

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Agape WIC	437	0.7	98.9	0.0	0.0	0.5	0.0
Albion Cli	216	67.6	29.2	3.2	0.0	0.0	0.0
Alcona	53	*	*	*	*	*	*
Algonac Cl	122	96.7	0.8	0.8	1.6	0.0	0.0
Allegan	105	92.4	2.9	4.8	0.0	0.0	0.0
Alma WIC	50	*	*	*	*	*	*
Alpena Cou	184	100.0	0.0	0.0	0.0	0.0	0.0
Antrim - B	61	*	*	*	*	*	*
Arab/Chald	168	92.9	6.5	0.0	0.0	0.6	0.0
Arenac Cou	96	*	*	*	*	*	*
Associated	8	*	*	*	*	*	*
Baldwin	149	61.1	23.5	14.1	0.0	1.3	0.0
Bangor	419	58.2	8.4	33.4	0.0	0.0	0.0
Baraga Cl	45	*	*	*	*	*	*
Barry-Eato	316	98.1	0.0	1.6	0.3	0.0	0.0
Barry-Eato	407	97.3	1.0	1.2	0.0	0.5	0.0
Battle Cre	938	69.9	21.6	7.4	0.2	0.9	0.0
Baxter Com	17	*	*	*	*	*	*
Bay Cl	693	96.5	1.9	1.2	0.0	0.4	0.0
Bayside WI	6	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Beaver Isl	2	*	*	*	*	*	*
Belding Cl	76	*	*	*	*	*	*
Bell Memor	85	*	*	*	*	*	*
Bendle/Bur	37	*	*	*	*	*	*
Benton Har	690	40.0	57.2	1.4	0.1	1.2	0.0
Benzie Cli	106	92.5	0.0	6.6	0.9	0.0	0.0
Berrien -	269	82.5	13.4	3.3	0.0	0.7	0.0
Bldhd Migr	7	*	*	*	*	*	*
Booth Clin	1	*	*	*	*	*	*
Branch-Hil	304	67.8	1.0	31.3	0.0	0.0	0.0
Branch-Hil	348	94.5	0.3	4.9	0.0	0.3	0.0
Branch-Hil	270	98.1	0.7	1.1	0.0	0.0	0.0
Browning-C	20	*	*	*	*	*	*
Bryant Com	20	*	*	*	*	*	*
Burton	1,287	83.2	12.2	3.3	0.5	0.7	0.0
Calumet WI	86	*	*	*	*	*	*
Capac Clin	42	*	*	*	*	*	*
Caro WIC S	3	*	*	*	*	*	*
Cedar Spri	243	94.7	0.8	4.1	0.4	0.0	0.0
Charlevoix	166	95.8	0.6	0.6	1.8	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Chass Heal	1,107	12.8	5.1	81.8	0.1	0.2	0.0
Cheboygan	181	98.3	0.0	0.6	0.0	1.1	0.0
Cherry Str	349	14.0	19.5	63.6	0.6	2.3	0.0
Chesaning	12	*	*	*	*	*	*
Clare Coun	213	99.5	0.5	0.0	0.0	0.0	0.0
Clinica Sa	327	0.6	0.0	99.4	0.0	0.0	0.0
Comstock C	0	*	*	*	*	*	*
Corner Hea	200	52.5	42.5	3.5	0.5	1.0	0.0
Crawford	110	99.1	0.9	0.0	0.0	0.0	0.0
Dewitt WIC	64	*	*	*	*	*	*
Dickinson-	194	100.0	0.0	0.0	0.0	0.0	0.0
Dickinson-	86	*	*	*	*	*	*
DMC Prim	14	*	*	*	*	*	*
DMC Prim	47	*	*	*	*	*	*
DMC Prim	15	*	*	*	*	*	*
Dorr Outre	99	*	*	*	*	*	*
Douglas He	686	12.4	53.5	26.4	0.4	7.3	0.0
Dowagiac	373	84.7	6.7	7.2	0.8	0.5	0.0
Durand WIC	94	*	*	*	*	*	*
East Area	322	2.2	96.9	0.6	0.0	0.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Eastside H	486	4.5	93.2	0.4	1.0	0.8	0.0
Eau Claire	195	1.5	0.5	97.9	0.0	0.0	0.0
Emmet Coun	248	92.3	0.0	2.0	4.4	1.2	0.0
Escanaba C	281	98.9	0.0	0.0	0.4	0.7	0.0
Family Com	356	38.2	48.6	5.6	0.0	7.6	0.0
Family Hlt	258	45.7	34.9	17.1	0.4	1.9	0.0
Family Roa	304	5.3	90.8	2.0	0.0	2.0	0.0
Farmington	104	71.2	25.0	1.0	0.0	2.9	0.0
Fife Lake	0	*	*	*	*	*	*
First Pres	0	*	*	*	*	*	*
Flint Fam	446	39.0	56.1	4.3	0.2	0.4	0.0
Fruitport	0	*	*	*	*	*	*
Gateway	417	0.7	98.6	0.0	0.2	0.5	0.0
Genesee -	12	*	*	*	*	*	*
Gladwin Co	155	100.0	0.0	0.0	0.0	0.0	0.0
Gogebic Cl	95	*	*	*	*	*	*
Golden	32	*	*	*	*	*	*
Grace Prof	120	0.8	98.3	0.8	0.0	0.0	0.0
Grace Ross	870	3.0	96.7	0.1	0.2	0.0	0.0
Grand Have	254	92.1	1.2	5.5	0.8	0.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Grand Trav	0	*	*	*	*	*	*
Grant	116	72.4	0.9	26.7	0.0	0.0	0.0
Gratiot	572	3.7	90.7	0.5	0.0	5.1	0.0
Greenville	127	99.2	0.0	0.8	0.0	0.0	0.0
Hamtramck	8	*	*	*	*	*	*
Hancock Cl	175	92.6	2.9	0.0	0.0	4.6	0.0
Hannahvil	13	*	*	*	*	*	*
Harper Gra	1,105	1.9	97.6	0.3	0.2	0.1	0.0
Haslett Cl	51	*	*	*	*	*	*
Haslett St	7	*	*	*	*	*	*
Herman Kie	1,288	2.9	95.7	0.2	0.1	1.2	0.0
Highland P	4	*	*	*	*	*	*
Holland Cl	825	49.2	3.5	40.6	0.0	6.7	0.0
Holly WIC	60	*	*	*	*	*	*
Holton	0	*	*	*	*	*	*
Howard Cit	54	*	*	*	*	*	*
Hudsonv	192	94.3	1.6	3.1	0.0	1.0	0.0
Hurley Cli	2	*	*	*	*	*	*
Huron Coun	226	98.2	0.4	1.3	0.0	0.0	0.0
Icchn-Spar	95	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Ingham Cou	1,517	61.4	23.1	11.5	0.4	3.6	0.0
Intervale	119	0.0	99.2	0.8	0.0	0.0	0.0
Ionia Coun	285	89.1	1.1	9.5	0.0	0.4	0.0
Iosco	150	98.0	1.3	0.0	0.7	0.0	0.0
Ironwood	0	*	*	*	*	*	*
Isabella C	283	95.1	2.1	1.1	0.7	1.1	0.0
Ithaca Cli	138	94.9	2.9	2.2	0.0	0.0	0.0
Ithaca WIC	0	*	*	*	*	*	*
Jackson Co	1,241	80.4	13.5	5.2	0.6	0.3	0.0
Jackson M	0	*	*	*	*	*	*
Janes St A	389	14.4	68.9	16.2	0.3	0.3	0.0
Joy-Greenf	733	33.8	63.0	3.0	0.0	0.1	0.0
Kalamazoo	384	30.2	53.4	14.1	0.5	1.8	0.0
Kalamazoo	776	80.3	17.1	1.8	0.0	0.8	0.0
Kalkaska	148	100.0	0.0	0.0	0.0	0.0	0.0
Kent	0	*	*	*	*	*	*
Kent Count	1,318	59.6	25.3	11.2	0.8	3.2	0.0
Kentwood	635	51.5	27.7	9.1	0.8	10.9	0.0
Keweenaw	25	*	*	*	*	*	*
Kingsley	8	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Lac Vieux	11	*	*	*	*	*	*
Lake Co	98	*	*	*	*	*	*
Lake Odess	45	*	*	*	*	*	*
Lapeer	489	98.0	0.6	1.2	0.0	0.2	0.0
Leelanu Cl	55	*	*	*	*	*	*
Lenawee	0	*	*	*	*	*	*
Leslie-Sto	18	*	*	*	*	*	*
Lincoln Pa	429	72.0	20.5	7.2	0.0	0.2	0.0
Livingston	358	96.9	0.6	2.2	0.0	0.3	0.0
Lowell	25	*	*	*	*	*	*
M-72 Migra	15	*	*	*	*	*	*
Mancelona	66	*	*	*	*	*	*
Manchester	8	*	*	*	*	*	*
Manchester	4	*	*	*	*	*	*
Manistee C	156	88.5	0.0	5.8	4.5	1.3	0.0
Manistique	63	*	*	*	*	*	*
Mansitee M	12	*	*	*	*	*	*
Marion Bra	39	*	*	*	*	*	*
Marquette	138	96.4	0.0	1.4	2.2	0.0	0.0
Marshall C	78	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Mason Clin	51	*	*	*	*	*	*
Mason Co	203	88.2	0.5	9.4	1.0	1.0	0.0
Mccree Hea	1,587	37.9	60.1	1.9	0.0	0.1	0.0
Mcnamara H	63	*	*	*	*	*	*
Mecosta Co	304	94.7	3.3	1.3	0.0	0.7	0.0
Medical Pr	7	*	*	*	*	*	*
Medical Pr	8	*	*	*	*	*	*
Menominee	136	99.3	0.7	0.0	0.0	0.0	0.0
Mercy	35	*	*	*	*	*	*
Mesick	1	*	*	*	*	*	*
Metropolit	0	*	*	*	*	*	*
Michigan S	80	*	*	*	*	*	*
Midland Cl	422	97.2	1.4	0.2	0.2	0.9	0.0
Midwest	425	17.6	36.0	43.1	0.7	2.6	0.0
Migrant H	104	1.0	0.0	99.0	0.0	0.0	0.0
Missaukee	66	*	*	*	*	*	*
Moms & Bab	449	6.0	61.2	32.1	0.0	0.7	0.0
Monroe Cou	758	92.3	4.5	2.9	0.1	0.1	0.0
Montague/	0	*	*	*	*	*	*
Montcalm	8	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Montmorenc	59	*	*	*	*	*	*
Mt Clemens	859	78.1	14.8	4.5	0.0	2.6	0.0
Munising C	37	*	*	*	*	*	*
Muskegon C	1,599	65.9	25.5	7.7	0.7	0.3	0.0
New Center	6	*	*	*	*	*	*
New Haven	74	*	*	*	*	*	*
New Hope	1	*	*	*	*	*	*
Newaygo	271	95.6	0.4	4.1	0.0	0.0	0.0
Newberry C	61	*	*	*	*	*	*
North Camp	128	35.9	5.5	7.8	0.0	50.8	0.0
North Midl	0	*	*	*	*	*	*
Northeast	926	2.9	87.5	0.4	0.0	9.2	0.0
Northern M	44	*	*	*	*	*	*
Northwest	215	1.4	98.6	0.0	0.0	0.0	0.0
Oakland Co	1,564	49.3	27.0	20.6	0.1	2.9	0.0
Oceana Co	188	84.6	0.5	14.4	0.5	0.0	0.0
Ogemaw	159	98.7	0.6	0.6	0.0	0.0	0.0
Olhsa Star	257	81.3	13.6	1.6	0.4	3.1	0.0
Olhsa WIC	65	*	*	*	*	*	*
Omni Care	26	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Omicare	0	*	*	*	*	*	*
Omicare-G	0	*	*	*	*	*	*
Omicare-M	27	*	*	*	*	*	*
Ontonagon	37	*	*	*	*	*	*
Oscoda	61	*	*	*	*	*	*
Otsego Co	204	99.5	0.0	0.0	0.0	0.5	0.0
Park Clin	0	*	*	*	*	*	*
Park Medic	14	*	*	*	*	*	*
Park Medic	20	*	*	*	*	*	*
Park Medic	0	*	*	*	*	*	*
Paw Paw Cl	132	90.9	2.3	6.1	0.0	0.8	0.0
PCOC	64	*	*	*	*	*	*
Pinconning	42	*	*	*	*	*	*
Port Huron	811	85.2	7.4	5.2	1.7	0.5	0.0
Portland	34	*	*	*	*	*	*
Presque Is	66	*	*	*	*	*	*
Profession	21	*	*	*	*	*	*
Pullman	253	59.3	4.0	36.4	0.0	0.4	0.0
Ravenna	0	*	*	*	*	*	*
Reed City	167	99.4	0.6	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Region II	595	76.1	1.3	22.2	0.0	0.3	0.0
Riverview	33	*	*	*	*	*	*
Romulus Fi	1	*	*	*	*	*	*
Roosevelt-	63	*	*	*	*	*	*
Roscommon	145	97.2	0.0	0.0	0.0	2.8	0.0
S & S Medi	7	*	*	*	*	*	*
Saginaw Co	1,093	54.8	34.9	10.2	0.0	0.0	0.0
Saginaw-Ch	14	*	*	*	*	*	*
Sanilac Co	296	99.0	0.7	0.3	0.0	0.0	0.0
Sault Ste	223	69.5	1.3	2.2	26.0	0.9	0.0
Sawyer Afb	74	*	*	*	*	*	*
School Bas	0	*	*	*	*	*	*
Schoolcraf	491	0.8	97.8	0.6	0.6	0.2	0.0
Seton Cent	28	*	*	*	*	*	*
Shaftsborg	40	*	*	*	*	*	*
Shelby M	116	1.7	0.0	98.3	0.0	0.0	0.0
Sheldon Co	595	10.1	52.4	37.1	0.0	0.3	0.0
Shiawassee	370	98.1	0.3	0.8	0.3	0.5	0.0
Southfield	931	51.3	45.4	1.3	0.0	1.9	0.0
Sparta Pub	61	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
St Charles	67	*	*	*	*	*	*
St Ignace	53	*	*	*	*	*	*
St Johns W	126	96.8	0.8	2.4	0.0	0.0	0.0
St Louis C	65	*	*	*	*	*	*
St.Clair S	343	86.9	9.6	1.5	0.0	2.0	0.0
Stanton Cl	252	98.4	0.8	0.8	0.0	0.0	0.0
Stockbridg	15	*	*	*	*	*	*
Sumpter He	379	74.9	24.3	0.8	0.0	0.0	0.0
Taylor	1,000	82.1	17.4	0.5	0.0	0.0	0.0
Teen Healt	1	*	*	*	*	*	*
Temperance	5	*	*	*	*	*	*
Thompsonv	10	*	*	*	*	*	*
Three Oaks	45	*	*	*	*	*	*
Three Rive	309	91.9	6.1	1.6	0.0	0.3	0.0
Traverse C	510	92.0	0.8	3.3	1.2	2.7	0.0
Tri-Cnty H	173	99.4	0.0	0.0	0.6	0.0	0.0
Tuscola	440	99.5	0.2	0.0	0.0	0.2	0.0
U Of M-Ped	0	*	*	*	*	*	*
Vcms River	35	*	*	*	*	*	*
Vicksburg	33	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Walled Lak	409	83.9	6.8	7.8	0.0	1.5	0.0
Warren Cli	536	82.8	10.8	0.9	0.2	5.2	0.0
Washtenaw	28	*	*	*	*	*	*
Wayne-Dear	866	98.3	0.8	0.6	0.0	0.3	0.0
Webbervill	16	*	*	*	*	*	*
West Genes	198	29.8	47.0	16.2	0.0	7.1	0.0
Western Wa	583	80.4	18.0	1.0	0.0	0.5	0.0
Westland C	719	67.5	28.8	3.1	0.1	0.6	0.0
Westside C	530	40.9	4.3	53.8	0.2	0.8	0.0
Westwood	28	*	*	*	*	*	*
Wexford	317	98.1	0.6	0.3	0.3	0.6	0.0
Wheeler WI	12	*	*	*	*	*	*
Whitmore L	16	*	*	*	*	*	*
Willow	5	*	*	*	*	*	*
Woodside	65	*	*	*	*	*	*
Wyoming Cl	656	73.5	8.8	14.2	0.6	2.9	0.0
Yale Clini	43	*	*	*	*	*	*
Ypsilanti	923	45.9	42.3	9.2	0.0	2.6	0.0
Michigan	61,040	59.8	28.1	10.1	0.5	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Nation (Prior Year)	727,815	49.8	24.9	20.1	1.2	2.8	1.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Agape WIC	437	0.7	6.2	10.5	53.1	27.7	1.6	0.2
Albion Cli	216	0.0	12.0	17.6	57.4	12.5	0.5	0.0
Alcona	53	*	*	*	*	*	*	*
Algonac Cl	122	0.0	5.7	15.6	59.0	17.2	2.5	0.0
Allegan	105	0.0	7.6	15.2	59.0	16.2	1.9	0.0
Alma WIC	50	*	*	*	*	*	*	*
Alpena Cou	184	0.0	4.9	13.0	69.0	12.0	1.1	0.0
Antrim - B	61	*	*	*	*	*	*	*
Arab/Chald	168	0.0	0.0	5.4	54.2	36.9	3.6	0.0
Arenac Cou	96	*	*	*	*	*	*	*
Associated	8	*	*	*	*	*	*	*
Baldwin	149	0.0	4.7	16.1	61.1	17.4	0.7	0.0
Bangor	419	0.2	8.1	16.0	57.3	17.4	1.0	0.0
Baraga Cl	45	*	*	*	*	*	*	*
Barry-Eato	316	0.0	8.2	15.8	55.7	19.0	1.3	0.0
Barry-Eato	407	0.0	5.9	17.4	61.9	13.5	1.2	0.0
Battle Cre	938	0.6	9.4	14.2	62.0	13.1	0.6	0.0
Baxter Com	17	*	*	*	*	*	*	*
Bay Cl	693	0.0	7.8	13.9	62.5	14.9	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Bayside WI	6	*	*	*	*	*	*	*
Beaver Isl	2	*	*	*	*	*	*	*
Belding Cl	76	*	*	*	*	*	*	*
Bell Memor	85	*	*	*	*	*	*	*
Bendle/Bur	37	*	*	*	*	*	*	*
Benton Har	690	0.9	10.6	15.7	56.8	14.8	1.3	0.0
Benzie Cli	106	0.0	4.7	17.9	57.5	17.9	1.9	0.0
Berrien -	269	0.0	9.3	14.5	62.8	13.4	0.0	0.0
Bldhd Migr	7	*	*	*	*	*	*	*
Booth Clin	1	*	*	*	*	*	*	*
Branch-Hil	304	0.3	7.9	11.8	61.5	17.4	1.0	0.0
Branch-Hil	348	0.6	7.5	16.1	62.6	12.6	0.6	0.0
Branch-Hil	270	0.4	6.3	21.1	60.0	11.9	0.4	0.0
Browning-C	20	*	*	*	*	*	*	*
Bryant Com	20	*	*	*	*	*	*	*
Burton	1,287	0.5	5.4	10.8	64.5	18.1	0.6	0.0
Calumet WI	86	*	*	*	*	*	*	*
Capac Clin	42	*	*	*	*	*	*	*
Caro WIC S	3	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cedar Spri	243	0.0	9.1	18.5	57.2	14.0	1.2	0.0
Charlevoix	166	0.0	8.4	13.9	60.2	16.9	0.6	0.0
Chass Heal	1,107	0.3	5.8	10.4	61.2	20.9	1.5	0.0
Cheboygan	181	0.0	5.5	15.5	58.6	19.9	0.6	0.0
Cherry Str	349	0.9	6.3	10.6	63.3	17.2	1.7	0.0
Chesaning	12	*	*	*	*	*	*	*
Clare Coun	213	1.4	9.9	16.4	58.7	11.7	1.9	0.0
Clinica Sa	327	0.3	6.1	8.3	63.6	19.9	1.8	0.0
Comstock C	0	*	*	*	*	*	*	*
Corner Hea	200	0.0	21.0	40.0	39.0	0.0	0.0	0.0
Crawford	110	0.9	9.1	20.0	57.3	10.9	1.8	0.0
Dewitt WIC	64	*	*	*	*	*	*	*
Dickinson-	194	0.0	8.2	12.4	62.9	15.5	1.0	0.0
Dickinson-	86	*	*	*	*	*	*	*
DMC Prim	14	*	*	*	*	*	*	*
DMC Prim	47	*	*	*	*	*	*	*
DMC Prim	15	*	*	*	*	*	*	*
Dorr Outre	99	*	*	*	*	*	*	*
Douglas He	686	0.4	5.8	10.6	61.4	20.6	1.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dowagiac	373	0.0	7.8	14.5	64.6	11.8	1.3	0.0
Durand WIC	94	*	*	*	*	*	*	*
East Area	322	1.2	11.5	13.4	52.8	19.3	1.9	0.0
Eastside H	486	0.4	6.6	14.6	55.8	21.4	1.2	0.0
Eau Claire	195	0.0	6.2	10.8	62.1	20.0	1.0	0.0
Emmet Coun	248	0.0	7.3	14.1	57.3	19.8	1.6	0.0
Eschanaba C	281	0.0	5.0	12.1	65.8	16.4	0.7	0.0
Family Com	356	0.0	3.9	7.9	60.4	25.8	2.0	0.0
Family Hlt	258	1.6	8.9	16.3	55.4	17.4	0.4	0.0
Family Roa	304	1.3	6.3	11.2	57.2	22.4	1.6	0.0
Farmington	104	0.0	5.8	3.8	66.3	23.1	1.0	0.0
Fife Lake	0	*	*	*	*	*	*	*
First Pres	0	*	*	*	*	*	*	*
Flint Fam	446	2.7	10.5	14.8	56.7	13.0	2.2	0.0
Fruitport	0	*	*	*	*	*	*	*
Gateway	417	1.7	13.7	14.6	51.3	17.0	1.7	0.0
Genesee -	12	*	*	*	*	*	*	*
Gladwin Co	155	0.0	9.7	12.9	57.4	19.4	0.6	0.0
Gogebic Cl	95	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Golden	32	*	*	*	*	*	*	*
Grace Prof	120	1.7	13.3	14.2	48.3	22.5	0.0	0.0
Grace Ross	870	0.5	4.7	14.9	56.2	22.1	1.6	0.0
Grand Have	254	0.0	5.1	11.0	63.8	19.3	0.8	0.0
Grand Trav	0	*	*	*	*	*	*	*
Grant	116	0.0	8.6	21.6	52.6	17.2	0.0	0.0
Gratiot	572	1.2	10.3	15.4	50.2	21.5	1.4	0.0
Greenville	127	0.0	9.4	13.4	66.9	9.4	0.8	0.0
Hamtramck	8	*	*	*	*	*	*	*
Hancock Cl	175	1.1	1.7	7.4	66.3	21.1	2.3	0.0
Hannahvil	13	*	*	*	*	*	*	*
Harper Gra	1,105	1.0	7.1	14.8	55.7	19.9	1.4	0.0
Haslett Cl	51	*	*	*	*	*	*	*
Haslett St	7	*	*	*	*	*	*	*
Herman Kie	1,288	0.3	5.9	11.9	55.8	24.2	1.9	0.0
Highland P	4	*	*	*	*	*	*	*
Holland Cl	825	0.2	8.7	14.5	61.7	14.2	0.6	0.0
Holly WIC	60	*	*	*	*	*	*	*
Holton	0	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Howard Cit	54	*	*	*	*	*	*	*
Hudsonv	192	1.0	5.7	11.5	57.3	22.9	1.6	0.0
Hurley Cli	2	*	*	*	*	*	*	*
Huron Coun	226	0.0	6.2	9.7	62.4	20.8	0.9	0.0
Icchn-Spar	95	*	*	*	*	*	*	*
Ingham Cou	1,517	0.2	7.2	12.5	64.5	14.5	1.1	0.0
Intervale	119	0.8	10.1	11.8	49.6	25.2	2.5	0.0
Ionia Coun	285	0.0	9.8	15.1	62.5	11.6	1.1	0.0
Iosco	150	0.0	4.7	16.7	65.3	12.7	0.7	0.0
Ironwood	0	*	*	*	*	*	*	*
Isabella C	283	0.0	4.9	15.5	66.4	13.1	0.0	0.0
Ithaca Cli	138	0.0	2.2	16.7	66.7	14.5	0.0	0.0
Ithaca WIC	0	*	*	*	*	*	*	*
Jackson Co	1,241	0.6	10.5	15.2	61.2	11.4	1.0	0.0
Jackson M	0	*	*	*	*	*	*	*
Janes St A	389	1.0	13.9	11.1	57.6	15.4	1.0	0.0
Joy-Greenf	733	0.4	4.5	10.9	59.8	23.2	1.2	0.0
Kalamazoo	384	0.3	10.4	14.1	60.4	13.5	1.3	0.0
Kalamazoo	776	0.5	5.7	13.5	64.9	14.3	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Kalkaska	148	0.7	4.7	12.2	69.6	10.1	2.7	0.0
Kent	0	*	*	*	*	*	*	*
Kent Count	1,318	0.5	5.9	13.7	61.1	17.5	1.3	0.0
Kentwood	635	0.5	4.6	11.2	61.6	20.6	1.6	0.0
Keweenaw	25	*	*	*	*	*	*	*
Kingsley	8	*	*	*	*	*	*	*
Lac Vieux	11	*	*	*	*	*	*	*
Lake Co	98	*	*	*	*	*	*	*
Lake Odess	45	*	*	*	*	*	*	*
Lapeer	489	0.2	5.9	16.4	58.1	18.6	0.8	0.0
Leelanu Cl	55	*	*	*	*	*	*	*
Lenawee	0	*	*	*	*	*	*	*
Leslie-Sto	18	*	*	*	*	*	*	*
Lincoln Pa	429	0.5	7.0	16.1	60.8	14.2	1.4	0.0
Livingston	358	0.0	7.3	13.7	59.2	19.3	0.6	0.0
Lowell	25	*	*	*	*	*	*	*
M-72 Migra	15	*	*	*	*	*	*	*
Mancelona	66	*	*	*	*	*	*	*
Manchester	8	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Manchester	4	*	*	*	*	*	*	*
Manistee C	156	1.3	7.1	10.9	64.7	14.7	1.3	0.0
Manistique	63	*	*	*	*	*	*	*
Mansitee M	12	*	*	*	*	*	*	*
Marion Bra	39	*	*	*	*	*	*	*
Marquette	138	0.0	5.8	6.5	72.5	12.3	2.9	0.0
Marshall C	78	*	*	*	*	*	*	*
Mason Clin	51	*	*	*	*	*	*	*
Mason Co	203	0.5	9.9	8.9	62.6	16.7	1.5	0.0
Mccree Hea	1,587	0.3	7.9	14.1	61.9	15.1	0.6	0.0
Mcnamara H	63	*	*	*	*	*	*	*
Mecosta Co	304	0.3	5.3	12.8	63.5	18.1	0.0	0.0
Medical Pr	7	*	*	*	*	*	*	*
Medical Pr	8	*	*	*	*	*	*	*
Menominee	136	0.0	8.1	18.4	58.1	14.0	1.5	0.0
Mercy	35	*	*	*	*	*	*	*
Mesick	1	*	*	*	*	*	*	*
Metropolit	0	*	*	*	*	*	*	*
Michigan S	80	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Midland Cl	422	0.5	5.9	14.7	63.3	14.2	1.4	0.0
Midwest	425	0.7	8.9	10.1	55.1	23.3	1.9	0.0
Migrant H	104	0.0	3.8	11.5	60.6	20.2	3.8	0.0
Missaukee	66	*	*	*	*	*	*	*
Moms & Bab	449	3.1	27.6	10.0	43.7	14.3	1.3	0.0
Monroe Cou	758	0.1	7.4	16.1	60.2	15.6	0.7	0.0
Montague/	0	*	*	*	*	*	*	*
Montcalm	8	*	*	*	*	*	*	*
Montmorenc	59	*	*	*	*	*	*	*
Mt Clemens	859	0.1	3.3	12.9	63.0	19.2	1.5	0.0
Munising C	37	*	*	*	*	*	*	*
Muskegon C	1,599	0.4	7.6	14.1	62.9	14.4	0.6	0.0
New Center	6	*	*	*	*	*	*	*
New Haven	74	*	*	*	*	*	*	*
New Hope	1	*	*	*	*	*	*	*
Newaygo	271	0.7	10.0	16.2	59.4	10.7	3.0	0.0
Newberry C	61	*	*	*	*	*	*	*
North Camp	128	0.0	0.0	1.6	60.2	38.3	0.0	0.0
North Midl	0	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Northeast	926	0.2	7.7	12.3	55.6	22.7	1.5	0.0
Northern M	44	*	*	*	*	*	*	*
Northwest	215	0.0	12.6	8.8	56.7	20.0	1.9	0.0
Oakland Co	1,564	0.4	7.2	11.3	62.2	18.2	0.7	0.0
Oceana Co	188	0.0	8.5	13.8	60.6	16.0	1.1	0.0
Ogemaw	159	0.0	6.3	11.3	64.2	17.6	0.6	0.0
Olhse Star	257	0.0	3.5	8.9	61.1	24.1	2.3	0.0
Olhse WIC	65	*	*	*	*	*	*	*
Omni Care	26	*	*	*	*	*	*	*
Omnicare	0	*	*	*	*	*	*	*
Omnicare-G	0	*	*	*	*	*	*	*
Omnicare-M	27	*	*	*	*	*	*	*
Ontonagon	37	*	*	*	*	*	*	*
Oscoda	61	*	*	*	*	*	*	*
Otsego Co	204	0.5	3.9	15.2	61.8	17.2	1.5	0.0
Park Clin	0	*	*	*	*	*	*	*
Park Medic	14	*	*	*	*	*	*	*
Park Medic	20	*	*	*	*	*	*	*
Park Medic	0	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Paw Paw Cl	132	0.0	9.1	14.4	63.6	12.1	0.8	0.0
PCOC	64	*	*	*	*	*	*	*
Pinconning	42	*	*	*	*	*	*	*
Port Huron	811	0.5	7.5	17.1	60.3	14.1	0.5	0.0
Portland	34	*	*	*	*	*	*	*
Presque Is	66	*	*	*	*	*	*	*
Profession	21	*	*	*	*	*	*	*
Pullman	253	1.6	10.3	15.4	56.9	13.8	2.0	0.0
Ravenna	0	*	*	*	*	*	*	*
Reed City	167	0.0	7.2	19.2	56.9	16.2	0.6	0.0
Region II	595	0.2	7.2	13.8	62.7	14.8	1.3	0.0
Riverview	33	*	*	*	*	*	*	*
Romulus Fi	1	*	*	*	*	*	*	*
Roosevelt-	63	*	*	*	*	*	*	*
Roscommon	145	0.0	9.7	17.9	57.2	13.8	1.4	0.0
S & S Medi	7	*	*	*	*	*	*	*
Saginaw Co	1,093	0.5	8.4	12.4	59.6	17.7	1.6	0.0
Saginaw-Ch	14	*	*	*	*	*	*	*
Sanilac Co	296	0.0	6.8	11.5	64.9	14.9	2.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sault Ste	223	0.0	5.4	14.8	61.0	17.9	0.9	0.0
Sawyer Afb	74	*	*	*	*	*	*	*
School Bas	0	*	*	*	*	*	*	*
Schoolcraf	491	0.2	5.9	11.4	59.3	21.4	1.8	0.0
Seton Cent	28	*	*	*	*	*	*	*
Shaftsburg	40	*	*	*	*	*	*	*
Shelby M	116	0.0	7.8	7.8	65.5	14.7	4.3	0.0
Sheldon Co	595	1.5	10.4	15.0	59.0	13.8	0.3	0.0
Shiawassee	370	0.0	7.8	15.1	62.2	14.6	0.3	0.0
Southfield	931	0.0	4.2	10.3	57.4	26.4	1.7	0.0
Sparta Pub	61	*	*	*	*	*	*	*
St Charles	67	*	*	*	*	*	*	*
St Ignace	53	*	*	*	*	*	*	*
St Johns W	126	0.0	6.3	13.5	64.3	13.5	2.4	0.0
St Louis C	65	*	*	*	*	*	*	*
St.Clair S	343	0.0	5.2	12.0	60.9	21.0	0.9	0.0
Stanton Cl	252	0.0	9.5	17.5	60.7	9.9	2.4	0.0
Stockbridg	15	*	*	*	*	*	*	*
Sumpter He	379	0.5	4.7	13.5	61.2	19.0	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Taylor	1,000	0.4	5.7	14.5	61.9	16.2	1.3	0.0
Teen Healt	1	*	*	*	*	*	*	*
Temperance	5	*	*	*	*	*	*	*
Thompsonv	10	*	*	*	*	*	*	*
Three Oaks	45	*	*	*	*	*	*	*
Three Rive	309	0.6	9.7	21.0	59.2	8.4	1.0	0.0
Traverse C	510	0.2	5.9	15.7	58.4	17.8	2.0	0.0
Tri-Cnty H	173	0.0	3.5	15.0	65.9	15.0	0.6	0.0
Tuscola	440	0.0	6.4	14.5	63.0	15.5	0.7	0.0
U Of M-Ped	0	*	*	*	*	*	*	*
Vcms River	35	*	*	*	*	*	*	*
Vicksburg	33	*	*	*	*	*	*	*
Walled Lak	409	0.2	4.6	14.9	59.4	19.3	1.5	0.0
Warren Cli	536	0.0	4.9	12.7	60.3	20.3	1.9	0.0
Washtenaw	28	*	*	*	*	*	*	*
Wayne-Dear	866	0.2	2.3	9.2	61.2	24.6	2.4	0.0
Webbervill	16	*	*	*	*	*	*	*
West Genes	198	0.5	11.1	7.6	64.1	15.7	1.0	0.0
Western Wa	583	0.5	4.5	10.6	62.8	19.4	2.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Westland C	719	0.3	6.1	11.5	60.8	21.1	0.1	0.0
Westside C	530	0.6	9.4	14.5	59.1	15.5	0.9	0.0
Westwood	28	*	*	*	*	*	*	*
Wexford	317	0.6	11.4	14.8	58.7	13.9	0.6	0.0
Wheeler WI	12	*	*	*	*	*	*	*
Whitmore L	16	*	*	*	*	*	*	*
Willow	5	*	*	*	*	*	*	*
Woodside	65	*	*	*	*	*	*	*
Wyoming Cl	656	0.5	6.3	11.6	62.5	18.1	0.9	0.2
Yale Clini	43	*	*	*	*	*	*	*
Ypsilanti	923	0.2	3.9	8.7	66.4	19.8	1.0	0.0
Michigan	61,040	0.4	7.2	13.4	60.3	17.5	1.2	0.0
Nation (Prior Year)	727,815	0.5	7.3	14.1	59.6	17.3	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Agape WIC	418	11.0 (43)	50.7 (111)	383	33.2 (88)	45.4 (55)	101	37.6 (22)	79	* *	
Albion Cli	202	10.4 (33)	47.5 (87)	173	26.0 (56)	42.8 (40)	33	* *	36	* *	
Alcona	51	* *	* *	43	* *	* *	2	* *	13	* *	
Algonac Cl	118	21.2 (122)	34.7 (8)	98	* *	* *	22	* *	71	* *	
Allegan	90	* *	* *	78	* *	* *	13	* *	18	* *	
Alma WIC	46	* *	* *	39	* *	* *	9	* *	11	* *	
Alpena Cou	172	11.6 (58)	44.8 (63)	156	22.4 (30)	50.6 (94)	20	* *	58	* *	
Antrim - B	54	* *	* *	47	* *	* *	3	* *	19	* *	
Arab/Chald	156	6.4 (4)	39.1 (24)	127	66.9 (113)	15.0 (2)	17	* *	13	* *	
Arenac Cou	86	* *	* *	37	* *	* *	12	* *	10	* *	
Associated	8	* *	* *	7	* *	* *	2	* *	3	* *	
Baldwin	134	9.7 (19)	38.1 (18)	98	* *	* *	32	* *	40	* *	
Bangor	376	14.1 (96)	42.0 (39)	326	34.0 (90)	38.0 (23)	59	* *	80	* *	
Baraga Cl	44	* *	* *	40	* *	* *	5	* *	5	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Barry-Eato	296	12.2 (65)	45.9 (75)	239	32.6 (87)	39.7 (26)	39	* *	73	* *	
Barry-Eato	384	11.2 (50)	42.4 (41)	329	24.3 (48)	49.2 (87)	66	* *	84	* *	
Battle Cre	863	12.6 (73)	45.5 (70)	755	24.0 (42)	48.2 (76)	177	25.4 (8)	142	39.4 (26)	
Baxter Com	17	* *	* *	11	* *	* *	2	* *	2	* *	
Bay Cl	630	11.0 (43)	47.1 (84)	532	24.1 (45)	46.6 (66)	85	* *	134	24.6 (9)	
Bayside WI	5	* *	* *	2	* *	* *	0	* *	1	* *	
Beaver Isl	1	* *	* *	1	* *	* *	0	* *	0	* *	
Belding Cl	76	* *	* *	65	* *	* *	15	* *	13	* *	
Bell Memor	77	* *	* *	65	* *	* *	6	* *	15	* *	
Bendle/Bur	35	* *	* *	27	* *	* *	8	* *	3	* *	
Benton Har	650	12.2 (65)	48.3 (94)	545	27.3 (60)	45.1 (53)	132	44.7 (27)	251	43.0 (31)	
Benzie Cli	99	* *	* *	87	* *	* *	14	* *	24	* *	
Berrien -	254	14.2 (98)	42.5 (44)	212	23.6 (39)	47.6 (71)	45	* *	74	* *	
Bldhd Migr	5	* *	* *	2	* *	* *	1	* *	0	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Booth Clin	1	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Branch-Hil	266	12.0 (62)	44.4 (60)		227	27.3 (60)	42.3 (37)		50	* *	51	* *
Branch-Hil	306	16.7 (114)	35.9 (10)		246	30.5 (81)	39.4 (25)		59	* *	51	* *
Branch-Hil	245	12.7 (77)	38.4 (21)		217	27.6 (63)	45.6 (56)		48	* *	62	* *
Browning-C	20	* *	* *	* *	16	* *	* *	* *	3	* *	2	* *
Bryant Com	20	* *	* *	* *	17	* *	* *	* *	2	* *	6	* *
Burton	1,129	12.7 (77)	45.2 (69)		974	19.4 (9)	51.8 (101)		221	36.2 (20)	264	25.8 (11)
Calumet WI	77	* *	* *	* *	62	* *	* *	* *	4	* *	27	* *
Capac Clin	41	* *	* *	* *	34	* *	* *	* *	7	* *	16	* *
Caro WIC S	1	* *	* *	* *	1	* *	* *	* *	0	* *	1	* *
Cedar Spri	220	17.3 (116)	34.1 (6)		198	23.2 (37)	46.0 (60)		33	* *	60	* *
Charlevoix	146	11.6 (58)	43.8 (57)		130	36.2 (94)	41.5 (30)		29	* *	64	* *
Chass Heal	1,034	10.8 (38)	36.9 (13)		889	36.1 (93)	35.4 (15)		158	23.4 (5)	536	21.3 (6)
Cheboygan	174	12.6 (73)	47.7 (88)		150	25.3 (53)	44.7 (51)		29	* *	37	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)										
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cherry Str	340	8.5 (13)	44.7 (61)	288	29.2 (71)	41.3 (29)	30	* *	28	* *
Chesaning	11	* *	* *	10	* *	* *	2	* *	2	* *
Clare Coun	199	10.1 (30)	48.2 (93)	173	22.5 (32)	48.0 (74)	26	* *	57	* *
Clinica Sa	298	3.4 (1)	49.7 (101)	271	62.0 (111)	16.2 (3)	31	* *	104	68.3 (40)
Comstock C	0	* *	* *	0	* *	* *	0	* *	0	* *
Corner Hea	194	16.0 (113)	33.5 (5)	171	19.9 (13)	49.1 (86)	12	* *	23	* *
Crawford	105	10.5 (34)	45.7 (73)	92	* *	* *	10	* *	29	* *
Dewitt WIC	54	* *	* *	48	* *	* *	12	* *	19	* *
Dickinson-	175	7.4 (8)	45.1 (68)	149	14.1 (2)	59.1 (115)	23	* *	44	* *
Dickinson-	82	* *	* *	70	* *	* *	9	* *	20	* *
DMC Prim	13	* *	* *	11	* *	* *	2	* *	4	* *
DMC Prim	45	* *	* *	45	* *	* *	10	* *	9	* *
DMC Prim	15	* *	* *	14	* *	* *	2	* *	3	* *
Dorr Outre	90	* *	* *	79	* *	* *	20	* *	20	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Douglas He	539	10.0 (25)	50.5 (109)	475	31.6 (85)	39.8 (27)	83	* *	116	33.6 (19)	
Dowagiac	342	14.3 (104)	43.3 (50)	278	50.4 (107)	32.4 (11)	70	* *	74	* *	
Durand WIC	76	* *	* *	64	* *	* *	8	* *	23	* *	
East Area	296	9.1 (17)	55.1 (121)	234	42.3 (101)	37.2 (18)	29	* *	23	* *	
Eastside H	463	11.4 (53)	50.1 (104)	411	25.3 (53)	44.3 (47)	104	47.1 (29)	175	49.7 (36)	
Eau Claire	145	4.1 (3)	56.6 (122)	117	38.5 (97)	29.1 (8)	20	* *	43	* *	
Emmet Coun	237	11.4 (53)	46.8 (82)	209	27.8 (65)	41.6 (31)	31	* *	48	* *	
Esanaba C	256	9.8 (22)	40.6 (30)	219	24.2 (47)	42.0 (35)	41	* *	43	* *	
Family Com	341	11.1 (45)	41.6 (35)	294	54.1 (109)	26.2 (7)	59	* *	50	* *	
Family Hlt	233	10.7 (36)	45.5 (70)	185	30.8 (84)	44.3 (47)	34	* *	31	* *	
Family Roa	298	9.7 (19)	50.0 (102)	252	27.0 (58)	44.8 (52)	69	* *	2	* *	
Farmington	96	* *	* *	77	* *	* *	25	* *	28	* *	
Fife Lake	0	* *	* *	0	* *	* *	0	* *	0	* *	
First Pres	0	* *	* *	0	* *	* *	0	* *	0	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Flint Fam	426	12.4 (68)	43.9 (58)	366	24.3 (48)	47.5 (70)	103	54.4 (32)	64	* *	
Fruitport	0	* *	* *	0	* *	* *	0	* *	0	* *	
Gateway	388	11.1 (45)	53.6 (117)	308	37.7 (96)	42.2 (36)	50	* *	50	* *	
Genesee -	11	* *	* *	0	* *	* *	0	* *	0	* *	
Gladwin Co	148	14.2 (98)	49.3 (100)	133	24.8 (52)	53.4 (106)	16	* *	22	* *	
Gogebic Cl	83	* *	* *	69	* *	* *	14	* *	26	* *	
Golden	31	* *	* *	28	* *	* *	5	* *	8	* *	
Grace Prof	119	10.9 (41)	50.4 (108)	105	21.9 (26)	54.3 (110)	23	* *	15	* *	
Grace Ross	837	6.6 (5)	53.8 (118)	671	29.8 (76)	48.6 (82)	185	33.5 (15)	165	49.1 (35)	
Grand Have	241	14.1 (96)	43.2 (49)	208	21.6 (23)	47.6 (71)	38	* *	65	* *	
Grand Trav	0	* *	* *	0	* *	* *	0	* *	0	* *	
Grant	106	6.6 (5)	48.1 (91)	97	* *	* *	16	* *	16	* *	
Gratiot	531	13.0 (86)	44.8 (63)	482	30.7 (82)	46.1 (62)	88	* *	256	44.9 (33)	
Greenville	118	17.8 (119)	42.4 (41)	106	29.2 (71)	46.2 (63)	21	* *	35	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hamtramck	6	* *	* *	* *	0	* *	* *	* *	2	* *	0	* *
Hancock Cl	157	15.3 (108)	30.6 (2)		122	75.4 (115)	10.7 (1)		23	* *	50	* *
Hannahvil	10	* *	* *	* *	9	* *	* *	* *	0	* *	3	* *
Harper Gra	1,047	9.7 (19)	48.6 (97)		914	41.7 (100)	37.7 (20)		198	34.8 (17)	169	41.4 (30)
Haslett Cl	47	* *	* *	* *	43	* *	* *	* *	8	* *	19	* *
Haslett St	7	* *	* *	* *	6	* *	* *	* *	1	* *	0	* *
Herman Kie	1,212	11.1 (45)	50.2 (105)		1,056	44.3 (103)	35.8 (16)		262	31.3 (14)	215	33.5 (17)
Highland P	4	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Holland Cl	755	13.2 (90)	41.9 (37)		661	53.4 (108)	26.0 (6)		128	34.4 (16)	254	25.6 (10)
Holly WIC	58	* *	* *	* *	44	* *	* *	* *	16	* *	12	* *
Holton	0	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Howard Cit	51	* *	* *	* *	44	* *	* *	* *	9	* *	19	* *
Hudsonv	174	11.5 (57)	36.8 (12)		162	44.4 (104)	30.9 (10)		31	* *	71	* *
Hurley Cli	2	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Huron Coun	213	9.9 (24)	52.6 (116)	188	29.3 (73)	44.1 (45)	28	* *	33	* *	
Icchn-Spar	63	* *	* *	44	* *	* *	8	* *	14	* *	
Ingham Cou	1,375	12.6 (73)	43.1 (48)	1,227	45.5 (106)	34.1 (14)	272	26.8 (10)	496	47.8 (34)	
Intervale	110	10.0 (25)	57.3 (123)	101	23.8 (41)	53.5 (107)	10	* *	25	* *	
Ionia Coun	269	10.8 (38)	49.1 (99)	227	19.4 (9)	48.9 (84)	50	* *	72	* *	
Iosco	147	13.6 (93)	41.5 (33)	128	18.8 (8)	48.4 (79)	21	* *	21	* *	
Ironwood	0	* *	* *	0	* *	* *	0	* *	0	* *	
Isabella C	255	14.9 (105)	42.7 (45)	185	54.6 (110)	30.3 (9)	50	* *	51	* *	
Ithaca Cli	120	11.7 (60)	52.5 (115)	102	21.6 (23)	52.9 (104)	19	* *	26	* *	
Ithaca WIC	0	* *	* *	0	* *	* *	0	* *	0	* *	
Jackson Co	1,119	13.0 (86)	41.4 (32)	1,013	19.6 (11)	51.7 (99)	176	23.9 (6)	155	59.4 (39)	
Jackson M	0	* *	* *	0	* *	* *	0	* *	0	* *	
Janes St A	377	9.0 (16)	51.2 (113)	322	30.1 (78)	37.3 (19)	60	* *	183	32.8 (16)	
Joy-Greenf	689	8.7 (15)	47.9 (89)	587	24.0 (42)	47.2 (67)	114	55.3 (33)	80	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Kalamazoo	350	11.1 (45)	47.1 (84)	279	37.6 (95)	39.8 (27)	58	* *	48	* *	
Kalamazoo	732	12.6 (73)	42.3 (40)	628	33.8 (89)	42.5 (39)	134	37.3 (21)	142	43.7 (32)	
Kalkaska	140	10.0 (25)	48.6 (97)	130	13.1 (1)	54.6 (113)	17	* *	48	* *	
Kent	0	* *	* *	0	* *	* *	0	* *	0	* *	
Kent Count	1,228	11.3 (51)	43.6 (54)	1,048	22.1 (27)	48.1 (75)	166	13.9 (1)	253	16.6 (4)	
Kentwood	587	15.7 (112)	39.7 (26)	525	22.3 (28)	44.4 (49)	122	13.9 (1)	138	13.0 (2)	
Keweenaw	25	* *	* *	23	* *	* *	5	* *	2	* *	
Kingsley	8	* *	* *	8	* *	* *	0	* *	2	* *	
Lac Vieux	11	* *	* *	11	* *	* *	2	* *	2	* *	
Lake Co	94	* *	* *	76	* *	* *	20	* *	23	* *	
Lake Odess	42	* *	* *	34	* *	* *	5	* *	8	* *	
Lapeer	450	12.0 (62)	44.2 (59)	386	20.2 (17)	49.7 (91)	76	* *	142	50.7 (37)	
Leelanu Cl	52	* *	* *	47	* *	* *	3	* *	11	* *	
Lenawee	0	* *	* *	0	* *	* *	0	* *	0	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Leslie-Sto	16	* *	* *	* *	12	* *	* *	* *	1	* *	6	* *
Lincoln Pa	396	12.9 (83)	44.7 (61)		270	21.1 (19)	53.3 (105)		105	25.7 (9)	73	* *
Livingston	306	17.6 (118)	33.0 (4)		271	15.5 (3)	53.5 (107)		75	* *	133	12.0 (1)
Lowell	24	* *	* *	* *	18	* *	* *	* *	6	* *	3	* *
M-72 Migra	7	* *	* *	* *	4	* *	* *	* *	1	* *	1	* *
Mancelona	63	* *	* *	* *	55	* *	* *	* *	5	* *	18	* *
Manchester	8	* *	* *	* *	6	* *	* *	* *	1	* *	2	* *
Manchester	2	* *	* *	* *	1	* *	* *	* *	0	* *	1	* *
Manistee C	149	12.8 (80)	45.0 (66)		128	21.1 (19)	48.4 (79)		27	* *	28	* *
Manistique	57	* *	* *	* *	54	* *	* *	* *	2	* *	11	* *
Mansitee M	6	* *	* *	* *	4	* *	* *	* *	1	* *	0	* *
Marion Bra	37	* *	* *	* *	35	* *	* *	* *	10	* *	16	* *
Marquette	129	7.8 (10)	35.7 (9)		113	17.7 (6)	47.8 (73)		13	* *	22	* *
Marshall C	76	* *	* *	* *	64	* *	* *	* *	12	* *	16	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mason Clin	44	* *	* *	* *	41	* *	* *	* *	10	* *	13	* *
Mason Co	196	12.8 (80)	46.4 (80)		171	23.4 (38)	51.5 (97)		29	* *	54	* *
Mccree Hea	1,458	12.7 (77)	47.4 (86)		1,266	24.3 (48)	48.4 (79)		322	46.0 (28)	307	35.5 (20)
Mcnamara H	60	* *	* *	* *	50	* *	* *	* *	14	* *	19	* *
Mecosta Co	282	14.2 (98)	43.6 (54)		249	64.3 (112)	16.5 (4)		30	* *	69	* *
Medical Pr	7	* *	* *	* *	6	* *	* *	* *	1	* *	2	* *
Medical Pr	8	* *	* *	* *	8	* *	* *	* *	2	* *	1	* *
Menominee	129	12.4 (68)	40.3 (29)		108	21.3 (21)	50.9 (95)		28	* *	29	* *
Mercy	35	* *	* *	* *	31	* *	* *	* *	9	* *	8	* *
Mesick	1	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Metropolit	0	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Michigan S	76	* *	* *	* *	67	* *	* *	* *	19	* *	36	* *
Midland Cl	380	12.4 (68)	48.4 (95)		317	21.8 (25)	51.7 (99)		62	* *	92	* *
Midwest	399	10.0 (25)	48.1 (91)		324	70.1 (114)	19.1 (5)		40	* *	26	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight	< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Migrant H	100	4.0 (2)	54.0 (120)	83	* *	* *	12	* *	18	* *	
Missaukee	62	* *	* *	57	* *	* *	11	* *	14	* *	
Moms & Bab	405	10.9 (41)	39.5 (25)	344	29.9 (77)	41.6 (31)	79	* *	65	* *	
Monroe Cou	702	14.2 (98)	37.7 (14)	614	24.1 (45)	48.2 (76)	119	20.2 (3)	98	* *	
Montague/	0	* *	* *	0	* *	* *	0	* *	0	* *	
Montcalm	6	* *	* *	5	* *	* *	0	* *	4	* *	
Montmorenc	56	* *	* *	51	* *	* *	5	* *	17	* *	
Mt Clemens	809	13.8 (94)	36.6 (11)	276	29.0 (69)	42.4 (38)	221	27.1 (12)	169	27.8 (12)	
Munising C	36	* *	* *	29	* *	* *	3	* *	11	* *	
Muskegon C	1,525	10.5 (34)	46.0 (78)	1,365	23.7 (40)	49.3 (88)	231	38.5 (24)	121	38.8 (25)	
New Center	5	* *	* *	5	* *	* *	0	* *	2	* *	
New Haven	68	* *	* *	33	* *	* *	10	* *	15	* *	
New Hope	1	* *	* *	1	* *	* *	1	* *	0	* *	
Newaygo	248	15.3 (108)	43.5 (53)	207	28.5 (67)	48.8 (83)	27	* *	84	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Newberry C	56	* *	* *	* *	46	* *	* *	* *	3	* *	12	* *
North Camp	121	28.9 (123)	16.5 (1)		95	* *	* *	* *	31	* *	40	* *
North Midl	0	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Northeast	872	10.7 (36)	50.0 (102)		718	45.1 (105)	33.6 (13)		99	* *	200	33.5 (17)
Northern M	42	* *	* *	* *	37	* *	* *	* *	5	* *	7	* *
Northwest	208	12.5 (72)	46.6 (81)		171	32.2 (86)	45.6 (56)		22	* *	14	* *
Oakland Co	1,444	11.1 (45)	48.0 (90)		1,264	27.0 (58)	46.0 (60)		301	38.9 (25)	402	30.6 (15)
Oceana Co	174	8.0 (12)	51.7 (114)		148	22.3 (28)	49.3 (88)		21	* *	37	* *
Ogemaw	148	14.2 (98)	41.9 (37)		122	23.0 (35)	51.6 (98)		12	* *	36	* *
Olhsa Star	231	15.6 (111)	32.5 (3)		179	34.1 (91)	43.0 (41)		60	* *	50	* *
Olhsa WIC	59	* *	* *	* *	45	* *	* *	* *	15	* *	8	* *
Omni Care	26	* *	* *	* *	24	* *	* *	* *	4	* *	0	* *
Omnicare	0	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Omnicare-G	0	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight		< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)		Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Omnicare-M	25	* * *	* *		20	* * *	* *		7	* *	7	* *
Ontonagon	29	* * *	* *		24	* * *	* *		5	* *	6	* *
Oscoda	54	* * *	* *		45	* * *	* *		5	* *	13	* *
Otsego Co	191	17.3 (116)	37.7 (14)		165	16.4 (4)	50.3 (92)		23	* *	42	* *
Park Clin	0	* * *	* *		0	* * *	* *		0	* *	0	* *
Park Medic	13	* * *	* *		12	* * *	* *		3	* *	1	* *
Park Medic	20	* * *	* *		20	* * *	* *		4	* *	6	* *
Park Medic	0	* * *	* *		0	* * *	* *		0	* *	0	* *
Paw Paw Cl	125	15.2 (107)	44.8 (63)		114	28.9 (68)	45.6 (56)		24	* *	23	* *
PCOC	60	* * *	* *		48	* * *	* *		7	* *	23	* *
Pinconning	38	* * *	* *		28	* * *	* *		5	* *	6	* *
Port Huron	775	14.2 (98)	42.7 (45)		691	22.4 (30)	46.2 (63)		134	35.8 (19)	414	41.1 (29)
Portland	29	* * *	* *		27	* * *	* *		6	* *	10	* *
Presque Is	61	* * *	* *		56	* * *	* *		9	* *	24	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight		< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Profession	20	* *	* *	* *	17	* *	* *	* *	2	* *	6	* *
Pullman	220	7.7 (9)	50.5 (109)		175	39.4 (98)	37.7 (20)		35	* *	31	* *
Ravenna	0	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Reed City	158	18.4 (121)	46.2 (79)		136	29.4 (74)	41.9 (33)		27	* *	41	* *
Region II	484	13.2 (90)	45.5 (70)		441	24.3 (48)	45.1 (53)		96	* *	125	37.6 (21)
Riverview	31	* *	* *	* *	29	* *	* *	* *	10	* *	12	* *
Romulus Fi	1	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Roosevelt-	59	* *	* *	* *	51	* *	* *	* *	11	* *	31	* *
Roscommon	139	12.9 (83)	38.1 (18)		120	20.0 (14)	55.8 (114)		23	* *	40	* *
S & S Medi	7	* *	* *	* *	5	* *	* *	* *	1	* *	1	* *
Saginaw Co	1,041	6.7 (7)	50.9 (112)		884	20.1 (15)	50.3 (92)		170	28.8 (13)	196	40.8 (28)
Saginaw-Ch	12	* *	* *	* *	8	* *	* *	* *	4	* *	5	* *
Sanilac Co	280	12.9 (83)	45.0 (66)		250	24.0 (42)	47.2 (67)		47	* *	96	* *
Sault Ste	213	8.5 (13)	42.7 (45)		188	17.6 (5)	54.3 (110)		27	* *	32	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)
Sawyer Afb	74	* *	* *		61	* *	* *		7	* *	11	* *
School Bas	0	* *	* *		0	* *	* *		0	* *	0	* *
Schoolcraf	465	10.1 (30)	50.3 (106)		418	21.5 (22)	48.3 (78)		109	48.6 (31)	114	38.6 (24)
Seton Cent	27	* *	* *		12	* *	* *		1	* *	1	* *
Shaftsbu	36	* *	* *		30	* *	* *		9	* *	9	* *
Shelby M	85	* *	* *		63	* *	* *		5	* *	12	* *
Sheldon Co	536	7.8 (10)	45.9 (75)		463	30.7 (82)	43.2 (42)		94	* *	89	* *
Shiawassee	141	11.3 (51)	53.9 (119)		119	22.7 (33)	52.1 (102)		42	* *	74	* *
Southfield	859	14.0 (95)	41.8 (36)		777	29.6 (75)	45.8 (59)		212	35.4 (18)	283	28.6 (13)
Sparta Pub	54	* *	* *		45	* *	* *		11	* *	11	* *
St Charles	63	* *	* *		59	* *	* *		16	* *	15	* *
St Ignace	52	* *	* *		46	* *	* *		6	* *	5	* *
St Johns W	116	18.1 (120)	37.9 (16)		98	* *	* *		15	* *	30	* *
St Louis C	60	* *	* *		48	* *	* *		9	* *	13	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
St.Clair S	321	12.1 (64)	38.3 (20)	97	* *	* *	84	* *	54	* *	
Stanton Cl	235	12.3 (67)	40.0 (27)	206	27.7 (64)	44.2 (46)	38	* *	59	* *	
Stockbridg	11	* *	* *	10	* *	* *	3	* *	3	* *	
Sumpter He	342	11.4 (53)	41.5 (33)	285	30.2 (79)	44.6 (50)	64	* *	101	23.8 (8)	
Taylor	855	15.0 (106)	40.2 (28)	716	44.1 (102)	38.0 (23)	221	24.0 (7)	222	22.1 (7)	
Teen Healt	1	* *	* *	1	* *	* *	0	* *	1	* *	
Temperance	5	* *	* *	5	* *	* *	1	* *	1	* *	
Thompsonv	10	* *	* *	8	* *	* *	1	* *	2	* *	
Three Oaks	42	* *	* *	41	* *	* *	14	* *	17	* *	
Three Rive	297	16.8 (115)	38.0 (17)	259	40.2 (99)	36.3 (17)	56	* *	59	* *	
Traverse C	465	13.1 (89)	38.7 (23)	407	23.1 (36)	47.4 (69)	67	* *	134	59.0 (38)	
Tri-Cnty H	153	9.8 (22)	50.3 (106)	136	25.7 (55)	49.3 (88)	29	* *	29	* *	
Tuscola	377	13.5 (92)	48.5 (96)	325	22.8 (34)	46.2 (63)	52	* *	89	* *	
U Of M-Ped	0	* *	* *	0	* *	* *	0	* *	0	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Vcms River	35	* *	* *	* *	33	* *	* *	* *	8	* *	6	* *
Vicksburg	30	* *	* *	* *	25	* *	* *	* *	1	* *	7	* *
Walled Lak	358	15.4 (110)	38.5 (22)		313	19.8 (12)	54.3 (110)		85	* *	133	21.1 (5)
Warren Cli	509	12.8 (80)	43.4 (51)		159	20.1 (15)	54.1 (109)		134	26.9 (11)	84	* *
Washtenaw	24	* *	* *	* *	22	* *	* *	* *	4	* *	5	* *
Wayne-Dear	766	9.5 (18)	34.1 (6)		573	35.8 (92)	32.6 (12)		149	47.7 (30)	207	39.6 (27)
Webbervill	12	* *	* *	* *	9	* *	* *	* *	1	* *	5	* *
West Genes	184	13.0 (86)	45.7 (73)		160	26.3 (57)	41.9 (33)		45	* *	41	* *
Western Wa	545	10.3 (32)	42.4 (41)		443	27.5 (62)	44.0 (44)		119	43.7 (26)	107	38.3 (23)
Westland C	670	12.4 (68)	43.4 (51)		470	28.3 (66)	48.9 (84)		191	20.4 (4)	173	28.9 (14)
Westside C	489	10.0 (25)	47.0 (83)		444	30.2 (79)	37.8 (22)		68	* *	86	* *
Westwood	27	* *	* *	* *	23	* *	* *	* *	3	* *	6	* *
Wexford	307	11.7 (60)	43.6 (54)		269	20.4 (18)	50.9 (95)		35	* *	89	* *
Wheeler WI	11	* *	* *	* *	10	* *	* *	* *	2	* *	2	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Whitmore L	14	* *	* *	* *	11	* *	* *	* *	4	* *	4	* *
Willow	4	* *	* *	* *	3	* *	* *	* *	1	* *	2	* *
Woodside	64	* *	* *	* *	55	* *	* *	* *	8	* *	12	* *
Wyoming Cl	621	10.8 (38)	40.7 (31)		570	18.6 (7)	52.5 (103)		97	* *	154	14.3 (3)
Yale Clini	39	* *	* *	* *	38	* *	* *	* *	12	* *	25	* *
Ypsilanti	814	11.4 (53)	45.9 (75)		674	29.1 (70)	43.5 (43)		198	37.9 (23)	155	38.1 (22)
Michigan	56,040	11.8	44.6		47,059	30.1	43.4		10,111	30.1	13,318	35.4
Nation (Prior Year)	658,420	12.7	42.2		558,067	25.6	44.8		100,071	30.6	241,000	28.1

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Agape WIC	413	79.4 (45)	2.4 (6)	419	10.0(125)
Albion Cli	193	67.9 (95)	20.2(110)	208	40.4 (40)
Alcona	46	* *	* *	52	* *
Algonac Cl	115	78.3 (57)	13.0 (90)	117	28.2 (87)
Allegan	85	* *	* *	99	* *
Alma WIC	46	* *	* *	49	* *
Alpena Cou	167	87.4 (9)	4.2 (23)	172	38.4 (50)
Antrim - B	55	* *	* *	58	* *
Arab/Chald	158	81.6 (34)	6.3 (49)	158	50.0 (11)
Arenac Cou	79	* *	* *	91	* *
Associated	8	* *	* *	8	* *
Baldwin	125	74.4 (75)	4.0 (20)	139	20.1(106)
Bangor	332	81.9 (30)	3.3 (10)	399	37.3 (55)
Baraga Cl	43	* *	* *	45	* *
Barry-Eato	273	78.8 (50)	11.4 (83)	300	46.7 (22)
Barry-Eato	344	84.9 (19)	5.8 (45)	384	33.3 (74)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Battle Cre	828	69.1 (93)	16.5(102)	893	35.2 (64)
Baxter Com	13	* *	* *	17	* *
Bay Cl	610	75.4 (70)	12.5 (87)	674	47.8 (19)
Bayside WI	3	* *	* *	6	* *
Beaver Isl	1	* *	* *	1	* *
Belding Cl	69	* *	* *	76	* *
Bell Memor	75	* *	* *	80	* *
Bendle/Bur	33	* *	* *	35	* *
Benton Har	611	66.9(100)	13.6 (93)	666	34.8 (69)
Benzie Cli	87	* *	* *	102	53.9 (3)
Berrien -	231	86.1 (11)	3.5 (14)	258	33.7 (73)
Bldhd Migr	4	* *	* *	7	* *
Booth Clin	0	* *	* *	1	* *
Branch-Hil	259	74.5 (74)	13.9 (95)	293	33.1 (75)
Branch-Hil	291	71.5 (84)	9.6 (72)	323	33.1 (75)
Branch-Hil	235	82.6 (27)	7.2 (58)	257	30.4 (79)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Browning-C	18	* *	* *	20	* *
Bryant Com	18	* *	* *	19	* *
Burton	1,144	77.4 (63)	8.8 (69)	1,208	28.3 (85)
Calumet WI	81	* *	* *	84	* *
Capac Clin	36	* *	* *	42	* *
Caro WIC S	2	* *	* *	3	* *
Cedar Spri	225	81.8 (31)	11.6 (84)	229	34.1 (70)
Charlevoix	147	78.2 (58)	13.6 (93)	162	41.4 (37)
Chass Heal	728	60.3(114)	21.8(111)	1,057	28.9 (83)
Cheboygan	163	89.0 (5)	4.9 (30)	176	46.6 (23)
Cherry Str	310	33.2(120)	49.7(119)	343	50.7 (8)
Chesaning	11	* *	* *	12	* *
Clare Coun	190	85.3 (14)	4.7 (29)	208	46.6 (23)
Clinica Sa	272	42.3(118)	16.2(101)	322	35.1 (66)
Comstock C	0	* *	* *	0	* *
Corner Hea	185	76.8 (64)	4.9 (30)	197	38.6 (47)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Crawford	97	* *	* *	106	50.9 (7)
Dewitt WIC	56	* *	* *	60	* *
Dickinson-	163	85.9 (12)	7.4 (61)	187	47.1 (21)
Dickinson-	75	* *	* *	86	* *
DMC Prim	12	* *	* *	12	* *
DMC Prim	45	* *	* *	46	* *
DMC Prim	14	* *	* *	14	* *
Dorr Outre	91	* *	* *	92	* *
Douglas He	551	64.4 (109)	10.7 (80)	663	25.6 (93)
Dowagiac	317	81.4 (37)	5.4 (42)	362	36.7 (57)
Durand WIC	81	* *	* *	87	* *
East Area	279	67.0 (99)	5.0 (34)	307	17.3(113)
Eastside H	452	67.5 (97)	4.2 (23)	467	13.7(121)
Eau Claire	87	* *	* *	189	46.6 (23)
Emmet Coun	223	84.3 (21)	9.0 (70)	241	46.1 (27)
Escanaba C	244	78.7 (52)	15.6 (99)	269	50.6 (9)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Family Com	322	76.7 (66)	5.0 (34)	350	24.3 (97)
Family Hlt	217	74.2 (76)	11.1 (82)	251	39.4 (45)
Family Roa	276	65.6(106)	3.3 (10)	299	11.7(122)
Farmington	96	* *	* *	94	* *
Fife Lake	0	* *	* *	0	* *
First Pres	0	* *	* *	0	* *
Flint Fam	413	66.6(102)	4.6 (28)	427	21.5(104)
Fruitport	0	* *	* *	0	* *
Gateway	368	66.0(103)	5.2 (38)	403	17.6(112)
Genesee -	1	* *	* *	12	* *
Gladwin Co	143	74.8 (73)	15.4 (98)	153	43.8 (30)
Gogebic Cl	81	* *	* *	94	* *
Golden	30	* *	* *	31	* *
Grace Prof	112	58.9(115)	5.4 (42)	119	15.1(119)
Grace Ross	783	73.4 (78)	3.4 (12)	843	16.5(115)
Grand Have	218	92.2 (1)	2.8 (8)	246	35.4 (62)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Grand Trav	0	* *	* *	0	* *
Grant	102	85.3 (14)	3.9 (19)	110	36.4 (59)
Gratiot	526	66.7(101)	5.3 (39)	541	10.2(123)
Greenville	118	83.1 (25)	10.2 (75)	120	36.7 (57)
Hamtramck	0	* *	* *	8	* *
Hancock Cl	139	87.1 (10)	9.4 (71)	162	50.6 (9)
Hannahvil	10	* *	* *	12	* *
Harper Gra	948	65.8(104)	6.5 (50)	1,069	14.4(120)
Haslett Cl	45	* *	* *	49	* *
Haslett St	6	* *	* *	6	* *
Herman Kie	1,160	67.8 (96)	6.9 (54)	1,239	17.3(113)
Highland P	0	* *	* *	4	* *
Holland Cl	716	85.1 (18)	1.7 (3)	786	34.9 (68)
Holly WIC	53	* *	* *	58	* *
Holton	0	* *	* *	0	* *
Howard Cit	49	* *	* *	51	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Hudsonv	172	89.5 (4)	1.2 (2)	178	29.2 (82)
Hurley Cli	0	* *	* *	2	* *
Huron Coun	209	84.2 (22)	5.7 (44)	218	47.7 (20)
Icchn-Spar	47	* *	* *	83	* *
Ingham Cou	1,385	79.0 (47)	3.5 (14)	1,427	16.0(116)
Intervale	112	72.3 (81)	7.1 (56)	113	15.9(117)
Ionia Coun	258	73.6 (77)	18.2(107)	275	37.8 (53)
Iosco	134	79.9 (43)	4.5 (26)	146	52.1 (4)
Ironwood	0	* *	* *	0	* *
Isabella C	265	77.7 (60)	12.5 (87)	277	31.4 (78)
Ithaca Cli	120	71.7 (82)	20.0(109)	133	39.8 (42)
Ithaca WIC	0	* *	* *	0	* *
Jackson Co	1,084	85.2 (17)	4.1 (22)	1,197	39.0 (46)
Jackson M	0	* *	* *	0	* *
Janes St A	339	56.3(117)	22.1(112)	383	51.4 (6)
Joy-Greenf	662	69.6 (91)	10.3 (76)	714	28.7 (84)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Clinic	Medical Care			WIC Enrollment		
	1st Trimester		No Care	1st Trimester		
	Number	% (Rank)	%	(Rank)	Number	% (Rank)
Kalamazoo	311	71.4 (85)	8.4	(64)	375	39.7 (43)
Kalamazoo	614	83.6 (24)	6.2	(48)	748	32.8 (77)
Kalkaska	131	80.9 (40)	8.4	(64)	143	40.6 (38)
Kent	0	* *	* *		0	* *
Kent Count	1,179	71.4 (85)	18.2	(107)	1,256	40.5 (39)
Kentwood	571	79.9 (43)	10.5	(78)	604	27.3 (89)
Keweenaw	24	* *	* *		25	* *
Kingsley	8	* *	* *		8	* *
Lac Vieux	11	* *	* *		10	* *
Lake Co	80	* *	* *		97	* *
Lake Odess	39	* *	* *		44	* *
Lapeer	441	67.1 (98)	24.7	(114)	474	42.2 (35)
Leelanu Cl	51	* *	* *		53	* *
Lenawee	0	* *	* *		0	* *
Leslie-Sto	15	* *	* *		17	* *
Lincoln Pa	381	73.2 (79)	8.7	(67)	411	21.9(103)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Livingston	324	81.8 (31)	8.6 (66)	326	25.5 (94)
Lowell	21	* *	* *	24	* *
M-72 Migra	3	* *	* *	13	* *
Mancelona	61	* *	* *	65	* *
Manchester	7	* *	* *	8	* *
Manchester	3	* *	* *	3	* *
Manistee C	125	76.8 (64)	14.4 (96)	154	35.1 (66)
Manistique	58	* *	* *	58	* *
Mansitee M	4	* *	* *	11	* *
Marion Bra	38	* *	* *	39	* *
Marquette	124	82.3 (28)	10.5 (78)	133	55.6 (2)
Marshall C	70	* *	* *	75	* *
Mason Clin	49	* *	* *	46	* *
Mason Co	174	75.3 (71)	14.4 (96)	200	48.5 (17)
Mccree Hea	1,392	71.0 (88)	8.2 (63)	1,538	22.3(101)
Mcnamara H	60	* *	* *	60	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Clinic	Medical Care			WIC Enrollment		
	1st Trimester		No Care	1st Trimester		
	Number	% (Rank)	%	(Rank)	Number	% (Rank)
Mecosta Co	272	85.3 (14)	3.7	(18)	291	35.4 (62)
Medical Pr	7	* *	*	*	7	* *
Medical Pr	8	* *	*	*	8	* *
Menominee	116	61.2(113)	30.2	(118)	134	35.8 (61)
Mercy	34	* *	*	*	35	* *
Mesick	0	* *	*	*	1	* *
Metropolit	0	* *	*	*	0	* *
Michigan S	70	* *	*	*	77	* *
Midland Cl	369	87.8 (8)	3.5	(14)	410	44.4 (29)
Midwest	376	75.8 (68)	4.0	(20)	404	23.5 (99)
Migrant H	93	* *	*	*	102	66.7 (1)
Missaukee	63	* *	*	*	65	* *
Moms & Bab	358	63.1(110)	5.3	(39)	424	18.2(111)
Monroe Cou	674	71.4 (85)	17.7	(106)	727	34.0 (71)
Montague/	0	* *	*	*	0	* *
Montcalm	6	* *	*	*	8	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Clinic	Medical Care				WIC Enrollment	
	1st Trimester		No Care		1st Trimester	
	Number	% (Rank)	% (Rank)		Number	% (Rank)
Montmorenc	57	* *	* *		55	* *
Mt Clemens	694	82.0 (29)	4.9 (30)		833	24.0 (98)
Munising C	32	* *	* *		36	* *
Muskegon C	1,481	75.8 (68)	12.5 (87)		1,566	38.3 (52)
New Center	6	* *	* *		6	* *
New Haven	63	* *	* *		69	* *
New Hope	1	* *	* *		1	* *
Newaygo	224	83.0 (26)	7.1 (56)		264	44.7 (28)
Newberry C	53	* *	* *		58	* *
North Camp	106	90.6 (2)	0.9 (1)		120	30.0 (81)
North Midl	0	* *	* *		0	* *
Northeast	838	69.3 (92)	8.0 (62)		890	19.1(108)
Northern M	41	* *	* *		43	* *
Northwest	198	65.2(107)	3.5 (14)		211	19.9(107)
Oakland Co	1,382	75.2 (72)	5.1 (37)		1,486	22.2(102)
Oceana Co	160	63.1(110)	28.1(115)		182	49.5 (13)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Ogemaw	137	80.3 (42)	7.3 (60)	153	51.6 (5)
Olhsa Star	220	85.5 (13)	2.3 (5)	240	26.3 (92)
Olhsa WIC	56	* *	* *	61	* *
Omni Care	25	* *	* *	26	* *
Omnicare	0	* *	* *	0	* *
Omnicare-G	0	* *	* *	0	* *
Omnicare-M	22	* *	* *	25	* *
Ontonagon	33	* *	* *	34	* *
Oscoda	51	* *	* *	58	* *
Otsego Co	187	78.6 (55)	12.3 (86)	196	49.0 (14)
Park Clin	0	* *	* *	0	* *
Park Medic	12	* *	* *	14	* *
Park Medic	20	* *	* *	20	* *
Park Medic	0	* *	* *	0	* *
Paw Paw Cl	117	78.6 (55)	6.8 (52)	126	37.3 (55)
PCOC	50	* *	* *	61	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Clinic	Medical Care				WIC Enrollment	
	1st Trimester		No Care		1st Trimester	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)
Pinconning	37	* *	* *	* *	39	* *
Port Huron	748	58.4 (116)	29.8 (117)	29.8 (117)	789	42.5 (34)
Portland	32	* *	* *	* *	31	* *
Presque Is	64	* *	* *	* *	61	* *
Profession	20	* *	* *	* *	20	* *
Pullman	197	79.2 (46)	6.6 (51)	6.6 (51)	245	39.6 (44)
Ravenna	0	* *	* *	* *	0	* *
Reed City	157	81.5 (36)	9.6 (72)	9.6 (72)	159	38.4 (50)
Region II	441	88.2 (7)	3.4 (12)	3.4 (12)	568	28.3 (85)
Riverview	31	* *	* *	* *	31	* *
Romulus Fi	0	* *	* *	* *	1	* *
Roosevelt-	57	* *	* *	* *	63	* *
Roscommon	127	81.1 (39)	8.7 (67)	8.7 (67)	140	38.6 (47)
S & S Medi	6	* *	* *	* *	7	* *
Saginaw Co	958	71.7 (82)	16.7 (104)	16.7 (104)	1,053	43.7 (31)
Saginaw-Ch	13	* *	* *	* *	13	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Sanilac Co	268	78.7 (52)	9.7 (74)	281	30.2 (80)
Sault Ste	194	78.9 (48)	13.4 (91)	217	46.5 (26)
Sawyer Afb	63	* *	* *	74	* *
School Bas	0	* *	* *	0	* *
Schoolcraf	457	68.5 (94)	4.2 (23)	474	10.1(124)
Seton Cent	16	* *	* *	28	* *
Shaftsburg	36	* *	* *	39	* *
Shelby M	72	* *	* *	109	48.6 (16)
Sheldon Co	536	64.6(108)	17.4(105)	581	36.3 (60)
Shiawassee	322	33.9(119)	63.4(120)	359	48.5 (17)
Southfield	819	77.5 (61)	3.1 (9)	879	18.5(110)
Sparta Pub	54	* *	* *	58	* *
St Charles	64	* *	* *	66	* *
St Ignace	48	* *	* *	53	* *
St Johns W	112	90.2 (3)	4.5 (26)	122	37.7 (54)
St Louis C	55	* *	* *	61	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
St.Clair S	296	83.8 (23)	6.1 (47)	331	26.6 (90)
Stanton Cl	222	80.6 (41)	10.8 (81)	242	42.6 (33)
Stockbridg	15	* *	* *	12	* *
Sumpter He	340	72.9 (80)	12.1 (85)	354	24.6 (96)
Taylor	836	70.7 (90)	13.5 (92)	958	23.3(100)
Teen Healt	1	* *	* *	1	* *
Temperance	5	* *	* *	5	* *
Thompsonv	9	* *	* *	10	* *
Three Oaks	45	* *	* *	42	* *
Three Rive	287	81.2 (38)	7.0 (55)	301	35.2 (64)
Traverse C	458	65.7(105)	24.2(113)	472	43.4 (32)
Tri-Cnty H	159	88.7 (6)	2.5 (7)	165	40.0 (41)
Tuscola	398	84.4 (20)	6.0 (46)	419	41.5 (36)
U Of M-Ped	0	* *	* *	0	* *
Vcms River	32	* *	* *	34	* *
Vicksburg	21	* *	* *	30	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Walled Lak	370	78.9 (48)	4.9 (30)	371	18.6(109)
Warren Cli	458	81.7 (33)	6.8 (52)	522	26.4 (91)
Washtenaw	27	* *	* *	27	* *
Wayne-Dear	742	76.1 (67)	10.4 (77)	834	38.5 (49)
Webbervill	14	* *	* *	12	* *
West Genes	179	81.6 (34)	2.2 (4)	194	15.5(118)
Western Wa	550	77.5 (61)	5.3 (39)	554	25.5 (94)
Westland C	627	78.8 (50)	7.2 (58)	687	21.1(105)
Westside C	495	62.0(112)	28.1(115)	519	49.7 (12)
Westwood	20	* *	* *	27	* *
Wexford	285	70.9 (89)	16.1(100)	310	49.0 (14)
Wheeler WI	11	* *	* *	12	* *
Whitmore L	14	* *	* *	16	* *
Willow	4	* *	* *	4	* *
Woodside	58	* *	* *	65	* *
Wyoming Cl	614	78.0 (59)	16.6(103)	628	33.9 (72)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Yale Clini	41	* *	* *	42	* *
Ypsilanti	822	78.7 (52)	5.0 (34)	886	27.9 (88)
Michigan	53,408	74.3	10.4	58,592	31.7
Nation (Prior Year)	437,601	75.9	9.6	666,694	27.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agape WIC	381	12.3 (8)	337	5.0 (5)	24	* *	0	* *	0	* *
Albion Cli	208	51.9 (88)	180	34.4 (101)	78	* *	0	* *	0	* *
Alcona	49	* *	39	* *	13	* *	0	* *	0	* *
Algonac Cl	122	53.3 (97)	102	36.3 (106)	48	* *	0	* *	0	* *
Allegan	94	* *	81	* *	34	* *	0	* *	0	* *
Alma WIC	41	* *	40	* *	11	* *	0	* *	0	* *
Alpena Cou	173	57.2 (108)	162	34.6 (102)	74	* *	0	* *	0	* *
Antrim - B	59	* *	51	* *	22	* *	0	* *	0	* *
Arab/Chald	137	4.4 (5)	135	4.4 (4)	5	* *	0	* *	0	* *
Arenac Cou	88	* *	42	* *	14	* *	0	* *	0	* *
Associated	7	* *	6	* *	1	* *	0	* *	0	* *
Baldwin	130	34.6 (34)	98	* *	27	* *	0	* *	0	* *
Bangor	384	35.9 (38)	345	20.3 (38)	96	* *	0	* *	0	* *
Baraga Cl	44	* *	40	* *	17	* *	0	* *	0	* *
Barry-Eato	259	57.9 (114)	216	33.3 (95)	80	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barry-Eato	384	52.3 (90)	324	28.4 (74)	123	23.6 (33)	0	* *	0	* *
Battle Cre	841	49.6 (77)	708	31.9 (91)	285	23.9 (32)	0	* *	0	* *
Baxter Com	17	* *	11	* *	4	* *	0	* *	0	* *
Bay Cl	567	57.3 (109)	503	28.0 (71)	189	26.5 (26)	0	* *	0	* *
Bayside WI	6	* *	2	* *	0	* *	0	* *	0	* *
Beaver Isl	2	* *	2	* *	0	* *	0	* *	0	* *
Belding Cl	74	* *	65	* *	29	* *	0	* *	0	* *
Bell Memor	78	* *	71	* *	33	* *	0	* *	0	* *
Bendle/Bur	37	* *	28	* *	11	* *	0	* *	0	* *
Benton Har	507	43.0 (57)	508	22.0 (46)	137	27.0 (25)	0	* *	0	* *
Benzie Cli	101	53.5 (101)	83	* *	37	* *	0	* *	0	* *
Berrien -	253	52.6 (93)	205	29.3 (77)	95	* *	0	* *	0	* *
Bldhd Migr	5	* *	2	* *	0	* *	0	* *	0	* *
Booth Clin	1	* *	0	* *	0	* *	0	* *	0	* *
Branch-Hil	266	33.1 (32)	205	23.9 (52)	55	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Branch-Hil	334	47.9 (70)	255	31.8 (90)	90	* *	0	* *	0	* *
Branch-Hil	255	53.3 (97)	231	34.6 (102)	96	* *	0	* *	0	* *
Browning-C	19	* *	15	* *	5	* *	0	* *	0	* *
Bryant Com	20	* *	16	* *	2	* *	0	* *	0	* *
Burton	1,087	52.3 (90)	930	31.3 (88)	324	26.5 (26)	0	* *	0	* *
Calumet WI	81	* *	68	* *	27	* *	0	* *	0	* *
Capac Clin	41	* *	35	* *	16	* *	0	* *	0	* *
Caro WIC S	3	* *	2	* *	0	* *	0	* *	0	* *
Cedar Spri	230	42.2 (54)	206	21.8 (43)	72	* *	0	* *	0	* *
Charlevoix	148	56.1 (105)	135	29.6 (79)	60	* *	0	* *	0	* *
Chass Heal	284	21.5 (16)	337	8.6 (9)	17	* *	0	* *	0	* *
Cheboygan	178	58.4 (116)	156	33.3 (95)	77	* *	0	* *	0	* *
Cherry Str	321	21.2 (15)	271	11.1 (14)	53	* *	0	* *	0	* *
Chesaning	11	* *	9	* *	2	* *	0	* *	0	* *
Clare Coun	144	82.6 (121)	94	* *	61	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Clinica Sa	312	1.0 (1)	288	0.0 (1)	3	* *	0	* *	0	* *
Comstock C	0	* *	0	* *	0	* *	0	* *	0	* *
Corner Hea	195	50.8 (83)	172	24.4 (55)	84	* *	0	* *	0	* *
Crawford	107	54.2 (102)	96	* *	45	* *	0	* *	0	* *
Dewitt WIC	54	* *	52	* *	19	* *	0	* *	0	* *
Dickinson-	179	54.7 (104)	151	38.4 (113)	76	* *	0	* *	0	* *
Dickinson-	83	* *	73	* *	30	* *	0	* *	0	* *
DMC Prim	12	* *	10	* *	1	* *	0	* *	0	* *
DMC Prim	41	* *	38	* *	5	* *	0	* *	0	* *
DMC Prim	14	* *	12	* *	3	* *	0	* *	0	* *
Dorr Outre	69	* *	57	* *	25	* *	0	* *	0	* *
Douglas He	538	15.8 (12)	589	6.1 (7)	56	* *	0	* *	0	* *
Dowagiac	322	49.1 (73)	254	32.3 (93)	107	26.2 (29)	0	* *	0	* *
Durand WIC	82	* *	81	* *	30	* *	0	* *	0	* *
East Area	291	20.3 (14)	217	12.4 (18)	31	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Eastside H	251	33.1 (32)	234	19.7 (36)	47	* *	0	* *	0	* *
Eau Claire	102	2.0 (3)	129	3.9 (3)	2	* *	0	* *	0	* *
Emmet Coun	237	49.4 (76)	177	32.8 (94)	85	* *	0	* *	0	* *
Escanaba C	258	56.6 (106)	233	35.6 (104)	106	21.7 (35)	0	* *	0	* *
Family Com	334	11.7 (7)	287	8.4 (8)	26	* *	0	* *	0	* *
Family Hlt	209	45.9 (66)	203	27.6 (69)	67	* *	0	* *	0	* *
Family Roa	299	24.1 (22)	263	12.9 (20)	45	* *	0	* *	0	* *
Farmington	91	* *	73	* *	10	* *	0	* *	0	* *
Fife Lake	0	* *	0	* *	0	* *	0	* *	0	* *
First Pres	0	* *	0	* *	0	* *	0	* *	0	* *
Flint Fam	437	39.4 (45)	340	25.3 (58)	114	22.8 (34)	0	* *	0	* *
Fruitport	0	* *	0	* *	0	* *	0	* *	0	* *
Gateway	387	23.8 (21)	304	14.5 (23)	35	* *	0	* *	0	* *
Genesee -	11	* *	0	* *	0	* *	0	* *	0	* *
Gladwin Co	154	52.6 (93)	138	30.4 (83)	66	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gogebic Cl	85	* *	80	* *	30	* *	0	* *	0	* *
Golden	26	* *	22	* *	4	* *	0	* *	0	* *
Grace Prof	95	* *	83	* *	11	* *	0	* *	0	* *
Grace Ross	780	26.3 (24)	609	11.2 (15)	122	51.6 (1)	0	* *	0	* *
Grand Have	244	41.4 (51)	216	23.1 (49)	72	* *	0	* *	0	* *
Grand Trav	0	* *	0	* *	0	* *	0	* *	0	* *
Grant	108	36.1 (39)	105	27.6 (69)	31	* *	0	* *	0	* *
Gratiot	544	18.2 (13)	503	9.5 (10)	49	* *	0	* *	0	* *
Greenville	104	57.7 (111)	105	30.5 (84)	48	* *	0	* *	0	* *
Hamtramck	8	* *	0	* *	0	* *	0	* *	0	* *
Hancock Cl	165	45.5 (62)	133	19.5 (34)	47	* *	0	* *	0	* *
Hannahvil	11	* *	11	* *	6	* *	0	* *	0	* *
Harper Gra	670	22.8 (17)	473	19.7 (36)	44	* *	0	* *	0	* *
Haslett Cl	45	* *	45	* *	13	* *	0	* *	0	* *
Haslett St	7	* *	6	* *	3	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Herman Kie	1,025	23.6 (20)	959	12.2 (17)	119	32.8 (10)	0	* *	0	* *
Highland P	4	* *	0	* *	0	* *	0	* *	0	* *
Holland Cl	783	29.0 (27)	682	13.8 (22)	157	47.8 (2)	0	* *	0	* *
Holly WIC	52	* *	43	* *	20	* *	0	* *	0	* *
Holton	0	* *	0	* *	0	* *	0	* *	0	* *
Howard Cit	42	* *	41	* *	14	* *	0	* *	0	* *
Hudsonv	182	37.4 (40)	172	16.3 (26)	49	* *	0	* *	0	* *
Hurley Cli	2	* *	0	* *	0	* *	0	* *	0	* *
Huron Coun	214	42.1 (53)	195	26.2 (63)	66	* *	0	* *	0	* *
Icchn-Spar	61	* *	44	* *	0	* *	0	* *	0	* *
Ingham Cou	1,324	42.7 (56)	1,172	25.7 (60)	322	27.6 (23)	0	* *	0	* *
Intervale	82	* *	72	* *	7	* *	0	* *	0	* *
Ionia Coun	276	46.0 (67)	238	30.3 (82)	95	* *	0	* *	0	* *
Iosco	141	63.8 (119)	122	36.9 (109)	67	* *	0	* *	0	* *
Ironwood	0	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Isabella C	252	46.4 (68)	134	15.7 (24)	35	* *	0	* *	0	* *
Ithaca Cli	100	64.0 (120)	101	30.7 (85)	36	* *	0	* *	0	* *
Ithaca WIC	0	* *	0	* *	0	* *	0	* *	0	* *
Jackson Co	1,212	51.3 (85)	1,052	32.0 (92)	433	28.2 (22)	0	* *	0	* *
Jackson M	0	* *	0	* *	0	* *	0	* *	0	* *
Janes St A	384	32.0 (31)	329	19.5 (34)	92	* *	0	* *	0	* *
Joy-Greenf	669	14.8 (10)	565	10.4 (13)	56	* *	0	* *	0	* *
Kalamazoo	311	40.8 (48)	298	24.2 (53)	86	* *	0	* *	0	* *
Kalamazoo	453	50.1 (79)	492	31.3 (88)	132	29.5 (18)	0	* *	0	* *
Kalkaska	143	54.5 (103)	135	38.5 (114)	64	* *	0	* *	0	* *
Kent	0	* *	0	* *	0	* *	0	* *	0	* *
Kent Count	1,148	42.3 (55)	1,020	20.6 (39)	284	31.7 (13)	0	* *	0	* *
Kentwood	560	30.5 (30)	528	15.7 (24)	110	46.4 (3)	0	* *	0	* *
Keweenaw	25	* *	23	* *	16	* *	0	* *	0	* *
Kingsley	8	* *	7	* *	6	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lac Vieux	11	* *	11	* *	7	* *	0	* *	0	* *
Lake Co	91	* *	69	* *	34	* *	0	* *	0	* *
Lake Odess	43	* *	35	* *	13	* *	0	* *	0	* *
Lapeer	460	51.5 (87)	405	29.9 (80)	167	29.3 (19)	0	* *	0	* *
Leelanu Cl	53	* *	47	* *	23	* *	0	* *	0	* *
Lenawee	0	* *	0	* *	0	* *	0	* *	0	* *
Leslie-Sto	16	* *	12	* *	3	* *	0	* *	0	* *
Lincoln Pa	399	45.1 (61)	284	28.2 (72)	95	* *	0	* *	0	* *
Livingston	328	53.0 (96)	292	29.5 (78)	109	31.2 (14)	0	* *	0	* *
Lowell	24	* *	19	* *	7	* *	0	* *	0	* *
M-72 Migra	6	* *	7	* *	0	* *	0	* *	0	* *
Mancelona	64	* *	54	* *	24	* *	0	* *	0	* *
Manchester	8	* *	6	* *	2	* *	0	* *	0	* *
Manchester	4	* *	3	* *	0	* *	0	* *	0	* *
Manistee C	144	49.3 (74)	127	28.3 (73)	57	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Manistique	61	* *	58	* *	28	* *	0	* *	0	* *
Mansitee M	6	* *	4	* *	0	* *	0	* *	0	* *
Marion Bra	34	* *	35	* *	20	* *	0	* *	0	* *
Marquette	130	52.3 (90)	120	33.3 (95)	56	* *	0	* *	0	* *
Marshall C	75	* *	62	* *	22	* *	0	* *	0	* *
Mason Clin	48	* *	43	* *	10	* *	0	* *	0	* *
Mason Co	199	47.2 (69)	176	23.3 (50)	71	* *	0	* *	0	* *
Mccree Hea	1,382	39.3 (44)	1,232	24.4 (55)	351	28.8 (20)	0	* *	0	* *
Mcnamara H	63	* *	48	* *	12	* *	0	* *	0	* *
Mecosta Co	288	49.3 (74)	260	26.5 (65)	97	* *	0	* *	0	* *
Medical Pr	6	* *	5	* *	1	* *	0	* *	0	* *
Medical Pr	4	* *	4	* *	0	* *	0	* *	0	* *
Menominee	116	57.8 (112)	114	36.8 (108)	49	* *	0	* *	0	* *
Mercy	31	* *	27	* *	3	* *	0	* *	0	* *
Mesick	1	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Metropolit	0	* *	0	* *	0	* *	0	* *	0	* *
Michigan S	69	* *	62	* *	3	* *	0	* *	0	* *
Midland Cl	407	44.0 (58)	346	22.8 (47)	134	29.9 (17)	0	* *	0	* *
Midwest	391	15.6 (11)	335	9.6 (11)	40	* *	0	* *	0	* *
Migrant H	102	2.0 (3)	84	* *	2	* *	0	* *	0	* *
Missaukee	63	* *	60	* *	22	* *	0	* *	0	* *
Moms & Bab	371	12.9 (9)	354	5.9 (6)	27	* *	0	* *	0	* *
Monroe Cou	730	50.8 (83)	636	33.6 (98)	262	26.0 (30)	0	* *	0	* *
Montague/	0	* *	0	* *	0	* *	0	* *	0	* *
Montcalm	6	* *	5	* *	0	* *	0	* *	0	* *
Montmorenc	54	* *	48	* *	27	* *	0	* *	0	* *
Mt Clemens	778	41.4 (51)	253	18.2 (32)	60	* *	0	* *	0	* *
Munising C	37	* *	28	* *	12	* *	0	* *	0	* *
Muskegon C	1,371	50.1 (79)	1,294	30.8 (86)	490	26.3 (28)	0	* *	0	* *
New Center	5	* *	5	* *	1	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
New Haven	69	* *	29	* *	2	* *	0	* *	0	* *
New Hope	1	* *	1	* *	0	* *	0	* *	0	* *
Newaygo	247	49.0 (71)	209	25.4 (59)	80	* *	0	* *	0	* *
Newberry C	58	* *	51	* *	26	* *	0	* *	0	* *
North Camp	116	1.7 (2)	95	* *	1	* *	0	* *	0	* *
North Midl	0	* *	0	* *	0	* *	0	* *	0	* *
Northeast	571	24.2 (23)	515	12.8 (19)	55	* *	0	* *	0	* *
Northern M	43	* *	39	* *	21	* *	0	* *	0	* *
Northwest	201	23.4 (18)	155	18.7 (33)	32	* *	0	* *	0	* *
Oakland Co	1,450	35.2 (37)	1,298	20.6 (39)	335	33.4 (8)	0	* *	0	* *
Oceana Co	177	45.8 (65)	154	23.4 (51)	61	* *	0	* *	0	* *
Ogemaw	153	56.9 (107)	126	30.2 (81)	64	* *	0	* *	0	* *
Olhsa Star	235	34.9 (36)	182	20.9 (42)	43	* *	0	* *	0	* *
Olhsa WIC	60	* *	45	* *	9	* *	0	* *	0	* *
Omni Care	15	* *	13	* *	2	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Omicare	0	* *	0	* *	0	* *	0	* *	0	* *
Omicare-G	0	* *	0	* *	0	* *	0	* *	0	* *
Omicare-M	23	* *	19	* *	2	* *	0	* *	0	* *
Ontonagon	34	* *	28	* *	10	* *	0	* *	0	* *
Oscoda	57	* *	51	* *	28	* *	0	* *	0	* *
Otsego Co	199	53.3 (97)	169	21.9 (44)	78	* *	0	* *	0	* *
Park Clin	0	* *	0	* *	0	* *	0	* *	0	* *
Park Medic	13	* *	12	* *	1	* *	0	* *	0	* *
Park Medic	15	* *	14	* *	2	* *	0	* *	0	* *
Park Medic	0	* *	0	* *	0	* *	0	* *	0	* *
Paw Paw Cl	128	57.8 (112)	120	36.7 (107)	55	* *	0	* *	0	* *
PCOC	41	* *	39	* *	10	* *	0	* *	0	* *
Pinconning	35	* *	27	* *	8	* *	0	* *	0	* *
Port Huron	791	58.9 (117)	718	35.7 (105)	366	30.6 (15)	0	* *	0	* *
Portland	32	* *	31	* *	13	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Presque Is	62	* *	60	* *	20	* *	0	* *	0	* *
Profession	15	* *	13	* *	1	* *	0	* *	0	* *
Pullman	231	38.1 (42)	187	21.9 (44)	53	* *	0	* *	0	* *
Ravenna	0	* *	0	* *	0	* *	0	* *	0	* *
Reed City	162	57.4 (110)	140	37.1 (110)	67	* *	0	* *	0	* *
Region II	408	39.7 (47)	490	26.3 (64)	109	30.3 (16)	0	* *	0	* *
Riverview	28	* *	23	* *	4	* *	0	* *	0	* *
Romulus Fi	1	* *	0	* *	0	* *	0	* *	0	* *
Roosevelt-	62	* *	51	* *	13	* *	0	* *	0	* *
Roscommon	141	61.0 (118)	120	38.3 (112)	61	* *	0	* *	0	* *
S & S Medi	6	* *	4	* *	0	* *	0	* *	0	* *
Saginaw Co	1,058	29.3 (28)	802	20.7 (41)	185	31.9 (12)	0	* *	0	* *
Saginaw-Ch	13	* *	8	* *	2	* *	0	* *	0	* *
Sanilac Co	291	45.7 (63)	264	26.1 (62)	99	* *	0	* *	0	* *
Sault Ste	213	50.2 (81)	175	27.4 (68)	75	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sawyer Afb	74	* *	60	* *	27	* *	0	* *	0	* *
School Bas	0	* *	0	* *	0	* *	0	* *	0	* *
Schoolcraf	452	27.2 (25)	398	11.6 (16)	72	* *	0	* *	0	* *
Seton Cent	16	* *	10	* *	0	* *	0	* *	0	* *
Shaftsburg	37	* *	32	* *	15	* *	0	* *	0	* *
Shelby M	94	* *	76	* *	3	* *	0	* *	0	* *
Sheldon Co	559	23.4 (18)	467	13.1 (21)	74	* *	0	* *	0	* *
Shiawassee	146	50.7 (82)	310	31.0 (87)	44	* *	0	* *	0	* *
Southfield	613	34.6 (34)	621	16.6 (27)	131	27.5 (24)	0	* *	0	* *
Sparta Pub	59	* *	49	* *	17	* *	0	* *	0	* *
St Charles	66	* *	57	* *	21	* *	0	* *	0	* *
St Ignace	50	* *	46	* *	20	* *	0	* *	0	* *
St Johns W	117	51.3 (85)	101	25.7 (60)	40	* *	0	* *	0	* *
St Louis C	47	* *	49	* *	22	* *	0	* *	0	* *
St.Clair S	322	44.4 (60)	92	* *	20	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Stanton Cl	213	58.2 (115)	208	34.1 (100)	96	* *	0	* *	0	* *
Stockbridg	14	* *	14	* *	4	* *	0	* *	0	* *
Sumpter He	353	40.8 (48)	317	23.0 (48)	81	* *	0	* *	0	* *
Taylor	753	52.1 (89)	723	28.9 (76)	267	32.2 (11)	0	* *	0	* *
Teen Healt	1	* *	1	* *	1	* *	0	* *	0	* *
Temperance	5	* *	5	* *	2	* *	0	* *	0	* *
Thompsonv	10	* *	8	* *	4	* *	0	* *	0	* *
Three Oaks	43	* *	44	* *	21	* *	0	* *	0	* *
Three Rive	293	45.7 (63)	253	33.6 (98)	91	* *	0	* *	0	* *
Traverse C	492	49.0 (71)	436	24.3 (54)	171	40.4 (5)	0	* *	0	* *
Tri-Cnty H	97	* *	63	* *	39	* *	0	* *	0	* *
Tuscola	392	53.3 (97)	365	27.1 (67)	132	35.6 (7)	0	* *	0	* *
U Of M-Ped	0	* *	0	* *	0	* *	0	* *	0	* *
Vcms River	34	* *	32	* *	4	* *	0	* *	0	* *
Vicksburg	14	* *	20	* *	3	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)		Number	% (Rank)	Number	% (Rank)	
Walled Lak	375	44.3 (59)	355	26.8 (66)	94	* *	0	* *	0	* *
Warren Cli	485	39.4 (45)	144	9.7 (12)	25	* *	0	* *	0	* *
Washtenaw	25	* *	23	* *	4	* *	0	* *	0	* *
Wayne-Dear	777	8.9 (6)	611	2.0 (2)	34	* *	0	* *	0	* *
Webbervill	16	* *	11	* *	5	* *	0	* *	0	* *
West Genes	177	37.9 (41)	164	17.7 (31)	42	* *	0	* *	0	* *
Western Wa	545	41.1 (50)	454	28.4 (74)	125	24.8 (31)	0	* *	0	* *
Westland C	564	49.6 (77)	387	25.1 (57)	129	33.3 (9)	0	* *	0	* *
Westside C	504	28.6 (26)	458	17.2 (29)	104	28.8 (20)	0	* *	0	* *
Westwood	19	* *	20	* *	2	* *	0	* *	0	* *
Wexford	296	52.7 (95)	234	37.2 (111)	99	* *	0	* *	0	* *
Wheeler WI	8	* *	8	* *	3	* *	0	* *	0	* *
Whitmore L	16	* *	12	* *	7	* *	0	* *	0	* *
Willow	5	* *	4	* *	0	* *	0	* *	0	* *
Woodside	63	* *	53	* *	28	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wyoming Cl	619	38.6 (43)	580	17.4 (30)	165	42.4 (4)	0	* *	0	* *
Yale Clini	40	* *	42	* *	23	* *	0	* *	0	* *
Ypsilanti	834	30.3 (29)	715	17.1 (28)	154	36.4 (6)	0	* *	0	* *
Michigan	52,553	39.6	45,315	23.1	13,107	30.5	0	*	0	*
Nation (Prior Year)	591,104	30.8	423,755	20.3	107,528	31.1	305,692	27.7	340,900	24.0

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agape WIC	391	9.0 (96)	7.7 (43)	394	0.5 (106)	386	9.1 (97)	378	40.7 (96)
Albion Cli	184	8.2 (89)	6.5 (22)	188	0.0 (1)	184	8.2 (87)	182	39.6 (101)
Alcona	23	* *	* *	45	* *	23	* *	43	* *
Algonac Cl	103	5.8 (38)	9.7 (74)	102	0.0 (1)	102	5.9 (39)	103	43.7 (89)
Allegan	76	* *	* *	90	* *	76	* *	84	* *
Alma WIC	36	* *	* *	40	* *	36	* *	40	* *
Alpena Cou	166	7.2 (68)	10.8 (90)	166	0.6 (110)	164	6.7 (54)	167	57.5 (36)
Antrim - B	45	* *	* *	53	* *	45	* *	51	* *
Arab/Chald	118	5.1 (25)	9.3 (69)	139	0.0 (1)	118	5.1 (23)	138	52.9 (59)
Arenac Cou	41	* *	* *	43	* *	40	* *	41	* *
Associated	6	* *	* *	6	* *	6	* *	6	* *
Baldwin	106	10.4 (107)	6.6 (24)	112	0.9 (117)	106	10.4 (106)	112	61.6 (25)
Bangor	222	8.1 (88)	7.7 (43)	346	0.0 (1)	220	8.2 (87)	330	52.7 (60)
Baraga Cl	40	* *	* *	41	* *	40	* *	40	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Barry-Eato	234	6.0 (45)	7.7 (43)	252	0.8 (116)	231	6.1 (44)	243	46.5 (79)
Barry-Eato	313	7.7 (79)	10.9 (91)	344	0.6 (110)	308	7.8 (82)	326	59.2 (32)
Battle Cre	792	5.9 (42)	8.5 (52)	813	0.0 (1)	790	5.9 (39)	755	40.3 (98)
Baxter Com	11	* *	* *	11	* *	11	* *	10	* *
Bay Cl	561	5.3 (28)	11.6 (99)	577	0.0 (1)	559	5.4 (29)	565	51.2 (64)
Bayside Wl	2	* *	* *	2	* *	2	* *	1	* *
Beaver Isl	2	* *	* *	2	* *	2	* *	2	* *
Belding Cl	65	* *	* *	65	* *	65	* *	63	* *
Bell Memor	70	* *	* *	70	* *	70	* *	65	* *
Bendle/Bur	29	* *	* *	29	* *	29	* *	29	* *
Benton Har	575	9.0 (96)	7.3 (34)	578	0.0 (1)	575	9.0 (96)	558	41.4 (94)
Benzie Cli	90	* *	* *	91	* *	90	* *	84	* *
Berrien -	218	6.9 (60)	11.0 (94)	226	0.0 (1)	218	6.9 (61)	214	53.3 (57)
Bldhd Migr	1	* *	* *	4	* *	1	* *	3	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Booth Clin	0	* *	* *		0	* *	0	* *	0	* *
Branch-Hil	196	5.6 (35)	5.1 (15)		253	0.4 (100)	195	5.1 (23)	232	60.8 (27)
Branch-Hil	261	7.7 (79)	13.0 (108)		283	0.0 (1)	261	7.7 (78)	281	48.8 (73)
Branch-Hil	237	7.2 (68)	4.6 (5)		237	0.0 (1)	237	7.2 (69)	226	54.9 (47)
Browning-C	15	* *	* *		16	* *	15	* *	12	* *
Bryant Com	17	* *	* *		16	* *	16	* *	17	* *
Burton	953	5.6 (35)	9.1 (65)		1,116	0.2 (81)	945	5.6 (35)	1,091	40.1 (99)
Calumet WI	64	* *	* *		70	* *	64	* *	66	* *
Capac Clin	34	* *	* *		35	* *	34	* *	34	* *
Caro WIC S	2	* *	* *		2	* *	2	* *	2	* *
Cedar Spri	214	2.3 (2)	13.1 (110)		215	0.0 (1)	214	2.3 (2)	198	63.6 (22)
Charlevoix	139	5.0 (23)	5.0 (13)		144	0.0 (1)	138	5.1 (23)	125	69.6 (13)
Chass Heal	885	5.9 (42)	11.2 (98)		934	0.1 (74)	878	5.9 (39)	907	51.6 (62)
Cheboygan	155	4.5 (16)	12.9 (106)		155	0.0 (1)	155	4.5 (15)	153	60.8 (27)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Cherry Str	284	6.7 (54)	8.1 (50)	283	0.0 (1)	282	6.7 (54)	280	64.6 (20)
Chesaning	11	* *	* *	11	* *	11	* *	11	* *
Clare Coun	171	8.2 (89)	7.6 (40)	179	0.6 (110)	169	8.3 (90)	169	43.8 (88)
Clinica Sa	286	4.5 (16)	7.0 (30)	286	0.0 (1)	286	4.5 (15)	273	94.5 (3)
Comstock C	0	* *	* *	0	* *	0	* *	0	* *
Corner Hea	174	9.2 (100)	8.6 (54)	177	0.0 (1)	174	9.2 (99)	157	44.6 (84)
Crawford	94	* *	* *	94	* *	94	* *	85	* *
Dewitt WIC	41	* *	* *	56	* *	41	* *	53	* *
Dickinson-	162	4.3 (14)	13.6 (115)	164	0.0 (1)	162	4.3 (13)	163	53.4 (56)
Dickinson-	70	* *	* *	71	* *	70	* *	69	* *
DMC Prim	12	* *	* *	11	* *	11	* *	12	* *
DMC Prim	45	* *	* *	47	* *	45	* *	46	* *
DMC Prim	13	* *	* *	13	* *	13	* *	13	* *
Dorr Outre	86	* *	* *	88	* *	86	* *	87	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Douglas He	586	6.5 (49)	9.7 (74)	601	0.3 (92)	584	6.5 (48)	587	48.7 (74)
Dowagiac	294	8.2 (89)	7.1 (32)	310	0.0 (1)	293	8.2 (87)	262	55.3 (45)
Durand WIC	81	* *	* *	82	* *	81	* *	75	* *
East Area	175	9.1 (98)	6.3 (20)	246	0.4 (100)	174	8.6 (92)	239	21.3 (118)
Eastside H	430	12.6 (112)	4.0 (2)	432	0.0 (1)	427	12.6 (111)	320	39.7 (100)
Eau Claire	136	3.7 (8)	12.5 (104)	153	0.7 (114)	135	3.7 (8)	150	78.0 (8)
Emmet Coun	186	5.4 (30)	4.8 (9)	216	0.0 (1)	185	5.4 (29)	183	78.7 (7)
Esanaba C	243	5.8 (38)	12.3 (103)	244	0.0 (1)	243	5.8 (37)	231	50.2 (71)
Family Com	261	6.5 (49)	7.3 (34)	308	0.3 (92)	260	6.5 (48)	300	45.0 (82)
Family Hlt	199	7.5 (74)	7.5 (37)	209	0.0 (1)	199	7.5 (73)	197	50.3 (70)
Family Roa	263	14.8 (117)	7.2 (33)	263	0.0 (1)	263	14.8 (116)	252	15.9 (121)
Farmington	81	* *	* *	83	* *	80	* *	83	* *
Fife Lake	0	* *	* *	0	* *	0	* *	0	* *
First Pres	0	* *	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Flint Fam	362	6.6 (52)	6.9 (27)	383	0.0 (1)	361	6.6 (52)	382	33.5 (109)
Fruitport	0	* *	* *	0	* *	0	* *	0	* *
Gateway	240	14.6 (116)	5.0 (13)	328	0.0 (1)	240	14.6 (115)	309	19.4 (119)
Genesee -	0	* *	* *	0	* *	0	* *	0	* *
Gladwin Co	127	3.1 (5)	11.8 (100)	140	0.0 (1)	127	3.1 (4)	139	61.2 (26)
Gogebic Cl	77	* *	* *	77	* *	77	* *	72	* *
Golden	29	* *	* *	29	* *	29	* *	27	* *
Grace Prof	103	4.9 (20)	6.8 (25)	102	0.0 (1)	102	4.9 (19)	103	18.4 (120)
Grace Ross	682	10.6 (108)	4.7 (7)	699	0.0 (1)	682	10.6 (107)	679	44.3 (86)
Grand Have	209	7.7 (79)	10.0 (79)	219	0.0 (1)	209	7.7 (78)	212	54.7 (50)
Grand Trav	0	* *	* *	0	* *	0	* *	0	* *
Grant	101	6.9 (60)	10.9 (91)	105	0.0 (1)	100	7.0 (65)	105	53.3 (57)
Gratiot	510	12.9 (114)	7.6 (40)	515	1.4 (121)	501	13.0 (113)	511	30.5 (114)
Greenville	112	8.0 (87)	8.9 (63)	111	0.0 (1)	111	8.1 (86)	104	54.8 (49)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
	Number	% (Rank)	% (Rank)							
Hamtramck	0	* * *	* *	0	* *	0	* *	0	* *	
Hancock Cl	135	3.7 (8)	13.3 (112)	141	0.0 (1)	135	3.7 (8)	137	75.2 (11)	
Hannahvil	11	* * *	* *	11	* *	11	* *	11	* *	
Harper Gra	866	11.9 (111)	6.4 (21)	959	0.3 (92)	859	12.0 (110)	811	33.4 (110)	
Haslett Cl	42	* * *	* *	47	* *	42	* *	42	* *	
Haslett St	6	* * *	* *	6	* *	6	* *	5	* *	
Herman Kie	1,045	11.4 (110)	4.9 (10)	1,117	0.1 (74)	1,041	11.4 (109)	1,077	31.1 (112)	
Highland P	0	* * *	* *	0	* *	0	* *	0	* *	
Holland Cl	540	6.7 (54)	8.7 (55)	710	0.3 (92)	536	6.3 (46)	679	67.0 (17)	
Holly WIC	44	* * *	* *	45	* *	43	* *	46	* *	
Holton	0	* * *	* *	0	* *	0	* *	0	* *	
Howard Cit	46	* * *	* *	46	* *	46	* *	44	* *	
Hudsonv	130	7.7 (79)	7.7 (43)	174	0.0 (1)	130	7.7 (78)	166	62.7 (24)	
Hurley Cli	0	* * *	* *	0	* *	0	* *	0	* *	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Huron Coun	196	6.6 (52)	13.3 (112)		197	0.0 (1)	195	6.7 (54)	187	43.9 (87)
Icchn-Spar	56	* *	* *		64	* *	56	* *	64	* *
Ingham Cou	1,238	7.2 (68)	9.1 (65)		1,352	0.2 (81)	1,234	7.2 (69)	1,159	65.1 (19)
Intervale	109	12.8(113)	9.2 (68)		110	0.0 (1)	109	12.8(112)	108	25.9(115)
Ionia Coun	240	7.5 (74)	8.8 (57)		242	0.0 (1)	240	7.5 (73)	239	51.0 (65)
Iosco	83	* *	* *		131	0.0 (1)	83	* *	125	36.0(105)
Ironwood	0	* *	* *		0	* *	0	* *	0	* *
Isabella C	204	4.9 (20)	5.9 (18)		204	0.0 (1)	203	4.9 (19)	199	75.9 (10)
Ithaca Cli	95	* *	* *		114	0.0 (1)	94	* *	110	53.6 (55)
Ithaca WIC	0	* *	* *		0	* *	0	* *	0	* *
Jackson Co	1,050	6.7 (54)	9.0 (64)		1,072	0.1 (74)	1,045	6.6 (52)	1,057	51.5 (63)
Jackson M	0	* *	* *		0	* *	0	* *	0	* *
Janes St A	324	9.9(106)	4.3 (4)		324	0.3 (92)	323	9.9(105)	293	25.9(115)
Joy-Greenf	603	7.5 (74)	9.5 (73)		622	0.3 (92)	600	7.5 (73)	608	56.1 (43)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kalamazoo	291	8.6 (93)	9.3 (69)	304	0.0 (1)	289	8.7 (94)	269	48.3 (76)
Kalamazoo	625	7.8 (85)	10.6 (88)	660	0.2 (81)	623	7.9 (85)	589	45.5 (80)
Kalkaska	134	4.5 (16)	6.0 (19)	135	0.0 (1)	134	4.5 (15)	128	66.4 (18)
Kent	0	* *	* *	0	* *	0	* *	0	* *
Kent Count	1,080	7.1 (64)	9.7 (74)	1,112	0.2 (81)	1,075	7.2 (69)	1,040	57.2 (37)
Kentwood	557	7.4 (73)	9.7 (74)	569	0.2 (81)	556	7.4 (72)	542	56.6 (38)
Keweenaw	23	* *	* *	23	* *	23	* *	23	* *
Kingsley	8	* *	* *	8	* *	8	* *	8	* *
Lac Vieux	10	* *	* *	9	* *	9	* *	10	* *
Lake Co	76	* *	* *	78	* *	76	* *	77	* *
Lake Odess	35	* *	* *	36	* *	35	* *	36	* *
Lapeer	414	5.6 (35)	11.1 (96)	420	0.2 (81)	413	5.6 (35)	407	50.4 (69)
Leelanu CI	44	* *	* *	49	* *	44	* *	47	* *
Lenawee	0	* *	* *	0	* *	0	* *	0	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Leslie-Sto	12	* * * *			14	* *			12	* *
Lincoln Pa	291	5.5 (32)	12.0 (101)		300	1.0 (118)			289	5.5 (31)
Livingston	310	7.7 (79)	16.8 (117)		317	0.3 (92)			309	7.8 (82)
Lowell	17	* * * *			19	* *			17	* *
M-72 Migra	7	* * * *			12	* *			7	* *
Mancelona	51	* * * *			53	* *			50	* *
Manchester	6	* * * *			6	* *			6	* *
Manchester	3	* * * *			3	* *			3	* *
Manistee C	131	9.2 (100)	9.9 (78)		132	0.0 (1)			130	9.2 (99)
Manistique	59	* * * *			59	* *			59	* *
Mansitee M	6	* * * *			8	* *			6	* *
Marion Bra	19	* * * *			37	* *			19	* *
Marquette	117	4.3 (14)	11.1 (96)		121	0.0 (1)			116	4.3 (13)
Marshall C	60	* * * *			65	* *			60	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Mason Clin	43	* * * *	* * * *	47	* *	42	* *	41	* *	
Mason Co	174	4.0 (11)	10.9 (91)	174	0.0 (1)	174	4.0 (10)	168	50.6 (67)	
Mccree Hea	1,328	9.6 (103)	5.4 (16)	1,373	0.3 (92)	1,324	9.5 (101)	1,212	35.7 (108)	
Mcnamara H	54	* * * *	* * * *	54	* *	54	* *	46	* *	
Mecosta Co	261	4.6 (19)	13.0 (108)	266	0.0 (1)	261	4.6 (18)	263	59.3 (31)	
Medical Pr	6	* * * *	* * * *	6	* *	6	* *	6	* *	
Medical Pr	8	* * * *	* * * *	8	* *	8	* *	8	* *	
Menominee	113	7.1 (64)	8.0 (48)	114	0.0 (1)	113	7.1 (67)	100	31.0 (113)	
Mercy	28	* * * *	* * * *	30	* *	27	* *	29	* *	
Mesick	0	* * * *	* * * *	0	* *	0	* *	0	* *	
Metropolit	0	* * * *	* * * *	0	* *	0	* *	0	* *	
Michigan S	64	* * * *	* * * *	70	* *	64	* *	67	* *	
Midland Cl	354	4.2 (13)	10.2 (83)	355	0.0 (1)	354	4.2 (12)	310	67.1 (16)	
Midwest	288	9.4 (102)	7.6 (40)	358	1.1 (120)	282	9.6 (102)	354	35.9 (106)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)												
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed			
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%		
	Number	% (Rank)	% (Rank)	(Rank)							(Rank)	(Rank)
Migrant H	85	* *	* *		85	* *		85	* *		84	* *
Missaukee	59	* *	* *		60	* *		59	* *		56	* *
Moms & Bab	355	8.7 (95)	6.5 (22)		382	0.0 (1)		353	8.8 (95)		370	35.9 (106)
Monroe Cou	640	7.5 (74)	8.8 (57)		656	0.2 (81)		638	7.5 (73)		621	44.4 (85)
Montague/	0	* *	* *		0	* *		0	* *		0	* *
Montcalm	6	* *	* *		7	* *		6	* *		7	* *
Montmorenc	54	* *	* *		56	* *		54	* *		56	* *
Mt Clemens	300	7.0 (63)	9.3 (69)		313	0.0 (1)		300	7.0 (65)		309	95.5 (2)
Munising C	30	* *	* *		30	* *		30	* *		29	* *
Muskegon C	1,375	5.9 (42)	8.8 (57)		1,407	0.1 (74)		1,373	5.9 (39)		1,318	39.4 (102)
New Center	6	* *	* *		6	* *		6	* *		6	* *
New Haven	37	* *	* *		37	* *		36	* *		38	* *
New Hope	1	* *	* *		1	* *		1	* *		1	* *
Newaygo	219	9.1 (98)	11.0 (94)		226	0.0 (1)		219	9.1 (97)		221	56.6 (38)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)											
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed		
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%	
	Number	% (Rank)	% (Rank)	(Rank)							(Rank)
Newberry C	48	* * * *	* * * *	50	* *	* *	48	* *	* *	48	* *
North Camp	100	3.0 (4)	10.0 (79)	101	0.0 (1)		99	* *		101	92.1 (4)
North Midl	0	* * * *	* * * *	0	* *	* *	0	* *	* *	0	* *
Northeast	664	10.7(109)	4.1 (3)	758	0.1 (74)		661	10.7(108)		715	32.4(111)
Northern M	39	* * * *	* * * *	39	* *	* *	39	* *	* *	37	* *
Northwest	131	13.7(115)	6.9 (27)	174	0.0 (1)		130	13.8(114)		174	25.3(117)
Oakland Co	1,345	7.5 (74)	8.8 (57)	1,371	0.2 (81)		1,339	7.5 (73)		1,348	43.1 (92)
Oceana Co	157	9.6(103)	7.0 (30)	158	0.0 (1)		157	9.6(102)		151	52.3 (61)
Ogemaw	125	3.2 (7)	8.8 (57)	132	0.0 (1)		125	3.2 (6)		128	46.9 (77)
Olhsa Star	192	7.3 (72)	4.7 (7)	201	0.5 (106)		191	6.8 (58)		196	51.0 (65)
Olhsa WIC	47	* * * *	* * * *	51	* *	* *	47	* *	* *	49	* *
Omni Care	24	* * * *	* * * *	24	* *	* *	24	* *	* *	24	* *
Omnicare	0	* * * *	* * * *	0	* *	* *	0	* *	* *	0	* *
Omnicare-G	0	* * * *	* * * *	0	* *	* *	0	* *	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Omnicare-M	20	* *	* *	* *	22	* *	20	* *	22	* *
Ontonagon	29	* *	* *	* *	31	* *	29	* *	29	* *
Oscoda	51	* *	* *	* *	52	* *	51	* *	49	* *
Otsego Co	164	6.7 (54)	9.1 (65)		172	0.0 (1)	163	6.7 (54)	143	72.7 (12)
Park Clin	0	* *	* *	* *	0	* *	0	* *	0	* *
Park Medic	12	* *	* *	* *	13	* *	12	* *	13	* *
Park Medic	20	* *	* *	* *	20	* *	20	* *	20	* *
Park Medic	0	* *	* *	* *	0	* *	0	* *	0	* *
Paw Paw Cl	67	* *	* *	* *	118	0.0 (1)	67	* *	116	44.8 (83)
PCOC	47	* *	* *	* *	51	* *	47	* *	42	* *
Pinconning	31	* *	* *	* *	32	* *	31	* *	31	* *
Port Huron	711	6.8 (58)	4.9 (10)		722	0.1 (74)	710	6.8 (58)	712	40.9 (95)
Portland	31	* *	* *	* *	31	* *	31	* *	31	* *
Presque Is	60	* *	* *	* *	60	* *	60	* *	60	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Profession	19	* *	* *		19	* *	19	* *	19	* *
Pullman	160	1.9 (1)	6.9 (27)		202	1.0 (118)	159	1.9 (1)	193	54.9 (47)
Ravenna	0	* *	* *		0	* *	0	* *	0	* *
Reed City	137	5.8 (38)	13.1 (110)		143	0.7 (114)	136	5.9 (39)	133	63.9 (21)
Region II	527	5.5 (32)	4.6 (5)		529	0.4 (100)	525	5.5 (31)	525	45.1 (81)
Riverview	30	* *	* *		30	* *	29	* *	31	* *
Romulus Fi	0	* *	* *		0	* *	0	* *	0	* *
Roosevelt-	55	* *	* *		55	* *	55	* *	51	* *
Roscommon	120	5.0 (23)	7.5 (37)		123	0.0 (1)	120	5.0 (22)	112	56.3 (42)
S & S Medi	5	* *	* *		5	* *	5	* *	5	* *
Saginaw Co	915	7.7 (79)	8.4 (51)		918	0.1 (74)	912	7.7 (78)	844	36.4 (104)
Saginaw-Ch	9	* *	* *		9	* *	9	* *	9	* *
Sanilac Co	265	6.8 (58)	4.9 (10)		267	0.0 (1)	264	6.8 (58)	245	53.9 (52)
Sault Ste	193	3.1 (5)	13.5 (114)		193	0.0 (1)	192	3.1 (4)	187	59.9 (29)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Sawyer Afb	61	* * * *	* * * *	61	* *	61	* *	53	* *	
School Bas	0	* * * *	* * * *	0	* *	0	* *	0	* *	
Schoolcraf	437	7.8 (85)	3.2 (1)	440	0.0 (1)	437	7.8 (82)	375	42.9 (93)	
Seton Cent	12	* * * *	* * * *	13	* *	12	* *	13	* *	
Shaftsburg	33	* * * *	* * * *	34	* *	32	* *	32	* *	
Shelby M	81	* * * *	* * * *	87	* *	81	* *	87	* *	
Sheldon Co	484	6.4 (47)	6.8 (25)	502	0.0 (1)	483	6.4 (47)	464	48.7 (74)	
Shiawassee	308	5.8 (38)	10.7 (89)	308	0.0 (1)	308	5.8 (37)	289	56.4 (41)	
Southfield	834	8.6 (93)	9.4 (72)	846	0.4 (100)	831	8.5 (91)	822	50.5 (68)	
Sparta Pub	53	* * * *	* * * *	53	* *	53	* *	46	* *	
St Charles	61	* * * *	* * * *	62	* *	61	* *	60	* *	
St Ignace	47	* * * *	* * * *	47	* *	47	* *	46	* *	
St Johns W	90	* * * *	* * * *	107	0.0 (1)	90	* *	106	50.0 (72)	
St Louis C	48	* * * *	* * * *	51	* *	47	* *	50	* *	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
St.Clair S	109	5.5 (32)	13.8 (116)		112	0.0 (1)	109	5.5 (31)	110	97.3 (1)
Stanton Cl	216	5.1 (25)	5.6 (17)		217	0.0 (1)	215	5.1 (23)	203	54.7 (50)
Stockbridg	14	* *	* *		14	* *	14	* *	12	* *
Sumpter He	307	8.5 (92)	8.8 (57)		319	0.6 (110)	301	8.6 (92)	284	43.7 (89)
Taylor	780	7.2 (68)	10.0 (79)		846	0.2 (81)	777	7.1 (67)	787	38.6 (103)
Teen Healt	1	* *	* *		1	* *	1	* *	1	* *
Temperance	5	* *	* *		5	* *	5	* *	4	* *
Thompsonv	8	* *	* *		8	* *	8	* *	7	* *
Three Oaks	43	* *	* *		44	* *	43	* *	40	* *
Three Rive	265	3.8 (10)	7.5 (37)		266	0.4 (100)	263	3.4 (7)	241	43.2 (91)
Traverse C	434	6.5 (49)	12.2 (102)		440	0.5 (106)	431	6.5 (48)	425	76.5 (9)
Tri-Cnty H	151	2.6 (3)	12.6 (105)		152	0.0 (1)	151	2.6 (3)	151	58.9 (33)
Tuscola	373	6.4 (47)	10.5 (87)		382	0.5 (106)	372	6.5 (48)	361	56.0 (44)
U Of M-Ped	0	* *	* *		0	* *	0	* *	0	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)											
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed		
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%	
	Number	% (Rank)	% (Rank)	(Rank)							(Rank)
Vcms River	31	* *	* *		33	* *		31	* *	32	* *
Vicksburg	25	* *	* *		26	* *		25	* *	19	* *
Walled Lak	358	5.3 (28)	8.7 (55)		359	0.0 (1)		356	5.3 (28)	361	56.5 (40)
Warren Cli	171	4.1 (12)	12.9 (106)		178	0.0 (1)		171	4.1 (11)	175	91.4 (5)
Washtenaw	22	* *	* *		25	* *		22	* *	24	* *
Wayne-Dear	569	5.4 (30)	7.9 (47)		649	0.0 (1)		567	5.5 (31)	630	81.1 (6)
Webbervill	12	* *	* *		13	* *		12	* *	11	* *
West Genes	164	6.1 (46)	8.5 (52)		171	0.0 (1)		163	6.1 (44)	141	53.9 (52)
Western Wa	467	7.1 (64)	7.3 (34)		473	0.2 (81)		464	6.9 (61)	450	40.4 (97)
Westland C	501	9.6 (103)	10.2 (83)		513	0.0 (1)		500	9.6 (102)	434	53.7 (54)
Westside C	451	4.9 (20)	10.4 (86)		471	0.0 (1)		450	4.9 (19)	454	58.1 (35)
Westwood	23	* *	* *		24	* *		23	* *	21	* *
Wexford	274	5.1 (25)	10.2 (83)		276	0.4 (100)		272	5.1 (23)	258	63.6 (22)
Wheeler WI	11	* *	* *		11	* *		11	* *	11	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Whitmore L	12	* * * *			13	* *			12	* *
Willow	3	* * * *			4	* *			3	* *
Woodside	53	* * * *			54	* *			53	* *
Wyoming Cl	594	7.1 (64)	10.1 (82)		596	0.2 (81)			591	6.9 (61)
Yale Clini	41	* * * *			41	* *			41	* *
Ypsilanti	728	6.9 (60)	8.0 (48)		765	0.0 (1)			725	6.9 (61)
Michigan	48,280	7.3	8.7		50,986	0.2			48,077	7.3
Nation (Prior Year)	572,813	8.0	7.5		559,297	11.2			478,431	4.0

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Agape WIC	1,371	0.7	98.8	0.1	0.1	0.3	0.0
Albion Cli	579	71.0	25.9	2.9	0.0	0.2	0.0
Alcona	147	98.6	0.0	1.4	0.0	0.0	0.0
Algonac Cl	328	98.2	0.3	0.6	0.9	0.0	0.0
Allegan	307	93.8	2.0	3.9	0.3	0.0	0.0
Alma WIC	202	91.1	0.5	5.4	0.5	2.5	0.0
Alpena Cou	573	99.8	0.0	0.0	0.2	0.0	0.0
Antrim - B	189	95.2	1.1	2.6	1.1	0.0	0.0
Arab/Chald	457	84.2	15.3	0.0	0.0	0.4	0.0
Arenac Cou	281	99.6	0.0	0.0	0.0	0.4	0.0
Associated	28	*	*	*	*	*	*
Baldwin	197	61.9	22.8	13.2	0.0	2.0	0.0
Bangor	1,320	58.6	9.5	31.0	0.5	0.5	0.0
Baraga Cl	123	100.0	0.0	0.0	0.0	0.0	0.0
Barry-Eato	924	98.4	0.3	1.0	0.3	0.0	0.0
Barry-Eato	1,213	97.2	1.8	0.7	0.0	0.2	0.0
Battle Cre	2,705	69.4	20.6	9.3	0.3	0.4	0.0
Baxter Com	65	*	*	*	*	*	*
Bay Cl	2,173	95.7	1.9	2.2	0.0	0.3	0.0
Bayside WI	39	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Beaver Isl	6	*	*	*	*	*	*
Belding Cl	261	96.2	0.4	1.5	0.0	1.9	0.0
Bell Memor	274	99.3	0.4	0.0	0.4	0.0	0.0
Bendle/Bur	121	95.0	1.7	1.7	1.7	0.0	0.0
Benton Har	2,115	37.6	60.3	0.9	0.2	1.0	0.0
Benzie Cli	309	94.8	0.0	4.2	1.0	0.0	0.0
Berrien -	918	82.4	13.8	2.5	0.5	0.8	0.0
Bldhd Migr	17	*	*	*	*	*	*
Booth Clin	12	*	*	*	*	*	*
Branch-Hil	946	93.8	0.6	5.4	0.1	0.1	0.0
Branch-Hil	824	99.2	0.4	0.5	0.0	0.0	0.0
Branch-Hil	859	70.1	1.0	28.9	0.0	0.0	0.0
Browning-C	70	*	*	*	*	*	*
Bryant Com	47	*	*	*	*	*	*
Burton	1,850	83.1	12.6	3.2	0.4	0.5	0.0
Calumet WI	263	100.0	0.0	0.0	0.0	0.0	0.0
Capac Clin	105	94.3	0.0	5.7	0.0	0.0	0.0
Caro WIC S	15	*	*	*	*	*	*
Cedar Spri	552	94.2	1.1	4.0	0.5	0.2	0.0
Charlevoix	472	95.1	1.3	1.5	1.5	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Chass Heal	3,472	9.1	6.5	84.2	0.1	0.1	0.0
Cheboygan	566	98.2	0.5	0.4	0.5	0.4	0.0
Cherry Str	883	12.3	20.4	64.0	0.5	2.8	0.0
Chesaning	95	*	*	*	*	*	*
Clare Coun	666	99.8	0.2	0.0	0.0	0.0	0.0
Clinica Sa	962	0.3	0.3	99.4	0.0	0.0	0.0
Comstock C	0	*	*	*	*	*	*
Corner Hea	612	53.3	44.3	1.5	0.2	0.8	0.0
Crawford	315	99.0	1.0	0.0	0.0	0.0	0.0
Dewitt WIC	173	97.1	2.3	0.6	0.0	0.0	0.0
Dickinson-	241	97.5	1.2	0.4	0.4	0.4	0.0
Dickinson-	561	99.8	0.0	0.2	0.0	0.0	0.0
DMC Prim	26	*	*	*	*	*	*
DMC Prim	134	2.2	97.0	0.7	0.0	0.0	0.0
DMC Prim	41	*	*	*	*	*	*
Dorr Outre	282	98.9	0.4	0.7	0.0	0.0	0.0
Douglas He	2,285	12.2	60.6	24.1	0.2	3.0	0.0
Dowagiac	1,093	85.0	7.5	6.8	0.4	0.4	0.0
Durand WIC	301	98.7	0.0	1.3	0.0	0.0	0.0
East Area	909	1.4	98.0	0.4	0.0	0.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Eastside H	1,655	3.0	95.2	0.5	0.3	1.0	0.0
Eau Claire	615	3.3	1.1	95.6	0.0	0.0	0.0
Emmet Coun	674	93.6	0.7	1.2	3.9	0.6	0.0
Escanaba C	852	99.2	0.1	0.2	0.1	0.4	0.0
Family Com	704	25.3	66.9	3.6	0.1	4.1	0.0
Family Hlt	688	50.3	32.4	15.7	0.3	1.3	0.0
Family Roa	973	6.0	89.4	2.9	0.1	1.6	0.0
Farmington	142	74.6	21.1	2.1	0.0	2.1	0.0
Fife Lake	2	*	*	*	*	*	*
First Pres	0	*	*	*	*	*	*
Flint Fam	876	39.0	57.2	3.3	0.1	0.3	0.0
Fruitport	21	*	*	*	*	*	*
Gateway	1,079	0.7	98.3	0.4	0.2	0.4	0.0
Genesee -	3,320	62.0	31.3	5.8	0.2	0.7	0.0
Gladwin Co	479	99.6	0.0	0.2	0.2	0.0	0.0
Gogebic Cl	268	98.1	0.4	0.7	0.7	0.0	0.0
Golden	104	1.0	98.1	0.0	0.0	1.0	0.0
Grace Prof	369	0.8	98.9	0.3	0.0	0.0	0.0
Grace Ross	2,614	3.4	96.1	0.3	0.1	0.1	0.0
Grand Have	740	92.8	0.5	6.1	0.4	0.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Grand Trav	13	*	*	*	*	*	*
Grant	345	71.6	0.3	28.1	0.0	0.0	0.0
Gratiot	1,837	3.1	90.4	0.5	0.1	5.9	0.0
Greenville	373	98.1	0.3	1.3	0.0	0.3	0.0
Hamtramck	440	68.9	26.8	1.8	0.2	2.3	0.0
Hancock Cl	467	94.2	1.5	0.2	0.0	4.1	0.0
Hannahvil	47	*	*	*	*	*	*
Harper Gra	3,330	1.9	97.3	0.6	0.2	0.1	0.0
Haslett Cl	154	81.8	7.1	7.1	0.0	3.9	0.0
Haslett St	29	*	*	*	*	*	*
Herman Kie	3,034	2.1	96.3	0.5	0.1	0.9	0.0
Highland P	333	5.7	94.0	0.3	0.0	0.0	0.0
Holland Cl	2,653	46.4	3.0	44.3	0.1	6.2	0.0
Holly WIC	100	95.0	4.0	1.0	0.0	0.0	0.0
Holton	82	*	*	*	*	*	*
Howard Cit	162	98.8	0.6	0.6	0.0	0.0	0.0
Hudsonv	571	93.9	1.4	3.7	0.0	1.1	0.0
Hurley Cli	567	34.4	60.7	4.4	0.5	0.0	0.0
Huron Coun	681	96.2	0.3	3.2	0.0	0.3	0.0
Icchn-Spar	331	0.6	0.0	99.4	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Ingham Cou	4,326	60.6	24.4	11.6	0.2	3.2	0.0
Intervale	269	0.4	98.9	0.7	0.0	0.0	0.0
Ionia Coun	842	92.2	0.6	6.9	0.0	0.4	0.0
Iosco	512	99.2	0.6	0.0	0.2	0.0	0.0
Ironwood	0	*	*	*	*	*	*
Isabella C	962	93.6	3.0	1.7	0.7	1.0	0.0
Ithaca Cli	381	93.7	1.0	5.2	0.0	0.0	0.0
Ithaca WIC	0	*	*	*	*	*	*
Jackson Co	3,529	80.3	13.7	5.3	0.5	0.2	0.0
Jackson M	0	*	*	*	*	*	*
Janes St A	1,166	13.6	69.8	16.1	0.1	0.3	0.0
Joy-Greenf	2,288	34.5	62.1	3.3	0.0	0.1	0.0
Kalamazoo	1,022	26.8	58.0	13.8	0.5	0.9	0.0
Kalamazoo	2,543	81.1	15.4	2.4	0.1	1.0	0.0
Kalkaska	448	99.6	0.0	0.2	0.2	0.0	0.0
Kent	0	*	*	*	*	*	*
Kent Count	3,956	60.2	25.5	10.8	0.9	2.6	0.0
Kentwood	1,854	54.4	26.5	9.3	0.3	9.5	0.0
Keweenaw	75	*	*	*	*	*	*
Kingsley	40	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Lac Vieux	25	*	*	*	*	*	*
Lake Co	242	76.4	16.1	7.0	0.0	0.4	0.0
Lake Odess	65	*	*	*	*	*	*
Lapeer	1,501	97.8	0.7	1.4	0.0	0.1	0.0
Leelanu Cl	182	58.8	0.5	14.3	25.8	0.5	0.0
Lenawee	0	*	*	*	*	*	*
Leslie-Sto	81	*	*	*	*	*	*
Lincoln Pa	1,096	74.4	21.8	3.7	0.0	0.1	0.0
Livingston	1,030	96.5	0.7	1.9	0.1	0.8	0.0
Lowell	139	93.5	0.0	5.0	1.4	0.0	0.0
M-72 Migra	34	*	*	*	*	*	*
Mancelona	231	99.1	0.4	0.0	0.4	0.0	0.0
Manchester	27	*	*	*	*	*	*
Manchester	8	*	*	*	*	*	*
Manistee C	433	90.1	0.0	4.6	3.7	1.6	0.0
Manistique	197	91.9	0.5	0.0	7.6	0.0	0.0
Mansitee M	48	*	*	*	*	*	*
Marion Bra	116	100.0	0.0	0.0	0.0	0.0	0.0
Marquette	438	97.3	0.9	0.7	0.9	0.2	0.0
Marshall C	312	98.1	1.3	0.6	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Mason Clin	112	100.0	0.0	0.0	0.0	0.0	0.0
Mason Co	580	87.1	0.7	10.9	0.9	0.5	0.0
Mccree Hea	3,512	35.4	62.3	1.8	0.0	0.4	0.0
Mcnamara H	176	95.5	0.6	3.4	0.0	0.6	0.0
Mecosta Co	915	92.9	5.5	0.8	0.1	0.8	0.0
Medical Pr	40	*	*	*	*	*	*
Medical Pr	87	*	*	*	*	*	*
Menominee	424	99.8	0.2	0.0	0.0	0.0	0.0
Mercy	120	0.0	100.0	0.0	0.0	0.0	0.0
Mesick	69	*	*	*	*	*	*
Metropolit	2	*	*	*	*	*	*
Michigan S	233	28.8	18.9	18.0	0.9	33.5	0.0
Midland Cl	1,264	97.6	1.3	0.7	0.1	0.3	0.0
Midwest	1,191	20.8	38.5	38.8	0.3	1.5	0.0
Migrant H	262	3.1	0.4	96.6	0.0	0.0	0.0
Missaukee	225	98.2	0.4	0.4	0.9	0.0	0.0
Moms & Bab	1,076	6.9	59.9	32.2	0.3	0.8	0.0
Monroe Cou	2,197	92.9	4.2	2.5	0.0	0.2	0.0
Montague/	100	92.0	3.0	2.0	2.0	1.0	0.0
Montcalm	25	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Montmorenc	182	98.4	0.0	0.5	0.0	1.1	0.0
Mt Clemens	2,663	78.9	14.7	4.5	0.1	1.8	0.0
Munising C	165	97.6	0.6	0.0	1.8	0.0	0.0
Muskegon C	4,545	63.7	28.2	7.2	0.6	0.4	0.0
New Center	24	*	*	*	*	*	*
New Haven	278	64.7	4.7	29.9	0.4	0.4	0.0
New Hope	13	*	*	*	*	*	*
Newaygo	781	96.8	0.4	2.8	0.0	0.0	0.0
Newberry C	180	95.6	0.6	0.6	3.3	0.0	0.0
North Camp	360	36.1	6.9	7.8	0.0	49.2	0.0
North Midl	0	*	*	*	*	*	*
Northeast	2,938	3.2	88.7	0.6	0.1	7.2	0.0
Northern M	134	93.3	0.7	0.0	5.2	0.7	0.0
Northwest	510	1.0	99.0	0.0	0.0	0.0	0.0
Oakland Co	4,556	51.5	26.8	19.4	0.1	2.2	0.0
Oceana Co	566	88.9	0.2	10.8	0.2	0.0	0.0
Ogemaw	501	98.8	0.6	0.6	0.0	0.0	0.0
Olhsa Star	373	81.2	13.9	1.1	0.5	3.2	0.0
Olhsa WIC	138	40.6	45.7	11.6	0.7	1.4	0.0
Omni Care	38	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Omicare	4	*	*	*	*	*	*
Omicare-G	0	*	*	*	*	*	*
Omicare-M	88	*	*	*	*	*	*
Ontonagon	97	*	*	*	*	*	*
Oscoda	176	96.0	0.0	2.8	0.6	0.6	0.0
Otsego Co	599	99.3	0.0	0.0	0.2	0.5	0.0
Park Clin	2	*	*	*	*	*	*
Park Medic	33	*	*	*	*	*	*
Park Medic	57	*	*	*	*	*	*
Park Medic	11	*	*	*	*	*	*
Paw Paw Cl	428	90.9	1.9	6.5	0.2	0.5	0.0
PCOC	127	88.2	8.7	2.4	0.0	0.8	0.0
Pinconning	118	99.2	0.0	0.0	0.0	0.8	0.0
Port Huron	2,365	89.1	6.2	3.3	0.9	0.5	0.0
Portland	94	*	*	*	*	*	*
Presque Is	206	100.0	0.0	0.0	0.0	0.0	0.0
Profession	77	*	*	*	*	*	*
Pullman	742	57.1	3.0	39.8	0.0	0.1	0.0
Ravenna	33	*	*	*	*	*	*
Reed City	447	99.6	0.2	0.0	0.2	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Region II	1,779	78.5	1.7	19.6	0.0	0.2	0.0
Riverview	90	*	*	*	*	*	*
Romulus Fi	287	67.9	29.3	1.0	0.0	1.7	0.0
Roosevelt-	154	20.8	61.7	16.9	0.6	0.0	0.0
Roscommon	440	98.0	0.0	0.5	0.2	1.4	0.0
S & S Medi	31	*	*	*	*	*	*
Saginaw Co	3,565	56.0	34.8	9.1	0.0	0.1	0.0
Saginaw-Ch	56	*	*	*	*	*	*
Sanilac Co	858	98.5	0.6	0.8	0.0	0.1	0.0
Sault Ste	655	68.2	1.1	1.8	28.2	0.6	0.0
Sawyer Afb	181	95.0	1.7	1.1	2.2	0.0	0.0
School Bas	0	*	*	*	*	*	*
Schoolcraf	1,512	1.1	97.4	0.9	0.3	0.3	0.0
Seton Cent	881	0.6	98.8	0.5	0.2	0.0	0.0
Shaftsborg	83	*	*	*	*	*	*
Shelby M	369	1.4	0.0	98.6	0.0	0.0	0.0
Sheldon Co	1,906	9.5	52.3	37.4	0.0	0.9	0.0
Shiawassee	1,131	98.2	0.4	1.0	0.3	0.2	0.0
Southfield	2,795	55.9	40.1	2.0	0.0	2.0	0.0
Sparta Pub	332	87.7	1.5	9.0	0.3	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
St Charles	164	97.6	1.2	1.2	0.0	0.0	0.0
St Ignace	176	74.4	1.1	1.7	22.7	0.0	0.0
St Johns W	391	97.7	1.3	1.0	0.0	0.0	0.0
St Louis C	206	93.7	1.9	4.4	0.0	0.0	0.0
St.Clair S	1,087	89.1	7.8	1.3	0.1	1.7	0.0
Stanton Cl	734	98.4	0.5	1.0	0.1	0.0	0.0
Stockbridg	67	*	*	*	*	*	*
Sumpter He	1,020	76.3	22.4	1.2	0.0	0.2	0.0
Taylor	2,940	84.3	15.2	0.5	0.0	0.0	0.0
Teen Healt	13	*	*	*	*	*	*
Temperance	77	*	*	*	*	*	*
Thompsonv	13	*	*	*	*	*	*
Three Oaks	137	92.7	4.4	1.5	0.0	1.5	0.0
Three Rive	849	91.8	6.7	1.2	0.0	0.4	0.0
Traverse C	1,428	91.1	1.3	4.5	1.7	1.5	0.0
Tri-Cnty H	556	98.2	0.9	0.5	0.2	0.2	0.0
Tuscola	1,242	99.3	0.4	0.2	0.0	0.1	0.0
U Of M-Ped	0	*	*	*	*	*	*
Vcms River	107	0.0	100.0	0.0	0.0	0.0	0.0
Vicksburg	112	99.1	0.0	0.9	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaskan Native	Asian/Pacific Islander	All Other/Unknown
		%	%	%	%	%	%
Walled Lak	1,035	85.4	5.2	7.2	0.3	1.8	0.0
Warren Cli	1,663	81.7	11.1	1.3	0.3	5.6	0.0
Washtenaw	91	*	*	*	*	*	*
Wayne-Dear	2,517	98.6	0.6	0.6	0.0	0.2	0.0
Webbervill	51	*	*	*	*	*	*
West Genes	603	29.5	48.1	17.7	0.0	4.6	0.0
Western Wa	1,239	81.6	16.6	1.4	0.0	0.4	0.0
Westland C	2,029	65.2	31.7	2.5	0.2	0.4	0.0
Westside C	1,604	46.6	4.3	47.4	0.8	0.9	0.0
Westwood	64	*	*	*	*	*	*
Wexford	843	99.1	0.2	0.2	0.2	0.2	0.0
Wheeler WI	35	*	*	*	*	*	*
Whitmore L	52	*	*	*	*	*	*
Willow	40	*	*	*	*	*	*
Woodside	185	67.0	1.1	0.5	29.7	1.6	0.0
Wyoming Cl	1,628	76.8	8.0	11.6	0.6	3.0	0.0
Yale Clini	111	97.3	0.9	1.8	0.0	0.0	0.0
Ypsilanti	2,906	47.7	41.7	7.6	0.1	2.9	0.0
Michigan	181,396	59.4	28.7	10.0	0.5	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Agape WIC	1,371	0.6	5.0	10.6	54.9	27.2	1.5	0.2
Albion Cli	579	0.9	11.7	17.8	55.8	13.0	0.9	0.0
Alcona	147	0.0	4.8	12.2	53.1	28.6	1.4	0.0
Algonac Cl	328	0.0	7.9	13.4	58.5	18.9	1.2	0.0
Allegan	307	0.0	8.8	13.4	63.2	13.7	1.0	0.0
Alma WIC	202	0.0	5.9	14.9	67.3	10.4	1.5	0.0
Alpena Cou	573	0.2	4.5	14.3	65.3	15.0	0.7	0.0
Antrim - B	189	0.5	6.9	13.2	61.4	16.4	1.6	0.0
Arab/Chald	457	0.2	0.4	5.7	54.9	33.9	4.8	0.0
Arenac Cou	281	0.4	5.7	17.8	61.2	13.5	1.4	0.0
Associated	28	*	*	*	*	*	*	*
Baldwin	197	0.0	3.6	14.7	62.4	18.8	0.5	0.0
Bangor	1,320	0.5	8.6	17.3	56.7	15.2	1.6	0.0
Baraga Cl	123	1.6	4.9	18.7	48.8	22.8	3.3	0.0
Barry-Eato	924	0.2	9.1	13.7	60.0	15.7	1.3	0.0
Barry-Eato	1,213	0.2	7.2	17.1	60.5	13.9	0.8	0.2
Battle Cre	2,705	0.9	9.2	14.0	62.8	12.2	0.9	0.0
Baxter Com	65	*	*	*	*	*	*	*
Bay Cl	2,173	0.1	7.7	12.8	62.7	15.3	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Bayside WI	39	*	*	*	*	*	*	*
Beaver Isl	6	*	*	*	*	*	*	*
Belding Cl	261	0.0	6.1	22.2	60.9	10.3	0.4	0.0
Bell Memor	274	0.4	5.8	16.4	54.7	20.8	1.8	0.0
Bendle/Bur	121	0.0	1.7	17.4	67.8	13.2	0.0	0.0
Benton Har	2,115	1.3	11.1	16.3	56.6	13.6	0.9	0.1
Benzie Cli	309	0.6	8.7	16.5	55.3	16.8	1.9	0.0
Berrien -	918	0.3	9.4	15.8	59.4	15.0	0.1	0.0
Bldhd Migr	17	*	*	*	*	*	*	*
Booth Clin	12	*	*	*	*	*	*	*
Branch-Hil	946	0.8	8.5	17.9	59.3	13.0	0.5	0.0
Branch-Hil	824	0.2	7.4	19.3	61.3	10.7	1.1	0.0
Branch-Hil	859	0.3	7.0	15.0	60.8	15.4	1.5	0.0
Browning-C	70	*	*	*	*	*	*	*
Bryant Com	47	*	*	*	*	*	*	*
Burton	1,850	0.4	5.2	11.9	64.9	16.8	0.7	0.0
Calumet WI	263	0.0	4.6	9.5	58.2	24.3	3.4	0.0
Capac Clin	105	1.0	4.8	21.9	61.0	11.4	0.0	0.0
Caro WIC S	15	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cedar Spri	552	0.0	8.7	16.3	58.0	15.9	1.1	0.0
Charlevoix	472	0.2	7.6	13.8	62.1	15.5	0.8	0.0
Chass Heal	3,472	0.4	6.5	10.5	61.1	20.0	1.6	0.0
Cheboygan	566	0.0	6.7	14.0	61.1	17.3	0.9	0.0
Cherry Str	883	1.0	7.2	11.8	61.5	17.0	1.5	0.0
Chesaning	95	*	*	*	*	*	*	*
Clare Coun	666	0.8	8.4	17.0	60.7	12.5	0.8	0.0
Clinica Sa	962	0.1	7.6	11.0	63.3	16.8	1.1	0.0
Comstock C	0	*	*	*	*	*	*	*
Corner Hea	612	1.0	20.4	42.6	35.9	0.0	0.0	0.0
Crawford	315	0.3	10.5	16.5	59.0	12.7	1.0	0.0
Dewitt WIC	173	0.6	6.9	14.5	58.4	19.7	0.0	0.0
Dickinson-	241	0.4	6.2	11.2	61.4	18.7	2.1	0.0
Dickinson-	561	0.2	7.1	15.2	60.4	16.2	0.9	0.0
DMC Prim	26	*	*	*	*	*	*	*
DMC Prim	134	0.0	6.0	14.2	55.2	23.1	1.5	0.0
DMC Prim	41	*	*	*	*	*	*	*
Dorr Outre	282	0.0	9.9	17.7	59.2	12.8	0.4	0.0
Douglas He	2,285	0.6	6.7	10.9	60.8	19.8	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dowagiac	1,093	0.1	7.0	15.5	63.2	12.7	1.5	0.0
Durand WIC	301	0.0	6.0	15.9	60.5	15.6	2.0	0.0
East Area	909	1.7	14.0	14.3	51.2	17.7	1.1	0.1
Eastside H	1,655	0.4	8.8	13.7	58.5	17.2	1.4	0.0
Eau Claire	615	0.7	6.8	12.4	62.4	16.7	1.0	0.0
Emmet Coun	674	0.3	6.4	12.9	59.8	19.1	1.5	0.0
Escanaba C	852	0.0	5.2	12.9	61.6	19.2	1.1	0.0
Family Com	704	0.4	6.0	8.0	62.5	21.6	1.6	0.0
Family Hlt	688	0.9	9.0	15.6	57.0	16.9	0.7	0.0
Family Roa	973	0.7	6.7	12.2	58.2	21.3	0.9	0.0
Farmington	142	0.0	4.2	5.6	65.5	21.8	2.8	0.0
Fife Lake	2	*	*	*	*	*	*	*
First Pres	0	*	*	*	*	*	*	*
Flint Fam	876	2.1	10.4	13.6	56.8	15.2	1.9	0.0
Fruitport	21	*	*	*	*	*	*	*
Gateway	1,079	1.8	11.6	13.5	53.8	17.6	1.7	0.0
Genesee -	3,320	0.5	7.4	15.8	60.1	15.2	1.0	0.0
Gladwin Co	479	0.2	7.7	16.5	57.4	17.5	0.6	0.0
Gogebic Cl	268	0.4	6.3	11.2	61.9	19.4	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Golden	104	1.9	5.8	12.5	54.8	23.1	1.9	0.0
Grace Prof	369	1.9	11.9	10.3	52.8	22.5	0.5	0.0
Grace Ross	2,614	0.7	6.5	14.3	55.7	21.6	1.1	0.0
Grand Have	740	0.1	5.5	13.1	63.5	16.6	1.1	0.0
Grand Trav	13	*	*	*	*	*	*	*
Grant	345	0.3	12.2	18.0	51.9	16.8	0.9	0.0
Gratiot	1,837	0.9	10.7	14.3	52.5	20.4	1.2	0.0
Greenville	373	0.3	8.0	17.2	63.0	10.7	0.8	0.0
Hamtramck	440	0.0	2.7	9.8	58.9	26.1	2.5	0.0
Hancock Cl	467	0.4	3.0	9.6	64.5	20.3	2.1	0.0
Hannahvil	47	*	*	*	*	*	*	*
Harper Gra	3,330	0.9	8.7	13.8	56.8	18.4	1.4	0.0
Haslett Cl	154	0.0	2.6	11.0	64.3	21.4	0.6	0.0
Haslett St	29	*	*	*	*	*	*	*
Herman Kie	3,034	0.7	6.2	11.6	56.6	23.4	1.5	0.0
Highland P	333	0.6	5.7	14.4	58.6	18.3	1.8	0.6
Holland Cl	2,653	0.6	8.8	14.6	61.4	13.9	0.7	0.0
Holly WIC	100	0.0	7.0	17.0	45.0	30.0	1.0	0.0
Holton	82	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Howard Cit	162	0.6	11.7	10.5	61.1	13.6	2.5	0.0
Hudsonv	571	0.5	7.0	12.8	57.8	20.8	1.1	0.0
Hurley Cli	567	0.9	9.5	14.5	57.3	16.2	1.6	0.0
Huron Coun	681	0.0	4.8	10.4	61.8	22.2	0.7	0.0
Icchn-Spar	331	0.6	10.0	8.8	60.4	19.0	1.2	0.0
Ingham Cou	4,326	0.3	7.5	14.0	62.6	14.8	0.9	0.0
Intervale	269	0.7	9.3	12.3	53.5	22.3	1.9	0.0
Ionia Coun	842	0.2	8.9	17.2	61.3	11.6	0.7	0.0
Iosco	512	0.4	7.6	16.6	59.4	14.6	1.4	0.0
Ironwood	0	*	*	*	*	*	*	*
Isabella C	962	0.2	4.6	16.0	65.3	13.5	0.4	0.0
Ithaca Cli	381	0.0	2.6	15.0	67.7	14.2	0.5	0.0
Ithaca WIC	0	*	*	*	*	*	*	*
Jackson Co	3,529	0.7	10.0	16.5	60.2	11.8	0.8	0.0
Jackson M	0	*	*	*	*	*	*	*
Janes St A	1,166	1.5	11.8	13.2	59.1	13.2	1.2	0.0
Joy-Greenf	2,288	0.3	4.5	11.6	59.2	22.7	1.6	0.0
Kalamazoo	1,022	0.4	11.0	14.5	58.8	14.0	1.4	0.0
Kalamazoo	2,543	0.4	6.7	13.3	62.7	15.8	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Kalkaska	448	0.7	6.0	13.2	64.1	13.6	2.5	0.0
Kent	0	*	*	*	*	*	*	*
Kent Count	3,956	0.4	7.0	13.2	61.5	16.5	1.5	0.0
Kentwood	1,854	0.4	4.7	11.7	62.7	19.6	1.0	0.0
Keweenaw	75	*	*	*	*	*	*	*
Kingsley	40	*	*	*	*	*	*	*
Lac Vieux	25	*	*	*	*	*	*	*
Lake Co	242	0.4	8.3	14.5	63.6	12.8	0.4	0.0
Lake Odess	65	*	*	*	*	*	*	*
Lapeer	1,501	0.1	6.3	16.4	58.8	17.2	1.3	0.0
Leelanu Cl	182	0.5	8.2	10.4	59.9	19.8	1.1	0.0
Lenawee	0	*	*	*	*	*	*	*
Leslie-Sto	81	*	*	*	*	*	*	*
Lincoln Pa	1,096	0.6	7.7	16.3	59.6	15.0	0.8	0.0
Livingston	1,030	0.1	5.9	15.9	58.4	18.3	1.3	0.0
Lowell	139	0.7	6.5	13.7	55.4	22.3	1.4	0.0
M-72 Migra	34	*	*	*	*	*	*	*
Mancelona	231	0.4	7.8	22.5	58.9	9.5	0.9	0.0
Manchester	27	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Manchester	8	*	*	*	*	*	*	*
Manistee C	433	1.2	5.3	14.5	60.3	17.3	1.4	0.0
Manistique	197	0.0	8.1	12.2	53.3	25.4	1.0	0.0
Mansitee M	48	*	*	*	*	*	*	*
Marion Bra	116	0.0	7.8	16.4	66.4	9.5	0.0	0.0
Marquette	438	0.0	4.6	12.3	66.4	15.3	1.4	0.0
Marshall C	312	0.0	7.7	11.5	67.0	12.8	1.0	0.0
Mason Clin	112	0.9	7.1	13.4	58.0	19.6	0.9	0.0
Mason Co	580	0.9	8.4	13.3	61.7	14.7	1.0	0.0
Mccree Hea	3,512	0.5	8.0	14.3	62.4	14.0	0.8	0.0
Mcnamara H	176	0.0	4.0	13.1	61.9	20.5	0.6	0.0
Mecosta Co	915	0.3	7.3	13.6	62.6	15.6	0.4	0.1
Medical Pr	40	*	*	*	*	*	*	*
Medical Pr	87	*	*	*	*	*	*	*
Menominee	424	0.0	9.0	17.7	55.0	17.2	1.2	0.0
Mercy	120	0.8	5.8	11.7	58.3	23.3	0.0	0.0
Mesick	69	*	*	*	*	*	*	*
Metropolit	2	*	*	*	*	*	*	*
Michigan S	233	0.0	0.9	2.1	57.1	39.1	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Midland Cl	1,264	0.5	5.7	15.7	62.9	14.2	1.1	0.0
Midwest	1,191	0.7	6.9	10.6	58.8	21.7	1.4	0.0
Migrant H	262	0.4	6.5	11.8	61.1	17.9	2.3	0.0
Missaukee	225	0.0	6.2	15.1	66.7	11.6	0.4	0.0
Moms & Bab	1,076	3.4	22.7	10.4	45.2	17.3	1.0	0.0
Monroe Cou	2,197	0.2	7.7	16.2	60.3	14.7	0.9	0.0
Montague/	100	0.0	11.0	21.0	50.0	18.0	0.0	0.0
Montcalm	25	*	*	*	*	*	*	*
Montmorenc	182	0.0	6.6	15.9	54.9	22.0	0.5	0.0
Mt Clemens	2,663	0.3	4.2	12.6	62.5	19.1	1.4	0.0
Munising C	165	0.0	4.8	13.9	59.4	20.0	1.8	0.0
Muskegon C	4,545	0.5	8.7	15.8	60.1	14.3	0.7	0.0
New Center	24	*	*	*	*	*	*	*
New Haven	278	0.0	4.7	11.5	65.8	16.2	1.8	0.0
New Hope	13	*	*	*	*	*	*	*
Newaygo	781	0.5	8.6	16.1	60.7	12.4	1.7	0.0
Newberry C	180	0.0	7.2	16.1	62.8	12.8	1.1	0.0
North Camp	360	0.0	0.0	2.5	59.4	37.2	0.8	0.0
North Midl	0	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Northeast	2,938	0.6	7.8	12.3	57.0	20.8	1.5	0.0
Northern M	134	0.7	3.7	9.0	69.4	17.2	0.0	0.0
Northwest	510	0.4	13.1	10.6	54.1	20.4	1.4	0.0
Oakland Co	4,556	0.4	6.9	13.9	60.5	17.3	0.9	0.0
Oceana Co	566	0.0	8.0	14.0	61.7	15.0	1.4	0.0
Ogemaw	501	0.0	6.0	14.2	62.9	16.4	0.6	0.0
Olhse Star	373	0.3	2.9	9.1	60.6	24.9	2.1	0.0
Olhse WIC	138	0.0	10.9	13.0	58.7	16.7	0.7	0.0
Omni Care	38	*	*	*	*	*	*	*
Omnicare	4	*	*	*	*	*	*	*
Omnicare-G	0	*	*	*	*	*	*	*
Omnicare-M	88	*	*	*	*	*	*	*
Ontonagon	97	*	*	*	*	*	*	*
Oscoda	176	0.0	16.5	17.0	51.7	13.1	1.7	0.0
Otsego Co	599	0.3	4.8	14.7	61.6	17.7	0.8	0.0
Park Clin	2	*	*	*	*	*	*	*
Park Medic	33	*	*	*	*	*	*	*
Park Medic	57	*	*	*	*	*	*	*
Park Medic	11	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Paw Paw Cl	428	0.2	8.6	15.9	61.9	12.6	0.7	0.0
PCOC	127	0.0	2.4	12.6	67.7	17.3	0.0	0.0
Pinconning	118	0.0	5.9	11.9	62.7	19.5	0.0	0.0
Port Huron	2,365	0.5	7.9	15.9	60.3	14.9	0.6	0.0
Portland	94	*	*	*	*	*	*	*
Presque Is	206	0.5	5.8	9.2	66.0	17.5	1.0	0.0
Profession	77	*	*	*	*	*	*	*
Pullman	742	0.8	9.7	15.1	58.5	14.4	1.5	0.0
Ravenna	33	*	*	*	*	*	*	*
Reed City	447	0.2	8.3	15.9	59.5	15.4	0.7	0.0
Region II	1,779	0.3	8.3	15.1	61.0	13.8	1.5	0.0
Riverview	90	*	*	*	*	*	*	*
Romulus Fi	287	0.3	5.6	15.0	64.1	14.6	0.3	0.0
Roosevelt-	154	2.6	13.6	11.7	53.9	17.5	0.6	0.0
Roscommon	440	0.0	6.6	15.0	58.2	18.6	1.6	0.0
S & S Medi	31	*	*	*	*	*	*	*
Saginaw Co	3,565	0.6	8.4	13.3	58.2	18.3	1.1	0.0
Saginaw-Ch	56	*	*	*	*	*	*	*
Sanilac Co	858	0.2	6.3	12.5	63.6	15.9	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Sault Ste	655	0.0	6.4	15.9	60.0	16.9	0.8	0.0
Sawyer Afb	181	1.7	3.9	10.5	66.3	16.0	1.7	0.0
School Bas	0	*	*	*	*	*	*	*
Schoolcraf	1,512	0.4	6.3	12.4	60.1	19.6	1.3	0.0
Seton Cent	881	0.6	7.4	11.8	56.5	21.8	1.9	0.0
Shaftsburg	83	*	*	*	*	*	*	*
Shelby M	369	0.0	9.8	15.2	57.2	15.4	2.4	0.0
Sheldon Co	1,906	1.2	12.4	14.5	56.4	14.7	0.8	0.0
Shiawassee	1,131	0.4	8.3	15.0	60.6	15.1	0.6	0.0
Southfield	2,795	0.1	3.9	10.6	56.9	26.4	2.2	0.0
Sparta Pub	332	0.0	8.4	17.2	59.6	14.2	0.6	0.0
St Charles	164	1.8	7.3	11.0	53.0	24.4	2.4	0.0
St Ignace	176	0.6	5.1	10.8	70.5	11.9	1.1	0.0
St Johns W	391	0.0	6.4	17.1	65.5	10.2	0.8	0.0
St Louis C	206	1.5	6.3	19.4	59.7	12.1	1.0	0.0
St.Clair S	1,087	0.1	5.4	14.3	59.2	20.0	1.1	0.0
Stanton Cl	734	0.0	9.3	17.7	61.2	10.8	1.1	0.0
Stockbridg	67	*	*	*	*	*	*	*
Sumpter He	1,020	0.5	4.7	12.7	63.2	17.5	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Taylor	2,940	0.2	6.2	14.5	63.3	14.8	0.9	0.0
Teen Healt	13	*	*	*	*	*	*	*
Temperance	77	*	*	*	*	*	*	*
Thompsonv	13	*	*	*	*	*	*	*
Three Oaks	137	0.0	8.0	18.2	57.7	13.9	2.2	0.0
Three Rive	849	0.6	9.2	18.1	59.0	12.0	1.1	0.0
Traverse C	1,428	0.3	6.1	16.0	59.7	16.2	1.7	0.0
Tri-Cnty H	556	0.2	6.8	12.9	64.0	15.6	0.4	0.0
Tuscola	1,242	0.1	6.7	15.3	61.5	15.9	0.6	0.0
U Of M-Ped	0	*	*	*	*	*	*	*
Vcms River	107	2.8	12.1	12.1	57.0	15.0	0.9	0.0
Vicksburg	112	0.0	8.0	17.9	51.8	21.4	0.9	0.0
Walled Lak	1,035	0.2	4.6	13.9	60.7	19.1	1.4	0.0
Warren Cli	1,663	0.2	5.1	11.1	62.5	19.6	1.6	0.0
Washtenaw	91	*	*	*	*	*	*	*
Wayne-Dear	2,517	0.1	2.3	9.3	59.4	26.0	2.9	0.0
Webbervill	51	*	*	*	*	*	*	*
West Genes	603	0.5	9.8	11.1	60.7	16.4	1.5	0.0
Western Wa	1,239	0.4	4.8	10.5	64.2	19.0	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Westland C	2,029	0.3	6.4	13.4	59.8	19.7	0.5	0.0
Westside C	1,604	0.5	9.6	14.6	57.9	16.2	1.2	0.0
Westwood	64	*	*	*	*	*	*	*
Wexford	843	0.5	8.2	17.0	59.3	14.4	0.7	0.0
Wheeler WI	35	*	*	*	*	*	*	*
Whitmore L	52	*	*	*	*	*	*	*
Willow	40	*	*	*	*	*	*	*
Woodside	185	0.5	8.1	17.8	57.8	15.7	0.0	0.0
Wyoming Cl	1,628	0.3	6.9	12.7	61.9	17.0	1.0	0.1
Yale Clini	111	0.9	8.1	16.2	54.1	20.7	0.0	0.0
Ypsilanti	2,906	0.2	3.2	7.9	68.0	19.5	1.2	0.0
Michigan	181,396	0.5	7.4	13.8	60.0	17.2	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Agape WIC	1,306	7.8 (8)	50.4 (169)	1,132	32.2 (109)	44.6 (97)	299	36.1 (60)	220	43.2 (69)	
Albion Cli	547	15.5 (164)	44.4 (100)	472	26.5 (71)	41.1 (67)	90	* *	100	45.0 (74)	
Alcona	140	12.1 (96)	47.9 (146)	123	19.5 (10)	52.0 (165)	14	* *	39	* *	
Algonac Cl	309	17.2 (175)	37.2 (23)	269	17.8 (5)	51.3 (160)	48	* *	174	33.9 (47)	
Allegan	273	11.4 (68)	41.8 (60)	232	40.1 (141)	37.9 (46)	45	* *	49	* *	
Alma WIC	163	17.8 (177)	32.5 (6)	143	31.5 (105)	39.2 (56)	27	* *	49	* *	
Alpena Cou	548	11.5 (71)	43.6 (89)	493	20.5 (18)	50.3 (156)	72	* *	147	17.7 (9)	
Antrim - B	170	9.4 (28)	40.6 (44)	146	37.0 (133)	27.4 (15)	19	* *	51	* *	
Arab/Chald	433	7.9 (10)	39.3 (34)	371	66.0 (169)	14.8 (5)	62	* *	42	* *	
Arenac Cou	259	17.8 (177)	49.8 (164)	101	44.6 (154)	33.7 (33)	35	* *	39	* *	
Associated	27	* *	* *	25	* *	* *	7	* *	7	* *	
Baldwin	174	12.1 (96)	36.2 (19)	129	45.0 (155)	33.3 (29)	40	* *	48	* *	
Bangor	1,158	11.6 (77)	45.1 (112)	975	34.1 (117)	40.1 (62)	175	24.0 (23)	213	54.5 (92)	
Baraga Cl	114	13.2 (122)	33.3 (10)	95	* *	* *	10	* *	18	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)										
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barry-Eato	866	14.1 (141)	42.4 (74)	717	36.8 (131)	38.4 (53)	120	29.2 (35)	221	51.1 (89)
Barry-Eato	1,132	12.2 (100)	40.7 (46)	957	25.6 (64)	48.0 (132)	204	17.2 (13)	235	28.5 (28)
Battle Cre	2,516	12.2 (100)	43.5 (88)	2,156	22.9 (42)	49.0 (141)	473	30.2 (39)	392	46.4 (78)
Baxter Com	61	* *	* *	45	* *	* *	12	* *	7	* *
Bay Cl	1,760	13.4 (125)	46.1 (125)	1,473	26.6 (74)	43.9 (92)	274	15.3 (9)	358	22.9 (18)
Bayside WI	28	* *	* *	15	* *	* *	5	* *	6	* *
Beaver Isl	5	* *	* *	5	* *	* *	1	* *	3	* *
Belding Cl	254	13.0 (117)	45.3 (115)	213	21.1 (26)	46.9 (118)	48	* *	63	* *
Bell Memor	259	12.0 (94)	44.0 (97)	217	24.4 (58)	46.1 (111)	32	* *	41	* *
Bendle/Bur	117	21.4 (181)	43.6 (89)	94	* *	* *	24	* *	19	* *
Benton Har	2,010	11.3 (63)	48.5 (151)	1,693	26.0 (68)	45.8 (107)	384	51.6 (77)	564	48.0 (84)
Benzie Cli	291	12.0 (94)	40.9 (49)	250	28.0 (86)	41.6 (73)	34	* *	68	* *
Berrien -	856	15.5 (164)	40.7 (46)	733	21.6 (31)	49.2 (146)	144	20.1 (16)	293	60.4 (97)
Bldhd Migr	12	* *	* *	8	* *	* *	2	* *	2	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Booth Clin	11	* *	* *	* *	0	* *	* *	* *	0	* *	0	* *
Branch-Hil	820	14.3 (147)	41.8 (60)		672	27.8 (84)	47.2 (121)		126	23.0 (18)	162	34.0 (48)
Branch-Hil	753	14.7 (153)	38.9 (32)		665	29.5 (92)	44.2 (94)		135	19.3 (15)	203	21.7 (15)
Branch-Hil	728	11.7 (80)	45.1 (112)		624	29.3 (91)	41.0 (65)		136	36.0 (59)	179	43.0 (68)
Browning-C	68	* *	* *	* *	53	* *	* *	* *	7	* *	11	* *
Bryant Com	47	* *	* *	* *	44	* *	* *	* *	7	* *	13	* *
Burton	1,651	12.9 (113)	45.5 (118)		1,450	20.7 (19)	49.4 (149)		287	35.9 (58)	404	27.7 (26)
Calumet WI	239	10.5 (43)	38.1 (26)		195	76.9 (175)	13.3 (2)		19	* *	60	* *
Capac Clin	99	* *	* *	* *	82	* *	* *	* *	21	* *	44	* *
Caro WIC S	8	* *	* *	* *	6	* *	* *	* *	0	* *	4	* *
Cedar Spri	491	13.6 (128)	40.5 (43)		437	19.9 (14)	47.4 (123)		81	* *	138	9.4 (1)
Charlevoix	409	9.3 (25)	40.3 (40)		357	33.6 (116)	41.7 (75)		73	* *	156	66.7(100)
Chass Heal	3,212	11.7 (80)	35.8 (16)		2,784	34.3 (120)	34.8 (36)		500	23.0 (18)	1,726	23.5 (19)
Cheboygan	533	11.3 (63)	45.2 (114)		468	23.5 (48)	47.0 (119)		75	* *	121	22.3 (17)

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)										
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cherry Str	856	10.4 (41)	39.8 (37)	733	36.2 (128)	36.8 (43)	100	31.0 (43)	88	* *
Chesaning	89	* *	* *	79	* *	* *	16	* *	14	* *
Clare Coun	622	14.0 (134)	43.6 (89)	509	35.2 (123)	38.3 (52)	92	* *	152	32.9 (44)
Clinica Sa	882	6.2 (4)	44.6 (103)	781	67.0 (170)	13.8 (3)	102	28.4 (33)	300	67.0(101)
Comstock C	0	* *	* *	0	* *	* *	0	* *	0	* *
Corner Hea	578	22.3 (182)	32.0 (5)	481	18.3 (8)	50.1 (153)	57	* *	82	* *
Crawford	296	10.5 (43)	44.6 (103)	262	19.8 (12)	53.1 (170)	29	* *	94	* *
Dewitt WIC	156	11.5 (71)	46.8 (132)	141	24.8 (60)	48.2 (134)	28	* *	58	* *
Dickinson-	221	6.8 (6)	41.6 (56)	194	21.6 (31)	43.3 (87)	21	* *	48	* *
Dickinson-	515	12.6 (107)	42.3 (71)	451	17.5 (4)	54.1 (173)	58	* *	105	21.9 (16)
DMC Prim	25	* *	* *	20	* *	* *	4	* *	7	* *
DMC Prim	128	7.0 (7)	59.4 (184)	113	38.9 (138)	31.9 (25)	24	* *	26	* *
DMC Prim	40	* *	* *	37	* *	* *	3	* *	14	* *
Dorr Outre	265	15.5 (164)	36.6 (20)	239	20.9 (25)	48.5 (136)	39	* *	67	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Douglas He	2,018	9.8 (32)	48.9 (154)	1,762	30.7 (101)	43.0 (84)	320	30.3 (40)	356	32.6 (42)	
Dowagiac	983	13.7 (130)	41.0 (51)	816	48.8 (163)	31.1 (23)	157	24.2 (24)	254	42.9 (67)	
Durand WIC	265	15.5 (164)	38.5 (28)	228	22.8 (41)	45.2 (102)	46	* *	61	* *	
East Area	849	9.0 (21)	52.7 (176)	642	40.3 (143)	37.9 (46)	70	* *	57	* *	
Eastside H	1,583	12.1 (96)	47.6 (141)	1,405	27.4 (79)	43.0 (84)	376	42.8 (72)	406	47.8 (83)	
Eau Claire	451	5.1 (2)	54.3 (180)	338	46.4 (159)	24.9 (11)	62	* *	120	55.0 (94)	
Emmet Coun	626	10.9 (54)	42.7 (77)	549	29.0 (89)	41.0 (65)	87	* *	147	61.9 (98)	
Esanaba C	740	11.5 (71)	43.2 (84)	635	62.7 (168)	21.3 (8)	110	14.5 (7)	148	24.3 (21)	
Family Com	678	11.7 (80)	42.2 (70)	584	41.6 (146)	33.2 (28)	135	31.1 (44)	115	41.7 (65)	
Family Hlt	581	10.7 (48)	33.7 (11)	500	33.0 (114)	39.8 (60)	93	* *	146	39.7 (58)	
Family Roa	956	10.1 (37)	49.1 (158)	811	26.3 (69)	46.0 (109)	228	46.1 (76)	46	* *	
Farmington	127	11.8 (87)	35.4 (15)	100	44.0 (153)	26.0 (13)	33	* *	41	* *	
Fife Lake	2	* *	* *	2	* *	* *	0	* *	0	* *	
First Pres	0	* *	* *	0	* *	* *	0	* *	0	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Flint Fam	826	12.3 (102)	45.8 (122)	715	25.2 (62)	47.8 (128)	192	54.7 (79)	138	44.9 (73)	
Fruitport	19	* *	* *	18	* *	* *	3	* *	3	* *	
Gateway	1,016	9.4 (28)	53.8 (178)	820	35.6 (126)	41.6 (73)	194	30.4 (42)	164	45.1 (75)	
Genesee -	3,103	15.7 (169)	43.4 (87)	2,532	23.1 (43)	47.2 (121)	543	34.4 (56)	549	29.5 (30)	
Gladwin Co	432	11.6 (77)	44.9 (107)	400	24.3 (57)	49.0 (141)	51	* *	82	* *	
Gogebic Cl	235	12.3 (102)	40.4 (42)	202	19.8 (12)	49.0 (141)	45	* *	59	* *	
Golden	102	7.8 (8)	53.9 (179)	94	* *	* *	25	* *	30	* *	
Grace Prof	362	9.9 (33)	51.1 (171)	320	21.6 (31)	55.0 (174)	84	* *	51	* *	
Grace Ross	2,518	8.3 (14)	50.3 (168)	1,981	31.5 (105)	45.0 (100)	658	34.2 (55)	629	51.5 (90)	
Grand Have	612	13.1 (119)	43.1 (82)	531	22.6 (38)	47.1 (120)	106	29.2 (35)	211	49.8 (86)	
Grand Trav	13	* *	* *	10	* *	* *	2	* *	3	* *	
Grant	320	9.1 (23)	45.0 (109)	274	25.9 (67)	46.4 (114)	58	* *	56	* *	
Gratiot	1,717	10.4 (41)	47.4 (138)	1,543	27.5 (83)	46.5 (115)	304	41.1 (67)	641	45.4 (76)	
Greenville	352	18.5 (179)	42.6 (76)	315	41.6 (146)	36.5 (40)	63	* *	123	16.3 (7)	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	% (Rank)
Hamtramck	406	14.8 (155)	29.1 (3)	287	48.1 (162)	24.0 (9)	95	* *	95	* *	
Hancock Cl	422	15.2 (161)	33.2 (8)	340	70.6 (172)	14.4 (4)	52	* *	112	41.1 (63)	
Hannahvil	44	* *	* *	38	* *	* *	6	* *	11	* *	
Harper Gra	3,148	10.8 (52)	47.7 (143)	2,701	54.2 (166)	29.1 (18)	689	41.7 (71)	513	44.6 (72)	
Haslett Cl	137	14.6 (152)	43.8 (95)	123	34.1 (117)	43.1 (86)	29	* *	51	* *	
Haslett St	28	* *	* *	27	* *	* *	7	* *	8	* *	
Herman Kie	2,857	11.1 (57)	49.5 (163)	2,474	36.9 (132)	41.4 (71)	611	36.7 (61)	453	35.8 (52)	
Highland P	298	11.7 (80)	52.0 (173)	245	30.6 (100)	42.9 (83)	86	* *	37	* *	
Holland Cl	2,419	11.8 (87)	41.8 (60)	2,102	45.3 (157)	29.3 (19)	415	31.8 (47)	744	29.6 (31)	
Holly WIC	91	* *	* *	65	* *	* *	24	* *	20	* *	
Holton	79	* *	* *	68	* *	* *	10	* *	13	* *	
Howard Cit	150	8.0 (11)	45.3 (115)	131	42.0 (150)	33.6 (31)	30	* *	58	* *	
Hudsonv	509	11.8 (87)	38.5 (28)	461	37.5 (135)	33.4 (30)	96	* *	197	31.5 (37)	
Hurley Cli	539	15.2 (161)	41.0 (51)	466	23.2 (45)	45.5 (104)	131	45.0 (74)	94	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Huron Coun	648	12.8 (109)	45.7 (120)	573	25.7 (65)	44.0 (93)	80	* *	75	* *	
Icchn-Spar	211	5.2 (3)	50.7 (170)	148	45.3 (157)	25.7 (12)	23	* *	56	* *	
Ingham Cou	3,913	13.2 (122)	42.3 (71)	3,458	36.6 (129)	39.8 (60)	763	26.9 (28)	1,220	49.9 (87)	
Intervale	252	10.7 (48)	57.1 (183)	232	26.7 (76)	50.0 (152)	36	* *	53	* *	
Ionia Coun	788	12.1 (96)	46.8 (132)	684	20.8 (21)	49.6 (150)	159	13.8 (5)	237	47.7 (81)	
losco	492	12.8 (109)	41.9 (64)	426	21.6 (31)	50.2 (154)	70	* *	103	34.0 (48)	
Ironwood	0	* *	* *	0	* *	* *	0	* *	0	* *	
Isabella C	886	11.5 (71)	44.7 (106)	738	30.9 (102)	43.5 (89)	178	29.8 (37)	280	30.4 (32)	
Ithaca Cli	310	11.9 (93)	48.7 (153)	276	31.9 (107)	43.5 (89)	61	* *	88	* *	
Ithaca WIC	0	* *	* *	0	* *	* *	0	* *	0	* *	
Jackson Co	3,119	14.1 (141)	41.3 (54)	2,849	21.6 (31)	48.4 (135)	528	25.2 (25)	493	46.7 (79)	
Jackson M	0	* *	* *	0	* *	* *	0	* *	0	* *	
Janes St A	1,099	8.9 (20)	52.0 (173)	920	29.2 (90)	37.9 (46)	164	41.5 (69)	555	34.4 (51)	
Joy-Greenf	2,141	10.0 (34)	45.3 (115)	1,837	32.2 (109)	39.0 (55)	381	38.6 (62)	237	51.5 (90)	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Kalamazoo	749	11.2 (60)	47.7 (143)	618	41.6 (146)	38.0 (49)	138	53.6 (78)	226	54.9 (93)	
Kalamazoo	2,405	14.0 (134)	41.8 (60)	2,062	52.9 (165)	29.8 (20)	290	25.9 (26)	876	37.0 (54)	
Kalkaska	424	10.6 (46)	45.5 (118)	382	18.6 (9)	51.3 (160)	55	* *	149	47.7 (81)	
Kent	0	* *	* *	0	* *	* *	0	* *	0	* *	
Kent Count	3,663	11.2 (60)	42.3 (71)	3,119	25.8 (66)	46.0 (109)	550	17.3 (14)	816	15.6 (4)	
Kentwood	1,700	14.1 (141)	40.3 (40)	1,517	26.3 (69)	43.4 (88)	294	12.9 (3)	441	15.6 (4)	
Keweenaw	75	* *	* *	67	* *	* *	13	* *	13	* *	
Kingsley	39	* *	* *	33	* *	* *	3	* *	11	* *	
Lac Vieux	25	* *	* *	22	* *	* *	5	* *	5	* *	
Lake Co	227	9.3 (25)	49.8 (164)	181	26.5 (71)	48.6 (137)	35	* *	46	* *	
Lake Odess	60	* *	* *	51	* *	* *	10	* *	12	* *	
Lapeer	1,380	12.3 (102)	42.1 (69)	1,181	18.2 (7)	51.5 (162)	232	33.2 (50)	378	39.7 (58)	
Leelanu Cl	169	10.7 (48)	52.1 (175)	145	20.7 (19)	51.7 (163)	20	* *	43	* *	
Lenawee	0	* *	* *	0	* *	* *	0	* *	0	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight		< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	Number	% (Rank)	
Leslie-Sto	77	* *	* *	* *	69	* *	* *	* *	13	* *	28	* *
Lincoln Pa	1,029	13.1 (119)	45.7 (120)	589	23.9 (54)	49.2 (146)	279	27.6 (30)	152	31.6 (40)		
Livingston	901	17.1 (173)	35.1 (14)	793	17.0 (3)	53.8 (172)	212	9.4 (1)	374	9.4 (1)		
Lowell	120	10.8 (52)	34.2 (12)	105	23.8 (52)	55.2 (175)	21	* *	38	* *		
M-72 Migra	19	* *	* *	7	* *	* *	3	* *	3	* *		
Mancelona	223	11.7 (80)	48.0 (148)	193	39.9 (139)	35.2 (37)	29	* *	56	* *		
Manchester	24	* *	* *	22	* *	* *	4	* *	6	* *		
Manchester	5	* *	* *	3	* *	* *	0	* *	1	* *		
Manistee C	412	13.8 (131)	46.4 (127)	356	20.2 (16)	48.9 (140)	69	* *	88	* *		
Manistique	183	6.6 (5)	46.4 (127)	164	45.1 (156)	32.3 (26)	13	* *	47	* *		
Mansitee M	24	* *	* *	15	* *	* *	3	* *	3	* *		
Marion Bra	110	10.0 (34)	44.5 (101)	103	31.1 (104)	42.7 (80)	20	* *	59	* *		
Marquette	410	8.3 (14)	40.0 (38)	354	22.0 (36)	45.5 (104)	43	* *	72	* *		
Marshall C	302	10.6 (46)	46.7 (130)	256	26.6 (74)	47.7 (126)	36	* *	62	* *		

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight		< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	Number	% (Rank)	
Mason Clin	97	* *	* *	* *	88	* *	* *	* *	19	* *	31	* *
Mason Co	548	11.7 (80)	44.5 (101)		469	23.9 (54)	49.3 (148)		81	* *	126	48.4 (85)
Mccree Hea	3,025	11.5 (71)	47.6 (141)		2,663	24.2 (56)	47.8 (128)		732	41.5 (69)	740	33.2 (45)
Mcnamara H	168	11.3 (63)	49.4 (162)		144	18.1 (6)	47.9 (131)		29	* *	45	* *
Mecosta Co	855	14.0 (134)	40.1 (39)		753	71.3 (174)	12.1 (1)		123	33.3 (51)	166	39.2 (57)
Medical Pr	35	* *	* *	* *	31	* *	* *	* *	9	* *	7	* *
Medical Pr	83	* *	* *	* *	79	* *	* *	* *	13	* *	20	* *
Menominee	391	12.8 (109)	39.4 (35)		339	26.5 (71)	46.6 (116)		68	* *	91	* *
Mercy	117	11.1 (57)	55.6 (181)		106	58.5 (167)	24.5 (10)		37	* *	33	* *
Mesick	66	* *	* *	* *	60	* *	* *	* *	13	* *	30	* *
Metropolit	0	* *	* *	* *	0	* *	* *	* *	1	* *	0	* *
Michigan S	215	24.7 (183)	16.7 (2)		190	34.7 (122)	27.4 (15)		37	* *	95	* *
Midland Cl	994	13.8 (131)	46.7 (130)		845	27.9 (85)	44.9 (99)		187	33.7 (54)	283	38.9 (55)
Midwest	1,142	10.1 (37)	48.1 (149)		874	70.6 (172)	18.2 (6)		96	* *	125	50.4 (88)

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Migrant H	251	3.6 (1)	49.0 (157)	210	28.6 (88)	32.9 (27)	24	* *	47	* *	
Missaukee	210	8.1 (12)	50.0 (166)	190	27.4 (79)	44.2 (94)	29	* *	70	* *	
Moms & Bab	974	11.8 (87)	42.4 (74)	822	30.0 (96)	41.7 (75)	225	11.6 (2)	220	78.6(102)	
Monroe Cou	2,041	14.9 (159)	40.7 (46)	1,759	22.6 (38)	47.8 (128)	361	26.9 (28)	296	32.8 (43)	
Montague/	95	* *	* *	89	* *	* *	14	* *	12	* *	
Montcalm	17	* *	* *	12	* *	* *	3	* *	7	* *	
Montmorenc	171	12.9 (113)	48.5 (151)	143	47.6 (161)	33.6 (31)	13	* *	47	* *	
Mt Clemens	2,527	14.2 (145)	36.1 (17)	887	29.7 (94)	42.8 (82)	630	31.6 (45)	522	28.7 (29)	
Munising C	156	14.7 (153)	37.8 (25)	136	43.4 (152)	28.7 (17)	15	* *	48	* *	
Muskegon C	4,330	11.7 (80)	45.0 (109)	3,864	23.1 (43)	48.8 (139)	652	40.6 (66)	696	40.2 (61)	
New Center	23	* *	* *	22	* *	* *	5	* *	10	* *	
New Haven	262	13.4 (125)	42.0 (67)	117	38.5 (137)	42.7 (80)	47	* *	67	* *	
New Hope	11	* *	* *	10	* *	* *	2	* *	4	* *	
Newaygo	726	14.3 (147)	44.1 (98)	620	27.4 (79)	47.7 (126)	89	* *	212	32.5 (41)	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	% (Rank)
Newberry C	159	8.2 (13)	47.8 (145)	132	35.6 (126)	36.4 (39)	14	* *	35	* *	
North Camp	328	29.6 (184)	16.5 (1)	254	32.3 (111)	27.2 (14)	68	* *	101	19.8 (12)	
North Midl	0	* *	* *	0	* *	* *	0	* *	0	* *	
Northeast	2,777	10.9 (54)	49.3 (161)	2,321	40.5 (144)	36.5 (40)	413	32.9 (48)	671	33.2 (45)	
Northern M	129	15.5 (164)	41.9 (64)	116	14.7 (1)	49.1 (144)	18	* *	21	* *	
Northwest	493	11.6 (77)	47.9 (146)	398	30.2 (98)	46.2 (113)	86	* *	41	* *	
Oakland Co	4,248	11.4 (68)	45.0 (109)	3,050	34.1 (117)	40.2 (63)	963	35.8 (57)	1,014	31.5 (37)	
Oceana Co	526	8.7 (19)	51.7 (172)	452	23.7 (50)	50.9 (158)	68	* *	113	39.8 (60)	
Ogemaw	474	12.4 (105)	42.8 (78)	413	21.5 (29)	51.8 (164)	56	* *	119	27.7 (26)	
Olhsa Star	332	14.2 (145)	34.3 (13)	247	47.0 (160)	34.4 (35)	87	* *	75	* *	
Olhsa WIC	119	12.6 (107)	39.5 (36)	84	* *	* *	29	* *	19	* *	
Omni Care	38	* *	* *	36	* *	* *	6	* *	2	* *	
Omnicare	4	* *	* *	4	* *	* *	0	* *	0	* *	
Omnicare-G	0	* *	* *	0	* *	* *	0	* *	0	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Omnicare-M	85	* *	* *	* *	77	* *	* *	* *	23	* *	25	* *
Ontonagon	81	* *	* *	* *	54	* *	* *	* *	16	* *	13	* *
Oscoda	160	20.0 (180)	31.3 (4)		130	21.5 (29)	45.4 (103)		15	* *	33	* *
Otsego Co	560	14.5 (149)	41.6 (56)		501	20.8 (21)	48.1 (133)		68	* *	204	66.2 (99)
Park Clin	2	* *	* *	* *	2	* *	* *	* *	1	* *	1	* *
Park Medic	31	* *	* *	* *	28	* *	* *	* *	6	* *	6	* *
Park Medic	55	* *	* *	* *	52	* *	* *	* *	11	* *	17	* *
Park Medic	11	* *	* *	* *	11	* *	* *	* *	4	* *	2	* *
Paw Paw Cl	406	14.8 (155)	44.3 (99)		355	30.4 (99)	41.4 (71)		60	* *	78	* *
PCOC	119	16.8 (172)	32.8 (7)		96	* *	* *	* *	12	* *	42	* *
Pinconning	100	9.0 (21)	47.0 (136)		80	* *	* *	* *	15	* *	19	* *
Port Huron	2,243	14.5 (149)	43.6 (89)		1,960	23.6 (49)	46.1 (111)		366	31.7 (46)	1,093	43.3 (70)
Portland	84	* *	* *	* *	80	* *	* *	* *	18	* *	31	* *
Presque Is	196	10.7 (48)	45.9 (123)		175	24.6 (59)	44.6 (97)		29	* *	59	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Profession	76	* *	* *	* *	66	* *	* *	* *	10	* *	17	* *
Pullman	647	9.3 (25)	47.4 (138)		525	37.1 (134)	38.7 (54)		97	* *	91	* *
Ravenna	29	* *	* *	* *	26	* *	* *	* *	9	* *	2	* *
Reed City	421	16.4 (171)	44.9 (107)		343	33.5 (115)	38.2 (50)		74	* *	100	31.0 (34)
Region II	1,288	14.8 (155)	43.9 (96)		1,148	29.5 (92)	40.8 (64)		326	13.8 (5)	387	31.0 (34)
Riverview	86	* *	* *	* *	83	* *	* *	* *	26	* *	26	* *
Romulus Fi	264	14.0 (134)	38.3 (27)		103	27.2 (78)	45.6 (106)		74	* *	49	* *
Roosevelt-	149	11.4 (68)	53.7 (177)		124	29.8 (95)	44.4 (96)		21	* *	78	* *
Roscommon	424	15.3 (163)	41.7 (58)		383	22.5 (37)	52.7 (167)		57	* *	149	24.8 (22)
S & S Medi	30	* *	* *	* *	27	* *	* *	* *	4	* *	9	* *
Saginaw Co	3,424	8.5 (18)	49.1 (158)		2,911	21.2 (27)	49.1 (144)		590	28.8 (34)	598	44.0 (71)
Saginaw-Ch	48	* *	* *	* *	27	* *	* *	* *	7	* *	17	* *
Sanilac Co	815	13.6 (128)	42.8 (78)		727	27.4 (79)	41.7 (75)		141	16.3 (11)	308	20.5 (13)
Sault Ste	608	11.8 (87)	42.9 (80)		538	20.8 (21)	52.8 (168)		75	* *	121	16.5 (8)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Sawyer Afb	175	10.3 (40)	46.9 (134)	155	22.6 (38)	45.8 (107)	13	* *	30	* *	
School Bas	0	* *	* *	0	* *	* *	0	* *	0	* *	
Schoolcraf	1,426	9.6 (31)	50.1 (167)	1,279	23.4 (47)	47.4 (123)	369	43.1 (73)	329	45.9 (77)	
Seton Cent	850	10.5 (43)	49.2 (160)	746	69.7 (171)	18.2 (6)	215	41.4 (68)	132	59.1 (96)	
Shaftsburg	75	* *	* *	66	* *	* *	17	* *	22	* *	
Shelby M	262	9.2 (24)	48.9 (154)	200	41.0 (145)	30.0 (21)	21	* *	64	* *	
Sheldon Co	1,713	9.4 (28)	46.4 (127)	1,486	30.9 (102)	39.5 (58)	287	23.7 (22)	248	26.2 (25)	
Shiawassee	788	12.9 (113)	46.2 (126)	684	20.3 (17)	49.6 (150)	154	29.9 (38)	223	35.9 (53)	
Southfield	2,590	13.3 (124)	41.7 (58)	1,972	40.0 (140)	37.7 (44)	634	30.3 (40)	722	30.6 (33)	
Sparta Pub	301	13.0 (117)	40.9 (49)	266	19.5 (10)	53.4 (171)	56	* *	76	* *	
St Charles	155	8.4 (16)	48.4 (150)	142	16.2 (2)	50.7 (157)	32	* *	36	* *	
St Ignace	166	8.4 (16)	47.0 (136)	147	36.7 (130)	39.5 (58)	17	* *	26	* *	
St Johns W	360	17.5 (176)	41.4 (55)	307	23.8 (52)	50.2 (154)	45	* *	95	* *	
St Louis C	171	14.0 (134)	43.3 (86)	153	35.3 (125)	42.5 (79)	22	* *	61	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
St.Clair S	1,019	14.1 (141)	36.8 (22)	329	27.1 (77)	43.5 (89)	281	33.5 (53)	184	25.5 (24)	
Stanton Cl	689	12.8 (109)	43.7 (93)	611	42.1 (151)	33.7 (33)	110	16.4 (12)	220	20.5 (13)	
Stockbridg	55	* *	* *	45	* *	* *	15	* *	21	* *	
Sumpter He	914	12.5 (106)	43.1 (82)	762	35.2 (123)	37.8 (45)	192	28.1 (32)	248	25.4 (23)	
Taylor	2,555	15.7 (169)	39.0 (33)	1,999	50.5 (164)	31.6 (24)	630	27.9 (31)	643	23.6 (20)	
Teen Healt	13	* *	* *	9	* *	* *	4	* *	5	* *	
Temperance	73	* *	* *	66	* *	* *	15	* *	16	* *	
Thompsonv	13	* *	* *	11	* *	* *	2	* *	3	* *	
Three Oaks	122	14.8 (155)	36.1 (17)	112	25.0 (61)	41.1 (67)	30	* *	52	* *	
Three Rive	793	14.0 (134)	42.0 (67)	689	37.7 (136)	38.2 (50)	145	23.4 (21)	143	42.0 (66)	
Traverse C	1,316	14.0 (134)	38.8 (30)	1,160	21.4 (28)	47.6 (125)	165	40.0 (63)	412	47.3 (80)	
Tri-Cnty H	509	11.2 (60)	48.9 (154)	449	25.2 (62)	48.6 (137)	95	* *	90	* *	
Tuscola	1,104	13.9 (133)	43.2 (84)	968	23.7 (50)	46.6 (116)	180	15.6 (10)	244	19.3 (11)	
U Of M-Ped	0	* *	* *	0	* *	* *	0	* *	0	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)											
Clinic	Prepregnancy BMI (4)			Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight	< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Vcms River	106	11.3 (63)	55.7 (182)	96	* *	* *	21	* *	32	* *	
Vicksburg	102	11.8 (87)	37.3 (24)	89	* *	* *	4	* *	47	* *	
Walled Lak	941	17.1 (173)	36.6 (20)	706	23.2 (45)	52.1 (166)	225	23.1 (20)	306	18.6 (10)	
Warren Cli	1,593	12.9 (113)	41.0 (51)	463	28.1 (87)	41.3 (70)	403	26.6 (27)	260	31.5 (37)	
Washtenaw	83	* *	* *	68	* *	* *	15	* *	21	* *	
Wayne-Dear	2,283	10.0 (34)	33.2 (8)	1,472	40.2 (142)	30.0 (21)	427	40.0 (63)	534	39.0 (56)	
Webbervill	44	* *	* *	40	* *	* *	9	* *	21	* *	
West Genes	551	11.1 (57)	47.4 (138)	483	31.9 (107)	39.3 (57)	133	33.1 (49)	110	55.5 (95)	
Western Wa	1,146	10.9 (54)	43.7 (93)	863	41.8 (149)	35.9 (38)	264	45.1 (75)	265	41.5 (64)	
Westland C	1,886	13.4 (125)	41.9 (64)	1,182	32.6 (113)	42.4 (78)	547	21.2 (17)	394	34.0 (48)	
Westside C	1,480	10.2 (39)	46.9 (134)	1,319	34.3 (120)	36.7 (42)	186	12.9 (3)	340	15.6 (4)	
Westwood	58	* *	* *	47	* *	* *	4	* *	15	* *	
Wexford	808	13.1 (119)	40.6 (44)	703	20.8 (21)	52.9 (169)	102	33.3 (51)	240	31.3 (36)	
Wheeler WI	29	* *	* *	28	* *	* *	5	* *	10	* *	

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
 - (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)												
Clinic	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)
Whitmore L	47	* *	* *	* *	38	* *	* *	* *	9	* *	18	* *
Willow	38	* *	* *	* *	31	* *	* *	* *	5	* *	14	* *
Woodside	172	14.5 (149)	43.0 (81)		145	20.0 (15)	51.0 (159)		22	* *	30	* *
Wyoming Cl	1,519	11.5 (71)	38.8 (30)		1,364	30.0 (96)	45.1 (101)		228	14.9 (8)	359	15.0 (3)
Yale Clini	100	15.0 (160)	46.0 (124)		97	* *	* *	* *	16	* *	49	* *
Ypsilanti	2,549	11.3 (63)	44.6 (103)		2,097	32.4 (112)	41.2 (69)		530	40.4 (65)	476	40.3 (62)
Michigan	166,099	12.2	43.9		137,167	31.8	41.7		30,317	30.3	39,910	36.0

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Agape WIC	1,254	75.9 (99)	3.7 (20)	1,326	11.5(181)
Albion Cli	516	67.2(149)	19.6(165)	563	39.3 (77)
Alcona	132	87.9 (6)	4.5 (35)	143	46.2 (36)
Algonac Cl	311	78.8 (73)	11.9(120)	315	27.3(136)
Allegan	252	86.5 (13)	2.8 (11)	293	42.0 (61)
Alma WIC	164	69.5(139)	17.7(153)	196	48.5 (22)
Alpena Cou	531	87.8 (7)	4.3 (29)	547	42.6 (55)
Antrim - B	166	86.7 (12)	6.0 (58)	178	48.9 (19)
Arab/Chald	423	81.1 (53)	4.7 (41)	435	39.8 (74)
Arenac Cou	252	75.0(108)	13.9(134)	273	45.1 (44)
Associated	27	* *	* *	28	* *
Baldwin	167	74.3(112)	4.2 (28)	181	17.1(166)
Bangor	1,051	78.0 (80)	4.5 (35)	1,250	39.7 (75)
Baraga Cl	115	78.3 (76)	13.9(134)	117	46.2 (36)
Barry-Eato	814	76.0 (98)	13.4(130)	881	41.8 (62)
Barry-Eato	1,035	80.7 (54)	9.0 (95)	1,152	34.6(103)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Battle Cre	2,405	67.9(144)	19.5(163)	2,584	38.7 (84)
Baxter Com	57	* *	* *	63	* *
Bay Cl	1,899	68.3(142)	21.9(170)	2,112	49.6 (13)
Bayside WI	21	* *	* *	35	* *
Beaver Isl	5	* *	* *	5	* *
Belding Cl	233	73.8(116)	16.7(148)	259	39.0 (81)
Bell Memor	236	82.6 (38)	10.6(104)	266	49.6 (13)
Bendle/Bur	109	82.6 (38)	1.8 (3)	118	31.4(116)
Benton Har	1,830	69.5(139)	11.3(113)	2,043	33.2(112)
Benzie Cli	269	75.1(107)	12.3(124)	297	57.2 (4)
Berrien -	810	85.8 (17)	2.2 (7)	875	29.9(126)
Bldhd Migr	10	* *	* *	15	* *
Booth Clin	0	* *	* *	12	* *
Branch-Hil	786	71.8(127)	10.6(104)	881	33.7(108)
Branch-Hil	713	81.6 (48)	7.3 (80)	784	33.4(110)
Branch-Hil	743	67.7(145)	15.6(142)	828	30.6(121)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Browning-C	63	* *	* *	70	* *
Bryant Com	45	* *	* *	46	* *
Burton	1,679	75.6(102)	10.1(102)	1,752	30.9(118)
Calumet WI	232	81.5 (49)	11.2(111)	256	48.8 (21)
Capac Clin	93	* *	* *	101	27.7(134)
Caro WIC S	9	* *	* *	14	* *
Cedar Spri	498	80.7 (54)	11.8(119)	522	34.3(106)
Charlevoix	411	78.3 (76)	12.4(125)	455	40.7 (68)
Chass Heal	2,384	61.3(172)	19.3(162)	3,289	25.7(141)
Cheboygan	517	86.3 (15)	5.8 (55)	544	47.8 (27)
Cherry Str	800	50.5(180)	21.8(169)	867	41.4 (65)
Chesaning	85	* *	* *	90	* *
Clare Coun	588	77.4 (87)	7.7 (87)	645	43.7 (51)
Clinica Sa	754	28.4(181)	15.9(144)	945	29.4(129)
Comstock C	0	* *	* *	0	* *
Corner Hea	545	72.1(124)	6.2 (61)	601	41.4 (65)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Crawford	274	85.4 (21)	7.3 (80)	303	46.2 (36)
Dewitt WIC	155	82.6 (38)	4.5 (35)	162	29.6(127)
Dickinson-	213	87.3 (9)	4.7 (41)	231	48.1 (25)
Dickinson-	490	85.5 (20)	8.4 (92)	535	44.7 (46)
DMC Prim	24	* *	* *	23	* *
DMC Prim	123	67.5(147)	4.9 (45)	129	20.9(153)
DMC Prim	38	* *	* *	38	* *
Dorr Outre	262	85.9 (16)	4.6 (39)	267	32.6(114)
Douglas He	1,948	66.1(155)	9.3 (96)	2,221	26.0(140)
Dowagiac	912	81.5 (49)	4.4 (32)	1,036	33.4(110)
Durand WIC	263	80.2 (61)	12.9(128)	283	34.3(106)
East Area	815	64.7(163)	5.2 (50)	886	19.0(159)
Eastside H	1,551	64.4(164)	4.4 (32)	1,596	11.6(180)
Eau Claire	329	78.4 (75)	6.1 (59)	579	44.2 (49)
Emmet Coun	615	82.4 (42)	9.8(101)	650	43.8 (50)
Escanaba C	639	74.6(109)	18.3(158)	821	47.6 (29)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Family Com	639	74.2(113)	3.1 (15)	687	19.7(155)
Family Hlt	462	65.4(160)	19.5(163)	666	40.4 (69)
Family Roa	886	64.8(161)	2.9 (12)	961	11.1(183)
Farmington	128	79.7 (64)	3.9 (23)	128	19.5(156)
Fife Lake	2	* *	* *	2	* *
First Pres	0	* *	* *	0	* *
Flint Fam	807	64.3(166)	6.1 (59)	839	20.0(154)
Fruitport	20	* *	* *	20	* *
Gateway	988	64.8(161)	3.6 (19)	1,043	18.3(161)
Genesee -	2,890	72.5(122)	11.0(108)	3,180	31.7(115)
Gladwin Co	444	70.5(136)	20.7(167)	470	43.6 (53)
Gogebic Cl	237	63.7(167)	27.0(175)	264	48.1 (25)
Golden	98	* *	* *	101	16.8(171)
Grace Prof	345	62.9(171)	7.2 (75)	364	14.3(175)
Grace Ross	2,320	71.7(131)	4.1 (25)	2,558	12.5(178)
Grand Have	587	86.4 (14)	7.2 (75)	712	40.4 (69)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Clinic	Medical Care				WIC Enrollment	
	1st Trimester		No Care		1st Trimester	
	Number	% (Rank)	% (Rank)		Number	% (Rank)
Grand Trav	12	* *	* *		13	* *
Grant	305	80.7 (54)	5.6 (52)		328	34.5(104)
Gratiot	1,672	64.4(164)	4.5 (35)		1,742	10.3(186)
Greenville	342	82.5 (41)	7.3 (80)		355	31.3(117)
Hamtramck	389	65.6(156)	6.9 (71)		419	15.5(174)
Hancock Cl	391	84.7 (26)	11.0(108)		435	49.2 (16)
Hannahvil	40	* *	* *		46	* *
Harper Gra	2,818	66.9(150)	6.8 (68)		3,213	13.0(177)
Haslett Cl	139	89.9 (2)	2.9 (12)		143	21.7(149)
Haslett St	26	* *	* *		27	* *
Herman Kie	2,691	67.7(145)	6.8 (68)		2,912	15.8(173)
Highland P	278	56.1(177)	10.1(102)		321	17.1(166)
Holland Cl	2,273	84.9 (23)	2.3 (8)		2,528	36.3 (99)
Holly WIC	92	* *	* *		92	* *
Holton	73	* *	* *		81	* *
Howard Cit	147	85.7 (18)	6.8 (68)		151	30.5(122)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Hudsonv	502	89.0 (4)	2.0 (5)	533	30.4(124)
Hurley Cli	505	73.5(118)	4.6 (39)	544	21.7(149)
Huron Coun	623	87.3 (9)	6.3 (63)	658	48.5 (22)
Icchn-Spar	164	78.7 (74)	2.4 (9)	296	36.8 (94)
Ingham Cou	3,924	79.4 (65)	3.4 (17)	4,075	17.8(164)
Intervale	254	66.5(152)	7.5 (84)	257	13.6(176)
Ionia Coun	760	75.8(100)	15.8(143)	806	38.2 (86)
Iosco	459	85.2 (22)	4.1 (25)	492	42.5 (57)
Ironwood	0	* *	* *	0	* *
Isabella C	880	80.0 (62)	7.2 (75)	923	30.1(125)
Ithaca Cli	307	72.0(125)	17.9(156)	372	37.4 (92)
Ithaca WIC	0	* *	* *	0	* *
Jackson Co	3,098	82.4 (42)	5.7 (53)	3,390	38.7 (84)
Jackson M	0	* *	* *	0	* *
Janes St A	1,013	57.0(176)	23.7(171)	1,147	49.1 (17)
Joy-Greenf	2,038	71.2(129)	7.7 (87)	2,219	24.8(142)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Kalamazoo	643	67.5(147)	13.8(133)	999	42.1 (60)
Kalamazoo	1,778	77.7 (83)	13.9(134)	2,441	36.7 (95)
Kalkaska	371	79.2 (67)	9.4 (97)	433	41.3 (67)
Kent	0	* *	* *	0	* *
Kent Count	3,515	70.8(133)	18.0(157)	3,765	38.2 (86)
Kentwood	1,676	79.1 (69)	11.3(113)	1,756	28.4(132)
Keweenaw	70	* *	* *	74	* *
Kingsley	36	* *	* *	38	* *
Lac Vieux	22	* *	* *	24	* *
Lake Co	206	72.8(120)	11.2(111)	235	46.0 (40)
Lake Odess	58	* *	* *	64	* *
Lapeer	1,301	73.9(114)	17.7(153)	1,427	42.3 (58)
Leelanu Cl	161	76.4 (96)	16.1(146)	175	45.7 (41)
Lenawee	0	* *	* *	0	* *
Leslie-Sto	73	* *	* *	80	* *
Lincoln Pa	977	71.8(127)	10.7(107)	1,067	24.1(143)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Livingston	901	80.5 (58)	7.0 (72)	945	27.6(135)
Lowell	125	85.6 (19)	4.8 (43)	130	30.8(120)
M-72 Migra	7	* *	* *	29	* *
Mancelona	208	70.7(134)	18.3(158)	226	47.8 (27)
Manchester	24	* *	* *	25	* *
Manchester	6	* *	* *	7	* *
Manistee C	364	71.2(129)	19.8(166)	427	36.5 (96)
Manistique	178	92.7 (1)	1.1 (2)	187	62.0 (3)
Mansitee M	17	* *	* *	41	* *
Marion Bra	113	72.6(121)	16.8(150)	115	36.5 (96)
Marquette	388	80.4 (60)	12.1(121)	424	49.3 (15)
Marshall C	278	75.5(104)	18.7(160)	303	45.5 (43)
Mason Clin	106	84.9 (23)	0.0 (1)	102	10.8(185)
Mason Co	491	78.0 (80)	13.6(131)	567	46.2 (36)
Mccree Hea	2,983	66.2(153)	13.3(129)	3,384	23.1(147)
Mcnamara H	164	80.5 (58)	12.2(123)	170	35.9(100)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Mecosta Co	824	84.8 (25)	4.1 (25)	888	37.0 (93)
Medical Pr	38	* *	* *	38	* *
Medical Pr	78	* *	* *	84	* *
Menominee	366	69.4(141)	26.0(174)	409	40.3 (72)
Mercy	115	69.6(138)	4.3 (29)	118	16.9(168)
Mesick	62	* *	* *	67	* *
Metropolit	0	* *	* *	2	* *
Michigan S	210	83.8 (30)	1.9 (4)	219	17.8(164)
Midland Cl	1,123	82.9 (37)	9.5 (99)	1,227	44.3 (48)
Midwest	1,066	76.5 (95)	3.8 (22)	1,150	29.6(127)
Migrant H	227	50.7(179)	42.3(181)	257	68.9 (1)
Missaukee	203	73.4(119)	17.7(153)	220	50.5 (9)
Moms & Bab	913	65.6(156)	5.7 (53)	1,006	18.3(161)
Monroe Cou	1,951	70.7(134)	17.3(152)	2,093	34.9(102)
Montague/	97	* *	* *	97	* *
Montcalm	17	* *	* *	22	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Montmorenc	161	88.2 (5)	3.7 (20)	168	53.0 (7)
Mt Clemens	2,208	79.0 (72)	7.2 (75)	2,590	27.3(136)
Munising C	152	77.6 (84)	15.1(141)	158	45.6 (42)
Muskegon C	4,171	73.7(117)	13.7(132)	4,453	39.0 (81)
New Center	24	* *	* *	24	* *
New Haven	241	77.6 (84)	6.6 (66)	269	30.9(118)
New Hope	11	* *	* *	13	* *
Newaygo	683	83.6 (32)	7.2 (75)	753	43.7 (51)
Newberry C	156	59.6(174)	32.7(178)	172	57.0 (5)
North Camp	305	89.2 (3)	3.3 (16)	333	35.4(101)
North Midl	0	* *	* *	0	* *
Northeast	2,660	66.2(153)	9.5 (99)	2,820	16.6(172)
Northern M	124	83.9 (29)	6.5 (65)	130	46.9 (33)
Northwest	482	65.6(156)	3.5 (18)	502	16.9(168)
Oakland Co	4,005	75.7(101)	5.0 (46)	4,369	23.2(146)
Oceana Co	488	70.9(132)	20.7(167)	542	46.7 (34)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Ogemaw	450	84.7 (26)	5.1 (48)	482	50.2 (11)
Olhse Star	328	83.5 (33)	2.1 (6)	346	23.7(145)
Olhse WIC	126	74.6(109)	7.1 (74)	126	21.4(151)
Omni Care	37	* *	* *	38	* *
Omnicare	4	* *	* *	4	* *
Omnicare-G	0	* *	* *	0	* *
Omnicare-M	81	* *	* *	84	* *
Ontonagon	86	* *	* *	90	* *
Oscoda	139	81.3 (52)	9.4 (97)	167	49.1 (17)
Otsego Co	554	79.4 (65)	12.5(126)	576	50.0 (12)
Park Clin	2	* *	* *	2	* *
Park Medic	31	* *	* *	33	* *
Park Medic	55	* *	* *	56	* *
Park Medic	11	* *	* *	11	* *
Paw Paw Cl	375	82.1 (44)	4.8 (43)	411	37.7 (91)
PCOC	98	* *	* *	122	42.6 (55)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Clinic	Medical Care			WIC Enrollment		
	1st Trimester		No Care	1st Trimester		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	
Pinconning	106	77.4 (87)	14.2(137)	112	47.3 (31)	
Port Huron	2,153	59.9(173)	30.6(177)	2,291	44.5 (47)	
Portland	89	* *	* *	88	* *	
Presque Is	190	82.1 (44)	11.1(110)	200	41.5 (64)	
Profession	74	* *	* *	76	* *	
Pullman	588	79.9 (63)	6.3 (63)	711	40.4 (69)	
Ravenna	32	* *	* *	32	* *	
Reed City	412	83.0 (36)	8.3 (91)	427	37.9 (89)	
Region II	1,200	83.8 (30)	5.8 (55)	1,676	29.2(130)	
Riverview	86	* *	* *	86	* *	
Romulus Fi	236	72.0(125)	10.6(104)	279	21.1(152)	
Roosevelt-	137	77.4 (87)	8.8 (94)	153	48.4 (24)	
Roscommon	401	79.1 (69)	11.7(116)	430	42.3 (58)	
S & S Medi	29	* *	* *	30	* *	
Saginaw Co	3,194	69.8(137)	16.7(148)	3,456	41.8 (62)	
Saginaw-Ch	44	* *	* *	53	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Sanilac Co	782	76.7 (94)	14.3(138)	821	39.2 (78)
Sault Ste	590	75.4(105)	16.6(147)	629	50.9 (8)
Sawyer Afb	164	83.5 (33)	8.5 (93)	178	44.9 (45)
School Bas	0	* *	* *	0	* *
Schoolcraf	1,408	63.6(169)	5.9 (57)	1,443	11.3(182)
Seton Cent	816	68.0(143)	4.3 (29)	860	10.9(184)
Shaftsburg	77	* *	* *	81	* *
Shelby M	213	52.1(178)	37.6(180)	353	43.6 (53)
Sheldon Co	1,712	65.5(159)	16.0(145)	1,850	34.4(105)
Shiawassee	1,001	59.3(175)	34.6(179)	1,102	47.5 (30)
Southfield	2,423	79.2 (67)	3.0 (14)	2,639	19.2(158)
Sparta Pub	305	80.7 (54)	11.5(115)	318	32.7(113)
St Charles	157	75.2(106)	14.6(139)	159	39.0 (81)
St Ignace	159	81.8 (47)	4.4 (32)	175	64.0 (2)
St Johns W	341	86.8 (11)	5.0 (46)	374	36.4 (98)
St Louis C	164	74.4(111)	18.9(161)	195	39.5 (76)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
St.Clair S	930	81.4 (51)	7.3 (80)	1,054	28.6(131)
Stanton Cl	663	83.1 (35)	8.0 (89)	700	39.1 (79)
Stockbridg	57	* *	* *	60	* *
Sumpter He	907	76.2 (97)	8.0 (89)	950	26.4(139)
Taylor	2,507	72.4(123)	12.1(121)	2,811	24.1(143)
Teen Healt	11	* *	* *	13	* *
Temperance	71	* *	* *	73	* *
Thompsonv	11	* *	* *	13	* *
Three Oaks	128	87.5 (8)	3.9 (23)	124	16.9(168)
Three Rive	771	76.8 (93)	11.7(116)	814	38.0 (88)
Traverse C	1,272	66.6(151)	24.0(172)	1,332	46.6 (35)
Tri-Cnty H	507	84.0 (28)	5.1 (48)	533	40.3 (72)
Tuscola	1,140	82.0 (46)	7.6 (85)	1,198	39.1 (79)
U Of M-Ped	0	* *	* *	0	* *
Vcms River	97	* *	* *	104	12.5(178)
Vicksburg	73	* *	* *	104	53.8 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Walled Lak	927	77.2 (90)	7.6 (85)	964	19.4(157)
Warren Cli	1,423	79.1 (69)	7.0 (72)	1,618	26.7(138)
Washtenaw	81	* *	* *	88	* *
Wayne-Dear	2,154	75.6(102)	11.7(116)	2,443	37.9 (89)
Webbervill	48	* *	* *	44	* *
West Genes	543	77.2 (90)	2.4 (9)	575	18.3(161)
Western Wa	1,165	77.9 (82)	6.6 (66)	1,177	23.0(148)
Westland C	1,763	78.2 (78)	6.2 (61)	1,965	19.0(159)
Westside C	1,456	63.7(167)	25.3(173)	1,563	47.0 (32)
Westwood	39	* *	* *	62	* *
Wexford	729	77.5 (86)	12.6(127)	823	50.3 (10)
Wheeler WI	29	* *	* *	34	* *
Whitmore L	43	* *	* *	50	* *
Willow	35	* *	* *	39	* *
Woodside	161	73.9(114)	16.8(150)	182	48.9 (19)
Wyoming Cl	1,502	77.0 (92)	14.6(139)	1,555	33.6(109)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)					
Clinic	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	% (Rank)	% (Rank)	Number	% (Rank)
Yale Clini	107	63.6(169)	27.1(176)	105	30.5(122)
Ypsilanti	2,556	78.2 (78)	5.2 (50)	2,769	28.3(133)
Michigan	157,725	73.6	10.6	174,143	31.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agape WIC	825	21.1 (17)	744	9.0 (13)	61	* *	0	* *	0	* *
Albion Cli	559	52.1 (116)	475	34.7 (137)	198	21.7 (99)	0	* *	0	* *
Alcona	135	57.0 (150)	110	41.8 (163)	42	* *	0	* *	0	* *
Algonac Cl	320	55.9 (143)	285	35.1 (142)	138	27.5 (74)	0	* *	0	* *
Allegan	279	48.0 (85)	230	32.6 (122)	97	* *	0	* *	0	* *
Alma WIC	145	48.3 (89)	150	34.0 (135)	42	* *	0	* *	0	* *
Alpena Cou	549	58.5 (162)	503	34.8 (139)	245	31.8 (37)	0	* *	0	* *
Antrim - B	179	51.4 (109)	154	27.9 (91)	64	* *	0	* *	0	* *
Arab/Chald	377	7.2 (7)	382	6.0 (9)	13	* *	0	* *	0	* *
Arenac Cou	263	51.7 (112)	107	18.7 (42)	39	* *	0	* *	0	* *
Associated	21	* *	20	* *	2	* *	0	* *	0	* *
Baldwin	167	35.9 (48)	128	24.2 (64)	32	* *	0	* *	0	* *
Bangor	1,183	38.2 (53)	1,012	23.6 (62)	310	30.6 (47)	0	* *	0	* *
Baraga Cl	120	52.5 (120)	102	26.5 (75)	47	* *	0	* *	0	* *
Barry-Eato	778	57.2 (152)	634	35.6 (146)	241	19.9 (107)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Barry-Eato	1,132	53.0 (123)	930	30.6 (109)	377	27.9 (71)	0	* *	0	* *
Battle Cre	2,516	48.0 (85)	2,070	30.6 (109)	825	25.5 (87)	0	* *	0	* *
Baxter Com	64	* *	46	* *	14	* *	0	* *	0	* *
Bay Cl	1,666	55.1 (138)	1,637	32.0 (119)	588	27.6 (73)	0	* *	0	* *
Bayside WI	36	* *	23	* *	2	* *	0	* *	0	* *
Beaver Isl	5	* *	6	* *	3	* *	0	* *	0	* *
Belding Cl	254	56.3 (145)	217	42.9 (165)	110	20.0 (106)	0	* *	0	* *
Bell Memor	260	58.5 (162)	222	35.1 (142)	102	28.4 (66)	0	* *	0	* *
Bendle/Bur	119	57.1 (151)	99	* *	40	* *	0	* *	0	* *
Benton Har	1,136	58.8 (166)	1,096	30.9 (113)	329	21.6 (100)	0	* *	0	* *
Benzie Cli	282	58.2 (159)	226	38.9 (158)	113	20.4 (104)	0	* *	0	* *
Berrien -	865	51.6 (111)	741	29.6 (99)	315	33.7 (26)	0	* *	0	* *
Bldhd Migr	12	* *	8	* *	0	* *	0	* *	0	* *
Booth Clin	11	* *	0	* *	0	* *	0	* *	0	* *
Branch-Hil	850	51.3 (106)	680	33.8 (133)	244	20.1 (105)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Branch-Hil	775	56.0 (144)	703	35.3 (145)	307	24.8 (92)	0	* *	0	* *
Branch-Hil	789	36.4 (50)	636	25.2 (68)	175	18.9 (108)	0	* *	0	* *
Browning-C	65	* *	49	* *	14	* *	0	* *	0	* *
Bryant Com	46	* *	43	* *	5	* *	0	* *	0	* *
Burton	1,606	52.2 (118)	1,363	33.4 (129)	502	24.9 (90)	0	* *	0	* *
Calumet WI	248	50.4 (99)	210	27.1 (85)	84	* *	0	* *	0	* *
Capac Clin	103	57.3 (153)	87	* *	38	* *	0	* *	0	* *
Caro WIC S	14	* *	11	* *	0	* *	0	* *	0	* *
Cedar Spri	513	47.8 (84)	468	27.1 (85)	174	35.6 (13)	0	* *	0	* *
Charlevoix	380	57.6 (154)	359	26.7 (78)	149	35.6 (13)	0	* *	0	* *
Chass Heal	961	22.1 (18)	1,012	10.7 (16)	61	* *	0	* *	0	* *
Cheboygan	548	56.9 (149)	491	34.8 (139)	236	29.7 (52)	0	* *	0	* *
Cherry Str	803	19.6 (16)	688	10.6 (15)	114	27.2 (77)	0	* *	0	* *
Chesaning	80	* *	76	* *	19	* *	0	* *	0	* *
Clare Coun	486	79.6 (180)	303	60.4 (171)	166	6.0 (109)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Clinica Sa	912	0.7 (2)	831	0.1 (3)	5	* *	0	* *	0	* *
Comstock C	0	* *	0	* *	0	* *	0	* *	0	* *
Corner Hea	589	51.8 (114)	485	26.8 (79)	225	34.7 (19)	0	* *	0	* *
Crawford	300	58.3 (161)	275	35.6 (146)	126	25.4 (89)	0	* *	0	* *
Dewitt WIC	154	51.3 (106)	145	28.3 (95)	50	* *	0	* *	0	* *
Dickinson-	227	56.4 (147)	207	36.7 (153)	76	* *	0	* *	0	* *
Dickinson-	529	50.9 (103)	456	32.9 (125)	201	26.4 (83)	0	* *	0	* *
DMC Prim	23	* *	19	* *	6	* *	0	* *	0	* *
DMC Prim	111	24.3 (22)	96	* *	18	* *	0	* *	0	* *
DMC Prim	33	* *	32	* *	7	* *	0	* *	0	* *
Dorr Outre	184	66.3 (177)	160	42.5 (164)	64	* *	0	* *	0	* *
Douglas He	1,302	28.3 (35)	1,532	11.4 (19)	180	28.9 (62)	0	* *	0	* *
Dowagiac	981	49.7 (95)	807	31.6 (115)	297	27.3 (76)	0	* *	0	* *
Durand WIC	279	60.9 (168)	256	44.1 (167)	116	31.0 (42)	0	* *	0	* *
East Area	839	22.6 (21)	591	14.2 (26)	98	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Eastside H	1,200	27.3 (30)	1,074	17.2 (38)	165	23.6 (96)	0	* *	0	* *
Eau Claire	402	5.7 (6)	364	3.6 (8)	15	* *	0	* *	0	* *
Emmet Coun	629	53.7 (130)	497	31.8 (116)	235	32.8 (31)	0	* *	0	* *
Escanaba C	719	55.6 (140)	651	33.2 (128)	287	24.7 (93)	0	* *	0	* *
Family Com	665	18.0 (13)	578	11.6 (20)	82	* *	0	* *	0	* *
Family Hlt	396	42.9 (67)	552	26.8 (79)	107	29.0 (57)	0	* *	0	* *
Family Roa	948	25.8 (27)	821	13.8 (24)	140	35.7 (12)	0	* *	0	* *
Farmington	119	27.7 (32)	90	* *	14	* *	0	* *	0	* *
Fife Lake	2	* *	2	* *	2	* *	0	* *	0	* *
First Pres	0	* *	0	* *	0	* *	0	* *	0	* *
Flint Fam	842	37.9 (52)	705	22.6 (59)	210	26.7 (79)	0	* *	0	* *
Fruitport	16	* *	19	* *	9	* *	0	* *	0	* *
Gateway	1,022	28.1 (34)	799	19.5 (47)	145	29.0 (57)	0	* *	0	* *
Genesee -	2,989	46.7 (80)	2,510	29.6 (99)	889	25.5 (87)	0	* *	0	* *
Gladwin Co	462	52.4 (119)	434	29.7 (103)	197	31.0 (42)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Gogebic Cl	242	54.1 (133)	224	33.5 (130)	92	* *	0	* *	0	* *
Golden	86	* *	83	* *	11	* *	0	* *	0	* *
Grace Prof	304	25.7 (26)	259	15.1 (31)	45	* *	0	* *	0	* *
Grace Ross	2,125	27.8 (33)	1,670	11.6 (20)	312	49.0 (1)	0	* *	0	* *
Grand Have	619	44.9 (72)	621	25.3 (69)	202	39.1 (9)	0	* *	0	* *
Grand Trav	13	* *	10	* *	3	* *	0	* *	0	* *
Grant	332	40.1 (61)	291	27.1 (85)	100	27.0 (78)	0	* *	0	* *
Gratiot	1,747	19.5 (15)	1,597	10.5 (14)	189	34.9 (17)	0	* *	0	* *
Greenville	296	62.5 (172)	288	35.1 (142)	130	30.8 (44)	0	* *	0	* *
Hamtramck	406	17.5 (11)	297	8.1 (12)	27	* *	0	* *	0	* *
Hancock Cl	450	46.9 (82)	366	21.3 (54)	140	41.4 (5)	0	* *	0	* *
Hannahvil	44	* *	40	* *	27	* *	0	* *	0	* *
Harper Gra	2,630	18.1 (14)	2,120	13.2 (22)	184	28.3 (68)	0	* *	0	* *
Haslett Cl	139	33.8 (43)	124	14.5 (28)	35	* *	0	* *	0	* *
Haslett St	27	* *	23	* *	11	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Herman Kie	1,981	30.7 (39)	1,909	16.0 (35)	267	28.5 (64)	0	* *	0	* *
Highland P	298	26.5 (28)	265	10.9 (17)	40	* *	0	* *	0	* *
Holland Cl	2,491	28.7 (36)	2,182	13.9 (25)	508	42.3 (4)	0	* *	0	* *
Holly WIC	84	* *	61	* *	26	* *	0	* *	0	* *
Holton	73	* *	69	* *	26	* *	0	* *	0	* *
Howard Cit	126	53.2 (126)	122	32.8 (123)	43	* *	0	* *	0	* *
Hudsonv	536	38.8 (56)	495	18.2 (41)	144	43.8 (3)	0	* *	0	* *
Hurley Cli	415	41.2 (63)	317	26.5 (75)	69	* *	0	* *	0	* *
Huron Coun	658	46.5 (79)	590	28.0 (92)	240	27.9 (71)	0	* *	0	* *
Icchn-Spar	215	0.5 (1)	161	0.0 (1)	1	* *	0	* *	0	* *
Ingham Cou	3,751	45.0 (73)	3,270	28.8 (96)	949	26.0 (86)	0	* *	0	* *
Intervale	184	22.3 (19)	160	15.0 (30)	20	* *	0	* *	0	* *
Ionia Coun	817	49.7 (95)	722	31.9 (118)	305	30.2 (50)	0	* *	0	* *
Iosco	467	61.9 (171)	410	38.5 (157)	203	20.7 (103)	0	* *	0	* *
Ironwood	0	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Isabella C	904	49.0 (91)	687	26.3 (73)	253	40.7 (6)	0	* *	0	* *
Ithaca Cli	271	57.6 (154)	289	31.8 (116)	92	* *	0	* *	0	* *
Ithaca WIC	0	* *	0	* *	0	* *	0	* *	0	* *
Jackson Co	3,398	52.7 (122)	2,895	33.5 (130)	1,196	24.2 (94)	0	* *	0	* *
Jackson M	0	* *	0	* *	0	* *	0	* *	0	* *
Janes St A	1,110	37.3 (51)	948	21.3 (54)	299	28.1 (69)	0	* *	0	* *
Joy-Greenf	1,307	26.9 (29)	1,274	15.1 (31)	170	29.4 (54)	0	* *	0	* *
Kalamazoo	607	38.6 (55)	761	26.0 (71)	148	22.3 (98)	0	* *	0	* *
Kalamazoo	1,084	46.8 (81)	1,260	26.6 (77)	245	33.9 (23)	0	* *	0	* *
Kalkaska	432	54.2 (134)	394	35.8 (148)	191	26.2 (84)	0	* *	0	* *
Kent	0	* *	0	* *	0	* *	0	* *	0	* *
Kent Count	3,244	45.3 (75)	2,824	22.4 (57)	868	33.8 (25)	0	* *	0	* *
Kentwood	1,631	33.8 (43)	1,512	17.3 (39)	352	40.6 (7)	0	* *	0	* *
Keweenaw	75	* *	67	* *	42	* *	0	* *	0	* *
Kingsley	40	* *	32	* *	17	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lac Vieux	25	* *	22	* *	12	* *	0	* *	0	* *
Lake Co	226	60.6 (167)	171	47.4 (169)	89	* *	0	* *	0	* *
Lake Odess	61	* *	54	* *	19	* *	0	* *	0	* *
Lapeer	1,435	53.8 (131)	1,258	32.0 (119)	551	31.6 (38)	0	* *	0	* *
Leelanu Cl	169	53.8 (131)	145	35.9 (149)	64	* *	0	* *	0	* *
Lenawee	0	* *	0	* *	0	* *	0	* *	0	* *
Leslie-Sto	75	* *	66	* *	27	* *	0	* *	0	* *
Lincoln Pa	1,031	44.6 (70)	590	26.1 (72)	212	33.5 (28)	0	* *	0	* *
Livingston	954	58.2 (159)	860	33.5 (130)	355	30.1 (51)	0	* *	0	* *
Lowell	91	* *	77	* *	25	* *	0	* *	0	* *
M-72 Migra	18	* *	10	* *	1	* *	0	* *	0	* *
Mancelona	223	51.1 (104)	192	30.7 (111)	89	* *	0	* *	0	* *
Manchester	25	* *	21	* *	7	* *	0	* *	0	* *
Manchester	8	* *	6	* *	0	* *	0	* *	0	* *
Manistee C	406	53.4 (129)	357	32.5 (121)	168	33.9 (23)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Manistique	189	55.0 (137)	160	33.1 (127)	82	* *	0	* *	0	* *
Mansitee M	28	* *	20	* *	1	* *	0	* *	0	* *
Marion Bra	106	55.7 (141)	98	* *	48	* *	0	* *	0	* *
Marquette	405	53.1 (124)	365	25.5 (70)	158	39.2 (8)	0	* *	0	* *
Marshall C	300	45.3 (75)	250	22.8 (60)	91	* *	0	* *	0	* *
Mason Clin	103	44.7 (71)	92	* *	24	* *	0	* *	0	* *
Mason Co	555	46.1 (78)	488	20.7 (51)	189	35.4 (16)	0	* *	0	* *
Mccree Hea	2,935	39.9 (60)	2,800	24.6 (66)	780	30.6 (47)	0	* *	0	* *
Mcnamara H	152	36.2 (49)	140	19.3 (46)	31	* *	0	* *	0	* *
Mecosta Co	865	49.7 (95)	784	27.7 (89)	318	31.4 (39)	0	* *	0	* *
Medical Pr	27	* *	26	* *	2	* *	0	* *	0	* *
Medical Pr	64	* *	52	* *	10	* *	0	* *	0	* *
Menominee	355	64.2 (174)	325	43.1 (166)	146	21.2 (101)	0	* *	0	* *
Mercy	107	22.4 (20)	98	* *	13	* *	0	* *	0	* *
Mesick	66	* *	60	* *	20	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Metropolit	1	* *	0	* *	0	* *	0	* *	0	* *
Michigan S	211	7.6 (8)	184	2.2 (6)	9	* *	0	* *	0	* *
Midland Cl	1,198	49.0 (91)	1,043	26.8 (79)	419	33.7 (26)	0	* *	0	* *
Midwest	1,113	17.7 (12)	893	11.1 (18)	122	31.1 (41)	0	* *	0	* *
Migrant H	255	2.4 (3)	212	0.9 (5)	5	* *	0	* *	0	* *
Missaukee	213	50.7 (102)	203	29.6 (99)	88	* *	0	* *	0	* *
Moms & Bab	917	16.0 (10)	877	7.3 (11)	92	* *	0	* *	0	* *
Monroe Cou	2,131	53.1 (124)	1,840	36.0 (150)	799	26.2 (84)	0	* *	0	* *
Montague/	91	* *	93	* *	40	* *	0	* *	0	* *
Montcalm	22	* *	16	* *	0	* *	0	* *	0	* *
Montmorenc	169	66.9 (178)	143	44.1 (167)	74	* *	0	* *	0	* *
Mt Clemens	2,482	42.4 (65)	816	16.3 (37)	194	37.1 (11)	0	* *	0	* *
Munising C	163	55.8 (142)	128	28.9 (97)	56	* *	0	* *	0	* *
Muskegon C	4,049	48.3 (89)	3,850	29.6 (99)	1,467	28.5 (64)	0	* *	0	* *
New Center	21	* *	20	* *	4	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
New Haven	265	34.7 (47)	110	6.4 (10)	11	* *	0	* *	0	* *
New Hope	10	* *	11	* *	1	* *	0	* *	0	* *
Newaygo	736	50.3 (98)	629	29.4 (98)	264	29.5 (53)	0	* *	0	* *
Newberry C	159	64.8 (175)	147	36.7 (153)	80	* *	0	* *	0	* *
North Camp	305	5.6 (5)	265	0.8 (4)	14	* *	0	* *	0	* *
North Midl	0	* *	0	* *	0	* *	0	* *	0	* *
Northeast	2,037	24.7 (23)	1,903	14.2 (26)	218	33.0 (30)	0	* *	0	* *
Northern M	127	49.6 (94)	118	20.3 (49)	45	* *	0	* *	0	* *
Northwest	471	25.1 (25)	370	15.9 (33)	70	* *	0	* *	0	* *
Oakland Co	4,060	39.3 (58)	2,763	21.2 (53)	726	32.2 (35)	0	* *	0	* *
Oceana Co	536	45.7 (77)	464	24.4 (65)	177	30.5 (49)	0	* *	0	* *
Ogemaw	490	56.3 (145)	415	30.4 (107)	205	28.8 (63)	0	* *	0	* *
Olhsa Star	318	34.0 (46)	234	19.2 (44)	51	* *	0	* *	0	* *
Olhsa WIC	124	33.9 (45)	76	* *	12	* *	0	* *	0	* *
Omni Care	24	* *	22	* *	2	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Omicare	4	* *	3	* *	0	* *	0	* *	0	* *
Omicare-G	0	* *	0	* *	0	* *	0	* *	0	* *
Omicare-M	71	* *	62	* *	8	* *	0	* *	0	* *
Ontonagon	87	* *	59	* *	19	* *	0	* *	0	* *
Oscoda	163	57.7 (157)	140	30.7 (111)	65	* *	0	* *	0	* *
Otsego Co	587	52.1 (116)	507	27.0 (83)	233	38.2 (10)	0	* *	0	* *
Park Clin	2	* *	2	* *	0	* *	0	* *	0	* *
Park Medic	26	* *	26	* *	2	* *	0	* *	0	* *
Park Medic	39	* *	38	* *	7	* *	0	* *	0	* *
Park Medic	11	* *	9	* *	0	* *	0	* *	0	* *
Paw Paw Cl	408	52.5 (120)	360	35.0 (141)	158	26.6 (80)	0	* *	0	* *
PCOC	70	* *	66	* *	14	* *	0	* *	0	* *
Pinconning	97	* *	86	* *	31	* *	0	* *	0	* *
Port Huron	2,299	58.7 (165)	2,057	36.0 (150)	1,042	29.0 (57)	0	* *	0	* *
Portland	90	* *	87	* *	46	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Presque Is	198	51.5 (110)	177	32.8 (123)	78	* *	0	* *	0	* *
Profession	63	* *	56	* *	4	* *	0	* *	0	* *
Pullman	672	33.6 (41)	550	22.2 (56)	149	23.5 (97)	0	* *	0	* *
Ravenna	30	* *	28	* *	11	* *	0	* *	0	* *
Reed City	436	57.6 (154)	362	39.0 (159)	168	23.8 (95)	0	* *	0	* *
Region II	1,121	40.2 (62)	1,343	24.7 (67)	276	29.3 (55)	0	* *	0	* *
Riverview	76	* *	72	* *	18	* *	0	* *	0	* *
Romulus Fi	263	51.7 (112)	117	28.2 (93)	33	* *	0	* *	0	* *
Roosevelt-	147	29.9 (37)	125	16.0 (35)	33	* *	0	* *	0	* *
Roscommon	425	60.9 (168)	390	37.7 (156)	199	26.6 (80)	0	* *	0	* *
S & S Medi	25	* *	23	* *	2	* *	0	* *	0	* *
Saginaw Co	2,915	30.5 (38)	2,754	20.8 (52)	567	34.7 (19)	0	* *	0	* *
Saginaw-Ch	51	* *	31	* *	6	* *	0	* *	0	* *
Sanilac Co	840	51.3 (106)	763	30.0 (106)	333	34.5 (22)	0	* *	0	* *
Sault Ste	623	53.3 (127)	532	31.0 (114)	242	28.1 (69)	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sawyer Afb	177	56.5 (148)	158	29.7 (103)	75	* *	0	* *	0	* *
School Bas	0	* *	0	* *	0	* *	0	* *	0	* *
Schoolcraf	1,375	27.6 (31)	1,231	13.5 (23)	228	46.1 (2)	0	* *	0	* *
Seton Cent	391	43.2 (68)	371	19.7 (48)	53	* *	0	* *	0	* *
Shaftsburg	76	* *	71	* *	30	* *	0	* *	0	* *
Shelby M	267	4.5 (4)	235	0.0 (1)	9	* *	0	* *	0	* *
Sheldon Co	1,769	25.0 (24)	1,416	15.9 (33)	272	29.0 (57)	0	* *	0	* *
Shiawassee	801	55.3 (139)	975	32.9 (125)	328	29.0 (57)	0	* *	0	* *
Southfield	1,843	38.4 (54)	1,373	17.8 (40)	269	24.9 (90)	0	* *	0	* *
Sparta Pub	315	49.5 (93)	286	27.3 (88)	106	34.9 (17)	0	* *	0	* *
St Charles	141	39.0 (57)	136	33.8 (133)	37	* *	0	* *	0	* *
St Ignace	157	58.6 (164)	153	39.2 (160)	73	* *	0	* *	0	* *
St Johns W	358	54.2 (134)	309	29.8 (105)	123	32.5 (34)	0	* *	0	* *
St Louis C	139	50.4 (99)	167	34.7 (137)	51	* *	0	* *	0	* *
St.Clair S	1,029	48.1 (87)	313	19.2 (44)	90	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Stanton Cl	604	61.1 (170)	592	37.0 (155)	278	26.6 (80)	0	* *	0	* *
Stockbridg	61	* *	53	* *	24	* *	0	* *	0	* *
Sumpter He	939	43.6 (69)	845	26.3 (73)	248	31.9 (36)	0	* *	0	* *
Taylor	2,515	51.8 (114)	2,159	27.8 (90)	799	30.8 (44)	0	* *	0	* *
Teen Healt	13	* *	9	* *	6	* *	0	* *	0	* *
Temperance	77	* *	69	* *	26	* *	0	* *	0	* *
Thompsonv	13	* *	11	* *	5	* *	0	* *	0	* *
Three Oaks	132	62.9 (173)	122	40.2 (162)	56	* *	0	* *	0	* *
Three Rive	803	50.4 (99)	684	36.0 (150)	269	21.2 (101)	0	* *	0	* *
Traverse C	1,382	51.1 (104)	1,212	28.2 (93)	521	33.4 (29)	0	* *	0	* *
Tri-Cnty H	336	78.3 (179)	233	60.9 (172)	133	6.0 (109)	0	* *	0	* *
Tuscola	1,086	54.7 (136)	1,029	30.4 (107)	414	29.2 (56)	0	* *	0	* *
U Of M-Ped	0	* *	0	* *	0	* *	0	* *	0	* *
Vcms River	87	* *	78	* *	12	* *	0	* *	0	* *
Vicksburg	40	* *	59	* *	5	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Walled Lak	962	47.5 (83)	745	27.0 (83)	213	34.7 (19)	0	* *	0	* *
Warren Cli	1,535	39.3 (58)	407	14.7 (29)	86	* *	0	* *	0	* *
Washtenaw	85	* *	71	* *	22	* *	0	* *	0	* *
Wayne-Dear	2,257	9.3 (9)	1,557	3.0 (7)	87	* *	0	* *	0	* *
Webbervill	48	* *	41	* *	11	* *	0	* *	0	* *
West Genes	464	42.2 (64)	431	23.9 (63)	110	32.7 (32)	0	* *	0	* *
Western Wa	1,154	42.4 (65)	909	26.8 (79)	260	31.2 (40)	0	* *	0	* *
Westland C	1,613	48.2 (88)	1,004	22.5 (58)	341	30.8 (44)	0	* *	0	* *
Westside C	1,512	33.6 (41)	1,341	20.4 (50)	370	28.4 (66)	0	* *	0	* *
Westwood	32	* *	34	* *	3	* *	0	* *	0	* *
Wexford	798	53.3 (127)	675	34.1 (136)	303	27.4 (75)	0	* *	0	* *
Wheeler WI	23	* *	26	* *	9	* *	0	* *	0	* *
Whitmore L	49	* *	41	* *	18	* *	0	* *	0	* *
Willow	35	* *	25	* *	4	* *	0	* *	0	* *
Woodside	171	65.5 (176)	147	48.3 (170)	80	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wyoming Cl	1,503	45.2 (74)	1,393	23.3 (61)	447	35.6 (13)	0	* *	0	* *
Yale Clini	105	58.1 (158)	107	39.3 (161)	51	* *	0	* *	0	* *
Ypsilanti	2,671	32.3 (40)	2,261	19.0 (43)	505	32.7 (32)	0	* *	0	* *
Michigan	152,542	41.5	130,034	24.3	38,846	29.8	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Agape WIC	1,140	8.9(151)	7.2 (32)		1,177	0.4 (155)	1,132	8.9(151)	1,019	42.3(141)
Albion Cli	485	7.6(122)	9.5 (90)		492	0.0 (1)	485	7.6(122)	481	35.8(155)
Alcona	63	* * * *			128	0.8 (179)	63	* *	122	49.2(113)
Algonac Cl	281	7.1(103)	8.2 (51)		284	0.0 (1)	279	6.8 (96)	222	53.2 (87)
Allegan	238	4.6 (30)	12.2 (147)		257	0.0 (1)	235	4.7 (32)	244	54.5 (73)
Alma WIC	162	3.1 (5)	9.3 (86)		170	0.0 (1)	162	3.1 (5)	167	44.3(131)
Alpena Cou	509	5.1 (41)	13.2 (162)		510	0.2 (120)	506	4.9 (37)	504	56.0 (62)
Antrim - B	126	11.1(163)	6.3 (21)		163	0.0 (1)	126	11.1(162)	151	81.5 (10)
Arab/Chald	169	6.5 (87)	7.7 (38)		393	0.3 (143)	169	6.5 (86)	388	52.1 (96)
Arenac Cou	110	10.9(160)	5.5 (12)		113	0.9 (180)	108	11.1(162)	110	72.7 (18)
Associated	23	* * * *			23	* *	23	* *	23	* *
Baldwin	139	9.4(153)	5.8 (16)		150	0.7 (172)	139	9.4(153)	148	58.1 (51)
Bangor	794	7.1(103)	8.2 (51)		1,076	0.2 (120)	789	7.0(104)	1,032	51.1(102)
Baraga Cl	102	6.9(96)	16.7 (174)		103	0.0 (1)	102	6.9 (98)	102	51.0(103)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Barry-Eato	715	5.7 (65)	8.0 (47)	761	0.5 (163)	710	5.5 (58)	733	52.5 (94)
Barry-Eato	954	6.2 (77)	10.3 (112)	1,017	0.2 (120)	947	6.1 (75)	978	56.9 (56)
Battle Cre	2,256	5.5 (58)	9.4 (88)	2,306	0.0 (1)	2,248	5.5 (58)	2,127	41.1(144)
Baxter Com	48	* *	* *	48	* *	47	* *	46	* *
Bay Cl	1,773	5.3 (49)	11.8 (143)	1,814	0.1 (97)	1,765	5.3 (48)	1,740	50.0(106)
Bayside Wl	21	* *	* *	21	* *	21	* *	20	* *
Beaver Isl	6	* *	* *	6	* *	6	* *	5	* *
Belding Cl	218	7.8(126)	8.3 (55)	219	0.0 (1)	217	7.8(125)	216	39.8(147)
Bell Memor	223	5.4 (54)	11.7 (142)	227	0.0 (1)	223	5.4 (54)	208	55.3 (70)
Bendle/Bur	90	* *	* *	101	0.0 (1)	90	* *	98	* *
Benton Har	1,770	8.6(148)	6.7 (29)	1,775	0.0 (1)	1,766	8.6(148)	1,705	41.4(143)
Benzie Cli	255	8.2(139)	10.6 (120)	263	0.0 (1)	255	8.2(139)	248	70.6 (21)
Berrien -	766	6.0 (72)	11.2 (133)	784	0.1 (97)	765	5.9 (71)	751	52.1 (96)
Bldhd Migr	7	* *	* *	12	* *	7	* *	11	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Booth Clin	0	* *	* *		0	* *	0	* *	0	* *
Branch-Hil	739	7.4(117)	10.7 (124)		784	0.0 (1)	736	7.3 (114)	779	46.6(124)
Branch-Hil	718	6.3 (79)	9.5 (90)		723	0.0 (1)	718	6.3 (80)	684	45.6(127)
Branch-Hil	643	5.1 (41)	7.6 (36)		729	0.4 (155)	641	4.8 (34)	693	55.6 (65)
Browning-C	50	* *	* *		52	* *	50	* *	47	* *
Bryant Com	44	* *	* *		43	* *	43	* *	42	* *
Burton	1,402	5.4 (54)	9.7 (98)		1,632	0.2 (120)	1,393	5.5 (58)	1,577	41.0(145)
Calumet WI	197	4.6 (30)	10.7 (124)		214	0.0 (1)	196	4.6 (30)	203	59.6 (44)
Capac Clin	85	* *	* *		88	* *	85	* *	66	* *
Caro WIC S	10	* *	* *		12	* *	10	* *	12	* *
Cedar Spri	487	2.5 (3)	12.5 (153)		489	0.0 (1)	487	2.5 (3)	455	60.4 (39)
Charlevoix	372	5.6 (61)	9.9 (106)		405	0.7 (172)	369	5.4 (54)	366	65.0 (28)
Chass Heal	2,856	5.8 (68)	10.4 (115)		2,988	0.3 (143)	2,838	5.8 (67)	2,916	53.2 (87)
Cheboygan	488	4.9 (36)	10.5 (116)		491	0.0 (1)	488	4.9 (37)	481	59.3 (46)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Cherry Str	743	5.2 (47)	8.9 (72)	741	0.0 (1)	738	5.3 (48)	726	64.2 (29)
Chesaning	84	* *	* *	85	* *	84	* *	81	* *
Clare Coun	521	7.1(103)	7.3 (34)	536	0.7 (172)	514	7.2(110)	519	42.6(138)
Clinica Sa	827	4.0(17)	7.7 (38)	828	0.0 (1)	826	4.0 (17)	801	95.9 (3)
Comstock C	0	* *	* *	0	* *	0	* *	0	* *
Corner Hea	495	6.9(96)	7.7 (38)	497	0.0 (1)	494	6.9 (98)	458	44.8(130)
Crawford	265	3.8(14)	13.6 (166)	269	0.7 (172)	262	3.8 (14)	244	59.8 (42)
Dewitt WIC	121	3.3 (7)	6.6 (26)	156	0.6 (170)	121	3.3 (8)	152	53.9 (83)
Dickinson-	201	4.0(17)	15.4 (172)	205	0.0 (1)	199	4.0 (17)	200	57.0 (55)
Dickinson-	477	4.0(17)	14.7 (171)	485	0.0 (1)	475	4.0 (17)	484	54.5 (73)
DMC Prim	21	* *	* *	20	* *	20	* *	21	* *
DMC Prim	110	8.2(139)	4.5 (3)	117	0.0 (1)	110	8.2(139)	112	25.0(170)
DMC Prim	33	* *	* *	36	* *	33	* *	34	* *
Dorr Outre	254	4.3(25)	11.4 (137)	255	0.0 (1)	253	4.3 (25)	251	58.2 (49)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Douglas He	1,938	7.2(110)	8.0 (47)	1,990	0.2 (120)	1,934	7.2(110)	1,945	44.3(131)
Dowagiac	887	9.1(152)	9.0 (75)	911	0.2 (120)	881	9.1 (152)	701	61.1 (38)
Durand WIC	255	9.8(156)	9.4 (88)	261	0.0 (1)	255	9.8(156)	249	49.0(115)
East Area	589	8.7(150)	5.6 (14)	672	0.3 (143)	585	8.5(146)	650	20.0(175)
Eastside H	1,454	11.6(167)	4.0 (1)	1,468	0.2 (120)	1,447	11.6(166)	1,281	32.2(159)
Eau Claire	396	4.5 (28)	11.1 (132)	441	0.2 (120)	394	4.6 (30)	429	82.1 (9)
Emmet Coun	490	7.1(103)	8.4 (58)	584	0.5 (163)	489	7.2(110)	500	77.0 (15)
Esanaba C	733	5.0 (39)	13.5 (164)	741	0.0 (1)	733	5.0 (41)	506	65.4 (26)
Family Com	553	8.1(136)	7.2 (32)	611	0.7 (172)	549	8.0(132)	596	37.4(152)
Family Hlt	555	7.6(122)	7.9 (44)	590	0.0 (1)	554	7.6(122)	543	49.5(110)
Family Roa	825	11.8(169)	7.6 (36)	828	0.0 (1)	825	11.8(169)	770	25.6(169)
Farmington	102	2.9 (4)	12.7 (156)	112	0.0 (1)	101	3.0 (4)	112	61.6 (36)
Fife Lake	2	* *	* *	2	* *	2	* *	2	* *
First Pres	0	* *	* *	0	* *	0	* *	0	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Flint Fam	730	7.9(132)	7.4 (35)	757	0.0 (1)	729	8.0(132)	745	34.2(158)
Fruitport	18	* *	* *	18	* *	18	* *	17	* *
Gateway	726	13.1(174)	4.5 (3)	862	0.0 (1)	723	13.1(174)	826	21.4(173)
Genesee -	2,474	8.4(144)	7.9 (44)	2,701	0.2 (120)	2,450	8.4(144)	2,531	37.3(153)
Gladwin Co	409	5.4 (54)	11.2 (133)	444	0.0 (1)	409	5.4 (54)	432	60.2 (41)
Gogebic Cl	216	5.1 (41)	8.3 (55)	223	0.0 (1)	216	5.1 (43)	209	54.5 (73)
Golden	96	* *	* *	96	* *	96	* *	93	* *
Grace Prof	321	5.9 (71)	5.9 (19)	321	0.0 (1)	320	5.9 (71)	310	23.2(172)
Grace Ross	2,009	10.2(157)	5.6 (14)	2,056	0.0 (1)	2,005	10.2(157)	1,975	42.0(142)
Grand Have	602	6.5 (87)	10.6 (120)	640	0.3 (143)	599	6.5 (86)	617	53.6 (85)
Grand Trav	9	* *	* *	10	* *	9	* *	10	* *
Grant	283	6.0 (72)	12.7 (156)	292	0.0 (1)	282	6.0 (73)	286	54.2 (78)
Gratiot	1,618	11.5(166)	6.1 (20)	1,635	0.7 (172)	1,601	11.6(166)	1,616	27.9(166)
Greenville	331	8.2(139)	10.6 (120)	330	0.0 (1)	330	8.2(139)	314	48.1(120)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Hamtramck	274	8.0(134)	9.5 (90)	312	0.3 (143)	272	8.1 (136)	303	56.8 (57)
Hancock Cl	368	4.3 (25)	14.1 (169)	380	0.3 (143)	366	4.4 (26)	368	71.7 (19)
Hannahvil	39	* * * *		39	* *	39	* *	38	* *
Harper Gra	2,524	11.6(167)	5.8 (16)	2,850	0.5 (163)	2,505	11.7(168)	2,296	28.9(165)
Haslett Cl	119	7.6(122)	11.8 (143)	135	0.0 (1)	118	7.6(122)	127	77.2 (14)
Haslett St	26	* * * *		28	* *	26	* *	24	* *
Herman Kie	2,474	11.2(165)	5.4 (11)	2,616	0.1 (97)	2,464	11.2(165)	2,457	31.2(160)
Highland P	250	8.4(144)	6.4 (22)	269	0.0 (1)	249	8.4(144)	255	31.0(161)
Holland Cl	1,858	6.8 (95)	8.7 (66)	2,288	0.1 (97)	1,850	6.7 (94)	2,186	63.8 (30)
Holly WIC	69	* * * *		72	* *	68	* *	71	* *
Holton	67	* * * *		67	* *	67	* *	61	* *
Howard Cit	140	3.6 (13)	10.7 (124)	140	0.0 (1)	140	3.6 (13)	134	59.0 (47)
Hudsonv	438	5.5 (58)	12.6 (155)	506	0.0 (1)	438	5.5 (58)	488	57.4 (53)
Hurley Cli	432	10.4(159)	7.9 (44)	490	0.2 (120)	428	10.3(158)	481	29.1(164)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Huron Coun	587	5.3 (49)	13.5 (164)	591	0.0 (1)	586	5.3 (48)	571	42.4 (140)	
Icchn-Spar	214	6.1 (75)	8.9 (72)	239	0.4 (155)	213	6.1 (75)	233	81.1 (11)	
Ingham Cou	3,202	8.0 (134)	8.7 (66)	3,819	0.2 (120)	3,185	8.0 (132)	3,325	63.0 (32)	
Intervale	245	11.8 (169)	9.0 (75)	247	0.0 (1)	245	11.8 (169)	235	26.0 (168)	
Ionia Coun	733	7.4 (117)	8.5 (60)	735	0.0 (1)	732	7.4 (119)	722	45.3 (129)	
Iosco	323	5.3 (49)	11.5 (140)	443	0.5 (163)	320	5.3 (48)	426	40.6 (146)	
Ironwood	0	* * *	* *	0	* *	0	* *	0	* *	
Isabella C	781	6.4 (82)	9.9 (106)	796	0.4 (155)	777	6.4 (83)	770	66.6 (24)	
Ithaca Cli	292	4.8 (33)	8.6 (64)	328	0.0 (1)	291	4.5 (28)	316	49.1 (114)	
Ithaca WIC	0	* * *	* *	0	* *	0	* *	0	* *	
Jackson Co	2,980	7.2 (110)	9.0 (75)	3,031	0.1 (97)	2,967	7.1 (106)	2,963	52.0 (99)	
Jackson M	0	* * *	* *	0	* *	0	* *	0	* *	
Janes St A	947	11.0 (162)	5.8 (16)	951	0.1 (97)	946	11.0 (161)	878	27.9 (166)	
Joy-Greenf	1,885	8.1 (136)	8.3 (55)	1,958	0.2 (120)	1,879	8.1 (136)	1,915	49.5 (110)	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kalamazoo	784	7.8(126)	8.0 (47)	825	0.1 (97)	780	7.8(125)	634	52.7 (91)
Kalamazoo	2,097	7.1(103)	10.5 (116)	2,155	0.1 (97)	2,094	7.1 (106)	1,948	43.4(135)
Kalkaska	392	6.4 (82)	10.5 (116)	396	0.0 (1)	392	6.4 (83)	374	69.8 (22)
Kent	0	* * * *		0	* *	0	* *	0	* *
Kent Count	3,218	7.3(114)	10.1 (109)	3,325	0.1 (97)	3,207	7.3(114)	3,096	56.0 (62)
Kentwood	1,615	7.1(103)	9.8 (100)	1,652	0.1 (97)	1,611	7.1 (106)	1,573	56.1 (60)
Keweenaw	67	* * * *		67	* *	66	* *	65	* *
Kingsley	31	* * * *		33	* *	31	* *	33	* *
Lac Vieux	21	* * * *		20	* *	20	* *	21	* *
Lake Co	189	7.9(132)	10.1 (109)	189	0.0 (1)	187	8.0(132)	188	43.1(137)
Lake Odess	54	* * * *		55	* *	54	* *	55	* *
Lapeer	1,275	5.5 (58)	11.8 (143)	1,290	0.2 (120)	1,272	5.5 (58)	1,246	51.0(103)
Leelanu CI	135	4.4 (27)	15.6 (173)	155	0.0 (1)	135	4.4 (26)	153	77.8 (13)
Lenawee	0	* * * *		0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Leslie-Sto	64	* * * *		71	* *	64	* *	63	* *	
Lincoln Pa	621	3.9 (15)	9.5 (90)	637	0.5 (163)	618	3.9 (15)	596	45.6 (127)	
Livingston	889	6.1 (75)	13.7 (167)	904	0.1 (97)	888	6.1 (75)	817	65.4 (26)	
Lowell	116	10.3 (158)	7.8 (41)	121	0.0 (1)	116	10.3 (158)	112	54.5 (73)	
M-72 Migra	10	* * * *		19	* *	10	* *	17	* *	
Mancelona	183	3.3 (7)	8.7 (66)	189	0.0 (1)	182	3.3 (8)	181	61.9 (35)	
Manchester	25	* * * *		25	* *	25	* *	23	* *	
Manchester	6	* * * *		6	* *	6	* *	6	* *	
Manistee C	364	7.1 (103)	9.1 (79)	370	0.3 (143)	362	6.9 (98)	351	55.3 (70)	
Manistique	165	4.2 (23)	9.7 (98)	174	0.6 (170)	165	4.2 (23)	167	46.7 (123)	
Mansitee M	20	* * * *		32	* *	19	* *	31	* *	
Marion Bra	51	* * * *		105	0.0 (1)	51	* *	103	51.5 (100)	
Marquette	362	3.9 (15)	13.0 (159)	378	0.0 (1)	361	3.9 (15)	341	66.0 (25)	
Marshall C	250	5.2 (47)	9.2 (84)	265	0.0 (1)	250	5.2 (46)	258	49.6 (109)	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Mason Clin	90	* *	* *		102	0.0 (1)	89	* *	92	* *
Mason Co	480	6.0 (72)	11.0 (130)		489	0.0 (1)	480	6.0 (73)	472	56.1 (60)
Mccree Hea	2,980	9.5(154)	5.3 (8)		3,067	0.2 (120)	2,972	9.5(154)	2,586	34.4(157)
Mcnamara H	154	7.8(126)	11.0 (130)		153	0.0 (1)	153	7.8(125)	137	48.9(116)
Mecosta Co	795	4.8 (33)	12.2 (147)		804	0.2 (120)	795	4.8 (34)	787	57.1 (54)
Medical Pr	36	* *	* *		36	* *	36	* *	35	* *
Medical Pr	77	* *	* *		81	* *	77	* *	74	* *
Menominee	333	6.3 (79)	10.2 (111)		368	0.0 (1)	333	6.3 (80)	243	52.3 (95)
Mercy	99	* *	* *		108	0.9 (180)	98	* *	104	17.3(176)
Mesick	61	* *	* *		61	* *	61	* *	57	* *
Metropolit	0	* *	* *		0	* *	0	* *	0	* *
Michigan S	176	5.1 (41)	9.1 (79)		204	0.0 (1)	175	5.1 (43)	199	95.5 (4)
Midland Cl	1,079	6.6 (90)	8.8 (70)		1,085	0.0 (1)	1,079	6.6 (90)	819	78.9 (12)
Midwest	881	8.2(139)	8.2 (51)		962	0.5 (163)	870	8.2(139)	948	34.5(156)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Migrant H	215	3.3 (7)	9.8 (100)	215	0.0 (1)	215	3.3 (8)	211	74.4 (17)
Missaukee	202	4.0 (17)	14.4 (170)	202	0.0 (1)	201	4.0 (17)	187	63.6 (31)
Moms & Bab	867	8.4 (144)	7.0 (31)	918	0.3 (143)	862	8.5 (146)	894	37.1 (154)
Monroe Cou	1,840	6.9 (96)	7.8 (41)	1,882	0.1 (97)	1,832	6.9 (98)	1,808	43.9 (134)
Montague/	88	* *	* *	88	* *	88	* *	79	* *
Montcalm	17	* *	* *	18	* *	17	* *	18	* *
Montmorenc	153	7.8 (126)	8.5 (60)	155	0.0 (1)	151	7.9 (131)	151	55.6 (65)
Mt Clemens	961	6.6 (90)	11.2 (133)	985	0.2 (120)	958	6.6 (90)	972	96.1 (2)
Munising C	140	6.4 (82)	11.4 (137)	144	0.7 (172)	138	6.5 (86)	141	48.9 (116)
Muskegon C	3,911	7.3 (114)	8.5 (60)	3,975	0.1 (97)	3,904	7.3 (114)	3,735	38.5 (151)
New Center	21	* *	* *	22	* *	21	* *	22	* *
New Haven	124	8.1 (136)	13.7 (167)	129	0.0 (1)	123	8.1 (136)	130	95.4 (5)
New Hope	11	* *	* *	12	* *	11	* *	11	* *
Newaygo	644	6.7 (93)	9.8 (100)	664	0.0 (1)	642	6.7 (94)	629	60.3 (40)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Newberry C	147	4.8 (33)	8.8 (70)	152	0.0 (1)	147	4.8 (34)	146	56.2 (59)
North Camp	263	3.4 (10)	9.1 (79)	281	0.4 (155)	257	3.1 (5)	280	91.8 (6)
North Midl	0	* *	* *	0	* *	0	* *	0	* *
Northeast	2,227	11.1(163)	4.8 (6)	2,442	0.2 (120)	2,212	11.1(162)	2,302	30.9(162)
Northern M	120	1.7 (1)	12.5 (153)	121	0.0 (1)	120	1.7 (1)	106	67.9 (23)
Northwest	364	12.1(171)	5.5 (12)	411	0.0 (1)	363	12.1(171)	398	24.9(171)
Oakland Co	3,199	7.8(126)	8.6 (64)	3,314	0.1 (97)	3,189	7.8(125)	3,254	48.6(118)
Oceana Co	467	4.7 (32)	10.3 (112)	480	0.0 (1)	466	4.7 (32)	462	53.5 (86)
Ogemaw	357	5.3 (49)	9.2 (84)	435	0.0 (1)	356	5.3 (48)	422	49.5(110)
Olhsa Star	249	7.2(110)	6.4 (22)	283	0.4 (155)	248	6.9 (98)	277	48.4(119)
Olhsa WIC	88	* *	* *	103	0.0 (1)	88	* *	99	* *
Omni Care	35	* *	* *	36	* *	35	* *	34	* *
Omnicare	4	* *	* *	4	* *	4	* *	4	* *
Omnicare-G	0	* *	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
Omnicare-M	71	* *	* *		78	* *	70	* *	74	* *
Ontonagon	64	* *	* *		66	* *	64	* *	64	* *
Oscoda	130	6.2 (77)	12.3 (149)		144	0.0 (1)	130	6.2 (79)	139	59.7 (43)
Otsego Co	488	5.1 (41)	9.8 (100)		524	0.4 (155)	484	5.2 (46)	436	71.6 (20)
Park Clin	2	* *	* *		2	* *	2	* *	2	* *
Park Medic	28	* *	* *		29	* *	28	* *	28	* *
Park Medic	55	* *	* *		55	* *	55	* *	52	* *
Park Medic	10	* *	* *		11	* *	10	* *	11	* *
Paw Paw Cl	265	4.2 (23)	10.9 (129)		369	0.0 (1)	264	4.2 (23)	362	49.7 (107)
PCOC	96	* *	* *		101	0.0 (1)	96	* *	89	* *
Pinconning	93	* *	* *		96	* *	93	* *	87	* *
Port Huron	2,042	7.0 (101)	5.3 (8)		2,066	0.0 (1)	2,040	7.0 (104)	1,640	47.6 (121)
Portland	89	* *	* *		89	* *	89	* *	88	* *
Presque Is	180	5.6 (61)	8.9 (72)		183	0.0 (1)	180	5.6 (63)	179	56.4 (58)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
	Number	% (Rank)	% (Rank)							
Profession	68	* *	* *	68	* *	68	* *	62	* *	
Pullman	529	4.0 (17)	5.3 (8)	594	0.3 (143)	528	4.0 (17)	569	54.0 (81)	
Ravenna	27	* *	* *	28	* *	27	* *	24	* *	
Reed City	350	5.4 (54)	13.1 (161)	367	0.3 (143)	349	5.4 (54)	323	59.4 (45)	
Region II	1,528	5.8 (68)	6.5 (24)	1,554	0.1 (97)	1,524	5.8 (67)	1,538	46.4 (125)	
Riverview	83	* *	* *	86	* *	82	* *	86	* *	
Romulus Fi	116	8.6 (148)	6.9 (30)	125	0.0 (1)	116	8.6 (148)	124	55.6 (65)	
Roosevelt-	128	12.5 (172)	4.7 (5)	129	0.0 (1)	128	12.5 (172)	120	20.8 (174)	
Roscommon	385	5.7 (65)	10.6 (120)	390	0.0 (1)	385	5.7 (65)	372	52.7 (91)	
S & S Medi	27	* *	* *	28	* *	27	* *	28	* *	
Saginaw Co	3,002	7.3 (114)	8.2 (51)	3,010	0.2 (120)	2,990	7.3 (114)	2,840	38.7 (150)	
Saginaw-Ch	31	* *	* *	32	* *	31	* *	31	* *	
Sanilac Co	761	7.5 (121)	6.6 (26)	766	0.0 (1)	757	7.5 (121)	711	54.4 (77)	
Sault Ste	554	3.1 (5)	13.4 (163)	569	0.2 (120)	552	3.1 (5)	544	62.3 (34)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Sawyer Afb	156	10.9(160)	9.6 (95)	160	0.0 (1)	156	10.9(160)	146	58.2 (49)	
School Bas	0	* * *	* *	0	* *	0	* *	0	* *	
Schoolcraf	1,330	9.5(154)	4.1 (2)	1,352	0.1 (97)	1,328	9.5(154)	1,209	39.1(148)	
Seton Cent	706	12.9(173)	5.2 (7)	765	0.1 (97)	701	12.8(173)	736	30.6(163)	
Shaftsburg	73	* * *	* *	74	* *	72	* *	71	* *	
Shelby M	233	1.7 (1)	11.6 (141)	270	0.0 (1)	233	1.7 (1)	269	55.8 (64)	
Sheldon Co	1,595	7.4(117)	6.5 (24)	1,642	0.0 (1)	1,594	7.4(119)	1,542	44.3(131)	
Shiawassee	954	4.9 (36)	10.0 (108)	967	0.1 (97)	950	4.9 (37)	937	53.0 (89)	
Southfield	1,965	7.8(126)	9.3 (86)	2,151	0.3 (143)	1,958	7.8(125)	2,093	55.4 (69)	
Sparta Pub	296	5.1 (41)	10.8 (128)	296	0.0 (1)	296	5.1 (43)	262	58.4 (48)	
St Charles	145	8.3(143)	12.4 (152)	148	0.0 (1)	145	8.3(143)	141	47.5(122)	
St Ignace	155	4.5 (28)	12.9 (158)	155	0.0 (1)	155	4.5 (28)	151	55.6 (65)	
St Johns W	277	4.0 (17)	9.0 (75)	331	0.0 (1)	276	4.0 (17)	328	54.0 (81)	
St Louis C	171	7.0(101)	12.3 (149)	178	0.0 (1)	170	7.1(106)	175	49.7(107)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
St.Clair S	371	6.5 (87)	10.5 (116)	378	0.0 (1)	371	6.5 (86)	374	97.3 (1)
Stanton Cl	644	6.4 (82)	8.5 (60)	642	0.0 (1)	638	6.4 (83)	612	52.6 (93)
Stockbridg	51	* *	* *	56	* *	51	* *	49	* *
Sumpter He	817	8.4(144)	9.5 (90)	858	0.3 (143)	807	8.6(148)	779	45.8(126)
Taylor	2,186	7.4(117)	9.6 (95)	2,344	0.5 (163)	2,173	7.3(114)	2,184	39.0(149)
Teen Healt	9	* *	* *	9	* *	9	* *	6	* *
Temperance	69	* *	* *	70	* *	68	* *	67	* *
Thompsonv	11	* *	* *	11	* *	11	* *	9	* *
Three Oaks	121	5.8 (68)	6.6 (26)	126	0.0 (1)	121	5.8 (67)	119	52.9 (90)
Three Rive	719	6.4 (82)	7.8 (41)	725	0.1 (97)	717	6.3 (80)	669	43.3(136)
Traverse C	1,170	6.6 (90)	11.8 (143)	1,229	0.2 (120)	1,162	6.6 (90)	1,187	75.1 (16)
Tri-Cnty H	483	3.5 (12)	13.0 (159)	488	0.0 (1)	483	3.5 (12)	484	51.2(101)
Tuscola	1,061	5.6 (61)	9.8 (100)	1,087	0.2 (120)	1,057	5.6 (63)	1,046	54.2 (78)
U Of M-Ped	0	* *	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)											
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed		
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%	
	Number	% (Rank)	% (Rank)	(Rank)							(Rank)
Vcms River	91	* *	* *		96	* *		91	* *	94	* *
Vicksburg	93	* *	* *		95	* *		93	* *	84	* *
Walled Lak	789	5.3 (49)	9.6 (95)		794	0.0 (1)		787	5.3 (48)	792	61.2 (37)
Warren Cli	487	5.7 (65)	11.3 (136)		507	0.0 (1)		486	5.8 (67)	501	91.4 (7)
Washtenaw	67	* *	* *		74	* *		67	* *	69	* *
Wayne-Dear	1,437	5.6 (61)	8.1 (50)		1,628	0.1 (97)		1,431	5.7 (65)	1,579	87.1 (8)
Webbervill	42	* *	* *		47	* *		42	* *	42	* *
West Genes	471	7.2(110)	9.1 (79)		530	0.2 (120)		469	7.2(110)	434	52.1 (96)
Western Wa	943	6.3 (79)	8.7 (66)		954	0.4 (155)		931	6.1 (75)	918	42.5(139)
Westland C	1,254	7.7(125)	9.1 (79)		1,291	0.2 (120)		1,251	7.8(125)	1,125	53.7 (84)
Westside C	1,357	4.9 (36)	10.3 (112)		1,408	0.0 (1)		1,355	4.9 (37)	1,329	57.9 (52)
Westwood	48	* *	* *		51	* *		47	* *	46	* *
Wexford	726	5.0 (39)	12.3 (149)		730	0.1 (97)		724	5.0 (41)	686	62.4 (33)
Wheeler WI	33	* *	* *		34	* *		33	* *	33	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2001 - 2003 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Whitmore L	38	* *	* *	41	* *	36	* *	40	* *
Willow	25	* *	* *	33	* *	25	* *	30	* *
Woodside	149	3.4 (10)	11.4 (137)	153	0.0 (1)	149	3.4 (11)	147	50.3 (105)
Wyoming Cl	1,435	6.9 (96)	10.7 (124)	1,446	0.1 (97)	1,432	6.8 (96)	1,306	54.8 (72)
Yale Clini	102	6.9 (96)	9.8 (100)	107	0.0 (1)	102	6.9 (98)	94	* *
Ypsilanti	2,306	6.7 (93)	8.4 (58)	2,394	0.1 (97)	2,292	6.6 (90)	2,275	54.2 (78)
Michigan	141,396	7.3	8.8	149,132	0.2	140,821	7.3	139,580	49.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

