



3 0000 004 809 707

LIBRARY OF MICHIGAN

JOINT DOCUME

render Michigan what she ought to be, rich, prosperous and happy.

I commit the whole subject to your deliberate consideration, gentlemen, and pray that this communication may be considered as for the use of both houses.

WILLIAM WOODBRIDGE.

Executive Office, January 27, 1841.

Resolutions.

At a general assembly of the state of Connecticut, holden at New Haven, in said state, on the first Wednesday of May, in the year of our Lord one thousand eight hundred and forty:

Resolved by this Assembly, That the public lands belonging to the United States, are the common property of all the states, and not of the particular states in which they lie; and that neither said lands or the proceeds thereof, ought to be appropriated for any purpose except for the benefit of all the states.

Resolved, That the proceeds of the public lands, not required for the payment of the debts of the government, or for other public purposes connected with the administration thereof, should of right be divided among the several states of this Union.

Resolved, That this assembly earnestly protest against any reduction in the prices of the public lands as now fixed by law, regarding such a measure as unnecessary and unjust.

Resolved, That this assembly do request their senators and representatives in congress, to use their exertions to sustain the principles contained in these resolutions.

Resolved, That his excellency the governor, be requested to forward copies of the foregoing resolutions to the senators and representatives in congress from this state, and also to the executives of the several states, that they may be communicated to the legislatures of their states respectively.

State of Connecticut, Secretary's Office, ss:

I hereby certify that the above is a true copy of record in this office. In testimony of which, I have hereto set the seal of this state, and signed the same.

ROYAL R. HINMAN,
Secretary of State.

[No. 11.]

Annual report of the state geologist.

OFFICE OF STATE GEOLOGIST, }
 Detroit, February 1, 1841. }

To the Senate and House of Representatives:

In conformity with instructions contained in the act authorizing a geological survey of the state, I have the honor to lay before you, an outline of the operations of the department over which I have been placed, together with the general progress towards completion of the whole work.

It is a matter of regret to me, that the sufferings and hardships to which I have been exposed in conducting the field work over the wilderness portions of our state, have so far impaired my health, as to render it impossible for me to enter into so minute details as had been anticipated. I regret this the more, since it leaves many wide spaces in portions of the present report, which are of much consequence to a proper understanding of the whole. But since the annual reports are intended to refer rather to the progress of the work than to its results, and since the whole will be embraced in a more perfect form hereafter, this defect is of less importance than it otherwise would be.

My individual labors, during the past season, have been chiefly devoted to surveys connected with the northern slope of the upper peninsula, and to this district, the chief observations in this report, will be directed.

UPPER OR NORTHERN PENINSULA.

General description and Topographical features.

The published maps of that portion of the state of Michigan usually known as the upper peninsula, are so defective, not only in details, but also in general outline, that the task of giving a description of any portion, in such a manner as to render it intelligible to any person who has not actually traveled over it, is exceedingly difficult. The extent of these geographical inaccuracies is much greater than would at first be supposed, for scarcely a single feature of the interior is given as it actually exists; mere brooks are magnified to rivers, and again those streams justly deserving the name of rivers, are either wholly omitted or scarcely noted, while the courses of the streams, as laid down, are almost invariably as far from the truth as could be conceived. Nor do the inaccuracies stop here, for even the coast maps of the great lakes, by which our upper peninsula is in part surrounded, are usually so defective

as scarcely to be recognized, except in their most general outlines.*

I have already, in a previous report, referred in general terms, to that portion of the upper peninsula bounded by lakes Huron and Michigan, for which reason my remarks at this time will be mostly confined to the northerly portion of the peninsula, or that portion bordering upon Lake Superior.

So little is known of the extent of country constituting the upper peninsula, that it may not be misplaced to make some reference to its dimensions, though at this time most of the estimates must be regarded as mere approximations. The most extreme length of the district is embraced between Point de Tour, of Lake Huron, on the east, and the mouth of Montreal river, of Lake Superior, on the west. From Point de Tour, the direction of the mouth of Montreal river is very nearly north, 80° 30' west, and the direct distance does not vary far from 316 miles. This estimate, it will be perceived, does not include Drummond's Island, which, if included, would add some 20 miles to the length as already given.

The easterly portion of the peninsula is narrow, and its width, for a distance of 130 miles west from Point de Tour, varies from 30 to 50 miles; west from this, the peninsula widens rapidly, though its width is exceedingly irregular. I am unable, at this time, to state with very much accuracy, the extreme width of the upper peninsula, but the area of the whole may be estimated at very nearly two-thirds that of the lower, or southern peninsula.

The topography and general features of the upper and lower peninsulas differ so widely from each other, that, with the simple exception of a part of the easterly extremity of the upper peninsula, they scarcely admit of a comparison. The wide contrast exhibited by the two districts, is wholly dependent upon geological differences, and these are so strongly marked, that they could not fail to attract the notice of the most superficial observer.

In the last report I had the honor to lay before you, some general references were made to the topography of the southern slope of the upper peninsula, which embraced simply those portions bordering upon lakes Huron and Michigan, and extending from Point de Tour to Menominee river.

Although the rocks of the district extending from Point de Tour to Chocolate river, upon the northerly or Lake Superior slope, belong to an older series than those lying south, and are different in composition, the general features of the two dis-

* I am happy in being able to except from this otherwise universal charge of inaccuracy, the coast map of Lake Superior published by the society for the diffusion of useful knowledge. This map was reduced from the surveys of Capt. Bayfield, R. N., as returned to the British admiralty office, and so far as the British coast is concerned, the map is minutely correct. The American coast upon this map is faithfully delineated in its general outline, but in minutiae it is frequently deficient.

districts, nevertheless, bear a close resemblance. Easterly from Point Iroquois, the country is for the most part flat, or but slightly elevated, and the near approach of the rock to the surface so far prevents the descent of the waters, as to give rise to extensive districts of wet and swaly land. Westerly from Point Iroquois to Chocolate river, the country is more elevated, and has a much smaller proportion of wet land. A range of hills, having an elevation varying from 300 to 600 feet above Lake Superior, commences a little easterly from Point Iroquois, and stretches very nearly west, or but a few degrees north of west, until the western escapement again appears upon the coast, giving rise to the elevated hills of which the Pictured rocks and Grand island form a part. The outline of this range of hills has the most perfect regularity, being unbroken and uniformly covered with a dense growth of timber.

West from Chocolate river, to our boundary line at Montreal river, the physical character of the country is widely different from that of the district before referred to. This country is made up of a series of irregular, knobby ranges of hills, that have a general easterly and westerly direction, with intervening valleys of flat or gently rolling land. These hills not unfrequently rise to a height of from 600 to 900 feet, very near to the immediate coast of Lake Superior, and at a distance of 15 to 20 miles south from the coast, portions of some of the ranges rise to a height of 1,200 to 1,300 feet above the level of that lake. The ragged and broken outline which this district presents, when viewed in detail, from the lake, contrasts in a striking manner, with that of the country lying east from Chocolate river; for, instead of the regular and unbroken range of hills, uniformly covered with a dense forest, that occur in the latter district, we have a series of ranges of broken hills, with knobs not unfrequently nearly or quite destitute of timber. The escapements of these hills are sometimes so abrupt as to render them difficult of ascent.

The only exception to the general easterly and westerly direction of these ranges of hills, occurs in that range constituting the Porcupine mountains. These mountains rise somewhat abruptly almost upon the immediate coast of Lake Superior, at a point 37 miles north-easterly from the mouth of Montreal river, and from this point they stretch inland, in a direction which, for the first 30 miles, is very nearly south-south-west, after which their course is more westerly, and in the direction of the sources of the Wisconsin river. The most elevated points of the Porcupine mountains, near to Lake Superior, attain an altitude of very nearly 950 feet, but several of the knobs, at a distance inland, rise from 1,000 to 1,300 feet above the level of the waters of that lake.

The valleys, before referred to as separating these ranges

of hills, are uniformly heavily timbered, and by far the largest proportion of this timber is beech and maple.

The length of the hilly or mountainous district, estimating in a direct line west from Chocolate river to the boundary line on Montreal river, is very nearly one hundred and sixty miles, and it does not probably extend, at any point, more than 20 to 25 miles south from this line. Estimating this hilly district to extend regularly 20 miles south of a line drawn from the points before mentioned, the greatest width of the district would be opposite Keweenaw point, which extends 67 miles north from this line, making the total width at this point 87 geographical miles. The very great irregularities of the coast, with the numerous deep bays and projecting points upon the north, together with the irregularities of the ranges of hills upon the south, cause so great variations in the width of the district, that it is impossible, with the present information upon this subject, to estimate the width of the district with any great degree of accuracy. Keweenaw bay, of Lake Superior stretches 60 miles, estimating from the extremity of Keweenaw point, into this hilly or mountainous country.

South from the range of hilly country alluded to, and extending to Green bay, the country at first becomes more level and finally flat, though with several regular and unbroken ranges of hills. In topography and general character, it more nearly resembles that district of country which lies east from Chocolate river.

Of the district of country lying between the hilly country and Green bay, less is known than of any other portion of the upper peninsula. The extent of my duties did not permit me to extend my examinations very far into it, nor was I enabled to obtain any information as to its general character.

The streams which discharge their waters into Lake Superior upon its south shore, are invariably short, and with very few exceptions, the quantity of water they discharge is small. This remark, in fact, may apply to the whole of the region of country surrounding that lake, for this immense body of water is completely surrounded by hills that, at no great distance from the lake, fall away more or less rapidly. Thus, while many of the streams discharging their waters into Lake Michigan, Green bay and the Mississippi river, have their sources near to the south shore of Lake Superior, so also, many of those streams which discharge their waters northerly into Hudson's bay, have their sources near to the north coast of the lake. The near approach of the summit of the ranges of hills surrounding the lake, to the immediate coast, leaves the

area of country draining into Lake Superior, comparatively small.

The most important of the streams entering Lake Superior upon its south shore and within the limits of our state, (commencing near the foot of the lake and enumerating westwardly,) are the Tequoimenon, Train, Chocolate, Death, Yellow dog, Huron, Portage, Fire Steel, Ontonagon, Iron, Presque Isle, Black and Montreal rivers. Besides these, there are innumerable creeks, which are usually known to voyageurs as rivers, for this term is applied indiscriminately to all. The waters of most of these streams are remarkably transparent and pure, with brisk currents and numerous cascades, and they almost invariably contain an abundance of the brook trout, a circumstance which I mention, from the fact that this fish is scarcely known in the streams of the southern peninsula.

The Tequoimenon river, which is the only stream east from Chocolate river, that in reality breaks through the range of sandstone hills, before mentioned as extending westerly from Point Iroquois, has its embouchure about 18 miles south from Whitefish point, and near the foot of the lake. The discharge is through loose sands, and there is an average of 4 to 4½ feet water over the bar. Having passed the bar the water for a distance of 7 to 8 miles varies from 10 to 15 feet.

Some of the sources of this stream approach very nearly to Lake Michigan, being directly at the base of the range of lime rock hills, referred to in my third annual report.

The Tequoimenon river, with the exception of a distance of some four to five miles, while passing through the range of sandstone hills before alluded to, is, through its whole course, a sluggish stream, though at many points having a strong, deep current. The character of the river, in its passage through the range of hills referred to, is totally changed, for it has there numerous chutes and falls, with almost continuous rapids. At one point, the whole body of water contained in the stream is precipitated by a single leap, from a height of forty-six feet, and the effect of this fall is much heightened by the elevated and overhanging rocks that bound the river upon either side.

Most of the small streams, discharging into Lake Superior between the foot of the lake and Chocolate river, have their sources to the north of the elevated range of hills mentioned, or minor branches only descend from those hills.

Chocolate river, which discharges its waters into Lake Superior at a point 146 miles very nearly due west from the Saut de Ste Marie, is a stream of considerable magnitude, though in consequence of the loose sands at its mouth, it is difficult of entrance at ordinary stages of water, even with barges of mo-

derate draught, but when once the bar is passed, the stream is found to be deep, and for several miles has a width varying from 80 to 150 feet.

This stream will be made a point of reference in the strictly geological portions of this report, for it winds along near the line of junction of two widely distinct geological districts; the general features of one being characterized by its ragged and broken hills, while the other is not less marked by its generally level or regularly undulating surface.

Chocolate river takes its name from the dark color of its waters.

Those streams which occur between Chocolate river and Keweenaw bay, are, with the exception of Huron river, small; though were we to refer to the published maps of this district, we would suppose that some of these streams were of very considerable length. But with the exception of Huron river and River Des Morts, I believe they all have their sources in small lakes lying along the bases of the elevated hills already described. These hills rarely recede farther than three to five miles from the coast, and the length of the streams, forming the outlets of the lakes referred to, is governed wholly by these features.

Ontonagon river, which is one of the most important of the rivers discharging its waters into Lake Superior, upon its south shore, has its embouchure very nearly fifty-five miles east, or rather north-easterly, from the western boundary of the state, at the mouth of Montreal river. Ontonagon river has its sources in a very great number of mountain lakes, situated in part upon the south-easterly spurs of the Porcupine mountains, and in part in the hilly district formed by the easterly and westerly ranges of hills before described; which ranges, upon this portion of the coast, curve very considerably to the south. Some of the sources of the Ontonagon river approach very near to the sources of the Chippewa river of the Mississippi. The great number of small tributaries of the Ontonagon, which are simply the outlets of the small lakes referred to, are concentrated into two principally branches that finally unite and form the principal river, at a distance of about eighteen miles from the coast of Lake Superior. The smaller tributaries are mostly mere torrents, with frequent perpendicular falls, and high banks, sometimes, of precipitous rock. The main stream, from the junction of the two principal branches to within five or six miles of the lake, is rapid and shoal, but below this, the stream is comparatively still, and with a good depth of water. The mouth of the Ontonagon river is obstructed by a bar of sand,

over which there is usually, at low stages, about six feet of water.

The principal rivers west from Ontonagon river and within the limits of our state, have already been stated to be Iron, Presque Isle, Black and Montreal rivers. These streams are all short, and the amount of water discharged by each separately, is comparatively small. Their waters descend from the elevated mountain region immediately south from the coast, and since the whole streams are concentrated before passing from these elevations, their waters are discharged in body and they descend with very great rapidity. A greater variety of grand and beautiful scenery than that presented by some of these streams in their descent to the lake, taken in connection with the rugged and wild character of the country, can scarcely be conceived. I was particularly struck with the great variety of picturesque views furnished by Black river, in its descent from the elevated country on the west side of the Porcupine mountains to Lake Superior. The stream was estimated to fall about five hundred feet in a fraction over four miles, and this descent is made up by a constant succession of falls, chutes and rapids, which continue with so little interruption that the waters for the whole distance may be said to be constantly white foam. The stream is bounded upon either side by banks elevated from one hundred to three hundred feet, sometimes sloping away from the stream, somewhat gently, and again rising in mural precipices of rock, separated from each other by so short distances as to appear scarcely sufficient to permit the passage of the waters of the river. The most considerable fall does not exceed fifty feet, and they are usually from ten to thirty feet in height, but their constant succession and variety, add much to its interest.

Montreal river is a comparatively small stream, made up of numerous small tributaries, that rise among the ranges of hills to the south-west and south-east of its mouth. The passage of the river through the range of hills near the lake, gives rise to several very considerable water-falls, as also to much rugged and wild scenery. Almost directly at the place of embouchure into Lake Superior, there is a perpendicular fall of about forty feet. This stream, it will be recollected, forms a portion of the boundary between Michigan and Wisconsin.

By the act admitting Michigan as a state into the confederacy, and in which her boundaries are defined, it does not appear to have been the intention to include within her limits any portion of territory lying upon the north shore of Lake Superior, but in consequence of the peculiar shape of the coast at that point where the *national* boundary line "last touches Lake

Superior," at the mouth of Pigeon river, a direct line to the mouth of the Montreal river, if followed literally, would throw within the state of Michigan several small rocky islands, together with a few miles of the south cape of Pigeon bay, situate upon the north coast. This boundary leaves in Wisconsin the whole of the Apostles' group of islands, near to the south coast, while it includes within Michigan, Isle Royale, situate near to the north coast of the lake.

Isle Royale is little less than an island of rock, rising abruptly from the lowest depth of the lake, in irregular hills, to a height varying from 100 to 450 feet above the level of the lake. The island has a length of a fraction over 45 miles from north-east to south-west, and a breadth varying from $3\frac{1}{2}$ to 8 miles. The most northerly point of the island is very nearly in latitude $48^{\circ} 12' 30''$ north, and the parallel of longitude 89° west from Greenwich, crosses the island a little east from its centre. Its nearest approach to the main land is near its north-westerly end, where it is separated from a point of the north coast, a few miles east from Pigeon river, by a distance of a fraction less than 13 miles. Isle Royale is separated from Keweenaw point, of the south coast, by a distance of 44 miles, and the elevated hills of this point may be distinctly seen from Isle Royale, when the atmosphere is clear.

Nearly the whole of the north-westerly side of Isle Royale is a continuous, elevated, rocky cliff, which will scarcely admit of a landing, but the south-easterly side, together with the easterly and westerly ends, are deeply indented with bays, which form secure harbors. The north-easterly end is made up of a series of elevated, rocky spits, with intervening bays. These spits of rock continue for a length varying from 10 to 12 miles, with a width scarcely exceeding half a mile, and altogether, they may not inaptly be compared to the hand with the fingers half spread. The bays have a sufficient depth of water to admit vessels of the largest class to enter nearly one third the whole length of the island.

Much of Isle Royale is absolutely destitute of soil, and the island has a most desolate appearance; but notwithstanding this, it is of immense value for its fisheries, which are as yet scarcely appreciated.

Though not within the limits of our state, I will briefly refer to the general character of a portion of the country west from Pigeon river, on the north coast. That district of country upon the immediate coast, extending from our national boundary, at Pigeon river, to Fond du Lac, is more decidedly and abruptly mountainous than any portion of the south coast of the lake. The hills rise in broad and somewhat knobby steppes or plateaus, to heights varying from 400 to 1,200 feet above the lake,

and the summits of these hills are usually not farther inland than from 10 to 20 miles. The rocks of the hills are very frequently bare over considerable areas, and the valleys containing arable soil, are few and very narrow.

The route of the fur trade to the north-west, via Rainy lakes, Lake of the Woods and Lake Winnepic, was formerly wholly carried on by passing over these hills, from a point a few miles west from the mouth of Pigeon river. The trail or portage path passes over a low portion of the range, and finally falls upon Pigeon river, which is ascended to its source, from which, by a series of portages, the sources of the streams flowing north-westerly are reached. The hilly portion of the country, though of exceeding interest in a geological point of view, is the most desolate that could be conceived.

GENERAL GEOLOGY OF THE UPPER PENINSULA.

The geology of the upper or northern peninsula of Michigan, when compared with that of the southern or lower peninsula, bears a striking contrast; for while that of the district last referred to is uniformly regular, with rocks which, though rarely exposed to view, are few and for the most part but little disturbed, over large areas of country, the upper peninsula embraces a much greater number of rocks, distributed over a somewhat smaller district of country, and a portion of which are so much disturbed, as to render their delineation exceedingly complex and difficult.

The widely different topographical features of the easterly and westerly portions of the northerly part of the upper peninsula would lead the most casual observer to infer that the geological features of the different districts would be equally distinct, and in this he would not be disappointed.

I have already referred to the rugged and broken character of the country, extending westerly from Chocolate river to our boundary at Montreal river, and have also attempted to define its general length and breadth. This district, which is essentially made up of primary, trap and metamorphic rocks, with intervening sedimentary rocks, usually occupying the valleys and out boundaries, may be estimated to cover an area equal to a little more than one-fourth of the whole of the upper peninsula. To the east and south of this district the rocks are wholly sedimentary, consisting of a series of sandstones, limestones and shales.

With a view of rendering the local details of the separate formations more intelligible, I will first describe, in general terms, the rocks occurring in so much of the peninsula as has been examined, together with their general extent. This will necessarily involve a repetition of a very small portion of the

report last made, upon the subject of the limestones of the south and east portions of the peninsula, but since the examinations of the past year have enabled me to add another member to the limestone group, and to define with more certainty its outline, this may not be devoid of interest.

1. *Primary Rocks.*—The rocks constituting what may be considered as the true primary group of this region, are chiefly granite, syenite and syenitic granites. The members of the group are first seen upon the south coast of Lake Superior, constituting a rocky point known as Little Presque Isle, a little south-east from Riviere Des Morts. These rocks frequently appear upon the coast north-westerly, nearly as far as Huron river, and the Huron islands off the mouth of Huron river belong to the same group. West from Huron Islands, no rock appears upon the coast which, in a strict sense, I should regard as primary. The rocks of this group arise upon or near to the coast, in irregular and broken ranges of hills, to a height varying from 300 to 700 feet above the waters of the lake, and these hills or ranges of hills are continued in a south-westerly direction. The precise limit of the primary rocks to the westward, has not yet been determined, but they are known to extend nearly or quite to the sources of the Wisconsin river.

A portion of the south-westerly prolongation of the Porcupine mountain range, is made up of rocks belonging to the primary group, but its precise limit here has not yet been determined.

2. *Trap Rocks.*—Flanking the primary rocks already described on the north and north-west, are a series of ranges of hills, stretching in a direction generally south-westerly and north-easterly, which attain an altitude of from three to nine hundred feet above the lake. They are more regular, or rather less broken in outline than the primary hills; a change, however, which in the transition is noticed to take place gradually, from one group to the other, or in other words, the knobbed character of the ranges becomes less and less apparent as we cross them in a north-westerly direction, or from the primary range. These hills are composed of rocks differing at first but slightly from those of the primary group, but gradually the difference becomes more and more apparent, as we proceed northerly. The rocks of those hills nearest the primary range may possibly be regarded simply as rocks of that group, more or less altered, though the rocks of the outer ranges are plainly trap. The range of these rocks which may be said to commence at the very extremity of Keweenaw point, extends, after a slight curve to the north, in a general south-westerly direction, gradually receding from the coast, until at the cross-

ing of the Ontonagon river, it is nearly 25 miles inland. West-erly from Ontonagon river, the range becomes confounded with the northerly portions of the Porcupine mountains; while west from these mountains, a portion of what may be considered the same range of rock, has taken a more westerly course and approaches the coast, until, at the crossing of the Montreal river, it is but a few miles distant from Lake Superior. West from the Porcupine mountains, a second range of trap is continued, at a distance of from fifteen to twenty miles inland. The trap range of Keweenaw point may be estimated to compose one third the entire width of the point, and the south-easterly portions of the range are made up of compact greenstone, while those portions to the north-west are amygdaloid.

The ranges of hills constituting the north-westerly part of Isle Royale, and extending its whole length, are of similar rocks, and single knobs of well defined trap rocks occasionally occur, in the very midst of the primary region before referred to, upon the south coast; the proofs of the character of which will be shown as we advance.

3. *Metamorphic Rocks.*—Flanking the primary rocks on the south, is a series of stratified rocks, consisting of talcose, mica and clay slates, slaty hornblende rock, and quartz rock; the latter rock constituting by far the largest proportion of the whole group. In traversing the country south-easterly from little Presque Isle, the point referred to as the most south-easterly prolongation of the granite, this last rock passes almost insensibly into a serpentine rock, which has a regular jointed structure, sometimes approaching to stratification; continuing in the same direction, we find a series of hornblende slates, talcose, mica and clay slates, resting against the serpentine rocks, and still farther to the south-east the rock becomes almost uniformly quartz. The rocks of this group dip irregularly to the south and south-east, while the cleavage of the slates is very uniformly to the north.

The rocks of the metamorphic group stretch into the interior, in a westerly or rather south-westerly direction, forming the south-easterly part of the hilly region.

Rocks referable to this group also occur upon the north coast of Lake Superior.

4. *Conglomerate.*—The rock to which I shall restrict this term, does not occur, well characterized, at any point east from the district referred to, as the commencement of the trap group, nor has it been noticed resting upon any of either the primary metamorphic rocks, but is invariably seen resting upon the trap rocks. Commencing upon the north side of the trap, at the extremity of Keweenaw point, the conglomerate

flanks the trap upon its northerly side, as far west as the boundary of our state; nor does it stop here, for the same rock is seen at intervals, as far west as the head of Lake Superior. A similar rock also rests upon the trap of Isle Royale, facing the south-east.

In the course of the range of conglomerate upon the south shore, it forms a nearly continuous range of hills, with somewhat steep escapements, but with a generally rounded outline. These hills sometimes rise to a height of from three to five hundred feet above the level of the lake.

The conglomerate attains a very great thickness, being greatest at its westerly prolongation, and it gradually thins out as we proceed north-easterly; but the irregularity in thickness is so very considerable that variations of several hundred feet are not uncommon within the space of a few miles.

The conglomerate rock of the south coast, dips in mass irregularly to the north and north-west, while that of Isle Royale dips to the south-east.

5. *Mixed Conglomerate and Sandstone.*—The rock or rocks to which I have fixed the above name consists of an alternating series of coarse conglomerates and red sandstones, resting conformably upon the conglomerate rock before described. In strictness, these rocks should probably be considered as a member of the conglomerate itself, but for the sake of convenience in description, I have deemed it desirable to separate them.

This mixed rock was only noticed, as before stated, resting upon the conglomerate, and this only between Point Keweenaw and Montreal river. Its thickness immediately west from and upon the flanks of the Porcupine mountains, is very considerable, but it wedges out rapidly both easterly and westerly, and on the east near the extremity of Keweenaw point, it wholly disappears.

The mixed rock dips regularly to the north and north-west.

6. *Lower, or Red Sandstone and Shales.*—The red sandstone, with its accompanying red and gray shales, occupies a much larger extent of the country bordering upon Lake Superior than any other single rock or group of rocks. It rests upon the primary and metamorphic rocks, immediately west from Chocolate river; upon the conglomerate and mixed rocks from near Eagle river, of Keweenaw point, west to the head of Lake Superior; upon the primary trap, metamorphic and conglomerate rocks of the north shore of the lake, and upon the conglomerate rock of Isle Royale. It is this rock which forms the basis of the level plateaus or valleys, occupying the spaces between the several ranges of hills south from Lake Superior,

and west from Chocolate river. In these last situations this rock is frequently seen undisturbed to surround the bases of isolated knobs of granite, though when near to or in contact with knobs or trap, there are invariable evidences of very great disturbance.

The rocks of this group attain their greatest thickness at their westerly prolongation, gradually thinning out as we proceed easterly.

With the exception of that portion of the coast extending from Point Iroquois, at the foot of the lake, to Grand Island, the predominating rock upon the immediate coast, both on the south and north shore, is this red sandstone; for even the primary trap and conglomerate rocks are almost invariably skirted with a band of it. It is also over this rock that the waters of Lake Superior are discharged at the Saut de Ste Marie.

The sandrock forms the chief portion of the group, the shales occurring rather as beds than otherwise, as will be hereafter described.

The red sandstone both upon the north and south shores of Lake Superior, invariably dips into the basin of that lake, which may therefore be regarded as a synclinal axis.

7. *Upper or Gray Sandstone.*—Upon the south shore of Lake Superior, and extending from Point Iroquois to Grand Island, a sandstone occurs differing widely in its appearance from that before described. This sandstone rests *unconformably* upon the red sandstone, the former dipping gently to the south or south-east, while the latter dips very considerably to the north or north-west.

The elevated range of hills before described, as commencing a little easterly from Point Iroquois and extending to the Pictured rocks, are composed of this rock. From the Pictured rocks, the range of hills curves more to the south, stretching very far to the south-west, but its precise limit is not yet determined.

In its easterly prolongation the grey sandstone thins out rapidly. It is last seen at the Neebesh rapids of the Riviere Ste Marie, on the east, at which point, in consequence of not having been sufficiently examined farther westerly, it was confounded with the red sandrock in the last report which I had the honor to lay before you.

8. *Sandy Lime Rock.**—Resting immediately upon this upper or grey sandstone is a sandy limerock, which, although nearly wanting at the very easterly extremity of the peninsula, as we proceed westerly, occupies a more important place.

* The names which have been affixed to the several sand and lime rocks are regarded as merely temporary, and are introduced, for the present, barely to facilitate description.

This rock, which, as its name implies, is intermediate between a sandstone and limerock, may be seen on Sailor's encampment island of the Riviere Ste Marie, as also at several points in the vicinity of Monusco bay, from whence it stretches westerly, occupying nearly the central portion of the peninsula, for a distance of at least sixty miles; from which its precise range and limit has not yet been determined. The outcropping edge of this rock appears at a level very considerably below that both of the sandstone to the north, and of the limestones to the south. Its width, for the distance mentioned, varies from ten to fifteen miles, and it dips uniformly to the south-south-east.

The sandy limerock has not yet been examined with sufficient care to admit of accurate description. It contains but few fossils, but those few are sufficiently characteristic, were there no other considerations, to separate it from the lower limerocks and shales.

Upon the sandy limerock, to which reference is above made, rests the lower limerock and shales, and upon this last the upper limerock, both uniformly dipping to the south or south-east. These limerocks were described in general terms, in my third annual report, and although many additional facts have been gathered respecting their character, range and extent, it is, perhaps, unnecessary to lay them before you at this time. I will barely add, with respect to them, that the suggestion there mentioned, that a more careful examination of these limestones would render a farther division of the groups necessary, has been fully confirmed.

I had hoped to lay before you a profile section of the rocks of the upper peninsula, but the impossibility of having it engraved in time to accompany this report, has led me to defer it. I regret this the more, since many of the facts connected with a full understanding of the economical portion of this report, are so intimately dependent upon the general geology of the country, that, in the absence of correct maps, and without a profile section of the rocks, I fear it will be impossible for me to render the most important portion, so far as regards the prosperity of the state, intelligible.

As it is, I can only, in the place of this, lay before you a general section of the rocks of the upper peninsula, together with their thickness, so far as the same has been satisfactorily determined. This section is intended simply to represent the order in which the several rocks rest upon each other.

Having already described in general terms, the range and extent of the rocks of the upper peninsula, so far as the same have been examined, the limits of the present report will admit of nothing more than a general description of the charac-

ters of these rocks, and I shall not attempt a minute description of any members of the series, except such as are more or less connected with subjects which are supposed to be of immediate practical importance.

Section illustrative of the order of super-position of the Rocks of the Upper Peninsula.

	Thickness in feet.
9. Tertiary Clays and Sands.	
8. Upper Limerock Group, (embracing as members, the Drummond Island and Mackinaw limestones.)	
7. Lower Limerock and Shales.	
6. Sandy or Intermediate Limestone.	
5. Upper or Grey Sandstone,	mean 700 feet.
4. Lower or Red Sandrock and Shales,	extreme 6,500.
3. Mixed Conglomerate and Sandrock,	extreme 4,200.
2. Conglomerate rock,	extreme 5,260.
1. Metamorphic, Trap and Primary rocks.	

PRIMARY ROCKS.

Although the usual ternary compound of quartz, feldspar and mica, occurs but rarely in the primary, in the vicinity of the coast of Lake Superior, and in fact but rarely in any portion of the range which I have visited, nevertheless, the great mass of rocks included within this range, may, in a broad sense, be called granite. The compound above referred to, is more common in the westerly than in the easterly portion of the

range. The more common rock is made up of quartz, feldspar and hornblende, giving rise to a very dark colored syenite; occasionally mica enters sufficiently into the compound to form syenitic granite, and sometimes the place of the hornblende in the syenite, is supplied by schorl or tourmaline, thus giving rise to a schorl rock.

The rocks of the south-easterly portion of the primary range of hills are more clearly defined as granite rocks, than those situated more northerly, for they are more distinctly and largely crystalline in structure, and quartz enters much more largely as a constituent into their composition. As we proceed north-westerly, from the south-east boundary of the primary, over the several broken ranges of hills, we find the character of the rocks in mass almost imperceptibly changing. The quartz as a mineral gradually forms a less important part, and it finally almost wholly disappears, leaving a binary compound of feldspar and hornblende, which then assumes a granular structure, constituting greenstone. The intermediate rock, between the syenite and greenstone ranges, may not inappropriately be called a syenitic greenstone.

The primary rocks which appear in the vicinity of Lake Superior, in the several ranges of hills extending from a point opposite little Presque Isle,* to Huron river, are essentially either syenite or syenitic granite. The rock, as a whole, is extremely compact, and the constituent minerals are mostly in small crystals, though occasionally the feldspar assumes a more largely crystalline form.

The granitic rocks, so far as the range has been examined, in a south-westerly direction, are largely traversed by dykes, that are almost without exception made up of materials in all respects identical with the greenstone, before alluded to, as forming the more north-westerly ranges of hills. The courses of these dykes or veins are invariably marked by striking changes in the character of the rock traversed, and in the larger dykes, the evidences of the changes produced by the heat of the injected matter, extend to several hundred feet upon either side of the dyke itself. The connection between the rocky matter composing these dykes and the ranges of greenstone, lying north-west, is clearly identified, not only by the perfect similarity in the mineral character, but also from the fact, that as we proceed in the direction of the ranges of greenstone, the dykes become much more frequent, until at length it becomes difficult to determine which of the rocks predominate in quantity.

These facts serve to throw much light upon the relative ages of the several ranges of hills, or in other words, serve to show the order in which they were severally uplifted; facts

* A little south-east from Riviere Des Mortis.

which will be more fully shown when we come to consider the present position of the overlying sedimentary rocks. These facts are not only important, to enable us to understand the many changes which have taken place, with regard to the relative position of the land and water, but they are rendered of practical importance for the reason, which I think may be satisfactorily shown, that the mineral region of the upper peninsula, to be hereafter described, is strictly confined to only the outer portion of the rocks of a single epoch.

The veins and dykes of greenstone, referred to as traversing the granite rocks, do not in this portion of the group, appear to have any regular magnetic bearing, for they traverse the rock in all directions. Veins of any other matter are very rarely seen traversing the granite. In a single instance, what was regarded as a true vein of porphyry, having a width of nearly three feet, was noticed, which vein is crossed, at angles of 53° and 107° , by a vein of greenstone, having a width somewhat less than that of the porphyry. In this instance, the greenstone is clearly the most recent vein.

The veins of greenstone traversing the granite, vary from a mere line, to 50 or 60 feet in width. The intimate blending of the material composing these veins, together with the chemical differences, causes them to disintegrate or waste away more rapidly than the rock they traverse; the result of which is, that deep grooves are frequently left in the granite, the simple result of the wasting away of these dykes or veins. This is peculiarly the case upon the coast of the lake, where the rocks are subject to the action of the waves, which have, in some instances, so removed the debris as to leave long and narrow bays, with high perpendicular walls, occupying simply the space once occupied by the dyke. The Huron Islands, which are simple elevated granite knobs, appear, upon first examination, as a mass of rocks, completely rent in many places, with portions separated from each other by narrow clefts, having perpendicular walls of great height. While these rents are of sufficient width to admit of being traversed by small boats, the perpendicular walls are so little varied in their elevation as scarcely to leave a point, in these narrow passages, where a landing can be effected. A careful examination of these passages shows them to be simply the spaces once occupied by dykes or veins of greenstone, which having disintegrated, and the detrital matter having been removed by the action of the waves, has left the walls of the more enduring granite rock, unbroken and almost untouched.

Upon the north coast of Lake Superior, well defined granite and syenite, or syenitic granite, occasionally appear upon the immediate coast of the lake, but more frequently these rocks are flanked on the south by greenstone, with occasional nar-

row bands of sandstone; thus precisely reversing the magnetic order of those rocks upon the south coast.

TRAP ROCKS.

Were we to consider the rocks of the district under consideration, strictly in their chronological order, those rocks which I propose to treat as trap rocks, would undoubtedly follow those slates and quartz rocks which are considered as metamorphic, and which may be regarded as identical in time of uplift with those rocks before alluded to, as being intermediate between the granitic and trap rocks. The almost insensible gradations by which the granitic rocks pass into the greenstone of the trap formation, and the near analogy of the whole of the rocks of both formations, to each other, renders it more convenient, at the same time that it is more simple to follow the arrangement or order that I have adopted.

I have already stated that in passing from the granitic region on the south side of Lake Superior, in a direction north-westerly, we cross a series of ranges of hills, varying in height from three to nine hundred feet above the lake, and that in pursuing this course, we observe that the character of the rocks gradually and almost insensibly change, until at length they become well defined greenstone.*

The rocks of the outer or north-western range of hills, which were clearly the last of the series of uplifts, bears more unequivocally the evidences of igneous origin, than either of the outer ranges alluded to. The rock upon the south flank of these hills, is invariably very compact greenstone, while upon the north-westerly line it is almost equally invariably an amygdaloid, or at least, has an amygdaloidal structure. The causes of this difference of structure of the rock, upon the opposite sides of this range of hills, when carefully examined upon the ground, are very apparent; for it is evident, as will hereafter be shown, that the uplift of the rocks of this range of hills was wholly upon the south-easterly side, and while the rocks of this portion were in a solidified state, or, in other words, that a point in Lake Superior may be regarded as the fixed axis of the uplifted mass. That this was the case, is shown by the fact that the sedimentary rocks to the south or south-east are

* In the present report, I use the term *greenstone* in its generic sense, applying it to all the compact rocks, of a granulated structure, belonging to the trap range. By far the larger proportion of these rocks are greenstone, in its most restricted sense, or in other words are composed of feldspar and hornblende; but the term is also used to include rocks which in a strict sense would be considered as altered syenite, syenitic granite, hornblende rock and angitic rock.

The term *amygdaloid*, I apply, as it is usually applied to that portion of the rock having a difference of form simply, without any reference to the constituents of the rock. This generic use of terms is employed for the reason that the limits of the present report will not allow any thing more than a very general consideration of the subject. The term *trap* is used in such a sense as to include both the greenstone and amygdaloid, though it may sometimes prove that the amygdaloid has had its origin from the fusion of the lower portions of the sedimentary rocks.

scarcely disturbed, so far as regards this range of hills, while the sedimentary rocks on the north and north-westerly side, are invariably tilted to a high angle, near the range of hills, which angle gradually decreases as we pass farther and farther from the hills themselves. These sedimentary rocks, which upon the north side always dip from the range of trap hills, are in their close proximity to the trap, inclined at angles varying from 45° to 85° . Dykes of from fifty to four or five hundred feet are of frequent occurrence, traversing these sedimentary rocks, but the widest of these have invariably been protruded between the strata of the sedimentary rocks, and consequently have the same general inclination. The result of these frequent dykes, which occur at comparatively short distances from the main body of trap, is, that the sedimentary rocks frequently so far lose their original character as scarcely to be recognized.

The rocks of the complete north-western escarpment of this range of hills, were evidently in an intense state of ignition while in contact with the sedimentary rocks, as is clearly shown by the very great changes which have taken place in the rocks last alluded to. In fact, I am disposed to refer the origin of much of the amygdaloid rock to the fusion of the lower portions of the sedimentary rocks referred to, for the reason, that as we pass south from this junction, the amygdaloid rocks wholly disappear, their place being supplied by greenstone; and again so intimately are they blended, that it is frequently impossible to determine where the amygdaloid ceases, and the upper sedimentary rocks commence. Fragments of the sedimentary rocks, the characters of which can be clearly recognized, are not of rare occurrence, imbedded in the amygdaloid rock, a circumstance which although by no means conclusive, should not be overlooked in considering this subject.

I would not wish to convey the idea that the amygdaloid rocks have their origin exclusively from the altered sedimentary rocks, but simply that the change in the structure of the trap, from greenstone to amygdaloid, may and no doubt does depend upon the proximity of the sedimentary rocks to the trap, while the latter was in a state of ignition.

I have been compelled to tread upon grounds which may, perhaps, be considered theoretical, but it would appear to be necessary in order to convey a proper idea of the condition of the rocks composing the range of hills under consideration. These views, however, would not have been alluded to at this time, had it not been for the fact, that an understanding of all that relates to the mineral resources of this portion of our state, is more or less intimately connected with this portion of the subject.

Although the general range of the trap hills has been alrea-

dy given, I will define, as nearly as is in my power, the line of junction between the trap and sedimentary rocks, upon the north escarpment, premising that the elevation at which this junction takes place, is usually at a height of from 100 to 500 feet above the lake, and only in a single instance does this line reach the coast of the lake. Commencing almost directly at the extremity of Keweenaw point, this line passes in a south-westerly direction, gradually receding from the coast; it crosses Sturgeon or Portage lake near its centre, after which it recedes still more rapidly from the coast, until finally it is seen to cross the upper forks of the Ontonagon river, and soon after the whole is apparently lost in the range of the Porcupine mountains; which last range has a course so much to the south-west as probably to completely intersect the first range mentioned. On the west side of the Porcupine mountains, the range of hills and the line of junction appear again, but many miles farther north than they would have been looked for; from thence the true line gradually approaches the coast, until, at its point of crossing the Montreal river, it is but about 2 miles above the mouth of that stream.

To the north and north-west, through the whole of the distance described, this trap is bounded by hills of conglomerate and sandstone, more or less elevated, but usually not exceeding four hundred feet. To the north-west of these hills of sedimentary rocks, a dyke of trap is seen to extend for many miles along the line of coast of Keweenaw point, and so great is the width of the dyke, that, unless carefully examined, its character might easily be misunderstood. It lies in a plane parallel to the stratification of the sedimentary rock by which it is embraced, and with that rock dips to the north-west. The dyke is chiefly made up of greenstone, but not unfrequently large portions of the mass consist of amygdaloid, in which the amygdules are filled or composed of quartz, chalcedony, agate, calc-spar, zeolite, &c.

The dykes just referred to, so far as their relation to the amygdaloidal portion of the trap is concerned, as also the many others similarly situated with respect to the superincumbent sedimentary rocks, will be regarded in the same light as contemporaneous veins, though they are only contemporaneous with the uplift of the strata, and not with their deposit. But there is still another class of veins which not only traverse a portion of the trap rocks, but also the upper sedimentary rocks, and which may be regarded as true veins. These last mentioned veins traverse the rocks at a high angle with the line of bearing of the sedimentary rocks, as also with the line of junction of those last mentioned with the trap rocks. The composition of these veins is widely different from that of the contemporaneous veins or dykes before referred to. As this subject will be

treated more at length in a succeeding portion of this report, I deem it unnecessary at this time to refer more particularly to the subject.

A single knob of trap appears under circumstances which add very much to its interest, at what is usually known as Presque Isle, an elevated rocky point immediately north-west from Riviere Des Morts, and almost directly within the granitic region. This point of land has its origin from the simple elevation of a mass of trap rock, which rises on the north in abrupt cliffs, varying from twenty to sixty feet in height. The trap is mostly greenstone, though portions of it are so largely impregnated with a dark colored, almost black serpentine, as to deserve the name of serpentine rock.

The knob of trap under consideration is possessed of additional interest, from the unequivocal evidence of uplift, as also from the manner in which these evidences are exhibited. The cliffs of trap occupy the very extremity of the point, while the neck and central portions are made up of conglomerate or trap-tuff and sandrock, resting upon the trap. These upper rocks also appear upon the immediate coast, in cliffs of from twenty to sixty feet in height, and in many places they are seen resting directly upon the trap. The stratification of these sedimentary rocks has been very much disturbed, and they invariably dip, at a high angle, in all directions from the trap itself. The character of both rocks, at the immediate line of junction, is almost completely lost, and the evidences of change most unequivocally marked. But the most curious feature of the whole is, that the sedimentary rocks, for a distance of several hundred feet, have been completely shattered or broken into minute fragments, which, having retained their original position, were again cemented by the injection of calcareous matter. This injection has filled the most minute fissures, and so perfect is it, that, in looking upon the face of a mural cliff of these rocks, the veins may be easily seen at a distance of many rods, forming, as it were, a complete net work over the cliff, and so minute is it, that a single hand specimen frequently contains many hundreds of these veins.

This knob of trap, like the rock before described, is also traversed by veins, of a date subsequent to the uplift of the rock.

The whole of the north-western portion of Isle Royale is made up of trap, and in truth that rock constitutes, by far, the largest proportion of the rocky mass of the island. The two northerly ranges of hills, already alluded to as traversing the island, in its greatest length, are wholly trap. The most northerly range of hills, is composed almost exclusively of

greenstone, while the rock of the south or south-easterly range, becomes more decidedly amygdaloidal in its structure; thus reversing the order which these portions of the rock bear to each other upon the south shore of the lake.

The ranges of hills immediately bounding Lake Superior upon its north coast, are almost invariably either well defined trap or altered syenite, while the decidedly primary rocks usually appear in ranges of hills to the north of these; thus following the reversed order of the rocks upon the south coast.

The character of the trap rocks of Lake Superior has perhaps been sufficiently described, to answer the purpose for which this hasty sketch is intended; and I will only add, that they are usually distinctly jointed, and where they approximate to the sedimentary rocks, there is not unfrequently so distinct a cleavage, opposed to the joints in direction, as to give the appearance of stratification. The jointed structure of the trap rocks sometimes, though rarely, passes to what may be termed a rudely columnar structure. Upon one of the long rocky points forming the north-easterly extremity of Isle Royale, this rock assumes the columnar form, and the columns are tolerably well defined, having a height of from eighty to ninety feet. The columns are also seen, but less perfectly developed, forming the coast of a small rocky island, two or three miles south from the point last alluded to. These are the only points in the trap of Lake Superior, where I have noticed the rock to assume this form.

METAMORPHIC ROCKS.

The general direction of the rocks composing this group, has already been described, and they are confined exclusively to the range of hills lying upon the south-east side of the granitic rocks. The general direction of these hills is south-west and north-east.

The outline of the hills of the metamorphic group is less broken than either the granitic or trap ranges, but these rocks sometimes rise in abrupt conical peaks, closely resembling those of the granitic rocks.

The area of country occupied by rocks of this group, is less than that of either the primary or trap, the general average width not exceeding six to eight miles. The precise limit of the group in a south-westerly direction, is not known.

It has already been stated that Chocolate river is the boundary, on the south-east, between these and the sedimentary rocks, and that they extend in a north-westerly direction from this stream to the granite, against which they rest. The group is made up of an alternating series of talcose and mica slates, sometimes graduating into clay slates, with quartz and serpen-

tine rocks, the quartz rock constituting by far the larger proportion of the whole mass. Since it would be nearly impossible to describe the alternations of these several rocks, in such a manner as to be understood, without the aid of a diagram, or section, no attempt will be made to do so.

The cleavage of all these rocks is usually north or north 10° west, at an angle, which, in the main, varies but little from 80°, but the mass of the group appears to dip regularly to the south or south-west. The talcose slates and quartz rocks alternate frequently with each other; and with the rock which has been called serpentine rock, less frequently.

The quartz rock is usually distinctly granular, though it is sometimes compact, with a conchoidal fracture. It usually separates by cleavage into masses, or strata, having a considerable degree of regularity, and varying from a few inches to several feet in thickness. The rock is usually more or less regularly jointed.

That rock which, for the sake of convenience, I have denominated serpentine rock, bears a close resemblance to greenstone, being essentially composed of granular feldspar and hornblende, with which serpentine is intimately blended. This rock only occurs in the talcose slate as we approach the granitic region, and possibly a more close examination may show it to be a simple series of dykes, lying parallel to the line of cleavage of the slate rocks.

The metamorphic rocks are occasionally traversed by trap dykes. The group of rocks under consideration has been comparatively little examined, and the more minute details connected with it, will be taken up at some future time.

CONGLOMERATE ROCK.

The lower of the sedimentary rocks, to which I have attached this name, appears to be invariably connected with, or to rest upon, the trap rock, nor has it been noticed, to any extent, in connection with either of the other lower rocks, for it wholly disappears as we approach the granitic and metamorphic groups. Of all the sedimentary rocks, this is the most variable in thickness, and not unfrequently does a few miles make a difference of several hundred feet. The conglomerate rock may, without doubt, be considered as a trap-tuff, which was gradually deposited or accumulated around the several conical knobs of trap, during their gradual elevation, and which would necessarily occupy the complete spaces or valleys between the several irregular ranges of knobs or hills.

If we regard this conglomerate rock in this light, we will at once perceive why the rock should be variable and irregular in its thickness.

The pebbles of which the mass of the rock is composed, consist of rounded masses of greenstone and amygdaloidal trap, of which the former make up by far the larger proportion, and scarcely a pebble of any other rock than trap, enters into its composition. These pebbles vary in size from that of a pea, to several pounds weight, but the average size may be stated at 1½ to 2 inches in diameter. The pebbles are usually united by a mixed calcareous and argillaceous cement, more or less colored by iron, and so firm is this union, that the most compact and tough of the greenstone pebbles, will frequently break through as freely as the cement, and crevices and narrow veins are frequently seen passing indiscriminately across the pebbles and cement. This fact is the more worthy of notice, since the pebbles are almost without exception, made up of the hardest and most indestructible portions of the trap rock.

The conglomerate rock can scarcely be said to occur in such form as to be well defined, in any portion of the country, excepting upon the northern flank of the outer trap range, before referred to. On the outer or northern side of Keweenaw point, the conglomerate commences near the extremity of the point, and extends several miles westwardly, forming a series of abrupt and precipitous cliffs upon the immediate shore, as also a range of well defined hills, a little in the interior; which hills have an elevation varying from 200 to 300 feet. After appearing for a few miles upon the coast, this rock gradually stretches into the interior, following the line before described as the most northerly boundary of the outer trap range of hills, and invariably occupying a place to the north of this range, and it may be observed, nearly or quite continuously, as far as Montreal river, which stream it crosses at a short distance above its mouth, thus making its complete length, within the limits of Michigan, computing its southerly curve, something over one hundred and forty miles; but the rock does not cease at Montreal river, for it may be seen at short intervals in the interior, as far westwardly as the head of Lake Superior.

At the trap knob of Presque Isle, the conglomerate is imperfectly developed, but on the south-westerly side of Isle Royale, it is more perfectly developed, flanking the hills of trap upon the southerly side.

The conglomerate rock is imperfectly stratified, in masses of immense thickness, and it dips, upon the south shore of Lake Superior, regularly to the north, and north-west,* usually at high angles, varying from 30° to 85°, while upon Isle Royale and the north shore, the dip is reversed, being south

* This variation in the dip is in conformity with the variation in the direction of the trap hills.

and south-easterly, or in other words, the rock upon all sides dips in the direction of the lake basin.

Upon the south shore of the lake, the thickness of this rock was not estimated at any point west from Montreal river, a little east from which it attains its greatest thickness, being, as estimated, 5,260 feet. In addition to the great variations in thickness, over comparatively small districts, the formation wedges out as we pass easterly along the range, and so rapid is this change, that near its easterly prolongation the thickness was estimated at something less than 1,000 feet.

The greatest estimated thickness of the rock upon the north coast, is a fraction less than 2,300 feet.

I have already stated that this rock is frequently traversed by dykes of trap, which are usually parallel to the line of stratification and dip of the rock. These dykes, which have sometimes a thickness of 50 to 60 feet, and even several hundred feet, are sometimes continuous for many miles, and are many times repeated. In addition to the dykes just alluded to, the rock is frequently traversed by veins of a more recent date, which traverse alike the trap and conglomerate rocks, always at very high angles with the line of bearing of the conglomerate. These veins, which are usually more perfectly developed near the line of junction of the two rocks, or for a distance of a few thousand feet upon each side of the junction, are clearly true veins, and since, with a few unimportant exceptions, they are the only veins belonging to this range which are metalliferous, they will be considered more fully under a separate head.

MIXED CONGLOMERATE AND SAND ROCK.

This rock formation is made up of an alternating series of conglomerate and red sandstones, which rest conformably upon the conglomerate rock last described, dipping with that rock, into the bed of Lake Superior. The mixed rock was not noticed upon the north side of the lake, or upon Isle Royale, but upon the south shore the rock was traced continuously for a distance of about one hundred and thirty miles, extending from a few miles westerly from the extremity of Keweenaw point, to Montreal river. It follows the line of the conglomerate before described, stretching from Keweenaw point, in a south-westerly direction, and again curving to the north-west, forming as it were, a crescent between the points before mentioned, the result of which is, that the rock only appears for a very limited distance upon the coast of the lake, at Keweenaw point.

From a point about eighteen miles easterly from Montreal river, the rock wedges out rapidly as we proceed westerly

from that point, and as we continue towards the head of Lake Superior, the rock wholly disappears, or becomes merged in the conglomerate rock below, and the sand rock above. The greatest observed thickness of this rock, is 4,200.

The conglomerate portion of the mixed rock consists of strata of conglomerate, varying from a few feet to several hundred feet in thickness, and it is composed of materials in all respects resembling those constituting the conglomerate rock before described, and these materials are united by a similar cement.

The sandstone portion of the formation occurs in strata of very nearly corresponding thickness, and the two rocks may be said to form nearly equal portions of the complete mass. But the material of which this sandstone is composed, differs widely from that of the true sandrock lying above, for while the latter is chiefly made up of quartzose materials, the former is composed of materials bearing a close analogy in composition to those of the conglomerate rock itself; or in other words the sand consists chiefly of greenstone, so much comminuted as, when cemented, to compose a coarse sandstone. It will thus be seen that the members of this formation differ only in the degrees and fineness of the material, and the character of this material will explain sufficiently why the true conglomerate, and the mixed rocks are referable to the same origin, for the materials of the several members of the group have their origin from the trap rock, and as a whole, may, perhaps, be regarded as a trap-tuff.

The coarser conglomerate of the formation is scarcely separated by lines of stratification, and the strata appears usually in mass, embraced between the strata of sandstone, but the stratification of the latter rock is perfect, and it bears evidence of having been deposited in shoal water, in the very abundant, perfectly defined ripple marks which it exhibits through its complete range.

No fossils were noticed in connection with either the mixed rock, or the conglomerate lying below it.

Dykes of greenstone occasionally appear in the mixed rock, but less frequently than in the rock below. These dykes almost invariably occupy places between the strata of the rock, and correspond in position to the direction and dip of the rocks by which they are embraced, or in other words, the rocky matter composing the dykes appears to have been injected in a plane corresponding with that of the stratification of the embracing rock. As in the conglomerate below, these dykes have produced very great changes in the color and structure of the mixed rock bounding them upon either side.

In addition to these, the mixed rock is occasionally, (though less frequently than the rock below,) traversed by veins or

cross courses of a more recent origin than the dykes, (which latter they usually cross at a high angle,) their course usually being at an angle of at least sixty degrees, opposed to the line of bearing of the mixed rock. These cross veins are usually made up of calcareous spar or a sub-granular limestone, and more rarely of some variety of quartz, and imperfect trap rock, the latter of which is usually of the amygdaloid variety.

RED SANDSTONE AND SHALES.

That rock to which I have applied the name of red sandstone, is emphatically the chief rock that appears upon the immediate coast of the south shore of Lake Superior, and the same remark will apply, in a more limited degree, to the complete coast of the lake. A traveler proceeding westerly along the coast, from Grand Island to the head of the lake, would imagine he had seen little else than red sandstone, and in fact, were he to confine his examinations to the immediate coast, he would see no other rock for nineteen-twentieths of the distance. From Grand Island westerly to the mouth of Chocolate river, no other rock is seen in place, and from Chocolate river to Keweenaw point, embracing the complete width of the primary, metamorphic and trap ranges, the hills forming these groups are almost invariably surrounded or flanked at their bases, by this sand rock, so that even along this portion, the hills are, for a long proportion of the distance, cut off from the lake by a narrow belt of the rock under consideration, and westerly from Keweenaw point to the head of Lake Superior, no other rocks appear upon the coast, if we except several trap dykes in the vicinity of the Porcupine mountains, and a series of more recent deposits of clay and sand, that appear west from Keweenaw point. This sand rock also occurs upon the southerly side of Isle Royale.

The material of which the red sandrock is composed, differs widely from that of the sedimentary rocks before described, for while the rocks last referred to are made up of materials clearly of trappean origin, and in which the material is very rarely quartz, the rock under consideration is composed of materials, the predominating portions of which are clearly derived from the granitic and metamorphic rocks, and in which quartz occurs abundantly, though with this, there is usually associated more or less sand, that has all the characters of the comminuted trap, constituting that portion of the mixed rock before referred to. Magnetic iron sand, sometimes becomes a constituent of the red sandrock, and occasionally continuous strata of several inches thickness, are almost wholly composed of this material. The material composing this rock is usually cemented by calcareous matter highly colored by the per oxyd

of iron, though not unfrequently these are associated with argillaceous matter.

While the chief mass of the rock is a coarse grained, somewhat compact, sandrock, there are portions of the formation where there are well formed red and grey flags, and red and green shales, forming, as it were, beds of a very considerable thickness, and occupying large districts of country. These red and green shales are more largely developed in that district extending from Granite point westerly to Keweenaw bay, and upon the south side of Keweenaw point, extending from the head of the bay to near the extremity of the point, they are particularly largely developed. These shales more usually occur in alternating bands of deep red and green colors, the red usually largely predominating, and they are made up of argillaceous matter, with sand, the whole of the materials being of extreme fineness.

On the south-east side of Keweenaw bay, near its head, an argillaceous rock appears, and extends for a short distance along the coast, which is an anomaly. The rock is evidently embraced in, or rather may be said to constitute a member of the sandstone series, but it differs widely from any other rock seen in connection with it. This argillaceous rock sometimes appears in the form of a slate, though its most usual form is that of a compact strata, frequently of several inches thickness, and which closely resembles indurated clay. A peculiar appearance is given to this rock by the innumerable layers or very thin strata, which compose the mass, being of different colors, sometimes red, grey and dark brown, alternating in the same hard specimen.

The material of which this argillaceous rock is composed, possesses an extreme degree of fineness, and is so soft as readily to be cut with the knife, which qualities render it a fit material for the manufacture of pipes, to which purpose the Indians of the country have long applied it. It has also been applied to use in sharpening tools, but its softness is a serious objection to its use for that purpose.

A similar argillaceous rock also appears at several other points in the interior, or southerly from that already described, but as yet I have been unable to determine its thickness at any point. The finely represented bands or zones, which may fairly be supposed to represent the original lines of deposition, are very much contorted, and in such a manner as to lead to the conclusion that this change must have taken place very soon after the deposition of the rocks, and while they were still in an unindurated state.

The rocks belonging to the red sandstone formation, bear the evidence of having been almost universally deposited in shoal water, for ripple marks occur abundantly at all points

where the rock takes on the decided character of sandrock, and these ripple marks may frequently be seen, for many rods together, as distinctly and clearly defined as they are at the present day in the loose sands forming the bottom of some of the shoal bays of Lake Superior. Fossils are rare in the red sandstone, and in fact, I have never seen any other than *fucoides*, of which there are three species, that are tolerably well defined.

The red sandrock is less frequently traversed by dykes of trap than either of the rocks before described, though dykes were sometimes noticed traversing the whole of the several rock formations, up to and including the red sandstone. Upon portions of the north coast, where the conglomerate and mixed rocks are more frequently wanting, and where the red sandrock is brought more nearly in contact with the trap, these dykes are of more frequent occurrence. It is deserving of remark, that where the lower rocks are either wholly or in part wanting, the red sandstone usually becomes of a deep brown color, and the material of which the sand is composed, gradually changes from that before described to greenstone.

I have already stated that the sandrock, at its westerly prolongation, attains its greatest thickness, which was estimated at 6,500 feet, but as the rock continues easterly, it gradually and quite regularly diminishes in thickness, and beyond Saut de Ste Marie, the thickness is very inconsiderable. The average rate of diminution which takes place in the thickness of the rock as we proceed easterly, was shown by a great number of observations, upon the south-westerly portions of the coast of Lake Superior, to be a fraction over fifteen feet to the mile, but this rate of decrease could not be satisfactorily estimated upon the lower or easterly half of the coast. The red sandrock thins out as we proceed southerly or inland from the coast, at a still more rapid rate, as was most satisfactorily shown, where it is connected with the several primary, metamorphic and trap ranges of hills, for all or nearly all the valleys, after passing the outer or northerly range of trap hills, are based upon this sandrock, and since we have every reason to believe that this sandrock was deposited in part, during the gradual elevation of the several chains of hills, it would follow, that over those districts which were last elevated, the rock would attain its greatest thickness. I have already alluded to the order in which the several ranges of hills appear to have been uplifted, and since more particular reference will be made to this hereafter, I leave the subject for the present.

The red sandrock south from Lake Superior, as well as upon the immediate coast, dips regularly northerly, while that upon the north coast dips invariably southerly, or, as has already been said of the lower rocks, this rock dips, upon all

sides, regularly into the basin of the lake. The quantity of dip is exceedingly variable, being always very much increased as it approaches the trap, and diminishing as it approaches the primary and metamorphic ranges.

The line of cleavage of some of the members of the lower sandrock and shales is frequently irregular, and opposed to the true stratification of the rock.

UPPER OR GRAY SAND ROCK.

The only remaining rock which separates the red sandrock from the limestones lying to the south, is a gray, or brownish sandrock, that is almost wholly composed of grains of quartz, usually feebly cemented with calcareous matter. The composition of this rock differs from that of the lower sandrock, in being more exclusively quartz, while in epoch of deposition, the rock under consideration should not be confounded with that of the red sandstone. It has already been stated that the red sandrock of the south coast, dips regularly northerly, while the upper or gray sandstone dips equally regularly south or south-easterly, in which respect the last mentioned rock conforms to the limestones resting upon it, while it rests itself upon the uplifted edge of the red sandrock, below.

I have already stated that this rock was first noticed, rising in hills, at a point not far distant from Riviere Ste Marie, and south-east from Point Iroquois; from this point, it stretches westerly in an elevated and very regular chain of hills, that are upon the coast, as far as Tequoimenon bay; westerly from which the shape of the coast is such that these hills do not again appear upon it, until we reach that precipitous portion of the lake coast known as the Pictured rocks, where the fury of the waves, aided by frost, has acted upon the feebly cemented material of which the rock is composed, to such an extent as to leave large portions of what was originally the northern escarpment of these hills, along this coast, in high mural and overhanging precipices. Westerly from the Pictured rocks, the ranges of hills, which are composed or made up of this rock, stretch in a south-westerly direction, passing completely to the south of the primary, trap and metamorphic regions. The westerly prolongation of this rock has not yet been determined.

The upper sandrock, like the lower, abounds in clearly defined ripple marks, and its line of cleavage is very irregular, frequently being opposed to the line of stratification over very considerable districts of country. Two indistinct species of *fucoides* were all the fossils noticed in connection with it.

I was unable to obtain any observation upon the thickness of the upper sandrock, which were satisfactory, but from the

imperfect observations which were obtained, I was led to conclude that the average thickness as far westerly as the Pictured rocks, does not vary very far from 700 feet. The upper sandrock, like the rocks before mentioned, wedges out as we proceed, in an easterly direction.

TERTIARY CLAYS AND SANDS.

As in the lower, so in the upper peninsula, the older rocks are more or less covered by deposits that may be severally arranged under the above head. To these deposits it is my intention, at the present time, barely to allude.

Stratified clays and sands, similar to those skirting the borders of the lower peninsula, are seen at many points, and continue for long distances upon the coast of Lake Superior; and they are also largely developed at many points in the interior of the country. These deposits sometimes attain a thickness of from 200 to 300 feet, and they are spread over the less elevated portions of the district. The character of these clays and sands bear a close resemblance to those described in a previous report, as occurring upon the lower peninsula.

ECONOMICAL GEOLOGY.

Rocks.

The series of limerocks resting upon the sand rock last described, were noticed in the report which was laid before you at a previous session, and the limits of the present report will not permit me to refer to them, more fully, at this time. My observations will, therefore, only include those rocks which lie below the limestones. It will be borne in mind that the whole of the group of limestones are embraced in the southerly portion of the upper peninsula, and that their outcropping edges do not reach within many miles of the coast of Lake Superior. This is an important fact, for it shows the whole of the northern part of the upper peninsula to be deficient in materials for the manufacture of lime, which are, in truth, wholly wanting.

Materials adapted to the purposes of building, abound, throughout the district of country under consideration, and though they vary exceedingly in value for that purpose, yet no portion of the country can be said to be without a supply.

Among the most valuable of the materials for this purpose, the syenites and syenitic granites deservedly rank first, and they occur of a quality which may be advantageously worked at various points in the primary range. Some of the syenites near the coast of the lake are so situated as to be readily quarried, and they may be made to furnish a beautiful and durable material for building. The color of these syenites is usually a

very dark gray, from the predominance of hornblende in the composition, but this is by no means invariably the case.

The metamorphic group scarcely furnishes a fit material for use as a building stone, for the structure of its schists would be an effectual bar against their use, and the difficulties of working the quartz rock will probably prevent that rock being applied to that purpose.

Some of the compact greenstones and altered syenites of the trap range, may be made to furnish an excellent building stone, which, although in powers of resisting the action of disintegrating agents, may be less than that of the unchanged syenite, nevertheless possess a very great degree of durability. The greenstone ranges of hills, frequently for very considerable distances, are made up of rock in which the jointed structure is so perfectly developed, that regular blocks, of a convenient size for building may be obtained, with comparatively little labor.

The conglomerate rock is scarcely applicable to use for purposes of building.

A very good building stone may be obtained from many portions of the lower, or red sandstone formation, and though the cement of this rock is usually not very perfect, yet, frequently, such changes have taken place in the rock, that it has almost taken on the character of granular quartz rock, in which cases, its durability is very much increased. The strata of this rock are usually of a convenient thickness to admit of being easily quarried, and they are so regular that the stone will require but little dressing.

The upper, or gray sandrock, being almost uniformly but feebly cemented, and sometimes decidedly friable, is of less value as a building stone than either of the rocks before mentioned. Those portions of the upper sandrock where the calcareous cement is perfect, but not sufficiently hardened, might be rendered much more capable of resisting the action of the elements if allowed to remain under shelter a sufficient length of time to allow this change to take place.

The value of the limestones of the southern part of the peninsula, for the purposes of building, as also for the manufacture of lime, was mentioned in a previous report upon the geology of that district of country. As has already been stated, these limerocks do not reach within many miles of the coast of Lake Superior, and it is certainly to be regretted that the shore of the northern portion of the peninsula is destitute of this important material. Nor have I seen any marls of sufficient extent, in the district, to admit of application to any of the purposes to which it is applicable, or to supply, even in part the deficiency in limestone. All the lime which would appear to be capable of being applied to practical purposes, is

that of the calcareous spar, composing the veins traversing the sandrock, and these are not only rare, but they are also of very limited extent.

MINERALS AND MINERAL VEINS.

In considering this portion of the subject, I propose to treat the minerals of the different formations separately, so far as the same can be done, and although this method will necessarily cause some repetition, it will enable me to show, more perfectly than could otherwise be done, the connection between those minerals that may be regarded as of practical value, and the rocks to which they belong.

As a whole, the rocks of the upper peninsula are deficient in *number* of minerals, though some few individual *species* occur abundantly.

Minerals of the Primary Rocks.

The following list can by no means be regarded as perfect, but it will serve, at least, to convey an idea of the small number of minerals which are found in connection with the rocks of this group.

Schorl,	Mica,
Tourmaline,	Feldspar,
Hornblende,	“ red,
Actynolite,	Quartz.

Minerals of the Metamorphic group of Rocks.

Quartz, common,	Iron, scaly red oxide of,
“ milky,	“ hæmatite,
“ greasy,	“ pyritous,
“ tabular,	Steatite,
Serpentine, common,	Novaculite.

Of the minerals enumerated as occurring in the metamorphic rocks, the milky variety of quartz is abundant, sometimes composing almost entire ranges of hills. The novaculite is also abundant, but of a coarse variety. This last is associated with the talcose slates. The remaining minerals appear either disseminated, or forming druses in the quartz rock, though sometimes they occur in thin beds or veins, in the talcose slate, which beds conform to the line of cleavage of that rock. Although the hæmatite is abundantly disseminated through all the rocks of the metamorphic group, it does not appear in sufficient quantity, at any one point that has been examined, to be of practical importance.

Minerals of the Trap Rocks.

Quartz, common,	Steatite, common,
“ smoky,	Asbestos,
“ milky,	Amianthus,
“ greasy,	Calcareous spar,
“ radiated,	Copper, native,
“ mamillary,	“ pyritous,
“ drusy,	“ black,
“ amethystine,	“ red oxyd of,
Chalcedony,	“ azure carbonate of,
Carnelian,	“ green carbonate of,
Jasper,	“ “ ferruginous,
Agate, common,	Lead, sulphuret of,
“ fortification,	“ carbonate of,
Augite,	Iron, pyritous,
Actynolite,	“ red oxyd of,
Serpentine,	“ hydrate of,
“ pseudomorphous,	“ silicate of,
Chlorite, common,	Manganese, ferruginous ox. of,
“ earthy,	Silver, native, (very rare.)

Since a consideration of the minerals contained in the trap, will also involve a portion of those embraced in the conglomerate, the mixed rock, and red sandrock and shales, I will, before referring minutely to those of the trap rocks, lay before you a list of those which occur most frequently in the sedimentary rocks last mentioned. The fact that veins of mineral matter, traversing the trap, are frequently continued across the several sedimentary rocks, and that dykes are of frequent occurrence in these latter rocks, would lead to the inference, that there would be a considerable degree of resemblance in the character of the minerals embraced in these dykes and veins, in both the trap and sedimentary rocks, and to a certain extent, this inference would be true; but it should be borne in mind, as has already been stated, that the veins, in traversing the several upper rocks, undergo very great changes in mineral character.

Minerals of the Conglomerate, Mixed Rock and Red Sand Rock.

Calcareous spar,	Copper, native,†
Quartz, common,	“ pyritous,†
“ milky,	“ blue carbonate of,†
“ drusy,	“ green carbonate of,†
Chalcedony,*	“ earthy green carb. of,†
Carnelian,*	“ black,†

*Occasionally occurring among the pebbles constituting the mass of the conglomerate.
† Chiefly in those portions of the veins traversing the conglomerate.

Jaspar,*
Agate,*

Zinc, silicious oxyd of,
" carbonate of,
Iron, pyritous,
" black oxyd of, (cemented iron sand,)
" red oxyd of,
" hydrate of,
" silicate of.

Mineral veins of the Trap, Conglomerate, &c.

In order to render the subject of the mineral veins traversing the above rock, so far intelligible as may be in my power, I have already been particular to define, as far as could be done without maps and sections, the relation which the trap rocks, together with the superincumbent conglomerate, mixed sand and conglomerate and red sandrock bear to each other, and it will be necessary, in considering the mineral contents of these rocks and the veins traversing them, to keep this relation constantly and clearly in view.

It will be recollected, that the north-westerly range of hills, commencing at the extremity of Keweenaw point, and stretching from thence in a south-westerly direction into the interior, were referred to as being more clearly of trapose origin than either of the other ranges, and that the rock of the southerly portion of this range is either compact greenstone or altered syenite, while that of the northerly flank is almost invariably either an amygdaloid or a rock approaching to toadstone.

The several ranges of hills to the south of that last alluded to, are either well formed, compact greenstones, altered syenite, or, (as we approach the primary range,) imperfectly formed granites. So far as the several ranges of hills, lying south from the northerly range, are concerned, they would appear to be, as a whole, deficient in minerals, and the rocks are not apparently traversed by veins or dykes of any more recent date than that of the uplift of the northerly trap hills.

Veins clearly of a date posterior to the uplift of that portion of the trap rock last mentioned, are of frequent occurrence, and these veins not only traverse a portion of the trap range, but also pass into the conglomerate, and sometimes completely across the three sedimentary rocks, immediately above the trap, thus having an unbroken length of several miles. The class of veins to which I now allude, where they occur in a connected or continuous portion of the range, rarely vary more than 12° to 15° from a right angle to the line of bearing of the sedimentary rocks, and in pursuing this course, they necessarily cut across the dykes of trap before alluded to as frequent-

*Occasionally occurring among the pebbles constituting the mass of the conglomerate.

ly appearing between the strata, and conforming to the dip of the lower sedimentary rocks.

That the veins under consideration belong to a single epoch, is inferred from the fact, that none have been noticed with other veins crossing them, as also for the reason that none have ever been noticed with dislocations, heaves or disturbance of any kind, save what may be referred to causes connected with their immediate origin.

That these veins must be regarded in the strictest sense as true veins, cannot be doubted, and that their origin or source, over the extended district alluded to, has been the same, is inferred from the perfect identity of their mineral contents; for a description of one of these true veins may be said to be essentially a description of the whole. Thus, while the mineral contents of the different portions of the same vein change as the rock traversed changes, the corresponding portions of different veins almost invariably bear a striking and close resemblance to each other.

These veins, as has already been stated, where they traverse connected ranges of the trap, are regular in course and direction, but when they are connected with a single uplifted knob of that rock, they are irregular and can scarcely be defined, appearing, in the latter instance, rather as matter injected into the fissures of a shattered mass of rock, than as connected veins.

The importance of carefully studying the relation which these veins bear to the rocks which they traverse, as also the relation which they bear to the numerous trap dykes, together with the few cotemporaneous veins noticed in the trap, is very much increased by the circumstance, that these veins are more or less connected with, or rather contain, metallic materials, which, it may be fairly inferred, will hereafter become of very considerable practical importance. In fact, so far as we may be enabled to judge from the examinations already made in this district of country, it is confidently believed that most, if not all the metalliferous veins of the upper peninsula, belong to veins of the epoch of those under consideration. It is true that native metals, more particularly copper, are sometimes found, in place, occupying the joints or natural septæ of the greenstone, but in these instances, the amount of metal is always comparatively small, and, with one or two exceptions, I have invariably been able to establish some connection between the native metal occupying these joints and the termination of some metalliferous vein that traverses other portions of the rock not far distant, and it is believed that the metal filling these joints has invariably resulted from the action of causes precisely analagous to those which have placed similar metals in the veins to which I have alluded.

The earliest as well as all travelers, who have visited the district of country under consideration, have not failed to make frequent allusion to the loose masses of native copper that have been occasionally found scattered over it, nor has any one failed to allude to the large boulder or loose mass of that metal upon the Ontonagon river. Almost invariably, the opinion has been expressed, from the frequent occurrences of these masses, that the metal must be abundant in the country. But, after all, the true sources from which these masses had their origin, or the relation which they held to the rocks of the district, would appear to have never been understood; and all, or nearly all, that was known of their true relations, was left to conjecture. The result of this has been, that while some have excessively magnified every thing connected with a subject of which, in truth, nothing was known, another class, equally far from what is really true, have regarded these masses of native copper as boulders transported from high northern latitudes.*

As far back as 1831 and 1832, I had occasion to pass, no less than three times, along the south coast of Lake Superior, as also to ascend several of the important tributaries of that lake, and during these years, I passed by three different routes, widely separated from each other, completely across to the Mississippi river. It is true that these journeys, made through a complete wilderness, uninhabited except by savages, were necessarily made under circumstances that admitted of only very general observations; but the result of these previous examinations have proved of immense service to me, in aiding the labors of the past season. I allude to these journeys and examinations at this time, in order to show you the difficulties by which a full understanding of the subject under consideration is surrounded, for I became satisfied at that time, not only that the subject was not understood by the mass of those who had traversed the country, but that even the natives of the country had no knowledge of the true sources from which the transported masses of copper had their origin.

During the time of the examinations referred to, a bare glimmer of light was thrown upon the subject by an examination of some small masses of copper, found occupying the joints of the greenstone; as also by the examination of a single vein in the conglomerate, containing the ores of copper, which has

*The vast area of country over which the boulders of native copper, from the district under consideration, (together with its westerly prolongation,) have been transported, is worthy of remark. They are not of unfrequent occurrence in the sand and gravel of the southern peninsula of Michigan, and since the commencement of the geological survey, many of these masses have been met, some of which weigh from seven to eight pounds. In the vicinity of Green bay, a mass was discovered, some ten years ago, which weighed 140 pounds, if my memory serves me correctly. Loose masses of a similar character have been met with in various other portions of Wisconsin, as also at various points in Illinois, Indiana and Ohio. In these cases, the occurrence of these masses of native copper are no more indications of the existence of veins of the metal in the immediate vicinity, than are the immense numbers of primary boulders scattered over the southern peninsula of Michigan indications of the existence of primary rock in place, in the district where they are found.

since been found to be the termination of a vein that is somewhat obscurely continued from the trap region. While these examinations were sufficient to enable me to draw the inference that the masses of native copper came chiefly, if not wholly, from the trap, and more rarely from those sedimentary rocks resting immediately upon it, it was supposed that this occurrence would follow the general law, and that it, together with the other ores of the metal, would occur in greatest abundance near the line of junction of this rock, with the overlaying sedimentary rocks. Nothing, or at least very little, was known of the true extent or range of the trap rocks, and the very great inaccuracies in the published maps of the country, rendered it almost impossible to apply even the data on hand to such purpose as to relieve the embarrassment.

With a full knowledge of these difficulties, I determined, during the past season, to endeavor to surmount them by so far adding to our geographical knowledge of the coast of the lake and its immediate vicinity, as to enable me to place whatever geological observations of importance might be made, in such condition that the relations of the several parts might be understood. Having sufficiently accomplished this, I proceeded to a very minute examination of the several rocks overlaying or resting against the trap, together with a determination of the thickness of the several members and their rate of decrease or wedging to the east. With these data I was enabled, by noting the dip of the rock upon the coast, to determine, with sufficient accuracy for the purposes to which the rule was to be applied, the line of junction between the trap and conglomerate rocks. This rule, when put in practice, enabled me to decide, with a very considerable degree of certainty, this line of junction, when the rocks were covered with a very considerable thickness of detrital matter, and when so covered, I was enabled, by traversing the country, on the line of bearing of the upper rocks, the more readily to gain access to such points as would admit of examination.

These observations soon showed me that this line of junction between the trap rock, and the south edge of the conglomerate, instead of pursuing a course parallel to the coast, only continued its parallelism for a few miles westerly from the extremity of Keweenaw point, after which, for a long distance, it recedes from the coast rapidly. These facts served to explain in part, why the subject of the origin of the masses of copper had remained a mystery, for the country through which this line passes is hardly ever passed over, even by the Indians, and probably large portions of it have never been passed over by whites, but in addition to this, the obscure character of the metalliferous veins is such, that they would scarcely attract the observation of the traveler whose attention was not called

especially to the subject, for many of the richest ores are so far from having the appearance of the pure metal, that they would be the last suspected to contain it in any form.

That the connection of these ores with the containing rocks was not understood by the English mining company, whose attention was turned to this subject, at an early day, is to be inferred from the fact that they commenced their operations at Miners' river, where the rock is the upper or grey sandstone, which has never been observed to contain mineral veins; and also on Ontonagon river, near the mass of native copper, at which point a shaft was commenced and carried about 40 feet through a reddish clay, at which point the red sandrock was reached. Now, although the metalliferous veins sometimes pass from the trap across the red sandstone, these veins in the red sandrock have never been noticed to contain any other ores than those of zinc and iron, unless it be at the immediate point where the vein crossing comes in contact with a dyke of trap, which condition does not exist at the point alluded to, on Ontonagon river. What indications could have induced these Quixotic trials at the points where they were commenced, is more than I have been able to divine, and as might have been anticipated, the attempts resulted in a failure to find the object sought.

Having thus, in a general manner, set forth the obscurity by which the subject of the true source of the transported masses of native copper has been surrounded, together with some of the reasons which have served to prevent its being fairly understood, I will now proceed to a general sketch of the metalliferous veins of the district, so far as the same have been examined; premising, that our knowledge of them is still deficient in very many important particulars, which can only be supplied by a careful and continued examination of the subject, which, in fact, can only be said to be just commenced.

I have had occasion to refer to the outer or northerly range of hills, or those from which the metalliferous veins may be said to spring, as being composed of trap rock, and lest what has been said may not be fairly understood, I will repeat, that the more southerly part of the range is uniformly composed of compact greenstone, under which head I not only include true greenstone, but also those forms of altered granular gneiss and gneissoid granite, which sometimes are associated with it, while the outer or northerly portion of the same range is usually composed of an amygdaloidal form of trap. The cells of the amygdaloid are usually filled with the different varieties of quartz, carnelian, chalcedony and agate, and sometimes, though more rarely, with native copper, or with calcareous spar,

though they are sometimes entirely empty, constituting a perfect toadstone.

The metalliferous veins cross this range of trap, usually very nearly at right angles to the prolongation of the hills, and are frequently continued in the same course, across the upper or sedimentary rocks, thus crossing the latter at an angle varying but little from their line of bearing. While the continuity, of course, of the vein, may remain perfect in its complete passage from the greenstone across the several members of the conglomerate, mixed and red sandstone rocks, the character and mineral contents of the vein undergoes essential change, and not only does the vein appear to be influenced in its mineral contents, but also in its width, for, as a general rule, the width of the vein increases as we proceed northerly, or from the greenstone. Thus, a vein which may appear of only a few inches in width, or as a bare line in the southerly or greenstone portion of the range, increases in width rapidly as it approaches and passes across the amygdaloid, and at or near the line of junction between the amygdaloid and the sedimentary rocks it will frequently be found to have attained a thickness of several feet, while in its passage across the sedimentary rocks it is usually either still further increased in width, or becomes so blended with the rock itself, as to render it difficult to define its boundaries.

These metalliferous veins, like those which occur under similar circumstances in other portions of the globe, do not continue uninterruptedly of any given width, for great distances, nor is their width increased regularly, for they frequently ramify or branch off in strings, that pursue a course generally somewhat parallel to the general direction of the main vein, and which eventually again unite with it. Sometimes these ramifications or branches destroy, as it were, for a considerable distance, the whole vein; but they at length unite again, and the main vein is, after their junction, as perfectly developed as before.

While traversing the most compact, southerly portions of the greenstone, the veins are most frequently made up of a very compact and finely granulated greenstone, sometimes associated with steatitic minerals and silicate of iron, under which circumstances they usually are destitute of any other metallic mineral, but occasionally, instead of the materials above mentioned, their place is supplied by native copper, without vein stone or matrix, and usually free from nearly all earthly impurities, but almost invariably incrustated with oxyd, or carbonate of the metal. Those portions of the vein traversing the greenstone, in which native copper occurs, under the circumstances above mentioned, are invariably thin, rarely exceeding three to four inches in thickness, and usually conside-

rably less, and they are liable to very considerable variation in width, from the divergence caused by the vein traversing the joints of the rock, where these joints produce the same character of change as is produced by the ordinary ramification of a vein.

As these metalliferous veins traverse the northerly portion of the range, or approach the sedimentary rocks, they undergo a gradual change in width as well as in mineral character, and it has been noticed that where the amygdaloid is most largely developed, the vein, as a general rule, has not only a greater width, but also has its mineral contents more perfectly developed, a circumstance which might fairly have been inferred from the fact that those points where the amygdaloid occurs most largely, may be regarded to have been so many centres of intensity of action, at the time of the original uplift of the range, from which circumstance they would remain in a softened state, or in such condition as to admit of the more perfect formation of these cross veins for a longer space of time after that condition had been passed at other points.

In the outer or amygdaloid portion of the rock, the vein is almost invariably accompanied by a veinstone of quartz, involving all the varieties before mentioned, as associated with the trap rocks, which quartz, though occasionally it occurs massive, of several feet in width, usually appears in the shape of a series of irregularly ramifying and branching minor veins, that may be said to constitute the main vein. These subordinate veins of quartz, which may be stated as the true veinstone, vary from a mere line to several inches in thickness, and in the aggregate they may be said to constitute from one-third to one-half the total thickness of the vein. In their branches and ramifications, they sometimes include portions of the rock which they traverse, at other times they embrace imperfectly formed steatite, with silicate, carbonate and red oxyd of iron,* and occasionally, though more rarely, it is associated with carbonate of lime, usually assuming the form of an opaque rhombic spar.

As the main vein traverses the conglomerate and overlaying rocks to, and including the red sandstone, these veins, as a general rule, undergo still farther changes, for very soon after entering the conglomerate, the veinstone changes from its quartzose character, and is made up, either wholly, of calcareous matter, mostly rhomb spar, or of this mineral with occasional ramifications of quartz. The whole usually including, and sometimes investing fragments of the conglomerate or the pebbles of that rock, separated.

As the vein is continued still farther in the direction of, and into the red sandstone, these changes are still noticed, and

* The latter closely resembling the Gossau, of the Cornish miners.

eventually the vein is found to be composed either entirely or mostly of calcareous spar, and eventually so completely is its metalliferous character lost, that it would not if examined singly, be suspected to be any portion of a metalliferous vein.

The metalliferous character of these veins is most largely developed almost directly at or near to the line of junction of the trap and sedimentary rocks, and they rarely continue, without considerable change, for a greater distance than one-fourth to one-third of a mile, on either side of the line, though a few veins were noticed in which, in the southerly or trap extension, the character of the vein continued for a distance of over a mile, nearly unchanged, while in its passage through the conglomerate, for half that distance, its character was also perfectly preserved.

The mineral character of the veins is somewhat varied in those having different degrees of thickness, though it is difficult if not impossible, to lay down any rule which would characterize this change. The different veins vary very greatly in width, ranging from a mere line to 14 or 15 feet—the greatest observed width of any single vein.

In the descriptions of the veins given above, I only intend to include those which are most perfectly developed, for, in addition to these, there are also many which are imperfectly formed and short, and in which many of the above characters are in part or entirely wanting. These latter are usually of little practical importance, and thus far have been comparatively little examined.

Of the metallic minerals occurring in those portions of the *true* veins which traverse the trap rocks, together with that portion of the conglomerate immediately resting upon or against the trap, by far the most important, consists of the several ores of copper, with which iron occurs, disseminated in the forms before described, and occasionally, though very rarely, native silver has been detected, associated in the same vein. After as minute an examination of the subject, as circumstances will permit, I am led to the conclusion, that the only ores of the metallic minerals, occurring in those portions of the veins, which traverse the rocks last alluded to, which can reasonably be hoped to be turned to practical account, are those of copper.

In these portions of the veins, the metal referred to, occurs very frequently in the form of native copper, with which are associated the red oxyd, azure carbonate, green carbonate, and more rarely what may be denominated copper black, and still more rarely, pyritous copper. *None* of these have been noticed in a crystalline form.

It must not be imagined that these several minerals make up the whole or even any very considerable portion of the entire length and breadth of the veins, in which they occur, for they

are distributed in bunches, strings, and comparatively narrow sub-veins, in a manner precisely analagous to that in which these ores are usually distributed, in similar rocks, in other portions of the globe. The quartz veinstone, before described, has always so much of the green tinge communicated by the carbonate of copper, that it cannot fail to be detected; but the presence of disseminated native copper, in this veinstone, would, at first, hardly be suspected, and it is not until a fresh fracture has been made, and the mineral closely examined, that the numerous dark points and minute threads are discovered to be copper in a native state. Large portions of this quartz veinstone, (when the included metal can scarcely be detected by the naked eye,) when examined with a glass, are found to contain very delicate threads of native copper, that traverse the quartz in every possible direction, and so completely is this latter mineral bound together, that it is fractured with difficulty, and its toughness is very greatly increased.

The specific gravity of this veinstone is very considerably above that of ordinary quartz, and usually, the difference is so considerable, even in those masses where the copper can scarcely be detected by the naked eye, as to be apparent to even the most careless observer. But in addition this finely disseminated condition of the native copper in the veinstone, it is also disseminated in a similar manner through the rocky matter embraced by the veinstone and in the amygdaloid and conglomerate portions of the rocks, it sometimes extends, for a distance of from two to three feet into the rocky matter on either side of the veins, sometimes completely, or in part, filling the cells of the amygdaloid rock.

The conditions above described refer to the main portions of the veins only, while there are other portions in which the copper appears to be concentrated in larger masses, constituting bunches and strings, and in which places the sides or walls of the veins are sometimes wholly made up of thin plates of native copper. In these portions of the metalliferous veins where the metal appears, as it were, to be concentrated, it also occurs, much in the form before described, except that the masses of metal vary from the merest speck to that of several pounds weight. In opening one of these veins, at a concentrated point, the observer, unless he had previously examined other portions of the vein, would be led to erroneous conclusions as to its richness, a source of error which cannot be too strongly guarded against; for while the vein, for a short distance, may be found to be exceedingly rich in mineral, the mineral in another portion of the vein may either wholly or in part disappear, a condition which is similar to that observed in those veins of copper that have been extensively worked and

found to be most productive, on the continent of Europe and the Island of Great Britain.

The excess of native copper, (compared with the other ores,) which occurs, in these portions of the veins, is a peculiar feature, for it may be said, in truth, that other ores are of rare occurrence. In those portions of the veins traversing the trap, and where other ores do occur, it is usually under such circumstances as to favor the presumption that their origin is chiefly from that which was previously in a native form; for the carbonates and oxyds, almost invariably appear either investing the native copper, or intimately associated with it, though they sometimes appear in distinct sub-veins. Pyritious copper is so rare, in connection with the trappean portions of the veins, as scarcely to deserve notice.

I have already stated that native silver, occasionally, though very rarely, occurs in the trappean portions of these veins, intimately associated with the copper, but it is in so minute quantities as to render it probable that it will not prove of any practical importance. Other mixed compounds of this metal occur so rarely as scarcely to deserve notice.

Leaving the trap rock, the character of these veins, as they traverse the conglomerate, undergoes important changes; for not only does the veinstone become gradually changed, from quartz to calcareous spar, but the amount of native copper diminishes, and its place is either supplied wholly or in part by ores of zinc and calcareous spar, or wholly by this latter mineral. There are, however, occasional exceptions to this general rule, for occasionally the place of the native copper in the veins, in their passage through the conglomerate, is supplied by a variety of complex compounds of the same metal, which compounds are of exceeding interest; but this change would appear always to be intimately connected with, or to bear some relation to, the dykes of trap which traverse the conglomerate rock. Several instances of this kind were noticed upon the northerly side of Keweenaw point, either directly upon or near to the coast, as also at several other places in the interior, westerly from Keweenaw point. A vein, which may without doubt be referred to as one of this character, (though in consequence of intervening bays and lakes between it and the ranges to the south, its connection with the main range has not been seen,) will serve to illustrate the character referred to.

This vein, which reaches the immediate coast of the lake, upon the easterly cape of the bay known to the voyageurs as the Grande Marrais of Keweenaw point, terminates, so far as examinations can be made, in the coarse conglomerate rock. The coast of the lake, for many miles on either side, is made up of abrupt cliffs of a similar rock, the rock as usual, being

made up of coarse rolled pebbles of trap, chiefly cemented with calcareous matter, which is usually associated, more or less, with the red oxyd of iron. Immediately south of the coast a heavy dyke of trap traverses the conglomerate, which dyke corresponds in position with line of bearing and dip of the conglomerate rock.

The vein, which, at its termination upon the immediate coast of the lake, has an extreme width of about ten feet, may be traced, in the bed of the lake, in a direction north 5° east, for a distance of several rods, after which, in consequence of the depth of water, it is completely lost. This vein, at the point where it appears upon the coast, may be said to be in a concentrated state, or in a condition analagous to that before described, where the native copper occurs in the condition of bunches and strings, though the condition in which the metallic minerals occur, is essentially different from that in the trap; for, instead of native copper, we have several mixed forms of the green and blue carbonates of copper and copper black, more or less intimately associated with calcareous spar, and in the adjoining rock, and in small ramifying veins, occasional small specks and masses of native copper, weighing from 1 to 3 oz., occur, but these are by no means abundant. No quartz occurs as the veinstone, and none of the ores have been noticed in a crystalline form.

It has already been stated, that these true veins, in traversing the conglomerate, frequently almost lose their character, and it becomes difficult to define their absolute width, or in other words, it would appear as if, at the time of the formation of the veins, the conglomerate had not been perfectly cemented, the result of which would be, that the mineral matter, which, under other circumstances, would constitute a perfect vein, would frequently appear in only an imperfect one, or the mineral which would, under other circumstances, make up the vein itself, may have been injected laterally through the interstices of the rolled masses constituting the conglomerate, in which case the mineral would, in fact, take the place of the ordinary cement, thus simply investing the pebbles of the conglomerate. Now, although at the point under consideration, a wide and remarkably distinct vein is developed, the rock, for many feet on either side, has the interstices between the pebbles filled wholly, or in part, with various mixed and irregular forms of the ores, accompanied by calcareous matter, as before stated, and with occasional specks and small masses of native copper.

Those veins traversing the conglomerate take on a similar character, to a greater or less extent, rather frequently, but the place of the copper is more usually supplied by the siliceous oxyd, and more rarely by the carbonate of zinc, which com-

pounds, sometimes may be seen forming a perfect or partial cement to the rock, for considerable distances on either side of the main vein. These ores of zinc, like those of copper, are uniformly amorphous, and almost invariably more or less associated with some form of carbonate of lime, with which they may, under some circumstances, unless closely examined, be confounded.

Although these copper and zinc ores occasionally appear in considerable quantities, in those portions of the veins traversing the conglomerate, they usually embrace or simply encrust portions of the rocky matter; or rather the rocky matter and those ores appear to be coarsely and mechanically mixed. These veins furnish beautiful cabinet specimens of the blue and green carbonates of copper, and more rarely of pyritous copper, together with the other varieties mentioned.

Having already devoted a larger space to the consideration of these veins than had been intended, I will simply add, that in pursuing their course northerly, across the mixed rock and the red sandrock, their mineral character is nearly or quite lost, the veins as before stated, being made up either entirely of calcareous spar or of that material containing very meagre ores of zinc.

The district of country to which these veins have been referred, thus far, only comprises the ranges of hills south of Lake Superior, but veins of a very similar character, and of similar mineral contents, also occur upon Isle Royale. The order and changes in the character of the veins upon Isle Royale is necessarily reversed, or in other words, the southerly point of the vein corresponds to that of the north point in the district south of Lake Superior. The mineral veins of Isle Royale have not been examined with sufficient care to enable me to determine with much certainty, their average width or value. Those examined were mostly narrow, the widest not exceeding eighteen inches, but in these the mineral contents are essentially the same as in those upon the south side of the lake.

Native copper, in very thin plates was occasionally noticed occupying irregularly the joints of the compact greenstone of Isle Royale, but invariably in comparatively small quantities. It should, however, be noticed of Isle Royale, that the veins, so far as examined, are less perfectly developed in their passage across the conglomerate, and that they very rarely contain any traces of zinc.

Upon the north shore of the lake, no attention was given to the subject of mineral veins, but from the character of the geology of that district, it may be inferred that they will also be found in portions of it, and that, where they do occur, they will

be uniformly either directly upon or not far from the coast of the lake.

In addition to the *regular* veins already described, irregular veins frequently occur, traversing the whole, or portions, of the outliers of trap, or those knobs which appear to have been elevated singly; and although these veins may without doubt, be referred to the same epoch as the regular veins before described they nevertheless frequently differ considerably in mineral contents.

The limits of the present report will not permit a separate description of these several distinct trap knobs. I will therefore confine my remarks to that already referred to, as occurring upon the south coast of Lake Superior, immediate northwest from Riviere Des Morts, and which forms the promontory known as Presque Isle.

In nearly all those portions of this knob, where the trap, conglomerate and sandstone, are exposed in such a manner as to permit examination, each of the rocks are seen to be traversed by innumerable irregular ramifying veins, which in the sandstones are made up of quartzose and calcareous matter; but many of which, near the junction of the igneous and sedimentary rocks, are metalliferous, and this metalliferous character is more fully developed as the veins are extended into the trap rock.

The metalliferous portions of these veins, rarely exceed three to four inches in width, and they ramify in such a manner that the mineral uniformly occupies situations similar to bunches or strings, at the junction of the ramifications. The minerals contained in the metalliferous portions of the veins, are sulphuret and carbonate of lead, earthy, green carbonate of copper, pyritous iron, and more rarely, pyritous copper. Occasionally there is a quartzose, or mixed quartzose and calcareous veinstone; but more usually the several metallic minerals are blended in a base of rocky matter. The sulphuret of lead is distributed in the form of small cubic crystals, while the other metallic minerals are usually distributed either in irregular masses, or investing portions of the rocky matter. These associations are referred to, as showing the character which these irregular veins assume, rather than from any supposed value which they may possess for practical purposes.

In addition to the minerals referred to, the trap of Presque Isle occasionally contains asbestos, common serpentine and imperfect agates; the two former minerals usually occupying the narrow joints of the rock.

Before referring to the economical considerations connected with the veins which have been described, I will briefly refer to

another situation in which the ores of copper have been observed in intimate connection with the trap range of rocks.

The southerly side, or greenstone portion of the trap range, appears to have been elevated in such a manner as to have caused but little disturbance to the sand rock lying between that and the range of simply altered rocks lying still further to the south; but near to the junction of the sandrock and greenstone there is usually a red slate resting against the trap, and which may be said to fill up, in a measure, the irregularities in the ranges of hills. This slate, which is sometimes seen of 100 to 200 feet in thickness, though usually it appears as a mere band, is traversed by irregular and imperfect veins, of what may be denominated a ferruginous steatite, containing placentiform masses of greasy and milkish quartz, that sometimes contain more or less of the ores of copper. The earthy carbonates of copper are also sometimes so intimately connected with these veins of steatitic matter as at first to be scarcely recognized. More rarely, distinct, very thin veins of green carbonate of copper occurs, well characterized, in this red slate, though these veins are never of any great length. The red shale extends, more or less perfectly, along the whole length of the trap range, skirting that range of hills upon the south, but I have not yet been enabled to devote sufficient time to its examination to enable me to determine whether any portions of these veins can be regarded as of practical importance. The examinations which have been made would lead me to look unfavorably upon these veins, and I regard them as having an origin completely distinct from that of the veins which traverse the northerly escarpment of the trap rock.

Having thus considered all the general circumstances under which the several ores of copper, zinc, lead, iron, manganese and silver have been noticed, in connection with the trap rock and the sedimentary rocks, immediately resting upon it, it becomes important to consider how far inferences may be drawn from these examinations, as to their occurrence in such quantities as to be of practical importance. I have already stated that so far as regards the ores of lead, iron, manganese and silver, I am led to conclude that at none of the points examined do they occur in veins, or otherwise, sufficiently developed to warrant favorable conclusions as to their existence in sufficient quantities to be made available, and from all that is now known of the country, I am led to infer that neither of these, unless it be iron, will be so found.*

The examinations which have thus far been made of those portions of the veins containing ores of zinc, have not been extended sufficiently to enable me to determine with much satis-

* These remarks are intended to apply directly to the trap region. Beds of bog iron ore occur, east from Chocolate river, which probably may at some future day be profitably worked.

faction, their extent as a whole. At several points in the veins, these ores are sufficiently abundant to admit of being profitably worked, but I would be unwilling, from an examination of a few points, to attempt to determine the character of the whole.

In considering the practical value of the copper ores of the upper peninsula of Michigan, where we are as yet compelled to judge from our examination, of what may be said to be the simply superficial portions of the veins, we can arrive at no safe conclusions, except by comparison of the district with those districts similarly situated, which have been extensively worked in other portions of the globe. Comparisons of this character, to be really useful, must necessarily be sufficiently minute to enable us to understand the relations which the ores in the districts compared, bear to each other, in all respects, which circumstances renders it necessary that a degree of minute information should be at hand, that is not at all times to be obtained. As the information on hand, with respect to the copper and tin veins of Cornwall, England, is more minute than that of any mineral district known, I propose, in order to avoid confusion, to confine my comparison to this district, simply premising, that however closely the two districts may resemble each other in character, it does not follow, as an axiom, that because the district with which we compare our own has been largely and profitably productive, that of Michigan must necessarily be so too, for it will be seen, as the subject is pursued, that there are not only several points in which it is impossible with our present knowledge of that of Michigan, to institute comparisons, but there are also some points on which there is a considerable degree of discrepancy.

The comparison instituted, in the main, is intended to refer rather to the character and contents of the mineral veins of the two districts than to the geology, although some general reference becomes necessary to the geology of the districts, to render the comparison perfect. The topography of the Cornish district bears a close resemblance to that of Michigan; both districts being marked by their irregular and broken outline, and by the occurrence of more or less frequent, nearly insulated knobs, rising to a considerable height above the elevation of the general ranges.

Although the older rock of Cornwall, or that from which the metalliferous veins of the district may be said to have their origin, is more distinctly granitic than that of the metalliferous region upon Lake Superior, the elements of which the rocks are composed, may be regarded as essentially bearing a very close resemblance; a resemblance, which it is conceived, would have been still more perfect had the granitic rocks of Cornwall been subject to the action of secondary causes similar to those

of the region under consideration. The rocks resting upon or against the granitic rocks of Cornwall, consist of clay slates, hornblende rocks, &c., which bear little real analogy to the rocks resting directly upon the trap of Lake Superior, but it is conceived that the composition of these upper rocks has little bearing upon the origin of the metalliferous veins, and may be regarded as in a measure unimportant; and however much these rocks may differ, they are traversed alike by the metalliferous veins of the lower rocks in such a manner, that the close resemblance cannot be mistaken.

It is a matter of history that the ores of tin have been, more or less, extensively raised in the mineral district of Cornwall, from the earliest settlement of the Island of Great Britain, but the working of the veins of copper at an early day, does not appear to have been carried on to any very considerable extent. The great importance to which the produce of copper from the Cornish veins, (in a district which, compared with the mineral district of our own state, is of very small dimensions,) has arisen, will be shown from the accompanying table, which I have reduced from the official returns, included in the several years, and which table, it will be seen, shows for a series of years, the average annual amount of copper produced from the ore, the average amount for which it sold, together with the amount per cent of copper contained in the ore, and the average value of the copper, per pound, at the smelting house. This table, which has been drawn with great care, from data that can scarcely lead to incorrect results, will not only serve to show the large aggregate amount of the metal produced, but it also shows, from the low average per cent of metal contained in the ores, (if we had no further knowledge upon the subject, that much capital must be required for, and a large amount of labor applied to the raising and smelting of these ores; a circumstance which should be carefully borne in mind, in all that relates to the mineral district of Michigan.

Table showing the average annual produce of the Copper Mines of the County of Cornwall, England, from 1771 to 1822.

YEARS.	Average No. of tons of ore per year.	Av. No. of tons cop- per produ- ced per year.	Av. amount per year for which sold.	Av. per ct. of copper produced from the ore.	Av. val. of the copper per lb.
1771 to 1775—5 years,	28,749	3,449	\$846,283	12	^{c. m.} 10 9
1776 to 1780 5 “	27,580	3,309	826,609	12	11 1
1781 to 1786 6 “	34,354	4,122	962,380	12	10 4
1796 to 1802 7 “	51,843	5,195	2,125,046	10	18 2
1803 to 1807 5 “	70,923	6,160	3,174,725	8	23
1808 to 1812 5 “	70,434	6,498	2,886,835	9	12 9
1813 to 1817 5 “	82,610	7,272	2,878,723	8 8	17 6
1818 to 1822 5 “	94,391	7,757	3,111,811	8 2	17 9

The general resemblance in the mineral contents of the copper veins of Cornwall and those of Michigan, is, for the most part, very great, though in some respects there is a considerable discrepancy. It should, however, be remarked that some difficulty exists in comparing the mineral veins of Cornwall, where several of them have been worked to depths, varying from 1,000 to 1,500 feet, with those of Michigan, where the examinations are nearly superficial.

In making these deep excavations, not only in the county of Cornwall, but also in the copper districts of Bohemia, Hungary, Silesia, Transylvania, Saxony, &c., (some of the veins in the latter districts having been explored to a depth very considerably greater than those of Cornwall,) an immense mass of facts has been accumulated, with respect to the general formation and mineral character of veins, or lodes of copper, which facts have led to an understanding of many of the contingencies connected with its associations, so universal, that when applied to this mineral, they may be regarded as general laws, that may fairly be inferred to govern, with more or less certainty, all those lodes or veins which have similar geological relations. Though a general consideration of those relations of the veins of other countries, may perhaps be regarded as somewhat foreign to the present report, I deem it more advisable to refer to these general laws in such a manner as to leave the reader to judge, by comparison, the condition in which the ores of Michigan may be fairly inferred to occur, rather than to draw conclusions directly; and in so doing, it will also become necessary to refer to some of the characters of mineral veins, or lodes, in general.

Veins are usually divided into two general orders, viz: “*cotemporaneous veins*, or those which were formed at the same

time as the containing rock, and *true veins*, whose formation is supposed to be subsequent to that of the rocks which are contiguous to them.” A *true vein* may be defined to be “the mineral contents of a vertical or inclined fissure, nearly straight, and of indefinite length and depth.”* The contents of a true vein, as a general rule, differ widely from the character of the rocks which it intersects, though this does not invariably hold good, and the vein also, as a general rule, has well defined walls.

The contents of cotemporaneous veins, bear a much closer resemblance to the rocks which embrace them, and as a general rule, they are shorter, more crooked, and less perfectly defined, than true veins.

The metalliferous veins being contained under the head of true veins, it is to these that the whole of my remarks will be directed.

Metallic veins are the repositories of most of the metals excepting iron, manganese and chrome, which occur more frequently and abundantly in beds than in veins. The thickness of metallic veins varies from a few inches to many feet, and the same vein also varies in thickness in different parts of its course, sometimes contracting to a narrow string of ore, and then expanding again to a width of many feet. The deposits of metal in the veins are as irregular as the widths of them, and so much so as to render the profits of mining proverbially uncertain. Ore is generally found to occupy certain portions of the veins only, differing constantly in extent, whether the length or depth on the course of the vein be considered, or the portion of its width which is filled up by it. No veins occur which are regularly impregnated with metal to any great extent, and when ore is found, it is in what the miners aptly term bunches or shoots, or in interspersed grains and strings, which are more or less connected with, or embraced in, veinstone, that, according to the rock which the veins intersect, will be fluor spar, calcareous spar, quartz, &c. The unproductive parts of veins, even in the most profitable mines, generally far exceed in extent the productive parts, but that mine is considered to be rich which has either frequent or extensive shoots of ore, and the great art of the miner consists in tracing and working the valuable accumulations of the metals, with as little waste of labor and expense on the poorer portions of the veins as possible. “In the mines of Cornwall, the ores of copper and tin commonly occur in detached masses, which are called bunches of ore; and the other parts of the vein, being unproductive, are called *deads*.”

The depth to which metallic veins descend is unknown, for

* Carne, on the mineral veins of Cornwall.

we believe no instance has occurred of a *considerable vein being worked out in depth*, though it may sink too deep to render the operation of the miner profitable, or it may branch off in a number of strings which are too much intermixed with the rock to be worked to advantage.* Some veins appear to grow wider, while others contract as they descend.

The superficial part of a vein generally contains the ore in a decomposing state, and it frequently happens that the ores in the upper and lower parts of a vein are different; thus, "in Cornwall, blende or sulphuret of zinc often occupies the *uppermost* part of the vein, to which succeeds, tinstone, and at a greater depth, copper pyrites." When a metallic vein, in its descent, passes through different kinds of rock, it is frequently observed that the products of the vein vary in each bed, and when it passes through regularly stratified beds of the same rock, there are particular strata in which the vein is always found most productive. This change in the productiveness of mineral veins, is more particularly noticed at or near to the transition from unstratified to stratified rocks; thus granite, syenite and those rocks which have a graniti-form structure, are frequently noticed to contain metals at or near their junction with stratified formations. On the other hand, the veins which traverse stratified rocks are, as a general law, more metalliferous near such junctions, than in other portions.†

Where a rock is crossed and penetrated by a great number of small veins in every direction, the whole mass is sometimes worked as an ore, and is called by the Germans, a "stockwerke." Where the ore is disseminated in particles through the rock, such rocks are also worked for the ore, when it exists in sufficient quantity.

As a general rule, those metals which are oxydable at ordinary temperatures, or which readily combine with sulphur, *rarely occur in a metallic state*, but are usually found in combination either with sulphur, oxygen or acids. The chief ore of copper raised from the mines of Cornwall, is the yellow sulphuret, though the blue and green carbonates and arseniate are more or less distributed; native copper and the oxyds are also, though more rarely, found.

By a comparison of what has been said upon the character and mineral contents of metallic veins in general, I trust a just view of the real condition in which the ores of copper are invariably found, will have been conveyed, and that, by the aid of this, we will be enabled to examine, without undue expectations, those mineral veins which occur within the limits of our own state. In the main, the resemblance between the character and contents of the copper veins of Cornwall and Michi-

* Koenig.

† Lyell. Necker.

gan, so far as can be determined, is close; the veinstones, (with the exception of fluor, which I have never observed in the latter,) are essentially the same; but in instituting this comparison, it should be borne in mind that the metallic veins of Cornwall have been in progress of exploration for centuries, and that shafts and galleries have been carried to great depths, while of those of Michigan, simply superficial examinations have as yet been made, and these in a wilderness country, under circumstances of the utmost embarrassment, and attended with the most excessive labor, privation and suffering.

In respect to the character of the ores which occur in the two districts, there are important differences, for while pyritous copper is the most important workable ore, not only in the Cornish mines, but also in those of other portions of our globe, it is comparatively of rare occurrence in the mineral district of upper Michigan; for, as I have already mentioned, the mineral of the trappean portions of the veins in the latter district, is essentially made up of strings, specks and bunches of native copper, with which more or less of the oxyds and carbonates are associated; while those portions of the veins traversing the conglomerate are characterized by the occurrence of the oxyds and carbonates, with occasional metallic and pyritous copper, or the places of all these are supplied by ores of zinc, associated with more or less calcareous matter. In the thin mineral veins of Presque Isle, pyritous copper is more abundant, where it is associated with sulphuret of lead, as before mentioned.

The occurrence of this native copper in the veins, and the manner in which it is associated with the veinstones, in all respects corresponds with the ordinary association of the other forms of ores, in those veins that have been extensively worked in other portions of the globe; but I confess that the preponderance of native to the other forms of copper, was regarded as an unfavorable indication, at least until this had been found to be more or less universal with respect to all the veins. It should, however, be remarked, that in those portions of the veins where the quartz of the vein and the accompanying rock are very compact, the native form is much more common than in those portions where the veinstone and accompanying rock are more or less cellular and soft.

The worked copper veins of Cornwall, are stated by Mr. Carne, to average from three to four feet in width, and to have a length, as yet, undetermined. But few have been traced for a greater distance than one to one and a half miles, and but one has been traced for a distance of three miles.

The veins which I have examined in the mineral district of Michigan, exceed the average of those last mentioned, but the imperfect examinations which have been made, render it diffi-

cult to determine this with certainty. I have traced no one vein for a further distance than one mile, and usually for distances considerably less. It was not, however, supposed that these veins terminated at the points where they were left, but the further examinations were abandoned at these points, in consequence of physical difficulties connected with the present condition of the country.

The native copper is frequently free from all foreign matter, and is as completely malleable as the most perfectly refined copper, but it more usually contains disseminated particles of earthy minerals, chiefly quartz. I have not been able to detect the alloy of any other metal, in a single instance.

The fatigues and exposures of the past season, have so far impaired my health, that, as yet, I have been unable to analyze as carefully, as could have been wished, the several ores furnished by the mineral veins of the upper peninsula, but sufficient has been done to show satisfactorily that the copper ores are not only of superior quality, but also that their associations are such as to render them easily reduced. Of those which have been examined, embracing nearly the whole, (and not including the native copper,) the per cent of pure metal ranges from 9.5 to 51.72, and the average may be stated at 21.10. Associated with some of these ores, I have detected a metal, the character of which remains, as yet, undetermined.

Were the analysis of the several ores of copper sufficiently perfected, I should deem it unnecessary to lay them before you at this time, for with what is now known of the district, it is conceived, the result would lead to erroneous, rather than to correct conclusions. The analysis of separate masses of ore, no matter how much care may be taken to select the poor as well as the richer ores, for the examinations, will usually be far from giving the average per cent of what would be the product when reduced to practice. I have, in order to arrive at safe conclusions, not only analyzed, but also assayed many of them, but when we come to consider what constitutes the true value of a vein of copper ore, we will perceive why it is unsafe to judge of the whole by the analysis of small portions.

By reference to the previous statistical table of the product of the copper mines of Cornwall, it will be seen, that the average produce of the ores since 1771, has never exceeded 12 per cent of the metal, and that from 1818 to 1822, it was only 8.2. This shows the aggregate, and it is well known that while many of the productive veins are considerably below this, the largest average per cent of any single vein, in that district, it is believed, has never been over 20 per cent, and it should be borne in mind that this average is taken after the ores have been carefully freed from all the rocky and other impurities, which can be separated by breaking and picking.

The value of a vein may be said to depend upon the abundance of the ore, and the ease with which it can be raised and smelted, rather than upon its purity or richness. Upon this point, with respect to our own mineral region, public opinion would perhaps be more in error than upon any other, and most certainly we could hardly look for a mineral district where the character of the ores were more liable to disseminate and keep alive such errors. The occurrence of masses of native metal, either transported or in place, are liable to excite, with those who have not reflected upon the subject, expectations which can never be realized, for while, in truth, the former show nothing but their own bare existence, the latter may be, as is frequently the case, simply imbedded masses, perfectly separated from all other minerals, or they may be associated in a vein where every comparison would lead to unfavorable conclusions, as to the existence of copper, in any considerable quantities. I have frequently noticed very considerable masses of native copper, occupying the joints of compact greenstone, under such circumstances as I conceive, might readily excite in many minds, high expectations, but a little reflection would satisfy the most careless observer of the uselessness of exploring these joints, under the expectation or hope of finding them a valuable repository of the metal. Again, not only native, but also the other ores of copper occur in veins, either so narrow as to render it useless to pursue them, or so associated as to render it probable that exploration would not be attended with success.

While I am fully satisfied that the mineral district of our state will prove a source of eventual and steadily increasing wealth to our people, I cannot fail to have before me the fear that it may prove the ruin of hundreds of adventurers, who will visit it with expectations never to be realized. The true resources have as yet been but little examined or developed, and even under the most favorable circumstances, we cannot expect to see this done but by the most judicious and economical expenditure of capital, at those points where the prospects of success are most favorable. It has been said of the Cornish district, in respect to the supposed large aggregate profits, that "a fair estimate of the expenditure and the return from all the mines that have been working for the last twenty or thirty years, if the necessary documents could be obtained from those who are interested in withholding them, would dispel the delusion which prevails on this subject, as well as check that ruinous spirit of gambling adventure which has been productive of so much misery."* And if these remarks will apply to a comparatively small district, which has been explored and extensively worked for centuries, with how much more

* Hawkins on the tin of Cornwall.

force must they apply to the mineral district of our own state. I would by no means desire to throw obstacles in the way of those who might wish to engage in the business of mining this ore, at such time as our government may see fit to permit it, but I would simply caution those persons who would engage in this business in the hope of accumulating wealth suddenly and without patient industry and capital, to look closely before the step is taken, which will most certainly end in disappointment and ruin.

The extreme length of what I have denominated the mineral district, (within the limits of Michigan,) may be estimated at a fraction over 135 miles, and it has a width varying from one to six miles; but it must not be imagined that mineral veins occur equally through all portions of it, for sometimes, for many miles together, none have been noticed, and the situation of the country is such as to render it probable they never will be. The range and course of the mineral district has been so far defined as to render it unnecessary to say more upon this subject, to enable such persons as may wish to examine, to pass directly along its complete length.

I have thus far omitted to allude particularly to the large mass of native copper, which has been so long known to exist in the bed of Ontonagon river, lest perhaps this isolated mass might be confounded with the products of the veins of the mineral district. That this mass has once occupied a place in some of these veins is quite certain, but it is now perfectly separated from its original connection, and appears simply as a loose transported boulder.

The attention of the earliest travelers was called to this mass of metallic copper by the natives of the country, and it has been repeatedly described by those who have visited it. The mass now lies in the bed of the westerly fork of the Ontonagon river, at a distance which may be estimated at 26 miles, by the stream, from its mouth. The rugged character of the country is such, that it is but rarely visited, in proof of which I may state, that upon my visit to it, during the last year, I found broken chisels, where I had left them on a previous visit, nine years before, and even a mass of the copper, which at that time had been partially detached, but which, for the want of sufficient implements, I was compelled to abandon, was found, after that interval, in precisely the same situation in which it had been left.

The copper in this boulder, is associated with rocky matter, which, in all respects, resembles that associated with that metal in some portions of the veins before described, the rocky matter being bound together by innumerable strings of metal; but a very considerable proportion of the whole is copper, in

a state of purity. The weight of copper is estimated at from three to four tons.

While the mass of native copper upon Ontonagon river cannot fail to excite much interest, from its great size and purity, it must be borne in mind, that it is a perfectly isolated mass, having no connection whatever with any other, nor does the character of the country lead to the inference that veins of the metal occur in the immediate vicinity, though, as before stated, the mineral district crosses the country at a distance of but a few miles.

The occurrence of carnelian, chalcedony, agate and amethystine quartz, in the amygdaloidal portion of the trap, has already been noticed, and these minerals are considerably abundant. They frequently possess very great beauty and perfection, and when ground and polished, they may be used for all the purposes to which those minerals are usually applied.

Minerals of the Upper or Grey Sandstone.

Though the upper sandrock is largely exposed along that portion of the lake coast known as the Pictured rocks, rising to a very considerable height in precipitous cliffs, there have, nevertheless, been no minerals noticed in connection with it, excepting iron pyrites. Along a portion of the distance, however, the rock of the cliffs is frequently colored by broad vertical bands, having a variety of tints, (which have given name to this portion of the coast,) and these bands have been, by some travelers, supposed to indicate the existence of important minerals in the rock; but the coloring matter of these bands is merely superficial. It chiefly consists of the oxyd and carbonate of iron, with occasional feint traces of carbonate of copper, both having been deposited from waters while trickling down the cliffs, the same having previously percolated the rock.

No mineral veins have been noticed in connection with this rock.

SOIL AND TIMBER OF THE UPPER PENINSULA.

The impressions which have gone abroad with respect to the character of this region for purposes of agriculture, are, in many respects, exceedingly erroneous, for which reason, I am the more solicitous to call attention to the true character of the country, as it regards the natural productions of its soil and its capabilities for cultivation. It has generally been supposed that the whole country is wild and sterile in the extreme,

and that, both from its high northern latitude and the rugged and broken character of its surface, it could never admit of the successful application of agriculture. This impression is, in great part, a mistaken one; for, while much of the country is, as has been supposed, extremely rugged, often presenting a rocky and sterile surface, and is besides exposed to the long and bleak winters of a high northern latitude, and to the cold winds of a vast and boisterous lake, yet, as I have already shown, a large part of the upper peninsula is far from presenting a rough and mountainous aspect; and much of the interior, at some distance from the lake shore, presents situations that are not only sheltered from the severity of the winter, but in soil and timber are wholly of a different character from what has been represented.

Much of the wrong impression which has been received, with regard to the timber and agricultural character of the country alluded to, has, no doubt, arisen from the circumstance that a judgment has been formed of the whole district of country, from the appearance of that part of it which lies more immediately upon the lake shore; and in fact, were we to form our estimate of the whole, from that only which is seen by the voyageur, in coasting along the shore, it would scarcely be possible to form any other than a very unfavorable opinion of the value of the country, in an agricultural point of view. Along that portion of the district which lies upon Lake Superior, about one-third of the entire coast, westwardly from the Ste Marie, is generally low and sandy, and thickly timbered with evergreens and white birch, which give to it a somewhat gloomy and forbidding aspect. Still further westerly, and extending to the extremely westerly boundary of our state, the coast presents little else than an almost unbroken succession of rocks, bare, and worn by the fury of the lake, and the country, as seen from the water, appears to be occupied by an almost countless succession of irregular knobs, either nearly destitute of timber, or producing only a few stunted yellow pines and firs, or a growth of worthless poplars; while the southern shores of the upper peninsula, upon Lake Michigan and Green Bay, though differing in character, present a scarcely less forbidding aspect, in general, when viewed from the lake. But these unfavorable impressions are almost wholly, or in great part, removed when we penetrate into the interior, beyond these local influencing causes, and become acquainted with the real condition and character of the districts described.

The general aspect of the surface, over the whole of the extensive district under consideration, has been already laid before you, under the head of "topography of the upper peninsula." It may be remarked, in general, that sand is by far the predominant soil throughout the entire district. A soil of this des-

cription prevails over the north-easterly or sandstone portion of the upper peninsula. This district which, as has been already described, consists of extensive level plateaus or steppes, with scarcely sufficient irregularities of surface to prevent the formation of numerous marshes, may be said to be timbered, in the largest proportion, with the several varieties of evergreens, among which hemlock, cedar and firs greatly abound. Considerable Norway or pitch pine is interspersed, with occasionally large white pines, though in limited quantity. This region, nevertheless, comprehends many extensive tracts of the sugar maple, lying in body, and these trees are frequently of large size. Several species of oaks are also occasionally met with. Upon the whole, much of this portion of the peninsula is better adapted to the wants of settlers for agricultural purposes than might, at first view, be supposed, and may be safely relied upon as capable of producing those crops which are of the most importance to the settler. Wheat, in small quantity, is said to have been raised upon Grand Island, in a spot exposed to the utmost rigor of that northern climate, and some species of Indian corn may, no doubt, be successfully cultivated in the most sheltered situations.

The south-easterly portion of the upper peninsula, embracing the lower limestone district, has a soil more nearly approaching to gravelly, and the pebbles composing which are chiefly derived from the northerly outcropping edges of the limerocks. This soil, in consequence, contains much more calcareous matter than that above described, which adds greatly to its fertility. Clays occur to a very limited extent, and clay soils may be said to be in general, rare throughout the district under consideration. Beach and sugar maple are abundant throughout the portion of the district described, mixed with hemlock and birch.

The hilly district, referred to as embracing the whole of that portion of the upper peninsula which lies west of the mouth of Chocolate river, though broken by ranges of knobby and often barren hills, is very far from being wholly or even generally sterile; for the broad and gently undulating valleys, described as occupying the intervening spaces between these ranges of rocky knobs, have in general a soil of dark, rich and deep loam, and in many places are covered with large bodies of sugar maples of unusual size. With this timber is frequently intermixed oaks and large hemlocks, and extensive bodies of the latter timber occur, together with occasional pines. The streams of this district, where they wind through the bottom lands, between the ranges of hills which inclose the valleys, are frequently densely wooded with all the varieties of hard wood timber, and their banks exhibit deep alluvial loams, which, when once brought under proper cultivation, will be unexcelled for fertility even by the rich plains of our southern peninsula.

These loams were sometimes observed to be underlaid by a red clay. As a whole, the soils of the hilly portion of the upper peninsula, may be said to be generally superior to those of the extensive easterly portion of the peninsula, and marshes are of a less frequent occurrence.

For purposes of lumber, the upper peninsula of our state cannot be said to hold out such inducements as, from its situation, might be imagined. White pine, though sometimes met with in considerable quantity, was not in any instance observed to have obtained to more than medium size, and is not generally abundant. The Norway pine is found in much greater abundance and of fair size, but this species of pine is of comparatively little value, as an article of lumber.

FURS, FISH AND HARBORS OF LAKE SUPERIOR.

In the general view I have attempted to give of the character and resources of that portion of our state bordering on Lake Superior, full justice cannot be done to the subject without adverting, however briefly, to the fur and fish trade of that upper country.

It is well known that the American fur company has for a long series of years, occupied posts, at convenient points upon the lake, as well as throughout the vast territory of the northwest, for the trade with the natives in the furs of the country. Since the year 1835, the general depot of the north-west trade in furs, has been established at La Pointe, one of the Apostles' Islands, near the western boundary line of upper Michigan, and in addition to this, other posts, of a more temporary and minor character, have from time to time, been occupied at various points in the same region. These all formed parts, or, as it were, links of that extended and connected chain, with which this company had been enabled to bind to its interests such an immense extent of territory, and to draw into its store-houses those treasures, in furs, with which this whole region originally abounded. The character of the company, its immense resources, the perfect system that characterized its operations, as well as its distinguished success, are so well known, and have so often been adverted to by travelers and historians, that I allude to the subject here only for the sake of exhibiting more perfectly the present condition of upper Michigan, and to show what is the extent of the inducements held out to individual enterprise in that region. Nearly the whole of this trade is now, as it has been for a series of years, in the hands of the company referred to, and it has been secured to them by a course of judicious management, as well as by a large command of capital, and so powerful continues to be the influence exerted by this company, as to leave little chance of

success to individual opposition, and the trade must continue to flow, mainly in its present channel.

The trade in furs is, however, very far from being of that importance which it was formerly, for the amount of furs has for many years past, been constantly diminishing, as the country becomes exhausted of the game, and so rapid has been the falling off, that at the present time, the amount of furs packed by the company, *at this station*, is scarcely half that which it was five years ago. At the same time, many important changes have been introduced by the company in the manner of conducting the traffic; the system formerly pursued, of granting credits to the natives on goods, sold at extravagant prices, a course not only unjust in itself, but destructive to habits of industry in the natives, has been entirely abandoned; goods are now sold at moderate prices, anticipating only reasonable profits, and an equally fair compensation is allowed for the furs. The company has, furthermore, entered fully into the measures of the government to prohibit the introduction of ardent spirits into the Indian country, and none is now employed in the trade; thus cutting off one of the chiefest sources of misery to the Indian population. I could not fail to notice that, since my last visit to the country, in 1832, the moral and social condition of the Indian population has greatly improved. This is owing, in great part, to the honorable and active measures adopted by the company, by which the means of subsistence among the Indians has been much increased, many evils that formerly existed have been removed, and an entire change has been brought about in their condition, and also to the exertions of the several very excellent and worthy missionaries, who have not only both by precept and example, been the means of introducing among them the meliorating influences of christianity, but have laboriously devoted their life to this "work and labor of love."

The American fur company also, at an early day, turned its attention to the fish of Lake Superior, and they have since engaged largely in the business of fishing. Many half-breeds and Indians are employed in the business, and this has also operated favorably upon the natives, and no doubt contributed largely to the improvement observable in their condition. Fair, if not high, wages are paid the Indians, which operates as a stimulus to industry, nor can we award too high praise to the equitable course pursued by the company in this matter.

Lake Superior abounds in trout, white fish and Siskowit,* the two former of which are larger and better flavored than any that are taken in the more southern waters, and the latter fish has been rarely taken at any other station on the lakes;

* An undescribed species of the genus *Salmo*.

the consequence of which facts is, that the barrels of this company sell in the market at a higher rate than those of the fisheries farther south; but, at the same time, great disadvantage results to the company from the necessity of keeping vessels constantly upon the lake, for the express purpose of the fish trade, thus greatly increasing the expenses and risks attending the business.

During the past season minor companies have been formed and have commenced the business of fishing upon this lake. Two vessels have been hauled around the rapids at the Saut, for that purpose, and as the fish are abundant, there is reasonable prospect that the fishing trade will eventually prove of great importance.

But cut off as Lake Superior is, from direct communication with the lower lakes, by an impassable rapid, this trade must continue to be carried on under great disadvantages, nor can the other resources of the upper country be fully developed, so long as this barrier exists. The construction of the Saut Ste Marie canal, which has been long projected, but the necessity for which seems to be not yet fully appreciated, would remove this obstacle to the growing importance of that great region; and when this shall have been done, we may expect to see all the resources of our upper peninsula fully appreciated and made available.

Directly connected with this subject, is that of the *harbors* on the southern shore of Lake Superior; for while such a field for enterprize is opened in that region, it is of essential importance to know what are the facilities it affords for the safe navigation of the lake. Six vessels are already navigating its waters, in the prosecution of the fur and fish trade, and although hitherto, but few harbors have been known to exist, which may be sought as places of refuge, the extreme breadth of the lake does not fail to afford an extraordinary facility in its navigation, by allowing a great extent of sea room, and thus enabling vessels, in case of storm, to "run before the wind."

The harbor afforded by the group of Apostles' Islands, and which is taken advantage of by the American fur company at its present station, is deep and completely "land-locked," and for safety and convenience could not be excelled. The same may be said of the natural harbor afforded by Grand Island and the neighboring shore. Both these harbors have long been known and appreciated, and more convenient and safe retreats could not be wished for vessels driven by stress of weather to "make a port;" nor could harbors be desired affording greater conveniences for permanent stations.

I was enabled, during the past season, to effect a partial tri-

angular survey of several points upon the coast, which seemed to promise advantages as harbors for vessels; and the depths of the water were uniformly taken, also, at the mouths of streams. The result shows that, although the number of places offering advantages for secure harbors is somewhat limited along a part of the coast, yet that portion of the coast of our state lying upon Lake Superior, may be said to hold out as great facilities, in this respect as any equal extent of coast upon the lakes. It might be supposed that Keweenaw point, from its extreme projection northerly, as well as from the "rock-bound" character of its shores, would serve to add to the dangers of the navigation, but along this rocky coast are found occasional inlets into large bays, stretching behind the extreme outer barrier of rock, and these are almost uniformly deep and completely sheltered, and at some future time, may, with little or no expense, be converted into a series of complete and permanent harbors. One of these bays or inlets, distinguished on the map as Copper Harbor, was surveyed by a series of triangulations and soundings, and was found to afford a bay, convenient for anchorage, about two and a half miles in length, stretching parallel to the coast, and having a depth of from five to seven fathoms, and the entrance to which has an uninterrupted breadth of three quarters of a mile, with a mean depth over the bar of twenty feet, and a maximum depth of thirty feet. These harbors, from their location with respect to the mineral district of the upper peninsula, as well as from the ease of access they will allow to vessels which are exposed "off the point," must eventually become of very great importance.

A natural harbor, of similar form to that above described, occurs upon a part of the coast east from the Pictured rocks, and in which the place of the cliffs of rock is supplied by cliffs or spits of sand. The bay here formed, behind the outer bar of sand and gravel, has a length of two miles, with an average depth exceeding six fathoms, and the water deepens so rapidly from the shore, that in most parts of it a vessel of the largest draught may approach so closely to the beach as that a landing might be effected without aid from the boats. The entrance into this bay has a width of three-fourths of a mile, but a bar of sand stretches across it, which would, at times, in its present condition, render the passage into the harbor dangerous, for the average depth of water over this bar does not exceed six feet, though there is a depth in the immediate channel, which is comparatively narrow, of ten feet.

Safe anchorage for vessels may also be found in Keweenaw bay, a deep expanse of water, on the easterly side of Keweenaw point, and which stretches for many miles inland.

PROGRESS AND CONDITION OF THE SURVEY, &c.

Notwithstanding the very many physical difficulties by which the geological survey of the upper peninsula of our state is surrounded, we have, nevertheless, been enabled to accomplish a much larger amount of the work than reasonably could have been hoped; but there still remains much to be done, before its geology and mineralogy can be fairly understood. Comparatively little has heretofore been known of the range and extent of the several rock formations, and, while the labor of the past season has shown the most interesting of these to have a much larger area than we had previously been led to infer; it has also shown, that the amount of work required, to enable us fairly to understand the geology and mineralogy of that interesting region, was considerably more than we had reason to look for.

The reports of the several assistants, will exhibit to you the progress that has been made, during the past season, in the surveys of the southern peninsula. Messrs. B. Hubbard and C. C. Douglass were engaged with me, during the early part of the season, in the upper peninsula; after which, they returned to carry forward the geological and topographical surveys of the lower peninsula. I was also accompanied, during a small portion of the season, by Mr. Frederick Hubbard, who acted as special assistant, and who has embodied a small part of his numerous observations in the form of a report, which is hereto appended.

The survey of the lower peninsula is mainly completed, but there are some few spaces, both in the geological and topographical portions of the work, which require to be filled up before the results can be fully laid before the public.

The drafting of the topographical portion of the survey, has advanced steadily towards completion, and the several county maps are in progress of publication, in conformity to your instructions.

While we had hoped to have been able to bring the survey to a close within the time originally contemplated, from the above statement of the progress and condition of the work, it will be seen, that some further time will be necessary for its final completion; but while this time will be essential to reach the object sought to be attained by our state, no farther appropriation will be necessary for that purpose.

In closing this report, I feel it a pleasure to refer to the very many acts of hospitality and kindness which have been extended to us by the citizens of Mackinac, Saut de Ste Marie, and

La Pointe. By the aids received at these several places, we have been enabled to accomplish much that otherwise could not have been done, while we have, at the same time, received much that has ministered to our comfort.

To the agents of the American fur company in Lake Superior, I feel very deeply indebted; for, through their polite attentions, I have been enabled to examine districts of country which otherwise could not have been reached, and to them and the several mission families, are we indebted for all that it was within their power to do, to aid us in our laborious duties, or to render our situation comfortable.

DOUGLASS HOUGHTON,
State Geologist.

Report of Frederick Hubbard, Special Assistant.

Utica, N. Y., November 20, 1840.

To DOUGLASS HOUGHTON, *State Geologist:*

DEAR SIR—I submit to you, the results of a portion of the observations for the determination of latitudes, magnetic variations, &c., made under your direction, during the recent expedition of the state geological corps to the upper peninsula and southern shore of Lake Superior.

The subject of latitudes was made a matter of particular attention, no regular survey ever having been made by the general government of that part of the lake lying within the boundary of our state. By means of instruments, with which the expedition was furnished, the positions of the most prominent points of the coast have been fixed by celestial observations, and with sufficient care to furnish, connected with the running meander, an accurate outline. Unfortunately, we were not furnished with the proper instruments for the determination of longitudes. This problem, always one of great difficulty, for our purpose, would have required a long series of the most careful observations, with the most delicate chronometers, and a greater devotion of time, than the circumstances of the expedition allowed. The ordinary lunar method, as practiced by navigators, is not susceptible of sufficient accuracy to be of service in a case of nicety.

Several rude attempts were made, however, with such means as we possessed, using a similar method, to fix the longitudes of a few points; but I do not place sufficient confidence in the results to deem it of importance to lay them before you.

The latitudes in the following table were mostly obtained by measured altitudes of the sun, Jupiter and the polar star.

Table of Latitudes.

<i>Ste Marie Riviere.</i>	
W. side Drummond Island, (encam'pt, May 27,)	45° 58' 26" N
E. side St. Joseph Island, May 28,)	46° 16' 45"
Pt. aux Pins, Little Lake George,	46° 32' 29"
Saut de Ste Marie, (head of rapids,	46° 31' 8"
<i>Lake Superior.</i>	
White fish point,	46° 45' 18"
Grand Marrais harbor,	46° 40' 5"
Mouth of Train river,	46° 25' 34"
Mouth of Chocolate river,	46° 29' 33"
Mouth La Riviere des Morts,	46° 35' 4"
Mouth Riviere Bay du Gres,	47° 21' 36"
Rock harbor, extremity of Keweenaw point,	47° 24' 32"
Mouth of Ontonagon river,	46° 57' 5"
Mouth of Montreal river,	46° 41' 19"
Village of La Pointe, Madaline Island,	46° 44' 31"

Variation of the compass.—The results, in the subjoined table, were deduced from a comparison of the observed magnetic azimuth of the sun with his true azimuth, as calculated for the measured altitude and place of observation. This method is one of great accuracy, and is almost the only one upon which dependence can be placed, for results which shall not contain an error of more than two or three minutes, due to inaccuracy of observation. The angles were measured with a theodolite, containing a needle of great delicacy.

A small correction to the table is necessary, on account of the diurnal variation. This may amount to some 8' or 10' and possibly may be greater in some instances, when the observations were made during the warmest seasons. As I have not the results of any experiments on this subject, now before me, I have thought proper to insert the date of each observation, and the hour of the day, that the correction may be made when the date can be obtained, with reference also to the metrological table kept during the survey.

By plating these variations upon the map of the coast, it will appear that there is a constant increase in the amount of deflection in passing westward, and at the same time that *the increment is not in proportion to the westing* but is in a decreasing ratio, the distance between the lines, connecting the points of the same variations, being greater as we recede from the line where the direction of the needle is due north.

From the observations in Riviere Ste Marie, it appears that the line of no variation passes through that strait, crossing

Drummond Island near its western, and St. Joseph through its central or eastern part, with a course about north by west.

The determination of its exact position is a matter of considerable importance, and I regret that circumstances did not allow us, when in that vicinity, to spend more time in the investigation of this interesting subject. The tracing of its supposed irregularities, and the ascertaining of the laws which govern its changes, have for many years attracted the attention of the scientific world.

By marking upon the map the points of 1, 2, 3, &c., degrees of variation, and drawing through them lines parallel to the line of no variation, it will be perceived that there are, in a few cases, important deviations from the general regularity in which the deflections are found to increase. These differences are by far too great to be attributed either to errors in observation, to diurnal variation, or to the effect of atmospheric disturbing causes. I know not what influence the general topography of a country may have upon the directions of the magnetic needle, but it appears highly probable that something may be due to the outline of coast, to the unequal distribution of land and water, and to the influence of an open extent of sea, on the one hand, and of a mountain range upon the other. It might be suspected that these irregularities were the result of local attraction of metallic veins, or of some unknown disturbances in the more immediate vicinity, did they not always occur where they seem to have an obvious connection with the circumstances I have mentioned. In all these cases there is a deflection of the needle towards the open lake, tending, when the land lies to the west of the place of observation to increase the amount of easterly variations, and the contrary. Thus we find, about the Riviere des Morts, where the trend of the shore is northerly, at the several points to the east of the Keweenaw peninsula, and at the village of La Pointe, lying to the eastward of a high range of hills upon the mainland, a too great deflection towards the east, as if the needle were actually affected by some repulsive influence existing in the land, or a contrary principle, in the water. One or two instances occur, where the attraction is to the west apparently from a similar cause.

Whether this is the true explanation of the difficulty, I will not pretend to decide. The subject is one that merits investigation, as having some bearing upon the science of terrestrial magnetism, many of whose principles are as yet enveloped in mystery.