

868. *Mycena acicula* Fr.

Epicrisis, 1836.

Illustrations: Fries, Icones, Pl. 85, Fig. 3.

Cooke, Ill., Pl. 190.

Patouillard, Tab. Analyt., No. 108.

Atkinson, Mushrooms, Fig. 98, p. 98, 1900.

PILEUS 2-4 mm. broad, campanulate-convex, sometimes papillate, glabrous, glaucous, striatulate on margin, *vermillion, reddish-orange or yellowish* with red center. FLESH very thin, membranaceous. GILLS adnexed, ascending, ventricose, rather broad, *subdistant to distant*, yellow, yellowish, creamy white, or white, edge minutely crenulate. STEM 2-5 cm. long, filiform, equal, toughish, hollow, glabrous or minutely pulverulent at first, *pellucid-yellowish or yellow*, more or less rooting. SPORES 7-9 x 2.5-4 micr., *narrow, fusiform or narrowly subovate*, smooth, white. CYSTIDIA none. ODOR none.

Gregarious or scattered. On rotten wood, or among leaves and grass, in woods, meadows, thickets, etc. Ann Arbor, Detroit, etc. May-June and September-October. Frequent.

This is a pretty little *Mycena* and one of the earliest to appear. It is not by any means limited to the woods or to growing on wood or twigs as most authors remark, but may be found among grass on the ground in low, moist meadows in spring. The spores are quite characteristic and help not a little in its positive identification. I suspect, in fact, that it has been reported as *M. adonis* Fr. when occurring on the ground in grassy places, but no spore measurements of that species seem to have been printed. On decayed wood the stem is rooting and hairy along the root, whereas on the ground it has few hairs and is scarcely at all rooting. Other minute *Mycenas* with rosy or red caps have been described by Fries from Europe: *M. stipularis* and *M. juncicola* have non-rooting stems inserted on stipules of fallen leaves; *M. pterigenus* has a bulbillose stem attached to roots and leaves of ferns, etc.

*Omphalia* Fr.

(From the Greek, *omphalos*, an umbilicus.)

White-spored. Stem *cartilaginous*, slender, usually hollow or loosely stuffed, widened above with the pileus in trumpet-form.

Gills *decurrent*, or adnate-decurrent. Pileus more or less *membranaceous*, its margin at first either incurved or straight-appressed.

Epiphytal, lignicolous or terrestrial; putrescent plants, with few exceptions small or minute. Very closely related to *Collybia* when the margin of the pileus is at first incurved; and to *Mycena* when it is straight; differing from these by the decurrent gills. Toughish, reviving species are referred to *Marasmius*. They differ from *Clitocybe* by the cartilaginous character of the stem.

The PILEUS varies conical, hemispherical, convex or campanulate, often quite expanded in the *Collybiariae*, and usually marked by a distinct umbilicus which becomes widened in some species to infundibuliform; some species show the umbilicate character only in occasional specimens, e. g., *O. gracillima*, while the other specimens are obtuse or somewhat papillate. Its color is often pure white; it may be gray, fuscous, brown, yellow, orange or reddish. It is generally hygrophanous and then striate when moist. The surface is usually glabrous or with a very minute pubescence which is rarely glandular. The GILLS are decurrent, sometimes not strongly so, most often running far down as the cap expands. Their width and spacing are used to separate the species into subdivisions, although this is not always sharply marked. Peck in his monograph of the New York species (Rep. 45, 1892) states that in his opinion this grouping is unsatisfactory, but as no better is offered, it is adhered to in this paper. CYSTIDIA are lacking in most species, and when present are rather few and inconspicuous. The STEM is usually slender to filiform, and when dry its cartilaginous cortex is like that of *Mycena*. The surface is glabrous, horny and shining, or varying to pruinose or hairy at base. The SPORES are smooth. As in *Mycena* the immature spores are loosened in a microscopic mount; so the same precaution must be observed as when studying the spores of *Mycena* in order that one may get the correct measurements of the mature spores. Very few *Omphalias* have a distinguishing ODOR, and none of the following are nitrous or fragrant. Their EDIBILITY is of no interest because of their thin texture and small size; none are known to be poisonous, but probably few have been tested.

They are found from early spring to late autumn, during wet weather and in low or damp, shaded places. With the exception of *O. campanella* and *O. fibula*, they occur sporadically, few of them are found in quantities, many are but rarely found. A rather large number of species are described, Peck has named about twenty-five.

## Key to the Species

- (a) Plant wholly white.
  - (b) Gills broad.
    - (c) Stem tubular; plant snow-white. 879. *O. gracillima* Fr.
    - (cc) Stem solid; plant soon dingy-white. 878. *O. albidula* Pk.
  - (bb) Gills rather narrow.
    - (c) Gills crowded; plant pure white. 869. *O. scyphoides* Fr.
    - (cc) Gills moderately close; plant dull white. 870. *O. scyphiformis* Fr.
- (aa) Plant not wholly white.
  - (b) Pileus viscid.
    - (c) Gills lilac; pileus greenish-yellowish. 872. *O. lilacifolia* Pk.
    - (cc) Gills whitish; pileus pale grayish-brown. (See 834. *Mycena vulgaris* Fr.)
  - (bb) Pileus not viscid.
    - (c) Pileus yellowish, orange or reddish.
      - (d) Stem date-brown, horny; on coniferous wood; gills very veiny. 883. *O. campanella* Fr.
      - (dd) Stem not date-brown and horny.
        - (e) Gills white or cream-colored; on mosses.
          - (f) Pileus 10-20 mm. broad; convex. 882. *O. fibuloides* Pk.
          - (ff) Pileus 3-8 mm. broad, cucullate. 880. *O. fibula* Fr.
        - (ee) Gills not white.
          - (f) Pileus brick-red to reddish-brown. 873. *O. pyxidata* Fr.
          - (ff) Pileus olivaceous-yellowish; stem dusky yellowish. 871. *O. olivaria* Pk.
    - (cc) Pileus with grayish, fuscous or brown shades.
      - (d) Pileus 3-7 mm. broad, whitish except the fuscous-brown center. 881. *O. schwartzii* Fr.
    - (dd) Pileus usually larger.
      - (e) Gills broad, crowded; pileus and stem dark, umber-fuscous to blackish. 884. *O. umbratilis* Fr.
      - (ee) Gills rather narrow.
        - (f) Pileus dotted with scurfy-blackish points; on sphagnum. 877. *O. gerardiana* Pk.
        - (ff) Not with blackish points.
          - (g) On decaying wood, logs, stumps, etc.
          - (h) Pileus rugose, brown; flesh emitting watery juice when cut. 874. *O. rugosodisca* Pk.
          - (hh) Pileus even, ashy to smoky, non-hygrophanous. 875. *O. epichyrium* Fr.
      - (gg) On the ground, pileus hygrophanous, substrate; gills subdistant. 876. *O. onisca* Fr.

Section I. *Collybiariae*. Margin of pileus at first incurved.

\*Gills narrow; crowded, close or subdistant.

869. *Omphalia scyphoides* Fr.

Syst. Myc., 1821.

Illustrations: Fries, Icones, Pl. 75, Fig. 2.

Patouillard, Tab. Analyt., No. 419.

"PILEUS 4-8 mm. broad, umbilicate then infundibuliform, often

irregular, pure white, even, silky. FLESH submembranaceous. GILLS decurrent, narrow, crowded, white. STEM 8-16 mm. long, short, stuffed, subvillose, white. SPORES 6 x 2 micr. (Sacc., Masee); 6 x 4-5 micr. (Pk.)

"Gregarious, in mossy, grassy places on the ground. (Dead leaves in woods—Longyear.)

"Variable, often flexuous and eccentric, commonly small, not hygrophanous, yellowish on drying, stem woolly at least at base, pileus sometimes often an inch broad."

Reported by Longyear. Greenville. July. Rare.

870. *Omphalia scyphiformis* Fr.

Epicrisis, 1836.

Illustration: Fries, Icones, Pl. 75, Fig. 3.

PILEUS 5-9 mm. broad, convex, then umbilicate or infundibuliform, glabrous, sometimes silky, dull white, margin obsolete striatulate. FLESH membranaceous. GILLS decurrent, thin, narrowly triangular, whitish, close, moderately broad in middle, attenuate at ends, edge entire, trama of interwoven hyphae. STEM 1.5-2.5 cm. long, equal, even, somewhat hollow, whitish, glabrous or loosely pubescent by flexuous longish hairs, base attached to moss by radiating hairs. SPORES 6-7 x 3.5-4 micr., ovate or pip-shaped, pointed at one end, smooth, white. CYSTIDIA none. ODOR none.

Gregarious on mosses or on moist earth. Ann Arbor. August. Rare.

This species is closely related to *O. scyphoides*. It differs in the dull white (albidus) color of the whole plant, whereas *M. scyphoides* is said to be pure white (candidus); the gills are not as narrow and crowded as in that species, and probably the spores differ. Our plants approached *M. scyphoides* in the pubescence toward the base of the stem, which Fries says is glabrous. On moss the silky-webby mycelium sometimes spreads some distance.

871. *Omphalia olivaria* Pk.

N. Y. State Mus. Rep. 25, 1873.

PILEUS 6-15 mm. broad, convex, subclavate, with decurved margin, deeply umbilicate; glabrous, pale yellowish to olive-green, even or obscurely striatulate. FLESH thin. GILLS long decurrent,

subdistant, rather narrow, yellow when fresh, edge entire. STEM 2-2.5 cm. long, 1 mm. thick, tough, equal, even, tubular, dusky yellowish, minutely pubescent, glabrescent. SPORES 6-7 x 5 micr., subglobose or broadly elliptical, smooth, white.

On decayed logs. Ann Arbor. September. Rare.

The plant revives somewhat after being moistened. The pubescence of the stem is white after the escape of the moisture. The trama of the gills is composed of interwoven, narrow hyphae, quite compactly arranged. *O. luteola* Pk. differs in having a brown and solid stem. *O. subclavata* Pk. has yellow gills, but the pileus is grayish-brown and stem white.

#### 872. *Omphalia lilacifolia* Pk.

N. Y. State Mus. Rep. 24, 1872 (as *O. lilacina*).

N. Y. State Mus. Rep. 29, 1878.

Illustrations: Peck, N. Y. State Mus. Rep. 24, Pl. 1, Fig. 10-13.

"PILEUS 1-2 cm. broad, convex, deeply umbilicate, glabrous, viscid, hygrophanous, dingy yellow with greenish tinge and striatulate when moist, bright sulphur-yellow when dry. GILLS decurrent, arcuate, close, narrow, pale lilac. STEM 1-2.5 cm. long, 1-2 mm. thick, equal, glabrous, hollow, viscid, yellowish, with a pale lilac-colored mycelium at the base. SPORES subelliptical, 5-6 x 3 micr."

Scattered or solitary, rarely subcaespitose, on decaying trunks of hemlock, in the coniferous regions of the State. Munising, Houghton, Huron Mountains, Bay View, New Richmond. July-September. Infrequent.

The colors and viscosity make this an easily recognized species. *O. lilacina* was described from Lapland, and anteceded Peck's first name; it is said to be violaceous except for the yellow gills; its cap is not viscid.

#### 873. *Omphalia pyxidata* Fr.

Syst. Myc., 1821.

Illustrations: Cooke, Ill., Pl. 194.

Patouillard, Tab. Analyt., No. 636.

"PILEUS 1-2 cm. broad, convex, umbilicate to infundibuliform,

glabrous, hygrophanous, brick red or reddish-brown and radiate-striate when moist, paler and silky when dry. FLESH membranaceous. GILLS decurrent, narrow, subdistant, tinged with flesh color then gilvus. STEM 2-3 cm. long, 2 mm. thick, equal, tough, glabrous, even, stuffed or hollow, pallid rufescent. SPORES ovoid, smooth, white, 6-8 x 5 micr. (Patouillard.)

"On the ground, roadsides, etc." East Lansing, on the campus. Reported by Longyear.

Evidently rare. Peck says he found it but once. Not much reliance can be placed on the spore-measurements given; Stevenson quotes three different sizes, one of which is followed by Saccardo. The pileus is said by Fries to fade strongly, even to become whitish. Some think that *O. muralis* Fr. and *O. hepatica* Fr. are to be considered as varieties of *O. pyxidata*. (See Barbier, Bull. 1, Soc. Myc. de France, Vol. XX, p. 105, 1904.)

#### 874. *Omphalia rugosodisca* Pk.

N. Y. State Mus. Rep. 26, 1874.

PILEUS 1-2.5 cm. broad, broadly convex or nearly plane, depressed or umbilicate, sometimes obtuse or slightly umbonate, rugose, hygrophanous, watery cinnamon-brown and striatulate (moist), paler when dry. FLESH thin. GILLS short-decurrent, narrow, close, whitish. STEM 2-3 cm. long, 1-2 mm. thick, glabrous, hollow, cartilaginous, concolor, paler at apex. SPORES elliptical, 6-7 x 4-5 micr., smooth, white.

On decaying prostrate trunks of hemlock. Bay View. July.

Known by the rugose cap, the brown color and the fact that every part of the plant emits a watery juice when cut or bruised. It seems to be limited to decaying wood of coniferous trees. Peck referred this species to *Collybia* in the 31st New York Museum Report, but later returned it to *Omphalia*. Saccardo placed it in the genus *Mycena* without giving any reason for doing so. At times the odor is said to be slightly of radish.

875. *Omphalia epichysium* Fr.

Syst. Myc., 1821.

Illustrations: Atkinson, Fig. 191, p. 101, 1900.

Plate CLXXI of this Report.

PILEUS 1-4 cm. broad, convex then expanded and umbilicate or umbilicate-infundibuliform, margin arcuate with decurved edge, *striatulate and dark cinereous to smoky-brown* (moist), even, silky and light gray to whitish (dry). FLESH thin, soft. GILLS acuminate-subdecurrent, ascending-arcuate, *narrow*, close, thin, *whitish-cinereous*, edge entire. STEM 1.5-3 cm. long, 1-3 mm. thick, equal, glabrous, *almost solid* or subfistulose, *smoky-cinereous*, concolor within. SPORES 7.5 x 4 micr., pip-shaped, smooth, white. CYSTIDIA none.

On decaying logs or remains of decayed wood in frondose or mixed woods. Throughout the State. Ann Arbor, New Richmond, Marquette and Houghton. July-September. Infrequent.

The pileus is often fibrillose-floccose on the umbilicate center. The plants are rather soft and watery and the stem soon shrivels. *O. onisca* Fr. differs in the hygrophamous pileus which is entirely glabrous, the smaller size and large spores; the gills are somewhat different in shape and spacing.

876. *Omphalia onisca* Fr.

Syst. Myc., 1821.

Illustrations: Fries, Icones, Pl. 75, Fig. 3.

Cooke, Ill., Pl. 209.

PILEUS 5-15 mm. broad, convex, soon plane and cyathiform-umbilicate, *glabrous*, flaccid, *hygrophamous*, smoky-fuscous or ashy-brown and striate on margin (moist), pale cinereous and hoary (dry), the umbilicus darker. FLESH thin, concolor when moist, pale grayish-white when dry. GILLS *short*, *decurrent*, broadish in middle, *subdistant*, thickish, cinereous-fuscous, sometimes forked, edge entire. STEM short, 1-1.5 cm. long, .5-1 mm. thick, slender, equal, toughish, firm, solid, *concolor*, glabrous, becoming pale within. SPORES 9-11 x 5 micr., distinctly *ovate*, apiculus curved, smooth, white. CYSTIDIA and sterile cells none. BASIDIA

clavate, 30-35 x 4-5 micr., 4-spored, sterigmata slender, 6-7 micr. long. ODOR and TASTE none.

Scattered or gregarious on the ground, in low, wet places in woods. Ann Arbor. May and September. Rare.

Differs from *O. epichysium* by its habitat on the ground, its glabrous, hygrophamous pileus and by its gills and spores. The hymenium occasionally has sterile basidia intermixed, bearing only a single long abortive sterigma. Authors disagree as to the size of the spores. Britzelmayr gives them as 12 x 7-8 micr. and Massee as 6 x 5 micr. It differs from *O. umbratilis* in the rather distant gills and larger spores.

877. *Omphalia gerardiana* Pk.N. Y. State Mus. Rep. 26, 1874 (as *Clitocybe*).

PILEUS 10-20 mm. broad, nearly plane or soon infundibuliform, hygrophamous, fragile, *grayish-brown to brownish-ashy and striatulate* (moist), paler when dry, *the surface is dotted by scurf-like scattered points which become blackish*. FLESH thin, concolor. GILLS decurrent, *narrow*, subdistant, whitish then tinged with ashy or obscure yellowish, sometimes forked. STEM 3-5 cm. long, 1-2 mm. thick, *cartilaginous*, equal, even, hollow by a narrow tubule, glabrous or pruinose-villose toward base, *at length darker than pileus*. SPORES 7-11 x 3.5-4 micr., variable in size, maturing slowly, oblong-ovate, narrow, smooth, white. CYSTIDIA none. ODOR none.

On sphagnum in cedar and tamarack swamps. Ann Arbor, Houghton. May, July. Local.

Differs from *O. sphagnophila* Berk., in color and spores, according to the description. Cooke (Ill., Pl. 289) gives figures which are very like our plants. *O. sphagnophila* is said to have a dingy pale ochre-colored pileus and gills, is tough, and has smaller spores. Peck vacillates in deciding whether this is an *Omphalia* or *Clitocybe*; it was originally described as a *Clitocybe*, then in the 45th Report it was placed among the *Omphalias*, finally in the list of species described by Peck (N. Y. State Mus. Bull. 131), it was referred back to *Clitocybe*. My collections indicate that it has mostly a distinctly cartilaginous stem, hence it is placed here. The color varies considerably during its development; in old specimens the whole plant becomes dingy dark brown. The scurfy points on the cap are somewhat as in that of *Clitocybe ectypoides* Pk.

\*\*Gills broad, distant.

878. *Omphalia albidula* Pk.

N. Y. State Mus. Rep. 49, 1896.

PILEUS 3-8 mm. broad, *convex-hemispherical*, at times papillate, with or without an umbilicus, glabrous, striatulate, at first pure white, then dingy. FLESH membranaceous. GILLS decurrent, *distant, broad*, white, then dingy. STEM *very slender*, 1-4 cm. long, .5 mm. thick, *toughish*, glabrous above, white, "solid," attached below to leaves, etc., by a very strigose base. SPORES 8-10 x 2-4 micr., elongated-oblong, on slender sterigmata, 4-5 micr. long, smooth, white.

On leaves and bark in frondose or mixed woods. Bay View. July. Rare.

Related to *O. stellata* Fr. which differs mainly in its fragile stem, diaphanous pileus and different spores. *O. papillata* Pk. seems to be very similar, but with a conic or campanulate cap, few gills and different spores; it is said to be pure white.

*Section II. Mycenariae.* Margin of pileus at first straight, appressed to stem.

879. *Omphalia gracillima* Fr.

Epicrisis, 1836.

Illustrations: Fries, Icones, Pl. 75, Fig. 5.

Cooke, Ill., Pl. 252.

Gillet, Champignons de France, No. 502.

Plate CLXXI of this Report.

PILEUS 3-10 mm. broad, at first campanulate, becoming *hemispherical*, sometimes *papillate*, sometimes depressed, *snowy-white*, pellucid-striate, soon sulcate, glabrous, subflocculose when dry, fragile. FLESH membranaceous. GILLS decurrent, *triangular, broad*, subdistant to distant, thin, pure white, edge fimbriate. STEM filiform, 2-5 cm. long, .5 mm. thick, equal, tubular, minutely subpruinose, *white* like pileus, *toughish*, flaccid, attached by floccose base or almost inserted. SPORES 6-8 x 3-4 micr., *oval-lanceolate*, apiculate, smooth, white. CYSTIDIA and *sterile cells* none. BASIDIA 24 x 6 micr., 4-spored.

Gregarious among and on fallen leaves and grass in frondose woods. Ann Arbor, Marquette. Spring and autumn. Infrequent.

A very pretty little plant as its snowy-white color shows against a background of leaves or grass in open woods. The sulcate character of the margin of the cap appears as the plant loses moisture; sometimes a slight papilla is present. The gills are quite distant at times, with shorter gills alternating. Under the microscope the stem is seen to be covered with very short hair-like cells which under a hand-lens appear only as a slight pruinosity. The trama of the gills consists of interwoven hyphae. Our plants were not attached by floccose hairs as shown in Cooke's figures, but were almost free of them.

880. *Omphalia fibula* Fr.

Syst. Myc., 1821 (c. syn.).

Illustrations: Cooke, Ill., Pl. 274.

Gillet, Champignons de France, No. 500.

Patouillard, Tab. Analyt., No. 110.

Hard, Mushrooms, Fig. 99, p. 134, 1908.

PILEUS 3-8 mm. broad, subhemispherical or *cupulate*, often not expanded, *subumbilicate*, obscurely striatulate, nonhygrophanous, *pallid ochraceous-orange*, fading, even when dry, minutely pubescent under lens. FLESH thin, pallid. GILLS arcuate-decurrent, *narrow*, close (or subdistant after expansion of pileus), rarely forked, subpruinose, whitish or creamy-yellowish, edge entire. STEM 2-5 cm. long, *scarcely 1 mm. thick*, equal, flexuous, at length subtubular, *toughish, whitish or with tinge of yellowish* or straw color, scarcely pubescent under a lens, cartilaginous when dry. SPORES elongated oblong, 4-6 x 2 micr., smooth, white. CYSTIDIA scattered on sides and edge of gills, narrowly fusiform-acuminate, sometimes capitate, 3-4.5 x 7-9 micr. ODOR and TASTE none.

On *mosses*, in low woods or moist places. Throughout the State. May-October. Common.

This is our commonest *Omphalia*, although often only several specimens are found in one place. Its cap is at the very first almost cylindrical, then the margin turns out like the brim of a hat, so that it has much the shape of a man's "high silk hat;" hence it is said to be *cupulate*. Under the microscope the stem and cap are

found covered with hyaline hairs, 50-60 micr. long. The gills are narrow and deeply decurrent, running up the inside of the unexpanded cap and down below it on the stem. The cap often becomes top-shaped instead of cucullate. Stevenson (British Fungi) and Massee (Fungus Flora) say the gills are broad, a statement which is clearly an error.

881. *Omphalia schwartzii* Fr.

Epicrisis, 1836 (as var. of *O. fibula*).

Illustrations: Patouillard, Tab. Analyt., No. 420.

Gillet, Champignons de France, No. 504.

PILEUS 4-7 mm. broad, rather firm, soon campanulate, obtuse or at first umbonate, at length slightly umbilicate, even or striatulate, subpruinose, whitish with a slight fuscous tinge, disk fuscous-brownish. FLESH thin, pale fuscous. GILLS adnate-decurrent to strongly decurrent, arcuate, moderately broad, subdistant, whitish. STEM 2-3 cm. long, filiform, hollow, equal, pallid below, tinged violaceous above, pruinose, even. SPORES oblong-elliptical, 5-6 x 2.5-3 micr., smooth, white. CYSTIDIA on sides and edge of gills, scattered, subventricose, subcapitate at apex, some cylindrical above, 40-45 x 9-12 micr.

Gregarious on moss (*Mnium*, etc.) in hemlock woods. New Richmond. June. Infrequent.

*O. schwartzii* is constantly distinct from *O. fibula* to which Fries joined it as a variety. Peck (45th Rep. N. Y. State Mus.) considered it an independent species. Patouillard (Tab. Analyt.) had already pointed out that it was distinct from Bulliard's *O. fibula*, giving figures of spores and cystidia as evidence. It seems to be limited to regions of coniferous forests.

882. *Omphalia fibuloides* Pk.

N. Y. State Mus. Rep. 24, 1872.

"PILEUS 1-2 cm. broad, convex, deeply umbilicate, glabrous, hygrophanous, dull orange and striatulate when moist, paler when dry. FLESH thin. GILLS strongly decurrent, arcuate, rather close, white, the interspaces venose. STEM 3-5 cm. long, scarcely 2 mm. thick, equal, glabrous, hollow, colored nearly like the pileus. SPORES elliptical, 7.5 x 5 micr."

On moss. Jackson County. June.

Reported by Longyear. It is unknown to me. It is said to occur on burned, mossy ground like *O. fibula*, "which it resembles in color, but from which it may easily be distinguished by its much larger size, more robust habit and venose interspaces; the spores also are larger than in that species."

883. *Omphalia campanella* Fr.

Syst. Myc., 1821.

Illustrations: Cooke, Ill., Pl. 273.

Michael, Führer f. Pilzfreunde, Vol. III, No. 67 (as *O. fragilis*).

Hard, Mushrooms, Pl. 17, p. 131.

White, Conn. State Geol. & Nat. Hist. Surv., Bull. 3, Pl. 16, p. 36.

Murrill, Mycologia, Vol. 4, Pl. 8, Fig. 10.

Plate CLXXII of this Report.

PILEUS 8-22 mm. broad, convex-campanulate, expanded, umbilicate, glabrous, striatulate to the umbilicus, dull orange-yellow or tinged reddish, watery when moist, paler when dry. FLESH thin, membranaceous, yellow. GILLS adnate-decurrent to deeply decurrent, arcuate, thick, very veiny, narrow, tapering to front, subdistant to distant, pruinose. STEM 1-4 cm. long, 1 mm. or less in thickness, horny-cartilaginous, minutely tubular, curved or straight, even, dark rufous-bay to date-brown, yellowish above, concolor within, glabrous above, pruinose elsewhere and with fulvous strigose hairs at the slightly thickened base. SPORES oblong, 6-7.5 x 3-3.5 micr., smooth, white. CYSTIDIA widely scattered on sides of gills, more abundant on the edge, subcylindrical or sublanceolate with obtuse apex. 50-55 micr. long. BASIDIA 30-32 x 4-5 micr. ODOR mild. In dense and spreading clusters on hemlock and pine logs, stumps, etc.; also on tamarack logs, stumps or their remains in the southern part of the State; sometimes on debris on the ground.

Throughout the State; collected from Detroit to Isle Royale in Lake Superior. June-November. (Earliest collection May 6.) Common on coniferous wood.

The Bell-omphalia is a striking and easily recognized plant because of its habit of forming extensive clusters, by its colors, and by its horny stem and veined gills. In the non-coniferous regions it

appears on tamarack and perhaps always on remains of wood of coniferous origin. Here it shows a marked variation from the typical plant as it appears on pine and hemlock; the gills are distant, always pruinose, and the stem is also pruinose and often solid. The typical stem of the plant from coniferous regions is usually attenuated below, but in the plant of the non-coniferous regions the stem is equal. The plants of non-coniferous regions are larger and the surface of the cap somewhat rivulose. Hard has illustrated a species (Mushrooms, Figure 98, p. 123, 1900) which he refers to *O. caespitosa* Bolt. It seems probable that this is a var. of *O. campanella*, perhaps var. *terrestris* Quel. Peck says its mycelium is regarded as destructive to the wood of coniferous trees.

884. *Omphalia umbratilis* Fr.

Syst. Myc., 1821.

Illustrations: Fries, Icones, Pl. 77, Fig. 3.

Cooke, Ill., Pl. 274.

"PILEUS 2-2.5 cm. broad, campanulate to convex, then *umbilicate*, glabrous, hygrophanous, *umber-fuscous* (moist), hoary when dry, margin somewhat striate. FLESH submembranaceous. GILLS adnato-decurrent, *broad, crowded*, becoming fuscous. STEM 2.5 cm. long, 2 mm. thick, equal, *tough*, stuffed then tubular, glabrous, *dark fuscous to blackish*. SPORES 6.7 x 4.5 micr." (Britz.)

Reported by Longyear. Chandlers. June. Rare.

Said to be gregarious on the ground, and imitating in color the blackish species of *Collybia*, like *C. atratus* and *C. ambustus*.

## MUSHROOM POISONING

BY O. E. FISCHER, M. D.\*

No one who has not followed the development of the study of mycology in its scientific or popular fields during the last twenty years can realize its development and the changes in our views that have taken place. Within this time a number of popular works have appeared, keeping pace with the great volume of general nature literature. Mycological and toadstool clubs have been organized in many cities in order to interest people in this fascinating branch of botany. Such clubs have easily won the interest of people at large because of their appeal on the score of mycophagy, the eating of fungi. In the purely botanical and technical field, colleges, universities and state herbaria have given more and more attention to the scientific and economic aspects of mycological study and have brought about the publication of journals devoted exclusively to studies made in these specialties. For centuries almost every community has had its enthusiastic amateur botanist who has collected and sought to name—to classify—the higher plants of his hunting ground. Now cometh also the humble collector of toadstools and mushrooms who finds that his hobby meets with sympathy and interest. Thus "The Spectator" in Outlook (January 13, 1915) gives a charming account of his initiation into the accuracies and delights of mycology.

It has been the purpose of Dr. Kauffman to supply in these volumes a manual for the use of both the amateur and advanced student of the Agaricaceae or gilled fungi of Michigan. It is similarly the purpose of this chapter on mushroom poisoning to place before layman, mycologist, mycophagist and physician an account of the present state of our knowledge of the subject. Having in mind such a varied class of readers, the author must include matter which will seem hopelessly technical to the one, and matter that may seem superfluously simple to the other. It is not the purpose of the paper to present much that is either new or original but it does

\*The author would acknowledge his thanks to the libraries of Parke, Davis & Co., to the University of Michigan and to the Wayne County Medical Society, as well as to Professor Kauffman and to numerous private correspondents and co-workers whose services have aided in the preparation of this paper. Our knowledge of mushroom poisoning is still far from complete and further aid from any source will be welcomed.  
Detroit, Michigan, 507 Field Ave., March, 1915.

seem possible, in the light of recent advances, to record more vital information about deleterious and suspected species than is to be found in similar articles in all the text books combined. The strictly technical literature of the subject is very large. A bibliography might easily include one hundred books and papers in French, German and English. Most of these are quite inaccessible to the average student and an attempt is made herein to present part of the matter they contain. No popular handbook covers a tenth of the field. The earliest views, still popularly held, regarded the Agarics as a large group of poisonous plants. Then, under the influence of teaching-mushroom-clubs and the invaluable published results of Peck and of McIlvaine and Macadam, the impression that there were but few deleterious and hundreds of edible species, has led to a reckless mycophagy which resulted in the discovery of new harmful species. These can no longer "be counted on the fingers" (Plowright), nor will Bagnall's "dozen in over a thousand edible" include all the toxic species. The bibliography, *On Mushroom Poisoning*, will guide the reader to the sources found most useful in the preparation of the paper.

A number of factors give myco-toxicology or mushroom intoxication increasing importance. The daily press endeavors to keep alive the fear of "mistaking a toadstool for a mushroom" in its frequent reports of the disastrous consequences of such a blunder if the toadstool is eaten, but the Sunday paper is not so consistent inasmuch as it prints and reprints irresponsible articles or quotes unreliable and dangerous rules and tests to apply, which, if followed, will lead the man of newspaper education into real danger. The enormous influx of foreigners from southern Europe, accustomed to seeing in the markets and gathering and eating certain species of higher fungi at home, gives us individuals who mistake some American deleterious species for an edible European one which resembles it. These people furnish most of the cases of poisoning which occur in the United States. It is affirmed that nearly all of more than thirty deaths from mushroom poisoning in and near New York City in 1911 occurred among them<sup>29,\*</sup> Through the growth of nature and mycological clubs, the sale and use of the several excellent popular books and bulletins, and the offering of wild species for sale in our markets, the use of fungi for food is rapidly increasing. This necessarily means that a larger number of poisonings of both major and minor importance will come about, since insufficiently trained, self-constituted "experts" may blunder

\* For numerical references see bibliography: *On Mushroom Poisoning*.

or fall into minor error. Some reasons for even the well trained student going a bit wrong will appear in other paragraphs. If he will but report his error in the manner indicated below much, or even all, may be forgiven or even approved.

Mushroom poisoning must have been fairly frequent in early times since it is well known that the Romans employed fungi in great quantity both as delicacies and as daily food. Paulet, in 1793, records their collection in Russia, China, Hungary, Italy and especially in Tuscany, and their public sale in Peking, Petrograd and Florence. Thus they have numbered among their victims the family of the Greek poet Euripedes, a wife, two sons and a daughter, Pope Clement VII, Emperor Jovian, Emperor Claudius, King Charles VI of France, and Czar Alexsis of Russia<sup>1</sup>. The Princess of Conti nearly lost her life through mistaking *Amanita muscaria* for *A. caesarea*. Definite knowledge of the number of fatalities from mushrooms begins with Paulet who states that from the year 1749 to 1788 there were a hundred deaths in the environs of Paris alone. More recently (1883) Bardy reported 60 cases in 6 years in Les Vosges, and Guillard (1885) estimated 100 deaths annually in southwestern France. Falck collected 53 cases in Germany with 40 deaths and Inoko in Japan reports 481 cases of mushroom intoxication in 8 years (1889). In this country Palmer, of Boston, collected 33 cases with 21 deaths and Forster, of Charlestown, 44 cases with 14 deaths<sup>1</sup>. Bagnall<sup>2</sup> quotes Clark and Smith to the effect that in one ten-day period (September, 1911), 22 deaths occurred in New York City and vicinity, 15 in 1906 and 30 cases with 12 deaths in 1905. In 1913 there were 26 cases of poisoning in Hartford in a few weeks. In 1900 Gillot found over 200 authentic cases of mushroom poisoning mostly in France (123 fatal due to *Amanita phalloides*) and Ford<sup>3</sup> added nearly as many more recorded in the German, English and French literature since 1900. Sartory, in France, records for the summer of 1912, 249 cases of fungus poisoning with 153 deaths. Of these 90% occurred in 15 days<sup>4</sup>. Ford is convinced that the majority of cases do not find their way into medical literature. I do not believe that 10% do. Thus in one summer there were unreported 2 cases (not fatal) in Baltimore, 2 deaths in Cleveland, 9 poisoned in Fostoria, Ohio with several deaths, and 10 in Toronto with 2 deaths. Murrill<sup>7</sup> estimates the annual deaths in the United States as probably 50 or more, as many are not reported. My own records, by no means complete, for southeastern Michigan only, for 10 years show 77 cases with 16 deaths. None reported medically. Most cases undoubtedly escape publicity.

## POISONING BY WHITE-SPORED AGARICS

*AMANITA PHALLOIDES*(See *A. verna*, *virosa* and *bisporiger*.)

*Amanita phalloides* is by long odds the most important of all poisonous mushrooms. It is widely distributed, common throughout most of the season and often exceedingly abundant. It is innocent in appearance, of delicious taste and of extreme toxicity. In considering it we may regard *A. verna*, *A. virosa* and *A. bisporiger* as included under the term "*A. phalloides*." Bulliard a hundred years ago gave it the common name "Destroying Angel." It is also known as the Death Cup, White or Deadly Amanita. The earlier species, named in Europe, such as *Amanita bulbosa* and its varieties, *alba*, *citrina*, *virescens* and *olivacea*, *Agaricus bulbosus*, *Amanita viridis*, *A. venenosa* and a number of others are without doubt identical. In older French literature it is known as *Vorange cigue*, *Vorange souris*, *Vorange blanche ou citronee*, *Vorange cigue jaunâtre* and *Vargaric bulbeux* and in German as *Giftwulstling* and *Knollenblaetterschwamm*.<sup>1</sup> Its identification is comparatively simple even in its disguises and no one who does not know this fungus well should dare to eat wild mushrooms or to recommend a portion of such to his neighbor. The possibility of its presence in a collection intended for the table should always be rigorously excluded. The 153 deaths of 1912, above mentioned, were due chiefly to *A. phalloides*. Gillet's thesis<sup>4</sup> maintains that all fatal cases are due to this Agaric. This is not strictly true. The statement that nine-tenths of all fatal cases are due to it, seems conservative enough. In 1845 Orfia reported 8 ill and 4 dead from its ingestion; Bock Ziemssen, 11 fatal cases; Mautner (1861), 4 cases with one death; Handford (Lancet, 1886), 2 fatal; Palmer, 16 cases, 7 fatal; Tappeiner, 5 cases, 2 fatal; Pfromm, 4 Italians, all died; Plowright, 6 cases, 4 fatal; Bulletin of French Mycological Society, 18 deaths 1900-07; in October, 1884, eleven children died in an orphanage in 5 days. Incomplete records for southeastern Michigan (10 years) show 16 deaths in 44 due to *phalloides* illnesses. In 22 of this 44 the white the Amanita was surely to blame, in 14 probably, in 5 presumably.

Surprisingly small quantities may bring on fatal consequences

and there are numerous deaths on record from eating one or two good-sized specimens.<sup>1</sup> Plowright<sup>2</sup> has reported the death of a child of ten years from one-third of the top of a small plant eaten raw; Pfromm, that of two children after taking a bit of juice soaked into bread.

The mortality from *A. phalloides* intoxication is extremely high, varying from 60 to 100 per cent and is dependent largely upon the amount ingested and probably somewhat upon the treatment.<sup>1</sup> Sixty to one hundred percent seems too high for adults, judging from local cases. One-half this would be more nearly correct, unless much is eaten or several children are included in each group. Recovery may be regarded as rare but not impossible. It may follow eight to twenty-one days of extreme suffering. Practically all deaths from mushrooms are attributed to this one species.

The cruelty of the poison and the horrible suffering it causes its victim may be faintly realized from a perusal of the clinical histories of the more fully reported cases. One gets the impression that scarcely another agent bears equal power to torture. A few typical, though varied, cases may be briefly quoted. Thus an Italian family at 6 p. m. on Sunday ate heartily of a cooking of fungi. By midnight vomiting had begun, attended by violent abdominal pain, headache and extreme thirst. A doctor was summoned the next morning. The father was cyanotic and twitching and delirious. The pupils were contracted. He improved slightly for a few hours. Periodical remissions and exacerbations of symptoms continued for eight days when coma and death supervened. The mother presented similar symptoms, with thirst and vomiting more violent, miscarried at five months and also died on the eighth day. Both children died in 58 hours. (Pfromm.<sup>3</sup>)

The Deep Valley, Pennsylvania, cases occurring in August, 1907, are of especial interest.<sup>8</sup> About 7 p. m. on Sunday a physician, three others and the man-of-all-work ate one quart of mixed fungi fried in flour and butter. Before 2 a. m. all began to suffer from excessive vomiting and violent diarrhea which continued all day Monday. Atropine, narcotics and an oil purgative were given. The gastrointestinal symptoms continued three days accompanied by subnormal temperature, more or less delirium (no salivation or urinary suppression) and in case of Dr. D., severe muscular cramps of limbs and abdominal muscles. His death occurred on Thursday morning. By Saturday the man-of-all-work was up and about, the three others still abed. In these, vomiting and diarrhea had ceased and an enlarged liver, distended gall bladder and jaundice were appear-

ing. The man who had gathered the fungi conducted the investigator to the place and indicated the varieties gathered and what had constituted most of the lot. These were *Amanita phalloides*, with smaller numbers of *Cantharellus*, *Amanitopsis vaginata* and a very few *Russula emetica*.

The following group of cases is very typical. In September, 1911, six persons were poisoned, two fatally, in Cleveland. The children, aged four and six, had a little gravy and recovered after nausea, vomiting and diarrhea. Mr. C., aged 67, ate some at supper, felt bad during the night but ate more for breakfast! About noon violent illness began with intense pain in the epigastrium, vomiting and diarrhea with loss of control, clonic spasms and great prostration. Urinary suppression was obstinate and lasted till death, three days after the first meal. Mrs. C., aged 65, ate one forkful at the first meal but did not like the taste. Profuse vomiting and diarrhea with great prostration began one hour after Mr. C.'s symptoms. She recovered rapidly after two days. The daughter-in-law, aged 40, ate at the second meal and, though feeling hot and feverish, ate more at noon! Eight hours later she had exactly the same symptoms as Mr. C. The physician arrived early on the next day. He gave oil, began stimulations with strychnia, nitroglycerin, aromatic spirits of ammonia and, after removal to the hospital, saline solution continuously and oxygen. Some improvement was noted except that the heart action was weak and intermittent and the extremities could not be kept warm. Hiccough for two days, great agony and unconsciousness preceded death on the seventh day. A son, aged 19, ate the second meal—breakfast. Though feeling bad, he worked until 4 p. m. By 9:30 he presented the same symptoms as mother and grand-father. After ten days of apparently as grave illness as theirs under "most terrific stimulation" (nitroglycerin, strychnia, oxygen and salines) he was reported out of danger though "looking like a corpse." "The fungi were gathered from a shady hillside. Some were over six inches across and white inside and out; others were yellow as saffron through and through and about four inches across. Others were white outside and brown under; some were small, white on top and pink under." The brief case histories leave no doubt as to *Amanita phalloides* being the offender. The botanical notes admit of considerable speculation. In every respect this report is pregnant with meaning—and full of food for reflection—for the student of toadstool poisoning.

Grouping the clinical histories from numerous sources the symp-

tomatology of *phalloides* intoxication may be described. When due to the deadly *Amanita* alone the clinical symptoms are practically always the same and are characteristic. Nevertheless IN EVERY CASE OF POISONING the etiology, i. e., WHAT FUNGUS HAS BEEN EATEN should by all means be determined at the earliest possible moment! Why? Not only for its scientific importance to myco-toxicology but more especially for guidance as to what treatment is indicated and required and very especially what the outlook may be for the patient. If *A. phalloides* can be ruled out the prognosis at once becomes very much better and useless fears may be allayed. The sufferer is entitled to this. Uneaten fungi should be submitted to competent authority, more should be gathered from the sources whence the suspected species were derived, and the opinion of the patient or of some one who gathered them with or for him as to their identity with those eaten, obtained.—Returning to symptomatology: After ingestion there is a prodromal stage of from six to fifteen hours—generally over ten—in which little or no discomfort is felt. Then follows a sudden seizure of extreme abdominal pain, cramp-like in character, accompanied by vomiting and diarrhea of undigested food, with blood and mucus. Discharges soon become cholera-like (serous) or rice-water in character. There is burning, consuming thirst. Anuria is usual; constipation rare. Prostration and sleeplessness, with the great nervous restlessness of weakness and suffering, are conspicuous. Muscular spasms in various groups are frequent, accompanied by cries or screams of pain. Loss of strength is rapid and excessive. Periods of pain and vomiting alternate with remissions and ameliorated symptoms. Haemoglobinuria does not occur. Within a few days jaundice, cyanosis and coldness of the skin and extremities develop followed by profound coma from which the patient does not rally. Ocular symptoms, the pupils varying, and convulsions are rare but may occur. Convulsions are often terminal, and death is due to cardiac failure. The course of the disease requires from four to six days in children and eight to ten in adults but death may occur within 48 hours if large quantities of the fungi have been eaten or they have not been thoroughly cooked. These points should be weighed in considering prognosis. The resemblance of the clinical picture to that of cholera and to acute yellow atrophy of the liver has often been remarked.

Atypical features occur especially in cases where *Amanita phalloides* was not proven to have been the sole etiological factor. Such cases may show dilated pupils, clear cerebration, albuminuria.<sup>2</sup>

Schuerer's six cases—most thoroughly studied and reported<sup>10</sup>—showed cramps in calf of leg, arms and other muscles, the left arm escaping in one. Many days of pains in the legs persisted. Recovery was more or less rapid according to age but the youngest (5) died in thirty-five hours after violent convulsions and coma. If recovery takes place the liver and spleen enlarge about the third day, after which day, according to Maass,<sup>12</sup> the prognosis becomes better. "In clear-cut cases the physician can diagnose the variety of toadstool from the typical symptoms."<sup>10</sup>

Reports on the post mortem findings in man in fatalities due to *A. phalloides* are not overly satisfactory. There is little to be found to account for the violent paroxysms of pain, vomiting and diarrhea. Schuerer found, in a child, colitis, pleural haemorrhages and fatty degeneration of liver, heart and kidneys. Microscopically there were "Very wide-spread and obviously severe lesions of the cell elements of the central nervous system, as heretofore hardly known in this form and to this extent"—regressive changes like those seen in the septic deliria. Harmsen, Maass and Kobert liken the postmortem findings to those of phosphorous poisoning. Thus, the normal liver contains from 8 to 25 per cent of fat, that of phosphorus and of alcohol poisoning 50 to 70 per cent and that of *Amanita*-toxin (2 cases) 53 and 69 per cent. Death seems to be due to this extreme fatty degeneration of the liver. (Ford, Schuerer.) Medico-legally, such a liver, with the addition of the other findings, makes the postmortem picture pathognomonic of *Amanita phalloides*.<sup>10</sup>

#### Treatment of Poisoning by *A. phalloides*

All authorities agree unanimously that therapeutic measures in these grave emergencies are almost useless. Case histories show that often the cause is not recognized, or the gravity of the cases not appreciated. There is no antidotal drug for *Amanita*-toxin and the treatment is that of poisoning and septic intoxication in general. Competent medical advice should be obtained as soon as possible. Active emetics (ipecac, mustard, apomorphine), assisted perhaps by the stomach-tube, purgatives (castor oil being preferable to the salines) should be administered at once and every effort thereby made to reduce further absorption of the poison to a minimum. By the time symptoms from *A. phalloides* have begun, the toxin is already in the circulation. High enemata to empty the lower bowel may be used early. Later normal saline solution should be given thus, or hypodermically, or even intra-

venously to supply the body's need for fluid and to ease the torturing thirst. Narcotics and anodynes in large doses are necessary to relieve the intense pain and to quiet convulsive movements. Nitroglycerine and strychnia frequently, and up to the limit of tolerance, are of great value. Cyanosis calls for oxygen inhalations. Atropine may be of use as a stimulant and a corrective with morphine *but it has no antidotal value here*. Milk, raw or boiled, may be regarded as a mild natural antidote.<sup>18</sup> Alcohol should probably not be given in any form. Strong coffee is indicated as are hot dry applications to the body. Digitalis may be used but requires from six to ten hours before effects are seen. Camphor<sup>10, 30</sup> in sterile oil given subcutaneously every hour is valuable. Suprarenal extract is mentioned. Large draughts of hot water, flaxseed tea, slippery elm or starch water<sup>14</sup> may be used, as well as tannigen, bismuth subcarbonate and opium to quiet excessive diarrhea and vomiting. Supportive measures and good nursing are of the greatest importance. Transfusion of blood would seem worthy of trial in graver, slower cases.

Ford, finding that protective and curative sera were theoretically possible, worked for three years on the serum-therapy. He was able to immunize animals to the aqueous extract up to five or six times the fatal dose<sup>15</sup> but efforts to manufacture a curative serum have thus far been unsuccessful.

The Fochier treatment by abscess of fixation<sup>53</sup> has been applied by Dr. A. Pic of the University of Lyon in 23 cases of which 9 died. The conclusion that "it is a therapeutic agent of the first order in those terrible intoxications due to *Amanita phalloides*" seems to have made no impression on the medical profession. Judging from Michigan cases—37 illnesses with 16 deaths—this is not a remarkably low mortality, 39 per cent versus 42. Pic and Martin contrast it with "the usual 86.8," based on 38 cases with 33 fatalities in France, 1913.

#### Poisonous Constituents of *Amanita phalloides*

The first attempt to obtain the active principle or poison of *A. phalloides* is probably that of Letellier, who in 1826 obtained a heat-resistant substance from a number of fungi. This was termed amanitin. Later he found two substances, one of an irritating nature, acting upon the mucous membranes of the alimentary canal and another characterized as a glucosidal alkaloid—the *Amanitia*. Boudier in 1866 ascribed the poisonous action to an

alkaloid which he named bulbosine, but was never able to isolate. In 1877 Oré concluded, on biological grounds alone, that *Amanita phalloides* must contain an alkaloid and this hypothetical poison he named phalloidin. These names are no longer employed. Kobert (1891) established the important fact that extracts of *A. phalloides* contain a substance which lyses or dissolves the red blood corpuscles of many animals and of man. With this "hemolysin" we shall have much to do in the remainder of this paper. Ford and his co-workers have investigated it most satisfactorily in their epoch-making labors which have been fully reported. This hemolysin is not the active principle—for we shall see that it is very easily destroyed by heat, much less than is usually employed in cooking, and that the digestive juices break it up as a rule. Furthermore, individuals dying of *A. phalloides* intoxication do not show symptoms which are to be ascribed to this kind of poison. Kobert gave this blood-dissolving hemolysin the name of phallin, regarded it as the essential poison, and gave it undeserved importance. He placed it in the group of protein-like poisons known as toxalbumins because of its susceptibility to destruction by heat. Beside the hemolysin, and more constantly present, Kobert found later an alcohol-soluble substance which was extremely poisonous to animals. This he regarded (1900) as an alkaloid, soluble in alcohol, which would not produce fatty degenerations. A toxalbumin (near thujon and pulegon), was held responsible for these.

Frey,<sup>16</sup> in 1912, comments, "The whole study of mushroom poisoning still lies very much in the dark. It is on the same plane as thirty years ago." He says that studies on muscarin and phallin show old results and theories to be wrong, but otherwise there is no progress. The publication of results of recent American investigators seems to have been unknown to him, for progress has been made, and a basis for further results established. Murrill<sup>17</sup> comments (1910) that it is remarkable how little is really known, and that the practical importance of the subject is vastly increasing. The important work of recent American investigators began with the proof (Ford<sup>18</sup>) that extracts of *Amanita phalloides* contain the hemolytic material described by Kobert and in addition a heat-resistant body which will reproduce in animals the majority of the lesions described in fatal cases of *A. phalloides* intoxication in man. These two substances were named by him the Amanita-hemolysin and the Amanita-toxin. The further chemical study upon the plant was carried out by Abel and Ford,<sup>19</sup> by Schlesinger and Ford,<sup>20</sup> and Ford and Prouty.<sup>21</sup> In these papers it was shown

that the hemolytic agent is not proteid (toxalbumin) but an easily-decomposed glucoside, insoluble in alcohol, extremely sensitive to heat, to small traces of acid, to pepsin and pancreatin, and that it can therefore play no role in poisoning in man when the fungi are cooked. It may be a factor if large quantities are eaten raw or insufficiently cooked, or if through deranged digestive action the hemolysin escapes destruction.<sup>15</sup> It is present in such great amount that under such circumstances the possibility of its having a poisonous action cannot be eliminated. The Amanita-toxin has so little in common with alkaloids that they hesitate to class it with them.<sup>15</sup> Amanita-toxin is the alcohol-soluble active principle, *the essential poison*, resisting the action of heat, of drying and of the digestive juices, and reproduces in animals the lesions found in *phalloides* intoxication in man. Chemically it cannot be characterized definitely, but the purest preparations do not give the reactions of either proteins, glucosides or alkaloids. Rabbits are not affected by various extracts by mouth, both the hemolysin and Amanita-toxin being quite innocuous to them, when one-fortieth of the amount was fatal when given subcutaneously. Dogs and cats are poisoned by the cooked fungus in the same degree as human beings. The raw hemolysin given subcutaneously has pronounced blood-dissolving properties, giving the picture of a hemolytic intoxication with extreme haemaglobinuria and pigmentation of the spleen.<sup>13</sup> [These are the properties assigned to the European *Helvella (Gyromitra) esculenta*.] Even when made from dried specimens of *Amanita phalloides* hemolysin will dissolve the red blood cells of guinea pigs, rabbits, fowls, pigeons, dogs, goats and man. Swine, sheep and beef bloods are not susceptible. The blood of the guinea pig is most susceptible and that of the goat least. When this hemolysin is heated to 140° F. it loses some of its activity, and 150° maintained for one-half hour, suspends it. (Hence the term "thermo-labile.") It may be classed with the bacterial hemolysins. Injection experiments on animals show its extreme toxicity. Within a few hours the fur ruffles and they refuse to eat. There is rapid loss of weight and strength, death occurring within one to three days under great dyspnoea. The heart stops last. In smaller doses a chronic intoxication is produced lasting three, four or six weeks. Convulsions are unusual and there is no salivation or gastro-intestinal disturbance—in contrast to muscarin poisoning.<sup>18</sup> Frequently, even in *A. phalloides*, the hemolysin is present only in small amounts and it may be absent whereas the edible *A. solitaria* and *A. rubescens* contain

it in great abundance. "It is probably a food and certainly harmless,"<sup>23</sup> i. e., when cooked and eaten by man.

The Amanita-toxin is the more active and acutely fatal, producing approximately the lesions seen in man from the whole cooked plant, ulcers in the stomach and intestine, serious hemorrhages, and in other organs, especially liver and kidney, cell necrosis and fatty degeneration.<sup>13</sup> It loses potency somewhat but not greatly on boiling. In a later report<sup>31</sup> Ford and Brush say that *Amanita phalloides* var. *citrina* gathered in France corresponds in all particulars to the *A. phalloides* gathered in America, and has identical properties and contains the same poisonous substances.

#### OTHER AMANITAS

*Amanita verna* and *A. virosa* have been already mentioned as included in the above section. *A. spreata* is recognized to be deadly poisonous. A group in which Amanita-toxin is present in small quantities includes *A. porphyria*, *strobiliformis*, *radicata*, *chlorinosma*, *mappa*, *morrisii*, *citrina*, and *crenulata*. The first four of this latter group are devoid of haemolysins but owe their toxicity to small amounts of Amanita-toxin. In *A. spreata* the hemolysin is small. The extract caused both acute and chronic intoxication in guinea pigs but not in rabbits. Poisonous, Boston Mycological Club, and Atkinson.

Ford and Sherrick<sup>41</sup> found in *Amanita mappa* a small amount of thermo-labile hemolysin, a chronic intoxication of guinea pigs closely resembling that of *Amanita phalloides*. No muscarin. Rabe says *A. mappa* has the same poisons as *A. phalloides* but in much smaller amount. It should be classed as perhaps less dangerous than *A. phalloides*. Other Amanitas may be grouped here by Ford's reports.<sup>22</sup> The agglutinin will receive attention in the account of *Amanita muscaria*. The chronic intoxication is shown by a progressive emaciation and death in 18 to 20 days.

*Amanita citrina* (of Europe) a yellow variety of *A. phalloides* (Kobert). No hemolysin or agglutinin. Poisonous to guinea pigs and rabbits by both acute and chronic intoxication. Seldom, if ever, toxic to man.<sup>18</sup> A distinct species from *Amanita phalloides* var. *citrina*.

*Amanita crenulata*—No hemolysin or agglutinin. Chronic intoxication in guinea pigs and rabbits. Extract made after one year of drying was fatal by chronic action after an acute. Poison,

small in amount, similar to Amanita-toxin. McIlvaine records it as edible.

*A. morrisii*—Small amount of hemolysin destroyed at 60° C. Poisonous to guinea pigs and rabbits. Should be grouped with the deadly species. Edibility not tested.

*A. chlorinosma* is probably seriously poisonous. *A. strobiliformis* acts like *phalloides* on frog's heart.

The species of the genus *Amanitopsis* as a whole are regarded as edible. McIlvaine warns against confusing *A. spreata* with these species.

*Amanitopsis volvata* may be grouped with *phalloides*. No hemolysin or agglutinin, fatal in 7 to 22 days to guinea pigs and rabbits, the intoxication resembling Amanitas. McIlvaine pronounces it edible, but it should be avoided. *Amanitopsis vaginata* is easily learned and is edible.

*Amanita junquillea*—rare and unimportant—free from poisonous properties.

*Amanita solitaria*—difficult to recognize. Contains small amount of hemolysin. Edible, McIlvaine. Ford<sup>32</sup> reports it almost free from poisonous action on rabbits and guinea pigs, but large doses produced a salivation in the latter. *Muscarin is more widely distributed in fungi than was originally supposed.*

*Amanita rubescens*—commonly known as "The Blusher," Red Amanita. Non-toxic to animals and man. Free from Amanita-toxin but has a powerful hemolysin. European authorities differ but our American form is a well-known edible species.

*Amanita frostiana*—difficult to identify, is closely related botanically to *A. muscaria*, of which it has been regarded as a small or depauperate form. It contains a small amount of a thermo-labile hemolysin, but no muscarin. Its extracts have no effect on animals. Not tested but probably edible.<sup>22</sup> CAUTION! lest *A. muscaria* be used.

#### AMANITA MUSCARIA

*Amanita muscaria*, the Fly Agaric, is a most interesting fungus. It is also called "the false orange" and "Fliegen Schwamm." It is less common and less toxic than the group of *A. phalloides* but is widely distributed over the world. In importance it ranks next to the white Amanitas. It is subject to great variations in color, size and markings, but is easily learned so that it may be distinguished from the famous edible *Amanita caesarea*. Ford<sup>3, 18</sup> and Michael

and others agree that its taste is bitter and unpleasant and this factor may save people from serious accident. Occasionally the bitter taste is absent, more is eaten and quick fatality may result.

Through the publication<sup>23</sup> of Circular No. 13, U. S. Dept. of Agriculture and of Prentiss' account<sup>24</sup> the fatal poisoning of Count de Vecchii in November, 1897, has become classic. He bought from a countryman a quantity of *Amanita muscaria*, picked in Virginia, seven miles from the capital. The Count was familiar with mushrooms and took these to be the Royal Agaric, *Amanita caesarea*. At breakfast, which was finished at 8:30, he ate two dozen and pronounced the taste particularly good. Dr. K. ate one dozen. By 9 a. m. the Count was lying on his bed in a state of collapse, filled with a sense of impending death, and soon lost consciousness. Blindness came on before this, as did rigid spasm of the lower jaw, and difficulty in swallowing. Convulsions were so violent as to break down the bed. Emetics were given and apomorphine and atropine subcutaneously. He became continually worse and died without regaining consciousness on the evening of the next day. Dr. K. went by car to his office. While sitting on a chair, about 9 a. m., he gradually passed into unconsciousness without feeling any premonitory pain or distress, though half-stupid and very restless just before. He noted, about 9:10, uncertain eyesight and double vision, without nausea. A prominent early symptom was sudden jerking back of the head. He remained unconscious for five hours; at one time his life was almost despaired of. He did not suffer the least pain but on the contrary was in a comfortable dreamy state. By 7 p. m. he was out of danger. Cold sweats were a prominent symptom. A total of one-tenth grain of atropine was given in 24 hours. Apomorphine produced no emesis, vomiting not occurring until evening. Castor oil and sweet oil were given about noon.

**THE CLINICAL FEATURES** of poisoning by *Amanita muscaria* are quite as characteristic and distinctive as those in *Amanita phalloides* intoxication and should enable physicians to distinguish clearly between the two conditions—when either fungus is eaten alone. So often a mixed lot of different varieties is used that the symptoms in patients point to the combined action of different toxic principles. In *A. muscaria* poisoning there is usually a very short interval between ingestion and first symptoms, one-half to one hour or at most three hours. If small amounts are eaten even five or six hours may elapse. This feature is of greatest value

in deciding upon the kind of intoxication which the cases present. Severe ones show excessive salivation and perspiration, a flow of tears, a feeling of laryngeal constriction, nausea, retching, vomiting and watery diarrhea. The last named almost always occur. The pulse is usually slow and irregular. There is no fever; pupils are small. Respirations are accelerated and the patients dyspnoeic, the bronchii being filled with mucus. (The action of atropine is the opposite of this, point for point.) Mental symptoms are also present, particularly giddiness with confusion of ideas and rarely hallucinations. All these symptoms may vary in intensity, at some times the gastro-intestinal predominating, and at other times the mental. In light cases only salivation or perspiration may be noticed, with uneasiness in stomach and bowels, for a few hours. In severe cases the vomiting and diarrhea may rapidly rid the alimentary canal of the offending material and the nervous symptoms then become predominant—delirium, violent convulsions and loss of consciousness developing in rapid succession and the patient's sinking into a deep coma. Rarely, consciousness is retained till the end, death resulting from paralysis of the respiration. Finally, in many cases, after the vomiting and diarrhea, the patients sink into a deep sleep, awakening later profoundly prostrated but on the road to recovery. Normal health reappears rapidly—two or three days. There are no late effects in *muscaria* intoxication as in that of *Amanita phalloides* with its degenerative changes in the internal organs. The prognosis is always good if the patient recovers from the preliminary symptoms. When, rarely, the nervous symptoms dominate the alimentary, excitement and hallucinations simulate alcoholic intoxication. (Quoted freely from Ford and Clark.<sup>25</sup>) The delirium is occasionally followed by loss or impairment of memory. The pupils dilate as death approaches. The action of muscarin is almost identical with that of pilocarpin.

Post-mortem examination reveals surprisingly little. The pathology of *Amanita phalloides* is absent, particularly the lesions of the liver. In general the findings point to the action of a profound nerve poison.<sup>3</sup> Medico-legally, remains of fungi in the alimentary canal would be of great importance.

#### Poisonous Constituents of *Amanita muscaria*

Schmiedeberg and Koppe, in 1869, showed by the most careful work, both chemical and pharmacological, that *Amanita muscaria* contains an active principle which they called muscarin. At first regarded as an alkaloid of the same general nature as strychnine

and morphine, later work has shown that it is probably a complex ammonia derivative. Muscarin is an extremely active substance, well known from the attention it has received in all works on toxicology and materia medica and therapeutics. In the latter field it can well be spared on account of its variability and unreliability and because we have better drugs of similar action. Muscarin is near pilocarpin and nicotine in action, exciting smooth muscle and stimulating all glands. At almost every point in its action it is the direct antagonist of atropine (from belladonna) but is far less powerful. It is present in the fungus in but small amounts but is nevertheless able to exert its characteristic effects, frequently with fatal outcome. In producing paralysis of heart and respiration it does so by stimulating the inhibitory nerve endings of the vagus. Atropin has a depressing action upon the same nerves which muscarin stimulates. The muscarin excitement, remarkably, does not pass over into a paralysis, its curare action (that of arrow-poison) being slight. Muscarin has been synthetically prepared by the oxidation of cholin but does not keep as well as the natural product, and differs materially in its action upon animals.<sup>18</sup> A ptomaine muscarin is also known.

But poisoning by *Amanita muscaria* and muscarin poisoning are by no means identical (Harmsen<sup>20</sup>). Kobert says the fly-agaric drunk (Fliegenschwamm Rausch) is by no means a pure muscarin "jag" but resembles haschisch (*Cannabis indica*). Harmsen found that he could extract from 100 g. of fresh *muscaria* 16 mg. of a fairly pure muscarin. This was twice as deadly to cats as to frogs. That it is not the sole factor in poisoning is shown from the following: (1) with the lethal dose of muscarin at 0.525 g., it would require 4 kg. (8.8 lbs.) of the fresh fly-fungus to produce a fatal outcome<sup>20</sup>; (2) when the action of the muscarin-part of an entire extract is physiologically neutralized by atropin, the animals nevertheless die; (3) the extract is deadly even when the muscarin is removed. He has also shown<sup>20a</sup> that the entire extracts of *A. muscaria* are twice as toxic, weight for weight, as pure muscarin and contain a poison which produces in animals continued convulsions with fatal outcome, not prevented by atropin. He therefore assumes the presence of at least one other substance which he names "Pilz-toxin." This pilz-toxin must be very unstable since it loses potency on drying, and is sensitive to heat (thermolabile). It does not appear in the urine. (Compare intoxication of the Koraks.) His work casts doubt over the value of atropin as an antidote and is in accord with clinical experience. In 1910 Ford<sup>22</sup> said *Amanita*

*muscaria* acts in all animal experiments as a convulsant and no other agaric shows similar action, not even the closely related *A. frostiana*.

#### Treatment for *Amanita muscaria* Poisoning

Just as in cases of *A. phalloides* intoxication, it is the important duty of physician and friends to get all the information possible as to the exact nature of the toadstool eaten and the amount ingested. *It will be shown below that a number of species of fungi, mildly toxic or simply deleterious and unwholesome, can produce a more or less typical picture of muscarin intoxication.* Confirmation of the species will therefore be of great value in determining prognosis and in giving a clue to antidotal treatment. The outlook in poisoning by the fly agaric is more hopeful than when the Destroying Angel (*A. phalloides*) has been ingested, the mortality runs much lower, the illness is briefer and the suffering less cruel—though bad enough. We do not have here the chronic and degenerative lesions produced by the white *Amanita* which defer death or prolong convalescence. The *muscaria* intoxication is acute, comes on soon after eating the fungus, develops rapidly and is amenable to treatment. Recovery often occurs without untoward symptoms. Lachrymation, salivation, contracted pupils, delirium, hallucinations, and coma call for atropin in large doses subcutaneously or intravenously. Even though the vomiting and diarrhea are pronounced, the stomach and bowels should be further emptied by the free use of emetics and purgatives, for parts of the fungus are often found in the canal post-mortem when profuse emptying seemed to have taken place. On account of coma, apomorphine subcutaneously is less apt to work, and other means of emptying the stomach should be begun early (stomach tube, mustard, zinc sulphate, sulphate of copper). In cases with bad heart action, respiratory distress and coma, atropin (intravenously) offers the only hope, though many other measures, as mentioned under *A. phalloides* treatment, should not be neglected. Absolute recumbent rest is enjoined. Sustain the heart. Give nitroglycerin for cold skin and extremities, and dry heat.<sup>24</sup> Atropin is not medically indicated in every case, and good nursing may easily be of greater importance to tide over periods of weakness and depression. Nourishment should be concentrated. Tannic acid is useless; acidulated water bad. Transfusion of blood, oxygen and galvanism, have been suggested.

The fly *Amanita* possesses interest in several other respects. It

is eaten in the Erzgebirge of Saxony and Bohemia.<sup>11</sup> Treated and untreated it has been eaten without bad results. A colored woman in Washington recited in detail how she was in the habit of cooking it. Rejecting gills and peeling the cap, specimens were boiled in salt water and then steeped in vinegar, then washed and cooked and served with steak, the whole process a rational process to remove poisons (?). Michael<sup>27</sup> worked up to eating a thick medium-sized cap (cooked) and "properly peeled." It tasted ill but did no harm. Then he ate a specimen prepared as salad which tasted worse. On this ground he classes it as "inedible." For reasons like this we are loath to take any one man's testimony in the great field of mycophagy. Peck has repeatedly received reports from various people who eat it.<sup>54</sup> He also records the eating of the fine variety *formosa* of *A. muscaria* by a sheep, but Ford<sup>22</sup> suggests that the herbivora are (at least, by mouth) immune to this toxin as well as to others. There seem to be *seasonable and local variations in the toxicity of Amanita muscaria*<sup>55</sup> and of other species.

One-tenth of a raw *A. muscaria* has produced in a man of thirty-seven years, eleven days of illness, with typical *muscaria* symptoms, but accompanied by fever.<sup>56</sup>

The use of *Amanita muscaria* simply and purely for producing drunkenness is well known, but has not been satisfactorily explained. Krashennikoff, who travelled in Siberia and Kamchatka for ten years (1733), reports that the Koraks used the fly Amanita—three or four for a moderate dose, and ten for a thorough drunk. Langsdorff (1803) confirms this and Kennan<sup>28</sup> describes it in some detail in his first Siberian journey. The natives call the fungus "muk-a-moor." Its sale has been made a penal offense by Russian law but "prohibition does not prohibit." One fungus may sell for \$20 worth of furs, and supply does not equal demand. The dried cap is used; a duly flavored decoction is made from them or pieces are swallowed whole. First effects come on rapidly and make the candidate cheerful and merry, then drowsy and sleepy for ten or twelve hours and he awakes in a state of exhaustion. During the stage of excitement there is a horrible kind of delirium and the experience of visions of varied character. The intoxication is prolonged or passed on (among the lowest and most degraded Koraks) by drinking the renal excretion and thus a spree may be economically kept up for a week.<sup>24</sup> Evidently the muscarin is excreted unchanged. (See Ziemssen, *Fungus Poisoning*, Vol. 17.) Toleration develops, though death from an orgy is not uncommon. The meat of animals dead of muscarin poisoning has a pronounced

poisonous action if eaten by others (Stellar & Erman). In regard to the use of *Amanita muscaria* as a fly poison, D. R. Sumstine (Penn.) reports that the apparently dead flies revive fully in about two hours. One of our mycologists has seen them recover after two days. Tappeiner<sup>36</sup> states that the fly poison is easily destroyed.

#### Toxic Principles of Amanitas

Harmsen's "Pilz-toxin" was never confirmed. Ford<sup>22</sup> agrees that *A. muscaria* owes its action to muscarin but in place of the second poison hypothesized by Harmsen it contains also an hemolysin (as in *A. phalloides*) soluble in alcohol and a *constantly-present agglutinin* belonging to the glucosides. Agglutinins are bodies capable of causing groupings, coherence or agglutination of blood corpuscles when brought in contact with them. They act directly on the blood cells. Given subcutaneously the agglutinin of *muscaria* always caused death in typical convulsions. Violent cooking of the plant, deadly without boiling, was shown to destroy both the muscarin and agglutinin. Subsequent studies of other fungi were based upon a search for the actions of the four active agents thus far enumerated. We have seen from the foregoing consideration of two deadly toadstools that *Amanita phalloides* contains two poisons, (1) an hemolysin which is thermolabile and also easily destroyed by the digestive juices and (2) an Amanita-toxin which is the very definite and powerful poison of the species. Now in *Amanita muscaria* we have (1) muscarin, a poison with its characteristic and individual physiological action, (2) hemolysin in small amount and (3) an agglutinin. Agglutinins are not common in plants. Out of ninety-nine examined they were present in four non-poisonous Papilionaceae and in six Daturas. Among 40 fungi they were present in one-quarter, thermolabile in some, in others heat-resistant. They resisted drying of the fungi better than did the hemolysins and were found to last for years in dried *A. muscaria*.<sup>22</sup>

*Amanita pantherina*, though rare or lacking in America, may be associated with *A. muscaria* since it is said to be used in Japan to produce mushroom drunkenness. Muscarin has been isolated from it as from the Siberian fungus. Delirium, dilated pupils and hallucinations with visions of beautiful red, yellow and brown objects predominate over the gastro-intestinal symptoms. *A. pantherina* is also used as a fly poison. Poisoning from it shows the usual alimentary irritation coming on within a few hours, great excitement,

delirium and convulsive seizures. Ocular symptoms, loss of memory and syncope are frequent. Gillot has collected thirty cases with two deaths and Inoko, in Japan, a series of thirty-two with one fatality. Recovery is usually rapid but occasionally convalescence requires fourteen days. Atkinson's *Amanita cothurnata* may be the American representative of *A. pantherina*, hence both of these should be avoided as esculents. *A. cothurnata* will poison flies. *A. pantherina* extracts were without effect on animals but only a few plants were tested.<sup>41</sup>

There has been in recent years a tendency to explain away too many cases of minor poisoning as due to indigestion, decomposition of the abundant proteid of mushrooms, or to the possible insect-infection of good fungi—and to refer too many of the cases to “probably *phalloides* or *muscaria*.” Now the rich labors of Ford and his co-workers, both in the field and in the laboratory, and the results of Clark, Smith and Kantor have verified certain clinical experiences and shown us that the list of more or less poisonous species must be considerably extended. *Amanita phalloides* and its few congeners still stand quite alone, head and shoulders above all others, for extreme toxicity. They are, most fortunately, not likely to have any rivals for dangerous qualities. They have retained their place easily at the head of the list of noxious species, but the minor and less poisonous list has been somewhat increased. These nearly all belong, in a way, to a *muscaria* group. It will be the problem of pharmacological and biologic chemistry to show why they cause such a variety of clinical disturbances,—by no means explainable by “indigestion,”—and yet resemble the action of muscarin.

#### THE GENUS *LEPIOTA*

In contrast to the genus *Amanita* with its very dangerous species and its few safe edible forms we have in the equally large genus *Lepiota* a number of highly prized edibles. *Amanita* requires close discrimination to distinguish its species; *Lepiota*, for the mycophagist, principally, that he shall not confuse its *L. naucina* with *Amanita phalloides* and that he shall not mistake the black sheep of the section, *Lepiota morgani* or green-spored *Lepiota* for *Lepiota procera*, “The Parasol.” *L. morgani* is an enticing plant and probably the largest Agaric in the world. It is distinctly American. This fine fungus shows very consistent partiality in selecting its victims for it always poisons certain individuals who try to eat it and never distresses others of the same family. It is credited with at

least one death and many serious illnesses. Significant it is that heating destroys the greater part of its toxic properties.

Dr. Blount (Illinois) says<sup>22</sup>: “One day last month the Man of Science of our house came home with a fine specimen of large white mushroom which he took to the library and identified as ‘horse mushroom.’” (If you do not care to discriminate between white-spored and purple-brown-spored Agarics, mycophagy is a dangerous field for you!) “So a few were collected and prepared for dinner. The Man of Science ate a small piece raw at 2 p. m. At 5 p. m., feeling well, he tried another piece, raw, as large as the little finger. At 6 p. m. he felt generally ill and ate no supper. In half an hour he began to have profuse, painless watery bowel movements, but blamed a dentist and his drugs for this.” Dr. B. took two small portions each as large as a pea about 6 p. m. The after-taste created loathing. Discomfort was immediate, and consisted of a warm heavy sensation, slight pharyngeal spasm and difficulty of swallowing. By 7 p. m. vomiting had begun, became very violent and continued every five to ten minutes. Diarrhea began and lasted all of the next day. Intense burning pain in the stomach alternated with intervals of lassitude and exhaustion.

At 9 p. m. hypodermic medication (strychnia 1/30, atropine 1/100, morph. sulph. 1/4) and cocaine produced relief and slumber came on at 11 p. m. Pulse was weak and rapid; perspiration free. The Man of Science vomited three times and had diarrhea all night. He felt as usual the next day. He ate most and suffered least. The action of the poison suggests muscarin.

V. K. Chestnut<sup>24</sup> records that the president of the Chicago Mycological Society mistook *L. morgani* for *L. procera*. Prof. Miller (Terre Haute) eats *L. morgani* and tells of six families that do so. One or two members of each family are made sick, though two families have eaten it repeatedly without trouble resulting. “The meat is simply delicious.” Galveston and Milwaukee record seven cases of illness and V. K. Chestnut adds twenty beside. Detroit might add four. The symptoms are as above described, apparently also from cooked specimens. The fatal case was that of a two-year old child who died in convulsions in seventeen hours after eating of a raw plant. Poisoning has resulted after every variety of cooking and after soaking in salt water. Webster<sup>25</sup> tells of a New England mycophagist who removed to Missouri, identified *L. morgani* as *L. procera* from pictures (!) and paid the penalty within two hours. He draws the moral, “Eat only what you KNOW!” McIlvaine<sup>28</sup> (p. 711) reports another case from Wisconsin of violent illness from

raw (?) *L. morgani*, mistaken for *L. procera* and eaten in very small amount, presenting all symptoms above recorded but with the addition of temporary blindness. Warren (Port Huron) records that in a family of five who ate it two girls were made ill. Symptoms came on early the next morning—seven hours after eating—and were “almost as bad as from Gyromitra poisoning as it is known in Port Huron.”

#### THE GENUS TRICHOLOMA

In this genus McIlvaine agrees to label *T. saponaceum* and *T. sulfureum* as inedible on account of taste. I had for years regarded the entire genus as safe, but in August, 1908, we had a group of seven cases of rather violent poisoning from an innocent-appearing Tricholoma. Good specimens were at once sent to Atkinson who described them as a new species which he named *Tricholoma venenatum*. This agaric has not been found again nor further tested. Of the lot eaten many were badly infested by insects when examined two days later. The symptoms came on one hour after supper and consisted of vomiting, sometimes bloody, retching and considerable prostration in three individuals. All recovered. Surprises like this will continue to occur as long as fungi are eaten. It may be years after some varieties of poisoning occur before the etiology is satisfactorily settled: whether due to a new deleterious species; a known inedible variety not recognized by the consumer; a personal physiologic sensitiveness of the individual; decayed fungi of good species or some infested by acrid insects; the rare presence of a minor toxin in some generally-esteemed edible variety; or simple acute indigestion—perhaps due to gluttony. The observer should endeavor to fix positively the responsibility on the one real cause.

#### THE GENUS CLITOCYBE

Like Lepiota, this large genus has for years been credited with but one deleterious species. Within a few years two others have been added. All three show muscarin symptoms in variety.

*Clitocybe illudens*, known as the “Jack o’ Lantern” because of its phosphorescent glow, or the “Deceiving Clitocybe,” is mistaken every year, in Detroit, by our foreign residents for *Armillaria mellea* or for the European Chanterelle. And on such annual fall occasions it holds high carnival and breaks into the newspapers. The attending physician has a busy night or a few busy hours—

and is credited in the daily press with having saved lives in toadstool poisoning. The mycological investigators visit the family and usually find its members up and about their usual occupations. This is the impression one gets of *Clitocybe illudens* from twenty-nine Detroit cases. The remnants of the feast are usually found to be large half-cooked tough masses. McIlvaine reports a saponaceous taste—and the ability to retain the fungus when eaten. It is possible to make it comparatively harmless by heating it in salt water for half-hour, then taking it out and frying it in butter.<sup>39</sup> Farlow<sup>36</sup> reports illness of four persons. They found the fried flavor excellent. Within two hours all had free vomiting lasting all afternoon, no depression, no intestinal disturbance. No emetics were used since the Jack o’ Lantern carries this property with it and may thus ward off more serious results. At a New York state institution eight teachers and children, after terrible nausea, recovered. No fatalities have been recorded. Diarrhea and prostration may occur. Clark and Smith<sup>37</sup> found that extracts of the plant would stop a frog’s heart which would recover under atropine. Similar results were obtained on the creature when paralyzed by the extract. They conclude that *Clitocybe illudens* exerts a characteristic muscarin effect on exposed hearts which effect is completely overcome by atropin. (Not the case in extracts of *Amanita muscaria*.) Ford finds no hemolysin but the power to produce an acute intoxication in guinea pigs, fatal in one to seven days or a chronic intoxication lasting fifteen days. No lesions postmortem. Rabbits unaffected. After one year of drying boiling for half-hour destroyed the toxicity.

The characteristic American *Clitocybe illudens* has its phosphorescent and related European correspondent in *Agaricus* (or *Pleurotus*) *olearius*, which, mistaken for the Chanterelle, caused illness of the *illudens* type in France.<sup>44</sup>

Fabre writes, “The soft light of *Agaricus olearius* has confounded our ideas of optics; it does not reflect, it does not form an image when passed through a lens, it does not affect ordinary photographic plates.” (Fabre, Poet of Science—LeGros.)

*Clitocybe dealbata* var. *sudorifica* or *Clitocybe sudorifica*, the sudorific Clitocybe, is an interesting little toxic toadstool recently added to the black list. It is often found among “fairy rings” (*Marasmius oreades*). I believe it has been picked with the latter and thus caused trouble, though easily distinguished. The flavor is good. Minneapolis has a record of two cases of poisoning. Peck himself tested it, eating eight caps slightly fried, and got the usual

reaction, i. e., some five hours of profuse perspiration beginning on the forehead and spreading over the body. This may be attended by increased nasal and salivary secretion, hiccough and discomfort, though there are no other ill effects.<sup>42</sup> The original lot was tested on animals.<sup>40, 41</sup> In a rabbit the watery extract produces profuse salivation in a few minutes with weakness and sickness, increased renal activity and activity of the bowels, followed by gradual improvement. Fatal to guinea pig in one quarter hour. Even the boiled extract paralyzed the respiration in seven minutes. Autopsy negative. One rabbit died with slightly contracted pupils. In a third guinea pig there was salivation, lachrymation, etc., increased respiration and then respiratory paralysis. Drops in the eye contracted the pupil for four hours. Its action therefore is that of the muscarin-pilocarpin series. The little *Clitocybe* seemed more poisonous than *muscaria* extract tested side by side with it for it killed rabbits that withstood larger doses of *muscaria* extract. A frog's heart could be stopped for one hour with it and then revived with atropine. *Clitocybe dealbata* should likewise be avoided for *C. sudorifica* has been mistaken for it by a well-trained mycologist.

*Clitocybe morbifera* is similar in habitat and appearance to the preceding and is closely related to it.<sup>42</sup> In four cases in Middleville, Michigan, which have come to my attention, the symptoms were more severe and serious than those of *C. sudorifica*. There was more discomfort and the attending physician recognized the likeness of the clinical picture to muscarin disturbance and used atropine. Four people ate, and three were made ill. The one that suffered most had over-taxed her digestive powers the day before—a factor that seems to predispose to mushroom poisoning. Symptoms came on two hours after eating and were abdominal pain, vomiting of food including entire specimens of tough "Fairy Ring" fungi, purging, sweating, cold extremities and collapse. In one case there was some blindness. All were fairly well the next day. Animal tests have not been made. These must henceforth be regarded as a necessary part of the record.

*Clitocybe nebularis* which made Cordier ill, and is reported as poisonous when raw (Bertillon), is legally allowed among the thirty varieties permitted in the markets of Munich.<sup>11</sup> Here legal enactments, duly enforced, have reduced the number of poisoning cases.

### THE GENUS *HYGROPHORUS*

Experimentally in man no *Hygrophorus* is known to be inedible and there are many fine esculents. *Hygrophorus conicus* used to be forbidden and Demange has attributed a serious outbreak of poisoning to it. It is fatal to guinea pigs by chronic intoxication—as are many perfectly safe fungi. Cooke and McIlvaine say it is all right. *H. pratensis* var. *cinereus* is toxic to guinea pigs.<sup>22</sup> It is edible. Var. *albus* contained a heat-resistant agglutinin and hemolysin: toxic to guinea pigs. Edible. *H. marginatus* similar and edible. See Bibliography, reference 22, for several others. The genus is either devoid of action or poisonous by chronic intoxication only to guinea pigs. An excellent record, so far. The species are clean, beautiful and inviting.

### THE GENUS *LACTARIUS*

*Lactarius* contains some well known edible mushrooms—*L. deliciosus*, *volemus*, *corrugis*, being well liked. McIlvaine says that not a single species retains its pepperiness after cooking. Some of the genus tasted raw are horrible. *L. torminosus* "the griping milky," is charged with having caused fatal illness.<sup>45</sup> In Germany it is known as the "Birken"—or "Gift-reizker." Eleven were poisoned. Three children ate it fried, the youngest, aged two, died in twenty-four hours. Eight Polish laborers, including two women who ate most, prepared it. The women died after six day's illness, treatment coming late in their case. Symptoms came on in about five hours and consisted of nausea, headache, abdominal cramps, vomiting prolonged and even bloody; diarrhea was synchronous, violent and profuse and accompanied by tenesmus. Anuria and albuminuria followed. Skin dry and later jaundiced; pupils dilated; heart negative but weakening; respiration rapid, shallow, irregular and finally Cheyne-Stokes. Temperature normal. Liver somewhat enlarged. The remainder of the description, as well as the postmortem findings in the adults, suggests very strongly that *A. phalloides* was the cause and not *L. torminosus*. Atropin proved without effect in the therapy and the invaluable hypodermoclysis of normal saline solution was not used. No account of why or how *L. torminosus* was settled upon as the cause is given. Hockauf<sup>11</sup> doubts the diagnosis and says many authors (six are named) say *L. torminosus* is harmless when cooked. Krombolz has eaten it though the taste

was bad. Huseman's two cases referred to the species fit *Amanita muscaria*. Great mycologists give conflicting reports. Kunkel says in Sweden it is used cooked and is poisonous only when raw, and this agrees with Ford's results.<sup>22</sup>—Ford found that its hemolysin and agglutinin were destroyed at 150° F. Though acutely fatal to both guinea pigs and rabbits which showed convulsive-like movements, with retraction of the head—a little like *A. muscaria* intoxication but with more somnolence—these toxic effects were not obtained when the extract was cooked one-half hour. Its safety or danger perhaps depends entirely upon the cooking. Maass<sup>12</sup> alludes to the presence in some fungi of drastically purgative resinous acids which may be decomposed by cooking processes and become foods. The milk of the Lactarii seems to be such a substance. "Insects eat both *L. torminosus* and *L. deliciosus*. They pronounce excellent what we find poisonous and vice versa" (Fabre). Hockauf regards *L. torminosus* and *L. zonarius* as poisonous and *L. plumbeus*, *L. chrysorrhoeus*, *L. vellereus*, *L. insulsus*, *L. pubescens*, *L. pyrogalus*, *L. fuliginosus* and *L. violescens* as suspicious or inedible. McIlvaine reports *L. insulsus* and *L. vellereus* as edible, as good as *L. deliciosus*. Murrill's list of forbidden mushrooms is headed by *L. rufus* and includes *L. torminosus*, *L. fuliginosus*, *L. vellereus*, *L. pyrogalus* and *L. theiogalus*—perhaps all condemned only on account of their taste when raw. Fabre's household finds *L. deliciosus* overrated, coarse and difficult to digest.

*Lactarius uvidus* extract was acutely poisonous to guinea pigs, fatal in forty-eight hours, but had no effect on rabbits. No hemolysin or agglutinin. Several authors rank it poisonous; the Boston Mycological Club pronounce it deleterious. With *L. torminosus* and *Clitocybe illudens* it is ranked as a violent gastro-intestinal irritant.

#### THE GENUS RUSSULA

Russula, one of the most difficult genera for reliable specific distinctions, appeals to the mycophagist because of the attractiveness, tenderness and abundance of its species. Members of the Detroit Mycological Club and the Institute of Science have for years eaten all the bright colored and peppery Russulas indiscriminately and believe that *Russula emetica* is a safe fungus to eat in Michigan. Hockauf says of the European *R. emetica* which is so often condemned, that our knowledge is insufficient and that exceptions can justly be taken to reports in the literature. Krapf

was made ill by it (?) before 1800 and its bad name has followed in all subsequent reports which are based on this almost exclusively. Hockauf would take reports of many bad sorts with reserve. McIlvaine is very emphatic that about all Russulas are good, even *R. emetica*, identified by Peck. *R. foetens* smells ill, tastes worse and made Krombholz slightly ill. In 1817 ten deaths in Bohemia were credited to Russula. Murrill<sup>7</sup> includes *R. emetica* in his poisonous list and credits it with cholin, pilztropin and muscarin and puts down *R. foetens*, *R. nitida* and *R. fragilis* as mildly poisonous or suspicious. Warren (Port Huron) says, "I have eaten every kind of Russula I have gathered except *R. foetens* and no one would care to eat that. Never any bad effects. Greatest fault is that they are liable to be wormy. *R. vesca*, *R. virescens*, *R. cyanoxantha*, and *R. alutacea* are permitted in the Munich markets. The "fraglos giftig" *R. emetica* is eaten in the Baltic province Esthonia after parboiling. (Maass.<sup>12</sup>)

Frey<sup>18</sup> says that poisoning by Russula should be classed among the greatest rarities. He reports two fatal cases, studied in the greatest detail and from every angle, with thorough autopsy. The clinical picture was not unlike that of *A. phalloides* intoxication, with gastro-intestinal symptoms dominant. The two boys that died (aged twelve and fourteen) ate the soup which they had prepared, on Sunday evening and Monday morning, were ill Monday night, attended school on Tuesday and became very ill that night, as did the father. They died on Thursday. Postmortem, the liver was not that of *A. phalloides* fatality and the gastro-intestinal hemorrhages and appearances were regarded as characteristic of the irritant action of Russula poison. It is assumed, in these two cases, that an essential change (spoilage) took place in the soup between the first and second meals. An official investigation of the abundant remnants of the fungi ruled out *A. phalloides* and placed the blame on spoiled Russula varieties.

#### THE GENUS MARASMIUS

Marasmius, the family of the internationally famous "fairy ring mushroom," has long been credited with having the poisonous *M. wrens* and the doubtful *peronatus*. McIlvaine would clear both of suspicion. We have no data but we would again warn of the danger of getting *Clitocybe sudorifica* and *Clitocybe morbifera* cooked with *M. oreades*. The latter, moreover, has been found tough, leathery

and entire in the vomited matter after a mushroom feast—illness *ex abusu*, a common form of spurious mushroom poisoning.

#### Conclusions on White-Spored Genera

Of some 50 families of Agarics about 23 are white-spored. Moreover, more than half of all species one finds belong to this section. Though the most dangerous toadstools belong to the Leucosporae, there are so many fine edibles that we do not wish to discard all the white-spored species. If we are to eat fungi at all we must expect to exercise discriminating observation on every specimen intended for the table. In *Amanita*, the edible *A. rubescens* is no harder to distinguish from the dangerous *Amanitas* than are *Lepiota naucina* and many others. Mixed lots of many varieties are a menace and should be used only by the trained student who knows the qualities of each species in the collection. The number of fatalities from fungi gathered by children tells its own story. Cases among students of mycology have all been due to the milder species, and have had the saving grace of adding real discoveries or valuable information to our knowledge. If such cases are duly published a real service and a duty are rendered to science. A synopsis of white-spored species which are definitely deleterious shows: About thirteen white *Amanitas* and a few nearer *A. muscaria* in their physiological action; one each in *Amanitopsis*, *Lepiota* and *Tricholoma*; three *Clitocybes*; at least one (and perhaps a half-dozen) in *Lactarius*; *Russula* uncertain. *Lactarius* and *Russula* are closely related genera, and will require much more investigation, both by eating and by laboratory studies, before the properties of the species will be known. Species closely related botanically are often widely separated toxicologically, and vice versa. This is seen in the contrasting qualities of *Amanita phalloides* versus *A. solitaria* and *A. rubescens*; *A. muscaria* vs. *A. frostiana* and *A. caesarea*; *Lepiota morgani* vs. *L. procera*; *Tricholoma venenatum* vs. *T. terreum* and others; *Clitocybe illudens* vs. *C. multiceps* and others; *Clitocybe dealbata* vs. *C. sudorifica*; *Lactarius torminosus* vs. *L. vellereus* and others.

#### PINK-SPORED AGARICS

The number of genera is small and some mycologists would avoid eating all species of the section. The common favorite *Pluteus cervinus*, or "fawn mushroom," has caused disturbance several times, attended by numbness and tingling in the extremities, mild general discomfort and an urticarial rash. Dr. Whetstone (Minnesota) has a record of the case of poisoning of an Iowa physician, attended by abdominal pain, nausea, and vomiting coming on three hours after eating. Cases like this should make one hesitate to recommend almost any species to the uninitiated.

#### THE GENUS *ENTOLOMA*

All species of *Entoloma* should be avoided by the mycophagist. They are seldom used. Warren says that you cannot cook the raw taste out of them. Vomiting, diarrhea, tenesmus, mental and physical depression are credited to them but no deaths.<sup>22</sup> Six species examined by Ford act identically, producing fatal chronic intoxication in guinea pigs or rabbits—sometimes in both. They vary somewhat in agglutinins and have no hemolysins. (*E. salmoneum*, *E. strictius*, *E. cuspidatum*, *E. nidorosum*, *E. rhodopolium*, *E. sinuatum* or *E. fertile*.) *E. grande* is under suspicion.<sup>17</sup> *E. modestum* and *E. subtruncatum* were negative.<sup>41</sup>

*E. fertile* (*sinuatum*)— $\frac{1}{4}$  oz. nearly killed W. G. Smith (Stevenson, Vol I). It "harbors a virulent poison." The genus may have its own poison, as *Amanita*. According to a recent collection of cases by Sartory<sup>46</sup> in France *E. lividum* is an extremely dangerous fungus, causing severe illness, and occasionally death. He believes that *E. lividum* is nearly as poisonous as some forms of *A. phalloides*. Butignot refers four cases of violent illness to it,<sup>52</sup> though but few specimens were in the mixture eaten. Vomiting and abdominal pain, sweating and a vile diarrhea were the result.

## BROWN OR OCHRE-SPORED AGARICS

These are not usually regarded as poisonous.

THE GENUS *PHOLIOTA*

"I have nothing but praise for the entire genus." (McIlvaine.)

Recently *P. autumnalis* has arisen to claim high rank as a toxic fungus.<sup>39</sup> In 1911 a mother and two children ate heartily of it. The children died. Severe poisoning of three individuals is also reported from Minnesota. Animal tests by Ford and Sherrick<sup>41</sup> on the Minnesota lot were negative on guinea pigs, rabbits and the frog heart, but a New York lot, although negative on blood corpuscles, was acutely poisonous to guinea pigs and rabbits even after heating. Atropin did not neutralize the dilating effect on the heart. Postmortem appearances resembled those of *A. phalloides* and the extracts were quite as poisonous. It should be grouped with the deadly poisonous Agarics, with the nature of the poison unknown.<sup>25</sup> *P. mutabilis* is approved in Munich.

THE GENUS *INOCYBE*

Absolutely negligible and uninviting as food, this genus has likewise recently taken an important rank toxicologically from laboratory studies. The trouble began when Dr. Deming (once Vice-President of the New York Mycological Club) knowingly gathered *Inocybe infida* and mixed it, for cooking, with *Panaeolus papilionaceus* which he knew to be non-poisonous. The chance taken was one in a thousand—but he lost. (See 37, 47 and 48 in Bibliography.) Five people were made ill. Symptoms, which came on soon, were a sense of fullness in the head and a rapid pulse—as if nitroglycerin had been taken. Sweating and warmth, no nausea or prostration; slight confusion, pressure and pain in the lower bowel. Some patients vomited, some had diarrhea. Recovery was complete in a few hours under simple treatment. Conclusions<sup>37, 48</sup> are that *I. infida* contains a poison of the type of muscarin, acting more particularly on the nervous system and similar to the nar-

cotic of *Inocybe infelix*. (See below.) Atropin acts as antidote. The relationship of the toxins of *I. infida* and *I. infelix* to those of *A. muscaria* is not yet clear.

*Inocybe infelix*, one of the most common Inocybes, closely related to the preceding, has not been tested for edibility, nor thus far been reported as toxic to man, but its poison seems definite and powerful. Ford and Sherrick<sup>40</sup> found it to contain a definite poison which resists dessication and boiling. Small doses produced a deep sleep in guinea pigs and rabbits from which they awoke well. A profound acute intoxication and coma, quickly fatal, followed large doses. The intoxication was such as is seen only with *Lactarius torminosus*, a somnolence with retracted head (rabbits), passing off in five hours. The action was not characteristically that of *A. muscaria*, though not inconsistent with muscarin, but that of a narcotic of some sort. Further work is promised. Autopsy on guinea pig showed hemorrhagic spots and perforating gastric ulcer. Generally the examination was negative. *Inocybe decipiens* has likewise no clinical record. Though its agglutinin is destroyed by heat, the heated extract in 2 to 4 cc. doses nevertheless kills guinea pigs acutely—even in 20 minutes—due to dilated heart. Smaller doses bring on lachrymation, salivation and nasal discharge, with labored respirations. These symptoms last a few hours and disappear, but the animals die in a day. Occasionally hemorrhage into the stomach is found postmortem. Dropped into the eye of rabbits, the pupil contracts—resembling *A. muscaria* and muscarin. Boiling the extract does not change its action. This significant record entitles this Inocybe likewise to a place among the more dangerous toadstools.<sup>41, 1</sup>

*Inocybe* sp. agrees in biologic pharmacology very closely with the preceding and the same conclusions are justified in this newest addition to a bad family. Agglutinin powerful, but thermolabile. A muscarin-pilocarpine poison.<sup>50</sup>

THE GENUS *HEBELOMA*

Hebeloma is closely related to Inocybe. The generic distinctions may be learned in order that the mycophagist may reject both genera as inedible. They never will be missed, anyway. Little is known about them. *Hebeloma sinapizans* is suspected (eaten with a gay mixture which included *Amanita jaune citrone*).<sup>51</sup> *Hebeloma fastibile* is related to Inocybe.<sup>17</sup> Kobert would class *I.*

*rimosa* and *H. fastibile* in the muscarin group.<sup>1</sup> Ford<sup>50</sup> reports favorably on *H. crustuliniforme* and another closely related species, even though the former is called "poison pie" in England.

#### Summary of Brown-Spored Genera

In summarizing the brown-spored group we have to take strong exception to the recent idea that none are known to be poisonous. *Pholiota* has some edibles worth while but the importance that *P. autumnalis* (perhaps identical with *P. marginata*) has recently achieved as a dangerous species is an unexpected warning on the family. *Flammula* has fair edibles and is free from suspicion at present. Kauffman's long and careful studies on the glorious genus *Cortinarius* will now enable us to begin to record the qualities of its species with some hope of accuracy. Past experience on its species warrants the statement that they are pretty safe esculents. The remaining brown-spored families should be rejected.

#### PURPLE-BROWN-SPORED AGARICS

*Hypholoma sublateritium* is regarded as poisonous in Europe. It is sometimes bitter, and on this account alone, like many other fungi, has probably been wrongly labeled. Our Club members have occasionally found the "Bricktops" and others inedible, on account of taste. Kobert states that the "Falscher Stockschwamm," *H. fasciculare*, is not edible. Kunkel says it may be poisonous but not very. *H. instratum* and *Psilocybe cernua* both produce acute intoxication in guinea pigs, fatal in three days. Internal and subserous hemorrhages and enlarged glands were found postmortem. (*Morchella esculenta*, the Morel, produces similar findings). Edible properties are not recorded. No fatalities in man have been referred to *Hypholoma* and there are many edible species.

Agaricus or Psalliota—the meadow mushroom family—contains the most famous and most sought edibles, a number of species. Any one who knows of the woods-inhabiting species in the genus would probably know and avoid the deadly *Amanitas* growing among them. Though they do occur, it is very rare for the dangerous white *Amanitas* to get out into the open grassy haunts of *Agaricus campestris* and *Lepiota naucina*. But no one should rely on usual habitat as his safeguard. He should know well the appearances in detail of the plant he may safely use. We know that two of our Michigan fatalities were due to children wandering into the woods and adding *Amanita phalloides* to "the meadows" they had collected, and three other Michigan deaths were caused by mistaking the "Destroying Angel" for *Lepiota naucina*.

## BLACK-SPORED AGARICS

Ford, in 1907, stated that no cases of poisoning have ever resulted from the use of any of the purple-spored or black-spored Agarics. I am of the opinion that a good many cases of the milder type have been caused by both *Coprinus* and *Panaeolus*. It will come as a shock to the lovers of the old reliable "inkies" to find them candidates for the increasing cohort of poisonous fungi. Their reputation has been as fair as their spores are black. That of the "shaggy mane" has been traced back to Pliny.

In "Good Housekeeping" (October, 1910) Dr. Cleghorn tells of ten people in four families accustomed to using ink caps, being made ill on three different dates by *Coprinus comatus*. The appearance was as of one intoxicated. There was failure of muscular coordination, standing being difficult and walking impossible. Drowsiness, loss of emotional control, bloodshot eyes, enlarged pupils, incoherent or inappropriate speech were the symptoms coming on in a few minutes or hours after eating. There was no prostration and heart and lung action were strong and regular. One patient complained of the apparent bending and swaying of the furniture. One had a temporary complete paralysis of the left arm. Practically no food had been eaten but the ink-caps. Prof. John Dearness<sup>58</sup> suggests *Panaeolus campanulatus* as the cause in these cases but in view of the circumstances reported this hardly seems likely. Detroit cases of unpleasant effects—more than an acute indigestion—have been reported from taking beer with a meal of *C. comatus*. I have also known of four cases in which flushed face, bloodshot eyes and rapid and distressing heart action followed the eating of *C. atramentarius*. In two of these cases no alcoholic beverage had been taken. In the other two, only a very small amount to which the individual was accustomed. Further reports concerning the inkies are desirable.

Ford<sup>22</sup> has examined *Panaeolus retirugis* only. It is regarded as edible by all authorities but is similar to *P. papilionaceus* which though edible, has been known to produce a peculiar intoxication. He found no hemolysin or agglutinin. Fatal to guinea pigs and postmortem negative. *P. campanulatus*, classed as poisonous by Murrill, is eaten by McIlvaine. Its bad reputation goes back to

1816 and has not been taken seriously. A Minneapolis report says that two rather delicate ladies ate of it—two tablespoonsful of stew. Drowsiness came on quickly; a sensation of intoxication, dizziness, staggering, trembling, numbness, contraction of the jaw, stricture of the throat, precordial distress, headache with sensation of fullness, face flushed and eyes injected, no nausea. Delusions of sight accompanied insomnia—the patients saw big red automobiles in the room or queer figures on the wall paper. The eyelids in one case were temporarily paralyzed. Mild but irritant diarrhea. In one case the heart was intermittent for a week. Recovery was not prompt. A third and more vigorous patient only tasted the stew. Two hours later she complained of dizziness, ringing in the ears and dry throat.

*P. papilionaceus*, "The Butterfly," has flitted into and out of the questionable list. McIlvaine has seen it produce hilarity and other mild symptoms of intoxication, soon over. Moderate quantities have no effect. Murrill does not rank it as certainly bad. It is a small uncommon Agaric and may therefore be easily spared. In nine years experience in cultivating the gardener's mushroom I have not seen the Coprini or Panaeoli coming on the beds in amount sufficient to warrant the picking of them, but know that *P. subbalteatus* has thus occurred.

## BOLETUS

Though these volumes deal only with the Agarics, or gilled fungi, a paper on mushroom poisoning would not be satisfactory did it fail to include some matter on the Boleti and on Gyromitra, especially since we have some positive data to report.

Ford<sup>22</sup> says that the definitely poisonous Boleti are not many, and that even the toxic, by reason of their bad taste or emetic or purgative action, protect the user from great harm. But few deaths have been traced to Polypores. Among the important esculents are *B. edulis*, *B. scaber* and *B. granulatus*. The majority are edible, but bitter and wormy varieties are common, and others produce vomiting and diarrhea. McIlvaine regards the genus as very safe. On the other hand, a gentleman at Walloon Lake, Michigan, after spending some weeks testing Boleti, said he had not found one variety that did not make him sick! Warren (Port Huron) says, "I never eat them and I tell others to let them alone. There are too many good kinds to bother with wormy Boletus." *B. satanus* and *B. luridus* are everywhere called poisonous, though the toxic principle is little known. Kobert found muscarin in the latter, but conservative Michael says it is edible. *B. clintonianus*, *B. cavipes*, *B. paluster*, *B. chrysenteron* var. *sphagnetum* were all found<sup>22</sup> free from muscarin or definite poisonous action on guinea pigs and rabbits. Variety *sphagnetum* has not been reported edible but *B. chrysenteron* and the other three are approved. McIlvaine, after years of testing by many people, is very positive that both *B. satanus* and *B. luridus* are edible. *Boletus felleus* is free from hemolysins and agglutinins and muscarin, but produces chronic intoxication in rabbits and guinea pigs, fatal in two or three weeks. Extract from the dried plant produced a steady emaciation in rabbits and progressive cachexia in guinea pigs. Probably to be classed as poisonous.<sup>40</sup> Very bitter and inedible.

*B. chromapes*: No hemolysin, agglutinin nor muscarin. Poisonous only to guinea pigs. Decision deferred. Edible (McIlvaine).

*B. affinis* and *ornatipes*: A thermolabile agglutinin destroyed at 150° F. No definite action. Edible (McIlvaine).

*B. bicolor*: An agglutinin; negative on hemolysin and muscarin. Non-toxic. One of the very best esculents (McIlvaine).

*B. separans*: An agglutinin requiring boiling for its destruction, non-toxic. Edible, McIlvaine and Hard.

*B. ravanelli* seems to be safe but not tested by actual use.

*B. roxanae*, similar to *B. separans*. Dietic properties unknown.

*B. miniato-olivaceus* should be regarded with suspicion because of the report on its var. *sensibilis* (below). Ford<sup>40</sup> finds it to contain a heat-resistant agglutinin and to be poisonous to guinea pigs by chronic emaciation. Rabbits were not affected. No evidence of muscarin. (Compare *Clitocybe dealbata* versus *C. sudorifica*.)

*B. pachypus* has a bitter taste and a bed-bug odor.<sup>21</sup> A case of poisoning which Hockauf would refer to cheese, has been credited to it. Murrill adds *B. ferruginatus*, *B. eastwoodiae*, *B. frostii* (edible, Peck), *B. morrissi*, and *B. rubinellus* to the uncertain or suspected. Fabre, in a chapter on insects and mushrooms, (Life of the Fly), records that his peasants eat *B. satanus* and other doubtful species after boiling them in salt water and rinsing.

*Boletus miniato-olivaceus* var. *sensibilis*. It will be seen from the above that our opinions on some species of *Boletus* are much at variance. The following case illustrates how effectively a student of mycology can add to our knowledge by following up thoroughly, and reporting cases of poisoning. Collins<sup>59</sup> records that a certain *Boletus*—found to agree with the erroneously-figured edible *B. subtomentosus* in Palmer's "Mushrooms of America"—was broiled and eaten for breakfast. Three persons ate sparingly and two ate freely. In two hours vomiting and then purging, with collapse calling for brandy and ether subcutaneously in one case, with narrowing or closing of the field of vision, coldness and helplessness, came on. There was no vertigo, headache nor acute pain. The action was mostly that of an irritant. Coffee was the principal stimulant used. One patient did not recover fully for several weeks. Fresh *Boletus* specimens were soon secured from the original spot and again one year later. These were identified by Peck as *B. miniato-olivaceus* var. *sensibilis*.

## GYROMITRA ESCULENTA

This fungus is also known as *Helvella esculenta*, the false Morel, and the Lorchel. Our Michigan species may include *G. brunnea*. Dispute still rages around this fungus and this is characteristic of a species that contains a minor poison or an inconstant one or one that affects only a few individuals and these only

at certain times. *G. esculenta* has a long criminal record in Europe. Nevertheless, it is not everywhere under the ban even there, for its sale, dried or fresh, is permitted in Berlin and Munich though forbidden in Austria. Dried, dusty, wormy (Anobium and Tinea), specimens are sold in the shops; old and inferior fresh ones at reduced prices in the markets. Several American authors say that only old specimens are dangerous. This is not true. The poison is very soluble in hot water and hence parboiling and rinsing may render the mushroom safe. Kobert says that all the Morchellas are safe but that the False Morel furnishes a record of over one hundred and sixty cases of poisoning. Hockauf<sup>60</sup> reports four cases with one fatality (girl of nine years) in April, 1905. Loevegren also has five cases with a fatality, in a girl of five years. Vomiting, colic, weakness, irregular respiration, tonic cramps of voluntary muscles, dilated pupils, jaundice and prolonged unconsciousness were the chief symptoms. Death may occur on the first day or in five days. The active principle has long been known as helvellic acid and has a true blood-dissolving action shown by the hemaglobinuria, icterus, and the pigmentation of the spleen. Nephritis and fatty degeneration of the liver are also found at autopsy. *Gyromitra esculenta* stands alone in producing a true hemolytic set of post-mortem appearances. Frey says that this form of poisoning seems to have become very rare. *Amanita phalloides* does not produce it. Animal tests following Hockauf's cases were negative. Kobert says the fresh extract is very variable. Allen (California) reports *G. esculenta* plentiful there and one of the best edibles, but that it should be let alone. In Michigan it begins to appear on the edge of melting snow banks even as early as mid-March and I have seen a small fall crop in northern Ontario in September. It is common about Port Huron and is eagerly sought and even sold in the markets. Warren and Peck (letters) and Dearness<sup>58</sup> report a number of illnesses and at least one fatality due to *Gyromitra*. In the cases of Dearness the family were made ill after eating of the warmed-over cooking. Coma and death of one adult came on the fourth day. In the Michigan cases there were two groups of nine people each who ate, with two illnesses in each group. Symptoms came on in about six hours. Very violent vomiting and diarrhea, with much weakness and fear were the chief symptoms. Heart action was good. The acute illness lasted thirty-six hours, inability to take food three days, and recovery required about five days. "No other ill effects except that they don't want any more Gyromi-

tras." The fungi were fresh and prepared as often before—and since. Warren says, "I know they may contain some kind of poison that affects some people at certain times."

Ford<sup>41</sup> has examined specimens of *G. esculenta* from Massachusetts. He found them entirely negative and harmless in every method of testing on guinea pigs, rabbits and the frog's heart.

#### CONCLUSIONS AND SUMMARY

Because of the growth of popular interest in the study of the mushrooms and toadstools, both as a "fad" and as a scientific past-time, and because of the influx of foreigners accustomed to use fungi for food, the subject of mushroom poisoning is assuming increasing importance in America.

The white Amanita or Death-cup (*Amanita phalloides*) and its few closely related species are responsible for at least nine-tenths of all fatal cases of mushroom poisoning. In illness caused by this fungus the mortality runs very high. Symptoms are six to ten hours in coming on. Suffering is extreme, and death often does not occur until a week or more has elapsed, though the course is quicker in children. There is no antidotal treatment. Clinical course and post-mortem findings are characteristic.

The white Amanita group contains a toxin found in no other fungi. It is a poison which causes profound degenerative changes in the internal organs and in the cells of the central nervous system.

The white Amanitas are easily recognized and avoided.

In all cases of mushroom intoxication, it is the duty of physicians and of friends to make every effort to learn whether or not *Amanita phalloides* has been eaten or whether some less dangerous toadstool is causing the sickness. Such determination is important both in determining treatment and especially prognosis.

The yellow Amanita or "Fly Fungus," *Amanita muscaria*, is second in importance. It is much less poisonous. It produces characteristic symptoms unlike those of *A. phalloides*, coming on in three hours or less, showing prompt disturbance of the nerve centers, and a disease of shorter course and lower mortality. The degenerative changes seen in *A. phalloides* intoxications, do not occur. If fatal results occur, this outcome may be expected early.

A large part of the disturbances produced by *Amanita muscaria* are due to the muscarin constituent of the fungus. This poison

can be counteracted by the drug atropin. Hence *A. muscaria* intoxication is somewhat amenable to treatment.

Physicians should be able to distinguish between these two forms of poisoning. Symptoms are not always definite enough to be relied upon, and specimens of the mushroom which has been eaten should be identified. Local or near-by botanical centers are always glad to be of service in such problems.

A number of minor poisonous species of mushrooms (about twenty) produce symptoms, when eaten, *which resemble the action of muscarin*. These species have not been given the importance and attention which they deserve. They usually also have an emetic action which prevents fatal consequences. Some of them are violent gastro-intestinal irritants and may thus add gravity to the illness. Deaths from them are almost unknown in healthy adults. Determination of the species is of great practical and scientific importance, since these minor cases are more numerous than is generally supposed.

Physicians and students of mycology should report cases in medical or botanical journals. Cases reported in the newspapers should be investigated.

Species closely related botanically may differ very widely in poisonous quality, though this is not usual. No variety should be eaten until its specific name has been determined by competent authority. If its edible qualities are not known or are in dispute, testing should proceed cautiously. The eating of mixed lots is to be condemned unless the user positively knows the reputation of each species to be good. Parboiling is a partial safeguard. None but clean, fresh specimens should be used, and these should be thoroughly cooked and indulged in only in moderation. Most mushrooms are not easily digested. Warmed-over portions are occasionally found to have developed toxic principles.

Good and abundant edible varieties are common, and the lover of fungi need take no chances. He can easily acquire a personal edible list, and can add new species to it as his knowledge and experience grow. Scientific mycology should precede mycophagy and increase one's pleasure in eating fungi.

## BIBLIOGRAPHY

### (A) BOOKS CONSULTED

- ATKINSON, GEO. F., *Mushrooms, Edible, Poisonous, etc.*, 1900. Third ed. 1911. Very useful to the beginner; beautifully illustrated with photographs.
- BARLA, J. B., *Les Champignons de la prov. de Nice*, 1859. With 48 colored plates of the common edible and poisonous species, with full descriptions. *Flore Mycologique illustree Les Champignons des Alper-Maritimes*, 1889-92. With 64 large, colored plates of species of *Amanita*, *Lepiota*, *Armillaria*, *Clitocybe* and *Tricholoma*.
- BERKELEY, REV. M. J., *Outlines of British Fungology*. 1860. A manual, with 24 colored plates of many species.
- BOUDIER, E., *Icones Mycologicae ou Iconograph, d. Champignons de France*, 1904. A series of the most accurate and most beautiful colored plates of fungi yet published; a relatively small number belonging to the Agarics. It is very expensive.
- BRESADOLA, ABBE J., *Fungi Tridentina*, 2 vol., 1881-1900. Illustrated with 217 colored plates; a critical study of the rarer or confused species; invaluable to the specialist. *I. Funghi mangerecci e velenosi*, 1906. Illustrated with 120 colored plates and descriptions in Italian of the more common edible and poisonous species.
- CLEMENTS, F. C., *Genera of Fungi*. This is a useful book of keys to the genera, largely taken from Saccardo and adapted to those who cannot read Latin. It contains no descriptions.
- COOKE, M. C., *Handbook of British Fungi*, 2 vol., 1871. With full descriptions of species. Not as helpful as either Masee or Stevenson. Illustrations of British Fungi. Eight volumes of colored plates illustrating practically all the Agarics of Great Britain; 1,198 plates in all. Necessary to the specialist.
- COOKE ET QUELET. *Clavis Synoptica Hymenomycetum Europaeorum*. A useful little book with brief Latin descriptions of the Hymenomycetes of Europe, arranged in the form of a pocket manual.
- CONSTANTIN ET DUFOUR. *Nouvelle Flore des Champignons*. A useful French book for beginners arranged in the form of keys, illustrated by 4,265, diagramatic line drawings. Mostly Hymenomycetes which are known in France.
- "ENGLER AND PRANTL", *Die Naturliche Pflanzen-familien*. The volume containing the Basidiomycetes; one of many volumes covering an outline of the whole plant kingdom to the genera; profusely illustrated with woodcuts. Indispensable to the mycologist.
- FARLOW, W. G., *Bibliographical Index of North American Fungi*, 1905. Giving the references to the literature for each species; only Vol. I, part I, has appeared to date.
- FRIES, ELIAS MAGNUS. *Systema Mycologicum*, 3 vol. Vol. I. contains the Basidiomycetes, 1821. This volume and its date was made the starting point for the nomenclature of the Agarics by the International Congress meeting at Brussels, 1910. *Epicrisis Systematis Mycologici*, 1836-38. An extension of Vol. I of *Syst. Myc.* with changes and additional species. *Monographia*, 1857-63. Being a series of monographs of the genera of the Agaricaceae; nearly all the species mentioned in this work are illustrated by colored drawings which are now deposited in the Royal Museum at Stockholm. It includes only Swedish species, but the descriptions are

- followed by commentaries and are invaluable to the specialist. *Hymenomyces Europaei*, 1874. This is one of the most important and fundamental works on the Basidiomycetes ever published. It represents the ripe experience and best judgment of the master. *Sveriges ätliga och giftiga Svam par.*, 1862-69. A volume of large plates with colored figures of the edible and poisonous species of Scandinavia. *Icones*, 1867-1884. Two volumes of 200 large colored plates of the Basidiomycetes, selected from the large number of drawings deposited at the Museum at Stockholm.
- GIBSON, W. H., *Our Edible Toadstools and Mushrooms*, 1895. (Ed. new, 1903.) A book written in a delightful literary vein, containing popular descriptions and discussions of some forty mushrooms; well illustrated with colored drawings.
- GILLET, C. C., *Les Hymenomyces*, 1874. A manual with full descriptions and artificial keys to all the Basidiomycetes growing in France. Very valuable. *Les Champignons de France*. A set of 712 colored plates accompanying the preceding manual.
- HARD, M. E., *Mushrooms Edible and Otherwise*, 1908. A popular book for the beginner, profusely illustrated with photographs.
- HERBST, WM., *Fungal Flora, of the Lehigh Valley, Pa.*, 1899. This book contains descriptions and 25 plates.
- HUSSEY, *Illustrations of British Mycology*, 1847-55.
- KARSTEN, P. A., *Mycologia Fennica*, 4 vol. Vol. 3—Basidiomycetes. A manual of the flora of Finland.
- LINDBLAD, M. A., *Svambok*, 1902, a popular manual in Swedish.
- MARSHALL, NINA L., *The Mushroom Book*, 1905. "A popular guide to the identification and study of our commoner fungi, with special emphasis on the edible varieties." With photographs, many colored.
- MASSE, GEO., *British Fungous Flora*, 4 vol. 1892-95. A useful manual for the English reader, but deals only with British forms. *European Fungus Flora; Agaricaceae*, 1902. A compilation of all European species of Agarics with only the essential characters given in brief description. *Text book of Fungi*, 1906. A general discussion of matters of interest and importance to the student of fungi, mostly of others than Agarics.
- McILVAINE, CHARLES, *One Thousand American Fungi*, 1900. New ed., 1912. A valuable compilation of descriptions of fungi formerly inaccessible to many. The last edition is well illustrated. An additional value of this book is in the data concerning the edibility of mushrooms as tested by the author.
- MICHAEL, EDMUND, *Führer für Pilzefreunde*. 3 Vol., 1903-05. Containing in all 307 beautifully colored plates of mushrooms, unusually true to nature cheap and very helpful to the beginner.
- MURRILL, W. A., *Agaricaceae*. *North Amer. Flora*, Vol. 9 et al., 1910.
- OUDEMANS, C. A. J. A., *Revision des Champignons dans les Pays-Bas*, 1893. Vol. I contains synopses and descriptions of Hymenomyces of Holland; in French.
- PATOUILLARD, N., *Tabulae Analyticae Fungorum*, 1883-89. A critical study of many species, with numerous colored figures; invaluable to the specialist. In French. *Les Hymenomyces d'Europe*, 1887. Deals with the general anatomy and classification of the higher fungi.
- PERSOON, C. H., *Synopsis Methodica Fungorum*, 1801. One of the best early efforts to place the classification of fungi on a scientific basis; in some groups it is considered by some as more fundamental than the works of Fries.
- QUELET, L., *Euchiridion Fungorum*, 1886. A manual. in Latin, of the species found in Middle Europe and France. *Les Champignons de Jura et des Vosges*, 1872. Classification and descriptions of some 700 species found in the mountainous regions of eastern France.
- RICKEN, A., *Die Blätterpilze (Agaricaceae)*. A manual, in German, of the Flora of Germany, with excellent modern descriptions, and profusely illustrated by colored plates. Appearing in parts; 1910-16. One of the most useful and complete works of its kind, giving the microscopic characters of most species. Invaluable to the student.
- SACCARDO, P. A., *Sylloge Fungorum*. A work of huge scope including descriptions, in Latin, of all known species of the world, 1882—. It comprises to the present 22 volumes; Vol. V, and parts of later volumes, deal with Agarics.
- SCHROETER, J., *Die Pilze Schlesiens*, 1889. A manual with full descriptions of the Agarics indigenous to Silesia.
- SCHWEINITZ, L. de, *Synopsis Fungorum*.
- SECRETAN, L., *Monographie Suisse*, 1833, 3 vol. Descriptions and classification of fungi growing in Switzerland.
- SMITH, W. G., *Synopsis of the British Basidiomycetes*. Brief diagnoses and sketches of all Agarics of Great Britain; spore-size not given.
- STEVENSON, REV. JOHN, *British Fungi*, 2 vol., 1886. A manual of the British Hymenomyces, whose descriptions are largely those of Fries. Very useful to the student who reads only English. The same descriptions of many species are also found in McIlvaine.
- SWANTON, E. W., *Fungi and How to Know Them*, 1909. A book for the beginner dealing in a popular way with all fungi, as well as Agarics; profusely illustrated with colored figures and cuts in black and white. The beginner will find it useful.
- TAYLOR, THOMAS, *Student Hand-book of Mushrooms of America*, 1897. A series of fine pamphlets bound in book-form, with 14 colored plates dealing with a few common edible and poisonous mushrooms.
- UNDERWOOD, L. M., *Moulds, Mildews and Mushrooms*, 1899. A small, compact text-book on the classification of fungi, with keys to families and genera, good bibliographies and explanatory notes on families; invaluable to the student who reads only English.
- VITTADINI, *Descriz. dei Funghi Mangerecci e velenosi d'Italia*. Milan, 1835. A carefully written work of the mushrooms of Italy.
- WINTER, GEO., *Die Pilze Deutschlands, Oesterreichs und der Schweiz*. A manual of the flora of the Basidiomycetes of Germany, Austria and Switzerland, with full descriptions; it is Vol. I, Part I, of Rabenhorst's *Kryptogamen Flora*.

## (B) JOURNALS OF MYCOLOGY CONSULTED

(For special papers in other journals, see (C) and text)

- ANNALES MYCOLOGICI. Entirely devoted to fungi; articles are printed in various languages.
- BULLETIN DE LA SOCIETE MYCOLOGIQUE DE FRANCE. Entirely devoted to fungi; in French. Started in 1885.
- BULLETIN OF THE TORREY BOTANICAL CLUB. Containing some papers on fungi, especially the descriptions of new species by Dr. Peck of such as were sent to him from outside of New York State; an American Journal.
- GREVILLEA. Published in England from 1892 to 1894. Devoted to Cryptogamic Botany and its literature; contains descriptions of the species in Cooke's Illustrations. Includes many references to American species.
- HEDWIGIA. A German journal dealing with Cryptogamic Botany; 1852 to the present.
- JOURNAL OF MYCOLOGY. An American journal devoted to fungi, superseded by *Mycologia*. 1885-1908.
- MYCOLOGIA. Started in 1909; devoted to mycology; published by the New York Botanical Garden.
- MYCOLOGICAL BULLETIN. A popular journal for the beginner which had a brief existence during four and one-half volumes.
- MYCOLOGICAL NOTES. Published privately by C. G. Lloyd since 1898. Devoted to critical notes and synopses of the higher fungi.
- REVUE MYCOLOGIQUE. A French journal for mycologists, started in 1879.
- RHODORA. Published by the New England Botanical Club. Contains articles on fungi of New England.
- TORREYA. Published by the Torrey Botanical Club, New York City. Containing occasional papers on Agarics, beginning 1901.

## (C) STATE REPORTS, FLORAS, KEYS, LISTS, ETC.

- ATKINSON, G. F., and STONEMAN, BERTHA, Key to the Genera of Hymenomyces. Ithaca, N. Y.
- BRADY, W. A., A Partial List of the Fungi of Wisconsin: Geol. of Wis., Vol. I, p. 396, 1883. (Includes 218 species of Basidiomycetes.)
- BURLINGHAM, GERTRUDE S., Some Lactarii from Wyndham Co., Vt.: Bull. Torr. Bot. Club, Vol. 34, 1907. (Contains a synopsis, list and new species.)
- BURT, E. A., Key to the Genera of the Basidiomycetes of Vermont: Contrib. to the Bot. of Vt., VI, 1899.
- CLEMENTS, F. C., Minnesota Mushrooms: Geol. and Nat. Hist. Surv. of Minn. Minn. Plant Studies, IV, 1910. (Published in book form for use in schools with descriptions and keys; profusely illustrated with photographs.)
- CURTIS, M. A., Catalogue of the plants of North Carolina, Raleigh, 1867.
- EARLE, F. S., (collaborator), Plant Life of Alabama: U. S. Dept. of Agr., Nat. Herbarium, Vol. 6, p. 150, 1901, (including a full list of the fungi reported from Alabama). The Genera of North American Gill Fungi: Bull. New York Botanical Garden, Vol. 5, p. 373, 1908. (Keys and descriptions of genera, classified according to the rule of priority.) Keys to species of various genera: Torrey, Vol. 2, (Russula, Hygrophorus, etc.).
- FARLOW, W. G., List of Fungi found in the vicinity of Boston: Bul. Bussey Institution, I, p. 430, 1876; II, p. 224, 1878.
- FROST, C. C., Further enumeration of New England Fungi: Proceed. Soc. Nat. Hist. Boston, Vol. XII, p. 77, 1869.
- GLATFELTER, N. M., Preliminary list of higher Fungi, etc.: Acad. Sci. of St. Louis, Vol. XVI, No. 4, p. 33, 1906.
- HESS AND VANDAERT, Basidiomycetes of Central Iowa: Proceed. Iowa Acad. Sci., Vol. 7, p. 133.
- KAUFFMAN, C. H., Michigan Fungi: Seventh Ann. Rep. Mich. Acad. Sci., p. 64, 1905. Unreported Michigan Fungi: Eighth Ann. Rep. Mich. Acad. Sci., p. 26, 1906. Ibid: Ninth Ann. Rep., p. 83, 1907. Ibid: Tenth Ann. Rep., p. 63, 1908. Ibid: Eleventh Ann. Rep., p. 55, 1909. Ibid: Twelfth Ann. Rep., p. 99, 1910. Ibid: Thirteenth Ann. Rep., p. 215, 1911.
- LONGYEAR, B. O., A preliminary list of the Saprophytic Fleshy Fungi known to occur in Michigan: Fourth Rep. Mich. Acad. Sci., p. 113, 1904. List of saprophytic Fungi: Rep. of Sec. to State Board Agr., 1898.
- MACBRIDE, T. H., The saprophytic fungi of Eastern Iowa: Bull. Tab. Nat. Hist. State Univ. of Iowa, Vol. I, p. 30, 1888. Ibid: p. 181, 1890. Ibid: p. 196, 1890.
- MOFFATT, W. S., The higher fungi of the Chicago Region: Nat. Hist. Surv. of Chicago Acad. Sci., Bull. VII, part 1, p. 1, 1909. Descriptions and 24 photo plates; including 360 species of Basidiomycetes.
- McKAY, A. H., Fungi of Nova Scotia; a preliminary list: Proceed. and Trans. Nova Scot. Inst. Sci., Vol. XI, part 1, p. 125, 1904. Includes 284 species of Basidiomycetes.
- MILLSPAUGH and NUTTALL, Flora of West Virginia: Field Columb. Mus., Bot. Ser. I, p. 60, 1896.
- MORGAN, A. P., The Mycologic Flora of the Miami Valley, Ohio: Journ. Cinn. Soc. Nat. Hist., Vol. 6, p. 54-81, 97-115, 173-199. With descriptions and six plates.
- PECK, CHAS. H., New York State Mus. Reports, published in the Annual Reports of the New York State Museum of Nat. Hist., and also separately as Reports 27-54 consecutively, afterwards as Bulletins until 1912 when Dr. Peck retired. The reports of Peck as State Botanist from 1870-1912, include plants of all groups, but the greater portion is devoted to lists of fungi found in New York State, to descriptions of new species, and to synopses of many of the genera of Agarics. See also bibliography of each genus. Fungi of Maryland: New York State Mus. Rep. 44, 1891. A list prepared by Mary E. Banning of Baltimore and including about 100 Agarics.
- REDDICK, D., A preliminary list of the Hymenomyces of Indiana: Dept. of Geol. and Nat. Resources of Ind., Rep. 32, p. 1193, 1907. A list including 88 Agarics.

- SPRAGUE, C. J., Contributions to New England Mycology: Proceed. Bost. Soc. Nat. Hist., Vol. 5, p. 325, 1856; Vol. 6, p. 315, 1859.
- STOVER, W. G., The Agaricaceae of Ohio: Proceed. Ohio State Acad. Sci., Vol. V, part IX, p. 462, 1912. Keys to the genera and species; comprising 540 species without descriptions.
- UNDERWOOD, L. M., Report of the Botanical Division of the Indiana State Biological Survey: Proceed. Ind. Acad. Sci., p. 13, 1894; p. 144, 1895; p. 171, 1896.
- VAN HOOK, J. M., Indiana Fungi: Proceed. Ind. Acad. Sci., p. 205, 1910; p. 347, 1911. Includes list of 347 Basidiomycetes.
- WEBBER, H. J., Catalogue of the Flora of Nebraska: Rep. Neb. State Board Agric., p. 37, 1890.
- WHITE, E. A., Hymeniales of Connecticut: Conn. State Geol. and Nat. Hist. Surv. Bull. 3, 1905; 15, 1910. Contain lists, keys to species and a few descriptions; illustrated with photographs of 28 and 39 plates respectively. Reference to 433 species of Basidiomycetes.

## (D) MISCELLANEOUS BOOKS, PAMPHLETS, PAPERS, ETC.

- ARTHUR, J. C., Three edible toadstools: Purdue Univ. Agr. Exp. Sta., Bull. No. 98, 1904.
- ATKINSON, G. F., and SHORE, R. B., Mushroom growing for amateurs: Cornell Univ. Agr. Exp. Sta., Bull. 227, 1905. Mushrooms: Ibid, Bull. 163, 1899.
- BADHAM, C. D., A Treatise on esculent funguses, 1847. (A British book, with 21 colored plates.)
- BATES, Mushrooms—poisonous Amanitas, harmless Agarics (directions for cultivation): Amer. Florist, Vol. II, 1896.
- BEACH, S. A., Mushrooms as a Greenhouse Crop: N. Y. State Agr. Exp. Sta., Bull. 88, 1895.
- BIFFEN, R. H., On the biology of *Collybia velutipes*: Jour. Linn. Soc., Vol. 34, p. 147.
- BRITZELMAYR, M., Materialien zur Beschreibung der Hymenomyceten: Bot. Centralblatt, Vol. 53-54, p. 33, 65 and 97, 1893; Vol. 62-63, p. 273, 1895; Vol. 71-72, p. 49, 1897. (Brief critical notes on European Agarics, with spore size.) Revision der Diagnosen zu den von Britzelmayr, aufgestellten Hymen. Arten., Bot. Centralblatt, Vol. 75-76, p. 129, 169, 203, 1898; Vol. 77-78, p. 356, 395, 433, 1899; Vol. 79-80, p. 57, 116, 1899. (Full description of the species described as new.)
- BULLER, A. H. K., Researches on Fungi, etc., 1909, (a valuable study of spore expulsion in Coprinus, etc.).
- BULLETIN, Boston Mycolog. Club, 1897, \* \* \* (small pamphlets giving directions to amateurs, with brief descriptions of the species of some genera).
- BURECK, A., Notes on enemies of mushrooms and on experiments with Remedies: U. S. Dept. of Agr., Entomol., 32-35.
- BURT, E. A., Collecting and preparing fleshy fungi for the herbarium: Bot. Gaz., Vol. 25, p. 172, 1898.
- COOKE, M. C., British edible fungi and how to distinguish and how to cook them. Edible and poisonous mushrooms; what to eat and what to avoid. (With 48 colored plates.)
- CORDIER, F. S., Les Champignons, Paris, 1876, 4th ed. with 60 colored plates.
- DAVIS, SIMON, Fleshy fungi of Stow, Mass.: Rhodora, Vol. 13, p. 57, 1911; Vol. 16, p. 45, 1914 (critical notes).
- DUGGAR, B. M., The principles of mushroom growing and mushroom spawn-making: U. S. Dept. Agr., Bureau of Plant Ind., Bull. 85, 1905. (The best modern discussion of the subject.) The cultivation of mushrooms: Farmer's Bull., No. 204, U. S. Dept. Agr., 1904.
- FALCONER, WM., How to grow mushrooms: Farmer's Bull., U. S. Dept. Agr., No. 53, 1897.
- FAYOD, V. M., Prodrome d'une Histoire Naturelle des Agricenes: Ann. des Nat. Botanique, 7 series, Vol. 9, p. 181.

- FARLOW, W. G., Some edible and poisonous fungi: U. S. Dept. Agr., Bull. No. 15, 1898. Notes for mushroom eaters: Reprinted from Garden and Forest Magazine.
- GAUTIER, L. M., Champignons, Paris, 1884.
- GOWDY, J. K., Mushroom culture in France: U. S. Consular Rep. 66, p. 259, 537, 1902.
- HARKNESS and MOORE, Catalogue of the Pacific Coast Fungi: Calif. Acad. Sci., 1881.
- HASTINGS, SOMERVILLE, Toadstools at home: Gowan's Nature Series, No. 7, London, 1908. (60 photographic reproductions of fungi in their natural surroundings; excellent little primer.)
- HAY, WM. D., Fungi hunter's guide, London, 1887.
- HENDERSON, L. P., Mushrooms or toadstools, a natural food-product: Idaho Agr. Exp. Sta., Bull. 27, p. 27.
- HESS, ALICE W., A few common fleshy fungi of Ames, Iowa: Iowa Agr. Exp. Sta., Bull. 61, p. 148.
- HOGG, ROBERT and JOHNSON, G. W., A selection of the edible funguses of Great Britain, London.
- HYAMS, C., Edible fungi of North Carolina: N. Car. Exp. Sta., Bull. 177, p. 27.
- JONES, C. H., Composition of mushrooms: Vermont Agr. Exp. Sta. Bull., p. 196, 1903.
- KAUFFMAN, C. H., Mushrooms: Nature Study Review, Vol. 8, p. 1.
- LANGWORTHY, C. F., Mushrooms as Food: Plant World, Vol. 8, p. 134, 1899.
- LONGNECKER, A. M., Mushrooms: Plant World, Vol. 5, p. 213.
- LONGYEAR, B. O., Remarks concerning saprophytic fungi, etc.: Mich. Acad. Sci., Rep. 1, p. 97, 1897. Notes on Michigan Saprophytic Fungi: Mich. Acad. Sci., Rep. 3, 1901.
- MARCEAU, L., Flore des champignons superior du Department de Saone-et-Loire, 1898. (A manual with extensive keys.)
- MANGET, C., Tableaux synoptiques des champignons comestible et venenaux, Paris, 1903.
- MAY, W. J., Mushroom culture for amateurs, 1897.
- MENDEL, L. B., Chemical composition and nutritive value of some edible American Fungi: Amer. Jour. Phys. 1, p. 225.
- MOYEN, J., Les Champignons, Paris, 1889.
- PALMER, JULIUS A., Mushrooms of America, Boston, 1885. About Mushrooms, Boston, 1894.
- PECK, C. H., Descriptions of new species of fungi: Bull. Soc. Nat. Sci. of Buffalo, 1. p. 41, 1873. New species of fungi, (see Bot. Gaz). New species of fungi, (see Torr. Bot. Club Bull., Vol. 22-38). Mushrooms and their uses: Pub. by Cambridge Bot. Supply Co., 1897. The smooth Agaric: Country Gentleman, 1886, p. 833. Edible and poisonous fungi: N. Y. State Mus. Rep. 48, p. 203, 26 Plates, 1895. Report of the State Botanist on the edible fungi of New York State: N. Y. State Mus. Memoir, No. 4, p. 113, 24 Plates, 1900. Lists of all species described and named by Dr. Peck to 1908, see N. Y. State Mus. Bull. 131, p. 59, 1909. List of figures and illustrations of mushrooms published by Dr. Peck, see N. Y. State Mus. Bull. 139, p. 78, 1910.
- RIDGWAY, ROBERT, Color standards and color nomenclature, Washington, D. C. 1912. (A series of 1115 named colors, invaluable for the future standardizing of the names of mushroom colors and tints.)
- ROBINSON, W., Mushroom culture, London, 1870. (A full account of the older methods.)
- SHEAR, C. L., Some common autumnal species of edible Fungi: Asa Gray Bull. 7, p. 93, 1899.
- SICARD, G., Histoire naturelle des Champignons, Paris, 1883 (with colored plates).
- SMITH, W. G., Clavis Agaricorum, London, 1870 (Descriptions of families and genera with keys.)
- STURGES, W. C., Edible and poisonous fungi: Rep. Sec. Connect. Board of Agr., 1895.

- SYDOW, P., Taschenbuch der wichtigeren essbaren u. giftigen Pilze Deutschlands u. der Schweiz. (Pocket size, with colored figures.)
- TAYLOR, EMMA L., Guide to the mushrooms. (Small primer, 200 pages and 66 plates.)
- TORREY, JR., J., Raising mushrooms in a cellar: Rhodora, Vol. 3, p. 57.
- TRELEASE, WM., Edible and poisonous mushrooms: Missouri State Hort. Soc. Rep., 1900, p. 224.
- UNDERWOOD, L. N., Edible Fungi—A great vegetable food in Indiana: Trans. Ind. Hort. Soc., 1893, p. 62. Edible Fungi: Alabama Agr. Exp. Sta. Bull. 73, p. 337. Suggestions to collectors of fleshy fungi: Ala. Agr. Exp. Sta. Bull. 80, p. 271.
- VIERMEYER, T. J., The Mycogone disease of mushrooms and its control: U. S. Dep. Agr. Bull. No. 127, 1915.
- WEBSTER, HOLLIS, See Rhodora, Vols. 1, 2, 4.
- WEEMS AND HESS, A study of the food value of some of the edible fungi of Ames, Iowa: Proceed. Ia. Soc. Prom. Agr. Sci., 1902, p. 105.
- WILLIAMS, E. M., Common edible and poisonous Amanitas: Asa Gray Bull. Vol. 6, p. 80.
- WINDISCH, E., Mushroom culture on an extensive scale, 1897.
- WINDOR, D. K., The Mushrooms of Canada, 1871 (a little primer).
- ZEGA, A., Composition of some edible Fungi: Chem. Zeitung, Vol. 26, p. 10, 1902. Mushrooms as food: U. S. Dept. Agr., Farmer's Bull. 79. Analyses of mushrooms: N. Y. State Agr. Exp. Sta. Rep. 1894, p. 134. Preliminary list of the edible fungi of Ohio: Ohio State Univ. Bull. 4, p. 30.

## (E) ON MUSHROOM POISONING

BY O. E. FISCHER, M. D.

1. Ford & Clark—A Consideration of the properties of poisonous fungi: Mycologia, July, 1914, Vol. VI, No. 4.
2. Bagnall—Mushroom poisoning, with atypical cases: Boston Med. & Surg. Jour., July 16, 1914.
3. Ford—A clin. study of mushroom intoxication: Johns Hopkins Hosp. Bull., Vol. 18, 1907.
4. Gillot—Etude med. sur. l'empoisonnement par les Champignons, Lyon, 1900.
5. Plowright—Brit. Med. Jour., Sept. 9, 1905, Vol. II, p. 541.
6. Pfromm—Med. Bull. Phila., Nov., 1905, p. 401.
7. Murrill—Poisonous mushrooms: Mycologia, Nov., 1910, p. 255.
8. Mycol. Bull., Vol. VI, No. 86, Feb., 1908, quoted from O. E. Jennings in Jour. of Mycol., Sept., 1907.
9. Barr—Letter. Kindness of Minneapolis Mycol. Club.
10. Schuerer—Kasuistischer Beitrag zur Kenntniss der Pilzvergiftungen: Deut. Med. Woch. Schr. XXXVIII, No. 12, Mar. 21, 1912.
11. Hockauf—Zur Kritik der Pilzvergiftungen: Wien Klin. Woch. Schr., 1904, p. 731.
12. Maass—Berl. Klin. Woch., June 26, 1905, p. 814.
13. Ford—Jour. Infect. Dis., Vol. 5, 1908, p. 116.
14. Jour. A. M. A.—Vol. LX, 1913, p. 1154.
15. Abel & Ford—Arch. f. Exper. Path. u. Pharm. Schmiedeberg Festschrift, 1908, p. 8.
16. Frey—Zeit. Sch. Klin. Med., 1912, p. 455.
17. Same as No. 7.
18. Ford—Jour. Inf. Dis., 1906, Vol. III, p. 191.
19. Abel & Ford—Poisons of *Amanita phalloides*: Jour. Biol. Chem., Vol. II, 1907, p. 273.
20. Schlesinger & Ford—Chem. Prop. of *A. phal.*: same as above, Vol. III, 1907, p. 279.
21. Ford & Prouty—Note on Amanita-toxin: Jour. Pharm. & Exp. Ther., Vol. I, 1907, p. 389.
22. Ford—Distribution of hemolysins, agglutinins & poisons in fungi: Jour. Pharm. & Exper. Ther., Vol. II, 1910, p. 285.

23. Coville—Observations on recent cases of mushroom poisoning in the District of Columbia: *Circ. 13, Div. of Botany, U. S. Dept. Agriculture, 1898.*
24. Prentiss—*Phil. Med. Jour.*, 1898, II, p. 607.
25. Ford & Clark—*Mycologia*, July, 1914, Vol. VI, No. 4.
26. Harmsen—*Zur Toxicologie des Fliegenschwammes: Archiv. f. Pathol. u. Pharmacol. Schmiedeberg, 1903.*
- 26a. *Ibid* —. Bd. 50, p. 361, 1903.
27. Michael—*Führer für Pilzfreunde. Zwickau, Sa., 1903.*
28. Kennan—*Tent Life in Siberia, New Ed., 1910, p. 198.*
29. Sumstine—*Mycol. Bull.*, Vol. IV, 56, Apr., 1906.
30. Tappeiner—*Munch. Med., Wochenschr.*, Feb. 12, 1895, p. 133, J. 42, No. 7.
31. Ford & Brush—*Note on properties of fungi from France: Jour. Pharm. & Exp. Ther.*, Vol. 6, No. 2, Nov., 1914.
32. Same—*Action of Amanitas upon the frog's heart. Same as above.*
33. Blount—*A personal experience with mushroom poisoning, Med. Record, Nov. 23, 1901, p. 815.*
34. Chestnut—*Asa Gray Bull.*, Oct., 1900.
35. Webster—*Rhodora, Jan., 1915, Vol. 17, p. 30.*
36. Farlow—*Poisoning by Ag. illudens: Rhodora, March, 1899, p. 43.*
37. Clark & Smith—*Toxicological studies on Clit. illudens & Inocybe infida: Mycologia, 1913, p. 224.*
38. McIlvaine & Macadam—*One thousand American fungi. Revised Ed.*
39. Peck—*Report of State Botanist, 1911, N. Y., Bull. 157.*
40. Ford & Sherrick—*Jour. Pharm. & Exp. Ther.*, 1911, II, p. 549.
41. Same—*On Clit. sudorifica, etc., etc. Same as above, Vol. IV, 1913, p. 321.*
42. Peck—*As above, Bull. 150, p. 44.*
43. Rabe—*Zeitscher. f. exp. Phar. u. Ther.*, Sept., 1911, p. 352.
44. *Bull. Soc. Mycol. de France, 1906, p. 22.*
45. Goldman—*Agaricus torminosus: Wien. Klin. Wochenschr.*, Mar. 21, 1901.
46. Sartory—*Les empois. par les Cham., (Etc., 1912) Paris, 1912.*
47. Murrill—*A new poisonous mushroom: Mycologia, 1909, p. 211.*
48. Clark & Kantor—*Toxicological experiments with higher fungi: Mycologia, 1911, p. 175.*
50. Ford—*Further observations on fungi: Journ. Pharm. & Exp. Ther.*, Vol. VI, No. 2, Nov., 1914.
51. *Bull. Soc. Mycol. de France, 1912, p. 28.*
52. Same—*1909, p. 25 and 1906, p. 22.*
53. *Jour. A. M. A., LXI, p. 1388.*
54. Peck *Report, 1895, p. 212. Same, 1901, p. 961.*
55. Clark & Smith—*Mycologia, 1913, p. 224.*
56. Von Wedekind—*Med. Record, Dec. 25, 1897, p. 919.*
57. Step—*Toadstools & mushrooms of the (British) country side.*
58. Dearness—*The personal factor in mushroom poisoning: Mycologia, Vol. 3, 1911, p. 75.*
59. Collins—*Rhodora, Vol. I, No. 2, Feb., 1899, p. 21.*
60. Hockauf—*Wien. Klin. Wochenschr, 1905, p. 1058.*

## (F) MONOGRAPHS, ETC., ARRANGED BY GENERA

AGARICUS. (See Psalliota).

AMANITA. Peck, N. Y. State Mus. Rep. 33, p. 38, 1880. Lloyd, compilation of the Volvae of U. S., p. 2, 1898. Boudier, *Obs. sur.—especies d'Amanitas, Bull. de la Soc. Mycol. de France, Vol. 18, p. 251, 1902.* Quelet et Bataille, *Flora Monographie des Amanites et des Lepiotas, Paris, 1902.* Beardslee, *Notes on the Amanitas of the Southern Appalachians, Lloyd's Mycological Notes, Vol. I. See also: Fries, Saccardo, Gillet, Bresadola, Barla, Stevenson, Masee, Atkinson, McIlvaine, Morgan, Ricken, etc.*

AMANITOPSIS. Peck, N. Y. State Mus. Rep. 33, p. 38 (under Amanita). Lloyd, compilation of Volvae of U. S., p. 8, 1898. Quelet et Bataille, *Flora monographie des Amanites et des Lepiotas, p. 42, 1902, (under Amanita). See also: Fries, Saccardo, etc. (under Amanita in the older books).*

- ANNULARIA. (See Chamaeota.)
- ARMILLARIA. Peck, N. Y. State Mus. Rep. 43 (Bot. ed.), p. 44, 1899. See also: McIlvaine, Morgan, Fries, Saccardo, Stevenson, Bresadola, Gillet, Stevenson, Barla, etc.
- BOLBITIUS. See Fries, Gillet, Ricken, etc.
- CANTHERELLUS. Peck, N. Y. State Mus. Bull. 2 (Bot. ed.), p. 34, 1887. See also: Saccardo, Fries, Gillet, Stevenson, Ricken, Morgan, Moffatt, McIlvaine, etc.
- CLAUDOPUS. Peck, N. Y. State Mus. Rep. 39, p. 67, 1886. See also: Saccardo, Fries, Gillet, Masee, Stevenson, Ricken, Patouillard, etc.
- CLITOCYBE. Peck, N. Y. State Mus. Bull. 157, (Bot. ed.), p. 59, 1912. See also: Barla, McIlvaine, Lloyd, Saccardo, Fries, Bresadola, Gillet, Stevenson, Masee, Morgan, Moffatt, etc.
- CLITOPILUS. Peck, N. Y. State Mus. Rep. 42, (Bot. ed.), p. 39, 1880. See also: Saccardo, Fries, Gillet, Stevenson, Masee, McIlvaine, Morgan, etc.
- COLLYBIA. Peck, N. Y. State Mus. Rep. 49, p. 32 (Bot. ed.), 1896, Lloyd, *Mycolog. Notes, Vol. I, p. 33, 1900.* See also: Saccardo, Fries, Gillet, Stevenson, Masee, Ricken, McIlvaine, Moffatt, etc.
- COPRINUS. Masee, *Revision of the Genus Coprinus: Ann. Bot., Vol. 10, p. 123, 1896.* Daniels, E. A., *Coprinus: Boston Mycolog. Club Bull. No. 15 and 16.* See also: Saccardo, Fries, Stevenson, Patouillard, Masee, Gillet, Ricken, Atkinson, Peck's Reports, Moffatt, etc.
- CORTINARIUS. Fries, *Monographia Cortinariorum Sueciae, Upsala, 1851.* Bataille, *Flore Monographique des Cortinaires d'Europe, Besancon, 1911.* Kauffman, *The Genus Cortinarius, Bull. Torr. Bot. Club, Vol. 32, pp. 301, 1905; The Genus Cortinarius with Key to Species, Jour. Mycol., Vol. 13, p. 32, 1907; Cortinarius as a Mycorrhiza-producing Fungus, Bot. Gaz., Vol. 42, p. 208, 1906.* Earle, *A Key to the North American Species of Cortinarius, Torrey, Vol. 2, pp. 169 and 180.* Britzelmayer, M., *Revision der Diagnosen zu den von Britzelmayer aufgestellten Hymenomyceten-arten, Bot. Centralblatt, Vol. 73, p. 129, 169 and 203, 1898.* Maire, R., *Notes critiques \* \* \* Bul. de la Soc. Myc. de France, Vol. 27, p. 423, 1911; Notes critiques \* \* \* ibid, Vol. 26, p. 176, 1910.* See also: Saccardo, Gillet, Quelet, Stevenson, Masee, Ricken, Boudier, Karsten, Michael, Peck (N. Y. State Reports) Hard, etc.
- CREPIDOTUS. Peck, N. Y. State Mus. Rep. 39, p. 69, 1886. See also: Saccardo, Fries, Gillet, Patouillard, Ricken, Masee, Stevenson, Moffatt, etc.
- DECONICA. See Psilocybe.
- ECCILIA. See: Saccardo, Fries, Ricken, Moffatt, Peck's Reports, Atkinson (Preliminary Notes on Some New Species: *Jour. Mycol., Vol. 8, p. 113, 1902*), etc.
- ENTOLOMA. Peck, N. Y. State Mus. Bull. 131, p. 47, 1909. See also: Saccardo, Fries, Gillet, Stevenson, Ricken, McIlvaine, Hard, Atkinson, Moffatt, Morgan, Davis, etc.
- FLAMMULA. Peck, N. Y. Mus. Rep. 50, p. 133, 1897. See also: Saccardo, Fries, Gillet, Stevenson, Ricken, Atkinson, Moffat, etc.
- GALERA. Peck, N. Y. State Mus. Rep. 46, p. 61, (Bot. ed.) 1893. See also: Saccardo, Fries, Gillet, Ricken, Stevenson, etc.
- GOMPHIDIUS. See: Saccardo, Fries, Ricken, Peck's Reports, etc.
- HEBELOMA. Peck, N. Y. State Mus. Bull. 139, p. 67 (Bot. ed.), 1910. See also: Saccardo, Fries, Ricken, Stevenson, Masee, etc.
- HYGROPHORUS. Peck, N. Y. State Mus. Bull. 116, p. 45, (Bot. ed.), 1907. Bataille, *Flora Monographique des Hygrophores, Besancon, 1910.* See also: Saccardo, Fries, Gillet, Schroeter, Ricken, Stevenson, Masee, Hard, McIlvaine, Atkinson, Moffat, Davis, etc.

- HYPHOLOMA.** Peck, N. Y. State Mus. Bull. 150, p. 73 (Bot. ed.), 1911. Harper, Species of Hypholoma \* \* \* \* Trans. Wis. Acad. Sci. Arts. and Let., Vol. 17, p. 1142, 1914. Morgan, North American Species of Agaricaceae; Hypholoma, Jour. Myc., Vol. 14, p. 27, 1908. See also: Saccardo, Fries, Ricken, Atkinson, Moffatt, etc.
- INOCYBE.** Peck, N. Y. State Mus. Bull. 139, p. 48 (Bot. ed.), 1910. Masee, Monograph of the Genus Inocybe, Ann. Bot., Vol. 18, p. 459, 1904. Bresadola, Fungi Tridentini. See also: Patouillard, Saccardo, Gillet, Ricken, Hard, Moffatt, Davis, etc.
- LACTARIUS.** Peck, N. Y. State Mus. Rep. 38, p. 111, 1885. Burlingham, Lactariae of the United States, Mem. Torr. Bot. Club, Vol. 14, p. 1, 1908. See also: Saccardo, Fries, Gillet, Bresadola, Ricken, Masee, Stevenson, McIlvaine, Hard, Atkinson, Moffatt, Morgan, Michael, etc.
- LENTINUS.** Peck, N. Y. State Mus. Bull. 131, p. 42, 1909. See also: Saccardo, Fries, Stevenson, Bresadola, Patouillard, McIlvaine, Hard, Moffatt, etc.
- LEPIOTA.** Peck, N. Y. State Mus. Rep. 35, p. 150, 1884. Quelet et Bataille, Flore Monographique des Amanites et des Lépiotes, Paris, 1902. Morgan, North American Species of Lepiota: Jour. Myc., Vol. 12, p. 154, 195 and 242, 1906; Vol. 13, p. 1, 1907. Lloyd, Myc. Notes, Vol. I, p. 4, 1898. Beardslee, The Lepiotes of Sweden: Jour. Myc., Vol. 13, p. 26, 1907. See also: Saccardo, Fries, Gillet, Barla, Bresadola, Ricken, Schroeter, Stevenson, Masee, McIlvaine, Hard, Atkinson, Moffatt, Michael, etc.
- LEPTONIA.** See: Saccardo, Ricken, Stevenson, etc.
- MARASMIUS.** Morgan, North American Species of Marasmius: Jour. Mycol., Vol. II, p. 201, 1905. Pennington, N. Y. State Mus. Bull., 1915. See also: Saccardo, Fries, Stevenson, Masee, Patouillard, Ricken, Hard, Moffatt, etc.
- MYCENA.** Peck, N. Y. State Cab. Rep. 23, p. 80 (Bot. ed.) 1872. See also: Saccardo, Fries, Masee, Stevenson, Gillet, Ricken, Bresadola, Patouillard, Atkinson, Hard, Moffatt, etc.
- NAUCORIA.** Peck, N. Y. State Cab. Rep. 23, p. 91 (Bot. ed.), 1872. See also: Saccardo, Fries, Ricken, Moffatt, etc.
- NOLANEA.** See: Saccardo, Fries, Gillet, Ricken, Stevenson, etc.
- OMPHALIA.** Peck, N. Y. State Mus. Rep. 45, p. 32, (Bot. ed.), 1893. See also: Saccardo, Fries, Stevenson, Ricken, Patouillard, Hard, Moffatt, etc.
- PANOEOLUS.** Morgan, North American Species of Agaricaceae; Panoeolus: Jour. Myc., Vol. 13, p. 59, 1907. See also: Saccardo, Fries, Ricken, Atkinson, Hard, Peck's Reports, etc.
- PANUS.** Forster, Agarics of United States Panus: Jour. Mycol., Vol. 4, p. 21, 1888. See also: Saccardo, Fries, Stevenson, McIlvaine, Hard, etc.
- PAXILLUS.** See Atkinson, Hard, Saccardo, etc.
- PHOLIOTA.** Peck, N. Y. State Mus. Bull. 122, p. 141, 1908. Harper, Species of Pholiota \* \* \* \* Trans. Wis. Acad. Sci. Arts and Let., Vol. 17, p. 470. See also: Saccardo, Fries, Gillet, Ricken, Hard, Atkinson, McIlvaine, Moffatt, etc.
- PLEUROTUS.** Peck, N. Y. State Mus. Rep. 39, p. 58, 1886. Webster, Bull. Boston Myc. Club, No. 8, 1898. See also: Saccardo, Fries, Gillet, Stevenson, Masee, Ricken, Patouillard, Atkinson, McIlvaine, Hard, Moffatt, etc.
- PLUTEOLUS.** Peck, N. Y. State Mus. Rep. 46, p. 58, 1893. See also: Saccardo.
- PLUTEUS.** Peck, N. Y. State Mus. Rep. 38, p. 133, 1885. Lloyd, Myc. Notes, Vol. I, p. 12. See also: Saccardo, Fries, Stevenson, Ricken, Patouillard, Atkinson, Hard, etc.
- PSALLIOTA.** Peck, N. Y. State Mus. Rep. 36, p. 41, 1884. Smith, Rhodora, Vol. 1, p. 161. Lloyd, Myc. Notes, Vol. 1, p. 25, 1899. (These

- are given as Agaricus.) See also: (under Agaricus), Saccardo, Fries, Gillet, Ricken, Stevenson, Atkinson, McIlvaine, Hard, Moffatt, etc.
- PSATHYRA.** Peck, N. Y. State Mus. Bull. 150, p. 84 (Bot. ed.), 1911. Morgan, Jour. Myc., Vol. 13, p. 147, 1907. See also: Saccardo, Fries, Ricken, etc.
- PSATHYRELLA.** Peck, N. Y. State Cab. Rep. 23, p. 102 (Bot. ed.), 1872. Morgan, Jour. Myc., Vol. 13, p. 54, 1907. See also: Saccardo, Fries, Ricken, etc.
- PSILOCYBE.** Peck, N. Y. State Mus. Bull. 157, p. 94 (Bot. ed.), 1912. Morgan, Jour. Myc., Vol. 13, p. 246, 1907. See also: Saccardo, Fries, Ricken, etc.
- RUSSULA.** Peck, N. Y. State Mus. Bull. 116, p. 67, 1907. Kauffman, C. H., Mich. Acad. Sci. Rep. 11, p. 57, 1909; Ibid, Rep. 13, p. 220, 1911. Kaufmann, F., Die bei Elbing gefundenen essbaren u. giftigen Täublinge (Russula): Schriften der Naturforschenden Ges. in Danzig, N. F., Vol. 8, p. 21, 1894. Maire, Les bases de la classification dans le genre Russula, Bull. de la Soc. Myc. de France, Vol. 26, p. 49, 1910. Notes critiques \* \* \* \* Ibid., p. 167. Peltereaux, Etudes et Observations sur les Russules: Bull. de la Soc. Myc. de France, Vol. 24, p. 95, 1908. Bataille, Flore monographique des Asterospores, 1907. Barbier, Encore les Russules: Bull. de la Soc. Myc. de France, Vol. 24, p. 230, 1908. Romell, Lindblads Svampbok, p. 62. Hymenomycetes of Lapland: Arkiv för Botanik, Vol. II, No. 3, p. 5, footnotes. Denniston, Trans. Wis. Acad. Sci. Arts & Let., Vol. 15, p. 71. Macadam, Jour. Mycol., Vol. 5, p. 58 and 135, 1889. See also: Saccardo, Fries, Gillet, Quelet, Masee, Stevenson, Ricken, McIlvaine, Burlingham, etc.
- STROPHARIA.** Harper, Species \* \* \* \* in the Region of the Great Lakes: Trans. Wis. Acad. Sci. Arts and Let., Vol. 17, p. 1014, 1913. Morgan, North American Species of Agaricaceae: Jour. Myc., Vol. 14, p. 57, 1908. See also: Peck's Reports, Saccardo, Fries, Gillet, Stevenson, Ricken, Hard, Atkinson, etc.
- TRICHOLOMA.** Peck, N. Y. State Mus. Rep. 44, p. 150, 1891. See also: Saccardo, Fries, Gillet, Barla, Ricken, Stevenson, Masee, McIlvaine, Hard, Moffatt, Morgan, etc.
- TUBARIA.** See: Saccardo, Fries, Stevenson, etc.
- VOLVARIA.** Lloyd, Volvae of United States, p. 10, 1898. See also: Saccardo, Fries, Patouillard, Ricken, Peck's Reports, Hard, etc.

AUTHORITIES AND ABBREVIATIONS

The binomial botanical name of each plant is followed by an abbreviation, e. g. Fr., which refers to the person who named the species. According to the rule established by the International Congress at Brussels in 1910, no names are to be considered valid in the case of the Agarics, earlier than those published in the *Systema Mycologica* of Fries, 1821-1832. In case the name was used for the same species by someone before Fries, reference may be made to it thus: Fr. (ex. Pers.).

AUTHORS OF AGARICS

A. & S.	Albertini and Schweinitz.
Atk.	Atkinson, Geo. F. (U. S.).
B. & C.	Berkeley and Curtis.
B. & Br.	Berkeley and Broome.
Bann	Banning, Mary E. (U. S.).
Barla	Barla, J. B. (France).
Batsch	Batsch, Augustus (German).
Beards	*Beardslee, H. C. (U. S.).
Berk	Berkeley, Rev. J. M. (England).
Bolt	Bolton, James (Canada).
Boud	*Boudier, E. (France).
Bosc	Bosc, Louis (U. S.).
Bres	*Bresadola, Abbe J. (Austria).
Britz	*Britzelmayr, Max (Germany).
Bull	Bulliard, Pierre (France).
Burl	*Burlingham, Gertrude S. (U. S.).
Cke	Cooke, M. C. (England).
Clem	*Clements, F. C. (U. S.).
Curt	Curtis, Rev. M. A. (U. S.).
D. C.	DeCandolle, Augustin P. (Switzerland).
E. & E.	Ellis and Everhart.
Earle	*Earle, F. S. (U. S.).
Ell	Ellis, J. B. (U. S.).
Fr	Fries, Elias Magnus (Sweden).
Frost	Frost, Charles C. (U. S.).
Gill	Gillet, C. C. (France).
Henn	Hennings, Paul (Germany).
Herbst	Herbst, Wm. (U. S.).
Kalchb	Kalchbreuner, Karoly (Hungary).
Karst	*Karsten, P. A. (Finland).
Kauff	*Kauffman, C. H. (U. S.).
Kromb	Krombholz (Germany).
L. or Linn	Linneaus, Carl von (Sweden).
Lev	Leveille, Joseph H. (France).
Lindb	Lindblad, M. A. (Sweden).
Lloyd	*Lloyd, C. G. (U. S.).
Longyear	*Longyear, B. O. (U. S.).
Maire	*Maire, René (France).

Mass	Massee, Geo. (England).
Mont	Montagne, Camille (France).
Morg	Morgan, A. P. (U. S.).
Murr	*Murrill, Wm. E. (U. S.).
Pat	*Patouillard, N. (France).
Pers	Persoon, Christian Hendrick (Europe).
Pk	Peck, Charles Horton (U. S.).
Q. & S.	Quelet and Schulzer.
Quel	Quelet, L. (France).
Rav	Ravenel, W. H. (U. S.).
Ricken	*Ricken, Rev. Adelbert (Germany).
Rom	*Romell, Lars (Sweden).
Roze	Roze, Ernest (France).
Sacc	*Saccardo, P. A. (Italy).
Schaeff	Schaeffer, Jacobi C. (Germany).
Schroet	Schroeter, Julius (Germany).
Schw	Schweinitz, Rev. Louis David de (U. S.).
Schulz	Schulzer, von Muggenberg (Europe).
Scop	Scopoli, Giovanni A. (Italy).
Sec	Secretan (Switzerland).
Smith	Smith, W. G. (England).
Sow	Sowerby, James (England).
Vahl	Vahl, Martin (Norway).
Vitt	Vittadini, Carlo (Italy).

Those names which are starred are living mycologists.

MISCELLANEOUS ABBREVIATIONS

Acad	Academy.
Bot	Botanical.
Bull	Bulletin; a publication.
Cab	Cabinet.
Cm	Centimetre.
Fig	Figure, referring to an illustration.
Gaz	Gazette.
Hist	History.
Jour	Journal.
Mem	Memoir, a publication.
Micr	Micron, one thousandth of a millimeter.
Mm	Millimetre.
Mus	Museum.
Myc	Mycology, Mycological.
Nat	Natural, Nature.
No	Number of a figure or plate.
Op	Opposite (page).
p	Page.
Pl	Plate; referring to illustration.
Rep	Report; a publication.
Ridg	Ridgway's Color Standards, 1912.
Sci	Science.
Soc	Society.
Sp. nov.	New Species; described for first time by the writer.
Var	Variety.

## GLOSSARY

- ABERRANT, differing from a certain species, genus, etc, in some respects, but not easily placed in another species, genus, etc.
- ABNORMAL, (of a specimen), not properly developed.
- ABORTIVE, (of a fruit-body or its parts), not perfect or entirely lacking.
- ABRUPT, (of a stem), terminating suddenly.
- ABRUPTLY-BULBOUS, (of the bulb of a stem), not rounded above.
- ACICULAR, (of a stem or cystidia), bristle-shaped, very slender.
- ACICULATE, same as acicular.
- ACRID, (of the taste of a mushroom or its juices), biting on the tongue.
- ACUMINATE, (of cystidia, or the ends of a lamella), gradually narrowed to a point.
- ACUTE, (of cystidia or the edge of the gills), pointed; less than a right-angle; sharp-edged; not prolonged.
- ADNATE, (of gills), see Fig. 1, 2; also (of the pellicle, scales, etc.), not capable of being peeled off or easily detached.
- ADNEXED, (of gills), see Fig. 1, 4, narrowly attached to the stem.
- AFFINITY, (of a species, genus, etc.), closely related by natural characters.
- AERUGINOSE, (color), verdigris-green.
- AERUGINOUS, same as aeruginose.
- AGGLUTINATE, (of fibrils, hairs, etc.), as if, glued together in tufts.
- AGGREGATE, crowded close together.
- ALLANTOID, (of spores), sausage-shaped.
- ALLIACEOUS, (odor), like onions or garlic.
- ALUTACEOUS, (color), light leather colored; isabelline; pale tan.
- ALVEOLATE, (of the surface of pileus or stem), deeply pitted.
- AMBIGUOUS, (of a species, genus, etc.), doubtful as to its place in classification.
- AMYGDALINE, (odor or taste), like that of peach or cherry stones, cherry-bark, etc.
- ANALOGOUS, similar in form, structure or appearance, but not necessarily related to.
- ANASTOMOSING, (of gills, ridges, wrinkles, etc.), connecting crosswise, so as to form angular areas or pits bounded by the connecting gills, etc.

- ANGULAR, (of spores), not regular in outline, not rounded; (of scales or pileus), when formed by cracking of cuticle, etc.
- ANNULATE, (of stem), bearing an annulus.
- ANNULAR, (of remains of veil on stem), resembling a ring.
- ANNULUS, the encircling band or curtain on the stem, resulting from the loosening of the inner veil from the margin of the pileus. See Fig. 2, 5.
- ANOMALOUS, deviating from the general rule.
- ANTERIOR, (of gills), the end of the gills at the margin of the pileus; in front.
- APICAL, (of stem), the portion near the pileus; referring to the apex.
- APICULUS, (of spores), the short, often sharp papilla at one end of a spore, by which it was attached to the sterigma.
- APICULATE, provided with an apiculus.
- APPENDICULATE, (of margin of pileus), hung with fragments of the veil.
- APPLANATE, (of pileus), flattened out or horizontally expanded.
- APPRESSED, (of scales, fibrils, hairs, etc.), closely flattened down; same as adpressed.
- APPROXIMATE, (of gills), free from but approaching the stem; closely; not remote.
- ARACHNOID, (of the partial veil), cobweb-like.
- ARCUATE, (of gills or margin of pileus), curved like a bow.
- ARCUATE-DECURRENT, (of gills), extending down the stem.
- AREOLATE, (of surface of pileus or stem), arranged in little areas.
- ARGILLACEOUS, (color), clay color, resembling ochraceous-cinnamon-brown.
- ARID, (of gills), dry, somewhat parchment-like.
- AROMATIC, (odor), of an agreeable aroma, reminding of drugs.
- ASCENDING, (of gills), in the case of a conical-shaped or unexpanded pileus.
- ASTRINGENT, (taste), causing more or less contraction or "pucker" of mouth membranes.
- ATOMATE, (surface of pileus or stem), covered with minute, shining, point-like particles.
- ATTENUATE, (of stem), gradually narrowed and thinner.
- AURANTIACOUS, (color), of an orange color.
- AUREOUS, (color), golden-yellow, reddish-yellow.
- AXIS, (of stem), the central, interior portion.
- AZONATE, (of surface of pileus), not zoned.
- BASAL, (of stem), at the lower end.

- BASIDIOMYCETES, see page 26.
- BASIDIUM, (of gills), one of the large cells which collectively compose the hymenium and which bear each four spores.
- BEADED, (of gills), applied to the row of drops exuding from the edge of gills.
- BEHIND, (of gills), toward the stem.
- BI—, of two, or twice.
- BIBULOUS, (of surface of pileus), capable of absorbing moisture.
- BIFURCATE, (of gills), forking by two's.
- BISTRE, (color), blackish-brown.
- BIOLOGY, the science of living organisms.
- BIOLOGICAL, concerning the life of plants or animals.
- BLOOM, (of pileus or stem), a minutely velvety surface.
- BROAD, (of gills), a relative term, opposed to narrow; determined by experience.
- BUFF, (color), pale creamy-gray.
- BULBOUS, (of stem), enlarged at base. (See also "abruptly-bulbous", clavate-bulbous and round-bulbous.)
- BULBILLATE, (of stem), provided with a small or obscure bulb.
- BULLATE, (of pileus), with a rounded knob.
- BYSSOID, (of mycelium), the condition when fine filaments spread from the base of the stem or fruit-body over the substratum.
- CAESIOUS, (color), pale bluish-gray.
- CAESPITOSE, aggregated in tufts but not grown together.
- CAMPANULATE, (of pileus), bell-shaped or similar.
- CANALICULATE, (of stem), furrowed or fluted.
- CANDIDOUS, (color), shining-white.
- CANESCENT, (surface), covered with hoary down.
- CAP, the pileus.
- CAPILLARY, (of stem), hair-like.
- CAPITATE, (of cystidia), with a minute knob at the tapering apex.
- CARBONACEOUS, (of tissue), of the texture of charcoal.
- CARINATE, (of spores), furnished with a keel, boat-shaped.
- CARNEOUS, (of trama), fleshy.
- CARTILAGINOUS, (of stem, cortex or cuticle), tough-brittle, breaking with a snap.
- CAULICOLOUS, growing on herbaceous stems.
- CELL, (of fungi), the living, protoplasmic units into which the mycelium and hyphae are divided.
- CERACEOUS, waxy.
- CEREBROSE, (of surface of pileus), convoluted like a brain.
- CHLAMYDOSPORES, (see secondary spores), thick-walled spores developed from hyphae but not on basidia.

CINEREOUS, (color), ashy-gray.  
 CINNABAR, (color), vermilion, red.  
 CINNAMON, (color), cinnamon-brown.  
 CIRCUMSCISSILE, (of volva). See page 593.  
 CITRINE, lemon-yellow.  
 CLAVATE, (of stem), thickened toward base, like a club; (of basidia and cystidia), thickened at apex, club-shaped.  
 CLAVATE-BULBOUS, (of stem), with a bulb which tapers gradually upwards.  
 CLAY-COLOR, argillaceous.  
 CLOSE, (of gills), halfway between crowded and subdistant; a relative term.  
 COBWEBBY, (of veil), composed of threads fine as those of a cobweb.  
 COCHLEATE, (of pileus), twisted like a shell.  
 COERULEOUS, (color), sky-blue.  
 COHERENT, (of stems), grown together.  
 COMPRESSED, (of stem), flattened lengthwise.  
 CONCAVE, (of pileus), round-depressed like a bowl.  
 CONCENTRIC, (of zonation, etc.), rings or zones within one another in a series.  
 CONCHATE, (of pileus), resembling an oyster shell in shape.  
 CONCOLOR, (of gills and stem), when of the same color as the pileus.  
 CONCOLOROUS, same as concolor.  
 CONFLUENT, (of flesh of stem), continuous with trama of pileus and of similar texture.  
 CONGLOBATE, (of base of stems), collected into a fleshy mass.  
 CONIDIUM, (see secondary spores), thin-walled spores developed on mycelium or on the hyphae of the fruit-body.  
 CONIDIA, plural of conidium.  
 CONIDIAL, relating to conidia.  
 CONIDIOPHORE, the specialized hypha bearing a conidium.  
 CONIFER, mostly evergreen trees bearing cones.  
 CONIFEROUS, said of forests or wood of conifer trees.  
 CONNATE, (of stems), grown together.  
 CONNIVENT, (of margin of pileus), converging on the stem.  
 CONSISTENCY, the firmness, density or solidity of the tissues which compose the parts of the fruit-body.  
 CONTEXT, the trama.  
 CONTINUOUS, (of stem), same as confluent.  
 CONVEX, (of pileus), regularly rounded, broadly obtuse, etc.

CONVEXO-PLANE, (of pileus), changing from convex when younger to flat when expanded.  
 CONVEX-EXPANDED, (of pileus), changing from convex and tending towards plane; the margin often remaining decurved.  
 CONVERGENT, (of trama of gills), in section the hyphae are seen to turn inwards to a median line.  
 CONVOLUTE, same as cerebrose.  
 CORIACEOUS, of a leathery texture.  
 CORNEOUS, of a horny texture.  
 CORRUGATE, (surface), coarsely wrinkled.  
 CORTEX, (of stem), the outer, denser rind.  
 CORTICAL, referring to the cortex.  
 CORTICATE, possessing a cortex.  
 CORTINA, the inner or partial veil in some genera of Agarics, the structure of which is cobwebby.  
 CORTINATE, provided with a cortina, or (of stem) covered by the threads of the cortina.  
 COSTATE, (of gills, etc.), veined or ribbed.  
 COTTONY, (of surface), covered by a soft cotton-like substance.  
 CRENATE, (of edge of gills or margin of pileus), scalloped.  
 CRENULATE, very finely crenate.  
 CRETACEOUS, (of color or consistency), like chalk.  
 CRISPED, (of gills), finely wavy.  
 CROWDED, (of gills), almost touching one another.  
 CRUCIATE, (of spores), having the general form of a cross.  
 CRYPTOGAMS, the group of plants which reproduce by spores and which include the fungi.  
 CRYPTOGAMIC, relating to cryptogams.  
 CUCULLATE, (of pileus), shape of a "high hat."  
 CUNEATE, (of pileus), wedge-shape.  
 CUSPIDATE, (of pileus or cystidia), tipped with a prominent sharp protuberance.  
 CUTICLE, (of pileus or stem), a differentiated thin layer of hyphae on the surface; same as pellicle.  
 CYATHIFORM, (of pileus), cup-shaped or bowl-shaped, flaring above.  
 CYLINDRICAL, (of stem or spores), of the same diameter throughout its length.  
 CYSTIDIUM, (of hymenium of gills), mixed with the basidia and usually projecting beyond them; large sterile cells.  
 CYSTIDIA, plural of cystidium.  
 DECORTICATED, of dead wood destitute of the bark.

DECUMBENT, (of stem), with the lower end lying against the substratum.

DECURRENT, (of gills), descending on the stem, see Fig. 1, (3).

DECURVED, (of margin of pileus), bent down.

DEBRIS, the mixture of fallen leaves, twigs, wood, etc., covering a forest floor.

DEFLEXED, same as decurved.

DELIQUESCENT, (of gills), absorbing water and dissolving at maturity.

DENTATE, (of gills), toothed on the edge.

DENTICULATE, (of gills), finely dentate.

DENUDED, (of pileus and stem), naked or glabrous by removal of the scales, flocci, etc.

DEPAUPERATE, undeveloped because of lack of favorable conditions.

DEPRESSED, (of pileus), central portion lower than margin.

DETERMINATE, having a fixed, definite limit.

DETERMINATION, assigning a plant to its correct place in the classification.

DIAGNOSIS, a distinctive description of a plant.

DIAPHANOUS, transparent or nearly so.

DICHOTOMOUS, (of gills), repeatedly forking in pairs.

DIFFERENTIATED, applied to portions or tissues of different character, all derived from a homogeneous tissue.

DIFFORMED, irregular in form.

DILATED, (of stem), enlarged.

DILUTE, (of color), reduced in strength.

DIMIDIATE, (of pileus), semicircular in outline, (of gills), that reach only half-way to stem.

DISK, (of pileus), the central portion of the surface.

DISCOID, (of pileus), with a noticeably marked, flattened disk.

DISCRETE, (of veil, scales, etc.), separate, not grown fast to and continuous with the surface.

DISTANT, (of gills), set far apart, especially toward the margin of the pileus; a relative term.

DIVERGENT, (of trama of gills), in section, the hyphae are seen to turn outwards from a median line.

DORSAL, (of pileus), the upper, back side.

DOWNY, (of pileus and stem), composed of fine hairiness.

DRY, not viscid nor hygrophanous.

EBENEIOUS, (color), black as ebony.

EBURNEOUS, (color), white like ivory.

ECCENTRIC, (of stem), not attached in the center.

ECHINATE, (of scales, etc.), sharply pointed spines.

ECHINULATE, (of spores, etc.), with minute and finely pointed spines.

ELEVATED, (of pileus), raised up at the margin.

ELLIPTICAL, (of spores, young pileus, gills, etc.), longer than broad, usually more than twice as long as broad and curved in outline.

ELLIPTIC, ELLIPSOID and ELLIPSOIDAL, similar to elliptical.

EMARGINATE, (of gills), notched near the stem. See Fig. 1, (6).

ENTIRE, (of gills), edge not toothed, etc.

EPIDERMIS, see cuticle.

EPISPORE, the outer wall of a spore.

EPIPHYTAL, growing on leaves.

EQUAL, (of stem), of uniform diameter; (of gills), alike in length.

ERODED, (of gills), edge as if gnawed.

EROSE, same as eroded.

ESCULENT, edible, can be eaten.

EVANESCENT, (of veil, annulus, scales, etc), but slightly developed and soon disappearing.

EVEN, (of surface of pileus, stem, spores), without striations, elevations, depressions or unevennesses of any kind. Compare glabrous and smooth.

EXCENTRIC, see eccentric.

EXOTIC, foreign, not native.

EXPANDED, (of pileus), the opening out of the cap while maturing or ageing.

EXSICCATI, dried specimens kept in herbaria, often in sets.

"FAIRY RINGS," mushrooms appearing in circles. See page 4.

FAMILY, a term in classification, each family includes related genera; the scientific ending of a family name is *aceae*.

FALCATE, (of spores), sickle-shaped.

FARINACEOUS, (odor and taste), like fresh meal; (of pileus and stem), covered by mealy particles.

FARINOSE, like farinaceous.

FASCIATED, (of stems, pilei, etc.), grown together so that tissues are intimately continuous.

FASCICULATE, (of fibrils, scales, stems, etc.), crowded in bundles.

FERRUGINOUS, (color), rusty-red.

FIBRILLOSE, (of surface of cap and stem), provided with fibrils or clusters of small fibres composed of hyphae.

FIBROUS, (of flesh of stem), composed of toughish, string-like tissue.

FILAMENT, a thread, applied to the separate threads of the mycelium.

FILAMENTOUS, composed of filaments.

FILIFORM, (of stem), slender as a thread.

FIMBRIATE, (of gills), with the edge minutely fringed, due to presence of cystidia or sterile cells.

FISTULOSE, (of stem), tubular.

FLABELLIFORM, (of pileus), fan-shaped.

FLACCID, flabby; soft and limber; without firmness or elasticity.

FLARING, (of volva or annulus), spreading away from stem at upper margin.

FLAVESCENT, (color), becoming yellowish.

FLAVUS, (color), of Saccardo's Color Key; a light cadmium-yellow.

FLESH, the trama of the mushroom, especially of the pileus and gills.

FLESHY, of rather soft consistency, putrescent; as opposed to leathery, corky, woody, membranous, etc., referring to the consistency of the trama of most of the Agarics.

FLEXUOUS, (of stem), bent in an undulate manner.

FLOCCI, (of pileus or stem), small points or tufts resembling cotton.

FLOCCOSE, (of pileus or stem), provided with cottony substance on the surface.

FLOCCULOSE, finely floccose.

FLOCCULOSE-CRENULATE, (of gills), edge with minute flocculose decoration.

FOETID, (odor), ill-smelling, nauseating.

FRIABLE, easily crumbled or breaking into powder.

FREE, (of gills), not attached to the stem at any time.

FRONDOSE, said of a forest or the wood of broad-leaved trees.

FRONT, (of gills), the end toward the margin of the pileus; anterior.

FRUIT-BODY, the term applied to the mushroom as opposed to the mycelium.

FRUCTIFICATION, the fruit-body.

FUGACIOUS, disappearing early or quickly or (of color) fading soon.

FULIGINOUS, (color), smoky, sooty.

FULVESCANT, (color), becoming fulvous.

FULVOUS, (color), of Saccardo's Color Key; reddish-cinnamon-brown.

FUNGUS, applied to the individuals of a group of plants which lack the green chlorophyll and hence subsist on other plants,

plant-remains or animals; they vegetate in the form of mycelium, and their fruit-bodies are also composed of mycelioid tissue.

FUNGI, the plural of fungus.

FURCATE, (of gills), forked.

FURFURACEOUS, (of pileus or stem), covered with bran-like particles; scurfy.

FUSCESENT, (color), becoming fuscous.

FUSCOUS, (color), a smoky drab; see Ridgway's Color Standards (1912). The term has been used in this report in a wider sense, including paler shades with more brown in them.

FUSIFORM, (of stem and spores), spindle-shaped.

GELATINOUS, jelly-like, applied to tissue whose hyphae become partially dissolved and glutinous in wet weather and when mounted in water under the microscope appear more transparent and wider, loosening from one another.

GENERIC, of the rank of a genus.

GENUS, a term in classification; each genus includes certain related species; the two names, viz. of its genus and its species, compose the binomial by which a plant is known in science, e. g. *Psalliota campestris*.

GIBBOUS, (of pileus), with an unsymmetrical convexity or umbo, or with convexity on one side.

GILLS, the knife-blade-like structures on the underside of the pileus; lamellae; collectively, the hymenophore.

GILL-TRAMA, the tissue of a gill between the two hymenial layers.

GILVOUS, (color), yellowish leather colored.

GLABRESCANT, becoming glabrous.

GLABROUS, (of pileus and stem), surface destitute of scales, hairs, etc., smooth.

GLANDULAR, with sticky drops or glands.

GLAUCOUS, (of pileus), covered with fine white bloom, easily rubbed off.

GLOBOSE, spherical or almost so.

GLUTEN, (of cuticle of pileus or stem, of universal veil), the dissolved gelatinous hyphae of certain tissues; very sticky and toughish.

GLUTINOUS, provided with gluten.

GRANULAR, (of pileus or stem), covered with granule-like substance.

GRANULOSE, same as granular.

GREGARIOUS, growing in company, scattered closely over a small area.

GROUP, a general term, applied indefinitely to a large or small number of plants whether classified or not.

GUTTATE, (of pileus), spotted as if by drops of liquid.

GUTTULATE, (of spores), containing an oily globule.

HABIT, the manner of growth of a plant.

HABITAT, the natural place of growth of a plant.

HAIRY, (of pileus), covered by an arrangement of fibrils resembling hairs.

HERBACEOUS, said of those flowering plants which perish annually down to the roots.

HERBARIUM, a collection of dried plants arranged systematically.

HETEROGENEOUS, applied to a structure composed of unlike tissues.

HIRSUTE, (of pileus), covered with rather long stiff fibres or hairs.

HISPID, (of pileus), covered with stiff bristle-like hairs.

HOARY, (of pileus or stem), covered with dense silky down; canescent.

HOMOGENEOUS, applied to structures composed of uniform tissues.

HOST, the plant or animal on or in which a parasitic fungus exists.

HUE, (of color), used here indiscriminately for "tint" or "shade." See Ridgway's Color Standards (1912) page 17, for correct usage.

HUMUS, the mixture of decayed vegetation and soil in the forest.

HYALINE, (of spores, gluten, etc.), colorless; transparent.

HYGROPHANOUS, (of flesh of mushrooms, or surface of pileus), watery in appearance, like the "water-core" of an apple, moisture disappearing rapidly accompanied by change in color, usually by fading.

HYGROSCOPIC, readily absorbing moisture from the atmosphere.

HYMENIUM, aggregation of the basidia in a continuous layer mixed with cystidia or sterile cells when present; the spore-bearing layer.

HYMENOMYCETES, the group of fungi possessing a hymenium composed of basidia which are exposed.

HYMENOPHORE, the portion of the fruit-body bearing the hymenium.

HYPHAE, plural of hypha; same as mycelium, composing also the fruit-body.

ICONES, colored plates illustrating fungi.

IDENTIFICATION, the study of the characters of a plant in order to determine its name.

IMBRICATE, (of pilei), overlapping one another, like the shingles of a roof.

INCARNATE, (color), flesh-colored.

INCISED, (of margin of pileus), as if cut into.

INCOMPLETE, (of annulus), forming a partial ring.

INCRASSATE, (of stem), thickened.

INCURVED, (of margin of pileus), same as inflexed.

INDIGENOUS, native, not foreign.

INFERIOR, (of annulus), below the middle of the stem.

INFLATED, (of cystidia), swollen like a bladder.

INFUNDIBULIFORM, (of pileus), funnel-shaped.

INNATE, (of scales, fibrils, etc.), a part of the surface tissue, not superficial.

INSERTED, (of base of stem), attached directly without "roots" or fibrils; instititious.

INSTITITIOUS, same as inserted.

INTERSPACES, (of gills), spaces between the attachment of the gills to the pileus.

INTERVENESE, (of gills), veined in the interspaces.

INTERWOVEN, (of trama), intermingled arrangement of hyphae, not parallel, convergent nor divergent.

INTRODUCED, brought from another country and growing spontaneously.

INVOLUTE, (of margin of pileus), rolled in, especially when young.

ISABELLINE, same as alutaceous; pale tan-color.

LABYRINTHIFORM, of sinuous lines; like a labyrinth.

LACERATE, (of annulus, scales, pileus, etc.), as if torn.

LACINIATE, (of margin of annulus or pileus), cut coarser than fimbriate; slashed.

LACTIFEROUS, (of hyphae of trama), bearing a milky juice.

LACUNOSE, (of pileus or stem), covered with pits or indentations.

LAMELLAE, plural of lamella; same as gills.

LANATE, same as woolly.

LANCEOLATE, (of spores, cystidia or gills), lance shaped; many times longer than broad, and tapering.

LATEX, a juice, usually of milky color, but also applied to other colors.

LATERAL, (of stem), attached to one side of the pileus.

LENS, a hand magnifying glass.

LIGNATILE, growing on wood.

LIGNICOLOUS, same as lignatile.

LINGULATE, (of pileus), tongue-shaped.

LIVID, (color), like that of a bruise; bluish-black.

LOBED, (of pileus), with rather large, rounded divisions on the margin.

LUCID, clear to the understanding; transparent.  
 LURID, (color), smoky-reddish, sordid.  
 LUTEOUS, (color), dull egg-yellow; see Saccardo's Color Key.  
 LUTESCENT, (color), becoming luteous.  
 MACROSCOPIC, visible without magnification.  
 MACULATE, spotted.  
 MAMMIFORM, (of umbo), breast-shaped.  
 MARGINATE, (of pileus), with a distinctly marked border; (of bulb), with a circular ridge on the exterior upper angle where the universal veil was attached.  
 MARGINATE-DEPRESSED, (of bulb), provided with a narrow, circular, horizontal platform on the upper side.  
 MAST, the fruit of forest trees like acorns and nuts, often used of a heap of nuts.  
 MATRIX, the substance on or in which a fungus grows.  
 MILKY, of the color of milk.  
 MEDIAL, (of annulus), situated at or near the middle of the stem.  
 MEMBRANOUS, (of pileus, annulus, etc.), thin and pliant like a membrane; applied when the trama of pileus is quite thin.  
 MEMBRANACEOUS, same as membranous.  
 MICACEOUS, (of pileus), covered with glistening mica-like particles.  
 MIXED, referring to forests containing both conifer and broad-leaved trees.  
 MILD, (odor and taste), not with a distinctly marked peculiarity.  
 MICROSCOPIC, of a size requiring the use of a microscope to see clearly.  
 MICROSCOPICAL, same as microscopic.  
 MICRON, (measure), of the length of one-thousandth part of a millimetre; used to designate size as measured by the use of a microscope.  
 MICROMETER, a disc of glass ruled with lines forming a metric scale for measuring objects under the microscope in microns.  
 MICRO-CHEMICAL, referring to tests with chemicals on microscopic objects.  
 MOLDS or MOULDS, certain fungi whose vegetative growth appears mouldy.  
 MONSTROSITY, applied to a specimen of a very abnormal appearance.  
 MORPHOLOGICAL, pertaining to form and structure, often used in a phylogenetic sense.  
 MOVABLE, (annulus), that can be moved more or less easily up and down the stem.

MUCILAGINOUS, slimy.  
 MUCCOUS, slime.  
 MUSHROOM, a general term applied to the fleshy Agarics and fleshy species of other fungi; a mushroom may be edible, poisonous, unpalatable, tough, etc., but popular usage applies the term only to edible ones. See toadstool.  
 MYC., MYCET., MYCETO., MYCO., prefixes signifying fungus.  
 MYCELIUM, came as hyphae; the vegetative part of a fungus consisting of microscopic threads usually with cross-walls to form the cells which contain the living protoplasm.  
 MYCELEOID, (of base of stem), provided with a white mouldy growth of mycelium.  
 MYCOLOGICAL, relating to mycology.  
 MYCOLOGY, the term applied to the science dealing with fungi.  
 MYCOPHAGIST, one who eats mushrooms; an epicure concerning mushrooms.  
 MYCOLOGIST, one who is versed in mycology; a specialist in the study of fungi.  
 MYCORHIZA, the stunted rootlets of trees, when such rootlets are covered or permeated by the mycelium of fungi.  
 NAKED, (of pileus or stem), entirely devoid of fibrils, scales or other covering.  
 NARROW, (of gills), a relative term, the opposite of broad; determined by experience.  
 NIGRESCENT, (color), turning blackish.  
 NUCLEATE, (of spores), containing microscopically visible, oil-like globules.  
 OBCLAVATE, (of cystidia, spores, stem), a reversal of clavate.  
 OBCONIC, reversal of conic.  
 OBLONG, (of spores), twice as long as wide.  
 OBOVATE, (of spores, etc.), reversal of ovate.  
 OBSOLETE, (of annulus, scales, etc.), very imperfectly developed, hardly perceptible; (of terms), no longer in use.  
 OBTUSE, (of pileus, cystidia, spores), rounded or blunt; greater than a right angle.  
 OCHRACEOUS, (color), dingy ochre-yellowish.  
 OCHREATE, (of volva), sheathing the stem at base like a stocking.  
 OLIVACEOUS, (color), with an olive shade.  
 OPAQUE, dull, not shining.  
 ORBICULAR, (of pileus), circular in outline.  
 OVAL, (of young pileus, spores), having the shape of an egg.  
 OVATE, similar to oval but rather pointed at the narrower end.  
 OVOID, same as oval.

- PALLID, (color), of an indefinite pale or whitish appearance.
- PAPILLA, a small, nipple-shaped elevation.
- PAPILLATE, (of pileus or spores), provided on surface with papillae.
- PARALLEL, (of trama of gills), in section the hyphae lie continuously side by side.
- PARASITE, an organism living upon another live organism and deriving food from it, with or without fatal effect.
- PARTIAL VEIL, the inner veil, extending from the margin of the pileus to the stem. See page 4.
- PATCHES, (of scales or remnants of the universal veil), flat, closely applied pieces.
- PECTINATE, (of margin of pileus), resembling the teeth of a comb.
- PEDICEL, (of cystidia) a slender stalk.
- PELLICLE, same as cuticle, sometimes thought of as thinner and more definite.
- PELLICULOSE, provided with a pellicle.
- PELLUCID, translucent.
- PELLUCID-STRIATE, (of pileus), when as the result of the thinness of the pileus the gills become visible thru it and appear as striae.
- PENTAGONAL, (of spores), angular and five-sided.
- PERIPHERY, the outer boundary or surface.
- PERONATE, (of stem), bootied; sheathed by the volva or universal veil.
- PERONATE-SCALY, (of stem), when the sheath of a peronate stem is broken up and the parts persist.
- PERSISTENT, retaining its place, shape or structure, not disappearing.
- PETALOID, (of pileus), shaped like the petal of a flower, narrowed somewhat at base.
- PHYLOGENETIC, pertaining to phylogeny.
- PHYLOGENY, the history of the evolution of the group or race to which a species belongs.
- PILEATE, possessing a pileus.
- PILEI, the plural of pileus.
- PILEUS, the cap or that structure of an Agaric which bears the gills on its under side.
- PILOSE, (of pileus), covered with long, soft, hairy filaments.
- PIPSHAPED, (of spores), shape of an apple seed.
- PITTED, (of pileus or stem), similar to lacunose; with little depressions.

- PITH, (of stem), the soft tissue in the interior, which often disappears so that the stem becomes hollow.
- PLANE, (of pileus), with a flat surface.
- PLIANT, not rigid nor firm; easily bent.
- PLICATE, (of pileus), plaited; folded like a fan.
- PLUMBEOUS, (color), like lead.
- PLUMOSE, finely feathery.
- POROSE, (of hymenophore), approaching the condition of possessing pores.
- POSTERIOR, (of gills), behind, toward the stem.
- PRAEMORSE, (of the "root" or base of stem), as if broken off abruptly.
- PROLIFEROUS, (of stem), producing other stems on itself near the base.
- PROTEAN, exceedingly variable.
- PROTOPLASM, the living semifluid substance of the cells which is the basis of life.
- PRUINATE, same as pruinose.
- PRUINOSE, (on the surface), as if finely powdered.
- PSEUDOPARENCHYMA, the tissue of fungi when its cellular structure imitates the parenchyma of higher plants.
- PSEUDOPROSENCHYMA, tissue of fine elongated hyphae, somewhat resembling the prosenchyma of higher plants.
- PUBESCENCE, a covering of short, soft, downy hairs.
- PUBESCENT, provided with pubescence.
- PULVERULENT, covered as if with powder.
- PUNCTATE, (of pileus or stem), dotted with minute scales or other substance.
- PUTRESCENT, soon decaying and becoming soft and mushy.
- PYRIFORM, (of cystidia or cells), pear-shaped.
- PYRAMIDAL, (of scales), pyramid-shaped.
- QUADRATE, (of spores), angularly four-sided.
- RADICATING, (of stem), imitating a root.
- RAMIFICATION, branching.
- REFLEXED, (of margin of pileus), turned up or back.
- REMOTE, (of gills), free and at some distance from the stem; (of annulus), at some distance from apex of stem.
- RENIFORM, (of pileus), kidney-shaped.
- REPAND, (of pileus), wavy on margin and turned back or elevated.
- RESUPINATE, (of pileus), with the upper surface reclining on the substratum, the gills facing outward.
- RESUPINATE-REFLEXED, (of pileus), attached for some dis-

- tance by the back surface, the other portion extending out like a shelf.
- RETICULATE**, (of pileus or stem), marked by lines, veins or ridges which cross one another as in a net.
- REVIVING**, said of a fruit-body which shrivels in dry weather and takes on its natural shape when wet.
- REVOLUTE**, (of margin of pileus), rolled back or up.
- RHIZOIDS**, radiating hyphae extending into substratum from base of stem.
- RHIZOMORPHS**, visible strands or cords of compacted mycelium, often dark colored, penetrating a soft substratum or between portions of it, as between bark and wood, etc.
- RIMOSE**, (of pileus), cracked.
- RIND**, same as cortex.
- RING**, same as annulus.
- RIVULOSE**, (of pileus and stem), marked with lines like a river-system on a map.
- ROOTING**, (of stem), an attenuated prolongation into the soil or substratum.
- ROUND-BULBOUS**, of a bulb not marginate.
- RUFESCENT**, (color), becoming reddish.
- RUFOUS**, (color), reddish, dull red.
- RUGOSE**, coarsely wrinkled.
- RUGULOSE**, finely wrinkled.
- SACCATE**, (of cystidia), shape of a meal-bag.
- SANGUINEOUS**, (color), blood-red.
- SAPROPHYTE**, a plant which lives on dead vegetable or animal matter.
- SCABROUS**, (of pileus), rough with short, rigid projections.
- SCALES**, applied to various decorations on the pileus and stem; torn portions of the cuticle or of the universal veil or of the volva; they may be membranous, fibrillose, hairy, floccose, hard, erect, flat, patch-like, etc.; often an important feature for identification.
- SCALY**, provided with scales.
- SCLEROTIA**, resting-bodies of small size, composed of a hardened mass of hyphae, from which fruit-bodies may develop.
- SCISSILE**, (of flesh of pileus), capable of being pulled into horizontal layers; this condition is most marked in a hygrophanous pileus.
- SECEDE**, (of gills), when at first attached to stem, i. e. adnate or adnexed, but separating from it later.
- SECONDARY SPORES**, not borne on basidia; conidia, chlamy-

- dospores, etc.; formed directly on the mycelium or on hyphae of the fruit body.
- SEPARABLE**, (of cuticle, pellicle, etc.), not adnate.
- SEPARATING**, see secede.
- SERICEOUS**, silky.
- SERRATE**, (of gills), with saw-tooth-like edge.
- SERRULATE**, minutely serrate.
- SERRATULATE**, same as serrulate.
- SESSILE**, (of pileus), without a stem.
- SETACEOUS**, (of stem), bristle-form.
- SHAGGY**, rough with long compact fibrils.
- SILKY**, covered with shining, close-fitting fibrils.
- SINUATE**, (of gills), a concave indentation of its edge near the stem.
- SINUOUS**, wavy, serpentine.
- SLENDER**, (of stem), very long as compared to its thickness; relative to stout.
- SMOOTH**, (of spores), not spiny, tuberculate, rough, nor angular, etc; (of pilus and stem), see glabrous.
- SOLITARY**, not growing in the immediate neighborhood of other individuals.
- SOLID**, (of stem), not hollow nor stuffed; of a texture in its central axis similar to that found in the rest of a cross-section.
- SORDID**, (color), dirty or dingy.
- SPADICEOUS**, (color), date-brown.
- SPATHULATE**, (of pileus), spatula-shaped; oblong with attenuated base.
- SPECIES**, the lowest term in classification; a group of individuals agreeing in certain characters which appear again in their progeny; one species differs from another in several marked characters agreed upon as sufficiently specific by tradition or by specialists in the group; a species is therefore a judgment, and has limitations imposed by an agreement of the judgments of scientific men. One or more species with certain common characters constitute a genus.
- SPECIFIC**, referring to characters which are used in designating or distinguishing species.
- SPHAGNUM**, a genus of mosses; bog-moss.
- SPINY**, (of spores), strongly echinulate.
- SPHOEROID**, (of spores), nearly spherical; similar to spherical.
- SPONGY**, (of stem), soft and tending to be water-soaked.
- SPONGY-STUFFED**, (of stem), with a spongy pith.

**SPORE**, the reproductive cells in Agarics borne four on each basidium; more accurately called basidio-spores. In other cryptogams the term is applied to reproductive cells or bodies of a great variety of kinds. The basidio-spores when they germinate give rise to mycelium.

**SPOROPHORE**, fruit-body.

**SPURIOUS**, false.

**SQUAMOSE**, (of pileus or stem), covered with scales.

**SQUAMULOSE**, minutely squamose.

**SQUAMULE**, scale.

**SQUARROSE**, (of pileus and stem), covered with recurved scales.

**STAINED**, said of any part which appears as if some coloring matter had been spilled on it and spread on the surface.

**STALK**, an indefinite term for stem, pedicel, etc.

**STELLATE**, (of spores, scales), with extensions like that of a star.

**STERIGMA**, the tiny spicule-like extension at the apex of a basidium on which the spores develop.

**STERILE**, said of a fruit-body or hymenium which is immature or produces no spores; or simply, without spores.

**STERILE CELLS**, term applied in this report to the slender cells on the edge of gills which bear no spores and which cause the fimbriate appearance of the edge as shown under a lens.

**STIPE**, technical term for the stem of mushrooms; see stem.

**STIPITATE**, possessing a stem.

**STOUT**, (of stem), relative to slender; not so many times longer than thick.

**STRAIGHT**, (margin of pileus), when not incurved.

**STRAMINEOUS**, straw-color.

**STRIATE**, (of margin of pileus), radiating minute furrows or lines; (of stem), longitudinal lines or minute furrows.

**STRIAE**, the lines or furrows when striate.

**STRIGOSE**, (of pileus or stem), with coarse or thick, long, rather stiff hairs.

**STUFFED**, (of stem), when the axis is filled with a differentiated pith which usually disappears in age leaving it hollow.

**SUB—**, prefix signifying "almost," "somewhat" or "under."

**SUBGENUS**, a term in classification; a grouping under a genus and containing groups of related species. The subgenera of the mycologists of one generation are often raised to the rank of genera by later students.

**SUBICULUM**, a more or less dense felt of hyphae covering the substratum, from which the fruit-bodies arise.

**SUBDISTANT**, (of gills), the spacing halfway between close and distant.

**SUBDECURRENT**, (of gills), when the attachment extends slightly farther down the stem than when adnate.

**SUBHYMENIUM**, a differentiated tissue just beneath the hymenium.

**SUBSTRATUM**, the substance in or on which the fungus grows, as soil, humus, fallen leaves, dung, wood, animal remains, etc. A better term than matrix.

**SUBULATE**, awl-shaped.

**SULCATE**, (of pileus and stem), grooved, more extreme than striate, less so than plicate.

**SUPERFICIAL**, (of scales, flocci, etc.), on the surface and easily removable.

**SUPERIOR**, (of annulus), attached above the middle of the stem.

**SYNONYM**, the name or names of a species or genus no longer tenable, either because of error in naming, rearrangement of the classification or as a result of "rules" promulgated by scientific men acting in agreement. Many of the long known plants have a number of such synonyms.

**TAN**, (color), leather-colored, similar to undressed leather; isabelline.

**TENACIOUS**, tough.

**TERETE**, (of stem), round like a broom-handle, not irregular.

**TERRESTRIAL**, growing on the ground.

**TESTACEOUS**, (color), brick-red.

**TEXTURE**, the arrangement of the components of the different tissues, as compact, loose, etc.

**TINGED**, with a tint of a color.

**TISSUE**, an aggregate of similar cells or hyphae.

**TOADSTOOL**, same as mushroom; popularly applied to those about which the user of the term often has no knowledge and which he therefore considers poisonous; a large number of so-called "toadstools" are edible.

**TOMENTOSE**, (of pileus or stem), densely covered with a matted wooliness or tomentum.

**TOMENTUM**, composed of long, soft, entangled or matted fibrils.

**TOXIC**, poisonous.

**TOOTH**, (of gills), decurrent by a tooth; see uncinatate.

**TRAMA**, the fleshy portion of pileus or gills composed of hyphae.

**TRANSLUCENT**, capable of transmitting light without being transparent.

TRANSVERSE, cross-wise.  
 TREMELLOID, of a gelatinous consistency.  
 TRUNCATE, an enlarged portion ending as if cut off.  
 TUBERCLE, any wart-like or knob-like excrescence.  
 TUBERCULATE, (of spores), covered with minute tubercles.  
 TUBERCULAR-STRIATE, (of pileus), when the striae are roughened by small tubercles.  
 TUMID, (of stem), swollen, inflated.  
 TURBINATE, (of pileus), top-shaped.  
 TYPE, the original specimen or specimens from which the species was described and named.  
 TYPICAL, agreeing with the descriptions of the type or with the type itself.  
 UMBER, (color), almost tobacco-colored; see Saccardo's Color Key.  
 UMBILICATE, (of pileus), with a central, naval-like depression.  
 UMBILICUS, a naval-like depression.  
 UMBO, (of pileus), a raised, conical to convex knob or mound on the center.  
 UMBONATE, (of pileus), provided with an umbo.  
 UNCINATE, (of gills), provided with a narrow, decurrent extension at the stem.  
 UNDULATE, same as wavy.  
 UNEQUAL, (of gills), of different length, some reaching the stem, others shorter.  
 UNEVEN, (of pileus or stem), said of surfaces with striations, reticulations, tubercles, etc.; not even.  
 UNICOLOROUS, of the same color throughout.  
 UNIVERSAL VEIL, sometimes used for volva.  
 VAGINATE, (of stem), provided with a long volva or sheath at the lower end.  
 VARIABILITY, the state of being variable.  
 VARIABLE, capable of taking on a number of different shapes, forms, colors or other characters, while retaining its specific identity.  
 VARIEGATED, marked with a variety of colors, which are intermingled.  
 VARIETY, (a). Here used to refer to a form which a species constantly assumes, under definite conditions, e. g. climate, soil, artificial culture, etc. Such forms are often given names, as *Psalliota compestris* var. *hortensis*.  
 (b). Also used to designate forms which are not typical, but which are not sufficiently known to be designated by a specific

name; such are given the name of the nearest species in this report followed by the abbreviation "var."  
 (c). When a variety is found to be constant in its characters and always distinct in some such characters from other species, it should eventually be given a specific name, but in rare varieties such constancy is not easily proven.  
 VEIL, see "partial veil," "universal veil," "cortina" and "volva."  
 VEINED, (of gills), with vein-like wrinkles or raised lines on the surfaces.  
 VELUM, see veil.  
 VENOSE, same as veined.  
 VENTRAL, on the under side of, opposed to dorsal.  
 VENTRICOSE, (of stem), swollen or enlarged in the middle.  
 VERRUCOSE, warty.  
 VERRUCULOSE, minutely warty.  
 VESICULOSE, referring to the trama of the Lactariae. See page 83.  
 VILLOSE, covered with long, soft, weak hairs.  
 VINACEOUS, (color), of the color of red wine.  
 VIOLACEOUS, (color), of some violet hue.  
 VIRESCENT, (color), becoming greenish.  
 VIRGATE, (of pileus), streaked, usually by differently colored fibrils.  
 VISCID, sticky.  
 VISCOUS, gluey.  
 VITELLINE, (color), egg-yellow.  
 VOLVA, the universal veil of certain genera. See pages 4 and 593.  
 WARTY, (of pileus, spores, etc.), covered by small wart-like excrescences.  
 WAVY, (of margin of pileus), alternately raised and depressed like waves.  
 WAXY, (of gills), of a consistency that can be partially or wholly moulded or compressed into balls.  
 ZONATE, (of pileus), marked with concentric bands of color.  
 ZONED, same as zonate.

---

---

INDEX

---

---

# INDEX

A.

	Page
Abbreviations .....	876
Agaricaceae .....	27
Agaricales .....	26
Agariceae .....	205
Agarics, black-spored .....	30, 205
poisoning by .....	858
classification of .....	23
collecting and preserving of .....	16
definition of .....	3, 4, 5
distribution of in Michigan .....	13
genera of .....	27
habit and growth conditions of .....	13
ochre-spored .....	28, 205
poisoning by .....	854
pink-spored .....	29, 205
poisoning by .....	853
poisoning by .....	825
preserving of .....	16
purple-spored .....	29, 205
poisoning by .....	857
white-spored .....	27, 205
poisoning by .....	828, 852
Amanita, genus .....	4, 27, 593
Amanita key to species .....	597
Amanita abrupta Pk. ....	599
bisporiger Atk. ....	603
bulbosa Bull. ....	601
caesarea Fr. ....	15, 600
candida Pk. ....	599
chlorinosma Pk. ....	615
poisoning by .....	837
chrysolema Atk. ....	613
cinera Bres. ....	606
cinereoconia Atk. ....	598
citrina, poisoning by .....	836
coccola (Scop.) Fr. ....	605
cothurnata Atk. ....	612
crenulata Pk. ....	599
poisoning by .....	836
excelsa Fr. ....	599
flavoconia Atk. ....	619
flavorubescens Atk. ....	619
frostiana Pk. ....	612
poisoning by .....	837
junquillea Quel. ....	617
poisoning by .....	837
mappa Fr. ....	609
poisoning by .....	836
muscaria Fr. ....	610
poisoning by .....	837, 863
poisoning by, clinical features of .....	838, 843
treatment for .....	841
poisonous constituents .....	839
var. alba .....	611
var. formosa .....	611
var. puellaris .....	611
var. regalis .....	611
ovoidea Fr. ....	605
pantherina Fr. ....	599
poisoning by .....	843
peckiana Kauff. ....	604
phalloides Fr. ....	600
poisoning by .....	827, 828, 863
treatment of .....	832
poisonous constituents of .....	833

	Page
<i>Amanita porphyria</i> Fr.	604
<i>radicata</i> Pk.	616
<i>recutita</i> Fr.	608
<i>rubescens</i> Fr.	617
<i>russuloides</i> Pk.	15, 616
<i>solitaria</i> Fr.	614
poisoning by	843
<i>spissa</i> Fr.	620
<i>spretia</i> Fr.	15, 606
poisoning by	836
<i>strobiliformis</i> Vitt.	615
<i>tomentella</i> Kromb.	607
<i>velatipes</i> Atk.	599
<i>verna</i> Fr.	602
poisoning by	846
<i>virosa</i> Fr.	603
poisoning by	836
<i>Amanita</i> , toxic principles of	843
<i>Amanitopsis</i> , genus	4, 27, 612
<i>Amanitopsis</i> , key to species	622
<i>Amanitopsis</i> <i>adnata</i> Smith	622
<i>agglutinata</i> B. & C.	623
<i>albocreata</i> Atk.	622
<i>farinosa</i> Schw.	622
<i>inaurata</i> Gill.	624
<i>parcivolvata</i> Pk.	622
<i>pusilla</i> Pk.	622
<i>strangulata</i> Fr.	624
<i>vaginata</i> Fr.	623
poisoning by	830
<i>var. alba</i>	623
<i>var. fulva</i>	623
<i>var. livida</i>	623
<i>volvata</i> Fr.	622
poisoning by	837
<i>Amaurospora</i>	205, 232
<i>Annulus</i> , definition of	5
<i>Armillaria</i> , genus	4, 27, 642
<i>Armillaria</i> , key to species	649
<i>Armillaria</i> <i>appendiculatus</i> Pk.	649
<i>aurantia</i> Fr.	651
<i>bulbiger</i> Fr.	649
<i>caligata</i> Vitt-Bres.	650
<i>corticata</i> Fr.-Pat.	654
<i>dryina</i> Fr.-Pat.	654
<i>focalis</i> Fr.	652
<i>macrospora</i> Pk.	648
<i>magnivelaris</i> Pk.	649
<i>mellea</i> Fr.	653
<i>mucida</i> Fr.	649
<i>nardosmia</i> Ell.	648, 649
<i>ponderosa</i> Pk.	649
<i>rhagadisma</i> Fr.	650
<i>robusta</i> Fr.	649
<i>viscidipes</i> Pk.	648, 649
B.	
<i>Basidia</i> , definition of	7
<i>Basidiomycetes</i>	26
<i>Bibliography</i> , general	865
<i>Bibliography</i> of monographs of genera	872
of mushroom poisoning	871
<i>Bolbitius</i> , genus	29, 502
<i>Bolbitius</i> , key to species	502
<i>Bolbitius</i> <i>fragilis</i> Fr.	503
<i>radicans</i> Morg.	506
<i>tener</i> Berk.	503
<i>vitellinus</i> Fr.	504
<i>Boletus</i> , poisoning by	860
C.	
<i>Cantherelleae</i>	31
<i>Cantherellus</i> , genus	27, 32
<i>Cantherellus</i> , key to species	33
<i>Cantherellus</i> <i>aurantiacus</i> Fr.	40
<i>cibarius</i> Fr.	35
<i>cinnabarinus</i> Schw.	36
<i>clavatus</i> Fr.	34
<i>dichotomus</i> Pk.	39
<i>focosus</i> Schw.	34
<i>friesii</i> Quel.	37
<i>infundibuliformis</i> Fr.	37
<i>lutescens</i> Fr.	38

	Page
<i>Cantherellus neurophyllum</i>	34
<i>tubaeformis</i> Fr.	38
<i>umbonatus</i> Fr.	39
<i>Carpophore</i> , definition of	4, 28, 533
<i>Chamaeota</i> , genus	4, 28, 533
<i>Chamaeota</i> <i>fenzlii</i> Fr.	534
<i>mampillata</i> (Longyear) Murr.	533
<i>sphaerospora</i> (Pk.)	534
<i>Claudopus</i> , genus	29, 590
<i>Claudopus</i> , key to species	591
<i>Claudopus</i> <i>byssisedus</i> Fr.	592
<i>depluens</i> Fr.	592
<i>griegensis</i> Pk.	593
<i>indulans</i> Fr.	591
<i>variabilis</i> Fr.	592
<i>Clitocybe</i> , genus	28, 715
<i>Clitocybe</i> , key to species	716
<i>Clitocybe</i> <i>adirondackensis</i> Pk.	741
<i>albida</i> Pk.	742
<i>albissima</i> Pk.	731
<i>angustissima</i> Fr.	746
<i>anisearia</i> Pk.	729
<i>caespitosa</i> Pk.	743
<i>candicans</i> Fr.	729
<i>candida</i> Bres.	720
<i>cartilaginea</i> Burl-Bres.	734
<i>carosior</i> Pk.	727
<i>catina</i> Fr.	735
<i>cerussata</i> Fr.	731
<i>clavipes</i> Fr.	726
<i>compressipes</i> Pk.	746
<i>connexa</i> Pk.	731
<i>cyathiforme</i> Fr.	739
<i>dealbata</i> Fr.	730
<i>dealbata</i> <i>var. deformata</i>	730
<i>var. sudorifica</i> , poisoning by	847
<i>decora</i> Fr.	724
<i>ditopoda</i> Fr.	730
<i>eccentrica</i> Pk.	741
<i>ectypoides</i> Pk.	740
<i>expallens</i> Fr.	740
<i>flaccida</i> Fr.	736
<i>gallinacea</i> Fr.	737
<i>geotropia</i> Fr.	736
<i>gigantea</i> Fr.	719
<i>illudens</i> Schw.	732
poisoning by	846
<i>incilis</i> Fr.	725
<i>infundibuliformis</i> Fr.	736
<i>laccata</i> Fr.	747
<i>var. amethystina</i> Bolt	747
<i>var. pallidifolia</i> Pk.	747
<i>var. striatula</i> Pk.	747
<i>leptoloma</i> Pk.	732, 742
<i>maxima</i> Fr.	721
<i>media</i> Pk.	727
<i>metachroa</i> Fr.	734
<i>monadelpha</i> Morg.	723
<i>morbifera</i> Pk.	745
poisoning by	848
<i>multiceps</i> Pk.	733
<i>nebularis</i> Fr.	725
poisoning by	848
<i>ochropurpurea</i> Berk.	748
<i>odora</i> Fr.	728
<i>paralis</i> Fr.	737
<i>peltigerina</i> Pk.	744
<i>phylliphila</i> Fr.	735
<i>piceina</i> Pk.	720
<i>pinophila</i> Pk.	737
<i>pithyophila</i> Fr.	737
<i>praecox</i> sp. nov.	724
<i>pulcherrima</i> Pk.	720
<i>sinopica</i> Fr.	738
<i>spinulosa</i> Smith	739
<i>subconcaeva</i> Pk.	740
<i>subcyathiforme</i> Pk.	740
<i>sulphurea</i> Pk.	739
<i>tabescens</i> Bres.	723
<i>torilis</i> Fr.	748
<i>truncicola</i> Pk.	732
<i>vilscens</i> Pk.	727
<i>viridis</i> Fr.	729
<i>Clitopilus</i> , genus	29, 562
<i>Clitopilus</i> , key to species	563

	Page
<i>Clitopilus abortivus</i> B. & C. ....	564
<i>albogriseus</i> Pk. ....	567
<i>caespitosus</i> Pk. ....	570
<i>micropus</i> Pk. ....	566
<i>novaboracensis</i> Pk. ....	569
<i>orella</i> Fr. ....	569
<i>popinalis</i> Fr. ....	570
<i>prunulus</i> Fr. ....	569
<i>subplanus</i> Pk. ....	567
<i>subvilis</i> Pk. ....	565
<i>undatus</i> Fr. ....	566
<i>woodianus</i> Pk. ....	564
<i>Collybia</i> , genus. ....	28, 749
<i>Collybia</i> , key to species. ....	751
<i>Collybia abundans</i> Pk. ....	764
<i>acervata</i> Fr. ....	759
<i>alba</i> Pk. ....	752
<i>albiflavida</i> (Pk.) ....	764
<i>albipilata</i> Pk. ....	772
<i>alcalinolens</i> Pk. ....	756
<i>amabilipes</i> Pk. ....	751
<i>aquosa</i> Fr. ....	758
<i>atrata</i> Fr. ....	761
<i>atratoidea</i> Pk. ....	763
<i>butyracea</i> Fr. ....	753
<i>campanella</i> Pk. ....	777
<i>cirrata</i> Fr. ....	773
<i>colorea</i> Pk. ....	758
<i>confluens</i> Fr. ....	775
<i>conigena</i> Fr.-Bres. ....	772
<i>conigenoides</i> Ellis. ....	772
<i>delicatella</i> Pk. ....	752
<i>dryophila</i> Fr. ....	754
<i>esculenta</i> Fr.-Bres. ....	772
<i>esculentoides</i> Pk. ....	752
<i>estensis</i> Morg. ....	755, 756
<i>expallens</i> Pk. ....	763
<i>famalia</i> Pk. ....	757
<i>floccipes</i> Fr. ....	771
<i>fuliginella</i> Pk. ....	752
<i>hariolorum</i> Fr. ....	774
<i>hygrophoroides</i> Pk. ....	760
<i>lacerata</i> Fr. ....	757
<i>lachnophylla</i> Berk. ....	750
<i>lacunosa</i> Pk. ....	777
<i>lentiginoides</i> Pk. ....	755
<i>longipes</i> Fr. ....	770
<i>maculata</i> A. & S. ....	768
<i>myriadophylla</i> Pk. ....	761
<i>platyphylla</i> Fr. ....	767
<i>plexipes</i> Fr. ....	762
<i>radicata</i> Fr. ....	766
<i>radicata</i> var. <i>furfuracea</i> Pk. ....	767
var. <i>pusilla</i> Pk. ....	767
<i>spinulifera</i> Pk. ....	751
<i>stipitaria</i> Fr. ....	776
<i>stipitaria</i> var. <i>setipes</i> Pk. ....	777
<i>strictipes</i> Pk. ....	755
<i>succinea</i> Fr. ....	765
<i>succosa</i> Pk. ....	770
<i>szorconera</i> Fr. ....	768
<i>tuberosa</i> Fr. ....	773
<i>velutipes</i> Fr. ....	769
<i>zonata</i> Pk. ....	776
<i>Coprinus</i> , genus. ....	15, 30, 206
<i>Coprinus</i> , key to species. ....	207
<i>Coprinus angulatus</i> Lloyd. ....	225
<i>arenatus</i> Pk. ....	217
<i>atramentarius</i> Fr. ....	212
<i>boudieri</i> Quel. ....	225
<i>brassicae</i> Pk. ....	222
<i>bulbilosus</i> Pat. ....	220
<i>comatus</i> Fr. ....	209
poisoning by. ....	858
<i>domesticus</i> Fr. ....	218
<i>ebulbosus</i> Pk. ....	213
<i>ephemerus</i> Fr. ....	224
<i>finetarius</i> Fr. ....	215
<i>finetarius</i> var. <i>macrohiza</i> ....	215
<i>friesii</i> Quel. ....	222
<i>granulosus</i> Clem. ....	219
<i>insignis</i> Pk. ....	212
<i>ioneshi</i> Pk. ....	216
<i>lagopides</i> Karst. ....	216
<i>lagopus</i> Fr. ....	197

	Page
<i>Coprinus laniger</i> Fr. ....	214
<i>macrosperma</i> Pk. ....	211
<i>micaceus</i> Fr. ....	218
<i>micaceus</i> var. <i>conicus</i> Pk. ....	219
<i>narcoticus</i> Fr. ....	222
<i>niveus</i> Fr. ....	217
<i>oblectus</i> Fr. ....	211
<i>ovatus</i> Fr. ....	210
<i>patouillardii</i> Quel. ....	223
<i>plicatilis</i> Fr. ....	226
<i>plicatilloides</i> Buller. ....	224
<i>pulcherrifolius</i> Pk. ....	219
<i>quadridius</i> Pk. ....	213
<i>radiatus</i> Fr. ....	219
<i>radiatus</i> Fr. ....	223
<i>sclerotigenus</i> E. & E. ....	221
<i>semilanatus</i> Pk. ....	217
<i>silvaticus</i> Pk. ....	225
<i>spraguei</i> B. & C. ....	224
<i>stenocoleus</i> Lindbl. ....	211
<i>stercorarius</i> Fr. ....	220
<i>sterquillinus</i> Fr. ....	210
<i>tigrinellus</i> Boud. ....	222
<i>tomentosus</i> Fr. ....	215
<i>tuberosus</i> Quel. ....	221
<i>Cortinarius</i> , genus. ....	4, 15, 28, 314
<i>Cortinarius</i> , key to species. ....	319
<i>Cortinarius</i> acutoidea Pk. ....	441
<i>acutus</i> Fr. ....	442
<i>adustus</i> Pk. ....	410
<i>aggregatus</i> sp. nov. ....	346
<i>albidifolius</i> Pk. ....	388, 393
<i>albidipes</i> Pk. ....	364
<i>albidus</i> Pk. ....	359
<i>alboviolaceus</i> Fr. ....	370
<i>aleuriosmus</i> Malre. ....	351
<i>amarus</i> Pk. ....	333
<i>anfractus</i> Fr. ....	366
<i>annulatus</i> Pk. ....	376
<i>anomalus</i> Fr. ....	390, 410
<i>arenatus</i> Fr. ....	373
<i>argentatus</i> Fr. ....	381
<i>armeniacus</i> Fr. ....	429
<i>armillatus</i> Fr. ....	13, 413
<i>arquatus</i> Fr. ....	338
<i>asper</i> Pk. ....	373
<i>atkinsonianus</i> Kauff. ....	338
<i>aureifolius</i> Pk. ....	398
<i>autumnalis</i> Pk. ....	386
<i>azureus</i> Fr. ....	384
<i>badius</i> Pk. ....	417
<i>badius</i> var. ....	419
<i>basalis</i> Pk. ....	395
<i>berlesianus</i> (Pk.) Sacc. ....	333
<i>bicolor</i> Cke. ....	427
<i>bolaris</i> Fr. ....	375
<i>braendlei</i> Pk. ....	384
<i>brevissimus</i> Pk. ....	393
<i>brunneofulvus</i> Fr. ....	420
<i>brunneus</i> Fr. ....	420
<i>bulliardii</i> Fr. ....	413
<i>caerulescens</i> Fr. ....	349
<i>caesiocyaneus</i> Britz. ....	342
<i>caesius</i> Clem. ....	350
<i>caespitosus</i> Pk. ....	387
<i>callisteus</i> Fr. ....	385
<i>calochrous</i> Fr. ....	339
<i>camphoratus</i> Fr. ....	342
<i>candelaris</i> Fr. ....	431
<i>canescens</i> Pk. ....	379
<i>caninus</i> Fr. ....	390
<i>castaneus</i> Pk. ....	395
<i>castaneoides</i> Pk. ....	417
<i>castaneus</i> Fr. ....	428
<i>catskillensis</i> Pk. ....	386
<i>cinnabarinus</i> Fr. ....	401
<i>cinnamomeus</i> Fr. ....	396
<i>cinnamomeus</i> var. <i>croceus</i> . ....	397
<i>claricolor</i> Fr. ....	363
<i>clintonianus</i> Pk. ....	385
<i>collintus</i> Fr. ....	329
<i>coloratus</i> Pk. ....	368
<i>colus</i> Fr. ....	413
<i>communis</i> Pk. ....	369
<i>copakensis</i> Pk. ....	363

	Page
Cortinarius corrugatus Pk.	356
croceocolor Kauff.	378
croceoconus	397
croceofolius Pk.	399
crystallinus Fr.	334
cumatilis Fr.	360
cyanopus Fr.	347
cylindripes Kauff.	330
deceptivus Kauff.	409
decipiens Fr.	438
decoloratus Fr.	365
delibutus Fr.	333
distans Pk.	421
dolobratius Fr.	431
duracinus Fr.	430
elator Fr.	407
elator pallidifolius Pk.	332
elegantior Fr. var.	355
elegantoides sp. nov.	344
emollitus Fr.	334
erraticus Pk.	374
erugatus Fr.	431
erythrinus Fr.	438
evernius Fr.	406
flaviofolius Pk.	377
flexipes var. minor Fr.	411
fulgens Fr.	354
fulmineus	355
fulmineus Fr. var. sulphureus var. nov.	354
furfurellus Fr.	422
fuscoviolaceus Pk.	437
glabrellus Kauff.	432
glandicolor Fr.	422, 435
glaucopus Fr.	352
glutinosus Pk.	367
gracilis Pk.	388
griseus Pk.	410
haematochelis Fr.	414
heliotropicus Pk.	414
hemitrichus Fr.	336
herpiticus Fr.	424
hinnuleus Fr.	340
iliopodius Fr.	416
imbutus Fr.	418
impenis Fr.	426
impolitus sp. nov.	371
infractus Pers.-Bres.	419
intrusus Pk.	365
iodes B. & C.	358
iodioides sp. nov.	335
juberinus Fr.	335
lanatipes Pk.	436
lapidophilus Pk.	262
lepidopus Cke.	363
leucopus Fr.	390
lignarius Pk.	439
lilacinus Pk.	440
limoneus Fr.	381
livor Fr.	378
longipes Pk.	428
lucorum Fr.	366
luteo-fuscus Fr.	408
lutescens Pk.	367
lutens Pk.	377
maculipes Pk.	398
maculipes Pk.	361
malachius Fr.	371
mallicorius Fr.	399
mammosus sp. nov.	415
michiganensis sp. nov.	350
modestus Pk.	388
morrisii Pk.	414
muclifluus Fr.	328
multiformis Fr.	357
musciogenus Pk.	331
newfieldtensis Ellis.	377
nigrillis Pk.	422
obliquus Pk.	382
obtusus Fr.	439
ochraceus Pk.	378
ochroleucus Fr.	394
olivaceo-stramineus Kauff.	341
olivaceus Pk.	366
ophiopus Pk.	369
paleaceus Fr.	425
paludosus Pk.	415

	Page
Cortinarius penicillatus Fr.	376
pholideus Fr.	372
phyllophilus Pk.	354
plumiger Fr.	405
praepallens Pk.	436
prasinus Fr.	353
privignus Fr.	433
pulchrifolius Pk.	383
punctata Fr.	419
purpurascens Fr.	345
purpureophyllus sp. nov.	348
raphanoides Fr.	403
rigens Fr.	440
rigidus Fr.	423
rigidus (Scop.) Ricken.	423
rimosus Pk.	383
robustus Pk.	387
rubens sp. nov.	343
rubricosus Fr.	412, 413
rubripes Kauff.	384
rubrocinerereus Pk.	343, 357
rufolivaceus Fr.	368
saginus Fr.	335
salor Fr.	402
sanguineus Fr.	427
saturinus Fr.	439
scandens Fr.	341, 353
scaurus Fr.	409
sciophyllus Fr.	408
scutulatus Fr.	400
semisanguineus (Fr.)	394
sericipes Pk.	391
sinnulans Pk.	347
sphaerosperma sp. nov.	362
sphagnophilus Pk.	332
sphaerosporus Pk.	331
splendidus Pk.	391
spilomeus Fr.	373
squamulosus Pk.	379
squarrosus Clem.	334
sterilis Kauff.	427
subferruginea Fr.	411
subflexipes Pk.	415
sublanatus Fr.	357
sublateritius Pk.	332
submarginalis Pk.	371
subpulchrifolius sp. nov.	346
subpurpurascens Fr.	433
subrigens sp. nov.	366
subsimile (Pers.)	392
subtabularis sp. nov.	344
sulfurinus Quel.	343, 357
testaceus Cke.	376
tophaceus Fr.	404
torvus Fr.	360
triumphans Fr.	407
umidicola Kauff.	435
urascens Fr.	403, 415
valgus Fr.	360
varicolor Fr.	339
vellicopia sp. nov.	333
vibratilis Fr.	380
violaceus Fr.	353
virentophyllus sp. nov.	387
whitel Pk.	28, 516
Crepidotus, genus	517
Crepidotus, key to species	519
abidus E. & E.	524
Crepidotus applanatus Fr.	522
calolepis	520
cinnabarinus Pk.	518
crocotinctus Pk.	525
crocophyllus Berk.	518
distans Pk.	525
dorsalis Pk.	521
fulvotomentosus Pk.	518
haerens Pk.	519
herbarum Pk.	518, 519
latifolius Pk.	525
haustellaris Fr.	523
malachius B. & C.	519
mollis Fr.	523
putrigenus B. & C.	525
rafsii B. & Br.	525
seplarius Pk.	521

	Page
Crepidotus stipitatus sp. nov.	524
tillophila Pk.	525
versutus Pk.	520
Cystidia, definition of.	10
D.	
Dictyolus retrugis.	31
E.	
Eccilia, genus.	29, 586
Eccilia, key to species.	587
Eccilia apiculatus Fr.	586
Eccilia atrides Fr.	588
carneogrisea B. & Br.	587
griseo-rubella Fr.	588
mordax Atk.	590
parkensis Fr.	587
pentagonospora Atk.	589
pirinoides sp. nov.	589
polita Fr.	587
rhodocyclioides Atk.	587
rhodocylix Fr.	587
roseoalbictrina Atk.	587
rusticoides Gill.	587
sphagnicola Pk.	587
vilis Fr.	587
Entoloma, genus.	29, 545
Entoloma, key to species.	547
Entoloma clypeatum Fr.	552
Entoloma cuspidatum Pk.	561
cyanum Pk.	551
dysthales Pk.	580
grande Pk.	548
grayanum Pk.	558
griseum Pk.	555
griseocyanum Fr.	551
jubatum Pk.	551
lividum Fr.	548
luteum Pk.	548
murinum Pk.	561
niderosum Fr. var.	554
pecktanum Burt.	560
prunuloides Fr.	547
rhodopodium Fr.	553
salmonium Pk.	560
scabrinellum Pk.	549
scabrosa Fr.	550
sericatum Britz.	554
sericellum Fr.	550
sericeum Fr.	556
sinuatum Fr.	549
speculum Fr.	550
strictius Pk.	557
variabilia Pk.	548
Entoloma, poisoning by.	853
F.	
"Fairy rings"	4
Families, key to.	26
Field, notes.	19
Fammula, genus.	29, 483
Fammula, key to species.	484
Fammula alnicola Fr.	491
carbonaria Fr.	488
flavida Fr.	490
fusa Fr.	488
gummosa Fr.	490
highlandensis Pk.	488
lenta Fr.	487
linulata Fr.	513
lubrica Fr.	486
parvifructa Pk.	487
polychora Berk.	484
sapinea Fr.	492
spumosa Fr.	489
Fruit body, definition of.	4
parts of.	4
Fungi, classes of.	26
outline of.	26
Galera, genus.	29, 492
Galera, key to species.	494

	Page
Galera antipus Lasch.	495
bulbifera sp. nov.	496
capillaripes Pk.	500
crispa Longyear.	498
cyanopes sp. nov.	500
hypnorum Fr.	502
laterifera (Lasch).	495
mniophila (Lasch).	500
ovalis Fr.	493, 494
paludosa Fr.	494
pellucida Fr.	494
plicatellus Pk.	501
pubescans Gill.	497
pygmaea-affinis Fr.	497
rufipes Pk.	494
sp.	497
spartea Fr.	499
stagnina Fr.	494
tenera Fr.	498
teneroides Pk.	499
Genera, key to.	27
Gills, definition of.	4, 7
Glossary of terms.	879
Gomphidius, genus.	30, 169
Gomphidius, key to species.	170
Gomphidius flavipes Pk.	172
furcatus Pk.	171
glutinosus Fr.	169
gracilis Berk.	171
maculatus Fr.	170
nigricand Pk.	170, 171
vinicolor Pk.	169
viscidus Fr.	169
Gyromitra esculenta, poisoning by.	861
H.	
Hebeloma, genus.	29, 468
Hebeloma, key to species.	470
Hebeloma albidulum Pk.	479
album Pk.	481
colvini Pk.	481
crustuliniforme Fr.	476
fastibite Fr.	472
firmum Fr.	470
glutinosum Fr.	471, 487
gregarium Pk.	473
hiemale Bres.	477
longicaudum Fr.	478
magnitimum Fr.	482
mesophaeum Fr.	473
neurophyllum Atk.	471, 479
pascuense Pk.	474
sarcophyllum Pk.	480
simile sp. nov.	479
sinapizans Fr.	475
sinuosum Fr.	476
sociale Pk.	471
syrjense Karst.	482
testaceum Fr.	471, 482
velatum Pk.	471
Hebeloma, poisoning by.	855
Heliomyces, genus.	28, 81
Heliomyces foetans Pat.	73
nigripes (Schw.) Morg.	81
pruinosisipes Pk. var. cerebrosus var. nov.	82
Herbarium, notes.	18
specimens.	18
Hygrophoreae.	169
Hygrophorus, genus.	27, 172
Hygrophorus, key to species.	175
Hygrophorus agathosmus Fr.	174
albipes Pk.	177
arbustus Fr.	186
aurens Fr.	182
basidiosus Pk.	177
borealis Pk.	192
var. subborealis var. nov.	192
bresadolae Quel.	183
burnhami Pk.	177
caerulescens B. & C.	195
ceraceus Fr.	193
chlorophanus Fr.	198
chrysodon Fr.	178
coccineus Fr.	196

	Page
Hygrophorus conicus Fr.	193
colemannianus Blox.	183
coloratus Pk.	174
cossus Fr.	178
eberneus Fr.	179
var. decipiens Pk.	177
var. unicolor Pk.	180
elegantulus Pk.	184
flavodiscus Frost.	189
foetens Phil.	201
fornicatus Fr.	183, 203
friesii Sacc.	187
fuliginus Frost.	183
fusco-albus var. occidentalis var. nov.	183
glutinifer Fr.	177
hypothecus Fr.	210
immutabilis Pk.	177
lactus Fr.	179
laricinus Pk.	188
laurae Morg.	177
leporinus Fr.	199
luridus B. & C.	177
marginatus Pk.	195
metapodius Pk.	195
miniatus Fr.	196
var. cantherellus Schw.	177
var. sphagnophilus Pk.	200
minutulus Pk.	174
nitidus B. & C. (non Fr.)	191
nitratus Fr.	184
niveus Fr.	194
olivaceo-albus Fr.	181
pallidus Pk.	194
paludosus Pk.	202
peckianus Howe.	190
peckii Atk.	190
pratensis Fr.	190
var. cinereus	202
var. pallidus	186
psitticinus Fr.	197
pudorinus Fr.	197
puniceus Fr.	177
var. flavescens Kauff.	179
ruber Pk.	185
rubropunctus Pk.	177
Russula (Fr.) nov. comb.	187
serotinus Pk.	181
sordidus Pk.	177
speciosus Pk.	177
sphaerosporus Pk.	195
subrufescens Pk.	203
subviolaceus Pk.	177
unquinosus Fr.	191
virgatulus Pk.	849
virginus Fr.	30, 254
Hygrophorus, poisoning by	255
Hypholoma, genus	259
Hypholoma, key to species	264
Hypholoma aggregatum Pk.	264
appendiculatum Fr.	255
candolleianum Fr.	265
capnoides Fr.	278
conatum Fr.	256
elongatum Pers.	255
epixanthum Fr.	266
fasciculare Fr.	266
hydrophilum Fr.-Rick.	263
hydrophilum Fr.-Sacc.	264
incertum Pk.	267
incertum var. sylvestrus var. nov.	259
instratum Britz.	258
lachrymabundum Fr.-Quel.	237
lactinum sp. nov.	261
peckianum Pk.	262
perplexum Pk.	267
populinum Britz.	279
rugocepalum Atk.	256
saccharinophilum Pk.	260
sarcocepalum Fr.	261
sublateritium Fr.	7
velutinum Fr.-Quel.	3
vinosum sp. nov.	29, 442
Hymenium, definition of	445
Hyphae, definition of	
Inocybe, genus	
Inocybe, key to species	

	Page
Inocybe albodisca Pk.	462
asterospora Quel.	462
bongardii Weim.	444
caesariata Fr.	448
calamistrata Fr.	447
calospora Quel.	450
carpta Bres.	461
cincinnati Fr.	443
comatella Pk.	465
cookei Bres.	458
curreyi Berk.	458
decipientoides Pk.	453
destricta Fr.	456
dulcamera Schw.	446
euthelus Fr.	460
eutheloides Pk.	459
fallax Pk.	462, 465
fastigiata Bres.	457
fibrosa Bres.	461
flocculosa Berk.	453
frumentacea Bres.	454
geophila Fr.	465
glaber sp. nov.	468
hirsuta Lasch.	443
hystrix Fr.	447
incarnata Bres.	444
infelix Pk.	452
poisoning by	855
infida Pk.	443, 465
poisoning by	854
lacera Fr.	452
lanatodisca sp. nov.	459
lanuginella Schroeter	449
lanuginosa Fr.-Bres.	449
leptophylla Atk.	449
lilacina Fr.	466
pyriodora Fr.	450
radiata Pk.	460
repanda Bres.	463
rigidipes Pk.	450
rimosa Fr.	455
scabella Fr.	466
scabella var. rufa	466
scaber Fr.	451
sindonia Fr.	464
squamosodisca Pk.	449
subdecurrans E. & E.	449
subochracea Pk.	464
subtomentosa Pk.	449
trechispora Berk.	467
trinti Wein.	444
umbratica Bres.	462
umbrina Bres.	461
unicolor Pk.	449
violaceifolia Pk.	445
Inocybe poisoning by	854
	K.
Keys, their use	23
	L.
Lamellae, definition of	4
Lactariae	83
Lactarius, genus	27, 83
Lactarius, key to species	86
Lactarius affinis Pk.	99
atroviridis Pk.	89
aurantiacus Fr.	110
boughtoni Pk.	108
brevipes Longyear	98
brevis Pk.	98
camphoratus Fr.	117
chrysorheus Fr.	96
cilicoides Fr.	91
cinereus Pk.	108
colorascens Pk.	110
controversus Fr.	94
corrugis Pk.	113
croceus Burl.	110
deceptivus Pk.	103
deliciosus Fr.	103
distanus Pk.	114
foetidus Pk.	115

	Page
Lactarius fuliginosus Fr.	105
gerardii Pk.	105
glycosmus Fr.	87
griseus Pk.	108
indigo Schw.	104
insulsus Fr.	98
isabellinus Burl.	111
helvus Fr.	106
helvus var. aquifolius Pk.	107
hygrophoroides B. & C.	114
hygginus Fr.	100
lignyotus Fr.	106
luteolus Pk.	115
maculatus Pk.	102
mucidus Burl.	109
oculatus Pk.-Burl.	116
parvus Pk.	111
perganenus Fr.	95
platyphyllus Pk.	99
piperatus Fr.	95
var. fragrans Burl.	95
pubescens Fr.	94
pyrogalus Fr.	96
regalis Pk.	99
resimus Fr.	99
rimosellus Pk.	117
rufus Fr.	107
scrobiculatus Fr.	90
sordidus Pk.	89
subdulcis Fr.	115
subpurpureus Pk.	102
theiogalus Fr.	97
tormentosus Fr.	91
poisoning by	849
trivialis Fr.	100
var. gracilis Pk.	101
var. maculatus Pk.	101
var. viridilactus var. nov.	101
turpis Fr.	88
uvidus Fr.	101
poisoning by	850
varius Pk.	112
vellerius Fr.	92
vietus Fr.	109
volemus Fr.	112
zonarius Fr.	99
Lactarius, poisoning by	845
Lentinus, genus	28, 49
Lentinus, key to species	51
Lentinus americana Pk.	51
chrysopelos B. & C.	50, 778
cochleatus Fr.	55
haematopus Berk.	54
lecomtei Schw.	46
lepideus Fr.	13, 53
microsperma Pk.	55
obconicus Pk.	51
omphalodes Fr.	54
spretus Pk.	51, 53
suavissimus Pk.	51
sulcatus Berk.	51
tigrinus Fr.	51
umbilicatus Pk.	54
underwoodii Pk.	51
ursinus Fr.	56
vulpinus Fr.	56
Lentodium squamulosum Morg.	52
Lepiota, genus	4, 27, 625
Lepiota, key to species	626
Lepiota acerina Pk.	635
acutaesquamosa Fr.	633
adnatifolia Pk.	637
alluvium Pk.	642
americana Pk.	645
amianthina Fr.	628, 637
angustana Britz.	642
arenicola Pk.	628
asperula Atk.	635
badhami Berk.	646
boudieri Bres.	636
caepestipes Fr.	640
caerulescens Pk.	627
calocephalus Atk.	628
charcarias Fr.	626
clypeolaria Fr.	631

	Page
Lepiota cristata Fr.	641
cristatataellus Pk.	643
cyanozonata Longyear	627
delicata Fr.	631
ecitodora Atk.	627
eritophora Pk.	628
excoriata Fr.	647
felina Fr.	633
fischerii sp. nov.	630
friesii Lasch.	634
geniculospora Atk.	677
glodermata Fr.	629
gracilis Pk.	628
granulosa Fr.	638
granosa Morg.	636
haematosperma Fr.-Bres.	262
helvola Bres.	626
lenticularis Fr.	630
illimita Fr.	629
incarnata Clem.	631
metulaespora	632
miamensis Morg.	642
morganii Pk.	15, 644
poisoning by	844
naucina Fr.	15, 646
naucinoides Pk.	647
oblita Pk.	631
persoonii Fr.	630
procera Fr.	643
pulveracea Pk.	638
purpureoconia Atk.	627
pusillomyces Pk.	639
rubrotincta Pk.	640
rufescens Morg.	631
rugoso-reticulata Lor.	636
rugulosa Pk.	627
seminuda Fr.	639
Leptonia, genus	29, 571
Leptonia, key to species	572
Leptonia aeruginosa Pk.	572
anatina Fr.	579
asprella Fr.	578
euchroa Fr.	577
formosa Fr.	576
grisea Pk.	572
incana Fr.	572
lampropoda Fr.	574
placida Fr.	573
rosea Longyear	575
serrulata Fr.	577
seticeps Atk.	575
solsitialis Fr.	572
subserulata Pk.	572
transformata Pk.	572
undulatella Pk.	572
Lucosporae	205
M.	
Marasmiaceae	41
Marasmius, genus	28, 57
Marasmius, key to species	59
Marasmius alliaceus Fr.	68
androsaceus Fr.	79
bellipes Morg.	66
calopus Fr.	72
campanulatus Pk.	77
candidus Fr.	74
capillaris Morg.	80
caricicola Kauff.	74
chordalis Fr.	76
cohaerens Fr.-Bres.	75
decurrans Pk.	71
delectans Morg.	67
elongatipes Pk.	75
epiphyllus Fr.	80
erythropus Fr.	70
fagineus Morg.	65
felix Morg.	77
foetidus Fr.	73
fuscoauripurea Fr.	69
glabellus Pk.	66
graminum Libert.	79
instititus Fr.	80
leptopus Pk.	73

	Page
Marasmius, longipes Pk	76
magnisporus Murr	74
olneyi B. & C	73
oreades Fr	15, 61
papillatus Pk	76
perforans	80, 78
peronatus Fr	62
polyphyllus Pk	68
prasiomius Fr	68
pulcherripes Pk	66
resinosus Pk	71
rotula Fr	78
scorodonius Fr	72
semihirtipes Pk	67
siccus Schw	65
spongiosus B. & C	65
subnudus Ellis-Pk	64
urens Fr	63
varicosus Fr	69
velutipes B. & C	70
viticola B. & C	64, 65
Marasmius, poisoning by	851
Melanosporae	205, 206
Mushrooms, collecting of	16
cultivation	21
edibility of	17
parts of	4
poisoning by	825, 863
preserving of	16
Mycelium, definition of	3
Mycena, genus	28, 778
Mycena, key to species	780
Mycena acicula Fr	812
adonis Fr	812
alcalina Fr	803
ammoniaca Fr	804
atroalba Fr	807
atroalboides Pk	808
atrocyaneus Fr	808
bryophila Vogl	802
calorhiza Bres	803
capillaripes Pk	792
clavicularis Fr	788
var. alba Pk	788
var. filipes Pk	788
var. luteipes var. nov	788
cohaerens Fr	75
collariata Fr	810
consimilis Cke	806
constans Pk	804, 806
corticola Fr	793
crystallina Pk	789
cyaneobasis Pk	802
cyanothrix Atk	810
debilis Fr	783
denticulata Pk	792
dissiliens Fr	806
echinipes Fr	790
epipterygia Fr	787
excisa Fr	799
filopes Fr	783
galericulata Fr	797
var. calopus Fr	797
haematopa Fr	783
hemispherica Pk	807
hiemalis Fr	793
immaculata Pk	796
inclinata Fr	798
iris Berk	803
juncicola Fr	812
lasiosperma Bres	801
leajana Berk	785
leptocephala Fr	806
marginella Fr	803
metata Fr	805
minutula Pk	795
parabolica Fr	800
pellanthina Fr	790
polygramma Fr, var. albida Kauf	801
praelonga Pk	809
prolifera Fr	799
pterigenus Fr	812
pulcherrima Pk	811
pura Fr	794
purpureofusca Fr	792

	Page
Mycena, raeborhiza (Lasch.)	802
receptibilis Britz	802
rosella Fr	791
rubromarginata Fr	792
sanguinolenta Fr	784
setosa Fr	794
stannea Fr	807
stipularis Fr	812
stylobates Fr	789
subincarnata Pk	811
sudora Fr	801
viscosa Maire	788
vitilis Fr	783
vulgaris Fr	786
Mycorrhiza, definition of	10
N.	
Naucoria, genus	29, 508
Naucoria, key to species	509
Naucoria bellula Pk	513
Naucoria centuncula Fr	511
cidaris Fr	511
horizontalis Fr	511
lignicola Pk	512
nimbosa Fr	510
pediades Fr	515
platysperma Pk	514
semiorbicularis Fr	513
siparia Fr	515
tabacina Fr	515
triscopoda Fr	512
verracti Fr	514
Nolanea, genus	29, 579
Nolanea, key to species	580
Nolanea babingtonii Berk	583
Nolanea caelestina var. violacea Kauf	585
conica Pk	584
cruenta Quel	586
dysthales Pk-Atk	580
fuscogrisella Pk	585
mammosa Fr	583
nodospora Atk	580
papillata Bres	584
pascua Fr	581
stauospora Bres	582
versatilis Fr	582
Nomenclature	24
Nyctalis, genus	27, 31
Nyctalis asterosphora	32
O.	
Ochrospora	205
Omphalia, genus	28, 812
Omphalia, key to species	814
Omphalia albidula Pk	820
Omphalia caespitosa Bolt	824
caespitosa Bolt	823
campanella Fr	818
epichysium Fr	821
fibula Fr	822
fibuloides Pk	819
gerardiana Pk	820
gracillima Fr	817
hepatica Fr	816
lilacifolia Pk	816
lilacina Laest	816
luteola Pk	817
muralis Fr	815
olivaria Pk	818
anisa Fr	820
papillata Pk	816
pyxidata Fr	817
rugodisca Pk	778
scabriuscula Pk	822
schwartzii Fr	815
scyphiformis Fr	814
scyphoides Fr	819
sphagnophila Berk	820
stellata Fr	816
subclavata Pk	824
umbratilis Fr	824

## P.

	Page
Panaeolus, genus	30, 228
Panaeolus campanulatus Fr.	230
poisoning by	858
epimyces Pk.	253
papilionaceus Fr.	231
poisoning by	859
retirugis Fr.	229
poisoning by	858
solidipes Pk.	228
sp.	231
sphinctrinus Fr.	232
Panus, genus	28, 43
Panus, key to species	44
Panus albotomentosus Cke. & Mass.	45
angustatus Berk.	48
betulinus Pk.	45
conchatus Fr.	447
dealbatus Pk.	45
dorsalis Bosc.	45
laevis B. & C.	44, 46
nigrifolius Pk.	45
operculatus B. & C.	44
rudis Fr.	46
salicinus Pk.	49
stipticus Fr.	48
strigosus B. & C.	45
torulosus Fr.	47
Partial veil, definition of	4, 7
Paxillus, genus	28, 284
Paxillus, key to species	285
Paxillus atrotomentosus Fr.	287
corrugatus Atk.	288
extrenuatus Fr.	721
involutus Fr.	286
lepista Fr.	284
panuoides Fr.	288
rhodoxanthus Schw.	285
Phollota, genus	4, 28, 289
Phollota, key to species	289
Phollota acericola Pk.	309
adiposa Fr.	305
aegerita Fr.	296
aeruginosa Pk.	308
aggregata Pk.	297
aggericola Pk.	297
var. retirugis Pk.	298
albocrenulata Pk.	301
autumnalis Pk.	311, 312
poisoning by	854
blattaria Fr.	313
caperata Fr.	294
cerasina Pk.	291
comosa Fr.	291
confragosa Fr.	310
curvipes Fr.	302
destruens Fr. -Bres.	298
detersibilis Pk.	313
discolor Pk.	310
dura Bolt.	293, 294
duroides Pk.	292
erinacella Pk.	303
flammans Fr.	306
fulvosquamosa Pk.	302
heteroclita Fr.	291
howeana Pk.	297
indecens Pk.	298
johnsoniana Pk. -Atk.	295
limonella Pk.	290, 307
lucifera Bres.	307
lutea Pk.	291
luteofolia Pk.	307
marginata Fr.	311
marginella Pk.	312
minima Pk.	291
muricata Fr.	304
mutabilis Fr.	291
mycenoides Fr.	314
ornella Pk.	291
praecox Fr.	293
var. sylvestris Pk.	293
radicosa Fr.	249
rugosa Pk.	312
spectabilis Fr.	304
squamosa Fr.	292, 300

	Page
Phollota squarrosoides Pk.	299
temnophylla Pk.	294
togularis Fr.	292, 313
tuberculosa Fr.	303
unicolor (Fl. D.) Fr.	311
vermiflua Pk.	294
verruculosa (Lasch) Fr.	300
Phollota, poisoning by	854
Photography of mushrooms	20
Pileus, definition of	4, 5
Pilosace algeriensis Fr. -Quel.	254
Pleurotus, genus	25, 656
Pleurotus, key to species	658
Pleurotus acerosus Fr.	670
albolanatus Pk. sp. nov.	45, 672
applicatus Fr.	674
atrocaeruleus Fr. var. griseus Pk.	673
atropellitus Pk.	675
borealis	657
candidissimus B. & C.	670
circinatus Fr.	666
corticatus	657
craspedius Fr.	659
cyphellaeformis Berk.	658
dryinus	657
elongatipes Pk.	661
euosmus	657
fimbriatus var. regularis var. nov.	667
lignatilis Fr.	666
limpidus Fr.	670
mastrucatus Fr.	673
mitis Fr.	670
nephretus Ell.	671
niger Schw.	656, 659, 675
ostreatus Fr.	663
petaloides Fr.	49, 669
porrigens Fr.	671
sapidus Kalchb.	665
septicus Fr.	671
serotinus Fr.	668
spatulatus Fr. -Pk.	669
striatulus Fr.	656, 659, 675
subareolatus Fr.	664
subpalmatius Fr.	662
sulfuroides Pk.	661
tremulus Fr.	659
ulmarius Fr.	659
umbonatus Pk.	658
Plicatura alni (Pk.)	42
Pluteolus, genus	29, 505
Pluteolus, key to species	505
Pluteolus aleuratus var. gracilis Pk.	506
coprophilus Pk.	506
expansus Pk.	507
var. terrestris Pk.	507
reticulatus Fr.	502, 508
Pluteus, genus	29, 535
Pluteus, key to species	536
Pluteus admirabilis Pk.	544
calocephalus Atk.	543
cervinus Fr.	537
var. alba Pk.	538
var. petastatus Fr.	538
var. viscosus Lloyd	538
chrysophaeus Fr.	537
ephebius Fr. var.	539
var. drephanophyllus Schultz	540
flavofulgineus Atk.	536
granularis Pk.	541
var. umbrosellus Atk.	542
intermedius var. nov.	542
leoninus Pk.	545
longistriatus Pk.	543
nanus Fr.	542
var. lutescens Fr.	543
pellitus Fr.	539
phleboporus	536
roseocandidus Atk.	541
salicinus Fr. var.	539
stercorarius Pk.	536
sterilomarginatus Pk.	536
tomentosulus Pk.	540
umbrosus Fr.	538
Poisoning by mushrooms, chapter on	825
statistics on	827

	Page
Poisonous mushrooms	828
<i>Amanita bisporiger</i>	828
<i>citrina</i>	836
<i>chlorinosoma</i>	837
<i>cothurnata</i>	844
<i>crenulata</i>	836
<i>frostiana</i>	837
<i>mappa</i>	836
<i>morrisii</i>	837
<i>muscaria</i>	837
constituents of poison of	839
poisoning by	837
clinical features of	838
treatment of	841
<i>pantherina</i>	843
<i>phalloides</i>	828
constituents of poison of	833
poisoning by	828
treatment of	832
<i>porphyria</i>	836
<i>radicata</i>	836
<i>spretia</i>	836
<i>strobiliformis</i>	836
<i>verna</i>	828
<i>virosa</i>	828
<i>Boletus luridus</i>	860
<i>satanus</i>	860
spp.	860-61
<i>Clitocybe illudens</i>	846
<i>morbifera</i>	848
<i>sudorifica</i>	847
<i>Coprinus</i> spp.	858
<i>Entoloma cuspidatum</i>	853
<i>lividum</i>	853
<i>nidorosum</i>	853
<i>rhodopodium</i>	853
<i>salmonium</i>	853
<i>sinuatum</i>	853
<i>strictius</i>	853
<i>Gyromitra esculenta</i>	861-62
<i>Hebeloma fastibile</i>	856
<i>sinapizans</i>	855
<i>Hygrophorus conicus</i>	849
<i>Hypholoma instratum</i>	857
<i>Inocybe decipiens</i>	855
<i>infelix</i>	855
<i>infida</i>	854
<i>rimosa</i>	856
<i>Lactarius</i> spp.	850
<i>torminosus</i>	849
<i>Lepiota morgani</i>	944
<i>Panaeolus</i> spp.	858
<i>Pholiota autumnalis</i>	854
<i>Pleurotus olearius</i>	847
<i>Psilocybe cernua</i>	857
<i>Russula emetica</i>	850
spp.	851
<i>Tricholoma venenatum</i>	846
Poisonous principles of <i>Amanitas</i>	843
<i>Psalliota</i> , genus	20, 232
<i>Psalliota</i> , key to species	233
<i>Psalliota abruptibulba</i> Pk.	237
<i>arvensis</i> Fr.	13, 15, 236
<i>augusta</i> Fr.	239
<i>campestris</i> Fr.	13, 15, 240
<i>comtula</i> Fr.	244
<i>cretacea</i> Fr.	234
<i>cretacea</i> Atk.	234
<i>diminutiva</i> Pk.	245
<i>echinata</i> Fr.	245, 262
<i>haemorrhodaria</i> Fr.	243
<i>halophila</i> Pk.	243
<i>micromegetha</i> Pk.	243
<i>perrara</i> Bres.	239, 242
<i>placomycetes</i> Pk.	238
<i>pusilla</i> Pk.	244
<i>rodmani</i> Pk.	235
<i>russiophylla</i> Lasch.	244
<i>sagata</i> Fr.	244
<i>subrufescens</i> Pk.	239
<i>sylvatica</i> Fr.	242
<i>villatica</i> Fr.-Bres.	242
<i>Psathyra</i> , genus	30, 268
<i>Psathyra</i> , key to species	268

	Page
<i>Psathyra conopilea</i> Fr.	269
<i>currugis</i> Fr.	269
<i>elata</i> Mass.	271
<i>microsperma</i> Pk.	270
<i>obtusata</i> Fr.	270
<i>persimplex</i> Britz.	271
<i>semivestita</i> B. & Br.	269
<i>umbonata</i> Pk.	271
<i>vestita</i> Pk.	271
<i>Psathyrella</i> , genus	30, 226
<i>Psathyrella crenata</i> Fr.	227
<i>disseminata</i> Fr.	269
<i>subatrata</i> Fr.	272
<i>Psilocybe</i> , genus	30, 272
<i>Psilocybe</i> , key to species	281
<i>Psilocybe agrariella</i> Atk.	281
<i>ammophila</i> Mont.	281
<i>arenulina</i> Pk.	277
<i>atrobrunnea</i> Fr.	275
<i>atrorufa</i> Fr.	275
<i>bullacea</i> Fr.	276
<i>canofaciens</i> Cke.	280
<i>cernua</i> Fr.	284
<i>conissans</i> Pk.	274
<i>coprinophila</i> Fr.	278
<i>ericaea</i> Fr.	15, 283
<i>foeniseii</i> Fr.	279
<i>larga</i> sp. nov.	274
<i>merdaria</i> Fr.	280
<i>murcida</i> Fr.	275
<i>physaloides</i> Fr.	267
<i>polycephala</i> Paul.-Sacc.	267, 279
<i>spadicea</i> Fr.	282
<i>submaculata</i> Atk.	275
<i>subviscida</i> Pk.	277
<i>uda</i> Fr.-Batt.	277
R.	
Rhizomorph, definition of	3
Rhodosporae	205
<i>Russula</i> , genus	27, 118
<i>Russula</i> , key to species	121
<i>Russula abietina</i> Pk.	166
<i>adusta</i> Fr.	128
<i>aeruginea</i> Lindb (non-Fr.)	144
<i>albella</i> Pk.	154
<i>albida</i> Pk.	158
<i>albidula</i> Pk.	154
<i>albonigra</i> (Kromb.)	129
<i>alutacea</i> Fr.	150
<i>amygdaloides</i> sp. nov.	162
<i>anomala</i> Pk.	159
<i>atropurpurea</i> Kromb-Maire (non Pk.)	142
<i>atropurpurea</i> Pk.	145
<i>aurata</i> Fr.	124
<i>aurantialutea</i> Kauff.	157
<i>azurea</i> Bres.	141
<i>batlae</i> Quel.	163
<i>borealis</i> Kauff.	150
<i>brevipes</i> Pk.	125
<i>caerulea</i> Pers.	165
<i>chamaeleontina</i> Fr.	165
<i>chloroides</i> Bres.	126
<i>claroflava</i> Grove	158
<i>compacta</i> Frost	129
<i>consobrina</i> Fr.	138
<i>constans</i> Karst.	148
<i>crustosa</i> Pk.	132
<i>cutefracta</i> Cke.	149
<i>cyanoxantha</i> Fr.	141
<i>decolorans</i> Fr.	146
var. <i>rubriceps</i> Kauff.	147
<i>delica</i> Fr.	15, 93, 125
var. <i>brevipes</i> Pk.	125
<i>densifolia</i> Secr.	127
<i>depallens</i> Fr.	147
<i>drimeia</i> Cke.	151
<i>emetica</i> Fr.	830, 850
poisoning by	850
var. <i>gregaria</i> Kauff.	152
<i>expallens</i> Fr.	143
<i>fallax</i> Cke.	154
<i>fellea</i> Pk.	158
<i>flava</i> Rom.	147

Russula flavida Frost	Page
flaviceps Pk	123
foetens Fr.	167
foetentula Pk	136
fragilis Fr.	136
furcata Fr.	153
graminicolor Quel	140
granulata Pk	144
granulosa Cke	133
graveolens Rom.	133, 158
heterophylla Fr.	145
incarnata Morg.	144
integra Fr.	130
lactea Fr.	161
lepida Fr. (non Bres.)	126
liacea Quel	133
linnaei Fr.	122
lutea Fr.	143
magnifica Pk	167
mariae Pk	121
melliolens Quel	143
migriscentipes Pk	162
nigricans Fr.	148
nitida Fr.	32, 126
obscura Rom.	163
ochracea Fr.	148
ochroleuca Fr.	158
ochroleucoides sp. nov.	133
ochrophylla Pk	132
olivacea Fr.	124, 150, 151
olivascens Fr.	145
palustris Fr.	144
pectinata Fr.	157
pectinatoides Pk	137
psuedointegra A. & G.	137
puellaris Fr.	160
pulverulenta Pk	169
punctata Gill	134
purpurea Gill	139
purpurina Q. & S.	143
quelletii Fr.	159
romellii Maire	143
rosacea Fr.	162
roselipes Sec.-Bres	155
rubra Fr.	163
rugulosa Pk	142
rubescens Beards.	152
sanguinea Fr.	149
sardonis Fr.	155
sericeo-nitens Kauff.	156
simillima Pk	161
sordida Pk	158
sororia Fr.	129
sphagnophila Kauff.	138
squalida Pk	165
subdepallens Pk	145
subfoetens Smith	159
subpunctata sp. nov.	137
subsordida Pk	139
tenuiceps Kauff.	129
uncialis Pk	156
variata Bann. & Pk	160
vesca Fr.-Bres	140
veternosa Fr.	138
vinosa Lindb.	155
virescens Fr.	148
vitellina Gill	131
xerampelina Fr.	167
Russula, poisoning by	144
	850

S.

Schizophyllum, genus	28, 42
Schizophyllum commune Fr.	43
Sclerotia, definition of	3
"Spawn," definition of	3
Species, arrangement of	24
Spores, definition of	7
Sterigma, definition of	7
Stipe, definition of	4
Stropharia, genus	4, 29, 246
Stropharia, key to species	247

Stropharia aeruginosa Fr.	Page
albonitens Fr.	247
battaraea Fr.	250
bilemellata Pk	247
caputmedusae Fr.	251
coprinophila Atk.	247
coronilla Bres	253
cotonea Fr.	250
depiata Fr.	259
epimyces Pk.-Atk.	249
johsoniana Pk.	253
melasperma Fr.	295
paradoxa P. Henn.	251
scobinaceum Fr.	253
semiglobata Fr.	247
stercoraria Fr.	252
umbonatescens Pk	251
ventricosa Mass.	252
	248

T.

Toxic principles of Amanitas	843
Tricholoma, genus	28, 675
Tricholoma, key to species	679
Tricholoma acerbum Fr.	705
acre Pk	694
albellum Fr.	680
albiflavidum Pk	764
album Fr.	704
aurantia Fr.	651
boreale Fr.	704
brevipes Fr.	679, 712
carneum Fr.	702
cerinum Fr.	681
chrysenteroides Pk	701
cinerascens Fr.	711
columbeta Fr.	692
conglobatum Fr.	704
coryphaeum Fr.	684
davisiae Pk	681
decorosum Pk	682
equestre Fr.	682
fallax Pk	681
flavescens Pk	680
flavobrunneum Fr.	679, 687
frumentaceum Fr.	687
fuliginum Pk	698
fumescens Pk	697
fumosiluteum Pk	707
gambosum Fr.	682
grammopodium Fr.	679, 712
grande Pk	680
grave Pk	681
humile Pk	680
imbricatum Fr.	693
intermedium Pk	679, 684
ionides Fr.	681, 709
lascivium Fr.	680
laterarium Pk	706
laticeps sp. nov.	700
leucocephaloides Pk	713
leucocephalum Fr.	706
loricatum Fr.	734
lugubre Pk	700
melaleucum Fr.	713
multipunctum Pk	724
murinaeum Fr.	695
nobile Pk	691
nudum Fr.	709
odorum Pk	702
paeonium Pk	703
pallidum Pk	699
panoecium var. caespitosum Bres	711
patulum Fr.	682
peckii Howe	651
personatum Fr.	707
pessundatum Fr.	688
portentosum Fr.	684
var. centrale Pk	685
putidum Fr.	679
radicatum Pk	681
rancidulum Bann	712
resplendens Fr.	187, 686
rimosum Pk	679
Russula Fr.	185

	Page
Tricholoma rutilans Fr. ....	689
saponaceum Fr. ....	699
scalpturatum Fr. ....	696
sejunctum Fr. ....	683
serratifolium Pk. ....	680, 692
silvaticum Pk. ....	680
sordidum Fr. ....	714
squarulosum Bres. ....	696
subacutum Pk. ....	681
subluteum Pk. ....	681
submaculatum Pk. ....	682, 768
sulphurascens Bres. ....	701
sulphureum Pk. ....	681
terreolens Pk. ....	680
terreum Fr. ....	696
terriferum Pk. ....	685
transmutans Fr. ....	687
tricolor Pk. ....	694
tumidum Fr. ....	710
unifacium Pk. ....	703
ustale Fr. ....	688
vaccinum Fr. ....	693
variegatum Fr. ....	690, 725
venenata Atk. ....	690
virgatum Fr. ....	681
viscosum Pk. ....	679
Tricholoma, poisoning by. ....	846
Trogia, genus. ....	28, 41
Trogia alni Pk. ....	42
crispa Fr. ....	41
Tubaria genus. ....	29
U.	
Universal veil, definition of. ....	4
V.	
Volva, definition of. ....	4
Volvaria, genus. ....	4, 29, 526
Volvaria, key to species. ....	527
Volvaria bombycina Fr. ....	527
gloiocephala Fr. ....	529
hypopithys Fr. ....	531
loveriana Berk. ....	32, 726
murinella Quel. ....	527, 531
parvula Fr. ....	532
pubescentipes Pk. ....	530
pusilla Pers.-Fr. ....	531
speciosa Fr. ....	528
striatula Pk. ....	527
umbonata Pk. ....	530
volvacea Fr. ....	527