

with sub-metallic lustre, which color is due to a large proportion of martite granules mingled with the mass. Next above this rock another belt of a finer grained and lighter colored reddish quartzite follows, equalling the former in thickness, and then a high bluff of dark green chlorito-hydro-micaceous schists inclosing irregular concretionary seams of a siliceous lime-rock, ascends closely adjoining the quartzite beds, and, as it appears, conformably superimposed on them. It forms the south part of the summit elevation, which is formed of a large granite belt adjoining the schists, and occupying the centre and the north slope of this ridge. The thickness of the chloritic schistose belt amounts to about 70 feet. The next following hill is formed of chloritic and dioritic schists, which have a northern dip; farther on, the line descends to a somewhat lower level, and passes over several knobs of crystalline diorite; but arrived at the quarter-post of the north line we find the flaggy beds of the iron formation beneath our feet. A very short distance east or south-east of this spot are the so-called old Tilden mines, and north-west of the place we find the Foster mines; both of them are at present abandoned. The above-described exposure of the quartzite formation, wedged in between two granite belts and overlaid by schists, which, according to their lithological character, belong to a lower horizon, must present the ledges in an overturned position. We should expect, according to the structural arrangement in the Gribben mines, to see the iron formation represented in this exposure between the upper ledges of the quartzite and the chloritic schists inclosing calcareous bands, which latter are very similar to the schists forming the foot-wall of the ore-formation in the Gribben mines, but I could not discover a sign of its being there; and north of this locality, where the iron-formation is very extensively exposed, directly reposing on rock-beds similar to the chloritic schists of the bluff, we cannot find a trace of the quartzite formation incumbent on the ores. Farther west, no more quartzite is seen within the examined district, which could with propriety be considered as a continuation of this large body of rocks, followed without interruption from the shore of Lake Superior, near Harvey's furnace, to the western town-line of R. 26, Town. 47.

CHAPTER IV.

IRON GROUP.

BEFORE I entered the special description of the different groups into which I subdivided the Huronian rock-series, I made the remark that, owing to the circumstance that I confined myself to the next environs of Marquette when I began my investigations, I was led into a misconception of the true order of sequence between the beds by the local conditions existing there, but that subsequent examination of the more western districts convinced me of being in error when I considered the quartzite formation as older than the iron formation.

The first reposes near Marquette, and also in the Teal Lake range, in apparent conformity on the dioritic rock-series, without intervention of any deposits which could be taken as an analogon or equivalent of the iron-ore group, developed a very short distance south of the Teal Lake range in so great a thickness that one can hardly imagine how so large a succession of beds should at once be eliminated from the succession in the series in a locality so near to the other. This fact, supported by several rock-exposures, in which the ore-bearing beds seem to be actually incumbent on the Teal Lake quartzites, if the conformable dip of the strata is taken as a proof of their regular succession, induced me to consider the ore-formation of the Negaunee mines as much younger than the Teal Lake quartzites, and, as a matter of course, the quartzites visibly reposing on the ore-beds in all the different mines of the district, I then thought to be still younger deposits totally different from the other quartzites, in accordance with the opinions of my predecessor in the examination of this district, Major T. B. Brooks.

According to the information subsequently acquired by me,

during the study of other localities, I had to change my views; and I hope to prove in the following pages their correctness—that is to say, I will demonstrate the identity of the so-called upper and lower quartzite formations, and the older age of the ore-bearing rock-series than these deposits have.

The conformable superposition of the quartzite formation on the iron-bearing rock-belt of the Cascade range, and the deposition of the latter on schistose beds analogous with those of the dioritic group, which on their part rest on the granite formation, is sufficiently described in the previous chapter, and needs only to be remembered here.

Major Brooks could not help making the same observation; but prepossessed by the idea that the ore-belt of the Negaunee mines was a much later deposit than the quartzite formation of the Teal Lake range, and at the same time acknowledging the identity of these quartzites with the quartzites overlying the Cascade ores, he solved the dilemma by the assumption of two ore-bearing horizons, one below, the other above the quartzite, although he found it strange to see the Cascade ore-belt not represented on the north side of the Teal Lake range, where it should be expected, according to the synclinal arrangement of the rock-crust.

The surface rock of the environs of Negaunee and Ishpeming is almost exclusively formed of the iron-bearing rock-series, but much of it is covered by drift deposits.

The strata are in an extremely disturbed condition, folded and distorted in every possible way, usually without causing a rupture of the beds; but in some other instances the laminated banded seams composing the thicker ledges have ruptured, and the ends often came in a faulted position to each other, and were so recemented by the siliceous ground-mass, or else the cracks of the shattered rock were filled out with hydrated oxide of iron in various form, from the compact aphanitic porcelain-like condition, to the coarsely crystalline, or the fibrous structure of grape-ore, or in porous ochraceous masses. These disturbed beds lie, in every instance, directly, but very often inconformably on chlorito-hydro-micaceous schists, or on crystalline dioritic masses, which are constant associates of these chloritic schists, or sometimes dioritic schists, as hornblende and chlorite substitute each other, or are both components of them.

Overlooking the extremely plicated and corrugated condition of the strata, they form, considered in their totality, a synclinal basin hemmed in between dioritic ridges. A northern ridge of diorite hills commences north of the last houses of Negaunee village, in the centre of the N.W. quarter of Sect. 6, Town. 47, R. 26, and extends in due western direction to the west line of Sect. 2, Town. 47, R. 27; the summit of this narrow ridge is cut up into rounded knobs, severed from each other by drift-covered intervals.

The diorite hills forming the south border of the synclinal, much higher than the northern, are not connected into one continuous ridge; they are parts of a large dioritic mountain mass, covering over 8 square miles of surface on the south side of Negaunee and Ishpeming, and cut up by deep valleys and ravines into a large number of smaller sub-parallel ridges and knobs, which trend from east-south-east to west-north-west, and many isolated diorite knobs, found dispersed over the inner part of the synclinal trough, actually are parts of this large southern body of diorite.

The structure of the ore-formation in the vicinity of Negaunee is well exhibited by the Jackson mine, which principally opens the upper part of the formation, while the lower beds, occupying the slope of the high diorite hills on the south side, are the mining ground for a large number of other companies, which have riddled the surface with hundreds of test-pits, besides the numerous larger excavations from which the ore is taken. The Jackson mines are situated on the north side of a short isolated ridge of from 100 to 150 feet elevation, which extends from east to west across the eastern two thirds of the north part of the south half of Sect. 1, Town. 47, R. 27. This ridge is completely formed of thick ledges of a banded, very compact rock of quartzose-ferruginous composition. The banded appearance of the rock is produced by an alternation of narrow, inseparably united seams, of which one kind consists of a jaspery quartz-mass, generally bright red colored by iron oxide; the other likewise quartzose seams of the rock are dark blackish, or else of a metallic steel color, by impregnation with a large proportion of granules of martite, or with scaly micaceous ore particles of metallic lustre. Oftentimes these ferruginous seams have the quality of a rich ore-mass, with very little siliceous matter in it. Interlaminated with these mixed rock-ledges are belts of pure iron-ore, in compact granular masses, or in the

schistose form of the so-called specular slate-ore, amounting to greater thickness, available for the miner, but found to be quite irregular in their extension, widening locally into large pockets, twenty, thirty, and more feet wide, or wedging out in another place to almost total disappearance, and some distance farther on perhaps expanding again into a seam of valuable thickness. Associated with these ore-seams are usually belts of argillitic schists called soapstone by the miner for its talcy lubricity; and the occurrence of these schists in a test-pit is considered by him as a sure indication of an ore-deposit somewhere near by. They consist of a fine scaly or granular soft aluminous silicate, readily cut with a knife, which, colorless in itself, is tinged by red oxide of iron, either uniformly and in various degrees of intensity, or the pale yellowish ground-mass is dotted with irregular large streaky blotches of red, or other beds of red color are found regularly speckled by white dots of the size of a lentil, which I cannot better describe than by comparing their aspect with the dotted plumage of a Guinea-hen. Such peculiarly speckled schists are met with in many other mines besides the Jackson. Other schists of similar argillitic and partly hydro-micaceous composition, of a gray sub-metallic color, are more or less densely disseminated with small octahedric crystals of martite, which, however, are also not missing in the red-colored layers. The ore of the Jackson mines is taken from different belts, of which nearly every one furnishes a different kind of ore, but the chaotically plicated and dislocated condition of the rock-beds rarely allows recognition of the exact relative position which one belt holds to the other; the miner pursuing a belt often finds it wedging out or abruptly cut off, and has to take his chances by drifting ahead in some direction to meet the other end again, or to find another different seam.

A very rich seam of ore is almost invariably found on top of this jasper-banded rock-series, immediately beneath the quartzites which form the terminal strata in all these exposures. This upper ore-belt is almost regularly brecciated in its upper part, and the same is true of the lower quartzite beds, which often are a mixture of ore-fragments with quartzite pieces held together by an arenaceous cement. As this is the case in nearly all the mines of the district, we must suggest that great disturbances, of not only a local extent, must have occurred at the end of this era of iron

sediments. At the Jackson mine these quartzites, terminating the succession of rock-beds, are only represented by a small series of dark iron-colored ledges. The general dip of the formation is northward, and the south side of the described ridge is formed of vertical rock-walls exhibiting the edges of the steeply inclined beds. Near the west end of the Jackson mines several small knobs of diorite, associated with schists belonging to this rock-group, project, closely surrounded by the jasper-banded rock-beds in a vertical position, and evidently corrugated by the intrusion of these masses. In the valley south of the bluffs of Jackson mine hill, and on the slope of the high diorite ridge on the south side of it, exposures of more thinly laminated ferruginous siliceous rock-beds follow, amounting to a belt of at least a thousand feet in thickness. The largest portion of this rock-series is made up of regularly alternating bands, from a quarter of an inch to one inch in thickness, of a paler or darker drab-colored jasper, and of similar seams of ferruginous matter in different degrees of purity, and of different molecular structure. Sometimes these seams, intimately coherent with the jasper-seams, are an almost pure granular martite of metallic lustre, but usually a dull blackish red impure siliceous iron-ore with a sandy fracture composes them, or else they are formed of a bluish black aphanitic oxide, hard, but quite brittle, breaking with very smooth conchoidal fracture.

Another part of this series, interstratified with the jasper-banded beds, consists of argillitic rock-seams, all impregnated with considerable proportions of red oxide of iron; but there are also rich ore-seams intercalated at different horizons of this large succession of beds, which make it one of the most productive fields for the miner. These ore-deposits are not regular sedimentary layers, originally formed of iron oxide in this state of purity, but are evidently the product of decomposition of the impurer mixed ferruginous ledges by percolating water, leaching out the siliceous matter, and replacing it by deposition of oxide of iron held in solution; the ore-seams are therefore very irregular, narrow in one place and widening in another into large pockets 40 and 50 feet in diameter, or even larger. They consist generally of a crummy, porous, dark brown or blackish mass of dull earthy aspect, readily dug out with the pickaxe and shovel, composed partly of not-hydrated, partly of hydrated oxide. Within these seams are more

compact concretionary masses, partly of small nodular size, partly in large bulky seams, so-called hog-backs, consisting almost exclusively of hydrated oxide of the radiated fibrous structure of grape-ore, or in the granular crystalline form of goethite, or in an amorphous compact condition. In other parts of the mines the ore is a yellowish-brown earthy ochraceous mass. All those ores contain a certain variable percentage of manganese, and a part of the concretionary masses consists of pure pyrolusite in brightly shining fibrous crystal clusters; more rare is the occurrence of concretions of the pale rose-colored sparry carbonate of manganese and lime. Associated with these concretionary hydrated ores are seams of milk-white glassy quartz, and nests of fine crystals of heavy-spar; occasionally occur also cubical pseudo-morphous crystals of hydrated iron oxide an inch and more in diameter; but particularly interesting is the abundant intermixture of snow-white nodular masses of a soft, very lubricous, minutely scaly aluminous silicate, termed by the miner *magnesia*, and mentioned in Major Brooks' Report under the name kaolinite; it is somewhat surprising how such concretions, totally inclosed within the intensely-colored ore-masses, could form without the least contamination with oxide of iron. The material of the so-called soapstone of the mines is chemically the same as that of the kaolinite, only in a harder, more compact, granular, instead of scaly condition. This lower thin-bedded series is, like the beds of the Jackson mine, bent into numerous sharp folds, many of which curvatures are splendidly exposed in the mining-pits; we can observe in the same places the contiguity of these rock-beds with dioritic rock-masses, which squarely broke through such arches of ledges, and pushed their severed ends in faulted positions to each other on the two sides. The discordance existing between the dioritic and the iron-bearing rock-groups is obvious in the majority of natural or artificial exposures, although it often occurs that they adjoin each other in parallelism.

On the higher part of the hill-slope south of the Jackson mine ridge, very close to the south line of Sect. 1, and not far from the margins of the ore-formation where it abuts against the dioritic summit part of the hill, a large doleritic dyke 70 or 80 feet wide intersects it in a direction from east to west. The external portions of the dyke-mass, in contiguity with the ferruginous beds,

are in a rotten, friable condition; the central parts of it are very hard and compact; the grain of the rock is coarse enough to distinguish with the naked eye its composition of greenish-brown augite crystals, and of glassy white feldspar in about an equal proportion. The feldspar, in the weathered, decomposing portions of the rock, is milky white, and some of it totally transformed into kaolin.

The above-described section across the ore-formation, from the Jackson mine southward, comprises Groups XII., XIII., and XIV. of Major Brooks' subdivisions. The quartzites constituting Group XIV. are, according to my views, analogous to the Teal Lake quartzites, which are in Major Brooks' arrangement placed with his Group V. These strata are in the Jackson mining location only poorly represented; we find them much better developed in the mines farther west. All the ore-bearing rock-beds of the Negaunee district I left together in one group; they form an uninterrupted succession, and are composed of similar material; moreover, these lower beds, constituting Group XII., are restricted to the vicinity of Negaunee and Ishpeming, while in the western mines no equivalent for them is found. I consider them as a locally developed series of deposits, not deserving to be ranked as a separate formation. The beds of hydrated ores are not peculiar to the lower strata alone; similar seams of hydrated oxides in the shape of grape-ore and non-crystalline earthy masses occur in the Jackson mines, closely adjoining the belts of specular-ore. In one of such seams of the Jackson mine, cellulose earthy hydrated ores occur, whose cavities are filled with a soft, bright green mealy mineral, which, on chemical examination, proves to be silicate of chromoxide.


All other parts of the hill-side situated north of the pits of the Jackson mine, and north of the Houghton and Ontonagon Railroad track, are formed of the same red jasper-banded ferruginous rocks as those composing the higher part of the ridge; they likewise inclose valuable belts of specular ore, which is mined on the west side of the slope. In the adjoining valley, and on the slope of the diorite range on its opposite side, the rock-beds are deeply buried under drift-deposits, with exception of a patch on the hill-slope, about 150 or 200 steps in width, where the red jasper banded lean ores are seen to repose with southern dip directly on

the diorite. The ledges exposed amount to a belt of 75 or 80 feet in thickness, and the ferruginous bands of the rock are rich in bright ore particles, but no ore-seam of any value has been found, although explorers have left vestiges of their work at this spot. East of the Jackson mine the ore-bearing beds soon disappear under the drift. About 300 steps from the most eastern pit, close to the Houghton and Ontonagon Railroad depot, a productive belt of soft hydrated iron-ores has been discovered by sinking a shaft; but farther east, across the creek passing Negaunee village, the ore-formation disappears beneath the dark-colored sandy flagstones of the fifth group. Some test-pits near the base of the knobs east of the Negaunee iron-furnace have uncovered red-colored siliceous flags of a ferruginous character, which are the last exposure of the ore-formation seen in this direction. The course of the above-mentioned creek, a tributary of the Escanaba, forms near Negaunee a natural line of demarcation for the different rock-formations. All the hill-slopes on the west side of the valley are formed of the lower, thinly-laminated part of the ore-bearing rock-series, and all on the east side, leaving drift deposits out of consideration, are found to be arenaceous and slaty rock-beds representing Group V. of the subdivisions adopted by me for the Huronian rock-series.

The description of the lower part of the iron formation south of the Jackson mines applies also to the more eastern exposures on the south side of the creek. The McOmber mine, in the N.W. quarter of the N.W. quarter of Sect. 7, Town. 47, R. 26, next adjoining the Jackson mine property, is one of the most productive mines of soft hydrated and generally more or less manganiferous ores, which the miner, ignoring their partially hydrated condition, calls hematite ores. East of the McOmber mine is Schade's mine, still richer in manganese ores than the former. It is situated in the N.E. quarter of the N.W. quarter of Sect. 7, high up on the hill, in a depression between two ridges formed on the summit of diorite and dioritic schists, but surrounded completely by the ore-bearing beds, which in their extremely folded condition dip in every direction, sometimes toward the diorite, other times falling away from it. Lower down on the slope of the northern ridge several other large mining-pits are opened in the S.E. quarter of the S.W. quarter of Sect. 6, and recently rich deposits of hematite

ore have been discovered in the S.E. quarter of the N.W. quarter of Sect. 7.

Many other mines in the same rock-belt are found scattered along the east slope of the hills in the S.E. quarter of Sect. 7, and in the N.E. quarter of Sect. 18. In the latter localities the dip of the ore-bearing beds is generally directed toward the hill, and on a higher level we find in some of these localities a series of dark blackish-colored ferrugino-arenaceous flagstones, or thicker quartzose ledges incumbent on them, which often contain large proportions of granules of magnetite. Good exposures of these quartzose magnetic strata can be seen near the S.E. corner of Sect. 7, where they lie on the side of a diorite hill near its summit, dipping toward it, while at the base the hematitic rock-beds crop out, dipping in the same direction as the black magnetic rock-ledges. The last mining-pit on this eastern slope of the range has been opened near the quarter-post on the east line of Sect. 18, but no merchantable ore was found; farther south, the above-mentioned arenaceous-ferruginous flagstones form the surface-rock, which I consider as representatives of the fifth subdivision of the Huronian series. Very extensive exposures of the lower iron-bearing rock-beds are in the N.W. quarter of Sect. 8, and in the adjoining parts of Sect. 7, where they surround an isolated short ridge of diorite, partly leaning on its sides, partly dipping toward it; they inclose valuable seams of the so-called hematite ores. In direct continuity with these outcrops is a cut of the Escanaba Railroad, made across



a bubble-like hillock of this formation, which forms a basal spur of a higher hill-range composed of massive quartzites and of slaty and flaggy arenaceous-ferruginous rock-beds. The strata in the cut form two successive anticlinal arches, which are in two places transversally intersected by wedge-like masses of chloritic schists

intruded from below. The upper portion of the ledges is formed of the banded alternating beds of pale brownish jasper, and of siliceous ore-seams like the jasper-banded rocks of the McOmber mines; beneath them are intensely red-colored argillite strata, in alternation with a few thick seams of dark green chloritic rock of very fine, almost aphanitic grain, and with siliceous flagstones richly impregnated with specular ore particles. The quartzites composing the higher part of the hill are not seen in direct contact with the ore-bearing beds, on account of heavy drift-deposits on the place, but from the general position of the ledges, and from the circumstance that masses of schist analogous with the schists of the dioritic group have been forced into the ore-belt from below, I infer that this could not be, if the very massive quartzites closely adjoining were the underlying rock; they should have first been pushed to the surface and been wedged into the ore-beds. In the mining-pits, not 50 steps from the place, several other such anticlinal bubbles of iron-rock, intersected by chloritic schists, can be seen denuded; and, moreover, we observe in these pits the direct superposition of the ore-beds on the diorite. I believe, therefore, that the quartzites forming the higher part of the hill are the incumbent, and not the underlying rock in this exposure. The absence of the upper ore-bearing, red jasper-banded rock-series, with inclosed seams of hard specular ore, and of the quartzites incumbent on it, from the exposures of the lower part of this group on the east side of Negaunee, is remarkable, because younger strata, which elsewhere have their place above these eliminated beds, directly succeed the others. These younger rock-beds cover most of the surface east of this place for many square miles, and no more of the iron-formation can be discovered in that direction. West of Negaunee and of the Jackson mines we find the ore-formation expanding over a large portion of the surface for a distance of three miles; much of it is covered up with drift; but the miner, diligently searching over all this space by sinking shafts and with the diamond drill, has demonstrated its uninterrupted extension beneath these drift-masses as a fact. Scattered over this area are numerous insular hills of dioritic rock, which pierces the incumbent ore-bearing layers, and stands out in high precipitous cliffs, or in lower hillocks rounded off by drift action. Mines are opened in the red-banded jaspery lean

ores in the S.E. corner of Sect. 2, Town. 48, R. 27, close to the base of vertical bluffs of diorite, which furnish principally soft hematite ores. Farther west, some hillocks in Sect. 11, south of the Houghton and Ontonagon Railroad track, are composed of a dioritic nucleus incased within a girdle of the jaspery lean ores, and next ahead of them a large knob, over which the west line of Sect. 11 runs, is totally composed of an anticlinal arch of these beds, with inclosed valuable seams of specular ore. In a western prolongation of this knob the Cleveland Mining Company has worked some of these seams; on the south side of the knob are the Marquette mines. North of this hill are the main openings of the Cleveland Company and the New York mines, which work the same body of ore-deposits. North and east of the mines are large diorite hills, on which the ore-bearing strata lean with southern dip, in direct contact with the schistose beds of this rock-group.

The ore-bearing rock-beds, after falling away from the diorites under a steep angle, bend up and down again to form several more shallow successive folds, which in places have been totally undermined, and the synclinal troughs excavated in that way, with a heavy hanging wall suspended over them by frail distant pillars, look very dangerous, and will sooner or later break down, as has already happened in some parts of the mines.

The quartzites covering the ore-bearing ledges are compact, dark-colored by hematitic pigment, and by intermingled granules of martite; the inferior strata are almost regularly brecciated and intermingled with ore-fragments. In the pits of the Cleveland mine, south of the Houghton and Ontonagon Railroad, these brecciated quartzites are immediately underlaid by a fine seam of specular ore, which itself is on its surface of brecciated structure. This ore-bed is underlaid by a series of light greenish or grayish-colored silky-shining hydro-micaceous schists, impregnated with small martite crystals, which on their part repose on the red jasper-banded lean ores, with inclosed seams of granular and of slaty specular ore. In the pits north of the railroad the quartzite is generally underlaid by chloritic schists inclosing locally quite large octahedrons of martite; below them usually follows a valuable seam of specular ore, schistose chloritic beds again, and then mixed jasper-banded lean ores, with other interstratified seams of ore, partly in the granular form, partly as a slate-ore; lowest, resting

on dioritic schists, are, in the New York mines, fine-grained silky-shining dark gray-colored argillites, charged with minute granules of martite more or less abundantly. The lower beds analogous to the strata of the McOmber mines are not found here.

Outcrops of the iron formation continue all along the hill-side from the Marquette mine west to the Lake Superior mine, situated on the north slope of an isolated diorite hill, which forms the terminal western spur of the range of diorite hills commencing south of Negaunee. By boring with the diamond drill it has been ascertained that the whole swamp valley on which Ishpeming is located is underlaid by the iron formation.

The ore-bearing beds in the Lake Superior mine lie in steep inclination, with northern dip directly on the diorite, or on schistose beds belonging to this group, but in other parts of the mine the strata are seen to be bent and folded repetitiously, and to dip in the most irregular way—much more so than in the Cleveland and New York mines. Uppermost in the succession we find heavy compact beds of a light-colored whitish or reddish quartzite, which are particularly well exposed in the pits of the Barnum mine, which naturally forms an integrant part of the Lake Superior mine, although owned by different parties. Beneath the quartzite succeeds a series of hydro-micaceous and argillitic schists, inter-laminated with seams of quartzite. The variety in the aspect of the schists is quite manifold, but closer examination shows them all to be composed of a fine scaly hydro-micaceous, or of a more granular crystalline argillitic ground-mass, modified by the admixture of different proportions of hematite or chlorite, or of martite in smaller or larger octahedric crystals, or in the form of micaceous iron oxide, which latter kind of schists is by gradations in the proportion of the micaceous iron particles directly allied with the specular slate-ores which occur in this upper horizon. These schists are underlaid by a fine-grained chloritic rock-seam, which overlies a seam of rich specular ore, and lower, a large succession of red jasper-banded mixed ores follows, which incloses other valuable ore-seams. In the old pits, mined out, we see large belts of the so-called soapstone interstratified with the jasper-banded lean ores, and adjoined sometimes by dioritic schists, protruding from beneath, and, as it appears, wedged in between the ore-bearing belt during the uplift and dislocation of the strata. On the east

side of the hill, and south of the jasper-banded specular lean ores, soft hydrated iron-ores similar to the Negaunee hematites are mined by the same company; the deposits visibly underlie the jaspery massive rock-ledges, and repose in a much-corrugated and shattered condition on the schists of the diorite group. These lower beds extend southward along the hill-slope, and are in all probability in direct connection with the soft ore-beds of the Lake Angeline mine. The Lake Angeline mines, situated in the north part of the N.W. quarter of Sect. 15, close to the shore of the west end of Lake Angeline and at the foot of a steep-sided ridge of diorite, principally work the lower hematitic ore-belt, which closely adjoins in an almost vertical position the dioritic rock-walls. Some distance farther east on the slope the higher red jasper-banded ledges, with inclosed belts of hard specular ore, are found developed, and some mining work is done in these hard ore-seams. On the south side of these same diorite hills are the Salisbury mines, in which locality only the lower strata of the ore-formation are represented. A large portion of the ledges exposed there are siliceous flag-ores which are not rich enough to be mined; the main product are hydrated ores, similar to those of the Negaunee hematite mines. In the Salisbury mines a large quantity of waste rock is hoisted and used as a bedding for the railroad track, not only there, but also on the road-bed of the North-Western Railroad, where it runs through the swamp lands near Ishpeming. The peculiar structure of the rock excites particular interest. It is a siliceous rock, richly impregnated with hydrated oxide of iron of a brownish, and to a great extent of a bright yellow ochraceous color. A part of it is compact, but most of it has a very open, cellulose, and to a certain extent laminated structure. The compact portions are comparable to an accumulation of layers of branching fucoidal stems, with club-shaped ends, which were imbedded in a solid siliceous ground-mass, which siliceous skeleton, after destruction of the material of the stems, had the remaining cylindrical branching and anastomosing cavities replenished with a porous silico-ochraceous substance. In the rock of cellulose structure, which seems to be a partially decomposed form of the first-mentioned compact rock, the compact interstitial siliceous skeleton seems to be removed by being leached out, and the porous, stem-like parts are the preserved portion.

Other portions of the rock of porous or cellulose structure consist of a succession of delicate laminæ, with a hollow interval between each of the layers, which are connected and support each other by their regular wavelike curvation, causing their alternate approximation and ankylosis, and the subsequent divergence of the laminæ. These last-mentioned rocks have a striking resemblance to the so-called eozone, but I have not the least idea of considering either them or the eozone as of organic origin, although I made the comparison, which was done simply to point out the occurrence of such resemblance to organic forms by a *lusus naturæ*, as it may be termed, because I am not prepared to give an explanation of the mode in which such singularly shaped rocks were formed.

A half mile south-east of the Salisbury mine a rich belt of specular ore has been discovered on the south side of a diorite knob in the N.W. quarter of the S.E. quarter of Sect. 16; the strata dip north toward the diorite, and were there covered deeply with drift-masses. Lower down the slope of this hill-side the strata come to the surface in natural outcrops at the location of the Parsons mine. On the N.W. side of the mining pits is a cluster of small rocky knobs, of which the largest, most western, consists of thick-bedded massive quartzite ledges in a succession amounting to about 200 feet. The highest of the strata, which dip under an angle of 45° west-north-west, are of a light whitish color; they exhibit on their side faces a striation of sedimentary deposition in discordant layers, which is so often noticeable in the sandstone deposits of Silurian age, and in all the more recent formations. Below the light-colored quartzites follow similar very compact ledges of dark purplish color, and of a rather coarse sandy grain, but a vitreous fracture; the dark color of the rock is due to abundant intermixture of martite granules with the mass. The lower strata of this dark quartzite are of brecciated structure, and inclose fragments of siliceous iron-ore. Underneath this rock-series, which composes the western of the knobs, follows a heavy belt of a breccia, composed almost exclusively of specular ore fragments cemented by siliceous matter, which rock-belt forms for itself the second row of lower knobs bordering the narrow valley of a small creek, beyond which, at a distance of a very few steps from the bluffs of the ore breccia, the diorite rises in steep walls at the base

of a high ridge composed of this rock. By artificial exposure in mining, the ore breccia is found to be underlaid by a thick belt of solid specular ore of a rich quality, and beneath it a large succession of red jasper-banded mixed ores follows, which inclose several other valuable ore-seams. Fissures in the ore-belt of the Parsons mine are sometimes lined with heavy spar crystals of exquisite beauty in hexagonal tabular form, with sharpened edges, some of brilliant lustre and translucent, others dull opaque. Associated with the spar occur crystals of quartz, large bunches of micaceous iron, and a soft, nacreous, scaly, white kaolinite.

South of the Parsons mines a high ridge trends westward across the centre of the N.W. quarter of Sect. 21. On its north slope the naturally and partly artificially denuded red jasper-banded ore-beds are conspicuous from a distance. The strata dip north under an angle of 60° or 70°, and are overlaid by the quartzite formation which composes the lower undulating hills farther north. The south side of these hills of ore-bearing beds is mostly covered by drift, but in the western part, opposite the New England mines, good exposures are found in which the ledges of the jaspery lean ores are brecciated and inclose seams of a rich specular slate-ore, besides more siliceous beds of slaty structure. These are in descending order conformably succeeded by a large series of thin-bedded ferruginous ledges, corresponding with the Negaunee hematite ore-belt, which contain rich seams of soft hydrated ore, mined on the New England location, where the strata lay on the north side of large diorite bluffs, in close, but, as it appears, discordant contact with them. Also the rich hematite ores of the Winthrop and Shenango mines are mined from the eastern continuation of the same rock-belt in the N.W. quarter of the S.E. quarter of Sect. 21, which is there in similar contiguity with the dioritic knobs bordering it on the south side.

From the New England mines the ore-formation trends west and little north, and can be traced by outcrops and in test-pits to connect directly with the ore-belt of the Saginaw mines and Goodrich mines, everywhere overlaid on the north side by the quartzite formation, and on the south side in contact with dioritic rocks.

South of the diorite knobs, at the Shenango mine, is another large drift-covered hill, composed of jasper-banded ferruginous ledges, but it contains, as far as known, no valuable ore-seams.

East of it, on the opposite side of the swampy valley of a tributary of the Escanaba River, commences a high hill-range, trending eastward, regularly followed by the south section line of Sects. 22 and 23, which is totally composed of siliceous flag-ores, generally not rich enough to be valuable. The strata dip under a high angle northward, reposing on dioritic rocks, which form a chain of lower hills on the south side of this ridge, which is on its north side likewise followed by a parallel row of diorite hills severed from it by the valley of a creek. The Tilden mines and Foster mines are located on the eastern part of this ridge of lean ores, both near the south quarter-post of Sect. 23—the first south of it, the second north-west of it.

In the abandoned pits of the old Tilden mine, and in several neighboring natural exposures, the ore-formation is found to repose on the diorite or on the schists belonging to this group, but as it seems always in discordance. It consists of a large succession of thin-bedded arenaceo-ferruginous layers, dark brownish red on the surface, and blackish gray with some metallic lustre on fractured faces. All the rock contains martite granules and amorphous hematitic oxide as constituents, but a good portion of the layers is so richly impregnated with these oxides as to constitute ores of from 45 to 50 per cent metal assay. These richer beds of flag-ore were mined once, but did not prove to be salable in the market. In the Foster mines, on the north side of the hill-range, the ore-formation is not altogether formed of flaggy ledges, but a great part of the very thick rock-series is formed of thicker seams, resembling the jasper-banded ores of the Ishpeming mines or Negaunee mines. The productive ore-belt of the Foster mine is not composed of flag-ores, as in the Tilden mine; it is a secondary accumulation of hydrated oxides in the form of a brecciated, very open cellulose, partly soft earthy, partly harder compact rock-mass; the cavities of the amorphous cellulose harder oxidic rock are incrustated with a thin minutely warty coating of grape-ore, and often with brilliant small quartz crystals, and in addition the porous ores contain frequently a brightly cinnabar red oxide of iron in pulverulent masses. The percentage of these ores in iron is about 50; the mines have long been abandoned, and most of the ore taken out of them lies there unused.

On the slope of the high plateau, a quarter mile south-east of

the old Tilden mine, is an exposure of a very large succession of ferruginous beds, dipping north under a high angle, which likewise resemble more the red jasper-banded rock-belt of the northern mines than the flaggy ore-bearing beds of the neighboring Tilden mines. At the base of the slope we find the granite at the surface.

The large succession of ore-bearing rock-beds surrounding the base of the diorite hills in the S.W. corner of Sect. 13, and the adjoining exposures in the S.E. quarter of Sect. 14, which were formerly mined by several companies, but are abandoned now, are perfectly analogous deposits with those of the old Tilden and of the Foster mine, and the same is the case with another outcrop on the north side of Lake Hall, in the N.E. quarter of the N.W. quarter of Sect. 14, which almost connects with the ore-deposits of the Lake Angeline mine.

Every one of these localities differs somewhat from the other in the character of its layers, but the unity of all these deposits as co-ordinate members of one formation is plainly obvious. The iron range on which the Tilden and Foster mines are located, and the deposits in Sects. 13 and 14, unquestionably represent the lower horizons of the ore-bearing series, but it is a remarkable fact that in the case of superposition of other strata on these beds we do not find them to be the quartzites next in the order of succession, but arenaceo-ferruginous beds identifiable with the younger group overlying the quartzite formation, which same observation I made before, when describing the exposures of the lower ore-bearing strata on the south-east side of Negaunee. A glance on the accompanying geological map will show a direct continuity existing between the beds incumbent on the ore-formation in the two mentioned places, and also the intercalation of the quartzite formation between these beds and the ore-bearing rock-belt of the Cascade mines will be noticed by looking more southward on the map.

Before I proceed to describe the ore-formation, as exhibited in the more western mining districts, I have yet to consider very interesting parts of the Negaunee and Ishpeming iron-bearing rock-belt on the north side of these villages. North-west of the Jackson mine we see the chain of diorite hills previously described interrupted by a depression through which a side-track of the North-Western Railroad Company is laid, which leads to the so-called

Teal Lake mines, situated on the north side of the aforesaid diorite range, and south of the west part of Teal Lake. The surface of this depression is covered with drift, but by examining the numerous test-pits opened on this space one can readily convince himself of the uninterrupted continuation of the iron-bearing rock-series across this gap from one side of the diorite range to the other. We found, as previously stated, a patch of the upper red jasper-banded beds of the iron formation leaning with southern dip on this diorite ridge, a few hundred steps east of this gap, but here we find the lower thin-bedded flaggy series, dipping likewise southward, on one side of the range as well as on the other; therefore, while on the south side the strata lie on the diorite, they seem to dip under it on the north side. The flaggy layers are rich in granular martite; some of them contain as much as 65 per cent of the oxide, and have been mined in the S.W. quarter of Sect. 36, Town. 48, R. 27, but the mines are now abandoned. Farther west, in the S.E. quarter of Sect. 35, other mines are opened in the brown jasper-banded layers, similar to those of the McOmber mine, which contain a rich belt of the soft hydrated ores, the belt is traced from there to the south-west corner of the section, south of which, in Sect. 2, Town. 47, R. 27, this soft ore-belt has a very great width and is of excellent quality. The soft ore in that locality, lying right under the surface, is taken out with the shovel like gravel, but as the excavations become deeper and the loose rock-masses cannot be timbered up, these mines are rather dangerous places to work in. The lower series of the ore-formation in the Teal Lake mines has a very great thickness, which cannot accurately be estimated, as the strata are much folded, but they amount at all events to six or eight hundred feet in thickness. By the excavations, various bubble-like dioritic masses have been uncovered, which by their intrusion into the stratified series have caused considerable corrugation of the adjoining rock-ledges.

This locality has caused me much perplexity with regard to the recognition of the true order in which the different groups of strata succeed each other. The ore-formation dips in this place, as previously stated, south, toward high cliffs of diorite, and is on its north side in close contact with slaty argillites, interstratified with sand-rock belts; still farther north follows a series of banded

ferrugino-siliceous beds, charged in certain seams with considerable magnetite, all of which strata dip south in apparent conformity with the ore-formation; then comes the bed of Teal Lake, and on its other side, the quartzite formation rises with its ledges from the water into high slanting bluffs, which likewise dip to the south. When I first saw this I felt great satisfaction, and considered this place as one of the rare examples in which the succession of the rock-formations could be plainly observed, and for a long time afterward I took it as an established fact that the ore-formation is the youngest of the three succeeding different rock-belts, and the quartzite-formation the oldest; but after a while I came across a good many other exposures, which did not harmonize with this supposed order in the succession, and finally, by inference from a multitude of facts observed, I became convinced that the seeming conformity of these beds is not a proof of their actual order of succession, and that their present relative position is the result of an overturn of the strata.

I must confess to have not yet clearly conceived the exact mode of dislocation which would bring the beds in their present overturned position; and I find it strange to see this ore-bearing series not at all developed on the north side of the quartzite range, where one naturally would expect it; but considering the other side of the question, I find a slaty and arenaceous rock-belt identical with the one adjoining the ore-formation of the Teal Lake mines on the north side, largely developed on the south side of the quartzite range east of Negaunee, which leans with southern dip on the quartzite, just as in the other place, but a mile south of the range, this rock-belt is reverted into a northern dip, and the ore-bearing strata, exposed on its south side, appear to underlie it conformably.

The previously described cut of the Escanaba Railroad, near the centre of Sect. 8, Town. 47, R. 26, offers another example of the superposition of this arenaceous slate formation on the iron-ore group. The most forcible argument, however, for the lower position of the ore-formation is the circumstance that we find it in all the mining locations of the Negaunee district, never underlaid by quartzites or slaty rock-beds of the above-mentioned class, but always resting on the dioritic formation, and in several localities a large belt of quartzite reposes on the ore-group, which, by its light

whitish color and great compactness, is so similar to the Teal Lake quartzite that hand-specimens cannot be distinguished. A quartzite of this description is naturally exposed in the village of Ishpeming on the railroad track a short distance west of the North-Western Railroad depot; the heavy strata dip south under a high angle. A drill-hole was sunk two years ago through this quartzite to the depth of 400 feet, when a valuable seam of specular ore was struck, after the drill had gone for most of its length through quartzite, and then through a breccia of quartz and ore fragments. Several other holes were drilled west of this place, on the high ground south of Lake Bancroft, all with the success of striking an ore-belt about 400 feet below the surface. The Barnum Mining Company is at present engaged in sinking a shaft in one of the spots explored by the drill. One of the most instructive exposures of the ore-formation, and of the incumbent and succumbent rocks, is found west of Lake Bancroft, on the other side of Carp River, in the S.W. quarter of Sect. 4. The red jasper-banded mixed ores are on the top of a terrace-like extension of the hills, in several places well denuded from the drift-masses which cover the greatest part of the surface; according to the distribution of the exposures they must amount to a belt of considerable thickness. North of this red-banded jasper rock follows a series of brownish jasper-banded ferruginous beds, which project in a row of low cliffs on the north edge of the terrace land. On its south side, and several hundred steps west of its exposures, a large belt of light-colored heavy quartzite beds crops out in low cliffs; the almost vertical strata dip south like the jasper ores, on which they lean conformably. On the slope directly south of the quartzites, which amount to about 200 feet in thickness, is another undulation formed by an alternating series of dark-colored slates with sand-rock seams which dip south, in conformity with the quartzite belt; they correspond in position and lithological character to the slate and sand-rock belt north of the Teal Lake mines, but are here indubitably deposited above the ore-formation. In this locality the superintendent of the Barnum mines, Mr. Sedgwick, sunk a hole with the diamond drill to the depth of 476 feet, and had the kindness to give me a complete set of cores in the order as the boring went on. The boring commenced in the quartzite. At a depth of 150 feet, sunk through

quartzite, a 50-foot-wide seam of hydro-micaceous schists, interspersed with an abundance of martite crystals, was pierced; then came again quartzites, part of them light-colored, others dark, highly ferruginous, and in a degree chloritic, to the depth of 320 feet from the surface; at 324 feet a seam of a very coarse breccia of quartz and ore fragments, only one foot wide, was bored through; then came an 18-foot seam of micaceous argillites, rich in martite granules; and again 5 feet of a brecciated ore; argillites 2 feet, same as before; brecciated ore, 35 feet; from 383 feet below the surface to 414 feet occur hydro-micaceo-chloritic schists, crowded with blackish octahedric crystals of martite; and the last specimen received from the drill-hole, at a depth of 476 feet, is again a brecciated siliceous iron-ore.

North-west of these exposures of quartzite, on the west line of Sect. 4, rises, at a distance of a few hundred steps, a steep crest of dioritic composition. The dioritic nucleus is surrounded with schistose beds dipping almost upright in position to the south; the beds next to the diorite are dark green, colored by chlorite, and consist of a granular feldspathic ground-mass in intermixture with considerable sparry carbonate of lime; the higher schistose strata are hydro-micaceous, richly disseminated with good-sized octahedric crystals of martite, and perfectly correspond with the rock found in the drill-hole at a depth of 400 feet. West of this diorite hill, a row of other, likewise dioritic, hills is seen to trend across the centre of the south half of Sect. 5. We find on their south slope a large belt of the just-mentioned hydro-mica schists, and on them follows a seam of brecciated specular iron-ore, more or less contaminated with quartzose matter, which does not naturally crop out, but is artificially denuded on the location of the Excelsior mine, situated on the west line of Sect. 5, 500 steps north of the south-west corner.

In the mine, which is on the west side of the diorite hills, at the base of a rounded drift-covered slope of the highlands, the belt of brecciated ore is overlaid by a compact, fine-grained sub-crystalline argillitic rock, which exhibits a banded sedimentary structure oblique to its imperfect schistose cleavage; a part of it is paler grayish green, with sedimentary stripes of a copper color imparted to it by ferruginous, silky-shining scales, which I consider to be an altered chlorite, by higher oxidation of its iron component, as

the green color of the rock is due to chlorite and insensibly merges into the red streaks. Other portions of the rock are dark green, and agree with the chloritic rock, which forms the hanging wall of the ore in the Lake Superior mine; it is in this form richly impregnated with octahedric crystals of martite. With the brecciated ore, which is always somewhat siliceous, a seam of a very pure coarsely crystalline ore occurs, which resembles the coarse-grained magnetite of the Champion mine, but is not at all magnetic. Next above the chloritic argillites, which amount to about 25 feet in thickness, follow light-colored, white and reddish quartzites, completely similar to the Teal Lake quartzite, in a thick belt, which is for about 50 feet uncovered, but south of there disappears under the drift.

The higher hills, ascending about 300 steps north of the ore-pits, are superficially covered with drift, but by natural outcrops and by numerous test-pits they are known to consist of a large succession of ferruginous sandy flagstones or jaspery mixed ores, erected into an almost vertical position, while the strata in the mine dip under an angle of only 45° to the south. Farther north argillitic slaty strata and seams of quartzite are found in the test-pits, also angular fragments and larger blocks of slate-rock are scattered abundantly over the surface; these slates have likewise a southern dip. The large series of banded lean ores are equivalent with the hematite ore-belt of the Negaunee mines; the slates and quartzites found north of the ore-formation are not underlying but younger deposits, which constitute the fifth group of Huronian deposits.

On the high lands west of the Excelsior mine the ore-bearing jaspery rock-series is across the centre part of Sect. 6, often seen in natural outcrops, and found at no great depth under the surface by test-pits, in which very fair seams of specular ore have lately been discovered, but they are not large enough to be profitably mined. The strata dip south. South of the exposures is a small knob of quartzite, situated a few hundred steps north of the south quarter-post of Sect. 6.

On the west line of the section, along its south half, are several knobs composed of the jasper-banded beds of the iron-formation; west of them is the swampy valley of the head-waters of Carp River, and no more is seen of this group in that direction for more

than a mile; but recent explorations have led to the discovery of a very valuable ore-belt in the S.W. quarter of Sect. 32, Town. 48, R. 28, which is the location of the newly-opened Boston mine. Intermediate between this mine and the before-mentioned exposures on the west line of Sect. 6, Town. 47, R. 27, the quartzite is seen in outcrops near the quarter-post on the line between Sects. 3 and 4, Town. 47, R. 28, and in the north half of the N.E. quarter of Sect. 5, and the ore-bearing jaspery rock-series is found on the north side of the quartzite outcrops in every one of these localities well exposed.

The Boston mine is opened on the north side of a hillock situated in the midst of a swamp; across the swamp, 500 steps northward of the mine, are high granite bluffs; south of the swamp are high drift-covered hills underlaid by slaty rocks of a blackish color, which belong to the upper horizon of the Huronian series subsequently to be described. The ore-belt of the mine is from 5 to 10 feet wide, stands nearly vertical, and is, without the intervention of many other rock ledges, conformably overlaid by a thick belt of light-colored compact quartzites, which have a dip to the south. The foot-wall of the ore is formed by a large succession of the red jasper-banded mixed ores, which inclose seams of fine-grained sub-crystalline argillitic schists analogous to the soapstones of the Ishpeming mines. The ore itself is a fine-grained, very compact laminated specular ore of a bluish gray color, with dull metallic lustre, but also seams of brightly shining slate-ore are found in the mine associated with the other kind.

The ore-formation has been traced for some distance west of the Boston mines, but I have not personally examined that district.

Returning to the south side of the synclinal trough of ore-bearing deposits, I have yet to describe the Saginaw and Goodrich mining locations, which, as we learned by previous remarks, are in direct continuity with the deposits of the New England mines.

The north slope of the hills, on top of which these mines are located, is largely covered with drift deposits; but in various places the higher arenaceous and slaty strata of the fifth group come to the surface, and in the swamp at the base of the hill, a little more than a quarter of a mile east of the Saginaw Railroad depot, a knoll of diorite protrudes. Next north of the mines we find the quartzite formation in a belt of great thickness; the upper

strata are light-colored compact ledges, the lower beds are conglomeratic and brecciated, full of ore fragments; and next to the productive ore-belt these brecciated very massive layers are almost totally composed of pure ore and of red jasper-banded mixed ore fragments. The productive ore-belt of the mine is interstratified between jasper-banded ledges, which amount to a great thickness, and consist partly of a bright metallic granular specular ore, partly of slate-ore, and in association with it are argillitic and hydro-micaceous schists in all shades of color, from white to red, yellow, brown, green, or gray, by impregnation with hematitic or ochraceous iron oxide; or with chlorite, and with granular martite. Some have an earthy aspect, others are silky-shining. We find also among these the peculiarly white and red dotted argillites, whose color I previously compared with the dotted plumage of a Guinea-hen. The strata dip all to the north. South of the mines is a high rocky knob in close proximity, which consists of a large succession of banded, thinly-laminated arenaceous beds, richly impregnated with iron oxide and in part with magnetite; frequently also the sandy rock-seams alternate with seams of actinolite, or the fibrous crystals of this mineral are scattered through the whole mass. The position of these beds is almost vertical, they succeed the ore-formation conformably, but the surface is too much incumbered with drift to allow us to see the junction between the contiguous rock-belts. Drift-deposits cover also, south of this knob for quite a distance, all the surface, with exception of limited exposures of granite, on the lower levels, a half mile distant from the knob, or, to indicate their position exactly, they are found in the S.W. quarter of the S.E. quarter of Sect. 19, Town. 47, R. 27. West of the Goodrich mines, located in the N.W. quarter of the N.W. quarter of Sect. 19, next adjoining to the Saginaw mines, the ore-formation disappears under the drift, and is not seen again for a distance of six miles, where it is uncovered by test-pits opened in the S.W. quarter of the S.W. quarter of Sect. 7, Town. 47, R. 28, adjoined on the north side by the quartzite formation and by a large series of higher arenaceous-chloritic rock-beds, all dipping conformably with the ore-belt to the south. The ore found there is partly a granular non-magnetic specular ore, partly a specular slate-ore, and is associated with gray hydro-mica schists, disseminated with an abundance of small martite crystals.

On the south side of the ore-belt we find the higher part of the ridge composed of banded arenaceous-ferruginous actinolitic rock-masses, similar to the strata composing the knob south of the Saginaw mines; the dip of these rock-beds is likewise southward, all these different strata must therefore be in an overturned position. A mile north-west of this locality we find in a continuation of this same hill-range the Washington mines and the Edwards mine. These mines surround in a semicircle the slope of a ridge isolated from the main body of the range by ravines. The new Washington mine and the Edwards mine occupy the north slope; the old Washington mine and the pits of the abandoned Hungerford mine are on the north-west and west side of the ridge.

North of them is the swampy valley of the Escanaba River. The rock-beds dip under a high angle to the north in the eastern pits, and curve into a western dip in the western; but considerable local deviations from this rule occur, as the strata are much folded. Quartzites of various character form the top layers of the formation; uppermost are light-colored, compact, thick-bedded ledges, which are well exposed near the railroad depot, a short distance up the hill-side; and still better exposures are half a mile farther west, on the north side of the railroad track to the Republic mines, above the cut it makes through the lower conglomeratic hydro-mica schist. The lower portion also of these quartzite ledges constitutes a coarse conglomerate of rounded pebbles of quartzite of different color, of siliceous ore-fragments and of schistose pieces of the ore-bearing series. Next under them follows in the latter locality the just-mentioned conglomeratic hydro-mica schist, which amounts to considerable thickness. The quartz pebbles in it are sometimes much crowded, at other times only distantly scattered through the silky-shining schist, gray-colored by minute scales of micaceous specular iron oxide. South of these conglomeratic schists, which compose a separate undulation of the surface, are other quartzose hydro-mica schists of very bright metallic lustre from intermingled scaly ore particles, and inclosed within these a large seam of excellent slate-ore occurs, which is merely a modification of the schists by total replacement of the hydro-mica by micaceous iron oxide; all gradations of the transition from the leanest schist to the rich ore can be seen in the mining-pits opened lately on the south side of the railroad cut,

and it requires some experience to make the distinction between the valuable ore and the less rich waste rock. Farther south the surface is covered by drift-masses, but not far off from the schist massive blackish crystalline rocks with brightly glistening fracture come to the surface, which consist of an intimate intermixture of white feldspar with black hornblende in thin-bladed, somewhat fibrous crystals. This rock, in composition a diorite, has an entirely different aspect from the ordinary diorites associated with the ore-formation in the eastern mines. In the Hungerford pits, south-west of the western part of the old Washington, the slate-ore is much more compact and harder than in the adjoining before-mentioned pits. East of the rock-belt inclosing the slate-ore occurs in the Hungerford location a peculiar kind of hard siliceous hematitic ore, which forms a part of the actinolite schist series which composes the higher part of the ridge. In the Franklin pit, on the west side of the Edwards mine, the following section through the ore-formation is observable. Uppermost of the strata exposed is a banded rock-belt of considerable thickness, composed of alternating thin seams of white granular quartzite interspersed with magnetite granules, and of other seams intensely impregnated with magnetite, so as to constitute a more or less siliceous ore-mass. Under it follows a pale yellowish-colored silky-shining hydro-mica schist in a seam of eight or ten feet in thickness; then comes a belt of siliceous specular slate-ore several feet wide, and again another seam of hydro-mica schist similar to the former. Next below are again banded quartzose magnetite ores like the upper rock-belt, which inclose a 15-feet wide seam of a high-graded pure magnetic ore.

Associated with this ore-belt is a dark blackish-colored glistening schist of micaceo-chloritic composition. Lowest in the mine follow actinolitic and quartzose schists, richly impregnated with granular magnetite, and in certain seams abundantly interspersed with brownish small garnet crystals. On the south side of these ore-bearing beds, all of which dip under an angle of about 70° to the north, a massive belt of a dark dioritic rock follows, and then the higher main body of the ridge is found to be composed of an actinolitic quartzoso-ferruginous banded rock-belt of immense thickness, which dips the same way and incloses chloritic seams full of brown garnets, and is from time to time interrupted by

parallel dioritic belts, of which it is uncertain whether they are intrusive or not, but I think they are. All these diorites, next to the ore-belt and the others farther south, are very similar to each other, but differ as before mentioned from the diorites in contact with the ore-belts of the eastern mines.

In the eastern pits of the Washington mines a good opportunity is offered for the study of the beds incumbent on the ore-belt by a long tunnel driven into the hill-side. The quartzite formation has in that locality a great thickness. Its upper beds are light-colored; the lower ones are dark, by intermingled ore granules and hematitic pigment, or by intermixture of chlorite. Interstratified with the quartzites occur chloritic seams and larger belts of micaceous quartz schists; next under the quartzose rock-series succeeds a dark blackish-green colored massive crystalline rock, in connection with other distinctly stratified sedimentary beds of similar mineral composition, which by gradation into the crystalline rock show its origin from such sediments, greatly altered by metamorphic agents. The massive part of the rock-belt resembles a dark, very coarsely crystalline diorite; it consists of black, rather hard, large tabular crystals, with a brightly shining micaceous cleavage, which I identify with ottrelite, and of a lighter-colored minutely granular feldspathic, or other times hydro-micaceous interstitial mass, serving as a scanty cement, or prevailing over the crystal blades, which finally entirely disappear in the sub-schistose laminated layers of the same rock-belt. Generally these rocks contain considerable proportions of minutely granular magnetite equally distributed through the mass, besides coarser octahedric magnetite crystals in clusters, or disseminated singly. Portions of this belt have an amygdaloid structure, by segregation of nodular rounded lumps of light greenish-colored earthy substance within the rock-mass, which by exposure of the rock readily becomes washed out, and leaves the exposed surface portions full of small cavities, from the size of a pin's head to that of a pea. Immediately beneath this rock series is a rich belt of magnetic ore, of a thickness from 15 to 25 feet, which is associated with narrow seams of hydro-micaceous and chloritic schist. The ore-belt reposes on a large succession of quartz-banded mixed magnetic ore-seams too low in percentage of the metal to be of value for the miner. In the presently considered eastern mining-pits of this

location, the strata of the ore-bearing group are seen to be transversally intersected by a doleritic dyke of about ten feet in width in one place, and much narrower in another; it divides into several branch seams. South of the mixed magnetic lean ore belt, which forms the foot-wall of the mine, an immense body of well-laminated, banded, but firmly coherent rock-strata succeeds in apparent conformity, but a direct superposition of the ore-belt on them is not observable. This large rock-belt, a quarter of a mile wide, consists of an endless alternation of thin laminar seams of arenaceous character, with seams of actinolite and with ferruginous, or jaspery, or chloritic bands, all united into clumsy masses like a crystalline non-stratified rock, and exhibiting great distortion and corrugation.

The actinolite forms partly seams exclusively composed of the stellate fibres of this mineral in longer asbestine crystals, or as a magma of minute needles, which are generally interspersed with magnetite granules and with small cinnamon-colored garnets; but also the other arenaceous, ferruginous, or jaspery seams of the rock are very often full of actinolite needles mingled with their substance. In the same way garnets are to be found in nearly all of the seams; but in the chloritic seams, which usually are larger than the others, the garnets are most abundant, and form clusters of sometimes quite large crystals of brown color, and not translucent. Narrow seams of this rock-belt are almost completely formed of magnetite; some others are composed of hematitic ore of a more or less siliceous character; the before-mentioned ore-seams, uncovered in the old pits of the Hungerford mine, belong to this class of sediments.

Four miles west of the Washington mines are the Keystone mines, and west of them the Champion mines. North of the Keystone, near the railroad track leading to the Champion mines, are exposures of a large belt of schistose arenaceous-chloritic and micaceous strata, interlaminated with coarsely conglomeratic seams, containing pebbles of granite, diorite, quartzite, banded jaspery rock-fragments, besides pieces of schists and slate-rocks. The not-conglomeratic seams are well-laminated striped rocks by alternation of lighter and darker colored layers in the composition of the strata. Certain seams rich in chlorite or black mica contain garnet crystals in abundance, and are disseminated with clusters of

actinolite crystals. The dip of this rock-belt is northward under a high angle; south of it, separated by a ravine, follows conformably another large rock-belt of light-colored quartzite, partly in heavy compact ledges, partly in more thinly stratified beds; the lower portion of the series has a schistose structure from copious intermixture of white mica scales with the quartzose mass, and a still lower succession of beds, amounting to 12 or 15 feet, is formed of fine-grained silky-shining mica schists, full of decomposed brownish earthy garnet crystals. These repose on a granular quartzite, intensely impregnated with magnetic granules, so as to impart to it the quality of a low-graded siliceous ore; it incloses, not far from its upper limits, a four-feet-wide belt of a pure rich granular magnetic ore, and in association with it chloritic-micaceous schists occur, densely crowded with dark reddish garnet crystals of the size of a pea and larger. The mixed siliceous lean ores continue on the south side of the productive ore-seam, and form a quite large belt, which is partially denuded in the mining-pits. South of the pits rise abrupt rock-walls, composed of the banded actinolite rock in compact masses, which form the crest part of the hill-range, and continue southward for about 800 feet, where a belt of diorite and then granite succeed.

Very similar is the succession of rock-beds in the Champion mines. Their dip is likewise northward. The schistose conglomeratic rock-belt, succeeded upward by black slate-rocks, and above them by siliceous and ferrugino-argillitic beds, compose the north slope of the hill-side, which layers are well exposed on the side of the road leading from the mines to the railroad depot. The underlying quartzite belt is well denuded in the mining-pits; its upper ledges are light-colored, thick, and compact; the lower are thinner and micaceous, like those of the Keystone mine, alternating with light-colored, argillitic, and hydro-micaceous, silky-shining schists. Beneath them follows a series of chloritic beds, interlaminated with quartzose seams, all of which are more or less intensely impregnated with magnetite granules. The chloritic rock is partly fine-grained, almost aphanitic, partly very coarsely scaly and glistening; it incloses an abundance of large-leaved black mica, and often white mica, which appears black from chloritic substance deposited between its laminae, besides large crystals of iron garnet, some of which have a diameter of three

inches, and all are complete; with them usually occur also octahedric magnetite crystals larger than a mustard-seed, and not rarely radiated bunches of long needle-shaped crystals of black tourmaline and cubes of iron pyrites are seen imbedded within the chloritic mass. The chloritic rock-belt forms the hanging wall of a large seam of granular magnetic ore, which is associated with silky-shining hydro-micaceous schists; some of them are even-bedded, dark gray colored by copious intermixture of granules of magnetite; others have a very rough brecciated structure from chloritic rock-fragments, inclosed by the much-corrugated hydro-micaceous ground-mass, or else large-leaved micaceous blades of black color copiously mingled with the whitish fatty schistose substance, compose these peculiar schistose beds. In some of the pits of the Champion mine a band of specular slate-ore occurs, which seems to have its position not far apart from the magnetic ore-belt. The foot-wall of the productive ore-seam is formed of a thick series of banded quartzite ledges, charged with large proportions of granular magnetite, unequally distributed in the mass in seams, with the magnetite prevailing over the quartz, and in others more sparingly dispersed. This ferruginous quartz-rock is on its south side in contact with a parallel belt of dioritic rock of partially schistose, partially massive crystalline structure; south of it are no more rock-beds exposed, for a distance of about 400 steps, which space is occupied by a swampy depression, on the other side of which granite, with inclosed hornblendic and micaceous-chloritic seams, comes to the surface and projects in a row of cliffs. In the eastern part of the mining location the granite disappears under drift deposits, and in its line of strike a diorite hill projects, which on its south side is in contiguity with actinolitic schists in irregular, nearly vertical position, much corrugated and twisted in some places, in others not. A broad strip of the surface is underlaid by a continued succession of these actinolitic beds; and high rock-knobs situated some distance farther south are found to consist of the same actinolite rock; still farther south are granite hills.

Eight miles south of the Champion mines we find the Republic mines, located on the S.E. side of Smith's Bay, a lake-like dilatation of Michigamee River. The iron formation surrounds this bay in a horseshoe-like arch, dipping with steeply erected beds from

all sides toward it. The inner circle of rock-beds is formed of compact, reddish-colored, in part brecciated quartzites, with a micaceous-schistose cement. Under the quartzite follows a seam of magnetic ore about 10 feet wide, inclosed within a series of quartzose beds impregnated with ore particles. Next below comes a narrow belt of silky-shining hydro-mica schists, some of which layers are charged with scales of brightly metallic, specular iron oxide, and beneath these occurs a thick seam of granular, coarse-grained, specular ore of great lustre. This ore-seam is succeeded by a large series of red jasper-banded mixed ores, which incloses other valuable ore-seams in connection with schistose bands of micaceous-argillitic character. The jasper-banded mixed ores exposed on the south-west side of Smith's Bay contain large seams of quartzose slate-ores of high percentage in iron, which are not regarded as being worth mining; but recently valuable specular ore has been discovered there in connection with the leaner beds. Above the jasper-banded strata of that locality are conglomeratic and brecciated layers, composed of larger and smaller pebbles and fragments of quartz of red and of white color, cemented by a siliceous mass of specular ore. The outer circumference of this jaspery rock-belt, which amounts to about 300 feet, is in contact with a large body of diorite of blackish color, similar to the diorite beneath the ore-belt of the Washington mine, but unlike the diorites of the Negaunee district. Portions of this diorite are much richer in feldspar than others, and from the red color of the spar I infer its being, in part at least, orthoclase. On the other side of this diorite belt adjoin it dark-colored, blackish ferruginous flaggy rock-beds, banded with lighter-colored narrow linear siliceous seams, in a nearly vertical position, which are succeeded by dark blackish-green colored micaceous and chloritic schists, full of good-sized brown garnet crystals; they are much corrugated and folded. South of them the whole slope of the hill-side is composed of an endless succession of banded ferruginous-siliceous actinolite schists, united into bulky compact masses, which are here and there interrupted by intrusive diorite belts of short local extension, and not, as represented by Major Brooks, in regular continuous bands encircling the whole side of the mountain. All these rock-beds dip under a high angle toward the hill-side, in apparent conformity with the direction of the ore-bearing rock-beds. At

the base of the slope, south of the actinolitic rock, a separate undulation of the surface is formed of a fine-grained quartzite of sand-rock structure, partly white, partly brownish-colored by admixture of small garnet crystals. I consider this quartzite merely as a local modification of the actinolite schists. A few steps farther, across a narrow strip of swamp, another hill chain rises, which is composed of gneissoid laminated rocks and of massive granite, both very rich in glistening white and blackish mica scales; the gneissoid ledges dip irregularly. In another locality farther east, the actinolite schists are in close contact with the granite, and dykes of it intersect them. At the Kloman location, north-west of the Republic mine, on the other side of Michigamee River, the quartzite formation and the ore-bearing rock-beds have a vertical position, and are seen in immediate contact with a dioritic rock-belt in a similar upright position, which is, on its other side, adjoined by ferrugino-quartzose banded rock-ledges, identical with those adjoining the same diorite belt on the south side at the Republic mine.

North-west of the Kloman mine are several other mines, close to Michigamee River—the Chippewa, Cannon, and Magnetic iron mines—which I have not examined, to my great regret, as the ore deposits in these localities, represented to be lower than those of the Republic mine, are not far off from the great body of stauroliteiferous mica-schists, which form the youngest strata of the Huronian series, and quite probably important facts regarding the succession of rock-beds could be observed in the surrounding country.*

Two other mines on the north side of Lake Michigamee—the Spurr and Michigamee mines—are yet to be described, not only for their economical importance, but as being of the highest scientific interest by the regularity and completeness of the geological sections offered there to the observer. The two mines are situated on the south slope of high diorite hills, which form a separate range in front of a large body of granite hills on their north side. The strata of the ore-formation, directly incumbent on the dioritic rocks, dip south under angles from 50° to 60° . These so-called diorites differ from the ordinary dioritic rocks by an abundance of

* While these pages were going through the press I had occasion to examine all these mines carefully, but it was too late to insert the results of my examinations.

quartz in their composition, and by the comparative scarcity of feldspar. Chlorite is one of the principal constituents of the rock-mass in intermixture with granular quartz, and usually with some proportion of carbonate of lime, which is readily discovered by the lively effervescence of the rock when put in muriatic acid. Within this chloritic ground-mass are dark-colored, somewhat fibrous actinolite-like hornblende crystals, more or less copiously dispersed, often without any visible admixture of feldspar; in other cases, crystals of this mineral can be distinguished. The generally massive non-stratified rock exhibits sometimes obscure traces of former stratification. Immediately on the diorite leans, in steep inclination, a large succession of banded ferruginous strata, which consist of an alternation of seams of a red-colored granular sand-rock-like quartz, with brightly metallic-shining seams of scaly specular ore; frequently these alternate seams are irregular, and represented by interrupted lenticular bands, wedging out on both ends. These are, in the Michigamee mines, succeeded by an equally large succession of banded quartzose beds of sand-rock structure, which differ from the former by the white color of the quartzose bands, and by the magnetic quality of the interlaminated alternating ore-seams, which have a more dull, blackish color, and are sometimes a soft friable mass. The ledges of this ore-bearing sand-rock belt are often brecciated, or else intersected by a dense network of shrinkage cracks, which are filled with ore particles. In the Spurr mines we find, between this and the lower red-banded series of ore-bearing beds, a belt of garnetiferous chlorite rock intercalated, which is associated with a massive crystalline diorite of somewhat different character from the other dioritic rocks at the base of the formation, as this one contains a good proportion of feldspar crystals, or else a granular feldspathic interstitial ground-mass. Above the quartzose rock-belt, banded with seams of magnetic ore, follow some layers of chloritic schist interstratified with quartzose seams, on which the productive magnetic ore-beds, from 20 to 40 feet in thickness, are deposited, and above them succeed other chloritic strata containing large crystals of well-formed iron garnets, and smaller octahedric crystals of magnetite, needles of actinolite, or larger stellate clusters of it in connection with smaller seams of magnetic ore, intermingled with larger masses of actinolite and with seams of quartzite. On top of them lies a thick belt

of heavy compact quartzite ledges, green-colored by chloritic scales, or light gray-colored. Part of these layers is conglomeratic, and incloses larger fragmental masses of the lower red-banded specular mixed ore-belt. This quartzite formation is overlaid by several hundred feet of chloritic and actinolitic quartz schists, which are succeeded by a very large succession of ferrugino-arenaceous banded actinolite schists, in appearance almost identical with the actinolite schists south of the Washington, Champion, and Republic mines.

Certain chloritic seams of the actinolitic rock-series are permeated with leaf-like agglomerations of crystal blades of margarite, which on the weather-worn surfaces of the rock project in high relief with a nacreous splendor; the same beds contain also black tourmaline crystals. Some of the actinolite seams contain a large percentage of magnetite, and in all of them small brown garnet crystals abound. Directly on the actinolitic series follow other banded, more slaty, and micaceous beds, dark green-colored by chlorite, or blackish by magnetite granules, which in some of the seams are also crowded with small garnet crystals; they are well exposed on the north side of the railroad track between the Michigamee and Spurr mines, and correspond to all appearances with the strata north of the quartzite belt of the Keystone mines; they are also intimately allied with the mica schist formation on the south-side of Lake Michigamee, opposite Michigamee village, which are doubtless the next succeeding deposits of the Huronian group.

The iron formation is traceable without interruption for six miles west of Michigamee to the vicinity of the three lakes; but I have only superficially examined these localities, and am not prepared to give a description. The ore-deposits of the L'Anse district I had not the time to examine.

NOTE.—These localities have been examined since, but I have to desist from their description at present as the printing of this report has advanced too far.

CHAPTER V.

ARENACEOUS SLATE GROUP.

I HAVE designated this large series of deposits with the name prefixed as a heading, because a large proportion of them consists of sandy siliceous layers in alternation with slaty argillitic rock-beds; but if we compare the strata representing this group in different localities, remote from each other, we will often find them to differ considerably. It is therefore impossible to select a name which would give a definite conception of the nature of all these multitudinous kinds of sediments. We find them incumbent on the quartzite formation, but sometimes also resting on the ore-bearing rock-group, and quite often in direct contact with the dioritic series.

The most eastern exposures of rocks identifiable with this group are found near the centre of the S.E. quarter of Sect. 6, Town. 47, R. 25. A thick belt of black, fine-grained slaty rock-beds, interlaminated with siliceous sandy seams, forms there with its vertically erected ledges the bed of Carp River, and its southern embankments; next south of the slates are high walls of massive light-colored quartzites in direct conformable contact with them, and also on the north side, at some distance from the river-bed, are large hills composed of the quartzite formation. In all the outcrops south or east of this place the quartzite formation is found to be overlaid with rocks representing the Huronian limestone formation, as we know from the previously given descriptions. Following from here the course of the river up stream, in a north-western direction, we find at intervals similar dark blackish-colored slates and lighter-colored arenaceo-ferruginous banded rock-ledges, exposed in the embankment and on the adjoining hill-slopes, but most of the surface is covered deeply with drift. Near

the N.E. corner of Sect. 1, Town. 47, R. 26, on the north side of the river, commences a conspicuous row of drift-hills, trending north-west, which terminates in the south part of Sect. 35, a half mile east of Eagle mills. The central nucleus of these hills is very probably formed of the banded silico-ferruginous layers of this group, as plenty of angular fragments of such rock are mingled with the drift-masses, and in one place an actual outcrop of the ledges is observable. A half mile north of the drift-hills are the limestone knobs of the Morgan furnace location; but as all the interval is covered with drift, we cannot inform ourselves of the relative position of the two rock-groups to each other. South of the drift-ridge is the swampy valley of Carp River, bordered on the south side of the river channel by highlands, slowly rising in several successive undulations until they culminate about a mile from the river in a row of knobs, formed of thick compact ledges of a light-colored quartzite, which dip under a high angle northward. The undulating, and to a great extent drift-covered north-slope is composed of a large succession of thin-bedded arenaceous-ferruginous flagstones, of slaty argillitic layers and of banded fine-grained siliceous ledges, interstratified with several larger belts of a granular sand-rock-like quartzite of whitish color, while the other rock-beds have a dark brownish or dirty greenish tint. This series has in part a conformable northern dip with the quartzites, but frequently also a southern dip is observed; the direct contact of the two formations is nowhere visible, on account of the covering drift-masses. The ferrugino-arenaceous schists, in considerably contorted corrugated condition, and dipping to the south, are well exposed in a rounded knob, a little distance north of the S.E. corner of Sect. 3, Town. 47, R. 26. A short distance west of this knob are other knobs composed of diorite, which are on their north side in close contact with the ferruginous flagstones and the associated beds. Examining the south slope of the most eastern of the before-mentioned quartzite knobs, over which the west line of Sect. 12 runs, we find it very brisk, full of cliffs; at the base is a swampy creek-bed running south-west, and across it, just at the quarter-post, rises another much lower rocky hill, formed of limestone. Taking from here a due west course, we pass over several low knobs of quartzite, which make part of the high ridge farther north; in the centre of Sect. 11 we descend into a broad swamp

valley, in which we remain until we come to the quarter-post on the west side of Sect. 11, close to the bed of a creek. Very few steps farther west we are at the base of a high ridge, the summit of which is composed of diorite; it extends from the west end of Goose Lake northward through the centre of Sect. 10, and terminates at the north end of the section. The slope of this diorite range is in all its circumference formed of the banded silico-ferruginous flagstones of the fifth group. We find them particularly well exposed at the south end of the ridge, near the west end of Goose Lake; the layers are in close contiguity with the diorite, and surround the base of high vertical walls of the diorite.

In the same locality the intersection of the diorite by a doleritic dyke is observable. A low hill, in front of the dioritic bluffs, which faces the Escanaba River near its entrance into Goose Lake with high vertical walls, exhibits a fine section through more than 200 feet of strata identical with those in contact with the diorite, and on the south side of the river channel an equally fine section, through the same banded silico-ferruginous beds, is seen in a cut of the Escanaba Railroad. The strata dip west, or north of west, under a moderate angle; east of these exposures the road-bed is cut through a large succession of lighter or darker gray-colored clay slates, with interstratified arenaceous seams which dip in the same direction, but under a much steeper angle than the former; and these are succeeded downward by a large belt of massive beds of light-colored compact quartzites, which form a high knob bordering the west side of Goose Lake, and are well laid open in a quarry close to the railroad. The upper, banded, flaggy rock-series amounts here to considerable thickness, as south of the railroad cut for a quarter of a mile all the surface is underlaid by rocks of this kind, somewhat differing from the others by lighter, more variegated colors, and representing another horizon of the series. Going from the exposures on the railroad westward on the track, we see the strata rapidly sink under the surface of marsh-lands adjoining the river channel on the west side. Farther on, after crossing the river a second time, the road passes for nearly two miles along the valley of this stream, which suddenly widens into marshy plains, bordered on the north side by a range of high hills composed of sand-rock and slate-beds analogous to the rock-belt exposed in the cut near Goose Lake, although they have not as much

similarity with each other as to allow an identification by the rock-character alone. South of the valley we observe an isolated knob of diorite at the place where it begins to widen, about 400 steps south of the N.W. corner of Sect. 15; west of it is a drift-covered ridge separated from the high quartzite range on its south side by a swampy depression; the slope of the quartzite range is deeply covered with drift. Farther west, beyond Partridge Creek, the valley is bordered by a high ridge of diorite; the west line of Sect. 16 runs over its highest summit knob. The slopes of this ridge, in its whole circumference, are composed of the banded arenaceo-ferruginous flags of the fifth group, but a large portion of the surface is covered with drift.

Near the west end of the ridge the lower hematitic rock-beds of the iron-formation come close to its base, but the contact between them and the flagstones of the fifth group is not seen. The high undulating plateau-lands on the south side of the ridge are underlaid by the arenaceous slate-group, which extends from there southward in a belt, hemmed in on the west side by dioritic rock-crests, on the east side leaning at the base of quartzite hills. In the centre part of Sect. 20 this rock-belt bends and is continued in a western direction, covering all the surface of Sect. 19 and the eastern diagonal half of Sect. 24 in the adjoining township, where it runs out into a point at the south-west corner of this section, close to the old Tilden mines. It reposes there on the ore-bearing rock-belt, as I infer from the relative position of the outcrops, although I have not observed the direct contact of the two rock-groups. The largest part of the rock-beds, representing the fifth group in the presently considered localities, consists of sandy, somewhat micaceous flagstones, or of finer-grained banded siliceous rock-beds, all more or less impregnated with protoxide of iron as a coloring matter, and with granules of magnetite. Some of the layers are almost rich enough to be used as iron-ores; such richer beds, dark blackish-colored by magnetite, are well exposed on the side of the railroad track near the N.E. corner of Sect. 24. The strata lie on the side of a diorite hill, dipping under a low angle toward it in a northern direction, and a few hundred steps farther east another body of these ledges lies in a slanting position on the diorite. Identical beds are observable near the N.E. corner of Sect. 18, Town. 47, R. 26; likewise in contact with diorite and

dipping toward it. In this latter locality they are conformably underlaid by a compact dark greenish colored rock, which by weathering turns rusty brown, and becomes full of small cavities; it effervesces with muriatic acid, which dissolves the small crystals of sparry carbonate of iron abundantly disseminated through the mass; the green color is due to chloritic scales. By boiling the rock with muriatic acid, it decolorizes, and a glistening cavernous white mass of micaceo-feldspathic composition remains. Good outcrops of this rock are observable a short distance south of the mentioned locality, in which its contiguity with the magnetic lean ores is not so well exposed; near the centre of the S.E. quarter of Sect. 17 a similar rock is found in connection with the flagstones of the fifth group.

In the above-mentioned exposures, near the N.E. corner of Sect. 18, we find, on a lower level of the hill-slope, the hematitic flags of the ore-formation uncovered in test-pits, which likewise dip toward the hill in conformity with the black magnetic ledges on the higher part of the slope. The just-described rock-series representing the fifth group, is, in the south part of Sect. 19, underlaid by a large body of massive layers of a granular, partly conglomeratic quartz-rock, with a chloritic cement; and therefore dark blackish-green colored, and from a distance is easily misapprehended for a dioritic rock; it is most favorably exposed for observation in the N.E. quarter of Sect. 30, Town. 47, R. 26, and is seen there to repose in apparent conformity on the altered granite-like quartzites, which compose the previously described rock-bluffs in the centre of Sect. 30. Equivalent, but less altered conglomeratic quartzites form a half mile farther south the hanging wall of the ore-belt in the Palmer mines.

Returning from here to the north margin of the examined belt of the fifth group, we find deposits of this kind to extend northward along the narrow valley of a creek which takes its head-waters from the S.W. quarter of Sect. 7, and passes near the south quarter-post of Sect. 18. The exposures in the S.W. quarter of Sect. 7 are particularly fine; the siliceous flags richly impregnated with magnetite, and amounting to a belt of considerable thickness, lean with western dip in steep inclination on the side of bubble-shaped hillocks of diorite. We are here on the top of a high plateau crowned with numerous dioritic ridges in parallel rows trend

ing from south-east to north-west ; and in every valley intervening between these ridges we meet again with these ferruginous flagstones. Exposures of this kind are found in the N.E. quarter of the S.W. quarter of Sect. 7, which continue into the S.W. quarter of the N.W. quarter of the same section, and into the S.E. quarter of the N.E. quarter of Sect. 12 in the next adjoining township, from whence the strata can be traced almost without an interruption to the N.W. quarter of the N.W. quarter of this section into close proximity with outcrops of the lower hematitic beds of the iron-ore group.

Other large deposits of this group are found in an adjoining parallel valley, which sends a creek westward into the small lake at the quarter-post of the west line of Sect. 12. The strata lean on the north side of the valley, steeply inclined on the hill-slope, in apparent conformity with the underlying dioritic rocks which compose the body of the ridge. On the south side, which is likewise bordered by a dioritic ridge, the strata, retaining the same direction as on the north side, dip toward the hill whose slope they form. Going across this ridge, we find its south slope again surrounded by a belt of these ferruginous flagstones. They are well exposed near the N.E. corner of Sect. 14, on the side of a wagon-road leading to the Ogden and Tilden mines, and farther east in the N.W. quarter of Sect. 13, up on the higher part of the hill-slope, and down near the shore of Lake Miller. As we pass along on this road to the south end of Lake Ogden we leave the area invested with deposits of the fifth group, and descend unto the older sediments of the iron-ore group, which are here well exposed in the now abandoned pits of the Ogden, Tilden, and Iron Mountain mines, of which I have spoken in the previous chapter.

We have followed in the preceding exposition the exposures of the fifth group southward and westward, after we left the track of the Escanaba Railroad, which was the starting-point, but we find the much larger display of this formation on the north side of the road. It covers, in Town. 47, R. 26, in one continuous body, almost the entire surface of Sects. 2, 3, 4, 5, and 9, besides the east halves of Sects. 6 and 8, and in the adjoining northern Town. 48, R. 26, the south halves of Sects. 31, 32, 33, and 34 have the same rock-series as a surface rock.

A large portion of the deposits in this northern part of the

formation consists of clay slates with interstratified larger belts of light-colored quartzites of a granular sand-rock structure, sometimes quite compact, of a vitreous fracture, other times porous ; but the silico-ferruginous banded rock-beds and flagstones seen in the before-described localities are also represented in this part.

The chain of hills which commences at the previously mentioned cut of the Escanaba Railroad through a bubble of the lower beds of the ore-formation, and trends from there eastward across the north part of Sects. 8 and 9, is formed of a belt of white quartzite in association with clay slates of a more or less arenaceous character. In some of the outcrops the quartzite is rather porous, and incloses an abundance of angular slate fragments ; its ledges are much corrugated. In other exposures we find the rock intermingled with martite granules, and in blotches tinted red by hematitic pigment, again in other places the rock is compact and has a glassy fracture. The hill north of this ridge, close to the north line of Sect. 8, consists of siliceous sandy flagstones, blackish colored by an abundance of magnetite granules ; a few hillocks farther north, in the S.W. quarter of the S.E. quarter of Sect. 5, are composed of clay slates in a nearly vertical position. By crossing over this large body of hills, which on its north side is bordered by a broad belt of swamp, we see a constant alternation of rock-beds of the described kind, without any observable regularity in the succession, a large portion of the surface is covered with drift, and in all probability numerous plications cause a repetition of the strata, which I infer from the much-corrugated condition of the rock-beds in many of the exposures. A short distance north of the quarter-post on the east line of Sect. 5, the section line crosses a narrow rock-ridge about 60 feet high, which is isolated from the other hill-lands in all its circumference by a strip of swamp. It consists of nearly vertical ledges of sand-rock and clay slate, with some intermingled ferruginous seams. East of it are other good exposures of light-colored massive quartzite beds in alternation with clay slates ; the strata there dip north. The north half of the S.E. quarter of Sect. 4 is occupied by an insular cluster of high diorite hills, surrounded on all sides by the arenaceous beds of the fifth group ; on the north side of these hills, which are quite steep, we find the sandy flagstones dipping north under a high angle. At the coaling kilns, situated on the

east line of Sect. 4, the arenaceous flags are dark blackish-colored and contain a large proportion of magnetite; a part of the beds is more argillitic, and merges with the clay slates which occur in association with the flags.

The nature of the eastern continuation of this rock-series in Sects. 3 and 2 has been described in a previous page of this chapter. I proceed, therefore, from the coal-kilns northward across a strip of swamp, and find on the north line of the N.E. quarter of Sect. 4, on the summit of an insular body of high land, three small knobs of a coarsely crystalline diorite, surrounded on all sides by the sand and slate rock-ledges of the fifth group, and in close contiguity with them. The old State road to Marquette passes close by the knobs, and the nearly upright sandstone and slate beds dipping north, are well exposed in the road-bed, and in some knobby protrusions north of the road. The greatest part of the hill-side sloping down toward the bed of Carp River is covered with drift, but on the opposite side of the river-bed the sand-rock and clay slate beds come again to the surface, and form an isolated chain of hills completely surrounded by marsh-lands; the strata have likewise a northern dip under a high angle. The west line of Sect. 34 runs over the highest part of this ridge, and the quarter-post is at the base of the north side of the hills, not many steps from the Houghton and Ontonagon Railroad, which leads through the marsh-lands. Following the railroad westward, we find, near the quarter-post of the west line of Sect. 33, another isolated knob in the marsh, consisting of the same sandstone and slate beds as the ridge east of it, and dipping north as they do. A few hundred steps north of this knob rises the high quartzite range which extends westward to the north side of Teal Lake, and has been described in a previous chapter; its rock-ledges have a southern dip, as will be remembered. Along the south side of this ridge extends, from this point westward, a broad belt of drift-covered undulating plateau-lands, from which in various places the underlying rock-beds of the fifth group project in low crests and knobs over the surface. In the exposures next to the quartzite ridge the strata have a southern dip conformable with it, but between the outcrops of the two rock-groups is always an interval several hundred feet in width, covered with soil and loose material, and their contact is not seen. The very large belt

of this younger rock-series consists, in the proximity of the quartzite range, of an alternation of clay slates with belts of sand-rock, which often has a density and compactness little inferior to the glassy quartzites of the high ridge on their north side; the clay slates are in part light gray-colored, others are dark blackish or lighter and darker sedimentary streaks alternate with each other in the slaty layers, whose cleavage is always contrary to this sedimentary striation; some of the slaty layers contain an abundance of small mica scales. Very good exposures of the clay slates are observable along the railroad where it intersects the west part of Sect. 32; others are found south of the centre of Sect. 31, near the wagon-road leading across the hill-range to Carp River. On the plateau we noticed, near the centre of the S.E. quarter of the N.E. quarter of Sect. 6, an old test-shaft with a large pile of rock thrown out of it, which consists in part of the ordinary clay slates, in part of banded fine-grained silico-ferruginous rock-beds, impregnated with magnetite granules, and of other silico-argillaceous rock-beds brightly red-colored by hematite, and also of dark green-colored chloritic schists. I am informed by the parties who dug this shaft that the strata had a northern dip, and that the hematitic rocks and chloritic schists were the lowest. We have here an instance of a direct superposition of the arenaceous slate-group on beds which evidently represent the lower strata of the iron-ore group, and the schistose chloritic beds, on which they are generally found to repose in the neighboring hematite mines on the other side of the valley, and from this we also may infer a considerable thinning out of the iron-bearing series, as the shaft, which in all is not over 300 feet deep, went clear through it and struck the usual foot-wall of the formation. South-west of the test-shaft, near the edge of the plateau, are several knobs formed of highly ferruginous, dark-colored siliceous flags, and of thicker banded ledges, which are identified with the large series of rock-beds intersected by the Escanaba Railroad near the west end of Goose Lake. They dip under a high angle northward, and seem to represent the upturned ends of the lowest beds in this rock-belt, which do not appear at the surface near the foot of the quartzite chain, and probably would be found to fill out the covered interval which, as I stated before, always exists between the slate exposures and the quartzites of the high ridge. The same strata are seen

exposed farther east, near the quarter-post on the west line of Sect. 5. West of the mentioned knobs, in the centre of Sect. 6, is an isolated rounded hillock composed of dioritic rocks, partly schistose, partly massive, crystalline, and about a quarter of a mile north-west of it, the previously mentioned ridge of diorite commences, which extends in a straight western direction to the west line of Sect. 2, 300 steps south of the north-west corner, near the Teal Lake hematite mines.

The occurrence of the slates and sand-rocks of the fifth group in that locality, and their contiguity with the hematitic strata of the iron-ore formation, is previously described. We find there the banded silico-ferruginous flaggy layers farthest north, close to the shore of the lake, and evidently holding a lower position than the slate and sand-rock layers. This rock-series extends in all probability across the south part of Sect. 34 to the north part of Sects. 4 and 5, but in Sect. 34 no outcrops of it are seen, on account of the heavy drift-deposits, while in 4 and 5 they occasionally are on the surface; we find them exposed in the N.W. quarter of the N.W. quarter of Sect. 4, in the S.W. quarter of the N.E. quarter of Sect. 5, and in test-pits in the N.W. quarter of the N.W. quarter of Sect. 5. The sandy ferruginous flags which compose the hills north of the Excelsior mine, and those leaning on the north side of the diorite knobs in that locality, I consider members of this rock-group.

Before I follow these deposits farther westward, I have to return to some more eastern exposures which I had omitted to describe, in order to adhere to the adopted system of giving the descriptions by guiding the reader in a certain direction from place to place without deviations.

On the south side of the large cluster of diorite knobs, north of the New York mines, we see generally the ore-formation in direct contiguity with the diorite; but on the higher part of one of the rock-bluffs another kind of stratified rock is found to repose on the diorite in seemingly discordant position. It is quite a large succession of well-laminated, banded, rather thin ledges of a dark rusty color, which consist of a radiated agglomeration of minute needles of actinolite, with intermixture of some proportion of granular magnetite; the ledges are intersected by numerous narrow asbestine seams vertical to the lamination or more irregular, and

are in some places of the outcrop considerably plicated and corrugated, or else shattered into irregular fragments. Portions of the exposed rock-faces are covered with a thin crust of malachite recently deposited there from decomposing nodules of copper pyrites inclosed within the rock-mass.

A mile west of this locality we find analogous beds on the slope of the diorite hills north of Lake Bancroft. Close to the north end of the lake are some lower hills formed of slate-rocks of dark blackish-brown color, very tough, of great specific gravity from a large proportion of magnetite disseminated through the mass. The cleavage of the beds is in discordance with the sedimentary lamination of the rock, which is well observable, as the slabs of slate are banded in that direction with streaks of different shades of color. The strata dip southward under an angle of about 60° ; they are intersected by seams of long-fibred asbestos in intermixture with quartz; some of the asbestos is flexible, but most of it rigid, wood-like. Next under them follows a massive, very tough, dark greenish-gray rock of crystalline structure, which contains a large proportion of carbonate of lime in bladed, acicular crystals, which on weathered surfaces of the rock become washed out, and leave in their place linear slits, intersecting each other in every direction. This rock is inseparably allied with the dioritic rocks, of which the main body of the ridge is composed, and consists of a granular feldspathic ground-mass in intermixture with delicate, partly chloritic, partly micaceous scales, and with a fibrous hornblendic mineral, in which mass larger bladed crystals of feldspar are dispersed. Another undulation of the hill-side next north of the slate-hill is composed of banded ferruginous and actinolitic rock-beds, which cleave with very even surfaces in accordance with the sedimentary lamination, but parts of the ledges are bent and twisted into serpentine curves; their dip is southward, like that of the slates, and the same massive diorite-like rocks underlie them. These actinolitic beds, richly impregnated with magnetite, I considered as analogous with the banded silico-ferruginous rock-beds exposed in the railroad cut near Goose Lake, and to those composing the knobs north-east of the furnace of Negaunee; but later observations convinced me of a much greater analogy existing between them and the actinolite schists below the ore-formation of the Saginaw, Washington, and Republic mines; still as actinolite

also occurs sometimes in the strata of the fifth group, I am not positive whether they represent these upper or the lower schists.

South of the exposures of ferruginous slates and actinolitic flagstones on Lake Bancroft, are knobs of diorite on both sides of the lake, which are not incumbered with these ledges; not far from their base the ore-formation is struck by test-pits and by the diamond drill. The diorite forming the nucleus of the knob on the west side of the lake is coarsely crystalline, and consists, besides the hornblende and feldspar, of a good proportion of epidote; with the massive rock occurs a large body of schistose beds of chloritic and hydro-micaceous character, some of which are crowded with red feldspar crystals dispersed through the mass, with a copious intermixture of granular octahedric crystals of martite. The knobs on the east side of the lake are composed of a different sort of crystalline rock, which consists of red feldspar crystals imbedded within a chloritic cement mass abundantly interspersed with granules of iron pyrites, which causes its rapid decomposition and disintegration on exposure; in intimate connection with the crystalline masses are well-laminated chloritic schistose strata.

Proceeding with the examination westward, we find the fifth group as a surface rock of a very extensive area, but frequently so much covered with drift-masses that the accurate relative position of certain rock-belts of the series can rarely be ascertained for want of connected exposures.

The occurrence of arenaceous and slaty deposits representing this group in the S.W. corner of Sect. 4 is mentioned; analogous beds underlie the drift-covered hills west of the Lake Superior mines, in the S.W. quarter of Sect. 9, and cover all the surface of Sects. 7, 8, 17, 18, and the west half of Sect. 16 in Town. 47, R. 27. If we start from the Parsons mines and follow the railroad which connects the mines with the main road, we pass the quartzite knobs west of the mine, and through a swampy valley densely strewn with large angular blocks of the same quartzite, we come to a crossing with the wagon-road to the Saginaw mines, and then take that road. Ascending the hill we soon see outcrops of thick ledges of white quartzite, which dip in a north-western direction; above them follows a large series of flaggy, ferruginous, and micaceous sand-rock ledges, which occupy the whole hill-side west-

ward, and are particularly well exposed in a cut of the railroad to the Saginaw mines, near the quarter-post on the west line of Sect. 16, where the strata have an almost southern dip, and are likewise seen to repose on a thick belt of a white compact quartzite. As we approach on the wagon-road the summit of the hill, we see it diagonally intersected by a thick belt of a bluish gray, dark-colored, banded siliceous rock similar to the banded siliceous beds of the fifth group in exposures of other localities. Farther on the strata disappear under the drift, and are not seen again until we come to the first houses on the Saginaw mining location, where the quartzite formation is well exposed on the roadside in massive compact ledges. The actinolitic silico-ferruginous rock-beds composing the knob on the south side of the Saginaw mines, considering their great lithological affinity with the sediments of the fifth group, I at first erroneously identified with this group, but I am fully convinced at present of their conformable succession below the ore-formation.

In the north part of the above-indicated area, over which the fifth group extends, I could not find any outcrops within the limits of Sect. 8; the drift-masses contain there an abundance of large boulders of brecciated specular ore, similar to the ore of the Excelsior mines. In Sect. 7 the larger portion of the surface is covered with drift, but we find exposures in various parts of it.

An outcrop of fine-grained black slates, which dip under a high angle southward, is observable on the south line of the section a quarter mile west of the east corner; other quite extensive exposures of dark blackish, siliceous, even-bedded flagstones, which carry a large proportion of magnetite, are seen farther on toward the west corner of the section, near which locality a large belt of a dark-colored, compact quartzite has been denuded in test-pits in association with the magnetic flagstones; the dip of the nearly vertical rock-beds is also here southward. In the north part of the S.W. quarter is a row of hills composed of these black magnetic flags interlaminated with slaty and arenaceous seams, which present themselves well denuded in cliffs. West of them is another rocky hillock, in the N.E. quarter of the N.E. quarter of Sect. 12, of the adjoining town, range 28, which consists of lighter greenish-gray colored, somewhat variegated, and much-corrugated seams

of micaceo-argillitic, ferruginous, and arenaceous character, all firmly united into a bulky banded rock-mass.

A chain of hills continues along the section line from the S.W. corner of Sect. 7 to the S.W. corner of Sect. 12, which consists of an alternation of dark slaty seams with thick belts of compact granular quartzites of various darker or lighter shades of color; with them occur also brecciated layers composed of slate and quartzite fragments. The strata dip south, but are nearly vertical. I have to mention yet a chain of rock-exposures along the north line of Sect. 7, which consists of peculiar schists and other more massive beds, some of which have an amygdaloid structure; within a dark gray or greenish-colored chlorito-argillitic or feldspathic ground-mass are flattened nodules of the size of a lentil, or somewhat larger, copiously imbedded, which in part consist of calcspar or sparry carbonate of iron; others are a granular white quartz, and still others are filled with radiated blades and fibres of a soft talcy, green-colored mineral. In the not-amygdaloidal beds the carbonates of lime and of iron are found abundantly disseminated through the mass in small crystals. The amygdaloid beds are best observable in the hills south of the Excelsior mine, in the N.E. corner of the section; farther west, near the quarter-post of the north line, the outcrops are much larger, but the amygdaloid structure of the rock is not so well developed. The dip of the strata is south; a short distance north of the quarter-post is a small quartzite knob, with ledges dipping in the same direction, and north of the knob succeed the jaspery beds of the iron-formation which have been mentioned in the previous chapter. From this relative position of the different rock-seams I infer the age of the schist to be the younger, and the other arenaceous and ferruginous strata in the south part of the section seem to follow them conformably in the ascending order; but south of these, in the north part of Sect. 18, and of Sect. 13 next to it, we meet again with a very large body of rocks resting in apparent conformity on the others, which, although not exactly identical with the amygdaloidic schists, resemble them so much that I consider them as analogous. The amygdaloid rocks in these exposures have a more perfect crystalline structure, and contain more mica in their composition; intimately associated with them are massive rocks of brecciated structure, composed of a similar dark-colored chlorito-feldspathic mass,

intermingled with calcspar or carbonate of iron, and not rarely with small crystals of black tourmaline; the brecciated rock is not always formed of intermingled fragments of different kinds of rock, but often consists only of an agglomeration of nodular lumps of the shattered rock, recemented by the same substance. Rocks of this description form the large bluffs north of the Saginaw Railroad depot, across the bed of Carp River. Half a mile farther west the analogous beds exposed close to the railroad track inclose an abundance of foreign rock-fragments, of slaty argillites, or of banded siliceous ledges, united with the mass in a mode which suggests to me its semi-fluid condition from the effects of heat, at the time the rock-fragments became imbedded in it; but the very plain sedimentary striation exhibited by the conglomeratic bluffs is, on the other hand, a positive proof of the origin of these sub-crystalline, much-altered rock-masses from sedimentary deposits.

On the line between Sects. 13 and 18, if we follow it from the railroad northward, we find, after ascending the front part of the hill-side, which is formed of the dark-colored, diorite-like, massive brecciated rocks, a second undulation of the range, formed of lighter-colored, more distinctly stratified beds, which are almost rich enough in sparry carbonate of iron and lime to pass for a siliceous limestone; the decomposed rock forms bright orange-colored, ochraceous, cellulose masses of soft friable condition. These rocks, of the conglomeratic and amygdaloid kind, are in the N.W. quarter of Sect. 13 in immediate contact with black-colored, very even-bedded flaggy layers in a vertical position, which form a small knob, partially intersected by the railroad. South and west of this knob only crystalline diorites are found in the exposures. On the north side of the railroad, near the last western bridge across the river, are fine exposures of the conglomeratic rocks, partly in massive, bulky form, partly schistose and rich in micaceous scales. The hills composed of them are on the west side closely adjoined by other hills, composed of dioritic schists and of massive coarsely crystalline diorites, which two sorts of rock are denuded in almost uninterrupted continuity on the south slope of this row of hills, and perfectly merge into each other, evincing in this case the original deposition of these altered sediments on the diorite, and not a merely accidental contiguity resulting from the

dislocation of the strata, although they are in neighboring localities, to all appearances, reposing on the quartzites above the iron formation. In Sect. 11, Town. 47, R. 28, the south-east corner is located on the terminal knob of the quartzite ridge, which follows the south line of Sect. 12. About 500 steps north of this knob, across the bed of Carp River, emerges an isolated knob above the drift-covered rolling lands, composed of black slates in an upright position, which are at least 300 feet in thickness; the south-west corner of this section, near the Greenwood furnace, is situated on a hill composed of the same black slate interstratified with quartzose seams; north of it, for three quarters of a mile, no outcrops occur, but in the north half of the N.W. quarter of Sect. 11 the black slates are again largely exposed in a row of hills bordering the north side of the wagon-road leading to the Greenwood and Clarksburg furnaces. The slates dip south under a high angle. North of the slates rises another higher hillock, consisting of banded ferrugino-arenaceous rock-seams firmly united into a bulky solid rock-mass, which exhibits an extreme degree of corrugation and torsion, made very conspicuous by the banded variegated structure of the rock. The iron formation must pass not a great distance north of this hill in its western extension toward the Boston mine, but there are no outcrops.

At the Clarksburg station the railroad passes a knob, composed of the altered sub-crystalline and occasionally amygdaloid rocks, encountered before on the side of the railroad in Sects. 18 and 13; they are here not conglomeratic, quite compact, partly of fine homogeneous grain, partly of coarser, crystalline structure. The carbonates of lime and iron are regular constituents of the rock, largely composed of a feldspathic ground-mass, with micaceous and chloritic scales dispersed through it; and in some of the collected specimens clusters of actinolite fibres are interspersed. Opposite this knob, toward the furnace buildings, the banded arenaceous-ferruginous beds, in a much corrugated condition, can be seen in close contact with the just-mentioned rocks. The hills north of the station are formed of a large succession of sandy flags and schistose, more or less chloritic and ferruginous rock-beds, representing the fifth group; they dip northward. Farther north a broad belt of surface is occupied by black slates, but the denuda-

tions are very imperfect. North of them, separated by a swamp, is the Boston mine. On the south side of Clarksburg are likewise the slaty and banded siliceous rock-strata of the fifth group—the general surface rock; they dip north, and beneath them the ore-formation is found, as previously stated; but on the higher part of the hill-range the same strata dip south, and above them succeeds a large belt of quartzite; next above it is a belt of specular iron ore partly granular, partly of slaty structure, in connection with hydro-mica schists, more or less impregnated with magnetite granules.

The summit of the ridge and its south slope are composed of an immensely thick series of banded but very tough and compact actinolite schists which on their part come in direct contact with diorite hills farther south.

All the described succession of rock-beds dips south, and as it represents the regular succession observed in other localities, but in inverted order, we have here before us a large overturn of the formations exhibited.

From Clarksburg to the Washington mines the rock-beds of the fifth group occupy both sides of the valley, presenting continued large exposures of this remarkably great succession of strata. The finer-grained schistose sediments prevail here over the coarser quartzite rock-belts, which, east of the Greenwood furnace, and generally in the eastern exposures of this formation, compose the greater bulk of it. At the Washington mine no rocks of this kind are visible in the hill-slope, but in the east half of Sect. 1 the outcrops of a large succession of micaceo-chloritic schists, dipping under a high angle southward, commence near the bed of the Escanaba River, and the strata continue to be exposed in a number of undulating hillocks south of there, without much change in their character; nearer the high hill-range where the dip of the strata is reverted into a northern, fine-grained arenaceous beds, with some magnetite granules and mica scales disseminated through the mass, begin to alternate with the schistose seams, and arriving at the base of the highest part of the hill-range, we find a large body of very even-bedded banded granular quartzose beds richly impregnated with granules of magnetite, leaning with steeply inclined beds on the side of the ridge, whose summit is formed of a diorite similar to, if not identical with, the

diorite cropping out on the south side of the Franklin pits of the old Washington mine. This hill is separated from the hill of the Washington mine by a deep ravine; close to the ravine a tunnel is driven on the mining property into the hill across a thick succession of micaceous quartz schists and of quartzite layers; no trace of these, or of the ore-formation, is observable on the east side of the ravine. Major Brooks considers, therefore, the chloritic schists and the banded ferruginous beds above mentioned as older than the iron-ore group; but the direct connection of these schists with the schists on the north side of the Escanaba valley, which are indubitably higher than the quartzite and the ore-formation, and their lithological resemblance with rock-beds of the fifth group in other localities, are sufficient proof for me to take them as such. The hills on the north side of Escanaba River, north-east of the Washington mines, are a direct continuation of those near Clarksburg station, and consist, like these, partly of schistose micaceo-argillitic beds, partly of the harder sub-crystalline and frequently conglomeratic rocks, of chlorito-feldspathic composition, with a good proportion of the carbonates of lime and iron. On the rolling drift-covered lands, north of the first-encountered rock-bluffs near the river, which are formed of the just-mentioned kind of rock-beds, dipping south under a high angle, we find numerous limited exposures of slaty micaceous and chloritic argillites and of banded siliceous rock-beds, most all of which inclose some granules of magnetite. These beds, amounting to a great thickness, are often considerably corrugated or bent into larger serpentine curves. Farther north succeeds a very large belt of fine-grained black slates, colored by carbonaceous matter, but only the general local distribution of all these deposits is observable, and about their order in the succession, their thickness and dip, nothing definite can be determined in this part of the country, where drift hides the rock ledges almost completely. West of the Washington mines we find, by following the railroad toward the Champion mines, the hills on both sides of the valley consisting of the same kind of rocks which compose the rock-ridges north of the railroad, between Clarksburg and the Washington mines; the black carbonaceous slates are in several places denuded on the road track. Near the Keystone mines the superposition of a schistose conglomeratic rock-belt, on the quartz-

ites above the ore-formation, is observable in the most unequivocal distinctness. These outcrops, in connection with the ore-bearing rock-series of this locality, have already been described. The identity of this conglomerate rock with the conglomeratic rocks encountered all along on our westward course, and first met with in the bluffs north of the railroad depot of Saginaw station, is not only indicated by a great similarity of the structure and of the material of these strata, but an almost direct continuity in the extension of this group of deposits can be traced.

In this locality we find for the first time, in this particular horizon of the rock-series, actinolite and garnets as constituents, which minerals are very abundant farther west, not only in the layers identifiable with these, but they are found to be equally common throughout the whole series of the ore-formation of the Spurr and Michigamee mines. In the higher strata of the mica schist group, garnets are likewise characteristic, and actinolite enters into the composition of some of the rock-seams. For this reason I suggest that at one time within the concerned area local conditions existed which were favorable for the formation of such minerals from the sedimentary material, while in other localities the same sediments were not altered in this way.

The large series of actinolitic rock-beds found south of the Washington mines, the Keystone, Champion, and Republic mines, in a position which evinces their older age than the ore-formation, has been previously described in connection with the ore-formation of these localities. In consideration of the total absence of such rock-beds beneath the ore-formation in the Michigamee and Spurr mines, and in most of the other mining localities, and of the occurrence of a rock-belt almost identical with these strata, resting in conformity on the ore-formation and on the quartzites in the Michigamee mine. I for a time believed in the equivalency of these similar deposits, and placed them all above the ore and quartzite formation, which position is unquestionably held by the actinolite beds of the Michigamee mine; but I have fully satisfied myself by subsequent careful examinations that the actinolite schists form two different belts, one above the ore-formation and another below it.

It remains yet for me to say that the schistose rock-ledges, analogous with those composing the north slope of the hills at the

Keystone mine, form also the opposite-hill slope near the Champion furnace, and continue westward to the north side of Lake Michigamee, where they are for a while interrupted by older rock-beds coming up to the shore; but they appear again on the west side of the Michigamee mines, and hold there a position next above the upper actinolitic rock-belt. The next succeeding strata, exposed on the south shore of the lake, will be described in another chapter as a separate group.

The Dead River, while it flows within the limits of Sect. 14, Town. 48, R. 26, has a narrow, ravine-like valley, bordered on the north side by a high ridge of granite, on the south side by steep drift-bluffs which form the edge of a high plateau. At its exit from the eastern limits of the section it rushes in small cascades and rapids over granitic rocks, which at this place form a narrow gorge with vertical rock-walls on both sides. Higher up stream the granitic rock-bluffs recede some distance from the channel on the north side, and are on the south side replaced by the mentioned drift-bluffs. Near the middle of the section the granite bluffs recede still farther north, and the river channel is, to the west end of the section, bordered by lower hills, which form a belt on the slope of the granitic summits. These hills are composed of quartzites and arenaceous flagstones, interstratified with argillaceous slaty rock-beds, which dip north toward the granite. The quartzites, of a purplish color, are partly thick-bedded, compact, with glossy fracture, but have a distinctly granular structure; other beds are more thinly laminated, finer-grained, and more porous, with a scaly micaceous or chloritic cement, or are more of an argillitic nature and merge into a slate-rock, with a cleavage discordant with the sedimentary stratification; these slaty and thin-bedded arenaceous beds have a dark gray color.

Farther up the river, in Sect. 15, we find in its embankment, and on the slope of the drift-covered terrace-lands, intervening between the river-bed and the granitic summit part of the hills, exposures of a large succession of black carbonaceous rather hard slate-rocks, interlaminated with more massive fine-grained silico-ferruginous layers of a banded structure, which is particularly obvious on weathered surfaces of the rock, as the darker and lighter-colored alternating seams contrast more than on fresh fractures. The strata dip north. We can follow from here

for two miles up the stream continued outcrops of similar black slates and banded siliceous ledges in the steep bluffs which border the river on both sides. Near the S.W. corner of Sect. 9 we arrive at the base of a cascade, in which the river leaps down over walls 90 feet high, formed of these dark carbonaceous slate-rocks, lifted up in all degrees of inclination, and bent into serpentine curves. This excessively disturbed condition is caused by a large doleritic dyke 42 feet wide, which here intersects the rock-ledges transversally. The outer part of the dyke is an amorphous black-colored homogeneous mass, which by degrees assumes a more and more perfect crystalline structure toward the centre of the rock-belt, consisting there of a magma of well-formed crystals of a white glassy feldspar and of a dark brown augite. The extensive exposures of slaty strata at the falls and in the hills south of the river give a conception of the great thickness of this rock-belt, but the folded condition of the strata allows no exact estimate of it; we see local dips in any direction, but generally a northern dip is observed in the outcrops.

South of the falls the slates continue for the distance of half a mile to form the surface rock, until we approach, in the south half of Sect. 17, the north slope of a granite-range which intersects the centre of Town. 48, R. 26, from east to west; here succeeds them conformably a series of arenaceous flagstones full of mica scales, brownish or reddish-colored by iron oxide; under them follow thick-bedded compact quartzites, part of which is conglomeratic.

These beds form the slope of the granite ridge, on which they lean with northern dip. A similar succession of rock-beds is exposed in numerous places, if we follow the slope of this granite range westward. The lower thick-bedded quartzites are not always found, and in their absence the next higher micaceous sand-rock flags lie directly on the granite, or sometimes on the schists of the dioritic group, which, in a part of the range, cover the granitic nucleus. The black slates above the sandy flagstones have a very great surface extension; they compose a broad belt of undulating hill-lands on the south side of Dead River, which extends from the falls without interruption to the west line of R. 26 of Town. 48, and spreads in the next Town., R. 27, still farther, occupying almost the whole north half of the township. We

find them often interstratified with arenaceous rock-seams; it must be remarked, however, that drift deposits are ignored in this consideration, or else the slaty rock-beds would have an inferior claim to the configuration of the surface.

A mile north of the falls of Dead River, in the S.E. corner of Sect. 5, Town. 48, R. 26, I found an isolated mass of banded silico-ferruginous rock-beds, consisting of lighter-colored jaspery seams, containing a small proportion of granules of magnetite, in alternation with darker-colored ones intensely impregnated with ore-particles, which are perfectly similar to the banded siliceous rock-beds of the fifth group in the environs of Negaunee. This succession of ledges, amounting to more than 40 or 50 feet in thickness, is wedged in between dioritic crystalline rock-masses, of which all the surrounding country is formed. It is, in all probability, a detached body of layers which, during the general upheaval, became inclosed within a fold of the dioritic rocks, and so escaped destruction during the drift period, or at any other time in which erosive forces acted on the surface. Another instance of the intercalation of still larger masses of ferrugino-siliceous banded rock-beds, inclosing narrow seams of magnetic flag-ore between the dioritic rock-series, occurs at the location of the Holyoke mines, in the N.W. quarter of Sect. 1, Town. 48, R. 27. They are found on the middle of the slope of a high ridge of dioritic rocks on the north side of Dead River, irregularly wedged in between the vertically erected beds of fine-grained dark-greenish gray schists, with a glossy surface and an almost perfect slaty cleavage, which are intimately connected with crystalline dioritic masses. A tunnel has been driven into the base of this hill-side through these schists, and the flaggy, ferruginous beds were there encountered. These were, however, not the object searched for by the miner. The dioritic rock-masses are here intersected by large veins of milk quartz, which carries some argentiferous galena, zincblende, copper and iron pyrites, in association with the sparry carbonates of iron and lime; but the amount of valuable ore is not large enough to be profitably mined, and the location, on which at one time large sums of money have been expended to develop it, is now abandoned.

All the hill-lands south of the Dead River, near the mining location, are composed of the black slate-rock. A large tributary

enters the river at this spot, which comes from the south-west; on the east side of this valley is a very broken, densely-timbered country, which continually rises as it recedes from the river, until it culminates a little over a mile's distance in high knobs with very steep slopes, formed of highly inclined layers of black slates dipping northward, which amount to an enormous thickness. On the south side of these knobs, the largest of which is situated near the centre of Sect. 12, and others in the S.E. quarter of Sect. 11, the arenaceous flagstones underlying the slate-formation come to exposure. On the west side of the mentioned tributary river the surface is less broken, rising first slowly into a drift-covered, undulating terrace, sparingly covered with pine, and partly clear of all timber, and then a more brisk ascent brings us to the summit of this body of hill-land, which is found to be isolated in its whole circumference. East and north are the valleys of Dead River and of its tributary branch; west and south a broad belt of marsh surrounds it. Nearly all the surface of this insular surface-elevation is covered with drift, but outcrops of the underlying slaty rock-beds are frequent enough in all parts of it to show the uniform structure of its nucleus of this kind of sediments. On the north-western slope of these highlands, in the N.E. quarter of Sect. 4, Town. 48, R. 27, are exposures of such slates with a fair-even cleavage, which induced the owners of this land to open various test-pits in search of a good quality of roofing slate, but, so far, they have not had success in finding any beds well qualified for this purpose, and as far as I examined the locality I can give no encouragement for continuing the explorations. West from here, on the other side of the broad marsh-belt surrounding this place, is another body of drift-covered hill-lands, which occupies the west half of Sect. 5, all of Sect. 6, the north half of Sect. 7, and the adjoining Sects. 1 and 12 of the next Town. R. 28. These are underlaid by identical slaty and arenaceous rock-beds, but outcrops are comparatively rare; farther south, in the western part of Sects. 13 and 24, frequent exposures of the black slates, underlaid by ferruginous and somewhat micaceous sandy flagstones, are observable, and the superposition of the latter on thick-bedded compact quartzites, similar to the quartzites of the Teal Lake range, is seen in various localities; as, for instance, near the N.W. corner of Sect. 24, which is located on the

ferrugino-arenaceous rock-belt, well exposed in this locality; and the higher part of the hills, situated in the adjoining Sects. 14 and 23, consists of quartzites of a coarse-grained structure, and partially in a brecciated form; other strata of the quartzitic rock-belt are sub-schistose by abundant intermixture of the granular quartz-mass with pale yellowish, scaly, hydro-mica of a silky lustre. Going from the N.W. corner of Sect. 24 southward on the section line we find the ferrugino-arenaceous flags and slaty rock-beds again well exposed near the quarter-post. Near the south-west corner of the section dioritic schistose rocks come to the surface. Taking from here an eastern course, we remain on diorite hills until we come to the quarter-post on the south line, where the quartzite formation crops out again in massive, partly conglomeratic ledges; farther on, we pass, before we reach the south-east corner, high vertical cliffs of quartzite, dipping with almost upright beds to the north. Ascending these cliffs and going north, we find them overlaid by a large succession of arenaceous flagstones, which continue northward until we reach the edge of the large marsh surrounding the head-waters of several tributaries of Dead River. A low undulation of the surface in the midst of the marsh, not many steps west of the N.E. corner of Sect. 19, is composed of steeply erected black slates interstratified with sand-rock seams; similar blackish slate and sandstone ledges underlie the drift-covered highlands in the south part of the section and in the west half of Sect. 20. In the N.W. quarter of the S.W. quarter of the latter section, a thick belt of massive conglomeratic quartzite ledges, with a chloritic cement, is exposed on top of a small knob, a short distance south of which the dioritic rocks are at the surface near the south-west corner of the section. In the N.E. part of Sect. 20, along the east section line, is a row of high knobs, with precipitous slopes rising above the marsh-lands, formed of the black slates and of interlaminated sand-rock beds, which dip north-west and are seen on the east side to repose directly on the quartzite formation. The drift-covered hills in the east part of Sect. 17 and in the northern two thirds of Sect. 16 are all formed of the black slates and connected quartzose seams; they are well-denuded in one of the parallel undulating ridges which intersects the central part of Sect. 16 in an east and west direction; their dip is always north, and on their south side follows another

parallel undulation formed of quartzite, in thick, light-colored, compact ledges. The same slaty rock-series, reposing on the south side, on the quartzite formation, composes the hills in the south half of Sect. 15, and in the S.W. quarter of Sect. 14. In the N.E. quarter of Sect. 15 is a row of hills composed of a peculiar variety of crystalline diorite, which extends into the N.W. quarter of Sect. 14, Town. 48, R. 27; all the surrounding country is formed of the slaty and arenaceous rock-beds of the fifth group; their contact with the diorite is not observable. I have, in the previous description of the topographical extent and the lithological character of this northern belt of deposits, representing the fifth group, only attempted to give the general features, and refer the reader to the accompanying geological map for further details, in order to avoid a too frequent repetition of similar statements, which is necessarily involved with the description of a rock-formation, as it occurs in a number of different remote localities. In the third volume of the Reports of the Geological Survey of Michigan I gave a short description of a cursory tour into the slate district of L'Anse and of Huron Bay, on which occasion I fell into an error regarding the dip of the strata and the cleavage of the slates, which I intend to rectify. At the same time I remarked that my investigations were not far enough advanced to enable me to give an opinion of the relative age of the slates, compared with the other sediments of the Huronian rock-series. While I examined the sedimentary series representing the fifth subdivision of the Huronian group in the environs of Negaunee, and particularly during the examination of the more northern exposures in the valley of Dead River, the similarity of these deposits with the slate-formation of L'Anse and Huron Bay made me suggest the analogy of all these compared rock-beds in the different localities, and a subsequent visit of Huron Bay fully convinced me of the correctness of my supposition. As the visit was a short one, and I had made no preparations to extend my explorations to any great distance from the shore, I confined myself to the examination of the beds of several creeks entering the bay, which in their descent from the hill-sides have carved deep ravines into them, and laid open splendid sections across the upheaved strata, amounting to an immense thickness. Clay slates in all shades of color, from whitish-gray, yellow, red, brown, to the darkest black, and in all degrees of hardness, from the fragile, absorbent, earthy condition of a shale, to the hard and compact structure of a roofing slate, are to be seen here, in endless repetition, often for many hundred feet not interrupted by inter-

position of any other kind of rock-seam, other times in frequent alternation with arenaceous layers and with belts of compact granular quartzites. The cleavage of the slates, which in uniformly colored layers cannot be discriminated from a sedimentary lamination, is in other sedimentary belts composed of an alternation of differently colored seams, obviously distinct from it, and intersects these laminæ under a more or less acute angle, partly under a right angle. In the above-mentioned essay I had the uniformly colored roofing slates under consideration, and I stated then that their cleavage was coincident with the sedimentary lamination, which, as I now see, was an error. This discordance between cleavage plains and sedimentary lamination makes it often very difficult to recognize the actual position of such rock-seams with regard to others, and to determine the direction in which the strata dip, which contrariety in the dip of the sedimentary laminæ with the cleavage plains, caused me also in the above case to assert the observation of a southern dip of the beds in the slate quarries, which according to the sedimentary striation of the beds is northward. It requires more time than I have devoted to the examination of this very large succession of slaty and arenaceous sediments to recognize the existing order in the structure of this rock-belt, and to form an estimate of its thickness. The lighter colored variegated slates, which are so well exposed in the bed of Slate Creek and along the shore of L'Anse Bay, appear to be the upper strata; the alternation of banded slaty beds with arenaceous seams, interstratified with larger compact quartzite belts of light color, and partly of dark blackish color, seem to represent the middle series, which in rock character corresponds most with the arenaceous slate-formation in the Negaunee district. These strata are best exposed in the bed of Silver Creek, near the bridge. The roofing slates form the lower horizon of the series; they are intersected by volcanic dykes, and are nearest in position to the granitic rocks, which compose the higher mountains southward, but I have not seen the slates in close contiguity with these rocks.*

* At the time the proof-sheets passed through my hands, I had made further examinations of the horizon of the arenaceous slate group in the Marquette region, and found that large iron-ore deposits, partly recently discovered, partly known for a good while, form a part of this group, occupying a position above the black slates which in these iron districts are of a graphitic nature. Of this class are the ore deposits of the Taylor mine in the L'Anse district, of the Northampton and D'Alaby mines, north of the Champion mine; likewise those of the S. C. Smith mine and of other localities not yet developed. At the same time I will mention that these deposits are contemporaneous and equivalent with the ore deposits of the Commonwealth mine in Wisconsin.

CHAPTER VI.

MICA SCHIST GROUP.

ON the west side of the Michigamee mines, near the railroad, are outcrops of dark blackish gray rock-beds, partly of slate structure, partly in well-laminated, banded, more compact seams, which succession of beds follows immediately above the actinolitic rock-series, dipping in conformity with that southward. These rock-beds consist of a sub-porous ground-mass, formed of very minute granules of white translucent quartz, in intermixture with a large proportion of brightly glistening black mica scales, and not rarely also with chlorite. In the softer, quite fissile schistose or slaty beds, the mica overbalances the granular quartz, and they have a silky lustre. In the compact banded ledges the quartzose ground-mass prevails, and their aspect is dull. Certain seams inclose an abundance of brown garnet crystals, from the size of a mustard-seed to that of a pea. These beds I consider as representatives of the upper horizon of the fifth group; they correspond with the micaceous schists, inclosing garnets exposed near the railroad track on the north side of the Keystone mines, which in that locality are succeeded on the north side by black slates of carbonaceous character, and by ferrugino-argillitic and siliceous beds. At the Michigamee location no exposures analogous to these are observable, as the hill slope on the north side of Michigamee Lake is deeply covered with boulder drift. On the south shore of the lake, opposite Michigamee village, the rock-beds come to the surface again near the water's edge; we find there silvery-shining gray-colored mica schists, some smooth, even-bedded, others much corrugated, which essentially consist of the same minutely granular quartzose ground-mass intermingled with lighter-colored mica scales, which composes the schists on the west side of

the Michigamee mines. This great similarity in the sedimentary material is an evidence of the close connection between this mica schist group and the arenaceous slate group, which proves the immediate succession of the first to the other more reliably than it is done by the southern dip of the strata, conformable with those of the Michigamee mine. These mica schists, interlaminated with large belts of a more compact, minutely granular, sub-porous quartz rock, not so rich in the micaceous constituent, but often mingled with a large proportion of hornblende in curved fibrous actinolite-like crystals of a dark green color, compose all the hills on the south side and west side of the shore of Michigamee Lake, and all the islands in the north part of it, as far as I examined them. The dip of the nearly vertical rock-beds is almost uniformly to the south, but the succession of beds is so immensely large that there must be suggested a frequent doubling up of the strata, which in a belt of several miles in width retain, from one end to the other, almost the same general rock character. We meet with a great many modifications of rock, differing in structure, or in the relative proportions of the constituents, or by admixture of accessory minerals, but the large bulk of all of them is formed of the same granular quartzose ground-mass mingled with mica scales. On Goat Island are very good exposures of this series, comprising there four or five hundred feet of strata. Some of the schists, composed of microscopically fine scales of a dark mica which loses its color by exposure to the weather, have a silvery-metallic lustre, made more conspicuous by the corrugated condition in which the schists frequently are found. Other schists are darker colored, less fissile, breaking crossways with a dull earthy fracture, but within the mass are scattered irregularly larger blades of a white mica, which on those dull fractured surfaces show with great brilliancy. Still others inclose quite abundantly small but very perfect translucent garnet crystals of the color of amethyst or ruby. The silvery variety of schists often contains a large number of good-sized twin crystals of staurolite dispersed through their substance, but usually the crystals are rough, with rounded angles, and always milky opaque. In the same kind of schists are irregularly digitated or cruciform clusters of nacreous margarite crystals associated with the staurolite.

Irregular fissure-seams in the schistose rocks are filled with milky

quartz in association with agglomerated coarse blades of a pale whitish mica, which often appears to be dark green, from chloritic substance filling the interstices between the blades, and imbedded within these mica-masses occur not rarely large square columns of rose-colored andalusite, from one to two inches in diameter, besides well-formed green crystals of apatite, some of which polygonal columns, with squarely truncated ends, are often one inch in diameter. These fissure-seams observable on Goat Island are larger and richer in the named crystalline minerals on another island, a mile farther south, called Silver Island, very probably on account of the silvery lustre of the schists composing it, as no silver has ever been discovered there.

The thick belts of more massive ledges, interstratified with the schists of Goat Island, are, as above remarked, composed of the same porous granular quartzose ground-mass as the schists, but contain less mica. Their color is either uniformly darker or lighter gray, or minutely speckled; a distinct sedimentary striation is observable in most of the beds. Other strata, in which hornblende is an additional component, have the fibrous actinolite-like green-colored crystals thinly disseminated within a white granular quartzose ground-mass, containing little or no mica, or the ground-mass is dark colored from intermingled mica. Elsewhere the bundles of curved, fibrous, actinolitic hornblende crystals compose almost exclusively the rock-mass, with very little of a dark micaceous cement in the interstices; these blackish-green colored beds resemble a coarsely crystalline diorite, but they never occur in belts of any great thickness, and always exhibit some trace of their sedimentary origin. How far southward this rock-series, amounting to a surprising thickness, extends, and what kind of rock-beds come there in contact with this formation, I have not yet examined. We will, in the subsequent description of the Menominee iron region, become acquainted with similar mica schists, believed to be identical with the just-considered group of sediments; they may even be in direct continuity with these, as they occur in the lower part of the Michigamee River valley, about forty miles south of Michigamee Lake; but this interval is not yet sufficiently examined to prove or disprove this suggestion.

CHAPTER VII.

SERPENTINE GROUP.

PRESQUE ISLE was for a long time the only locality in the Marquette district where serpentine and kindred magnesian rocks were known to occur.

During the summer of 1878, while engaged in the examination of the environs of Negaunee, Mr. Ropes, Postmaster of Ishpeming, informed me he had discovered large outcrops of marble and of serpentine some distance west of Deer Lake, in Sects. 29 and 30 of Town. 48, R. 27, and accompanied me to the indicated localities, where I saw his statements fully verified. Subsequently, when I came to extend my explorations over this district, I found the extent of the serpentine group much larger than it was previously supposed. A broad belt of such rocks, commencing with a narrow end in the centre of Sect. 27, Town. 48, R. 27, continues westward through Sects. 28, 29, and 30; thence, taking a south-western direction, it intersects diagonally Sect. 31 and Sect. 36 of the next adjoining Town. R. 28. Farther west I have not examined. The distance between the outcrops of Presque Isle and those in the centre of Sect. 27 is 13 miles, on which intervening space not a trace of serpentine or other magnesian rocks could be discovered.

On inquiring which position the serpentine formation holds in relation to the other six groups, I find it difficult to give a satisfactory answer.

The outcrops of Presque Isle are completely surrounded by the granite formation, which comes close to the serpentine, but the contact between these rocks is not seen in that locality. The large western belt of serpentines is bordered on both sides by dioritic rocks, and is often seen in intimate contact with them.

In two localities the quartzite formation is found to repose on the serpentine, leaning on the two opposite slopes of a large body of serpentine hills, and dipping south on the southern slope, north on the northern. The quartzite on the south side is an isolated strip of the formation, about a mile long in an east and west direction, and in thickness representing about four or five hundred feet of strata, mostly in heavy compact beds of a white or reddish color; their position is almost vertical. South of the quartzites rises another high serpentine ridge close by, so that this outlier of the formation is completely surrounded by the serpentine.

The intimate connection of the serpentines with the dioritic formation, and on the other hand the superposition of the quartzite formation on this rock-belt, might lead to the inference of its intermediate position between these two groups, as a link of the sedimentary series; but if this was the case I could not well conceive how a rock-series of so large a bulk, and so different in composition from the succumbent and incumbent layers, should be in some places sporadically developed, and in next adjoining localities be totally missing, in which the concerned strata, supposed to be above and below, are well represented and exposed. Another hypothesis, considering the serpentine formation to be the product of local metamorphic influences on the dioritic rock-series, is very improbable, as the lithological character of the two contiguous rock-species changes abruptly, which could not be if the differences depended upon a secondary transformation of the same, or at least a similar sort of sediments; we would in this case doubtless observe intermediate stages of the transformation.

The rocks belonging to the serpentine formation occur generally in bulky non-stratified masses, which, if they ever originated from mechanical sedimentary deposits, are by chemical action so completely transformed as to efface all traces of their former detrital structure. They resemble a volcanic eruptive rock, forced to the surface in a soft, plastic condition, and most likely heat was one of the prime agents in their formation, or else transformation, in combination with aqueous vapors, which is suggested by the hydrated composition of the serpentine.

The cliffs at the north-east end of Presque Isle are formed of a black, semi-crystalline massive rock, consisting of a dark greenish-brown amorphous ground-mass of serpentine, semi-transparent in

thin splinters, in intimate intermixture with a crystalline foliated mineral of a brassy metallic lustre, which seems to me identical with the schillerspar of the Harz Mountains. In addition, the rock contains a large proportion of protoxide of iron as a coloring matter, and about 2 per cent of chrome iron in small octahedric crystals, readily attracted by the magnet.

Boiling muriatic acid dissolves about half the weight of the rock; the solution contains a large proportion of iron and magnesia, besides small quantities of alumina and lime. The discolored white crystalline residue of pearly lustre, after its separation from the chrome iron by washing, is on analysis found to be composed of silicate of magnesia and alumina, with some lime; it represents the discolored crystals of the schillerspar in the rock-mass.

Associated with the black serpentine are lighter-colored greenish or reddish or variegated rocks, intersected in all directions by seams of calcspar, or by asbestine or chrysotile bands of silky lustre with the parallel fibres transverse to the walls inclosing the bands. The material of these lighter-colored rocks is frequently identical with the black rock, but in a half-decomposed friable condition, with the protoxide of iron changed into hematitic oxide. In the more compact rocks of this seamy character often a system of capillary fibres pervades the rock in a rectangular position to these larger chrysotile seams, which are comparable to the capillary channels in a block of slowly melting ice; they consist also of chrysotile, tinged sometimes with bright red hematitic pigment. There are also rocks, found in the same association of feldspathic composition, resembling a porphyry by intermixture of larger crystals of feldspar with the ground-mass. The principal bulk of the upper beds, which are much corrugated and folded, or shattered by the upheaval, is formed of dolomites in various modifications. Some beds are compact saccharoidal, white or reddish-colored, but the larger portion of the dolomitic rocks is a mixture of pale green milky serpentine, in concretionary nodular masses, with fine-grained dolomite. Fissures in the rock exhibiting a sliding dislocation of the walls are filled with long-fibred flexible asbestus, or with rigid wood-like masses of it, and druse cavities lined with brilliantly shining rhombic crystals of dolomite spar are quite common. The rock incloses also concretions of iron and copper

pyrites, of zincblende, galena, etc., which have induced some mining experiments, but the quantity of these minerals is too small to be of practical value.

The inconformable superposition of Silurian sand-rock on the serpentine formation of Presque Isle is previously mentioned.

A rock identical with the black serpentine of Presque Isle composes an isolated knob near the centre of Sect. 27, Town. 48, R. 27, which is the first outcrop of the serpentine group observed west of Presque Isle. Next adjoining on the east side is another knob, consisting of a hard feldspathic rock of porphyritic structure by a copious intermixture of feldspar crystals with an almost aphanitic brown-colored ground-mass. Another portion of the rock is completely crystalline, granite-like; it contains small hornblende crystals sparingly intermingled with the feldspar crystals. Adjoining outcrops on the south and west side of the serpentine knob are schistose dioritic strata, which show their affinity to the feldspar porphyry by insensible gradations into it. North of the serpentine knob, on the opposite side of the marshy valley of a creek which intervenes, other outcrops of magnesian rocks occur in intimate association with diorites and dioritic schists. A part of these has a schistose structure, and consists of a dark green amorphous serpentine, intermingled with an abundance of pearly scales of talc, of which mineral, large bunches of beautifully greenish white color occur in the schists in association with the sparry carbonates of lime, magnesia and iron. Another portion of the magnesian rocks is perfectly crystalline, and resembles the diorites of neighboring exposures, but by closer inspection they are found to be composed of pearly shining crystals of diallage, incorporated with an amorphous interstitial mass of dark green serpentine; the rock is partially soluble in warm muriatic acid, which solution gives a strong reaction indicating magnesia and iron, besides small proportions of alumina and lime. The adjoining diorite is not acted upon by the acid, and decomposed by fusion with alkali, it is found to contain only 3 per cent of magnesia and 5 of lime; the other constituents have not been determined by weight. All outcrops further north are formed of dioritic rocks. Westward, and yet within the limits of Sect. 27, we find a short ridge on the south side of Carp River, formed of bulky masses of a dark lead-colored steatite, which contains a good proportion of sparry car-

bonates (of lime, magnesia, and iron), besides an abundance of magnetite granules, and invariably some chrome iron. With the steatite occurs a belt of granular dolomite and sparry seams banded with streaky laminæ of pale green serpentine. South of this steatite ridge, and in close connection with it, are several other parallel undulating ridges, which consist of dioritic rocks of fine grain, cleaving into rhomboidal fragments.

On the north side of Carp River, opposite this place, we find the lower part of the slope of a high range of mountains deeply covered with drift; higher up are cliffs of serpentine and dolomitic rock, and the crest part is again formed of diorite. Farther west, within Sect. 28, the entire body of the continuation of this range is composed of magnesian rocks belonging to the serpentine group, and a part of the same range, extending over the S.W. quarter of Sect. 21, and the S.E. quarter of Sect. 20. In the latter localities the whole mountain crest is formed of bold cliffs with vertical walls. In Sect. 21 is the before-mentioned locality in which the quartzites are seen to repose with a northern dip on the basal slope of this extremely broken body of serpentine hills. East from there, the novaculitic schists underlying the quartzite formation are observable in close proximity to the serpentine, and, further east, the dioritic schists come in contact with it. This large serpentine belt, over a half mile in width, exhibits a remarkable uniformity in its composition, although numberless varieties of rock occur, dependent upon differences in structure, color, or the admixture of accessory minerals to these specifically magnesian masses, which are partly silicates, partly carbonates, or an intermixture of both chemical combinations. The recognition of an order in the arrangement or succession of these rocks is scarcely possible, owing to their greatly disturbed upheaved position, and to their massive non-stratified structure. A large proportion of the rock material composing the presently considered group of hills consists of light grayish green and, in thin splinters, semi-transparent serpentines, which usually have a granular, subcrystalline, glistening fracture, as the crystalline mineral *marmolite* is substituted for the amorphous serpentine, which amorphous form, with a waxy fracture, is however abundant enough, and forms masses of considerable bulk. Some of it is the so-called precious serpentine, of bright yellowish green color and semi-transparent;

its adaptation for ornamental work is much impaired by nodular clusters of magnetite crystals, abundantly disseminated through the mass. The common serpentine has usually a much paler grayish green color, and is likewise full of magnetite granules, contains always some chrome iron, and not rarely, *nikelglanz*.

Other large masses of the rock, blackish green or dark lead-colored from copiously intermingled magnetite granules, and from a dark green-colored interstitial mass of amorphous serpentine, have a crystalline foliated or a stellato-fibrose structure, and are principally composed of crystal blades of *marmolite*, or, in the other case, of *picrolite* or *chrysotile* in intermixture with the above-mentioned cement mass. All these rocks exhibit on polished surfaces a peculiarly cellulose structure, from a network of capillary seams pervading them; the magnetite in the darker rock is particularly concentrated into these seams. In other instances they are formed of *picrolite* fibres. Most all contain a certain proportion of carbonates, which make them effervesce by immersion in warm muriatic acid. The cellulose reticulated structure of the rock, caused by intersecting capillary seams mainly composed of magnetite granules, is particularly plain in rock-specimens collected on the north line of Sect. 28, a quarter mile east of the N.W. corner; at right angles with these finer anastomosing filaments, the rock is banded by other larger seams of *chrysotile* or of *asbestos* in intermixture with *calcspar*, which usually are parallel with each other, or the bands composed of vertical fibres split into smaller seams, which occasionally may reunite and inclose lenticular segments of the rock-mass within the loops. It is difficult to give a plausible explanation of the mode in which such transverse fissures in the rock could form and fill up with these silky *chrysotile* and *asbestos* bands; it could not be done by aqueous infiltration, as there is no indication of a successive deposition of the substance. It appears more as if it were an exudation from the capillary pores of the rock into these fissures, while it was in a plastic semi-fluid condition, which same process formed probably the bands of *epidote* in the diorites, which also consist of fibres transverse to the bands or the fissure walls.

Besides the serpentine and other magnesian silicates, magnesian limestones have an important share in the composition of these hills. Such limestones occur in the S.W. quarter of the S.E.

quarter of Sect. 20 on the south-west slope of the range. A part of the rock on the surface is in a decayed, soft, porous condition, and has a brown ochraceous color; the not decayed, undecomposed rock is whitish, compact, of fine-grained crystalline structure. It contains, besides the carbonates of lime, magnesia, and iron, minute scales of talc, and is densely interwoven with fine seams of serpentine.

Still larger exposures of limestone occur in the N.E. quarter of the N.W. quarter of Sect. 29, on the south side of an isolated ridge surrounded by a swamp. The main mass composing this ridge is a dark-colored serpentine of subcrystalline marmolitic structure, which incloses seams of talc in intermixture with the sparry carbonates of lime and of iron; other sparry seams contain long-fibred asbestos, permeating the spar in distant bundles, or its densely crowded fibres form the prevailing material of the rock-seam, and the spar fills out the narrow interstices between them. The talc of pale green color, partly in granular masses, partly in large-leaved micaceous form, is very curiously mingled with the spar, which forms parallel columnar stems of the thickness of a finger, down to that of a lead-pencil, more or less remote from each other, and the interstices between these columns are solidly filled out with the talc. I have observed this singular columnar structure of sparry rock seams associated with talc in several localities.

The limestone belt forming part of this ridge is best exposed on the south-east side of the slope; it amounts to considerable bulk, and is partly composed of coarsely crystalline sparry masses of white color on fresh fractured surfaces, but rusty brown on exposed weathered surfaces, from oxidation of a certain proportion of carbonate of iron combined with the limy carbonate; other portions of this limerock are gray-colored, fine-grained, with a conchoidal fracture. Both the fine and the coarser-grained limerocks are mingled with linear seams of talc and contain scattered columnar crystal blades of bright lustre which I have not chemically examined. South from there, on the other side of the swampy valley, rises a high mountain mass, cut up into numerous parallel summit ridges, which is formed of the serpentine group. We meet also here with calcareous rock-belts of a more impure mixed character. Some of these rocks, consisting of a reticulated intermixture of dark green amorphous serpentine with white spar-

seams, take a good polish and are very ornamental. The serpentine, which forms the largest portion of the rock-mass of these hills, varies in color from dark blackish green to pale greenish gray; most of it occurs in massive non-stratified belts; sometimes an obscurely laminated schistose structure is observed.

On the east side of this hill-group, near the centre of Sect. 29, are large bluffs in which the various modifications of the magnesian rocks can be studied to advantage. We find there quite large veins of rigid asbestos resembling petrified wood.

By following these bluffs southward we come into the previously mentioned belt of quartzite which borders the south slope of the serpentine range, trending in south-western direction along the north side of the valley of a creek flowing to the Carp River.

The corner of Sect. 29 is on the other side of the creek, and south of the quartzite belt, at the base of another range of serpentine hills, in the north part of Sect. 31.

The high bluffs on the summit part of them are composed of a dark-colored steatite, which incloses seams of white calcspar, with cleavage plains large enough to split off rhombohedric blocks 8 inches in diameter. Associated with the spar occurs a pale green pure talc of pearly lustre, in bunches larger than a man's fist.

Passing over these hills next to the valley-westward, nearly all the before-described kinds of magnesian rocks are met with. The dark green-colored serpentines, intersected by a network of sparry seams, are locally well qualified to be cut and polished for ornamental trinkets, but not for larger-sized works of art. Light-colored serpentine is not so common as in some of the before-examined localities; but dark lead-colored subcrystalline rocks, composed of pearly shining marmolite crystals imbedded in a green amorphous serpentine mass, and richly impregnated with magnetite granules, belong to the commonest in the exposures. Dolomite also occurs in fine-grained compact massive belts, dark blackish-colored by magnetite granules, and chrome iron is discovered in it by the blow-pipe. The serpentine formation composes all the hills on both sides of the creek from our starting point near the S.W. corner of Sect. 29 to the west line of Sect. 31. The quarter-post of the west line is situated on a hill of serpentine, but on the south side of this creek only the first row of hills adjoining the valley is formed of rocks representing the serpentine

group; the large body of hills south of there is all composed of dioritic rocks. On the north side of the creek the serpentine group covers not only all the surface of Sect. 31, which is on that side, but also the southern three quarters of the east half of Sect. 30; and in the S.W. quarter of the S.W. quarter of Sect. 30 are again several serpentine knobs, on one of which the corner of the section is located. North and west of the corner are only dioritic rocks, but south-west of it the serpentine continues diagonally across the N.E. quarter of Sect. 36, and forms the surface of the whole S.W. quarter of that section, entering into Sect. 35, but further the examination was not extended. In the S.W. quarter of Sect. 36 the exposures are very extensive and easily accessible, as the forest is cleared away from the greatest part of the section. A large portion of the outcrop is formed of crystalline marmolitic rocks of finer or coarser grain; they have a dark gray color from intermingled granules of magnetite, which always is associated with small quantities of chrome iron. The bladed crystals of marmolite, imbedded in an amorphous ground-mass of serpentine, giving it a silvery glistening lustre, often occur in the rock-mass also red, copper-colored blades and scales, with the silvery ones. Polished specimens of this kind of rock are very beautiful on closer inspection, but disastrously for their value in the market, this same polished rock, seen from a greater distance, has a dull gray unpleasing color, because the eye cannot distinguish the small scaly molecules which constitute its beauty. Similar aventurine-like rocks of a brown color, with thickly disseminated copper-colored scales, are found in the S.E. quarter of the N.W. quarter of Sect. 29, Town. 48, R. 27, which polish well and look very fine held near enough to the eye, but their aspect from a distance is also not attractive. Occasionally narrow seams of precious semi-transparent serpentine are found disseminated with such scales of metallic lustre, these being polished are, also seen from a distance, very beautiful.

An industrious collector can gather a great variety of such specimens large enough to make paper-weights and other small ornamental trinkets; but if I were asked for advice whether it would be a paying business to quarry these rocks, and send them to the market in sawed slabs and blocks, I would certainly warn one from this enterprise, because I know that not enough of the

finest varieties could be found to make the success certain, and probably not enough to prevent a total failure and disappointment. In the S.W. quarter of Sect. 36 are good exposures of the contact between the serpentine and the dioritic rocks; the serpentine leans with a southern dip on the diorite, both rocks being in an almost vertical position; the diorite, in contiguity with the serpentine, is full of fissures filled with epidote, red feldspar, quartz, and chlorite; but both kinds of rock are well defined, and no transition from one into the other is observable.

CHAPTER VIII.

ERUPTIVE DYKES.

THE previously described series of rocks, representing the Huronian group, considered to be a succession of sedimentary deposits in a more or less altered condition, is frequently found intersected by rock-seams of a different kind, which must have come into the position held by them subsequent to the formation of the stratified sedimentary beds, and even subsequent to their transformation into rocks of crystalline structure and to their great dislocation. There are two essentially different sorts of these transverse rock-belts. One represents, evidently, fissures in the mother rock, replenished with various crystalline minerals, and partially with detrital fragments, by intervention of water as a transporting and solvent medium. These so-called fissure veins have nothing in common with the other kind of rock-seams formed of a homogeneous solid rock-mass, which evidently represents a cooled lava stream forced from the interior through the ruptured superficial crust of the globe; this latter kind only shall be considered in this chapter.

Various lithologically different lava-masses can be distinguished, which represent different periods of volcanic action, as one kind is found to intersect the other whenever they come in contact, and the belt intersected must necessarily be the older of the two.

The oldest of the volcanic dyke-rocks in this district is a truly dioritic rock, consisting of a whitish or greenish anorthic feldspar and of hornblende. As accidental additional constituents, granules of iron pyrites and of magnetite are usually found sparingly intermingled with the mass; less common is the intermixture of brightly yellowish green small clusters of epidote crystals. The color of the rock is lighter or darker greenish gray; in the smaller dykes, which vary in width from less than a foot to 50 or 60 feet, or even more, the rock-mass is usually fine-grained, homogeneous

for the naked eye, quite compact, with a smooth conchoidal fracture; the wider dykes have generally a coarser crystalline grain, in which the component minerals can be recognized with the naked eye, or at least with a small magnifying power. The dyke-mass generally adheres firmly to the inclosing walls, but seems to have exerted little or no altering influence on them. Dykes are frequently seen to divide into numerous side-branches, which often are not wider than a few inches, and may extend in a geniculated course for quite a distance from the main belt. These thin seams are generally of a homogeneous aphanitic structure, owing to the rapid refrigeration of such narrow seams of molten substance.

Another younger kind of dyke rock, much darker than the other, and often intensely black-colored, has the composition of a dolerite; it either forms a homogeneous basalt-like mass of aphanitic grain, or consists of a magma of crystals of dark brown augite, with translucent glassy crystals of colorless or yellowish anorthic feldspar; nearly always a certain proportion of magnetite is mingled with the mass, which is often large enough to impart to the rock strong magnetic properties. These augitic dykes, if coming in contact with the dioritic kind of transverse rock-belts, invariably intersect them, and hereby prove their younger age.

Doleritic dykes frequently intersect the iron formation and higher Huronian rock-beds, while the dioritic dykes seem to be restricted to the lower horizons of the Huronian group, if we are not inclined to consider the larger belts of diorite which protrude in the central part of the synclinal rock-basin, from the midst of the upper Huronian strata, as analogous eruptive masses, coeval with the smaller transverse belts. The often perfect similarity of the dioritic rocks of the larger belts, parallel to the trend of the formations with these transverse, decidedly volcanic dyke-masses, proves, at all events, the transformation of the former sedimentary rock-ledges into a crystalline diorite, by the action of heat principally, or else the structure of these diorites could not be so absolutely similar with the dyke-rocks, whose igneous origin is unquestionable. Both kinds of dykes, the dioritic and the doleritic, are abundantly represented in the Marquette district. We can observe a dioritic dyke well exposed in the quarries at the Lighthouse Point of Marquette, while it intersects a large belt of nearly vertical well-stratified ledges of dioritic schists under an angle of 45° . This


fine-grained crystalline transverse seam, about 12 feet in diameter, differs in its lithological character so little from the substance of the compact schistose ledges intersected by it, that it requires a careful observer to make a distinction in hand-specimens. In the same locality another larger dyke of doleritic composition is to be seen, which is parallel with the schists in its general course, but intersects the dioritic dyke. This doleritic rock-belt, of a coarsely crystalline structure and about 50 feet wide, extends into the Lighthouse Point, composing the knob on which the turret for the light is built. East of the building it divides into several branches, which inclose within their anastomosing loops, schistose masses, whose broken ends correspond with the ends of other segments of schist, inclosed within another adjoining loop in front of that one, and obviously the grain of these smaller branches of the dyke is finer, more homogeneous than that of the thick main belt, in which the crystals of augite and feldspar are plainly discernible. A third sort of dykes is disclosed in the Lighthouse quarries, smaller than the others, not over two feet in width, and sometimes only a few inches. It intersects the strata in an irregular geniculated course, and disappears for a while under the accumulations of lake-sand in this place, but reappears again in the sub-schistose dioritic bluffs on the road near the Marquette saw-mill. The dark olive green, perfectly amorphous, smooth-fracturing dyke-mass differs from any other I have observed in this district, but may perhaps be only a modification of the dioritic dykes; it does not intersect the other dykes in the place. The large augitic dyke of the Lighthouse Point appears to continue for three miles westward in one uninterrupted belt, parallel with the stratified rock-series. We meet with outcrops of the same rock in Michigan Street, east of Front Street, and again in the quarries at the west end of Arch Street, in which latter locality the broad main belt gives off several smaller side-arms which intersect the strata diagonally. Here it can be plainly observed how the more or less rapid cooling of the lava prevented the crystallization completely or partially. The narrowest side-seams and the outside of the larger belt in contact with the wall-rock, consist of a black, homogeneous, fine-grained mass, which toward the centre of the large belt gradually becomes a crystalline structure, and forms in the centre of the dyke a magma of well-formed crystals of white

glassy feldspar with augite. Other narrower augitic dykes, intersecting a bulky crystalline diorite mass, are observable two blocks farther north in Vermont Street. West of the Marquette cemetery, not far from the south quarter-post on the north line of Sect. 22, the supposed continuation of this large doleritic dyke is found again largely denuded, and exhibits the homogeneous aphanitic condition of the rock, as well as the coarsely crystalline form, of a black and white speckled appearance, and of bright lustre on fractured faces. This dyke intersects in this vicinity, on Hudson's Hill, another dyke of dioritic composition, rather fine-grained, and copiously mingled with iron pyrites; fissures in the mass are filled with epidote and quartz crystals.

Pursuing the doleritic dyke farther west, we find some low hillocks in the north-east corner of Sect. 21 composed of it, and a row of knobs already described in a previous chapter, trending along with the south line of Sect. 16 to its west end, is formed of the same augitic rock, which differs only from the eastern exposures, and the otherwise fully corresponding mass, by the admixture of a good proportion of flesh-red feldspar crystals. Farther west this rock-belt disappears under drift-accumulations.

It seems to me the most expedient way to describe the different kinds of eruptive dykes promiscuously as they occur in certain localities. I therefore lead the reader from the south line of Sect. 16 to a before-examined locality in the south-east quarter of this section, in which we have seen the repeated alternation of granite belts with the schists of the dioritic group. Here we see the schists and granites transversely intersected by various dioritic and doleritic dykes. One of the dioritic dykes in its main belt, about 25 to 30 feet wide, consists of a uniformly green-colored magma of hornblende crystals with feldspar, in which larger white feldspar crystals have segregated, which give the rock a spotted appearance; narrower side-branches of this same dyke are a non-crystalline, homogeneous, compact mass of a pale green color. Iron pyrites is the hardly ever missing accessory constituent of all these dyke-rocks. Doleritic dykes of black color, in belts a few feet in width, are likewise seen here to intersect the granite and the schists. Another narrow dyke, only a few feet wide, and dividing into numerous narrower tortuous seams, is seen to intersect the dioritic schists of this locality; it consists of a reddish

gray fine-grained ground-mass, principally formed of red feldspar, in which small dark green-colored blades of hornblende are distributed in a gneissoid laminated arrangement, which structure is, however, not in all parts of the dyke equally plain. A dyke of similar nature is found in the exposures at Stone's mills in the S.W. quarter of Sect. 7, Town. 48, R. 25, where it intersects the granite and associated schistose beds. Most likely also a dyke belongs to the same class, which intersects the granite in the N.W. quarter of the N.W. quarter of Sect. 17; though it contains a larger proportion of hornblende, and is of a coarser crystalline grain. In this association belongs probably also a dyke four or five feet wide, intersecting the dioritic rock-mass of the Picnic Islands, which I have previously mentioned in order to give an example of a laminated gneissoid structure, exhibited by an evidently eruptive rock-belt. Before leaving the S.E. quarter of Sect. 16, which is the same locality that offered us the natural section through the lower part of the dioritic series described in the second chapter, I must give a description of a most instructive exposure in the bed of the creek which follows the east line of Sec. 16. An augitic dyke, dividing in several branches and



loops, is seen here to intersect a large series of dioritic schist, interlaminated with several belts of granite, in the high vertical bluffs, which present a section through these ledges in steeply erected position. One arm of the dyke, about three feet wide, coming to the surface in the bed of the creek, cuts diagonally across the granite and schist ledges exposed in the rock-wall, causing a lateral dislocation of the disrupted corresponding ends of the beds, amounting to about eight inches. Another arm of the same dyke forced its way parallel with the bedding between two granite seams, and divides afterward into reuniting branches, which inclose within these loops fragmental portions of the granite belt. The above sketch shows the exposure.

North of here, in the granite hills intermediate between the valleys of Dead River and of Campo Creek, dykes are very abundant. I have mentioned the occurrence of a row of diorite hills, parallel to the trend of the formation, and on both sides surrounded by granitic rock, whose eruptive nature and relationship with the smaller transverse dioritic rock-seams is suggested on that occasion. Of the smaller dykes a great many occur in close vicinity to these larger eruptive masses; they exhibit a good many minor variations in external aspect of the rock-mass in each dyke, but these depend only from a more or less perfect crystallization, and all of them can with propriety be arranged under the two classes, the dioritic and the doleritic. In the south half of Sects. 5 and 6, several dioritic dykes, from 30 to 50 feet in diameter, exhibiting all gradations of structure from the aphanitic to the coarsely crystalline, can be observed in the granite, and in a few places their intersection by doleritic dykes is seen. A large doleritic dyke in these localities, which sends out a number of smaller side-branches, exhibits the contrast of the quickly cooled aphanitic masses, with those slowly refrigerated, well-crystallized, and all the intermediate gradations, particularly well; the narrowest seams and the wall portions of the larger are a homogeneous black mass of conchoidal fracture; in other portions of the dyke the aphanitic black material is seen disseminated with scattered white, translucent feldspar crystals of good size, and the central part of the largest seams is formed of a magma of augite and feldspar crystals, without a trace of an amorphous interstitial ground-mass. Some of these augitic dyke-masses are strongly magnetic; others, of exactly the same appearance, exert no influence on the compass needle. Partridge Island and the Gull Islands, east of Presque Isle, are other favorable localities for the observation of dykes of both kinds; particularly the Gull Islands, perfectly denuded of soil, exhibit the intersection of the dioritic by the doleritic dykes most plainly. We notice there also the basalt-like, columnar cleavage of the dykes of doleritic nature, which is transverse, from wall to wall, of the fissures filled with the amorphous lava-mass.

Leaving the granitic districts, and examining the area of the dioritic rocks, the dykes occur not near as frequent, and there the doleritic dykes are much oftener seen than the dioritic, which may, however, in part be due to an oversight of the dioritic seams, which

look so perfectly similar to the indigenous masses inclosing them, while the doleritic are at once discernible from them.

Still less common are dykes in the districts covered by Huronian rocks, younger than the dioritic series, and the doleritic kind is the only one observable. The intersection of the iron-formation by doleritic rock-seams on the north side of Negaunee, in the hematite mines on the Jackson mining property, at the Cascade mines in several of the different locations, and in the eastern pits of the Washington mine, has incidentally been mentioned in the previous pages. The intersection of the novaculitic rock-series beneath the quartzite formation by a doleritic dyke, is observable in the N.E. quarter of the S.W. quarter of Sect. 21, Town. 48, R. 27; this is the only locality in which I observed an appreciable altering influence of a dyke-mass on the inclosing walls, but it does not extend much over an inch sideways into the wall-rock.

The intersection of the black slates, representing the fifth group by a large doleritic dyke at the falls of Dead River, is previously mentioned, and in the surrounding country several other doleritic rock-seams can be observed cutting across the slate-rock.

The only instance of the intersection of the serpentine group by a doleritic dyke, occurred to me in the N.W. quarter of Sect. 31, Town. 48, R. 27. In the south part of the examined district, dioritic and doleritic dykes are equally common, as in the north part, within the area of granitic rock-exposures, but on the large space covered by the quartzite formation, and by the arenaceous slate-group, no dykes were observed, if we except the large insular protrusions of dioritic rocks which occur there.

A narrow branching dyke intersects the roofing slates in the quarries of Huron Bay, which in its rock character is unlike any of the before-described dyke-masses; it consists of a compact amorphous light-gray colored ground-mass, sparingly disseminated with small pale green columnar hornblende crystals.

In connection with the volcanic dykes, I have to describe other narrow rock-seams of a partially schistose structure, which transversally intersect the granitic and dioritic rocks in the islands and on the shore of the main-land west of Presque Isle, and in several other localities more remote from there. The partially slaty structure of these transverse rock-seams induced me at first to consider them as sedimentary strata which became accidentally inclosed within a fold of the upheaved granitic masses, which case evidently happened sometimes. A slaty novaculitic rock-seam intersecting the quartzite formation near the east end of Teal

Lake is mentioned by Major Brooks under the name of a slate dyke, which belongs to this class; but the above-mentioned transverse rock-belts have a different origin; there is no rock-formation similar to them developed in these localities which could have been inclosed, except those narrow seams which in several instances are seen to intersect other dykes, of the dioritic and the doleritic kind, while it had to be expected that these lava-belts, erupted after the general large upheavals, should intersect them, if they are sedimentary layers wedged into a fold of the rock-masses. It is also impossible to suppose sedimentary deposits, holding a position as they do, could form subsequent to the eruption of the doleritic lava-streams. In fact, the laminated structure of the schists is totally unlike a sedimentary lamination, and is the result of sliding pressure on a semi-plastic mass; moreover, the larger proportion of the substance of these rock-belts has a massive sub-crystalline structure closely similar to igneous eruptive rocks. The character of the rock-mass in dykes of this kind is not altogether alike; a number of them occur in close vicinity in the N.E. part of Town. 48, R. 25, and almost every one of them varies some from the others. Some are richly impregnated with granular magnetite, and exert a powerful influence on the magnetic needle; others of the dykes are not, or only faintly, magnetic.

On a naked granite island, situated near the imaginary north line of the N.W. quarter of Sect. 2, Town. 48, R. 25, a narrow dyke about five feet wide intersects the granite in a north and south direction. The exterior portions of the dyke-rock, intimately coherent with the granitic side-walls, have a laminated schistose structure; the central portions are massive, not laminated. The schistose parts consist of a minutely granular pale greenish-colored feldspathic ground-mass, laminated by interrupted sub-parallel linear seams of dark green-colored fibroscaly blades arranged in a similar mode as the micaceous laminae of a gneiss. The lamination is in the vertical sense parallel with the walls of the fissure, but oblique to the walls from one side to the other; the schistose parts of the dyke, as well as the central massive portion of it, contain a large proportion of carbonate of lime, causing a lively effervescence if the rock is immersed in muriatic acid; usually also a certain quantity of magnetite and iron pyrites granules is found intermingled with the schistose part of the rock-belt, but in the central massive part of the dyke magnetite is very abundant, giving the rock a dark lead color, and imparting to it a powerful influence on the magnetic needle. This massive central part of the dyke is sub-crystal-

line, but earthy, absorbent, or porous ; it consists of the same granular feldspathic ground-mass as the outer, and is copiously intermingled with small flattened, lenticular bits of a green-colored fatty, talcose mineral, which becomes more visible if the surface of the rock is moistened ; pearly mica scales are an additional component, and some harder portions of the dyke-mass also contain radiated clusters of actinolite of pale green color. A similar narrow transverse rock-seam, richly impregnated with magnetite, occurs on the shore of the main-land near the quarter-post on the north line of Sect. 4 ; the rock-mass of this is harder, and actinolite is one of its principal constituents. Other magnetic and schistose dykes are found in the N.E. quarter of the S.E. quarter of Sect. 4, and one near the quarter-post on the south line of the same section, which latter intersects a dioritic dyke. Perfectly similar dykes can be observed in the N.W. quarter of the N.W. quarter of Sect. 9, in the S.W. quarter of the N.E. quarter of Sect. 10, at the N.W. corner of Sect. 18, the N.E. quarter of the S.W. quarter of Sect. 20 ; and in Michigan and Arch Streets of Marquette a similar dyke, not much over one foot wide, occurs. Also in Town. 48, R. 26, a dyke of this kind intersects the granite in the S.W. quarter of the N.W. quarter of Sect. 23, and in Town. 48, R. 27, in the S.W. quarter of the N.W. quarter of Sect. 19.

On the west side of Partridge Island is another dyke of this group, which consists principally of dark green silky-shining schists, which insensibly merge into rigid masses of asbestos ; the schists and the asbestos are in an extremely contorted, corrugated condition, and intermingled with siliceous seams and with a large proportion of sparry carbonate of lime as filling material of the interstices. With these schists occurs a seam of a dark basalt-like rock of aphanitic structure, and a narrow band of a compact rock richly impregnated with magnetite granules, and with ferruginous calcspar crystals imbedded within a micaceous-feldspathic and partly actinolitic ground-mass. This dyke, about six feet wide, intersects a dark black basalt-like dolerite dyke of aphanitic grain, not far off from the water-line of the shore. A larger dyke identical with this one is on the north side of the island, which contains, besides the asbestine and actinolitic masses, thick seams of saccharoidal calcspar interwoven with asbestine fibres, and with well-formed large epidote crystals ; these rocks have some resemblance with certain seams of the serpentine formation, but contain no serpentine and no chrome-iron, which is a constant associate with the latter group.

CHAPTER IX.

FISSURE VEINS.

ESSENTIALLY distinct from the described narrow transverse rock-seams, are these clefts in the rock-masses, subsequently replenished with crystalline mineral substances deposited there successively by percolating aqueous solutions. We find them in all parts of the Huronian series, and in the granites, retaining in general the same character, whether they occur in the higher or lower horizons. The commonest are veins of white milky quartz, which sometimes fills the fissures solidly without the intermixture of any other mineral. In other cases the quartz is associated with calcspar, or with a mixture of calcspar and sparry carbonate of iron. Not rarely red feldspar crystals and epidote crystals intermingle with the quartz and calcspar ; the quartz, epidote, and calcspar compose together fibrous bands with fibres transverse to the walls of the fissures, but this occurs only in narrower seams, not wider than a few inches. Frequently nodular concretions of iron and copper pyrites are intermingled with the quartz veins, and chlorite is almost constantly present as the last-formed deposit filling the interstices left between the other minerals. A more rare mineral is tourmaline in the quartz veins ; it occurs in a black, fibrous, radiated masses. Quite common is large-leaved micaceous iron oxide in such fissures, and in a few instances botryoidal incrustations of the fissure walls with banded chalcedony occur. In several localities larger metalliferous quartz-veins have been discovered, which, besides the before-enumerated minerals, contain argentiferous galena associated with zincblende and copper pyrites. The Holyoke mines were opened in such veins, which, after expenditure of large sums of money, had to be abandoned again, as not a sufficient quantity of the valuable

ores could be found. A similar costly experiment, which likewise proved to be a failure, was made in the S.W. quarter of Sect. 27, Town. 48, R. 26, at the so-called Sedgwick mine, where a quartz vein carries a small quantity of copper pyrites. Considerable test-pitting has also been done in examination of a quartz-vein containing carbonate of iron and some copper pyrites in Sect. 25 of the same township, a short distance north of the Morgan furnace, but with no better success. A good many other localities could be enumerated in which explorers spent weeks and months of hard labor to follow such quartz veins, in which they discovered a few nodules of copper ore or galena. The most promising veins of argentiferous galena have been discovered in the granite hills near the head-waters of the Chocolate River. I have seen there, some years ago, many tons of pure lead-ore piled up on the side of the test-pits, but it seems as if these also are abandoned at present. It can safely be asserted, from precedent experiences, that the Marquette district is, with exception of its immense wealth in iron-ore, rather barren of metallic ores of another kind.

PART II.

MENOMENEE IRON REGION.