

Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
0101	212	92.9	0.0	2.4	0.0	0.0	4.7	0.0
0102	606	91.4	1.2	4.3	0.0	0.0	3.1	0.0
0103	617	94.2	0.0	2.3	0.0	0.0	3.6	0.0
0104	210	95.2	0.0	1.0	0.0	0.0	3.8	0.0
0250	266	82.3	0.0	0.4	1.5	0.0	15.8	0.0
0251	220	87.7	0.0	0.5	0.0	0.5	11.4	0.0
0252	256	44.1	1.2	3.5	3.1	0.4	47.7	0.0
0253	272	66.2	0.0	1.5	4.4	0.0	27.9	0.0
0401	875	92.5	0.6	2.1	0.0	0.0	4.9	0.0
0402	729	82.9	0.1	2.5	0.0	0.0	14.5	0.0
0403	235	97.4	0.0	0.4	0.4	0.0	1.7	0.0
0404	267	91.0	0.0	1.9	0.0	0.0	7.1	0.0
0501	870	94.1	0.5	1.1	0.1	0.0	4.1	0.0
0502	6	*	*	*	*	*	*	*
0503	267	91.8	0.0	3.7	0.0	0.0	4.1	0.4
0504	986	88.2	0.2	1.9	2.6	0.7	6.3	0.0
0505	636	86.2	2.0	3.0	0.6	0.2	8.0	0.0
0506	345	92.5	0.0	4.1	0.0	0.0	3.2	0.3
0650	1,679	87.1	2.7	3.7	0.5	0.5	5.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
0651	1,055	88.5	0.3	4.1	0.2	0.5	6.4	0.0
0652	594	93.8	0.0	1.9	0.0	0.0	4.4	0.0
0653	783	98.0	0.3	1.0	0.0	0.0	0.8	0.0
0654	604	92.2	0.8	2.2	0.5	0.0	4.3	0.0
0655	623	94.1	0.0	1.3	0.2	0.5	4.0	0.0
0657	148	93.9	0.0	1.4	0.0	0.0	4.7	0.0
0701	592	93.2	0.7	0.7	0.5	3.0	1.9	0.0
0702	162	87.7	0.0	0.0	0.0	0.0	12.3	0.0
0703	378	89.7	0.5	1.9	0.5	0.0	7.4	0.0
0704	155	92.9	0.6	3.9	0.0	0.0	2.6	0.0
0705	312	95.2	0.0	1.6	0.3	0.0	2.9	0.0
0850	1,837	82.5	3.9	7.1	0.0	0.5	5.9	0.1
0851	1,525	91.9	0.8	4.3	0.1	0.2	2.8	0.1
0901	2,784	73.9	2.0	14.0	0.0	0.2	9.8	0.1
0902	125	96.0	0.0	2.4	0.0	0.0	1.6	0.0
1051	2,182	87.5	0.7	5.8	0.3	0.3	5.3	0.0
1101	2,912	30.0	58.0	4.9	0.1	0.6	6.3	0.1
1102	1,435	68.3	11.3	6.3	0.2	0.4	13.4	0.0
1106	178	85.4	1.7	6.7	0.0	0.0	6.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
1250	1,833	80.6	0.5	14.8	0.0	1.3	2.8	0.1
1251	1,461	93.9	0.2	2.7	0.0	0.0	3.1	0.0
1252	894	52.8	0.7	43.3	0.0	0.1	3.1	0.0
1253	1,724	76.9	4.5	13.2	0.0	0.2	5.1	0.1
1350	3,859	54.9	19.0	11.5	0.1	1.9	12.5	0.1
1351	313	88.5	0.0	4.5	0.0	0.6	6.4	0.0
1352	731	56.6	25.3	5.9	0.0	0.3	11.8	0.1
1401	409	87.3	0.7	6.4	0.2	0.0	5.4	0.0
1402	293	45.7	0.3	34.5	7.8	0.7	10.9	0.0
1403	63	*	*	*	*	*	*	*
1404	39	*	*	*	*	*	*	*
1501	1,994	38.1	6.5	49.1	0.2	0.0	6.0	0.2
1503	2,230	51.1	5.4	35.3	0.3	0.5	7.0	0.4
1504	4,199	40.4	3.7	48.1	0.0	3.7	4.0	0.0
1505	1,608	85.0	1.8	8.6	0.0	0.2	4.2	0.1
1506	1,067	75.5	1.7	17.1	0.0	0.0	5.7	0.0
1507	1,136	87.5	0.4	5.3	0.0	0.8	6.0	0.0
1508	254	0.4	0.0	98.4	0.0	0.0	0.0	1.2
1509	1,265	47.5	1.6	48.7	0.1	0.1	1.8	0.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
1510	161	78.9	6.2	12.4	0.0	0.0	2.5	0.0
1511	1,135	88.5	1.3	5.8	0.1	0.1	4.0	0.2
1512	538	90.1	0.6	5.6	0.0	0.0	3.3	0.4
1701	1,012	46.5	0.7	3.0	22.4	1.5	25.9	0.0
1712	174	48.3	0.0	1.7	24.7	0.6	24.7	0.0
1950	1,114	90.7	0.1	6.6	0.1	0.3	2.2	0.0
1952	271	92.3	0.0	4.1	0.7	0.0	3.0	0.0
1953	536	90.3	0.2	5.8	0.0	0.2	3.5	0.0
1954	1,075	81.8	0.1	15.9	0.0	0.0	2.2	0.0
1955	285	80.0	0.0	16.1	0.0	0.0	3.9	0.0
1957	738	84.6	1.2	9.6	0.3	0.4	3.9	0.0
1959	303	77.6	5.0	8.9	0.0	0.3	8.3	0.0
2150	1,116	87.8	0.1	2.7	1.1	0.1	8.2	0.0
2151	422	92.2	0.0	3.6	0.0	0.0	4.3	0.0
2251	736	89.9	0.1	3.3	0.0	0.4	6.0	0.3
2252	362	88.4	0.0	4.1	0.0	0.0	7.5	0.0
2551	5,676	26.2	66.9	2.3	0.0	0.1	4.6	0.0
2554	222	89.2	6.3	2.7	0.0	0.0	1.8	0.0
2555	6,465	70.9	20.4	4.1	0.1	0.4	4.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
3001	1,690	9.1	66.5	20.3	0.1	0.5	3.6	0.0
3002	465	13.5	51.8	30.5	0.0	0.0	4.1	0.0
3003	21	*	*	*	*	*	*	*
3004	356	0.3	0.0	99.7	0.0	0.0	0.0	0.0
3007	75	*	*	*	*	*	*	*
3009	23	*	*	*	*	*	*	*
3010	5	*	*	*	*	*	*	*
3201	996	91.3	0.4	5.6	0.2	0.2	2.3	0.0
3301	5,977	33.7	28.7	22.8	0.1	3.0	11.6	0.0
3302	576	14.4	43.8	25.7	0.0	4.7	11.5	0.0
3304	237	32.1	23.2	8.9	0.0	30.8	5.1	0.0
3305	210	63.8	7.6	16.7	0.0	3.3	8.6	0.0
3306	410	85.4	0.7	6.1	0.0	0.2	7.3	0.2
3308	108	85.2	0.0	7.4	0.0	0.0	7.4	0.0
3309	84	*	*	*	*	*	*	*
3550	148	2.0	0.0	2.0	8.1	0.0	87.8	0.0
3551	49	*	*	*	*	*	*	*
3552	77	*	*	*	*	*	*	*
3701	1,833	83.4	0.3	12.5	0.1	0.1	3.7	0.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
3805	5,315	66.1	11.8	9.3	0.1	0.4	12.2	0.0
3950	3,146	54.8	18.0	11.1	0.0	1.3	14.7	0.0
3952	61	*	*	*	*	*	*	*
3953	69	*	*	*	*	*	*	*
3954	192	74.0	2.1	8.9	0.0	5.2	9.9	0.0
4001	1,818	18.1	58.0	11.3	0.2	0.8	11.6	0.0
4002	1,675	38.3	28.2	19.0	0.1	1.0	13.5	0.1
4103	211	6.2	71.6	10.4	0.0	1.9	10.0	0.0
4106	1,890	6.2	45.8	42.5	0.1	0.1	5.3	0.1
4107	4,987	41.0	26.6	19.4	0.1	1.1	11.9	0.0
4109	2,234	82.1	1.4	9.4	0.0	0.2	6.8	0.0
4110	2,976	24.4	8.4	58.3	0.0	0.3	8.6	0.0
4112	1,183	0.2	0.2	99.7	0.0	0.0	0.0	0.0
4113	1,180	3.0	13.3	80.5	0.0	0.4	2.8	0.0
4120	3,630	46.7	7.8	34.4	0.1	2.1	8.9	0.0
4160	3,084	34.0	33.6	16.1	0.1	6.0	10.1	0.0
4351	1,047	88.2	0.8	6.7	0.2	0.0	4.1	0.1
4352	354	63.8	13.3	3.4	0.0	0.0	19.5	0.0
4353	869	67.5	0.3	27.6	0.0	0.1	4.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
4354	1,289	86.3	3.3	2.4	0.0	0.1	7.9	0.0
4355	574	74.6	0.0	10.6	0.2	0.2	14.5	0.0
4356	949	75.0	0.1	15.5	0.1	0.4	8.9	0.0
4357	9	*	*	*	*	*	*	*
4358	413	1.9	0.0	97.6	0.0	0.0	0.2	0.2
4359	964	63.0	0.0	34.8	0.1	0.0	2.2	0.0
4360	281	84.7	0.0	11.4	0.0	0.0	3.9	0.0
4361	369	91.6	1.4	1.6	0.0	0.0	5.4	0.0
4362	570	90.9	0.2	3.3	0.0	0.2	5.4	0.0
4363	1,469	88.5	0.3	5.0	0.0	0.3	5.9	0.0
4401	2,234	83.6	0.8	7.4	0.0	0.1	8.1	0.0
4501	1,607	90.7	0.4	6.4	0.1	0.1	2.2	0.0
4601	2,369	67.6	0.9	24.7	0.1	0.1	6.5	0.0
4701	1,910	90.6	0.7	4.9	0.2	1.1	2.5	0.0
5050	2,133	62.0	24.3	2.7	0.1	1.1	9.8	0.0
5051	3,899	55.8	25.6	5.8	0.1	2.2	10.5	0.0
5052	3,178	59.8	19.3	3.2	0.1	7.6	10.1	0.0
5101	3,256	65.7	8.4	14.8	0.0	1.2	9.8	0.0
5202	933	87.4	0.2	2.1	0.4	0.1	9.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
5203	327	80.4	1.5	5.8	0.0	0.3	11.9	0.0
5204	249	83.5	0.4	2.8	0.8	0.0	12.4	0.0
5601	1,874	92.4	1.0	1.7	0.0	0.1	4.7	0.1
5801	3,264	79.9	4.3	7.7	0.0	0.3	7.4	0.3
5803	65	*	*	*	*	*	*	*
5804	81	*	*	*	*	*	*	*
6101	136	89.0	2.2	3.7	0.0	0.0	5.1	0.0
6102	447	84.3	2.2	4.0	0.0	0.4	8.9	0.0
6103	6,247	51.1	25.1	13.2	0.2	0.3	10.2	0.0
6104	275	10.5	73.8	5.1	0.0	0.4	10.2	0.0
6105	145	85.5	2.8	11.7	0.0	0.0	0.0	0.0
6106	38	*	*	*	*	*	*	*
6301	6,911	40.2	25.7	26.1	0.1	2.3	5.6	0.0
6312	3,794	41.5	47.7	2.8	0.2	1.7	6.2	0.0
6316	2,033	67.9	12.2	11.8	0.1	2.1	5.9	0.0
6320	472	65.7	24.8	1.9	0.0	3.0	4.7	0.0
6325	257	89.5	3.1	4.3	0.0	0.8	2.3	0.0
6330	655	20.3	50.2	20.6	0.3	0.5	8.1	0.0
6335	980	76.9	11.0	2.4	0.0	4.2	5.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
6340	271	33.6	18.1	39.1	0.0	4.4	4.8	0.0
6345	69	*	*	*	*	*	*	*
6350	440	58.2	33.6	0.7	0.0	1.4	6.1	0.0
7304	423	85.6	0.0	11.1	0.0	0.5	2.6	0.2
7306	4,092	39.8	33.8	19.7	0.0	0.3	6.3	0.0
7308	352	87.8	1.7	6.0	0.0	0.0	4.5	0.0
7310	69	*	*	*	*	*	*	*
7401	3,430	73.0	5.0	8.7	0.1	0.4	12.7	0.0
7402	329	90.6	0.0	2.7	0.0	0.0	6.7	0.0
7403	113	80.5	0.0	16.8	0.0	0.0	2.7	0.0
7404	105	88.6	0.0	4.8	0.0	0.0	6.7	0.0
7601	1,186	90.0	0.3	7.1	0.1	0.1	2.5	0.0
7850	1,557	87.2	0.2	6.4	0.1	0.0	6.2	0.0
7851	262	90.8	0.8	6.1	0.0	0.0	2.3	0.0
7852	178	93.3	0.0	0.6	0.0	0.0	6.2	0.0
8150	151	45.0	31.1	11.9	0.0	3.3	8.6	0.0
8151	4,188	34.6	40.7	12.2	0.5	1.7	10.2	0.0
8152	521	26.5	47.8	6.9	0.0	0.0	18.8	0.0
8153	65	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
8154	488	36.1	21.3	16.2	0.2	22.7	3.5	0.0
8155	142	52.1	30.3	6.3	0.0	3.5	7.7	0.0
8156	69	*	*	*	*	*	*	*
8157	48	*	*	*	*	*	*	*
8201	2,901	53.5	23.7	8.5	0.0	2.7	11.5	0.0
8202	4,108	88.5	5.1	3.8	0.2	0.5	1.9	0.0
8203	461	68.8	22.8	2.8	0.2	0.7	4.8	0.0
8250	813	41.9	45.0	4.8	0.4	1.6	6.3	0.0
8255	3,432	60.3	24.1	7.5	0.0	0.9	7.2	0.0
8257	909	47.9	23.2	0.6	0.0	25.0	3.4	0.0
8258	2,348	58.3	26.1	7.4	0.0	0.8	7.4	0.0
8260	716	45.3	29.9	2.9	0.0	18.9	3.1	0.0
8901	1,606	0.2	95.8	0.9	0.1	0.6	2.4	0.0
8902	1,016	2.4	92.7	0.9	0.0	0.5	3.5	0.0
8903	1,873	1.9	95.1	0.7	0.1	0.6	1.6	0.0
8904	1,496	4.8	78.0	0.5	0.3	12.2	4.2	0.0
8905	4,801	36.4	24.6	28.9	0.1	5.7	4.3	0.0
9004	15	*	*	*	*	*	*	*
9012	24	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
9013	84	*	*	*	*	*	*	*
9015	20	*	*	*	*	*	*	*
9016	92	*	*	*	*	*	*	*
9017	33	*	*	*	*	*	*	*
9018	82	*	*	*	*	*	*	*
9019	23	*	*	*	*	*	*	*
9021	26	*	*	*	*	*	*	*
9040	22	*	*	*	*	*	*	*
9041	942	0.2	97.6	0.5	0.0	0.1	1.6	0.0
9045	2,187	6.4	41.3	50.3	0.0	0.4	1.5	0.0
9050	3,082	14.9	49.3	34.3	0.1	0.7	0.8	0.0
9051	3,380	5.1	9.6	84.4	0.0	0.0	0.9	0.0
9053	3,519	1.1	96.1	0.9	0.0	0.7	1.2	0.0
9055	233	1.3	96.6	1.3	0.0	0.4	0.4	0.0
9058	1,165	76.7	15.8	0.4	0.0	6.3	0.9	0.0
9066	3,207	0.6	98.6	0.1	0.1	0.2	0.4	0.0
9068	2,056	2.1	96.1	0.4	0.0	0.4	1.0	0.0
9069	2,876	1.0	95.8	1.1	0.0	0.1	2.0	0.0
9070	2,674	1.4	86.9	0.3	0.1	9.1	2.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 4A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
9072	2,265	1.1	93.6	0.4	0.0	2.7	2.3	0.0
9074	2,177	1.7	93.7	0.6	0.0	0.3	3.7	0.0
9080	3,888	29.0	65.2	2.6	0.0	0.2	2.9	0.0
9151	4	*	*	*	*	*	*	*
Unknown Clinics	134	31.3	10.4	7.5	12.7	9.7	28.4	0.0
Michigan	260,806	51.9	25.5	14.4	0.3	1.4	6.5	0.0
Nation (Prior Year)	8,164,612	31.7	18.8	41.0	1.0	2.6	2.1	2.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0101	212	34.9	20.8	44.3
0102	606	27.9	26.1	46.0
0103	617	31.9	20.4	47.6
0104	210	31.9	16.2	51.9
0250	266	23.3	29.3	47.4
0251	220	29.1	21.8	49.1
0252	256	18.8	24.2	57.0
0253	272	29.0	22.8	48.2
0401	875	27.3	25.9	46.7
0402	729	24.0	26.6	49.4
0403	235	22.6	26.0	51.5
0404	267	23.2	23.2	53.6
0501	870	26.8	24.1	49.1
0502	6	*	*	*
0503	267	34.1	19.9	46.1
0504	986	27.9	24.4	47.7
0505	636	28.3	20.4	51.3
0506	345	20.9	26.7	52.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0650	1,679	34.5	21.9	43.7
0651	1,055	30.8	22.1	47.1
0652	594	24.1	22.4	53.5
0653	783	27.5	21.6	51.0
0654	604	32.9	20.2	46.9
0655	623	29.7	23.9	46.4
0657	148	23.6	17.6	58.8
0701	592	31.9	20.6	47.5
0702	162	32.1	16.0	51.9
0703	378	28.6	23.5	47.9
0704	155	23.2	22.6	54.2
0705	312	31.1	18.3	50.6
0850	1,837	34.3	22.0	43.7
0851	1,525	31.2	20.7	48.1
0901	2,784	32.3	21.1	46.6
0902	125	29.6	24.8	45.6
1051	2,182	27.5	25.1	47.4
1101	2,912	33.6	22.6	43.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1102	1,435	31.5	21.6	46.9
1106	178	31.5	20.2	48.3
1250	1,833	33.5	17.2	49.3
1251	1,461	31.1	19.0	49.8
1252	894	26.6	20.6	52.8
1253	1,724	32.3	22.7	45.1
1350	3,859	33.2	21.5	45.3
1351	313	36.1	19.2	44.7
1352	731	28.6	21.5	49.9
1401	409	27.9	23.5	48.7
1402	293	28.3	19.5	52.2
1403	63	*	*	*
1404	39	*	*	*
1501	1,994	28.3	20.5	51.2
1503	2,230	29.5	21.0	49.5
1504	4,199	25.6	26.1	48.3
1505	1,608	26.6	24.1	49.3
1506	1,067	30.4	23.1	46.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1507	1,136	28.4	23.6	48.0
1508	254	22.0	24.0	53.9
1509	1,265	29.2	17.5	53.3
1510	161	29.8	24.2	46.0
1511	1,135	32.8	21.1	46.1
1512	538	28.3	19.1	52.6
1701	1,012	31.1	23.8	45.1
1712	174	25.3	13.8	60.9
1950	1,114	33.8	19.7	46.4
1952	271	25.8	24.7	49.4
1953	536	27.4	22.8	49.8
1954	1,075	31.6	21.2	47.2
1955	285	26.7	18.6	54.7
1957	738	35.6	21.5	42.8
1959	303	30.4	21.8	47.9
2150	1,116	30.3	22.3	47.4
2151	422	33.2	21.6	45.3
2251	736	25.1	22.6	52.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
2252	362	33.7	21.5	44.8
2551	5,676	34.4	23.1	42.4
2554	222	24.8	23.4	51.8
2555	6,465	35.7	21.8	42.4
3001	1,690	29.5	20.1	50.4
3002	465	27.5	18.3	54.2
3003	21	*	*	*
3004	356	19.9	21.1	59.0
3007	75	*	*	*
3009	23	*	*	*
3010	5	*	*	*
3201	996	30.9	20.3	48.8
3301	5,977	36.6	20.4	43.1
3302	576	28.3	20.3	51.4
3304	237	24.5	23.6	51.9
3305	210	28.6	25.7	45.7
3306	410	34.1	23.7	42.2
3308	108	32.4	21.3	46.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
3309	84	*	*	*
3550	148	24.3	27.0	48.6
3551	49	*	*	*
3552	77	*	*	*
3701	1,833	29.7	23.3	46.9
3805	5,315	31.2	23.0	45.8
3950	3,146	33.8	22.1	44.1
3952	61	*	*	*
3953	69	*	*	*
3954	192	20.3	26.6	53.1
4001	1,818	31.1	20.9	48.0
4002	1,675	31.2	21.6	47.3
4103	211	21.8	25.6	52.6
4106	1,890	30.1	19.0	50.8
4107	4,987	37.0	19.6	43.4
4109	2,234	30.3	20.9	48.7
4110	2,976	30.3	20.2	49.5
4112	1,183	26.2	21.4	52.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
4113	1,180	31.7	20.8	47.5
4120	3,630	31.1	20.1	48.8
4160	3,084	31.9	21.8	46.3
4351	1,047	26.8	23.1	50.0
4352	354	25.4	24.9	49.7
4353	869	27.4	24.6	48.0
4354	1,289	31.2	21.8	47.0
4355	574	25.4	26.3	48.3
4356	949	26.7	24.4	48.9
4357	9	*	*	*
4358	413	21.8	22.3	55.9
4359	964	24.6	24.6	50.8
4360	281	26.3	24.2	49.5
4361	369	32.0	26.3	41.7
4362	570	28.9	25.6	45.4
4363	1,469	30.0	22.8	47.2
4401	2,234	32.8	22.8	44.4
4501	1,607	28.9	23.5	47.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
4601	2,369	35.0	21.4	43.6
4701	1,910	30.4	25.9	43.7
5050	2,133	40.1	26.2	33.8
5051	3,899	39.6	25.7	34.7
5052	3,178	40.5	25.3	34.2
5101	3,256	35.0	23.7	41.2
5202	933	30.7	23.8	45.6
5203	327	26.6	24.2	49.2
5204	249	29.3	22.5	48.2
5601	1,874	30.0	25.0	45.0
5801	3,264	35.0	22.8	42.2
5803	65	*	*	*
5804	81	*	*	*
6101	136	22.1	23.5	54.4
6102	447	21.9	28.2	49.9
6103	6,247	31.3	23.2	45.5
6104	275	22.9	22.2	54.9
6105	145	17.2	22.8	60.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
6106	38	*	*	*
6301	6,911	35.7	21.6	42.7
6312	3,794	37.3	20.5	42.3
6316	2,033	38.8	19.7	41.6
6320	472	29.0	19.1	51.9
6325	257	34.6	19.5	45.9
6330	655	37.3	19.5	43.2
6335	980	31.9	21.3	46.7
6340	271	28.0	17.7	54.2
6345	69	*	*	*
6350	440	36.1	22.0	41.8
7304	423	28.8	20.6	50.6
7306	4,092	32.1	21.7	46.3
7308	352	29.5	19.6	50.9
7310	69	*	*	*
7401	3,430	30.3	23.7	45.9
7402	329	25.2	31.9	42.9
7403	113	24.8	28.3	46.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
7404	105	21.9	33.3	44.8
7601	1,186	32.5	21.0	46.5
7850	1,557	27.0	24.8	48.2
7851	262	24.8	22.5	52.7
7852	178	28.1	25.3	46.6
8150	151	21.9	19.9	58.3
8151	4,188	37.5	19.9	42.6
8152	521	45.1	23.6	31.3
8153	65	*	*	*
8154	488	29.3	23.2	47.5
8155	142	21.8	23.9	54.2
8156	69	*	*	*
8157	48	*	*	*
8201	2,901	48.3	26.7	25.0
8202	4,108	31.2	25.4	43.4
8203	461	45.3	27.1	27.5
8250	813	62.0	12.9	25.1
8255	3,432	41.8	31.8	26.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
8257	909	30.5	22.1	47.4
8258	2,348	32.6	30.3	37.1
8260	716	29.7	24.2	46.1
8901	1,606	31.6	22.3	46.1
8902	1,016	35.0	19.3	45.7
8903	1,873	33.8	21.9	44.3
8904	1,496	33.0	21.1	45.9
8905	4,801	33.6	19.7	46.8
9004	15	*	*	*
9012	24	*	*	*
9013	84	*	*	*
9015	20	*	*	*
9016	92	*	*	*
9017	33	*	*	*
9018	82	*	*	*
9019	23	*	*	*
9021	26	*	*	*
9040	22	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
9041	942	26.3	25.1	48.6
9045	2,187	27.1	22.6	50.3
9050	3,082	34.3	20.7	45.0
9051	3,380	32.0	24.6	43.5
9053	3,519	38.8	21.4	39.9
9055	233	100.0	0.0	0.0
9058	1,165	28.3	18.3	53.4
9066	3,207	36.5	24.5	39.0
9068	2,056	34.7	22.2	43.1
9069	2,876	43.6	21.1	35.3
9070	2,674	34.5	24.5	41.0
9072	2,265	35.6	22.6	41.8
9074	2,177	37.9	20.5	41.7
9080	3,888	28.0	23.0	48.9
9151	4	*	*	*
Unknown Clinics	134	29.1	33.6	37.3
Michigan	260,806	32.9	22.5	44.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Nation (Prior Year)	8,164,612	35.8	23.0	41.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0101	49	* *	* *	205	2.9 (11)	2.9 (76)	10.2 (33)	93	* *	* *	119	7.6 (46)
0102	135	8.1 (46)	7.4 (66)	591	5.9(113)	5.2(154)	11.8 (69)	275	20.0(142)	11.3 (34)	384	15.4(119)
0103	146	4.1 (2)	15.8(119)	603	4.6 (51)	2.3 (40)	16.3(161)	282	18.1(114)	19.1(153)	391	10.5 (74)
0104	47	* *	* *	209	7.2(160)	2.9 (76)	13.4(114)	111	18.0(111)	15.3(113)	134	6.0 (33)
0250	51	* *	* *	265	5.7(105)	4.5(131)	14.3(132)	124	25.0(162)	16.1(129)	190	14.2(113)
0251	46	* *	* *	219	4.1 (42)	2.3 (40)	11.0 (48)	110	22.7(156)	16.4(136)	144	10.4 (73)
0252	44	* *	* *	247	3.6 (26)	2.0 (26)	14.6(140)	139	18.7(122)	17.3(143)	176	14.8(116)
0253	55	* *	* *	261	2.7 (8)	11.9(179)	6.1 (3)	122	11.5 (8)	4.9 (2)	182	13.2(103)
0401	207	12.6(105)	10.6(110)	852	7.7(171)	3.6(105)	12.2 (75)	403	19.9(140)	14.4 (91)	538	10.2 (70)
0402	145	4.8 (4)	11.7(117)	721	5.0 (70)	3.6(105)	15.3(147)	359	14.5 (33)	20.1(155)	518	10.2 (70)
0403	48	* *	* *	231	6.5(137)	6.5(173)	16.5(165)	119	13.4 (24)	24.4(160)	168	6.0 (33)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0404	51	* *	* *	261	2.3 (5)	3.8(110)	11.5 (59)	142	16.2 (58)	14.8(104)	166	9.6 (62)
0501	174	6.9 (27)	10.9(112)	845	5.0 (70)	3.2 (89)	12.5 (86)	420	17.9(109)	12.9 (57)	595	6.2 (38)
0502	0	* *	* *	6	* *	* *	* *	4	* *	* *	6	* *
0503	75	* *	* *	260	5.8(111)	5.4(159)	10.0 (29)	123	12.2 (13)	7.3 (8)	157	5.7 (30)
0504	202	8.9 (61)	10.4(108)	953	7.1(157)	2.5 (53)	13.1(105)	472	20.1(143)	12.5 (47)	674	3.9 (9)
0505	119	5.9 (10)	10.9(112)	614	5.2 (79)	2.4 (49)	12.2 (75)	318	17.0 (81)	10.7 (27)	451	3.1 (6)
0506	57	* *	* *	337	5.6 (99)	2.7 (67)	15.7(153)	180	17.2 (88)	14.4 (91)	251	3.2 (8)
0650	427	7.5 (34)	5.6 (27)	1,653	6.3(128)	2.5 (53)	13.4(114)	718	15.3 (41)	18.2(149)	1,056	6.4 (40)
0651	243	6.6 (20)	5.8 (32)	1,034	5.7(105)	3.1 (83)	13.8(129)	485	21.2(151)	13.2 (66)	706	7.4 (43)
0652	97	* *	* *	582	6.0(117)	3.3 (94)	14.3(132)	313	22.4(155)	16.3(134)	431	5.6 (29)
0653	160	10.0 (78)	8.1 (79)	772	6.1(121)	2.5 (53)	13.5(121)	390	12.3 (16)	16.2(130)	553	9.6 (62)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0654	146	10.3 (85)	10.3(107)	589	4.6 (51)	4.8(140)	8.8 (16)	278	12.9 (19)	12.2 (45)	405	23.0(154)
0655	133	6.8 (24)	6.0 (36)	613	7.2(160)	2.6 (59)	13.4(114)	283	17.0 (81)	14.5 (95)	416	8.9 (57)
0657	24	* *	* *	148	6.1(121)	2.7 (67)	13.5(121)	85	* *	* *	113	22.1(151)
0701	143	4.9 (5)	11.2(115)	578	6.6(141)	3.3 (94)	8.0 (9)	277	15.2 (40)	7.2 (7)	307	4.6 (20)
0702	28	* *	* *	159	7.5(167)	5.0(149)	15.1(146)	84	* *	* *	102	3.9 (9)
0703	78	* *	* *	372	6.7(145)	2.4 (49)	11.0 (48)	174	13.8 (26)	14.4 (91)	258	0.8 (1)
0704	27	* *	* *	153	6.5(137)	2.6 (59)	17.0(171)	81	* *	* *	102	3.9 (9)
0705	59	* *	* *	306	2.9 (11)	2.3 (40)	8.8 (16)	156	17.3 (94)	9.6 (15)	172	5.8 (32)
0850	425	5.9 (10)	8.7 (91)	1,814	4.7 (56)	3.9(114)	12.5 (86)	790	19.2(131)	13.4 (73)	1,181	4.2 (16)
0851	315	6.7 (22)	7.3 (64)	1,482	3.6 (26)	2.8 (72)	9.0 (20)	729	17.6(100)	10.3 (23)	1,019	4.7 (23)
0901	634	7.9 (38)	9.6(100)	2,752	6.3(128)	1.9 (20)	12.9 (97)	1,277	17.3 (94)	15.3(113)	1,922	11.0 (78)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0902	22	* *	* *	123	9.8(179)	3.3 (94)	13.0(102)	56	* *	* *	90	* *
1051	492	6.5 (18)	9.6(100)	2,133	4.9 (62)	6.2(169)	12.8 (93)	1,027	16.5 (67)	13.6 (76)	1,359	9.3 (61)
1101	727	8.1 (46)	5.4 (25)	2,871	6.3(128)	2.1 (30)	11.7 (66)	1,249	17.0 (81)	12.8 (54)	1,887	22.0(150)
1102	333	6.6 (20)	7.5 (67)	1,416	5.6 (99)	2.5 (53)	13.1(105)	657	17.8(105)	13.1 (64)	965	8.4 (55)
1106	37	* *	* *	175	4.0 (39)	1.1 (4)	12.6 (89)	83	* *	* *	123	10.6 (75)
1250	409	5.6 (8)	8.3 (84)	1,809	6.5(137)	3.3 (94)	14.9(142)	880	19.1(130)	13.9 (84)	1,173	8.1 (52)
1251	297	6.4 (15)	6.4 (44)	1,451	5.9(113)	3.6(105)	12.3 (79)	725	19.9(140)	14.5 (95)	1,035	4.0 (14)
1252	114	7.9 (38)	8.8 (93)	879	5.1 (76)	1.9 (20)	20.7(177)	471	22.3(154)	26.1(162)	638	11.0 (78)
1253	413	8.0 (42)	6.8 (53)	1,702	6.4(132)	4.2(122)	15.0(144)	750	17.2 (88)	17.9(148)	1,192	11.4 (85)
1350	876	8.0 (42)	8.0 (77)	3,807	6.3(128)	3.1 (83)	14.8(141)	1,728	17.7(103)	15.4(116)	2,613	12.5 (97)
1351	73	* *	* *	304	4.3 (44)	3.0 (81)	17.8(172)	132	12.9 (19)	16.7(137)	215	7.4 (43)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
1352	147	9.5 (71)	9.5 (99)	713	4.3 (44)	3.1 (83)	15.8(154)	353	14.2 (28)	15.9(125)	524	22.1(151)
1401	84	* *	* *	407	4.9 (62)	4.9(146)	12.0 (72)	201	12.4 (17)	14.9(105)	262	4.2 (16)
1402	63	* *	* *	286	3.1 (18)	4.2(122)	16.1(157)	150	18.0(111)	15.3(113)	200	5.0 (25)
1403	6	* *	* *	63	* *	* *	* *	41	* *	* *	47	* *
1404	6	* *	* *	38	* *	* *	* *	18	* *	* *	32	* *
1501	389	6.7 (22)	9.3 (98)	1,952	4.5 (48)	2.9 (76)	16.0(156)	985	17.9(109)	14.1 (86)	1,412	28.3(163)
1503	454	6.8 (24)	8.6 (89)	2,187	5.5 (92)	4.7(138)	13.6(125)	1,062	13.3 (23)	13.7 (79)	1,543	29.2(166)
1504	823	8.3 (48)	6.7 (51)	4,112	3.7 (29)	3.9(114)	12.6 (89)	2,012	16.5 (67)	14.1 (86)	2,893	18.6(137)
1505	336	8.0 (42)	10.1(105)	1,577	3.2 (19)	3.5(103)	10.7 (42)	797	13.0 (22)	8.9 (13)	1,076	23.4(156)
1506	239	7.5 (34)	9.6(100)	1,048	5.2 (79)	5.6(162)	16.9(169)	496	20.2(144)	15.7(122)	735	20.1(145)
1507	229	7.0 (29)	8.3 (84)	1,122	2.9 (11)	3.8(110)	10.5 (39)	542	15.9 (56)	11.3 (34)	791	28.7(164)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
1508	31	* *	* *	249	3.6 (26)	4.8(140)	24.5(181)	139	18.7(122)	23.7(158)	164	20.1(145)
1509	265	6.8 (24)	6.8 (53)	1,247	3.8 (34)	3.2 (89)	14.4(136)	661	19.7(138)	13.0 (61)	865	14.5(114)
1510	15	* *	* *	161	3.7 (29)	1.9 (20)	13.0(102)	77	* *	* *	112	12.5 (97)
1511	266	10.2 (83)	7.9 (75)	1,117	2.2 (4)	3.3 (94)	12.8 (93)	510	19.4(135)	15.9(125)	739	21.0(149)
1512	94	* *	* *	526	4.0 (39)	2.7 (67)	14.1(131)	284	18.0(111)	17.3(143)	378	17.5(128)
1701	256	7.0 (29)	10.9(112)	994	5.3 (84)	2.8 (72)	12.8 (93)	444	18.2(117)	12.6 (48)	664	8.0 (51)
1712	27	* *	* *	171	7.6(170)	5.3(156)	12.9 (97)	103	10.7 (6)	14.6(100)	130	14.6(115)
1950	267	6.4 (15)	6.0 (36)	1,099	6.0(117)	2.5 (53)	11.4 (58)	510	19.2(131)	11.4 (37)	740	4.6 (20)
1952	45	* *	* *	268	5.6 (99)	2.2 (35)	13.4(114)	132	14.4 (32)	14.4 (91)	193	8.3 (53)
1953	102	10.8 (94)	5.9 (35)	532	5.5 (92)	1.5 (10)	10.7 (42)	265	18.9(126)	11.3 (34)	375	5.1 (26)
1954	222	9.5 (71)	7.2 (62)	1,062	6.4(132)	2.9 (76)	15.4(150)	502	19.7(138)	15.1(110)	669	6.0 (33)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
1955	48	* *	* *	282	4.3 (44)	1.8 (16)	15.6(152)	157	16.6 (73)	14.0 (85)	187	9.1 (59)
1957	189	7.4 (33)	12.2 (118)	731	6.7(145)	2.2 (35)	16.7(168)	314	16.2 (58)	17.8(147)	481	4.6 (20)
1959	53	* *	* *	301	7.0(155)	2.3 (40)	16.9(169)	147	14.3 (30)	15.0(109)	211	5.7 (30)
2150	262	8.8 (57)	8.8 (93)	1,099	6.7(145)	1.9 (20)	12.3 (79)	526	16.3 (61)	12.7 (51)	731	1.4 (2)
2151	105	9.5 (71)	10.5(109)	419	3.3 (21)	2.6 (59)	11.5 (59)	186	17.7(103)	15.1(110)	276	1.4 (2)
2251	160	9.4 (69)	8.1 (79)	731	6.2(126)	1.2 (7)	11.2 (53)	379	17.2 (88)	13.2 (66)	482	4.4 (18)
2252	92	* *	* *	355	5.4 (87)	3.4(100)	11.5 (59)	159	14.5 (33)	11.9 (40)	232	3.0 (5)
2551	1,415	12.7(108)	3.1 (3)	5,535	7.0(155)	3.9(114)	14.5(137)	2,361	17.4 (96)	13.8 (82)	3,643	24.4(159)
2554	29	* *	* *	218	3.7 (29)	2.3 (40)	19.3(175)	111	16.2 (58)	16.2(130)	156	20.5(148)
2555	1,678	9.1 (65)	6.6 (49)	6,354	5.3 (84)	4.4(130)	13.0(102)	2,697	15.6 (47)	13.6 (76)	4,150	18.6(137)
3001	360	12.5(103)	2.2 (1)	1,630	7.1(157)	2.0 (26)	18.7(173)	842	19.0(127)	16.7(137)	1,166	15.2(118)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
3002	95	* *	* *	451	6.9(151)	1.8 (16)	16.2(159)	245	20.8(149)	15.5(117)	320	18.4(133)
3003	2	* *	* *	21	* *	* *	* *	12	* *	* *	18	* *
3004	52	* *	* *	341	5.6 (99)	1.8 (16)	23.5(180)	204	21.6(152)	24.0(159)	265	6.0 (33)
3007	11	* *	* *	74	* *	* *	* *	40	* *	* *	54	* *
3009	6	* *	* *	22	* *	* *	* *	12	* *	* *	17	* *
3010	0	* *	* *	5	* *	* *	* *	3	* *	* *	5	* *
3201	209	5.3 (7)	9.1 (96)	969	3.8 (34)	2.6 (59)	11.6 (62)	463	16.8 (77)	13.4 (73)	599	11.2 (84)
3301	1,496	7.9 (38)	7.3 (64)	5,884	5.4 (87)	3.1 (83)	13.4(114)	2,549	16.4 (63)	14.9(105)	3,871	10.0 (67)
3302	107	6.5 (18)	5.6 (27)	565	7.3(162)	1.1 (4)	16.1(157)	285	20.7(146)	20.0(154)	402	16.7(120)
3304	41	* *	* *	233	3.0 (16)	4.3(127)	9.0 (20)	120	15.8 (54)	10.0 (21)	172	11.0 (78)
3305	43	* *	* *	207	4.3 (44)	3.9(114)	9.2 (23)	95	* *	* *	146	4.8 (24)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
3306	98	* *	* *	403	6.0(117)	4.2(122)	10.4 (36)	172	17.4 (96)	9.9 (19)	284	3.9 (9)
3308	26	* *	* *	105	2.9 (11)	4.8(140)	5.7 (1)	45	* *	* *	74	* *
3309	14	* *	* *	82	* *	* *	* *	46	* *	* *	62	* *
3550	32	* *	* *	144	5.6 (99)	2.1 (30)	20.1(176)	71	* *	* *	99	* *
3551	7	* *	* *	46	* *	* *	* *	23	* *	* *	39	* *
3552	16	* *	* *	76	* *	* *	* *	40	* *	* *	47	* *
3701	397	7.1 (31)	5.8 (32)	1,791	5.6 (99)	3.6(105)	11.6 (62)	851	20.8(149)	12.9 (57)	1,314	8.7 (56)
3805	1,181	8.5 (50)	6.3 (42)	5,216	4.8 (60)	5.5(161)	10.8 (45)	2,394	17.6(100)	12.6 (48)	2,546	9.8 (65)
3950	726	8.5 (50)	8.4 (88)	3,106	4.7 (56)	5.1(152)	13.5(121)	1,369	16.5 (67)	13.2 (66)	2,020	18.6(137)
3952	4	* *	* *	61	* *	* *	* *	35	* *	* *	47	* *
3953	3	* *	* *	69	* *	* *	* *	42	* *	* *	50	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)													
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)			
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number	% (Rank)
3954	22	* *	* *	190	5.8(111)	4.2(122)	13.2(111)	101	18.8(124)	14.9(105)		142	16.9(124)
4001	390	15.6(118)	3.3 (4)	1,766	5.7(105)	5.9(166)	14.3(132)	852	15.4 (42)	14.2 (88)		1,250	30.0(168)
4002	362	10.8 (94)	5.8 (32)	1,643	5.3 (84)	4.7(138)	16.3(161)	773	19.0(127)	17.3(143)		1,172	23.9(157)
4103	31	* *	* *	203	4.9 (62)	8.4(177)	6.4 (4)	106	14.2 (28)	7.5 (9)		157	19.1(141)
4106	394	8.9 (61)	5.3 (24)	1,862	5.0 (70)	2.6 (59)	14.5(137)	939	15.7 (53)	15.9(125)		1,331	16.8(121)
4107	1,204	8.8 (57)	7.8 (72)	4,933	7.1(157)	1.9 (20)	13.4(114)	2,161	16.1 (57)	13.6 (76)		3,218	14.1(112)
4109	456	10.1 (82)	9.9(104)	2,216	4.6 (51)	2.6 (59)	9.7 (27)	1,081	14.3 (30)	11.1 (33)		1,564	7.6 (46)
4110	611	7.9 (38)	7.7 (69)	2,930	5.7(105)	1.6 (12)	16.3(161)	1,440	18.3(119)	18.3(151)		2,045	12.5 (97)
4112	221	3.2 (1)	11.3(116)	1,157	6.9(151)	4.6(136)	22.2(178)	595	20.7(146)	25.0(161)		940	11.5 (87)
4113	225	8.0 (42)	6.2 (40)	1,165	2.7 (8)	2.1 (30)	15.5(151)	558	16.8 (77)	17.2(141)		862	7.0 (41)
4120	745	8.7 (56)	7.5 (67)	3,575	6.9(151)	1.7 (13)	15.3(147)	1,722	18.1(114)	14.3 (90)		2,545	14.9(117)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
4160	653	11.3 (98)	6.6 (49)	3,049	6.1(121)	2.2 (35)	13.1(105)	1,421	15.6 (47)	13.7 (79)	2,068	13.6(107)
4351	214	10.3 (85)	10.7(111)	1,021	5.2 (79)	3.1 (83)	11.9 (71)	517	17.6(100)	15.7(122)	745	12.5 (97)
4352	71	* *	* *	344	7.8(173)	2.3 (40)	14.5(137)	172	18.6(120)	12.8 (54)	247	18.6(137)
4353	170	5.9 (10)	7.1 (59)	850	5.1 (76)	2.8 (72)	12.9 (97)	407	19.4(135)	16.2(130)	613	13.5(106)
4354	314	5.7 (9)	8.3 (84)	1,274	4.9 (62)	3.2 (89)	9.5 (26)	598	16.4 (63)	9.7 (17)	861	11.8 (90)
4355	124	9.7 (77)	8.1 (79)	560	5.4 (87)	1.4 (9)	12.1 (74)	267	24.3(158)	14.6(100)	378	7.4 (43)
4356	212	8.5 (50)	8.0 (77)	922	6.8(148)	2.7 (67)	12.9 (97)	455	19.3(134)	16.3(134)	644	11.0 (78)
4357	3	* *	* *	8	* *	* *	* *	3	* *	* *	4	* *
4358	73	* *	* *	401	9.2(177)	0.7 (1)	22.4(179)	224	28.6(163)	28.1(163)	301	6.3 (39)
4359	192	4.2 (3)	7.8 (72)	924	7.3(162)	2.3 (40)	19.2(174)	472	16.7 (75)	16.7(137)	633	11.5 (87)
4360	53	* *	* *	279	2.9 (11)	0.7 (1)	9.3 (24)	135	15.6 (47)	11.9 (40)	199	18.1(132)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
4361	97	* *	* *	366	6.6(141)	1.1 (4)	10.4 (36)	151	24.5(160)	13.2 (66)	239	13.4(104)
4362	135	11.1 (96)	8.1 (79)	562	7.5(167)	3.0 (81)	8.7 (14)	259	15.4 (42)	10.8 (29)	397	17.4(127)
4363	305	9.2 (66)	9.2 (97)	1,446	5.0 (70)	2.3 (40)	11.0 (48)	690	17.2 (88)	14.6(100)	1,044	17.2(125)
4401	497	10.7 (93)	7.2 (62)	2,162	9.8(179)	3.5(103)	12.3 (79)	985	17.8(105)	13.7 (79)	1,387	12.4 (96)
4501	345	7.5 (34)	6.4 (44)	1,592	6.8(148)	3.3 (94)	8.7 (14)	759	16.5 (67)	9.9 (19)	1,152	6.1 (37)
4601	519	9.2 (66)	6.4 (44)	2,288	4.9 (62)	4.9(146)	13.2(111)	984	17.8(105)	16.7(137)	1,296	7.6 (46)
4701	449	7.3 (32)	10.2(106)	1,887	4.9 (62)	2.7 (67)	11.0 (48)	825	17.8(105)	15.8(124)	1,292	4.0 (14)
5050	652	10.3 (85)	5.7 (31)	2,104	6.1(121)	2.3 (40)	13.1(105)	710	14.9 (37)	12.4 (46)	1,305	17.8(130)
5051	1,159	10.3 (85)	6.5 (47)	3,830	6.8(148)	2.4 (49)	13.9(130)	1,344	15.6 (47)	10.9 (30)	2,417	19.8(143)
5052	961	10.0 (78)	5.0 (22)	3,129	5.4 (87)	1.7 (13)	13.7(127)	1,077	17.2 (88)	15.5(117)	1,943	18.4(133)
5101	869	7.7 (37)	5.5 (26)	3,220	4.9 (62)	2.0 (26)	11.7 (66)	1,316	16.5 (67)	13.2 (66)	2,045	12.7(102)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
5202	230	8.3 (48)	7.0 (58)	892	4.5 (48)	2.1 (30)	13.5(121)	414	22.9(157)	15.5(117)	530	9.1 (59)
5203	68	* *	* *	306	3.9 (37)	0.7 (1)	16.3(161)	154	17.5 (99)	13.0 (61)	193	5.2 (27)
5204	64	* *	* *	240	1.3 (1)	1.7 (13)	11.3 (55)	118	20.3(145)	12.7 (51)	139	10.1 (69)
5601	454	8.6 (55)	7.9 (75)	1,839	8.1(175)	3.2 (89)	16.6(166)	833	17.0 (81)	14.2 (88)	1,149	5.5 (28)
5801	813	8.5 (50)	8.6 (89)	3,211	7.3(162)	2.1 (30)	13.2(111)	1,362	17.0 (81)	14.7(103)	2,018	10.2 (70)
5803	10	* *	* *	65	* *	* *	* *	29	* *	* *	50	* *
5804	14	* *	* *	80	* *	* *	* *	36	* *	* *	62	* *
6101	21	* *	* *	132	6.1(121)	3.8(110)	9.1 (22)	70	* *	* *	102	9.8 (65)
6102	78	* *	* *	438	3.4 (23)	4.8(140)	10.0 (29)	221	18.6(120)	12.7 (51)	291	11.0 (78)
6103	1,517	9.2 (66)	6.7 (51)	6,092	4.2 (43)	4.6(136)	12.2 (75)	2,826	15.5 (45)	12.8 (54)	3,978	19.3(142)
6104	51	* *	* *	272	3.7 (29)	6.3(170)	14.3(132)	157	11.5 (8)	13.4 (73)	186	19.9(144)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)													
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)			
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number	% (Rank)
6105	20	* *	* *	142	2.8 (10)	5.6(162)	8.5 (12)	81	* *	* *		107	16.8(121)
6106	6	* *	* *	37	* *	* *	* *	21	* *	* *		29	* *
6301	1,704	9.5 (71)	6.9 (56)	6,799	5.7(105)	2.6 (59)	12.0 (72)	2,925	16.5 (67)	15.9(125)		4,586	13.4(104)
6312	970	10.2 (83)	5.6 (27)	3,738	6.4(132)	2.6 (59)	11.3 (55)	1,555	17.2 (88)	12.9 (57)		2,527	17.6(129)
6316	530	6.4 (15)	9.8(103)	2,004	4.7 (56)	2.4 (49)	8.9 (19)	819	16.4 (63)	11.0 (32)		1,345	8.3 (53)
6320	87	* *	* *	460	5.0 (70)	5.0(149)	11.3 (55)	243	10.7 (6)	13.2 (66)		305	7.9 (50)
6325	53	* *	* *	249	2.4 (7)	5.2(154)	11.6 (62)	115	24.3(158)	13.0 (61)		170	7.6 (46)
6330	177	11.9(100)	5.6 (27)	633	6.2(126)	3.2 (89)	13.4(114)	282	14.9 (37)	14.5 (95)		389	17.2(125)
6335	217	6.9 (27)	8.8 (93)	964	4.0 (39)	3.8(110)	9.3 (24)	455	14.1 (27)	8.6 (12)		633	9.6 (62)
6340	53	* *	* *	268	3.0 (16)	3.4(100)	11.2 (53)	150	9.3 (3)	10.7 (27)		177	11.9 (94)
6345	18	* *	* *	69	* *	* *	* *	31	* *	* *		43	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
6350	100	6.0 (14)	4.0 (12)	434	5.1 (76)	4.8(140)	12.7 (91)	175	16.6 (73)	14.9(105)	271	13.7 (109)
7304	75	* *	* *	407	4.9 (62)	3.9(114)	16.2(159)	210	24.8(161)	23.3(157)	311	3.9 (9)
7306	924	8.8 (57)	6.5 (47)	3,947	4.8 (60)	2.9 (76)	13.7(127)	1,853	19.2(131)	14.5 (95)	2,893	12.6(101)
7308	74	* *	* *	344	5.5 (92)	1.5 (10)	11.6 (62)	173	16.8 (77)	16.2(130)	254	11.4 (85)
7310	26	* *	* *	68	* *	* *	* *	28	* *	* *	37	* *
7401	833	8.9 (61)	7.1 (59)	3,342	5.5 (92)	3.4(100)	11.1 (52)	1,547	14.8 (35)	11.8 (39)	2,352	11.8 (90)
7402	49	* *	* *	324	7.7(171)	3.1 (83)	11.7 (66)	143	11.9 (11)	20.3(156)	244	11.1 (83)
7403	21	* *	* *	110	6.4(132)	4.5(131)	10.9 (46)	52	* *	* *	77	* *
7404	16	* *	* *	100	5.0 (70)	4.0(119)	10.0 (29)	50	* *	* *	80	* *
7601	263	4.9 (5)	3.8 (9)	1,163	3.9 (37)	5.8(165)	6.6 (5)	536	9.3 (3)	6.7 (4)	828	4.5 (19)
7850	339	8.8 (57)	6.2 (40)	1,515	3.4 (23)	5.3(156)	12.9 (97)	734	15.5 (45)	17.2(141)	1,063	3.1 (6)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
7851	47	* *	* *	258	6.6(141)	4.3(127)	13.6(125)	137	18.2(117)	18.2(149)	184	7.1 (42)
7852	38	* *	* *	170	3.5 (25)	5.9(166)	10.6 (40)	81	* *	* *	127	1.6 (4)
8150	19	* *	* *	144	2.1 (3)	4.2(122)	9.7 (27)	85	* *	* *	100	10.0 (67)
8151	1,047	9.6 (75)	7.7 (69)	4,100	4.5 (48)	4.0(119)	12.4 (83)	1,757	16.3 (61)	13.3 (72)	2,695	24.9(161)
8152	184	12.0(101)	8.2 (83)	496	3.8 (34)	6.3(170)	12.3 (79)	159	19.5(137)	13.8 (82)	284	34.2(171)
8153	10	* *	* *	64	* *	* *	* *	38	* *	* *	45	* *
8154	92	* *	* *	482	2.3 (5)	5.0(149)	6.0 (2)	234	12.8 (18)	8.1 (10)	291	8.9 (57)
8155	16	* *	* *	137	1.5 (2)	2.2 (35)	8.8 (16)	71	* *	* *	97	* *
8156	9	* *	* *	68	* *	* *	* *	32	* *	* *	47	* *
8157	9	* *	* *	48	* *	* *	* *	31	* *	* *	27	* *
8201	964	8.9 (61)	8.3 (84)	2,848	6.9(151)	2.5 (53)	11.8 (69)	714	17.1 (86)	10.6 (25)	1,589	11.6 (89)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
8202	720	8.5 (50)	6.1 (39)	4,056	3.7 (29)	2.8 (72)	12.4 (83)	1,790	15.6 (47)	13.1 (64)	2,639	11.8 (90)
8203	161	13.0 (109)	8.7 (91)	451	9.3 (178)	1.3 (8)	13.1 (105)	122	22.1 (153)	9.0 (14)	222	18.5 (136)
8250	459	10.5 (90)	6.3 (42)	798	6.0 (117)	5.4 (159)	6.8 (6)	198	13.6 (25)	11.6 (38)	331	26.0 (162)
8255	1,065	10.0 (78)	6.9 (56)	3,383	7.5 (167)	2.0 (26)	13.1 (105)	894	20.7 (146)	17.7 (146)	1,836	13.7 (109)
8257	184	10.3 (85)	4.3 (16)	893	5.5 (92)	4.5 (131)	12.7 (91)	427	12.2 (13)	12.9 (57)	569	23.0 (154)
8258	609	9.4 (69)	7.1 (59)	2,311	7.3 (162)	1.9 (20)	12.8 (93)	841	18.8 (124)	15.6 (120)	1,528	13.7 (109)
8260	151	10.6 (91)	3.3 (4)	703	5.5 (92)	5.3 (156)	8.3 (10)	322	14.9 (37)	7.1 (6)	469	24.1 (158)
8901	329	15.2 (115)	3.3 (4)	1,552	5.7 (105)	5.1 (152)	12.4 (83)	728	12.2 (13)	10.9 (30)	879	41.1 (172)
8902	241	11.2 (97)	4.6 (18)	958	5.2 (79)	7.5 (175)	12.5 (86)	432	11.6 (10)	12.0 (43)	594	29.1 (165)
8903	427	13.3 (110)	4.0 (12)	1,793	4.6 (51)	5.7 (164)	10.4 (36)	818	15.8 (54)	10.0 (21)	1,148	42.2 (174)
8904	325	13.5 (113)	4.0 (12)	1,435	4.6 (51)	7.6 (176)	10.7 (42)	676	14.8 (35)	12.1 (44)	957	41.3 (173)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
8905	1,159	9.6 (75)	6.8 (53)	4,622	5.2 (79)	7.4(174)	15.0(144)	2,178	16.8 (77)	15.6(120)	2,112	33.5 (170)
9004	15	* *	* *	10	* *	* *	* *	0	* *	* *	0	* *
9012	7	* *	* *	21	* *	* *	* *	8	* *	* *	4	* *
9013	32	* *	* *	81	* *	* *	* *	35	* *	* *	44	* *
9015	8	* *	* *	20	* *	* *	* *	7	* *	* *	6	* *
9016	27	* *	* *	90	* *	* *	* *	37	* *	* *	47	* *
9017	10	* *	* *	31	* *	* *	* *	13	* *	* *	4	* *
9018	38	* *	* *	78	* *	* *	* *	29	* *	* *	7	* *
9019	11	* *	* *	23	* *	* *	* *	12	* *	* *	2	* *
9021	13	* *	* *	25	* *	* *	* *	3	* *	* *	4	* *
9040	6	* *	* *	20	* *	* *	* *	11	* *	* *	11	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
9041	142	13.4 (111)	4.9 (21)	888	6.4(132)	4.5(131)	14.9(142)	432	17.1 (86)	10.4 (24)	263	29.7 (167)
9045	451	10.0 (78)	7.8 (72)	2,137	6.6(141)	2.2 (35)	16.6(166)	1,059	17.4 (96)	14.5 (95)	937	22.5 (153)
9050	738	12.6 (105)	6.0 (36)	3,000	7.4(166)	4.5(131)	15.3(147)	1,363	15.4 (42)	15.2(112)	1,878	13.6 (107)
9051	730	5.9 (10)	7.7 (69)	3,311	4.7 (56)	1.8 (16)	15.9(155)	1,419	19.0(127)	18.4(152)	2,218	10.9 (76)
9053	936	13.4 (111)	4.1 (15)	3,447	6.5(137)	4.8(140)	10.9 (46)	1,385	16.7 (75)	11.9 (40)	2,298	18.4 (133)
9055	232	12.1 (102)	5.2 (23)	215	9.8(179)	12.1(181)	7.9 (8)	0	* *	* *	1	* *
9058	235	10.6 (91)	4.7 (20)	1,147	3.2 (19)	4.1(121)	8.6 (13)	616	11.9 (11)	8.1 (10)	798	20.4 (147)
9066	833	13.7 (114)	3.7 (8)	3,118	5.9(113)	6.3(170)	10.3 (35)	1,220	10.2 (5)	6.9 (5)	1,819	11.9 (94)
9068	496	15.5 (117)	3.8 (9)	1,969	5.4 (87)	11.9(179)	7.4 (7)	835	4.4 (1)	4.1 (1)	1,308	32.3 (169)
9069	856	12.5 (103)	4.3 (16)	2,800	5.5 (92)	6.1(168)	10.2 (33)	1,000	16.4 (63)	12.6 (48)	1,556	11.8 (90)
9070	692	11.4 (99)	3.3 (4)	2,600	5.9(113)	8.5(178)	10.0 (29)	1,060	6.7 (2)	5.4 (3)	1,538	16.8 (121)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
9072	608	16.4 (119)	3.9 (11)	2,197	8.0(174)	3.6(105)	12.2 (75)	921	18.1(114)	9.6 (15)	1,419	10.9 (76)
9074	596	15.4 (116)	2.9 (2)	2,120	8.6(176)	4.3(127)	10.6 (40)	898	15.6 (47)	10.6 (25)	1,336	18.0 (131)
9080	842	12.6 (105)	4.6 (18)	3,848	3.3 (21)	4.9(146)	8.4 (11)	1,882	12.9 (19)	9.8 (18)	2,688	24.7 (160)
9151	0	* *	* *	1	* *	* *	* *	1	* *	* *	2	* *
Unknown Clinics	16	* *	* *	75	* *	* *	* *	28	* *	* *	87	* *
Michigan	61,144	9.4	6.8	255,398	5.6	3.6	12.8	114,489	16.6	13.7	166,887	15.0
Nation (Prior Year)	1,909,620	9.0	6.4	7,107,000	6.0	4.5	14.1	2,943,996	16.5	14.8	4,916,523	14.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)												
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)
0101	51	* *	49	* *	28	* *	39	* *	45	* *	91	* *
0102	138	54.3 (73)	82	* *	99	* *	47	* *	63	* *	253	83.8 (94)
0103	148	60.8 (47)	93	* *	97	* *	51	* *	64	* *	275	81.8(115)
0104	47	* *	30	* *	24	* *	24	* *	29	* *	96	* *
0250	51	* *	48	* *	60	* *	28	* *	41	* *	113	90.3 (29)
0251	46	* *	35	* *	33	* *	15	* *	19	* *	96	* *
0252	44	* *	27	* *	42	* *	22	* *	26	* *	133	89.5 (38)
0253	55	* *	46	* *	43	* *	30	* *	43	* *	120	86.7 (66)
0401	208	42.8(103)	147	19.7 (51)	138	10.9 (21)	115	17.4 (14)	135	8.9 (15)	361	95.6 (7)
0402	145	66.2 (19)	108	23.1 (25)	113	7.1 (57)	73	* *	74	* *	338	89.1 (42)
0403	48	* *	29	* *	38	* *	31	* *	29	* *	111	83.8 (94)
0404	51	* *	28	* *	40	* *	23	* *	14	* *	126	90.5 (25)
0501	187	67.4 (15)	143	21.7 (39)	132	10.6 (24)	95	* *	123	7.3 (24)	402	89.1 (42)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% (Rank)	Number	% (Rank)				
0502	0	* *	1	* *	2	* *	0	* *	1	* *	4	* *	6	* *
0503	76	* *	36	* *	28	* *	39	* *	31	* *	122	91.8 (16)	248	3.2 (16)
0504	212	75.0 (3)	119	22.7 (30)	125	10.4 (25)	63	* *	56	* *	436	91.3 (20)	854	5.6 (42)
0505	127	74.8 (4)	97	* *	81	* *	74	* *	83	* *	293	87.7 (61)	566	8.1 (89)
0506	57	* *	34	* *	57	* *	21	* *	23	* *	169	90.5 (25)	302	21.2 (178)
0650	428	64.0 (31)	333	24.0 (22)	298	8.7 (45)	226	14.2 (25)	266	7.9 (19)	783	85.2 (81)	1,675	10.1 (123)
0651	243	51.4 (82)	194	10.8 (98)	174	4.0 (96)	141	9.9 (52)	165	3.6 (67)	538	86.8 (65)	1,055	16.4 (170)
0652	98	* *	83	* *	91	* *	59	* *	62	* *	341	82.7(107)	590	16.9 (172)
0653	160	60.6 (49)	116	25.0 (19)	127	5.5 (79)	61	* *	85	* *	422	83.6 (97)	779	22.1 (179)
0654	147	69.4 (10)	100	17.0 (64)	102	7.8 (53)	83	* *	91	* *	302	87.7 (61)	604	13.9 (163)
0655	133	70.7 (8)	132	15.2 (76)	124	8.1 (51)	81	* *	108	1.9 (86)	313	88.5 (50)	621	13.2 (155)
0657	24	* *	18	* *	18	* *	10	* *	14	* *	93	* *	148	14.2 (164)
0701	145	78.6 (2)	106	37.7 (2)	91	* *	87	* *	96	* *	271	97.8 (3)	546	1.6 (8)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% (Rank)	Number	% (Rank)				
0702	26	* *	31	* *	21	* *	22	* *	29	* *	75	* *	141	8.5 (97)
0703	78	* *	64	* *	64	* *	43	* *	60	* *	174	93.1 (12)	337	8.3 (93)
0704	27	* *	16	* *	19	* *	10	* *	15	* *	75	* *	133	15.8 (169)
0705	59	* *	57	* *	43	* *	46	* *	52	* *	152	98.0 (1)	298	2.3 (10)
0850	434	69.4 (10)	371	25.9 (14)	333	9.6 (36)	299	15.4 (19)	344	7.0 (27)	858	86.2 (72)	1,835	5.4 (39)
0851	321	59.8 (54)	282	22.0 (33)	267	9.0 (40)	204	17.2 (15)	263	5.7 (44)	786	86.5 (67)	1,522	6.2 (55)
0901	633	53.6 (75)	517	15.3 (75)	429	5.1 (87)	386	10.9 (48)	393	3.6 (67)	1,226	90.0 (33)	2,577	9.7 (116)
0902	21	* *	21	* *	20	* *	9	* *	10	* *	59	* *	116	11.2 (139)
1051	495	82.2 (1)	360	34.4 (4)	348	16.4 (2)	246	26.8 (3)	322	10.2 (7)	934	88.3 (52)	1,961	5.1 (31)
1101	728	43.3(101)	471	12.3 (92)	481	6.9 (59)	300	8.3 (62)	404	3.0 (73)	1,349	72.3(143)	2,905	10.8 (132)
1102	334	47.6 (92)	282	21.6 (40)	258	12.4 (12)	199	13.1 (32)	254	6.3 (37)	716	82.0(112)	1,434	12.4 (154)
1106	37	* *	33	* *	33	* *	22	* *	28	* *	89	* *	178	13.5 (158)
1250	411	59.1 (57)	280	22.9 (28)	229	10.0 (30)	259	14.3 (24)	265	8.3 (17)	958	83.6 (97)	1,833	9.7 (116)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
1251	303	64.4 (28)	223	27.4 (11)	210	9.0 (40)	169	20.7 (5)	202	13.9 (1)	778	87.9 (58)	1,459	8.6 (100)
1252	117	63.2 (35)	132	15.2 (76)	131	5.3 (83)	90	* *	103	8.7 (16)	518	80.3(123)	894	6.4 (60)
1253	416	59.4 (56)	288	18.8 (57)	308	5.5 (79)	255	14.1 (27)	235	7.7 (22)	812	83.4(100)	1,724	8.6 (100)
1350	883	47.5 (94)	679	12.5 (91)	606	3.3 (100)	556	8.3 (62)	637	3.8 (66)	1,867	88.2 (54)	3,843	6.3 (58)
1351	73	* *	53	* *	51	* *	48	* *	52	* *	151	98.0 (1)	313	4.5 (25)
1352	147	51.0 (84)	102	10.8 (98)	123	4.9 (89)	94	* *	98	* *	387	83.2(102)	731	13.7 (161)
1401	84	* *	61	* *	51	* *	52	* *	54	* *	191	84.8 (85)	370	5.9 (47)
1402	63	* *	30	* *	43	* *	13	* *	15	* *	135	89.6 (36)	266	7.1 (73)
1403	6	* *	5	* *	10	* *	6	* *	5	* *	37	* *	58	* *
1404	6	* *	5	* *	4	* *	3	* *	5	* *	17	* *	37	* *
1501	403	66.0 (21)	305	28.5 (8)	314	8.3 (49)	218	14.2 (25)	236	6.8 (29)	1,053	84.7 (86)	1,980	7.4 (79)
1503	473	63.6 (32)	345	25.2 (17)	371	11.6 (17)	266	13.5 (29)	267	9.0 (14)	1,155	81.9(113)	2,219	8.2 (92)
1504	880	67.0 (16)	496	20.2 (48)	756	9.1 (39)	280	15.0 (21)	388	5.7 (44)	2,118	84.6 (87)	4,175	2.9 (13)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
1505	337	72.1 (6)	188	25.5 (16)	260	12.7 (9)	116	24.1 (4)	160	10.0 (8)	855	89.6 (36)	1,600	2.9 (13)
1506	241	68.0 (14)	187	21.9 (36)	191	6.8 (63)	116	17.2 (15)	145	12.4 (2)	535	88.2 (54)	1,066	6.8 (65)
1507	241	65.6 (23)	170	29.4 (7)	206	10.2 (27)	100	27.0 (2)	151	10.6 (6)	575	89.7 (35)	1,135	5.8 (45)
1508	39	* *	38	* *	43	* *	22	* *	33	* *	146	91.1 (21)	253	0.4 (1)
1509	270	61.9 (44)	182	18.1 (61)	185	8.1 (51)	151	11.3 (44)	152	6.6 (33)	679	82.9(105)	1,257	6.7 (63)
1510	16	* *	23	* *	25	* *	11	* *	21	* *	79	* *	160	8.8 (103)
1511	271	65.7 (22)	199	18.6 (60)	198	9.6 (36)	159	9.4 (54)	182	5.5 (47)	545	83.9 (92)	1,133	5.0 (28)
1512	96	* *	84	* *	79	* *	67	* *	78	* *	302	88.4 (51)	538	6.1 (50)
1701	255	61.2 (46)	169	21.9 (36)	165	8.5 (47)	116	18.1 (11)	125	6.4 (36)	395	88.9 (46)	877	8.3 (93)
1712	27	* *	26	* *	19	* *	17	* *	22	* *	89	* *	149	11.4 (141)
1950	266	62.0 (43)	190	15.8 (71)	163	6.7 (64)	122	13.1 (32)	162	8.0 (18)	546	86.3 (70)	1,114	11.4 (141)
1952	45	* *	47	* *	41	* *	28	* *	37	* *	143	89.5 (38)	270	9.6 (114)
1953	102	58.8 (58)	87	* *	83	* *	55	* *	69	* *	283	86.9 (63)	535	9.3 (110)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
1954	244	60.2 (52)	150	18.7 (59)	157	12.7 (9)	110	13.6 (28)	125	4.8 (57)	546	94.3 (9)	1,067	6.2 (55)
1955	55	* *	36	* *	34	* *	24	* *	24	* *	164	89.0 (44)	285	6.0 (48)
1957	190	66.8 (17)	154	22.1 (32)	124	12.9 (8)	108	20.4 (8)	138	5.1 (53)	341	86.2 (72)	734	5.4 (39)
1959	52	* *	57	* *	57	* *	33	* *	48	* *	157	87.9 (58)	303	5.3 (34)
2150	262	57.3 (64)	188	20.7 (46)	158	10.8 (22)	117	15.4 (19)	130	5.4 (48)	485	93.0 (13)	994	6.8 (65)
2151	105	47.6 (92)	81	* *	72	* *	71	* *	78	* *	158	88.0 (56)	357	5.3 (34)
2251	160	60.6 (49)	121	26.4 (13)	115	13.0 (7)	99	* *	110	10.9 (4)	320	90.9 (24)	626	5.0 (28)
2252	92	* *	70	* *	56	* *	52	* *	66	* *	133	91.7 (18)	308	7.1 (73)
2551	1,432	35.0(111)	1,206	6.1 (108)	928	1.7 (104)	899	4.7 (81)	1,038	1.8 (88)	2,258	76.7(133)	5,203	8.4 (96)
2554	29	* *	36	* *	47	* *	9	* *	32	* *	107	86.9 (63)	201	11.9 (150)
2555	1,719	49.4 (86)	1,273	12.6 (89)	1,050	4.8 (91)	929	9.1 (56)	970	3.9 (65)	2,588	84.9 (84)	5,945	6.0 (48)
3001	361	45.2 (98)	284	8.8 (103)	247	3.2 (102)	257	6.2 (72)	259	2.3 (78)	801	75.2(136)	1,547	13.5 (158)
3002	93	* *	68	* *	61	* *	54	* *	64	* *	237	81.9(113)	431	7.2 (78)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% (Rank)	Number	% (Rank)				
3003	2	* *	2	* *	1	* *	1	* *	2	* *	12	* *	20	* *
3004	53	* *	45	* *	46	* *	26	* *	31	* *	197	73.6(139)	326	1.5 (7)
3007	10	* *	11	* *	8	* *	8	* *	11	* *	37	* *	70	* *
3009	6	* *	2	* *	4	* *	2	* *	2	* *	12	* *	23	* *
3010	0	* *	1	* *	1	* *	0	* *	0	* *	3	* *	4	* *
3201	211	52.1 (79)	165	15.8 (71)	131	6.9 (59)	131	13.0 (35)	146	1.4 (94)	451	92.5 (14)	906	7.5 (81)
3301	1,516	60.6 (49)	1,131	21.6 (40)	960	10.3 (26)	835	6.6 (69)	1,015	3.1 (71)	2,719	80.1(125)	5,951	6.1 (50)
3302	108	56.5 (65)	82	* *	90	* *	56	* *	76	* *	287	72.5(142)	575	6.1 (50)
3304	41	* *	29	* *	38	* *	19	* *	27	* *	128	79.7(128)	237	0.8 (2)
3305	45	* *	26	* *	33	* *	17	* *	24	* *	104	85.6 (76)	210	2.9 (13)
3306	100	58.0 (62)	68	* *	76	* *	56	* *	63	* *	187	93.6 (11)	410	7.1 (73)
3308	26	* *	18	* *	17	* *	18	* *	16	* *	52	* *	105	7.6 (82)
3309	14	* *	9	* *	13	* *	8	* *	8	* *	51	* *	84	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)			Exclusive Breastfeeding (7)			≤2 Hours TV Viewing/Day (8)		Smoking in Household					
	Number	% Ever Breastfed (Rank)	Number Breastfed At Least 6 Months (Rank)	Number Breastfed At Least 12 Months (Rank)	Number Breastfed At Least 3 Months (Rank)	Number Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)				
3550	32	* *	25	* *	25	* *	20	* *	25	* *	74	* *	135	4.4 (23)
3551	7	* *	5	* *	12	* *	4	* *	5	* *	23	* *	41	* *
3552	16	* *	8	* *	14	* *	8	* *	8	* *	36	* *	67	* *
3701	395	63.5 (33)	270	14.4 (82)	327	6.7 (64)	156	20.5 (6)	228	7.9 (19)	921	96.2 (5)	1,831	9.3 (110)
3805	1,187	57.4 (63)	1,051	17.6 (62)	975	6.6 (68)	812	9.1 (56)	1,005	5.1 (53)	2,583	82.9(105)	5,295	8.5 (97)
3950	732	56.4 (66)	605	23.0 (27)	575	10.8 (22)	450	12.7 (36)	527	6.6 (33)	1,476	82.6(109)	3,145	6.3 (58)
3952	4	* *	11	* *	13	* *	8	* *	11	* *	37	* *	61	* *
3953	3	* *	7	* *	13	* *	5	* *	4	* *	43	* *	69	* *
3954	22	* *	24	* *	35	* *	19	* *	21	* *	114	90.4 (27)	192	2.1 (9)
4001	402	55.7 (68)	274	19.0 (56)	285	6.7 (64)	219	9.1 (56)	195	5.1 (53)	935	59.5(161)	1,817	13.4 (157)
4002	366	62.3 (39)	263	23.6 (23)	270	10.0 (30)	206	13.1 (32)	220	6.8 (29)	849	66.2(153)	1,671	9.6 (114)
4103	31	* *	20	* *	41	* *	15	* *	20	* *	115	69.6(148)	210	10.0 (122)
4106	396	54.3 (73)	304	22.0 (33)	277	8.7 (45)	260	5.0 (78)	273	5.1 (53)	1,011	80.5(122)	1,883	6.1 (50)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	%	Number	% (Rank)
4107	1,213	62.3 (39)	1,093	25.2 (17)	779	10.0 (30)	823	11.1 (47)	959	6.8 (29)	2,304	85.3 (80)	4,974	5.2 (33)
4109	455	59.6 (55)	378	22.0 (33)	348	7.8 (53)	286	10.8 (50)	300	9.7 (10)	1,170	88.3 (52)	2,230	6.5 (62)
4110	620	62.3 (39)	547	21.8 (38)	494	11.7 (16)	390	6.2 (72)	444	4.1 (62)	1,573	86.4 (68)	2,965	6.1 (50)
4112	223	63.2 (35)	134	38.8 (1)	188	16.0 (4)	108	3.7 (83)	122	4.1 (62)	670	94.6 (8)	1,181	4.3 (21)
4113	224	65.2 (24)	195	32.8 (5)	186	15.1 (5)	158	6.3 (71)	116	2.6 (76)	607	84.3 (88)	1,177	1.4 (6)
4120	750	64.8 (25)	662	20.8 (45)	555	9.0 (40)	500	8.8 (60)	578	4.7 (59)	1,901	86.3 (70)	3,621	3.7 (18)
4160	662	60.7 (48)	536	24.6 (21)	494	9.7 (34)	399	9.0 (59)	454	5.9 (42)	1,531	83.9 (92)	3,067	4.4 (23)
4351	214	68.7 (13)	127	22.8 (29)	159	10.1 (29)	78	* *	115	9.6 (12)	520	85.8 (74)	968	11.8 (148)
4352	71	* *	53	* *	50	* *	37	* *	51	* *	159	78.6(131)	315	18.7 (176)
4353	173	60.1 (53)	140	25.7 (15)	123	11.4 (18)	99	* *	104	11.5 (3)	407	75.9(135)	797	14.2 (164)
4354	315	62.9 (37)	218	21.6 (40)	207	9.2 (38)	178	19.1 (9)	194	9.8 (9)	564	91.8 (16)	1,184	9.9 (119)
4355	124	66.1 (20)	85	* *	95	* *	56	* *	69	* *	247	90.3 (29)	509	7.7 (84)
4356	213	58.7 (59)	136	16.2 (66)	169	8.9 (43)	83	* *	114	6.1 (38)	430	85.8 (74)	858	9.7 (116)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)												
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)
4357	3	* *	2	* *	1	* *	2	* *	2	* *	8	* *
4358	75	* *	54	* *	51	* *	41	* *	52	* *	220	85.0 (82) 384 1.3 (5)
4359	196	69.4 (10)	106	14.2 (83)	150	3.3 (100)	76	* *	98	* *	480	82.7(107) 875 9.0 (106)
4360	54	* *	37	* *	50	* *	30	* *	34	* *	134	88.8 (47) 261 11.9 (150)
4361	97	* *	63	* *	66	* *	29	* *	38	* *	143	90.2 (31) 330 10.9 (133)
4362	134	74.6 (5)	70	* *	101	8.9 (43)	25	* *	35	* *	240	90.4 (27) 520 15.4 (168)
4363	304	71.1 (7)	237	16.0 (68)	225	6.2 (70)	186	18.3 (10)	224	7.1 (25)	664	90.2 (31) 1,339 9.9 (119)
4401	495	64.4 (28)	402	19.7 (51)	336	11.3 (19)	315	12.7 (36)	355	5.9 (42)	891	90.0 (33) 1,907 9.4 (112)
4501	346	58.1 (61)	198	16.2 (66)	259	10.0 (30)	113	17.7 (13)	149	6.7 (32)	729	83.7 (96) 1,482 9.2 (109)
4601	591	61.4 (45)	472	20.1 (49)	404	7.2 (56)	354	11.3 (44)	422	3.1 (71)	1,046	85.5 (78) 2,248 5.6 (42)
4701	448	64.7 (26)	291	20.3 (47)	341	5.9 (73)	197	17.8 (12)	275	6.9 (28)	852	88.8 (47) 1,845 5.0 (28)
5050	649	49.0 (87)	398	19.6 (53)	376	5.3 (83)	302	13.2 (30)	326	7.1 (25)	716	81.8(115) 1,949 5.7 (44)
5051	1,157	52.5 (78)	704	19.5 (55)	718	6.5 (69)	589	9.7 (53)	583	4.3 (60)	1,343	81.8(115) 3,603 5.8 (45)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
5052	964	47.4 (95)	632	13.3 (84)	605	5.5 (79)	504	7.3 (67)	541	3.0 (73)	1,072	80.1(125)	2,951	6.8 (65)
5101	883	62.9 (37)	682	20.1 (49)	608	8.4 (48)	564	12.2 (41)	593	4.2 (61)	1,315	83.4(100)	3,104	5.3 (34)
5202	229	64.6 (27)	143	27.3 (12)	145	12.4 (12)	127	20.5 (6)	134	9.7 (10)	378	91.0 (22)	798	5.1 (31)
5203	69	* *	52	* *	44	* *	36	* *	48	* *	145	89.0 (44)	281	7.1 (73)
5204	64	* *	41	* *	35	* *	37	* *	40	* *	104	94.2 (10)	212	4.2 (20)
5601	452	70.4 (9)	271	24.7 (20)	300	9.7 (34)	155	29.7 (1)	243	9.1 (13)	848	89.3 (40)	1,769	5.4 (39)
5801	813	55.0 (70)	649	16.6 (65)	574	5.7 (77)	503	12.5 (39)	598	6.0 (40)	1,435	82.5(111)	3,189	9.9 (119)
5803	10	* *	12	* *	18	* *	9	* *	11	* *	32	* *	65	* *
5804	14	* *	19	* *	20	* *	13	* *	15	* *	40	* *	81	* *
6101	22	* *	20	* *	28	* *	16	* *	16	* *	77	* *	135	14.8 (167)
6102	79	* *	45	* *	85	* *	27	* *	38	* *	236	91.9 (15)	443	8.8 (103)
6103	1,521	47.9 (91)	985	12.9 (86)	1,088	5.0 (88)	760	9.2 (55)	840	2.6 (76)	3,001	82.6(109)	6,232	9.1 (107)
6104	51	* *	40	* *	46	* *	28	* *	39	* *	159	73.6(139)	274	16.4 (170)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)												
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)
6105	20	* *	11	* *	26	* *	7	* *	9	* *	88	* *
6106	6	* *	3	* *	7	* *	3	* *	3	* *	21	* *
6301	1,718	45.0 (99)	1,336	17.1 (63)	1,056	7.0 (58)	1,052	12.7 (36)	1,205	5.4 (48)	2,910	83.0(104)
6312	973	56.0 (67)	731	23.1 (25)	561	11.8 (15)	594	12.5 (39)	672	6.1 (38)	1,536	79.4(130)
6316	534	54.5 (72)	461	15.4 (74)	316	5.7 (77)	337	14.5 (22)	398	5.3 (50)	823	81.7(118)
6320	93	* *	83	* *	65	* *	67	* *	78	* *	225	88.0 (56)
6325	54	* *	49	* *	41	* *	48	* *	48	* *	103	86.4 (68)
6330	180	39.4(106)	124	11.3 (97)	82	* *	123	5.7 (76)	121	1.7 (90)	277	74.7(137)
6335	219	39.7(105)	164	14.6 (80)	142	12.7 (9)	151	13.2 (30)	154	6.5 (35)	456	87.9 (58)
6340	54	* *	37	* *	30	* *	32	* *	33	* *	143	91.6 (19)
6345	17	* *	16	* *	8	* *	11	* *	13	* *	31	* *
6350	104	37.5(107)	88	* *	65	* *	75	* *	78	* *	180	85.6 (76)
7304	78	* *	51	* *	46	* *	36	* *	40	* *	210	84.3 (88)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
7306	925	46.8 (96)	583	11.8 (94)	562	2.8 (103)	436	8.5 (61)	426	3.5 (69)	1,805	81.3(119)	3,708	8.8 (103)
7308	74	* *	38	* *	46	* *	30	* *	24	* *	167	89.2 (41)	314	11.5 (144)
7310	26	* *	11	* *	5	* *	10	* *	8	* *	24	* *	62	* *
7401	832	53.0 (77)	617	12.8 (88)	613	5.9 (73)	414	10.9 (48)	537	5.6 (46)	1,540	84.2 (91)	3,212	11.5 (144)
7402	49	* *	55	* *	78	* *	27	* *	40	* *	133	91.0 (22)	295	6.8 (65)
7403	21	* *	20	* *	24	* *	9	* *	16	* *	53	* *	103	10.7 (131)
7404	16	* *	21	* *	26	* *	7	* *	14	* *	50	* *	95	* *
7601	261	55.6 (69)	212	13.2 (85)	178	4.5 (92)	161	16.1 (17)	173	5.2 (52)	510	84.3 (88)	1,088	10.1 (123)
7850	340	58.5 (60)	194	12.9 (86)	282	6.7 (64)	112	8.0 (64)	175	1.7 (90)	778	88.7 (49)	1,549	10.1 (123)
7851	47	* *	18	* *	33	* *	9	* *	17	* *	143	96.5 (4)	258	8.5 (97)
7852	38	* *	17	* *	33	* *	7	* *	16	* *	90	* *	178	10.1 (123)
8150	19	* *	15	* *	22	* *	13	* *	11	* *	94	* *	151	8.6 (100)
8151	1,053	63.4 (34)	805	28.3 (9)	621	16.1 (3)	648	15.6 (18)	678	7.4 (23)	1,851	78.1(132)	4,119	10.2 (127)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
8152	185	54.6 (71)	124	14.5 (81)	97	* *	90	* *	112	0.9 (97)	170	83.5 (99)	509	22.4 (180)
8153	10	* *	5	* *	11	* *	10	* *	3	* *	41	* *	64	* *
8154	92	* *	70	* *	82	* *	37	* *	31	* *	239	96.2 (5)	456	1.1 (3)
8155	16	* *	20	* *	21	* *	9	* *	14	* *	82	* *	139	13.7 (161)
8156	10	* *	11	* *	9	* *	5	* *	6	* *	33	* *	69	* *
8157	9	* *	5	* *	4	* *	6	* *	5	* *	31	* *	47	* *
8201	968	51.9 (80)	737	15.7 (73)	600	8.2 (50)	592	11.8 (42)	686	3.4 (70)	769	85.4 (79)	2,767	5.3 (34)
8202	849	66.4 (18)	625	35.5 (3)	717	12.0 (14)	364	11.3 (44)	464	7.8 (21)	1,877	80.8(121)	3,917	5.3 (34)
8203	161	53.4 (76)	63	* *	86	* *	37	* *	38	* *	127	85.0 (82)	425	7.1 (73)
8250	465	41.5(104)	155	14.8 (78)	74	* *	181	7.2 (68)	146	4.8 (57)	209	72.7(141)	810	6.4 (60)
8255	1,080	49.0 (87)	580	16.0 (68)	740	6.1 (71)	391	10.5 (51)	530	5.3 (50)	972	79.6(129)	3,232	8.1 (89)
8257	207	50.2 (85)	113	32.7 (6)	127	10.2 (27)	84	* *	101	10.9 (4)	444	81.3(119)	877	7.9 (88)
8258	619	43.9(100)	284	6.3 (107)	487	5.3 (83)	163	8.0 (64)	228	1.3 (95)	896	83.1(103)	2,189	6.7 (63)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
8260	151	64.2 (30)	130	27.7 (10)	120	17.5 (1)	84	* *	124	4.0 (64)	323	60.7(159)	674	6.8 (65)
8901	330	23.9(120)	258	11.6 (96)	242	4.5 (92)	205	3.4 (85)	213	2.3 (78)	737	61.9(158)	1,523	17.6 (173)
8902	242	34.7(112)	183	14.8 (78)	144	6.9 (59)	147	4.1 (82)	154	0.6(101)	448	67.9(151)	964	12.0 (152)
8903	434	27.4(117)	300	10.0 (100)	275	4.0 (96)	260	2.3 (90)	265	1.5 (93)	846	59.6(160)	1,792	12.1 (153)
8904	328	43.0(102)	236	23.3 (24)	215	7.4 (55)	183	11.5 (43)	179	2.2 (80)	689	69.7(147)	1,421	11.8 (148)
8905	1,180	34.3(113)	782	21.6 (40)	635	5.8 (75)	644	3.7 (83)	599	3.0 (73)	2,246	74.3(138)	4,616	8.1 (89)
9004	15	* *	1	* *	0	* *	1	* *	1	* *	0	* *	14	* *
9012	9	* *	0	* *	4	* *	0	* *	0	* *	9	* *	23	* *
9013	32	* *	10	* *	12	* *	10	* *	10	* *	35	* *	84	* *
9015	8	* *	2	* *	2	* *	1	* *	2	* *	7	* *	20	* *
9016	27	* *	12	* *	15	* *	7	* *	12	* *	40	* *	92	* *
9017	10	* *	2	* *	6	* *	2	* *	2	* *	14	* *	33	* *
9018	38	* *	9	* *	6	* *	9	* *	8	* *	32	* *	82	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
9019	11	* *	0	* *	0	* *	0	* *	0	* *	12	* *	23	* *
9021	12	* *	5	* *	6	* *	3	* *	3	* *	3	* *	26	* *
9040	6	* *	2	* *	1	* *	1	* *	2	* *	13	* *	22	* *
9041	141	35.5(110)	152	6.6 (106)	160	4.4 (94)	67	* *	126	0.8 (98)	447	80.1(125)	872	11.1 (135)
9045	453	46.8 (96)	301	15.9 (70)	325	4.9 (89)	213	7.5 (66)	226	2.2 (80)	1,138	69.9(146)	2,116	7.4 (79)
9050	745	30.2(115)	531	18.8 (57)	463	6.9 (59)	396	3.0 (87)	443	2.0 (83)	1,430	71.6(144)	3,008	7.0 (72)
9051	784	51.3 (83)	627	20.9 (44)	554	5.8 (75)	421	14.5 (22)	521	6.0 (40)	1,488	80.2(124)	3,231	2.8 (11)
9053	939	37.3(108)	694	12.0 (93)	543	5.3 (83)	575	4.9 (80)	662	2.1 (82)	1,438	64.7(155)	3,403	14.3 (166)
9055	233	25.8(119)	55	* *	0	* *	72	* *	54	* *	0	* *	220	9.1 (107)
9058	244	62.3 (39)	190	22.6 (31)	172	15.1 (5)	149	2.7 (89)	183	1.1 (96)	635	68.2(149)	1,140	13.2 (155)
9066	843	26.7(118)	615	7.8 (104)	567	3.4 (99)	522	5.0 (78)	587	0.7(100)	1,303	76.1(134)	3,110	7.8 (86)
9068	497	35.6(109)	405	7.4 (105)	319	6.0 (72)	327	2.8 (88)	367	0.8 (98)	890	68.2(149)	1,974	13.6 (160)
9069	884	48.0 (90)	624	9.5 (101)	439	5.5 (79)	536	6.0 (75)	603	2.0 (83)	1,028	70.5(145)	2,787	11.0 (134)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
9070	706	33.1(114)	407	11.8 (94)	456	3.7 (98)	284	5.3 (77)	331	1.8 (88)	1,130	62.9(157)	2,555	8.3 (93)
9072	618	29.9(116)	336	8.9 (102)	362	1.4 (105)	245	6.5 (70)	309	1.9 (86)	962	66.1(154)	2,182	11.7 (147)
9074	597	48.2 (89)	404	12.6 (89)	311	4.2 (95)	346	6.1 (74)	386	1.6 (92)	895	66.3(152)	2,108	11.2 (139)
9080	851	51.8 (81)	592	19.6 (53)	652	11.2 (20)	482	3.1 (86)	554	2.0 (83)	1,936	64.3(156)	3,775	10.2 (127)
9151	0	* *	0	* *	1	* *	0	* *	0	* *	1	* *	2	* *
Unknown Clinics	19	* *	20	* *	24	* *	15	* *	20	* *	45	* *	108	18.5 (175)
Michigan	62,038	53.5	44,879	18.5	42,347	7.8	33,662	11.0	38,570	4.9	117,932	81.6	250,058	7.9
Nation (Prior Year)	1,719,083	62.0	848,761	26.9	1,038,203	19.1	144,421	12.9	206,416	7.2	744,697	74.8	3,277,352	10.9

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0101	132	7.6 (49)	8.3 (113)	609	4.8 (34)	3.1 (83)	12.3 (87)	270	15.2 (44)	11.9 (47)	375	9.3 (64)
0102	399	7.0 (36)	8.8 (118)	1,786	6.6 (131)	4.9 (161)	15.0 (164)	798	19.8 (169)	13.2 (77)	1,119	17.1 (145)
0103	428	6.8 (30)	11.7 (160)	1,829	6.6 (131)	2.3 (32)	15.2 (168)	808	20.4 (176)	16.3 (161)	1,188	10.6 (90)
0104	130	9.2 (106)	7.7 (91)	651	5.8 (76)	3.5 (100)	13.2 (117)	330	14.5 (31)	13.0 (73)	463	9.7 (74)
0250	166	6.0 (14)	9.6 (133)	806	4.7 (32)	3.6 (103)	13.5 (132)	379	23.5 (189)	15.6 (147)	537	13.0 (114)
0251	114	4.4 (5)	9.6 (133)	636	4.2 (21)	2.0 (19)	14.3 (151)	307	22.8 (186)	16.0 (156)	404	8.2 (50)
0252	137	8.8 (88)	10.2 (145)	728	3.7 (13)	2.1 (23)	14.1 (145)	380	20.5 (178)	17.9 (176)	508	10.4 (87)
0253	178	11.8 (150)	10.7 (154)	821	3.7 (13)	8.2 (198)	8.5 (11)	400	14.0 (23)	11.5 (41)	507	18.3 (153)
0401	553	9.9 (124)	7.8 (97)	2,430	7.6 (171)	2.9 (71)	15.5 (173)	1,101	18.0 (129)	14.5 (117)	1,576	11.8 (102)
0402	465	5.4 (8)	9.0 (121)	2,118	4.2 (21)	3.6 (103)	13.0 (108)	986	15.9 (65)	17.5 (174)	1,452	9.6 (70)
0403	123	8.9 (92)	7.3 (78)	684	6.7 (139)	4.1 (128)	17.4 (187)	347	17.3 (110)	21.3 (185)	474	6.1 (37)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0404	153	6.5 (20)	8.5 (115)	765	2.2 (1)	3.8 (117)	9.8 (25)	402	14.2 (25)	10.4 (20)	509	10.0 (79)
0501	428	6.8 (30)	10.0 (139)	2,369	4.9 (38)	3.4 (96)	12.5 (95)	1,201	17.9 (124)	13.2 (77)	1,729	4.7 (22)
0502	3	* *	* *	22	* *	* *	* *	11	* *	* *	16	* *
0503	169	13.0 (155)	10.1 (142)	743	5.0 (44)	5.4 (175)	11.6 (59)	329	13.4 (17)	10.6 (24)	496	4.2 (15)
0504	607	9.4 (112)	9.7 (135)	2,959	6.0 (97)	2.7 (58)	13.3 (122)	1,425	18.9 (157)	14.1 (104)	2,092	3.2 (7)
0505	371	7.8 (57)	7.8 (97)	1,868	5.8 (76)	3.8 (117)	13.3 (122)	937	17.8 (122)	12.3 (57)	1,354	4.1 (13)
0506	200	10.5 (132)	5.5 (29)	1,057	5.1 (49)	4.2 (135)	13.3 (122)	550	18.2 (133)	14.2 (107)	764	3.3 (8)
0650	1,077	8.3 (76)	7.5 (83)	4,604	7.6 (171)	3.0 (75)	11.6 (59)	1,928	16.6 (79)	14.7 (126)	2,898	6.9 (41)
0651	652	8.9 (92)	4.6 (18)	2,982	5.8 (76)	3.4 (96)	13.1 (113)	1,382	19.7 (168)	14.5 (117)	2,014	11.0 (95)
0652	322	7.1 (39)	9.0 (121)	1,735	6.7 (139)	3.1 (83)	14.8 (161)	894	20.9 (181)	17.1 (171)	1,200	4.8 (23)
0653	462	8.4 (80)	8.0 (102)	2,277	6.3 (113)	1.8 (10)	14.4 (154)	1,125	17.1 (99)	15.5 (144)	1,600	14.4 (125)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0654	400	9.0 (96)	11.8(162)	1,675	5.3 (56)	3.6(103)	11.0 (45)	763	15.5 (51)	14.2(107)	1,129	23.8(175)
0655	415	6.7 (26)	7.0 (64)	1,905	7.5(169)	2.5 (43)	13.2(117)	886	18.5(142)	15.5(144)	1,309	7.5 (47)
0657	72	* *	* *	475	6.1(100)	4.2(135)	12.2 (82)	257	17.1 (99)	13.6 (87)	364	16.2(138)
0701	422	5.5 (9)	10.9(156)	1,820	7.3(158)	3.0 (75)	10.5 (34)	822	17.5(116)	8.9 (7)	1,031	4.8 (23)
0702	99	* *	* *	471	7.4(164)	2.8 (65)	15.5(173)	225	16.9 (89)	18.2(178)	292	3.8 (11)
0703	264	8.0 (61)	9.8(137)	1,085	7.9(182)	2.2 (27)	10.4 (30)	486	18.5(142)	12.3 (57)	662	2.3 (3)
0704	85	* *	* *	458	7.0(147)	2.6 (48)	17.5(188)	233	15.9 (65)	16.7(165)	294	2.7 (5)
0705	189	3.7 (4)	9.0(121)	957	5.7 (74)	2.9 (71)	9.4 (17)	483	15.7 (57)	10.8 (27)	575	5.4 (28)
0850	1,212	7.7 (51)	9.5(129)	4,914	5.9 (88)	4.0(126)	11.4 (52)	2,042	18.5(142)	13.3 (82)	2,840	3.9 (12)
0851	877	6.8 (30)	8.1(107)	3,970	5.6 (65)	4.2(135)	9.4 (17)	1,796	15.8 (61)	11.1 (32)	2,414	5.6 (31)
0901	1,877	8.8 (88)	10.7(154)	8,032	7.3(158)	2.5 (43)	11.8 (70)	3,615	16.8 (85)	14.2(107)	5,378	13.0(114)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
0902	71	* *	* *	368	7.6(171)	3.0 (75)	12.5 (95)	172	20.3(172)	12.2 (54)	255	14.1(122)
1051	1,472	8.2 (68)	10.3(147)	6,507	5.9 (88)	5.3(172)	11.6 (59)	3,053	16.5 (76)	12.0 (49)	3,851	9.9 (77)
1055	0	* *	* *	1	* *	* *	* *	0	* *	* *	0	* *
1101	1,919	9.6(119)	5.8 (31)	7,979	7.0(147)	2.5 (43)	11.4 (52)	3,416	17.2(103)	11.8 (44)	5,190	22.0(172)
1102	945	7.4 (45)	8.9(120)	3,839	5.6 (65)	2.3 (32)	12.2 (82)	1,676	16.1 (70)	12.5 (63)	2,495	8.2 (50)
1106	112	6.3 (17)	6.3 (41)	481	5.6 (65)	1.0 (3)	11.6 (59)	198	12.1 (8)	16.2(160)	318	10.1 (81)
1250	1,089	6.3 (17)	8.1(107)	5,060	6.4(117)	3.6(103)	14.2(146)	2,374	17.3(110)	13.7 (92)	3,311	9.6 (70)
1251	721	8.2 (68)	7.1 (70)	3,897	6.4(117)	3.6(103)	13.0(108)	1,895	18.9(157)	16.4(162)	2,705	4.4 (17)
1252	602	8.3 (76)	6.5 (46)	3,542	6.2(107)	4.6(150)	20.0(193)	1,692	23.6(191)	26.2(190)	2,376	11.7(100)
1253	955	6.5 (20)	6.8 (53)	3,940	6.1(100)	4.8(157)	13.2(117)	1,749	16.8 (85)	16.4(162)	2,550	11.2 (96)
1350	2,691	8.9 (92)	7.9(101)	10,700	7.5(169)	3.4 (96)	13.7(136)	4,514	18.6(149)	15.0(137)	6,439	14.9(130)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
1351	169	6.5 (20)	5.9 (34)	741	6.6(131)	1.8 (10)	14.7(160)	334	15.6 (53)	14.7(126)	480	15.6(137)
1352	471	9.1(103)	7.6 (88)	2,110	7.4(164)	3.7(113)	15.4(171)	993	15.7 (57)	15.9(151)	1,320	19.8(165)
1401	244	9.4(112)	9.8(137)	1,215	5.6 (65)	4.4(143)	13.9(142)	590	14.7 (38)	16.1(158)	788	4.6 (20)
1402	176	6.8 (30)	13.6(166)	840	4.2 (21)	3.0 (75)	17.0(184)	429	20.7(180)	18.2(178)	588	8.2 (50)
1403	28	* *	* *	209	7.7(177)	5.7(180)	7.7 (8)	121	18.2(133)	5.8 (2)	151	2.6 (4)
1404	9	* *	* *	88	* *	* *	* *	44	* *	* *	75	* *
1501	1,080	7.6 (49)	7.5 (83)	5,369	5.8 (76)	3.3 (93)	16.5(183)	2,568	17.4(114)	14.9(132)	3,653	24.9(178)
1502	6	* *	* *	50	* *	* *	* *	28	* *	* *	34	* *
1503	1,342	8.2 (68)	9.0(121)	6,141	5.8 (76)	5.0(164)	13.2(117)	2,902	14.5 (31)	13.9 (99)	4,008	21.8(171)
1504	2,482	8.2 (68)	7.1 (70)	11,537	6.5(124)	3.6(103)	13.4(129)	5,298	16.6 (79)	13.9 (99)	7,442	18.5(154)
1505	933	7.4 (45)	9.5(129)	4,172	4.4 (29)	3.6(103)	10.6 (36)	2,000	12.1 (8)	9.0 (9)	2,686	18.2(151)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and >= 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
1506	657	8.2 (68)	10.2(145)	2,918	6.5(124)	4.6(150)	17.3(186)	1,325	17.7(121)	16.8(166)	1,869	18.5(154)
1507	703	6.7 (26)	11.1(158)	3,094	5.2 (52)	3.9(123)	12.1 (81)	1,402	18.3(137)	12.4 (61)	2,013	25.6(180)
1508	137	8.0 (61)	19.7(168)	814	12.5(200)	5.4(175)	23.2(197)	401	19.5(167)	23.2(187)	502	31.9(193)
1509	676	7.1 (39)	7.7 (91)	3,365	5.0 (44)	3.3 (93)	14.8(161)	1,701	18.3(137)	14.0(101)	2,288	15.4(135)
1510	182	9.3(108)	12.1(164)	830	5.4 (57)	3.1 (83)	13.7(136)	368	17.9(124)	13.0 (73)	505	8.7 (54)
1511	681	9.0 (96)	10.1(142)	2,851	4.4 (29)	4.4(143)	11.4 (52)	1,268	18.7(153)	14.6(122)	1,843	21.1(169)
1512	311	8.0 (61)	11.3(159)	1,538	6.0 (97)	2.7 (58)	13.3(122)	731	18.3(137)	14.6(122)	1,039	20.6(168)
1701	656	6.6 (25)	12.0(163)	2,758	5.8 (76)	2.7 (58)	13.0(108)	1,202	17.5(116)	14.0(101)	1,863	8.0 (49)
1712	88	* *	* *	605	8.3(189)	3.5(100)	13.1(113)	343	14.3 (26)	14.3(111)	446	15.2(133)
1950	702	7.0 (36)	7.0 (64)	3,134	6.2(107)	4.1(128)	10.7 (40)	1,433	15.6 (53)	9.1 (10)	2,059	4.4 (17)
1952	144	2.8 (1)	9.7(135)	809	4.8 (34)	3.2 (90)	12.4 (91)	391	16.6 (79)	13.3 (82)	573	7.7 (48)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
1953	332	7.5 (48)	9.3 (126)	1,632	5.5 (61)	3.7 (113)	11.6 (59)	763	16.5 (76)	10.9 (28)	1,088	5.1 (26)
1954	637	8.9 (92)	6.8 (53)	2,862	7.1 (151)	3.0 (75)	15.0 (164)	1,350	19.3 (163)	16.1 (158)	1,763	4.8 (23)
1955	144	3.5 (2)	13.9 (167)	818	6.4 (117)	2.4 (38)	14.2 (146)	406	17.5 (116)	13.8 (96)	530	5.8 (33)
1957	482	5.8 (10)	7.5 (83)	1,985	6.3 (113)	2.4 (38)	14.5 (155)	859	18.0 (129)	16.9 (168)	1,307	7.0 (42)
1959	226	8.4 (80)	10.6 (152)	998	7.9 (182)	2.6 (48)	13.8 (139)	439	18.5 (142)	12.8 (68)	643	5.3 (27)
1963	24	* *	* *	194	6.7 (139)	3.6 (103)	12.9 (106)	98	* *	* *	139	9.4 (66)
2150	726	7.2 (42)	10.1 (142)	3,217	6.9 (144)	2.2 (27)	11.8 (70)	1,524	18.6 (149)	14.5 (117)	2,119	2.1 (2)
2151	324	8.6 (86)	6.8 (53)	1,283	4.9 (38)	3.1 (83)	9.9 (26)	583	16.5 (76)	12.2 (54)	839	1.3 (1)
2251	503	7.8 (57)	10.3 (147)	2,197	7.1 (151)	2.0 (19)	12.2 (82)	1,034	17.2 (103)	13.5 (86)	1,439	4.6 (20)
2252	248	7.7 (51)	7.7 (91)	984	5.0 (44)	2.5 (43)	12.4 (91)	463	16.2 (72)	13.6 (87)	646	5.7 (32)
2551	4,351	13.4 (157)	3.9 (11)	16,656	8.9 (192)	3.6 (103)	13.4 (129)	6,834	16.6 (79)	12.7 (66)	10,467	28.2 (188)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
2552	0	* *	* *	3	* *	* *	* *	1	* *	* *	4	* *
2554	114	5.3 (7)	7.0 (64)	633	6.5(124)	3.8(117)	17.5(188)	311	12.5 (11)	17.0(169)	419	23.2(174)
2555	4,742	9.3(108)	6.7 (51)	17,974	7.0(147)	3.9(123)	12.7(101)	7,185	15.2 (44)	13.2 (77)	11,191	18.9(159)
3001	1,072	13.6(158)	3.2 (2)	4,790	6.6(131)	2.4 (38)	17.2(185)	2,422	16.8 (85)	15.9(151)	3,157	15.1(132)
3002	250	11.6(146)	6.4 (44)	1,301	5.6 (65)	2.8 (65)	15.1(166)	693	17.2(103)	14.0(101)	863	26.8(182)
3003	4	* *	* *	49	* *	* *	* *	31	* *	* *	43	* *
3004	189	5.8 (10)	4.8 (22)	1,165	4.9 (38)	1.3 (5)	22.2(195)	665	23.5(189)	23.9(189)	862	8.9 (57)
3007	36	* *	* *	222	3.2 (6)	2.3 (32)	23.9(198)	119	19.3(163)	23.5(188)	165	11.5 (97)
3009	14	* *	* *	75	* *	* *	* *	45	* *	* *	57	* *
3010	0	* *	* *	8	* *	* *	* *	6	* *	* *	8	* *
3201	611	6.2 (15)	10.6(152)	2,831	4.8 (34)	4.2(135)	11.1 (47)	1,322	16.9 (89)	12.0 (49)	1,759	8.7 (54)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
3301	3,114	8.7 (87)	7.3 (78)	15,354	6.2(107)	2.6 (48)	13.5(132)	6,781	16.9 (89)	14.3(111)	9,790	11.6 (99)
3302	260	11.5(144)	6.2 (36)	1,658	7.4(164)	2.0 (19)	14.5(155)	879	21.4(183)	15.9(151)	1,117	13.7 (119)
3303	8	* *	* *	85	* *	* *	* *	42	* *	* *	72	* *
3304	82	* *	* *	617	2.6 (4)	4.2(135)	8.6 (12)	315	13.0 (13)	8.9 (7)	444	10.4 (87)
3305	87	* *	* *	499	6.4(117)	2.2 (27)	8.4 (9)	231	16.9 (89)	6.9 (4)	346	8.7 (54)
3306	158	10.1(128)	9.5(129)	831	6.1(100)	3.2 (90)	10.7 (40)	365	19.2(161)	10.7 (26)	556	6.7 (39)
3308	48	* *	* *	270	4.1 (20)	4.8(157)	13.3(122)	119	18.5(142)	14.3(111)	177	7.3 (45)
3309	27	* *	* *	207	4.3 (26)	2.4 (38)	15.9(176)	105	18.1(132)	21.0(183)	149	5.4 (28)
3311	3	* *	* *	11	* *	* *	* *	2	* *	* *	7	* *
3550	84	* *	* *	393	5.9 (88)	2.8 (65)	21.4(194)	190	27.9(193)	21.6(186)	262	9.2 (62)
3551	36	* *	* *	139	2.2 (1)	2.2 (27)	15.8(175)	61	* *	* *	88	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
3552	58	* *	* *	260	5.8 (76)	2.7 (58)	25.0(201)	128	18.8(155)	20.3(182)	162	14.8 (128)
3701	1,048	7.7 (51)	6.6 (49)	4,514	6.5(124)	4.1(128)	11.9 (75)	2,049	19.4(165)	14.7(126)	3,093	10.9 (94)
3702	98	* *	* *	444	5.9 (88)	4.1(128)	14.2(146)	208	16.3 (73)	15.4(143)	293	9.2 (62)
3703	47	* *	* *	189	6.3(113)	4.8(157)	13.8(139)	68	* *	* *	125	7.2 (43)
3704	38	* *	* *	171	7.6(171)	3.5(100)	14.6(158)	79	* *	* *	111	11.7 (100)
3805	3,576	8.2 (68)	8.0(102)	14,141	5.5 (61)	4.6(150)	10.5 (34)	5,956	16.1 (70)	12.9 (71)	6,298	10.8 (92)
3950	2,271	9.3(108)	8.2(111)	9,125	5.8 (76)	5.3(172)	13.6(135)	4,011	17.2(103)	13.7 (92)	5,631	17.8(148)
3952	46	* *	* *	257	5.8 (76)	1.9 (15)	16.0(177)	129	20.9(181)	9.3 (14)	168	11.9(103)
3953	37	* *	* *	260	4.2 (21)	3.8(117)	12.3 (87)	130	10.8 (5)	9.2 (11)	173	16.8(141)
3954	135	8.1 (67)	6.7 (51)	674	4.0 (18)	5.5(178)	13.8(139)	297	18.9(157)	14.8(129)	439	18.5(154)
4001	1,123	12.6(152)	5.4 (27)	4,967	6.9(144)	5.3(172)	14.0(144)	2,329	15.8 (61)	15.5(144)	3,455	27.6(184)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
4002	1,001	10.5 (132)	7.2 (77)	4,381	7.4 (164)	5.0 (164)	15.1 (166)	1,985	17.5 (116)	15.0 (137)	2,983	27.2 (183)
4103	98	* *	* *	521	5.0 (44)	8.6 (199)	7.5 (6)	264	14.4 (28)	11.0 (30)	369	22.5 (173)
4106	1,078	9.5 (116)	4.7 (20)	5,546	5.9 (88)	2.6 (48)	13.5 (132)	2,755	17.1 (99)	14.3 (111)	3,877	18.7 (158)
4107	3,238	9.7 (121)	8.2 (111)	13,340	7.3 (158)	2.3 (32)	13.1 (113)	5,769	16.6 (79)	13.8 (96)	8,527	15.3 (134)
4109	1,342	10.2 (129)	10.0 (139)	5,958	5.4 (57)	3.0 (75)	9.4 (17)	2,725	14.4 (28)	11.2 (34)	4,032	7.2 (43)
4110	1,743	7.2 (42)	6.9 (60)	8,075	5.9 (88)	1.8 (10)	16.0 (177)	3,881	17.9 (124)	18.1 (177)	5,590	11.9 (103)
4112	711	3.5 (2)	11.0 (157)	3,301	8.6 (191)	3.7 (113)	22.8 (196)	1,590	20.6 (179)	26.5 (192)	2,509	9.8 (76)
4113	607	6.9 (34)	7.1 (70)	3,382	3.6 (9)	2.3 (32)	16.1 (179)	1,663	17.3 (110)	19.7 (181)	2,347	9.1 (59)
4120	2,089	8.4 (80)	9.0 (121)	9,914	7.3 (158)	1.8 (10)	15.2 (168)	4,636	17.9 (124)	14.9 (132)	6,753	14.1 (122)
4160	1,977	9.4 (112)	8.1 (107)	8,861	6.2 (107)	2.6 (48)	12.8 (102)	3,901	15.6 (53)	13.6 (87)	5,700	14.1 (122)
4351	673	9.1 (103)	11.7 (160)	3,120	6.2 (107)	3.1 (83)	11.5 (56)	1,472	16.8 (85)	13.0 (73)	2,100	12.3 (109)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
4352	205	8.8 (88)	7.3 (78)	944	7.2(156)	2.6 (48)	14.2(146)	467	16.3 (73)	14.3(111)	632	16.5(140)
4353	539	7.8 (57)	6.9 (60)	2,476	6.6(131)	2.1 (23)	12.0 (78)	1,157	18.2(133)	14.6(122)	1,693	9.6 (70)
4354	826	7.7 (51)	8.8(118)	3,603	4.7 (32)	3.0 (75)	9.2 (16)	1,673	16.3 (73)	9.6 (17)	2,380	11.5 (97)
4355	390	10.8(138)	8.5(115)	1,658	6.4(117)	2.6 (48)	11.7 (67)	759	21.6(184)	15.9(151)	1,092	5.8 (33)
4356	625	8.5 (83)	9.3(126)	2,670	6.5(124)	2.0 (19)	11.8 (70)	1,251	19.9(170)	14.9(132)	1,794	10.4 (87)
4357	17	* *	* *	49	* *	* *	* *	14	* *	* *	30	* *
4358	201	5.0 (6)	10.0(139)	1,197	7.9(182)	0.8 (1)	24.7(199)	672	25.6(192)	32.0(193)	897	4.3 (16)
4359	553	6.9 (34)	8.3(113)	2,430	9.3(196)	2.6 (48)	18.1(192)	1,169	16.9 (89)	16.8(166)	1,575	9.1 (59)
4360	189	5.8 (10)	9.5(129)	842	4.0 (18)	2.6 (48)	8.9 (15)	371	15.9 (65)	12.4 (61)	562	17.1(145)
4361	279	8.2 (68)	6.5 (46)	1,084	6.4(117)	1.9 (15)	11.4 (52)	456	23.0(187)	14.5(117)	694	10.8 (92)
4362	379	9.5(116)	7.7 (91)	1,604	5.9 (88)	2.9 (71)	9.4 (17)	702	14.1 (24)	11.8 (44)	1,086	13.3(116)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
4363	1,000	8.5 (83)	9.3 (126)	4,236	5.6 (65)	2.1 (23)	10.4 (30)	1,892	17.8(122)	13.3 (82)	2,847	16.9(142)
4401	1,422	9.0 (96)	8.6(117)	5,679	10.5(199)	3.1 (83)	12.5 (95)	2,477	17.2(103)	13.2 (77)	3,333	13.7 (119)
4501	1,042	8.3 (76)	7.8 (97)	4,762	5.8 (76)	4.1(128)	9.5 (21)	2,228	15.7 (57)	10.5 (21)	3,182	6.0 (35)
4502	0	* *	* *	1	* *	* *	* *	1	* *	* *	1	* *
4601	1,590	8.2 (68)	6.8 (53)	6,494	8.4(190)	4.5(148)	13.9(142)	2,736	17.9(124)	16.0(156)	3,533	10.0 (79)
4701	1,334	8.0 (61)	10.5(150)	5,261	5.4 (57)	3.8(117)	10.1 (27)	2,167	18.7(153)	15.0(137)	3,320	4.1 (13)
5050	1,772	10.5(132)	5.4 (27)	5,678	6.5(124)	2.7 (58)	13.1(113)	1,756	13.9 (22)	12.8 (68)	3,364	19.2(161)
5051	3,632	9.0 (96)	6.9 (60)	12,092	7.0(147)	2.3 (32)	14.3(151)	3,866	14.4 (28)	11.3 (38)	7,236	20.1(166)
5052	2,656	10.7(136)	5.8 (31)	8,557	6.4(117)	2.2 (27)	14.2(146)	2,726	16.9 (89)	14.9(132)	5,074	19.4(163)
5054	0	* *	* *	6	* *	* *	* *	2	* *	* *	4	* *
5101	1,777	9.2(106)	6.2 (36)	6,107	6.1(100)	2.1 (23)	12.2 (82)	2,476	17.2(103)	13.3 (82)	3,767	12.5(112)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
5202	637	7.8 (57)	8.0 (102)	2,409	5.6 (65)	1.8 (10)	12.5 (95)	1,025	20.3(172)	13.7 (92)	1,419	7.3 (45)
5203	214	6.5 (20)	12.6 (165)	954	5.2 (52)	0.9 (2)	14.9 (163)	479	19.4 (165)	12.3 (57)	596	4.4 (17)
5204	173	6.4 (19)	10.4 (149)	699	2.9 (5)	1.3 (5)	10.9 (44)	323	20.4 (176)	11.1 (32)	422	9.7 (74)
5205	21	* *	* *	165	9.1 (195)	1.2 (4)	7.3 (3)	85	* *	* *	111	2.7 (5)
5601	1,278	8.8 (88)	8.1 (107)	5,161	8.1 (187)	2.8 (65)	17.9 (191)	2,276	17.4 (114)	14.9 (132)	2,916	6.0 (35)
5801	2,311	7.1 (39)	7.0 (64)	8,132	7.4 (164)	2.7 (58)	11.8 (70)	2,966	17.2 (103)	13.2 (77)	4,630	9.9 (77)
5803	11	* *	* *	81	* *	* *	* *	38	* *	* *	59	* *
5804	14	* *	* *	80	* *	* *	* *	36	* *	* *	62	* *
6101	53	* *	* *	286	7.3 (158)	2.4 (38)	11.2 (48)	145	14.5 (31)	13.8 (96)	211	13.3 (116)
6102	183	7.7 (51)	7.7 (91)	941	5.2 (52)	4.6 (150)	10.3 (29)	474	19.2 (161)	12.2 (54)	620	12.1 (107)
6103	4,673	9.4 (112)	6.6 (49)	18,091	5.5 (61)	5.0 (164)	11.5 (56)	8,092	15.2 (44)	13.1 (76)	11,537	17.8 (148)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
6104	95	* *	* *	552	4.3 (26)	6.0(185)	12.5 (95)	327	12.2 (10)	12.8 (68)	381	27.8 (187)
6105	49	* *	* *	295	6.1(100)	4.1(128)	10.2 (28)	162	17.3(110)	10.5 (21)	210	19.0 (160)
6106	6	* *	* *	37	* *	* *	* *	21	* *	* *	29	* *
6301	4,997	9.9 (124)	6.5 (46)	19,655	6.1(100)	2.7 (58)	11.7 (67)	8,035	17.6(120)	15.8(150)	12,945	13.9 (121)
6312	2,795	10.7 (136)	6.8 (53)	10,960	6.2(107)	2.9 (71)	10.8 (42)	4,456	17.0 (95)	12.5 (63)	7,181	16.9 (142)
6316	1,421	7.4 (45)	8.0 (102)	5,273	4.9 (38)	2.6 (48)	9.7 (23)	2,114	15.6 (53)	12.0 (49)	3,415	10.1 (81)
6320	267	9.7 (121)	7.1 (70)	1,364	4.9 (38)	6.1(187)	10.6 (36)	669	14.8 (39)	12.0 (49)	901	9.3 (64)
6325	165	11.5 (144)	3.6 (6)	740	3.6 (9)	3.8(117)	12.4 (91)	359	20.3(172)	14.5(117)	470	9.4 (66)
6330	441	10.2 (129)	5.9 (34)	1,610	5.8 (76)	4.2(135)	12.8(102)	713	13.6 (19)	14.6(122)	972	16.4 (139)
6335	589	8.3 (76)	7.6 (88)	2,688	3.6 (9)	4.7(154)	10.6 (36)	1,184	14.9 (41)	10.9 (28)	1,737	9.5 (68)
6340	178	6.7 (26)	6.2 (36)	1,113	5.1 (49)	2.5 (43)	12.6(100)	548	17.0 (95)	14.1(104)	746	11.9 (103)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
6345	47	* *	* *	233	5.6 (65)	1.3 (5)	12.0 (78)	113	18.6(149)	10.6 (24)	146	8.9 (57)
6350	285	10.2 (129)	6.3 (41)	1,134	5.6 (65)	6.3(189)	12.9(106)	440	18.0(129)	14.8(129)	665	14.7 (127)
7304	250	11.6 (146)	7.6 (88)	1,214	4.9 (38)	3.1 (83)	16.1(179)	649	20.2(171)	21.1(184)	941	6.7 (39)
7306	2,881	11.0 (139)	6.8 (53)	11,122	5.1 (49)	3.2 (90)	12.8(102)	5,328	18.4(140)	13.6 (87)	8,296	14.8 (128)
7308	219	5.9 (13)	10.5(150)	957	6.8(142)	3.3 (93)	13.7(136)	510	15.3 (48)	15.7(149)	749	10.1 (81)
7310	33	* *	* *	93	* *	* *	* *	40	* *	* *	54	* *
7401	2,410	9.3 (108)	6.2 (36)	9,612	5.7 (74)	5.9(183)	10.8 (42)	4,208	15.1 (42)	11.4 (39)	6,487	14.6 (126)
7402	295	8.5 (83)	7.8 (97)	1,248	7.2(156)	5.0(164)	10.4 (30)	503	15.9 (65)	14.1(104)	831	12.2 (108)
7403	100	7.0 (36)	7.0 (64)	429	6.5(124)	4.4(143)	14.5(155)	189	13.2 (14)	17.5(174)	289	8.3 (53)
7404	113	6.2 (15)	6.2 (36)	442	5.2 (52)	8.1(197)	9.7 (23)	173	13.3 (16)	9.8 (18)	292	13.4 (118)
7601	789	7.7 (51)	4.3 (16)	3,447	3.9 (16)	5.1(168)	7.4 (4)	1,613	12.6 (12)	9.2 (11)	2,359	3.3 (8)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
7850	1,044	8.0 (61)	6.8 (53)	4,522	5.0 (44)	6.3(189)	11.9 (75)	2,036	15.9 (65)	15.9(151)	3,026	5.5 (30)
7851	187	8.0 (61)	7.0 (64)	985	7.1(151)	4.2(135)	13.4(129)	495	17.0 (95)	17.4(172)	690	6.4 (38)
7852	126	9.5 (116)	7.1 (70)	546	4.8 (34)	6.6(191)	12.3 (87)	268	19.0(160)	15.3(142)	380	3.7 (10)
8150	64	* *	* *	434	3.9 (16)	5.8(181)	11.8 (70)	219	14.6 (36)	11.9 (47)	291	12.4(111)
8151	2,604	9.6 (119)	8.0 (102)	10,553	5.9 (88)	4.7(154)	12.4 (91)	4,230	15.8 (61)	14.2(107)	6,414	25.1(179)
8152	544	11.2(141)	5.5 (29)	1,500	3.8 (15)	6.7(194)	11.3 (49)	412	18.4(140)	13.6 (87)	832	30.8(190)
8153	35	* *	* *	178	10.1(198)	6.2(188)	14.6(158)	89	* *	* *	116	10.3 (85)
8154	312	6.7 (26)	7.4 (82)	1,499	2.5 (3)	4.5(148)	7.4 (4)	697	15.4 (50)	10.5 (21)	901	10.3 (85)
8155	81	* *	* *	464	4.5 (31)	2.8 (65)	8.6 (12)	217	16.6 (79)	6.9 (4)	301	18.6(157)
8156	37	* *	* *	204	3.4 (8)	4.9(161)	12.3 (87)	98	* *	* *	125	9.6 (70)
8157	23	* *	* *	166	9.0(193)	3.6(103)	13.3(122)	112	10.7 (3)	15.2(140)	105	27.6(184)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
8201	2,719	9.7 (121)	7.3 (78)	8,149	6.8(142)	3.0 (75)	11.6 (59)	2,076	14.5 (31)	11.2 (34)	4,124	12.3(109)
8202	2,786	7.2 (42)	7.1 (70)	13,995	3.6 (9)	4.4(143)	11.9 (75)	6,426	15.7 (57)	12.7 (66)	8,467	9.5 (68)
8203	359	11.7(149)	7.5 (83)	997	9.8(197)	1.6 (8)	11.5 (56)	315	23.2(188)	11.7 (43)	532	17.9(150)
8250	459	10.5(132)	6.3 (41)	798	6.0 (97)	5.4(175)	6.8 (1)	198	13.6 (19)	11.6 (42)	331	26.0(181)
8255	3,406	9.0 (96)	7.5 (83)	10,723	7.8(180)	2.8 (65)	13.3(122)	3,145	20.3(172)	17.4(172)	5,590	10.1 (81)
8257	491	9.0 (96)	3.9 (11)	2,472	5.8 (76)	3.9(123)	13.0(108)	1,085	13.2 (14)	12.9 (71)	1,593	19.6(164)
8258	2,175	10.0(127)	7.1 (70)	7,138	8.0(186)	1.9 (15)	13.2(117)	2,590	18.6(149)	16.4(162)	4,367	17.0(144)
8260	447	11.2(141)	3.4 (5)	1,737	5.9 (88)	5.2(170)	8.8 (14)	800	14.3 (26)	9.5 (16)	1,119	27.7(186)
8901	1,042	14.9(164)	3.6 (6)	4,544	7.1(151)	5.8(181)	13.0(108)	2,136	14.8 (39)	11.8 (44)	2,330	37.8(199)
8902	608	12.8(153)	5.1 (24)	2,660	6.6(131)	7.8(196)	11.7 (67)	1,226	11.3 (6)	11.2 (34)	1,589	31.1(192)
8903	1,265	14.3(162)	3.7 (8)	5,231	7.1(151)	5.1(168)	12.2 (82)	2,391	15.1 (42)	11.0 (30)	2,818	37.7(198)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
8904	923	12.9 (154)	4.8 (22)	4,035	5.5 (61)	6.6(191)	12.0 (78)	1,914	13.6 (19)	12.5 (63)	2,483	33.5 (195)
8905	2,629	9.0 (96)	6.9 (60)	10,790	6.6(131)	7.7(195)	15.2(168)	4,926	17.1 (99)	15.2(140)	3,173	34.1 (197)
9004	140	20.0 (168)	2.9 (1)	972	9.0(193)	3.4 (96)	24.7(199)	406	18.5(142)	26.4(191)	683	33.5 (195)
9010	12	* *	* *	40	* *	* *	* *	14	* *	* *	26	* *
9012	7	* *	* *	21	* *	* *	* *	8	* *	* *	4	* *
9013	32	* *	* *	81	* *	* *	* *	35	* *	* *	44	* *
9015	8	* *	* *	20	* *	* *	* *	7	* *	* *	6	* *
9016	27	* *	* *	90	* *	* *	* *	37	* *	* *	47	* *
9017	10	* *	* *	31	* *	* *	* *	13	* *	* *	4	* *
9018	38	* *	* *	78	* *	* *	* *	29	* *	* *	7	* *
9019	11	* *	* *	23	* *	* *	* *	12	* *	* *	2	* *

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
9021	13	* *	* *	25	* *	* *	* *	3	* *	* *	4	* *
9023	6	* *	* *	49	* *	* *	* *	26	* *	* *	31	* *
9040	78	* *	* *	753	4.2 (21)	4.8(157)	11.0 (45)	401	14.5 (31)	9.2 (11)	557	33.0(194)
9041	640	14.7(163)	4.2 (13)	3,395	14.4(202)	5.9(183)	16.4(182)	1,588	18.5(142)	14.3(111)	1,166	24.7(177)
9045	1,256	11.0(139)	6.4 (44)	5,893	6.1(100)	4.9(161)	17.8(190)	2,820	18.2(133)	17.0(169)	1,288	21.2(170)
9050	2,021	11.4(143)	5.8 (31)	8,608	7.6(171)	4.0(126)	15.4(171)	3,866	15.5 (51)	15.6(147)	4,987	15.5(136)
9051	2,352	6.5 (20)	7.7 (91)	9,897	5.4 (57)	1.9 (15)	16.3(181)	4,174	18.8(155)	19.5(180)	4,189	10.6 (90)
9053	2,808	13.9(159)	4.2 (13)	10,550	7.6(171)	4.7(154)	11.6 (59)	4,297	15.2 (44)	12.0 (49)	6,911	19.2(161)
9054	51	* *	* *	258	6.6(131)	6.6(191)	12.8(102)	105	21.9(185)	11.4 (39)	151	24.5(176)
9055	455	9.9(124)	4.6 (18)	492	6.9(144)	12.8(202)	6.9 (2)	36	* *	* *	44	* *
9058	552	9.1(103)	4.5 (17)	2,968	3.2 (6)	5.2(170)	9.6 (22)	1,614	11.6 (7)	10.1 (19)	2,083	20.1(166)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
9066	2,402	13.9 (159)	3.8 (10)	8,809	7.7(177)	6.0(185)	10.4 (30)	3,448	10.7 (3)	8.4 (6)	4,474	12.0(106)
9068	1,421	13.9 (159)	4.2 (13)	5,706	6.3(113)	11.9(201)	7.5 (6)	2,384	5.2 (1)	4.4 (1)	3,541	30.8(190)
9069	2,105	13.0 (155)	4.7 (20)	7,382	7.3(158)	5.6(179)	11.3 (49)	2,669	17.0 (95)	13.7 (92)	4,066	12.9(113)
9070	1,955	12.0 (151)	3.3 (3)	7,552	7.7(177)	9.2(200)	10.6 (36)	2,949	6.4 (2)	5.8 (2)	4,352	15.0(131)
9072	1,641	15.1 (165)	3.7 (8)	6,137	7.8(180)	4.1(128)	11.3 (49)	2,508	15.3 (48)	11.2 (34)	3,482	9.1 (59)
9074	1,597	15.1 (165)	3.3 (3)	5,779	8.1(187)	3.7(113)	11.6 (59)	2,323	15.8 (61)	12.3 (57)	3,556	18.2(151)
9079	405	17.0 (167)	5.2 (25)	1,845	7.9(182)	4.4(143)	14.3(151)	742	14.6 (36)	14.8(129)	1,009	17.4(147)
9080	2,272	11.6 (146)	5.2 (25)	10,600	4.3 (26)	4.3(142)	8.4 (9)	5,160	13.5 (18)	9.3 (14)	7,047	28.7(189)
9151	2	* *	* *	33	* *	* *	* *	16	* *	* *	31	* *
Unknown Clinics	33	* *	* *	230	12.6(201)	1.7 (9)	27.0(202)	38	* *	* *	452	59.7(200)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
Michigan	174,742	9.5	7.1	726,369	6.4	3.8	12.8	315,917	16.5	13.8	449,245	15.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
0101	214	64.5 (39)	165	20.6 (35)	123	8.1 (124)	81	* *	75	* *	91	* *	189	11.6 (148)
0102	651	49.2(131)	483	11.4 (144)	391	7.4 (135)	163	11.0 (99)	152	0.7(133)	253	83.8 (92)	529	11.0 (136)
0103	657	52.1(119)	491	13.6 (126)	392	7.1 (140)	154	12.3 (88)	144	2.1(125)	275	81.8(115)	558	11.3 (144)
0104	193	49.2(131)	157	4.5 (179)	137	5.1 (161)	64	* *	58	* *	96	* *	179	24.0 (182)
0250	271	59.4 (77)	225	16.4 (90)	183	11.5 (50)	86	* *	85	* *	113	90.3 (29)	237	8.4 (97)
0251	210	44.8(152)	155	13.5 (127)	122	9.8 (86)	51	* *	50	* *	96	* *	193	5.7 (44)
0252	223	68.2 (20)	181	22.7 (27)	169	12.4 (42)	77	* *	75	* *	133	89.5 (38)	230	11.3 (144)
0253	291	50.2(127)	221	18.6 (55)	179	12.3 (43)	108	20.4 (27)	103	12.6 (4)	120	86.7 (66)	247	18.2 (175)
0401	921	45.9(147)	731	16.7 (86)	608	11.3 (52)	343	16.3 (56)	323	8.0 (35)	361	95.6 (7)	761	8.3 (90)
0402	731	63.7 (46)	564	20.4 (37)	463	12.3 (43)	192	19.3 (35)	171	5.8 (66)	338	89.1 (42)	648	7.3 (76)
0403	207	64.3 (41)	168	17.9 (65)	152	13.2 (33)	76	* *	69	* *	111	83.8 (92)	213	9.9 (125)
0404	227	59.0 (81)	185	19.5 (44)	147	10.9 (59)	51	* *	39	* *	126	90.5 (25)	228	21.1 (180)
0501	789	64.4 (40)	651	16.0 (96)	544	9.0 (105)	280	19.6 (31)	263	6.1 (62)	402	89.1 (42)	780	10.0 (127)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
0502	6	* *	7	* *	7	* *	3	* *	3	* *	4	* *	5	* *
0503	255	74.5 (10)	200	24.0 (23)	164	17.1 (18)	83	* *	67	* *	122	91.8 (16)	247	3.6 (18)
0504	968	68.4 (18)	725	19.2 (47)	635	12.9 (38)	208	19.7 (30)	184	8.2 (34)	436	91.3 (20)	847	5.4 (36)
0505	627	69.5 (15)	477	26.4 (14)	410	15.1 (26)	211	31.8 (3)	199	11.1 (11)	293	87.7 (60)	569	8.6 (100)
0506	334	65.6 (31)	257	15.2 (104)	240	9.6 (93)	93	* *	87	* *	169	90.5 (25)	304	20.7 (178)
0650	1,778	65.2 (35)	1,388	19.4 (46)	1,088	10.8 (61)	545	19.1 (37)	520	8.3 (33)	783	85.2 (80)	1,668	9.7 (118)
0651	1,036	49.1(133)	831	10.3 (156)	697	7.0 (141)	382	12.3 (88)	349	2.6(117)	538	86.8 (65)	1,053	16.6 (172)
0652	527	52.9(113)	417	18.7 (52)	366	11.2 (53)	165	12.1 (91)	141	4.3 (91)	341	82.7(106)	588	16.5 (170)
0653	733	58.9 (82)	576	13.7 (123)	513	10.3 (72)	186	17.7 (49)	177	5.6 (72)	422	83.6 (97)	774	21.2 (181)
0654	625	62.7 (48)	493	15.0 (108)	376	6.9 (142)	237	17.3 (51)	219	7.8 (39)	302	87.7 (60)	603	13.4 (159)
0655	665	61.7 (56)	542	15.9 (97)	457	9.2 (100)	232	15.9 (57)	223	5.8 (66)	313	88.5 (50)	621	12.9 (157)
0657	118	66.1 (26)	104	17.3 (78)	101	8.9 (107)	44	* *	42	* *	93	* *	150	16.7 (173)
0701	626	76.4 (6)	468	36.1 (3)	350	24.3 (4)	209	35.9 (1)	188	17.0 (1)	271	97.8 (3)	546	1.5 (7)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
0702	151	57.0 (95)	129	17.1 (81)	95	* *	58	* *	57	* *	75	* *	140	6.4 (61)
0703	401	60.3 (71)	295	17.6 (72)	209	10.5 (66)	136	21.3 (23)	132	6.8 (55)	174	93.1 (11)	336	8.3 (90)
0704	133	65.4 (33)	102	16.7 (86)	95	* *	40	* *	39	* *	75	* *	133	16.5 (170)
0705	307	66.1 (26)	249	30.9 (8)	185	16.8 (19)	114	28.9 (5)	103	11.7 (8)	152	98.0 (2)	298	2.3 (10)
0850	1,806	64.7 (38)	1,376	19.9 (42)	999	9.7 (89)	785	18.0 (46)	731	6.6 (58)	942	85.8 (73)	1,907	5.5 (38)
0851	1,358	54.4(104)	1,132	15.5 (101)	835	10.1 (80)	623	18.0 (46)	593	6.1 (62)	851	86.1 (71)	1,587	6.1 (54)
0901	2,830	52.0(122)	2,213	11.6 (140)	1,731	5.5 (157)	893	10.8 (100)	795	3.3(107)	1,226	90.0 (33)	2,577	10.3 (131)
0902	119	70.6 (13)	93	* *	70	* *	21	* *	18	* *	59	* *	116	11.2 (140)
1051	2,263	78.7 (4)	1,770	31.4 (7)	1,450	22.3 (6)	685	33.3 (2)	659	11.8 (6)	934	88.3 (51)	1,959	5.1 (28)
1055	0	* *	1	* *	0	* *	0	* *	0	* *	0	* *	0	* *
1101	3,013	43.8(155)	2,394	10.3 (156)	1,921	6.8 (143)	925	10.8 (100)	887	3.6(101)	1,350	72.3(142)	2,921	10.8 (135)
1102	1,434	51.8(123)	1,154	18.6 (55)	924	12.3 (43)	520	18.7 (40)	508	7.3 (46)	721	82.0(113)	1,440	11.9 (151)
1106	182	58.8 (83)	153	18.3 (59)	127	15.0 (27)	62	* *	59	* *	89	* *	180	13.9 (164)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
1250	1,696	59.1 (79)	1,262	18.0 (62)	1,012	10.4 (69)	723	18.4 (44)	630	7.5 (44)	981	83.3(102)	1,855	9.8 (119)
1251	1,261	61.2 (64)	1,056	18.4 (58)	827	9.1 (102)	553	26.8 (8)	512	13.1 (3)	811	86.9 (62)	1,482	8.6 (100)
1252	1,041	59.6 (74)	841	18.0 (62)	762	12.1 (48)	292	12.0 (93)	269	6.7 (56)	537	80.4(123)	899	6.9 (69)
1253	1,556	54.4(104)	1,064	11.6 (140)	872	5.0 (162)	578	12.1 (91)	502	5.2 (80)	842	83.7 (95)	1,760	8.9 (103)
1350	3,760	49.9(128)	2,737	10.7 (148)	2,069	4.9 (163)	1,509	8.3 (110)	1,359	3.5(104)	2,017	87.8 (59)	3,953	6.6 (64)
1351	284	64.1 (42)	200	17.5 (75)	152	11.2 (53)	138	18.8 (38)	120	5.8 (66)	164	98.2 (1)	323	4.6 (25)
1352	689	47.9(138)	519	8.5 (167)	410	3.4 (173)	315	7.9 (113)	281	2.5(120)	414	83.8 (92)	754	13.5 (160)
1401	403	67.7 (21)	305	23.3 (26)	262	10.7 (63)	143	25.2 (11)	128	11.7 (8)	191	84.8 (85)	377	6.4 (61)
1402	264	78.0 (5)	180	28.3 (11)	142	22.5 (5)	40	* *	40	* *	135	89.6 (37)	267	7.1 (72)
1403	55	* *	46	* *	39	* *	21	* *	18	* *	37	* *	56	* *
1404	31	* *	29	* *	23	* *	12	* *	12	* *	17	* *	35	* *
1501	1,783	61.1 (66)	1,368	20.1 (40)	1,157	10.4 (69)	578	15.7 (61)	515	7.4 (45)	1,078	84.8 (85)	1,983	8.0 (85)
1502	6	* *	7	* *	6	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
1503	2,138	61.8 (55)	1,659	19.2 (47)	1,327	12.7 (39)	697	15.8 (59)	614	8.8 (24)	1,199	82.2(112)	2,251	8.3 (90)
1504	4,070	62.4 (52)	2,928	17.6 (72)	2,652	10.5 (66)	1,173	21.6 (22)	1,106	7.2 (48)	2,195	84.5 (87)	4,263	2.8 (13)
1505	1,414	66.5 (24)	1,029	25.6 (17)	908	13.0 (37)	427	27.9 (7)	407	14.3 (2)	899	89.9 (35)	1,631	2.9 (14)
1506	1,049	61.2 (64)	775	16.6 (89)	654	8.9 (107)	356	18.8 (38)	328	7.6 (42)	554	88.1 (54)	1,086	6.8 (68)
1507	1,141	62.6 (49)	812	20.7 (33)	749	9.9 (82)	408	28.7 (6)	382	12.0 (5)	592	89.7 (36)	1,170	6.3 (58)
1508	261	60.5 (70)	218	32.1 (5)	190	13.2 (33)	75	* *	72	* *	149	90.6 (24)	256	0.0 (1)
1509	1,093	57.3 (93)	857	16.8 (85)	723	9.7 (89)	421	18.1 (45)	373	8.0 (35)	692	83.1(103)	1,272	6.7 (66)
1510	283	49.8(129)	218	14.2 (118)	177	10.7 (63)	77	* *	70	* *	83	* *	172	11.0 (136)
1511	1,060	61.4 (61)	792	16.2 (94)	657	9.1 (102)	440	17.0 (52)	403	5.2 (80)	562	83.6 (97)	1,155	5.2 (31)
1512	493	58.4 (86)	407	14.7 (115)	360	7.8 (131)	201	19.9 (29)	182	7.1 (49)	313	88.2 (52)	537	5.6 (42)
1701	1,011	61.3 (62)	747	20.5 (36)	625	12.6 (40)	281	19.6 (31)	252	7.1 (49)	395	88.9 (46)	869	8.1 (87)
1712	145	55.2(100)	128	17.2 (80)	107	11.2 (53)	48	* *	47	* *	89	* *	154	11.0 (136)
1950	1,128	57.4 (92)	856	15.0 (108)	675	8.9 (107)	385	16.9 (54)	360	8.6 (28)	546	86.3 (69)	1,108	11.2 (140)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
1952	270	65.6 (31)	210	20.0 (41)	172	8.1 (124)	98	* *	91	* *	143	89.5 (38)	270	9.3 (113)
1953	540	57.6 (90)	435	17.5 (75)	346	9.2 (100)	180	16.7 (55)	172	4.7 (88)	283	86.9 (62)	529	8.9 (103)
1954	1,034	61.7 (56)	788	20.7 (33)	604	13.2 (33)	343	21.9 (16)	317	7.6 (42)	546	94.3 (9)	1,058	6.2 (56)
1955	250	59.2 (78)	219	17.8 (68)	187	8.0 (127)	85	* *	78	* *	164	89.0 (44)	287	6.3 (58)
1957	762	62.6 (49)	605	20.2 (38)	435	14.7 (29)	294	21.8 (18)	278	7.9 (38)	341	86.2 (70)	734	5.4 (36)
1959	370	67.6 (22)	323	19.2 (47)	251	10.4 (69)	123	17.9 (48)	118	8.5 (30)	157	87.9 (57)	310	5.8 (46)
1963	10	* *	19	* *	22	* *	0	* *	0	* *	0	* *	0	* *
2150	1,125	57.5 (91)	832	18.0 (62)	682	9.8 (86)	290	12.4 (87)	270	4.4 (89)	485	93.0 (12)	994	6.0 (50)
2151	479	47.0(143)	327	10.4 (154)	284	6.7 (145)	168	13.7 (78)	150	3.3(107)	158	88.0 (55)	355	5.1 (28)
2251	735	60.1 (72)	593	21.6 (32)	489	10.2 (76)	257	23.7 (15)	242	8.7 (26)	320	90.9 (23)	629	5.1 (28)
2252	375	58.7 (84)	251	16.3 (93)	190	7.9 (129)	137	26.3 (9)	126	9.5 (19)	133	91.7 (18)	309	8.1 (87)
2551	6,551	34.0(171)	4,889	5.3 (177)	3,586	3.3 (174)	2,157	5.3 (129)	2,009	1.6(131)	2,258	76.7(133)	5,193	8.4 (97)
2552	0	* *	1	* *	2	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
2554	205	31.2(174)	154	5.2 (178)	144	4.2 (168)	67	* *	66	* *	107	86.9 (62)	198	11.6 (148)
2555	7,236	47.7(140)	5,299	10.3 (156)	3,897	6.2 (149)	2,104	9.6 (107)	1,900	3.6(101)	2,588	84.9 (84)	5,951	5.9 (48)
3001	1,645	43.2(159)	1,255	7.5 (173)	962	4.5 (167)	633	7.0 (119)	573	3.7 (98)	801	75.2(136)	1,545	13.3 (158)
3002	398	34.2(170)	318	2.8 (180)	244	2.0 (176)	158	3.2 (137)	148	0.0(135)	237	81.9(114)	428	7.7 (80)
3003	10	* *	11	* *	11	* *	6	* *	5	* *	12	* *	20	* *
3004	288	66.0 (28)	243	20.2 (38)	211	15.6 (24)	75	* *	70	* *	197	73.6(139)	329	1.5 (7)
3007	60	* *	47	* *	33	* *	22	* *	19	* *	37	* *	66	* *
3009	21	* *	12	* *	13	* *	6	* *	4	* *	12	* *	23	* *
3010	3	* *	3	* *	2	* *	1	* *	1	* *	3	* *	4	* *
3201	952	53.2(109)	722	12.9 (132)	555	9.7 (89)	357	18.5 (43)	319	5.3 (77)	451	92.5 (13)	901	7.5 (79)
3301	5,298	60.9 (67)	4,316	18.7 (52)	3,173	11.4 (51)	2,267	12.3 (88)	2,081	5.3 (77)	2,951	79.8(127)	6,211	6.0 (50)
3302	449	52.1(119)	369	12.7 (134)	303	8.9 (107)	191	9.9 (104)	183	4.4 (89)	303	71.9(143)	584	6.0 (50)
3303	4	* *	10	* *	9	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
3304	176	92.6 (1)	153	62.7 (1)	138	49.3 (1)	82	* *	85	* *	135	78.5(131)	235	1.7 (9)
3305	182	80.2 (3)	140	27.9 (12)	119	16.8 (19)	77	* *	74	* *	106	85.8 (73)	209	2.4 (11)
3306	337	67.1 (23)	271	15.1 (106)	213	9.9 (82)	151	11.9 (94)	137	5.8 (66)	198	92.4 (14)	414	7.7 (80)
3308	96	* *	82	* *	51	* *	50	* *	45	* *	53	* *	109	9.2 (111)
3309	66	* *	56	* *	47	* *	26	* *	23	* *	53	* *	85	* *
3311	1	* *	1	* *	0	* *	0	* *	0	* *	0	* *	0	* *
3550	139	48.2(137)	106	6.6 (174)	101	4.0 (172)	60	* *	57	* *	74	* *	135	5.2 (31)
3551	55	* *	39	* *	38	* *	19	* *	19	* *	23	* *	41	* *
3552	84	* *	72	* *	59	* *	28	* *	26	* *	36	* *	66	* *
3701	1,756	51.3(125)	1,285	13.3 (130)	1,082	8.1 (124)	583	18.7 (40)	561	10.5 (13)	921	96.2 (5)	1,834	9.2 (111)
3702	81	* *	99	* *	82	* *	0	* *	0	* *	0	* *	0	* *
3703	41	* *	49	* *	39	* *	0	* *	0	* *	0	* *	0	* *
3704	35	* *	44	* *	38	* *	0	* *	0	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
3805	5,095	54.2(107)	4,027	13.9 (120)	2,985	6.6 (147)	2,302	13.2 (83)	2,161	5.5 (75)	2,754	82.6(108)	5,417	8.5 (99)
3950	3,415	57.7 (87)	2,600	18.2 (60)	1,968	10.2 (76)	1,209	18.6 (42)	1,124	9.5 (19)	1,492	82.6(108)	3,131	5.8 (46)
3952	76	* *	63	* *	47	* *	25	* *	25	* *	38	* *	61	* *
3953	67	* *	67	* *	55	* *	25	* *	22	* *	43	* *	75	* *
3954	219	52.1(119)	193	15.0 (108)	165	10.3 (72)	78	* *	70	* *	114	90.4 (27)	192	1.0 (4)
4001	1,766	56.3 (96)	1,314	13.9 (120)	1,052	8.4 (118)	537	12.7 (86)	451	5.3 (77)	941	59.4(161)	1,830	13.8 (163)
4002	1,612	58.6 (85)	1,226	17.5 (75)	990	10.2 (76)	604	17.5 (50)	535	7.3 (46)	854	66.0(154)	1,671	9.3 (113)
4103	200	43.5(158)	153	13.1 (131)	134	6.7 (145)	81	* *	78	* *	122	67.2(151)	222	9.0 (106)
4106	1,668	53.2(109)	1,387	15.6 (100)	1,125	8.4 (118)	672	9.5 (108)	602	5.0 (83)	1,012	80.5(122)	1,889	6.2 (56)
4107	5,038	60.9 (67)	3,984	19.5 (44)	2,887	10.8 (61)	1,958	14.7 (69)	1,810	7.8 (39)	2,333	85.2 (80)	5,004	5.3 (34)
4109	2,042	60.9 (67)	1,702	16.4 (90)	1,339	8.8 (114)	697	15.9 (57)	625	8.8 (24)	1,187	88.2 (52)	2,258	6.5 (63)
4110	2,696	56.0 (97)	2,149	17.3 (78)	1,799	11.0 (57)	889	6.4 (124)	821	4.3 (91)	1,581	86.5 (67)	2,966	5.5 (38)
4112	1,088	73.2 (11)	833	25.8 (16)	731	15.7 (23)	380	4.2 (133)	350	2.6(117)	671	94.6 (8)	1,177	4.2 (22)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
4113	1,070	71.8 (12)	902	24.1 (22)	715	15.2 (25)	265	5.7 (127)	201	2.0(126)	607	84.3 (88)	1,172	1.3 (5)
4120	3,426	61.6 (59)	2,878	17.9 (65)	2,232	9.9 (82)	1,329	14.1 (74)	1,230	6.7 (56)	1,928	86.1 (71)	3,654	3.5 (17)
4160	3,132	59.6 (74)	2,659	18.2 (60)	1,989	10.9 (59)	1,044	13.0 (84)	964	7.0 (52)	1,547	83.6 (97)	3,079	4.5 (24)
4351	1,027	63.8 (44)	772	17.6 (72)	662	8.2 (122)	300	26.3 (9)	286	8.7 (26)	520	85.8 (73)	968	11.6 (148)
4352	339	52.2(118)	240	13.8 (122)	210	6.2 (149)	126	14.3 (73)	122	5.7 (70)	159	78.6(130)	316	19.9 (177)
4353	876	55.1(101)	661	17.9 (65)	557	9.3 (98)	278	15.8 (59)	252	9.5 (19)	407	75.9(135)	795	13.7 (161)
4354	1,291	61.3 (62)	940	17.1 (81)	786	9.7 (89)	456	21.9 (16)	408	11.8 (6)	564	91.8 (16)	1,179	9.8 (119)
4355	589	61.6 (59)	437	13.7 (123)	390	8.2 (122)	155	19.4 (34)	144	5.6 (72)	247	90.3 (29)	505	7.9 (83)
4356	934	53.2(109)	676	14.8 (114)	597	7.7 (133)	269	20.4 (27)	253	9.1 (23)	430	85.8 (73)	864	9.3 (113)
4357	24	* *	14	* *	13	* *	2	* *	2	* *	2	* *	7	* *
4358	346	75.7 (8)	263	31.6 (6)	243	19.8 (12)	128	13.3 (81)	121	4.1 (93)	220	85.0 (82)	384	1.3 (5)
4359	924	61.7 (56)	624	17.8 (68)	602	10.1 (80)	314	15.3 (66)	292	6.2 (61)	480	82.7(106)	869	9.0 (106)
4360	295	68.5 (17)	248	14.9 (111)	211	9.5 (96)	101	21.8 (18)	91	* *	134	88.8 (47)	259	12.7 (155)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
4361	398	64.8 (37)	277	12.6 (135)	238	8.4 (118)	77	* *	70	* *	143	90.2 (31)	333	10.2 (129)
4362	564	74.6 (9)	396	23.5 (25)	326	15.0 (27)	93	* *	88	* *	240	90.4 (27)	519	15.8 (168)
4363	1,558	70.2 (14)	1,166	16.9 (83)	918	9.3 (98)	540	23.9 (14)	501	8.4 (32)	664	90.2 (31)	1,330	9.8 (119)
4401	2,311	57.7 (87)	1,590	16.9 (83)	1,163	9.9 (82)	837	15.5 (64)	754	6.4 (59)	891	90.0 (33)	1,909	8.9 (103)
4501	1,643	60.1 (72)	1,203	15.1 (106)	986	8.9 (107)	444	21.8 (18)	423	9.7 (16)	729	83.7 (95)	1,482	9.4 (117)
4601	2,578	55.4 (99)	1,979	15.8 (98)	1,524	8.0 (127)	925	15.7 (61)	843	3.8 (97)	1,046	85.5 (78)	2,244	5.6 (42)
4701	2,128	66.4 (25)	1,409	18.5 (57)	1,218	10.3 (72)	647	21.3 (23)	613	7.7 (41)	852	88.8 (47)	1,843	4.9 (27)
5050	2,703	48.8(136)	1,823	13.5 (127)	1,456	6.4 (148)	844	14.5 (72)	749	5.6 (72)	716	81.8(115)	1,946	6.0 (50)
5051	5,337	49.1(133)	3,784	14.2 (118)	3,058	7.3 (137)	1,528	13.6 (79)	1,324	5.4 (76)	1,343	81.8(115)	3,603	5.9 (48)
5052	4,063	43.6(156)	2,825	10.7 (148)	2,225	5.9 (153)	1,291	9.9 (104)	1,151	3.4(105)	1,072	80.1(125)	2,957	6.7 (66)
5054	0	* *	1	* *	1	* *	0	* *	0	* *	0	* *	0	* *
5101	2,945	57.2 (94)	2,035	17.7 (70)	1,447	10.2 (76)	1,191	14.1 (74)	1,090	5.1 (82)	1,315	83.4(101)	3,097	5.5 (38)
5202	986	63.8 (44)	708	23.6 (24)	570	14.0 (32)	340	21.8 (18)	314	9.6 (18)	378	91.0 (21)	806	5.5 (38)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
5203	333	62.5 (51)	236	26.3 (15)	198	18.2 (16)	119	29.4 (4)	111	9.9 (15)	145	89.0 (44)	284	7.0 (70)
5204	245	68.6 (16)	185	27.0 (13)	140	21.4 (7)	83	* *	74	* *	104	94.2 (10)	202	4.0 (21)
5205	9	* *	20	* *	22	* *	0	* *	0	* *	0	* *	0	* *
5601	1,976	65.7 (30)	1,434	15.3 (103)	1,143	9.0 (105)	604	24.5 (12)	586	9.2 (22)	848	89.3 (40)	1,766	5.2 (31)
5801	3,617	52.9(113)	2,855	13.7 (123)	2,077	7.3 (137)	1,306	14.1 (74)	1,199	6.3 (60)	1,435	82.5(111)	3,189	10.5 (133)
5803	38	* *	32	* *	25	* *	23	* *	21	* *	32	* *	69	* *
5804	50	* *	39	* *	27	* *	39	* *	34	* *	40	* *	81	* *
6101	120	41.7(162)	98	* *	76	* *	55	* *	50	* *	77	* *	139	15.1 (167)
6102	376	64.1 (42)	256	22.7 (27)	224	10.7 (63)	139	24.5 (12)	130	10.0 (14)	236	91.9 (15)	445	8.3 (90)
6103	6,694	47.0(143)	4,714	11.3 (145)	4,211	6.8 (143)	2,148	11.9 (94)	1,939	4.0 (95)	3,029	82.6(108)	6,261	8.7 (102)
6104	173	31.2(174)	124	5.6 (176)	118	4.2 (168)	89	* *	84	* *	159	73.6(139)	276	15.9 (169)
6105	110	59.1 (79)	82	* *	77	* *	47	* *	43	* *	88	* *	143	9.8 (119)
6106	19	* *	13	* *	12	* *	14	* *	12	* *	21	* *	39	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
6301	7,728	45.9(147)	5,797	14.9 (111)	4,219	8.9 (107)	2,688	14.7 (69)	2,473	5.9 (65)	2,910	83.0(105)	6,517	4.7 (26)
6312	4,324	54.9(103)	3,300	19.7 (43)	2,315	11.2 (53)	1,524	19.2 (36)	1,396	8.5 (30)	1,536	79.4(129)	3,569	3.9 (19)
6316	2,226	51.8(123)	1,660	15.7 (99)	1,112	9.6 (93)	777	14.9 (68)	720	6.0 (64)	823	81.7(118)	1,951	3.3 (16)
6320	447	52.6(116)	360	22.5 (30)	265	20.0 (10)	168	15.5 (64)	154	9.7 (16)	225	88.0 (55)	429	0.9 (2)
6325	274	59.5 (76)	201	13.4 (129)	143	10.5 (66)	116	20.7 (25)	100	7.0 (52)	103	86.4 (68)	236	8.1 (87)
6330	675	43.9(154)	460	11.3 (145)	327	7.3 (137)	264	9.8 (106)	230	1.7(130)	277	74.7(137)	620	9.0 (106)
6335	994	46.0(146)	762	15.4 (102)	571	12.3 (43)	390	15.6 (63)	349	8.6 (28)	456	87.9 (57)	923	3.9 (19)
6340	275	47.3(141)	243	17.7 (70)	198	19.2 (14)	82	* *	74	* *	143	91.6 (19)	255	3.1 (15)
6345	79	* *	62	* *	39	* *	23	* *	21	* *	31	* *	66	* *
6350	486	42.2(161)	361	11.9 (138)	240	8.3 (121)	188	13.8 (77)	169	4.1 (93)	180	85.6 (77)	406	4.4 (23)
7304	393	47.3(141)	285	8.1 (170)	244	5.3 (160)	123	14.6 (71)	112	7.1 (49)	210	84.3 (88)	399	8.3 (90)
7306	4,294	45.0(149)	2,729	8.0 (171)	2,393	4.1 (170)	1,221	6.9 (121)	1,056	3.1(110)	1,805	81.3(119)	3,704	9.0 (106)
7308	336	57.7 (87)	196	12.8 (133)	184	4.9 (163)	82	* *	70	* *	167	89.2 (41)	308	11.0 (136)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)**

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
7310	51	* *	28	* *	21	* *	18	* *	15	* *	24	* *	64	* *
7401	3,603	45.0(149)	2,784	10.6 (151)	2,244	5.4 (158)	1,119	11.3 (98)	1,085	4.8 (86)	1,540	84.2 (91)	3,208	11.3 (144)
7402	449	49.7(130)	374	12.3 (137)	296	7.4 (135)	103	13.6 (79)	98	* *	133	91.0 (21)	284	6.3 (58)
7403	141	50.4(126)	125	10.4 (154)	101	2.0 (176)	33	* *	33	* *	53	* *	101	9.9 (125)
7404	158	44.9(151)	136	10.3 (156)	123	4.9 (163)	40	* *	40	* *	50	* *	103	7.8 (82)
7601	1,203	55.1(101)	915	11.1 (147)	690	6.1 (151)	372	15.1 (67)	350	3.7 (98)	510	84.3 (88)	1,083	9.8 (119)
7850	1,598	55.9 (98)	1,252	11.9 (138)	1,073	7.8 (131)	482	10.4 (102)	468	3.0(112)	778	88.7 (49)	1,545	9.8 (119)
7851	298	65.4 (33)	243	18.9 (51)	210	11.0 (57)	80	* *	77	* *	143	96.5 (4)	263	9.1 (110)
7852	191	66.0 (28)	145	14.5 (116)	123	9.8 (86)	55	* *	54	* *	90	* *	179	11.2 (140)
8150	136	68.4 (18)	116	25.0 (19)	103	19.4 (13)	38	* *	33	* *	94	* *	143	7.0 (70)
8151	4,538	62.4 (52)	3,327	24.9 (20)	2,339	16.3 (21)	1,574	20.7 (25)	1,405	11.7 (8)	1,851	78.1(132)	4,122	10.1 (128)
8152	737	54.3(106)	527	8.0 (171)	391	4.1 (170)	214	9.3 (109)	197	3.6(101)	170	83.5(100)	510	21.0 (179)
8153	58	* *	43	* *	31	* *	17	* *	10	* *	41	* *	63	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
8154	502	87.8 (2)	408	57.4 (2)	322	35.7 (2)	72	* *	61	* *	239	96.2 (5)	457	0.9 (2)
8155	155	76.1 (7)	127	34.6 (4)	103	32.0 (3)	33	* *	32	* *	82	* *	139	13.7 (161)
8156	78	* *	52	* *	46	* *	19	* *	17	* *	33	* *	69	* *
8157	35	* *	22	* *	22	* *	10	* *	7	* *	31	* *	48	* *
8201	4,229	52.9(113)	3,074	14.5 (116)	2,154	8.7 (115)	1,491	13.0 (84)	1,363	4.8 (86)	769	85.4 (79)	2,764	5.3 (34)
8202	4,900	65.0 (36)	3,297	29.2 (10)	2,961	21.3 (8)	1,268	11.9 (94)	1,199	7.0 (52)	1,877	80.8(121)	3,936	5.7 (44)
8203	573	53.1(112)	276	11.6 (140)	254	8.7 (115)	120	6.7 (122)	103	1.9(127)	127	85.0 (82)	422	6.6 (64)
8250	628	43.6(156)	259	16.2 (94)	136	10.3 (72)	305	10.2 (103)	247	5.7 (70)	209	72.7(141)	819	7.1 (72)
8255	5,299	47.9(138)	3,185	15.2 (104)	2,834	9.5 (96)	1,502	13.3 (81)	1,396	4.9 (85)	972	79.6(128)	3,237	7.9 (83)
8257	914	52.3(117)	617	25.4 (18)	588	17.9 (17)	276	19.6 (31)	258	10.9 (12)	444	81.3(119)	873	7.3 (76)
8258	3,261	40.1(164)	1,726	9.4 (163)	1,754	5.9 (153)	765	5.1 (131)	726	1.9(127)	896	83.1(103)	2,191	7.2 (74)
8260	743	61.9 (54)	512	30.9 (8)	409	20.0 (10)	277	6.5 (123)	262	5.0 (83)	323	60.7(159)	673	6.1 (54)
8901	1,656	26.9(180)	1,211	9.9 (161)	959	8.6 (117)	585	4.3 (132)	518	2.5(120)	737	61.9(158)	1,512	17.6 (174)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
8902	1,037	33.4(173)	762	16.7 (86)	555	14.1 (30)	368	3.8 (134)	332	1.2(132)	448	67.9(150)	966	12.0 (152)
8903	1,993	25.7(183)	1,453	10.7 (148)	1,072	8.9 (107)	700	3.4 (135)	620	2.4(123)	846	59.6(160)	1,794	12.1 (154)
8904	1,516	46.7(145)	1,030	24.7 (21)	853	20.6 (9)	491	11.4 (97)	430	3.7 (98)	689	69.7(147)	1,429	12.0 (152)
8905	4,268	34.6(169)	2,996	16.4 (90)	2,327	13.1 (36)	1,606	3.4 (135)	1,358	1.8(129)	2,246	74.3(138)	4,604	8.0 (85)
9004	168	28.0(179)	269	8.6 (166)	252	7.9 (129)	0	* *	0	* *	0	* *	13	* *
9010	11	* *	17	* *	13	* *	0	* *	0	* *	0	* *	0	* *
9012	11	* *	5	* *	4	* *	3	* *	3	* *	9	* *	23	* *
9013	52	* *	24	* *	18	* *	25	* *	23	* *	35	* *	86	* *
9015	12	* *	4	* *	2	* *	4	* *	4	* *	7	* *	19	* *
9016	52	* *	28	* *	21	* *	28	* *	28	* *	40	* *	88	* *
9017	19	* *	11	* *	9	* *	10	* *	10	* *	14	* *	34	* *
9018	49	* *	15	* *	9	* *	18	* *	14	* *	32	* *	78	* *
9019	11	* *	0	* *	0	* *	0	* *	0	* *	12	* *	23	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
9021	20	* *	10	* *	8	* *	7	* *	7	* *	3	* *	27	* *
9023	7	* *	9	* *	10	* *	0	* *	0	* *	0	* *	0	* *
9040	103	28.2(178)	138	11.6 (140)	146	12.3 (43)	4	* *	4	* *	13	* *	24	* *
9041	1,193	28.7(177)	872	9.2 (165)	803	9.1 (102)	304	5.3 (129)	300	0.7(133)	447	80.1(125)	877	10.6 (134)
9045	2,118	42.9(160)	1,534	14.9 (111)	1,368	11.6 (49)	627	8.1 (112)	561	2.5(120)	1,138	69.9(146)	2,118	7.3 (76)
9050	3,179	40.1(164)	2,290	19.2 (47)	1,824	14.1 (30)	1,074	7.9 (113)	954	3.9 (96)	1,430	71.6(144)	3,004	7.2 (74)
9051	3,810	54.2(107)	2,463	18.7 (52)	2,232	12.6 (40)	1,109	17.0 (52)	1,063	8.0 (35)	1,488	80.2(124)	3,228	2.7 (12)
9053	4,378	34.8(167)	2,933	10.5 (153)	2,339	7.7 (133)	1,522	7.0 (119)	1,405	3.1(110)	1,438	64.7(155)	3,398	14.2 (166)
9054	46	* *	31	* *	43	* *	0	* *	0	* *	0	* *	0	* *
9055	353	26.1(182)	57	* *	14	* *	67	* *	42	* *	0	* *	216	9.3 (113)
9058	915	63.6 (47)	713	21.9 (31)	595	15.8 (22)	393	6.4 (124)	371	3.2(109)	635	68.2(148)	1,137	12.8 (156)
9066	3,898	26.9(180)	2,331	8.3 (169)	2,083	5.4 (158)	1,419	8.3 (110)	1,289	2.9(113)	1,303	76.1(134)	3,113	8.3 (90)
9068	2,274	34.8(167)	1,713	10.6 (151)	1,352	6.0 (152)	833	5.9 (126)	776	3.4(105)	890	68.2(148)	1,986	14.0 (165)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic
Children Aged < 5 Years (3)

(2)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
Clinic	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
9069	3,439	41.7(162)	2,323	9.9 (161)	1,619	5.6 (156)	1,304	7.7 (116)	1,178	2.9(113)	1,028	70.5(145)	2,792	10.3 (131)
9070	3,133	33.6(172)	1,997	8.5 (167)	1,755	4.8 (166)	886	7.1 (118)	815	2.6(117)	1,130	62.9(157)	2,559	8.3 (90)
9072	2,588	29.2(176)	1,502	6.1 (175)	1,396	3.1 (175)	790	7.6 (117)	736	2.3(124)	962	66.1(153)	2,185	11.5 (147)
9074	2,483	44.7(153)	1,816	10.2 (160)	1,227	5.9 (153)	874	7.8 (115)	817	2.8(115)	895	66.3(152)	2,107	11.2 (140)
9079	338	39.1(166)	429	12.4 (136)	364	9.6 (93)	0	* *	0	* *	0	* *	0	* *
9080	3,700	48.9(135)	2,777	22.7 (27)	2,378	19.1 (15)	1,448	5.7 (127)	1,357	2.7(116)	1,936	64.3(156)	3,766	10.2 (129)
9151	6	* *	10	* *	11	* *	1	* *	1	* *	1	* *	3	* *
Unknown Clinics	339	8.3(184)	139	9.4 (163)	71	* *	55	* *	51	* *	45	* *	109	19.3 (176)
Michigan	274,527	52.0	201,805	15.8	161,280	9.7	92,303	13.6	84,586	5.8	119,260	81.6	251,189	7.9

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

