

Table 8: Demographic information on HIV infection cases currently living in Michigan, 2012

	REPORTED HIV INFECTION PREVALENCE										CENSUS 2010**	
	EST PREV*	HIV, non-stage 3		HIV, stage 3 (AIDS)		TOTAL		Late HIV diagnosis		Num	Percent	
	Num	Num	Percent	Num	Percent	Num	Percent	Rate per 100,000	Num			Percent of stage 3 cases
RACE/ETHNICITY[§]												
White	7,410	2,545	35%	3,121	36%	5,666	36%	75	1,409	45%	7,569,939	77%
Black	11,620	4,111	57%	4,777	56%	8,888	56%	642	1,895	40%	1,383,756	14%
Hispanic	1,000	337	5%	431	5%	768	5%	176	207	48%	436,358	4%
Asian/NH/OPI	130	45	1%	51	1%	96	1%	40	28	55%	238,660	2%
AI/AN	50	26	<1%	15	<1%	41	<1%	75	4	27%	54,665	1%
Multi/other/unk	380	124	2%	170	2%	294	2%	N/A	51	30%	200,262	2%
SEX & RACE												
Male	16,040	5,450	76%	6,819	80%	12,269	78%	253	2,952	43%	4,848,114	49%
White male	6,470	2,160	30%	2,784	33%	4,944	31%	133	1,292	46%	3,728,507	38%
Black male	8,360	2,894	40%	3,500	41%	6,394	41%	973	1,415	40%	657,181	7%
Hispanic male	790	258	4%	346	4%	604	4%	272	175	51%	221,913	2%
Other male	430	138	2%	189	2%	327	2%	136	70	37%	240,513	2%
Female	4,560	1,738	24%	1,746	20%	3,484	22%	69	642	37%	5,035,526	51%
White female	940	385	5%	337	4%	722	5%	19	117	35%	3,841,432	39%
Black female	3,260	1,217	17%	1,277	15%	2,494	16%	343	480	38%	726,575	7%
Hispanic female	210	79	1%	85	1%	164	1%	76	32	38%	214,445	2%
Other female	140	57	1%	47	1%	104	1%	41	13	28%	253,074	3%
RISK†												
Male-male sex (MSM)	10,160	3,451	48%	4,320	50%	7,771	49%	--	1,832	42%	--	--
Injection drug use (IDU)	2,010	593	8%	946	11%	1,539	10%	--	321	34%	--	--
MSM/IDU	910	294	4%	405	5%	699	4%	--	119	29%	--	--
Blood products	120	31	<1%	61	1%	92	1%	--	17	28%	--	--
Heterosexual contact (HC)	3,600	1,317	18%	1,437	17%	2,754	17%	--	534	37%	--	--
HCFR (male)	720	239	3%	313	4%	552	4%	--	123	39%	--	--
HCM (female)	2,880	1,078	15%	1,124	13%	2,202	14%	--	411	37%	--	--
Perinatal	230	108	2%	65	1%	173	1%	--	25	38%	--	--
Undetermined	3,560	1,394	19%	1,331	16%	2,725	17%	--	746	56%	--	--
AGE AT HIV DIAGNOSIS												
0 - 12 years	270	127	2%	76	1%	203	1%	--	23	30%	--	--
13 - 19 years	1,030	473	7%	311	4%	784	5%	--	58	19%	--	--
20 - 24 years	2,810	1,204	17%	947	11%	2,151	14%	--	198	21%	--	--
25 - 29 years	3,440	1,275	18%	1,356	16%	2,631	17%	--	401	30%	--	--
30 - 39 years	7,140	2,266	32%	3,193	37%	5,459	35%	--	1,340	42%	--	--
40 - 49 years	4,200	1,306	18%	1,905	22%	3,211	20%	--	1,078	57%	--	--
50 - 59 years	1,380	438	6%	620	7%	1,058	7%	--	385	62%	--	--
60 years and over	330	96	1%	157	2%	253	2%	--	111	71%	--	--
Unspecified	10	3	<1%	0	0%	3	<1%	--	--	--	--	--
AREA OF CURRENT RESIDENCE^{††}												
Detroit Metro	13,040	4,453	62%	5,466	64%	9,919	63%	232	2,325	43%	4,267,304	43%
Out-State	7,080	2,512	35%	2,877	34%	5,389	34%	96	1,213	42%	5,616,336	57%
Prison	370	174	2%	189	2%	363	2%	N/A	43	23%	N/A	N/A
Unknown ^{††}	110	49	1%	33	<1%	82	1%	N/A	13	39%	N/A	N/A
STATEWIDE TOTAL	20,600	7,188	100%	8,565	100%	15,753	100%	159	3,594	42%	9,883,640	100%

*See pages iv-v for descriptions of prevalence estimate calculations. NOTE: prevalence estimates throughout this document are based on the number of people currently living with HIV in Michigan as of January 2012. Prevalence estimates in other MDCH documents (such as quarterly stats) are based on the number of people living with HIV who were diagnosed in MI.

† See page vi of the Forward and Appendix 2 for risk category groupings. Risk categories used in Michigan are redefined as of January 2012. NOTE: Heterosexual contact for males includes only males whose sexual partners are known to be HIV infected or at high risk for HIV (HCFR). Heterosexual contact for females includes all females who have had sex with a male regardless of what is known about the male's HIV status or behaviors (HCM).

§ In this report, persons described as white, black, Asian/Native Hawaiian or Other Pacific Islander (Asian/NH/OPI), or American Indian/Alaskan Native (AI/AN) are all non-Hispanic; persons described as Hispanic may be of any race.

¶ The Detroit Metro Area consists of Lapeer, Macomb, Monroe, Oakland, St. Clair, and Wayne Counties. The remaining counties comprise the Out-State Area.

** Rates are not reported for risk categories and age at diagnosis because no reliable denominator data exist for these groups.

†† Unknown residence consists of 80 persons released from prison with unknown current location and two non-prisoners with no known residence.

Table 9: HIV infection cases currently living in Michigan by county of current residence, 2012

REPORTED HIV INFECTION PREVALENCE

COUNTY	EST PREV*	HIV, non-stage 3		HIV, stage 3 (AIDS)		TOTAL			Late diagnosis		CENSUS 2010	
		Num	Percent	Num	Percent	Num	Percent	Rate per 100,000	Num	Percent of stage 3 cases	Num	Percent
Alcona	10	0	0%	1	<1%	1	<1%	9	1	100%	10,942	<1%
Alger	10	1	<1%	4	<1%	5	<1%	52	0	0%	9,601	<1%
Allegan	140	38	1%	72	1%	110	1%	99	27	38%	111,408	1%
Alpena	10	1	<1%	9	<1%	10	<1%	34	2	22%	29,598	<1%
Antrim	10	6	<1%	5	<1%	11	<1%	47	3	60%	23,580	<1%
Arenac	10	3	<1%	4	<1%	7	<1%	44	2	50%	15,899	<1%
Baraga	10	1	<1%	4	<1%	5	<1%	56	3	75%	8,860	<1%
Barry	30	6	<1%	18	<1%	24	<1%	41	10	56%	59,173	1%
Bay	100	36	1%	39	<1%	75	<1%	70	15	38%	107,771	1%
Benzie	10	3	<1%	4	<1%	7	<1%	40	1	25%	17,525	<1%
Berrien	330	105	1%	148	2%	253	2%	161	61	41%	156,813	2%
Branch	20	8	<1%	4	<1%	12	<1%	27	2	50%	45,248	<1%
Calhoun	220	80	1%	84	1%	164	1%	120	21	25%	136,146	1%
Cass	40	15	<1%	18	<1%	33	<1%	63	7	39%	52,293	1%
Charlevoix	10	4	<1%	7	<1%	11	<1%	42	3	43%	25,949	<1%
Cheboygan	10	4	<1%	6	<1%	10	<1%	38	1	17%	26,152	<1%
Chippewa	20	11	<1%	8	<1%	19	<1%	49	3	38%	38,520	<1%
Clare	40	10	<1%	17	<1%	27	<1%	87	7	41%	30,926	<1%
Clinton	70	26	<1%	29	<1%	55	<1%	73	13	45%	75,382	1%
Crawford	10	3	<1%	4	<1%	7	<1%	50	3	75%	14,074	<1%
Delta	20	10	<1%	8	<1%	18	<1%	49	2	25%	37,069	<1%
Dickinson	10	1	<1%	5	<1%	6	<1%	23	3	60%	26,168	<1%
Eaton	130	47	1%	51	1%	98	1%	91	11	22%	107,759	1%
Emmet	20	5	<1%	7	<1%	12	<1%	37	4	57%	32,694	<1%
Genesee	720	261	4%	286	3%	547	3%	128	113	40%	425,790	4%
Gladwin	10	2	<1%	2	<1%	4	<1%	16	2	100%	25,692	<1%
Gogebic	10	2	<1%	2	<1%	4	<1%	24	1	50%	16,427	<1%
Grand Traverse	100	35	<1%	39	<1%	74	<1%	85	20	51%	86,986	1%
Gratiot	20	10	<1%	9	<1%	19	<1%	45	5	56%	42,476	<1%
Hillsdale	10	2	<1%	7	<1%	9	<1%	19	4	57%	46,688	<1%
Houghton	10	7	<1%	4	<1%	11	<1%	30	2	50%	36,628	<1%
Huron	10	3	<1%	2	<1%	5	<1%	15	1	50%	33,118	<1%
Ingham	600	227	3%	231	3%	458	3%	163	95	41%	280,895	3%
Ionia	50	21	<1%	18	<1%	39	<1%	61	7	39%	63,905	1%
Iosco	10	5	<1%	2	<1%	7	<1%	27	0	0%	25,887	<1%
Iron	10	1	<1%	1	<1%	2	<1%	17	1	100%	11,817	<1%
Isabella	70	22	<1%	28	<1%	50	<1%	71	10	36%	70,311	1%
Jackson	230	82	1%	96	1%	178	1%	111	35	36%	160,248	2%
Kalamazoo	430	161	2%	169	2%	330	2%	132	57	34%	250,331	3%
Kalkaska	10	3	<1%	0	0%	3	<1%	17	--	--	17,153	<1%
Kent	1,330	452	6%	559	7%	1,011	6%	168	250	45%	602,622	6%
Keweenaw	10	0	0%	0	0%	0	0%	0	--	--	2,156	<1%
Lake	20	5	<1%	7	<1%	12	<1%	104	4	57%	11,539	<1%
Lapeer	50	17	<1%	24	<1%	41	<1%	46	10	42%	88,319	1%
Leelanau	10	3	<1%	6	<1%	9	<1%	41	4	67%	21,708	<1%
Lenawee	90	35	<1%	30	<1%	65	<1%	65	16	53%	99,892	1%
Livingston	90	28	<1%	39	<1%	67	<1%	37	17	44%	180,967	2%
Luce	10	1	<1%	1	<1%	2	<1%	30	0	0%	6,631	<1%

Table 9: HIV infection cases currently living in Michigan by county of current residence, 2012 (continued)

COUNTY	EST PREV*	REPORTED HIV INFECTION PREVALENCE									CENSUS 2010	
		HIV, non-stage 3		HIV, stage 3 (AIDS)		TOTAL		Late diagnosis		Num	Percent	
		Num	Percent	Num	Percent	Num	Percent	Rate per 100,000†	Num			Percent of stage 3 cases
Mackinac	10	3	<1%	1	<1%	4	<1%	36	0	0%	11,113	<1%
Macomb	990	365	5%	391	5%	756	5%	90	189	48%	840,978	9%
Manistee	10	4	<1%	7	<1%	11	<1%	44	3	43%	24,733	<1%
Marquette	50	11	<1%	24	<1%	35	<1%	52	14	58%	67,077	1%
Mason	20	6	<1%	10	<1%	16	<1%	56	6	60%	28,705	<1%
Mecosta	20	7	<1%	8	<1%	15	<1%	35	4	50%	42,798	<1%
Menominee	10	4	<1%	1	<1%	5	<1%	21	1	100%	24,029	<1%
Midland	30	12	<1%	14	<1%	26	<1%	31	9	64%	83,629	1%
Missaukee	10	4	<1%	3	<1%	7	<1%	47	2	67%	14,849	<1%
Monroe	100	37	1%	42	<1%	79	1%	52	22	52%	152,021	2%
Montcalm	30	9	<1%	13	<1%	22	<1%	35	7	54%	63,342	1%
Montmorency	10	0	0%	3	<1%	3	<1%	31	3	100%	9,765	<1%
Muskegon	200	79	1%	72	1%	151	1%	88	34	47%	172,188	2%
Newaygo	20	5	<1%	9	<1%	14	<1%	29	1	11%	48,460	<1%
Oakland	2,400	865	12%	958	11%	1,823	12%	152	402	42%	1,202,362	12%
Oceana	10	5	<1%	2	<1%	7	<1%	26	1	50%	26,570	<1%
Ogemaw	10	1	<1%	2	<1%	3	<1%	14	1	50%	21,699	<1%
Ontonagon	10	0	0%	3	<1%	3	<1%	44	2	67%	6,780	<1%
Osceola	10	2	<1%	5	<1%	7	<1%	30	2	40%	23,528	<1%
Oscoda	10	2	<1%	1	<1%	3	<1%	35	0	0%	8,640	<1%
Otsego	20	4	<1%	9	<1%	13	<1%	54	5	56%	24,164	<1%
Ottawa	140	45	1%	62	1%	107	1%	41	34	55%	263,801	3%
Presque Isle	10	0	0%	2	<1%	2	<1%	15	1	50%	13,376	<1%
Roscommon	20	5	<1%	12	<1%	17	<1%	70	7	58%	24,449	<1%
Saginaw	320	119	2%	121	1%	240	2%	120	47	39%	200,169	2%
Sanilac	20	10	<1%	8	<1%	18	<1%	42	6	75%	43,114	<1%
Schoolcraft	10	1	<1%	0	0%	1	<1%	12	--	--	8,485	<1%
Shiawassee	40	11	<1%	16	<1%	27	<1%	38	9	56%	70,648	1%
St. Clair	150	51	1%	61	1%	112	1%	69	29	48%	163,040	2%
St. Joseph	40	13	<1%	20	<1%	33	<1%	54	9	45%	61,295	1%
Tuscola	10	5	<1%	4	<1%	9	<1%	16	3	75%	55,729	1%
Van Buren	90	36	1%	33	<1%	69	<1%	90	12	36%	76,258	1%
Washtenaw	820	313	4%	310	4%	623	4%	181	139	45%	344,791	3%
Wayne Total	9,340	3,118	43%	3,990	47%	7,108	45%	390	1,673	42%	1,820,584	18%
Wayne, excl. Detroit	2,040	672	9%	882	10%	1,554	10%	140	385	44%	1,106,807	11%
Detroit	7,300	2,446	34%	3,108	36%	5,554	35%	778	1,288	41%	713,777	7%
Wexford	20	4	<1%	8	<1%	12	<1%	37	2	25%	32,735	<1%
AREA †												
Detroit Metro	13,040	4,453	62%	5,466	64%	9,919	63%	232	2,325	43%	4,267,304	43%
Out-State	7,080	2,512	35%	2,877	34%	5,389	34%	96	1,213	42%	5,616,336	57%
Prison	370	174	2%	189	2%	363	2%	N/A	43	23%	N/A	N/A
Unknown§	110	49	1%	33	<1%	82	1%	N/A	13	39%	N/A	N/A
STATEWIDE TOTAL	20,600	7,188	100%	8,565	100%	15,753	100%	159	3,594	42%	9,883,640	100%

*See pages iv-v for descriptions of prevalence estimate calculations. NOTE: prevalence estimates throughout this document are based on the number of people currently living with HIV in Michigan as of January 2012. Prevalence estimates in other MDCH documents (such as quarterly stats) are based on the number of people living with HIV who were diagnosed in MI.

† The Detroit Metro Area consists of Lapeer, Macomb, Monroe, Oakland, St. Clair, and Wayne Counties. The remaining counties comprise the Out-State Area.

§ Unknown residence consists of 80 persons released from prison with unknown current location and two non-prisoners with no known residence.

Table 10: Risk transmission and exposure categories for HIV infection cases currently living in Michigan by sex, 2012

REPORTED HIV INFECTION PREVALENCE

	Male		Female		Overall	
	Num	Percent	Num	Percent	Num	Percent
RISK TRANSMISSION CATEGORIES (CDC Hierarchy) *§						
<i>(Mutually exclusive: one case is represented in ONLY one category)</i>						
Male-male sex (MSM)	7,771	63%	N/A	--	7,771	49%
Injection drug use (IDU)	904	7%	635	18%	1,539	10%
MSM/IDU	699	6%	N/A	--	699	4%
Blood products	77	1%	15	<1%	92	1%
Heterosexual contact (HC)	552	4%	2,202	63%	2,754	17%
HCFR (male)	552	4%	N/A	--	552	4%
HCM (female)	N/A	--	2,202	63%	2,202	14%
Perinatal	97	1%	76	2%	173	1%
Undetermined	2,169	18%	556	16%	2,725	17%

EXPOSURE CATEGORIES *†

<i>(Mutually exclusive: one case is represented in ONLY one category)</i>						
Male-male sex only	5,087	41%	N/A	--	5,087	32%
MSM & HC	2,637	21%	N/A	--	2,637	17%
MSM & IDU	306	2%	N/A	--	306	2%
MSM & blood products	25	<1%	N/A	--	25	<1%
MSM & HC & IDU	377	3%	N/A	--	377	2%
MSM & HC & blood products	22	<1%	N/A	--	22	<1%
MSM & IDU & blood products	4	<1%	N/A	--	4	<1%
MSM & HC & IDU & blood products	12	<1%	N/A	--	12	<1%
Heterosexual contact only	1,959	16%	2,506	72%	4,465	28%
HC & IDU	680	6%	555	16%	1,235	8%
HC & blood products	50	<1%	40	1%	90	1%
HC & IDU & blood products	22	<1%	17	<1%	39	<1%
Injection drug use only	201	2%	63	2%	264	2%
IDU & blood products	1	<1%	0	0%	1	<1%
Perinatal exposure	97	1%	76	2%	173	1%
Exposure to blood products only	41	<1%	4	<1%	45	<1%
Undetermined	748	6%	223	6%	971	6%
TOTAL	12,269	100%	3,484	100%	15,753	100%

SUMMARIZED EXPOSURE CATEGORIES*

<i>(NOT mutually exclusive: one case may be represented in multiple categories)</i>						
Any MSM	8,470	69%	N/A	--	8,470	54%
Behaviorally bisexual males	3,048	25%	N/A	--	3,048	19%
Any heterosexual contact	5,759	47%	3,118	89%	8,877	56%
Any IDU	1,603	13%	635	18%	2,238	14%

*See page ii for descriptions of risk transmission and exposure categories.

§ Risk transmission categories are grouped based on hierarchical categories determined by the CDC. Any one person with multiple risks is only represented in the highest category, with the exception of MSM/IDU (based on the hierarchical algorithm).

† Exposure categories are mutually exclusive and grouped to allow all possible combinations of exposures that any one person may have. NOTE: Heterosexual contact (HC) in exposure categories includes males and females who had heterosexual contact, regardless of what is known about their partners' risk or HIV status.

* Summarized exposure categories are NOT mutually exclusive, i.e. a case may be represented in multiple categories. These summarized categories are meant to give a broader picture of exposure and will NOT add up to the total number of persons living with HIV infection.

Table 11: Sex, race, and risk among HIV infection cases currently living in Michigan, 2012

MALE	White		Black		Hispanic		Other or unknown		All male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	3,702	75%	3,557	56%	332	55%	180	55%	7,771	63%
Injection drug use (IDU)	189	4%	634	10%	60	10%	21	6%	904	7%
MSM/IDU	313	6%	326	5%	35	6%	25	8%	699	6%
Blood products	59	1%	14	<1%	2	<1%	2	1%	77	1%
Heterosexual contact (HCFR)	109	2%	386	6%	43	7%	14	4%	552	4%
Perinatal	18	<1%	67	1%	4	1%	8	2%	97	1%
Undetermined	554	11%	1,410	22%	128	21%	77	24%	2,169	18%
Male Subtotal	4,944	40%	6,394	52%	604	5%	327	3%	12,269	100%

FEMALE	White		Black		Hispanic		Other or unknown		All female	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Injection drug use (IDU)	138	19%	454	18%	29	18%	14	13%	635	18%
Blood products	10	1%	4	<1%	1	1%	0	0%	15	<1%
Heterosexual contact (HCM)	467	65%	1,552	62%	114	70%	69	66%	2,202	63%
Perinatal	10	1%	52	2%	9	5%	5	5%	76	2%
Undetermined	97	13%	432	17%	11	7%	16	15%	556	16%
Female Subtotal	722	21%	2,494	72%	164	5%	104	3%	3,484	100%

ALL	White		Black		Hispanic		Other or unknown		Risk all	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	3,702	65%	3,557	40%	332	43%	180	42%	7,771	49%
Injection drug use (IDU)	327	6%	1,088	12%	89	12%	35	8%	1,539	10%
MSM/IDU	313	6%	326	4%	35	5%	25	6%	699	4%
Blood products	69	1%	18	<1%	3	<1%	2	<1%	92	1%
Heterosexual contact (HC)	576	10%	1,938	22%	157	20%	83	19%	2,754	17%
<i>HCFR (male)</i>	109	2%	386	4%	43	6%	14	3%	552	4%
<i>HCM (female)</i>	467	8%	1,552	17%	114	15%	69	16%	2,202	14%
Perinatal	28	<1%	119	1%	13	2%	13	3%	173	1%
Undetermined	651	11%	1,842	21%	139	18%	93	22%	2,725	17%
RACE ALL	5,666	36%	8,888	56%	768	5%	431	3%	15,753	100%

Table 12: Sex, race, and age at HIV diagnosis among HIV infection cases currently living in Michigan, 2012

<i>MALE</i>	White		Black		Hispanic		Other or unknown		All male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
0 - 12 years	32	1%	73	1%	4	1%	9	3%	118	1%
13 - 19 years	86	2%	454	7%	21	3%	16	5%	577	5%
20 - 24 years	467	9%	1,078	17%	78	13%	45	14%	1,668	14%
25 - 29 years	828	17%	1,025	16%	123	20%	66	20%	2,042	17%
30 - 39 years	1,928	39%	2,036	32%	220	36%	116	35%	4,300	35%
40 - 49 years	1,152	23%	1,242	19%	106	18%	56	17%	2,556	21%
50 - 59 years	353	7%	407	6%	33	5%	16	5%	809	7%
60 years and over	98	2%	77	1%	19	3%	3	1%	197	2%
Unknown	0	0%	2	<1%	0	0%	0	0%	2	<1%
Male Subtotal	4,944	40%	6,394	52%	604	5%	327	3%	12,269	100%

<i>FEMALE</i>	White		Black		Hispanic		Other or unknown		All female	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
0 - 12 years	12	2%	59	2%	9	5%	5	5%	85	2%
13 - 19 years	48	7%	144	6%	12	7%	3	3%	207	6%
20 - 24 years	131	18%	315	13%	24	15%	13	13%	483	14%
25 - 29 years	139	19%	402	16%	30	18%	18	17%	589	17%
30 - 39 years	226	31%	835	33%	55	34%	43	41%	1,159	33%
40 - 49 years	109	15%	510	20%	22	13%	14	13%	655	19%
50 - 59 years	47	7%	185	7%	9	5%	8	8%	249	7%
60 years and over	9	1%	44	2%	3	2%	0	0%	56	2%
Unknown	1	<1%	0	0%	0	0%	0	0%	1	<1%
Female Subtotal	722	21%	2,494	72%	164	5%	104	3%	3,484	100%

<i>ALL</i>	White		Black		Hispanic		Other or unknown		Age all	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
0 - 12 years	44	1%	132	1%	13	2%	14	3%	203	1%
13 - 19 years	134	2%	598	7%	33	4%	19	4%	784	5%
20 - 24 years	598	11%	1,393	16%	102	13%	58	13%	2,151	14%
25 - 29 years	967	17%	1,427	16%	153	20%	84	19%	2,631	17%
30 - 39 years	2,154	38%	2,871	32%	275	36%	159	37%	5,459	35%
40 - 49 years	1,261	22%	1,752	20%	128	17%	70	16%	3,211	20%
50 - 59 years	400	7%	592	7%	42	5%	24	6%	1,058	7%
60 years and over	107	2%	121	1%	22	3%	3	1%	253	2%
Unknown	1	<1%	2	<1%	0	0%	0	0%	3	<1%
RACE ALL	5,666	36%	8,888	56%	768	5%	431	3%	15,753	100%

Table 13: Sex, risk, and age at HIV diagnosis among HIV infection cases currently living in Michigan, 2012

MALE	0 - 12 years		13 - 19 years		20 - 24 years		25 - 29 years		30 - 39 years		40 - 49 years		50 - 59 years		60 years and over		All male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex	0	0%	436	76%	1,298	78%	1,439	70%	2,740	64%	1,372	54%	393	49%	93	47%	7,771	63%
Injection drug use	0	0%	5	1%	33	2%	85	4%	330	8%	338	13%	98	12%	14	7%	903	7%
MSM/IDU	0	0%	14	2%	81	5%	131	6%	293	7%	144	6%	34	4%	2	1%	699	6%
Blood products	14	12%	20	3%	11	1%	12	1%	15	<1%	4	<1%	1	<1%	0	0%	77	1%
Heterosexual contact (HCFR)	0	0%	10	2%	35	2%	96	5%	227	5%	125	5%	46	6%	13	7%	552	4%
Perinatal	93	79%	4	1%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	97	1%
Undetermined	11	9%	88	15%	210	13%	279	14%	695	16%	573	22%	237	29%	75	38%	2,168	18%
Male Subtotal*	118	1%	577	5%	1,668	14%	2,042	17%	4,300	35%	2,556	21%	809	7%	197	2%	12,267	100%
FEMALE	0 - 12 years		13 - 19 years		20 - 24 years		25 - 29 years		30 - 39 years		40 - 49 years		50 - 59 years		60 years and over		All female	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Injection drug use	0	0%	12	6%	63	13%	91	15%	250	22%	167	25%	44	18%	8	14%	635	18%
Blood products	0	0%	2	1%	2	<1%	0	0%	4	<1%	2	<1%	2	1%	3	5%	15	<1%
Heterosexual contact (HCM)	0	0%	162	78%	348	72%	396	67%	720	62%	388	59%	156	63%	32	57%	2,202	63%
Perinatal	76	89%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	76	2%
Undetermined	9	11%	31	15%	70	14%	102	17%	185	16%	98	15%	47	19%	13	23%	555	16%
Female Subtotal*	85	2%	207	6%	483	14%	589	17%	1,159	33%	655	19%	249	7%	56	2%	3,483	100%
ALL	0 - 12 years		13 - 19 years		20 - 24 years		25 - 29 years		30 - 39 years		40 - 49 years		50 - 59 years		60 years and over		Risk all	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex	0	0%	436	56%	1,298	60%	1,439	55%	2,740	50%	1,372	43%	393	37%	93	37%	7,771	49%
Injection drug use	0	0%	17	2%	96	4%	176	7%	580	11%	505	16%	142	13%	22	9%	1,538	10%
MSM/IDU	0	0%	14	2%	81	4%	131	5%	293	5%	144	4%	34	3%	2	1%	699	4%
Blood products	14	7%	22	3%	13	1%	12	<1%	19	<1%	6	<1%	3	<1%	3	1%	92	1%
Heterosexual contact (HC)	0	0%	172	22%	383	18%	492	19%	947	17%	513	16%	202	19%	45	18%	2,754	17%
HCFR (male)	0	0%	10	1%	35	2%	96	4%	227	4%	125	4%	46	4%	13	5%	552	4%
HCM (female)	0	0%	162	21%	348	16%	396	15%	720	13%	388	12%	156	15%	32	13%	2,202	14%
Perinatal	169	83%	4	1%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	173	1%
Undetermined	20	10%	119	15%	280	13%	381	14%	880	16%	671	21%	284	27%	88	35%	2,723	17%
AGE TOTAL*	203	1%	784	5%	2,151	14%	2,631	17%	5,459	35%	3,211	20%	1,058	7%	253	2%	15,750	100%

*Not included in this table are the following cases with unknown age at diagnosis: one male IDU, one male with unknown risk, and one female with unknown risk.

Table 14: Estimated number and rate of new HIV infections in Michigan and the U.S., 2006-2009

	2006				2007				2008				2009			
	Num*	Percent	Rate†	U.S. rate§	Num*	Percent	Rate†	U.S. rate§	Num*	Percent	Rate†	U.S. rate§	Num*	Percent	Rate†	U.S. rate§
SEX																
Male	548	79%	13.4	30.1	770	83%	18.9	34.9	517	77	12.7	29	531	74%	13.1	29.8
Female	150	21%	3.5	9.8	154	17	3.6	10.7	157	23	3.7	9.5	188	26%	4.4	8.6
RACE/ETHNICITY																
White	221	32%	3.3	9.8	327	35	5	11.2	174	26	2.6	8.7	287	40%	4.4	9.1
Black	373	54%	33	72.7	522	57	46.2	79.2	440	65	39	73.2	389	54%	34.6	69.9
AGE AT INFECTION																
13-29 years	239	34%	10.1	21.8	379	41	16	27.2	298	44	12.7	26.5	298	41%	12.8	25.8
30-39 years	184	26%	14	37	276	30	21.6	27.9	178	26	14.2	34.2	161	22%	13.1	32.2
40+ years	274	39%	5.9	N/A¶	269	29	5.7	N/A¶	198	29	4.2	N/A¶	261	36%	5.5	N/A¶
RISK																
Male-male sex (MSM)	466	67%	--	--	630	68	--	--	447	66	--	--	456	63%	--	--
Injection drug user (IDU)	N/A	N/A	--	--	104	11	--	--	108	16	--	--	144	20%	--	--
Heterosexual contact (HC)	154	22%	--	--	189	20	--	--	117	17	--	--	121	17%	--	--
TOTAL	697	100%	8.3	19.8	924	100%	11.1	22.5	674	100%	8.1	19.0	720	100%	8.6	19.0

*Numbers have been adjusted to account for reporting delay.

† Rate per 100,000 population for ages 13 and older, 2009 intercensal estimates.

§ U.S. Rates are from "Estimated HIV Incidence in the United States, 2006-2009" in the online journal, PLoS One, August 2011, Volume 6, Issue 8, e17502 (www.plosone.org).

¶ National data did not include a 40+ age group. Rates were reported for 40-49 and 50-99 age groups.

Table 15: Demographic characteristics of HIV-positive persons with met need compared to HIV-positive persons with unmet need in Michigan, as of November 2011

	<i>Met need</i>		<i>Unmet need</i>		<i>Total</i>		<i>Overall percent unmet need</i>
	<i>Num</i>	<i>Percent of total</i>	<i>Num</i>	<i>Percent of total</i>	<i>Num</i>	<i>Percent of total</i>	
STAGE OF INFECTION							
HIV, non-stage 3	3,691	38%	2,861	52%	6,552	43%	44%
HIV stage 3 (AIDS)	5,974	62%	2,670	48%	8,644	57%	31%
RACE/ETHNICITY*							
White	3,635	38%	1,867	34%	5,502	36%	34%
Black	5,393	56%	3,175	57%	8,568	56%	37%
Hispanic	366	4%	361	7%	727	5%	50%
Asian/NH/OPI	48	<1%	34	1%	82	1%	41%
AI/AN	20	<1%	17	<1%	37	<1%	46%
Multi/other/unk	203	2%	77	1%	280	2%	28%
SEX & RACE							
Male	7,526	78%	4,313	78%	11,839	78%	36%
<i>White male</i>	3,204	33%	1,616	29%	4,820	32%	34%
<i>Black male</i>	3,833	40%	2,318	42%	6,151	40%	38%
<i>Hispanic male</i>	288	3%	276	5%	564	4%	49%
<i>Other male</i>	201	2%	103	2%	304	2%	34%
Female	2,139	22%	1,218	22%	3,357	22%	36%
<i>White female</i>	431	4%	251	5%	682	4%	37%
<i>Black female</i>	1,560	16%	857	15%	2,417	16%	35%
<i>Hispanic female</i>	78	1%	85	2%	163	1%	52%
<i>Other female</i>	70	1%	25	<1%	95	1%	26%
RISK							
Male-male sex (MSM)	4,875	50%	2,601	47%	7,476	49%	35%
Injection drug use (IDU)	844	9%	782	14%	1,626	11%	48%
MSM/IDU	442	5%	275	5%	717	5%	38%
Blood recipient	59	1%	35	1%	94	1%	37%
Heterosexual contact (HC)†	1,825	19%	847	15%	2,672	18%	32%
Perinatal	130	1%	40	1%	170	1%	24%
Undetermined	1,490	15%	951	17%	2,441	16%	39%
AGE AT HIV DIAGNOSIS							
0 - 12 yrs	143	1%	44	1%	187	1%	24%
13 - 19 yrs	352	4%	244	4%	596	4%	41%
20 - 24 yrs	976	10%	803	15%	1,779	12%	45%
25 - 29 yrs	1,391	14%	1,011	18%	2,402	16%	42%
30 - 34 yrs	1,606	17%	1,100	20%	2,706	18%	41%
35 - 39 yrs	1,587	16%	938	17%	2,525	17%	37%
40 - 44 yrs	1,215	13%	701	13%	1,916	13%	37%
45 - 49 yrs	746	8%	402	7%	1,148	8%	35%
50 - 54 yrs	426	4%	223	4%	649	4%	34%
55 - 59 yrs	197	2%	90	2%	287	2%	31%
60 - 64 yrs	87	1%	48	1%	135	1%	36%
65 yrs and older	55	1%	32	1%	87	1%	37%
CURRENT RESIDENCE							
Detroit Metro Area	6,167	64%	3,379	61%	9,546	63%	35%
<i>Lapeer</i>	28	<1%	8	<1%	36	<1%	19%
<i>Macomb</i>	468	5%	228	4%	630	4%	33%
<i>Monroe</i>	49	1%	29	1%	75	<1%	43%
<i>Oakland</i>	1,080	11%	606	11%	1,551	10%	39%
<i>St Clair</i>	77	1%	29	1%	87	1%	33%
<i>Wayne, excl. Detroit</i>	935	10%	486	9%	1,421	9%	35%
<i>Detroit</i>	3,530	37%	1,993	36%	5,251	35%	39%
Out-State Michigan	3,236	33%	1,953	35%	5,189	34%	38%
<i>Washtenaw</i>	392	4%	192	3%	584	4%	33%
<i>Berrien</i>	132	1%	112	2%	244	2%	46%
<i>Genesee</i>	302	3%	229	4%	531	3%	43%
<i>Allegan, Kent, Muskegon and Ottawa</i>	834	9%	466	8%	1,300	9%	36%
<i>Jackson</i>	143	1%	102	2%	245	2%	42%
<i>Kalamazoo and Calhoun</i>	292	3%	168	3%	460	3%	37%
<i>Clinton, Eaton and Ingham</i>	363	4%	197	4%	560	4%	35%
<i>Saginaw, Bay and Midland</i>	193	2%	135	2%	328	2%	41%
<i>Other Out-State counties</i>	585	6%	352	6%	937	6%	38%
Other/unknown§	262	3%	199	4%	461	3%	43%
TOTAL	9,665	100%	5,531	100%	15,196	100%	36%

* In this report, persons described as white, black, Asian/Native Hawaiian or Other Pacific Islander (A/NH/OPI) or American Indian/Alaska Native (AI/AN) are all non-Hispanic. Persons described as Hispanic may be of any race.

† Heterosexual contact (HC) includes males who had sex with females with known risk for HIV (HCFR) and females who had sex with males, regardless of what was known about the male partners' risks (HCM).

§ Persons who are currently in prison are included in 'Other/Unknown' residence.

Table 16: Selected characteristics of HIV-positive persons with viral suppression (<=200 copies/ml) among persons living with HIV infection in Michigan as of 2009*

	Overall population		Persons with at least 1 VL test in 2009		Persons with VL <=200 ^{†§}	
	Num	Percent	Num	Percent	Num	Percent
SEX						
Male	10,547	78%	5,764	55%	3,954	69%
Female	3,056	22%	1,749	57%	1,096	63%
Missing/unknown	1	<1%	0	0%	--	--
AGE AS OF 12/31/2008						
13-24 years	736	5%	416	57%	152	37%
25-34 years	1,940	14%	1,025	53%	569	56%
35-44 years	4,381	32%	2,372	54%	1,596	67%
45-54 years	4,476	33%	2,509	56%	1,799	72%
55-64 years	1,709	13%	998	58%	779	78%
65 years and over	362	3%	193	53%	155	80%
RACE/ETHNICITY[¶]						
Black/African American	7,493	55%	4,199	56%	2,544	61%
Hispanic/Latino	669	5%	293	44%	217	74%
White	5,033	37%	2,778	55%	2,134	77%
Other	409	3%	243	59%	155	64%
RISK						
Male-male sex (MSM)	6,706	49%	3,854	57%	2,677	69%
Injection drug use (IDU) - males	864	6%	409	47%	272	67%
Injection drug use (IDU) - females	612	4%	300	49%	171	57%
MSM/IDU	677	5%	340	50%	219	64%
Heterosexual contact - males	517	4%	283	55%	190	67%
Heterosexual contact - females	1,259	9%	769	61%	502	65%
Other/unknown - males	1,783	13%	878	49%	596	68%
Other/unknown - females	1,185	9%	680	57%	423	62%
COUNTRY OF BIRTH						
U.S.	9,688	71%	5,531	57%	3,635	66%
U.S. dependency	77	1%	27	35%	19	70%
Foreign country	680	5%	318	47%	248	78%
Missing/unknown	3,159	23%	1,637	52%	1,148	70%
TOTAL	13,604	100%	7,513	55%	5,050	67%

*Monitored viral load is calculated based on laboratory testing data which has a longer lag than case reporting. For that reason, data from 2009 is the latest year viral load analyses can be conducted at this time. Analysis based on HIV surveillance data reported through 05/25/2012.

[†]Based on the most recent viral load test result from 01/01/2009 through 12/31/2009.

[§] Among persons with at least 1 VL test.

[¶] Persons described as white, black, and other are all non-Hispanic; persons described as Hispanic/Latino may be of any race. "Other" includes American Indian/Alaska Native, Asian/Native Hawaiian or Other Pacific Islander, multiple races, and unknown race.

[°] Heterosexual contact with a person known to have, or to have a known risk factor for, HIV infection.

Table 17: Gonorrhea, syphilis, and chlamydia cases by sex, race, and age group, Michigan, 2011

	<i>Gonorrhea</i>			<i>P&S syphilis*</i>			<i>Chlamydia</i>			<i>Census 2010</i>	
	Num	Percent	Rate [^]	Num	Percent	Rate [^]	Num	Percent	Rate [^]	Num	Percent
<i>RACE/ ETHNICITY</i>											
White	1,399	11%	18.5	93	34%	1.2	10,866	22%	143.5	7,569,939	77%
Black	6,382	49%	461.2	169	62%	12.2	17,912	36%	1294.4	1,383,756	14%
Hispanic	146	1%	33.5	4	1%	0.9	1,069	2%	245.0	436,358	4%
Other/multi	155	1%	31.4	3	1%	0.6	894	2%	181.1	493,587	5%
Unknown race	4,988	38%	N/A	4	1%	N/A	19,322	39%	N/A	N/A	N/A
<i>SEX & RACE</i>											
Male	5,343	41%	110.2	247	90%	5.1	13,221	26%	272.7	4,848,114	49%
<i>White male</i>	412	3%	11.0	88	32%	2.4	2,491	5%	66.8	3,728,507	38%
<i>Black male</i>	2,975	23%	452.7	149	54%	22.7	5,497	11%	836.5	657,181	7%
<i>Hispanic male</i>	39	<1%	17.6	4	1%	1.8	287	1%	129.3	221,913	2%
<i>Other male</i>	60	<1%	N/A	3	1%	1.2	224	<1%	N/A	240,513	2%
<i>Unknown male</i>	1,857	14%	N/A	3	1%	N/A	4,722	9%	N/A	N/A	N/A
Female	7,706	59%	153.0	26	9%	0.0	36,732	73%	729.5	5,035,526	51%
<i>White female</i>	981	8%	25.5	1	<1%	0.0	8,364	17%	217.7	3,841,432	39%
<i>Black female</i>	3,406	26%	468.8	20	7%	2.8	12,407	25%	1707.6	726,575	7%
<i>Hispanic female</i>	107	1%	49.9	0	0%	0.0	779	2%	363.3	214,445	2%
<i>Other female</i>	101	1%	39.9	0	0%	0.0	667	1%	263.6	253,074	3%
<i>Unknown female</i>	3,111	24%	N/A	1	<1%	N/A	14,515	29%	N/A	N/A	N/A
Unknown sex - all races	21	<1%	N/A	0	0%	N/A	110	<1%	N/A	N/A	N/A
<i>AGE</i>											
0-4 years	10	<1%	1.7	0	0%	0.0	11	<1%	1.8	596,286	6%
5-9 years	3	<1%	0.5	0	0%	0.0	10	<1%	1.6	637,784	6%
10-14 years	148	1%	21.9	0	0%	0.0	662	1%	98.0	675,216	7%
15-19 years	4,249	33%	574.5	14	5%	1.9	19,426	39%	2626.6	739,599	7%
20-24 years	4,473	34%	668.5	63	23%	9.4	18,877	38%	2821.4	669,072	7%
25-29 years	1,806	14%	306.3	44	16%	7.5	6,026	12%	1022.1	589,583	6%
30-34 years	925	7%	161.0	35	13%	6.1	2,556	5%	444.9	574,566	6%
35-39 years	550	4%	89.8	40	15%	6.5	1,185	2%	193.5	612,493	6%
40-44 years	348	3%	52.3	28	10%	4.2	583	1%	87.6	665,481	7%
45-54 years	368	3%	24.4	36	13%	2.4	450	1%	29.8	1,510,033	15%
55-64 years	113	1%	9.0	13	5%	1.0	107	<1%	8.5	1,251,997	13%
65 and over	27	<1%	2.0	1	<1%	0.1	31	<1%	2.3	1,361,530	14%
Unknown age	50	<1%	N/A	0	0%	N/A	139	<1%	N/A	N/A	N/A
<i>TOTAL</i>	13,070	100%	132.2	274	100%	2.8	50,063	100%	506.5	9,883,640	100%

*P&S: Primary and secondary syphilis.

[^]Rate per 100,000 population.

Table 18: Gonorrhea, syphilis, and chlamydia cases by area and local health department jurisdiction, 2011

<i>Local health department jurisdiction</i>	<i>Gonorrhea</i>		<i>P&S syphilis*</i>		<i>Chlamydia</i>		<i>Census 2010</i>
	<i>Num</i>	<i>Rate[^]</i>	<i>Num</i>	<i>Rate[^]</i>	<i>Num</i>	<i>Rate[^]</i>	
Allegan	31	27.8	1	0.9	268	240.6	111,408
Barry/Eaton	55	32.9	1	0.6	424	254.0	166,932
Bay	28	26.0	0	0.0	339	314.6	107,771
Benzie/Leelanau	1	2.5	0	0.0	83	211.6	39,233
Berrien	224	142.8	0	0.0	1,067	680.4	156,813
Br/Hills/St Joseph	25	16.3	0	0.0	299	195.1	153,231
Calhoun	125	91.8	2	1.5	924	678.7	136,146
Chippewa	-	0.0	0	0.0	88	228.5	38,520
Central MI Dist	54	28.3	4	2.1	474	248.4	190,805
Delta/Menominee	4	6.5	3	4.9	101	165.3	61,098
Dickinson/Iron	2	5.3	0	0.0	70	184.3	37,985
District #2	3	4.5	0	0.0	70	104.2	67,168
District #4	11	13.9	1	1.3	92	116.6	78,891
District #10	30	11.5	2	0.8	541	206.8	261,616
Genesee	875	205.5	5	1.2	3,192	749.7	425,790
Grand Traverse	5	5.7	1	1.1	255	293.2	86,986
Huron	-	0.0	0	0.0	47	141.9	33,118
Ingham	342	121.8	7	2.5	1,915	681.7	280,895
Ionia	7	11.0	0	0.0	128	200.3	63,905
Jackson	84	52.4	2	1.2	696	434.3	160,248
Kalamazoo	351	140.2	15	6.0	1,759	702.7	250,331
Kent	734	121.8	9	1.5	3,615	599.9	602,622
Lapeer	13	14.7	1	1.1	122	138.1	88,319
Lenawee	34	34.0	2	2.0	209	209.2	99,892
Livingston	12	6.6	1	0.6	251	138.7	180,967
LMAS District	5	18.3	0	0.0	38	139.0	27,345
Macomb	501	59.6	25	3.0	1,960	233.1	840,978
Marquette	10	14.9	0	0.0	142	211.7	67,077
Midland	14	16.7	1	1.2	191	228.4	83,629
Monroe	64	42.1	1	0.7	352	231.5	152,021
Muskegon	217	126.0	4	2.3	1,219	707.9	172,188
Mid-MI District	42	23.2	2	1.1	349	192.6	181,200
NW Michigan	21	19.7	0	0.0	216	203.0	106,387
Oakland	989	82.3	30	2.5	3,691	307.0	1,202,362
Ottawa	66	25.0	3	1.1	571	216.5	263,801
Saginaw	202	100.9	1	0.5	1,558	778.3	200,169
Sanilac	3	7.0	0	0.0	50	116.0	43,114
Shiawassee	18	25.5	1	1.4	163	230.7	70,648
St Clair	56	34.3	2	1.2	425	260.7	163,040
Tuscola	8	14.4	0	0.0	104	186.6	55,729
Van Buren/Cass	49	38.1	0	0.0	323	251.3	128,551
Washtenaw	237	68.7	11	3.2	1,392	403.7	344,791
Wayne excl Detroit	989	89.4	34	3.1	3,774	341.0	1,106,807
City of Detroit	6,521	913.6	102	14.3	16,414	2299.6	713,777
WestUpDist	3	4.2	0	0.0	91	128.4	70,851
Detroit Metro Area [#]	9,132	214.0	194	4.5	26,738	626.6	4,267,304
Out-State	3,937	70.1	79	1.4	26,738	476.1	5,616,336
TOTAL	13,070	132.2	274	2.8	50,063	506.5	9,883,640

[#]Detroit Metro Area includes Lapeer, Monroe, Macomb, Oakland, St. Clair, and Wayne counties.

* P&S: Primary and secondary syphilis.

[^] Rate per 100,000 population.

Table 19: Reported cases of acute and chronic hepatitis C by sex, race, and age group, Michigan, 2011

	<i>Acute hepatitis C</i>		<i>Chronic hepatitis C</i>			<i>Census 2010</i>	
	Num	Percent	Num	Percent	Rate*	Num	Percent
SEX							
Male	16	52%	4,426	63%	91	4,848,114	49%
Female	15	48%	2,543	36%	51	5,035,526	51%
Unknown	0	0%	22	<1%	N/A	N/A	N/A
RACE[†]							
White	20	65%	2,712	39%	35	7,803,120	79%
Black	3	10%	1,379	20%	98	1,400,362	14%
Asian	1	3%	20	<1%	8	238,199	2%
Native Hawaiian/Other Pacific Islander	0	0%	4	<1%	--	2,604	<1%
American Indian/Alaska Native	0	0%	43	1%	69	62,007	1%
Other	1	3%	82	1%	56	147,029	1%
Unknown race	6	19%	2,524	36%	N/A	N/A	N/A
Multiracial	0	0%	227	3%	99	230,319	2%
AGE							
0-4 years	1	3%	4	<1%	--	596,286	6%
5-9 years	0	0%	2	<1%	--	637,784	6%
10-14 years	0	0%	2	<1%	--	675,216	7%
15-19 years	1	3%	94	1%	13	739,599	7%
20-24 years	5	16%	414	6%	62	669,072	7%
25-29 years	7	23%	516	7%	88	589,583	6%
30-34 years	5	16%	387	6%	67	574,566	6%
35-39 years	3	10%	287	4%	47	612,493	6%
40-44 years	0	0%	411	6%	62	665,481	7%
45-49 years	3	10%	695	10%	93	744,581	8%
50-54 years	3	10%	1,268	18%	166	765,452	8%
55-64 years	2	6%	2,394	34%	191	1,251,997	13%
65 and over	1	3%	501	7%	37	1,361,530	14%
Unknown age	0	0%	16	<1%	N/A	N/A	N/A
TOTAL	31	100%	6,991	100%	71	9,883,640	100%

* Rates are not displayed for <10 cases.

† Hispanic ethnicity is not categorized due to incomplete data. Each race category includes both Hispanic and non-Hispanic persons.

Table 20: Sex, race, and risk among HIV-positive persons currently incarcerated in Michigan, 2012

<i>MALE</i>	White		Black		Hispanic		Other or unknown		All male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	24	41%	93	35%	2	22%	3	50%	122	36%
Injection drug use (IDU)	12	20%	40	15%	3	33%	0	0%	55	16%
MSM/IDU	8	14%	31	12%	1	11%	3	50%	43	13%
Blood products	1	2%	0	0%	0	0%	0	0%	1	<1%
Heterosexual contact (HCFR)	4	7%	30	11%	3	33%	0	0%	37	11%
Perinatal	0	0%	2	1%	0	0%	0	0%	2	1%
Undetermined	10	17%	71	27%	0	0%	0	0%	81	24%
Male Subtotal	59	17%	267	78%	9	3%	6	2%	341	100%

<i>FEMALE</i>	White		Black		Hispanic		Other or unknown		All female	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Injection drug use (IDU)	3	38%	6	50%	0	0%	0	0%	9	41%
Blood Products	0	0%	0	0%	0	0%	0	0%	0	0%
Heterosexual contact (HCM)	5	63%	6	50%	0	0%	1	0%	12	55%
Perinatal	0	0%	0	0%	0	0%	0	0%	0	0%
Undetermined	0	0%	0	0%	0	0%	1	0%	1	5%
Female Subtotal	8	36%	12	55%	0	0%	2	9%	22	100%

<i>All</i>	White		Black		Hispanic		Other or unknown		Risk all	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	24	36%	93	33%	2	22%	3	38%	122	34%
Injection drug use (IDU)	15	22%	46	16%	3	33%	0	0%	64	18%
MSM/IDU	8	12%	31	11%	1	11%	3	38%	43	12%
Blood products	1	1%	0	0%	0	0%	0	0%	1	<1%
Heterosexual contact (HC)	9	13%	36	13%	3	33%	1	13%	49	13%
<i>HCFR (male)</i>	4	6%	30	11%	3	33%	0	0%	37	10%
<i>HCM (female)</i>	5	7%	6	2%	0	0%	1	13%	12	3%
Perinatal	0	0%	2	1%	0	0%	0	0%	2	1%
Undetermined	10	15%	71	25%	0	0%	1	13%	82	23%
RACE ALL	67	18%	279	77%	9	2%	8	2%	363	100%

Table 21: Sex, race, and age at HIV diagnosis among HIV-positive persons currently incarcerated in Michigan, 2012

	White		Black		Hispanic		Other or unknown		All male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
0 - 12 years	0	0%	2	1%	0	0%	0	0%	2	1%
13 - 19 years	2	3%	16	6%	0	0%	0	0%	18	5%
20 - 24 years	11	19%	49	18%	2	22%	1	17%	63	18%
25 - 29 years	15	25%	59	22%	3	33%	1	17%	78	23%
30 - 39 years	22	37%	103	39%	3	33%	3	50%	131	38%
40 - 49 years	8	14%	30	11%	1	11%	1	17%	40	12%
50 - 59 years	1	2%	8	3%	0	0%	0	0%	9	3%
60 years and over	0	0%	0	0%	0	0%	0	0%	0	0%
Male Subtotal	59	17%	267	78%	9	3%	6	2%	341	100%

	White		Black		Hispanic		Other or unknown		All female	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
0 - 12 years	0	0%	0	0%	0	0%	0	0%	0	0%
13 - 19 years	1	13%	0	0%	0	0%	0	0%	1	5%
20 - 24 years	2	25%	3	25%	0	0%	0	0%	5	23%
25 - 29 years	2	25%	4	33%	0	0%	2	0%	8	36%
30 - 39 years	2	25%	3	25%	0	0%	0	0%	5	23%
40 - 49 years	1	13%	2	17%	0	0%	0	0%	3	14%
50 - 59 years	0	0%	0	0%	0	0%	0	0%	0	0%
60 years and over	0	0%	0	0%	0	0%	0	0%	0	0%
Female Subtotal	8	36%	12	55%	0	0%	2	9%	22	100%

	White		Black		Hispanic		Other or unknown		Age all	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
0 - 12 years	0	0%	2	1%	0	0%	0	0%	2	1%
13 - 19 years	3	4%	16	6%	0	0%	0	0%	19	5%
20 - 24 years	13	19%	52	19%	2	22%	1	13%	68	19%
25 - 29 years	17	25%	63	23%	3	33%	3	38%	86	24%
30 - 39 years	24	36%	106	38%	3	33%	3	38%	136	37%
40 - 49 years	9	13%	32	11%	1	11%	1	13%	43	12%
50 - 59 years	1	1%	8	3%	0	0%	0	0%	9	2%
60 years and over	0	0%	0	0%	0	0%	0	0%	0	0%
RACE ALL	67	18%	279	77%	9	2%	8	2%	363	100%

Table 22: Sex, risk, and age at HIV diagnosis among HIV-positive persons currently incarcerated in Michigan, 2012

MALE	0 - 12 years		13 - 19 years		20 - 24 years		25 - 29 years		30 - 39 years		40 - 49 years		50 - 59 years		60 years and over		All male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex	0	0%	10	56%	35	56%	31	40%	39	30%	6	15%	1	11%	0	0%	122	36%
Injection drug use	0	0%	0	0%	6	10%	7	9%	28	21%	11	28%	3	33%	0	0%	55	16%
MSM/IDU	0	0%	3	17%	6	10%	14	18%	15	11%	3	8%	2	22%	0	0%	43	13%
Blood products	0	0%	0	0%	1	2%	0	0%	0	0%	0	0%	0	0%	0	0%	1	0%
Heterosexual contact (HCFR)	0	0%	2	11%	3	5%	10	13%	15	11%	7	18%	0	0%	0	0%	37	11%
Perinatal	2	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	2	1%
Undetermined	0	0%	3	17%	12	19%	16	21%	34	26%	13	33%	3	33%	0	0%	81	24%
Male Subtotal	2	1%	18	5%	63	18%	78	23%	131	38%	40	12%	9	3%	0	0%	341	100%
FEMALE	0 - 12 years		13 - 19 years		20 - 24 years		25 - 29 years		30 - 39 years		40 - 49 years		50 - 59 years		60 years and over		All female	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Injection drug use	0	0%	0	0%	3	60%	4	50%	0	0%	2	67%	0	0%	0	0%	9	41%
Blood products	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
Heterosexual contact (HCM)	0	0%	1	100%	2	40%	3	38%	5	100%	1	33%	0	0%	0	0%	12	55%
Perinatal	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%
Undetermined	0	0%	0	0%	0	0%	1	13%	0	0%	0	0%	0	0%	0	0%	1	5%
Female Subtotal	0	0%	1	5%	5	23%	8	36%	5	23%	3	14%	0	0%	0	0%	22	100%
ALL	0 - 12 years		13 - 19 years		20 - 24 years		25 - 29 years		30 - 39 years		40 - 49 years		50 - 59 years		60 years and over		Age all	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex	0	0%	10	53%	35	51%	31	36%	39	29%	6	14%	1	11%	0	0%	122	34%
Injection drug use	0	0%	0	0%	9	13%	11	13%	28	21%	13	30%	3	33%	0	0%	64	18%
MSM/IDU	0	0%	3	16%	6	9%	14	16%	15	11%	3	7%	2	22%	0	0%	43	12%
Blood products	0	0%	0	0%	1	1%	0	0%	0	0%	0	0%	0	0%	0	0%	1	<1%
Heterosexual contact (HC)	0	0%	3	16%	5	7%	13	15%	20	15%	8	19%	0	0%	0	0%	49	13%
HCFR (male)	0	0%	2	11%	3	4%	10	12%	15	11%	7	16%	0	0%	0	0%	37	10%
HCM (female)	0	0%	1	5%	2	3%	3	3%	5	4%	1	2%	0	0%	0	0%	12	3%
Perinatal	2	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	0	0%	2	1%
Undetermined	0	0%	3	16%	12	18%	17	20%	34	25%	13	30%	3	33%	0	0%	82	23%
AGE ALL	2	1%	19	5%	68	19%	86	24%	136	37%	43	12%	9	2%	0	0%	363	100%

Table 23: Demographic information on Arab American HIV infection cases currently living in Michigan, 2012

	REPORTED PREVALENCE							
	HIV, non-stage 3		HIV, stage 3 (AIDS)		TOTAL		Late HIV diagnosis	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent of stage 3 cases
SEX								
Male	30	75%	47	90%	77	84%	23	44%
Female	10	25%	5	10%	15	16%	1	2%
RISK*								
Male-male sex (MSM)	16	40%	21	40%	37	40%	10	19%
Injection drug use (IDU)	2	5%	2	4%	4	4%	1	2%
MSM/IDU	1	3%	3	6%	4	4%	2	4%
Blood products	1	3%	0	0%	1	1%	--	--
Heterosexual contact (HC)	7	18%	10	19%	17	18%	1	2%
<i>HCFR (male)</i>	1	3%	5	10%	6	7%	0	0%
<i>HCM (female)</i>	6	15%	5	10%	11	12%	1	2%
Perinatal	1	3%	0	0%	1	1%	--	--
Undetermined	12	30%	16	31%	28	30%	10	19%
AGE AT HIV DIAGNOSIS								
0 - 12 years	1	3%	0	0%	1	1%	--	--
13 - 19 years	4	10%	0	0%	4	4%	--	--
20 - 24 years	2	5%	6	12%	8	9%	0	0%
25 - 29 years	13	33%	8	15%	21	23%	1	2%
30 - 39 years	11	28%	19	37%	30	33%	10	19%
40 - 49 years	6	15%	15	29%	21	23%	9	17%
50 - 59 years	2	5%	2	4%	4	4%	2	4%
60 and over	0	0%	2	4%	2	2%	2	4%
Unspecified	1	3%	0	0%	1	1%	--	--
AREA OF CURRENT RESIDENCE[†]								
Detroit Metro Area	37	93%	49	94%	86	93%	24	46%
Out-State	3	8%	3	6%	6	7%	0	0%
TOTAL	40	100%	52	100%	92	100%	24	46%

*See page vi of the Forward and Appendix 2 for risk category groupings. Risk categories used in Michigan are redefined as of January 2012. NOTE: Heterosexual contact for males includes only males whose sexual partners are known to be HIV infected or at high risk for HIV (HCFR). Heterosexual contact for females includes all females who have had sex with a male regardless of what is known about the male's HIV status or behaviors (HCM).

[†] The Detroit Metro Area consists of Lapeer, Macomb, Monroe, Oakland, St. Clair, and Wayne Counties. The remaining counties comprise the Out-State Area.

Table 24: Sex, risk, and age at HIV diagnosis among Arab American HIV infection cases currently living in Michigan, 2012

MALE	0 - 19 years		20 - 29 years		30 - 39 years		40 years and older		All male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	1	25%	13	68%	12	41%	11	44%	37	48%
Injection drug use (IDU)	0	0%	0	0%	1	3%	1	4%	2	3%
MSM/IDU	0	0%	0	0%	3	10%	1	4%	4	5%
Blood products	1	25%	0	0%	0	0%	0	0%	1	1%
Heterosexual contact (HCFR)	0	0%	1	5%	3	10%	2	8%	6	8%
Perinatal	0	0%	0	0%	0	0%	0	0%	0	0%
Undetermined	2	50%	5	26%	10	34%	10	40%	27	35%
Male Subtotal	4	5%	19	25%	29	38%	25	32%	77	100%

FEMALE	0 - 19 years		20 - 29 years		30 - 39 years		40 years and older		All female	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Injection drug use (IDU)	0	0%	1	10%	1	100%	0	0%	2	14%
Blood products	0	0%	0	0%	0	0%	0	0%	0	0%
Heterosexual contact (HCM)	0	0%	9	90%	0	0%	2	100%	11	79%
Perinatal	1	100%	0	0%	0	0%	0	0%	1	7%
Undetermined	0	0%	0	0%	0	0%	0	0%	0	0%
Female Subtotal*	1	7%	10	71%	1	7%	2	14%	14	100%

ALL	0 - 19 years		20 - 29 years		30 - 39 years		40 years and older		Risk all	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	1	20%	13	45%	12	40%	11	41%	37	41%
Injection drug use (IDU)	0	0%	1	3%	2	7%	1	4%	4	4%
MSM/IDU	0	0%	0	0%	3	10%	1	4%	4	4%
Blood products	1	20%	0	0%	0	0%	0	0%	1	1%
Heterosexual contact (HC)	0	0%	10	34%	3	10%	4	15%	17	19%
<i>HCFR (male)</i>	0	0%	1	3%	3	10%	2	7%	6	7%
<i>HCM (female)</i>	0	0%	9	31%	0	0%	2	7%	11	12%
Perinatal	1	20%	0	0%	0	0%	0	0%	1	1%
Undetermined	2	40%	5	17%	10	33%	10	37%	27	30%
AGE ALL*	5	5%	29	32%	30	33%	27	30%	91	100%

*Not included in this table are the following cases with unknown age at diagnosis: one female with unknown risk.

Table 25: Demographic information on Asian, Native Hawaiian, and Other Pacific Islander HIV infection cases currently living in Michigan, 2012

	REPORTED PREVALENCE							
	HIV, non-stage 3		HIV, stage 3 (AIDS)		TOTAL		Late HIV diagnosis	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent of stage 3 cases
SEX								
Male	29	64%	41	80%	70	73%	23	45%
Female	16	36%	10	20%	26	27%	5	10%
RISK*								
Male-male sex (MSM)	13	29%	15	29%	28	29%	12	24%
Injection drug use (IDU)	3	7%	3	6%	6	6%	1	2%
MSM/IDU	0	0%	1	2%	1	1%	0	0%
Blood products	1	2%	0	0%	1	1%	--	--
Heterosexual contact (HC)	8	18%	12	24%	20	21%	5	10%
<i>HCFR (male)</i>	0	0%	3	6%	3	3%	1	2%
<i>HCM (female)</i>	8	18%	9	18%	17	18%	4	8%
Perinatal	2	4%	0	0%	2	2%	--	--
Undetermined	19	42%	20	39%	39	41%	10	20%
AGE AT HIV DIAGNOSIS								
0 - 12 years	2	4%	0	0%	2	2%	--	--
13 - 19 years	1	2%	2	4%	3	3%	--	--
20 - 24 years	7	16%	4	8%	11	11%	2	4%
25 - 29 years	13	29%	16	31%	29	30%	7	14%
30 - 39 years	16	36%	15	29%	31	32%	11	22%
40 - 49 years	6	13%	12	24%	18	19%	8	16%
50 - 59 years	0	0%	2	4%	2	2%	0	0%
60 and over	0	0%	0	0%	0	0%	--	--
Unspecified	0	0%	0	0%	0	0%	--	--
AREA OF CURRENT RESIDENCE[†]								
Detroit Metro Area	21	47%	27	53%	48	50%	17	33%
Out-State	24	53%	22	43%	46	48%	9	18%
Prison or unknown	0	0%	3	6%	3	3%	2	4%
TOTAL	45	100%	51	100%	96	100%	28	55%

*See page vi of the Forward and Appendix 2 for risk category groupings. Risk categories used in Michigan are redefined as of January 2012. NOTE: Heterosexual contact for males includes only males whose sexual partners are known to be HIV infected or at high risk for HIV (HCFR). Heterosexual contact for females includes all females who have had sex with a male regardless of what is known about the male's HIV status or behaviors (HCM).

[†] The Detroit Metro Area consists of Lapeer, Macomb, Monroe, Oakland, St. Clair, and Wayne Counties. The remaining counties comprise the Out-State Area.

Table 26: Sex, Risk, and Age at HIV Diagnosis Among Asian, Native Hawaiian and Pacific Islander HIV Infection Cases Currently Living in Michigan, 2012

MALE	0 - 19 years		20 - 29 years		30 - 39 years		40 years and older		All male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	0	0%	12	41%	9	39%	7	47%	28	40%
Injection drug use (IDU)	0	0%	3	10%	2	9%	1	7%	6	9%
MSM/IDU	0	0%	1	3%	0	0%	0	0%	1	1%
Blood products	0	0%	0	0%	0	0%	0	0%	0	0%
Heterosexual contact (HCFR)	0	0%	2	7%	1	4%	0	0%	3	4%
Perinatal	1	33%	0	0%	0	0%	0	0%	1	1%
Undetermined	2	67%	11	38%	11	48%	7	47%	31	44%
Male Subtotal	3	4%	29	41%	23	33%	15	21%	70	100%

FEMALE	0 - 19 years		20 - 29 years		30 - 39 years		40 years and older		All female	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Injection drug use (IDU)	0	0%	0	0%	0	0%	0	0%	0	0%
Blood products	0	0%	0	0%	0	0%	0	0%	0	0%
Heterosexual contact (HCM)	1	50%	7	64%	4	50%	5	100%	17	65%
Perinatal	1	50%	0	0%	0	0%	0	0%	1	4%
Undetermined	0	0%	4	36%	4	50%	0	0%	8	31%
Female Subtotal*	2	8%	11	42%	8	31%	5	19%	26	100%

ALL	0 - 19 years		20 - 29 years		30 - 39 years		40 years and older		Risk all	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	0	0%	12	30%	9	29%	7	35%	28	29%
Injection drug use (IDU)	0	0%	3	8%	2	6%	1	5%	6	6%
MSM/IDU	0	0%	1	3%	0	0%	0	0%	1	1%
Blood products	0	0%	0	0%	0	0%	0	0%	0	0%
Heterosexual contact (HC)	1	20%	9	23%	5	16%	5	25%	20	21%
<i>HCFR (male)</i>	0	0%	2	5%	1	3%	0	0%	3	3%
<i>HCM (female)</i>	1	20%	7	18%	4	13%	5	25%	17	18%
Perinatal	2	40%	0	0%	0	0%	0	0%	2	2%
Undetermined	2	40%	15	38%	15	48%	7	35%	39	41%
AGE ALL*	5	5%	40	42%	31	32%	20	21%	96	100%

Table 27: Demographic information on American Indian and Alaska Native HIV infection cases currently living in Michigan, 2012

	REPORTED PREVALENCE							
	HIV, non-stage 3		HIV, stage 3 (AIDS)		TOTAL		Late HIV diagnosis	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent of stage 3 cases
SEX								
Male	20	77%	12	80%	32	78%	3	20%
Female	6	23%	3	20%	9	22%	1	7%
RISK*								
Male-male sex (MSM)	9	35%	6	40%	15	37%	2	13%
Injection drug use (IDU)	1	4%	0	0%	1	2%	--	--
MSM/IDU	3	12%	3	20%	6	15%	0	0%
Blood products	1	4%	0	0%	1	2%	--	--
Heterosexual contact (HC)	6	23%	4	27%	10	24%	1	7%
HCFR (male)	1	4%	1	7%	2	5%	0	0%
HCM (female)	5	19%	3	20%	8	20%	1	7%
Perinatal	1	4%	0	0%	1	2%	--	--
Undetermined	6	23%	2	13%	8	20%	1	7%
AGE AT HIV DIAGNOSIS								
0 - 12 years	1	4%	0	0%	1	2%	--	--
13 - 19 years	0	0%	0	0%	0	0%	--	--
20 - 24 years	8	31%	3	20%	11	27%	1	7%
25 - 29 years	1	4%	3	20%	4	10%	0	0%
30 - 39 years	11	42%	8	53%	19	46%	3	20%
40 - 49 years	4	15%	1	7%	5	12%	0	0%
50 - 59 years	0	0%	0	0%	0	0%	--	--
60 and over	1	4%	0	0%	1	2%	--	--
Unspecified	0	0%	0	0%	0	0%	--	--
AREA OF CURRENT RESIDENCE[†]								
Detroit Metro Area	11	42%	5	33%	16	39%	1	7%
Out-State	15	58%	9	60%	24	59%	3	20%
Prison	0	0%	1	7%	1	2%	0	0%
TOTAL	26	100%	15	100%	41	100%	4	27%

*See page vi of the Forward and Appendix 2 for risk category groupings. Risk categories used in Michigan are redefined as of January 2012. NOTE: Heterosexual contact for males includes only males whose sexual partners are known to be HIV infected or at high risk for HIV (HCFR). Heterosexual contact for females includes all females who have had sex with a male regardless of what is known about the male's HIV status or behaviors (HCM).

[†] The Detroit Metro Area consists of Lapeer, Macomb, Monroe, Oakland, St. Clair, and Wayne Counties. The remaining counties comprise the Out-State Area.

Table 28: Sex, risk, and age at HIV diagnosis among American Indian and Alaska Native HIV infection cases currently living in Michigan, 2012

MALE	0 - 19 years		20 - 29 years		30 - 39 years		40 years and older		All male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	0	0%	5	42%	8	50%	2	50%	15	47%
Injection drug use (IDU)	0	0%	0	0%	1	6%	0	0%	1	3%
MSM/IDU	0	0%	4	33%	2	13%	0	0%	6	19%
Blood products	0	0%	0	0%	0	0%	0	0%	0	0%
Heterosexual contact (HCFR)	0	0%	1	8%	0	0%	1	25%	2	6%
Perinatal	0	0%	0	0%	0	0%	0	0%	0	0%
Undetermined	0	0%	2	17%	5	31%	1	25%	8	25%
Male Subtotal	0	0%	12	38%	16	50%	4	13%	32	100%

FEMALE	0 - 19 years		20 - 29 years		30 - 39 years		40 years and older		All female	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Injection drug use (IDU)	0	0%	0	0%	0	0%	0	0%	0	0%
Blood products	0	0%	0	0%	0	0%	0	0%	0	0%
Heterosexual contact (HCM)	0	0%	3	100%	3	100%	2	100%	8	89%
Perinatal	1	100%	0	0%	0	0%	0	0%	1	11%
Undetermined	0	0%	0	0%	0	0%	0	0%	0	0%
Female Subtotal*	1	11%	3	33%	3	33%	2	22%	9	100%

ALL	0 - 19 years		20 - 29 years		30 - 39 years		40 years and older		Risk all	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
Male-male sex (MSM)	0	0%	5	33%	8	42%	2	33%	15	37%
Injection drug use (IDU)	0	0%	0	0%	1	5%	0	0%	1	2%
MSM/IDU	0	0%	4	27%	2	11%	0	0%	6	15%
Blood products	0	0%	0	0%	0	0%	0	0%	0	0%
Heterosexual contact (HC)	0	0%	4	27%	3	16%	3	50%	10	24%
<i>HCFR (male)</i>	0	0%	1	7%	0	0%	1	17%	2	5%
<i>HCM (female)</i>	0	0%	3	20%	3	16%	2	33%	8	20%
Perinatal	1	100%	0	0%	0	0%	0	0%	1	2%
Undetermined	0	0%	0	0%	0	0%	0	0%	8	20%
AGE ALL*	1	2%	15	37%	19	46%	6	15%	41	100%