

## **CDC Health Advisory: Interim Recommendations for the Use of Influenza Antiviral Medications in the Setting of Oseltamivir Resistance among Circulating Influenza A (H1N1) Viruses, 2008-09 Influenza Season**

Enclosed are CDC's interim recommendations for the use of influenza antiviral medications. Health care professionals should continue to refer to the CDC website at [www.cdc.gov/flu](http://www.cdc.gov/flu) for updated information.

### **New Labeling for Some Influenza Antivirals**

Influenza viruses change over time. Emergence of resistance mutations could decrease drug effectiveness. Other factors (for example, changes in viral virulence) might also diminish clinical benefit of antiviral drugs. Prescribers should consider available information on influenza drug susceptibility patterns and treatment effects when deciding whether to use RELENZA® (zanamivir) Inhalation Powder.

### **Indications and Usage**

- RELENZA is indicated for treatment of uncomplicated acute illness due to influenza A and B virus in adults and pediatric patients 7 years of age and older who have been symptomatic for no more than 2 days.
- RELENZA is indicated for prophylaxis of influenza in adults and pediatric patients 5 years of age and older.

### **Important Limitations on Use of RELENZA**

Not recommended for treatment or prophylaxis of influenza in:

- Individuals with underlying airways disease.

Not proven effective for:

- Treatment in individuals with underlying airways disease.
- Prophylaxis in nursing home residents.

Not a substitute for annual influenza vaccination.

Consider available information on influenza drug susceptibility patterns and treatment effects when deciding whether to use RELENZA.

### **Warnings and Precautions**

- Bronchospasm: Serious, sometimes fatal, cases have occurred. Not recommended in individuals with underlying airways disease. Discontinue RELENZA if bronchospasm or decline in respiratory function develops.
- Allergic Reactions: Discontinue RELENZA and initiate appropriate treatment if an allergic reaction occurs or is suspected.
- Neuropsychiatric Events: Patients with influenza, particularly pediatric patients, may be at an increased risk of seizures, confusion, or abnormal behavior early in their illness. Monitor for signs of abnormal behavior.
- High-risk Underlying Medical Conditions: Safety and effectiveness have not been demonstrated in these patients.

Clinical significance cannot be inferred from surveillance data

GlaxoSmithKline markets and distributes RELENZA, and is disseminating these interim recommendations at the company's expense.

*Please see accompanying complete Prescribing Information for RELENZA.*



**This is an official  
CDC HEALTH ADVISORY**

Distributed via Health Alert Network  
Friday, December 19, 2008, 11:50 EST (11:50 AM EST)  
CDCHAN-00279-2008-12-19-ADV-N

**CDC Issues Interim Recommendations for the Use  
of Influenza Antiviral Medications in the Setting of  
Oseltamivir Resistance among Circulating  
Influenza A (H1N1) Viruses,  
2008-09 Influenza Season**

Although influenza activity is low in the United States to date, preliminary data from a limited number of states indicate that the prevalence of influenza A (H1N1) virus strains resistant to the antiviral medication oseltamivir is high. Therefore, CDC is issuing interim recommendations for antiviral treatment and chemoprophylaxis of influenza during the 2008-09 influenza season. When influenza A (H1N1) virus infection or exposure is suspected, zanamivir or a combination of oseltamivir and rimantadine are more appropriate options than oseltamivir alone. Local influenza surveillance data and laboratory testing can help with physician decision-making regarding the choice of antiviral agents for their patients. The 2008-09 influenza vaccine is expected to be effective in preventing or reducing the severity of illness with currently circulating influenza viruses, including oseltamivir-resistant influenza A (H1N1) virus strains. Since influenza activity remains low and is expected to increase in the weeks and months to come, CDC recommends that influenza vaccination efforts continue.

**Background**

Influenza A viruses, including two subtypes (H1N1) and (H3N2), and influenza B viruses, currently circulate worldwide, but the prevalence of each can vary among communities and within a single community over the course of an influenza season. In the United States, four prescription antiviral medications (oseltamivir, zanamivir, amantadine and rimantadine) are approved for treatment and chemoprophylaxis of influenza. Since January 2006, the neuraminidase inhibitors (oseltamivir, zanamivir) have been the only recommended influenza antiviral drugs because of widespread resistance to the adamantanes (amantadine, rimantadine) among influenza A (H3N2) virus strains. The neuraminidase inhibitors have activity against influenza A and B viruses while the adamantanes have activity only against influenza A viruses. In 2007-08, a significant increase in the prevalence of oseltamivir resistance was reported among influenza A (H1N1) viruses worldwide. During the 2007-08 influenza season, 10.9% of H1N1 viruses tested in the U.S. were resistant to oseltamivir.

Influenza activity has been low thus far this season in the United States. As of December 19, 2008, a limited number of influenza viruses isolated in the U.S. since October 1 have been available for antiviral resistance testing at CDC. Of the 50 H1N1 viruses tested to date from 12 states, 98% were resistant to oseltamivir, and all were susceptible to zanamivir, amantadine and rimantadine. Preliminary data indicate that oseltamivir-resistant influenza A (H1N1) viruses do not cause different or more severe symptoms compared to oseltamivir sensitive influenza A (H1N1) viruses. Influenza A (H3N2) and B viruses remain susceptible to oseltamivir. The proportion of influenza A (H1N1) viruses among all influenza A and B viruses that will circulate during the 2008-09 season cannot be predicted, and will likely vary over the course of the season and among communities. Oseltamivir-resistant influenza A (H1N1) viruses are antigenically similar to the influenza A (H1N1) virus strain represented in 2008-09 influenza vaccine, and CDC recommends that influenza vaccination efforts continue as the primary method to prevent influenza.

Oseltamivir resistance among circulating influenza A (H1N1) virus strains presents challenges for the selection of antiviral medications for treatment and chemoprophylaxis of influenza, and provides additional reasons for clinicians to test patients for influenza virus infection and to consult surveillance data when evaluating persons with acute respiratory illnesses during influenza season. These interim guidelines provide options for treatment or chemoprophylaxis of influenza in the United States if oseltamivir-resistant H1N1 viruses are circulating widely in a community or if the prevalence of oseltamivir resistant H1N1 viruses is uncertain.

### **Interim Recommendations**

Persons providing medical care for patients with suspected influenza or persons who are candidates for chemoprophylaxis against influenza should consider the following guidance for assessing and treating patients during the 2008-09 influenza season (see Table below). Guidance Table):

- 1) Review local or state influenza virus surveillance data weekly during influenza season, to determine which types (A or B) and subtypes of influenza A virus (H3N2 or H1N1) are currently circulating in the area. For some communities, surveillance data might not be available or timely enough to provide information useful to clinicians.
- 2) Consider use of influenza tests that can distinguish influenza A from influenza B.
  - a. Patients testing positive for influenza B may be given either oseltamivir or zanamivir (no preference) if treatment is indicated.
  - b. At this time, if a patient tests positive for influenza A, use of zanamivir should be considered if treatment is indicated. Oseltamivir should be used alone only if recent local surveillance data indicate that circulating viruses are likely to be influenza A (H3N2) or influenza B viruses. Combination treatment with oseltamivir and rimantadine is an acceptable alternative, and might be necessary for patients that cannot receive zanamivir, (e.g., patient is <7 years old, has chronic underlying airways disease, or cannot use the zanamivir inhalation device), or zanamivir is unavailable. Amantadine can be substituted for rimantadine if rimantadine is unavailable.
  - c. If a patient tests negative for influenza, consider treatment options based on local influenza activity and clinical impression of the likelihood of influenza. Because rapid antigen tests may have low sensitivity, treatment should still be considered during periods of high influenza activity for persons with respiratory symptoms consistent with influenza who test negative and have no alternative diagnosis. Use of zanamivir should be considered if treatment is indicated. Combination treatment with oseltamivir and rimantadine (substitute amantadine if rimantadine unavailable) is an acceptable alternative. Oseltamivir should be used alone only if recent local surveillance data indicates that circulating viruses are likely to be influenza A(H3N2) or influenza B viruses.
  - d. If available, confirmatory testing with a diagnostic test capable of distinguishing influenza caused by influenza A (H1N1) virus from influenza caused by influenza A (H3N2) or influenza B virus can also be used to guide treatment. When treatment is indicated, influenza A (H3N2) and influenza B virus infections should be treated with oseltamivir or zanamivir (no preference). Influenza A (H1N1) virus infections should be treated with zanamivir or combination treatment with oseltamivir and rimantadine is an acceptable alternative.
- 3) Persons who are candidates for chemoprophylaxis (e.g., residents in an assisted living facility during an influenza outbreak, or persons who are at higher risk for influenza-related complications and have had recent household or other close contact with a person with laboratory confirmed influenza) should be provided with medications most likely to be effective against the influenza virus that is the cause of the outbreak, if known. Respiratory specimens from ill persons during institutional outbreaks should be obtained and sent for testing to determine the type and subtype of influenza A viruses associated with the outbreak and to guide antiviral therapy decisions. Persons whose need for chemoprophylaxis is due to potential exposure to a person with laboratory-confirmed influenza A (H3N2) or influenza B should receive oseltamivir or zanamivir (no preference). Zanamivir should be used when persons require chemoprophylaxis due to exposure to influenza A (H1N1) virus. Rimantadine can be used if zanamivir use is contraindicated.

Enhanced surveillance for influenza antiviral resistance is ongoing at CDC in collaboration with local and state health departments. Clinicians should remain alert for additional changes in recommendations that might occur as the 2008--09 influenza season progresses. Oseltamivir resistant influenza A (H1N1) viruses are antigenically similar to the influenza A(H1N1) viruses represented in the vaccine, and vaccination should continue to be considered the primary prevention strategy regardless of oseltamivir sensitivity. Information on antiviral resistance will be updated in weekly surveillance reports (available at <http://www.cdc.gov/flu/weekly/fluactivity.htm>)

For more information on antiviral medications and additional considerations related to antiviral use during the 2008-09 influenza season, visit <http://www.cdc.gov/flu/professionals/antivirals/index.htm>

TABLE

**Interim recommendations for the selection of antiviral treatment using**

Rapid antigen or other laboratory test	Predominant virus(es) in community	Preferred medication(s)	Alternative (combination antiviral treatment)
Not done or negative, but clinical suspicion for influenza	H1N1 or unknown	Zanamivir	Oseltamivir + Rimantadine*
Not done or negative, but clinical suspicion for influenza	H3N2 or B	Oseltamivir or Zanamivir	None
Positive A	H1N1 or unknown	Zanamivir	Oseltamivir + Rimantadine*
Positive A	H3N2 or B	Oseltamivir or Zanamivir	None
Positive B	Any	Oseltamivir or Zanamivir	None
Positive A+B**	H1N1 or unknown	Zanamivir	Oseltamivir + Rimantadine*
Positive A+B**	H3N2 or B	Oseltamivir or Zanamivir	None

**laboratory test results and viral surveillance data, United States, 2008-09 season<sup>‡</sup>**

\*Amantadine can be substituted for rimantadine but has increased risk of adverse events. Human data are lacking to support the benefits of combination antiviral treatment of influenza; however, these interim recommendations are intended to assist clinicians treating patients who might be infected with oseltamivir-resistant influenza A (H1N1) virus.

\*\*Positive A+B indicates a rapid antigen test that cannot distinguish between influenza and influenza B viruses

‡ Influenza antiviral medications used for treatment are most beneficial when initiated within the first two days of illness. Clinicians should consult the package insert of each antiviral medication for specific dosing information, approved indications and ages, contraindications/warnings/precautions, and adverse effects.

---

##This Message was distributed to State and Local Health Officers, Public Information Officers, Epidemiologists, State Laboratory Directors, PHEP/BT Coordinators and HAN Coordinators, as well as Public Health Associations and Clinician organizations##

**Categories of Health Alert Messages:**

---

<b>Health Alert</b>	Conveys the highest level of importance; warrants immediate action or attention.
<b>Health Advisory</b>	Provides important information for a specific incident or situation; may not require immediate action.
<b>Health Update</b>	Provides updated information regarding an incident or situation; unlikely to require immediate action.

---

You have received this message based upon the information contained within our emergency notification database. If you have a different or additional e-mail or fax address that you would like us to use, please contact the Health Alert Network program at your State or local health department.