

TABLE 1. Demographic Information on Prevalent HIV/AIDS Cases Living in KENT CO. at Diagnosis

	<i>EST PREV*</i>	<i>REPORTED PREVALENCE[†]</i>						<i>CENSUS 2008 ESTIMATES</i>	
		HIV, not AIDS		AIDS		Total		Rate per 100,000 [§]	Number Percent
	Number	Number	Percent	Number	Percent	Number	Percent		
RACE/ ETHNICITY[¶]									
White	510	180	(53%)	208	(48%)	388	(50%)	82.6	469,908 (78%)
Black	350	116	(34%)	150	(35%)	266	(35%)	503.3	52,852 (9%)
Hispanic	130	37	(11%)	61	(14%)	98	(13%)	169.9	57,691 (10%)
Asian/PI	10	<5	**	<5	**	7	(1%)	---	12,874 (2%)
Am Indian/AN	10	<5	**	<5	**	<5	**	**	2,550 (<1%)
Multi/Unk	10	<5	**	5	(1%)	8	(1%)	N/A	9,338 (2%)
SEX & RACE									
Males	780	246	(72%)	344	(80%)	590	(77%)	196.8	299,863 (50%)
White Males	460	156	(46%)	195	(45%)	351	(46%)	151.8	231,169 (38%)
Black Males	200	57	(17%)	93	(22%)	150	(19%)	590.8	25,388 (4%)
Hispanic Males	100	28	(8%)	50	(12%)	78	(10%)	247.2	31,554 (5%)
Other Males	10	5	(1%)	6	(1%)	11	(1%)	93.6	11,752 (2%)
Females	240	95	(28%)	86	(20%)	181	(23%)	59.3	305,350 (50%)
White Females	50	24	(7%)	13	(3%)	37	(5%)	15.5	238,739 (39%)
Black Females	150	59	(17%)	57	(13%)	116	(15%)	422.4	27,464 (5%)
Hispanic Fmls	30	9	(3%)	11	(3%)	20	(3%)	76.5	26,137 (4%)
Other Females	10	<5	**	5	(1%)	8	(1%)	---	13,010 (2%)
RISK*									
Male-Male Sex	530	172	(50%)	229	(53%)	401	(52%)	N/A	N/A N/A
Injection Drug Use	60	18	(5%)	25	(6%)	43	(6%)	N/A	N/A N/A
MSM/IDU	50	19	(6%)	16	(4%)	35	(5%)	N/A	N/A N/A
Blood Products	10	0	(0%)	<5	**	<5	**	N/A	N/A N/A
Heterosexual	220	80	(23%)	87	(20%)	167	(22%)	N/A	N/A N/A
HRH	160	57	(17%)	63	(15%)	120	(16%)	N/A	N/A N/A
PH-Female	60	23	(7%)	24	(6%)	47	(6%)	N/A	N/A N/A
Perinatal	10	<5	**	<5	**	7	(1%)	N/A	N/A N/A
Undetermined	150	48	(14%)	68	(16%)	116	(15%)	N/A	N/A N/A
PH-Male	110	28	(8%)	52	(12%)	80	(10%)	N/A	N/A N/A
Unknown	50	20	(6%)	16	(4%)	36	(5%)	N/A	N/A N/A
AGE AT HIV DIAGNOSIS									
0 - 12 years	10	6	(2%)	<5	**	9	(1%)	N/A	N/A N/A
13 - 19 years	50	14	(4%)	23	(5%)	37	(5%)	N/A	N/A N/A
20 - 24 years	130	56	(16%)	46	(11%)	102	(13%)	N/A	N/A N/A
25 - 29 years	180	68	(20%)	71	(17%)	139	(18%)	N/A	N/A N/A
30 - 39 years	360	110	(32%)	163	(38%)	273	(35%)	N/A	N/A N/A
40 - 49 years	200	68	(20%)	83	(19%)	151	(20%)	N/A	N/A N/A
50 - 59 years	70	17	(5%)	33	(8%)	50	(6%)	N/A	N/A N/A
60 years and over	10	<5	**	8	(2%)	10	(1%)	N/A	N/A N/A
Unspecified	0	0	(0%)	0	(0%)	0	(0%)	N/A	N/A N/A
TOTAL	1,020	341	(100%)	430	(100%)	771	(100%)	127	605,213 (100%)

*See page 4 for descriptions of prevalence estimate calculations and new risk category groupings. Risk categories used in Michigan are newly defined as of the July 2007 quarter.

[†] Includes reports that contain patient name or are otherwise unduplicated. <5 and "*" = 1, 2, 3, or 4 cases.

[§] To calculate "1 out of x" statements, divide the census number by total reported prevalence. For example, for non-Hispanic whites: 469,908/ 388 = 1,211. Thus, 1 out of every 1,211 non-Hispanic white persons in KENT CO. are living with HIV. Rates and "1 out of x" statements are not reliable for <10 cases. Thus, rates for <10 cases are shown as "----".

[¶] In this report, persons described as white, black, Asian/Pacific Islander (PI), or American Indian/Alaska Native (AN) are all non-Hispanic; persons described as Hispanic might be of any race.

TABLE 2. Sex, Race, and Risk Among Prevalent HIV/AIDS Cases in KENT CO.

MALES	White		Black		Hispanic		Other or Unknown		Male Subtotal	
Male-Male sex	288	(82%)	64	(43%)	45	(58%)	<5	**	401	(68%)
Injecting Drug Use	5	(1%)	13	(9%)	<5	**	0	(0%)	22	(4%)
Male-Male Sex/IDU	20	(6%)	11	(7%)	<5	**	<5	**	35	(6%)
Blood Products	<5	**	0	(0%)	0	(0%)	0	(0%)	<5	**
Heterosexual*	6	(2%)	17	(11%)	7	(9%)	<5	**	32	(5%)
Perinatal	<5	**	<5	**	0	(0%)	0	(0%)	<5	**
Undetermined	29	(8%)	42	(28%)	20	(26%)	<5	**	94	(16%)
<i>PH-Male</i>	21	(6%)	38	(25%)	18	(23%)	<5	**	80	(14%)
<i>Unknown</i>	8	(2%)	<5	**	<5	**	0	(0%)	14	(2%)
Male Subtotal	351	(59%)	150	(25%)	78	(13%)	11	(2%)	590	(100%)
FEMALES	White		Black		Hispanic		Other or Unknown		Female Subtotal	
Injecting Drug Use	<5	**	11	(9%)	<5	**	<5	**	21	(12%)
Blood Products	0	(0%)	0	(0%)	0	(0%)	0	(0%)	0	(0%)
Heterosexual	29	(78%)	87	(75%)	14	(70%)	5	(63%)	135	(75%)
<i>HRH</i>	27	(73%)	47	(41%)	12	(60%)	<5	**	88	(49%)
<i>PH-Female</i>	<5	**	40	(34%)	<5	**	<5	**	47	(26%)
Perinatal	0	(0%)	<5	**	<5	**	0	(0%)	<5	**
Undetermined*	5	(14%)	17	(15%)	0	(0%)	0	(0%)	22	(12%)
Female Subtotal	37	(20%)	116	(64%)	20	(11%)	8	(4%)	181	(100%)
TOTAL	White		Black		Hispanic		Other or Unknown		Risk Total	
Male-Male sex	288	(74%)	64	(24%)	45	(46%)	<5	**	401	(52%)
Injecting Drug Use	8	(2%)	24	(9%)	8	(8%)	<5	**	43	(6%)
Male-Male Sex/IDU	20	(5%)	11	(4%)	<5	**	<5	**	35	(5%)
Blood Products	<5	**	0	(0%)	0	(0%)	0	(0%)	<5	**
Heterosexual	35	(9%)	104	(39%)	21	(21%)	7	(37%)	167	(22%)
<i>HRH</i>	33	(9%)	64	(24%)	19	(19%)	<5	**	120	(16%)
<i>PH-Female</i>	<5	**	40	(15%)	<5	**	<5	**	47	(6%)
Perinatal	<5	**	<5	**	<5	**	0	(0%)	7	(1%)
Undetermined	34	(9%)	59	(22%)	20	(20%)	<5	**	116	(15%)
<i>PH-Male</i>	21	(5%)	38	(14%)	18	(18%)	<5	**	80	(10%)
<i>Unknown</i>	13	(3%)	21	(8%)	<5	**	0	(0%)	36	(5%)
RACE TOTAL	388	(50%)	266	(35%)	98	(13%)	19	(2%)	771	(100%)

*In the male subset all cases in the heterosexual category are HRH because the PH-Female category is not applicable to males and, likewise, in the female subset, all cases in the undetermined category are of unknown risk because the PH-Male category is not applicable to females. See page 4 for additional details about the changes to Michigan's risk categories.

TABLE 3. New Diagnoses, Deaths, and Prevalence of HIV/AIDS in KENT CO. by Year


Year	<i>HIV/AIDS</i>			<i>AIDS</i>		
	New HIV Diagnoses	Deaths	Prevalence	New AIDS Diagnoses	Deaths	Prevalence
1983	1	0	1	1	0	1
1984	3	1	3	2	1	2
1985	22	3	22	7	3	6
1986	25	2	45	3	2	7
1987	39	5	79	10	5	12
1988	47	10	116	18	9	21
1989	68	20	164	35	20	36
1990	65	16	213	33	16	53
1991	77	28	262	53	26	80
1992	59	29	292	67	28	119
1993	50	47	295	52	47	124
1994	49	43	301	52	43	133
1995	47	37	311	48	36	145
1996	43	33	321	36	30	151
1997	34	28	327	37	27	161
1998	34	18	343	23	17	167
1999	45	12	376	25	12	180
2000	55	14	417	32	12	200
2001	40	14	443	29	10	219
2002	52	9	486	27	9	237
2003	50	7	529	31	5	263
2004	53	8	574	31	6	288
2005	60	11	623	42	8	322
2006	52	13	662	45	10	357
2007	45	11	696	38	11	384
2008	36	6	726	24	6	402
2009	49	4	771	31	3	430
TOTAL	1,200	429		832	402	

The prevalence of HIV in Michigan has steadily increased, since persons with HIV are living longer. This is largely due to improved anti-retroviral therapy.

The increase in HIV prevalence is also reflected in Figure 1 on page 5, which shows that the number of persons diagnosed, while stable for the last several years, is greater than the number of deaths each year. This directly contributes to the increase in prevalence. The current reported prevalence of HIV/AIDS in KENT CO. is 771. The prevalence of AIDS, which is a subset of HIV/AIDS prevalence, is 430.

As implied, the HIV/AIDS section displays data on all persons with HIV, including those with AIDS, as well as those who have not been diagnosed with AIDS. Thus, persons represented in the AIDS section are also represented in the HIV/AIDS section. The number of reported deaths includes deaths directly attributable to presence of HIV/AIDS as well as deaths due to other causes.

NOTE: Reporting for recent years may not be complete. Data are not adjusted to account for reporting delays.

FIGURE 1. New Diagnoses, Deaths, and Prevalence of HIV/AIDS in KENT CO. by Year**TABLE 4. Comparison of HIV/AIDS Prevalence in KENT CO. to State**

Residence at Diagnosis	Estimated Prevalence	Reported Prevalence				Cumulative AIDS
		HIV, not AIDS	AIDS	Total	Rate	
Kent Co.	1,020	341	430	771	127	832
Michigan Total	18,800	6,695	7,676	14,371	144	16,678

ADDITIONAL NOTES

Prevalence Estimate Calculations

HIV prevalence estimates include all persons living in KENT CO. at diagnosis of HIV or AIDS and incorporates an approximation of those not reported or not yet diagnosed. All estimates are rounded to the nearest ten, and the minimum estimate given is 10. Prevalence estimates were most recently revised in January 2010, so the estimates in this report may vary in comparison to reports published before the revision.

Risk Groups

In June 2007, the CSTE issued a position statement proposing the creation of a "presumed heterosexual" category. As a result, the following changes were made to Michigan's HIV risk categories.

Heterosexual risk is made up of two sub-categories, (1) HRH [High Risk Heterosexual]: Males and females whose sexual partners are known to be HIV-infected or at high risk for HIV. The partners meet one of the following criteria: a history of sexual contact with bi-sexual males (for females), IDU, hemophiliacs, HIV+ transfusion recipients or other HIV+ persons of unknown risk and (2) PH [Presumed Heterosexual]-female: Females who are not injection drug users and whose only reported risk is heterosexual contact, but their male partners' risk and HIV status are unknown.

Undetermined risk is made up of two sub-categories, (1) PH [Presumed Heterosexual]-Male: Males whose only reported risk is heterosexual contact, and their female partners' risk and HIV status are unknown and (2) Unknown: Males and Females with no identified risk.



Michigan Department of Community Health
HIV/STD/VH/TB Epidemiology Section
Division of Communicable Disease
Bureau of Epidemiology

Jennifer M. Granholm, Governor
Janet Olszewski, Director
<http://www.michigan.gov/hivstd>

Lansing Office 201 Townsend St., 5th Floor
Lansing, MI 48913
517-335-8165

Detroit Office 1151 Taylor St., Room 211B
Detroit, MI 48202
313-876-0353