Nausea & Vomiting

Pre-Medical Control

MFR/EMT/SPECIALIST/PARAMEDIC

1. Follow **General Pre-hospital Care Protocol**.

SPECIALIST/PARAMEDIC

2. Administer NS IV/IO fluid bolus up to 1 liter, wide open.
3. Hypotensive patients should receive additional IV/IO fluid boluses, as indicated by hemodynamic state. Continue IV/IO fluid bolus to a maximum of 2 liters.

PARAMEDIC

4. Administer Ondansetron (Zofran) 4mg IV/IM.

Post-Medical Control

5. Repeat Ondansetron (Zofran) 4mg IV/IM.
6. For Pediatric Patients refer to **Pediatric Nausea/Vomiting Protocol.**


Follow **General Pre-hospital Care Protocol**
Administer NS IV/IO fluid bolus up to 1 liter, wide open. Hypotensive patients should receive additional IV/IO fluid boluses, as indicated by hemodynamic state. Continue IV/IO fluid boluses to a maximum of 2 liters."
Administer Ondansetron (Zofran) 4 mg IV/IM
Contact Medical Control
Repeat Ondansetron (Zofran) 4 mg IV/IM
For Pediatric Patients refer to **Pediatric Nausea/Vomiting Protocol.**