

TABLE 1. Demographic Information on Prevalent HIV/AIDS Cases Living in WAYNE CO. (excluding Detroit) at Diagnosis

	<i>EST PREV*</i>	<i>REPORTED PREVALENCE[†]</i>						<i>CENSUS 2008 ESTIMATES[§]</i>	
		<i>HIV, not AIDS</i>		<i>AIDS</i>		<i>Total</i>		<i>Number</i>	<i>Percent</i>
<i>RACE/ ETHNICITY[‡]</i>	<i>Number</i>	<i>Number</i>	<i>Percent</i>	<i>Number</i>	<i>Percent</i>	<i>Number</i>	<i>Percent</i>		
White	910	302	(48%)	398	(49%)	700	(48%)	77	905,033 (87%)
Black	880	295	(47%)	380	(46%)	675	(47%)	1,735	38,904 (4%)
Hispanic	40	14	(2%)	19	(2%)	33	(2%)	92	35,982 (3%)
Asian/PI	10	<5	**	7	(1%)	11	(1%)	28	39,056 (4%)
Am Indian/AN	0	0	(0%)	0	(0%)	0	(0%)	**	3,494 (<1%)
Multi/Unk	40	12	(2%)	15	(2%)	27	(2%)	N/A	15,398 (1%)
<i>SEX & RACE</i>									
Males	1,500	475	(76%)	670	(82%)	1,145	(79%)	223	512,762 (49%)
White Males	820	256	(41%)	369	(45%)	625	(43%)	N/A	N/A N/A
Black Males	610	196	(31%)	267	(33%)	463	(32%)	N/A	N/A N/A
Hispanic Males	30	11	(2%)	15	(2%)	26	(2%)	N/A	N/A N/A
Other Males	40	12	(2%)	19	(2%)	31	(2%)	N/A	N/A N/A
Females	390	152	(24%)	149	(18%)	301	(21%)	57	525,105 (51%)
White Females	100	46	(7%)	29	(4%)	75	(5%)	N/A	N/A N/A
Black Females	280	99	(16%)	113	(14%)	212	(15%)	N/A	N/A N/A
Hispanic Fmls	10	<5	**	<5	**	7	(<1%)	N/A	N/A N/A
Other Females	10	<5	**	<5	**	7	(<1%)	N/A	N/A N/A
<i>RISK*</i>									
Male-Male Sex	990	310	(49%)	446	(54%)	756	(52%)	N/A	N/A N/A
Injection Drug Use	180	55	(9%)	79	(10%)	134	(9%)	N/A	N/A N/A
MSM/IDU	80	19	(3%)	44	(5%)	63	(4%)	N/A	N/A N/A
Blood Products	10	<5	**	5	(1%)	9	(1%)	N/A	N/A N/A
Heterosexual	300	101	(16%)	126	(15%)	227	(16%)	N/A	N/A N/A
HRH	200	60	(10%)	92	(11%)	152	(11%)	N/A	N/A N/A
PH-Female	100	41	(7%)	34	(4%)	75	(5%)	N/A	N/A N/A
Perinatal	10	6	(1%)	<5	**	9	(1%)	N/A	N/A N/A
Undetermined	320	132	(21%)	116	(14%)	248	(17%)	N/A	N/A N/A
PH-Male	160	51	(8%)	68	(8%)	119	(8%)	N/A	N/A N/A
Unknown	170	81	(13%)	48	(6%)	129	(9%)	N/A	N/A N/A
<i>AGE AT HIV DIAGNOSIS</i>									
0 - 12 years	10	7	(1%)	<5	**	11	(1%)	N/A	N/A N/A
13 - 19 years	70	38	(6%)	17	(2%)	55	(4%)	N/A	N/A N/A
20 - 24 years	200	88	(14%)	68	(8%)	156	(11%)	N/A	N/A N/A
25 - 29 years	310	103	(16%)	135	(16%)	238	(16%)	N/A	N/A N/A
30 - 39 years	660	191	(30%)	317	(39%)	508	(35%)	N/A	N/A N/A
40 - 49 years	420	128	(20%)	196	(24%)	324	(22%)	N/A	N/A N/A
50 - 59 years	160	56	(9%)	70	(9%)	126	(9%)	N/A	N/A N/A
60 years and over	30	13	(2%)	12	(1%)	25	(2%)	N/A	N/A N/A
Unspecified	10	<5	**	0	(0%)	<5	**	N/A	N/A N/A
TOTAL	1,890	627 (100%)		819 (100%)		1,446 (100%)		139	1,037,867 (100%)

*See page 5 for descriptions of prevalence estimate calculations and new risk category groupings. Risk categories used in Michigan are newly defined as of the July 2007 quarter.

[†] Includes reports that contain patient name or are otherwise unduplicated. <5 and ** = 1, 2, 3, or 4 cases.

[§] Estimated census data for sex by race are not available for sub-county areas.

[¶] To calculate "1 out of x" statements, divide the census number by total reported prevalence. For example, for non-Hispanic whites: 905,033/ 700 = 1,293. Thus, 1 out of every 1,293 non-Hispanic white persons in WAYNE CO. (excl. Detroit) are living with HIV. Rates and "1 out of x" statements are not reliable for <10 cases. Thus, rates for <10 cases are shown as "----".

[‡] In this report, persons described as white, black, Asian/Pacific Islander (PI), or American Indian/Alaska Native (AN) are all non-Hispanic; persons described as Hispanic might be of any race.

TABLE 2. Sex, Race, and Risk Among Prevalent HIV/AIDS Cases in WAYNE CO. (excl Detroit)

MALES	White	Black	Hispanic	Other or Unknown	Male Subtotal
Male-Male sex	460 (74%)	262 (57%)	18 (69%)	16 (52%)	756 (66%)
Injecting Drug Use	34 (5%)	43 (9%)	0 (0%)	<5 **	78 (7%)
Male-Male Sex/IDU	32 (5%)	28 (6%)	<5 **	<5 **	63 (6%)
Blood Products	7 (1%)	0 (0%)	0 (0%)	0 (0%)	7 (1%)
Heterosexual*	20 (3%)	18 (4%)	<5 **	0 (0%)	39 (3%)
Perinatal	<5 **	5 (1%)	0 (0%)	0 (0%)	6 (1%)
Undetermined	71 (11%)	107 (23%)	6 (23%)	12 (39%)	196 (17%)
<i>PH-Male</i>	44 (7%)	67 (14%)	<5 **	5 (16%)	119 (10%)
<i>Unknown</i>	27 (4%)	40 (9%)	<5 **	7 (23%)	77 (7%)
Male Subtotal	625 (55%)	463 (40%)	26 (2%)	31 (3%)	1,145 (100%)
FEMALES	White	Black	Hispanic	Other or Unknown	Female Subtotal
Injecting Drug Use	15 (20%)	40 (19%)	0 (0%)	<5 **	56 (19%)
Blood Products	<5 **	0 (0%)	<5 **	0 (0%)	<5 **
Heterosexual	52 (69%)	125 (59%)	5 (71%)	6 (86%)	188 (62%)
<i>HRH</i>	36 (48%)	69 (33%)	<5 **	5 (71%)	113 (38%)
<i>PH-Female</i>	16 (21%)	56 (26%)	<5 **	<5 **	75 (25%)
Perinatal	0 (0%)	<5 **	<5 **	0 (0%)	<5 **
Undetermined*	7 (9%)	45 (21%)	0 (0%)	0 (0%)	52 (17%)
Female Subtotal	75 (25%)	212 (70%)	7 (2%)	7 (2%)	301 (100%)
TOTAL	White	Black	Hispanic	Other or Unknown	Risk Total
Male-Male sex	460 (66%)	262 (39%)	18 (55%)	16 (42%)	756 (52%)
Injecting Drug Use	49 (7%)	83 (12%)	0 (0%)	<5 **	134 (9%)
Male-Male Sex/IDU	32 (5%)	28 (4%)	<5 **	<5 **	63 (4%)
Blood Products	8 (1%)	0 (0%)	<5 **	0 (0%)	9 (1%)
Heterosexual	72 (10%)	143 (21%)	6 (18%)	6 (16%)	227 (16%)
<i>HRH</i>	56 (8%)	87 (13%)	<5 **	5 (13%)	152 (11%)
<i>PH-Female</i>	16 (2%)	56 (8%)	<5 **	<5 **	75 (5%)
Perinatal	<5 **	7 (1%)	<5 **	0 (0%)	9 (1%)
Undetermined	78 (11%)	152 (23%)	6 (18%)	12 (32%)	248 (17%)
<i>PH-Male</i>	44 (6%)	67 (10%)	<5 **	5 (13%)	119 (8%)
<i>Unknown</i>	34 (5%)	85 (13%)	<5 **	7 (18%)	129 (9%)
RACE TOTAL	700 (48%)	675 (47%)	33 (2%)	38 (3%)	1,446 (100%)

*In the male subset all cases in the heterosexual category are HRH because the PH-Female category is not applicable to males and, likewise, in the female subset, all cases in the undetermined category are of unknown risk because the PH-Male category is not applicable to females. See page 5 for additional details about the changes to Michigan's risk categories.

**TABLE 3. Prevalent HIV/AIDS Cases in WAYNE CO.
(excluding Detroit) According to City of Residence at
Diagnosis**

CITY OF RESIDENCE AT DIAGNOSIS	EST PREV*	REPORTED PREVALENCE [†]			
		HIV, not AIDS	AIDS	Total	
				Number	Percent
Allen Park	20	7	6	13	(1%)
Belleville	80	28	34	62	(4%)
Brownstown Township	10	<5	<5	<5	**
Canton	70	26	25	51	(4%)
Dearborn	120	43	49	92	(6%)
Dearborn Heights	70	27	30	57	(4%)
Ecorse	40	14	17	31	(2%)
Flat Rock	10	<5	5	6	(<1%)
Garden City	30	10	10	20	(1%)
Gibraltar	10	0	<5	<5	**
Grosse Ile	10	<5	<5	<5	**
Grosse Pointe	30	8	16	24	(2%)
Grosse Pointe Farms	10	<5	<5	5	(<1%)
Grosse Pointe Park	10	<5	0	<5	**
Grosse Pointe Woods	10	<5	<5	5	(<1%)
Hamtramck	100	34	45	79	(5%)
Harper Woods	30	10	13	23	(2%)
Highland Park	280	92	124	216	(15%)
Inkster	170	51	81	132	(9%)
Lincoln Park	60	20	24	44	(3%)
Livonia	80	28	37	65	(4%)
Melvindale	20	6	6	12	(1%)
New Boston	10	<5	<5	5	(<1%)
Northville	20	7	7	14	(1%)
Plymouth	20	10	7	17	(1%)
Redford Township	110	38	46	84	(6%)
River Rouge	40	15	13	28	(2%)
Riverview	10	<5	5	9	(1%)
Rockwood	10	0	<5	<5	**
Romulus	70	23	28	51	(4%)
Southgate	30	12	8	20	(1%)
Taylor	110	34	48	82	(6%)
Trenton	20	<5	13	14	(1%)
Wayne	40	<5	23	27	(2%)
Westland	150	53	62	115	(8%)
Woodhaven	10	<5	0	<5	**
Wyandotte	40	9	19	28	(2%)
UNKNOWN	10	<5	<5	<5	**
TOTAL	1,890	627	819	1,446	(100%)

*See page 5 for descriptions of prevalence estimate calculations.

[†] Includes reports that contain patient name or are otherwise unduplicated. <5 and ** = 1, 2, 3, or 4 cases.

TABLE 4. New Diagnoses, Deaths, and Prevalence of HIV/AIDS in WAYNE CO. (excluding Detroit) by Year

Year	<i>HIV/AIDS</i>			<i>AIDS</i>		
	New HIV Diagnoses	Deaths	Prevalence	New AIDS Diagnoses	Deaths	Prevalence
1981	1	1	0	1	1	0
1982	0	0	0	0	0	0
1983	3	0	3	3	0	3
1984	5	1	7	4	1	6
1985	36	4	39	11	4	13
1986	47	14	72	14	14	13
1987	61	16	117	25	15	23
1988	88	23	182	49	21	51
1989	128	41	269	78	41	88
1990	138	40	367	77	38	127
1991	126	50	443	94	50	171
1992	134	61	516	117	58	230
1993	107	84	539	102	79	253
1994	127	83	583	100	77	276
1995	135	101	617	112	93	295
1996	73	63	627	67	60	302
1997	102	40	689	67	33	336
1998	90	31	748	69	26	379
1999	74	31	791	50	26	403
2000	93	31	853	67	26	444
2001	94	35	912	53	27	470
2002	69	23	958	58	19	509
2003	81	21	1,018	58	19	548
2004	97	28	1,087	59	24	583
2005	82	24	1,145	63	20	626
2006	86	25	1,206	72	23	675
2007	108	22	1,292	74	21	728
2008	90	28	1,354	68	27	769
2009	87	17	1,424	57	17	809
2010 [†]	24	2	1,446	11	1	819
TOTAL	2,386	940		1,680	861	


[†] Reporting for 2010 is incomplete at this time.

The prevalence of HIV in Michigan has steadily increased, since persons with HIV are living longer. This is largely due to improved anti-retroviral therapy.

The increase in HIV prevalence is also reflected in Figure 1 on page 5, which shows that the number of persons diagnosed, while stable for the last several years, is greater than the number of deaths each year. This directly contributes to the increase in prevalence. The current reported prevalence of HIV/AIDS in WAYNE CO. is 1,446. The prevalence of AIDS, which is a subset of HIV/AIDS prevalence, is 819

As implied, the HIV/AIDS section displays data on all persons with HIV, including those with AIDS, as well as those who have not been diagnosed with AIDS. Thus, persons represented in the AIDS section are also represented in the HIV/AIDS section. The number of reported deaths includes deaths directly attributable to presence of HIV/AIDS as well as deaths due to other causes.

NOTE: Reporting for recent years may not be complete. Data are not adjusted to account for reporting delays.

FIGURE 1. New Diagnoses, Deaths, and Prevalence of HIV/AIDS in WAYNE CO. (excl Detroit) by Year**TABLE 5. Comparison of HIV/AIDS Prevalence in WAYNE CO. (excl Detroit) to State**

<i>Residence at Diagnosis</i>	<i>Estimated Prevalence</i>	<i>Reported Prevalence</i>				<i>Cumulative AIDS</i>
		<i>HIV, not AIDS</i>	<i>AIDS</i>	<i>Total</i>	<i>Rate</i>	
Wayne Co.	1,890	627	819	1,446	139	1,680
Michigan Total	18,800	6,776	7,798	14,574	146	16,811

ADDITIONAL NOTES

Prevalence Estimate Calculations

HIV prevalence estimates include all persons living in WAYNE CO. at diagnosis of HIV or AIDS and incorporates an approximation of those not reported or not yet diagnosed. All estimates are rounded to the nearest ten, and the minimum estimate given is 10. Prevalence estimates were most recently revised in January 2010, so the estimates in this report may vary in comparison to reports published before the revision.

Risk Groups

In June 2007, the CSTE issued a position statement proposing the creation of a "presumed heterosexual" category. As a result, the following changes were made to Michigan's HIV risk categories.

Heterosexual risk is made up of two sub-categories, (1) HRH [High Risk Heterosexual]: Males and females whose sexual partners are known to be HIV-infected or at high risk for HIV. The partners meet one of the following criteria: a history of sexual contact with bi-sexual males (for females), IDU, hemophiliacs, HIV+ transfusion recipients or other HIV+ persons of unknown risk and (2) PH [Presumed Heterosexual]-female: Females who are not injection drug users and whose only reported risk is heterosexual contact, but their male partners' risk and HIV status are unknown.

Undetermined risk is made up of two sub-categories, (1) PH [Presumed Heterosexual]-Male: Males whose only reported risk is heterosexual contact, and their female partners' risk and HIV status are unknown and (2) Unknown: Males and Females with no identified risk.



Michigan Department of Community Health
HIV/STD/VH/TB Epidemiology Section
Division of Communicable Disease
Bureau of Epidemiology

Jennifer M. Granholm, Governor
Janet Olszewski, Director
<http://www.michigan.gov/hivstd>

Lansing Office 201 Townsend St., 5th Floor
Lansing, MI 48913
517-335-8165

Detroit Office 1151 Taylor St., Room 211B
Detroit, MI 48202
313-876-0353