WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	39	*	*	*	*	*	*	*
0102	166	89.8	0.6	3.0	0.0	0.0	6.6	0.0
0103	156	91.7	1.3	3.8	0.0	0.0	3.2	0.0
0104	54	*	*	*	*	*	*	*
0250	51	*	*	*	*	*	*	*
0251	40	*	*	*	*	*	*	*
0252	43	*	*	*	*	*	*	*
0253	64	*	*	*	*	*	*	*
0401	188	92.0	1.1	3.2	0.0	0.5	2.7	0.5
0402	141	80.9	0.0	3.5	0.0	1.4	14.2	0.0
0403	49	*	*	*	*	*	*	*
0404	52	*	*	*	*	*	*	*
0501	192	94.8	0.0	2.1	0.0	1.6	1.6	0.0
0503	39	*	*	*	*	*	*	*
0504	176	86.4	1.7	1.1	1.1	0.6	9.1	0.0
0505	115	86.1	0.9	3.5	0.0	0.9	8.7	0.0
0506	68	*	*	*	*	*	*	*
0507	25	*	*	*	*	*	*	*
0508	16	*	*	*	*	*	*	*
0509	3	*	*	*	*	*	*	*
0650	396	79.5	4.0	4.3	2.3	3.8	6.1	0.0
0651	213	90.6	0.0	4.7	0.0	0.5	2.8	1.4
0652	96	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0653	145	94.5	0.0	2.1	0.0	0.0	3.4	0.0
0654	140	94.3	1.4	1.4	0.7	0.0	1.4	0.7
0655	120	90.8	0.8	3.3	0.0	0.0	4.2	0.8
0657	25	*	*	*	*	*	*	*
0701	140	89.3	2.1	0.7	0.0	5.0	2.9	0.0
0702	22	*	*	*	*	*	*	*
0703	86	*	*	*	*	*	*	*
0704	18	*	*	*	*	*	*	*
0705	52	*	*	*	*	*	*	*
0706	2	*	*	*	*	*	*	*
0850	310	84.8	5.5	5.5	0.0	1.0	2.9	0.3
0851	244	95.5	0.8	3.3	0.0	0.4	0.0	0.0
0901	600	79.7	3.8	9.5	0.0	0.8	5.3	0.8
0902	12	*	*	*	*	*	*	*
1051	503	86.1	1.4	6.2	1.0	0.8	4.4	0.2
1101	635	37.5	51.2	6.9	0.2	0.5	3.0	0.8
1102	271	76.8	11.1	6.3	0.4	0.4	4.8	0.4
1106	29	*	*	*	*	*	*	*
1250	337	87.5	0.3	10.7	0.0	0.3	1.2	0.0
1251	298	93.0	0.0	1.7	0.0	0.3	4.0	1.0
1252	201	63.2	0.0	34.3	0.0	0.5	2.0	0.0
1253	288	78.5	6.9	10.4	0.0	0.0	2.8	1.4
1350	828	55.7	21.4	7.9	0.2	7.6	6.5	0.7

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1351	43	*	*	*	*	*	*	*
1352	132	62.9	27.3	6.1	0.0	0.0	3.8	0.0
1401	81	*	*	*	*	*	*	*
1402	54	*	*	*	*	*	*	*
1403	6	*	*	*	*	*	*	*
1501	268	49.6	4.5	37.7	0.0	0.4	3.0	4.9
1503	410	61.2	6.3	24.1	0.2	0.7	6.6	0.7
1504	763	41.8	5.1	44.8	0.0	4.6	2.6	1.0
1505	144	84.7	3.5	8.3	0.0	0.7	1.4	1.4
1506	208	75.0	1.0	18.3	0.0	0.0	3.4	2.4
1507	219	87.2	1.4	5.9	0.0	1.4	3.7	0.5
1508	40	*	*	*	*	*	*	*
1509	149	48.3	3.4	40.9	0.0	0.0	3.4	4.0
1511	250	88.8	1.2	3.2	0.0	2.4	3.6	0.8
1512	123	91.9	0.0	1.6	1.6	0.8	3.3	0.8
1513	33	*	*	*	*	*	*	*
1514	55	*	*	*	*	*	*	*
1701	240	56.3	0.4	1.3	20.4	0.0	21.7	0.0
1950	208	93.3	0.0	3.8	0.5	0.5	1.0	1.0
1952	43	*	*	*	*	*	*	*
1953	103	91.3	0.0	4.9	0.0	0.0	2.9	1.0
1954	205	84.9	0.5	13.7	0.0	0.5	0.5	0.0
1955	28	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1957	176	83.5	0.6	14.2	0.0	0.0	1.7	0.0
1964	13	*	*	*	*	*	*	*
2150	242	86.8	0.4	2.9	0.4	1.2	7.9	0.4
2151	80	*	*	*	*	*	*	*
2251	144	93.1	0.0	2.8	1.4	0.0	2.8	0.0
2252	73	*	*	*	*	*	*	*
2554	29	*	*	*	*	*	*	*
2555	2,823	47.5	40.5	4.8	0.1	0.5	6.2	0.4
2556	77	*	*	*	*	*	*	*
2557	2	*	*	*	*	*	*	*
2558	2	*	*	*	*	*	*	*
2559	6	*	*	*	*	*	*	*
3001	378	13.0	67.5	14.0	0.3	1.3	3.4	0.5
3002	89	*	*	*	*	*	*	*
3004	49	*	*	*	*	*	*	*
3007	14	*	*	*	*	*	*	*
3009	3	*	*	*	*	*	*	*
3201	174	93.7	0.0	4.0	0.0	0.0	0.6	1.7
3301	1,479	38.9	30.0	15.1	0.2	7.0	8.6	0.3
3304	31	*	*	*	*	*	*	*
3305	48	*	*	*	*	*	*	*
3306	83	*	*	*	*	*	*	*
3308	25	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3309	17	*	*	*	*	*	*	*
3312	148	39.9	27.0	21.6	1.4	2.0	6.8	1.4
3550	30	*	*	*	*	*	*	*
3551	12	*	*	*	*	*	*	*
3552	19	*	*	*	*	*	*	*
3701	379	92.3	0.0	5.8	0.0	0.3	1.6	0.0
3850	1,077	70.7	16.2	5.9	0.2	0.7	6.2	0.1
3901	126	31.7	50.0	6.3	1.6	2.4	4.0	4.0
3902	65	*	*	*	*	*	*	*
3903	2	*	*	*	*	*	*	*
3950	653	60.8	23.9	7.7	0.0	1.4	5.5	0.8
4001	420	30.5	55.0	8.6	0.0	1.2	4.5	0.2
4002	269	49.4	26.8	16.7	0.0	0.7	6.3	0.0
4003	2	*	*	*	*	*	*	*
4004	4	*	*	*	*	*	*	*
4103	26	*	*	*	*	*	*	*
4106	810	16.2	43.8	33.6	0.1	0.7	4.7	0.9
4107	1,175	49.0	24.9	16.9	0.3	2.1	6.6	0.2
4109	348	85.3	1.1	8.6	0.0	1.4	3.4	0.0
4112	196	1.5	0.5	96.4	0.0	0.5	0.0	1.0
4113	227	4.8	19.4	74.0	0.0	0.4	1.3	0.0
4115	1,188	39.1	24.0	17.3	0.5	12.7	6.0	0.4
4351	213	91.1	0.0	5.6	0.0	0.5	2.3	0.5

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4352	63	*	*	*	*	*	*	*
4353	147	70.1	0.0	25.9	0.7	0.0	2.0	1.4
4354	248	89.9	4.0	2.8	0.0	0.0	3.2	0.0
4355	128	78.1	0.8	8.6	3.1	0.8	8.6	0.0
4356	178	89.3	0.6	7.9	0.0	0.0	2.2	0.0
4358	50	*	*	*	*	*	*	*
4359	158	69.0	0.0	27.2	0.0	0.6	1.3	1.9
4360	75	*	*	*	*	*	*	*
4361	100	95.0	0.0	0.0	1.0	0.0	3.0	1.0
4362	96	*	*	*	*	*	*	*
4363	330	93.6	0.0	3.6	0.0	0.0	2.1	0.6
4401	357	89.4	0.0	5.0	0.0	0.3	4.8	0.6
4501	331	92.7	1.5	3.3	0.0	0.6	1.8	0.0
4601	486	76.5	1.9	15.0	0.0	0.2	5.3	0.8
4602	13	*	*	*	*	*	*	*
4701	370	88.6	0.3	3.2	0.5	2.4	2.7	2.2
5050	613	51.9	38.7	3.3	0.0	1.8	3.9	0.5
5051	945	51.3	36.9	3.9	0.0	2.6	4.9	0.3
5052	951	43.0	35.2	1.4	0.1	14.8	4.8	0.5
5055	408	96.6	0.2	0.2	0.0	2.2	0.2	0.5
5101	672	69.3	10.0	14.9	0.0	1.5	4.3	0.0
5202	235	85.1	0.9	3.8	0.0	1.3	7.7	1.3
5203	4	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
5204	4	*	*	*	*	*	*	*
5601	407	93.9	0.5	1.7	0.2	0.7	2.5	0.5
5801	591	82.7	7.4	5.1	0.2	0.3	3.2	1.0
5803	45	*	*	*	*	*	*	*
5804	8	*	*	*	*	*	*	*
5806	11	*	*	*	*	*	*	*
6101	13	*	*	*	*	*	*	*
6102	37	*	*	*	*	*	*	*
6103	1,447	54.2	30.1	8.2	0.3	0.6	6.2	0.4
6105	8	*	*	*	*	*	*	*
6301	1,531	41.3	29.8	21.2	0.1	2.7	4.1	0.7
6312	862	33.5	57.5	3.7	0.1	2.2	2.7	0.2
6316	433	66.1	22.4	6.5	0.2	3.0	1.4	0.5
6320	35	*	*	*	*	*	*	*
6325	33	*	*	*	*	*	*	*
6330	114	24.6	54.4	18.4	0.0	0.9	1.8	0.0
6335	320	78.8	14.7	2.5	0.3	2.5	1.3	0.0
6340	14	*	*	*	*	*	*	*
6345	11	*	*	*	*	*	*	*
6360	110	21.8	62.7	10.0	0.0	1.8	3.6	0.0
7304	53	*	*	*	*	*	*	*
7306	907	44.1	35.0	14.8	0.0	0.6	4.7	0.9
7308	39	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
7310	10	*	*	*	*	*	*	*
7401	799	83.2	5.1	5.3	0.0	0.6	5.1	0.6
7601	230	94.3	0.4	3.5	0.0	0.0	1.7	0.0
7850	384	86.7	0.3	7.6	0.0	0.0	4.7	0.5
7852	10	*	*	*	*	*	*	*
8101	696	5.9	90.9	0.7	0.0	0.4	1.7	0.3
8102	155	8.4	9.0	79.4	0.0	0.0	2.6	0.6
8105	603	31.2	61.7	3.6	0.0	0.8	2.3	0.3
8106	153	23.5	66.7	4.6	0.7	0.0	4.6	0.0
8150	11	*	*	*	*	*	*	*
8151	1,156	37.1	39.0	9.5	0.1	4.2	9.6	0.4
8152	18	*	*	*	*	*	*	*
8153	2	*	*	*	*	*	*	*
8155	11	*	*	*	*	*	*	*
8157	3	*	*	*	*	*	*	*
8201	921	48.5	36.6	5.3	0.4	3.0	5.6	0.4
8202	1,341	88.6	7.9	1.7	0.0	0.7	0.7	0.4
8203	181	57.5	29.3	4.4	0.0	1.1	6.6	1.1
8250	285	23.5	64.9	3.9	0.0	1.1	6.7	0.0
8252	57	*	*	*	*	*	*	*
8255	876	55.8	29.0	7.2	0.1	3.4	3.9	0.6
8257	673	52.5	15.5	0.7	0.1	29.7	1.0	0.4
8263	261	68.6	16.1	9.6	0.0	1.9	3.4	0.4

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8264	252	63.5	27.0	3.6	0.0	2.4	2.8	0.8
8266	9	*	*	*	*	*	*	*
8267	316	58.9	18.7	17.1	0.3	0.0	3.8	0.6
8901	402	2.2	93.5	1.2	0.0	1.0	1.7	0.2
8902	489	3.5	92.4	2.0	0.0	0.2	1.8	0.0
8903	411	2.7	92.7	1.5	0.0	0.7	1.9	0.5
8904	840	1.9	93.7	1.1	0.1	0.7	2.4	0.1
8905	561	59.5	28.3	9.3	0.0	0.9	1.6	0.4
8908	18	*	*	*	*	*	*	*
8909	189	1.1	95.2	1.1	0.0	0.5	1.6	0.5
9016	7	*	*	*	*	*	*	*
9018	3	*	*	*	*	*	*	*
9045	598	9.5	27.8	60.0	0.0	0.2	2.2	0.3
9050	337	12.5	72.1	13.1	0.0	0.6	1.8	0.0
9051	639	8.6	7.7	81.7	0.0	0.5	1.6	0.0
9053	453	3.5	90.5	1.1	0.0	3.1	1.3	0.4
9055	13	*	*	*	*	*	*	*
9058	370	23.5	63.2	0.8	0.0	11.4	0.8	0.3
9066	425	1.9	97.6	0.0	0.0	0.2	0.2	0.0
9068	324	2.8	96.0	0.6	0.0	0.0	0.3	0.3
9069	1,007	1.0	97.5	0.3	0.0	0.4	0.6	0.2
9070	385	1.8	93.5	0.0	0.0	3.6	1.0	0.0
9072	670	2.1	95.1	0.3	0.0	0.3	1.9	0.3

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9080	1,043	21.3	75.6	1.6	0.0	0.0	1.2	0.2
9081	16	*	*	*	*	*	*	*
9082	55	*	*	*	*	*	*	*
9083	20	*	*	*	*	*	*	*
9084	13	*	*	*	*	*	*	*
9085	3	*	*	*	*	*	*	*
Michigan	**57,669**	**52.1**	**30.4**	**10.4**	**0.3**	**2.3**	**4.0**	**0.5**
National (Prior Year)	**120,297**	**55.2**	**27.3**	**9.5**	**1.5**	**2.3**	**3.9**	**0.3**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	39	*	*	*	*	*	*	*
0102	166	0.0	4.8	10.2	65.1	19.3	0.6	0.0
0103	156	0.0	1.9	7.7	66.7	22.4	1.3	0.0
0104	54	*	*	*	*	*	*	*
0250	51	*	*	*	*	*	*	*
0251	40	*	*	*	*	*	*	*
0252	43	*	*	*	*	*	*	*
0253	64	*	*	*	*	*	*	*
0401	188	0.0	2.1	11.2	64.4	21.3	1.1	0.0
0402	141	0.0	0.7	5.0	75.2	18.4	0.7	0.0
0403	49	*	*	*	*	*	*	*
0404	52	*	*	*	*	*	*	*
0501	192	0.0	3.1	8.9	64.1	23.4	0.5	0.0
0503	39	*	*	*	*	*	*	*
0504	176	0.0	1.1	12.5	63.6	22.2	0.6	0.0
0505	115	0.0	2.6	10.4	52.2	33.0	1.7	0.0
0506	68	*	*	*	*	*	*	*
0507	25	*	*	*	*	*	*	*
0508	16	*	*	*	*	*	*	*
0509	3	*	*	*	*	*	*	*
0650	396	0.0	2.8	7.6	65.7	23.0	0.8	0.3
0651	213	0.5	2.8	11.3	66.7	16.9	1.9	0.0
0652	96	*	*	*	*	*	*	*
0653	145	0.0	3.4	9.0	64.8	22.1	0.7	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0654	140	0.0	4.3	10.7	56.4	27.9	0.7	0.0
0655	120	0.0	3.3	12.5	63.3	20.8	0.0	0.0
0657	25	*	*	*	*	*	*	*
0701	140	0.0	2.1	9.3	58.6	28.6	1.4	0.0
0702	22	*	*	*	*	*	*	*
0703	86	*	*	*	*	*	*	*
0704	18	*	*	*	*	*	*	*
0705	52	*	*	*	*	*	*	*
0706	2	*	*	*	*	*	*	*
0850	310	0.0	1.6	7.7	62.6	26.5	1.0	0.6
0851	244	0.0	4.5	10.2	68.0	16.4	0.8	0.0
0901	600	0.2	2.5	7.8	66.8	21.0	1.7	0.0
0902	12	*	*	*	*	*	*	*
1051	503	0.0	2.8	7.2	61.8	25.8	1.8	0.6
1101	635	0.2	5.2	8.2	62.5	22.5	1.1	0.3
1102	271	0.4	4.1	9.6	65.7	17.7	2.2	0.4
1106	29	*	*	*	*	*	*	*
1250	337	0.0	2.4	8.0	62.0	24.3	3.0	0.3
1251	298	0.0	3.0	10.1	68.8	17.4	0.7	0.0
1252	201	0.0	4.5	15.9	57.7	20.9	1.0	0.0
1253	288	0.0	2.1	10.4	67.7	18.8	1.0	0.0
1350	828	0.1	3.5	8.8	63.5	22.6	1.4	0.0
1351	43	*	*	*	*	*	*	*
1352	132	0.0	3.0	7.6	70.5	16.7	2.3	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1401	81	*	*	*	*	*	*	*
1402	54	*	*	*	*	*	*	*
1403	6	*	*	*	*	*	*	*
1501	268	0.0	3.0	12.3	63.4	19.4	1.9	0.0
1503	410	0.0	5.9	8.3	60.7	23.9	1.2	0.0
1504	763	0.1	4.1	7.1	61.7	25.3	1.7	0.0
1505	144	0.0	0.7	7.6	63.2	27.8	0.7	0.0
1506	208	0.0	2.9	10.6	63.0	22.6	1.0	0.0
1507	219	0.0	1.8	6.4	64.8	25.1	1.8	0.0
1508	40	*	*	*	*	*	*	*
1509	149	0.0	4.7	9.4	57.7	22.8	5.4	0.0
1511	250	0.8	2.4	12.0	62.8	20.8	1.2	0.0
1512	123	0.0	1.6	6.5	66.7	25.2	0.0	0.0
1513	33	*	*	*	*	*	*	*
1514	55	*	*	*	*	*	*	*
1701	240	0.0	0.4	7.1	63.3	27.5	1.7	0.0
1950	208	0.0	2.4	12.0	67.3	17.3	1.0	0.0
1952	43	*	*	*	*	*	*	*
1953	103	1.0	1.0	9.7	70.9	17.5	0.0	0.0
1954	205	0.0	3.4	5.9	62.4	26.3	2.0	0.0
1955	28	*	*	*	*	*	*	*
1957	176	0.0	4.5	10.2	64.2	18.8	2.3	0.0
1964	13	*	*	*	*	*	*	*
2150	242	0.0	1.7	9.9	63.2	24.4	0.8	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2151	80	*	*	*	*	*	*	*
2251	144	0.0	3.5	8.3	56.9	28.5	2.1	0.7
2252	73	*	*	*	*	*	*	*
2554	29	*	*	*	*	*	*	*
2555	2,823	0.2	2.9	9.5	65.4	20.7	1.2	0.1
2556	77	*	*	*	*	*	*	*
2557	2	*	*	*	*	*	*	*
2558	2	*	*	*	*	*	*	*
2559	6	*	*	*	*	*	*	*
3001	378	0.8	5.0	10.1	64.3	18.0	1.6	0.3
3002	89	*	*	*	*	*	*	*
3004	49	*	*	*	*	*	*	*
3007	14	*	*	*	*	*	*	*
3009	3	*	*	*	*	*	*	*
3201	174	0.0	2.3	10.3	64.4	23.0	0.0	0.0
3301	1,479	0.3	3.0	7.6	62.4	24.5	1.9	0.2
3304	31	*	*	*	*	*	*	*
3305	48	*	*	*	*	*	*	*
3306	83	*	*	*	*	*	*	*
3308	25	*	*	*	*	*	*	*
3309	17	*	*	*	*	*	*	*
3312	148	0.0	2.0	3.4	62.2	29.7	2.7	0.0
3550	30	*	*	*	*	*	*	*
3551	12	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3552	19	*	*	*	*	*	*	*
3701	379	0.0	3.2	10.3	68.1	17.7	0.8	0.0
3850	1,077	0.2	3.5	9.3	67.3	18.4	1.1	0.2
3901	126	0.0	2.4	9.5	66.7	19.8	1.6	0.0
3902	65	*	*	*	*	*	*	*
3903	2	*	*	*	*	*	*	*
3950	653	0.0	3.5	9.5	61.4	23.7	1.8	0.0
4001	420	0.2	4.8	9.8	62.9	20.2	2.1	0.0
4002	269	0.0	5.2	8.6	59.1	25.7	1.5	0.0
4003	2	*	*	*	*	*	*	*
4004	4	*	*	*	*	*	*	*
4103	26	*	*	*	*	*	*	*
4106	810	0.6	4.7	10.0	59.1	23.3	2.1	0.1
4107	1,175	0.0	3.2	6.9	63.8	24.0	1.8	0.3
4109	348	0.3	3.2	6.6	63.8	23.9	2.3	0.0
4112	196	0.0	8.7	8.2	51.0	27.0	5.1	0.0
4113	227	0.0	3.1	7.5	51.1	32.2	1.3	4.8
4115	1,188	0.1	2.3	6.7	61.2	27.0	2.7	0.0
4351	213	0.0	2.8	14.6	64.8	17.4	0.5	0.0
4352	63	*	*	*	*	*	*	*
4353	147	0.0	3.4	8.8	68.7	17.7	1.4	0.0
4354	248	0.0	4.4	8.9	68.5	17.3	0.8	0.0
4355	128	0.0	7.0	15.6	56.3	19.5	0.8	0.8
4356	178	0.0	3.9	5.1	71.9	19.1	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4358	50	*	*	*	*	*	*	*
4359	158	0.0	3.2	6.3	59.5	26.6	4.4	0.0
4360	75	*	*	*	*	*	*	*
4361	100	0.0	1.0	9.0	61.0	29.0	0.0	0.0
4362	96	*	*	*	*	*	*	*
4363	330	0.0	1.8	8.8	68.5	20.6	0.3	0.0
4401	357	0.3	1.7	9.0	69.2	19.6	0.3	0.0
4501	331	0.3	2.1	8.8	64.4	21.8	2.4	0.3
4601	486	0.0	2.9	9.7	65.2	21.6	0.6	0.0
4602	13	*	*	*	*	*	*	*
4701	370	0.0	1.9	7.0	57.8	31.9	1.4	0.0
5050	613	0.0	1.3	6.2	62.5	29.4	0.7	0.0
5051	945	0.1	0.7	3.7	67.7	25.7	2.0	0.0
5052	951	0.1	2.9	5.6	64.6	24.8	2.0	0.0
5055	408	0.0	0.2	0.2	57.6	38.0	3.9	0.0
5101	672	0.0	0.7	6.1	61.9	28.7	2.5	0.0
5202	235	0.0	4.7	6.0	57.0	31.1	1.3	0.0
5203	4	*	*	*	*	*	*	*
5204	4	*	*	*	*	*	*	*
5601	407	0.0	1.5	9.6	64.1	23.3	1.2	0.2
5801	591	0.0	3.7	11.3	63.1	20.8	0.8	0.2
5803	45	*	*	*	*	*	*	*
5804	8	*	*	*	*	*	*	*
5806	11	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
6101	13	*	*	*	*	*	*	*
6102	37	*	*	*	*	*	*	*
6103	1,447	0.2	3.8	9.0	65.6	19.8	1.7	0.0
6105	8	*	*	*	*	*	*	*
6301	1,531	0.1	3.3	7.4	61.6	25.9	1.6	0.1
6312	862	0.0	1.6	5.2	58.4	32.4	2.4	0.0
6316	433	0.0	1.2	5.3	55.7	34.2	3.7	0.0
6320	35	*	*	*	*	*	*	*
6325	33	*	*	*	*	*	*	*
6330	114	0.0	1.8	9.6	61.4	27.2	0.0	0.0
6335	320	0.0	1.3	5.6	50.6	38.4	4.1	0.0
6340	14	*	*	*	*	*	*	*
6345	11	*	*	*	*	*	*	*
6360	110	0.0	0.9	8.2	73.6	16.4	0.9	0.0
7304	53	*	*	*	*	*	*	*
7306	907	0.2	2.6	7.2	67.5	20.9	1.5	0.0
7308	39	*	*	*	*	*	*	*
7310	10	*	*	*	*	*	*	*
7401	799	0.1	3.1	8.6	64.1	23.2	0.9	0.0
7601	230	0.0	6.1	7.8	60.9	25.2	0.0	0.0
7850	384	0.0	2.9	12.0	68.5	16.1	0.5	0.0
7852	10	*	*	*	*	*	*	*
8101	696	0.1	3.2	8.6	64.2	21.0	2.9	0.0
8102	155	0.0	2.6	3.9	58.1	31.6	3.2	0.6

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8105	603	0.2	2.7	5.3	60.2	29.2	2.5	0.0
8106	153	0.0	5.2	5.2	60.1	27.5	2.0	0.0
8150	11	*	*	*	*	*	*	*
8151	1,156	0.2	3.0	6.8	60.9	26.6	2.5	0.0
8152	18	*	*	*	*	*	*	*
8153	2	*	*	*	*	*	*	*
8155	11	*	*	*	*	*	*	*
8157	3	*	*	*	*	*	*	*
8201	921	0.2	3.1	6.8	64.1	24.2	1.5	0.0
8202	1,341	0.1	0.8	3.3	56.6	33.3	4.3	1.6
8203	181	0.6	2.2	8.3	58.0	28.2	2.8	0.0
8250	285	0.0	1.8	7.0	61.8	28.8	0.7	0.0
8252	57	*	*	*	*	*	*	*
8255	876	0.0	2.2	5.3	66.6	25.0	1.0	0.0
8257	673	0.1	1.5	4.2	58.8	30.6	4.6	0.1
8263	261	0.0	0.8	5.0	62.8	29.9	1.5	0.0
8264	252	0.0	2.0	7.9	63.9	24.2	2.0	0.0
8266	9	*	*	*	*	*	*	*
8267	316	0.0	3.2	5.1	63.9	25.6	2.2	0.0
8901	402	0.0	3.2	8.0	62.4	23.9	2.5	0.0
8902	489	0.6	1.6	9.4	63.2	22.7	2.2	0.2
8903	411	0.0	3.2	6.3	67.2	21.7	1.5	0.2
8904	840	0.0	3.8	8.0	68.5	18.0	1.8	0.0
8905	561	0.0	1.1	5.5	58.6	31.7	3.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8908	18	*	*	*	*	*	*	*
8909	189	0.5	3.7	8.5	60.3	25.9	1.1	0.0
9016	7	*	*	*	*	*	*	*
9018	3	*	*	*	*	*	*	*
9045	598	0.0	3.7	8.9	57.0	28.4	2.0	0.0
9050	337	0.6	2.7	8.0	61.4	24.0	3.3	0.0
9051	639	0.0	3.4	7.5	52.0	31.8	5.3	0.0
9053	453	0.0	2.9	5.1	71.7	18.1	2.2	0.0
9055	13	*	*	*	*	*	*	*
9058	370	0.0	1.4	3.5	58.4	33.0	3.8	0.0
9066	425	0.0	3.8	8.0	63.3	23.5	1.4	0.0
9068	324	0.0	2.5	10.5	63.6	21.6	1.9	0.0
9069	1,007	0.2	2.3	6.4	67.1	22.7	1.3	0.0
9070	385	0.5	4.2	7.5	64.4	21.8	1.6	0.0
9072	670	0.1	2.7	9.7	65.7	19.7	2.1	0.0
9080	1,043	0.1	1.9	6.3	63.3	26.2	2.2	0.0
9081	16	*	*	*	*	*	*	*
9082	55	*	*	*	*	*	*	*
9083	20	*	*	*	*	*	*	*
9084	13	*	*	*	*	*	*	*
9085	3	*	*	*	*	*	*	*
Michigan	**57,676**	**0.1**	**2.8**	**7.7**	**63.1**	**24.3**	**1.8**	**0.1**
National (Prior Year)	**120,303**	**0.2**	**3.4**	**9.1**	**63.1**	**22.7**	**1.5**	**0.1**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	29	*	*	6	*	*	4	*	7	*	35	*	35	*
0102	140	4.3 (61)	60.0 (61)	62	*	*	23	*	34	*	156	7.7 (82)	156	5.1 (6)
0103	124	3.2 (27)	61.3 (73)	88	*	*	17	*	39	*	148	10.1 (109)	148	14.9 (100)
0104	45	*	*	21	*	*	6	*	7	*	48	*	48	*
0250	48	*	*	21	*	*	5	*	10	*	50	*	50	*
0251	34	*	*	9	*	*	3	*	11	*	38	*	38	*
0252	43	*	*	38	*	*	3	*	10	*	40	*	40	*
0253	44	*	*	5	*	*	6	*	13	*	59	*	59	*
0401	175	2.9 (18)	55.4 (21)	140	17.9 (38)	57.9 (73)	25	*	47	*	172	4.1 (3)	172	7.6 (21)
0402	128	6.3 (105)	54.7 (16)	86	*	*	26	*	23	*	134	6.7 (47)	134	5.2 (7)
0403	48	*	*	33	*	*	9	*	12	*	47	*	47	*
0404	50	*	*	39	*	*	9	*	9	*	51	*	51	*
0501	106	2.8 (15)	63.2 (95)	40	*	*	28	*	32	*	180	3.3 (1)	180	7.8 (22)
0503	33	*	*	18	*	*	3	*	6	*	36	*	36	*
0504	154	5.2 (87)	48.7 (2)	72	*	*	31	*	24	*	167	5.4 (21)	167	9.0 (29)
0505	100	3.0 (21)	54.0 (12)	70	*	*	18	*	18	*	109	7.3 (70)	109	7.3 (19)
0506	48	*	*	16	*	*	3	*	12	*	65	*	65	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0507	17	*	*	10	*	*	3	*	6	*	25	*	25	*
0508	11	*	*	4	*	*	3	*	0	*	16	*	16	*
0509	3	*	*	3	*	*	0	*	0	*	3	*	3	*
0650	352	4.8 (77)	59.9 (57)	237	15.6 (18)	59.5 (77)	61	*	78	*	373	8.3 (89)	373	9.7 (37)
0651	203	4.9 (81)	64.0 (99)	172	19.2 (49)	54.1 (46)	35	*	29	*	196	7.1 (64)	196	6.6 (13)
0652	64	*	*	46	*	*	19	*	12	*	94	*	94	*
0653	130	2.3 (8)	60.0 (61)	82	*	*	12	*	27	*	138	4.3 (5)	139	10.8 (55)
0654	124	4.0 (54)	69.4 (111)	81	*	*	14	*	27	*	125	6.4 (39)	125	8.0 (23)
0655	110	7.3 (109)	53.6 (10)	41	*	*	14	*	23	*	112	7.1 (64)	113	12.4 (74)
0657	13	*	*	6	*	*	4	*	2	*	24	*	24	*
0701	79	*	*	34	*	*	14	*	32	*	132	5.3 (18)	132	9.1 (31)
0702	20	*	*	9	*	*	1	*	4	*	18	*	18	*
0703	51	*	*	28	*	*	14	*	15	*	73	*	73	*
0704	13	*	*	4	*	*	1	*	5	*	14	*	14	*
0705	32	*	*	8	*	*	6	*	15	*	46	*	46	*
0706	2	*	*	2	*	*	0	*	1	*	2	*	2	*
0850	195	5.6 (96)	55.4 (21)	137	16.8 (25)	51.1 (26)	60	*	69	*	290	11.4 (113)	291	17.5 (117)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0851	147	4.1 (58)	57.8 (40)	49	*	*	30	*	50	*	220	6.8 (52)	221	13.6 (87)
0901	482	5.0 (86)	60.0 (61)	363	22.6 (66)	51.5 (30)	79	*	86	*	565	7.1 (64)	565	10.8 (55)
0902	9	*	*	6	*	*	0	*	2	*	11	*	11	*
1051	463	5.4 (94)	50.1 (5)	381	16.5 (23)	52.8 (36)	77	*	68	*	485	4.5 (8)	486	10.3 (46)
1101	536	3.5 (36)	60.4 (65)	397	19.6 (50)	54.9 (55)	107	54.2 (19)	100	48.0 (29)	588	5.1 (16)	588	13.6 (87)
1102	225	2.7 (12)	58.7 (49)	169	13.0 (2)	61.5 (78)	34	*	63	*	247	4.9 (14)	247	4.5 (4)
1106	24	*	*	17	*	*	4	*	9	*	25	*	25	*
1250	247	4.0 (54)	59.9 (57)	93	*	*	48	*	59	*	310	10.3 (110)	310	6.5 (11)
1251	262	5.7 (100)	56.5 (29)	173	24.9 (73)	44.5 (6)	30	*	58	*	279	4.7 (10)	279	6.5 (11)
1252	145	2.8 (15)	60.7 (68)	72	*	*	22	*	35	*	187	6.4 (39)	187	17.1 (115)
1253	208	3.4 (29)	58.7 (49)	121	13.2 (3)	52.9 (38)	36	*	67	*	253	9.1 (102)	255	15.3 (104)
1350	701	2.1 (4)	61.2 (71)	383	17.8 (36)	54.6 (52)	108	48.1 (10)	144	44.4 (25)	765	6.4 (39)	761	9.6 (36)
1351	34	*	*	32	*	*	5	*	10	*	41	*	41	*
1352	65	*	*	34	*	*	12	*	27	*	120	8.3 (89)	120	4.2 (2)
1401	77	*	*	56	*	*	16	*	7	*	72	*	72	*
1402	50	*	*	38	*	*	12	*	6	*	49	*	49	*
1403	5	*	*	4	*	*	1	*	2	*	5	*	5	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1501	240	2.9 (18)	61.3 (73)	148	48.0 (77)	23.0 (2)	26	*	35	*	226	5.3 (18)	226	9.3 (33)
1503	352	2.8 (15)	60.5 (66)	209	15.3 (15)	56.0 (64)	36	*	64	*	374	4.3 (5)	374	9.4 (34)
1504	646	5.6 (96)	58.0 (43)	435	20.2 (55)	45.1 (8)	117	61.5 (27)	146	48.6 (31)	700	10.0 (108)	700	8.9 (27)
1505	116	3.4 (29)	57.8 (40)	70	*	*	15	*	40	*	133	5.3 (18)	133	10.5 (49)
1506	157	4.5 (68)	58.6 (48)	53	*	*	34	*	40	*	189	6.9 (57)	189	7.4 (20)
1507	159	3.8 (46)	54.1 (13)	79	*	*	24	*	37	*	200	8.5 (96)	200	13.5 (84)
1508	18	*	*	5	*	*	1	*	2	*	30	*	30	*
1509	111	0.9 (1)	65.8 (106)	62	*	*	15	*	18	*	121	9.1 (102)	121	13.2 (81)
1511	200	4.0 (54)	62.5 (88)	96	*	*	34	*	59	*	227	6.6 (44)	229	15.7 (110)
1512	90	*	*	27	*	*	18	*	22	*	114	7.0 (61)	114	15.8 (112)
1513	27	*	*	24	*	*	5	*	8	*	32	*	32	*
1514	48	*	*	36	*	*	8	*	24	*	53	*	53	*
1701	150	2.7 (12)	62.0 (83)	30	*	*	24	*	36	*	215	8.4 (95)	215	12.6 (76)
1950	195	1.0 (2)	64.1 (100)	159	23.3 (67)	54.1 (46)	22	*	43	*	189	7.9 (87)	190	14.7 (98)
1952	42	*	*	36	*	*	12	*	13	*	41	*	41	*
1953	94	*	*	85	*	*	14	*	29	*	101	11.9 (116)	101	14.9 (100)
1954	171	5.3 (91)	55.6 (23)	96	*	*	27	*	33	*	190	7.4 (73)	190	13.2 (81)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1955	25	*	*	12	*	*	6	*	4	*	25	*	25	*
1957	160	5.6 (96)	57.5 (35)	121	14.9 (9)	57.0 (69)	30	*	43	*	159	6.9 (57)	159	15.7 (110)
1964	12	*	*	9	*	*	3	*	4	*	13	*	13	*
2150	147	3.4 (29)	53.7 (11)	59	*	*	26	*	44	*	211	13.7 (118)	211	13.3 (83)
2151	36	*	*	22	*	*	9	*	18	*	68	*	68	*
2251	110	4.5 (68)	50.0 (3)	39	*	*	15	*	32	*	125	5.6 (25)	125	10.4 (47)
2252	44	*	*	9	*	*	12	*	11	*	61	*	61	*
2554	25	*	*	21	*	*	4	*	7	*	28	*	28	*
2555	2,308	5.5 (95)	57.3 (33)	1,206	18.8 (45)	52.8 (36)	468	48.5 (11)	451	36.6 (13)	2,656	7.5 (76)	2,651	15.5 (106)
2556	71	*	*	62	*	*	5	*	16	*	76	*	76	*
2557	0	*	*	0	*	*	2	*	0	*	1	*	1	*
2558	1	*	*	0	*	*	2	*	0	*	2	*	2	*
2559	4	*	*	2	*	*	2	*	0	*	6	*	6	*
3001	350	3.1 (22)	62.6 (90)	265	25.7 (74)	42.3 (4)	67	*	243	48.6 (31)	355	5.4 (21)	355	8.5 (25)
3002	85	*	*	65	*	*	19	*	64	*	82	*	82	*
3004	43	*	*	38	*	*	7	*	8	*	45	*	45	*
3007	11	*	*	7	*	*	0	*	3	*	9	*	9	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3009	2	*	*	0	*	*	0	*	0	*	0	*	0	*
3201	142	4.9 (81)	63.4 (96)	108	20.4 (58)	42.6 (5)	20	*	41	*	162	10.5 (111)	162	11.7 (69)
3301	1,284	3.9 (49)	58.7 (49)	1,037	19.0 (47)	51.2 (27)	268	44.0 (4)	244	38.9 (19)	1,404	6.3 (35)	1,404	9.5 (35)
3304	25	*	*	22	*	*	3	*	4	*	31	*	31	*
3305	38	*	*	31	*	*	5	*	4	*	46	*	46	*
3306	76	*	*	65	*	*	15	*	22	*	79	*	79	*
3308	24	*	*	21	*	*	1	*	8	*	24	*	24	*
3309	16	*	*	11	*	*	4	*	5	*	16	*	16	*
3312	124	7.3 (109)	64.5 (102)	103	17.5 (33)	51.5 (30)	35	*	26	*	142	12.0 (117)	142	9.9 (42)
3550	29	*	*	25	*	*	4	*	9	*	29	*	29	*
3551	11	*	*	7	*	*	0	*	8	*	12	*	12	*
3552	17	*	*	10	*	*	1	*	2	*	17	*	17	*
3701	243	5.3 (91)	62.6 (90)	168	13.7 (5)	54.2 (48)	50	*	90	*	351	6.8 (52)	351	14.0 (92)
3850	797	6.5 (107)	58.7 (49)	466	20.8 (62)	53.4 (42)	161	56.5 (22)	194	46.9 (27)	996	5.8 (29)	996	11.3 (63)
3901	96	*	*	71	*	*	16	*	40	*	118	5.9 (31)	118	4.2 (2)
3902	53	*	*	42	*	*	3	*	18	*	64	*	64	*
3903	2	*	*	1	*	*	0	*	1	*	2	*	2	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3950	490	3.1 (22)	58.0 (43)	279	16.8 (25)	56.6 (67)	89	*	172	31.4 (4)	584	9.6 (104)	584	10.4 (47)
4001	353	4.2 (60)	60.9 (69)	147	18.4 (41)	48.3 (11)	54	*	60	*	369	7.3 (70)	370	15.1 (103)
4002	234	5.6 (96)	62.4 (86)	132	15.9 (20)	59.1 (76)	37	*	38	*	253	8.3 (89)	253	7.1 (16)
4003	2	*	*	1	*	*	1	*	0	*	2	*	2	*
4004	4	*	*	3	*	*	0	*	0	*	4	*	4	*
4103	12	*	*	2	*	*	3	*	7	*	21	*	21	*
4106	599	4.3 (61)	61.8 (78)	352	19.9 (53)	48.9 (14)	144	59.0 (24)	175	45.7 (26)	735	5.0 (15)	736	10.1 (44)
4107	946	4.8 (77)	55.0 (17)	485	18.4 (41)	50.1 (20)	229	48.5 (11)	328	34.5 (8)	1,082	7.0 (61)	1,083	11.3 (63)
4109	162	7.4 (111)	50.0 (3)	60	*	*	48	*	93	*	330	6.7 (47)	331	13.9 (90)
4112	194	4.6 (71)	62.4 (86)	180	72.2 (78)	8.9 (1)	27	*	24	*	188	20.7 (119)	188	10.6 (50)
4113	198	2.5 (9)	65.7 (105)	39	*	*	37	*	58	*	218	9.6 (104)	217	11.5 (65)
4115	771	4.9 (81)	54.5 (15)	348	20.1 (54)	50.0 (19)	198	41.4 (1)	375	37.9 (18)	1,109	7.8 (84)	1,109	9.8 (40)
4351	207	3.4 (29)	66.2 (107)	162	23.5 (70)	52.5 (35)	23	*	42	*	203	7.4 (73)	203	10.8 (55)
4352	53	*	*	32	*	*	7	*	14	*	55	*	55	*
4353	122	2.5 (9)	64.8 (104)	78	*	*	18	*	29	*	132	8.3 (89)	132	17.4 (116)
4354	213	4.7 (75)	55.9 (25)	125	15.2 (13)	55.2 (57)	44	*	56	*	239	4.2 (4)	239	14.2 (95)
4355	114	4.4 (65)	56.1 (26)	37	*	*	20	*	30	*	116	11.2 (112)	116	6.9 (14)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4356	138	6.5 (107)	55.1 (18)	49	*	*	23	*	37	*	167	6.6 (44)	167	14.4 (96)
4358	39	*	*	31	*	*	6	*	8	*	42	*	42	*
4359	139	3.6 (39)	61.9 (80)	63	*	*	18	*	26	*	148	11.5 (114)	148	11.5 (65)
4360	71	*	*	56	*	*	9	*	14	*	73	*	73	*
4361	93	*	*	72	*	*	10	*	25	*	94	*	94	*
4362	86	*	*	67	*	*	13	*	30	*	93	*	93	*
4363	318	3.1 (22)	56.3 (27)	275	23.3 (67)	51.3 (28)	45	*	68	*	313	4.8 (11)	313	11.2 (62)
4401	348	4.9 (81)	58.3 (47)	299	15.1 (11)	55.5 (59)	63	*	82	*	337	4.5 (8)	337	7.1 (16)
4501	243	4.9 (81)	54.3 (14)	182	15.4 (16)	53.3 (41)	31	*	57	*	319	7.8 (84)	319	9.7 (37)
4601	362	3.6 (39)	57.7 (38)	97	*	*	98	*	97	*	447	6.7 (47)	447	13.0 (78)
4602	8	*	*	5	*	*	1	*	5	*	13	*	13	*
4701	327	5.2 (87)	56.3 (27)	210	15.2 (13)	55.7 (61)	65	*	124	13.7 (1)	331	11.5 (114)	332	13.0 (78)
5050	584	3.4 (29)	59.9 (57)	497	17.9 (38)	54.7 (54)	135	45.2 (5)	128	33.6 (7)	557	7.5 (76)	557	12.0 (71)
5051	914	4.7 (75)	57.2 (32)	779	15.5 (17)	56.5 (66)	163	45.4 (6)	232	36.6 (13)	872	8.6 (97)	872	11.5 (65)
5052	902	5.2 (87)	57.4 (34)	764	20.2 (55)	50.1 (20)	212	47.2 (8)	195	37.4 (17)	884	7.8 (84)	884	10.7 (52)
5055	192	2.1 (4)	46.4 (1)	89	*	*	31	*	103	41.7 (22)	391	6.9 (57)	393	4.8 (5)
5101	610	5.2 (87)	59.0 (55)	465	16.8 (25)	49.9 (18)	123	51.2 (15)	152	35.5 (11)	631	9.8 (107)	630	11.0 (58)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5202	203	3.9 (49)	55.2 (19)	141	19.1 (48)	44.7 (7)	28	*	59	*	207	6.8 (52)	210	6.2 (9)
5203	3	*	*	2	*	*	0	*	0	*	3	*	3	*
5204	4	*	*	2	*	*	1	*	0	*	4	*	4	*
5601	333	2.1 (4)	58.0 (43)	118	22.0 (64)	54.2 (48)	62	*	105	33.3 (5)	387	4.4 (7)	387	10.1 (44)
5801	371	4.6 (71)	56.9 (31)	174	17.8 (36)	54.6 (52)	104	51.0 (14)	121	34.7 (9)	546	6.4 (39)	546	12.6 (76)
5803	42	*	*	29	*	*	7	*	11	*	42	*	42	*
5804	1	*	*	1	*	*	0	*	1	*	8	*	8	*
5806	7	*	*	6	*	*	0	*	5	*	11	*	11	*
6101	13	*	*	10	*	*	0	*	3	*	12	*	12	*
6102	35	*	*	29	*	*	4	*	4	*	35	*	35	*
6103	1,360	3.8 (46)	59.9 (57)	1,027	20.3 (57)	50.3 (23)	213	47.9 (9)	138	31.2 (3)	1,341	8.7 (98)	1,343	12.1 (72)
6105	8	*	*	6	*	*	0	*	0	*	8	*	8	*
6301	1,424	3.6 (39)	57.7 (38)	1,096	20.6 (61)	51.0 (25)	299	49.5 (13)	316	37.0 (15)	1,435	5.4 (21)	1,435	12.2 (73)
6312	797	3.4 (29)	61.9 (80)	629	17.0 (29)	53.9 (45)	188	46.8 (7)	241	36.5 (12)	812	5.2 (17)	812	8.9 (27)
6316	416	4.8 (77)	59.1 (56)	354	13.6 (4)	55.6 (60)	83	*	114	33.3 (5)	404	6.9 (57)	404	11.1 (60)
6320	22	*	*	8	*	*	3	*	10	*	31	*	31	*
6325	18	*	*	7	*	*	3	*	14	*	32	*	32	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6330	62	*	*	16	*	*	14	*	15	*	103	6.8 (52)	104	15.4 (105)
6335	199	3.5 (36)	52.3 (8)	46	*	*	49	*	55	*	302	9.6 (104)	302	10.6 (50)
6340	10	*	*	3	*	*	1	*	3	*	13	*	13	*
6345	10	*	*	6	*	*	3	*	4	*	11	*	11	*
6360	67	*	*	21	*	*	12	*	18	*	102	3.9 (2)	101	14.9 (100)
7304	43	*	*	37	*	*	12	*	11	*	50	*	50	*
7306	729	3.7 (42)	64.1 (100)	560	20.5 (59)	50.7 (24)	141	51.8 (16)	128	43.0 (23)	840	6.8 (52)	839	11.1 (60)
7308	37	*	*	32	*	*	7	*	8	*	38	*	38	*
7310	3	*	*	2	*	*	2	*	3	*	10	*	10	*
7401	759	4.3 (61)	58.9 (54)	643	14.6 (8)	55.7 (61)	125	54.4 (20)	149	39.6 (21)	728	7.3 (70)	732	12.4 (74)
7601	107	3.7 (42)	64.5 (102)	31	*	*	32	*	44	*	212	6.6 (44)	212	13.7 (89)
7850	365	6.0 (103)	57.8 (40)	301	16.3 (22)	50.2 (22)	58	*	63	*	355	7.6 (80)	355	9.0 (29)
7852	10	*	*	8	*	*	3	*	1	*	10	*	10	*
8101	568	4.0 (54)	62.5 (88)	350	17.7 (35)	53.4 (42)	125	70.4 (29)	73	*	646	7.1 (64)	645	18.1 (118)
8102	113	4.4 (65)	63.7 (98)	46	*	*	22	*	9	*	141	5.7 (28)	141	13.5 (84)
8105	455	5.7 (100)	61.5 (75)	347	17.6 (34)	56.2 (65)	119	61.3 (25)	127	48.0 (29)	566	9.0 (100)	567	14.6 (97)
8106	89	*	*	62	*	*	28	*	32	*	146	4.8 (11)	146	11.0 (58)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8150	8	*	*	3	*	*	1	*	3	*	9	*	9	*
8151	1,045	3.9 (49)	57.5 (35)	663	21.6 (63)	49.6 (16)	212	61.3 (25)	277	39.4 (20)	1,045	7.4 (73)	1,044	13.1 (80)
8152	16	*	*	6	*	*	6	*	3	*	17	*	17	*
8153	2	*	*	2	*	*	0	*	1	*	2	*	2	*
8155	10	*	*	6	*	*	2	*	2	*	11	*	11	*
8157	3	*	*	1	*	*	0	*	1	*	3	*	3	*
8201	735	4.8 (77)	58.8 (53)	383	15.1 (11)	55.9 (63)	194	43.8 (3)	250	27.2 (2)	847	8.0 (88)	847	9.2 (32)
8202	793	5.7 (100)	50.1 (5)	427	22.0 (64)	45.2 (9)	220	52.3 (18)	279	43.4 (24)	1,283	7.5 (76)	1,281	4.0 (1)
8203	161	3.1 (22)	60.9 (69)	134	15.7 (19)	57.5 (71)	34	*	58	*	170	8.8 (99)	170	15.9 (113)
8250	235	2.1 (4)	61.7 (77)	179	14.5 (6)	57.0 (69)	75	*	74	*	264	8.3 (89)	264	18.2 (119)
8252	51	*	*	42	*	*	14	*	12	*	56	*	56	*
8255	724	4.3 (61)	58.1 (46)	460	11.5 (1)	55.4 (58)	204	43.6 (2)	262	37.0 (15)	815	6.3 (35)	816	8.7 (26)
8257	417	3.8 (46)	50.1 (5)	192	29.2 (76)	38.0 (3)	109	56.9 (23)	194	46.9 (27)	640	8.3 (89)	640	6.3 (10)
8263	240	2.9 (18)	62.9 (93)	187	16.0 (21)	55.1 (56)	48	*	113	35.4 (10)	244	9.0 (100)	244	14.8 (99)
8264	158	3.2 (27)	53.2 (9)	97	*	*	50	*	74	*	226	7.1 (64)	226	9.7 (37)
8266	4	*	*	2	*	*	0	*	2	*	9	*	9	*
8267	223	6.3 (105)	55.6 (23)	133	16.5 (23)	58.6 (74)	75	*	66	*	285	7.0 (61)	286	8.0 (23)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8901	343	3.5 (36)	61.8 (78)	239	18.0 (40)	51.9 (33)	63	*	54	*	373	6.7 (47)	374	15.5 (106)
8902	430	5.3 (91)	61.9 (80)	326	18.7 (43)	53.1 (39)	77	*	76	*	460	5.9 (31)	460	13.9 (90)
8903	352	1.4 (3)	63.6 (97)	224	20.5 (59)	53.1 (39)	62	*	63	*	393	6.4 (39)	392	16.3 (114)
8904	736	3.9 (49)	62.0 (83)	514	16.9 (28)	54.5 (51)	165	73.9 (30)	117	60.7 (34)	786	5.9 (31)	787	15.6 (108)
8905	405	3.7 (42)	56.8 (30)	219	19.6 (50)	49.8 (17)	91	*	87	*	525	6.7 (47)	525	10.7 (52)
8908	16	*	*	9	*	*	1	*	1	*	18	*	18	*
8909	165	6.1 (104)	69.1 (110)	117	17.1 (30)	59.0 (75)	41	*	21	*	181	7.7 (82)	181	11.6 (68)
9016	6	*	*	3	*	*	0	*	2	*	7	*	7	*
9018	3	*	*	3	*	*	0	*	0	*	3	*	3	*
9045	474	2.5 (9)	63.1 (94)	408	23.8 (72)	48.5 (13)	90	*	58	*	568	6.3 (35)	568	7.0 (15)
9050	290	3.4 (29)	60.0 (61)	230	26.5 (75)	49.1 (15)	55	*	34	*	322	7.5 (76)	321	10.0 (43)
9051	477	3.1 (22)	61.2 (71)	275	14.5 (6)	48.4 (12)	60	*	76	*	592	7.6 (80)	592	5.4 (8)
9053	434	4.4 (65)	61.5 (75)	387	15.0 (10)	57.6 (72)	75	*	53	*	437	4.8 (11)	437	14.0 (92)
9055	11	*	*	11	*	*	1	*	0	*	13	*	13	*
9058	281	4.6 (71)	62.6 (90)	212	19.8 (52)	46.7 (10)	65	*	53	*	349	7.2 (69)	350	11.7 (69)
9066	373	2.7 (12)	67.6 (109)	287	17.4 (32)	54.4 (50)	90	*	47	*	410	5.6 (25)	409	15.6 (108)
9068	321	3.7 (42)	66.4 (108)	284	23.6 (71)	51.4 (29)	56	*	22	*	305	5.6 (25)	305	7.2 (18)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9069	922	4.1 (58)	57.6 (37)	787	18.7 (43)	53.6 (44)	190	52.1 (17)	89	*	963	5.5 (24)	962	10.7 (52)
9070	284	3.9 (49)	55.3 (20)	209	17.2 (31)	51.7 (32)	67	*	30	*	369	6.0 (34)	370	14.1 (94)
9072	608	4.6 (71)	62.2 (85)	492	23.4 (69)	52.2 (34)	132	56.1 (21)	71	*	636	6.3 (35)	636	13.5 (84)
9080	821	4.5 (68)	60.5 (66)	613	18.8 (45)	56.6 (67)	206	65.5 (28)	150	48.7 (33)	1,001	5.8 (29)	1,001	9.8 (40)
9081	11	*	*	7	*	*	3	*	2	*	12	*	12	*
9082	53	*	*	34	*	*	12	*	5	*	51	*	52	*
9083	17	*	*	16	*	*	0	*	0	*	20	*	20	*
9084	12	*	*	12	*	*	0	*	1	*	13	*	13	*
9085	3	*	*	2	*	*	0	*	0	*	3	*	3	*
Michigan	**47,111**	**4.3**	**58.8**	**31,292**	**19.1**	**51.9**	**9,416**	**49.2**	**11,325**	**36.6**	**53,680**	**7.1**	**53,693**	**11.2**
National (Prior Year)	**106,629**	**4.4**	**58.2**	**62,805**	**21.4**	**50.4**	**35,452**	**17.4**	**20,899**	**27.9**	**60,377**	**6.9**	**60,411**	**10.7**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	36	*	*	39	*	36	*	31	*
0102	153	82.4 (68)	9.8 (80)	166	41.6 (24)	150	17.3 (73)	135	89.6 (49)
0103	150	75.3 (100)	14.7 (100)	154	32.5 (63)	137	21.2 (46)	126	91.3 (37)
0104	48	*	*	54	*	48	*	46	*
0250	50	*	*	50	*	34	*	38	*
0251	38	*	*	40	*	35	*	32	*
0252	40	*	*	42	*	43	*	37	*
0253	60	*	*	61	*	59	*	57	*
0401	173	85.0 (48)	9.2 (76)	184	32.6 (61)	172	30.8 (18)	152	95.4 (3)
0402	134	84.3 (55)	7.5 (66)	140	37.9 (43)	126	21.4 (43)	109	85.3 (79)
0403	47	*	*	48	*	46	*	45	*
0404	51	*	*	52	*	51	*	43	*
0501	182	96.2 (1)	0.0 (1)	190	34.2 (54)	177	23.2 (39)	155	87.7 (61)
0503	36	*	*	38	*	37	*	33	*
0504	166	88.6 (30)	7.2 (60)	175	32.6 (61)	163	30.1 (19)	140	87.1 (68)
0505	109	80.7 (80)	10.1 (81)	114	36.8 (49)	103	24.3 (32)	96	*
0506	65	*	*	68	*	58	*	52	*
0507	25	*	*	25	*	23	*	16	*
0508	15	*	*	15	*	15	*	10	*
0509	3	*	*	3	*	2	*	1	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	374	84.0 (60)	9.6 (77)	393	38.2 (41)	338	29.3 (23)	318	95.0 (4)
0651	195	79.5 (85)	14.4 (97)	211	45.5 (14)	191	12.6 (92)	181	91.2 (38)
0652	94	*	*	94	*	84	*	76	*
0653	139	72.7 (111)	20.1 (113)	144	49.3 (7)	136	29.4 (22)	124	87.9 (57)
0654	126	74.6 (103)	19.0 (111)	140	48.6 (8)	132	21.2 (46)	128	78.1 (105)
0655	112	81.3 (76)	7.1 (57)	117	44.4 (15)	109	23.9 (36)	97	*
0657	24	*	*	23	*	23	*	21	*
0701	132	79.5 (85)	16.7 (104)	131	38.2 (41)	125	54.4 (3)	113	92.0 (25)
0702	18	*	*	22	*	21	*	20	*
0703	71	*	*	84	*	76	*	65	*
0704	13	*	*	18	*	17	*	15	*
0705	46	*	*	52	*	48	*	43	*
0706	2	*	*	2	*	1	*	0	*
0850	289	88.9 (28)	2.1 (13)	308	20.5 (100)	269	23.0 (40)	224	95.5 (2)
0851	222	94.1 (3)	2.7 (20)	241	34.4 (53)	211	23.7 (37)	180	93.9 (10)
0901	558	72.4 (112)	21.9 (116)	594	43.9 (17)	538	16.2 (78)	489	81.4 (99)
0902	11	*	*	12	*	8	*	6	*
1051	482	73.7 (107)	17.0 (106)	494	32.0 (66)	438	25.1 (30)	343	89.2 (52)
1101	580	74.0 (106)	12.4 (90)	632	27.1 (78)	526	11.2 (99)	473	84.1 (90)
1102	246	72.0 (114)	19.9 (112)	266	40.2 (31)	217	14.7 (87)	215	94.9 (6)
1106	24	*	*	28	*	21	*	18	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1250	256	83.2 (64)	10.5 (85)	329	41.3 (27)	280	22.5 (41)	280	76.1 (108)
1251	278	79.5 (85)	10.1 (81)	293	41.6 (24)	264	19.7 (56)	236	84.3 (89)
1252	177	68.9 (116)	20.3 (114)	196	45.9 (11)	184	17.9 (70)	164	87.2 (65)
1253	247	72.1 (113)	20.6 (115)	283	35.7 (52)	237	14.8 (86)	214	90.2 (43)
1350	734	89.4 (22)	4.4 (38)	813	37.4 (46)	516	18.6 (62)	635	77.2 (107)
1351	41	*	*	43	*	37	*	33	*
1352	120	87.5 (33)	2.5 (16)	130	39.2 (34)	111	2.7 (112)	103	88.3 (56)
1401	72	*	*	79	*	44	*	57	*
1402	48	*	*	52	*	32	*	35	*
1403	5	*	*	6	*	4	*	5	*
1501	224	79.9 (81)	14.7 (100)	260	41.5 (26)	179	8.9 (102)	197	78.2 (104)
1503	367	89.1 (26)	6.8 (54)	399	45.9 (11)	346	47.4 (4)	334	73.7 (110)
1504	668	90.9 (12)	3.3 (30)	757	32.9 (59)	477	21.4 (43)	549	69.8 (111)
1505	127	91.3 (10)	5.5 (48)	142	30.3 (69)	100	29.0 (24)	93	*
1506	188	71.8 (115)	23.9 (117)	205	42.4 (21)	182	7.1 (109)	167	91.0 (40)
1507	196	91.8 (9)	4.6 (40)	213	37.1 (48)	160	28.8 (25)	158	92.4 (21)
1508	27	*	*	40	*	29	*	29	*
1509	118	90.7 (13)	5.1 (43)	147	50.3 (6)	100	37.0 (12)	124	82.3 (94)
1511	219	83.6 (61)	8.2 (73)	245	41.2 (29)	157	24.2 (34)	192	86.5 (72)
1512	111	89.2 (24)	8.1 (72)	122	32.0 (66)	73	*	95	*
1513	31	*	*	32	*	24	*	24	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1514	50	*	*	54	*	34	*	36	*
1701	218	85.3 (44)	13.8 (95)	230	52.6 (5)	203	57.6 (2)	200	86.0 (76)
1950	186	84.9 (49)	9.1 (75)	206	41.3 (27)	179	19.6 (59)	163	89.6 (49)
1952	41	*	*	40	*	35	*	32	*
1953	101	87.1 (36)	6.9 (55)	97	*	90	*	69	*
1954	188	85.1 (47)	10.1 (81)	201	36.8 (49)	179	28.5 (26)	164	92.1 (24)
1955	25	*	*	27	*	26	*	22	*
1957	160	86.3 (41)	6.3 (53)	173	31.2 (68)	146	20.5 (48)	132	92.4 (21)
1964	12	*	*	13	*	13	*	11	*
2150	212	88.2 (31)	5.2 (44)	239	46.0 (10)	218	24.3 (32)	203	92.6 (19)
2151	68	*	*	77	*	66	*	62	*
2251	125	90.4 (16)	7.2 (60)	141	41.1 (30)	131	34.4 (14)	112	91.1 (39)
2252	62	*	*	73	*	65	*	69	*
2554	28	*	*	29	*	23	*	23	*
2555	2,626	87.0 (37)	3.2 (26)	2,762	24.6 (86)	2,093	8.9 (102)	2,021	89.7 (48)
2556	75	*	*	73	*	58	*	14	*
2557	1	*	*	2	*	2	*	2	*
2558	2	*	*	2	*	2	*	2	*
2559	6	*	*	6	*	5	*	6	*
3001	356	75.3 (100)	9.6 (77)	376	36.2 (51)	340	5.6 (110)	315	81.9 (96)
3002	82	*	*	89	*	83	*	80	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3004	43	*	*	48	*	45	*	46	*
3007	12	*	*	14	*	12	*	11	*
3009	1	*	*	3	*	2	*	2	*
3201	159	81.8 (71)	13.8 (95)	172	45.9 (11)	142	38.7 (9)	136	85.3 (79)
3301	1,385	81.1 (77)	10.5 (85)	1,461	27.4 (76)	1,306	8.7 (106)	1,117	87.9 (57)
3304	31	*	*	29	*	30	*	24	*
3305	46	*	*	48	*	43	*	34	*
3306	78	*	*	82	*	80	*	60	*
3308	24	*	*	24	*	19	*	14	*
3309	16	*	*	17	*	15	*	14	*
3312	140	83.6 (61)	7.1 (57)	146	27.4 (76)	125	12.0 (96)	113	87.6 (62)
3550	29	*	*	30	*	28	*	28	*
3551	12	*	*	12	*	8	*	7	*
3552	17	*	*	19	*	16	*	16	*
3701	349	86.5 (39)	7.4 (64)	374	37.4 (46)	341	18.2 (65)	291	92.8 (17)
3850	985	84.7 (51)	8.2 (73)	1,062	38.6 (37)	916	14.4 (88)	882	87.2 (65)
3901	115	80.9 (79)	0.0 (1)	119	0.8 (118)	63	*	19	*
3902	64	*	*	61	*	36	*	4	*
3903	1	*	*	2	*	2	*	0	*
3950	575	78.1 (91)	11.7 (88)	645	33.3 (58)	572	18.0 (68)	493	84.6 (88)
4001	368	73.4 (108)	17.1 (107)	413	55.9 (2)	415	15.2 (82)	417	74.1 (109)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4002	254	74.8 (102)	18.1 (109)	269	53.9 (4)	265	20.0 (51)	269	84.8 (83)
4003	2	*	*	2	*	2	*	2	*
4004	4	*	*	4	*	4	*	4	*
4103	21	*	*	26	*	20	*	22	*
4106	725	81.8 (71)	7.2 (60)	798	30.3 (69)	711	8.9 (102)	652	86.7 (69)
4107	1,079	84.9 (49)	7.1 (57)	1,155	25.7 (81)	811	19.9 (52)	868	91.6 (34)
4109	329	81.8 (71)	12.8 (93)	345	30.1 (71)	305	19.7 (56)	248	94.4 (7)
4112	187	35.8 (119)	0.0 (1)	193	21.2 (96)	188	94.1 (1)	179	95.0 (4)
4113	213	89.2 (24)	0.0 (1)	225	24.0 (88)	158	20.3 (49)	208	86.1 (75)
4115	1,095	87.5 (33)	5.5 (48)	1,170	27.5 (75)	957	16.7 (76)	886	90.1 (45)
4351	203	92.1 (7)	3.4 (31)	212	43.4 (19)	85	*	175	94.3 (9)
4352	55	*	*	62	*	56	*	51	*
4353	129	81.4 (75)	12.4 (90)	145	38.6 (37)	128	24.2 (34)	114	83.3 (93)
4354	238	85.3 (44)	7.6 (68)	246	37.8 (44)	155	25.2 (29)	198	91.4 (36)
4355	116	88.8 (29)	7.8 (71)	125	32.8 (60)	115	23.5 (38)	103	93.2 (14)
4356	167	82.6 (66)	11.4 (87)	177	44.1 (16)	161	27.3 (27)	151	90.7 (41)
4358	40	*	*	48	*	35	*	31	*
4359	144	79.9 (81)	13.2 (94)	157	43.3 (20)	124	29.8 (21)	123	89.4 (51)
4360	72	*	*	72	*	48	*	54	*
4361	94	*	*	98	*	90	*	82	*
4362	93	*	*	96	*	89	*	82	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4363	309	74.1 (104)	18.8 (110)	319	42.0 (23)	209	32.5 (15)	250	84.8 (83)
4401	336	85.7 (43)	7.7 (70)	351	29.6 (73)	326	18.1 (67)	277	92.8 (17)
4501	317	74.1 (104)	17.4 (108)	326	40.2 (31)	301	29.9 (20)	254	91.7 (32)
4601	446	88.1 (32)	5.2 (44)	478	28.0 (74)	435	19.8 (54)	374	87.4 (63)
4602	13	*	*	13	*	7	*	5	*
4701	332	91.9 (8)	3.0 (23)	367	22.1 (93)	283	38.9 (8)	240	92.9 (16)
5050	552	90.2 (17)	3.1 (24)	599	21.9 (95)	385	19.7 (56)	435	91.7 (32)
5051	863	89.9 (18)	3.2 (26)	929	21.0 (97)	686	18.2 (65)	696	93.5 (13)
5052	871	84.7 (51)	4.5 (39)	938	21.0 (97)	625	17.4 (71)	709	91.8 (29)
5055	378	79.9 (81)	16.7 (104)	398	55.8 (3)	209	16.7 (76)	353	79.6 (102)
5101	618	89.3 (23)	5.3 (46)	664	24.5 (87)	521	25.5 (28)	491	93.9 (10)
5202	208	87.5 (33)	7.2 (60)	233	43.8 (18)	167	38.3 (10)	194	90.2 (43)
5203	3	*	*	4	*	4	*	4	*
5204	4	*	*	4	*	3	*	4	*
5601	382	84.3 (55)	12.6 (92)	398	33.7 (56)	372	30.9 (17)	313	97.4 (1)
5801	541	86.3 (41)	7.4 (64)	582	33.8 (55)	537	15.8 (81)	467	86.7 (69)
5803	41	*	*	44	*	40	*	28	*
5804	8	*	*	8	*	7	*	6	*
5806	11	*	*	10	*	4	*	1	*
6101	12	*	*	12	*	11	*	11	*
6102	35	*	*	34	*	30	*	26	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6103	1,321	84.6 (54)	7.0 (56)	1,404	38.4 (40)	1,263	15.0 (83)	1,159	78.8 (103)
6105	8	*	*	8	*	7	*	6	*
6301	1,433	84.1 (58)	5.6 (50)	1,521	25.1 (84)	1,247	12.1 (94)	1,127	89.2 (52)
6312	808	89.9 (18)	3.6 (34)	849	19.0 (105)	667	16.8 (74)	608	91.9 (27)
6316	401	89.0 (27)	3.2 (26)	424	26.7 (79)	356	19.9 (52)	294	92.2 (23)
6320	31	*	*	35	*	15	*	29	*
6325	32	*	*	33	*	18	*	21	*
6330	103	78.6 (88)	14.6 (99)	114	47.4 (9)	90	*	114	81.6 (97)
6335	303	86.5 (39)	10.2 (84)	316	42.1 (22)	91	*	270	87.8 (59)
6340	13	*	*	14	*	7	*	13	*
6345	11	*	*	11	*	7	*	9	*
6360	102	85.3 (44)	2.9 (22)	107	17.8 (110)	46	*	67	*
7304	49	*	*	53	*	48	*	42	*
7306	824	76.0 (98)	14.4 (97)	899	37.5 (45)	720	14.0 (89)	698	84.8 (83)
7308	38	*	*	38	*	34	*	29	*
7310	9	*	*	10	*	10	*	8	*
7401	731	67.4 (117)	24.4 (118)	793	39.7 (33)	709	16.2 (78)	658	86.5 (72)
7601	206	81.1 (77)	9.7 (79)	226	38.5 (39)	157	40.1 (7)	175	85.1 (82)
7850	344	76.2 (96)	15.4 (103)	380	39.2 (34)	347	15.0 (83)	299	86.6 (71)
7852	10	*	*	10	*	9	*	9	*
8101	637	78.0 (92)	2.5 (16)	684	19.0 (105)	458	15.9 (80)	480	90.4 (42)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8102	139	82.0 (70)	5.0 (42)	143	39.2 (34)	108	19.4 (60)	125	84.8 (83)
8105	566	84.1 (58)	4.2 (36)	594	23.6 (89)	539	18.6 (62)	453	92.5 (20)
8106	143	76.2 (96)	5.6 (50)	148	18.2 (107)	101	11.9 (97)	103	81.6 (97)
8150	9	*	*	11	*	9	*	9	*
8151	1,041	90.6 (14)	2.7 (20)	1,140	32.3 (65)	1,054	18.0 (68)	910	91.8 (29)
8152	17	*	*	16	*	17	*	17	*
8153	2	*	*	2	*	1	*	0	*
8155	11	*	*	10	*	10	*	9	*
8157	3	*	*	3	*	3	*	3	*
8201	847	91.0 (11)	1.4 (8)	909	18.2 (107)	666	37.4 (11)	652	93.7 (12)
8202	1,260	84.2 (57)	12.1 (89)	1,320	33.7 (56)	1,020	12.5 (93)	1,038	81.3 (101)
8203	170	94.1 (3)	0.0 (1)	181	19.9 (102)	109	22.0 (42)	129	93.0 (15)
8250	262	83.6 (61)	1.9 (12)	281	16.0 (115)	195	24.6 (31)	189	87.3 (64)
8252	55	*	*	55	*	45	*	45	*
8255	811	89.8 (20)	1.7 (11)	859	17.1 (113)	536	36.4 (13)	607	91.8 (29)
8257	635	95.7 (2)	0.3 (6)	658	26.6 (80)	601	19.8 (54)	522	85.2 (81)
8263	244	92.2 (6)	1.6 (9)	260	13.5 (117)	189	32.3 (16)	176	92.0 (25)
8264	224	92.9 (5)	3.1 (24)	251	20.7 (99)	219	17.4 (71)	196	94.4 (7)
8266	8	*	*	9	*	9	*	9	*
8267	284	90.5 (15)	2.5 (16)	313	19.8 (103)	272	41.2 (6)	229	90.0 (46)
8901	370	77.0 (95)	4.3 (37)	392	25.5 (82)	223	10.8 (100)	295	81.4 (99)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8902	449	82.4 (68)	2.2 (14)	480	22.5 (92)	305	18.4 (64)	342	85.7 (77)
8903	393	77.4 (93)	5.3 (46)	400	22.0 (94)	243	13.2 (90)	301	84.1 (90)
8904	782	79.8 (84)	4.0 (35)	817	20.0 (101)	470	16.8 (74)	610	87.2 (65)
8905	521	86.8 (38)	6.0 (52)	543	29.8 (72)	390	19.0 (61)	429	84.8 (83)
8908	18	*	*	18	*	15	*	18	*
8909	179	78.2 (90)	2.2 (14)	184	17.9 (109)	141	14.9 (85)	143	88.8 (54)
9016	7	*	*	7	*	7	*	7	*
9018	3	*	*	3	*	3	*	3	*
9045	570	72.8 (109)	15.3 (102)	581	32.4 (64)	108	46.3 (5)	452	86.5 (72)
9050	314	77.4 (93)	7.6 (68)	329	23.1 (90)	252	20.2 (50)	249	83.9 (92)
9051	561	62.0 (118)	29.2 (119)	615	56.3 (1)	520	13.1 (91)	577	65.7 (112)
9053	438	75.6 (99)	7.5 (66)	452	19.5 (104)	362	8.8 (105)	298	77.9 (106)
9055	13	*	*	12	*	8	*	3	*
9058	349	89.7 (21)	3.2 (26)	365	23.0 (91)	290	5.5 (111)	278	88.8 (54)
9066	360	84.7 (51)	0.8 (7)	417	17.3 (112)	302	11.3 (98)	297	91.9 (27)
9068	307	81.8 (71)	1.6 (9)	316	25.0 (85)	240	12.1 (94)	237	87.8 (59)
9069	955	78.3 (89)	3.4 (31)	998	17.4 (111)	610	8.7 (106)	668	90.0 (46)
9070	364	72.8 (109)	4.7 (41)	378	16.9 (114)	238	21.4 (43)	254	85.4 (78)
9072	630	82.5 (67)	3.5 (33)	664	16.0 (115)	529	7.6 (108)	456	82.2 (95)
9080	996	82.9 (65)	2.6 (19)	1,028	25.2 (83)	746	9.4 (101)	784	91.6 (34)
9081	12	*	*	16	*	15	*	16	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9082	52	*	*	55	*	48	*	35	*
9083	20	*	*	20	*	13	*	1	*
9084	13	*	*	13	*	4	*	0	*
9085	3	*	*	3	*	2	*	3	*
Michigan	**53,045**	**83.1**	**7.7**	**56,717**	**30.6**	**44,576**	**19.0**	**43,643**	**43643.0**
National (Prior Year)	**88,167**	**83.4**	**5.4**	**108,369**	**19.0**	**50,771**	**25.4**	**50,162**	**87.6**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS
Table 7A
2017 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	37	*	35	*	12	*	31	*	35	*
0102	150	49.3 (113)	156	34.0 (117)	64	*	135	25.9 (109)	156	8.3 (116)
0103	146	50.0 (114)	151	36.4 (119)	62	*	126	23.8 (106)	151	5.3 (107)
0104	48	*	48	*	23	*	46	*	48	*
0250	38	*	50	*	23	*	38	*	50	*
0251	35	*	38	*	20	*	32	*	38	*
0252	43	*	40	*	16	*	37	*	40	*
0253	59	*	59	*	29	*	57	*	59	*
0401	173	47.4 (112)	171	28.7 (110)	62	*	152	27.0 (110)	171	6.4 (112)
0402	126	57.1 (116)	134	30.6 (114)	59	*	109	22.9 (105)	134	3.0 (60)
0403	46	*	47	*	22	*	45	*	47	*
0404	51	*	51	*	20	*	43	*	51	*
0501	178	45.5 (107)	180	35.6 (118)	64	*	155	19.4 (99)	180	5.6 (109)
0503	37	*	36	*	16	*	33	*	36	*
0504	163	33.7 (70)	167	18.6 (73)	49	*	140	14.3 (77)	167	4.2 (87)
0505	103	38.8 (85)	108	21.3 (85)	35	*	96	*	109	9.2 (118)
0506	59	*	65	*	27	*	52	*	65	*
0507	23	*	25	*	6	*	16	*	25	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	15	*	16	*	1	*	10	*	16	*
0509	2	*	3	*	1	*	1	*	3	*
0650	366	41.5 (98)	372	23.9 (96)	127	37.8 (19)	318	14.5 (80)	373	2.4 (49)
0651	197	45.2 (105)	195	30.8 (115)	77	*	181	21.0 (102)	195	5.1 (106)
0652	84	*	94	*	41	*	76	*	94	*
0653	136	45.6 (109)	139	28.8 (111)	53	*	123	27.6 (111)	139	7.9 (115)
0654	134	45.5 (107)	125	26.4 (105)	54	*	128	22.7 (104)	125	4.0 (84)
0655	110	51.8 (115)	114	29.8 (113)	49	*	97	*	114	4.4 (93)
0657	22	*	24	*	8	*	21	*	24	*
0701	126	31.0 (60)	132	17.4 (65)	34	*	113	7.1 (24)	132	1.5 (25)
0702	21	*	18	*	6	*	20	*	18	*
0703	76	*	72	*	24	*	65	*	73	*
0704	17	*	14	*	4	*	15	*	14	*
0705	48	*	46	*	15	*	43	*	46	*
0706	1	*	2	*	0	*	0	*	2	*
0850	273	39.9 (92)	292	23.3 (94)	84	*	224	6.7 (21)	292	3.1 (66)
0851	219	31.1 (62)	223	22.4 (90)	55	*	180	11.1 (56)	223	1.8 (34)
0901	539	41.9 (99)	565	24.4 (98)	206	34.0 (28)	489	13.9 (75)	566	3.5 (74)
0902	8	*	11	*	2	*	6	*	11	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1051	438	37.0 (82)	486	22.2 (88)	124	37.1 (20)	343	9.0 (42)	486	3.9 (83)
1101	543	28.5 (52)	588	14.3 (53)	125	39.2 (18)	473	10.1 (50)	588	2.7 (56)
1102	239	31.0 (60)	247	18.6 (73)	58	*	215	13.5 (71)	247	2.8 (58)
1106	22	*	25	*	4	*	18	*	25	*
1250	315	34.9 (75)	311	19.0 (77)	96	*	280	12.9 (66)	311	4.8 (100)
1251	265	45.3 (106)	279	29.7 (112)	96	*	236	14.4 (78)	279	4.7 (97)
1252	185	35.7 (79)	187	17.1 (64)	47	*	164	15.9 (86)	187	3.7 (78)
1253	238	39.1 (88)	255	21.2 (84)	72	*	214	12.1 (63)	256	3.5 (74)
1350	717	30.4 (59)	760	17.6 (67)	195	29.7 (31)	644	13.2 (68)	763	3.1 (66)
1351	39	*	41	*	8	*	33	*	41	*
1352	113	43.4 (102)	120	20.8 (83)	42	*	103	1.9 (2)	120	0.8 (10)
1401	70	*	72	*	24	*	57	*	72	*
1402	45	*	49	*	10	*	35	*	49	*
1403	6	*	4	*	3	*	5	*	5	*
1501	220	23.6 (34)	227	9.7 (31)	40	*	197	6.1 (17)	225	1.8 (34)
1503	365	25.5 (45)	374	11.5 (42)	80	*	335	9.0 (42)	374	1.6 (28)
1504	657	21.3 (27)	699	9.6 (30)	129	46.5 (9)	559	6.8 (22)	701	2.3 (48)
1505	127	25.2 (42)	133	9.8 (32)	26	*	95	*	134	1.5 (25)
1506	187	24.6 (40)	189	14.8 (55)	39	*	167	4.2 (14)	189	1.6 (28)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1507	191	27.7 (50)	201	15.9 (59)	44	*	161	11.8 (61)	202	3.5 (74)
1508	32	*	30	*	2	*	29	*	30	*
1509	129	18.6 (21)	120	12.5 (49)	21	*	124	8.9 (41)	121	3.3 (71)
1511	217	37.3 (83)	229	20.5 (81)	75	*	192	13.5 (71)	229	2.6 (54)
1512	108	34.3 (71)	113	23.0 (91)	33	*	95	*	114	2.6 (54)
1513	26	*	32	*	4	*	24	*	32	*
1514	47	*	52	*	12	*	36	*	51	*
1701	221	42.5 (101)	215	31.2 (116)	80	*	201	28.4 (112)	214	17.8 (119)
1950	187	40.6 (94)	190	18.4 (72)	61	*	164	15.9 (86)	190	4.7 (97)
1952	37	*	41	*	12	*	32	*	41	*
1953	92	*	101	23.8 (95)	29	*	69	*	101	2.0 (44)
1954	186	44.1 (104)	190	27.9 (107)	69	*	164	13.4 (70)	190	1.1 (14)
1955	26	*	25	*	11	*	22	*	25	*
1957	158	32.3 (67)	160	20.6 (82)	45	*	133	6.8 (22)	160	1.3 (17)
1964	13	*	13	*	3	*	11	*	13	*
2150	221	38.9 (86)	211	23.2 (92)	66	*	203	9.9 (48)	211	1.9 (40)
2151	70	*	68	*	31	*	62	*	68	*
2251	138	42.0 (100)	125	25.6 (101)	38	*	112	3.6 (11)	126	1.6 (28)
2252	71	*	61	*	26	*	68	*	61	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2554	23	*	28	*	5	*	23	*	28	*
2555	2,383	29.6 (55)	2,663	16.2 (61)	619	36.2 (22)	2,023	10.2 (51)	2,667	2.7 (56)
2556	58	*	76	*	4	*	14	*	76	*
2557	2	*	1	*	0	*	2	*	1	*
2558	2	*	2	*	0	*	2	*	2	*
2559	6	*	6	*	1	*	6	*	6	*
3001	340	34.4 (72)	355	14.1 (52)	96	*	315	6.3 (18)	355	2.5 (51)
3002	84	*	82	*	19	*	80	*	82	*
3004	47	*	45	*	6	*	46	*	45	*
3007	12	*	9	*	1	*	11	*	9	*
3009	2	*	0	*	0	*	2	*	0	*
3201	164	32.3 (67)	162	16.0 (60)	45	*	137	16.1 (88)	162	2.5 (51)
3301	1,319	17.3 (15)	1,403	11.6 (44)	187	34.8 (26)	1,118	7.7 (31)	1,404	4.5 (95)
3304	30	*	31	*	0	*	24	*	31	*
3305	43	*	46	*	9	*	34	*	46	*
3306	80	*	79	*	9	*	60	*	79	*
3308	19	*	24	*	5	*	14	*	24	*
3309	15	*	16	*	3	*	14	*	16	*
3312	126	23.0 (31)	142	16.2 (61)	26	*	113	8.0 (36)	142	4.2 (87)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3550	28	*	29	*	13	*	28	*	29	*
3551	8	*	12	*	4	*	7	*	12	*
3552	16	*	15	*	11	*	16	*	17	*
3701	342	40.1 (93)	353	23.2 (92)	114	33.3 (29)	291	14.8 (82)	353	3.7 (78)
3850	966	40.7 (95)	995	26.4 (105)	339	28.9 (32)	882	16.3 (94)	994	3.0 (60)
3901	62	*	118	18.6 (73)	8	*	19	*	118	0.8 (10)
3902	36	*	64	*	1	*	4	*	64	*
3903	2	*	2	*	0	*	0	*	2	*
3950	573	32.1 (66)	585	17.9 (69)	144	35.4 (24)	493	12.0 (62)	585	2.2 (45)
4001	418	35.9 (80)	369	19.8 (80)	128	28.9 (32)	418	12.4 (65)	370	3.0 (60)
4002	269	35.3 (78)	253	17.4 (65)	90	*	269	10.8 (53)	252	1.6 (28)
4003	2	*	2	*	1	*	2	*	2	*
4004	4	*	4	*	2	*	4	*	4	*
4103	22	*	21	*	4	*	22	*	21	*
4106	720	23.6 (34)	736	11.3 (41)	127	46.5 (9)	652	3.8 (13)	736	1.8 (34)
4107	1,053	21.3 (27)	1,086	12.0 (47)	171	39.8 (16)	868	11.1 (56)	1,085	4.3 (90)
4109	307	30.3 (58)	332	15.7 (57)	77	*	248	8.5 (37)	332	1.2 (15)
4112	188	4.3 (3)	188	1.6 (3)	4	*	179	7.3 (27)	188	3.2 (69)
4113	210	7.6 (6)	218	2.8 (6)	17	*	208	0.5 (1)	219	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4115	1,047	18.3 (19)	1,108	8.6 (27)	153	45.1 (14)	888	3.4 (10)	1,110	0.7 (7)
4351	199	45.7 (111)	203	28.6 (108)	77	*	174	16.7 (95)	203	3.0 (60)
4352	56	*	54	*	24	*	51	*	54	*
4353	128	35.2 (77)	132	24.2 (97)	41	*	114	7.9 (34)	132	1.5 (25)
4354	226	43.4 (102)	239	25.9 (103)	79	*	198	16.2 (92)	239	4.6 (96)
4355	117	39.3 (90)	117	18.8 (76)	37	*	103	2.9 (6)	117	4.3 (90)
4356	162	33.3 (69)	167	16.2 (61)	44	*	151	7.9 (34)	166	1.8 (34)
4358	36	*	43	*	6	*	31	*	43	*
4359	136	30.1 (56)	148	17.6 (67)	37	*	123	9.8 (47)	148	1.4 (20)
4360	60	*	73	*	24	*	54	*	73	*
4361	93	*	94	*	34	*	82	*	94	*
4362	89	*	93	*	42	*	82	*	93	*
4363	283	39.6 (91)	313	25.6 (101)	92	*	250	15.6 (84)	313	4.8 (100)
4401	326	34.7 (74)	337	15.7 (57)	90	*	277	2.2 (3)	337	0.3 (2)
4501	302	41.1 (97)	319	22.3 (89)	102	34.3 (27)	254	16.1 (88)	319	4.1 (86)
4601	439	40.8 (96)	447	19.0 (77)	137	48.2 (8)	374	11.0 (55)	448	3.8 (81)
4602	7	*	13	*	2	*	5	*	13	*
4701	326	34.4 (72)	329	17.9 (69)	76	*	242	11.2 (58)	332	2.4 (49)
5050	524	24.4 (37)	558	11.5 (42)	102	50.0 (6)	435	7.1 (24)	558	1.4 (20)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5051	849	23.0 (31)	872	10.3 (34)	161	46.0 (12)	697	5.7 (16)	873	1.8 (34)
5052	861	22.6 (30)	884	11.2 (40)	150	36.7 (21)	712	8.7 (38)	884	1.4 (20)
5055	386	1.8 (1)	394	0.3 (1)	9	*	362	3.3 (9)	393	1.3 (17)
5101	607	24.4 (37)	628	11.9 (45)	124	46.0 (12)	494	3.6 (11)	630	1.9 (40)
5202	215	45.6 (109)	210	28.6 (108)	87	*	196	12.2 (64)	211	1.4 (20)
5203	4	*	3	*	3	*	4	*	3	*
5204	4	*	4	*	0	*	4	*	4	*
5601	372	29.3 (53)	387	18.1 (71)	91	*	312	9.6 (45)	387	1.0 (12)
5801	538	38.7 (84)	546	25.1 (100)	165	30.3 (30)	467	13.3 (69)	546	4.8 (100)
5803	40	*	42	*	8	*	28	*	42	*
5804	7	*	8	*	3	*	6	*	8	*
5806	4	*	11	*	0	*	1	*	11	*
6101	11	*	12	*	5	*	11	*	12	*
6102	30	*	35	*	8	*	26	*	35	*
6103	1,289	31.2 (63)	1,339	21.3 (85)	347	24.5 (35)	1,160	13.6 (73)	1,344	5.9 (110)
6105	7	*	8	*	1	*	6	*	8	*
6301	1,366	20.6 (25)	1,438	12.0 (47)	214	35.0 (25)	1,131	7.3 (27)	1,438	1.3 (17)
6312	794	6.7 (5)	812	2.0 (5)	38	*	610	2.6 (5)	810	0.5 (4)
6316	406	15.3 (13)	404	8.7 (28)	31	*	294	2.4 (4)	404	0.5 (4)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6320	33	*	31	*	3	*	29	*	31	*
6325	30	*	32	*	7	*	21	*	32	*
6330	114	21.9 (29)	104	14.4 (54)	23	*	114	8.8 (39)	104	1.9 (40)
6335	302	8.3 (8)	304	4.3 (11)	17	*	270	3.0 (7)	304	0.7 (7)
6340	14	*	13	*	2	*	13	*	13	*
6345	11	*	11	*	3	*	9	*	11	*
6360	84	*	101	11.9 (45)	14	*	67	*	102	1.0 (12)
7304	50	*	50	*	19	*	42	*	50	*
7306	790	36.6 (81)	841	13.4 (50)	249	49.8 (7)	699	14.6 (81)	841	3.0 (60)
7308	34	*	38	*	11	*	29	*	38	*
7310	10	*	10	*	4	*	8	*	10	*
7401	730	39.2 (89)	732	24.7 (99)	243	28.4 (34)	658	15.2 (83)	732	3.6 (77)
7601	200	35.0 (76)	210	26.2 (104)	70	*	183	14.2 (76)	212	4.7 (97)
7850	342	38.9 (86)	353	22.1 (87)	117	35.9 (23)	298	15.8 (85)	355	3.1 (66)
7852	9	*	10	*	1	*	9	*	10	*
8101	571	26.6 (47)	646	9.4 (29)	125	61.6 (2)	477	16.1 (88)	646	4.8 (100)
8102	134	16.4 (14)	141	7.1 (22)	18	*	125	4.8 (15)	141	0.7 (7)
8105	542	14.6 (12)	567	6.0 (19)	56	*	453	11.5 (59)	567	3.2 (69)
8106	129	24.8 (41)	145	10.3 (34)	28	*	102	7.8 (33)	146	1.4 (20)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8150	9	*	9	*	1	*	9	*	9	*
8151	1,054	24.4 (37)	1,045	10.4 (36)	194	43.8 (15)	909	7.7 (31)	1,044	1.8 (34)
8152	17	*	17	*	5	*	17	*	17	*
8153	1	*	2	*	0	*	0	*	2	*
8155	10	*	11	*	1	*	9	*	11	*
8157	3	*	3	*	0	*	3	*	3	*
8201	832	20.0 (23)	846	10.2 (33)	123	46.3 (11)	652	11.7 (60)	847	2.2 (45)
8202	1,228	5.0 (4)	1,284	1.9 (4)	53	*	1,041	6.3 (18)	1,282	2.5 (51)
8203	163	31.3 (64)	169	19.5 (79)	44	*	129	10.9 (54)	170	1.2 (15)
8250	244	31.6 (65)	264	10.6 (37)	54	*	189	13.8 (74)	264	3.4 (73)
8252	50	*	56	*	13	*	45	*	56	*
8255	752	30.1 (56)	814	15.5 (56)	180	39.4 (17)	606	9.2 (44)	815	1.7 (33)
8257	625	2.2 (2)	640	1.1 (2)	12	*	524	3.1 (8)	639	0.5 (4)
8263	226	27.0 (48)	244	11.1 (39)	49	*	176	7.4 (29)	244	1.6 (28)
8264	233	27.5 (49)	225	10.7 (38)	51	*	197	9.6 (45)	226	2.2 (45)
8266	9	*	9	*	1	*	9	*	9	*
8267	285	29.5 (54)	286	14.0 (51)	60	*	229	10.5 (52)	286	0.3 (2)
8901	327	23.5 (33)	373	6.2 (20)	65	*	296	16.2 (92)	373	3.8 (81)
8902	420	14.5 (11)	459	3.3 (9)	51	*	342	12.9 (66)	459	5.9 (110)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8903	340	25.3 (43)	393	7.6 (25)	80	*	302	19.5 (100)	393	7.1 (113)
8904	720	24.2 (36)	786	5.6 (17)	143	72.0 (1)	609	18.2 (98)	787	7.5 (114)
8905	505	17.6 (17)	525	8.0 (26)	76	*	429	6.3 (18)	525	3.0 (60)
8908	18	*	18	*	6	*	18	*	18	*
8909	169	20.7 (26)	181	2.8 (6)	31	*	143	16.1 (88)	181	5.5 (108)
9016	7	*	7	*	2	*	7	*	7	*
9018	3	*	3	*	1	*	3	*	3	*
9045	525	12.0 (10)	569	5.1 (15)	56	*	452	8.8 (39)	569	3.3 (71)
9050	284	28.2 (51)	322	4.7 (13)	78	*	251	19.9 (101)	321	3.7 (78)
9051	602	8.1 (7)	593	3.5 (10)	38	*	579	7.4 (29)	593	1.9 (40)
9053	378	18.0 (18)	438	5.7 (18)	56	*	299	14.4 (78)	438	5.0 (105)
9055	9	*	13	*	1	*	3	*	13	*
9058	314	9.2 (9)	350	3.1 (8)	22	*	278	7.2 (26)	349	4.0 (84)
9066	348	18.4 (20)	411	5.4 (16)	57	*	297	17.8 (97)	410	2.9 (59)
9068	266	25.9 (46)	307	4.6 (12)	58	*	237	17.7 (96)	307	4.9 (104)
9069	842	20.5 (24)	963	6.7 (21)	142	59.2 (3)	668	22.0 (103)	962	8.7 (117)
9070	320	17.5 (16)	370	4.9 (14)	46	*	253	25.7 (108)	368	4.3 (90)
9072	543	25.4 (44)	637	7.5 (24)	114	54.4 (5)	456	24.8 (107)	637	4.4 (93)
9080	898	19.5 (22)	1,000	7.2 (23)	149	58.4 (4)	784	9.9 (48)	1,001	4.2 (87)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9081	16	*	12	*	0	*	16	*	12	*
9082	49	*	52	*	10	*	35	*	52	*
9083	15	*	20	*	0	*	1	*	18	*
9084	10	*	13	*	0	*	0	*	13	*
9085	3	*	3	*	1	*	3	*	3	*
Michigan	**51,028**	**27.1**	**53,711**	**14.1**	**11,398**	**39.3**	**43,712**	**11.3**	**53,738**	**3.2**
National (Prior Year)	**112,214**	**29.5**	**114,607**	**15.2**	**15,803**	**35.0**	**75,630**	**14.7**	**88,858**	**11.7**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	32	*	*	33	*	27	*	32	*
0102	148	6.8 (17)	9.5 (89)	151	7.3 (4)	138	5.8 (94)	148	63.5 (81)
0103	146	7.5 (24)	7.5 (57)	147	8.8 (16)	132	4.5 (67)	146	66.4 (74)
0104	48	*	*	48	*	47	*	48	*
0250	45	*	*	48	*	40	*	45	*
0251	35	*	*	37	*	34	*	35	*
0252	38	*	*	38	*	35	*	38	*
0253	52	*	*	52	*	46	*	52	*
0401	159	11.3 (82)	13.2 (109)	164	9.8 (28)	143	5.6 (87)	158	68.4 (68)
0402	127	9.4 (56)	7.9 (65)	130	10.0 (29)	114	5.3 (78)	127	78.0 (33)
0403	46	*	*	46	*	41	*	46	*
0404	49	*	*	50	*	44	*	49	*
0501	126	9.5 (57)	13.5 (113)	170	10.6 (37)	111	2.7 (23)	153	77.8 (36)
0503	32	*	*	33	*	30	*	32	*
0504	156	5.8 (8)	8.3 (72)	162	6.8 (3)	145	3.4 (40)	159	79.9 (24)
0505	97	*	*	106	6.6 (2)	90	*	104	83.7 (11)
0506	58	*	*	62	*	54	*	59	*
0507	20	*	*	22	*	18	*	22	*
0508	14	*	*	15	*	12	*	15	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0509	3	*	*	3	*	3	*	3	*
0650	341	8.8 (49)	7.0 (52)	360	9.2 (22)	311	5.8 (94)	341	79.5 (29)
0651	180	6.1 (10)	9.4 (86)	188	9.0 (20)	165	3.0 (31)	180	67.8 (69)
0652	85	*	*	86	*	74	*	85	*
0653	126	10.3 (68)	8.7 (79)	134	9.7 (26)	114	6.1 (98)	127	84.3 (10)
0654	119	3.4 (1)	10.9 (100)	122	13.1 (84)	105	0.0 (1)	119	86.6 (3)
0655	106	7.5 (24)	7.5 (57)	108	12.0 (66)	92	*	107	72.0 (60)
0657	21	*	*	21	*	21	*	21	*
0701	118	3.4 (1)	13.6 (114)	121	8.3 (9)	102	1.0 (2)	118	85.6 (5)
0702	14	*	*	17	*	14	*	14	*
0703	71	*	*	71	*	59	*	71	*
0704	13	*	*	14	*	13	*	13	*
0705	42	*	*	45	*	39	*	42	*
0706	2	*	*	2	*	1	*	2	*
0850	282	9.9 (63)	7.1 (53)	292	11.3 (52)	249	2.0 (8)	286	84.6 (8)
0851	212	9.9 (63)	9.4 (86)	218	11.0 (44)	185	3.8 (54)	217	79.7 (25)
0901	534	6.9 (18)	9.7 (93)	540	10.2 (33)	479	3.5 (46)	534	65.4 (75)
0902	11	*	*	11	*	11	*	11	*
1051	455	8.4 (41)	9.7 (93)	468	8.3 (9)	415	4.1 (59)	455	86.6 (3)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1101	560	10.4 (71)	6.8 (48)	583	11.8 (60)	494	5.3 (78)	563	71.6 (61)
1102	230	5.2 (7)	10.0 (96)	240	9.6 (25)	208	3.8 (54)	230	76.1 (44)
1106	24	*	*	24	*	20	*	24	*
1250	298	4.7 (3)	10.1 (97)	299	9.0 (20)	266	2.3 (14)	300	83.3 (12)
1251	253	5.1 (6)	9.5 (89)	266	10.9 (41)	227	1.3 (4)	254	78.0 (33)
1252	172	7.6 (28)	8.1 (69)	176	11.9 (63)	148	2.7 (23)	173	85.0 (6)
1253	246	11.0 (80)	8.5 (76)	247	12.1 (67)	213	2.3 (14)	248	84.7 (7)
1350	713	10.9 (79)	6.5 (43)	735	12.9 (76)	615	3.4 (40)	714	75.4 (47)
1351	39	*	*	39	*	38	*	39	*
1352	115	9.6 (59)	12.2 (108)	118	11.9 (63)	103	5.8 (94)	115	62.6 (84)
1401	66	*	*	64	*	60	*	66	*
1402	44	*	*	47	*	38	*	44	*
1403	4	*	*	4	*	4	*	4	*
1501	209	6.2 (11)	9.1 (82)	211	10.0 (29)	183	2.2 (12)	210	65.2 (76)
1503	345	8.1 (36)	10.1 (97)	350	12.9 (76)	294	3.1 (34)	347	80.7 (18)
1504	667	8.5 (45)	9.1 (82)	686	11.8 (60)	585	2.7 (23)	675	75.7 (45)
1505	126	10.3 (68)	15.1 (115)	128	12.5 (71)	110	2.7 (23)	127	79.5 (29)
1506	181	5.0 (5)	13.3 (111)	185	10.3 (35)	162	2.5 (19)	182	74.7 (52)
1507	182	8.2 (38)	13.2 (109)	191	8.9 (18)	162	3.1 (34)	187	77.0 (40)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1508	29	*	*	29	*	24	*	29	*
1509	112	9.8 (61)	7.1 (53)	115	13.0 (82)	98	*	113	64.6 (77)
1511	214	7.5 (24)	11.7 (105)	215	10.7 (39)	189	1.1 (3)	217	79.7 (25)
1512	110	13.6 (106)	9.1 (82)	112	11.6 (56)	98	*	110	80.9 (16)
1513	30	*	*	31	*	27	*	31	*
1514	50	*	*	51	*	38	*	51	*
1701	205	10.7 (76)	11.7 (105)	200	12.5 (71)	171	4.1 (59)	206	69.9 (65)
1950	184	9.8 (61)	5.4 (28)	186	11.8 (60)	160	4.4 (64)	184	87.0 (1)
1952	40	*	*	38	*	35	*	40	*
1953	97	*	*	94	*	86	*	97	*
1954	182	6.6 (13)	10.4 (99)	183	8.7 (14)	164	3.0 (31)	182	80.8 (17)
1955	24	*	*	24	*	23	*	24	*
1957	150	10.0 (65)	11.3 (104)	151	8.6 (12)	135	4.4 (64)	151	86.8 (2)
1964	10	*	*	11	*	9	*	10	*
2150	196	6.6 (13)	8.2 (71)	202	10.9 (41)	175	1.7 (5)	196	75.0 (50)
2151	66	*	*	65	*	54	*	66	*
2251	119	8.4 (41)	5.9 (35)	122	7.4 (5)	109	2.8 (28)	119	77.3 (39)
2252	57	*	*	59	*	49	*	57	*
2554	27	*	*	28	*	23	*	27	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2555	2,573	11.9 (89)	4.9 (22)	2,596	13.6 (90)	2,192	5.7 (91)	2,584	57.9 (92)
2556	73	*	*	72	*	58	*	73	*
2557	1	*	*	1	*	0	*	1	*
2558	2	*	*	2	*	2	*	2	*
2559	6	*	*	6	*	5	*	6	*
3001	333	13.8 (108)	4.2 (20)	339	13.9 (94)	285	6.0 (97)	333	58.9 (89)
3002	78	*	*	81	*	71	*	78	*
3004	40	*	*	39	*	36	*	40	*
3007	9	*	*	9	*	8	*	9	*
3009	0	*	*	0	*	0	*	0	*
3201	156	7.7 (32)	10.9 (100)	158	13.3 (86)	135	2.2 (12)	157	74.5 (53)
3301	1,327	10.0 (65)	6.1 (36)	1,370	12.2 (68)	1,161	4.4 (64)	1,335	80.1 (21)
3304	28	*	*	29	*	26	*	29	*
3305	44	*	*	45	*	41	*	44	*
3306	74	*	*	76	*	67	*	74	*
3308	23	*	*	24	*	22	*	23	*
3309	16	*	*	16	*	14	*	16	*
3312	132	10.6 (75)	6.8 (48)	140	8.6 (12)	122	5.7 (91)	135	66.7 (72)
3550	26	*	*	29	*	24	*	26	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3551	12	*	*	12	*	11	*	12	*
3552	17	*	*	17	*	14	*	17	*
3701	337	9.2 (55)	6.2 (37)	343	13.4 (88)	290	3.4 (40)	338	79.0 (31)
3850	943	8.7 (48)	7.1 (53)	964	13.8 (92)	809	3.0 (31)	943	73.6 (58)
3901	106	12.3 (93)	2.8 (4)	110	10.9 (41)	90	*	107	73.8 (56)
3902	60	*	*	59	*	50	*	60	*
3903	2	*	*	2	*	2	*	2	*
3950	546	9.5 (57)	8.4 (74)	565	12.4 (69)	477	3.6 (49)	550	75.5 (46)
4001	348	11.5 (86)	5.2 (25)	354	14.1 (96)	294	5.4 (81)	349	70.8 (64)
4002	242	7.4 (21)	9.5 (89)	249	11.2 (49)	215	1.9 (7)	243	67.5 (71)
4003	2	*	*	2	*	2	*	2	*
4004	3	*	*	4	*	3	*	3	*
4103	20	*	*	21	*	16	*	20	*
4106	700	12.1 (91)	4.9 (22)	716	13.3 (86)	603	6.5 (101)	705	72.1 (59)
4107	1,024	11.3 (82)	8.1 (69)	1,060	13.2 (85)	886	4.7 (70)	1,028	76.7 (42)
4109	309	11.7 (88)	7.8 (64)	314	17.2 (113)	255	2.0 (8)	312	82.4 (15)
4112	185	6.5 (12)	6.5 (43)	185	11.4 (53)	163	1.8 (6)	185	29.7 (116)
4113	211	7.6 (28)	7.1 (53)	216	10.6 (37)	187	3.7 (51)	211	80.1 (21)
4115	1,048	9.0 (52)	7.7 (61)	1,079	13.0 (82)	907	2.5 (19)	1,053	82.7 (13)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4351	187	7.5 (24)	13.4 (112)	195	12.8 (75)	164	2.4 (18)	186	75.3 (48)
4352	53	*	*	54	*	41	*	53	*
4353	125	8.0 (33)	5.6 (31)	126	6.3 (1)	115	5.2 (77)	125	76.8 (41)
4354	233	7.3 (20)	6.4 (39)	232	8.2 (8)	213	4.2 (61)	233	75.1 (49)
4355	109	12.8 (98)	11.9 (107)	113	9.7 (26)	95	*	108	80.6 (19)
4356	163	8.6 (47)	8.0 (68)	164	11.6 (56)	143	4.2 (61)	163	77.9 (35)
4358	40	*	*	39	*	37	*	41	*
4359	134	6.7 (15)	7.5 (57)	135	7.4 (5)	123	3.3 (39)	134	70.9 (63)
4360	62	*	*	67	*	57	*	62	*
4361	88	*	*	89	*	79	*	88	*
4362	86	*	*	90	*	78	*	86	*
4363	291	6.9 (18)	7.9 (65)	296	9.5 (23)	256	3.9 (56)	292	82.5 (14)
4401	322	8.4 (41)	9.3 (85)	324	10.8 (40)	284	3.9 (56)	323	79.6 (28)
4501	309	8.4 (41)	11.0 (103)	310	11.0 (44)	272	3.7 (51)	309	73.8 (56)
4601	420	7.4 (21)	8.8 (80)	433	9.5 (23)	377	3.4 (40)	419	77.6 (38)
4602	12	*	*	12	*	12	*	12	*
4701	323	8.0 (33)	8.4 (74)	330	11.2 (49)	285	2.5 (19)	323	74.9 (51)
5050	549	10.4 (71)	5.5 (30)	546	13.7 (91)	465	3.4 (40)	549	58.3 (91)
5051	846	11.3 (82)	5.6 (31)	855	12.9 (76)	726	3.7 (51)	847	63.3 (82)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5052	865	11.6 (87)	4.0 (15)	868	13.8 (92)	738	5.1 (74)	865	59.9 (88)
5055	291	10.7 (76)	3.1 (6)	373	8.8 (16)	261	5.4 (81)	364	44.0 (109)
5101	609	6.7 (15)	7.7 (61)	620	10.5 (36)	543	2.6 (22)	610	71.6 (61)
5202	203	10.3 (68)	9.9 (95)	210	10.0 (29)	182	7.1 (104)	205	80.0 (23)
5203	3	*	*	3	*	3	*	3	*
5204	4	*	*	4	*	3	*	4	*
5601	368	6.0 (9)	10.9 (100)	372	8.9 (18)	334	2.1 (10)	369	74.5 (53)
5801	514	8.0 (33)	6.4 (39)	529	10.0 (29)	458	3.1 (34)	498	63.7 (78)
5803	40	*	*	40	*	37	*	39	*
5804	8	*	*	8	*	7	*	8	*
5806	11	*	*	10	*	8	*	11	*
6101	12	*	*	11	*	11	*	12	*
6102	34	*	*	31	*	27	*	34	*
6103	1,272	10.5 (73)	5.2 (25)	1,289	14.4 (101)	1,056	4.2 (61)	1,274	63.7 (78)
6105	8	*	*	8	*	6	*	8	*
6301	1,372	12.8 (98)	5.0 (24)	1,411	14.1 (96)	1,174	5.4 (81)	1,381	66.6 (73)
6312	782	11.4 (85)	5.2 (25)	796	13.9 (94)	671	3.9 (56)	786	77.7 (37)
6316	390	12.1 (91)	7.7 (61)	396	13.4 (88)	333	4.5 (67)	392	80.4 (20)
6320	29	*	*	30	*	26	*	30	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6325	25	*	*	31	*	24	*	29	*
6330	95	*	*	101	21.8 (117)	74	*	97	*
6335	289	9.0 (52)	6.6 (46)	295	10.2 (33)	259	5.4 (81)	295	63.1 (83)
6340	10	*	*	11	*	8	*	11	*
6345	9	*	*	9	*	8	*	9	*
6360	96	*	*	98	*	75	*	99	*
7304	47	*	*	47	*	42	*	47	*
7306	812	13.4 (103)	5.8 (34)	824	12.9 (76)	705	5.1 (74)	813	60.3 (87)
7308	38	*	*	37	*	35	*	38	*
7310	9	*	*	10	*	7	*	9	*
7401	691	8.1 (36)	6.4 (39)	703	11.1 (46)	613	2.9 (29)	690	69.7 (66)
7601	204	8.8 (49)	6.9 (50)	206	11.2 (49)	177	2.3 (14)	204	63.7 (78)
7850	341	7.6 (28)	7.9 (65)	346	8.7 (14)	309	3.2 (37)	341	73.9 (55)
7852	10	*	*	10	*	10	*	10	*
8101	583	16.6 (115)	3.1 (6)	629	20.3 (116)	457	6.3 (100)	610	47.4 (105)
8102	124	10.5 (73)	8.9 (81)	126	18.3 (114)	92	*	124	38.7 (115)
8105	537	8.9 (51)	6.5 (43)	553	11.6 (56)	468	3.6 (49)	541	67.7 (70)
8106	129	13.2 (102)	6.2 (37)	140	11.4 (53)	111	5.4 (81)	133	57.9 (92)
8150	8	*	*	9	*	8	*	8	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8151	994	10.1 (67)	5.6 (31)	1,015	14.2 (99)	851	4.5 (67)	954	79.7 (25)
8152	17	*	*	15	*	12	*	16	*
8153	2	*	*	2	*	2	*	2	*
8155	10	*	*	10	*	10	*	10	*
8157	2	*	*	2	*	2	*	2	*
8201	815	11.0 (80)	6.4 (39)	832	11.1 (46)	718	4.9 (71)	817	58.8 (90)
8202	1,108	8.3 (39)	6.7 (47)	1,230	8.5 (11)	1,002	5.0 (73)	1,220	48.1 (103)
8203	166	9.6 (59)	5.4 (28)	170	12.4 (69)	146	2.1 (10)	166	78.3 (32)
8250	252	11.9 (89)	4.0 (15)	255	16.1 (107)	208	5.3 (78)	253	68.8 (67)
8252	55	*	*	54	*	45	*	55	*
8255	786	8.5 (45)	7.5 (57)	796	11.9 (63)	685	3.5 (46)	789	60.5 (86)
8257	559	13.4 (103)	4.1 (19)	617	11.5 (55)	481	7.1 (104)	601	76.2 (43)
8263	240	8.3 (39)	9.6 (92)	243	7.8 (7)	220	3.2 (37)	239	62.3 (85)
8264	213	10.8 (78)	8.5 (76)	222	12.6 (74)	185	2.7 (23)	214	57.0 (94)
8266	7	*	*	9	*	6	*	9	*
8267	276	7.6 (28)	6.9 (50)	280	11.1 (46)	244	2.9 (29)	276	56.5 (95)
8901	350	14.9 (111)	4.0 (15)	363	16.8 (111)	288	5.6 (87)	352	47.7 (104)
8902	433	14.1 (109)	3.5 (11)	446	16.4 (108)	358	5.6 (87)	434	52.8 (99)
8903	363	13.5 (105)	3.0 (5)	376	15.7 (106)	301	7.0 (103)	367	40.9 (113)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8904	727	12.4 (96)	2.5 (3)	757	16.8 (111)	596	5.7 (91)	737	54.8 (97)
8905	447	7.4 (21)	8.3 (72)	497	12.5 (71)	381	3.4 (40)	486	51.6 (101)
8908	16	*	*	18	*	13	*	16	*
8909	173	15.0 (112)	2.3 (2)	175	20.0 (115)	136	6.6 (102)	169	55.0 (96)
9016	7	*	*	7	*	6	*	7	*
9018	3	*	*	3	*	3	*	3	*
9045	533	9.0 (52)	9.4 (86)	544	12.9 (76)	456	3.5 (46)	540	47.4 (105)
9050	300	12.3 (93)	4.0 (15)	312	16.7 (110)	246	4.9 (71)	304	52.3 (100)
9051	513	4.9 (4)	8.6 (78)	532	11.7 (59)	442	2.3 (14)	521	84.5 (9)
9053	407	12.5 (97)	4.7 (21)	427	14.1 (96)	353	6.2 (99)	407	45.5 (107)
9055	12	*	*	12	*	9	*	12	*
9058	325	12.3 (93)	3.4 (9)	338	14.2 (99)	277	5.1 (74)	334	53.0 (98)
9066	387	13.7 (107)	1.8 (1)	402	12.9 (76)	333	7.5 (106)	388	44.8 (108)
9068	287	14.6 (110)	3.5 (11)	295	15.6 (105)	238	8.0 (107)	288	43.1 (112)
9069	920	13.0 (100)	3.6 (13)	943	15.1 (103)	779	5.6 (87)	920	39.0 (114)
9070	354	15.8 (113)	3.4 (9)	362	16.6 (109)	289	9.0 (109)	356	43.8 (110)
9072	597	16.4 (114)	3.2 (8)	628	14.8 (102)	505	8.3 (108)	598	43.8 (110)
9080	955	13.0 (100)	3.6 (13)	973	15.4 (104)	800	5.4 (81)	959	48.6 (102)
9081	11	*	*	11	*	10	*	11	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9082	49	*	*	52	*	42	*	49	*
9083	17	*	*	20	*	16	*	18	*
9084	13	*	*	13	*	10	*	13	*
9085	3	*	*	3	*	3	*	3	*
Michigan	**50,526**	**10.2**	**6.7**	**52,025**	**12.4**	**43,785**	**4.3**	**51,003**	**66.7**
National (Prior Year)	**56,577**	**9.7**	**7.0**	**58,441**	**12.2**	**49,086**	**4.1**	**56,962**	**58.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	114	95.6	0.0	1.8	0.0	0.0	2.6	0.0
0102	527	91.7	0.4	2.7	0.2	0.0	4.7	0.4
0103	452	92.5	0.4	3.5	0.0	0.0	3.3	0.2
0104	169	95.3	0.6	2.4	0.0	0.6	0.6	0.6
0250	124	87.1	0.0	1.6	1.6	0.0	9.7	0.0
0251	102	91.2	0.0	0.0	3.9	0.0	4.9	0.0
0252	139	46.8	1.4	3.6	1.4	0.7	45.3	0.7
0253	179	79.9	0.6	0.6	7.8	0.0	11.2	0.0
0401	549	93.8	0.5	2.6	0.0	0.4	2.0	0.7
0402	412	83.3	1.0	2.4	0.0	1.0	12.4	0.0
0403	136	97.1	0.0	1.5	0.7	0.0	0.0	0.7
0404	140	93.6	0.7	0.7	0.0	0.7	4.3	0.0
0501	605	95.9	0.2	1.7	0.0	1.0	1.3	0.0
0503	115	92.2	0.0	4.3	0.0	0.0	2.6	0.9
0504	545	87.5	1.5	1.5	3.5	0.6	5.5	0.0
0505	342	84.2	0.6	4.7	0.3	0.9	9.4	0.0
0506	222	94.6	0.0	0.9	0.5	0.0	4.1	0.0
0507	88	*	*	*	*	*	*	*
0508	34	*	*	*	*	*	*	*
0509	11	*	*	*	*	*	*	*
0650	1,186	80.7	4.2	3.9	2.5	3.0	5.6	0.1
0651	668	91.6	0.0	4.2	0.1	0.3	2.4	1.3
0652	317	93.4	0.0	2.2	0.3	0.6	3.2	0.3

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0653	476	96.4	0.2	1.9	0.0	0.0	1.3	0.2
0654	385	94.5	1.0	2.1	0.5	0.0	1.3	0.5
0655	392	90.8	0.8	2.6	0.0	0.8	3.8	1.3
0657	77	*	*	*	*	*	*	*
0701	466	88.8	1.1	0.9	0.0	6.0	3.0	0.2
0702	85	*	*	*	*	*	*	*
0703	253	93.3	1.6	2.4	0.8	0.4	0.8	0.8
0704	68	*	*	*	*	*	*	*
0705	140	95.7	0.0	0.0	0.0	0.0	4.3	0.0
0706	2	*	*	*	*	*	*	*
0850	1,037	88.5	3.8	5.1	0.0	0.6	1.9	0.1
0851	821	94.8	0.2	3.0	0.1	0.6	1.0	0.1
0901	1,829	81.3	3.2	9.5	0.0	0.3	5.0	0.7
0902	43	*	*	*	*	*	*	*
1051	1,557	87.7	1.2	6.2	1.3	0.8	2.5	0.4
1101	1,948	38.5	50.1	6.9	0.2	0.5	2.6	1.3
1102	960	74.3	11.9	6.4	0.4	1.1	4.7	1.3
1106	96	*	*	*	*	*	*	*
1250	1,099	85.7	0.7	10.4	0.0	0.5	2.0	0.6
1251	937	91.5	0.2	2.9	0.0	0.1	3.5	1.8
1252	550	61.1	0.7	34.9	0.0	0.4	2.0	0.9
1253	1,006	81.0	6.0	8.8	0.1	0.2	3.1	0.8
1350	2,743	59.1	19.4	7.5	0.2	6.6	6.3	0.9
1351	168	92.9	0.6	4.2	0.0	0.6	1.8	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1352	382	64.7	25.4	5.5	0.3	0.0	3.7	0.5
1401	280	87.5	1.1	6.1	1.1	0.7	1.8	1.8
1402	167	54.5	1.8	22.2	9.0	1.2	8.4	3.0
1403	23	*	*	*	*	*	*	*
1404	5	*	*	*	*	*	*	*
1501	951	47.8	5.4	39.0	0.1	0.4	3.0	4.2
1503	1,269	61.5	6.0	24.7	0.3	0.7	5.4	1.3
1504	2,503	43.6	5.2	42.8	0.0	5.1	2.3	1.0
1505	569	85.6	2.3	8.6	0.0	0.5	2.3	0.7
1506	682	77.1	1.0	15.4	0.1	0.6	3.5	2.2
1507	662	86.1	1.2	5.3	0.3	2.0	4.5	0.6
1508	122	6.6	0.0	86.9	0.0	0.0	0.0	6.6
1509	539	48.2	2.4	43.0	0.0	0.2	2.0	4.1
1511	786	91.0	1.0	3.3	0.3	1.0	2.7	0.6
1512	391	91.3	0.5	2.3	0.5	0.8	4.1	0.5
1513	58	*	*	*	*	*	*	*
1514	108	86.1	0.0	9.3	0.0	1.9	2.8	0.0
1701	752	58.9	0.4	2.1	17.6	0.9	20.1	0.0
1950	666	93.1	0.3	4.2	0.2	0.2	1.5	0.6
1952	127	96.9	0.0	2.4	0.0	0.0	0.0	0.8
1953	339	90.9	0.3	5.3	0.0	0.0	3.2	0.3
1954	655	85.8	0.3	12.4	0.0	0.3	1.1	0.2
1955	81	*	*	*	*	*	*	*
1957	621	85.3	1.0	11.0	0.2	0.3	2.3	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1964	15	*	*	*	*	*	*	*
2150	773	85.4	0.4	2.7	0.6	1.2	9.2	0.5
2151	278	82.0	1.1	6.1	1.1	0.0	7.2	2.5
2251	475	92.2	0.0	2.1	0.4	0.6	3.2	1.5
2252	212	94.8	0.0	2.4	0.0	0.0	1.4	1.4
2554	104	91.3	4.8	2.9	0.0	0.0	1.0	0.0
2555	8,889	48.7	39.4	4.7	0.1	0.5	6.0	0.6
2556	85	*	*	*	*	*	*	*
2557	2	*	*	*	*	*	*	*
2558	2	*	*	*	*	*	*	*
2559	6	*	*	*	*	*	*	*
3001	1,156	16.5	65.7	13.7	0.1	1.0	2.7	0.3
3002	269	21.6	52.4	22.7	0.0	0.0	3.3	0.0
3004	139	12.2	0.0	85.6	0.0	0.0	0.7	1.4
3007	32	*	*	*	*	*	*	*
3009	8	*	*	*	*	*	*	*
3201	553	93.5	0.2	4.2	0.2	0.0	1.3	0.7
3301	4,547	40.5	29.7	15.0	0.2	6.4	7.5	0.7
3304	122	34.4	19.7	8.2	0.0	33.6	4.1	0.0
3305	162	56.8	16.7	9.9	0.0	10.5	4.9	1.2
3306	272	84.9	4.0	6.3	0.0	0.7	2.9	1.1
3308	78	*	*	*	*	*	*	*
3309	44	*	*	*	*	*	*	*
3312	450	37.8	28.2	24.0	1.1	2.0	6.2	0.7

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3550	79	*	*	*	*	*	*	*
3551	28	*	*	*	*	*	*	*
3552	58	*	*	*	*	*	*	*
3701	1,190	90.7	0.2	6.8	0.1	0.3	1.9	0.0
3850	3,383	73.7	14.0	5.6	0.2	0.6	5.6	0.3
3901	126	31.7	50.0	6.3	1.6	2.4	4.0	4.0
3902	65	*	*	*	*	*	*	*
3903	2	*	*	*	*	*	*	*
3950	2,209	61.9	22.8	7.4	0.2	1.3	5.6	0.8
3954	9	*	*	*	*	*	*	*
4001	1,468	32.1	54.4	8.0	0.1	1.1	4.1	0.3
4002	1,030	54.2	24.8	13.7	0.0	0.8	6.2	0.4
4003	8	*	*	*	*	*	*	*
4004	5	*	*	*	*	*	*	*
4103	67	*	*	*	*	*	*	*
4106	2,402	17.6	41.7	34.3	0.1	0.4	5.0	0.9
4107	3,619	52.6	23.4	16.2	0.3	1.9	5.6	0.1
4109	1,135	87.8	1.1	6.2	0.0	0.7	4.1	0.1
4112	614	1.0	0.5	97.1	0.0	0.7	0.2	0.7
4113	950	9.4	16.3	70.8	0.1	1.5	1.9	0.0
4115	3,696	41.1	24.1	17.7	0.4	10.7	5.4	0.5
4120	72	*	*	*	*	*	*	*
4160	80	*	*	*	*	*	*	*
4351	616	89.6	0.3	5.8	0.0	0.8	3.1	0.3

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4352	217	71.4	17.5	6.0	0.0	0.0	4.1	0.9
4353	506	66.2	0.4	29.1	0.2	0.0	3.2	1.0
4354	806	90.1	4.5	2.2	0.1	0.1	2.4	0.6
4355	398	79.6	1.8	6.8	1.8	0.5	9.3	0.3
4356	541	84.8	0.7	10.0	0.2	0.0	3.9	0.4
4358	162	26.5	0.0	68.5	0.0	0.0	1.9	3.1
4359	535	72.1	0.2	25.0	0.2	0.2	0.7	1.5
4360	237	95.4	0.0	3.8	0.4	0.0	0.4	0.0
4361	290	93.8	0.3	0.3	0.3	0.3	4.5	0.3
4362	302	94.4	0.0	1.7	0.3	0.7	3.0	0.0
4363	995	94.4	0.3	2.2	0.0	0.3	2.6	0.2
4401	1,237	88.2	0.4	6.0	0.1	0.3	4.4	0.6
4501	1,003	93.2	0.8	3.4	0.2	0.2	2.0	0.2
4601	1,665	75.4	1.6	17.2	0.0	0.4	4.3	0.9
4602	23	*	*	*	*	*	*	*
4701	1,271	89.5	0.7	4.8	0.2	1.5	2.4	0.9
5050	1,963	55.3	35.4	2.8	0.0	1.4	4.7	0.5
5051	2,970	53.4	35.1	3.5	0.0	2.5	5.1	0.4
5052	2,972	45.4	34.7	1.9	0.1	13.4	4.2	0.3
5055	1,235	96.1	0.4	0.1	0.0	2.8	0.1	0.5
5101	2,163	71.9	10.1	12.3	0.0	1.6	4.1	0.0
5202	821	85.4	0.6	2.9	1.1	1.2	7.9	0.9
5203	43	*	*	*	*	*	*	*
5204	30	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
5601	1,237	93.6	0.7	2.0	0.1	0.7	2.3	0.6
5801	1,937	83.2	6.6	5.5	0.2	0.4	3.4	0.7
5803	139	89.2	1.4	6.5	0.0	1.4	1.4	0.0
5804	31	*	*	*	*	*	*	*
5806	11	*	*	*	*	*	*	*
6101	32	*	*	*	*	*	*	*
6102	135	91.1	0.0	4.4	0.0	0.0	3.7	0.7
6103	4,357	57.2	27.3	8.4	0.3	0.4	6.0	0.4
6104	102	39.2	51.0	6.9	0.0	1.0	2.0	0.0
6105	30	*	*	*	*	*	*	*
6301	4,679	44.6	28.5	20.0	0.1	2.7	3.7	0.4
6312	2,644	35.2	55.1	3.3	0.2	2.7	3.3	0.2
6316	1,349	68.9	18.5	7.6	0.1	2.4	2.3	0.2
6320	139	62.6	23.7	4.3	0.0	5.0	3.6	0.7
6325	154	93.5	1.3	3.2	0.0	0.6	1.3	0.0
6330	721	25.1	54.0	16.8	0.0	1.2	2.9	0.0
6335	959	77.2	14.0	3.5	0.2	3.9	1.1	0.1
6340	66	*	*	*	*	*	*	*
6345	28	*	*	*	*	*	*	*
6350	18	*	*	*	*	*	*	*
6360	160	22.5	59.4	13.1	0.0	2.5	2.5	0.0
7304	161	92.5	0.0	5.0	0.0	0.0	1.9	0.6
7306	2,773	44.3	34.7	15.8	0.0	0.5	3.9	0.6
7308	143	83.9	1.4	10.5	0.0	0.0	4.2	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
7310	44	*	*	*	*	*	*	*
7401	2,567	82.8	5.8	5.5	0.2	0.6	4.7	0.4
7601	707	92.9	0.1	4.7	0.1	0.0	1.4	0.7
7850	1,241	87.9	0.6	5.6	0.1	0.4	4.8	0.4
7852	45	*	*	*	*	*	*	*
8101	2,026	6.0	90.5	0.7	0.1	0.4	2.2	0.1
8102	526	13.7	7.6	75.9	0.0	0.0	2.5	0.4
8105	1,915	32.4	60.3	2.9	0.1	1.0	3.0	0.3
8106	553	25.1	63.3	6.5	0.2	0.4	4.0	0.5
8150	50	*	*	*	*	*	*	*
8151	3,583	37.3	38.7	10.1	0.1	4.3	8.7	0.6
8152	159	39.6	44.0	8.8	0.0	0.6	6.9	0.0
8153	19	*	*	*	*	*	*	*
8155	48	*	*	*	*	*	*	*
8157	16	*	*	*	*	*	*	*
8201	3,061	50.0	36.3	4.8	0.2	2.5	5.5	0.7
8202	4,137	86.7	9.0	2.5	0.0	0.7	0.6	0.3
8203	554	62.1	27.6	3.6	0.2	0.9	4.9	0.7
8250	862	22.0	66.5	4.8	0.0	1.0	5.5	0.2
8252	150	1.3	94.7	0.0	0.0	1.3	2.0	0.7
8255	3,115	57.3	29.1	6.4	0.2	2.9	3.5	0.5
8257	1,713	51.2	14.9	0.8	0.1	31.5	0.9	0.6
8258	1	*	*	*	*	*	*	*
8263	877	70.4	12.4	11.7	0.0	1.5	3.4	0.6

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8264	747	58.9	29.9	4.3	0.1	2.5	4.0	0.3
8265	58	*	*	*	*	*	*	*
8266	46	*	*	*	*	*	*	*
8267	501	59.5	17.4	18.2	0.2	0.2	3.2	1.0
8901	1,212	1.4	95.0	1.0	0.1	0.5	1.9	0.1
8902	1,366	4.0	91.7	1.0	0.0	0.2	2.9	0.2
8903	1,252	2.2	94.7	0.8	0.0	0.4	1.6	0.3
8904	2,236	2.2	92.9	0.8	0.1	1.6	2.2	0.1
8905	1,690	58.3	28.2	10.8	0.1	0.9	1.5	0.2
8908	33	*	*	*	*	*	*	*
8909	266	0.8	94.0	1.9	0.0	0.4	2.3	0.8
9012	21	*	*	*	*	*	*	*
9013	7	*	*	*	*	*	*	*
9016	119	1.7	96.6	0.0	0.0	0.0	1.7	0.0
9017	22	*	*	*	*	*	*	*
9018	115	3.5	94.8	0.0	0.0	0.0	1.7	0.0
9041	9	*	*	*	*	*	*	*
9045	1,849	9.4	29.7	57.8	0.0	0.3	2.7	0.2
9050	1,128	10.5	69.9	16.9	0.0	0.7	1.9	0.1
9051	1,775	9.3	7.9	80.8	0.0	0.2	1.6	0.2
9053	1,665	3.1	92.0	0.5	0.1	3.2	1.0	0.2
9055	59	*	*	*	*	*	*	*
9058	1,103	26.7	60.0	1.3	0.1	10.6	1.0	0.3
9066	1,337	1.9	96.9	0.5	0.0	0.1	0.6	0.1

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 3A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9068	1,114	2.5	95.7	0.4	0.0	0.3	0.9	0.3
9069	3,328	1.5	97.1	0.6	0.0	0.2	0.4	0.1
9070	1,244	3.3	89.5	0.3	0.1	5.3	1.4	0.1
9072	2,169	1.8	94.4	1.0	0.0	0.7	1.5	0.5
9080	2,954	19.9	76.1	1.8	0.0	0.1	1.7	0.4
9081	84	*	*	*	*	*	*	*
9082	58	*	*	*	*	*	*	*
9083	20	*	*	*	*	*	*	*
9084	13	*	*	*	*	*	*	*
9085	3	*	*	*	*	*	*	*
Michigan	**180,927**	**53.2**	**29.4**	**10.4**	**0.3**	**2.2**	**3.9**	**0.6**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	114	0.0	2.6	8.8	63.2	24.6	0.9	0.0
0102	527	0.0	5.1	6.8	66.6	20.5	0.9	0.0
0103	452	0.0	3.1	8.0	69.0	18.8	1.1	0.0
0104	169	0.0	1.8	11.2	72.2	13.6	1.2	0.0
0250	124	0.0	4.0	3.2	73.4	19.4	0.0	0.0
0251	102	0.0	5.9	4.9	61.8	25.5	1.0	1.0
0252	139	0.0	2.2	2.9	66.2	28.1	0.7	0.0
0253	179	1.1	2.8	11.2	67.0	17.9	0.0	0.0
0401	549	0.2	3.6	9.5	63.4	22.0	1.3	0.0
0402	412	0.0	2.2	7.3	71.6	18.0	1.0	0.0
0403	136	0.0	2.9	8.8	69.9	18.4	0.0	0.0
0404	140	0.0	2.1	10.7	62.1	20.7	4.3	0.0
0501	605	0.2	3.8	9.1	62.8	23.3	0.8	0.0
0503	115	0.0	1.7	7.8	56.5	32.2	1.7	0.0
0504	545	0.0	1.3	9.4	67.3	21.5	0.6	0.0
0505	342	0.0	2.9	11.7	59.6	24.0	1.8	0.0
0506	222	0.0	1.8	10.8	67.6	18.9	0.9	0.0
0507	88	*	*	*	*	*	*	*
0508	34	*	*	*	*	*	*	*
0509	11	*	*	*	*	*	*	*
0650	1,186	0.0	2.7	8.0	66.0	22.5	0.7	0.1
0651	668	0.3	3.4	11.5	66.6	16.3	1.8	0.0
0652	317	0.0	4.1	10.4	64.4	21.1	0.0	0.0
0653	476	0.0	3.6	8.4	68.5	18.1	1.1	0.4

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0654	385	0.3	5.5	9.6	60.8	22.9	1.0	0.0
0655	392	0.3	3.1	9.9	65.3	20.7	0.8	0.0
0657	77	*	*	*	*	*	*	*
0701	466	0.0	1.3	8.8	62.2	26.4	1.3	0.0
0702	85	*	*	*	*	*	*	*
0703	253	0.0	3.6	9.1	64.0	22.5	0.8	0.0
0704	68	*	*	*	*	*	*	*
0705	140	0.0	4.3	8.6	54.3	30.0	2.9	0.0
0706	2	*	*	*	*	*	*	*
0850	1,037	0.2	2.0	8.8	63.5	24.0	1.2	0.3
0851	821	0.1	4.1	9.5	68.6	17.1	0.6	0.0
0901	1,829	0.1	3.2	8.9	64.4	22.6	0.9	0.0
0902	43	*	*	*	*	*	*	*
1051	1,557	0.0	2.7	7.1	62.9	25.5	1.3	0.4
1101	1,948	0.5	4.7	9.1	64.4	20.0	1.0	0.3
1102	960	0.6	3.3	10.2	63.0	20.9	1.6	0.3
1106	96	*	*	*	*	*	*	*
1250	1,099	0.1	3.9	9.2	62.9	21.5	2.4	0.1
1251	937	0.0	3.1	10.4	69.6	16.1	0.9	0.0
1252	550	0.0	4.2	12.7	59.3	22.7	1.1	0.0
1253	1,006	0.1	4.2	12.0	63.0	20.0	0.7	0.0
1350	2,743	0.1	3.3	9.6	64.7	20.7	1.2	0.3
1351	168	0.0	1.2	4.8	71.4	22.0	0.6	0.0
1352	382	0.0	3.7	7.1	70.9	16.8	1.6	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1401	280	0.0	4.3	7.1	67.5	18.9	1.8	0.4
1402	167	0.0	2.4	8.4	55.7	31.7	0.6	1.2
1403	23	*	*	*	*	*	*	*
1404	5	*	*	*	*	*	*	*
1501	951	0.1	4.1	10.2	62.4	20.8	2.4	0.0
1503	1,269	0.1	4.4	9.3	63.4	21.7	1.1	0.0
1504	2,503	0.1	3.4	7.7	63.7	23.2	1.8	0.0
1505	569	0.0	1.9	7.6	63.4	25.3	1.4	0.4
1506	682	0.1	3.5	7.8	63.5	23.5	1.6	0.0
1507	662	0.0	2.1	6.2	64.4	25.5	1.8	0.0
1508	122	0.0	5.7	5.7	46.7	35.2	5.7	0.8
1509	539	0.0	4.3	9.1	62.5	21.3	2.8	0.0
1511	786	0.4	3.8	10.9	64.2	19.3	1.3	0.0
1512	391	0.3	3.1	6.9	64.5	25.1	0.3	0.0
1513	58	*	*	*	*	*	*	*
1514	108	0.0	2.8	4.6	50.9	40.7	0.9	0.0
1701	752	0.0	2.0	8.8	63.7	24.3	1.2	0.0
1950	666	0.0	3.0	10.4	68.5	17.1	1.1	0.0
1952	127	0.8	3.1	3.9	69.3	20.5	2.4	0.0
1953	339	0.3	1.8	10.0	72.0	15.6	0.3	0.0
1954	655	0.2	2.6	8.9	65.5	21.2	1.7	0.0
1955	81	*	*	*	*	*	*	*
1957	621	0.0	2.4	8.5	63.0	24.3	1.8	0.0
1964	15	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2150	773	0.1	2.5	8.0	64.8	23.5	1.0	0.0
2151	278	0.0	4.3	14.4	56.8	23.4	0.7	0.4
2251	475	0.0	3.6	8.6	61.3	24.8	1.5	0.2
2252	212	0.0	3.3	9.9	58.5	26.4	1.9	0.0
2554	104	0.0	2.9	9.6	65.4	20.2	1.9	0.0
2555	8,889	0.2	3.0	9.9	66.4	19.4	1.1	0.0
2556	85	*	*	*	*	*	*	*
2557	2	*	*	*	*	*	*	*
2558	2	*	*	*	*	*	*	*
2559	6	*	*	*	*	*	*	*
3001	1,156	0.4	4.8	10.2	64.3	18.8	1.4	0.2
3002	269	0.4	4.1	10.4	62.1	22.3	0.7	0.0
3004	139	0.0	2.2	7.9	58.3	28.1	3.6	0.0
3007	32	*	*	*	*	*	*	*
3009	8	*	*	*	*	*	*	*
3201	553	0.0	2.2	10.1	63.1	23.7	0.7	0.2
3301	4,547	0.2	3.1	8.6	63.3	23.0	1.6	0.2
3304	122	0.0	0.0	2.5	50.8	45.1	1.6	0.0
3305	162	0.0	0.6	4.3	56.2	35.2	3.7	0.0
3306	272	0.4	7.7	7.0	56.6	28.3	0.0	0.0
3308	78	*	*	*	*	*	*	*
3309	44	*	*	*	*	*	*	*
3312	450	0.2	3.3	6.4	62.7	25.3	2.0	0.0
3550	79	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3551	28	*	*	*	*	*	*	*
3552	58	*	*	*	*	*	*	*
3701	1,190	0.2	3.7	10.9	67.6	17.1	0.6	0.0
3850	3,383	0.1	3.5	9.1	66.9	18.9	1.1	0.3
3901	126	0.0	2.4	9.5	66.7	19.8	1.6	0.0
3902	65	*	*	*	*	*	*	*
3903	2	*	*	*	*	*	*	*
3950	2,209	0.0	3.4	10.3	62.2	22.4	1.6	0.1
3954	9	*	*	*	*	*	*	*
4001	1,468	0.3	6.0	10.5	62.3	19.5	1.4	0.0
4002	1,030	0.1	4.7	9.1	60.5	24.3	1.3	0.1
4003	8	*	*	*	*	*	*	*
4004	5	*	*	*	*	*	*	*
4103	67	*	*	*	*	*	*	*
4106	2,402	0.4	4.8	9.3	60.5	23.2	1.7	0.1
4107	3,619	0.2	3.4	7.1	63.8	23.8	1.5	0.2
4109	1,135	0.2	3.6	7.1	63.0	24.0	2.1	0.0
4112	614	0.0	4.9	10.6	48.2	31.4	4.6	0.3
4113	950	0.4	4.4	8.3	44.2	32.8	1.9	7.9
4115	3,696	0.1	2.2	6.3	62.3	27.1	1.9	0.1
4120	72	*	*	*	*	*	*	*
4160	80	*	*	*	*	*	*	*
4351	616	0.2	4.7	10.2	65.7	18.2	1.0	0.0
4352	217	0.0	5.1	12.4	67.7	13.8	0.9	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4353	506	0.0	4.2	10.3	64.0	20.6	1.0	0.0
4354	806	0.0	3.6	10.0	68.2	17.0	1.1	0.0
4355	398	0.0	5.5	14.3	55.8	22.9	1.3	0.3
4356	541	0.2	3.5	8.3	70.8	16.3	0.9	0.0
4358	162	0.0	4.3	9.3	63.0	22.2	1.2	0.0
4359	535	0.2	2.8	9.3	59.6	24.7	3.4	0.0
4360	237	0.0	3.4	9.3	67.5	18.1	1.7	0.0
4361	290	0.0	3.8	10.3	63.1	22.4	0.3	0.0
4362	302	0.3	3.3	7.3	66.6	21.5	1.0	0.0
4363	995	0.2	3.1	9.2	67.5	19.7	0.2	0.0
4401	1,237	0.2	2.6	7.8	68.6	19.8	1.0	0.2
4501	1,003	0.2	3.2	8.7	66.4	20.1	1.3	0.1
4601	1,665	0.2	3.5	9.8	65.6	19.8	1.0	0.0
4602	23	*	*	*	*	*	*	*
4701	1,271	0.0	1.8	6.0	63.6	27.1	1.5	0.0
5050	1,963	0.1	1.9	8.7	61.0	26.7	1.7	0.0
5051	2,970	0.1	1.5	4.9	66.7	25.0	1.8	0.0
5052	2,972	0.1	2.4	6.1	63.5	25.9	2.0	0.0
5055	1,235	0.0	0.2	0.7	56.6	38.1	4.2	0.1
5101	2,163	0.1	1.2	5.7	62.5	27.9	2.5	0.0
5202	821	0.0	3.8	6.7	62.5	25.2	1.8	0.0
5203	43	*	*	*	*	*	*	*
5204	30	*	*	*	*	*	*	*
5601	1,237	0.2	2.5	7.4	66.8	21.9	1.1	0.1

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5801	1,937	0.1	4.1	10.5	64.0	20.2	1.1	0.1
5803	139	0.0	1.4	2.9	64.7	30.2	0.7	0.0
5804	31	*	*	*	*	*	*	*
5806	11	*	*	*	*	*	*	*
6101	32	*	*	*	*	*	*	*
6102	135	0.0	6.7	8.1	60.7	23.0	1.5	0.0
6103	4,357	0.2	3.9	8.9	65.8	20.0	1.1	0.0
6104	102	0.0	2.0	13.7	67.6	16.7	0.0	0.0
6105	30	*	*	*	*	*	*	*
6301	4,679	0.1	3.0	7.4	63.2	24.4	1.8	0.1
6312	2,644	0.0	1.7	5.4	58.6	31.9	2.4	0.0
6316	1,349	0.0	0.9	5.1	59.7	31.7	2.4	0.1
6320	139	0.0	0.0	0.7	51.1	40.3	7.2	0.7
6325	154	0.0	2.6	5.8	66.2	25.3	0.0	0.0
6330	721	0.4	3.6	10.5	66.0	18.0	1.4	0.0
6335	959	0.0	1.7	5.2	51.8	37.6	3.6	0.0
6340	66	*	*	*	*	*	*	*
6345	28	*	*	*	*	*	*	*
6350	18	*	*	*	*	*	*	*
6360	160	0.0	0.6	7.5	75.0	15.6	1.3	0.0
7304	161	0.0	1.9	5.6	67.1	25.5	0.0	0.0
7306	2,773	0.3	3.1	8.1	66.7	20.6	1.3	0.0
7308	143	0.0	2.8	14.7	61.5	21.0	0.0	0.0
7310	44	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
7401	2,567	0.1	3.3	9.5	64.8	21.0	1.3	0.0
7601	707	0.0	5.4	9.2	63.4	21.5	0.6	0.0
7850	1,241	0.2	3.3	10.3	66.8	18.3	1.0	0.0
7852	45	*	*	*	*	*	*	*
8101	2,026	0.1	3.6	9.1	64.6	20.9	1.8	0.0
8102	526	0.0	3.0	6.7	59.7	26.6	3.8	0.2
8105	1,915	0.1	2.1	6.0	61.4	28.4	2.0	0.1
8106	553	0.2	5.1	9.0	61.7	23.1	0.9	0.0
8150	50	*	*	*	*	*	*	*
8151	3,583	0.1	2.3	5.9	61.7	27.5	2.4	0.0
8152	159	0.0	3.1	17.0	73.6	5.7	0.6	0.0
8153	19	*	*	*	*	*	*	*
8155	48	*	*	*	*	*	*	*
8157	16	*	*	*	*	*	*	*
8201	3,061	0.1	2.3	7.3	63.9	24.5	1.6	0.3
8202	4,137	0.0	0.9	3.6	57.8	32.4	3.8	1.4
8203	554	0.5	1.3	8.5	63.0	24.4	2.2	0.2
8250	862	0.0	3.0	7.4	62.9	25.3	1.3	0.1
8252	150	0.0	4.0	11.3	65.3	18.0	1.3	0.0
8255	3,115	0.0	2.3	7.1	66.5	22.7	1.3	0.0
8257	1,713	0.1	1.3	4.6	57.7	31.5	4.4	0.4
8258	1	*	*	*	*	*	*	*
8263	877	0.1	2.2	6.0	63.9	26.0	1.7	0.1
8264	747	0.1	3.2	6.8	62.7	24.9	2.3	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8265	58	*	*	*	*	*	*	*
8266	46	*	*	*	*	*	*	*
8267	501	0.0	2.2	6.8	63.9	25.3	1.8	0.0
8901	1,212	0.2	3.5	8.9	65.2	20.5	1.7	0.1
8902	1,366	0.2	2.5	8.9	63.3	22.8	2.1	0.1
8903	1,252	0.2	4.6	7.9	64.7	20.8	1.7	0.2
8904	2,236	0.1	3.0	7.9	66.5	20.7	1.8	0.1
8905	1,690	0.0	1.3	5.1	60.7	30.3	2.5	0.1
8908	33	*	*	*	*	*	*	*
8909	266	0.4	3.4	8.6	58.3	27.8	1.5	0.0
9012	21	*	*	*	*	*	*	*
9013	7	*	*	*	*	*	*	*
9016	119	0.8	2.5	10.1	67.2	19.3	0.0	0.0
9017	22	*	*	*	*	*	*	*
9018	115	0.0	4.3	8.7	62.6	20.0	4.3	0.0
9041	9	*	*	*	*	*	*	*
9045	1,849	0.2	3.6	8.6	57.2	28.0	2.3	0.0
9050	1,128	0.4	3.5	8.2	61.9	23.0	2.9	0.0
9051	1,775	0.1	4.0	7.5	50.0	33.5	4.9	0.0
9053	1,665	0.1	2.8	7.2	67.9	20.2	1.9	0.0
9055	59	*	*	*	*	*	*	*
9058	1,103	0.0	2.4	4.4	59.9	29.5	3.8	0.0
9066	1,337	0.1	3.6	7.9	64.5	22.3	1.6	0.0
9068	1,114	0.0	3.8	9.7	63.6	21.3	1.5	0.1

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 4A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9069	3,328	0.1	3.2	7.5	66.0	21.2	1.9	0.1
9070	1,244	0.4	2.7	8.0	66.1	21.7	1.1	0.0
9072	2,169	0.0	4.1	8.1	67.3	18.9	1.6	0.0
9080	2,954	0.1	2.3	6.4	63.6	25.1	2.5	0.1
9081	84	*	*	*	*	*	*	*
9082	58	*	*	*	*	*	*	*
9083	20	*	*	*	*	*	*	*
9084	13	*	*	*	*	*	*	*
9085	3	*	*	*	*	*	*	*
Michigan	**180,927**	**0.1**	**3.0**	**8.0**	**63.4**	**23.6**	**1.7**	**0.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	102	5.9 (141)	67.6 (151)	55	*	*	16	*	27	*	104	4.8 (16)	104	9.6 (55)
0102	479	5.0 (110)	61.4 (119)	326	20.6 (98)	50.6 (53)	62	*	88	*	497	4.8 (16)	497	8.5 (32)
0103	398	3.0 (24)	59.5 (102)	279	16.8 (22)	52.7 (85)	47	*	86	*	420	8.3 (132)	420	12.6 (117)
0104	153	4.6 (96)	52.9 (21)	99	*	*	15	*	30	*	157	7.0 (91)	157	7.6 (22)
0250	121	2.5 (10)	67.8 (152)	76	*	*	14	*	23	*	121	9.9 (152)	121	9.1 (45)
0251	93	*	*	54	*	*	13	*	23	*	95	*	95	*
0252	138	1.4 (1)	53.6 (25)	121	23.1 (127)	52.1 (77)	15	*	24	*	124	5.6 (39)	124	14.5 (146)
0253	147	5.4 (128)	51.0 (14)	79	*	*	12	*	31	*	168	8.9 (140)	168	13.7 (134)
0401	523	3.8 (52)	56.6 (51)	462	20.1 (84)	51.1 (60)	81	*	113	23.0 (14)	512	3.1 (2)	511	7.4 (19)
0402	385	5.2 (120)	56.9 (55)	317	19.9 (81)	50.5 (52)	66	*	73	*	391	4.9 (19)	391	5.4 (7)
0403	130	6.2 (145)	55.4 (37)	104	18.3 (54)	48.1 (23)	20	*	29	*	126	7.1 (92)	126	11.1 (85)
0404	122	2.5 (10)	53.3 (24)	92	*	*	28	*	31	*	133	5.3 (29)	133	9.0 (42)
0501	459	4.4 (82)	57.5 (66)	318	18.6 (58)	52.2 (78)	98	*	101	36.6 (67)	571	4.2 (7)	570	9.6 (55)
0503	100	3.0 (24)	58.0 (74)	66	*	*	13	*	18	*	105	7.6 (109)	105	11.4 (91)
0504	458	3.9 (57)	50.7 (12)	266	20.7 (101)	51.1 (60)	89	*	65	*	512	5.1 (23)	513	8.6 (33)
0505	295	4.4 (82)	55.9 (39)	201	17.9 (42)	51.2 (64)	44	*	52	*	327	6.7 (78)	327	7.6 (22)
0506	145	9.0 (153)	57.2 (61)	72	*	*	19	*	34	*	204	5.4 (32)	205	6.3 (12)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0507	70	*	*	46	*	*	7	*	17	*	84	*	84	*
0508	20	*	*	11	*	*	4	*	3	*	34	*	34	*
0509	10	*	*	10	*	*	2	*	4	*	11	*	11	*
0650	1,106	3.7 (47)	58.7 (92)	812	18.7 (60)	51.1 (60)	169	32.5 (14)	233	30.0 (36)	1,108	7.6 (109)	1,108	8.4 (29)
0651	638	5.2 (120)	63.6 (140)	520	19.6 (76)	49.8 (42)	78	*	95	*	614	7.2 (97)	615	6.0 (10)
0652	261	3.4 (35)	58.2 (83)	199	18.1 (46)	52.8 (87)	43	*	45	*	306	8.2 (130)	306	10.5 (74)
0653	435	5.3 (123)	59.8 (106)	332	18.1 (46)	50.0 (46)	36	*	90	*	446	4.3 (10)	447	12.1 (107)
0654	350	4.3 (80)	62.0 (130)	270	18.1 (46)	54.4 (110)	42	*	93	*	355	5.6 (39)	355	8.7 (37)
0655	364	5.5 (129)	54.4 (30)	195	21.5 (111)	51.8 (71)	45	*	58	*	361	7.2 (97)	362	11.6 (97)
0657	56	*	*	40	*	*	9	*	12	*	69	*	70	*
0701	367	2.5 (10)	46.3 (1)	239	16.3 (16)	55.2 (120)	56	*	118	19.5 (6)	438	5.9 (52)	438	12.1 (107)
0702	75	*	*	54	*	*	9	*	21	*	77	*	77	*
0703	202	4.5 (89)	55.9 (39)	132	20.5 (94)	47.0 (17)	34	*	47	*	224	8.9 (140)	224	11.6 (97)
0704	62	*	*	37	*	*	4	*	16	*	60	*	60	*
0705	110	5.5 (129)	50.0 (6)	62	*	*	17	*	35	*	127	7.1 (92)	127	10.2 (68)
0706	2	*	*	2	*	*	0	*	1	*	2	*	2	*
0850	790	5.1 (115)	56.1 (43)	561	16.2 (14)	52.4 (82)	173	34.7 (18)	234	20.1 (8)	953	9.1 (145)	953	15.2 (153)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0851	627	4.9 (105)	58.2 (83)	369	14.1 (3)	59.1 (132)	104	30.8 (11)	157	16.6 (3)	756	6.9 (85)	754	13.3 (124)
0901	1,418	4.8 (102)	58.3 (85)	1,080	20.4 (90)	48.5 (25)	238	32.8 (15)	245	26.9 (21)	1,709	6.7 (78)	1,709	9.8 (60)
0902	30	*	*	18	*	*	2	*	9	*	41	*	41	*
1051	1,393	5.1 (115)	50.5 (10)	1,103	16.2 (14)	52.0 (75)	220	60.0 (70)	205	45.4 (87)	1,470	6.1 (60)	1,471	8.4 (29)
1101	1,697	4.0 (63)	58.6 (89)	1,291	18.7 (60)	52.6 (84)	314	47.5 (49)	267	39.0 (73)	1,776	5.7 (45)	1,777	13.4 (126)
1102	793	3.0 (24)	57.5 (66)	586	14.3 (4)	59.7 (133)	118	30.5 (10)	214	30.8 (42)	850	5.9 (52)	850	8.6 (33)
1106	80	*	*	65	*	*	14	*	21	*	88	*	88	*
1250	916	4.7 (98)	56.9 (55)	584	16.3 (16)	52.4 (82)	138	13.8 (1)	148	23.0 (14)	988	8.3 (132)	988	6.3 (12)
1251	801	5.7 (136)	57.4 (63)	559	18.2 (51)	48.5 (25)	112	32.1 (13)	157	23.6 (16)	856	4.2 (7)	857	6.5 (14)
1252	437	2.5 (10)	62.0 (130)	302	18.9 (66)	55.6 (124)	61	*	105	32.4 (53)	505	5.5 (37)	505	17.2 (158)
1253	805	3.2 (31)	60.0 (107)	586	17.1 (30)	54.9 (116)	130	41.5 (32)	176	29.0 (32)	911	6.6 (76)	914	12.0 (104)
1350	2,308	4.0 (63)	56.8 (54)	1,426	19.4 (71)	52.2 (78)	387	52.2 (58)	476	40.1 (76)	2,501	6.5 (72)	2,498	9.8 (60)
1351	143	5.6 (132)	53.1 (22)	111	18.9 (66)	51.4 (66)	26	*	33	*	161	6.8 (81)	161	7.5 (21)
1352	256	5.9 (141)	59.4 (101)	149	22.1 (121)	45.0 (9)	44	*	59	*	351	7.1 (92)	350	5.1 (6)
1401	269	3.0 (24)	58.7 (92)	219	23.7 (129)	53.0 (94)	28	*	25	*	254	3.9 (6)	255	5.5 (8)
1402	151	2.6 (17)	60.3 (111)	120	15.0 (8)	54.2 (108)	24	*	26	*	152	5.9 (52)	152	8.6 (33)
1403	22	*	*	19	*	*	4	*	5	*	22	*	22	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1404	2	*	*	0	*	*	0	*	0	*	0	*	0	*
1501	818	2.7 (18)	63.0 (139)	555	43.4 (135)	27.2 (2)	95	*	131	21.4 (13)	822	6.3 (67)	822	7.4 (19)
1503	1,015	4.0 (63)	58.4 (88)	671	18.6 (58)	50.8 (55)	147	43.5 (38)	158	31.0 (44)	1,129	5.3 (29)	1,126	9.9 (62)
1504	2,111	5.1 (115)	56.9 (55)	1,444	21.3 (109)	46.5 (14)	343	55.4 (65)	455	42.4 (81)	2,254	9.3 (148)	2,255	8.3 (28)
1505	483	2.3 (4)	54.0 (27)	324	14.5 (6)	54.9 (116)	72	*	131	30.5 (39)	534	6.9 (85)	534	9.7 (58)
1506	540	3.5 (38)	57.6 (69)	182	22.0 (120)	47.3 (19)	87	*	116	29.3 (34)	620	7.9 (119)	619	6.8 (17)
1507	516	3.1 (29)	56.0 (41)	284	17.3 (33)	50.7 (54)	83	*	112	33.9 (59)	610	8.0 (122)	608	13.5 (128)
1508	51	*	*	14	*	*	6	*	15	*	79	*	79	*
1509	437	2.1 (3)	65.4 (145)	218	25.2 (130)	47.2 (18)	52	*	70	*	435	9.9 (152)	436	7.6 (22)
1511	653	5.2 (120)	59.1 (97)	361	21.9 (118)	52.9 (92)	103	38.8 (28)	147	36.1 (65)	717	7.9 (119)	721	12.5 (115)
1512	285	4.2 (76)	59.3 (100)	118	22.9 (126)	47.5 (21)	44	*	82	*	368	4.3 (10)	368	13.6 (133)
1513	48	*	*	38	*	*	7	*	14	*	56	*	56	*
1514	95	*	*	68	*	*	19	*	43	*	103	6.8 (81)	103	11.7 (100)
1701	594	2.9 (22)	61.6 (123)	286	17.8 (40)	53.8 (105)	70	*	107	31.8 (49)	686	7.7 (113)	686	12.4 (113)
1950	629	4.0 (63)	59.1 (97)	532	21.6 (112)	52.8 (87)	86	*	161	16.8 (4)	618	6.8 (81)	620	13.5 (128)
1952	121	5.8 (139)	56.2 (45)	101	18.8 (64)	46.5 (14)	28	*	46	*	119	5.9 (52)	119	8.4 (29)
1953	309	5.5 (129)	50.8 (13)	264	22.3 (122)	44.7 (8)	42	*	88	*	320	8.4 (134)	320	10.3 (70)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1954	589	5.8 (139)	59.1 (97)	415	19.5 (74)	53.3 (97)	76	*	118	20.3 (9)	613	6.0 (58)	613	15.8 (155)
1955	72	*	*	46	*	*	13	*	15	*	77	*	77	*
1957	560	4.5 (89)	57.1 (59)	457	18.4 (56)	56.0 (126)	100	20.0 (3)	156	20.5 (10)	574	6.1 (60)	575	13.9 (138)
1964	14	*	*	11	*	*	3	*	5	*	14	*	15	*
2150	628	4.5 (89)	54.3 (29)	407	22.6 (124)	42.0 (5)	79	*	147	25.9 (18)	699	13.2 (155)	699	11.0 (82)
2151	197	3.0 (24)	48.2 (2)	119	20.2 (85)	52.9 (92)	35	*	48	*	234	5.6 (39)	234	8.1 (27)
2251	409	4.4 (82)	51.3 (16)	243	16.0 (13)	45.3 (11)	52	*	99	*	422	3.3 (3)	420	10.5 (74)
2252	167	2.4 (7)	56.3 (47)	72	*	*	22	*	33	*	187	5.3 (29)	188	14.9 (150)
2554	99	*	*	83	*	*	20	*	19	*	98	*	98	*
2555	7,725	5.3 (123)	55.8 (38)	4,658	17.9 (42)	53.3 (97)	1,531	46.4 (44)	1,450	32.6 (55)	8,300	7.4 (101)	8,296	13.5 (128)
2556	78	*	*	68	*	*	5	*	21	*	84	*	84	*
2557	0	*	*	0	*	*	2	*	0	*	1	*	1	*
2558	1	*	*	0	*	*	2	*	0	*	2	*	2	*
2559	4	*	*	2	*	*	2	*	0	*	6	*	6	*
3001	1,086	3.9 (57)	62.2 (135)	851	27.5 (132)	42.1 (7)	185	37.8 (26)	748	44.7 (85)	1,071	4.9 (19)	1,071	10.3 (70)
3002	238	2.5 (10)	75.2 (153)	179	22.3 (122)	46.9 (16)	39	*	178	37.6 (70)	252	3.6 (4)	252	11.5 (94)
3004	121	3.3 (33)	62.0 (130)	108	28.7 (133)	41.7 (4)	17	*	26	*	127	7.1 (92)	127	8.7 (37)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3007	26	*	*	16	*	*	1	*	8	*	21	*	21	*
3009	3	*	*	0	*	*	0	*	0	*	1	*	1	*
3201	467	4.1 (71)	61.9 (129)	381	17.8 (40)	49.9 (45)	55	*	104	21.2 (12)	511	9.2 (147)	509	12.8 (120)
3301	3,924	4.4 (82)	57.2 (61)	3,008	18.1 (46)	51.9 (74)	746	37.1 (25)	814	30.6 (41)	4,196	5.4 (32)	4,199	10.6 (77)
3304	98	*	*	77	*	*	14	*	16	*	117	13.7 (157)	117	2.6 (1)
3305	130	7.7 (151)	56.2 (45)	108	17.6 (38)	53.7 (104)	13	*	32	*	152	4.6 (13)	152	5.9 (9)
3306	247	6.9 (149)	56.3 (47)	209	12.9 (1)	61.2 (136)	42	*	55	*	256	7.8 (116)	256	9.0 (42)
3308	69	*	*	63	*	*	10	*	18	*	75	*	75	*
3309	39	*	*	32	*	*	6	*	16	*	43	*	42	*
3312	386	4.9 (105)	62.4 (137)	306	21.6 (112)	51.6 (68)	94	*	73	*	424	7.5 (105)	425	12.0 (104)
3550	78	*	*	57	*	*	7	*	18	*	70	*	70	*
3551	26	*	*	21	*	*	3	*	11	*	28	*	28	*
3552	52	*	*	38	*	*	4	*	14	*	55	*	55	*
3701	923	4.9 (105)	58.6 (89)	730	17.1 (30)	53.2 (96)	183	16.4 (2)	230	11.3 (1)	1,090	5.2 (25)	1,089	12.0 (104)
3850	2,516	5.9 (141)	58.0 (74)	1,617	18.1 (46)	51.2 (64)	479	49.9 (55)	653	40.0 (74)	3,109	6.1 (60)	3,110	10.0 (65)
3901	96	*	*	71	*	*	16	*	40	*	118	5.9 (52)	118	4.2 (4)
3902	53	*	*	42	*	*	3	*	18	*	64	*	64	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3903	2	*	*	1	*	*	0	*	1	*	2	*	2	*
3950	1,744	3.8 (52)	58.3 (85)	1,206	16.3 (16)	56.0 (126)	334	42.8 (35)	492	31.1 (45)	1,994	9.0 (144)	1,995	9.7 (58)
3954	7	*	*	5	*	*	1	*	2	*	8	*	8	*
4001	1,251	4.0 (63)	58.8 (95)	783	23.4 (128)	49.4 (36)	162	55.6 (66)	299	43.5 (83)	1,301	7.8 (116)	1,302	11.4 (91)
4002	940	4.9 (105)	59.5 (102)	668	19.5 (74)	55.7 (125)	126	40.5 (30)	214	26.2 (19)	952	8.9 (140)	952	7.8 (25)
4003	7	*	*	5	*	*	2	*	1	*	8	*	8	*
4004	5	*	*	3	*	*	0	*	0	*	5	*	5	*
4103	46	*	*	31	*	*	6	*	10	*	59	*	59	*
4106	1,867	3.6 (45)	60.0 (107)	1,206	21.1 (106)	48.5 (25)	371	54.2 (62)	521	41.7 (79)	2,179	5.5 (37)	2,182	9.2 (48)
4107	2,868	4.2 (76)	55.0 (35)	1,771	18.9 (66)	50.8 (55)	578	43.8 (39)	999	33.8 (57)	3,316	5.8 (48)	3,322	11.7 (100)
4109	496	7.9 (152)	50.6 (11)	189	14.3 (4)	54.5 (112)	140	29.3 (8)	317	18.3 (5)	1,060	5.8 (48)	1,063	14.8 (149)
4112	606	2.8 (20)	65.2 (143)	543	71.5 (136)	11.6 (1)	70	*	54	*	582	17.0 (158)	581	11.0 (82)
4113	870	3.1 (29)	61.6 (123)	397	20.9 (105)	45.3 (11)	128	23.4 (4)	190	31.6 (48)	874	11.0 (154)	873	11.0 (82)
4115	2,461	5.0 (110)	54.2 (28)	1,421	18.7 (60)	49.3 (34)	619	36.8 (24)	1,050	33.8 (57)	3,442	7.4 (101)	3,443	10.6 (77)
4120	68	*	*	42	*	*	0	*	8	*	59	*	59	*
4160	49	*	*	27	*	*	0	*	6	*	71	*	71	*
4351	594	4.7 (98)	61.6 (123)	465	19.8 (79)	50.1 (49)	72	*	113	27.4 (24)	580	9.1 (145)	581	13.8 (136)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4352	193	5.7 (136)	61.7 (127)	134	25.4 (131)	56.0 (126)	23	*	36	*	197	5.6 (39)	199	9.0 (42)
4353	433	3.7 (47)	64.0 (141)	340	17.4 (35)	53.8 (105)	53	*	71	*	453	7.3 (99)	453	13.7 (134)
4354	683	5.6 (132)	58.3 (85)	489	20.0 (82)	47.9 (22)	128	31.3 (12)	151	23.8 (17)	757	4.8 (16)	757	10.7 (79)
4355	357	5.6 (132)	56.6 (51)	240	18.8 (64)	53.3 (97)	70	*	101	27.7 (26)	362	7.7 (113)	362	9.9 (62)
4356	463	3.7 (47)	57.5 (66)	324	20.4 (90)	49.4 (36)	72	*	101	28.7 (30)	503	5.2 (25)	503	12.7 (119)
4358	128	2.3 (4)	65.6 (147)	100	20.0 (82)	42.0 (5)	17	*	23	*	136	8.1 (126)	136	9.6 (55)
4359	490	3.5 (38)	60.8 (115)	360	21.7 (115)	48.6 (28)	60	*	92	*	504	8.1 (126)	505	15.2 (153)
4360	222	2.7 (18)	57.7 (71)	186	20.4 (90)	50.0 (46)	32	*	53	*	223	5.8 (48)	223	9.4 (50)
4361	278	4.0 (63)	61.5 (121)	225	16.4 (20)	60.4 (135)	31	*	73	*	273	6.6 (76)	274	9.1 (45)
4362	285	3.5 (38)	51.6 (17)	230	19.6 (76)	54.3 (109)	40	*	81	*	286	8.4 (134)	286	6.6 (15)
4363	960	4.1 (71)	56.0 (41)	843	22.8 (125)	51.8 (71)	124	41.1 (31)	215	27.0 (22)	932	5.7 (45)	933	11.1 (85)
4401	1,210	4.8 (102)	56.4 (49)	1,046	15.2 (9)	55.3 (121)	192	34.9 (19)	304	30.9 (43)	1,148	4.7 (14)	1,149	7.1 (18)
4501	761	5.7 (136)	53.2 (23)	572	19.4 (71)	51.0 (57)	118	26.3 (6)	156	20.5 (10)	962	7.5 (105)	962	10.7 (79)
4601	1,375	3.9 (57)	58.0 (74)	782	21.2 (107)	49.4 (36)	313	35.8 (20)	334	29.0 (32)	1,526	6.9 (85)	1,528	12.4 (113)
4602	17	*	*	11	*	*	2	*	6	*	22	*	22	*
4701	1,195	5.9 (141)	52.0 (20)	896	14.8 (7)	54.5 (112)	219	25.6 (5)	411	16.5 (2)	1,154	8.0 (122)	1,157	11.3 (89)
5050	1,878	3.5 (38)	57.8 (73)	1,604	18.2 (51)	53.4 (100)	394	42.9 (36)	459	31.2 (46)	1,798	7.6 (109)	1,795	12.5 (115)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5051	2,849	4.4 (82)	58.1 (79)	2,430	15.6 (11)	55.1 (118)	563	42.5 (34)	755	34.4 (60)	2,715	8.1 (126)	2,718	11.1 (85)
5052	2,829	5.0 (110)	57.0 (58)	2,441	20.5 (94)	49.7 (41)	562	47.3 (48)	631	35.2 (64)	2,766	8.0 (122)	2,769	10.4 (72)
5055	576	2.8 (20)	50.0 (6)	249	20.5 (94)	51.8 (71)	110	49.1 (53)	289	37.4 (69)	1,184	8.0 (122)	1,189	3.6 (2)
5101	1,992	4.4 (82)	54.7 (33)	1,634	16.8 (22)	52.3 (80)	411	46.5 (45)	583	35.0 (62)	2,008	9.5 (150)	2,008	11.2 (88)
5202	721	4.2 (76)	53.7 (26)	548	18.2 (51)	48.7 (29)	105	27.6 (7)	225	19.6 (7)	720	6.5 (72)	723	10.5 (74)
5203	38	*	*	31	*	*	4	*	11	*	39	*	39	*
5204	27	*	*	24	*	*	2	*	10	*	30	*	30	*
5601	1,105	3.4 (35)	57.7 (71)	759	20.7 (101)	51.1 (60)	172	33.7 (16)	296	29.4 (35)	1,164	6.2 (65)	1,164	9.4 (50)
5801	1,511	5.1 (115)	55.1 (36)	952	20.6 (98)	52.0 (75)	346	43.4 (37)	382	31.9 (50)	1,762	6.7 (78)	1,764	12.9 (121)
5803	131	5.3 (123)	51.1 (15)	94	*	*	22	*	33	*	126	13.5 (156)	126	9.5 (54)
5804	21	*	*	16	*	*	3	*	5	*	30	*	30	*
5806	7	*	*	6	*	*	0	*	5	*	11	*	11	*
6101	30	*	*	24	*	*	2	*	3	*	28	*	28	*
6102	131	5.3 (123)	50.4 (9)	110	17.3 (33)	49.1 (33)	14	*	14	*	128	9.4 (149)	127	12.6 (117)
6103	4,046	4.2 (76)	59.5 (102)	3,225	20.8 (103)	50.2 (51)	582	41.8 (33)	403	28.8 (31)	4,046	8.5 (137)	4,047	13.4 (126)
6104	99	*	*	85	*	*	5	*	4	*	89	*	90	*
6105	28	*	*	24	*	*	2	*	5	*	30	*	30	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6301	4,336	4.3 (80)	57.4 (63)	3,481	19.6 (76)	51.0 (57)	913	46.0 (42)	983	32.9 (56)	4,349	6.4 (69)	4,349	11.6 (97)
6312	2,425	3.5 (38)	58.7 (92)	1,909	18.0 (45)	51.7 (70)	550	46.0 (42)	846	36.5 (66)	2,464	5.0 (22)	2,464	8.8 (40)
6316	1,309	3.8 (52)	56.1 (43)	1,126	17.4 (35)	52.7 (85)	241	36.5 (23)	387	30.5 (39)	1,251	6.9 (85)	1,251	8.7 (37)
6320	110	7.3 (150)	48.2 (2)	58	*	*	20	*	39	*	126	8.7 (138)	126	4.8 (5)
6325	112	4.5 (89)	58.0 (74)	70	*	*	24	*	59	*	150	7.3 (99)	150	14.7 (147)
6330	558	4.7 (98)	58.1 (79)	327	19.0 (69)	48.3 (24)	120	60.8 (71)	122	45.1 (86)	662	4.7 (14)	663	12.2 (112)
6335	700	3.9 (57)	51.9 (19)	297	16.8 (22)	53.5 (103)	146	47.9 (51)	174	27.6 (25)	905	8.8 (139)	905	8.8 (40)
6340	48	*	*	24	*	*	11	*	17	*	62	*	62	*
6345	25	*	*	19	*	*	4	*	11	*	26	*	26	*
6350	17	*	*	13	*	*	2	*	1	*	17	*	17	*
6360	101	4.0 (63)	54.5 (32)	35	*	*	24	*	30	*	147	6.8 (81)	146	14.4 (145)
7304	112	1.8 (2)	67.0 (150)	89	*	*	26	*	32	*	154	8.4 (134)	154	14.9 (150)
7306	2,109	3.7 (47)	61.4 (119)	1,565	21.7 (115)	49.8 (42)	445	53.7 (60)	376	37.8 (71)	2,557	6.5 (72)	2,557	12.1 (107)
7308	118	2.5 (10)	62.7 (138)	94	*	*	26	*	27	*	135	5.9 (52)	135	10.4 (72)
7310	15	*	*	9	*	*	7	*	12	*	42	*	42	*
7401	2,402	4.9 (105)	54.4 (30)	2,004	16.8 (22)	52.8 (87)	383	46.5 (45)	466	32.0 (51)	2,319	7.6 (109)	2,327	9.9 (62)
7601	543	6.4 (147)	57.6 (69)	302	17.9 (42)	50.0 (46)	97	*	143	26.6 (20)	648	6.0 (58)	649	10.2 (68)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7850	1,173	5.3 (123)	58.9 (96)	1,003	18.4 (56)	50.1 (49)	188	34.0 (17)	210	30.0 (36)	1,172	7.9 (119)	1,172	9.1 (45)
7852	42	*	*	34	*	*	8	*	7	*	44	*	44	*
8101	1,710	3.9 (57)	61.5 (121)	1,244	20.3 (87)	49.4 (36)	371	67.9 (76)	268	53.7 (91)	1,918	6.9 (85)	1,920	16.7 (157)
8102	436	3.7 (47)	65.1 (142)	255	21.6 (112)	45.5 (13)	62	*	48	*	478	5.6 (39)	477	8.6 (33)
8105	1,637	5.0 (110)	60.5 (114)	1,335	17.1 (30)	54.5 (112)	386	54.4 (64)	410	41.0 (77)	1,790	6.9 (85)	1,792	11.8 (102)
8106	339	3.8 (52)	54.9 (34)	244	15.6 (11)	56.6 (130)	110	52.7 (59)	115	38.3 (72)	517	4.3 (10)	516	9.3 (49)
8150	42	*	*	30	*	*	7	*	15	*	43	*	43	*
8151	3,268	4.0 (63)	58.1 (79)	2,228	19.8 (79)	49.0 (32)	604	47.8 (50)	919	35.1 (63)	3,228	7.8 (116)	3,228	11.9 (103)
8152	142	6.3 (146)	48.6 (4)	94	*	*	22	*	28	*	148	2.7 (1)	148	13.5 (128)
8153	14	*	*	13	*	*	2	*	4	*	14	*	14	*
8155	43	*	*	36	*	*	5	*	19	*	48	*	48	*
8157	13	*	*	8	*	*	1	*	4	*	15	*	15	*
8201	2,564	4.5 (89)	57.1 (59)	1,722	17.0 (28)	55.3 (121)	664	44.1 (40)	839	30.3 (38)	2,803	7.4 (101)	2,805	10.0 (65)
8202	2,432	5.6 (132)	48.7 (5)	1,303	20.4 (90)	47.4 (20)	700	52.0 (57)	902	41.6 (78)	3,935	7.4 (101)	3,934	3.6 (2)
8203	462	3.2 (31)	58.0 (74)	353	13.3 (2)	58.1 (131)	99	*	178	28.1 (28)	505	7.1 (92)	506	11.3 (89)
8250	666	3.5 (38)	60.2 (109)	465	17.0 (28)	55.1 (118)	187	46.5 (45)	190	40.0 (74)	798	6.4 (69)	798	13.2 (123)
8252	122	2.5 (10)	62.3 (136)	89	*	*	36	*	24	*	143	4.2 (7)	143	14.7 (147)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8255	2,617	4.6 (96)	56.6 (51)	1,911	15.3 (10)	54.4 (110)	667	38.5 (27)	874	32.0 (51)	2,887	6.3 (67)	2,891	9.4 (50)
8257	1,020	5.1 (115)	50.2 (8)	470	29.4 (134)	38.1 (3)	283	48.4 (52)	578	42.4 (81)	1,617	9.6 (151)	1,617	6.2 (11)
8258	0	*	*	0	*	*	0	*	0	*	1	*	1	*
8263	726	3.9 (57)	58.1 (79)	587	16.4 (20)	52.8 (87)	193	29.5 (9)	296	28.4 (29)	825	8.2 (130)	826	13.9 (138)
8264	547	2.9 (22)	51.7 (18)	409	16.9 (26)	53.1 (95)	151	35.8 (20)	213	27.2 (23)	692	6.2 (65)	693	11.5 (94)
8265	53	*	*	41	*	*	16	*	16	*	55	*	55	*
8266	27	*	*	11	*	*	5	*	12	*	42	*	42	*
8267	380	4.5 (89)	57.4 (63)	257	16.3 (16)	59.9 (134)	129	36.4 (22)	132	28.0 (27)	459	8.1 (126)	459	10.0 (65)
8901	1,039	3.8 (52)	60.8 (115)	738	20.3 (87)	49.5 (40)	219	68.0 (77)	154	59.7 (96)	1,142	5.1 (23)	1,143	14.3 (144)
8902	1,167	4.8 (102)	61.2 (118)	857	17.5 (37)	53.4 (100)	259	65.3 (74)	207	59.4 (95)	1,274	6.1 (60)	1,274	13.9 (138)
8903	1,092	2.3 (4)	65.7 (149)	766	20.8 (103)	53.4 (100)	211	70.1 (78)	148	61.5 (98)	1,183	5.2 (25)	1,183	15.8 (155)
8904	1,910	3.5 (38)	61.6 (123)	1,451	16.9 (26)	54.5 (112)	496	73.4 (79)	312	60.6 (97)	2,104	5.7 (45)	2,105	14.0 (142)
8905	1,339	3.3 (33)	56.4 (49)	912	20.6 (98)	49.3 (34)	268	54.1 (61)	276	44.2 (84)	1,596	7.7 (113)	1,598	9.4 (50)
8908	29	*	*	20	*	*	7	*	5	*	33	*	33	*
8909	234	6.4 (147)	65.4 (145)	164	18.3 (54)	56.1 (129)	58	*	36	*	254	7.5 (105)	254	11.4 (91)
9012	16	*	*	11	*	*	1	*	0	*	20	*	20	*
9013	5	*	*	5	*	*	0	*	0	*	7	*	7	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9016	93	*	*	61	*	*	15	*	12	*	104	3.8 (5)	104	13.5 (128)
9017	17	*	*	12	*	*	3	*	5	*	20	*	20	*
9018	92	*	*	68	*	*	5	*	4	*	107	5.6 (39)	107	12.1 (107)
9041	9	*	*	4	*	*	0	*	0	*	9	*	9	*
9045	1,397	2.4 (7)	62.0 (130)	1,031	21.8 (117)	48.8 (30)	287	44.3 (41)	198	31.3 (47)	1,763	6.4 (69)	1,762	7.9 (26)
9050	936	4.1 (71)	61.0 (117)	732	21.9 (118)	51.0 (57)	209	56.5 (67)	153	54.2 (92)	1,073	6.5 (72)	1,074	10.7 (79)
9051	1,350	2.4 (7)	61.8 (128)	869	19.1 (70)	45.2 (10)	178	40.4 (29)	361	32.4 (53)	1,636	8.9 (140)	1,639	6.7 (16)
9053	1,588	4.4 (82)	60.3 (111)	1,389	17.6 (38)	55.4 (123)	281	57.3 (69)	220	50.0 (90)	1,570	4.9 (19)	1,570	13.9 (138)
9055	56	*	*	53	*	*	2	*	0	*	59	*	59	*
9058	707	5.0 (110)	60.4 (113)	508	20.5 (94)	48.8 (30)	172	51.2 (56)	181	37.0 (68)	1,055	7.5 (105)	1,056	12.1 (107)
9066	1,226	3.6 (45)	65.3 (144)	1,027	20.3 (87)	49.8 (42)	253	61.3 (72)	164	54.3 (93)	1,280	5.4 (32)	1,277	14.2 (143)
9068	1,086	3.4 (35)	65.6 (147)	949	21.3 (109)	54.0 (107)	179	54.2 (62)	106	34.9 (61)	1,046	5.4 (32)	1,046	13.3 (124)
9069	3,042	4.1 (71)	59.6 (105)	2,559	20.2 (85)	52.8 (87)	677	49.8 (54)	357	41.7 (79)	3,177	5.4 (32)	3,178	13.0 (122)
9070	1,057	4.5 (89)	58.6 (89)	777	19.4 (71)	51.6 (68)	189	66.7 (75)	103	55.3 (94)	1,163	6.1 (60)	1,165	13.8 (136)
9072	1,959	4.1 (71)	62.1 (134)	1,528	21.2 (107)	51.5 (67)	439	57.2 (68)	250	48.0 (88)	2,048	5.2 (25)	2,047	15.0 (152)
9080	2,409	4.7 (98)	60.2 (109)	1,808	18.7 (60)	52.3 (80)	551	63.7 (73)	430	48.8 (89)	2,804	5.8 (48)	2,801	11.5 (94)
9081	77	*	*	58	*	*	12	*	6	*	74	*	74	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 5A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9082	56	*	*	36	*	*	12	*	6	*	54	*	55	*
9083	17	*	*	16	*	*	0	*	0	*	20	*	20	*
9084	12	*	*	12	*	*	0	*	1	*	13	*	13	*
9085	3	*	*	2	*	*	0	*	0	*	3	*	3	*
Michigan	**151,746**	**4.4**	**57.8**	**108,901**	**19.4**	**51.2**	**28,808**	**45.2**	**35,717**	**34.2**	**167,329**	**6.9**	**167,397**	**10.9**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	106	80.2 (94)	11.3 (100)	112	33.9 (101)	104	21.2 (73)	90	*
0102	493	82.8 (70)	11.2 (97)	517	41.2 (49)	476	18.1 (100)	430	88.8 (74)
0103	419	74.5 (131)	15.3 (128)	445	38.4 (71)	393	20.9 (78)	368	89.9 (55)
0104	157	77.1 (118)	11.5 (106)	167	41.3 (48)	147	23.8 (55)	133	91.0 (40)
0250	119	73.1 (137)	18.5 (137)	120	50.8 (8)	90	*	97	*
0251	95	*	*	101	48.5 (15)	84	*	81	*
0252	124	75.0 (127)	13.7 (121)	138	47.1 (18)	132	22.7 (61)	119	81.5 (128)
0253	169	88.2 (30)	11.2 (97)	172	65.7 (3)	165	32.1 (17)	155	87.1 (88)
0401	511	87.3 (37)	7.0 (66)	538	39.4 (63)	509	24.4 (48)	465	91.6 (30)
0402	391	83.4 (68)	9.7 (85)	404	37.4 (77)	376	21.0 (76)	328	89.9 (55)
0403	126	78.6 (102)	13.5 (119)	135	40.0 (58)	123	25.2 (39)	112	90.2 (48)
0404	133	82.0 (77)	11.3 (100)	139	40.3 (57)	135	25.9 (33)	118	93.2 (12)
0501	576	91.1 (11)	3.6 (33)	598	36.5 (85)	552	24.6 (45)	485	90.3 (47)
0503	105	81.0 (87)	8.6 (75)	111	36.0 (89)	111	28.8 (26)	97	*
0504	513	85.6 (52)	8.6 (75)	539	37.1 (82)	508	29.3 (25)	443	91.0 (40)
0505	328	82.6 (72)	11.3 (100)	337	46.6 (22)	312	25.0 (41)	286	87.4 (85)
0506	208	72.1 (139)	23.6 (152)	220	52.3 (6)	201	18.9 (92)	184	82.1 (124)
0507	85	*	*	85	*	76	*	53	*
0508	33	*	*	32	*	31	*	21	*
0509	11	*	*	11	*	10	*	5	*
0650	1,108	83.7 (66)	9.0 (80)	1,176	40.5 (56)	1,015	28.4 (27)	963	92.8 (13)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0651	608	68.8 (148)	25.0 (154)	659	46.7 (20)	588	14.1 (123)	550	87.1 (88)
0652	305	77.7 (113)	17.7 (134)	313	41.9 (46)	288	22.2 (65)	267	85.0 (114)
0653	450	73.3 (136)	22.0 (147)	475	50.7 (9)	433	24.2 (51)	403	88.8 (74)
0654	357	72.0 (140)	20.7 (143)	385	44.4 (28)	347	23.1 (60)	324	80.2 (132)
0655	361	81.4 (84)	8.6 (75)	383	48.6 (14)	351	25.6 (36)	328	90.2 (48)
0657	71	*	*	74	*	71	*	69	*
0701	441	78.5 (104)	17.5 (133)	449	37.2 (80)	430	39.8 (9)	384	89.8 (60)
0702	76	*	*	84	*	77	*	65	*
0703	217	91.7 (8)	2.3 (17)	248	39.9 (59)	228	33.8 (14)	199	91.0 (40)
0704	58	*	*	67	*	67	*	57	*
0705	127	83.5 (67)	11.0 (94)	140	36.4 (86)	128	41.4 (7)	112	93.8 (8)
0706	2	*	*	2	*	1	*	0	*
0850	935	90.6 (12)	2.5 (18)	1,031	29.9 (114)	809	24.8 (43)	787	91.5 (31)
0851	747	94.0 (2)	2.0 (11)	813	37.3 (78)	619	24.4 (48)	644	92.1 (21)
0901	1,675	71.3 (143)	21.6 (146)	1,810	46.7 (20)	1,683	20.0 (81)	1,553	84.2 (118)
0902	41	*	*	43	*	33	*	27	*
1051	1,461	67.8 (151)	22.8 (150)	1,526	34.9 (96)	1,393	25.9 (33)	1,123	89.7 (64)
1101	1,754	76.7 (121)	8.5 (74)	1,920	28.4 (120)	1,615	13.1 (126)	1,472	85.5 (110)
1102	844	80.2 (94)	10.2 (89)	942	38.2 (72)	685	23.9 (54)	735	93.9 (5)
1106	85	*	*	94	*	67	*	65	*
1250	818	78.4 (105)	12.2 (112)	1,074	44.6 (27)	931	21.7 (69)	923	80.9 (131)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1251	853	77.1 (118)	14.0 (122)	921	40.7 (54)	833	19.2 (89)	755	83.3 (122)
1252	490	72.4 (138)	17.8 (135)	540	40.9 (51)	501	20.2 (80)	453	89.4 (65)
1253	895	68.6 (150)	22.2 (148)	987	39.7 (61)	896	17.0 (104)	810	87.9 (83)
1350	2,425	88.7 (27)	4.7 (45)	2,697	37.2 (80)	1,666	19.4 (87)	2,150	79.3 (135)
1351	156	89.1 (24)	6.4 (61)	166	31.3 (111)	146	24.7 (44)	124	94.4 (4)
1352	345	84.9 (54)	7.8 (69)	376	42.8 (40)	330	5.8 (153)	310	87.4 (85)
1401	250	69.2 (147)	24.8 (153)	270	43.0 (37)	158	29.7 (24)	213	81.2 (129)
1402	151	66.2 (153)	21.2 (145)	161	31.7 (110)	112	25.9 (33)	117	76.1 (142)
1403	22	*	*	23	*	18	*	21	*
1404	0	*	*	3	*	1	*	2	*
1501	812	81.7 (79)	11.3 (100)	931	44.1 (33)	680	10.3 (141)	749	78.9 (136)
1503	1,104	87.0 (41)	8.0 (71)	1,233	44.2 (31)	1,021	45.9 (4)	1,029	78.1 (139)
1504	2,164	89.9 (18)	4.4 (41)	2,486	39.5 (62)	1,538	22.2 (65)	1,924	73.2 (144)
1505	519	89.2 (23)	6.2 (60)	562	36.3 (87)	403	33.3 (15)	425	89.2 (69)
1506	614	77.4 (114)	17.4 (132)	668	42.2 (45)	583	7.9 (147)	533	88.7 (77)
1507	607	92.6 (6)	4.0 (38)	651	40.9 (51)	476	27.5 (28)	519	89.8 (60)
1508	71	*	*	122	38.5 (70)	73	*	77	*
1509	424	77.4 (114)	17.0 (131)	534	50.2 (10)	345	23.5 (58)	439	71.3 (146)
1511	701	86.9 (42)	6.0 (59)	774	40.6 (55)	511	31.3 (20)	629	85.1 (113)
1512	363	85.7 (48)	9.6 (84)	385	36.1 (88)	223	32.3 (16)	300	86.0 (103)
1513	55	*	*	57	*	42	*	38	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1514	100	96.0 (1)	0.0 (1)	107	17.8 (153)	69	*	60	*
1701	689	88.2 (30)	10.0 (87)	727	50.1 (12)	648	64.4 (3)	639	86.7 (94)
1950	615	85.7 (48)	7.8 (69)	654	39.3 (64)	551	21.1 (75)	530	89.2 (69)
1952	118	87.3 (37)	5.1 (52)	121	29.8 (116)	103	15.5 (113)	98	*
1953	319	90.6 (12)	3.8 (36)	330	31.8 (109)	284	24.3 (50)	264	94.7 (2)
1954	609	83.9 (62)	11.5 (106)	645	43.1 (36)	582	25.3 (38)	535	89.0 (72)
1955	76	*	*	79	*	73	*	69	*
1957	573	89.0 (26)	4.7 (45)	613	36.9 (84)	519	24.5 (46)	481	90.4 (46)
1964	14	*	*	15	*	15	*	11	*
2150	696	89.5 (21)	5.5 (54)	762	45.8 (23)	701	23.7 (56)	637	91.8 (25)
2151	231	84.0 (61)	11.7 (109)	269	42.8 (40)	238	21.0 (76)	219	84.5 (116)
2251	420	88.1 (33)	9.3 (82)	462	44.2 (31)	424	26.4 (31)	381	89.8 (60)
2252	191	81.7 (79)	15.2 (126)	209	56.0 (5)	178	65.7 (2)	184	82.6 (123)
2554	98	*	*	102	37.3 (78)	86	*	80	*
2555	8,201	86.5 (43)	3.1 (27)	8,725	26.6 (125)	5,974	11.8 (133)	6,507	88.3 (80)
2556	83	*	*	81	*	63	*	14	*
2557	1	*	*	2	*	2	*	2	*
2558	2	*	*	2	*	2	*	2	*
2559	6	*	*	6	*	5	*	6	*
3001	1,073	73.9 (132)	10.0 (87)	1,149	38.6 (69)	1,033	6.5 (151)	975	79.5 (133)
3002	251	85.7 (48)	1.2 (6)	267	45.7 (25)	246	10.2 (142)	238	78.2 (138)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3004	124	89.5 (21)	4.0 (38)	137	48.9 (13)	127	15.0 (117)	125	68.0 (149)
3007	25	*	*	32	*	27	*	25	*
3009	2	*	*	8	*	4	*	4	*
3201	506	81.6 (82)	13.6 (120)	549	44.4 (28)	468	31.8 (18)	431	87.0 (90)
3301	4,151	77.8 (111)	12.4 (114)	4,500	29.7 (117)	3,894	12.2 (129)	3,498	87.6 (84)
3304	117	82.1 (75)	14.5 (125)	119	41.2 (49)	113	9.7 (144)	97	*
3305	152	80.9 (89)	11.2 (97)	160	35.6 (92)	143	10.5 (139)	117	92.3 (20)
3306	255	84.3 (59)	7.1 (67)	270	37.0 (83)	251	10.4 (140)	211	91.5 (31)
3308	75	*	*	76	*	65	*	53	*
3309	42	*	*	44	*	40	*	31	*
3312	417	79.4 (100)	10.3 (91)	440	27.3 (121)	332	16.6 (106)	359	88.6 (78)
3550	71	*	*	78	*	67	*	70	*
3551	28	*	*	28	*	21	*	20	*
3552	55	*	*	58	*	50	*	51	*
3701	1,082	81.0 (87)	10.2 (89)	1,178	38.9 (68)	1,086	19.2 (89)	959	91.7 (29)
3850	3,059	86.0 (46)	6.5 (63)	3,329	39.9 (59)	2,968	14.7 (120)	2,763	85.2 (112)
3901	115	80.9 (89)	0.0 (1)	119	0.8 (163)	63	*	19	*
3902	64	*	*	61	*	36	*	4	*
3903	1	*	*	2	*	2	*	0	*
3950	1,982	79.9 (97)	12.1 (111)	2,179	34.8 (98)	2,009	16.4 (107)	1,780	86.7 (94)
3954	8	*	*	9	*	9	*	8	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4001	1,296	69.8 (146)	19.0 (138)	1,441	48.4 (16)	1,365	13.1 (126)	1,290	72.5 (145)
4002	953	73.9 (132)	19.1 (139)	1,016	44.0 (34)	963	18.2 (99)	888	82.0 (125)
4003	8	*	*	8	*	7	*	6	*
4004	5	*	*	5	*	5	*	5	*
4103	58	*	*	66	*	56	*	54	*
4106	2,126	78.0 (109)	11.4 (105)	2,372	35.0 (95)	2,107	7.0 (149)	1,975	86.4 (100)
4107	3,312	83.9 (62)	8.3 (73)	3,570	30.5 (113)	2,592	19.2 (89)	2,784	89.9 (55)
4109	1,057	76.0 (123)	19.2 (140)	1,123	33.7 (103)	1,018	23.4 (59)	841	93.9 (5)
4112	580	56.7 (158)	0.3 (3)	602	27.2 (123)	574	92.5 (1)	562	92.0 (23)
4113	868	87.2 (39)	0.6 (4)	940	38.0 (73)	597	18.4 (97)	874	70.6 (147)
4115	3,423	84.7 (55)	8.0 (71)	3,647	28.6 (119)	3,044	18.9 (92)	2,838	91.0 (40)
4120	58	*	*	72	*	67	*	71	*
4160	70	*	*	80	*	71	*	79	*
4351	577	89.6 (20)	6.6 (65)	609	45.8 (23)	302	44.4 (5)	519	91.1 (37)
4352	195	71.3 (143)	17.9 (136)	213	50.2 (10)	190	15.8 (110)	187	86.1 (102)
4353	444	75.2 (125)	19.4 (141)	499	42.3 (44)	440	21.4 (71)	403	84.6 (115)
4354	755	80.7 (93)	12.3 (113)	794	39.0 (67)	345	43.8 (6)	652	90.2 (48)
4355	359	78.0 (109)	12.5 (115)	389	32.6 (107)	348	22.7 (61)	315	91.1 (37)
4356	499	75.2 (125)	16.8 (130)	534	44.0 (34)	490	24.1 (52)	450	89.3 (67)
4358	129	59.7 (157)	34.1 (158)	158	42.4 (43)	129	17.8 (102)	124	64.5 (150)
4359	501	78.8 (101)	14.4 (124)	523	42.8 (40)	431	25.1 (40)	422	91.2 (34)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4360	218	61.0 (156)	30.3 (157)	230	47.0 (19)	178	23.6 (57)	192	81.8 (127)
4361	273	82.8 (70)	11.0 (94)	287	47.4 (17)	252	21.8 (68)	241	90.0 (52)
4362	286	69.9 (145)	23.4 (151)	298	43.0 (37)	267	22.1 (67)	251	88.8 (74)
4363	931	71.6 (141)	20.9 (144)	967	44.3 (30)	671	31.3 (20)	797	86.6 (99)
4401	1,141	81.9 (78)	11.8 (110)	1,217	34.8 (98)	1,101	19.3 (88)	964	89.9 (55)
4501	954	67.0 (152)	22.6 (149)	995	41.9 (46)	921	25.4 (37)	805	91.8 (25)
4601	1,513	87.6 (36)	4.5 (43)	1,642	29.9 (114)	1,476	19.5 (86)	1,278	90.0 (52)
4602	23	*	*	23	*	16	*	11	*
4701	1,153	91.2 (10)	3.9 (37)	1,260	25.6 (129)	1,013	31.5 (19)	881	90.1 (51)
5050	1,777	89.1 (24)	2.5 (18)	1,926	21.6 (145)	1,189	24.0 (53)	1,395	92.5 (16)
5051	2,693	89.8 (19)	3.0 (25)	2,916	22.6 (143)	2,054	22.7 (61)	2,205	91.3 (33)
5052	2,737	85.7 (48)	4.3 (40)	2,928	23.1 (139)	1,908	18.7 (96)	2,195	91.8 (25)
5055	1,162	84.2 (60)	13.2 (117)	1,212	57.8 (4)	636	40.4 (8)	1,104	76.8 (141)
5101	1,987	88.2 (30)	5.5 (54)	2,127	24.9 (131)	1,582	26.2 (32)	1,588	92.1 (21)
5202	722	87.8 (34)	5.5 (54)	807	39.3 (64)	584	39.6 (10)	664	92.5 (16)
5203	39	*	*	41	*	27	*	36	*
5204	30	*	*	29	*	25	*	27	*
5601	1,154	81.2 (86)	14.0 (122)	1,219	37.6 (76)	1,145	30.7 (22)	995	96.2 (1)
5801	1,761	83.8 (64)	9.5 (83)	1,915	35.9 (91)	1,759	18.1 (100)	1,551	86.7 (94)
5803	125	80.8 (91)	10.4 (92)	136	33.8 (102)	124	15.3 (115)	98	*
5804	30	*	*	31	*	24	*	24	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5806	11	*	*	10	*	4	*	1	*
6101	28	*	*	31	*	28	*	25	*
6102	128	79.7 (99)	13.3 (118)	127	35.4 (93)	117	19.7 (84)	105	68.6 (148)
6103	3,954	81.3 (85)	9.8 (86)	4,277	40.8 (53)	3,839	15.7 (111)	3,522	77.5 (140)
6104	90	*	*	102	71.6 (2)	101	10.9 (137)	102	49.0 (151)
6105	30	*	*	30	*	24	*	20	*
6301	4,335	86.5 (43)	5.0 (51)	4,626	25.1 (130)	3,596	18.8 (95)	3,459	90.8 (44)
6312	2,452	90.4 (15)	2.7 (21)	2,611	20.4 (149)	2,026	19.8 (83)	1,906	92.6 (15)
6316	1,248	88.7 (27)	4.8 (48)	1,331	24.9 (131)	1,151	22.7 (61)	981	92.0 (23)
6320	124	92.7 (5)	3.2 (28)	138	31.9 (108)	49	*	102	88.2 (81)
6325	147	84.4 (58)	5.4 (53)	154	27.3 (121)	119	26.9 (29)	106	90.6 (45)
6330	661	78.1 (108)	8.9 (79)	709	33.3 (104)	560	15.4 (114)	616	86.0 (103)
6335	907	86.2 (45)	11.6 (108)	948	37.8 (74)	240	20.4 (79)	784	88.9 (73)
6340	62	*	*	66	*	40	*	49	*
6345	26	*	*	27	*	18	*	21	*
6350	17	*	*	18	*	1	*	18	*
6360	147	87.1 (40)	2.7 (21)	156	16.0 (158)	72	*	105	86.7 (94)
7304	151	74.8 (128)	15.2 (126)	159	35.2 (94)	143	18.9 (92)	124	93.5 (9)
7306	2,508	68.7 (149)	19.7 (142)	2,740	37.7 (75)	2,284	11.7 (134)	2,198	83.4 (121)
7308	130	80.8 (91)	10.8 (93)	141	26.2 (128)	131	8.4 (146)	113	91.2 (34)
7310	41	*	*	44	*	43	*	35	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7401	2,323	65.4 (154)	26.9 (155)	2,545	43.0 (37)	2,251	16.4 (107)	2,140	85.6 (106)
7601	636	83.3 (69)	8.8 (78)	693	34.9 (96)	428	36.0 (12)	553	83.9 (119)
7850	1,147	77.2 (117)	15.9 (129)	1,222	39.2 (66)	1,127	14.9 (118)	1,002	86.2 (101)
7852	43	*	*	45	*	41	*	40	*
8101	1,904	75.8 (124)	3.6 (33)	1,989	19.2 (152)	1,322	15.7 (111)	1,435	89.3 (67)
8102	476	81.7 (79)	6.5 (63)	501	45.5 (26)	355	15.2 (116)	441	78.7 (137)
8105	1,787	84.5 (56)	3.4 (30)	1,888	22.2 (144)	1,698	14.5 (121)	1,379	92.8 (13)
8106	510	78.4 (105)	3.5 (31)	544	20.2 (151)	389	9.8 (143)	399	86.7 (94)
8150	43	*	*	50	*	47	*	44	*
8151	3,219	88.6 (29)	4.8 (48)	3,525	32.8 (105)	3,262	18.3 (98)	2,814	89.8 (60)
8152	148	84.5 (56)	2.7 (21)	156	34.6 (100)	155	7.1 (148)	149	93.3 (11)
8153	14	*	*	19	*	16	*	13	*
8155	48	*	*	47	*	42	*	39	*
8157	15	*	*	16	*	14	*	14	*
8201	2,798	90.2 (16)	2.0 (11)	3,022	20.5 (148)	2,316	26.7 (30)	2,216	93.4 (10)
8202	3,848	85.4 (53)	11.0 (94)	4,069	36.0 (89)	3,249	11.3 (135)	3,310	81.1 (130)
8203	504	94.0 (2)	0.8 (5)	551	22.7 (142)	302	24.5 (46)	403	91.8 (25)
8250	786	85.8 (47)	2.0 (11)	850	22.9 (141)	527	34.9 (13)	633	87.2 (87)
8252	140	78.6 (102)	3.6 (33)	146	15.1 (161)	109	11.0 (136)	114	88.6 (78)
8255	2,870	90.6 (12)	1.6 (7)	3,078	21.4 (146)	2,051	30.7 (22)	2,285	92.5 (16)
8257	1,568	92.9 (4)	2.2 (15)	1,680	29.2 (118)	1,528	12.0 (131)	1,322	85.6 (106)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8258	1	*	*	1	*	1	*	1	*
8263	823	91.9 (7)	1.8 (8)	863	15.8 (159)	670	24.9 (42)	613	94.5 (3)
8264	688	91.3 (9)	2.9 (24)	738	16.8 (157)	586	20.0 (81)	526	93.9 (5)
8265	55	*	*	58	*	49	*	44	*
8266	40	*	*	46	*	40	*	40	*
8267	458	90.0 (17)	2.2 (15)	497	15.7 (160)	428	38.8 (11)	328	89.9 (55)
8901	1,130	76.9 (120)	3.5 (31)	1,189	23.3 (137)	666	14.9 (118)	917	85.6 (106)
8902	1,260	80.1 (96)	3.3 (29)	1,341	23.6 (136)	937	17.1 (103)	1,032	86.9 (91)
8903	1,165	76.7 (121)	4.8 (48)	1,223	23.3 (137)	751	11.9 (132)	931	85.9 (105)
8904	2,093	81.5 (83)	3.0 (25)	2,179	20.3 (150)	1,362	16.9 (105)	1,638	89.4 (65)
8905	1,588	83.8 (64)	7.2 (68)	1,647	31.3 (111)	1,234	21.2 (73)	1,298	85.4 (111)
8908	33	*	*	32	*	28	*	29	*
8909	252	77.8 (111)	2.0 (11)	261	13.8 (162)	199	16.1 (109)	184	89.1 (71)
9012	20	*	*	21	*	19	*	19	*
9013	7	*	*	7	*	7	*	7	*
9016	102	71.6 (141)	12.7 (116)	117	24.8 (133)	114	8.8 (145)	111	85.6 (106)
9017	20	*	*	22	*	20	*	19	*
9018	106	82.1 (75)	1.9 (9)	114	21.1 (147)	109	6.4 (152)	105	92.4 (19)
9041	9	*	*	9	*	8	*	8	*
9045	1,765	73.4 (135)	11.3 (100)	1,803	32.7 (106)	557	21.4 (71)	1,388	88.1 (82)
9050	1,056	73.9 (132)	9.2 (81)	1,111	27.1 (124)	884	19.6 (85)	872	84.3 (117)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 6A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9051	1,558	62.8 (155)	27.5 (156)	1,724	51.9 (7)	1,557	12.7 (128)	1,565	74.8 (143)
9053	1,567	74.7 (129)	6.4 (61)	1,655	23.0 (140)	1,409	13.4 (125)	1,227	79.5 (133)
9055	59	*	*	58	*	45	*	5	*
9058	1,055	87.8 (34)	5.5 (54)	1,081	26.6 (125)	866	4.6 (154)	829	82.0 (125)
9066	1,223	82.6 (72)	2.5 (18)	1,310	16.9 (156)	1,021	6.8 (150)	922	91.1 (37)
9068	1,050	78.2 (107)	4.6 (44)	1,100	26.4 (127)	725	14.2 (122)	851	86.8 (92)
9069	3,163	77.4 (114)	4.4 (41)	3,299	17.8 (153)	2,030	10.9 (137)	2,403	91.2 (34)
9070	1,148	74.7 (129)	5.9 (58)	1,214	24.0 (135)	793	21.7 (69)	913	83.6 (120)
9072	2,034	82.5 (74)	1.9 (9)	2,148	17.6 (155)	1,424	12.1 (130)	1,585	86.8 (92)
9080	2,789	79.8 (98)	4.7 (45)	2,912	24.2 (134)	1,879	13.7 (124)	2,202	90.0 (52)
9081	75	*	*	81	*	69	*	66	*
9082	55	*	*	58	*	49	*	35	*
9083	20	*	*	20	*	13	*	1	*
9084	13	*	*	13	*	4	*	0	*
9085	3	*	*	3	*	2	*	3	*
Michigan	**165,526**	**82.1**	**8.6**	**178,109**	**32.4**	**139,078**	**19.5**	**139,833**	**87.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	105	52.4 (155)	104	34.6 (155)	42	*	90	*	104	1.9 (32)
0102	477	50.7 (151)	497	32.0 (147)	205	30.2 (93)	430	19.1 (127)	497	4.6 (116)
0103	416	45.0 (133)	423	30.0 (140)	157	30.6 (91)	368	23.1 (139)	423	7.1 (144)
0104	150	52.0 (154)	158	35.4 (156)	66	*	133	19.5 (129)	158	7.0 (143)
0250	104	50.0 (148)	121	29.8 (139)	49	*	97	*	121	0.0 (1)
0251	90	*	95	*	51	*	82	*	95	*
0252	134	47.8 (143)	124	35.5 (157)	52	*	119	20.2 (132)	124	4.8 (121)
0253	165	44.2 (128)	168	31.0 (142)	65	*	155	20.6 (134)	168	7.1 (144)
0401	511	47.2 (141)	511	27.4 (130)	203	36.5 (64)	466	22.5 (136)	511	2.7 (66)
0402	378	47.1 (140)	391	27.6 (131)	149	39.6 (50)	328	25.0 (147)	391	4.1 (107)
0403	123	49.6 (147)	127	25.2 (121)	55	*	112	24.1 (143)	127	7.1 (144)
0404	135	43.0 (120)	133	24.8 (117)	49	*	118	22.9 (137)	133	7.5 (147)
0501	556	44.4 (130)	571	28.0 (133)	202	30.7 (90)	485	17.3 (114)	572	4.7 (118)
0503	111	42.3 (116)	105	32.4 (150)	38	*	97	*	105	5.7 (133)
0504	509	37.7 (98)	513	17.3 (77)	160	41.3 (42)	443	10.6 (61)	513	2.9 (71)
0505	312	39.4 (108)	326	20.9 (101)	111	42.3 (38)	286	15.4 (99)	327	3.4 (84)
0506	202	51.5 (152)	206	34.0 (153)	91	*	184	30.4 (150)	206	7.8 (150)
0507	76	*	84	*	17	*	53	*	84	*
0508	31	*	34	*	5	*	21	*	34	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0509	10	*	11	*	1	*	5	*	11	*
0650	1,105	38.6 (100)	1,107	22.7 (111)	359	33.4 (78)	961	14.7 (94)	1,106	3.2 (77)
0651	608	46.4 (138)	614	31.3 (146)	242	22.7 (110)	550	24.4 (144)	614	5.9 (135)
0652	289	51.9 (153)	306	24.8 (117)	135	37.0 (63)	266	17.7 (117)	306	6.9 (142)
0653	446	48.0 (144)	448	31.0 (142)	187	24.6 (109)	402	26.4 (148)	448	8.7 (154)
0654	355	44.8 (132)	355	28.2 (134)	132	26.5 (106)	324	18.5 (123)	355	3.9 (103)
0655	359	50.4 (150)	363	32.0 (147)	153	28.8 (98)	328	20.4 (133)	363	5.5 (128)
0657	71	*	70	*	23	*	69	*	70	*
0701	432	35.2 (88)	438	19.2 (91)	134	38.1 (60)	383	6.0 (17)	438	1.4 (18)
0702	77	*	78	*	22	*	65	*	78	*
0703	228	45.6 (136)	223	34.1 (154)	77	*	199	13.1 (80)	224	4.5 (113)
0704	67	*	60	*	23	*	57	*	60	*
0705	128	43.8 (125)	127	29.1 (137)	51	*	112	18.8 (126)	127	3.1 (76)
0706	1	*	2	*	0	*	0	*	2	*
0850	923	38.8 (101)	955	24.8 (117)	290	29.3 (95)	787	9.7 (49)	954	2.7 (66)
0851	751	32.4 (74)	758	20.4 (98)	203	31.0 (88)	645	11.9 (70)	759	1.8 (29)
0901	1,689	45.4 (135)	1,709	25.5 (124)	682	36.4 (66)	1,553	14.0 (87)	1,710	3.5 (90)
0902	34	*	41	*	10	*	27	*	41	*
1051	1,395	37.0 (94)	1,474	20.6 (99)	420	39.5 (51)	1,123	8.6 (38)	1,473	3.3 (80)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1101	1,673	31.6 (70)	1,778	14.1 (59)	425	43.8 (34)	1,470	13.0 (79)	1,780	3.7 (97)
1102	844	34.7 (85)	851	17.9 (80)	219	42.0 (39)	734	15.9 (102)	851	3.4 (84)
1106	79	*	88	*	20	*	65	*	88	*
1250	1,012	35.8 (90)	990	22.1 (107)	304	28.9 (97)	923	13.5 (83)	990	4.0 (104)
1251	838	46.4 (138)	856	31.0 (142)	322	22.4 (111)	755	17.9 (118)	858	4.5 (113)
1252	502	34.5 (84)	505	19.0 (89)	128	28.1 (101)	452	15.3 (97)	505	4.0 (104)
1253	899	40.8 (111)	914	23.9 (115)	307	32.6 (81)	810	15.3 (97)	915	3.8 (102)
1350	2,402	35.1 (87)	2,486	20.0 (95)	750	32.5 (82)	2,162	14.1 (88)	2,499	3.6 (94)
1351	150	34.7 (85)	161	18.6 (85)	42	*	124	7.3 (23)	161	1.9 (32)
1352	333	47.7 (142)	350	23.4 (112)	141	25.5 (107)	310	5.2 (12)	350	1.1 (14)
1401	247	49.4 (146)	255	32.2 (149)	108	32.4 (84)	214	20.1 (131)	255	5.5 (128)
1402	147	27.9 (56)	152	9.9 (38)	33	*	117	2.6 (3)	152	1.3 (15)
1403	23	*	21	*	12	*	21	*	22	*
1404	2	*	0	*	0	*	2	*	0	*
1501	816	21.8 (27)	825	10.8 (41)	153	28.1 (101)	751	4.9 (11)	823	1.9 (32)
1503	1,115	29.6 (64)	1,127	15.1 (63)	289	27.0 (104)	1,030	8.5 (37)	1,129	2.2 (47)
1504	2,204	25.7 (47)	2,251	10.7 (40)	520	47.5 (24)	1,962	6.6 (20)	2,256	2.3 (49)
1505	512	26.2 (49)	533	13.1 (55)	123	48.8 (21)	427	8.0 (31)	531	2.4 (53)
1506	605	25.6 (45)	620	15.6 (67)	130	29.2 (96)	534	5.2 (12)	619	2.1 (43)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1507	603	30.8 (68)	608	15.6 (67)	167	38.9 (53)	526	9.1 (41)	612	2.6 (60)
1508	88	*	79	*	4	*	76	*	79	*
1509	466	24.2 (36)	435	15.4 (65)	103	34.0 (75)	445	8.1 (32)	434	2.5 (58)
1511	708	39.3 (107)	716	22.3 (109)	252	32.5 (82)	631	13.8 (85)	719	4.3 (110)
1512	344	32.8 (76)	367	17.4 (78)	106	37.7 (61)	300	8.7 (39)	367	3.5 (90)
1513	45	*	56	*	7	*	38	*	56	*
1514	87	*	102	8.8 (29)	20	*	60	*	101	1.0 (12)
1701	701	42.8 (119)	687	33.5 (152)	255	17.6 (112)	641	31.8 (151)	686	19.5 (158)
1950	605	39.7 (109)	619	21.6 (102)	196	40.3 (48)	530	18.7 (124)	620	4.7 (118)
1952	113	42.5 (118)	119	28.6 (136)	37	*	97	*	119	5.0 (123)
1953	307	44.3 (129)	319	27.0 (129)	117	39.3 (52)	264	14.4 (93)	320	4.7 (118)
1954	605	43.1 (121)	612	25.2 (121)	223	36.3 (67)	535	12.9 (78)	613	3.4 (84)
1955	75	*	77	*	27	*	69	*	77	*
1957	570	35.6 (89)	575	18.8 (87)	172	44.8 (30)	482	10.0 (52)	576	1.7 (26)
1964	15	*	15	*	3	*	11	*	15	*
2150	706	44.1 (126)	698	29.4 (138)	248	25.4 (108)	637	12.2 (72)	699	2.0 (38)
2151	252	48.4 (145)	235	36.6 (158)	90	*	219	16.4 (107)	235	3.4 (84)
2251	446	44.4 (130)	423	26.7 (127)	155	34.2 (72)	381	6.8 (21)	424	0.9 (10)
2252	199	43.7 (124)	188	28.2 (134)	72	*	183	16.4 (107)	188	3.2 (77)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2554	86	*	98	*	22	*	80	*	98	*
2555	7,599	31.6 (70)	8,325	16.0 (71)	2,109	38.8 (54)	6,519	11.7 (69)	8,332	2.3 (49)
2556	63	*	84	*	4	*	14	*	84	*
2557	2	*	1	*	0	*	2	*	1	*
2558	2	*	2	*	0	*	2	*	2	*
2559	6	*	6	*	1	*	6	*	6	*
3001	1,033	37.6 (97)	1,073	18.5 (84)	335	42.4 (37)	975	12.6 (75)	1,074	2.9 (71)
3002	247	27.5 (54)	252	17.1 (75)	60	*	238	11.3 (65)	252	2.8 (70)
3004	129	10.1 (9)	127	3.1 (7)	11	*	125	4.8 (10)	126	0.8 (6)
3007	27	*	21	*	2	*	25	*	21	*
3009	4	*	1	*	0	*	4	*	1	*
3201	507	36.9 (93)	511	20.2 (97)	156	43.6 (35)	431	17.6 (116)	510	4.5 (113)
3301	4,087	25.1 (42)	4,197	14.1 (59)	830	35.7 (70)	3,499	11.1 (62)	4,204	5.1 (124)
3304	116	1.7 (1)	117	0.9 (3)	2	*	97	*	117	0.9 (10)
3305	145	20.0 (20)	152	6.6 (18)	24	*	117	4.3 (8)	152	2.0 (38)
3306	256	25.0 (40)	256	15.6 (67)	52	*	211	5.2 (12)	256	4.3 (110)
3308	66	*	75	*	16	*	53	*	75	*
3309	40	*	43	*	9	*	31	*	43	*
3312	392	30.6 (66)	425	15.3 (64)	104	37.5 (62)	359	9.2 (42)	425	5.2 (125)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3550	73	*	71	*	40	*	70	*	71	*
3551	21	*	28	*	13	*	20	*	28	*
3552	51	*	53	*	34	*	51	*	55	*
3701	1,089	44.1 (126)	1,094	24.9 (120)	394	33.0 (79)	958	16.7 (111)	1,094	2.6 (60)
3850	3,064	42.1 (115)	3,109	25.6 (125)	1,095	29.8 (94)	2,760	16.4 (107)	3,111	3.7 (97)
3901	62	*	118	18.6 (85)	8	*	19	*	118	0.8 (6)
3902	36	*	64	*	1	*	4	*	64	*
3903	2	*	2	*	0	*	0	*	2	*
3950	2,012	31.8 (73)	1,998	18.1 (81)	524	36.5 (64)	1,781	10.4 (55)	1,998	2.6 (60)
3954	9	*	8	*	2	*	8	*	8	*
4001	1,372	32.9 (77)	1,301	17.4 (78)	366	28.7 (100)	1,291	13.2 (81)	1,302	2.4 (53)
4002	976	31.7 (72)	952	18.1 (81)	262	30.5 (92)	888	10.5 (58)	950	2.4 (53)
4003	7	*	8	*	1	*	6	*	8	*
4004	5	*	5	*	2	*	5	*	5	*
4103	59	*	59	*	19	*	54	*	59	*
4106	2,144	24.9 (39)	2,184	11.8 (45)	425	41.4 (41)	1,973	7.9 (29)	2,182	2.1 (43)
4107	3,275	22.3 (28)	3,330	12.1 (47)	579	40.6 (44)	2,784	9.6 (46)	3,328	2.9 (71)
4109	1,027	33.0 (79)	1,064	16.8 (74)	279	40.5 (45)	840	7.9 (29)	1,065	1.6 (23)
4112	579	2.4 (2)	584	0.7 (2)	10	*	562	7.5 (26)	584	2.7 (66)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4113	896	11.7 (11)	880	5.6 (14)	95	*	872	4.5 (9)	885	1.0 (12)
4115	3,326	19.3 (19)	3,441	8.8 (29)	516	47.3 (25)	2,839	4.2 (7)	3,445	0.8 (6)
4120	72	*	59	*	21	*	72	*	59	*
4160	77	*	71	*	18	*	79	*	71	*
4351	575	41.6 (113)	582	23.5 (114)	203	33.5 (77)	518	14.3 (91)	581	3.3 (80)
4352	196	53.1 (157)	199	32.7 (151)	93	*	187	28.9 (149)	199	6.5 (140)
4353	443	32.5 (75)	454	19.8 (94)	127	32.3 (85)	403	15.1 (95)	455	2.6 (60)
4354	733	43.4 (123)	757	27.9 (132)	269	32.3 (85)	652	14.1 (88)	757	3.4 (84)
4355	353	45.6 (136)	363	22.6 (110)	137	38.7 (55)	315	14.3 (91)	363	3.6 (94)
4356	493	37.1 (95)	503	19.5 (92)	158	38.6 (57)	450	12.7 (77)	502	3.6 (94)
4358	131	15.3 (15)	137	9.5 (35)	18	*	124	2.4 (2)	137	0.7 (3)
4359	468	28.8 (61)	506	16.4 (72)	123	34.1 (73)	422	7.6 (28)	506	1.8 (29)
4360	207	36.2 (91)	223	20.6 (99)	70	*	192	16.7 (111)	223	2.7 (66)
4361	267	50.2 (149)	274	31.0 (142)	116	31.9 (87)	241	16.6 (110)	274	2.6 (60)
4362	281	53.0 (156)	286	30.8 (141)	132	34.1 (73)	251	24.7 (146)	286	5.9 (135)
4363	886	43.2 (122)	936	26.9 (128)	314	28.0 (103)	797	15.6 (100)	936	4.0 (104)
4401	1,134	37.1 (95)	1,149	19.7 (93)	343	38.2 (58)	961	6.2 (18)	1,148	1.5 (19)
4501	926	42.4 (117)	962	22.2 (108)	340	35.9 (68)	806	15.9 (102)	962	4.2 (109)
4601	1,490	34.4 (83)	1,534	17.1 (75)	399	44.1 (33)	1,279	10.2 (53)	1,535	2.9 (71)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4602	16	*	23	*	3	*	11	*	23	*
4701	1,143	39.2 (105)	1,154	19.0 (89)	327	45.6 (27)	885	10.4 (55)	1,157	2.5 (58)
5050	1,727	27.4 (53)	1,799	12.1 (47)	369	48.0 (22)	1,395	8.2 (33)	1,801	1.5 (19)
5051	2,664	25.2 (43)	2,714	12.0 (46)	540	44.4 (32)	2,205	7.3 (23)	2,721	2.1 (43)
5052	2,666	24.6 (37)	2,773	12.7 (51)	538	38.7 (55)	2,203	9.4 (43)	2,772	1.9 (32)
5055	1,191	2.4 (2)	1,192	0.5 (1)	29	*	1,120	5.6 (16)	1,188	1.3 (15)
5101	1,951	28.0 (57)	2,006	14.0 (58)	441	44.7 (31)	1,595	7.4 (25)	2,015	2.4 (53)
5202	751	39.0 (102)	722	25.3 (123)	258	34.5 (71)	663	12.1 (71)	727	1.8 (29)
5203	38	*	39	*	15	*	36	*	39	*
5204	27	*	30	*	13	*	27	*	30	*
5601	1,145	34.0 (82)	1,164	21.9 (106)	337	28.8 (98)	994	9.6 (46)	1,164	1.5 (19)
5801	1,776	39.2 (105)	1,766	23.4 (112)	557	33.8 (76)	1,552	15.6 (100)	1,766	4.6 (116)
5803	127	28.3 (58)	126	18.3 (83)	26	*	98	*	126	4.8 (121)
5804	25	*	30	*	11	*	24	*	30	*
5806	4	*	11	*	0	*	1	*	11	*
6101	28	*	28	*	12	*	25	*	28	*
6102	115	30.4 (65)	128	18.8 (87)	35	*	105	11.4 (67)	128	1.6 (23)
6103	3,901	33.6 (80)	4,030	21.6 (102)	1,145	26.7 (105)	3,520	13.9 (86)	4,051	5.5 (128)
6104	102	45.1 (134)	90	*	40	*	102	19.6 (130)	90	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6105	25	*	29	*	7	*	20	*	30	*
6301	4,178	20.1 (21)	4,354	11.0 (42)	636	40.4 (47)	3,462	7.5 (26)	4,351	1.6 (23)
6312	2,420	9.4 (7)	2,462	2.9 (6)	168	61.9 (7)	1,909	3.6 (6)	2,462	1.7 (26)
6316	1,264	20.9 (25)	1,251	10.4 (39)	182	47.8 (23)	980	12.6 (75)	1,250	5.6 (132)
6320	127	4.7 (5)	126	3.2 (8)	6	*	102	1.0 (1)	126	0.0 (1)
6325	141	39.0 (102)	150	20.0 (95)	47	*	106	16.0 (105)	150	3.3 (80)
6330	671	26.5 (50)	661	14.7 (61)	152	38.2 (58)	613	10.3 (54)	663	3.2 (77)
6335	910	10.1 (9)	908	4.8 (9)	72	*	783	3.3 (4)	909	0.8 (6)
6340	59	*	61	*	7	*	49	*	62	*
6345	26	*	26	*	6	*	21	*	26	*
6350	18	*	17	*	1	*	18	*	17	*
6360	129	20.9 (25)	146	13.0 (53)	21	*	105	11.4 (67)	147	0.7 (3)
7304	146	38.4 (99)	154	13.0 (53)	51	*	124	10.5 (58)	155	2.6 (60)
7306	2,476	36.4 (92)	2,563	13.9 (57)	763	51.6 (16)	2,204	15.1 (95)	2,569	3.4 (84)
7308	132	41.7 (114)	134	21.6 (102)	45	*	114	8.8 (40)	135	2.2 (47)
7310	44	*	42	*	16	*	35	*	42	*
7401	2,347	41.2 (112)	2,328	25.6 (125)	808	30.8 (89)	2,142	18.7 (124)	2,328	4.1 (107)
7601	625	39.0 (102)	646	24.0 (116)	217	35.9 (68)	564	14.2 (90)	649	3.7 (97)
7850	1,121	40.0 (110)	1,165	21.7 (105)	397	33.0 (79)	1,001	17.3 (114)	1,172	2.4 (53)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7852	41	*	44	*	17	*	40	*	44	*
8101	1,717	23.2 (32)	1,921	9.7 (36)	316	54.4 (13)	1,432	17.9 (118)	1,921	5.5 (128)
8102	470	17.0 (16)	478	9.0 (31)	70	*	441	5.2 (12)	478	1.9 (32)
8105	1,728	17.8 (18)	1,793	7.6 (24)	229	49.3 (19)	1,382	9.8 (51)	1,792	3.7 (97)
8106	475	25.7 (47)	515	12.6 (50)	101	41.6 (40)	398	12.3 (73)	517	1.5 (19)
8150	47	*	43	*	9	*	44	*	43	*
8151	3,267	25.6 (45)	3,233	11.3 (43)	649	45.1 (29)	2,810	8.2 (33)	3,228	1.7 (26)
8152	155	32.9 (77)	148	12.2 (49)	48	*	149	13.4 (82)	148	2.0 (38)
8153	16	*	14	*	3	*	13	*	14	*
8155	43	*	48	*	7	*	39	*	47	*
8157	15	*	15	*	1	*	14	*	15	*
8201	2,758	23.7 (35)	2,804	13.1 (55)	506	40.5 (45)	2,220	9.6 (46)	2,806	2.3 (49)
8202	3,829	8.3 (6)	3,936	2.5 (5)	268	63.4 (5)	3,321	8.3 (36)	3,933	2.3 (49)
8203	499	30.7 (67)	504	16.5 (73)	131	41.2 (43)	405	11.1 (62)	507	2.0 (38)
8250	740	28.5 (60)	797	11.7 (44)	165	54.5 (12)	629	13.5 (83)	796	2.9 (71)
8252	126	20.6 (24)	143	5.6 (14)	24	*	113	12.4 (74)	143	3.5 (90)
8255	2,749	29.2 (63)	2,889	15.6 (67)	642	40.3 (48)	2,285	9.7 (49)	2,884	2.0 (38)
8257	1,579	3.5 (4)	1,616	1.7 (4)	45	*	1,325	3.5 (5)	1,615	1.3 (15)
8258	1	*	1	*	0	*	1	*	1	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8263	781	28.4 (59)	827	14.9 (62)	173	45.7 (26)	610	8.2 (33)	827	1.9 (32)
8264	680	28.8 (61)	692	12.7 (51)	152	42.8 (36)	527	11.2 (64)	692	3.3 (80)
8265	52	*	56	*	20	*	44	*	56	*
8266	46	*	42	*	1	*	40	*	42	*
8267	449	31.0 (69)	460	15.4 (65)	95	*	327	10.4 (55)	460	0.7 (3)
8901	1,025	25.2 (43)	1,140	9.0 (31)	223	52.9 (15)	916	18.0 (121)	1,141	6.4 (138)
8902	1,202	20.1 (21)	1,274	5.5 (12)	201	67.2 (1)	1,032	15.9 (102)	1,272	6.4 (138)
8903	1,046	26.5 (50)	1,184	9.8 (37)	242	58.7 (10)	934	18.2 (122)	1,181	7.5 (147)
8904	1,955	22.8 (29)	2,104	7.1 (22)	373	65.1 (3)	1,637	17.9 (118)	2,105	7.8 (150)
8905	1,524	17.1 (17)	1,599	7.1 (22)	210	53.8 (14)	1,298	9.4 (43)	1,600	3.7 (97)
8908	32	*	33	*	9	*	29	*	33	*
8909	232	22.8 (29)	254	5.1 (11)	46	*	184	16.3 (106)	254	7.5 (147)
9012	19	*	20	*	5	*	19	*	20	*
9013	7	*	7	*	4	*	7	*	7	*
9016	112	33.9 (81)	104	8.7 (28)	33	*	110	23.6 (140)	104	5.8 (134)
9017	20	*	20	*	6	*	19	*	20	*
9018	111	23.4 (34)	107	9.3 (34)	22	*	105	10.5 (58)	107	8.4 (153)
9041	8	*	9	*	1	*	8	*	9	*
9045	1,633	13.1 (13)	1,765	6.2 (16)	163	45.4 (28)	1,388	9.5 (45)	1,764	4.4 (112)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 7A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9050	970	27.7 (55)	1,075	6.9 (20)	245	64.9 (4)	873	19.4 (128)	1,076	6.1 (137)
9051	1,652	12.8 (12)	1,642	5.0 (10)	163	49.7 (18)	1,567	7.0 (22)	1,641	2.1 (43)
9053	1,444	25.0 (40)	1,570	8.5 (26)	294	62.9 (6)	1,228	23.9 (142)	1,572	9.2 (155)
9055	47	*	59	*	1	*	5	*	59	*
9058	969	9.8 (8)	1,056	5.5 (12)	74	*	829	6.4 (19)	1,055	3.5 (90)
9066	1,086	15.1 (14)	1,282	6.7 (19)	145	49.0 (20)	921	16.9 (113)	1,281	5.4 (127)
9068	956	26.9 (52)	1,049	8.5 (26)	233	61.4 (8)	852	21.9 (135)	1,049	6.5 (140)
9069	2,861	23.2 (32)	3,181	7.0 (21)	551	66.1 (2)	2,401	23.7 (141)	3,178	10.4 (157)
9070	1,071	20.2 (23)	1,166	6.4 (17)	174	58.0 (11)	913	23.0 (138)	1,165	9.4 (156)
9072	1,833	24.6 (37)	2,052	9.2 (33)	384	50.5 (17)	1,585	24.4 (144)	2,054	7.9 (152)
9080	2,548	23.1 (31)	2,805	8.3 (25)	492	60.0 (9)	2,203	11.3 (65)	2,806	5.2 (125)
9081	73	*	74	*	4	*	66	*	74	*
9082	51	*	55	*	10	*	35	*	55	*
9083	15	*	20	*	0	*	1	*	18	*
9084	10	*	13	*	0	*	0	*	13	*
9085	3	*	3	*	1	*	3	*	3	*
Michigan	**161,648**	**28.8**	**167,486**	**15.0**	**38,754**	**39.4**	**139,966**	**12.2**	**167,578**	**3.6**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	94	*	*	98	*	80	*	94	*
0102	472	6.4 (21)	9.3 (117)	477	9.0 (19)	421	3.6 (67)	472	60.2 (116)
0103	405	7.2 (42)	7.9 (78)	408	9.1 (24)	365	4.1 (90)	402	67.2 (94)
0104	147	8.2 (72)	6.1 (45)	148	8.1 (7)	134	4.5 (106)	147	75.5 (62)
0250	113	3.5 (3)	13.3 (151)	114	9.6 (37)	99	*	113	77.0 (51)
0251	87	*	*	90	*	83	*	87	*
0252	118	6.8 (30)	7.6 (69)	121	11.6 (88)	104	1.0 (3)	118	83.1 (13)
0253	150	9.3 (93)	12.0 (147)	149	12.1 (103)	128	7.0 (143)	149	61.7 (113)
0401	475	8.6 (78)	11.6 (146)	490	10.6 (62)	418	3.8 (79)	475	62.1 (112)
0402	368	9.2 (91)	7.9 (78)	370	11.1 (75)	321	4.0 (87)	368	77.4 (48)
0403	123	8.9 (85)	8.9 (104)	125	9.6 (37)	110	3.6 (67)	123	65.9 (97)
0404	129	7.0 (38)	7.0 (62)	129	10.9 (69)	114	1.8 (9)	129	56.6 (130)
0501	448	6.0 (13)	12.1 (149)	545	8.3 (11)	408	2.9 (39)	514	74.5 (68)
0503	98	*	*	97	*	90	*	97	*
0504	485	6.8 (30)	10.9 (138)	502	7.0 (2)	448	3.8 (79)	492	84.3 (8)
0505	286	8.4 (75)	9.4 (119)	307	8.1 (7)	259	2.7 (26)	303	80.5 (29)
0506	166	6.0 (13)	6.0 (41)	195	7.2 (4)	153	2.6 (24)	188	78.2 (42)
0507	73	*	*	78	*	64	*	76	*
0508	32	*	*	33	*	28	*	33	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0509	11	*	*	11	*	10	*	10	*
0650	1,028	8.7 (81)	9.0 (106)	1,072	9.0 (19)	934	5.2 (123)	1,018	78.2 (42)
0651	560	5.9 (11)	6.4 (50)	580	9.8 (43)	504	2.4 (18)	556	65.1 (101)
0652	273	5.9 (11)	9.5 (120)	285	10.5 (56)	242	2.1 (15)	272	71.3 (85)
0653	419	8.8 (82)	7.9 (78)	434	11.1 (75)	375	3.5 (63)	420	79.3 (38)
0654	334	6.3 (17)	10.2 (128)	341	11.7 (91)	296	1.0 (3)	334	81.7 (21)
0655	332	8.1 (69)	6.0 (41)	343	12.2 (107)	287	4.2 (96)	333	73.9 (73)
0657	62	*	*	62	*	56	*	62	*
0701	399	6.5 (23)	11.0 (143)	409	10.3 (51)	347	1.7 (6)	398	87.4 (3)
0702	72	*	*	74	*	61	*	72	*
0703	217	8.8 (82)	5.1 (26)	217	15.7 (146)	179	2.8 (33)	216	68.5 (91)
0704	53	*	*	55	*	47	*	53	*
0705	117	6.0 (13)	10.3 (133)	125	7.2 (4)	108	2.8 (33)	119	79.8 (35)
0706	2	*	*	2	*	1	*	2	*
0850	870	8.4 (75)	6.4 (50)	945	10.9 (69)	773	2.7 (26)	908	81.5 (23)
0851	712	9.7 (106)	8.7 (101)	748	11.0 (73)	627	4.6 (109)	730	77.4 (48)
0901	1,617	7.9 (64)	10.2 (128)	1,641	10.4 (53)	1,440	3.6 (67)	1,619	63.7 (104)
0902	39	*	*	39	*	38	*	39	*
1051	1,393	7.8 (57)	9.9 (123)	1,423	9.2 (25)	1,250	3.4 (55)	1,391	85.8 (6)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1101	1,670	9.2 (91)	6.9 (60)	1,732	10.9 (69)	1,481	4.1 (90)	1,671	67.4 (93)
1102	796	6.7 (29)	9.2 (111)	824	10.3 (51)	708	3.2 (51)	795	73.2 (76)
1106	83	*	*	84	*	73	*	83	*
1250	925	6.5 (23)	9.0 (106)	939	9.6 (37)	818	3.1 (44)	922	80.6 (28)
1251	793	6.3 (17)	8.3 (90)	817	8.4 (12)	719	2.4 (18)	785	78.1 (44)
1252	468	6.6 (27)	8.5 (94)	480	10.6 (62)	411	2.7 (26)	471	82.2 (18)
1253	867	7.7 (55)	7.8 (75)	889	10.2 (48)	770	2.7 (26)	870	81.8 (20)
1350	2,364	9.4 (97)	7.6 (69)	2,419	10.2 (48)	2,105	4.2 (96)	2,369	72.8 (79)
1351	152	5.3 (5)	9.2 (111)	154	6.5 (1)	141	4.3 (101)	154	74.0 (71)
1352	333	9.6 (104)	7.8 (75)	340	12.9 (118)	291	5.5 (125)	334	60.2 (116)
1401	231	7.8 (57)	8.7 (101)	233	9.0 (19)	204	4.4 (104)	229	82.5 (17)
1402	139	5.8 (10)	14.4 (154)	145	11.7 (91)	121	1.7 (6)	139	89.9 (1)
1403	19	*	*	19	*	19	*	18	*
1404	0	*	*	0	*	0	*	0	*
1501	760	6.3 (17)	9.1 (108)	772	10.2 (48)	670	1.6 (5)	766	68.5 (91)
1503	1,055	7.4 (46)	10.0 (124)	1,062	11.7 (91)	911	3.3 (53)	1,060	79.8 (35)
1504	2,122	8.0 (67)	8.6 (96)	2,200	11.4 (82)	1,870	2.6 (24)	2,153	76.1 (58)
1505	498	6.4 (21)	14.3 (153)	519	10.4 (53)	444	1.8 (9)	501	84.0 (10)
1506	577	5.5 (6)	10.9 (138)	590	9.5 (34)	516	1.9 (12)	582	76.3 (54)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1507	561	7.0 (38)	10.9 (138)	591	9.6 (37)	502	2.2 (16)	574	78.6 (41)
1508	73	*	*	74	*	58	*	74	*
1509	397	8.3 (74)	7.3 (66)	422	13.5 (124)	345	4.1 (90)	411	71.0 (88)
1511	682	7.8 (57)	10.1 (127)	694	9.5 (34)	609	2.0 (13)	689	79.5 (37)
1512	345	10.7 (125)	9.6 (122)	356	14.6 (141)	291	3.1 (44)	352	77.8 (46)
1513	52	*	*	54	*	44	*	53	*
1514	98	*	*	100	23.0 (156)	75	*	99	*
1701	654	6.9 (35)	10.9 (138)	649	10.6 (62)	564	1.8 (9)	652	71.3 (85)
1950	583	7.2 (42)	6.9 (60)	600	11.5 (84)	511	3.1 (44)	583	83.5 (11)
1952	115	6.1 (16)	7.0 (62)	112	8.9 (17)	102	2.9 (39)	115	81.7 (21)
1953	311	9.0 (88)	10.0 (124)	309	12.0 (101)	268	3.7 (72)	311	83.3 (12)
1954	578	9.3 (93)	9.2 (111)	590	9.8 (43)	517	4.1 (90)	576	82.1 (19)
1955	75	*	*	74	*	67	*	75	*
1957	538	6.5 (23)	8.9 (104)	556	9.2 (25)	489	2.2 (16)	541	88.2 (2)
1964	12	*	*	13	*	10	*	12	*
2150	649	7.7 (55)	10.3 (133)	665	9.6 (37)	586	3.1 (44)	635	76.1 (58)
2151	225	10.2 (112)	5.8 (34)	225	11.6 (88)	196	4.6 (109)	226	59.3 (123)
2251	398	8.0 (67)	9.3 (117)	400	9.5 (34)	351	3.4 (55)	393	75.3 (63)
2252	172	7.6 (50)	7.6 (69)	174	12.6 (114)	150	4.0 (87)	172	80.2 (32)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2554	95	*	*	96	*	83	*	95	*
2555	8,089	11.4 (128)	5.2 (28)	8,154	13.8 (130)	6,894	5.0 (118)	8,092	59.7 (122)
2556	80	*	*	80	*	64	*	80	*
2557	1	*	*	1	*	0	*	1	*
2558	2	*	*	2	*	2	*	2	*
2559	6	*	*	6	*	5	*	6	*
3001	1,004	12.9 (138)	4.8 (24)	1,028	13.7 (126)	862	5.7 (130)	1,006	62.8 (108)
3002	230	10.4 (116)	6.1 (45)	240	13.8 (130)	202	5.0 (118)	230	50.0 (139)
3004	117	3.4 (2)	5.1 (26)	118	9.3 (27)	106	0.9 (2)	117	75.2 (64)
3007	20	*	*	20	*	17	*	20	*
3009	0	*	*	0	*	0	*	0	*
3201	490	6.5 (23)	11.4 (145)	497	11.1 (75)	436	2.5 (22)	490	71.0 (88)
3301	3,965	10.0 (111)	5.9 (38)	4,107	12.5 (112)	3,457	4.4 (104)	3,984	77.9 (45)
3304	107	5.6 (7)	6.5 (52)	113	7.1 (3)	98	*	113	86.7 (5)
3305	142	6.3 (17)	13.4 (152)	146	8.9 (17)	127	2.4 (18)	142	87.3 (4)
3306	236	8.1 (69)	10.2 (128)	245	9.4 (30)	213	2.8 (33)	238	81.1 (26)
3308	70	*	*	75	*	68	*	69	*
3309	42	*	*	43	*	36	*	43	*
3312	407	9.8 (109)	5.4 (29)	414	9.9 (46)	365	4.7 (112)	411	71.8 (83)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3550	65	*	*	70	*	59	*	65	*
3551	28	*	*	28	*	26	*	28	*
3552	49	*	*	54	*	41	*	49	*
3701	1,054	7.6 (50)	8.3 (90)	1,067	11.8 (97)	926	2.9 (39)	1,057	80.1 (33)
3850	2,953	9.3 (93)	7.4 (67)	2,994	12.9 (118)	2,545	3.2 (51)	2,946	73.4 (75)
3901	106	12.3 (132)	2.8 (5)	110	10.9 (69)	90	*	107	73.8 (74)
3902	60	*	*	59	*	50	*	60	*
3903	2	*	*	2	*	2	*	2	*
3950	1,897	8.6 (78)	10.6 (136)	1,943	12.0 (101)	1,666	3.5 (63)	1,903	74.9 (66)
3954	7	*	*	8	*	7	*	7	*
4001	1,197	10.4 (116)	5.8 (34)	1,239	12.3 (108)	1,035	4.5 (106)	1,202	65.0 (102)
4002	886	10.2 (112)	7.9 (78)	914	11.6 (88)	772	4.3 (101)	893	65.3 (99)
4003	8	*	*	8	*	7	*	7	*
4004	4	*	*	5	*	4	*	4	*
4103	55	*	*	58	*	43	*	55	*
4106	2,047	10.5 (120)	5.5 (32)	2,116	12.9 (118)	1,771	5.3 (124)	2,056	70.4 (90)
4107	3,153	10.6 (121)	8.1 (86)	3,240	13.1 (121)	2,725	4.1 (90)	3,157	74.7 (67)
4109	998	9.5 (103)	10.0 (124)	1,020	14.2 (138)	852	2.5 (22)	1,007	82.6 (16)
4112	560	5.7 (8)	8.6 (96)	560	10.5 (56)	497	2.4 (18)	557	31.8 (155)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4113	844	7.9 (64)	6.6 (55)	867	10.7 (65)	749	4.0 (87)	845	80.4 (30)
4115	3,256	9.4 (97)	8.0 (84)	3,346	13.2 (122)	2,814	3.0 (43)	3,271	81.1 (26)
4120	57	*	*	59	*	52	*	57	*
4160	71	*	*	71	*	67	*	71	*
4351	543	8.1 (69)	10.9 (138)	557	12.4 (110)	475	2.7 (26)	540	75.2 (64)
4352	186	7.5 (47)	5.4 (29)	188	12.8 (116)	159	3.8 (79)	186	71.5 (84)
4353	425	7.5 (47)	8.0 (84)	432	9.0 (19)	383	3.7 (72)	422	76.1 (58)
4354	721	7.1 (40)	7.9 (78)	728	9.3 (27)	648	3.4 (55)	717	76.2 (55)
4355	338	8.9 (85)	8.3 (90)	347	11.2 (79)	295	3.7 (72)	336	80.4 (30)
4356	486	8.8 (82)	7.6 (69)	488	12.1 (103)	422	4.7 (112)	485	77.3 (50)
4358	129	3.1 (1)	8.5 (94)	128	8.6 (14)	117	1.7 (6)	130	83.1 (13)
4359	461	7.6 (50)	10.2 (128)	466	8.4 (12)	415	3.9 (84)	455	71.2 (87)
4360	198	6.6 (27)	10.6 (136)	211	9.0 (19)	177	3.4 (55)	196	84.2 (9)
4361	258	9.3 (93)	12.0 (147)	262	10.7 (65)	228	3.1 (44)	257	82.9 (15)
4362	269	7.8 (57)	8.6 (96)	276	8.7 (16)	243	4.1 (90)	266	85.7 (7)
4363	874	7.2 (42)	8.8 (103)	880	10.1 (47)	767	3.3 (53)	874	78.9 (40)
4401	1,105	7.5 (47)	9.2 (111)	1,110	9.4 (30)	989	3.8 (79)	1,104	74.0 (71)
4501	910	7.8 (57)	8.2 (88)	933	11.8 (97)	796	2.8 (33)	909	74.5 (68)
4601	1,450	6.9 (35)	8.6 (96)	1,489	9.7 (42)	1,303	3.1 (44)	1,442	76.2 (55)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4602	22	*	*	22	*	20	*	22	*
4701	1,109	6.9 (35)	8.6 (96)	1,143	9.3 (27)	1,002	3.7 (72)	1,102	79.9 (34)
5050	1,756	9.7 (106)	5.8 (34)	1,760	12.7 (115)	1,508	3.4 (55)	1,758	59.8 (121)
5051	2,638	11.6 (130)	5.9 (38)	2,656	13.7 (126)	2,246	4.3 (101)	2,636	62.6 (109)
5052	2,699	11.6 (130)	5.4 (29)	2,705	13.4 (123)	2,306	5.6 (127)	2,697	58.6 (126)
5055	756	9.4 (97)	4.6 (19)	1,114	8.6 (14)	686	4.8 (114)	1,085	38.2 (152)
5101	1,961	8.4 (75)	8.1 (86)	1,970	11.0 (73)	1,729	3.7 (72)	1,959	72.7 (81)
5202	699	8.9 (85)	10.4 (135)	709	11.4 (82)	610	3.9 (84)	699	81.4 (25)
5203	36	*	*	36	*	31	*	36	*
5204	29	*	*	29	*	23	*	29	*
5601	1,106	7.1 (40)	11.2 (144)	1,129	10.5 (56)	988	2.7 (26)	1,103	76.2 (55)
5801	1,681	8.6 (78)	7.6 (69)	1,727	11.3 (80)	1,481	3.4 (55)	1,611	63.1 (106)
5803	122	5.7 (8)	8.2 (88)	122	10.7 (65)	108	2.8 (33)	118	56.8 (129)
5804	30	*	*	30	*	26	*	30	*
5806	11	*	*	10	*	8	*	11	*
6101	28	*	*	27	*	23	*	28	*
6102	118	6.8 (30)	10.2 (128)	114	10.5 (56)	101	2.0 (13)	116	76.7 (53)
6103	3,806	10.6 (121)	5.8 (34)	3,920	13.6 (125)	3,238	4.8 (114)	3,812	65.2 (100)
6104	80	*	*	87	*	68	*	80	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6105	30	*	*	30	*	27	*	30	*
6301	4,161	12.6 (136)	5.7 (33)	4,265	14.0 (135)	3,563	5.0 (118)	4,168	63.9 (103)
6312	2,344	10.3 (114)	6.5 (52)	2,407	12.3 (108)	2,048	3.5 (63)	2,343	73.1 (78)
6316	1,201	9.0 (88)	6.8 (58)	1,226	11.7 (91)	1,054	3.7 (72)	1,204	73.2 (76)
6320	120	10.8 (126)	9.2 (111)	122	11.5 (84)	105	4.8 (114)	119	79.0 (39)
6325	132	7.6 (50)	12.9 (150)	148	11.5 (84)	120	4.2 (96)	137	75.9 (61)
6330	607	11.4 (128)	4.9 (25)	648	14.0 (135)	522	5.7 (130)	627	60.1 (118)
6335	849	9.4 (97)	5.9 (38)	873	10.5 (56)	757	5.0 (118)	864	66.0 (96)
6340	58	*	*	60	*	50	*	60	*
6345	23	*	*	24	*	22	*	24	*
6350	17	*	*	17	*	16	*	17	*
6360	137	16.1 (153)	1.5 (1)	142	19.0 (152)	108	7.4 (147)	143	51.7 (137)
7304	147	6.8 (30)	9.5 (120)	148	8.1 (7)	133	3.8 (79)	147	62.6 (109)
7306	2,449	11.3 (127)	6.5 (52)	2,491	11.8 (97)	2,148	5.0 (118)	2,448	60.4 (115)
7308	133	3.8 (4)	8.3 (90)	132	7.6 (6)	121	0.8 (1)	132	55.3 (132)
7310	41	*	*	42	*	36	*	41	*
7401	2,201	7.9 (64)	7.1 (65)	2,251	11.5 (84)	1,937	2.9 (39)	2,197	65.9 (97)
7601	617	7.3 (45)	7.8 (75)	635	9.8 (43)	548	2.7 (26)	618	66.7 (95)
7850	1,112	8.2 (72)	9.1 (108)	1,132	9.4 (30)	1,000	3.4 (55)	1,111	72.8 (79)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7852	41	*	*	44	*	41	*	41	*
8101	1,764	16.7 (154)	2.9 (6)	1,866	20.6 (155)	1,380	6.4 (139)	1,804	46.6 (141)
8102	427	9.4 (97)	9.1 (108)	438	15.3 (142)	345	5.5 (125)	421	40.1 (150)
8105	1,722	10.3 (114)	6.0 (41)	1,756	12.1 (103)	1,495	3.5 (63)	1,727	63.1 (106)
8106	472	12.5 (134)	4.7 (23)	505	12.5 (112)	409	6.1 (135)	484	56.0 (131)
8150	42	*	*	43	*	39	*	41	*
8151	3,068	10.6 (121)	6.3 (48)	3,132	13.7 (126)	2,632	4.5 (106)	2,919	81.5 (23)
8152	144	10.4 (116)	6.3 (48)	144	10.4 (53)	126	7.1 (144)	135	77.0 (51)
8153	14	*	*	14	*	13	*	14	*
8155	42	*	*	45	*	37	*	41	*
8157	13	*	*	14	*	10	*	13	*
8201	2,685	9.4 (97)	6.7 (57)	2,766	10.5 (56)	2,390	4.2 (96)	2,655	61.5 (114)
8202	3,330	7.8 (57)	6.2 (47)	3,761	9.4 (30)	2,977	3.9 (84)	3,720	44.5 (143)
8203	480	10.6 (121)	4.6 (19)	504	11.7 (91)	424	3.1 (44)	485	72.6 (82)
8250	761	10.4 (116)	6.0 (41)	775	11.1 (75)	665	4.8 (114)	754	63.4 (105)
8252	139	15.1 (151)	4.3 (18)	138	13.8 (130)	117	8.5 (150)	140	57.9 (127)
8255	2,760	9.1 (90)	7.5 (68)	2,842	11.3 (80)	2,434	4.2 (96)	2,749	60.1 (118)
8257	1,392	12.4 (133)	4.1 (17)	1,553	10.8 (68)	1,219	7.5 (148)	1,471	74.5 (68)
8258	1	*	*	1	*	1	*	1	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8263	803	7.6 (50)	9.2 (111)	812	8.1 (7)	728	3.6 (67)	802	59.2 (124)
8264	656	9.6 (104)	7.0 (62)	676	11.7 (91)	574	3.7 (72)	655	62.3 (111)
8265	55	*	*	56	*	50	*	55	*
8266	35	*	*	41	*	29	*	41	*
8267	444	9.7 (106)	6.8 (58)	453	12.1 (103)	388	2.8 (33)	443	60.0 (120)
8901	1,067	14.5 (149)	2.5 (2)	1,112	19.0 (152)	855	6.3 (138)	1,066	42.0 (146)
8902	1,197	13.3 (142)	3.6 (14)	1,236	16.3 (147)	992	5.7 (130)	1,203	52.9 (135)
8903	1,106	13.0 (140)	3.6 (14)	1,146	17.5 (151)	898	6.2 (137)	1,112	38.4 (151)
8904	1,951	12.8 (137)	3.5 (13)	2,031	15.6 (144)	1,615	5.6 (127)	1,960	52.3 (136)
8905	1,416	7.8 (57)	6.6 (55)	1,524	12.8 (116)	1,211	3.6 (67)	1,490	53.1 (134)
8908	30	*	*	33	*	24	*	31	*
8909	244	16.0 (152)	2.9 (6)	247	19.8 (154)	193	7.3 (146)	241	57.7 (128)
9012	20	*	*	20	*	17	*	20	*
9013	6	*	*	7	*	4	*	6	*
9016	99	*	*	102	16.7 (149)	81	*	101	37.6 (153)
9017	20	*	*	20	*	16	*	20	*
9018	101	12.9 (138)	4.0 (16)	105	17.1 (150)	83	*	102	43.1 (145)
9041	7	*	*	8	*	7	*	7	*
9045	1,652	9.9 (110)	7.6 (69)	1,683	11.9 (100)	1,428	4.6 (109)	1,656	47.5 (140)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNNS & PedNSS

Table 8A

2015 - 2017 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9050	1,014	13.3 (142)	4.6 (19)	1,039	15.6 (144)	844	5.8 (134)	1,019	58.7 (125)
9051	1,456	6.8 (30)	7.9 (78)	1,501	12.4 (110)	1,252	3.4 (55)	1,482	77.7 (47)
9053	1,482	14.0 (144)	4.6 (19)	1,540	13.8 (130)	1,277	6.9 (142)	1,481	46.5 (142)
9055	50	*	*	58	*	45	*	50	*
9058	974	12.5 (134)	3.0 (8)	1,017	13.7 (126)	828	5.7 (130)	1,003	54.6 (133)
9066	1,212	14.2 (146)	2.7 (4)	1,250	14.4 (139)	1,022	7.1 (144)	1,201	36.1 (154)
9068	988	14.2 (146)	3.4 (11)	1,021	14.5 (140)	834	7.7 (149)	955	40.8 (149)
9069	3,053	14.0 (144)	3.0 (8)	3,121	13.8 (130)	2,615	6.5 (140)	3,040	41.4 (148)
9070	1,103	14.2 (146)	2.6 (3)	1,134	16.3 (147)	905	6.1 (135)	1,109	43.3 (144)
9072	1,952	14.7 (150)	3.3 (10)	2,024	14.1 (137)	1,668	6.6 (141)	1,916	41.5 (147)
9080	2,681	13.2 (141)	3.4 (11)	2,733	15.4 (143)	2,244	5.6 (127)	2,686	50.6 (138)
9081	69	*	*	70	*	60	*	69	*
9082	52	*	*	55	*	43	*	52	*
9083	17	*	*	20	*	16	*	18	*
9084	13	*	*	13	*	10	*	13	*
9085	3	*	*	3	*	3	*	3	*
Michigan	**157,373**	**9.9**	**6.9**	**162,286**	**12.2**	**136,810**	**4.2**	**158,346**	**58.1**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.