WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	43	*	*	*	*	*	*	*
0102	192	91.7	0.0	2.1	1.0	0.0	4.7	0.5
0103	139	90.6	0.0	2.9	0.7	0.0	4.3	1.4
0104	59	*	*	*	*	*	*	*
0105	7	*	*	*	*	*	*	*
0250	60	*	*	*	*	*	*	*
0251	28	*	*	*	*	*	*	*
0252	33	*	*	*	*	*	*	*
0253	47	*	*	*	*	*	*	*
0401	148	95.9	0.0	2.0	0.0	0.0	2.0	0.0
0402	128	88.3	0.8	2.3	0.0	0.0	8.6	0.0
0403	51	*	*	*	*	*	*	*
0404	49	*	*	*	*	*	*	*
0501	163	93.3	1.2	3.1	0.0	0.6	1.8	0.0
0503	41	*	*	*	*	*	*	*
0504	166	88.6	0.0	3.0	3.0	0.6	4.8	0.0
0505	118	79.7	1.7	5.9	0.8	0.0	11.9	0.0
0506	70	*	*	*	*	*	*	*
0507	31	*	*	*	*	*	*	*
0508	9	*	*	*	*	*	*	*
0509	2	*	*	*	*	*	*	*
0650	388	81.2	4.4	5.7	2.8	0.5	5.4	0.0
0651	198	93.9	0.5	3.5	0.0	0.0	2.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0652	117	95.7	0.0	2.6	0.0	0.0	1.7	0.0
0653	147	95.9	0.7	1.4	0.0	0.0	1.4	0.7
0654	127	93.7	1.6	0.8	0.8	0.0	3.1	0.0
0655	126	91.3	0.0	2.4	0.0	0.0	5.6	0.8
0657	17	*	*	*	*	*	*	*
0701	148	83.1	2.0	4.1	0.0	6.1	2.7	2.0
0702	22	*	*	*	*	*	*	*
0703	73	*	*	*	*	*	*	*
0704	15	*	*	*	*	*	*	*
0705	42	*	*	*	*	*	*	*
0706	3	*	*	*	*	*	*	*
0850	296	85.1	5.4	5.4	0.3	0.0	3.0	0.7
0851	273	93.0	0.4	4.8	0.0	0.4	1.5	0.0
0901	541	80.2	0.9	10.5	0.0	0.0	7.6	0.7
0902	13	*	*	*	*	*	*	*
1051	436	81.7	1.1	8.7	1.6	1.4	4.4	1.1
1101	625	35.5	50.2	8.2	0.3	1.3	2.9	1.6
1102	278	71.2	10.1	6.8	0.7	1.1	8.3	1.8
1106	21	*	*	*	*	*	*	*
1250	380	86.1	0.5	9.5	0.0	0.5	2.9	0.5
1251	266	94.0	0.8	2.3	0.0	0.4	1.9	0.8
1252	200	58.0	1.0	36.0	0.0	0.0	4.0	1.0
1253	274	78.1	3.3	10.6	0.0	0.4	5.8	1.8

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1350	849	61.2	18.3	6.7	0.0	5.9	7.4	0.5
1351	40	*	*	*	*	*	*	*
1352	124	66.1	23.4	7.3	0.0	0.0	2.4	0.8
1401	77	*	*	*	*	*	*	*
1402	42	*	*	*	*	*	*	*
1403	10	*	*	*	*	*	*	*
1501	266	49.6	3.8	34.6	0.0	0.4	6.4	5.3
1503	346	61.6	7.5	21.7	0.3	0.6	7.2	1.2
1504	711	40.1	4.9	44.4	0.0	3.7	5.8	1.1
1505	135	83.7	5.2	3.7	0.0	0.7	5.2	1.5
1506	187	78.1	0.5	15.5	1.1	0.5	3.2	1.1
1507	213	86.4	0.5	6.1	0.0	0.5	6.1	0.5
1508	30	*	*	*	*	*	*	*
1509	147	51.7	2.7	40.1	0.7	0.0	1.4	3.4
1511	265	87.9	2.6	4.5	0.0	0.8	3.4	0.8
1512	120	87.5	0.0	5.8	0.0	0.8	5.8	0.0
1513	21	*	*	*	*	*	*	*
1514	74	*	*	*	*	*	*	*
1701	243	57.6	0.4	2.9	16.5	0.4	21.4	0.8
1950	198	93.4	0.0	3.0	0.5	0.0	3.0	0.0
1952	47	*	*	*	*	*	*	*
1953	114	93.0	0.9	4.4	0.0	0.0	1.8	0.0
1954	193	87.0	0.0	11.4	0.0	0.0	1.6	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1955	34	*	*	*	*	*	*	*
1957	163	81.0	0.0	16.6	0.0	0.0	2.5	0.0
1963	11	*	*	*	*	*	*	*
1964	19	*	*	*	*	*	*	*
2150	220	87.7	0.0	0.9	0.0	0.9	10.5	0.0
2151	81	*	*	*	*	*	*	*
2251	144	93.8	0.7	1.4	0.0	0.0	4.2	0.0
2252	64	*	*	*	*	*	*	*
2554	23	*	*	*	*	*	*	*
2555	2,567	46.9	40.3	4.7	0.1	0.4	6.9	0.7
2556	119	35.3	52.1	5.0	0.0	0.8	6.7	0.0
2557	7	*	*	*	*	*	*	*
2558	2	*	*	*	*	*	*	*
2559	3	*	*	*	*	*	*	*
2560	2	*	*	*	*	*	*	*
2562	2	*	*	*	*	*	*	*
2563	5	*	*	*	*	*	*	*
3001	385	17.7	65.5	13.2	0.0	0.3	3.4	0.0
3002	91	*	*	*	*	*	*	*
3004	60	*	*	*	*	*	*	*
3007	7	*	*	*	*	*	*	*
3009	3	*	*	*	*	*	*	*
3201	136	95.6	0.0	2.2	0.7	0.0	1.5	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3301	1,425	38.9	30.0	16.6	0.2	5.3	8.7	0.4
3304	34	*	*	*	*	*	*	*
3305	49	*	*	*	*	*	*	*
3306	88	*	*	*	*	*	*	*
3308	27	*	*	*	*	*	*	*
3309	12	*	*	*	*	*	*	*
3312	123	31.7	31.7	29.3	0.0	0.8	6.5	0.0
3550	24	*	*	*	*	*	*	*
3551	10	*	*	*	*	*	*	*
3552	14	*	*	*	*	*	*	*
3701	358	89.1	0.3	7.0	0.0	0.0	3.6	0.0
3850	1,081	72.1	14.3	5.2	0.0	0.4	7.7	0.4
3901	454	33.5	53.5	6.4	0.0	0.7	5.1	0.9
3902	203	45.3	9.9	20.2	0.0	0.0	24.6	0.0
3903	8	*	*	*	*	*	*	*
3904	3	*	*	*	*	*	*	*
3905	2	*	*	*	*	*	*	*
3950	645	57.8	21.4	8.4	0.3	1.4	10.5	0.2
3954	2	*	*	*	*	*	*	*
4001	92	*	*	*	*	*	*	*
4002	58	*	*	*	*	*	*	*
4103	15	*	*	*	*	*	*	*
4106	889	15.6	43.4	34.2	0.3	0.7	4.8	0.8

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4107	1,064	42.7	27.2	18.9	0.4	1.9	8.6	0.4
4109	383	82.8	2.1	8.1	0.3	1.6	5.2	0.0
4112	189	2.6	1.1	94.7	0.0	1.6	0.0	0.0
4113	116	4.3	16.4	75.0	0.0	0.9	3.4	0.0
4115	1,169	35.9	26.7	19.8	0.3	11.7	4.7	0.8
4351	200	89.0	0.0	4.5	0.0	1.0	5.5	0.0
4352	48	*	*	*	*	*	*	*
4353	145	66.9	0.7	31.0	0.0	0.0	1.4	0.0
4354	276	87.0	2.5	3.6	0.7	0.7	4.7	0.7
4355	111	80.2	0.9	9.9	0.9	0.0	8.1	0.0
4356	145	83.4	2.1	11.7	0.0	0.0	2.8	0.0
4358	54	*	*	*	*	*	*	*
4359	153	69.3	0.7	26.1	0.0	1.3	1.3	1.3
4360	86	*	*	*	*	*	*	*
4361	76	*	*	*	*	*	*	*
4362	108	88.9	0.0	4.6	0.0	0.9	4.6	0.9
4363	295	91.9	0.0	3.1	0.0	0.0	4.7	0.3
4401	377	87.3	0.8	6.4	0.0	0.5	5.0	0.0
4501	357	90.2	0.0	5.6	0.0	0.0	3.9	0.3
4601	477	75.3	1.7	16.8	0.0	0.0	5.5	0.8
4602	9	*	*	*	*	*	*	*
4701	379	92.9	0.3	3.4	0.0	1.3	1.8	0.3
5050	579	52.2	37.8	2.6	0.0	1.0	6.2	0.2

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
5051	882	53.7	33.2	4.8	0.2	2.4	5.3	0.3
5052	1,024	40.1	37.0	2.2	0.3	14.7	5.0	0.6
5055	392	92.6	2.0	0.0	0.3	3.1	0.8	1.3
5101	618	68.4	12.3	12.5	0.0	1.1	5.7	0.0
5102	1	*	*	*	*	*	*	*
5202	294	85.0	0.3	4.1	1.7	0.0	8.5	0.3
5601	336	93.2	0.9	1.8	0.0	1.5	2.7	0.0
5801	577	80.8	5.2	7.6	0.0	0.5	5.4	0.5
5803	44	*	*	*	*	*	*	*
5804	1	*	*	*	*	*	*	*
5806	19	*	*	*	*	*	*	*
6101	11	*	*	*	*	*	*	*
6102	39	*	*	*	*	*	*	*
6103	1,386	54.0	29.0	9.1	0.1	0.4	7.1	0.2
6105	9	*	*	*	*	*	*	*
6106	17	*	*	*	*	*	*	*
6301	1,413	42.3	31.1	19.0	0.1	2.1	5.0	0.4
6312	702	31.6	58.3	3.0	0.0	1.1	5.0	0.9
6316	398	66.8	16.8	8.0	0.3	3.8	4.0	0.3
6320	41	*	*	*	*	*	*	*
6325	54	*	*	*	*	*	*	*
6330	4	*	*	*	*	*	*	*
6335	352	77.6	13.1	1.7	0.0	6.5	0.9	0.3

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6340	16	*	*	*	*	*	*	*
6345	12	*	*	*	*	*	*	*
6360	225	23.6	47.1	16.0	0.0	0.9	12.4	0.0
7304	29	*	*	*	*	*	*	*
7306	836	42.5	34.4	17.3	0.0	0.2	4.8	0.7
7308	36	*	*	*	*	*	*	*
7310	10	*	*	*	*	*	*	*
7401	772	82.1	5.4	6.1	0.0	0.1	5.6	0.6
7601	208	92.3	0.0	5.8	0.0	0.0	1.4	0.5
7850	381	89.8	0.3	5.0	0.3	0.3	4.5	0.0
7852	5	*	*	*	*	*	*	*
8101	660	6.2	90.5	0.2	0.2	0.2	2.7	0.2
8102	124	15.3	8.9	71.8	0.0	0.0	3.2	0.8
8105	511	33.5	58.5	2.9	0.4	0.4	3.9	0.4
8106	158	22.2	60.8	11.4	0.6	0.0	5.1	0.0
8150	10	*	*	*	*	*	*	*
8151	1,035	36.0	39.8	10.3	0.4	2.4	9.9	1.2
8152	23	*	*	*	*	*	*	*
8153	2	*	*	*	*	*	*	*
8155	5	*	*	*	*	*	*	*
8157	1	*	*	*	*	*	*	*
8201	893	45.8	40.3	4.0	0.1	3.8	5.6	0.3
8202	1,413	85.2	10.5	2.1	0.0	0.2	1.5	0.5

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8203	128	57.8	28.9	4.7	0.8	3.1	3.9	0.8
8250	285	19.3	67.4	5.6	0.4	0.7	6.3	0.4
8252	49	*	*	*	*	*	*	*
8255	875	53.4	32.3	7.3	0.3	2.1	4.1	0.5
8257	761	46.6	15.0	0.9	0.0	35.0	2.1	0.4
8263	221	70.6	13.6	7.7	0.5	2.3	5.0	0.5
8264	185	67.0	23.8	4.9	0.5	0.0	3.2	0.5
8267	323	56.7	16.4	19.8	0.0	0.3	6.2	0.6
8901	394	1.5	93.7	0.0	0.3	1.5	2.8	0.3
8902	512	4.9	90.8	1.0	0.0	0.0	2.9	0.4
8903	511	2.2	92.2	0.8	0.0	1.4	3.5	0.0
8904	820	2.2	91.7	1.2	0.4	1.0	3.3	0.2
8905	531	56.5	32.4	8.1	0.0	0.8	2.1	0.2
8909	216	2.3	94.4	0.5	0.0	0.5	1.9	0.5
9045	666	10.5	25.5	57.7	0.2	0.6	5.1	0.5
9050	302	10.6	72.5	15.6	0.0	0.0	1.3	0.0
9051	563	9.6	8.2	81.2	0.0	0.0	0.9	0.0
9053	217	3.2	86.2	1.4	0.0	6.5	2.8	0.0
9055	7	*	*	*	*	*	*	*
9058	345	22.0	65.5	1.7	0.0	8.1	2.6	0.0
9066	349	1.4	96.3	0.9	0.0	0.0	1.1	0.3
9068	339	1.8	96.2	0.6	0.0	0.3	0.9	0.3
9069	987	0.6	97.8	0.4	0.0	0.1	1.0	0.1

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9070	476	3.2	92.9	1.1	0.0	1.1	1.7	0.2
9072	585	1.9	95.9	0.3	0.0	0.3	1.4	0.2
9080	891	20.5	74.6	1.9	0.4	0.1	2.1	0.2
9082	65	*	*	*	*	*	*	*
9083	36	*	*	*	*	*	*	*
9084	12	*	*	*	*	*	*	*
9085	46	*	*	*	*	*	*	*
9086	36	*	*	*	*	*	*	*
Michigan	**55,407**	**51.7**	**29.9**	**10.5**	**0.4**	**2.1**	**4.9**	**0.5**
National (Prior Year)	**98,878**	**47.8**	**21.6**	**18.0**	**1.9**	**2.7**	**3.2**	**4.9**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	43	*	*	*	*	*	*	*
0102	192	0.0	4.2	9.9	63.5	21.9	0.5	0.0
0103	139	0.0	3.6	7.9	65.5	21.6	1.4	0.0
0104	59	*	*	*	*	*	*	*
0105	7	*	*	*	*	*	*	*
0250	60	*	*	*	*	*	*	*
0251	28	*	*	*	*	*	*	*
0252	33	*	*	*	*	*	*	*
0253	47	*	*	*	*	*	*	*
0401	148	0.0	6.1	9.5	62.8	19.6	2.0	0.0
0402	128	0.8	1.6	8.6	64.1	25.0	0.0	0.0
0403	51	*	*	*	*	*	*	*
0404	49	*	*	*	*	*	*	*
0501	163	0.0	2.5	9.8	65.0	22.1	0.6	0.0
0503	41	*	*	*	*	*	*	*
0504	166	0.0	0.0	6.6	63.9	28.3	1.2	0.0
0505	118	0.0	0.0	9.3	67.8	22.0	0.8	0.0
0506	70	*	*	*	*	*	*	*
0507	31	*	*	*	*	*	*	*
0508	9	*	*	*	*	*	*	*
0509	2	*	*	*	*	*	*	*
0650	388	0.0	3.1	9.0	61.9	24.7	1.0	0.3
0651	198	0.5	2.5	10.6	65.2	19.2	2.0	0.0
0652	117	0.0	1.7	5.1	65.8	23.1	4.3	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0653	147	0.0	2.0	10.9	70.7	16.3	0.0	0.0
0654	127	0.0	6.3	13.4	60.6	18.9	0.8	0.0
0655	126	0.0	3.2	11.1	61.9	23.0	0.8	0.0
0657	17	*	*	*	*	*	*	*
0701	148	0.0	2.7	4.7	62.2	25.0	4.7	0.7
0702	22	*	*	*	*	*	*	*
0703	73	*	*	*	*	*	*	*
0704	15	*	*	*	*	*	*	*
0705	42	*	*	*	*	*	*	*
0706	3	*	*	*	*	*	*	*
0850	296	0.0	0.7	12.5	61.1	24.3	0.7	0.7
0851	273	0.0	1.8	9.9	63.7	22.7	1.8	0.0
0901	541	0.0	2.2	10.7	64.0	21.3	1.8	0.0
0902	13	*	*	*	*	*	*	*
1051	436	0.0	3.0	7.3	62.4	24.8	2.5	0.0
1101	625	0.0	3.4	9.6	64.3	20.5	1.4	0.8
1102	278	0.0	4.7	11.5	60.4	21.9	1.4	0.0
1106	21	*	*	*	*	*	*	*
1250	380	0.0	2.4	8.9	62.6	23.4	2.6	0.0
1251	266	0.4	3.0	12.0	63.9	19.9	0.8	0.0
1252	200	0.0	1.0	11.0	68.0	17.0	3.0	0.0
1253	274	0.0	5.5	6.2	69.7	17.9	0.7	0.0
1350	849	0.0	3.1	10.4	62.3	22.7	1.3	0.2
1351	40	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1352	124	0.0	5.6	7.3	68.5	16.9	1.6	0.0
1401	77	*	*	*	*	*	*	*
1402	42	*	*	*	*	*	*	*
1403	10	*	*	*	*	*	*	*
1501	266	0.0	3.8	10.9	57.9	24.8	2.3	0.4
1503	346	0.6	1.7	8.7	63.9	22.8	2.3	0.0
1504	711	0.4	3.1	9.1	60.6	24.1	2.7	0.0
1505	135	0.0	1.5	5.9	62.2	28.9	0.7	0.7
1506	187	0.0	2.7	7.5	61.0	28.3	0.5	0.0
1507	213	0.0	0.5	5.2	69.0	23.5	1.9	0.0
1508	30	*	*	*	*	*	*	*
1509	147	0.0	8.2	6.1	62.6	21.1	2.0	0.0
1511	265	0.0	3.8	12.5	59.6	23.0	1.1	0.0
1512	120	0.0	2.5	6.7	61.7	28.3	0.8	0.0
1513	21	*	*	*	*	*	*	*
1514	74	*	*	*	*	*	*	*
1701	243	0.4	1.2	7.8	60.9	28.4	1.2	0.0
1950	198	0.0	5.6	14.1	64.6	14.6	1.0	0.0
1952	47	*	*	*	*	*	*	*
1953	114	0.0	0.9	7.9	73.7	15.8	1.8	0.0
1954	193	0.0	2.1	9.8	65.3	21.2	1.6	0.0
1955	34	*	*	*	*	*	*	*
1957	163	0.0	2.5	9.2	65.0	21.5	1.8	0.0
1963	11	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1964	19	*	*	*	*	*	*	*
2150	220	0.0	2.7	8.2	61.8	26.8	0.5	0.0
2151	81	*	*	*	*	*	*	*
2251	144	0.0	1.4	10.4	61.8	25.0	1.4	0.0
2252	64	*	*	*	*	*	*	*
2554	23	*	*	*	*	*	*	*
2555	2,567	0.2	2.4	7.9	64.6	23.5	1.4	0.0
2556	119	0.8	1.7	8.4	65.5	22.7	0.8	0.0
2557	7	*	*	*	*	*	*	*
2558	2	*	*	*	*	*	*	*
2559	3	*	*	*	*	*	*	*
2560	2	*	*	*	*	*	*	*
2562	2	*	*	*	*	*	*	*
2563	5	*	*	*	*	*	*	*
3001	385	0.5	4.7	9.1	64.7	19.7	1.3	0.0
3002	91	*	*	*	*	*	*	*
3004	60	*	*	*	*	*	*	*
3007	7	*	*	*	*	*	*	*
3009	3	*	*	*	*	*	*	*
3201	136	0.0	2.2	7.4	61.8	27.2	1.5	0.0
3301	1,425	0.3	2.9	8.0	62.7	23.9	2.1	0.1
3304	34	*	*	*	*	*	*	*
3305	49	*	*	*	*	*	*	*
3306	88	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3308	27	*	*	*	*	*	*	*
3309	12	*	*	*	*	*	*	*
3312	123	0.0	3.3	7.3	62.6	26.0	0.8	0.0
3550	24	*	*	*	*	*	*	*
3551	10	*	*	*	*	*	*	*
3552	14	*	*	*	*	*	*	*
3701	358	0.0	3.4	7.3	69.8	18.7	0.8	0.0
3850	1,081	0.1	3.6	8.9	67.3	19.2	0.5	0.4
3901	454	0.0	4.0	10.6	64.3	19.6	1.5	0.0
3902	203	0.0	3.9	10.3	57.1	26.6	2.0	0.0
3903	8	*	*	*	*	*	*	*
3904	3	*	*	*	*	*	*	*
3905	2	*	*	*	*	*	*	*
3950	645	0.2	2.5	7.6	65.3	22.5	1.9	0.2
3954	2	*	*	*	*	*	*	*
4001	92	*	*	*	*	*	*	*
4002	58	*	*	*	*	*	*	*
4103	15	*	*	*	*	*	*	*
4106	889	0.2	3.3	10.5	59.4	24.7	1.7	0.2
4107	1,064	0.2	2.6	9.7	60.2	25.6	1.6	0.1
4109	383	0.0	2.1	6.0	64.8	25.3	1.6	0.3
4112	189	0.0	3.2	6.3	54.5	33.3	2.6	0.0
4113	116	0.0	0.9	8.6	41.4	34.5	7.8	6.9
4115	1,169	0.2	2.2	6.3	58.9	30.4	1.8	0.2

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4351	200	0.0	2.0	8.0	73.0	16.5	0.5	0.0
4352	48	*	*	*	*	*	*	*
4353	145	0.0	4.1	7.6	65.5	20.7	2.1	0.0
4354	276	0.0	2.2	11.6	67.4	17.8	1.1	0.0
4355	111	0.0	2.7	10.8	58.6	23.4	4.5	0.0
4356	145	0.7	4.8	4.8	69.0	20.0	0.7	0.0
4358	54	*	*	*	*	*	*	*
4359	153	0.0	3.9	7.2	64.7	22.9	1.3	0.0
4360	86	*	*	*	*	*	*	*
4361	76	*	*	*	*	*	*	*
4362	108	0.0	0.9	7.4	75.0	16.7	0.0	0.0
4363	295	0.0	4.1	5.4	66.8	22.7	1.0	0.0
4401	377	0.0	4.0	8.0	61.5	24.4	1.9	0.3
4501	357	0.0	2.8	9.2	64.4	22.7	0.8	0.0
4601	477	0.0	2.5	10.3	64.6	22.0	0.6	0.0
4602	9	*	*	*	*	*	*	*
4701	379	0.0	1.6	7.4	61.2	28.2	1.6	0.0
5050	579	0.0	2.4	4.7	57.5	32.8	2.6	0.0
5051	882	0.0	0.7	4.6	64.9	28.3	1.5	0.0
5052	1,024	0.0	2.3	6.3	61.2	27.5	2.6	0.0
5055	392	0.0	0.3	1.3	51.3	42.6	4.6	0.0
5101	618	0.2	0.8	4.9	61.7	30.1	2.4	0.0
5102	1	*	*	*	*	*	*	*
5202	294	0.3	2.0	6.8	61.6	28.9	0.3	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5601	336	0.0	2.1	6.5	67.6	22.6	1.2	0.0
5801	577	0.0	2.8	9.9	63.1	22.4	1.9	0.0
5803	44	*	*	*	*	*	*	*
5804	1	*	*	*	*	*	*	*
5806	19	*	*	*	*	*	*	*
6101	11	*	*	*	*	*	*	*
6102	39	*	*	*	*	*	*	*
6103	1,386	0.4	3.5	9.2	66.1	19.6	1.3	0.0
6105	9	*	*	*	*	*	*	*
6106	17	*	*	*	*	*	*	*
6301	1,413	0.1	2.6	9.5	61.2	24.1	2.5	0.0
6312	702	0.0	0.9	4.6	58.7	32.9	3.0	0.0
6316	398	0.0	0.8	3.5	61.1	30.9	3.8	0.0
6320	41	*	*	*	*	*	*	*
6325	54	*	*	*	*	*	*	*
6330	4	*	*	*	*	*	*	*
6335	352	0.0	1.1	3.4	53.4	40.3	1.7	0.0
6340	16	*	*	*	*	*	*	*
6345	12	*	*	*	*	*	*	*
6360	225	0.4	1.3	7.1	61.8	26.2	2.7	0.4
7304	29	*	*	*	*	*	*	*
7306	836	0.1	2.6	9.8	64.4	21.2	1.9	0.0
7308	36	*	*	*	*	*	*	*
7310	10	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
7401	772	0.1	3.4	8.2	64.5	22.5	1.2	0.1
7601	208	0.0	1.9	7.7	61.1	27.4	1.9	0.0
7850	381	0.3	2.6	8.7	70.6	16.3	1.6	0.0
7852	5	*	*	*	*	*	*	*
8101	660	0.5	2.0	9.8	64.7	20.9	2.1	0.0
8102	124	0.0	1.6	4.0	61.3	33.1	0.0	0.0
8105	511	0.0	1.8	4.3	61.6	31.1	1.2	0.0
8106	158	0.0	1.3	5.1	69.0	20.3	4.4	0.0
8150	10	*	*	*	*	*	*	*
8151	1,035	0.2	2.4	5.5	60.1	28.6	3.0	0.2
8152	23	*	*	*	*	*	*	*
8153	2	*	*	*	*	*	*	*
8155	5	*	*	*	*	*	*	*
8157	1	*	*	*	*	*	*	*
8201	893	0.0	3.0	5.4	61.7	27.7	2.2	0.0
8202	1,413	0.1	1.3	3.5	56.5	33.8	4.2	0.6
8203	128	0.0	1.6	7.0	60.2	29.7	1.6	0.0
8250	285	0.7	2.1	9.1	56.5	29.8	1.8	0.0
8252	49	*	*	*	*	*	*	*
8255	875	0.0	2.1	7.1	64.8	24.8	1.3	0.0
8257	761	0.0	0.9	3.9	58.6	31.9	4.5	0.1
8263	221	0.0	0.5	5.4	66.5	26.7	0.9	0.0
8264	185	0.0	5.4	1.6	64.9	23.8	4.3	0.0
8267	323	0.0	4.3	7.4	59.1	28.5	0.6	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8901	394	0.3	2.5	8.4	67.0	20.1	1.8	0.0
8902	512	0.0	2.3	8.0	65.0	22.3	2.3	0.0
8903	511	0.0	2.5	9.4	61.4	25.4	1.2	0.0
8904	820	0.2	2.4	7.0	64.0	24.4	2.0	0.0
8905	531	0.0	1.5	4.3	58.9	32.2	3.0	0.0
8909	216	0.5	4.6	9.7	57.9	25.9	1.4	0.0
9045	666	0.2	3.2	7.2	58.1	28.4	2.7	0.3
9050	302	0.3	3.0	6.3	59.3	28.8	2.3	0.0
9051	563	0.2	3.0	8.2	50.6	30.7	7.3	0.0
9053	217	0.0	1.8	9.2	61.3	25.3	2.3	0.0
9055	7	*	*	*	*	*	*	*
9058	345	0.6	2.0	4.6	62.9	26.1	3.5	0.3
9066	349	0.3	2.3	6.9	67.3	20.9	2.3	0.0
9068	339	0.0	3.2	9.1	60.2	26.0	1.5	0.0
9069	987	0.0	3.5	6.6	63.5	23.9	2.4	0.0
9070	476	0.0	2.3	10.1	62.0	23.9	1.7	0.0
9072	585	0.2	2.6	9.1	67.4	20.0	0.9	0.0
9080	891	0.0	2.5	6.3	61.1	28.2	2.0	0.0
9082	65	*	*	*	*	*	*	*
9083	36	*	*	*	*	*	*	*
9084	12	*	*	*	*	*	*	*
9085	46	*	*	*	*	*	*	*
9086	36	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Michigan	**55,413**	**0.1**	**2.6**	**7.7**	**62.5**	**25.1**	**2.0**	**0.1**
National (Prior Year)	**103,413**	**0.1**	**2.7**	**7.5**	**61.6**	**26.0**	**2.0**	**0.1**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	23	*	*	4	*	*	8	*	12	*	42	*	42	*
0102	135	5.9 (99)	60.7 (62)	65	*	*	32	*	41	*	177	3.4 (5)	177	9.6 (33)
0103	121	2.5 (12)	59.5 (51)	72	*	*	20	*	28	*	125	8.0 (87)	125	8.0 (21)
0104	47	*	*	29	*	*	8	*	13	*	56	*	56	*
0105	7	*	*	6	*	*	1	*	0	*	7	*	7	*
0250	55	*	*	34	*	*	7	*	17	*	52	*	53	*
0251	17	*	*	6	*	*	1	*	8	*	28	*	27	*
0252	33	*	*	27	*	*	4	*	8	*	30	*	30	*
0253	25	*	*	7	*	*	4	*	14	*	45	*	45	*
0401	139	2.9 (23)	58.3 (38)	105	18.1 (37)	56.2 (64)	19	*	35	*	137	4.4 (15)	137	7.3 (14)
0402	121	6.6 (105)	58.7 (45)	60	*	*	23	*	14	*	121	10.7 (108)	121	9.9 (40)
0403	48	*	*	33	*	*	7	*	11	*	48	*	48	*
0404	45	*	*	36	*	*	5	*	11	*	47	*	47	*
0501	83	*	*	27	*	*	24	*	30	*	149	7.4 (68)	149	6.0 (5)
0503	38	*	*	24	*	*	5	*	6	*	39	*	39	*
0504	144	2.8 (20)	53.5 (12)	67	*	*	30	*	27	*	161	8.1 (89)	161	10.6 (47)
0505	91	*	*	44	*	*	21	*	16	*	107	6.5 (49)	108	8.3 (24)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	47	*	*	12	*	*	10	*	4	*	64	*	65	*
0507	16	*	*	8	*	*	1	*	8	*	31	*	31	*
0508	5	*	*	1	*	*	0	*	4	*	8	*	8	*
0509	2	*	*	1	*	*	0	*	1	*	2	*	2	*
0650	305	3.3 (32)	61.0 (66)	182	20.9 (57)	50.5 (33)	64	*	84	*	365	5.5 (32)	365	11.2 (60)
0651	189	4.8 (77)	60.3 (58)	153	20.9 (57)	48.4 (19)	29	*	26	*	184	3.3 (4)	184	7.6 (19)
0652	74	*	*	38	*	*	19	*	13	*	112	2.7 (1)	112	8.0 (21)
0653	125	4.0 (54)	57.6 (33)	94	*	*	23	*	20	*	141	4.3 (13)	141	13.5 (88)
0654	113	2.7 (17)	64.6 (94)	76	*	*	17	*	33	*	116	9.5 (103)	115	13.0 (81)
0655	115	3.5 (37)	66.1 (99)	73	*	*	14	*	32	*	118	7.6 (72)	119	15.1 (103)
0657	13	*	*	4	*	*	1	*	5	*	17	*	17	*
0701	103	8.7 (110)	41.7 (1)	45	*	*	32	*	52	*	136	4.4 (15)	137	11.7 (67)
0702	21	*	*	12	*	*	1	*	3	*	18	*	18	*
0703	62	*	*	26	*	*	9	*	19	*	62	*	62	*
0704	14	*	*	11	*	*	4	*	2	*	13	*	13	*
0705	12	*	*	6	*	*	8	*	9	*	40	*	40	*
0706	1	*	*	0	*	*	0	*	0	*	3	*	3	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	212	2.4 (10)	61.8 (74)	134	17.2 (29)	58.2 (74)	47	*	51	*	269	7.8 (79)	270	15.2 (106)
0851	154	7.1 (106)	51.9 (9)	45	*	*	35	*	60	*	250	8.0 (87)	250	11.2 (60)
0901	465	4.1 (56)	60.6 (61)	361	18.0 (36)	52.1 (44)	89	*	67	*	508	7.9 (83)	508	10.2 (42)
0902	13	*	*	11	*	*	0	*	2	*	13	*	13	*
1051	397	3.8 (45)	51.9 (9)	328	17.4 (32)	53.4 (52)	57	*	59	*	408	9.3 (102)	409	11.7 (67)
1101	532	3.8 (45)	62.4 (80)	403	17.9 (35)	55.6 (60)	116	53.4 (15)	86	*	572	5.6 (34)	571	11.9 (70)
1102	223	1.8 (3)	60.5 (59)	157	14.6 (9)	56.7 (67)	35	*	46	*	245	6.1 (43)	245	9.4 (32)
1106	19	*	*	15	*	*	1	*	8	*	18	*	19	*
1250	282	7.4 (107)	62.4 (80)	161	16.1 (23)	54.0 (55)	62	*	53	*	346	7.5 (69)	346	7.2 (13)
1251	252	5.2 (88)	58.3 (38)	187	15.0 (13)	55.6 (60)	39	*	49	*	243	4.1 (10)	243	7.4 (16)
1252	159	2.5 (12)	59.7 (55)	82	*	*	32	*	33	*	188	4.8 (20)	188	14.9 (100)
1253	212	5.7 (96)	57.5 (31)	114	18.4 (41)	49.1 (24)	43	*	48	*	244	5.3 (25)	244	11.9 (70)
1350	713	3.2 (29)	60.7 (62)	426	21.6 (63)	55.2 (57)	125	51.2 (12)	156	31.4 (5)	762	7.6 (72)	763	9.7 (38)
1351	35	*	*	30	*	*	3	*	11	*	40	*	40	*
1352	76	*	*	42	*	*	18	*	25	*	119	4.2 (12)	119	5.9 (4)
1401	67	*	*	50	*	*	11	*	15	*	73	*	73	*
1402	38	*	*	29	*	*	3	*	8	*	39	*	39	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1403	10	*	*	6	*	*	0	*	1	*	10	*	10	*
1501	234	4.3 (60)	67.5 (101)	176	50.6 (78)	21.6 (2)	27	*	42	*	232	7.8 (79)	232	9.1 (29)
1503	296	2.7 (17)	62.8 (86)	168	16.7 (26)	56.5 (66)	42	*	37	*	313	8.6 (97)	313	15.3 (107)
1504	589	3.9 (50)	61.8 (74)	423	17.3 (30)	48.0 (18)	103	56.3 (22)	198	44.9 (29)	653	8.4 (94)	655	9.6 (33)
1505	106	3.8 (45)	52.8 (11)	73	*	*	22	*	40	*	130	5.4 (29)	130	15.4 (110)
1506	152	3.3 (32)	62.5 (83)	51	*	*	27	*	32	*	173	6.9 (56)	173	8.7 (27)
1507	166	3.6 (42)	51.2 (7)	77	*	*	24	*	49	*	192	6.8 (54)	192	8.3 (24)
1508	13	*	*	4	*	*	2	*	3	*	23	*	23	*
1509	115	6.1 (101)	68.7 (108)	69	*	*	18	*	36	*	123	8.9 (100)	124	7.3 (14)
1511	207	3.9 (50)	58.9 (49)	100	17.0 (27)	47.0 (12)	41	*	62	*	250	7.6 (72)	251	13.9 (91)
1512	74	*	*	24	*	*	24	*	30	*	112	6.3 (46)	113	10.6 (47)
1513	19	*	*	7	*	*	0	*	4	*	17	*	17	*
1514	47	*	*	38	*	*	9	*	32	*	71	*	71	*
1701	175	4.0 (54)	62.9 (87)	92	*	*	38	*	38	*	220	8.2 (91)	219	10.5 (45)
1950	186	6.5 (104)	62.4 (80)	143	18.9 (44)	53.8 (54)	28	*	51	*	187	11.2 (110)	188	16.0 (112)
1952	42	*	*	35	*	*	5	*	15	*	45	*	45	*
1953	106	5.7 (96)	55.7 (23)	85	*	*	15	*	30	*	104	7.7 (76)	104	16.3 (114)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1954	174	7.5 (109)	58.6 (43)	132	12.9 (2)	61.4 (78)	29	*	40	*	179	11.7 (114)	179	14.5 (96)
1955	31	*	*	23	*	*	4	*	8	*	32	*	32	*
1957	148	4.7 (71)	68.2 (104)	112	15.2 (14)	52.7 (47)	16	*	45	*	156	8.3 (93)	156	12.2 (73)
1963	10	*	*	10	*	*	1	*	5	*	10	*	10	*
1964	18	*	*	16	*	*	3	*	4	*	16	*	16	*
2150	133	0.8 (1)	51.1 (6)	72	*	*	26	*	52	*	196	10.2 (105)	196	8.7 (27)
2151	50	*	*	32	*	*	17	*	22	*	67	*	67	*
2251	119	4.2 (58)	55.5 (20)	49	*	*	17	*	21	*	135	3.7 (7)	135	14.8 (98)
2252	45	*	*	7	*	*	7	*	12	*	53	*	53	*
2554	19	*	*	13	*	*	4	*	4	*	21	*	21	*
2555	2,007	5.2 (88)	56.7 (28)	865	17.3 (30)	51.1 (35)	496	54.8 (20)	391	35.0 (9)	2,381	8.2 (91)	2,385	16.0 (112)
2556	104	1.9 (6)	59.6 (53)	87	*	*	15	*	21	*	115	6.1 (43)	115	25.2 (119)
2557	3	*	*	0	*	*	3	*	1	*	6	*	6	*
2558	2	*	*	2	*	*	0	*	0	*	2	*	2	*
2559	3	*	*	2	*	*	1	*	0	*	3	*	3	*
2560	2	*	*	0	*	*	0	*	0	*	2	*	2	*
2562	1	*	*	0	*	*	2	*	1	*	1	*	1	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2563	5	*	*	2	*	*	0	*	1	*	5	*	5	*
3001	359	3.3 (32)	62.7 (85)	267	22.8 (69)	44.6 (8)	72	*	242	47.1 (33)	355	5.1 (23)	354	10.5 (45)
3002	88	*	*	70	*	*	13	*	64	*	88	*	88	*
3004	58	*	*	51	*	*	7	*	10	*	54	*	54	*
3007	6	*	*	2	*	*	1	*	2	*	5	*	5	*
3009	2	*	*	0	*	*	0	*	1	*	1	*	1	*
3201	106	3.8 (45)	67.9 (102)	87	*	*	17	*	37	*	131	7.6 (72)	131	18.3 (117)
3301	1,194	5.0 (83)	59.5 (51)	970	20.3 (53)	52.8 (48)	227	49.8 (10)	172	32.0 (7)	1,327	6.5 (49)	1,328	10.2 (42)
3304	25	*	*	20	*	*	3	*	6	*	32	*	32	*
3305	42	*	*	36	*	*	6	*	8	*	48	*	48	*
3306	76	*	*	65	*	*	16	*	24	*	87	*	87	*
3308	22	*	*	16	*	*	0	*	8	*	25	*	25	*
3309	12	*	*	9	*	*	1	*	4	*	11	*	11	*
3312	110	4.5 (68)	68.2 (104)	90	*	*	22	*	13	*	114	4.4 (15)	114	7.0 (11)
3550	19	*	*	3	*	*	3	*	6	*	22	*	22	*
3551	6	*	*	2	*	*	1	*	4	*	8	*	8	*
3552	9	*	*	0	*	*	1	*	3	*	11	*	11	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3701	206	4.9 (82)	57.8 (34)	129	15.5 (18)	55.8 (62)	65	*	102	20.6 (2)	327	7.0 (60)	326	17.2 (115)
3850	818	6.1 (101)	60.1 (57)	455	15.2 (14)	52.5 (46)	171	43.3 (4)	194	42.3 (24)	1,004	5.4 (29)	1,005	11.3 (62)
3901	351	4.6 (69)	57.8 (34)	155	20.0 (50)	47.1 (14)	70	*	93	*	421	5.2 (24)	421	10.7 (51)
3902	178	2.8 (20)	68.5 (107)	82	*	*	32	*	36	*	187	11.2 (110)	188	6.9 (10)
3903	7	*	*	7	*	*	0	*	5	*	8	*	8	*
3904	3	*	*	0	*	*	1	*	1	*	3	*	3	*
3905	2	*	*	2	*	*	0	*	0	*	2	*	2	*
3950	533	3.8 (45)	57.0 (29)	331	15.7 (20)	58.0 (71)	92	*	181	39.8 (18)	590	7.5 (69)	590	9.3 (31)
3954	1	*	*	1	*	*	0	*	1	*	2	*	2	*
4001	74	*	*	26	*	*	0	*	12	*	79	*	79	*
4002	50	*	*	27	*	*	0	*	10	*	48	*	48	*
4103	7	*	*	3	*	*	2	*	4	*	12	*	12	*
4106	656	2.9 (23)	61.7 (73)	308	20.5 (54)	47.1 (14)	145	54.5 (17)	280	45.0 (30)	820	7.3 (67)	819	11.0 (57)
4107	899	4.3 (60)	57.5 (31)	491	14.7 (10)	51.3 (38)	171	42.1 (3)	395	43.5 (27)	1,001	5.8 (38)	1,000	13.9 (91)
4109	154	5.2 (88)	49.4 (3)	58	*	*	50	*	89	*	346	8.1 (89)	350	15.1 (103)
4112	189	2.1 (7)	65.6 (98)	171	77.2 (79)	5.3 (1)	22	*	24	*	179	12.3 (117)	179	10.6 (47)
4113	102	1.0 (2)	73.5 (110)	25	*	*	15	*	32	*	112	12.5 (118)	112	8.0 (21)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4115	685	3.9 (50)	54.7 (19)	169	16.0 (22)	46.2 (9)	185	54.6 (19)	454	42.3 (24)	1,089	7.0 (60)	1,090	11.6 (66)
4351	189	2.6 (15)	58.7 (45)	138	26.1 (77)	43.5 (5)	36	*	38	*	188	6.9 (56)	188	9.6 (33)
4352	46	*	*	35	*	*	6	*	10	*	47	*	47	*
4353	126	2.4 (10)	61.1 (68)	82	*	*	23	*	32	*	135	5.9 (40)	135	9.6 (33)
4354	219	5.0 (83)	53.9 (14)	131	23.7 (72)	51.1 (35)	38	*	66	*	250	4.4 (15)	250	10.8 (53)
4355	88	*	*	20	*	*	13	*	32	*	108	3.7 (7)	108	11.1 (59)
4356	107	4.7 (71)	49.5 (4)	31	*	*	29	*	25	*	132	3.0 (2)	132	9.1 (29)
4358	47	*	*	37	*	*	4	*	6	*	47	*	47	*
4359	129	4.7 (71)	64.3 (92)	60	*	*	18	*	34	*	144	11.8 (115)	145	12.4 (75)
4360	82	*	*	66	*	*	10	*	16	*	81	*	81	*
4361	61	*	*	52	*	*	10	*	15	*	73	*	73	*
4362	95	*	*	72	*	*	25	*	24	*	99	*	99	*
4363	280	3.2 (29)	62.5 (83)	204	17.6 (34)	51.5 (41)	37	*	71	*	270	7.0 (60)	270	13.0 (81)
4401	373	5.6 (93)	54.2 (16)	326	14.7 (10)	55.2 (57)	61	*	117	31.6 (6)	355	3.4 (5)	355	3.9 (2)
4501	272	7.4 (107)	55.5 (20)	220	14.5 (8)	58.2 (74)	54	*	64	*	342	7.9 (83)	342	12.6 (78)
4601	415	4.3 (60)	54.2 (16)	113	23.0 (70)	44.2 (7)	95	*	121	42.1 (22)	433	7.2 (64)	434	14.1 (94)
4602	6	*	*	3	*	*	0	*	2	*	9	*	9	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4701	359	4.7 (71)	51.8 (8)	253	15.4 (16)	53.4 (52)	61	*	125	18.4 (1)	358	11.2 (110)	357	15.4 (110)
5050	561	5.0 (83)	60.8 (64)	492	16.3 (24)	55.5 (59)	120	53.3 (14)	156	37.2 (13)	540	8.7 (98)	538	13.6 (89)
5051	833	4.8 (77)	58.9 (49)	733	19.5 (47)	51.8 (43)	187	50.3 (11)	256	42.2 (23)	827	7.7 (76)	827	10.8 (53)
5052	970	4.8 (77)	63.3 (89)	824	21.5 (62)	47.0 (12)	188	48.9 (7)	246	39.0 (17)	954	8.7 (98)	954	11.0 (57)
5055	271	2.6 (15)	57.9 (36)	115	18.3 (39)	48.7 (21)	37	*	110	45.5 (31)	372	11.8 (115)	371	4.6 (3)
5101	577	4.3 (60)	58.4 (41)	462	14.3 (6)	51.3 (38)	118	39.0 (1)	137	36.5 (12)	569	10.5 (106)	570	13.7 (90)
5102	1	*	*	1	*	*	0	*	0	*	1	*	1	*
5202	258	3.5 (37)	50.8 (5)	196	22.4 (65)	43.9 (6)	38	*	80	*	267	4.1 (10)	268	19.0 (118)
5601	282	2.8 (20)	59.6 (53)	50	*	*	46	*	168	28.6 (3)	320	5.9 (40)	321	15.3 (107)
5801	458	4.1 (56)	57.9 (36)	271	23.6 (71)	46.9 (10)	93	*	108	40.7 (19)	516	7.8 (79)	516	12.2 (73)
5803	40	*	*	31	*	*	7	*	18	*	42	*	42	*
5804	1	*	*	0	*	*	0	*	0	*	0	*	0	*
5806	15	*	*	13	*	*	1	*	6	*	19	*	19	*
6101	9	*	*	8	*	*	0	*	5	*	11	*	11	*
6102	37	*	*	24	*	*	3	*	7	*	39	*	39	*
6103	1,282	3.9 (50)	61.4 (70)	927	20.7 (55)	48.8 (22)	167	40.7 (2)	163	32.5 (8)	1,269	7.2 (64)	1,272	13.3 (87)
6105	9	*	*	7	*	*	0	*	4	*	8	*	9	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6106	17	*	*	12	*	*	2	*	4	*	17	*	17	*
6301	1,290	4.3 (60)	58.8 (48)	880	18.5 (42)	49.3 (27)	270	48.5 (6)	289	37.4 (14)	1,309	4.9 (21)	1,310	8.6 (26)
6312	655	3.1 (26)	60.5 (59)	476	14.9 (12)	51.3 (38)	140	52.1 (13)	184	51.1 (35)	659	3.2 (3)	659	6.7 (8)
6316	383	3.4 (36)	53.8 (13)	325	15.4 (16)	59.1 (76)	76	*	115	35.7 (10)	368	7.9 (83)	368	10.6 (47)
6320	34	*	*	11	*	*	7	*	13	*	39	*	39	*
6325	39	*	*	12	*	*	6	*	20	*	53	*	53	*
6330	1	*	*	1	*	*	0	*	0	*	4	*	4	*
6335	227	3.1 (26)	47.6 (2)	77	*	*	45	*	72	*	334	7.5 (69)	334	7.5 (18)
6340	13	*	*	5	*	*	0	*	1	*	16	*	15	*
6345	9	*	*	7	*	*	0	*	5	*	12	*	12	*
6360	164	5.5 (92)	61.0 (66)	36	*	*	40	*	51	*	208	6.7 (53)	208	9.6 (33)
7304	22	*	*	11	*	*	1	*	6	*	27	*	27	*
7306	676	5.2 (88)	58.7 (45)	473	21.1 (60)	48.4 (19)	133	63.9 (26)	162	38.3 (16)	787	5.7 (36)	787	10.2 (42)
7308	32	*	*	21	*	*	4	*	1	*	29	*	29	*
7310	2	*	*	1	*	*	0	*	0	*	7	*	7	*
7401	734	6.1 (101)	56.4 (26)	623	18.3 (39)	57.3 (69)	91	*	180	36.1 (11)	702	5.3 (25)	704	13.1 (83)
7601	43	*	*	23	*	*	34	*	42	*	189	5.8 (38)	189	7.4 (16)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7850	367	4.4 (66)	60.8 (64)	303	22.4 (65)	46.9 (10)	53	*	85	*	357	8.4 (94)	357	13.2 (86)
7852	4	*	*	4	*	*	0	*	2	*	5	*	5	*
8101	561	3.2 (29)	64.9 (95)	407	19.7 (48)	49.6 (29)	101	65.3 (28)	96	*	621	6.6 (52)	623	12.7 (79)
8102	94	*	*	62	*	*	17	*	18	*	112	12.5 (118)	112	12.5 (76)
8105	381	4.2 (58)	65.1 (97)	225	22.7 (68)	52.9 (49)	98	*	122	41.0 (20)	472	10.8 (109)	472	14.8 (98)
8106	72	*	*	44	*	*	28	*	23	*	147	6.8 (54)	147	6.8 (9)
8150	7	*	*	2	*	*	0	*	1	*	8	*	8	*
8151	964	3.5 (37)	61.4 (70)	650	18.2 (38)	50.6 (34)	195	53.8 (16)	297	41.8 (21)	933	7.9 (83)	933	14.0 (93)
8152	23	*	*	17	*	*	5	*	10	*	22	*	22	*
8153	2	*	*	2	*	*	0	*	0	*	2	*	2	*
8155	4	*	*	3	*	*	0	*	2	*	4	*	4	*
8157	1	*	*	0	*	*	1	*	0	*	0	*	0	*
8201	671	6.0 (100)	56.5 (27)	399	19.0 (45)	50.4 (32)	174	48.9 (7)	254	29.5 (4)	826	6.9 (56)	826	11.3 (62)
8202	998	4.7 (71)	54.0 (15)	500	22.6 (67)	41.6 (4)	276	58.3 (24)	366	49.2 (34)	1,346	7.2 (64)	1,346	3.4 (1)
8203	112	1.8 (3)	61.6 (72)	100	17.0 (27)	49.0 (23)	25	*	41	*	122	5.7 (36)	122	13.1 (83)
8250	252	3.6 (42)	58.3 (38)	203	19.7 (48)	52.2 (45)	60	*	91	*	259	10.0 (104)	259	13.1 (83)
8252	43	*	*	29	*	*	6	*	11	*	44	*	44	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8255	718	5.6 (93)	56.3 (25)	475	13.9 (3)	57.3 (69)	182	48.9 (7)	275	37.8 (15)	805	7.8 (79)	805	11.3 (62)
8257	501	4.4 (66)	57.1 (30)	235	21.3 (61)	40.4 (3)	123	54.5 (17)	213	46.9 (32)	711	10.5 (106)	711	6.3 (7)
8263	182	2.7 (17)	54.4 (18)	131	22.1 (64)	47.3 (16)	50	*	64	*	206	9.2 (101)	206	10.7 (51)
8264	134	3.0 (25)	61.9 (76)	97	*	*	42	*	69	*	170	11.2 (110)	170	10.0 (41)
8267	191	5.8 (98)	55.5 (20)	119	14.3 (6)	56.3 (65)	74	*	93	*	290	6.9 (56)	290	7.6 (19)
8901	369	3.5 (37)	65.0 (96)	239	16.3 (24)	56.9 (68)	61	*	35	*	365	6.3 (46)	365	12.9 (80)
8902	432	5.6 (93)	58.6 (43)	242	17.4 (32)	55.8 (62)	82	*	80	*	472	4.0 (9)	472	12.5 (76)
8903	462	4.8 (77)	63.2 (88)	294	20.1 (51)	54.1 (56)	105	76.2 (29)	68	*	470	4.3 (13)	470	17.4 (116)
8904	685	3.1 (26)	61.2 (69)	482	20.7 (55)	51.2 (37)	155	63.2 (25)	127	57.5 (36)	764	5.5 (32)	764	14.1 (94)
8905	438	2.5 (12)	58.4 (41)	294	15.6 (19)	49.7 (30)	83	*	104	42.3 (24)	496	7.1 (63)	496	10.9 (56)
8909	185	4.3 (60)	64.3 (92)	119	21.0 (59)	58.0 (71)	43	*	31	*	202	5.0 (22)	203	15.3 (107)
9045	502	1.8 (3)	68.7 (108)	388	24.2 (76)	47.7 (17)	118	45.8 (5)	87	*	614	6.4 (48)	615	6.2 (6)
9050	253	4.7 (71)	64.0 (90)	116	24.1 (75)	49.1 (24)	47	*	39	*	290	4.5 (19)	290	11.7 (67)
9051	437	2.1 (7)	68.4 (106)	289	14.2 (5)	50.2 (31)	45	*	73	*	518	7.7 (76)	518	7.1 (12)
9053	190	2.1 (7)	64.2 (91)	128	14.1 (4)	59.4 (77)	36	*	22	*	202	5.4 (29)	202	11.4 (65)
9055	7	*	*	6	*	*	2	*	2	*	7	*	7	*
9058	209	4.8 (77)	62.2 (78)	146	18.5 (42)	49.3 (27)	56	*	72	*	333	8.4 (94)	333	15.0 (101)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9066	254	3.5 (37)	68.1 (103)	141	11.3 (1)	65.2 (79)	57	*	58	*	327	6.1 (43)	327	15.0 (101)
9068	328	3.7 (44)	66.8 (100)	244	23.8 (74)	53.3 (51)	60	*	33	*	320	5.9 (40)	320	9.7 (38)
9069	890	3.3 (32)	62.2 (78)	735	19.0 (45)	51.7 (42)	174	55.7 (21)	97	*	938	6.5 (49)	939	10.8 (53)
9070	351	4.6 (69)	55.8 (24)	228	23.7 (72)	53.1 (50)	85	*	69	*	453	5.3 (25)	453	14.6 (97)
9072	483	5.0 (83)	59.8 (56)	293	20.1 (51)	49.1 (24)	117	58.1 (23)	78	*	563	5.3 (25)	563	15.1 (103)
9080	760	5.0 (83)	62.1 (77)	510	15.7 (20)	58.0 (71)	186	64.5 (27)	183	43.7 (28)	853	5.6 (34)	853	12.1 (72)
9082	60	*	*	29	*	*	10	*	8	*	57	*	57	*
9083	36	*	*	34	*	*	3	*	1	*	35	*	35	*
9084	11	*	*	11	*	*	0	*	0	*	12	*	12	*
9085	45	*	*	15	*	*	22	*	4	*	42	*	42	*
9086	34	*	*	20	*	*	15	*	4	*	30	*	30	*
Michigan	**45,296**	**4.3**	**59.5**	**29,118**	**19.1**	**51.0**	**9,090**	**49.4**	**12,050**	**37.9**	**51,368**	**7.0**	**51,399**	**11.4**
National (Prior Year)	**81,973**	**4.2**	**58.2**	**63,088**	**20.7**	**48.2**	**19,941**	**34.5**	**19,251**	**29.8**	**66,742**	**9.3**	**65,904**	**11.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	42	*	*	43	*	40	*	32	*
0102	173	88.4 (34)	7.5 (73)	191	33.0 (47)	171	18.7 (57)	156	90.4 (41)
0103	123	81.3 (76)	14.6 (104)	135	43.0 (7)	121	24.8 (32)	117	89.7 (47)
0104	56	*	*	59	*	55	*	47	*
0105	7	*	*	7	*	5	*	5	*
0250	53	*	*	59	*	41	*	49	*
0251	28	*	*	28	*	22	*	21	*
0252	30	*	*	33	*	28	*	26	*
0253	45	*	*	47	*	44	*	42	*
0401	135	86.7 (45)	7.4 (71)	147	35.4 (41)	136	24.3 (36)	117	92.3 (26)
0402	121	85.1 (58)	7.4 (71)	127	32.3 (52)	108	29.6 (20)	101	92.1 (29)
0403	47	*	*	50	*	41	*	38	*
0404	47	*	*	48	*	43	*	38	*
0501	150	94.7 (2)	0.7 (3)	162	32.1 (56)	142	33.8 (15)	116	87.1 (63)
0503	40	*	*	40	*	36	*	28	*
0504	160	93.1 (6)	4.4 (42)	165	24.2 (89)	149	36.9 (12)	120	88.3 (55)
0505	109	87.2 (41)	8.3 (80)	117	29.1 (75)	113	33.6 (16)	97	*
0506	65	*	*	66	*	62	*	57	*
0507	31	*	*	31	*	29	*	20	*
0508	8	*	*	9	*	8	*	7	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0509	2	*	*	2	*	1	*	1	*
0650	364	85.7 (55)	7.1 (70)	383	36.6 (32)	325	28.6 (22)	314	91.7 (35)
0651	183	74.3 (108)	15.3 (108)	195	40.5 (16)	170	17.6 (64)	166	88.0 (56)
0652	110	77.3 (97)	15.5 (109)	116	31.9 (58)	103	28.2 (23)	93	*
0653	140	76.4 (101)	18.6 (112)	147	42.9 (8)	134	26.9 (26)	120	90.0 (44)
0654	116	75.0 (105)	19.8 (113)	126	42.1 (9)	114	22.8 (42)	106	77.4 (103)
0655	117	87.2 (41)	6.0 (59)	124	46.8 (4)	113	23.0 (39)	105	87.6 (60)
0657	17	*	*	17	*	13	*	13	*
0701	139	79.9 (81)	12.9 (94)	142	30.3 (69)	132	40.2 (7)	114	89.5 (48)
0702	18	*	*	22	*	21	*	17	*
0703	61	*	*	72	*	64	*	55	*
0704	13	*	*	15	*	13	*	12	*
0705	40	*	*	42	*	38	*	31	*
0706	3	*	*	3	*	2	*	2	*
0850	268	89.2 (29)	3.0 (25)	294	32.0 (57)	265	19.6 (53)	227	92.1 (29)
0851	251	86.5 (48)	9.6 (84)	268	35.1 (42)	255	22.0 (43)	221	94.6 (7)
0901	504	73.8 (109)	20.6 (116)	538	41.4 (12)	500	14.0 (88)	456	82.9 (92)
0902	13	*	*	13	*	13	*	12	*
1051	411	78.6 (89)	15.1 (107)	431	29.7 (72)	390	23.3 (38)	309	86.1 (71)
1101	558	73.5 (111)	9.9 (87)	618	27.7 (79)	529	13.0 (96)	456	87.3 (62)
1102	239	72.4 (113)	13.4 (97)	272	32.7 (49)	230	20.0 (51)	199	96.0 (3)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1106	19	*	*	21	*	19	*	16	*
1250	298	80.9 (77)	9.7 (85)	374	34.8 (44)	336	25.3 (29)	303	83.2 (89)
1251	242	78.1 (92)	12.4 (93)	262	35.9 (36)	226	15.5 (81)	198	83.8 (84)
1252	183	74.9 (106)	13.1 (95)	197	37.6 (26)	182	13.7 (92)	163	91.4 (36)
1253	235	75.3 (104)	14.0 (102)	267	31.5 (61)	246	18.7 (57)	218	88.5 (53)
1350	745	89.3 (28)	4.6 (46)	838	35.9 (36)	565	18.2 (62)	648	79.5 (99)
1351	40	*	*	40	*	32	*	25	*
1352	119	89.9 (23)	5.0 (49)	124	41.1 (14)	114	1.8 (112)	104	93.3 (11)
1401	72	*	*	77	*	48	*	58	*
1402	39	*	*	42	*	35	*	24	*
1403	10	*	*	8	*	4	*	6	*
1501	231	89.6 (26)	5.6 (56)	262	35.9 (36)	161	23.0 (39)	197	84.8 (80)
1503	304	83.2 (67)	7.6 (74)	339	39.8 (19)	275	40.0 (8)	281	79.4 (100)
1504	632	91.3 (13)	2.7 (22)	703	31.9 (58)	495	26.5 (27)	473	72.1 (107)
1505	128	89.1 (30)	4.7 (48)	132	25.0 (87)	96	*	86	*
1506	171	74.9 (106)	21.1 (117)	185	36.8 (30)	160	9.4 (104)	145	93.8 (9)
1507	190	87.4 (40)	6.3 (60)	212	35.8 (39)	154	24.0 (37)	158	89.9 (45)
1508	22	*	*	29	*	22	*	23	*
1509	122	85.2 (57)	6.6 (64)	144	40.3 (18)	116	19.8 (52)	116	81.9 (94)
1511	250	91.6 (12)	3.6 (32)	260	32.3 (52)	196	18.4 (60)	195	86.7 (67)
1512	114	88.6 (33)	7.0 (69)	120	22.5 (94)	85	*	84	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1513	18	*	*	21	*	21	*	21	*
1514	72	*	*	74	*	59	*	41	*
1701	222	86.9 (43)	8.1 (78)	233	46.8 (4)	207	45.9 (5)	198	82.8 (93)
1950	184	78.8 (87)	15.8 (110)	195	39.5 (20)	165	19.4 (54)	150	90.7 (38)
1952	45	*	*	46	*	42	*	36	*
1953	103	85.4 (56)	13.6 (100)	112	39.3 (21)	101	21.8 (45)	87	*
1954	179	78.8 (87)	14.0 (102)	191	36.1 (35)	178	25.3 (29)	156	92.9 (15)
1955	33	*	*	33	*	26	*	23	*
1957	156	86.5 (48)	5.1 (50)	159	31.4 (62)	134	27.6 (24)	114	84.2 (82)
1963	10	*	*	11	*	6	*	1	*
1964	16	*	*	19	*	16	*	13	*
2150	197	92.4 (8)	1.0 (4)	215	44.2 (6)	201	24.4 (35)	181	92.3 (26)
2151	65	*	*	79	*	59	*	56	*
2251	138	88.4 (34)	8.0 (77)	142	41.5 (11)	134	24.6 (33)	122	95.1 (6)
2252	57	*	*	64	*	56	*	57	*
2554	22	*	*	22	*	20	*	17	*
2555	2,372	85.8 (54)	4.5 (44)	2,521	23.7 (91)	2,130	11.0 (98)	1,856	89.4 (50)
2556	113	77.9 (94)	4.4 (42)	115	10.4 (120)	89	*	36	*
2557	5	*	*	7	*	5	*	6	*
2558	1	*	*	2	*	2	*	2	*
2559	3	*	*	3	*	3	*	3	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2560	2	*	*	2	*	2	*	2	*
2562	1	*	*	2	*	2	*	2	*
2563	5	*	*	3	*	0	*	0	*
3001	354	72.0 (115)	7.9 (76)	383	31.3 (64)	328	9.8 (101)	302	77.8 (102)
3002	88	*	*	91	*	81	*	76	*
3004	53	*	*	60	*	58	*	53	*
3007	5	*	*	7	*	6	*	6	*
3009	0	*	*	3	*	2	*	2	*
3201	130	80.0 (80)	13.1 (95)	136	37.5 (27)	119	38.7 (10)	99	*
3301	1,313	88.0 (37)	5.9 (58)	1,411	30.2 (71)	1,257	6.0 (109)	1,067	93.0 (14)
3304	32	*	*	33	*	32	*	23	*
3305	48	*	*	49	*	47	*	37	*
3306	85	*	*	88	*	77	*	66	*
3308	25	*	*	26	*	23	*	16	*
3309	11	*	*	11	*	10	*	8	*
3312	112	83.9 (64)	5.4 (52)	123	29.3 (73)	105	10.5 (99)	98	*
3550	22	*	*	23	*	20	*	19	*
3551	8	*	*	9	*	9	*	8	*
3552	11	*	*	14	*	12	*	12	*
3701	326	93.6 (5)	2.5 (19)	356	31.7 (60)	322	21.1 (47)	274	96.7 (1)
3850	997	88.1 (36)	5.7 (57)	1,070	40.6 (15)	954	14.7 (85)	879	85.4 (74)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3901	419	80.9 (77)	6.7 (65)	447	32.2 (54)	391	10.0 (100)	337	76.3 (104)
3902	184	79.9 (81)	12.0 (91)	202	48.5 (3)	201	13.9 (89)	199	79.9 (98)
3903	8	*	*	8	*	7	*	0	*
3904	3	*	*	3	*	3	*	3	*
3905	2	*	*	2	*	1	*	0	*
3950	581	83.1 (68)	9.8 (86)	634	33.4 (46)	572	19.1 (55)	479	86.6 (70)
3954	2	*	*	2	*	2	*	0	*
4001	77	*	*	91	*	91	*	92	*
4002	48	*	*	58	*	55	*	58	*
4103	12	*	*	13	*	15	*	15	*
4106	815	86.6 (46)	5.5 (55)	878	28.5 (76)	772	8.4 (106)	700	87.7 (58)
4107	996	82.2 (73)	8.1 (78)	1,053	27.7 (79)	731	17.2 (70)	790	92.5 (21)
4109	347	82.4 (71)	13.5 (98)	379	31.4 (62)	346	17.3 (67)	280	91.8 (34)
4112	180	49.4 (119)	0.0 (1)	185	26.5 (84)	178	90.4 (1)	165	90.3 (43)
4113	112	90.2 (19)	0.0 (1)	116	37.1 (29)	116	12.9 (97)	116	91.4 (36)
4115	1,085	87.6 (38)	4.6 (46)	1,159	25.7 (86)	1,023	15.8 (78)	863	92.5 (21)
4351	188	89.4 (27)	3.2 (27)	199	36.2 (33)	62	*	171	92.4 (25)
4352	47	*	*	48	*	42	*	42	*
4353	129	73.6 (110)	20.2 (114)	143	41.3 (13)	129	16.3 (76)	116	75.9 (105)
4354	249	90.0 (21)	4.0 (39)	266	35.7 (40)	217	27.6 (24)	202	90.6 (39)
4355	107	94.4 (4)	2.8 (23)	111	32.4 (51)	98	*	86	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4356	132	86.4 (50)	6.8 (66)	144	31.3 (64)	126	37.3 (11)	115	96.5 (2)
4358	44	*	*	50	*	44	*	39	*
4359	142	70.4 (116)	20.4 (115)	153	41.8 (10)	135	21.5 (46)	118	89.8 (46)
4360	82	*	*	86	*	57	*	71	*
4361	74	*	*	76	*	69	*	61	*
4362	99	*	*	105	39.0 (23)	93	*	88	*
4363	272	82.7 (70)	10.3 (88)	289	40.5 (16)	225	34.2 (14)	234	86.8 (65)
4401	355	80.6 (79)	14.9 (106)	369	32.2 (54)	343	15.2 (83)	283	92.9 (15)
4501	341	77.7 (96)	14.7 (105)	351	36.2 (33)	309	25.2 (31)	266	92.9 (15)
4601	433	84.5 (62)	7.6 (74)	466	27.0 (82)	408	25.7 (28)	344	84.9 (79)
4602	9	*	*	9	*	7	*	7	*
4701	356	90.4 (17)	3.7 (33)	371	22.1 (95)	266	35.0 (13)	247	93.1 (13)
5050	532	90.0 (21)	3.2 (27)	566	17.7 (111)	392	16.8 (73)	399	93.2 (12)
5051	819	90.5 (15)	3.8 (35)	867	20.6 (100)	646	17.8 (63)	582	92.1 (29)
5052	952	88.7 (32)	3.4 (31)	1,004	22.1 (95)	714	14.3 (86)	732	92.5 (21)
5055	362	85.9 (53)	12.2 (92)	386	51.8 (2)	182	16.5 (74)	317	80.4 (97)
5101	566	90.6 (14)	3.9 (37)	611	25.9 (85)	473	20.9 (48)	443	92.8 (19)
5102	1	*	*	1	*	1	*	0	*
5202	264	87.5 (39)	6.8 (66)	289	34.9 (43)	218	30.3 (19)	215	94.0 (8)
5601	323	84.8 (59)	11.1 (89)	332	32.8 (48)	296	31.8 (17)	256	95.7 (4)
5801	517	77.2 (98)	13.5 (98)	569	32.5 (50)	518	18.9 (56)	442	85.3 (75)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5803	42	*	*	41	*	39	*	25	*
5804	0	*	*	1	*	1	*	1	*
5806	19	*	*	19	*	11	*	3	*
6101	11	*	*	10	*	9	*	6	*
6102	39	*	*	39	*	33	*	28	*
6103	1,250	86.2 (52)	6.9 (68)	1,350	39.3 (21)	1,177	20.7 (49)	1,071	79.4 (100)
6105	9	*	*	9	*	8	*	6	*
6106	17	*	*	17	*	11	*	2	*
6301	1,292	86.8 (44)	5.4 (52)	1,398	27.0 (82)	1,139	9.6 (102)	1,059	86.7 (67)
6312	633	86.3 (51)	4.3 (41)	688	23.4 (92)	541	13.3 (93)	506	84.4 (81)
6316	365	86.6 (46)	3.0 (25)	392	24.5 (88)	329	24.6 (33)	283	92.9 (15)
6320	39	*	*	38	*	13	*	24	*
6325	52	*	*	52	*	39	*	30	*
6330	4	*	*	4	*	2	*	4	*
6335	332	90.4 (17)	8.4 (81)	344	38.1 (25)	54	*	260	91.9 (33)
6340	16	*	*	16	*	7	*	10	*
6345	11	*	*	12	*	7	*	6	*
6360	206	88.8 (31)	5.3 (51)	218	21.1 (99)	104	31.7 (18)	161	83.2 (89)
7304	26	*	*	28	*	24	*	23	*
7306	776	76.9 (100)	11.1 (89)	822	30.8 (66)	705	17.3 (67)	600	86.8 (65)
7308	28	*	*	35	*	34	*	33	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7310	8	*	*	10	*	9	*	9	*
7401	705	67.9 (118)	22.4 (118)	766	36.8 (30)	639	17.2 (70)	613	86.9 (64)
7601	182	82.4 (71)	9.3 (83)	206	37.4 (28)	151	51.0 (4)	161	83.9 (83)
7850	354	77.1 (99)	15.8 (110)	379	38.5 (24)	348	13.2 (95)	303	90.4 (41)
7852	5	*	*	5	*	5	*	4	*
8101	623	79.0 (86)	2.2 (16)	647	17.8 (110)	397	17.4 (65)	415	86.7 (67)
8102	108	81.5 (75)	3.7 (33)	116	29.3 (73)	60	*	88	*
8105	472	83.3 (66)	3.8 (35)	507	20.3 (103)	445	17.3 (67)	350	92.3 (26)
8106	144	72.2 (114)	6.3 (60)	156	20.5 (101)	122	4.1 (110)	113	85.0 (78)
8150	8	*	*	8	*	8	*	7	*
8151	932	90.2 (19)	2.5 (19)	1,029	30.8 (66)	926	16.5 (74)	790	90.6 (39)
8152	22	*	*	23	*	21	*	20	*
8153	2	*	*	2	*	1	*	1	*
8155	4	*	*	5	*	4	*	3	*
8157	0	*	*	1	*	1	*	1	*
8201	822	91.7 (10)	2.1 (14)	881	19.3 (105)	627	29.2 (21)	584	88.0 (56)
8202	1,330	92.5 (7)	5.4 (52)	1,394	30.3 (69)	929	13.9 (89)	1,092	83.6 (88)
8203	122	91.8 (9)	1.6 (9)	128	15.6 (119)	109	17.4 (65)	83	*
8250	258	79.5 (85)	3.9 (37)	280	16.8 (114)	236	22.0 (43)	198	88.4 (54)
8252	44	*	*	48	*	33	*	25	*
8255	798	89.7 (25)	1.1 (6)	860	18.0 (109)	559	42.4 (6)	581	92.8 (19)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8257	702	95.7 (1)	1.6 (9)	750	30.8 (66)	692	7.4 (108)	578	83.7 (85)
8263	205	94.6 (3)	1.0 (4)	216	16.2 (117)	101	39.6 (9)	147	95.2 (5)
8264	169	91.7 (10)	1.2 (7)	185	20.5 (101)	147	15.6 (80)	123	93.5 (10)
8267	289	84.8 (59)	4.5 (44)	321	18.1 (108)	278	51.8 (3)	225	85.8 (72)
8901	361	82.0 (74)	2.8 (23)	384	22.7 (93)	279	13.3 (93)	294	83.0 (91)
8902	469	84.6 (61)	2.1 (14)	503	21.9 (97)	338	15.1 (84)	359	87.7 (58)
8903	466	78.1 (92)	2.6 (21)	504	21.4 (98)	367	15.3 (82)	369	81.6 (95)
8904	752	84.0 (63)	2.0 (12)	805	16.4 (116)	428	18.5 (59)	539	89.2 (51)
8905	494	90.5 (15)	2.0 (12)	520	28.3 (77)	364	20.1 (50)	375	87.5 (61)
8909	199	83.9 (64)	1.5 (8)	213	18.3 (107)	166	16.9 (72)	143	89.5 (48)
9045	612	72.5 (112)	13.9 (101)	654	33.8 (45)	118	65.3 (2)	523	85.1 (77)
9050	285	75.4 (103)	6.3 (60)	294	23.8 (90)	239	23.0 (39)	208	81.3 (96)
9051	498	68.7 (117)	26.3 (119)	548	55.5 (1)	508	13.8 (91)	482	68.9 (108)
9053	201	76.1 (102)	8.5 (82)	217	27.6 (81)	165	15.8 (78)	166	75.3 (106)
9055	7	*	*	7	*	6	*	5	*
9058	332	89.8 (24)	4.2 (40)	340	19.7 (104)	245	2.9 (111)	240	88.8 (52)
9066	294	79.9 (81)	2.4 (18)	348	16.7 (115)	166	16.3 (76)	221	83.7 (85)
9068	320	77.8 (95)	1.9 (11)	337	28.2 (78)	272	14.3 (86)	260	85.8 (72)
9069	941	79.8 (84)	2.3 (17)	976	16.0 (118)	564	8.9 (105)	625	92.0 (32)
9070	448	78.3 (91)	6.3 (60)	468	17.1 (113)	283	18.4 (60)	311	85.2 (76)
9072	554	78.5 (90)	3.2 (27)	578	18.7 (106)	473	8.2 (107)	424	83.7 (85)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9080	852	83.0 (69)	3.3 (30)	875	17.4 (112)	578	9.5 (103)	610	92.5 (21)
9082	57	*	*	64	*	59	*	43	*
9083	35	*	*	36	*	28	*	7	*
9084	12	*	*	12	*	4	*	0	*
9085	40	*	*	45	*	42	*	45	*
9086	30	*	*	36	*	36	*	36	*
Michigan	**50,875**	**84.3**	**6.8**	**54,591**	**29.4**	**42,983**	**19.1**	**40,941**	**40941.0**
National (Prior Year)	**84,995**	**68.3**	**8.9**	**96,065**	**21.0**	**70,195**	**23.0**	**71,805**	**88.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	40	*	42	*	19	*	32	*	42	*
0102	172	54.7 (117)	177	39.5 (119)	80	*	156	23.1 (106)	177	2.3 (53)
0103	123	41.5 (103)	125	28.0 (112)	45	*	117	19.7 (101)	125	6.4 (110)
0104	56	*	56	*	24	*	47	*	56	*
0105	5	*	7	*	1	*	5	*	7	*
0250	54	*	53	*	17	*	49	*	53	*
0251	23	*	28	*	11	*	21	*	28	*
0252	29	*	30	*	10	*	26	*	30	*
0253	44	*	45	*	21	*	42	*	45	*
0401	138	49.3 (114)	137	25.5 (106)	54	*	117	12.0 (67)	137	2.2 (50)
0402	109	48.6 (113)	120	27.5 (111)	46	*	101	12.9 (73)	121	3.3 (78)
0403	41	*	49	*	21	*	38	*	49	*
0404	43	*	47	*	17	*	38	*	47	*
0501	143	42.0 (107)	149	24.2 (100)	42	*	116	18.1 (92)	149	9.4 (118)
0503	37	*	40	*	14	*	28	*	40	*
0504	149	36.2 (84)	161	18.0 (70)	45	*	120	5.8 (17)	161	1.2 (20)
0505	113	37.2 (86)	108	18.5 (73)	39	*	97	*	108	2.8 (65)
0506	63	*	65	*	29	*	57	*	65	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0507	29	*	31	*	9	*	20	*	31	*
0508	8	*	8	*	3	*	7	*	8	*
0509	1	*	2	*	1	*	1	*	2	*
0650	353	38.2 (87)	364	21.4 (89)	111	45.9 (9)	314	14.0 (80)	365	3.8 (92)
0651	180	49.4 (115)	184	35.3 (118)	77	*	166	18.1 (92)	184	4.9 (100)
0652	103	49.5 (116)	112	25.9 (107)	43	*	93	*	112	6.3 (108)
0653	136	41.9 (105)	141	27.0 (109)	49	*	120	15.0 (86)	141	2.8 (65)
0654	114	46.5 (111)	115	30.4 (115)	44	*	106	27.4 (109)	115	6.1 (107)
0655	115	40.9 (102)	119	31.1 (116)	41	*	105	19.0 (97)	119	8.4 (116)
0657	13	*	17	*	5	*	13	*	17	*
0701	134	25.4 (46)	137	18.2 (71)	31	*	113	6.2 (21)	137	0.7 (5)
0702	21	*	18	*	7	*	17	*	18	*
0703	63	*	62	*	24	*	55	*	62	*
0704	13	*	13	*	4	*	12	*	13	*
0705	40	*	40	*	11	*	31	*	40	*
0706	2	*	3	*	2	*	2	*	3	*
0850	265	40.8 (101)	270	18.9 (75)	86	*	227	9.7 (51)	270	1.5 (28)
0851	255	38.8 (92)	250	19.2 (78)	79	*	221	7.7 (34)	250	1.6 (32)
0901	499	41.9 (105)	509	25.1 (103)	193	32.6 (21)	456	12.3 (69)	509	3.1 (74)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0902	13	*	13	*	4	*	12	*	13	*
1051	391	33.8 (74)	410	21.5 (91)	114	31.6 (23)	309	11.0 (62)	410	3.2 (76)
1101	534	24.9 (44)	575	12.2 (46)	105	43.8 (13)	456	12.1 (68)	575	3.8 (92)
1102	233	33.0 (72)	245	19.2 (78)	59	*	199	16.6 (90)	245	3.3 (78)
1106	19	*	19	*	3	*	16	*	19	*
1250	345	31.6 (67)	346	21.1 (87)	88	*	303	12.9 (73)	346	2.3 (53)
1251	228	43.4 (109)	243	28.8 (113)	80	*	198	19.2 (98)	243	7.0 (112)
1252	183	32.2 (70)	188	20.2 (83)	51	*	163	13.5 (78)	188	3.2 (76)
1253	247	36.4 (85)	244	18.9 (75)	69	*	218	12.4 (71)	244	2.9 (69)
1350	738	32.1 (69)	760	19.6 (80)	201	23.4 (30)	648	11.0 (62)	765	2.4 (57)
1351	33	*	40	*	10	*	25	*	40	*
1352	114	34.2 (76)	119	14.3 (57)	35	*	104	1.9 (3)	119	0.8 (8)
1401	67	*	73	*	28	*	59	*	73	*
1402	38	*	39	*	6	*	24	*	39	*
1403	8	*	10	*	4	*	6	*	10	*
1501	224	19.2 (21)	232	11.2 (40)	41	*	197	3.0 (8)	231	1.7 (37)
1503	305	27.2 (56)	314	16.2 (63)	72	*	281	3.9 (11)	314	2.9 (69)
1504	601	21.0 (28)	655	9.8 (32)	108	50.0 (4)	478	4.0 (12)	654	0.9 (12)
1505	115	26.1 (50)	128	13.3 (50)	30	*	87	*	129	3.1 (74)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1506	168	25.0 (45)	173	17.3 (68)	38	*	145	2.8 (7)	173	1.2 (20)
1507	184	34.2 (76)	190	15.8 (59)	58	*	158	5.7 (16)	191	1.6 (32)
1508	23	*	23	*	0	*	23	*	23	*
1509	130	27.7 (57)	124	16.1 (61)	32	*	117	7.7 (34)	124	2.4 (57)
1511	225	29.3 (60)	250	17.2 (66)	55	*	196	11.7 (65)	251	1.6 (32)
1512	100	30.0 (64)	112	18.8 (74)	31	*	85	*	113	1.8 (39)
1513	21	*	17	*	9	*	21	*	17	*
1514	70	*	71	*	6	*	40	*	71	*
1701	223	42.2 (108)	219	33.8 (117)	81	*	201	25.9 (108)	218	18.3 (119)
1950	169	46.2 (110)	188	29.3 (114)	68	*	150	20.0 (102)	188	5.3 (102)
1952	43	*	45	*	16	*	36	*	45	*
1953	101	39.6 (96)	104	17.3 (68)	31	*	87	*	104	2.9 (69)
1954	179	40.2 (99)	179	21.8 (94)	60	*	156	7.1 (32)	179	1.1 (16)
1955	27	*	33	*	9	*	23	*	33	*
1957	143	39.2 (94)	156	19.9 (81)	45	*	114	10.5 (58)	157	1.3 (26)
1963	6	*	10	*	0	*	1	*	10	*
1964	17	*	16	*	5	*	13	*	16	*
2150	201	46.8 (112)	197	24.4 (101)	75	*	181	14.4 (83)	197	1.0 (14)
2151	70	*	67	*	22	*	56	*	67	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2251	136	40.4 (100)	135	27.4 (110)	49	*	122	9.8 (54)	136	1.5 (28)
2252	60	*	53	*	25	*	57	*	53	*
2554	20	*	21	*	6	*	17	*	21	*
2555	2,200	29.1 (58)	2,396	16.1 (61)	539	36.4 (18)	1,860	8.6 (40)	2,397	2.2 (50)
2556	90	*	115	16.5 (65)	6	*	35	*	114	3.5 (85)
2557	6	*	6	*	0	*	6	*	6	*
2558	2	*	2	*	0	*	2	*	2	*
2559	3	*	3	*	1	*	3	*	3	*
2560	2	*	2	*	1	*	2	*	2	*
2562	2	*	1	*	0	*	2	*	1	*
2563	1	*	5	*	0	*	0	*	5	*
3001	329	34.3 (78)	355	18.3 (72)	96	*	302	11.9 (66)	355	3.7 (88)
3002	81	*	88	*	21	*	76	*	88	*
3004	59	*	54	*	9	*	53	*	54	*
3007	6	*	5	*	1	*	6	*	5	*
3009	2	*	1	*	0	*	2	*	1	*
3201	128	34.4 (79)	131	23.7 (97)	39	*	100	14.0 (80)	131	4.6 (97)
3301	1,263	17.2 (15)	1,327	11.5 (42)	170	28.8 (27)	1,067	6.2 (21)	1,328	4.1 (95)
3304	32	*	32	*	0	*	23	*	32	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3305	47	*	48	*	4	*	37	*	48	*
3306	77	*	87	*	19	*	66	*	87	*
3308	23	*	25	*	2	*	16	*	25	*
3309	10	*	11	*	5	*	8	*	11	*
3312	105	17.1 (14)	114	20.2 (83)	16	*	98	*	114	3.5 (85)
3550	20	*	22	*	12	*	19	*	22	*
3551	9	*	8	*	4	*	8	*	8	*
3552	12	*	11	*	4	*	12	*	11	*
3701	322	38.2 (87)	328	20.7 (86)	103	30.1 (26)	274	9.1 (49)	328	2.1 (45)
3850	982	38.2 (87)	1,007	25.9 (107)	321	26.5 (29)	879	10.7 (61)	1,007	2.3 (53)
3901	396	25.5 (47)	421	14.0 (55)	79	*	337	8.3 (38)	421	1.2 (20)
3902	202	22.3 (34)	188	14.9 (58)	41	*	199	6.5 (26)	188	2.1 (45)
3903	7	*	8	*	0	*	0	*	8	*
3904	3	*	3	*	0	*	3	*	3	*
3905	1	*	2	*	0	*	0	*	2	*
3950	572	25.5 (47)	588	14.1 (56)	113	34.5 (19)	479	8.6 (40)	590	1.2 (20)
3954	2	*	2	*	0	*	0	*	2	*
4001	92	*	79	*	24	*	92	*	79	*
4002	58	*	48	*	13	*	58	*	48	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4103	15	*	12	*	3	*	15	*	12	*
4106	777	22.3 (34)	821	10.6 (36)	147	36.7 (17)	699	3.7 (10)	821	1.2 (20)
4107	932	21.2 (29)	1,001	10.6 (36)	144	43.8 (13)	790	8.6 (40)	1,001	2.5 (60)
4109	345	29.9 (63)	351	13.7 (53)	80	*	279	6.5 (26)	351	1.1 (16)
4112	177	1.7 (1)	180	1.1 (1)	3	*	164	7.3 (33)	180	3.3 (78)
4113	116	3.4 (3)	112	3.6 (7)	4	*	116	0.0 (1)	112	0.0 (1)
4115	1,036	17.9 (18)	1,088	7.3 (20)	148	47.3 (8)	864	6.0 (19)	1,090	1.9 (40)
4351	189	41.8 (104)	188	20.2 (83)	68	*	171	17.0 (91)	188	1.6 (32)
4352	43	*	47	*	27	*	42	*	47	*
4353	129	31.8 (68)	135	20.0 (82)	33	*	116	8.6 (40)	135	0.7 (5)
4354	239	38.9 (93)	250	24.0 (99)	77	*	202	15.8 (87)	250	4.0 (94)
4355	100	34.0 (75)	108	21.3 (88)	32	*	86	*	108	2.8 (65)
4356	128	29.7 (62)	134	22.4 (96)	32	*	115	8.7 (45)	134	3.7 (88)
4358	44	*	46	*	5	*	39	*	47	*
4359	138	26.8 (53)	145	12.4 (47)	28	*	118	10.2 (57)	145	0.7 (5)
4360	77	*	82	*	17	*	71	*	82	*
4361	70	*	75	*	28	*	61	*	75	*
4362	97	*	99	*	47	*	88	*	99	*
4363	266	35.3 (83)	270	25.2 (105)	78	*	234	12.8 (72)	270	3.7 (88)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4401	343	32.4 (71)	354	11.9 (44)	85	*	283	2.5 (6)	353	0.6 (3)
4501	313	35.1 (82)	342	21.6 (93)	93	*	266	13.2 (76)	343	2.6 (62)
4601	412	38.6 (90)	437	17.2 (66)	125	48.8 (6)	344	9.0 (48)	437	3.7 (88)
4602	7	*	9	*	5	*	7	*	9	*
4701	338	34.6 (80)	358	19.0 (77)	89	*	248	6.9 (30)	358	3.4 (82)
5050	499	26.9 (54)	539	13.5 (51)	102	48.0 (7)	400	9.8 (54)	540	2.6 (62)
5051	768	23.3 (37)	827	9.4 (30)	138	52.9 (3)	583	8.4 (39)	827	1.9 (40)
5052	898	22.0 (33)	953	12.5 (48)	146	40.4 (16)	732	8.7 (45)	954	1.9 (40)
5055	363	4.1 (4)	372	1.3 (2)	9	*	329	3.6 (9)	372	1.1 (16)
5101	534	24.2 (40)	570	10.4 (35)	101	43.6 (15)	444	6.3 (23)	569	1.9 (40)
5102	1	*	1	*	0	*	0	*	1	*
5202	265	40.0 (97)	267	25.1 (103)	86	*	216	9.7 (51)	266	1.9 (40)
5601	296	31.1 (66)	321	21.5 (91)	77	*	256	8.6 (40)	321	2.5 (60)
5801	520	39.2 (94)	518	24.9 (102)	153	30.7 (25)	442	13.8 (79)	519	1.7 (37)
5803	39	*	42	*	7	*	25	*	42	*
5804	1	*	0	*	0	*	1	*	0	*
5806	11	*	19	*	2	*	3	*	19	*
6101	9	*	11	*	3	*	6	*	11	*
6102	33	*	39	*	12	*	28	*	39	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6103	1,217	33.2 (73)	1,269	21.4 (89)	345	33.0 (20)	1,072	13.3 (77)	1,274	5.8 (105)
6105	8	*	9	*	0	*	6	*	9	*
6106	12	*	17	*	0	*	2	*	17	*
6301	1,269	17.3 (16)	1,310	7.6 (22)	167	44.3 (12)	1,060	5.0 (15)	1,310	0.8 (8)
6312	619	5.8 (5)	659	2.7 (5)	27	*	506	2.0 (5)	659	0.9 (12)
6316	376	19.9 (23)	369	7.9 (23)	52	*	284	4.2 (13)	369	1.1 (16)
6320	36	*	39	*	1	*	24	*	39	*
6325	45	*	53	*	5	*	30	*	53	*
6330	4	*	4	*	2	*	4	*	4	*
6335	331	6.0 (6)	334	3.3 (6)	16	*	261	1.1 (2)	334	0.6 (3)
6340	13	*	16	*	3	*	10	*	16	*
6345	11	*	12	*	2	*	6	*	12	*
6360	189	21.2 (29)	208	11.5 (42)	35	*	161	8.1 (37)	208	2.4 (57)
7304	24	*	26	*	13	*	23	*	27	*
7306	707	34.9 (81)	787	13.5 (51)	201	49.3 (5)	600	13.0 (75)	788	2.3 (53)
7308	34	*	29	*	7	*	33	*	29	*
7310	10	*	7	*	2	*	9	*	7	*
7401	678	40.1 (98)	707	23.8 (98)	225	32.4 (22)	613	19.6 (100)	707	5.5 (104)
7601	181	29.3 (60)	188	16.0 (60)	45	*	163	6.7 (28)	188	0.5 (2)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7850	350	38.6 (90)	358	22.3 (95)	118	31.4 (24)	303	18.5 (95)	357	3.4 (82)
7852	5	*	5	*	3	*	4	*	5	*
8101	534	20.0 (25)	624	7.5 (21)	78	*	414	14.7 (84)	623	5.1 (101)
8102	96	*	112	6.3 (16)	15	*	88	*	112	3.6 (87)
8105	451	20.0 (25)	472	9.1 (29)	60	*	350	9.7 (51)	472	3.0 (73)
8106	130	13.8 (11)	147	8.8 (27)	18	*	113	8.0 (36)	146	2.1 (45)
8150	8	*	8	*	1	*	7	*	8	*
8151	930	23.7 (39)	940	11.3 (41)	159	45.3 (10)	791	7.0 (31)	939	1.6 (32)
8152	21	*	21	*	5	*	20	*	22	*
8153	1	*	2	*	0	*	1	*	2	*
8155	4	*	4	*	1	*	3	*	4	*
8157	1	*	0	*	0	*	1	*	0	*
8201	772	18.7 (19)	826	11.9 (44)	108	28.7 (28)	583	9.9 (56)	827	2.2 (50)
8202	1,296	6.5 (7)	1,347	1.6 (3)	70	*	1,103	6.3 (23)	1,346	2.1 (45)
8203	116	31.0 (65)	122	16.4 (64)	25	*	83	*	122	0.8 (8)
8250	243	24.7 (43)	259	10.0 (33)	39	*	198	10.6 (59)	259	2.7 (64)
8252	35	*	44	*	7	*	25	*	44	*
8255	746	29.1 (58)	806	13.9 (54)	159	45.3 (10)	584	9.6 (50)	806	1.5 (28)
8257	699	2.4 (2)	711	1.7 (4)	14	*	580	1.9 (3)	711	0.8 (8)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8263	195	23.6 (38)	205	10.7 (39)	35	*	147	6.1 (20)	206	1.5 (28)
8264	161	24.2 (40)	170	10.6 (36)	26	*	123	10.6 (59)	170	1.2 (20)
8267	286	26.9 (54)	290	13.1 (49)	56	*	223	6.7 (28)	290	1.0 (14)
8901	326	26.1 (50)	365	7.9 (23)	77	*	293	19.5 (99)	365	6.3 (108)
8902	422	19.0 (20)	471	6.6 (18)	66	*	360	14.7 (84)	472	7.2 (113)
8903	441	25.6 (49)	472	8.7 (26)	89	*	369	21.4 (104)	471	7.9 (115)
8904	668	22.9 (36)	763	7.9 (23)	116	67.2 (1)	539	16.1 (88)	763	6.9 (111)
8905	459	13.9 (12)	495	6.5 (17)	48	*	375	6.4 (25)	496	1.4 (27)
8909	187	20.9 (27)	203	4.4 (11)	36	*	142	18.3 (94)	203	7.4 (114)
9045	593	12.1 (10)	619	4.0 (8)	58	*	527	8.9 (47)	619	2.9 (69)
9050	244	26.2 (52)	291	4.5 (12)	57	*	208	16.3 (89)	291	4.5 (96)
9051	521	9.8 (8)	517	4.3 (10)	42	*	483	5.8 (17)	518	2.1 (45)
9053	193	21.2 (29)	201	5.5 (13)	36	*	166	11.4 (64)	202	5.4 (103)
9055	6	*	7	*	1	*	5	*	7	*
9058	288	11.8 (9)	332	6.0 (15)	25	*	240	4.6 (14)	333	3.3 (78)
9066	273	24.2 (40)	326	9.5 (31)	56	*	223	18.8 (96)	327	3.4 (82)
9068	292	19.9 (23)	321	10.0 (33)	52	*	260	14.2 (82)	321	2.8 (65)
9069	793	15.9 (13)	941	6.7 (19)	95	*	625	21.0 (103)	941	9.1 (117)
9070	399	21.3 (32)	454	8.8 (27)	63	*	311	23.8 (107)	453	6.0 (106)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9072	487	19.7 (22)	564	4.1 (9)	83	*	424	22.9 (105)	564	4.6 (97)
9080	737	17.6 (17)	852	5.6 (14)	107	58.9 (2)	609	12.3 (69)	853	4.8 (99)
9082	61	*	57	*	8	*	43	*	57	*
9083	28	*	35	*	0	*	7	*	35	*
9084	7	*	12	*	0	*	0	*	12	*
9085	45	*	42	*	10	*	45	*	42	*
9086	36	*	30	*	0	*	36	*	30	*
Michigan	**48,601**	**26.1**	**51,441**	**13.9**	**10,331**	**38.9**	**40,995**	**10.4**	**51,462**	**3.0**
National (Prior Year)	**84,358**	**26.3**	**90,718**	**13.4**	**19,249**	**38.5**	**75,977**	**9.4**	**91,112**	**3.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	39	*	*	40	*	38	*	39	*
0102	172	7.6 (31)	8.1 (68)	174	9.8 (35)	155	3.2 (31)	172	69.2 (68)
0103	118	8.5 (50)	14.4 (114)	117	8.5 (17)	105	1.0 (3)	118	62.7 (84)
0104	53	*	*	53	*	49	*	53	*
0105	7	*	*	7	*	7	*	7	*
0250	53	*	*	52	*	47	*	53	*
0251	27	*	*	28	*	25	*	27	*
0252	30	*	*	30	*	28	*	30	*
0253	41	*	*	42	*	35	*	41	*
0401	133	6.0 (10)	9.8 (96)	135	8.1 (12)	123	1.6 (6)	133	72.2 (62)
0402	109	10.1 (71)	9.2 (90)	113	9.7 (33)	97	*	109	82.6 (13)
0403	46	*	*	46	*	38	*	46	*
0404	47	*	*	46	*	41	*	47	*
0501	114	8.8 (55)	6.1 (39)	145	6.2 (2)	106	3.8 (45)	141	83.7 (9)
0503	37	*	*	39	*	35	*	37	*
0504	151	9.9 (67)	10.6 (102)	157	8.3 (16)	138	5.8 (88)	154	82.5 (15)
0505	94	*	*	105	5.7 (1)	89	*	103	84.5 (6)
0506	57	*	*	59	*	49	*	62	*
0507	23	*	*	29	*	20	*	29	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	7	*	*	7	*	6	*	7	*
0509	2	*	*	2	*	2	*	2	*
0650	339	7.7 (34)	8.8 (80)	352	8.0 (10)	313	4.8 (76)	340	75.0 (53)
0651	175	6.9 (18)	11.4 (108)	177	9.0 (21)	158	3.8 (45)	175	70.3 (67)
0652	102	8.8 (55)	8.8 (80)	106	9.4 (29)	92	*	102	67.6 (73)
0653	130	7.7 (34)	10.8 (103)	137	10.2 (39)	116	3.4 (37)	130	82.3 (16)
0654	109	6.4 (11)	12.8 (113)	111	8.1 (12)	99	*	109	76.1 (48)
0655	115	8.7 (52)	9.6 (94)	116	7.8 (6)	107	4.7 (74)	115	75.7 (50)
0657	15	*	*	15	*	12	*	15	*
0701	124	4.0 (3)	8.9 (82)	130	6.9 (5)	110	1.8 (8)	126	88.1 (1)
0702	18	*	*	18	*	14	*	18	*
0703	60	*	*	60	*	54	*	60	*
0704	13	*	*	13	*	13	*	13	*
0705	36	*	*	37	*	33	*	35	*
0706	2	*	*	2	*	2	*	2	*
0850	256	5.5 (7)	10.2 (98)	268	9.0 (21)	237	1.7 (7)	263	84.4 (7)
0851	239	7.1 (23)	10.9 (104)	246	8.1 (12)	221	4.5 (67)	241	79.3 (29)
0901	482	7.1 (23)	9.1 (86)	498	9.4 (29)	438	2.3 (14)	482	72.4 (60)
0902	12	*	*	13	*	12	*	12	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1051	395	8.4 (48)	11.9 (111)	402	9.5 (31)	356	2.2 (10)	395	85.3 (3)
1101	541	8.7 (52)	7.2 (52)	561	9.3 (26)	490	4.3 (63)	541	66.4 (75)
1102	225	7.1 (23)	8.4 (75)	234	6.8 (4)	208	4.3 (63)	225	76.4 (45)
1106	19	*	*	19	*	17	*	19	*
1250	324	7.4 (28)	5.6 (31)	332	12.3 (68)	281	2.1 (9)	328	78.7 (34)
1251	225	5.8 (8)	6.2 (40)	238	10.5 (43)	200	3.0 (25)	225	79.6 (25)
1252	170	6.5 (12)	4.7 (21)	177	10.7 (48)	152	1.3 (5)	171	80.7 (18)
1253	230	7.8 (37)	9.1 (86)	236	9.3 (26)	209	3.8 (45)	230	85.7 (2)
1350	730	7.9 (40)	7.8 (63)	745	10.2 (39)	650	3.2 (31)	732	72.7 (58)
1351	38	*	*	38	*	33	*	38	*
1352	116	10.3 (77)	7.8 (63)	117	12.0 (63)	102	4.9 (79)	116	62.1 (87)
1401	70	*	*	72	*	59	*	70	*
1402	37	*	*	39	*	33	*	37	*
1403	10	*	*	9	*	9	*	10	*
1501	218	8.7 (52)	12.4 (112)	224	10.7 (48)	192	4.2 (59)	221	64.3 (81)
1503	290	6.9 (18)	7.2 (52)	295	13.6 (85)	247	2.4 (16)	290	78.3 (37)
1504	620	7.6 (31)	8.2 (70)	645	11.2 (53)	551	2.9 (22)	627	79.1 (31)
1505	125	12.8 (99)	11.2 (107)	128	7.8 (6)	115	8.7 (106)	126	80.2 (21)
1506	159	6.9 (18)	7.5 (56)	165	11.5 (55)	139	2.2 (10)	160	68.8 (70)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1507	175	13.1 (100)	14.9 (115)	185	16.2 (105)	147	3.4 (37)	176	79.0 (32)
1508	23	*	*	22	*	19	*	23	*
1509	121	5.8 (8)	6.6 (46)	119	10.1 (38)	106	2.8 (21)	122	72.1 (64)
1511	240	10.4 (78)	8.3 (72)	244	12.7 (75)	207	4.8 (76)	241	78.8 (33)
1512	107	9.3 (60)	10.3 (99)	112	18.8 (113)	86	*	107	85.0 (4)
1513	14	*	*	16	*	14	*	15	*
1514	70	*	*	70	*	60	*	70	*
1701	207	4.3 (4)	10.1 (97)	207	9.7 (33)	180	1.1 (4)	208	73.6 (57)
1950	180	6.7 (15)	8.9 (82)	182	9.3 (26)	160	2.5 (17)	181	77.9 (41)
1952	42	*	*	42	*	41	*	42	*
1953	95	*	*	100	13.0 (79)	84	*	95	*
1954	170	9.4 (61)	8.2 (70)	175	14.3 (92)	146	3.4 (37)	170	79.4 (28)
1955	30	*	*	32	*	27	*	30	*
1957	153	8.5 (50)	8.5 (76)	152	12.5 (71)	132	3.0 (25)	153	83.7 (9)
1963	10	*	*	10	*	6	*	10	*
1964	16	*	*	16	*	15	*	16	*
2150	181	6.6 (13)	11.0 (106)	186	9.1 (23)	164	3.0 (25)	181	82.9 (12)
2151	66	*	*	65	*	58	*	66	*
2251	130	6.9 (18)	8.5 (76)	132	9.1 (23)	116	2.6 (19)	130	72.3 (61)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2252	46	*	*	48	*	43	*	46	*
2554	21	*	*	21	*	19	*	21	*
2555	2,327	12.3 (96)	5.6 (31)	2,346	13.6 (85)	1,984	5.8 (88)	2,330	58.8 (92)
2556	112	20.5 (117)	2.7 (5)	111	19.8 (118)	86	*	112	50.9 (110)
2557	5	*	*	6	*	5	*	6	*
2558	2	*	*	2	*	2	*	2	*
2559	3	*	*	3	*	2	*	3	*
2560	2	*	*	2	*	2	*	2	*
2562	1	*	*	1	*	1	*	1	*
2563	4	*	*	3	*	2	*	4	*
3001	334	12.6 (97)	5.1 (24)	338	13.3 (83)	289	4.5 (67)	334	57.8 (95)
3002	86	*	*	88	*	73	*	86	*
3004	51	*	*	52	*	44	*	51	*
3007	4	*	*	4	*	4	*	4	*
3009	1	*	*	1	*	1	*	1	*
3201	121	6.6 (13)	16.5 (117)	125	8.0 (10)	112	3.6 (42)	121	80.2 (21)
3301	1,254	10.7 (85)	6.9 (48)	1,296	13.4 (84)	1,086	5.2 (83)	1,260	78.2 (39)
3304	29	*	*	30	*	25	*	30	*
3305	43	*	*	47	*	40	*	44	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3306	80	*	*	82	*	72	*	80	*
3308	22	*	*	22	*	21	*	22	*
3309	11	*	*	10	*	8	*	11	*
3312	109	11.0 (89)	7.3 (54)	113	15.0 (96)	93	*	109	65.1 (78)
3550	21	*	*	21	*	19	*	20	*
3551	6	*	*	8	*	6	*	6	*
3552	10	*	*	11	*	8	*	10	*
3701	316	9.2 (58)	5.4 (28)	321	11.8 (60)	279	2.2 (10)	317	84.2 (8)
3850	945	9.7 (64)	7.6 (58)	973	12.0 (63)	828	3.9 (49)	946	74.6 (54)
3901	392	9.9 (67)	7.4 (55)	404	13.6 (85)	336	3.9 (49)	398	70.9 (66)
3902	174	7.5 (30)	10.9 (104)	175	13.1 (80)	151	4.0 (56)	175	69.1 (69)
3903	7	*	*	8	*	7	*	7	*
3904	2	*	*	2	*	2	*	2	*
3905	2	*	*	2	*	2	*	2	*
3950	550	10.5 (82)	10.5 (101)	570	12.8 (76)	476	2.9 (22)	554	75.6 (51)
3954	2	*	*	2	*	2	*	2	*
4001	76	*	*	77	*	68	*	76	*
4002	46	*	*	48	*	40	*	46	*
4103	11	*	*	10	*	8	*	11	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4106	761	13.1 (100)	5.1 (24)	795	13.1 (80)	658	6.7 (96)	771	79.6 (25)
4107	943	11.6 (92)	9.1 (86)	979	14.6 (94)	803	4.4 (66)	944	76.4 (45)
4109	343	7.3 (26)	8.7 (79)	345	11.3 (54)	303	3.0 (25)	343	78.7 (34)
4112	175	3.4 (1)	6.3 (42)	172	9.9 (36)	155	0.6 (1)	175	22.9 (118)
4113	105	9.5 (62)	9.5 (93)	111	16.2 (105)	88	*	106	83.0 (11)
4115	1,042	10.6 (83)	7.1 (51)	1,065	12.6 (73)	906	4.1 (58)	1,042	80.3 (20)
4351	176	6.8 (16)	5.7 (33)	181	10.5 (43)	157	3.2 (31)	176	74.4 (55)
4352	42	*	*	43	*	35	*	42	*
4353	131	8.4 (48)	7.6 (58)	129	7.8 (6)	119	4.2 (59)	130	79.2 (30)
4354	230	8.3 (45)	7.8 (63)	236	14.8 (95)	192	3.1 (29)	230	79.6 (25)
4355	104	3.8 (2)	16.3 (116)	105	6.7 (3)	97	*	104	77.9 (41)
4356	128	10.2 (75)	3.9 (13)	132	18.9 (114)	104	2.9 (22)	128	78.1 (40)
4358	44	*	*	44	*	41	*	44	*
4359	138	8.0 (41)	8.0 (66)	140	15.0 (96)	118	3.4 (37)	138	79.7 (24)
4360	71	*	*	80	*	66	*	71	*
4361	65	*	*	74	*	53	*	65	*
4362	92	*	*	92	*	80	*	92	*
4363	257	8.2 (44)	8.6 (78)	261	11.5 (55)	224	3.1 (29)	257	79.8 (23)
4401	337	8.3 (45)	7.7 (60)	340	10.6 (45)	295	2.7 (20)	337	78.6 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4501	320	7.8 (37)	11.6 (109)	332	9.9 (36)	285	3.2 (31)	320	72.5 (59)
4601	415	5.3 (6)	9.2 (90)	424	8.5 (17)	374	3.7 (43)	417	67.9 (72)
4602	8	*	*	9	*	8	*	8	*
4701	346	10.1 (71)	11.6 (109)	351	12.0 (63)	305	4.3 (63)	346	78.3 (37)
5050	526	10.6 (83)	5.7 (33)	522	16.5 (108)	431	3.9 (49)	527	60.3 (90)
5051	805	10.4 (78)	6.7 (47)	809	12.2 (66)	697	3.7 (43)	806	67.5 (74)
5052	925	10.4 (78)	4.4 (20)	932	12.3 (68)	797	5.1 (82)	929	62.4 (85)
5055	326	7.4 (28)	2.1 (2)	342	8.2 (15)	298	4.7 (74)	335	38.5 (116)
5101	544	9.2 (58)	5.7 (33)	557	12.6 (73)	472	4.2 (59)	547	72.2 (62)
5102	1	*	*	1	*	1	*	1	*
5202	258	10.9 (88)	6.2 (40)	263	12.9 (77)	224	4.9 (79)	258	82.6 (13)
5601	303	6.9 (18)	8.3 (72)	314	11.1 (52)	267	2.2 (10)	303	76.9 (44)
5801	496	7.3 (26)	6.3 (42)	506	11.7 (58)	433	2.3 (14)	496	65.1 (78)
5803	38	*	*	40	*	33	*	38	*
5804	0	*	*	0	*	0	*	0	*
5806	18	*	*	19	*	16	*	18	*
6101	11	*	*	10	*	9	*	11	*
6102	38	*	*	38	*	35	*	38	*
6103	1,206	11.2 (91)	5.4 (28)	1,218	15.8 (102)	991	4.5 (67)	1,207	64.2 (82)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6105	9	*	*	9	*	8	*	9	*
6106	17	*	*	17	*	10	*	17	*
6301	1,247	11.6 (92)	5.8 (37)	1,274	12.9 (77)	1,082	5.7 (87)	1,253	66.3 (76)
6312	631	10.1 (71)	5.4 (28)	640	10.6 (45)	554	4.0 (56)	635	76.1 (48)
6316	361	8.3 (45)	6.9 (48)	362	12.2 (66)	314	2.5 (17)	361	77.6 (43)
6320	36	*	*	37	*	35	*	36	*
6325	52	*	*	52	*	50	*	51	*
6330	4	*	*	4	*	4	*	4	*
6335	316	9.8 (65)	4.1 (17)	318	8.5 (17)	285	5.3 (85)	317	76.3 (47)
6340	16	*	*	16	*	14	*	16	*
6345	12	*	*	12	*	9	*	12	*
6360	196	12.2 (95)	7.7 (60)	198	13.1 (80)	164	6.1 (91)	199	64.8 (80)
7304	26	*	*	26	*	25	*	26	*
7306	766	10.2 (75)	5.9 (38)	766	10.2 (39)	678	4.6 (71)	767	60.9 (88)
7308	29	*	*	29	*	27	*	29	*
7310	7	*	*	7	*	7	*	7	*
7401	673	7.6 (31)	8.9 (82)	687	10.8 (50)	598	3.3 (35)	673	71.6 (65)
7601	184	4.9 (5)	10.3 (99)	188	11.7 (58)	162	0.6 (1)	184	54.9 (101)
7850	338	8.0 (41)	8.0 (66)	343	7.9 (9)	311	3.5 (41)	338	75.4 (52)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7852	5	*	*	5	*	5	*	5	*
8101	563	15.1 (111)	4.3 (19)	607	18.9 (114)	450	6.7 (96)	586	47.1 (112)
8102	103	6.8 (16)	9.7 (95)	103	20.4 (119)	77	*	104	51.0 (109)
8105	460	12.0 (94)	5.2 (26)	464	11.6 (57)	404	4.2 (59)	460	63.3 (83)
8106	134	14.2 (108)	7.5 (56)	144	11.8 (60)	116	7.8 (104)	138	62.3 (86)
8150	6	*	*	5	*	5	*	6	*
8151	900	11.1 (90)	5.3 (27)	923	13.9 (90)	780	4.5 (67)	877	81.2 (17)
8152	22	*	*	22	*	20	*	22	*
8153	2	*	*	2	*	1	*	2	*
8155	4	*	*	4	*	4	*	4	*
8157	0	*	*	0	*	0	*	0	*
8201	805	10.4 (78)	7.0 (50)	813	13.9 (90)	691	3.9 (49)	805	60.6 (89)
8202	1,267	7.7 (34)	6.3 (42)	1,308	9.6 (32)	1,142	3.9 (49)	1,295	46.9 (113)
8203	119	12.6 (97)	9.2 (90)	120	12.5 (71)	104	3.8 (45)	119	84.9 (5)
8250	247	10.1 (71)	5.7 (33)	255	10.6 (45)	219	5.5 (86)	248	68.5 (71)
8252	44	*	*	43	*	37	*	44	*
8255	777	9.9 (67)	7.7 (60)	789	10.3 (42)	689	4.6 (71)	779	58.2 (94)
8257	633	10.7 (85)	3.9 (13)	685	11.8 (60)	554	6.3 (92)	656	80.5 (19)
8263	199	8.0 (41)	9.0 (85)	203	8.9 (20)	181	3.9 (49)	199	57.8 (95)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8264	156	9.6 (63)	8.3 (72)	165	9.1 (23)	142	6.3 (92)	157	54.1 (102)
8267	285	9.8 (65)	8.1 (68)	288	10.8 (50)	256	3.9 (49)	285	56.8 (99)
8901	342	14.0 (106)	1.8 (1)	352	17.3 (110)	274	7.7 (103)	355	58.6 (93)
8902	416	13.9 (105)	2.2 (4)	462	16.2 (105)	343	5.0 (81)	441	59.2 (91)
8903	430	17.2 (114)	2.1 (2)	461	19.1 (116)	342	8.8 (107)	444	52.9 (106)
8904	681	14.1 (107)	3.2 (8)	743	17.5 (111)	553	6.3 (92)	722	55.5 (100)
8905	452	10.8 (87)	6.4 (45)	477	13.8 (89)	385	5.2 (83)	469	65.7 (77)
8909	188	18.1 (116)	3.2 (8)	197	19.3 (117)	147	7.5 (100)	194	54.1 (102)
9045	577	9.0 (57)	4.9 (22)	595	14.3 (92)	486	3.3 (35)	585	57.4 (97)
9050	272	13.6 (102)	3.7 (12)	275	16.0 (103)	224	7.6 (102)	276	44.9 (115)
9051	462	7.8 (37)	9.1 (86)	472	12.3 (68)	395	4.6 (71)	472	73.7 (56)
9053	191	9.9 (67)	4.2 (18)	199	13.6 (85)	165	4.8 (76)	192	53.1 (105)
9055	7	*	*	7	*	7	*	7	*
9058	303	17.2 (114)	4.0 (16)	322	16.1 (104)	254	9.8 (108)	313	56.9 (98)
9066	307	16.0 (113)	4.9 (22)	326	17.5 (111)	252	7.5 (100)	311	52.1 (108)
9068	307	15.3 (112)	3.3 (10)	317	16.7 (109)	258	8.5 (105)	307	46.6 (114)
9069	891	14.9 (110)	3.1 (7)	921	15.5 (100)	746	6.0 (90)	892	35.1 (117)
9070	436	13.8 (104)	3.9 (13)	448	15.0 (96)	366	6.8 (98)	436	49.8 (111)
9072	526	14.4 (109)	3.0 (6)	555	15.3 (99)	446	6.3 (92)	527	53.3 (104)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9080	818	13.6 (102)	3.4 (11)	832	15.7 (101)	678	6.8 (98)	822	52.8 (107)
9082	55	*	*	56	*	45	*	55	*
9083	33	*	*	35	*	24	*	33	*
9084	10	*	*	12	*	10	*	10	*
9085	42	*	*	41	*	34	*	42	*
9086	30	*	*	30	*	26	*	30	*
Michigan	**48,594**	**10.2**	**6.7**	**49,968**	**12.5**	**42,176**	**4.5**	**48,971**	**67.7**
National (Prior Year)	**91,478**	**9.5**	**6.8**	**88,038**	**11.4**	**75,818**	**4.1**	**92,044**	**59.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	117	94.0	0.0	1.7	0.0	0.0	4.3	0.0
0102	540	90.9	0.4	2.8	0.4	0.0	5.4	0.2
0103	455	91.4	0.4	3.3	0.2	0.0	4.0	0.7
0104	161	92.5	0.0	3.1	0.0	0.6	3.1	0.6
0105	7	*	*	*	*	*	*	*
0250	146	83.6	0.0	2.1	0.7	0.0	13.7	0.0
0251	95	*	*	*	*	*	*	*
0252	122	46.7	1.6	3.3	1.6	0.0	46.7	0.0
0253	175	78.3	0.6	0.6	5.1	0.0	15.4	0.0
0401	513	94.0	0.6	2.5	0.0	0.4	1.9	0.6
0402	386	83.7	1.0	2.6	0.0	0.8	11.9	0.0
0403	140	96.4	0.0	2.1	0.7	0.0	0.0	0.7
0404	148	91.2	1.4	1.4	0.0	0.7	5.4	0.0
0501	551	94.6	0.5	2.2	0.0	0.9	1.8	0.0
0503	118	93.2	0.0	3.4	0.0	0.0	2.5	0.8
0504	516	87.6	1.4	2.1	2.3	0.4	6.2	0.0
0505	351	82.3	1.1	5.4	0.3	0.3	10.5	0.0
0506	220	95.0	0.5	0.5	0.5	0.0	3.6	0.0
0507	85	*	*	*	*	*	*	*
0508	32	*	*	*	*	*	*	*
0509	8	*	*	*	*	*	*	*
0650	1,171	81.0	4.3	4.5	2.5	2.3	5.3	0.1
0651	625	91.5	0.2	4.2	0.2	0.2	2.7	1.1

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0652	313	95.2	0.0	1.9	0.3	0.3	1.9	0.3
0653	451	96.0	0.2	2.0	0.0	0.0	1.6	0.2
0654	386	93.5	1.3	2.1	0.8	0.0	1.8	0.5
0655	393	90.6	0.5	2.8	0.0	0.8	4.3	1.0
0657	66	*	*	*	*	*	*	*
0701	437	87.2	1.6	2.1	0.0	5.9	2.5	0.7
0702	68	*	*	*	*	*	*	*
0703	233	93.1	1.3	1.7	0.0	0.4	2.1	1.3
0704	54	*	*	*	*	*	*	*
0705	135	95.6	0.0	0.0	0.0	0.0	3.7	0.7
0706	5	*	*	*	*	*	*	*
0850	945	86.8	4.7	5.4	0.1	0.3	2.4	0.3
0851	804	93.9	0.4	3.7	0.1	0.6	1.1	0.1
0901	1,775	80.4	2.5	10.2	0.0	0.3	5.7	0.9
0902	46	*	*	*	*	*	*	*
1051	1,454	85.6	1.1	7.2	1.2	1.0	3.2	0.6
1101	1,890	37.1	50.3	7.6	0.2	0.7	2.8	1.3
1102	888	73.4	10.7	6.6	0.6	1.1	6.2	1.4
1106	79	*	*	*	*	*	*	*
1250	1,083	86.1	0.4	10.2	0.0	0.5	2.4	0.5
1251	882	92.4	0.3	2.6	0.0	0.2	2.9	1.5
1252	585	60.9	0.7	34.9	0.0	0.2	2.7	0.7
1253	900	79.4	5.3	9.4	0.1	0.2	4.3	1.1
1350	2,631	59.8	19.2	7.0	0.2	6.6	6.7	0.6

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1351	143	92.3	1.4	4.2	0.0	0.0	2.1	0.0
1352	383	65.5	24.3	6.0	0.3	0.0	3.1	0.8
1401	255	88.2	0.4	6.3	0.8	0.8	2.0	1.6
1402	147	59.9	1.4	23.1	6.8	2.0	4.8	2.0
1403	28	*	*	*	*	*	*	*
1404	1	*	*	*	*	*	*	*
1501	841	49.8	4.6	36.7	0.0	0.4	3.9	4.5
1503	1,146	61.4	6.6	23.6	0.3	0.7	6.3	1.0
1504	2,322	42.2	4.7	44.2	0.0	4.7	3.2	1.0
1505	477	85.5	2.9	6.9	0.0	0.6	2.7	1.3
1506	626	75.9	1.0	16.8	0.5	0.6	3.2	2.1
1507	659	86.6	1.1	5.2	0.0	1.7	5.0	0.5
1508	114	8.8	0.0	86.0	0.0	0.0	0.0	5.3
1509	485	51.1	2.3	40.6	0.2	0.2	1.6	3.9
1511	763	89.0	1.6	3.5	0.1	1.3	3.7	0.8
1512	386	90.4	0.5	2.8	0.5	0.8	4.7	0.3
1513	78	*	*	*	*	*	*	*
1514	170	87.1	0.0	6.5	0.0	0.6	4.7	1.2
1701	712	57.9	0.4	2.2	17.3	0.6	21.3	0.3
1950	625	93.3	0.3	3.8	0.3	0.2	1.8	0.3
1952	139	95.7	0.0	2.2	0.0	0.0	1.4	0.7
1953	332	91.6	0.6	5.4	0.0	0.0	2.1	0.3
1954	601	86.4	0.3	11.6	0.0	0.3	1.2	0.2
1955	89	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1957	567	83.6	0.5	13.2	0.0	0.2	2.5	0.0
1963	11	*	*	*	*	*	*	*
1964	34	*	*	*	*	*	*	*
2150	705	85.4	0.1	2.4	0.4	1.0	10.2	0.4
2151	256	79.3	1.2	7.4	0.8	0.0	7.4	3.9
2251	464	92.7	0.2	1.7	0.4	0.6	3.4	0.9
2252	210	92.9	0.5	2.4	0.0	0.0	2.9	1.4
2554	85	*	*	*	*	*	*	*
2555	8,410	47.8	39.9	4.9	0.1	0.4	6.4	0.5
2556	204	34.3	51.5	5.9	0.0	0.5	7.4	0.5
2557	9	*	*	*	*	*	*	*
2558	4	*	*	*	*	*	*	*
2559	9	*	*	*	*	*	*	*
2560	2	*	*	*	*	*	*	*
2562	2	*	*	*	*	*	*	*
2563	5	*	*	*	*	*	*	*
3001	1,176	16.2	66.2	13.4	0.1	0.8	3.0	0.3
3002	264	21.2	50.0	24.2	0.0	0.0	4.5	0.0
3004	153	22.2	0.0	75.2	0.7	0.0	2.0	0.0
3007	30	*	*	*	*	*	*	*
3009	10	*	*	*	*	*	*	*
3201	502	93.6	0.2	4.0	0.4	0.0	1.2	0.6
3301	4,437	39.9	29.3	15.7	0.2	6.2	8.2	0.6
3304	112	42.9	15.2	5.4	0.0	33.0	3.6	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3305	148	65.5	12.8	9.5	0.0	6.1	5.4	0.7
3306	263	84.4	4.2	6.5	0.4	0.0	3.4	1.1
3308	76	*	*	*	*	*	*	*
3309	40	*	*	*	*	*	*	*
3312	422	38.9	26.8	24.9	0.7	1.4	6.9	0.5
3550	76	*	*	*	*	*	*	*
3551	29	*	*	*	*	*	*	*
3552	53	*	*	*	*	*	*	*
3701	1,142	90.2	0.1	6.8	0.0	0.2	2.7	0.0
3850	3,304	72.7	14.2	5.6	0.1	0.6	6.5	0.3
3901	580	33.1	52.8	6.4	0.3	1.0	4.8	1.6
3902	268	48.9	10.8	19.8	0.0	0.4	20.1	0.0
3903	10	*	*	*	*	*	*	*
3904	3	*	*	*	*	*	*	*
3905	2	*	*	*	*	*	*	*
3950	2,066	60.0	23.5	7.9	0.1	1.4	6.6	0.5
3954	5	*	*	*	*	*	*	*
4001	1,045	31.0	55.1	8.4	0.0	1.1	4.1	0.3
4002	720	51.5	25.4	15.1	0.1	1.0	6.5	0.3
4003	5	*	*	*	*	*	*	*
4004	5	*	*	*	*	*	*	*
4103	62	*	*	*	*	*	*	*
4106	2,477	16.4	42.1	35.0	0.2	0.6	4.8	0.9
4107	3,386	48.3	24.9	17.3	0.3	1.9	7.0	0.2

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4109	1,118	85.8	1.6	7.3	0.1	1.2	3.9	0.1
4112	582	1.9	0.9	95.5	0.0	1.0	0.2	0.5
4113	682	7.3	17.3	72.9	0.0	0.6	1.9	0.0
4115	3,649	39.0	24.6	19.0	0.4	11.0	5.2	0.7
4351	616	89.6	0.2	5.7	0.0	0.6	3.7	0.2
4352	178	69.1	16.9	7.3	0.0	0.0	6.2	0.6
4353	466	66.1	0.4	30.9	0.2	0.0	1.9	0.4
4354	800	88.8	3.6	2.9	0.4	0.4	3.6	0.4
4355	382	79.3	1.3	7.9	1.8	0.5	8.9	0.3
4356	496	85.1	0.8	10.7	0.0	0.0	3.4	0.0
4358	163	31.3	0.0	63.8	0.0	0.0	2.5	2.5
4359	495	72.3	0.2	24.4	0.2	0.6	1.0	1.2
4360	245	94.3	0.4	4.5	0.4	0.0	0.4	0.0
4361	252	94.4	0.0	0.8	0.4	0.4	3.6	0.4
4362	302	91.7	0.0	2.6	0.3	1.0	4.0	0.3
4363	944	93.2	0.1	2.6	0.0	0.3	3.4	0.3
4401	1,159	87.9	0.4	5.6	0.1	0.3	5.2	0.4
4501	1,027	92.3	0.5	4.5	0.0	0.2	2.3	0.2
4601	1,541	75.5	1.7	16.7	0.0	0.1	4.9	1.0
4602	30	*	*	*	*	*	*	*
4701	1,184	90.8	0.5	3.8	0.3	1.6	2.2	0.8
5050	1,845	54.0	36.7	2.8	0.0	1.3	4.8	0.3
5051	2,778	52.8	35.2	4.0	0.1	2.6	5.1	0.3
5052	3,006	42.4	36.0	1.8	0.2	14.5	4.7	0.5

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
5055	1,237	94.4	1.1	0.1	0.1	3.2	0.3	0.8
5101	2,011	69.3	11.3	12.8	0.0	1.5	5.0	0.0
5102	1	*	*	*	*	*	*	*
5202	834	84.9	0.6	3.7	1.2	0.5	8.4	0.7
5203	27	*	*	*	*	*	*	*
5204	14	*	*	*	*	*	*	*
5601	1,140	92.9	0.7	2.0	0.1	1.1	2.9	0.4
5801	1,797	81.3	6.8	6.6	0.1	0.4	4.1	0.7
5803	134	86.6	0.7	9.7	0.0	0.7	2.2	0.0
5804	19	*	*	*	*	*	*	*
5806	30	*	*	*	*	*	*	*
6101	31	*	*	*	*	*	*	*
6102	129	86.0	0.8	5.4	0.0	0.0	7.0	0.8
6103	4,346	55.3	28.6	8.6	0.3	0.4	6.5	0.3
6105	28	*	*	*	*	*	*	*
6106	17	*	*	*	*	*	*	*
6301	4,502	42.8	29.9	19.9	0.1	2.5	4.3	0.5
6312	2,399	33.7	56.4	3.3	0.0	2.1	4.0	0.4
6316	1,253	67.5	18.4	7.9	0.2	3.0	2.6	0.3
6320	127	64.6	22.0	5.5	0.0	1.6	5.5	0.8
6325	146	92.5	2.1	2.7	0.0	0.0	2.7	0.0
6330	365	27.1	53.2	15.9	0.0	1.1	2.7	0.0
6335	1,010	77.3	13.8	3.0	0.1	4.6	1.1	0.2
6340	58	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6345	29	*	*	*	*	*	*	*
6360	384	22.9	52.3	14.8	0.0	1.6	8.3	0.0
7304	128	88.3	0.0	7.0	0.0	0.0	3.9	0.8
7306	2,622	43.0	35.4	16.0	0.0	0.4	4.5	0.7
7308	129	87.6	0.8	7.8	0.0	0.0	3.1	0.8
7310	38	*	*	*	*	*	*	*
7401	2,470	82.0	5.7	5.9	0.1	0.4	5.3	0.6
7601	683	93.7	0.1	4.4	0.0	0.0	1.5	0.3
7850	1,170	88.3	0.4	6.2	0.2	0.3	4.4	0.3
7852	29	*	*	*	*	*	*	*
8101	2,053	5.6	90.8	0.6	0.1	0.3	2.4	0.1
8102	460	12.2	9.1	75.7	0.0	0.0	2.4	0.7
8105	1,780	32.2	59.9	3.0	0.2	1.0	3.4	0.3
8106	518	23.4	63.5	7.3	0.4	0.2	5.0	0.2
8150	42	*	*	*	*	*	*	*
8151	3,360	36.7	39.3	10.0	0.2	3.5	9.5	0.8
8152	111	45.9	36.9	9.0	0.0	0.9	7.2	0.0
8153	14	*	*	*	*	*	*	*
8155	39	*	*	*	*	*	*	*
8157	9	*	*	*	*	*	*	*
8201	2,826	46.9	39.0	4.8	0.2	2.9	5.8	0.4
8202	4,133	86.8	9.1	2.2	0.0	0.6	0.9	0.3
8203	481	61.5	26.8	4.0	0.4	1.5	5.2	0.6
8250	832	21.2	67.4	4.3	0.1	0.8	5.9	0.2

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8252	132	2.3	93.2	0.0	0.0	2.3	2.3	0.0
8255	2,797	55.6	30.5	6.7	0.3	2.6	3.7	0.6
8257	1,986	49.2	15.4	0.9	0.1	32.6	1.4	0.5
8263	758	70.3	12.8	9.9	0.1	1.8	4.4	0.7
8264	718	61.3	28.1	4.7	0.3	1.4	3.8	0.4
8265	21	*	*	*	*	*	*	*
8266	41	*	*	*	*	*	*	*
8267	804	58.1	17.3	18.8	0.1	0.2	4.4	0.9
8901	1,175	1.6	94.5	0.6	0.2	0.9	2.0	0.2
8902	1,414	4.3	91.4	1.3	0.0	0.1	2.5	0.3
8903	1,354	2.1	93.8	0.8	0.0	0.7	2.3	0.2
8904	2,487	2.1	92.6	1.0	0.2	1.3	2.5	0.2
8905	1,663	57.5	29.9	9.7	0.1	0.8	1.7	0.2
8908	31	*	*	*	*	*	*	*
8909	482	1.5	94.2	1.2	0.0	0.4	2.1	0.6
9012	7	*	*	*	*	*	*	*
9013	1	*	*	*	*	*	*	*
9016	52	*	*	*	*	*	*	*
9017	6	*	*	*	*	*	*	*
9018	55	*	*	*	*	*	*	*
9041	2	*	*	*	*	*	*	*
9045	1,872	10.0	26.4	59.3	0.1	0.4	3.5	0.4
9050	1,012	10.8	71.1	15.9	0.0	0.3	1.8	0.1
9051	1,762	9.4	7.6	81.0	0.0	0.2	1.5	0.1

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9053	1,164	3.4	90.1	0.8	0.0	4.0	1.5	0.2
9055	50	*	*	*	*	*	*	*
9058	1,099	23.7	63.8	1.5	0.1	9.6	1.2	0.3
9066	1,180	1.5	97.0	0.7	0.0	0.1	0.6	0.1
9068	1,020	2.3	96.1	0.4	0.0	0.3	0.8	0.2
9069	3,034	1.1	97.5	0.5	0.0	0.2	0.6	0.2
9070	1,204	2.7	92.4	0.4	0.0	2.7	1.6	0.2
9072	1,967	2.0	95.1	0.5	0.1	0.5	1.5	0.4
9080	2,862	20.4	75.5	1.8	0.1	0.1	1.7	0.3
9081	49	*	*	*	*	*	*	*
9082	123	3.3	95.1	0.8	0.0	0.0	0.8	0.0
9083	56	*	*	*	*	*	*	*
9084	25	*	*	*	*	*	*	*
9085	49	*	*	*	*	*	*	*
9086	36	*	*	*	*	*	*	*
Michigan	**173,232**	**52.4**	**29.7**	**10.5**	**0.3**	**2.2**	**4.3**	**0.5**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	117	0.0	1.7	9.4	59.8	28.2	0.9	0.0
0102	540	0.0	4.6	8.3	65.2	21.5	0.4	0.0
0103	455	0.0	2.6	7.7	68.4	20.0	1.3	0.0
0104	161	0.0	2.5	9.3	76.4	11.8	0.0	0.0
0105	7	*	*	*	*	*	*	*
0250	146	0.0	4.1	6.2	70.5	19.2	0.0	0.0
0251	95	*	*	*	*	*	*	*
0252	122	0.0	2.5	2.5	68.0	25.4	1.6	0.0
0253	175	0.6	2.9	9.1	65.1	22.3	0.0	0.0
0401	513	0.0	3.3	9.9	63.0	22.2	1.6	0.0
0402	386	0.3	1.3	6.2	71.0	20.7	0.5	0.0
0403	140	0.0	2.1	7.9	68.6	21.4	0.0	0.0
0404	148	0.0	2.0	9.5	64.2	20.9	3.4	0.0
0501	551	0.2	3.1	8.9	65.2	22.3	0.4	0.0
0503	118	0.0	1.7	6.8	59.3	30.5	1.7	0.0
0504	516	0.0	1.0	10.1	64.0	24.0	1.0	0.0
0505	351	0.0	1.4	10.8	62.1	24.2	1.4	0.0
0506	220	0.0	2.3	13.2	63.2	20.5	0.9	0.0
0507	85	*	*	*	*	*	*	*
0508	32	*	*	*	*	*	*	*
0509	8	*	*	*	*	*	*	*
0650	1,171	0.0	2.8	7.9	65.3	23.0	0.9	0.2
0651	625	0.3	2.7	11.2	67.4	16.6	1.8	0.0
0652	313	0.0	3.5	9.3	65.5	20.1	1.6	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0653	451	0.0	2.7	9.5	68.5	18.4	0.4	0.4
0654	386	0.3	5.4	11.4	59.1	22.5	1.3	0.0
0655	393	0.0	2.8	10.2	64.1	22.4	0.5	0.0
0657	66	*	*	*	*	*	*	*
0701	437	0.0	1.8	7.8	62.2	25.6	2.3	0.2
0702	68	*	*	*	*	*	*	*
0703	233	0.0	2.6	9.4	64.8	22.7	0.4	0.0
0704	54	*	*	*	*	*	*	*
0705	135	0.0	4.4	7.4	53.3	31.1	3.7	0.0
0706	5	*	*	*	*	*	*	*
0850	945	0.2	1.7	9.3	62.8	24.6	1.0	0.5
0851	804	0.1	3.5	9.8	66.0	19.4	1.1	0.0
0901	1,775	0.1	2.9	9.0	64.3	22.3	1.4	0.0
0902	46	*	*	*	*	*	*	*
1051	1,454	0.0	3.0	7.2	62.0	25.7	1.9	0.3
1101	1,890	0.3	4.4	9.2	63.7	20.8	1.0	0.5
1102	888	0.3	3.9	10.9	62.5	20.5	1.6	0.2
1106	79	*	*	*	*	*	*	*
1250	1,083	0.0	2.8	8.8	62.7	23.4	2.3	0.1
1251	882	0.1	3.2	10.7	66.9	18.3	0.9	0.0
1252	585	0.0	3.4	12.6	62.4	19.7	1.9	0.0
1253	900	0.1	4.4	9.6	66.4	18.4	1.0	0.0
1350	2,631	0.1	3.2	9.5	63.6	22.0	1.4	0.2
1351	143	0.0	2.1	6.3	60.8	30.1	0.7	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1352	383	0.0	4.4	7.3	69.5	16.7	2.1	0.0
1401	255	0.0	2.7	8.2	64.3	22.4	2.0	0.4
1402	147	0.0	1.4	7.5	59.9	28.6	1.4	1.4
1403	28	*	*	*	*	*	*	*
1404	1	*	*	*	*	*	*	*
1501	841	0.0	4.0	10.8	61.5	21.2	2.4	0.1
1503	1,146	0.2	3.5	9.2	62.2	23.2	1.7	0.0
1504	2,322	0.3	3.3	8.1	62.1	24.1	2.1	0.0
1505	477	0.0	1.5	5.5	64.6	27.3	1.0	0.2
1506	626	0.2	3.0	7.7	62.0	25.6	1.6	0.0
1507	659	0.0	1.5	5.5	66.2	24.9	2.0	0.0
1508	114	0.0	7.0	7.0	41.2	39.5	4.4	0.9
1509	485	0.0	5.2	7.6	63.1	21.2	2.9	0.0
1511	763	0.3	3.9	11.0	62.4	21.1	1.3	0.0
1512	386	0.3	3.1	6.0	63.2	26.9	0.5	0.0
1513	78	*	*	*	*	*	*	*
1514	170	0.0	2.4	4.1	49.4	42.9	1.2	0.0
1701	712	0.1	1.7	7.6	63.8	25.4	1.4	0.0
1950	625	0.0	3.4	12.5	68.0	15.0	1.1	0.0
1952	139	0.7	2.2	7.9	64.0	23.0	2.2	0.0
1953	332	0.3	1.2	10.5	70.5	16.9	0.6	0.0
1954	601	0.2	2.2	7.7	65.7	22.5	1.8	0.0
1955	89	*	*	*	*	*	*	*
1957	567	0.0	2.8	10.8	62.1	22.4	1.9	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1963	11	*	*	*	*	*	*	*
1964	34	*	*	*	*	*	*	*
2150	705	0.0	2.6	8.7	61.0	26.7	1.1	0.0
2151	256	0.0	4.7	10.9	59.8	23.0	1.6	0.0
2251	464	0.0	3.2	8.6	62.3	24.4	1.3	0.2
2252	210	0.0	1.9	10.5	59.5	25.7	2.4	0.0
2554	85	*	*	*	*	*	*	*
2555	8,410	0.2	2.6	9.0	65.9	21.1	1.2	0.0
2556	204	1.0	1.5	7.8	67.2	22.1	0.5	0.0
2557	9	*	*	*	*	*	*	*
2558	4	*	*	*	*	*	*	*
2559	9	*	*	*	*	*	*	*
2560	2	*	*	*	*	*	*	*
2562	2	*	*	*	*	*	*	*
2563	5	*	*	*	*	*	*	*
3001	1,176	0.5	4.6	9.7	64.6	19.0	1.4	0.1
3002	264	0.4	4.9	12.9	60.6	20.1	1.1	0.0
3004	153	0.0	0.7	6.5	62.1	27.5	3.3	0.0
3007	30	*	*	*	*	*	*	*
3009	10	*	*	*	*	*	*	*
3201	502	0.0	2.4	9.4	62.2	25.1	0.8	0.2
3301	4,437	0.3	3.1	8.1	62.6	24.0	1.9	0.1
3304	112	0.0	0.0	0.9	48.2	49.1	1.8	0.0
3305	148	0.0	2.7	5.4	48.0	40.5	3.4	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3306	263	0.4	9.5	5.7	58.2	25.9	0.4	0.0
3308	76	*	*	*	*	*	*	*
3309	40	*	*	*	*	*	*	*
3312	422	0.2	3.6	6.2	62.6	25.6	1.9	0.0
3550	76	*	*	*	*	*	*	*
3551	29	*	*	*	*	*	*	*
3552	53	*	*	*	*	*	*	*
3701	1,142	0.1	3.3	8.9	69.2	17.8	0.7	0.0
3850	3,304	0.2	3.5	9.1	66.9	19.3	0.8	0.2
3901	580	0.0	3.6	10.3	64.8	19.7	1.6	0.0
3902	268	0.0	3.0	9.3	60.8	24.6	2.2	0.0
3903	10	*	*	*	*	*	*	*
3904	3	*	*	*	*	*	*	*
3905	2	*	*	*	*	*	*	*
3950	2,066	0.0	3.0	9.6	62.7	22.7	1.8	0.1
3954	5	*	*	*	*	*	*	*
4001	1,045	0.6	5.5	11.3	61.7	19.5	1.4	0.0
4002	720	0.0	4.9	8.5	61.9	23.5	1.3	0.0
4003	5	*	*	*	*	*	*	*
4004	5	*	*	*	*	*	*	*
4103	62	*	*	*	*	*	*	*
4106	2,477	0.4	4.0	9.7	60.3	23.5	1.9	0.2
4107	3,386	0.1	3.2	7.7	62.8	24.4	1.7	0.1
4109	1,118	0.1	3.0	6.1	64.0	24.6	2.1	0.1

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4112	582	0.0	4.8	9.5	51.0	30.8	3.8	0.2
4113	682	0.4	3.7	7.8	45.6	32.6	2.6	7.3
4115	3,649	0.1	2.2	6.4	60.7	28.5	1.9	0.2
4351	616	0.2	3.2	9.4	68.3	17.9	1.0	0.0
4352	178	0.0	3.4	12.4	66.9	15.7	1.7	0.0
4353	466	0.0	3.6	8.6	66.7	19.5	1.5	0.0
4354	800	0.0	3.4	9.9	68.9	17.0	0.9	0.0
4355	382	0.0	5.0	13.4	58.6	20.7	2.1	0.3
4356	496	0.2	3.8	5.8	71.4	18.1	0.6	0.0
4358	163	0.6	3.1	6.7	65.0	23.3	1.2	0.0
4359	495	0.0	3.0	8.7	61.6	23.6	3.0	0.0
4360	245	0.0	3.3	9.4	68.2	16.7	2.4	0.0
4361	252	0.0	3.2	9.1	62.3	25.4	0.0	0.0
4362	302	0.0	2.3	6.3	71.9	18.9	0.7	0.0
4363	944	0.0	3.4	8.4	66.8	20.9	0.5	0.0
4401	1,159	0.2	2.9	8.4	65.3	21.8	1.3	0.1
4501	1,027	0.2	3.1	8.7	65.2	21.3	1.4	0.1
4601	1,541	0.0	3.2	10.3	63.6	21.9	1.0	0.0
4602	30	*	*	*	*	*	*	*
4701	1,184	0.0	1.6	6.1	62.1	28.7	1.5	0.0
5050	1,845	0.1	1.8	6.9	59.7	29.5	2.0	0.0
5051	2,778	0.0	1.2	4.8	65.8	26.5	1.7	0.0
5052	3,006	0.0	2.5	5.9	63.0	26.3	2.2	0.0
5055	1,237	0.0	0.2	0.8	54.7	40.1	4.1	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5101	2,011	0.1	0.8	5.6	61.8	28.9	2.7	0.0
5102	1	*	*	*	*	*	*	*
5202	834	0.1	4.0	7.0	60.3	27.6	1.1	0.0
5203	27	*	*	*	*	*	*	*
5204	14	*	*	*	*	*	*	*
5601	1,140	0.0	2.0	7.4	66.8	22.4	1.4	0.1
5801	1,797	0.0	3.6	10.8	62.9	21.4	1.3	0.1
5803	134	0.0	1.5	2.2	63.4	31.3	1.5	0.0
5804	19	*	*	*	*	*	*	*
5806	30	*	*	*	*	*	*	*
6101	31	*	*	*	*	*	*	*
6102	129	0.0	7.0	7.0	58.9	25.6	1.6	0.0
6103	4,346	0.3	3.6	9.2	65.5	20.0	1.4	0.0
6105	28	*	*	*	*	*	*	*
6106	17	*	*	*	*	*	*	*
6301	4,502	0.2	3.1	8.0	62.2	24.6	1.9	0.1
6312	2,399	0.0	1.4	5.0	58.4	32.8	2.5	0.0
6316	1,253	0.0	0.9	5.0	59.0	32.0	3.0	0.1
6320	127	0.0	0.0	0.8	52.8	40.2	6.3	0.0
6325	146	0.0	2.1	4.8	67.8	25.3	0.0	0.0
6330	365	0.3	3.0	10.4	63.8	21.6	0.8	0.0
6335	1,010	0.0	1.2	4.8	53.4	37.3	3.4	0.0
6340	58	*	*	*	*	*	*	*
6345	29	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
6360	384	0.3	1.0	7.0	67.4	21.9	2.1	0.3
7304	128	0.0	2.3	6.3	65.6	25.8	0.0	0.0
7306	2,622	0.2	3.1	8.2	65.6	21.4	1.4	0.0
7308	129	0.0	3.1	11.6	65.1	18.6	1.6	0.0
7310	38	*	*	*	*	*	*	*
7401	2,470	0.2	3.3	8.6	64.5	22.0	1.4	0.0
7601	683	0.0	5.0	8.6	61.3	24.2	0.9	0.0
7850	1,170	0.3	2.7	10.3	68.8	16.9	1.0	0.0
7852	29	*	*	*	*	*	*	*
8101	2,053	0.2	3.1	9.0	65.0	20.6	2.1	0.0
8102	460	0.0	3.5	5.7	59.1	28.5	3.0	0.2
8105	1,780	0.1	2.0	5.3	61.4	29.3	1.8	0.1
8106	518	0.2	4.1	7.1	65.3	21.0	2.3	0.0
8150	42	*	*	*	*	*	*	*
8151	3,360	0.1	2.6	6.0	60.6	27.9	2.7	0.1
8152	111	0.0	4.5	14.4	73.0	8.1	0.0	0.0
8153	14	*	*	*	*	*	*	*
8155	39	*	*	*	*	*	*	*
8157	9	*	*	*	*	*	*	*
8201	2,826	0.1	2.6	6.5	63.6	25.4	1.7	0.1
8202	4,133	0.0	1.1	3.2	57.2	33.3	4.1	1.1
8203	481	0.4	1.5	8.1	61.3	26.4	2.3	0.0
8250	832	0.2	2.8	7.1	60.7	27.4	1.8	0.0
8252	132	0.0	3.0	6.1	68.9	20.5	1.5	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8255	2,797	0.0	2.0	6.4	66.2	24.0	1.3	0.0
8257	1,986	0.1	1.1	4.2	58.7	31.2	4.5	0.3
8263	758	0.1	1.5	5.8	64.4	26.9	1.3	0.0
8264	718	0.1	4.0	5.7	61.0	26.2	2.9	0.0
8265	21	*	*	*	*	*	*	*
8266	41	*	*	*	*	*	*	*
8267	804	0.0	3.1	7.0	62.3	26.2	1.4	0.0
8901	1,175	0.3	3.4	8.2	64.8	21.5	1.9	0.0
8902	1,414	0.2	2.2	8.6	64.1	22.3	2.5	0.1
8903	1,354	0.0	3.1	7.9	64.8	22.5	1.6	0.1
8904	2,487	0.1	2.9	7.8	65.8	21.7	1.8	0.0
8905	1,663	0.0	1.3	5.1	59.7	31.0	2.8	0.1
8908	31	*	*	*	*	*	*	*
8909	482	0.4	3.9	9.1	58.1	27.0	1.5	0.0
9012	7	*	*	*	*	*	*	*
9013	1	*	*	*	*	*	*	*
9016	52	*	*	*	*	*	*	*
9017	6	*	*	*	*	*	*	*
9018	55	*	*	*	*	*	*	*
9041	2	*	*	*	*	*	*	*
9045	1,872	0.1	3.0	8.0	57.6	28.7	2.5	0.1
9050	1,012	0.5	3.1	6.8	61.1	25.9	2.7	0.0
9051	1,762	0.1	3.3	8.0	50.3	32.6	5.7	0.0
9053	1,164	0.0	2.7	6.4	67.6	21.0	2.3	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9055	50	*	*	*	*	*	*	*
9058	1,099	0.2	2.5	4.4	59.6	29.6	3.7	0.1
9066	1,180	0.1	3.0	7.3	65.9	22.0	1.7	0.0
9068	1,020	0.0	3.2	9.7	62.6	22.7	1.7	0.0
9069	3,034	0.1	2.9	6.8	65.4	22.8	2.0	0.0
9070	1,204	0.2	2.8	8.6	64.1	22.8	1.3	0.0
9072	1,967	0.1	3.4	8.2	67.0	19.6	1.7	0.0
9080	2,862	0.1	2.3	6.4	61.8	26.9	2.4	0.1
9081	49	*	*	*	*	*	*	*
9082	123	0.0	4.9	6.5	52.8	35.8	0.0	0.0
9083	56	*	*	*	*	*	*	*
9084	25	*	*	*	*	*	*	*
9085	49	*	*	*	*	*	*	*
9086	36	*	*	*	*	*	*	*
Michigan	**173,232**	**0.1**	**2.8**	**7.8**	**63.0**	**24.3**	**1.9**	**0.1**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	86	*	*	28	*	*	19	*	31	*	112	5.4 (25)	112	8.9 (38)
0102	445	4.9 (114)	60.7 (106)	239	17.6 (47)	51.5 (60)	75	*	103	36.9 (56)	508	4.7 (11)	508	7.9 (24)
0103	392	2.6 (9)	60.7 (106)	245	18.8 (75)	48.6 (24)	58	*	100	35.0 (49)	422	8.5 (134)	422	10.9 (72)
0104	138	2.9 (18)	58.0 (67)	73	*	*	17	*	31	*	151	7.9 (113)	151	8.6 (37)
0105	7	*	*	6	*	*	1	*	0	*	7	*	7	*
0250	138	3.6 (50)	69.6 (151)	73	*	*	16	*	33	*	136	7.4 (96)	137	10.2 (57)
0251	77	*	*	35	*	*	8	*	24	*	91	*	90	*
0252	122	0.8 (1)	59.0 (80)	107	25.2 (126)	50.5 (44)	13	*	23	*	113	8.0 (120)	113	15.0 (145)
0253	125	5.6 (136)	55.2 (31)	42	*	*	13	*	39	*	165	8.5 (134)	165	17.0 (156)
0401	481	3.5 (45)	57.4 (57)	398	20.6 (99)	51.8 (63)	73	*	106	29.2 (25)	473	4.2 (4)	473	7.2 (21)
0402	360	5.3 (129)	58.6 (75)	248	18.1 (57)	50.4 (43)	68	*	55	*	368	7.1 (84)	368	6.5 (11)
0403	131	7.6 (151)	54.2 (22)	93	*	*	21	*	30	*	129	6.2 (56)	129	11.6 (90)
0404	127	1.6 (4)	60.6 (102)	94	*	*	25	*	34	*	143	6.3 (60)	143	11.2 (80)
0501	342	4.4 (89)	56.1 (37)	166	17.5 (45)	51.8 (63)	90	*	95	*	510	5.1 (17)	510	8.2 (28)
0503	101	5.0 (118)	49.5 (5)	59	*	*	11	*	20	*	109	8.3 (130)	109	8.3 (31)
0504	447	3.8 (58)	51.7 (12)	221	19.0 (77)	48.9 (26)	88	*	72	*	489	6.3 (60)	490	9.4 (44)
0505	295	4.1 (71)	52.9 (16)	180	18.3 (64)	56.7 (120)	51	*	50	*	331	6.9 (77)	332	7.5 (22)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	145	8.3 (153)	57.9 (65)	51	*	*	22	*	30	*	202	5.9 (44)	204	6.9 (19)
0507	59	*	*	33	*	*	5	*	21	*	84	*	84	*
0508	19	*	*	7	*	*	3	*	6	*	31	*	31	*
0509	8	*	*	7	*	*	2	*	1	*	8	*	8	*
0650	1,021	3.7 (53)	59.8 (88)	648	18.1 (57)	52.2 (65)	179	34.1 (9)	242	27.7 (22)	1,095	7.0 (81)	1,095	9.6 (47)
0651	594	5.1 (122)	62.1 (124)	475	18.5 (67)	51.6 (62)	87	*	76	*	581	6.9 (77)	581	6.2 (7)
0652	222	4.5 (92)	50.9 (9)	139	16.5 (29)	52.5 (69)	52	*	40	*	301	6.0 (50)	301	10.0 (52)
0653	408	3.9 (61)	61.0 (114)	287	13.6 (3)	53.0 (79)	50	*	77	*	426	4.2 (4)	427	12.6 (111)
0654	345	4.1 (71)	63.5 (132)	237	18.1 (57)	57.0 (122)	45	*	87	*	349	8.3 (130)	348	10.1 (55)
0655	362	5.0 (118)	58.6 (75)	168	15.5 (19)	59.5 (130)	45	*	76	*	363	7.2 (86)	365	13.2 (125)
0657	44	*	*	21	*	*	8	*	12	*	63	*	63	*
0701	307	4.6 (97)	45.6 (1)	140	14.3 (7)	53.6 (93)	66	*	130	16.9 (3)	410	5.1 (17)	411	12.2 (102)
0702	63	*	*	36	*	*	7	*	13	*	59	*	59	*
0703	173	4.6 (97)	55.5 (32)	77	*	*	29	*	51	*	199	8.5 (134)	199	6.5 (11)
0704	47	*	*	23	*	*	6	*	11	*	46	*	46	*
0705	78	*	*	29	*	*	19	*	34	*	123	7.3 (91)	123	10.6 (64)
0706	3	*	*	2	*	*	0	*	1	*	5	*	5	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	648	4.3 (81)	57.6 (61)	416	15.9 (22)	54.6 (105)	163	38.7 (17)	210	22.4 (12)	872	8.9 (142)	874	15.8 (151)
0851	511	6.3 (143)	57.3 (55)	184	14.1 (6)	55.4 (110)	103	34.0 (8)	165	18.2 (6)	735	7.9 (113)	736	13.2 (125)
0901	1,469	4.3 (81)	58.7 (78)	1,101	19.7 (85)	50.6 (46)	258	37.6 (14)	228	24.1 (15)	1,659	7.7 (105)	1,659	10.1 (55)
0902	40	*	*	28	*	*	1	*	8	*	44	*	44	*
1051	1,312	4.6 (97)	51.8 (13)	1,033	16.9 (37)	52.5 (69)	208	60.6 (65)	209	45.5 (80)	1,381	6.9 (77)	1,383	9.5 (45)
1101	1,622	3.8 (58)	60.6 (102)	1,181	18.5 (67)	54.2 (98)	333	51.4 (48)	277	38.6 (64)	1,730	6.2 (56)	1,729	13.0 (119)
1102	752	2.8 (15)	57.8 (63)	516	14.7 (9)	58.1 (126)	115	37.4 (13)	201	32.8 (40)	776	6.1 (53)	776	8.2 (28)
1106	69	*	*	51	*	*	9	*	25	*	69	*	70	*
1250	842	5.3 (129)	60.9 (110)	450	15.1 (12)	54.4 (100)	155	12.9 (1)	154	23.4 (14)	978	7.9 (113)	978	6.5 (11)
1251	771	5.8 (139)	58.4 (72)	514	18.5 (67)	50.2 (38)	121	38.0 (16)	156	25.6 (16)	814	4.3 (7)	815	6.5 (11)
1252	458	2.8 (15)	61.6 (116)	271	18.5 (67)	58.3 (127)	77	*	94	*	542	5.4 (25)	542	16.4 (154)
1253	701	4.7 (105)	60.3 (96)	438	16.4 (27)	53.4 (90)	118	37.3 (12)	171	26.3 (19)	810	6.9 (77)	812	12.7 (113)
1350	2,221	3.1 (22)	59.0 (80)	1,240	19.8 (87)	53.2 (86)	370	52.4 (51)	474	38.4 (61)	2,394	6.8 (72)	2,393	9.6 (47)
1351	123	3.3 (30)	54.5 (26)	96	*	*	19	*	31	*	140	8.6 (137)	140	7.9 (24)
1352	228	6.6 (145)	56.1 (37)	106	23.6 (122)	50.9 (51)	45	*	65	*	354	6.2 (56)	353	4.0 (3)
1401	237	3.4 (40)	56.5 (42)	182	19.2 (80)	51.1 (53)	33	*	30	*	235	5.5 (30)	235	6.8 (18)
1402	135	1.5 (2)	60.0 (92)	100	17.0 (39)	58.0 (125)	21	*	25	*	134	5.2 (21)	134	6.7 (16)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1403	27	*	*	20	*	*	3	*	4	*	27	*	27	*
1404	1	*	*	0	*	*	0	*	0	*	0	*	0	*
1501	747	3.3 (30)	64.4 (136)	501	47.1 (130)	24.8 (2)	84	*	127	22.0 (11)	723	6.6 (68)	724	7.7 (23)
1503	972	3.5 (45)	60.3 (96)	579	17.6 (47)	53.2 (86)	135	44.4 (33)	145	29.7 (28)	1,033	6.3 (60)	1,032	11.8 (96)
1504	1,966	4.8 (110)	59.7 (87)	1,354	20.2 (94)	46.3 (11)	339	58.7 (62)	509	45.0 (78)	2,116	9.3 (147)	2,118	9.1 (40)
1505	392	3.3 (30)	54.8 (28)	248	15.3 (17)	53.6 (93)	58	*	123	33.3 (41)	445	5.6 (34)	445	11.5 (87)
1506	505	3.2 (27)	58.6 (75)	154	17.5 (45)	48.1 (20)	87	*	112	28.6 (23)	567	7.4 (96)	566	8.5 (35)
1507	513	3.9 (61)	53.0 (17)	241	14.9 (11)	51.0 (52)	76	*	117	36.8 (55)	598	7.5 (101)	597	11.2 (80)
1508	49	*	*	13	*	*	5	*	13	*	85	*	85	*
1509	378	3.2 (27)	65.9 (145)	197	20.3 (95)	51.3 (57)	50	*	81	*	394	10.2 (154)	395	8.1 (27)
1511	606	4.5 (92)	60.7 (106)	275	21.1 (105)	49.8 (34)	106	38.7 (17)	164	33.5 (43)	702	7.1 (84)	705	14.0 (134)
1512	272	3.3 (30)	56.3 (39)	89	*	*	52	*	85	*	361	5.0 (15)	362	13.0 (119)
1513	67	*	*	45	*	*	7	*	17	*	72	*	72	*
1514	131	3.1 (22)	56.5 (42)	96	*	*	28	*	69	*	162	8.0 (120)	162	11.7 (94)
1701	509	2.9 (18)	62.3 (126)	160	20.0 (90)	53.1 (83)	87	*	103	31.1 (34)	647	8.0 (120)	646	12.2 (102)
1950	589	4.1 (71)	60.3 (96)	475	21.3 (110)	53.5 (92)	82	*	153	17.0 (4)	579	8.8 (140)	582	14.8 (143)
1952	132	6.8 (149)	62.1 (124)	108	18.5 (67)	47.2 (14)	26	*	52	*	133	8.3 (130)	133	9.0 (39)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1953	309	5.8 (139)	51.5 (10)	253	21.3 (110)	47.4 (16)	45	*	85	*	309	9.1 (144)	309	11.3 (83)
1954	528	6.6 (145)	57.0 (51)	336	17.6 (47)	56.0 (115)	82	*	108	14.8 (2)	559	7.2 (86)	559	14.3 (140)
1955	79	*	*	48	*	*	14	*	19	*	83	*	83	*
1957	519	4.6 (97)	60.9 (110)	410	17.1 (41)	55.1 (108)	82	*	153	20.9 (9)	529	6.8 (72)	530	13.2 (125)
1963	10	*	*	10	*	*	1	*	5	*	10	*	10	*
1964	32	*	*	27	*	*	6	*	9	*	30	*	31	*
2150	494	4.0 (67)	54.0 (19)	253	20.6 (99)	45.5 (9)	79	*	145	30.3 (30)	630	13.5 (157)	630	11.1 (77)
2151	154	2.6 (9)	48.7 (2)	79	*	*	36	*	49	*	213	5.6 (34)	213	10.8 (70)
2251	386	4.1 (71)	51.6 (11)	176	13.1 (2)	48.9 (26)	52	*	94	*	416	3.4 (2)	415	12.3 (104)
2252	150	4.0 (67)	56.7 (45)	25	*	*	26	*	36	*	181	5.5 (30)	182	15.9 (152)
2554	76	*	*	58	*	*	17	*	20	*	80	*	80	*
2555	6,910	5.1 (122)	57.0 (51)	3,075	17.7 (54)	52.5 (69)	1,493	49.3 (45)	1,327	34.7 (48)	7,851	7.7 (105)	7,853	15.0 (145)
2556	182	3.3 (30)	54.9 (29)	155	16.8 (35)	56.1 (118)	20	*	42	*	199	7.5 (101)	199	19.6 (158)
2557	3	*	*	0	*	*	5	*	1	*	7	*	7	*
2558	3	*	*	2	*	*	2	*	0	*	4	*	4	*
2559	7	*	*	4	*	*	3	*	0	*	9	*	9	*
2560	2	*	*	0	*	*	0	*	0	*	2	*	2	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2562	1	*	*	0	*	*	2	*	1	*	1	*	1	*
2563	5	*	*	2	*	*	0	*	1	*	5	*	5	*
3001	1,102	3.5 (45)	62.0 (123)	815	26.0 (128)	43.4 (6)	199	48.7 (43)	748	46.9 (84)	1,089	5.2 (21)	1,088	9.7 (50)
3002	253	1.6 (4)	78.3 (153)	198	22.7 (118)	48.0 (19)	43	*	184	45.1 (79)	249	2.8 (1)	249	12.9 (116)
3004	139	2.9 (18)	64.0 (135)	124	28.2 (129)	40.3 (3)	20	*	21	*	139	5.8 (43)	139	7.9 (24)
3007	25	*	*	13	*	*	2	*	7	*	20	*	20	*
3009	5	*	*	0	*	*	0	*	1	*	2	*	2	*
3201	412	3.9 (61)	65.3 (143)	334	16.2 (25)	53.3 (89)	50	*	115	26.1 (18)	474	9.7 (150)	472	14.6 (142)
3301	3,784	4.7 (105)	58.4 (72)	2,878	18.5 (67)	52.7 (75)	727	42.9 (26)	748	32.4 (39)	4,112	5.7 (38)	4,114	10.0 (52)
3304	88	*	*	69	*	*	10	*	17	*	107	15.0 (158)	107	4.7 (5)
3305	120	6.7 (147)	60.0 (92)	96	*	*	15	*	29	*	142	7.7 (105)	142	6.3 (8)
3306	236	7.2 (150)	54.2 (22)	195	11.3 (1)	59.0 (129)	44	*	59	*	254	5.9 (44)	254	8.3 (31)
3308	66	*	*	56	*	*	6	*	20	*	72	*	72	*
3309	37	*	*	27	*	*	6	*	14	*	38	*	38	*
3312	372	5.4 (131)	64.8 (139)	293	20.8 (102)	52.6 (74)	89	*	68	*	398	7.0 (81)	398	10.8 (70)
3550	70	*	*	40	*	*	9	*	20	*	70	*	70	*
3551	23	*	*	15	*	*	3	*	12	*	27	*	27	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3552	44	*	*	21	*	*	4	*	12	*	47	*	47	*
3701	824	5.0 (118)	57.4 (57)	594	14.6 (8)	54.5 (104)	183	21.9 (2)	266	14.7 (1)	1,045	5.9 (44)	1,043	14.1 (135)
3850	2,450	5.9 (141)	59.4 (85)	1,364	16.5 (29)	52.5 (69)	484	49.6 (47)	610	42.8 (72)	3,041	5.9 (44)	3,041	10.9 (72)
3901	447	5.6 (136)	57.5 (59)	226	22.1 (114)	45.6 (10)	86	*	133	51.9 (87)	539	5.4 (25)	539	9.3 (43)
3902	231	2.6 (9)	66.2 (146)	124	13.7 (4)	54.0 (96)	35	*	54	*	251	10.4 (156)	252	8.3 (31)
3903	9	*	*	8	*	*	0	*	6	*	10	*	10	*
3904	3	*	*	0	*	*	1	*	1	*	3	*	3	*
3905	2	*	*	2	*	*	0	*	0	*	2	*	2	*
3950	1,651	3.6 (50)	58.4 (72)	1,028	16.1 (24)	57.4 (124)	302	44.0 (28)	510	35.1 (51)	1,870	8.7 (138)	1,871	9.6 (47)
3954	4	*	*	2	*	*	0	*	1	*	4	*	4	*
4001	887	3.7 (53)	59.1 (83)	461	22.1 (114)	49.5 (31)	107	52.3 (50)	179	43.6 (73)	932	7.4 (96)	933	11.7 (94)
4002	645	4.7 (105)	60.6 (102)	398	17.1 (41)	57.0 (122)	79	*	124	30.6 (31)	659	8.2 (127)	659	6.7 (16)
4003	4	*	*	2	*	*	2	*	1	*	5	*	5	*
4004	5	*	*	3	*	*	0	*	0	*	5	*	5	*
4103	36	*	*	17	*	*	8	*	12	*	53	*	53	*
4106	1,881	3.2 (27)	60.9 (110)	1,036	19.5 (84)	48.3 (22)	404	57.2 (60)	620	44.0 (75)	2,256	6.0 (50)	2,257	10.2 (57)
4107	2,783	4.1 (71)	56.8 (46)	1,509	17.6 (47)	50.6 (46)	569	44.3 (31)	1,076	38.8 (66)	3,133	5.9 (44)	3,136	12.9 (116)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4109	507	7.7 (152)	49.5 (5)	186	16.7 (33)	52.2 (65)	148	29.1 (4)	298	20.1 (7)	1,041	7.0 (81)	1,047	15.1 (147)
4112	577	3.3 (30)	63.1 (130)	516	76.4 (131)	6.8 (1)	69	*	63	*	552	15.0 (158)	551	9.8 (51)
4113	611	2.9 (18)	63.7 (133)	119	17.6 (47)	41.2 (5)	99	*	151	33.8 (44)	632	10.0 (153)	634	11.5 (87)
4115	2,350	4.6 (97)	54.4 (25)	1,016	17.6 (47)	50.2 (38)	602	44.2 (30)	1,177	38.5 (62)	3,396	7.3 (91)	3,397	10.7 (68)
4351	587	3.9 (61)	61.8 (120)	426	22.8 (119)	46.9 (13)	86	*	113	31.9 (35)	578	9.0 (143)	579	11.9 (98)
4352	160	4.4 (89)	60.6 (102)	106	19.8 (87)	52.8 (78)	20	*	32	*	163	4.3 (7)	164	10.4 (61)
4353	398	2.8 (15)	65.1 (141)	283	19.1 (79)	52.7 (75)	60	*	85	*	426	6.1 (53)	426	12.4 (107)
4354	678	4.3 (81)	57.7 (62)	424	20.8 (102)	50.7 (48)	126	31.0 (6)	173	20.2 (8)	749	4.4 (9)	749	11.2 (80)
4355	337	5.0 (118)	60.2 (94)	170	16.5 (29)	51.2 (54)	68	*	99	*	358	7.8 (110)	358	9.5 (45)
4356	408	5.1 (122)	55.1 (30)	207	19.8 (87)	47.8 (18)	75	*	94	*	457	4.2 (4)	457	12.3 (104)
4358	133	1.5 (2)	66.9 (147)	102	18.6 (73)	45.1 (8)	16	*	23	*	137	7.3 (91)	137	12.4 (107)
4359	433	3.7 (53)	63.3 (131)	246	21.5 (112)	51.2 (54)	61	*	95	*	466	9.2 (146)	468	14.1 (135)
4360	230	4.3 (81)	60.4 (100)	187	18.7 (74)	50.8 (50)	31	*	53	*	233	4.7 (11)	233	11.6 (90)
4361	227	4.8 (110)	59.9 (89)	177	16.4 (27)	62.1 (131)	27	*	57	*	237	5.1 (17)	238	9.2 (42)
4362	274	4.7 (105)	54.0 (19)	208	21.2 (108)	50.5 (44)	53	*	74	*	285	7.7 (105)	285	6.0 (6)
4363	905	3.4 (40)	57.8 (63)	749	21.1 (105)	52.5 (69)	124	41.9 (23)	204	26.5 (20)	883	5.7 (38)	883	12.1 (100)
4401	1,137	4.9 (114)	55.5 (32)	972	15.1 (12)	55.5 (112)	193	39.4 (19)	317	32.2 (38)	1,085	4.9 (13)	1,085	6.6 (15)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4501	786	6.7 (147)	54.5 (26)	573	16.9 (37)	53.8 (95)	136	27.9 (3)	176	21.6 (10)	981	8.1 (126)	981	11.0 (75)
4601	1,279	3.9 (61)	56.9 (48)	465	24.3 (124)	48.2 (21)	291	42.6 (25)	342	33.3 (41)	1,401	7.4 (96)	1,403	12.9 (116)
4602	21	*	*	12	*	*	1	*	7	*	30	*	30	*
4701	1,095	5.6 (136)	52.5 (14)	742	15.1 (12)	53.1 (83)	205	30.2 (5)	388	17.3 (5)	1,085	10.2 (154)	1,088	12.6 (111)
5050	1,774	4.1 (71)	59.9 (89)	1,498	18.3 (64)	54.4 (100)	382	45.5 (35)	438	34.5 (46)	1,692	7.9 (113)	1,689	13.3 (130)
5051	2,667	4.6 (97)	58.8 (79)	2,278	17.0 (39)	54.0 (96)	547	46.4 (36)	730	38.5 (62)	2,569	7.7 (105)	2,569	11.3 (83)
5052	2,858	4.7 (105)	59.4 (85)	2,411	21.1 (105)	49.1 (30)	573	49.0 (44)	651	37.8 (58)	2,788	7.9 (113)	2,789	10.6 (64)
5055	660	2.6 (9)	53.3 (18)	269	19.7 (85)	49.8 (34)	97	*	294	38.8 (66)	1,175	9.3 (147)	1,178	4.1 (4)
5101	1,847	4.6 (97)	56.4 (40)	1,465	16.2 (25)	51.3 (57)	374	47.3 (39)	462	38.3 (60)	1,861	9.5 (149)	1,864	12.0 (99)
5102	1	*	*	1	*	*	0	*	0	*	1	*	1	*
5202	737	3.8 (58)	52.6 (15)	540	20.0 (90)	46.7 (12)	109	35.8 (11)	231	22.9 (13)	736	6.3 (60)	738	12.5 (110)
5203	25	*	*	19	*	*	3	*	6	*	24	*	24	*
5204	12	*	*	10	*	*	2	*	3	*	14	*	14	*
5601	982	3.3 (30)	57.9 (65)	447	22.6 (116)	50.3 (41)	159	34.6 (10)	376	31.9 (35)	1,078	5.5 (30)	1,079	11.3 (83)
5801	1,307	4.5 (92)	56.4 (40)	671	21.6 (113)	49.6 (32)	311	48.2 (40)	354	36.7 (54)	1,644	7.2 (86)	1,644	12.3 (104)
5803	123	4.1 (71)	48.8 (3)	84	*	*	24	*	41	*	122	9.8 (152)	122	10.7 (68)
5804	11	*	*	6	*	*	0	*	3	*	18	*	18	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5806	22	*	*	19	*	*	1	*	11	*	30	*	30	*
6101	29	*	*	23	*	*	0	*	8	*	28	*	28	*
6102	124	4.0 (67)	50.8 (8)	93	*	*	10	*	15	*	123	7.3 (91)	122	13.9 (132)
6103	4,057	4.2 (79)	60.2 (94)	3,055	20.8 (102)	50.0 (36)	569	43.8 (27)	429	31.0 (33)	4,009	8.0 (120)	4,013	13.2 (125)
6105	27	*	*	22	*	*	1	*	6	*	27	*	28	*
6106	17	*	*	12	*	*	2	*	4	*	17	*	17	*
6301	4,174	4.4 (89)	58.2 (68)	3,091	19.4 (81)	50.7 (48)	907	47.0 (38)	921	35.0 (49)	4,186	5.9 (44)	4,185	11.0 (75)
6312	2,225	3.1 (22)	60.9 (110)	1,650	17.4 (44)	52.2 (65)	486	48.4 (41)	691	41.5 (69)	2,243	4.5 (10)	2,243	8.2 (28)
6316	1,207	3.6 (50)	56.9 (48)	1,019	15.6 (20)	55.3 (109)	228	41.2 (21)	335	34.0 (45)	1,165	7.9 (113)	1,165	10.0 (52)
6320	97	*	*	35	*	*	18	*	40	*	115	8.7 (138)	115	3.5 (1)
6325	96	*	*	39	*	*	23	*	60	*	143	5.6 (34)	143	16.1 (153)
6330	255	4.3 (81)	57.3 (55)	90	*	*	58	*	60	*	333	5.4 (25)	334	11.1 (77)
6335	687	4.1 (71)	49.3 (4)	190	24.7 (125)	47.4 (16)	146	48.6 (42)	191	30.9 (32)	948	8.2 (127)	948	8.5 (35)
6340	42	*	*	15	*	*	10	*	12	*	56	*	55	*
6345	24	*	*	17	*	*	4	*	10	*	28	*	28	*
6360	264	4.9 (114)	58.3 (71)	70	*	*	64	*	81	*	354	6.8 (72)	353	11.6 (90)
7304	96	*	*	68	*	*	19	*	27	*	121	9.1 (144)	121	17.4 (157)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7306	2,098	4.3 (81)	60.7 (106)	1,488	20.7 (101)	50.2 (38)	423	59.8 (63)	403	38.7 (65)	2,438	6.3 (60)	2,437	11.5 (87)
7308	116	4.3 (81)	63.8 (134)	84	*	*	20	*	18	*	115	7.8 (110)	115	13.9 (132)
7310	13	*	*	8	*	*	4	*	8	*	35	*	35	*
7401	2,330	5.5 (132)	55.7 (35)	1,939	16.7 (33)	54.8 (106)	356	51.4 (48)	495	36.0 (52)	2,226	6.8 (72)	2,236	12.1 (100)
7601	360	6.4 (144)	57.2 (54)	124	15.3 (17)	53.2 (86)	97	*	131	26.0 (17)	622	6.1 (53)	623	11.1 (77)
7850	1,124	5.1 (122)	59.9 (89)	935	19.0 (77)	50.3 (41)	179	39.7 (20)	219	37.4 (57)	1,100	8.0 (120)	1,100	10.5 (62)
7852	27	*	*	22	*	*	5	*	5	*	28	*	28	*
8101	1,736	3.4 (40)	63.0 (129)	1,193	20.5 (98)	49.7 (33)	354	68.9 (73)	271	55.0 (88)	1,928	6.5 (67)	1,931	15.4 (150)
8102	357	2.5 (8)	68.1 (150)	175	19.4 (81)	48.6 (24)	53	*	48	*	416	6.7 (70)	416	10.6 (64)
8105	1,409	4.8 (110)	61.7 (117)	1,012	18.4 (66)	56.0 (115)	354	57.6 (61)	413	43.8 (74)	1,654	8.4 (133)	1,655	13.4 (131)
8106	281	3.9 (61)	55.5 (32)	184	15.2 (16)	56.0 (115)	100	54.0 (55)	97	*	488	5.3 (23)	487	8.4 (34)
8150	35	*	*	23	*	*	4	*	10	*	37	*	37	*
8151	3,072	3.7 (53)	59.0 (80)	1,965	20.1 (93)	48.9 (26)	606	52.8 (52)	898	38.1 (59)	3,030	7.9 (113)	3,032	12.8 (115)
8152	102	5.9 (141)	55.9 (36)	59	*	*	18	*	23	*	102	3.9 (3)	102	12.7 (113)
8153	10	*	*	10	*	*	0	*	4	*	10	*	10	*
8155	35	*	*	27	*	*	3	*	15	*	38	*	38	*
8157	9	*	*	4	*	*	1	*	3	*	7	*	7	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8201	2,237	5.1 (122)	57.5 (59)	1,288	17.8 (55)	54.3 (99)	600	46.7 (37)	820	29.4 (26)	2,609	7.2 (86)	2,609	10.2 (57)
8202	2,595	5.5 (132)	50.4 (7)	1,226	22.6 (116)	43.7 (7)	713	54.6 (56)	937	44.5 (77)	3,936	7.4 (96)	3,935	3.5 (1)
8203	411	3.4 (40)	58.2 (68)	320	13.8 (5)	55.6 (113)	91	*	163	28.8 (24)	450	7.3 (91)	450	12.4 (107)
8250	696	3.3 (30)	59.1 (83)	502	17.1 (41)	56.2 (119)	194	44.3 (31)	229	46.7 (83)	770	7.5 (101)	770	15.1 (147)
8252	115	1.7 (6)	72.2 (152)	82	*	*	27	*	27	*	125	7.2 (86)	125	15.2 (149)
8255	2,321	5.1 (122)	57.1 (53)	1,514	14.7 (9)	55.4 (110)	587	44.1 (29)	841	36.0 (52)	2,584	6.7 (70)	2,586	10.2 (57)
8257	1,257	4.8 (110)	54.1 (21)	541	25.7 (127)	40.5 (4)	329	52.9 (53)	614	46.4 (82)	1,865	9.7 (150)	1,865	6.3 (8)
8263	672	3.1 (22)	58.2 (68)	529	17.6 (47)	51.4 (59)	162	31.5 (7)	270	30.0 (29)	712	8.8 (140)	713	13.2 (125)
8264	503	2.6 (9)	54.3 (24)	350	15.1 (12)	53.1 (83)	141	37.6 (14)	226	29.6 (27)	659	6.8 (72)	660	10.5 (62)
8265	18	*	*	9	*	*	2	*	3	*	19	*	19	*
8266	25	*	*	10	*	*	5	*	7	*	37	*	37	*
8267	552	5.1 (122)	56.5 (42)	358	15.9 (22)	58.4 (128)	201	42.3 (24)	214	27.6 (21)	729	7.5 (101)	729	9.1 (40)
8901	1,040	3.5 (45)	62.4 (127)	687	18.2 (61)	53.0 (79)	202	67.8 (71)	132	63.6 (95)	1,100	5.7 (38)	1,101	14.3 (140)
8902	1,211	5.5 (132)	60.3 (96)	793	18.2 (61)	54.4 (100)	247	68.0 (72)	216	57.9 (91)	1,313	5.0 (15)	1,312	13.0 (119)
8903	1,179	3.3 (30)	65.2 (142)	740	20.3 (95)	53.0 (79)	239	74.5 (75)	183	62.8 (94)	1,274	4.9 (13)	1,273	16.4 (154)
8904	2,135	3.5 (45)	61.8 (120)	1,546	18.0 (56)	53.0 (79)	530	72.1 (74)	350	58.3 (92)	2,331	5.4 (25)	2,332	14.2 (137)
8905	1,279	3.4 (40)	56.8 (46)	776	18.2 (61)	50.0 (36)	273	56.0 (57)	291	44.0 (75)	1,566	7.8 (110)	1,567	10.6 (64)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8908	27	*	*	18	*	*	6	*	4	*	31	*	31	*
8909	419	5.5 (132)	64.9 (140)	283	19.4 (81)	56.9 (121)	101	60.4 (64)	67	*	456	6.4 (66)	457	13.1 (123)
9012	6	*	*	4	*	*	0	*	0	*	6	*	6	*
9013	1	*	*	1	*	*	0	*	0	*	1	*	1	*
9016	46	*	*	32	*	*	5	*	6	*	47	*	47	*
9017	4	*	*	3	*	*	0	*	2	*	5	*	5	*
9018	49	*	*	36	*	*	3	*	2	*	51	*	51	*
9041	2	*	*	1	*	*	0	*	0	*	2	*	2	*
9045	1,436	2.2 (7)	64.5 (137)	1,059	23.3 (120)	48.3 (22)	300	45.0 (34)	220	34.5 (46)	1,755	6.6 (68)	1,756	7.1 (20)
9050	852	4.0 (67)	62.4 (127)	584	23.3 (120)	51.2 (54)	182	66.5 (70)	126	58.7 (93)	967	6.0 (50)	966	11.8 (96)
9051	1,358	2.6 (9)	64.7 (138)	814	15.8 (21)	47.2 (14)	176	41.5 (22)	237	32.1 (37)	1,626	8.2 (127)	1,627	6.3 (8)
9053	1,099	4.6 (97)	61.7 (117)	921	16.6 (32)	55.7 (114)	189	61.4 (67)	134	49.3 (86)	1,101	5.1 (17)	1,101	13.0 (119)
9055	47	*	*	45	*	*	3	*	2	*	50	*	50	*
9058	727	4.5 (92)	61.9 (122)	521	20.3 (95)	48.9 (26)	175	56.0 (57)	195	40.0 (68)	1,048	8.0 (120)	1,049	13.1 (123)
9066	1,001	3.1 (22)	67.2 (149)	731	16.8 (35)	54.9 (107)	213	60.6 (65)	161	57.1 (90)	1,127	6.2 (56)	1,126	14.2 (137)
9068	993	3.3 (30)	66.9 (147)	811	24.0 (123)	52.3 (68)	187	53.5 (54)	101	41.6 (70)	961	5.7 (38)	961	10.9 (72)
9069	2,762	3.7 (53)	60.5 (101)	2,306	18.9 (76)	52.7 (75)	573	49.4 (46)	296	42.6 (71)	2,906	5.5 (30)	2,907	11.6 (90)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9070	931	4.5 (92)	56.9 (48)	609	20.0 (90)	53.4 (90)	207	65.2 (69)	134	56.0 (89)	1,148	5.7 (38)	1,149	14.2 (137)
9072	1,729	4.2 (79)	61.7 (117)	1,241	21.2 (108)	51.5 (60)	382	56.0 (57)	230	46.1 (81)	1,871	5.3 (23)	1,871	14.8 (143)
9080	2,345	4.9 (114)	61.4 (115)	1,627	18.1 (57)	54.4 (100)	540	65.0 (68)	469	47.8 (85)	2,726	5.6 (34)	2,726	11.3 (83)
9081	43	*	*	28	*	*	7	*	4	*	42	*	42	*
9082	116	4.3 (81)	65.5 (144)	65	*	*	22	*	14	*	111	6.3 (60)	112	20.5 (159)
9083	53	*	*	50	*	*	3	*	1	*	55	*	55	*
9084	23	*	*	23	*	*	0	*	1	*	25	*	25	*
9085	48	*	*	17	*	*	22	*	4	*	45	*	45	*
9086	34	*	*	20	*	*	15	*	4	*	30	*	30	*
Michigan	**143,426**	**4.3**	**58.8**	**93,551**	**19.2**	**51.4**	**28,148**	**47.8**	**35,457**	**36.4**	**160,515**	**7.0**	**160,593**	**11.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	113	85.8 (61)	8.0 (80)	116	31.9 (97)	108	24.1 (54)	92	*
0102	501	86.4 (52)	8.2 (85)	532	38.3 (57)	486	20.0 (84)	439	90.4 (52)
0103	421	77.7 (120)	13.5 (123)	445	36.9 (64)	398	21.9 (68)	369	90.5 (51)
0104	151	82.8 (82)	7.9 (79)	159	35.8 (71)	142	23.2 (62)	128	90.6 (50)
0105	7	*	*	7	*	5	*	5	*
0250	137	75.2 (131)	19.0 (148)	142	45.8 (16)	100	33.0 (18)	112	91.1 (42)
0251	91	*	*	95	*	81	*	76	*
0252	113	69.9 (155)	20.4 (153)	121	41.3 (32)	112	20.5 (82)	100	83.0 (123)
0253	166	82.5 (86)	13.9 (125)	170	59.4 (1)	161	31.1 (26)	152	88.2 (81)
0401	470	86.2 (56)	8.5 (89)	505	38.2 (58)	474	27.2 (33)	424	92.9 (16)
0402	368	83.7 (71)	8.2 (85)	379	35.6 (74)	340	22.1 (66)	303	88.8 (77)
0403	128	82.8 (82)	7.0 (68)	138	33.3 (90)	121	24.8 (47)	111	91.0 (43)
0404	143	83.2 (77)	11.2 (114)	147	45.6 (17)	139	25.9 (42)	122	91.0 (43)
0501	514	92.6 (6)	1.6 (12)	545	33.8 (86)	497	26.6 (40)	424	88.9 (76)
0503	110	86.4 (52)	7.3 (72)	114	33.3 (90)	110	31.8 (20)	91	*
0504	490	88.8 (28)	6.7 (66)	513	31.6 (101)	473	31.3 (23)	401	90.0 (58)
0505	333	83.5 (73)	10.8 (112)	347	39.5 (47)	327	27.8 (32)	291	90.7 (47)
0506	206	82.0 (93)	14.1 (127)	215	48.4 (8)	195	23.1 (63)	175	80.6 (130)
0507	84	*	*	85	*	77	*	50	*
0508	30	*	*	31	*	29	*	21	*
0509	8	*	*	8	*	6	*	5	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	1,096	84.9 (66)	7.6 (76)	1,160	38.9 (51)	999	29.4 (31)	944	92.8 (17)
0651	576	73.4 (145)	18.1 (146)	615	43.7 (22)	544	16.4 (114)	521	89.4 (65)
0652	298	79.9 (111)	14.8 (130)	310	35.2 (76)	278	24.5 (52)	253	85.8 (103)
0653	427	74.5 (139)	20.4 (153)	450	47.1 (12)	411	26.8 (35)	381	89.2 (69)
0654	350	76.9 (125)	16.9 (141)	385	44.9 (19)	352	21.9 (68)	330	80.0 (133)
0655	361	82.5 (86)	8.0 (80)	385	47.0 (13)	348	23.9 (56)	325	89.2 (69)
0657	63	*	*	64	*	59	*	56	*
0701	414	78.7 (116)	16.7 (139)	420	35.7 (73)	398	43.2 (8)	357	89.4 (65)
0702	58	*	*	68	*	64	*	57	*
0703	196	92.9 (5)	1.5 (10)	229	35.8 (71)	206	31.1 (26)	178	93.3 (10)
0704	45	*	*	54	*	51	*	46	*
0705	123	87.0 (46)	8.9 (92)	135	30.4 (105)	122	63.9 (3)	106	91.5 (36)
0706	5	*	*	5	*	3	*	2	*
0850	858	90.2 (19)	2.3 (16)	939	28.6 (115)	815	21.3 (75)	708	93.2 (11)
0851	736	92.3 (7)	4.6 (46)	792	36.0 (68)	679	20.8 (80)	633	93.8 (7)
0901	1,621	74.2 (143)	19.6 (150)	1,756	44.0 (21)	1,635	17.4 (105)	1,496	83.8 (116)
0902	44	*	*	46	*	37	*	33	*
1051	1,377	73.1 (148)	18.1 (146)	1,432	30.4 (105)	1,283	23.9 (56)	1,000	88.2 (81)
1101	1,700	75.2 (131)	9.4 (97)	1,872	27.5 (119)	1,566	12.9 (134)	1,406	86.3 (100)
1102	764	74.5 (139)	14.1 (127)	871	36.9 (64)	710	18.5 (93)	673	95.1 (2)
1106	68	*	*	77	*	64	*	56	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1250	813	80.4 (109)	10.8 (112)	1,057	40.8 (36)	935	24.6 (50)	889	80.9 (128)
1251	809	76.5 (127)	13.3 (121)	867	39.1 (50)	776	18.3 (96)	695	83.5 (120)
1252	524	74.2 (143)	15.1 (131)	575	39.7 (45)	537	17.5 (103)	478	89.5 (63)
1253	791	71.7 (151)	20.0 (151)	880	36.6 (66)	791	17.2 (109)	714	88.0 (84)
1350	2,324	90.2 (19)	3.8 (40)	2,598	37.3 (63)	1,650	19.2 (90)	2,041	78.7 (137)
1351	137	87.6 (41)	8.0 (80)	142	28.9 (111)	124	21.8 (70)	103	92.2 (28)
1352	349	87.4 (45)	5.7 (59)	379	43.0 (23)	333	3.0 (157)	313	88.8 (77)
1401	231	77.1 (123)	16.5 (136)	250	33.2 (93)	143	30.1 (30)	185	83.8 (116)
1402	133	71.4 (152)	16.5 (136)	145	29.7 (108)	101	34.7 (15)	97	*
1403	27	*	*	26	*	17	*	22	*
1404	0	*	*	1	*	0	*	1	*
1501	717	83.8 (70)	11.2 (114)	822	41.4 (31)	579	13.3 (130)	647	80.4 (131)
1503	1,009	87.0 (46)	6.6 (65)	1,117	42.3 (25)	938	45.2 (6)	931	78.1 (139)
1504	2,038	90.8 (12)	3.4 (33)	2,301	35.1 (77)	1,509	23.0 (64)	1,672	71.8 (146)
1505	435	90.6 (14)	4.6 (46)	469	32.4 (94)	333	32.7 (19)	333	88.0 (84)
1506	562	74.4 (141)	21.4 (156)	618	40.8 (36)	541	7.8 (152)	492	90.4 (52)
1507	593	90.6 (14)	4.7 (49)	649	40.4 (40)	475	26.7 (37)	508	90.0 (58)
1508	77	*	*	113	46.9 (14)	79	*	79	*
1509	384	84.1 (69)	9.9 (105)	480	46.0 (15)	314	26.8 (35)	387	75.7 (144)
1511	691	87.8 (37)	5.5 (57)	749	36.4 (67)	520	25.0 (46)	579	86.7 (95)
1512	359	87.5 (43)	8.4 (87)	382	31.7 (99)	236	33.1 (17)	286	87.1 (91)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1513	72	*	*	77	*	62	*	59	*
1514	160	94.4 (2)	1.3 (7)	169	21.3 (139)	122	27.0 (34)	101	90.1 (57)
1701	650	87.8 (37)	9.7 (101)	686	48.5 (7)	617	56.4 (4)	599	86.1 (102)
1950	574	83.1 (80)	10.3 (107)	615	40.3 (42)	525	20.0 (84)	487	89.7 (62)
1952	133	85.0 (65)	9.8 (104)	135	33.3 (90)	117	21.4 (73)	108	92.6 (22)
1953	308	88.0 (35)	8.1 (83)	323	34.4 (81)	293	21.8 (70)	251	93.2 (11)
1954	555	82.7 (84)	12.1 (117)	592	40.9 (35)	540	26.7 (37)	492	89.4 (65)
1955	83	*	*	87	*	76	*	70	*
1957	528	87.5 (43)	5.1 (53)	558	33.5 (89)	480	23.3 (60)	426	89.0 (75)
1963	10	*	*	11	*	6	*	1	*
1964	30	*	*	34	*	31	*	24	*
2150	631	90.2 (19)	3.6 (38)	692	45.1 (18)	639	25.7 (43)	581	91.0 (43)
2151	209	86.1 (57)	9.6 (99)	249	38.6 (55)	201	18.4 (94)	189	83.6 (119)
2251	420	88.1 (32)	9.3 (94)	453	42.6 (24)	418	24.6 (50)	379	91.3 (39)
2252	188	80.9 (104)	17.0 (142)	210	57.6 (2)	182	64.8 (2)	187	85.0 (111)
2554	81	*	*	84	*	73	*	66	*
2555	7,764	86.4 (52)	3.4 (33)	8,256	25.4 (124)	6,181	10.9 (142)	6,095	89.2 (69)
2556	196	78.1 (119)	2.6 (19)	196	6.6 (161)	152	17.1 (110)	50	*
2557	6	*	*	9	*	7	*	8	*
2558	3	*	*	4	*	4	*	4	*
2559	9	*	*	9	*	8	*	9	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2560	2	*	*	2	*	2	*	2	*
2562	1	*	*	2	*	2	*	2	*
2563	5	*	*	3	*	0	*	0	*
3001	1,089	73.0 (149)	9.3 (94)	1,169	35.5 (75)	1,030	7.1 (154)	964	79.9 (134)
3002	249	85.9 (60)	1.2 (6)	262	40.1 (43)	241	9.5 (147)	231	81.0 (127)
3004	136	90.4 (17)	2.9 (21)	152	50.0 (6)	144	19.4 (89)	140	69.3 (148)
3007	24	*	*	30	*	26	*	25	*
3009	2	*	*	10	*	6	*	6	*
3201	469	81.0 (103)	13.6 (124)	499	41.7 (28)	425	35.3 (14)	380	87.9 (87)
3301	4,074	82.0 (93)	9.5 (98)	4,391	28.8 (114)	3,833	10.1 (143)	3,328	89.9 (60)
3304	107	83.2 (77)	11.2 (114)	108	40.7 (38)	106	12.3 (136)	86	*
3305	142	86.6 (48)	7.7 (77)	146	33.6 (87)	134	6.7 (155)	107	94.4 (4)
3306	251	82.5 (86)	9.6 (99)	262	38.2 (58)	242	7.9 (150)	205	92.7 (19)
3308	72	*	*	73	*	64	*	48	*
3309	37	*	*	39	*	35	*	28	*
3312	392	83.4 (75)	7.1 (70)	415	27.0 (120)	339	11.8 (140)	337	90.2 (56)
3550	71	*	*	74	*	65	*	64	*
3551	27	*	*	28	*	23	*	21	*
3552	47	*	*	53	*	47	*	47	*
3701	1,039	86.6 (48)	6.9 (67)	1,129	35.9 (69)	1,032	19.9 (86)	894	93.1 (13)
3850	3,009	85.8 (61)	7.5 (75)	3,254	40.7 (38)	2,899	14.9 (123)	2,710	85.4 (107)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3901	534	80.9 (104)	5.2 (54)	566	25.6 (121)	454	10.1 (143)	356	77.5 (141)
3902	248	81.5 (97)	9.7 (101)	263	37.6 (62)	237	15.2 (121)	203	79.8 (135)
3903	9	*	*	10	*	9	*	0	*
3904	3	*	*	3	*	3	*	3	*
3905	2	*	*	2	*	1	*	0	*
3950	1,850	81.1 (102)	10.7 (111)	2,042	34.6 (80)	1,859	17.9 (101)	1,619	86.2 (101)
3954	4	*	*	5	*	5	*	3	*
4001	924	73.3 (146)	17.2 (143)	1,021	52.4 (5)	980	14.6 (124)	949	74.0 (145)
4002	659	76.6 (126)	17.5 (145)	713	48.1 (9)	676	18.2 (97)	645	82.9 (124)
4003	5	*	*	5	*	5	*	5	*
4004	5	*	*	5	*	5	*	5	*
4103	53	*	*	59	*	53	*	54	*
4106	2,211	82.2 (90)	8.1 (83)	2,443	31.4 (103)	2,156	7.9 (150)	1,976	87.6 (89)
4107	3,127	83.5 (73)	7.8 (78)	3,341	27.7 (118)	2,404	18.9 (91)	2,535	91.8 (32)
4109	1,039	79.8 (112)	15.8 (134)	1,109	31.7 (99)	1,006	21.0 (79)	825	93.6 (8)
4112	552	52.4 (159)	0.4 (3)	571	28.9 (111)	551	91.5 (1)	524	92.0 (30)
4113	630	88.3 (30)	0.2 (2)	674	32.2 (95)	477	18.4 (94)	632	80.9 (128)
4115	3,376	85.8 (61)	7.0 (68)	3,605	28.1 (117)	3,007	18.1 (98)	2,769	91.2 (40)
4351	577	91.5 (8)	3.3 (30)	611	41.6 (29)	225	55.6 (5)	519	92.5 (23)
4352	164	72.0 (150)	16.5 (136)	175	48.0 (10)	156	15.4 (119)	151	86.8 (93)
4353	412	77.4 (121)	17.2 (143)	460	40.0 (44)	410	18.8 (92)	369	81.6 (126)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4354	747	84.5 (67)	8.8 (90)	782	37.7 (61)	481	31.8 (20)	627	90.3 (55)
4355	353	88.1 (32)	5.7 (59)	375	30.7 (104)	338	35.5 (13)	305	92.5 (23)
4356	455	80.2 (110)	13.0 (120)	491	39.3 (49)	444	30.2 (29)	410	91.2 (40)
4358	130	70.8 (154)	20.8 (155)	156	37.8 (60)	130	26.2 (41)	120	75.8 (143)
4359	463	76.2 (129)	16.4 (135)	487	41.3 (32)	416	23.8 (59)	386	89.1 (73)
4360	233	63.9 (157)	28.3 (159)	240	44.2 (20)	172	26.7 (37)	196	80.1 (132)
4361	238	86.6 (48)	8.8 (90)	249	47.4 (11)	222	24.8 (47)	208	90.4 (52)
4362	285	74.7 (137)	20.0 (151)	298	41.6 (29)	268	21.3 (75)	251	85.7 (106)
4363	882	75.9 (130)	16.7 (139)	916	42.0 (26)	654	31.2 (24)	744	85.8 (103)
4401	1,084	82.1 (92)	12.7 (119)	1,139	32.2 (95)	1,053	18.1 (98)	887	91.9 (31)
4501	973	71.4 (152)	19.3 (149)	1,014	38.8 (52)	917	25.5 (44)	792	91.5 (36)
4601	1,392	86.3 (55)	6.3 (63)	1,516	29.7 (108)	1,345	22.0 (67)	1,156	88.4 (80)
4602	30	*	*	30	*	22	*	17	*
4701	1,086	90.7 (13)	3.9 (41)	1,169	23.4 (131)	909	34.2 (16)	793	91.6 (35)
5050	1,673	89.4 (25)	2.5 (18)	1,807	19.7 (143)	1,161	21.4 (73)	1,289	92.7 (19)
5051	2,543	90.1 (22)	3.1 (26)	2,729	21.0 (140)	1,965	19.9 (86)	1,982	91.8 (32)
5052	2,764	86.5 (51)	3.9 (41)	2,951	22.3 (134)	1,955	17.4 (105)	2,194	92.3 (26)
5055	1,147	82.2 (90)	15.2 (132)	1,210	56.5 (3)	571	23.3 (60)	1,064	78.2 (138)
5101	1,842	89.6 (24)	4.9 (51)	1,977	25.2 (127)	1,518	25.4 (45)	1,454	93.1 (13)
5102	1	*	*	1	*	1	*	0	*
5202	732	88.5 (29)	5.9 (62)	819	38.8 (52)	593	36.8 (12)	650	92.8 (17)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5203	24	*	*	25	*	18	*	24	*
5204	14	*	*	13	*	10	*	12	*
5601	1,075	84.2 (68)	12.1 (117)	1,121	33.6 (87)	1,036	31.5 (22)	887	96.4 (1)
5801	1,638	82.7 (84)	10.0 (106)	1,775	34.1 (84)	1,622	17.3 (108)	1,413	86.7 (95)
5803	121	85.1 (64)	7.4 (74)	130	24.6 (128)	121	17.4 (105)	88	*
5804	18	*	*	19	*	14	*	14	*
5806	30	*	*	29	*	15	*	4	*
6101	28	*	*	29	*	26	*	21	*
6102	123	81.3 (98)	10.6 (110)	122	41.8 (27)	110	24.5 (52)	98	*
6103	3,952	83.2 (77)	8.4 (87)	4,244	39.4 (48)	3,761	16.5 (113)	3,447	79.6 (136)
6105	28	*	*	28	*	24	*	18	*
6106	17	*	*	17	*	11	*	2	*
6301	4,164	86.0 (59)	5.5 (57)	4,454	25.5 (122)	3,598	16.0 (117)	3,367	89.2 (69)
6312	2,207	89.4 (25)	3.3 (30)	2,364	20.7 (141)	1,868	18.0 (100)	1,720	90.9 (46)
6316	1,158	87.7 (40)	4.3 (44)	1,234	25.4 (124)	1,049	21.8 (70)	888	92.7 (19)
6320	115	91.3 (10)	5.2 (54)	123	30.1 (107)	45	*	88	*
6325	142	91.5 (8)	3.5 (36)	144	21.5 (138)	108	24.1 (54)	93	*
6330	333	78.7 (116)	10.5 (109)	361	40.4 (40)	288	12.5 (135)	331	83.1 (122)
6335	949	87.6 (41)	10.4 (108)	994	38.7 (54)	236	19.9 (86)	794	89.5 (63)
6340	56	*	*	58	*	35	*	42	*
6345	27	*	*	28	*	19	*	18	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6360	352	88.1 (32)	4.3 (44)	373	19.0 (147)	176	23.9 (56)	266	84.6 (112)
7304	119	74.8 (136)	14.3 (129)	127	31.5 (102)	113	24.8 (47)	101	93.1 (13)
7306	2,393	74.3 (142)	14.0 (126)	2,593	34.8 (79)	2,112	14.0 (126)	2,001	85.3 (108)
7308	113	80.5 (107)	9.7 (101)	126	28.6 (115)	119	13.4 (129)	106	93.4 (9)
7310	35	*	*	38	*	36	*	31	*
7401	2,234	66.3 (156)	24.9 (157)	2,449	39.6 (46)	2,124	16.9 (112)	2,015	86.4 (99)
7601	610	82.5 (86)	9.3 (94)	672	35.1 (77)	448	43.1 (9)	531	84.2 (114)
7850	1,079	77.0 (124)	15.5 (133)	1,158	38.4 (56)	1,061	14.1 (125)	932	87.8 (88)
7852	27	*	*	29	*	25	*	24	*
8101	1,919	77.2 (122)	3.1 (26)	2,017	18.5 (150)	1,289	15.7 (118)	1,403	88.7 (79)
8102	409	81.2 (100)	6.4 (64)	431	41.1 (34)	293	15.0 (122)	367	76.6 (142)
8105	1,652	82.9 (81)	4.2 (43)	1,758	21.8 (137)	1,575	17.6 (102)	1,264	92.3 (26)
8106	480	75.0 (134)	4.6 (46)	510	18.8 (149)	371	9.2 (148)	365	85.8 (103)
8150	37	*	*	40	*	38	*	36	*
8151	3,024	90.3 (18)	3.2 (28)	3,318	31.9 (97)	3,042	17.5 (103)	2,611	90.7 (47)
8152	102	89.2 (27)	0.0 (1)	108	34.3 (82)	106	7.5 (153)	103	94.2 (5)
8153	10	*	*	14	*	10	*	8	*
8155	38	*	*	38	*	34	*	30	*
8157	7	*	*	9	*	9	*	8	*
8201	2,603	90.6 (14)	1.8 (13)	2,791	19.7 (143)	2,089	30.5 (28)	1,985	91.8 (32)
8202	3,860	88.0 (35)	9.0 (93)	4,069	34.1 (84)	2,944	13.2 (131)	3,258	81.8 (125)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8203	449	93.8 (3)	0.7 (4)	480	19.0 (147)	301	21.3 (75)	331	94.0 (6)
8250	763	83.6 (72)	2.1 (15)	816	15.6 (158)	554	22.7 (65)	552	88.0 (84)
8252	124	75.0 (134)	4.8 (50)	128	17.2 (153)	97	*	92	*
8255	2,562	89.9 (23)	1.5 (10)	2,752	19.4 (145)	1,789	38.4 (11)	1,960	92.1 (29)
8257	1,843	95.0 (1)	1.3 (7)	1,953	29.4 (110)	1,775	12.1 (137)	1,530	85.2 (109)
8263	709	93.1 (4)	1.3 (7)	749	15.6 (158)	504	31.2 (24)	526	94.5 (3)
8264	656	91.2 (11)	3.0 (25)	711	20.4 (142)	587	16.4 (114)	515	92.4 (25)
8265	19	*	*	21	*	19	*	21	*
8266	36	*	*	41	*	38	*	40	*
8267	727	87.8 (37)	3.2 (28)	798	17.0 (154)	690	44.5 (7)	551	88.2 (81)
8901	1,087	79.2 (113)	3.3 (30)	1,148	24.2 (129)	721	13.0 (133)	881	83.5 (120)
8902	1,298	81.9 (96)	2.9 (21)	1,386	22.3 (134)	965	15.3 (120)	1,024	87.5 (90)
8903	1,261	78.5 (118)	3.5 (36)	1,326	22.5 (132)	857	13.9 (127)	988	83.7 (118)
8904	2,310	82.0 (93)	2.8 (20)	2,435	18.5 (150)	1,435	17.0 (111)	1,748	89.1 (73)
8905	1,559	86.1 (57)	5.4 (56)	1,620	28.9 (111)	1,177	21.3 (75)	1,226	86.6 (97)
8908	31	*	*	30	*	26	*	28	*
8909	451	80.5 (107)	1.8 (13)	474	15.8 (157)	365	16.4 (114)	327	89.3 (68)
9012	6	*	*	7	*	7	*	7	*
9013	1	*	*	1	*	1	*	1	*
9016	47	*	*	52	*	52	*	52	*
9017	5	*	*	6	*	6	*	6	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9018	50	*	*	55	*	54	*	54	*
9041	2	*	*	2	*	2	*	2	*
9045	1,755	73.3 (146)	13.3 (121)	1,824	34.2 (83)	400	40.0 (10)	1,450	87.0 (92)
9050	949	76.3 (128)	7.3 (72)	991	25.4 (124)	783	20.6 (81)	748	84.5 (113)
9051	1,556	63.9 (157)	27.7 (158)	1,701	52.7 (4)	1,533	13.8 (128)	1,538	70.2 (147)
9053	1,099	74.7 (137)	7.1 (70)	1,159	21.9 (136)	943	12.0 (138)	829	77.6 (140)
9055	50	*	*	49	*	39	*	9	*
9058	1,047	88.2 (31)	5.0 (52)	1,078	24.0 (130)	822	4.1 (156)	798	86.5 (98)
9066	1,038	83.3 (76)	2.3 (16)	1,163	16.6 (156)	771	8.9 (149)	781	89.9 (60)
9068	962	78.9 (115)	2.9 (21)	1,007	25.5 (122)	766	13.2 (131)	763	86.8 (93)
9069	2,897	79.0 (114)	3.4 (33)	3,005	17.0 (154)	1,796	9.6 (146)	2,045	91.4 (38)
9070	1,134	75.2 (131)	5.8 (61)	1,177	19.4 (145)	733	20.5 (82)	814	85.1 (110)
9072	1,855	80.9 (104)	2.9 (21)	1,945	17.9 (152)	1,451	9.7 (145)	1,404	84.2 (114)
9080	2,714	81.2 (100)	3.7 (39)	2,815	22.5 (132)	1,941	11.4 (141)	2,064	90.7 (47)
9081	43	*	*	49	*	39	*	39	*
9082	112	81.3 (98)	0.9 (5)	122	14.8 (160)	108	12.0 (138)	78	*
9083	55	*	*	56	*	41	*	8	*
9084	25	*	*	25	*	8	*	0	*
9085	43	*	*	48	*	44	*	48	*
9086	30	*	*	36	*	36	*	36	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Michigan	**158,822**	**83.2**	**7.7**	**170,497**	**30.9**	**133,582**	**19.4**	**131,162**	**87.3**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	109	50.5 (151)	112	32.1 (149)	45	*	92	*	112	3.6 (101)
0102	487	50.9 (153)	508	34.1 (155)	213	27.7 (94)	439	21.0 (135)	508	3.9 (112)
0103	412	46.4 (139)	425	31.5 (146)	161	28.6 (91)	369	21.7 (137)	425	6.8 (149)
0104	144	49.3 (149)	151	33.8 (154)	62	*	128	17.2 (118)	151	4.6 (129)
0105	5	*	7	*	1	*	5	*	7	*
0250	121	51.2 (154)	137	30.7 (142)	55	*	112	19.6 (129)	137	0.7 (7)
0251	82	*	91	*	45	*	76	*	91	*
0252	114	48.2 (146)	113	32.7 (152)	44	*	100	16.0 (109)	113	5.3 (135)
0253	161	46.0 (137)	165	31.5 (146)	67	*	152	17.1 (117)	165	4.8 (132)
0401	477	46.8 (141)	472	26.7 (131)	180	36.1 (60)	424	20.5 (131)	472	3.0 (83)
0402	341	48.4 (147)	367	28.3 (136)	141	38.3 (52)	303	20.5 (131)	368	3.8 (108)
0403	121	52.1 (155)	131	27.5 (134)	59	*	111	25.2 (145)	131	9.2 (156)
0404	139	45.3 (135)	143	25.9 (129)	54	*	122	22.1 (141)	143	6.3 (145)
0501	499	43.3 (129)	510	29.0 (138)	164	28.7 (89)	424	18.6 (125)	510	6.3 (145)
0503	111	49.5 (150)	110	37.3 (158)	43	*	91	*	110	7.3 (153)
0504	473	37.0 (104)	490	17.8 (88)	148	41.2 (42)	401	10.5 (67)	490	2.2 (54)
0505	327	39.1 (111)	331	20.8 (106)	116	40.5 (44)	291	13.1 (87)	332	4.2 (122)
0506	197	52.8 (157)	205	34.6 (157)	88	*	175	27.4 (147)	205	7.3 (153)
0507	77	*	84	*	20	*	50	*	84	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	29	*	31	*	6	*	21	*	31	*
0509	6	*	8	*	2	*	5	*	8	*
0650	1,079	38.5 (108)	1,093	21.9 (112)	342	38.6 (50)	944	12.6 (84)	1,094	3.0 (83)
0651	568	46.7 (140)	580	32.2 (150)	233	21.5 (105)	521	21.7 (137)	580	4.5 (125)
0652	278	50.7 (152)	301	24.6 (122)	124	35.5 (61)	252	20.6 (133)	301	6.0 (142)
0653	420	47.1 (143)	427	29.7 (140)	170	25.9 (99)	380	22.6 (142)	427	6.1 (143)
0654	357	45.9 (136)	348	27.6 (135)	135	30.4 (85)	330	21.8 (139)	348	4.3 (124)
0655	356	47.2 (144)	366	30.6 (141)	144	31.9 (79)	325	20.6 (133)	366	6.3 (145)
0657	58	*	63	*	18	*	56	*	63	*
0701	401	29.4 (70)	411	17.8 (88)	107	29.9 (87)	355	6.2 (21)	411	1.2 (19)
0702	64	*	59	*	22	*	57	*	59	*
0703	205	47.3 (145)	198	37.4 (159)	70	*	178	14.6 (96)	199	4.5 (125)
0704	51	*	46	*	16	*	46	*	46	*
0705	124	40.3 (116)	123	28.5 (137)	43	*	106	16.0 (109)	123	3.3 (93)
0706	3	*	5	*	2	*	2	*	5	*
0850	835	38.8 (109)	875	23.2 (116)	256	35.2 (63)	708	8.3 (42)	875	2.1 (48)
0851	734	33.0 (88)	738	20.5 (102)	198	31.3 (81)	631	9.5 (60)	738	1.4 (23)
0901	1,635	44.0 (133)	1,660	25.3 (126)	647	34.8 (65)	1,496	13.0 (86)	1,661	3.4 (99)
0902	38	*	44	*	14	*	33	*	44	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1051	1,286	35.9 (98)	1,385	21.9 (112)	375	34.7 (66)	1,000	8.6 (46)	1,385	3.2 (88)
1101	1,616	28.7 (68)	1,732	13.2 (56)	372	43.8 (36)	1,405	11.1 (73)	1,734	3.2 (88)
1102	765	32.7 (86)	776	17.4 (83)	189	39.2 (48)	673	15.2 (102)	776	3.1 (86)
1106	66	*	70	*	15	*	56	*	70	*
1250	995	33.1 (89)	979	21.3 (110)	275	28.7 (89)	889	13.2 (89)	979	3.8 (108)
1251	783	43.7 (131)	814	29.2 (139)	281	23.5 (103)	695	16.7 (115)	816	5.8 (140)
1252	539	33.8 (92)	542	19.2 (95)	140	30.0 (86)	478	15.1 (101)	542	3.7 (103)
1253	794	37.4 (106)	812	20.6 (103)	240	33.8 (71)	714	13.6 (90)	813	3.2 (88)
1350	2,299	32.5 (84)	2,384	19.0 (94)	653	29.9 (87)	2,051	12.6 (84)	2,395	3.3 (93)
1351	128	35.9 (98)	140	16.4 (78)	37	*	103	7.8 (36)	140	2.1 (48)
1352	335	41.2 (120)	354	17.8 (88)	123	27.6 (95)	313	3.2 (6)	354	1.1 (15)
1401	220	46.8 (141)	235	32.8 (153)	90	*	186	15.6 (106)	235	5.1 (133)
1402	129	29.5 (71)	134	9.0 (36)	29	*	97	*	134	1.5 (26)
1403	26	*	26	*	13	*	22	*	27	*
1404	1	*	0	*	0	*	1	*	0	*
1501	717	20.4 (26)	725	11.0 (43)	125	24.0 (102)	647	3.6 (8)	722	1.5 (26)
1503	1,010	26.8 (62)	1,034	14.9 (68)	236	24.2 (101)	932	6.8 (26)	1,034	1.8 (41)
1504	1,998	22.4 (39)	2,115	9.8 (39)	412	48.3 (20)	1,700	5.6 (18)	2,120	1.7 (33)
1505	419	25.8 (56)	443	14.7 (64)	98	*	335	7.5 (33)	444	2.7 (73)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1506	560	24.1 (48)	567	14.3 (62)	117	23.1 (104)	494	3.4 (7)	566	1.6 (31)
1507	590	30.8 (78)	596	15.3 (72)	161	41.6 (40)	513	7.4 (32)	599	2.0 (47)
1508	88	*	85	*	4	*	79	*	85	*
1509	418	24.4 (49)	394	15.5 (74)	89	*	392	7.9 (37)	394	2.5 (64)
1511	666	35.7 (97)	702	20.7 (105)	207	32.4 (77)	583	13.7 (91)	704	2.6 (69)
1512	330	31.5 (82)	360	18.6 (92)	102	33.3 (72)	286	3.8 (12)	362	1.7 (33)
1513	65	*	72	*	16	*	59	*	72	*
1514	150	19.3 (24)	161	8.7 (35)	26	*	100	3.0 (5)	160	0.0 (1)
1701	661	42.2 (127)	646	32.0 (148)	236	17.4 (106)	603	28.2 (148)	644	18.0 (159)
1950	555	43.4 (130)	582	24.9 (123)	200	37.5 (58)	488	19.9 (130)	582	4.5 (125)
1952	126	44.4 (134)	133	25.6 (127)	43	*	107	15.9 (107)	133	3.0 (83)
1953	302	42.1 (126)	309	23.0 (115)	102	45.1 (30)	251	12.0 (81)	309	3.9 (112)
1954	555	41.8 (124)	558	24.4 (121)	194	37.6 (57)	492	10.8 (71)	559	2.5 (64)
1955	78	*	84	*	28	*	70	*	84	*
1957	514	37.4 (106)	531	20.3 (101)	163	46.6 (26)	427	8.7 (47)	532	1.7 (33)
1963	6	*	10	*	0	*	1	*	10	*
1964	32	*	31	*	8	*	24	*	31	*
2150	644	43.9 (132)	631	26.6 (130)	221	27.1 (97)	581	13.1 (87)	631	1.9 (45)
2151	222	52.3 (156)	214	34.1 (155)	83	*	189	15.9 (107)	214	3.3 (93)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2251	436	41.3 (122)	417	24.9 (123)	144	31.3 (81)	379	7.1 (28)	419	1.2 (19)
2252	198	46.0 (137)	182	31.3 (144)	75	*	186	14.5 (95)	182	3.8 (108)
2554	73	*	80	*	20	*	66	*	80	*
2555	7,163	30.1 (73)	7,876	15.8 (75)	1,861	38.2 (54)	6,099	10.6 (69)	7,886	2.3 (58)
2556	153	17.6 (18)	199	16.1 (76)	10	*	49	*	198	5.6 (138)
2557	8	*	7	*	0	*	8	*	7	*
2558	4	*	4	*	0	*	4	*	4	*
2559	9	*	9	*	2	*	9	*	9	*
2560	2	*	2	*	1	*	2	*	2	*
2562	2	*	1	*	0	*	2	*	1	*
2563	1	*	5	*	0	*	0	*	5	*
3001	1,031	36.0 (101)	1,090	18.0 (91)	316	43.4 (37)	964	10.3 (66)	1,090	2.8 (75)
3002	242	26.0 (57)	249	13.7 (58)	56	*	231	9.1 (53)	249	1.6 (31)
3004	147	12.9 (13)	139	7.2 (22)	16	*	140	3.6 (8)	138	0.7 (7)
3007	26	*	20	*	3	*	25	*	20	*
3009	6	*	2	*	0	*	6	*	2	*
3201	467	34.0 (93)	474	20.0 (98)	139	44.6 (34)	382	16.2 (111)	474	3.6 (101)
3301	3,948	20.6 (31)	4,114	12.5 (54)	645	32.2 (78)	3,330	8.2 (41)	4,118	4.2 (122)
3304	107	0.9 (1)	107	0.0 (1)	1	*	86	*	107	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3305	135	19.3 (24)	142	7.7 (30)	20	*	107	3.7 (10)	142	2.1 (48)
3306	244	24.6 (51)	254	13.4 (57)	48	*	205	3.9 (13)	254	2.8 (75)
3308	64	*	72	*	13	*	48	*	72	*
3309	35	*	38	*	10	*	28	*	38	*
3312	368	28.3 (66)	398	17.6 (86)	95	*	337	8.3 (42)	398	4.0 (115)
3550	67	*	71	*	36	*	64	*	71	*
3551	23	*	27	*	12	*	21	*	27	*
3552	47	*	45	*	27	*	47	*	47	*
3701	1,035	41.7 (123)	1,049	23.5 (117)	353	33.1 (73)	893	14.1 (93)	1,049	2.8 (75)
3850	2,993	40.3 (116)	3,043	25.8 (128)	1,033	28.1 (93)	2,710	15.0 (100)	3,043	3.2 (88)
3901	458	26.0 (57)	539	15.0 (69)	87	*	356	8.7 (47)	539	1.1 (15)
3902	238	21.8 (36)	252	14.7 (64)	42	*	203	6.9 (27)	252	2.4 (62)
3903	9	*	10	*	0	*	0	*	10	*
3904	3	*	3	*	0	*	3	*	3	*
3905	1	*	2	*	0	*	0	*	2	*
3950	1,859	30.6 (77)	1,870	16.8 (79)	457	37.9 (56)	1,620	10.7 (70)	1,872	2.1 (48)
3954	5	*	4	*	0	*	3	*	4	*
4001	986	33.2 (90)	932	17.4 (83)	272	30.5 (84)	950	12.2 (82)	933	2.3 (58)
4002	687	32.6 (85)	659	17.3 (82)	188	30.9 (83)	645	11.0 (72)	659	2.3 (58)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4003	5	*	5	*	1	*	5	*	5	*
4004	5	*	5	*	2	*	5	*	5	*
4103	56	*	53	*	16	*	54	*	53	*
4106	2,173	22.6 (40)	2,261	11.3 (45)	397	39.8 (46)	1,974	4.8 (16)	2,260	1.5 (26)
4107	3,022	21.3 (34)	3,145	11.4 (46)	496	41.9 (38)	2,534	9.8 (63)	3,143	3.3 (93)
4109	1,012	30.2 (74)	1,049	14.8 (67)	248	46.0 (29)	824	7.3 (31)	1,049	1.1 (15)
4112	552	2.7 (2)	555	0.9 (3)	11	*	523	8.0 (39)	555	2.9 (80)
4113	647	9.4 (9)	638	4.2 (9)	55	*	631	2.4 (2)	640	0.6 (5)
4115	3,252	18.6 (21)	3,395	7.8 (32)	488	47.7 (21)	2,771	4.4 (15)	3,399	1.0 (13)
4351	576	42.7 (128)	580	23.8 (119)	209	34.0 (70)	518	17.0 (116)	580	2.8 (75)
4352	160	54.4 (159)	164	31.1 (143)	79	*	151	27.2 (146)	164	3.7 (103)
4353	412	30.8 (78)	426	20.0 (98)	110	28.2 (92)	369	11.7 (79)	427	2.1 (48)
4354	716	41.9 (125)	749	26.7 (131)	250	35.2 (63)	627	14.8 (97)	749	4.1 (118)
4355	345	41.2 (120)	359	20.9 (107)	126	32.5 (76)	305	11.5 (75)	359	2.8 (75)
4356	448	34.2 (94)	459	19.4 (96)	129	39.5 (47)	410	10.5 (67)	458	3.1 (86)
4358	131	13.0 (14)	137	7.3 (25)	14	*	120	5.0 (17)	138	0.0 (1)
4359	439	27.1 (63)	469	14.5 (63)	101	38.6 (50)	386	8.5 (45)	469	1.7 (33)
4360	212	33.5 (91)	234	17.1 (80)	67	*	196	17.9 (122)	234	3.8 (108)
4361	233	48.9 (148)	240	31.3 (144)	98	*	208	19.2 (127)	240	1.7 (33)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4362	279	53.8 (158)	285	32.6 (151)	132	32.6 (75)	251	21.5 (136)	285	4.6 (129)
4363	840	40.8 (119)	885	27.1 (133)	286	25.2 (100)	744	15.5 (103)	885	4.5 (125)
4401	1,056	34.9 (95)	1,084	16.3 (77)	289	40.5 (44)	887	3.7 (10)	1,083	0.9 (11)
4501	925	39.9 (115)	981	22.0 (114)	316	34.5 (67)	792	14.8 (97)	982	3.9 (112)
4601	1,354	36.3 (102)	1,411	17.6 (86)	386	44.6 (34)	1,157	9.7 (61)	1,412	3.3 (93)
4602	22	*	30	*	8	*	17	*	30	*
4701	1,061	35.9 (98)	1,085	17.5 (85)	269	47.6 (22)	798	8.8 (49)	1,088	2.5 (64)
5050	1,603	26.4 (60)	1,692	12.6 (55)	325	48.9 (19)	1,290	8.4 (44)	1,694	2.1 (48)
5051	2,468	23.7 (46)	2,567	10.7 (41)	466	46.8 (23)	1,983	7.1 (28)	2,570	1.9 (45)
5052	2,671	23.0 (43)	2,790	12.3 (50)	476	38.2 (54)	2,201	8.9 (51)	2,790	1.8 (41)
5055	1,175	2.8 (3)	1,182	0.6 (2)	28	*	1,086	4.3 (14)	1,179	0.8 (9)
5101	1,784	26.1 (59)	1,863	12.4 (51)	378	44.7 (32)	1,459	6.2 (21)	1,868	2.2 (54)
5102	1	*	1	*	0	*	0	*	1	*
5202	755	39.6 (114)	737	25.1 (125)	256	36.3 (59)	651	10.1 (65)	739	1.5 (26)
5203	23	*	24	*	10	*	24	*	24	*
5204	12	*	14	*	6	*	12	*	14	*
5601	1,036	31.3 (80)	1,079	20.9 (107)	274	26.6 (98)	886	7.9 (37)	1,079	1.5 (26)
5801	1,632	39.1 (111)	1,646	23.5 (117)	504	34.3 (68)	1,414	14.8 (97)	1,647	3.4 (99)
5803	123	28.5 (67)	122	19.7 (97)	22	*	88	*	122	4.1 (118)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5804	15	*	18	*	5	*	14	*	18	*
5806	15	*	30	*	2	*	4	*	30	*
6101	26	*	28	*	8	*	21	*	28	*
6102	109	30.3 (76)	123	17.1 (80)	32	*	98	*	123	4.1 (118)
6103	3,849	32.4 (83)	3,998	21.2 (109)	1,068	27.2 (96)	3,447	13.8 (92)	4,018	5.7 (139)
6105	24	*	27	*	4	*	18	*	28	*
6106	12	*	17	*	0	*	2	*	17	*
6301	4,030	18.9 (23)	4,190	10.7 (41)	582	38.3 (52)	3,372	6.4 (24)	4,189	1.2 (19)
6312	2,183	7.4 (7)	2,241	2.8 (7)	117	63.2 (6)	1,722	2.8 (3)	2,241	1.0 (13)
6316	1,182	17.9 (20)	1,166	8.5 (34)	137	41.6 (40)	889	5.7 (19)	1,166	1.1 (15)
6320	115	4.3 (5)	115	2.6 (6)	5	*	88	*	115	0.0 (1)
6325	131	31.3 (80)	143	15.4 (73)	33	*	93	*	143	1.4 (23)
6330	351	24.5 (50)	334	14.7 (64)	77	*	331	9.1 (53)	334	2.7 (73)
6335	952	8.5 (8)	952	4.6 (10)	61	*	794	2.1 (1)	952	0.6 (5)
6340	52	*	55	*	8	*	42	*	56	*
6345	27	*	28	*	5	*	18	*	28	*
6360	317	21.1 (33)	353	12.2 (49)	56	*	266	9.4 (59)	354	1.7 (33)
7304	116	40.5 (118)	120	15.0 (69)	45	*	101	11.9 (80)	121	1.7 (33)
7306	2,297	36.3 (102)	2,441	14.1 (60)	698	50.7 (18)	2,004	14.3 (94)	2,444	2.6 (69)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7308	119	37.0 (104)	114	20.2 (100)	35	*	107	9.3 (56)	115	2.6 (69)
7310	38	*	35	*	13	*	31	*	35	*
7401	2,223	39.2 (113)	2,240	24.1 (120)	720	31.4 (80)	2,016	17.3 (121)	2,240	4.0 (115)
7601	600	35.0 (96)	620	20.6 (103)	187	34.2 (69)	542	11.6 (78)	622	2.6 (69)
7850	1,056	38.8 (109)	1,094	21.3 (110)	362	32.9 (74)	931	18.3 (124)	1,100	2.5 (64)
7852	25	*	28	*	12	*	24	*	28	*
8101	1,704	22.8 (41)	1,933	9.1 (38)	305	55.7 (12)	1,399	16.4 (112)	1,932	5.3 (135)
8102	395	15.7 (15)	416	7.0 (20)	55	*	367	5.7 (19)	416	2.2 (54)
8105	1,590	16.4 (17)	1,656	7.5 (29)	186	46.2 (28)	1,265	11.5 (75)	1,656	3.2 (88)
8106	434	21.7 (35)	486	10.1 (40)	82	*	364	9.1 (53)	487	1.4 (23)
8150	38	*	37	*	7	*	36	*	37	*
8151	3,047	25.2 (53)	3,042	11.2 (44)	575	44.7 (32)	2,609	8.0 (39)	3,039	1.8 (41)
8152	106	30.2 (74)	101	13.9 (59)	30	*	103	9.7 (61)	102	2.9 (80)
8153	10	*	10	*	1	*	8	*	10	*
8155	35	*	38	*	5	*	30	*	38	*
8157	9	*	7	*	0	*	8	*	7	*
8201	2,508	20.5 (28)	2,608	11.9 (48)	391	39.1 (49)	1,983	9.9 (64)	2,612	2.3 (58)
8202	3,806	6.9 (6)	3,938	2.0 (5)	226	63.7 (4)	3,278	7.5 (33)	3,936	2.2 (54)
8203	429	32.9 (87)	448	18.8 (93)	113	35.4 (62)	331	11.5 (75)	450	1.3 (22)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8250	690	28.1 (65)	769	12.4 (51)	139	53.2 (17)	549	12.2 (82)	769	3.3 (93)
8252	108	23.1 (44)	125	4.0 (8)	23	*	91	*	125	4.0 (115)
8255	2,415	29.3 (69)	2,584	15.2 (71)	547	40.6 (43)	1,962	8.8 (49)	2,584	1.8 (41)
8257	1,827	2.9 (4)	1,865	1.4 (4)	44	*	1,536	2.8 (3)	1,864	0.9 (11)
8263	672	25.0 (52)	712	12.4 (51)	133	46.6 (26)	525	7.2 (30)	713	1.7 (33)
8264	650	25.5 (55)	659	11.4 (46)	127	44.9 (31)	516	9.3 (56)	660	2.4 (62)
8265	21	*	19	*	9	*	21	*	19	*
8266	41	*	37	*	1	*	40	*	37	*
8267	718	29.7 (72)	730	14.1 (60)	151	41.7 (39)	548	8.9 (51)	730	0.8 (9)
8901	976	25.2 (53)	1,098	8.1 (33)	218	58.3 (11)	879	16.4 (112)	1,099	5.4 (137)
8902	1,212	18.6 (21)	1,311	5.3 (14)	185	65.4 (2)	1,025	15.5 (103)	1,312	6.4 (148)
8903	1,139	26.4 (60)	1,275	9.0 (36)	249	58.6 (10)	991	19.1 (126)	1,273	7.2 (151)
8904	2,119	22.8 (41)	2,330	7.2 (22)	391	66.5 (1)	1,747	17.2 (118)	2,331	7.2 (151)
8905	1,475	16.2 (16)	1,567	7.0 (20)	191	55.5 (13)	1,226	7.6 (35)	1,569	2.9 (80)
8908	30	*	31	*	9	*	28	*	31	*
8909	419	22.0 (37)	457	4.8 (12)	82	*	326	17.2 (118)	457	7.4 (155)
9012	7	*	6	*	1	*	7	*	6	*
9013	1	*	1	*	0	*	1	*	1	*
9016	50	*	47	*	17	*	51	*	47	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9017	6	*	5	*	1	*	6	*	5	*
9018	54	*	51	*	9	*	54	*	51	*
9041	2	*	2	*	0	*	2	*	2	*
9045	1,655	12.3 (12)	1,762	5.3 (14)	167	46.7 (24)	1,454	9.3 (56)	1,762	3.7 (103)
9050	847	27.2 (64)	970	5.5 (16)	215	65.1 (3)	750	19.2 (127)	969	5.1 (133)
9051	1,641	10.8 (11)	1,627	4.6 (10)	137	46.7 (24)	1,541	6.7 (25)	1,626	2.5 (64)
9053	996	22.1 (38)	1,101	7.4 (26)	182	58.8 (9)	830	15.5 (103)	1,103	7.1 (150)
9055	41	*	50	*	2	*	9	*	50	*
9058	940	10.3 (10)	1,048	5.2 (13)	74	*	798	6.3 (23)	1,048	3.7 (103)
9066	934	17.6 (18)	1,127	6.9 (19)	141	54.6 (16)	782	16.6 (114)	1,127	4.1 (118)
9068	865	23.8 (47)	964	7.7 (30)	185	62.2 (7)	764	18.2 (123)	964	3.7 (103)
9069	2,523	20.4 (26)	2,911	6.7 (18)	422	63.7 (4)	2,044	21.9 (140)	2,909	9.3 (157)
9070	1,015	20.5 (28)	1,151	7.4 (26)	166	55.4 (14)	813	23.6 (143)	1,148	6.2 (144)
9072	1,632	23.2 (45)	1,873	7.2 (22)	318	55.0 (15)	1,404	24.1 (144)	1,874	5.9 (141)
9080	2,429	20.7 (32)	2,724	7.4 (26)	422	59.0 (8)	2,064	11.4 (74)	2,726	4.6 (129)
9081	41	*	42	*	1	*	39	*	42	*
9082	112	20.5 (28)	112	6.3 (17)	18	*	78	*	112	9.8 (158)
9083	43	*	55	*	0	*	8	*	53	*
9084	17	*	25	*	0	*	0	*	25	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9085	48	*	45	*	11	*	48	*	45	*
9086	36	*	30	*	0	*	36	*	30	*
Michigan	**153,367**	**27.3**	**160,682**	**14.4**	**34,505**	**39.2**	**131,305**	**11.2**	**160,776**	**3.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	103	9.7 (105)	11.7 (149)	107	13.1 (120)	92	*	103	74.8 (74)
0102	487	6.0 (15)	9.2 (114)	491	9.2 (25)	437	3.2 (49)	487	65.5 (107)
0103	403	8.2 (64)	10.2 (131)	405	8.9 (19)	361	3.6 (71)	402	65.4 (108)
0104	142	4.9 (7)	4.9 (23)	143	5.6 (3)	133	3.8 (80)	142	75.4 (71)
0105	7	*	*	7	*	7	*	7	*
0250	129	4.7 (6)	14.0 (157)	130	8.5 (13)	115	1.7 (5)	129	76.0 (68)
0251	85	*	*	89	*	80	*	85	*
0252	109	6.4 (22)	8.3 (90)	110	10.0 (42)	98	*	109	87.2 (6)
0253	148	13.5 (141)	10.8 (139)	149	13.4 (125)	126	8.7 (149)	147	62.6 (116)
0401	445	8.1 (58)	11.9 (151)	457	9.4 (29)	402	3.5 (64)	443	67.0 (100)
0402	339	9.4 (101)	7.7 (76)	344	9.0 (22)	302	5.3 (119)	339	79.4 (45)
0403	125	12.0 (132)	9.6 (122)	127	11.8 (93)	109	5.5 (124)	125	65.6 (106)
0404	139	3.6 (3)	11.5 (147)	139	9.4 (29)	125	1.6 (4)	139	63.3 (114)
0501	381	7.6 (48)	9.4 (121)	491	8.8 (16)	343	3.5 (64)	463	78.2 (53)
0503	103	6.8 (25)	7.8 (80)	104	4.8 (2)	95	*	103	80.6 (33)
0504	462	8.0 (54)	11.7 (149)	478	7.9 (9)	423	4.7 (106)	469	83.6 (12)
0505	295	7.5 (44)	8.1 (86)	319	6.9 (5)	272	2.9 (32)	312	83.3 (14)
0506	170	5.9 (14)	5.9 (41)	190	7.4 (6)	155	3.2 (49)	187	79.1 (47)
0507	70	*	*	79	*	63	*	78	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	28	*	*	29	*	23	*	29	*
0509	8	*	*	8	*	8	*	7	*
0650	1,009	8.2 (64)	8.8 (105)	1,056	8.1 (10)	926	5.4 (123)	1,009	77.9 (56)
0651	545	6.1 (19)	8.1 (86)	557	8.8 (16)	493	3.0 (37)	543	65.4 (108)
0652	274	8.0 (54)	9.1 (112)	284	9.5 (32)	246	3.7 (74)	273	68.9 (94)
0653	398	9.3 (94)	8.8 (105)	415	10.1 (47)	357	3.9 (83)	399	81.5 (23)
0654	329	5.8 (13)	10.9 (141)	335	11.3 (78)	292	2.1 (11)	329	81.2 (29)
0655	343	8.5 (73)	6.7 (51)	349	9.7 (36)	307	4.2 (90)	344	73.8 (78)
0657	56	*	*	56	*	49	*	56	*
0701	372	6.2 (20)	10.8 (139)	386	10.1 (47)	321	2.2 (15)	374	89.3 (3)
0702	53	*	*	56	*	44	*	53	*
0703	193	8.8 (79)	7.8 (80)	193	13.0 (116)	165	3.0 (37)	193	68.4 (96)
0704	44	*	*	46	*	40	*	44	*
0705	111	5.4 (12)	9.0 (109)	118	7.6 (7)	102	2.9 (32)	111	80.2 (39)
0706	4	*	*	4	*	3	*	4	*
0850	814	8.6 (75)	7.0 (59)	869	11.0 (73)	726	2.2 (15)	845	83.9 (11)
0851	697	9.0 (84)	9.3 (118)	725	10.2 (49)	620	4.4 (102)	716	78.1 (55)
0901	1,571	7.1 (34)	10.1 (129)	1,601	9.6 (34)	1,415	2.8 (27)	1,571	67.0 (100)
0902	42	*	*	43	*	42	*	42	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1051	1,314	8.4 (72)	10.9 (141)	1,342	9.2 (25)	1,185	3.2 (49)	1,313	86.2 (9)
1101	1,646	9.0 (84)	7.1 (63)	1,699	10.2 (49)	1,473	4.3 (100)	1,650	68.7 (95)
1102	718	6.5 (23)	9.2 (114)	747	9.6 (34)	645	3.7 (74)	718	76.2 (67)
1106	66	*	*	67	*	58	*	66	*
1250	913	6.0 (15)	7.9 (83)	931	10.2 (49)	802	2.2 (15)	924	79.8 (41)
1251	753	4.9 (7)	8.2 (88)	782	9.0 (22)	681	1.8 (6)	753	79.0 (48)
1252	499	7.0 (31)	7.0 (59)	515	10.9 (69)	440	2.3 (18)	503	81.1 (30)
1253	763	8.1 (58)	8.4 (92)	783	10.7 (63)	676	3.1 (44)	765	84.4 (10)
1350	2,273	9.3 (94)	7.6 (74)	2,324	10.5 (59)	2,021	3.9 (83)	2,276	73.2 (83)
1351	130	6.9 (27)	7.7 (76)	132	9.1 (24)	118	5.1 (116)	132	80.3 (38)
1352	340	9.1 (88)	8.2 (88)	345	11.3 (78)	302	5.6 (127)	341	61.0 (117)
1401	218	8.3 (68)	7.3 (67)	222	9.9 (39)	192	3.1 (44)	216	82.9 (16)
1402	122	6.6 (24)	13.9 (156)	129	10.9 (69)	107	0.9 (2)	122	89.3 (3)
1403	25	*	*	24	*	24	*	24	*
1404	0	*	*	0	*	0	*	0	*
1501	672	7.3 (37)	10.4 (133)	682	11.0 (73)	586	2.9 (32)	677	66.5 (102)
1503	958	7.4 (41)	8.7 (102)	967	13.1 (120)	814	2.8 (27)	961	79.7 (42)
1504	2,004	8.0 (54)	9.0 (109)	2,071	11.6 (85)	1,763	2.8 (27)	2,029	77.3 (60)
1505	422	8.8 (79)	14.7 (158)	433	8.8 (16)	384	3.9 (83)	423	81.8 (21)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1506	530	6.0 (15)	10.9 (141)	543	10.5 (59)	470	2.1 (11)	533	73.5 (81)
1507	549	9.7 (105)	12.9 (155)	576	12.0 (96)	479	3.1 (44)	559	78.5 (51)
1508	82	*	*	81	*	68	*	82	*
1509	368	7.9 (52)	7.6 (74)	383	12.0 (96)	323	4.3 (100)	378	69.3 (92)
1511	669	8.7 (77)	9.3 (118)	679	10.3 (55)	592	3.0 (37)	676	79.4 (45)
1512	341	10.6 (117)	10.3 (132)	352	15.9 (145)	284	3.5 (64)	345	80.0 (40)
1513	65	*	*	69	*	57	*	67	*
1514	156	16.0 (155)	9.6 (122)	158	17.7 (152)	128	4.7 (106)	157	90.4 (1)
1701	614	7.0 (31)	11.6 (148)	612	11.6 (85)	523	1.9 (7)	615	71.4 (89)
1950	556	7.6 (48)	6.8 (55)	564	10.5 (59)	490	3.3 (56)	557	82.6 (17)
1952	127	3.1 (1)	7.1 (63)	125	6.4 (4)	117	0.9 (2)	127	81.9 (20)
1953	294	7.5 (44)	9.9 (127)	296	11.1 (76)	258	3.1 (44)	294	83.3 (14)
1954	529	8.1 (58)	8.5 (97)	538	10.8 (67)	469	3.8 (80)	529	81.1 (30)
1955	79	*	*	81	*	75	*	79	*
1957	508	7.3 (37)	9.1 (112)	514	9.7 (36)	455	2.6 (22)	510	86.9 (7)
1963	10	*	*	10	*	6	*	10	*
1964	28	*	*	29	*	25	*	28	*
2150	582	8.1 (58)	10.5 (134)	600	10.2 (49)	523	3.6 (71)	570	77.7 (57)
2151	209	10.5 (114)	7.7 (76)	207	11.6 (85)	182	4.9 (111)	209	54.1 (139)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2251	397	6.8 (25)	8.8 (105)	400	9.3 (28)	352	2.6 (22)	396	74.5 (75)
2252	163	9.2 (92)	8.0 (84)	169	12.4 (103)	144	5.6 (127)	163	82.2 (19)
2554	78	*	*	80	*	68	*	78	*
2555	7,654	11.5 (127)	5.3 (29)	7,708	13.7 (129)	6,529	5.3 (119)	7,665	58.7 (131)
2556	192	18.8 (158)	2.1 (1)	191	18.8 (155)	150	10.0 (151)	192	53.1 (141)
2557	6	*	*	7	*	5	*	7	*
2558	4	*	*	4	*	4	*	4	*
2559	9	*	*	9	*	7	*	9	*
2560	2	*	*	2	*	2	*	2	*
2562	1	*	*	1	*	1	*	1	*
2563	4	*	*	3	*	2	*	4	*
3001	1,024	13.4 (140)	4.7 (21)	1,042	14.9 (140)	867	5.1 (116)	1,025	60.5 (120)
3002	238	12.2 (134)	6.7 (51)	244	13.1 (120)	208	4.8 (110)	238	50.4 (145)
3004	130	6.9 (27)	8.5 (97)	130	12.3 (100)	114	2.6 (22)	130	77.7 (57)
3007	19	*	*	19	*	17	*	19	*
3009	1	*	*	1	*	1	*	1	*
3201	452	6.9 (27)	12.6 (154)	460	10.0 (42)	408	2.9 (32)	452	75.0 (73)
3301	3,888	10.3 (111)	6.4 (44)	4,025	13.0 (116)	3,376	4.7 (106)	3,919	78.9 (49)
3304	98	*	*	101	10.9 (69)	86	*	102	89.2 (5)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3305	133	7.5 (44)	11.3 (145)	136	10.3 (55)	117	2.6 (22)	134	90.3 (2)
3306	234	8.1 (58)	12.0 (152)	241	10.4 (57)	209	1.9 (7)	235	83.4 (13)
3308	67	*	*	69	*	64	*	66	*
3309	38	*	*	37	*	32	*	38	*
3312	380	10.5 (114)	5.8 (38)	391	10.7 (63)	339	5.6 (127)	384	68.2 (97)
3550	64	*	*	69	*	59	*	63	*
3551	25	*	*	27	*	23	*	25	*
3552	42	*	*	47	*	35	*	42	*
3701	1,010	8.6 (75)	7.1 (63)	1,022	12.6 (105)	879	2.8 (27)	1,011	80.6 (33)
3850	2,880	9.0 (84)	7.4 (70)	2,934	12.6 (105)	2,492	3.3 (56)	2,879	74.3 (77)
3901	498	10.4 (112)	6.4 (44)	514	13.0 (116)	426	4.2 (90)	505	71.5 (88)
3902	234	7.3 (37)	11.1 (144)	234	12.8 (111)	201	3.5 (64)	235	70.2 (91)
3903	9	*	*	10	*	9	*	9	*
3904	2	*	*	2	*	2	*	2	*
3905	2	*	*	2	*	2	*	2	*
3950	1,763	9.3 (94)	10.7 (136)	1,818	12.2 (98)	1,547	3.3 (56)	1,773	76.5 (64)
3954	4	*	*	4	*	4	*	4	*
4001	862	10.0 (110)	5.5 (33)	885	11.9 (94)	746	4.7 (106)	867	67.8 (98)
4002	623	9.1 (88)	7.5 (72)	639	11.0 (73)	549	4.2 (90)	628	66.1 (104)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4003	5	*	*	5	*	4	*	4	*
4004	4	*	*	5	*	4	*	4	*
4103	49	*	*	50	*	37	*	49	*
4106	2,116	12.0 (132)	5.2 (28)	2,189	13.3 (124)	1,820	6.2 (134)	2,132	73.7 (80)
4107	2,968	11.0 (123)	8.3 (90)	3,071	13.7 (129)	2,549	4.2 (90)	2,975	75.9 (69)
4109	1,004	9.3 (94)	9.3 (118)	1,016	13.6 (127)	864	2.8 (27)	1,009	80.7 (32)
4112	540	5.2 (10)	6.7 (51)	537	10.2 (49)	481	2.1 (11)	539	27.1 (158)
4113	608	8.9 (82)	7.1 (63)	629	12.9 (114)	527	4.2 (90)	609	80.6 (33)
4115	3,226	9.5 (102)	7.7 (76)	3,311	12.8 (111)	2,799	3.2 (49)	3,238	81.3 (25)
4351	542	8.5 (73)	9.6 (122)	559	12.5 (104)	473	3.0 (37)	541	73.2 (83)
4352	153	10.5 (114)	3.9 (17)	156	17.3 (151)	123	4.9 (111)	153	73.2 (83)
4353	401	8.2 (64)	7.0 (59)	406	8.9 (19)	361	4.2 (90)	400	78.5 (51)
4354	707	7.4 (41)	7.8 (80)	710	10.7 (63)	621	3.4 (61)	706	77.1 (62)
4355	337	8.0 (54)	10.7 (136)	345	10.4 (57)	296	3.4 (61)	336	79.5 (43)
4356	445	9.2 (92)	5.8 (38)	449	13.6 (127)	382	4.2 (90)	445	78.2 (53)
4358	131	4.6 (5)	9.2 (114)	130	4.6 (1)	123	2.4 (19)	132	81.8 (21)
4359	431	7.2 (36)	8.4 (92)	434	9.9 (39)	383	3.1 (44)	428	73.8 (78)
4360	206	4.9 (7)	10.7 (136)	222	8.6 (14)	186	3.2 (49)	205	81.5 (23)
4361	220	9.5 (102)	11.4 (146)	231	13.0 (116)	189	3.7 (74)	220	80.5 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4362	268	7.8 (50)	7.5 (72)	272	9.9 (39)	240	3.3 (56)	267	86.5 (8)
4363	831	8.3 (68)	8.4 (92)	838	11.3 (78)	718	3.5 (64)	831	80.4 (37)
4401	1,038	7.5 (44)	9.2 (114)	1,045	10.0 (42)	920	3.2 (49)	1,037	76.8 (63)
4501	930	7.4 (41)	10.6 (135)	953	10.8 (67)	821	2.9 (32)	929	73.4 (82)
4601	1,329	6.2 (20)	9.0 (109)	1,363	9.5 (32)	1,194	3.4 (61)	1,326	72.9 (86)
4602	28	*	*	29	*	27	*	28	*
4701	1,047	8.1 (58)	9.9 (127)	1,074	10.5 (59)	934	3.7 (74)	1,043	78.9 (49)
5050	1,654	11.0 (123)	5.6 (35)	1,649	14.4 (133)	1,388	3.8 (80)	1,657	59.4 (125)
5051	2,501	11.4 (126)	6.2 (42)	2,517	13.5 (126)	2,134	4.2 (90)	2,505	63.8 (112)
5052	2,723	11.1 (125)	4.9 (23)	2,725	12.9 (114)	2,331	5.5 (124)	2,727	60.0 (123)
5055	883	9.1 (88)	3.6 (15)	1,098	8.4 (12)	800	5.0 (114)	1,073	38.0 (157)
5101	1,802	8.7 (77)	6.8 (55)	1,820	11.7 (91)	1,573	3.9 (83)	1,805	72.7 (87)
5102	1	*	*	1	*	1	*	1	*
5202	711	9.8 (109)	8.4 (92)	723	11.9 (94)	619	4.5 (103)	711	82.4 (18)
5203	22	*	*	21	*	17	*	22	*
5204	13	*	*	13	*	9	*	13	*
5601	1,020	7.1 (34)	10.1 (129)	1,041	10.9 (69)	904	2.5 (21)	1,020	75.7 (70)
5801	1,570	8.2 (64)	6.9 (58)	1,608	11.4 (83)	1,376	3.0 (37)	1,536	64.1 (111)
5803	116	6.0 (15)	8.6 (99)	118	10.2 (49)	103	1.9 (7)	114	60.5 (120)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5804	18	*	*	18	*	16	*	18	*
5806	29	*	*	29	*	24	*	29	*
6101	28	*	*	26	*	25	*	28	*
6102	113	8.8 (79)	9.7 (126)	110	10.0 (42)	98	*	112	81.3 (25)
6103	3,782	10.6 (117)	5.3 (29)	3,863	14.4 (133)	3,167	4.5 (103)	3,791	65.0 (110)
6105	28	*	*	28	*	25	*	28	*
6106	17	*	*	17	*	10	*	17	*
6301	3,988	12.5 (136)	5.6 (35)	4,088	14.0 (132)	3,412	5.3 (119)	3,999	65.8 (105)
6312	2,144	10.4 (112)	5.8 (38)	2,193	12.3 (100)	1,867	3.5 (64)	2,157	76.4 (66)
6316	1,120	9.3 (94)	7.4 (70)	1,140	12.6 (105)	969	3.0 (37)	1,125	77.7 (57)
6320	108	8.3 (68)	6.5 (46)	110	8.2 (11)	98	*	107	81.3 (25)
6325	132	4.5 (4)	12.1 (153)	141	7.8 (8)	124	2.4 (19)	136	77.2 (61)
6330	303	12.2 (134)	4.3 (19)	327	14.4 (133)	257	5.8 (131)	319	54.2 (137)
6335	895	9.3 (94)	5.5 (33)	914	10.0 (42)	801	5.0 (114)	910	70.8 (90)
6340	53	*	*	54	*	46	*	54	*
6345	26	*	*	26	*	22	*	26	*
6360	332	13.9 (146)	5.1 (27)	339	15.6 (144)	271	6.6 (138)	341	59.5 (124)
7304	116	5.2 (10)	8.6 (99)	117	9.4 (29)	105	1.9 (7)	116	63.8 (112)
7306	2,349	11.5 (127)	6.3 (43)	2,378	11.3 (78)	2,067	5.1 (116)	2,352	60.3 (122)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7308	113	3.5 (2)	8.8 (105)	113	9.7 (36)	101	0.0 (1)	113	53.1 (141)
7310	34	*	*	35	*	32	*	34	*
7401	2,124	7.8 (50)	8.0 (84)	2,167	11.2 (77)	1,877	3.0 (37)	2,123	69.2 (93)
7601	596	6.9 (27)	8.7 (102)	610	10.7 (63)	524	2.1 (11)	596	62.9 (115)
7850	1,047	8.3 (68)	8.4 (92)	1,062	8.9 (19)	948	3.5 (64)	1,046	74.5 (75)
7852	27	*	*	28	*	27	*	27	*
8101	1,760	16.5 (156)	3.3 (9)	1,884	19.9 (158)	1,393	6.9 (143)	1,823	46.3 (149)
8102	372	9.7 (105)	8.6 (99)	379	18.2 (154)	286	4.9 (111)	371	43.7 (154)
8105	1,592	10.9 (121)	5.7 (37)	1,622	12.6 (105)	1,376	3.7 (74)	1,597	67.1 (99)
8106	444	11.9 (131)	5.4 (31)	476	11.6 (85)	389	5.7 (130)	456	59.4 (125)
8150	34	*	*	34	*	32	*	33	*
8151	2,892	10.8 (119)	5.4 (31)	2,956	13.7 (129)	2,490	4.6 (105)	2,769	81.3 (25)
8152	102	10.8 (119)	4.9 (23)	99	*	87	*	96	*
8153	10	*	*	10	*	9	*	10	*
8155	33	*	*	35	*	30	*	32	*
8157	5	*	*	6	*	4	*	5	*
8201	2,514	9.6 (104)	6.8 (55)	2,568	11.6 (85)	2,210	4.1 (89)	2,503	60.6 (119)
8202	3,478	7.9 (52)	6.6 (49)	3,782	9.2 (25)	3,129	4.2 (90)	3,750	46.2 (150)
8203	432	10.9 (121)	6.5 (46)	447	12.3 (100)	379	2.6 (22)	434	75.1 (72)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8250	733	11.5 (127)	4.9 (23)	746	12.2 (98)	631	5.5 (124)	732	66.4 (103)
8252	124	15.3 (153)	6.5 (46)	121	13.2 (123)	105	8.6 (148)	124	56.5 (133)
8255	2,486	9.3 (94)	7.0 (59)	2,529	11.6 (85)	2,179	4.0 (88)	2,489	59.2 (127)
8257	1,621	11.8 (130)	4.3 (19)	1,801	11.3 (78)	1,415	6.6 (138)	1,715	76.5 (64)
8263	695	7.3 (37)	9.6 (122)	705	8.7 (15)	631	3.3 (56)	692	58.8 (129)
8264	614	9.1 (88)	7.3 (67)	641	11.4 (83)	541	3.9 (83)	617	59.0 (128)
8265	19	*	*	19	*	17	*	19	*
8266	32	*	*	36	*	26	*	36	*
8267	709	9.7 (105)	7.3 (67)	721	11.7 (91)	626	3.2 (49)	708	58.8 (129)
8901	1,033	14.1 (148)	2.7 (2)	1,066	19.1 (156)	819	6.3 (135)	1,040	48.0 (146)
8902	1,209	13.7 (143)	3.3 (9)	1,278	16.3 (149)	996	5.3 (119)	1,241	55.3 (136)
8903	1,185	14.9 (152)	3.2 (5)	1,236	18.1 (153)	956	7.2 (145)	1,204	44.9 (152)
8904	2,135	13.2 (139)	3.0 (3)	2,257	15.9 (145)	1,768	6.3 (135)	2,191	54.2 (137)
8905	1,378	8.9 (82)	6.7 (51)	1,497	12.8 (111)	1,181	4.2 (90)	1,466	57.1 (132)
8908	28	*	*	31	*	23	*	29	*
8909	432	16.9 (157)	3.0 (3)	444	19.6 (157)	340	7.4 (147)	435	56.1 (134)
9012	6	*	*	6	*	4	*	6	*
9013	1	*	*	1	*	1	*	1	*
9016	44	*	*	47	*	39	*	44	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9017	5	*	*	5	*	4	*	5	*
9018	48	*	*	51	*	44	*	49	*
9041	2	*	*	2	*	2	*	2	*
9045	1,655	9.0 (84)	6.6 (49)	1,682	12.6 (105)	1,416	3.7 (74)	1,670	51.7 (143)
9050	904	13.6 (142)	4.2 (18)	929	15.9 (145)	746	7.0 (144)	915	53.4 (140)
9051	1,434	7.0 (31)	8.7 (102)	1,470	12.6 (105)	1,219	3.6 (71)	1,460	79.5 (43)
9053	1,037	13.1 (137)	4.7 (21)	1,081	14.5 (137)	886	6.5 (137)	1,036	46.4 (147)
9055	44	*	*	49	*	39	*	44	*
9058	965	13.8 (145)	3.2 (5)	1,009	14.4 (133)	818	6.7 (140)	995	55.8 (135)
9066	1,060	14.4 (149)	3.4 (13)	1,107	15.4 (141)	888	7.2 (145)	1,065	42.4 (155)
9068	913	15.9 (154)	3.2 (5)	939	15.4 (141)	766	9.1 (150)	901	43.8 (153)
9069	2,773	13.7 (143)	3.2 (5)	2,848	14.7 (138)	2,352	5.9 (133)	2,770	38.5 (156)
9070	1,097	14.6 (151)	3.3 (9)	1,125	16.6 (150)	897	6.7 (140)	1,101	45.8 (151)
9072	1,760	14.5 (150)	3.4 (13)	1,845	14.7 (138)	1,495	6.8 (142)	1,751	46.4 (147)
9080	2,603	13.1 (137)	3.3 (9)	2,652	15.5 (143)	2,171	5.8 (131)	2,617	50.5 (144)
9081	39	*	*	41	*	36	*	39	*
9082	107	14.0 (147)	3.7 (16)	111	16.2 (148)	88	*	107	60.7 (118)
9083	50	*	*	55	*	40	*	51	*
9084	23	*	*	25	*	20	*	23	*

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2016 - 2018 Pregnancy Nutrition Surveillance (1)

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9085	45	*	*	44	*	37	*	45	*
9086	30	*	*	30	*	26	*	30	*
Michigan	**151,265**	**10.0**	**6.8**	**155,722**	**12.4**	**131,154**	**4.3**	**152,480**	**57.8**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.