

Quick Reference to Combination and/or Reconstituted Vaccines: Childhood

Highlight Vaccines in Your Refrigerator/Freezer and Post

Brand Name	Contains	Use for Ages	Use for Dose	Administration Tips ¹
Pentacel® SP	DTaP, IPV + Hib	6 weeks through 4 years	1, 2, 3 or 4 of DTaP, IPV or Hib	<ul style="list-style-type: none"> • DTaP/IPV liquid (diluent) packaged with vial • Draw up diluent; add to Hib vial; shake well • Administer within 30 minutes; give IM
Pediarix® GSK	DTaP, IPV, Hep B	6 weeks through 6 years	1, 2, or 3 of IPV or DTaP; any dose of hep B	<ul style="list-style-type: none"> • Premixed • Shake well before administering; give IM
Kinrix® GSK	DTaP, IPV	4 through 6 years	5 th dose of DTaP; 4 th (valid) dose of IPV ²	<ul style="list-style-type: none"> • Premixed • Shake well before administering; give IM
Quadracel™ SP	DTaP, IPV	4 through 6 years	5 th dose of DTaP; 4 th or 5 th dose of IPV	<ul style="list-style-type: none"> • Premixed • Shake well before administering; give IM
ProQuad® Merck	MMR, Var (MMRV)	If 1 st dose: ages 12-47 months, use separate MMR and Var If 1 st dose: ages 4-12 years, use MMRV If 2 nd dose: ages 15 months-12 years, use MMRV		<ul style="list-style-type: none"> • Draw up supplied diluent • Add diluent to MMRV vial; shake well • Administer within 30 minutes; give SC
M-M-R II® Merck	MMR	12 months and older	1 or 2 of MMR	<ul style="list-style-type: none"> • Draw up supplied diluent • Add diluent to MMR vial; shake well; give SC
ActHIB® SP	Hib	6 weeks through 4 years	Any dose of Hib	<ul style="list-style-type: none"> • Draw up diluent packaged with Hib vial • Add diluent to Hib vial; shake well; give IM
Hiberix® GSK	Hib	6 weeks through 4 years	Any dose of Hib	<ul style="list-style-type: none"> • Draw up diluent supplied with Hib vial • Add diluent to Hib vial; shake well; give IM
Varivax® Merck	Var	12 months and older	1 or 2 of Var	<ul style="list-style-type: none"> • Draw up diluent supplied with Varicella vial • Add diluent to Varicella vial; shake well • Administer within 30 minutes; give SC
Menveo® Novartis	MenACWY	2 months through 55 years	Any dose of MenACWY	<ul style="list-style-type: none"> • MenCYW liquid (diluent) packaged with vial • Draw up MenCYW diluent; add to MenA vial • Invert; shake well; give IM
Rotarix® GSK	RV1 (Rotavirus)	6 weeks through 7 months	Any dose of RV	<ul style="list-style-type: none"> • Diluent packaged with vial • Shake diluent in pre-filled oral applicator • Add to RV1 vial; shake; withdraw; give orally

¹Refer to the manufacturer's package insert for further details regarding reconstituting and/or administering these products.

²When used in combination with Pentacel (DTaP-IPV-Hib), Kinrix may be used for the 5th (4th valid) dose of the IPV series.

Avoid medication errors! Use only the diluent that is packaged or sent with each specific vaccine—don't use any other liquid.