


# COVID-19 Vaccination Schedule\*



Vaccine	0 month	1 month	2 month	3 month	4 month	5 month	6 month	7 month	8 month	9 month	10 month	11 month
<b>Pfizer-BioNTech (ages 5-11 years)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (3 weeks after 1 <sup>st</sup> dose)										
<b>Pfizer-BioNTech (ages 12 years and older)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose<sup>1</sup></b> (3-8 weeks after 1 <sup>st</sup> dose)					<b>Booster Dose<sup>2</sup></b> (at least 5 months after 2 <sup>nd</sup> dose)				<b>2<sup>nd</sup> Booster Dose<sup>3</sup></b> (See footnote)	
<b>Moderna (ages 18 years and older)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose<sup>1</sup></b> (4-8 weeks after 1 <sup>st</sup> dose)					<b>Booster Dose<sup>2</sup></b> (at least 5 months after 2 <sup>nd</sup> dose)				<b>2<sup>nd</sup> Booster Dose<sup>3</sup></b> (See footnote)	
<b>Janssen (ages 18 years and older)</b>	<b>1<sup>st</sup> Dose</b>			<b>Booster Dose<sup>2</sup></b> (at least 2 months after 1 <sup>st</sup> dose)				<b>2<sup>nd</sup> Booster Dose<sup>3</sup></b> (See footnote)				

Note: Timeline is approximate. Intervals of 3 months or fewer are converted into weeks per the formula “1 month = 4 weeks.” Intervals of 4 months or more are converted into calendar months.

\* See Guidance for COVID-19 Vaccination Schedule for People Who are Moderately or Severely Immunocompromised.

<sup>1</sup> An 8-week interval may be optimal for some people ages 12 years and older, especially for males ages 12 to 39 years. A shorter interval (3 weeks for Pfizer-BioNTech; 4 weeks for Moderna) between the first and second doses remains the recommended interval for: people who are moderately or severely immunocompromised; adults ages 65 years and older; and others who need rapid protection due to increased concern about community transmission or risk of severe disease. For more information, view [the CDC Interim Clinical Considerations](https://bit.ly/COVIDClinicalConsiderations) (bit.ly/COVIDClinicalConsiderations).

<sup>2</sup> An mRNA COVID-19 vaccine is preferred over the Janssen COVID-19 Vaccine for booster vaccination of people ages 18 years and older. For people ages 12 through 17 years, only Pfizer-BioNTech can be used. People ages 5 through 11 years should not receive a booster dose.

<sup>3</sup> People ages 18 through 49 years who received Janssen COVID-19 Vaccine as both their primary series dose and booster dose may receive an mRNA COVID-19 booster dose at least 4 months after the Janssen booster dose. People ages 50 years and older may choose to receive a second booster dose if it has been at least 4 months after the first booster dose.

