



# COVID-19 Vaccination Schedule

## For Those who are Moderately or Severely Immunocompromised

Vaccine	0 month	1 month	2 month	3 month	4 month	5 month	6 month	7 month	8 month	9 month
<b>Pfizer-BioNTech (ages 5-11 years)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (3 weeks after 1 <sup>st</sup> dose)	<b>3<sup>rd</sup> Dose</b> (At least 4 weeks after 2 <sup>nd</sup> dose)			<b>Booster Dose<sup>1</sup></b> (at least 3 months after 3 <sup>rd</sup> dose)				
<b>Pfizer-BioNTech (ages 12 years and older)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (3 weeks after 1 <sup>st</sup> dose)	<b>3<sup>rd</sup> Dose</b> (At least 4 weeks after 2 <sup>nd</sup> dose)			<b>Booster Dose<sup>1</sup></b> (at least 3 months after 3 <sup>rd</sup> dose)				<b>2<sup>nd</sup> Booster Dose<sup>3</sup></b> (See footnote)
<b>Moderna (ages 18 years and older)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (4 weeks after 1 <sup>st</sup> dose)	<b>3<sup>rd</sup> Dose</b> (At least 4 weeks after 2 <sup>nd</sup> dose)			<b>Booster Dose<sup>1</sup></b> (at least 3 months after 3 <sup>rd</sup> dose)				<b>2<sup>nd</sup> Booster Dose<sup>3</sup></b> (See footnote)
<b>Janssen (ages 18 years and older)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> (Additional) Dose<sup>2</sup> using an mRNA COVID-19 Vaccine</b> (At least 4 weeks after 1 <sup>st</sup> dose)			<b>Booster Dose<sup>1</sup></b> (at least 2 months after additional dose)			<b>2<sup>nd</sup> Booster Dose<sup>3</sup></b> (See footnote)		

Note: Timeline is approximate. Intervals of 3 months or fewer are converted into weeks per the formula “1 month = 4 weeks.” Intervals of 4 months or more are converted into calendar months.

<sup>1</sup> An mRNA COVID-19 vaccine is preferred over the Janssen COVID-19 Vaccine for booster vaccination of people ages 18 years and older. For people ages 5 through 17 years, only Pfizer-BioNTech can be used for the first booster dose.

<sup>2</sup> Only Pfizer-BioNTech or Moderna COVID-19 Vaccine should be used. Visit [the CDC Interim Clinical Considerations](https://www.cdc.gov/media/releases/2022/s0927-covid-19-vaccine-immunocompromised.html) ([bit.ly/COVIDClinicalConsiderations](https://bit.ly/COVIDClinicalConsiderations)) for more information on vaccinating people who are moderately or severely immunocompromised and who received Janssen COVID-19 vaccine for the primary series.

<sup>3</sup> People ages 12 years and older who are moderately or severely immunocompromised should receive a second booster dose using an mRNA COVID-19 vaccine if it has been at least 4 months after the first booster dose. For people ages 12 through 17 years, only Pfizer-BioNTech can be used for the second booster dose.