

ULTRA BLAZER®

For use on sugar beets for management of Pigweed, Palmer Amaranth, and glyphosate-resistant Waterhemp,

For distribution and use only in Michigan under Section 18 emergency exemption

This labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its label. Read entire Directions for Use and Disclaimer before using this product. All applicable directions, restrictions, and precautions on the EPA registered product label for Ultra Blazer (EPA Reg. No. 70506-60) as well as those on these directions for use must be followed.

Any adverse effects resulting from the use of Ultra Blazer under this Section 18 Emergency Exemption must be immediately reported to the Michigan Department of Agriculture.

Effective date: June 1, 2021. Expiration date July 31, 2021. No applications of Ultra Blazer may be made under the emergency exemption after the expiration date.

A maximum acreage of 48,000 acres of sugar beets may be treated in Michigan in the counties of Arenac, Bay, Clinton, Eaton, Genesee, Gladwin, Gratiot, Huron, Ionia, Iosco, Isabella, Lapeer, Macomb, Midland, Montcalm, Ogemaw, Saginaw, Sanilac, Shiawassee, St. Clair, and Tuscola.

Product: Ultra Blazer Herbicide (EPA Reg. No. 70506-60)

Firm Name: UPL NA Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
For product information call 1-800-438-6071

Crop/Site/Commodity: Sugar beets

File Symbol: 21-MI-02

Target Pest/Problem: Pigweed, Palmer Amaranth, glyphosate-resistant Waterhemp

Application Rate: 16 fl. oz./A/year Ultra Blazer (0.25 lb ai)

Application Method: Apply by ground equipment only, aerial application and chemigation are prohibited

Application Volume: 10-20 gallons of spray solution per broadcast acre

Sugar beet Application

Timing: Make one application at greater than 6 leaf stage of sugar beet. Application before 6 leaf sugar beet will result in crop injury and potential yield loss.

Weed

Application Timing: For control of these weeds applications should occur at 3-4 inch weed height. Treating weeds greater than 4 inches in height or with incomplete coverage will result in reduced weed control.

Additives: Ultra Blazer may be tank mixed with a non-ionic surfactant, or with Roundup PowerMax + AMS.

Other Requirements, Restrictions, and

Precautions: Applications of Ultra Blazer will result in visual crop injury to exposed leaf tissue. Under normal growing conditions new leaf tissue will emerge without symptoms. This product is only effective on emerged weeds. This product will not control Group 14 (PPO inhibitor) resistant waterhemp or Palmer amaranth. Do not make more than one application per year. Do not apply more than 16 fl. oz. Ultra Blazer (0.25 lb ai) per acre per year. Application before 6 leaf sugar beet will result in crop injury and potential yield loss. Use of crop oil or methylated seed oil will significantly increase the potential for visual crop injury. Do not tank mix with pesticides other than glyphosate. Do not apply by air or chemigation to sugar beet.

Restricted Entry

Interval (REI): 48 hours. See additional Worker Protection information on the Section 3 labeling.

Pre-Harvest Interval

(PHI): 45 days.

**FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY:
1-866-673-6671**

**FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC:
1-800-424-9300 or +1-703-527-3887.**

Ultra Blazer is a registered trademark of a UPL Corporation Limited Group Company.