



COVID-19 Vaccination Schedule

For Those Who are Moderately or Severely Immunocompromised

Vaccine	0 month	1 month	2 month	3 month	4 month	5 month	6 month	7 month
Pfizer-BioNTech (ages 6 months – 4 years)	1st Dose	2nd Dose (3 weeks after 1 st dose)		3rd Dose (At least 8 weeks after 2 nd dose)				
Moderna (ages 6 months – 5 years)	1st Dose	2nd Dose (4 weeks after 1 st dose)	3rd Dose (At least 4 weeks after 2 nd dose)					
Pfizer-BioNTech (ages 5 years and older)	1st Dose	2nd Dose (3 weeks after 1 st dose)	3rd Dose (At least 4 weeks after 2 nd dose)			Bivalent Booster Dose^{1 3 4} (At least 2 months after last dose)		
Moderna (ages 6 years and older)	1st Dose	2nd Dose (4 weeks after 1 st dose)	3rd Dose (At least 4 weeks after 2 nd dose)			Bivalent Booster Dose^{1 3 4} (At least 2 months after last dose)		
Novavax (ages 12 years and older)	1st Dose	2nd Dose (3 weeks after 1 st dose)		Bivalent Booster Dose^{1 3 4} (At least 2 months after last dose)				
Janssen (ages 18 years and older)	1st Dose	2nd (Additional) Dose² using an mRNA COVID-19 Vaccine (4 weeks after 1 st dose)		Bivalent Booster Dose^{1 3 4} (At least 2 months after last dose)				

Note: Timeline is approximate. Intervals of 3 months or fewer are converted into weeks per the formula “1 month = 4 weeks.” Intervals of 4 months or more are converted into calendar months.

¹ For people ages 5 years and older, an age-appropriate COVID-19 bivalent vaccine booster dose is recommended, regardless of the primary series given.

² Janssen COVID-19 Vaccine should only be used in certain limited situations.

³ The bivalent booster dose is administered at least 2 months after completion of the primary series. For people who previously received a monovalent booster dose(s), the bivalent booster dose is administered at least 2 months after the last monovalent booster dose.

⁴ A monovalent Novavax booster dose may be used in limited situations in people ages 18 years and older who completed a primary series using any COVID-19 vaccine, have not received any previous booster dose(s), and are unable or unwilling to receive an mRNA vaccine. The monovalent Novavax booster dose is administered at least 6 months after completion of a primary series.

