


COVID-19 Vaccination Schedule*



Vaccine	0 month	1 month	2 month	3 month	4 month	5 month	6 month	7 month	8 month	9 month	10 month	11 month
Pfizer-BioNTech (ages 6 months – 4 years)	1st Dose	2nd Dose¹ (3–8 weeks after 1 st dose)		3rd Dose (at least 8 weeks after 2 nd dose)								
Moderna (ages 6 months – 5 years)	1st Dose	2nd Dose¹ (4–8 weeks after 1 st dose)										
Pfizer-BioNTech (ages 5–11 years)	1st Dose	2nd Dose¹ (3–8 weeks after 1 st dose)					Booster Dose² (at least 5 months after 2 nd dose)					
Pfizer-BioNTech (ages 12 years and older)	1st Dose	2nd Dose¹ (3–8 weeks after 1 st dose)					Booster Dose² (at least 5 months after 2 nd dose)				2nd Booster Dose³ (See footnote)	
Moderna (ages 18 years and older)	1st Dose	2nd Dose¹ (4–8 weeks after 1 st dose)					Booster Dose² (at least 5 months after 2 nd dose)				2nd Booster Dose³ (See footnote)	
Janssen (ages 18 years and older)	1st Dose		Booster Dose² (at least 2 months after 1 st dose)				2nd Booster Dose³ (See footnote)					

Note: Timeline is approximate. Intervals of 3 months or fewer are converted into weeks per the formula “1 month = 4 weeks.” Intervals of 4 months or more are converted into calendar months.

* See Guidance for COVID-19 Vaccination Schedule for People Who are Moderately or Severely Immunocompromised.

¹ An 8-week interval may be optimal for some people ages 6 months–64 years, especially for males ages 12–39 years. A shorter interval (3 weeks for Pfizer-BioNTech; 4 weeks for Moderna) between the first and second doses remains the recommended interval for: people who are moderately or severely immunocompromised; adults ages 65 years and older; and others who need rapid protection due to increased concern about community transmission or risk of severe disease. For more information, view [the CDC Interim Clinical Considerations](https://bit.ly/COVIDClinicalConsiderations) (bit.ly/COVIDClinicalConsiderations).

² All people ages 5 years and older should receive an age appropriate COVID-19 vaccine booster dose, an mRNA (Pfizer or Moderna) COVID-19 vaccine is preferred. For people ages 5-17 years, Pfizer COVID-19 vaccine is currently FDA authorized for use as a booster dose in people who received Pfizer as their primary series. Moderna COVID-19 vaccine is not authorized for use as a booster dose in this age group. For people ages 18 years and older, Pfizer-BioNTech and Moderna can be used as a booster dose.

³ People ages 18–49 who received Janssen COVID-19 vaccine as both their primary dose and first booster dose may receive a second booster dose using an mRNA vaccine at least 4 months after the first booster dose. People ages 50 years and older should receive a second booster dose if it has been at least 4 months after the first booster dose. An mRNA vaccine must be used for the second booster.

