


COVID-19 Vaccination Schedule

For Those Who are Moderately or Severely Immunocompromised



Vaccine	0 month	1 month	2 month	3 month	4 month	5 month	6 month	7 month	8 month	9 month
Pfizer-BioNTech (ages 6 months – 4 years)	1st Dose	2nd Dose (3 weeks after 1 st dose)		3rd Dose (At least 8 weeks after 2 nd dose)						
Moderna (ages 6 months – 17 years)	1st Dose	2nd Dose (4 weeks after 1 st dose)	3rd Dose (At least 4 weeks after 2 nd dose)							
Pfizer-BioNTech (ages 5-11 years)	1st Dose	2nd Dose (3 weeks after 1 st dose)	3rd Dose (At least 4 weeks after 2 nd dose)				Booster Dose¹ (at least 3 months after 3 rd dose)			
Pfizer-BioNTech (ages 12 years and older)	1st Dose	2nd Dose (3 weeks after 1 st dose)	3rd Dose (At least 4 weeks after 2 nd dose)				Booster Dose¹ (at least 3 months after 3 rd dose)			2nd Booster Dose³ (at least 4 months after 4 th dose)
Moderna (ages 18 years and older)	1st Dose	2nd Dose (4 weeks after 1 st dose)	3rd Dose (At least 4 weeks after 2 nd dose)				Booster Dose¹ (at least 3 months after 3 rd dose)			2nd Booster Dose³ (at least 4 months after 4 th dose)
Janssen (ages 18 years and older)	1st Dose	2nd (Additional) Dose² using an mRNA COVID-19 Vaccine (At least 4 weeks after 1 st dose)		Booster Dose¹ (at least 2 months after additional dose)				2nd Booster Dose³ (at least 4 months after 4 th dose)		

Note: Timeline is approximate. Intervals of 3 months or fewer are converted into weeks per the formula “1 month = 4 weeks.” Intervals of 4 months or more are converted into calendar months.

¹ If eligible, people 5 years and older should receive an age appropriate COVID-19 vaccine booster dose, an mRNA (Pfizer or Moderna) COVID-19 vaccine is preferred. For people ages 5-17 years who received Pfizer COVID-19 vaccine as their primary series should receive an age-appropriate COVID-19 vaccine booster dose. Those 6 months through 17 years who received Moderna COVID-19 vaccine as their primary series are not recommended to receive a COVID-19 booster dose as a booster dose is not authorized for use in this age group. For people ages 18 years and older, either Pfizer-BioNTech or Moderna may be used as a booster dose.

² Only Pfizer-BioNTech or Moderna COVID-19 vaccine should be used. Visit appendix D in [the CDC's Interim Clinical Considerations](#) for more information on vaccinating people who are moderately or severely immunocompromised and who received Janssen COVID-19 vaccine for the primary series.

³ People ages 18–49 years who received Janssen COVID-19 vaccine as both their primary dose and first booster dose may receive a second booster dose using an mRNA (Pfizer or Moderna) COVID-19 vaccine at least 4 months after the first booster dose. People ages 12 and older who are moderately or severely immunocompromised should receive an age appropriate second booster dose at least 4 months after the first booster. An mRNA (Pfizer or Moderna) COVID-19 vaccine must be used for the second booster.

