


# COVID-19 Vaccination Schedule



## For Those Who are Moderately or Severely Immunocompromised

Vaccine	0 month	1 month	2 month	3 month	4 month	5 month	6 month	7 month	8 month	9 month
<b>Pfizer-BioNTech (ages 6 months – 4 years)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (3 weeks after 1 <sup>st</sup> dose)		<b>3<sup>rd</sup> Dose</b> (At least 8 weeks after 2 <sup>nd</sup> dose)						
<b>Moderna (ages 6 months – 17 years)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (4 weeks after 1 <sup>st</sup> dose)	<b>3<sup>rd</sup> Dose</b> (At least 4 weeks after 2 <sup>nd</sup> dose)							
<b>Pfizer-BioNTech (ages 5-11 years)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (3 weeks after 1 <sup>st</sup> dose)	<b>3<sup>rd</sup> Dose</b> (At least 4 weeks after 2 <sup>nd</sup> dose)			<b>Booster Dose<sup>1</sup></b> (at least 3 months after 3 <sup>rd</sup> dose)				
<b>Pfizer-BioNTech (ages 12 years and older)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (3 weeks after 1 <sup>st</sup> dose)	<b>3<sup>rd</sup> Dose</b> (At least 4 weeks after 2 <sup>nd</sup> dose)			<b>Booster Dose<sup>1</sup></b> (at least 3 months after 3 <sup>rd</sup> dose)				<b>2<sup>nd</sup> Booster Dose<sup>3</sup></b> (at least 4 months after 4 <sup>th</sup> dose)
<b>Moderna (ages 18 years and older)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (4 weeks after 1 <sup>st</sup> dose)	<b>3<sup>rd</sup> Dose</b> (At least 4 weeks after 2 <sup>nd</sup> dose)			<b>Booster Dose<sup>1</sup></b> (at least 3 months after 3 <sup>rd</sup> dose)				<b>2<sup>nd</sup> Booster Dose<sup>3</sup></b> (at least 4 months after 4 <sup>th</sup> dose)
<b>Janssen (ages 18 years and older)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> (Additional) Dose<sup>2</sup> using an mRNA COVID-19 Vaccine</b> (At least 4 weeks after 1 <sup>st</sup> dose)		<b>Booster Dose<sup>1</sup></b> (at least 2 months after additional dose)				<b>2<sup>nd</sup> Booster Dose<sup>3</sup></b> (at least 4 months after 4 <sup>th</sup> dose)		
<b>Novavax (ages 18 years and older)</b>	<b>1<sup>st</sup> Dose</b>	<b>2<sup>nd</sup> Dose</b> (3 weeks after 1 <sup>st</sup> dose)								

Note: Timeline is approximate. Intervals of 3 months or fewer are converted into weeks per the formula “1 month = 4 weeks.” Intervals of 4 months or more are converted into calendar months.

<sup>1</sup> If eligible, people 5 years and older should receive an age appropriate COVID-19 vaccine booster dose, an mRNA (Pfizer or Moderna) COVID-19 vaccine is preferred. For people ages 5-17 years who received Pfizer COVID-19 vaccine as their primary series should receive an age-appropriate COVID-19 vaccine booster dose. Currently, a booster dose using any COVID-19 vaccine is not authorized for people ages 6 months through 17 years who receive a Moderna primary series or people ages 18 years and older who receive a Novavax primary series. For people ages 18 years and older, either Pfizer-BioNTech or Moderna may be used as a booster dose.

<sup>2</sup> Only Pfizer-BioNTech or Moderna COVID-19 vaccine should be used. Visit appendix D in [the CDC's Interim Clinical Considerations](#) for more information on vaccinating people who are moderately or severely immunocompromised and who received Janssen COVID-19 vaccine for the primary series.

<sup>3</sup> People ages 18–49 years who received Janssen COVID-19 vaccine as both their primary dose and first booster dose may receive a second booster dose using an mRNA (Pfizer or Moderna) COVID-19 vaccine at least 4 months after the first booster dose. People ages 12 and older who are moderately or severely immunocompromised should receive an age appropriate second booster dose at least 4 months after the first booster. An mRNA (Pfizer or Moderna) COVID-19 vaccine must be used for the second booster.

