

MI COVID RESPONSE DATA AND MODELING UPDATE

September 27, 2022

Epidemiologic Surveillance: Key Messages

COVID-19 pandemic is plateauing in some parts of the globe; decreases are seen in parts of the United States

- However, case rates in Europe are increasing
- Within the U.S., case rates decreased 10.6% over the past week
- Midwestern states (region 5) are showing signs of declines or plateaus

COVID spread in Michigan is plateaued

- COVID spread is assessed from many different markers including CDC community levels and other surveillance systems
- As of September 22, 48% of Michigan counties are at medium or high COVID-19 community levels
 - 7 Michigan counties are classified as High this week according to CDC's Community Levels (8%). This represents 12% of the population.
 - 33 Michigan counties are currently at Medium level (40%). This represents 65% of the population.
- The R_t for Michigan is around 1 indicating COVID is plateaued
- The proportion of specimens sequenced and identified as BA.5 in the U.S. and Michigan continues to remain the most dominant
- 50% of SWEEP sites saw a decrease in the most recent week; 35% of SWEEP sites saw increases

COVID-19 hospital metrics in Michigan remain lower than past surges



- COVID-19 hospital admissions, hospital census, ICU census, and pediatric census showed decreases this week

Global and National Trends

Daily new confirmed COVID-19 cases per million people

7-day rolling average. Due to limited testing, the number of confirmed cases is lower than the true number of infections.

Our World
in Data



Globally, 615,185,298 cases and 6,537,269 deaths (Data* through 9/26/2022)

- Case rates for some European countries are increasing


United States: Reported cases (7-day average) have decreased 10.6% since the prior week[†]

- In the U.S., the case rate is 112.5 cases/100,000 in last 7 days (last week: 126.2 cases per/100,000)


Region 5 (Midwest) states are declining or plateaued

- Indiana and Minnesota have the lowest case rates *in Region 5* (9/22/2022)

Local Prevention Decisions Should Use Community Levels in Concert with Other Pandemic Indicators



As of Sep 22, 7 Michigan Counties at High COVID-19 Community Level



- In the US, 7% of counties have high risk for medically significant disease and healthcare strain
- In Michigan, 8% (7/83) of counties are at high risk. This represents 12% of the population
- 33 Michigan counties are currently at Medium level (40%). This represents 65% of the population
- 43 Michigan counties are currently at Low level (52%). This represents 23% of the population

Percent of Counties

	United States	Michigan	Percent of MI Population
Low	62%	52%	23%
Medium	31%	40%	65%
High	7%	8%	12%



Low	Medium	High
<ul style="list-style-type: none"> • Stay <u>up to date</u> with COVID-19 vaccines • <u>Get tested</u> if you have symptoms 	<ul style="list-style-type: none"> • If you are <u>at high risk for severe illness</u>, talk to your healthcare provider about whether you need to wear a mask and take other precautions • Stay <u>up to date</u> with COVID-19 vaccines • <u>Get tested</u> if you have symptoms 	<ul style="list-style-type: none"> • Wear a <u>mask</u> indoors in public • Stay <u>up to date</u> with COVID-19 vaccines • <u>Get tested</u> if you have symptoms • Additional precautions may be needed for people <u>at high risk for severe illness</u>

Source: CDC COVID-19 Community Levels https://covid.cdc.gov/covid-data-tracker/#county-view?list_select_state=all_states&list_select_county=all_counties&data-type=CommunityLevels

Michigan Trends of COVID-19 Community Levels

- As of September 22, 7 (8%) Michigan counties at high COVID-19 community level and another 33 Michigan counties are currently at Medium level (40%)
- The proportion of Michigan counties at medium and high is lower than last week
- Current number of counties at high and medium are the lowest since July 28

Weekly Percent of MI Counties by CDC COVID-19 Community Level




This metric uses three indicators for categorization: (1) new COVID-19 cases per 100,000 population in the last 7 days lagged 1 day behind the date the COVID-19 Community Level is calculated; (2) new COVID-19 hospital admissions per 100,000 population in the last 7 days; and (3) percent of staffed inpatient beds occupied by patients with confirmed COVID-19 (7-day average) lagged 1 day behind the 7-day case rate .

Recent statewide trends are plateaued

Statewide trends

— 7-day average ■ Daily values


Daily Positive Test Rate
MICHIGAN STATEWIDE



Positivity, %


Current: 15.8%
Last Week: 17.0%

Daily cases per million




Current: 183.9
Last Week: 180.6

Daily hospitalization rate, %



Current: 5.6%
Last Week: 5.8%


Deaths





Current: 1.2
Last Week: 1.1

MERC Regional breakdown: Positivity, cases, hospitalization rate, and deaths


Positivity: 7-day average positivity, %
Cases: 7-day average cases per million
Hosp. rate: 7-day average hospitalization rate, %
Deaths: 7-day average deaths per million



Cases are plateaued in Michigan




18 counties currently showing increases and 14 in elevated incidence plateaus (via mistartmap.info as of 9/23/22, data through 9/15/22).




Identified COVID-19 Cases Caused by Variants of Concern (VOC) in US and Michigan: Predominately BA.5 and BA.4 lineages

SARS-CoV-2 Variants Circulating in the United States, Jul 10– Sep 24 (NOWCAST)



* Enumerated lineages are US VOC and lineages circulating above 1% nationally in at least one week period. "Other" represents the aggregation of lineages which are circulating <1% nationally during all weeks displayed.
 ** These data include Nowcast estimates, which are modeled projections that may differ from weighted estimates generated at later dates
 # AY.1-AY.133 and their sublineages are aggregated with B.1.617.2. BA.1, BA.3 and their sublineages (except BA.1.1 and its sublineages) are aggregated with B.1.1.529. Except BA.2.12.1, BA.2.75 and their sublineages, BA.2 sublineages are aggregated with BA.2. Except BA.4.6, sublineages of BA.4 are aggregated to BA.4. Except BF.7, sublineages of BA.5 are aggregated to BA.5. Sublineages of BA.1.1 and BA.2.75 are aggregated to the parental BA.1.1 and BA.2.75 respectively. Previously, BA.2.75 was aggregated with BA.2, and BF.7 was aggregated with BA.5.





VOC Distribution in Michigan



- Since August 15, there have 2,512 VOC specimens sequenced
- 100% of specimens sequenced are Omicron, 90% of those are BA.5 lineage
 - Since August 15, 9% of specimens sequenced and reported (n=216) have been identified as BA.4; however, 70% of those specimens are BA.4.6 (n=152)

Scenario Hub projections suggest plateau/smaller fall surge + potential winter surge

- Explored scenarios with early/late boosters and potential new variant
- Fall: smaller surge or plateau
- Winter
 - If no new immune escape variant, suggests plateau through winter (left two plots)
 - If new variant, potential for larger winter surge (right two plots)
- Similar patterns for cases and deaths (see link below)




Plotted with 50% uncertainty interval

Source: [Round 15 Scenario Modeling Hub Projections](#)



Case rates by age, race, and ethnicity

Daily new confirmed and probable cases per million by age group (7-day rolling average)



- Case rates by onset date for all age groups are between 129.0 and 244.9 cases per million (through 9/16)
- Case counts and case rates are highest for 80+-year-olds this week, followed by 30-39-year-olds and the 70-79-year-old age groups
- There are early indicators of potential increases in the lag period with the start of academic year (red box)

Daily new confirmed and probable cases per million (7 day rolling average) by race & ethnicity category





- Case rates are highest for Asian/Pacific Islander populations (160.5 cases/million)
- Between 22-28% of cases in last 30 days have missing race/ethnicity data

Note: Case information sourced from MDHHS and reflects date of onset of symptoms
Source: MDHHS – Michigan Disease Surveillance System

COVID case rates among K-12 age individuals are increasing

K-12 age population summary:

- Overall case counts among school-aged populations are increasing (7-day average 240.6)
- 60% (↓2%) of school district areas have between 1-10 cases.
- 11 ISD areas have greater than 5 cases.



Interpreting Wastewater Should Be In Context with Other Indicators

- When levels of virus in wastewater are low, a modest increase overall in virus level can appear much larger as numbers are translated into percentages
- This does not necessarily mean we will see major increases in transmission in the community
- When increases are seen within one wastewater site, public health officials compare with neighboring communities and other data sources to understand potential of surges
- For example, the Ypsilanti WWTP saw increases in SARS-CoV-2 levels which correlated with increasing presence of Omicron BA.2 lineage and then followed by an increase in cases

Ypsilanti WWTP

The most recent sample concentration is higher than 86% of samples collected at this site, which puts it in the 81-100 percentile category. As of 7/27/2022, the change in viral concentration over the past 15 days is increasing.


Wastewater SARS-CoV-2 Levels and COVID-19 Cases



The blue line on the graph shows the levels of SARS-CoV-2, the virus that causes COVID-19, in the wastewater samples collected from Ypsilanti WWTP. Each data point is calculated by averaging the number of viral gene copies detected per 100mL of wastewater in the 3 most recent samples. The orange bars on the graph show the COVID-19 cases reported to MDHHS from the zip codes that the wastewater treatment plant serves (7-day average). Both the virus levels and COVID-19 cases are calculated per 100,000 people. Case data will not be shown on the graph when the average number of cases is fewer than 10 per 100,000 people to protect the confidentiality of individuals with infections. This will be represented by an orange dashed line with gray shading below.

Michigan COVID-19 SWEEP Sentinel Wastewater Dashboard

The map below shows 20 sewershed sites in Michigan where wastewater is being monitored for the presence of SARS-CoV-2, the virus that causes COVID-19. These sentinel sites serve as a subset of wastewater surveillance in Michigan distributed across the Michigan Economic Recovery Council (MERC) Regions. Click on each site on the map to see wastewater and clinical case data over time. To view wastewater data from previous weeks, please use the "Map - All Data" and "Trends - All Data" tabs.




Site	Population	Consecutive Weeks of Virus Detection	Trend As Of	15-Day Trend
Alma WWTP	8976	22	9/12/2022	↓
Battle Creek WWTP	51093	22	9/14/2022	↓
Bay City WWTP	34000	13	9/15/2022	↑
Delhi Township WWTP	22500	24	9/8/2022	↑
Escanaba WWTP	12600	20	9/14/2022	↓
GLWA Detroit River Interce..	492000	9	9/8/2022	↓
GLWA North Interceptor-Ea..	1482000	76	9/8/2022	↔
GLWA Oakwood-Northwest..	840600	99	9/8/2022	↓
Grand Rapids WWTP	265000	58	9/15/2022	↔
Holland WWTP North	45606	22	9/14/2022	↑
Holland WWTP South	36912	24	9/14/2022	↓
Jackson WWTP	90000	61	9/15/2022	↑
Kalamazoo WWTP	150000	25	9/15/2022	↑
Petoskey WWTP	7900	22	9/14/2022	↓
Portage Lake WWTP	14000	53	9/14/2022	↓
Saginaw Township WWTP	40000	23	9/15/2022	↓
Tecumseh WWTP	8680	36	9/16/2022	↑
Traverse City WWTP	45000	27	9/15/2022	↓
Warren WWTP	135000	21	9/8/2022	↑
Ypsilanti WWTP	330000	61	9/15/2022	↔

Abbreviations: GLWA - Great Lakes Water Authority; WWTP - Waste Water Treatment Plant

Definitions and descriptions of data calculations can be found in the "About" tab.

Current results reflect data that were uploaded to MDHHS as of 9/21/2022. Labs are required to report test results to local partners within 24 hours. Data is subject to change as additional wastewater data and case data are received.




SWEEP Summary


- 35% (7/20) of sentinel sites are showing increasing trends over last 15-days
- 15% (3/20) of sites have plateaued over the last 15 days
- 50% (10/20) of sentinel sites are showing declines in the previous 15-days

Cases Among Staff and Residents in Long Term Care Facilities

STATE OF MICHIGAN WEEKLY TOTAL CONFIRMED COVID-19 CASES IN
AFC/HFA RESIDENTS AND STAFF
09/22/2021 TO 09/21/2022



STATE OF MICHIGAN WEEKLY TOTAL CONFIRMED COVID-19 CASES IN
SNF
RESIDENTS AND STAFF
09/24/2021 TO 09/23/2022



- Case counts in residents increased in AFC/HFA (129 to 165), but decreased in SNFs (337 to 307) since last week
- Case counts in staff increased in AFC/HFA (145 to 155), however has been steady in SNFs (479) since last week
- **29%** of SNFs are reporting **nursing shortages** and **31%** of SNFs are reporting **aide shortages**, which is stable from last week

Abbreviations: AFC: Adult Foster Care; HFAs: Homes for the Aged; and SNF: Skilled Nursing Facilities

Reported Number of Clusters in Long Term Care Facilities

Number of AFC/HFAs with 3 or more Confirmed Cases




Number of SNFs with 3 or more Confirmed Cases



- The number of Long-Term Care Facilities reporting 3 or more cases within a single reporting period has increased over the past week
- This week, the number has increased in **AFC/HFAs** (15 to 19) and in **SNFs** (33 to 43) since the previous week

The data is from weekly reporting by facilities with bed occupancy of at least 13 beds.

Hospital admissions due to COVID-19 remain lower than past surges




- Trends for daily average hospital admissions decreased (-10%) compared to last week (vs. +2% prior week)
- Nearly all age groups of saw decreases or plateaus this week
- Those 60-69, 70-79, and 80+ are seeing between 20 and 40 daily hospital admissions

Source: CHECC & EM Resource


Statewide Hospitalization Trends: Total COVID+ Census

Hospitalization Trends 7/1/2021 – 9/26/2022
Confirmed Positive & Persons Under Investigation (PUI)




COVID+ census in hospitals has decreased 4% from last week. Overall census is currently 1,009 patients.

Hospitalized COVID Positive Long Term Trend (beginning March 2020)



Statewide Hospitalization Trends: Regional COVID+ Census

Hospitalization Trends 7/1/2021 – 9/26/2022
Confirmed Positive by Region




This week hospitalizations have decreased or remained flat in Regions 1, 2N, 2S, 5, 6, and 8. Hospitalizations have increased in Regions 3 and 7.

Regions 2N, 2S, and 3 have greater than 100 hospitalizations/M.

Region	COVID+ Hospitalizations (% Δ from last week)	COVID+ Hospitalizations / MM
Region 1	87 (0%)	80/M
Region 2N	247 (-5%)	112/M
Region 2S	281 (-2%)	126/M
Region 3	139 (11%)	123/M
Region 5	82 (-19%)	86/M
Region 6	126 (-13%)	86/M
Region 7	31 (7%)	62/M
Region 8	16 (-16%)	51/M

Statewide Hospitalization Trends: ICU COVID+ Census

Hospitalization Trends 7/1/2021 – 9/26/2022
Confirmed Positive in ICUs



Overall, the volume of COVID+ patients in ICUs has decreased by 4% from last week. There are 117 COVID+ patients in ICU beds across the state.

ICU occupancy is less than 85% in all regions except Region 3. All regions have fewer than 10% of ICU beds occupied by COVID+ patients.

Region	Adult COVID+ in ICU (% Δ from last week)	ICU Occupancy	% of ICU beds COVID+
Region 1	12 (9%)	81%	7%
Region 2N	21 (-22%)	66%	4%
Region 2S	36 (-8%)	77%	5%
Region 3	20 (11%)	85%	7%
Region 5	5 (0%)	66%	3%
Region 6	16 (14%)	78%	7%
Region 7	4 (-33%)	76%	3%
Region 8	3 (50%)	55%	5%

Statewide Hospitalization Trends: Pediatric COVID+ Census


Hospitalization Trends 1/1/2021 – 9/26/2022
Pediatric Hospitalizations, Confirmed + PUI



Percent of Inpatients with COVID is Similar to One Year Ago During the Delta Wave


- The percent of inpatients who are COVID+ remains lower than Alpha, Omicron, and holiday wave peaks
- Current hospital levels are plateaued and between levels seen during fall 2020 and fall 2021

7-day rolling average of percent of inpatients who are COVID positive




Average new deaths have plateaued for those over the age of 80

Daily COVID-19 deaths in confirmed and probable cases per million by age group (7 day rolling average)



Total COVID-19 deaths in confirmed and probable cases by age group (past 30 days, ending 9/16/2022)

- 9.9% of deaths below age sixty



- Through 9/16, the 7-day avg. death rate has plateaued (13.2 deaths per million people) for those over the age of 80
- In the past 30 days, there are fewer than 25 confirmed and probable COVID-19 deaths under the age of 50
- 30-day proportion of deaths among those under 60 years of age is 9.9%.

Harm Reduction: Key Messages

Empowering community members to make best choices for their individual circumstances and to be prepared by making a COVID plan



- Michiganders can take advantage of local, state, and national COVID-19 resources
- Get tested, and if positive, seek care with therapeutics (e.g., antibodies or antiviral medications)
 - Cumulative therapeutic availability and administration is plateaued for the last month and below all-time highs
 - Talk to your doctor or pharmacist about whether you should get antibody or antiviral treatment, and where you can find treatment
 - Therapeutics are authorized for people who meet select criteria
 - Additional public health, regulatory, and policy efforts might help decrease barriers to oral antiviral access, particularly in communities with high social vulnerability
- Vaccinations remain the best way to protect from COVID-19, especially from severe disease
 - COVID-19 vaccines are now available for ages 6 months and up
 - Everyone 6 months and older should also get an age-appropriate COVID-19 booster, when eligible
 - Data tracker now includes Novavax administration
 - Over 6.8 million Michiganders have received at least one dose (68.3%)
 - 56.6% of fully vaccinated Michiganders have received at least one booster
 - 36.3% of people in Michigan (793K+) with a first booster dose have received a second booster dose
 - Over 176,000 bivalent booster doses had been administered as of 9/20

Federal & Michigan websites assist COVID positive residents find treatment

COVID-19 resources available on federal website: [COVID.gov](https://www.covid.gov)

Test-to-Treat program simplifies access to COVID treatment: [Find a Test-to-Treat location near you](#)

- If you have COVID-19 symptoms, do not wait to get treated
- You must take oral COVID-19 medication within 5 days of your first COVID-19 symptoms
- Use the tool to find a location that is right for you



Source: Screen capture of Michigan Test-to-Treat sites from linked website

Therapeutic administration increased during Michigan's Spring Omicron surge. Supply limitations in January 2022 required strategic distribution and should not be compared directly.

Source: HHS – Tiberius


*Data is reported as a single patient course, except for Evusheld, which is reported as the number of 300mg doses administered. Data Updated September 19

^Federally supplied Bebtelovimab has concluded, and product has transitioned to the commercial marketplace

Vaccinations and Boosters

- Over 16.9 million COVID-19 vaccine doses have been administered in Michigan
 - Over 6.8 million Michiganders have received at least one dose (68.3%)
 - Over 6.1 million Michiganders have completed a primary series (61.2%)
 - Over 3.4 million additional/booster doses have been administered in Michigan
 - 56.6% of the fully vaccinated population has received a booster
 - 78.1% of the fully vaccinated population 65 years of age or older has received a booster
- Nearly 793,872 Michiganders 50 years of age or older who have received a first booster dose have received second booster (36.3%)

COVID-19 Vaccine Coverage by Age Group



Note: Cumulative data and population percentages now include those ages 6 months and older


Sources: Michigan Coronavirus Vaccine Dashboard; CDC COVID-19 Data Tracker: Vaccine Coverage Dashboard

Bivalent (Omicron) Pfizer and Moderna COVID-19 vaccines available for booster shots in Michigan

- The Moderna and Pfizer bivalent boosters target two strains of COVID-19: the original strain of the virus and the widely spread Omicron variants (BA.4 and BA.5, including BF.7)
- **Who is eligible to receive a single bivalent booster dose and when:**
 - Individuals 18 years of age and older are eligible for a single booster dose of the bivalent **Moderna** COVID-19 vaccine if it has been at least two months since they completed primary vaccination or received the most recent booster dose with any authorized or approved monovalent COVID-19 vaccine
 - Individuals 12 years of age and older are eligible for a single booster dose of the bivalent **Pfizer-BioNTech** COVID-19 vaccine if it has been at least two months since they completed primary vaccination or received the most recent booster dose with any authorized or approved monovalent COVID-19 vaccine
- Individuals may choose to receive either the Pfizer or Moderna bivalent booster, regardless of which primary series vaccine or original booster dose they had previously.
- Influenza vaccines, which are now available in Michigan, can also be co-administered with the COVID-19 bivalent booster doses

Bivalent Administration


- As of 9/20, 176,457 Michiganders had received their bivalent booster
- These data are updated every Wednesday on our COVID-19 vaccination Dashboard under Additional/Booster Administration Trends and then restricting the view to just Moderna and Pfizer bivalent doses
- Note: the data for the week ending 9/17 would have been incomplete on the date the dashboard was last refreshed (9/13)



Co-administration of Flu and COVID-19 Vaccines

- Coadministration of influenza and COVID-19 vaccines has proven to be safe and effective*¹, is recommended by the CDC², and provides an efficient way to immunize the population against two potentially serious illnesses
- During the 2021-2022 season (Sep 1, 2021 – Feb 12, 2022), 13.4% (413,101/3,075,658) of all flu vaccines administered in Michigan were co-administered with COVID-19 vaccines and highest among 18–49-year-olds
 - Pharmacies reported the most coadministrations of flu and COVID vaccines during this season
- Educational outreach to pharmacies, other health care providers, and the general public to co-administer flu and COVID vaccines can be an effective strategy to enhance immunizations

Age group	Flu-COVID Coadministrations (%)
5 – 11 years	283 (0.1%)
12 – 17 years	11,144 (2.5%)
18 – 49 years	161,949 (37.0%)
50 – 64 years	114,210 (26.1%)
65 years and above	150,025 (34.3%)




* Routine vaccinations for all persons aged ≥ 6 months who do not have contraindications

Sources: Michigan Department of Health and Human Services: Division of Immunization; **1.** Toback S, et al. Safety, immunogenicity, and efficacy of a COVID-19 vaccine (NVX-CoV2373) co-administered with seasonal influenza vaccines: an exploratory substudy of a randomised, observer-blinded, placebo-controlled, phase 3 trial. *Lancet Respir Med.* 2022 Feb;10(2):167-179. doi: 10.1016/S2213-2600(21)00409-4. Epub 2021 Nov 17. **2** Grohskopf LA et al. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices, United States, 2021–22 Influenza Season. *MMWR Recomm Rep* 2021;70(No. RR-5):1–28. DOI: <http://dx.doi.org/10.15585/mmwr.rr7005a1>

COVID-19 mRNA Vaccine is Safe Among Children Aged 6 Months–5 Years

- Approximately 1 million COVID-19 vaccine doses were administered to children 6 months to 5 years between June and August 2022, following ACIP and CDC recommendations
- Reports of serious adverse events were rare and there were 0 reports of myocarditis
- Everyone 6 months and older should receive COVID-19 vaccine



Data support the safety of COVID-19 vaccination for children ages 6 months to 5 years*


More than 1 million children were vaccinated; reports of serious adverse events were rare**

Most reported reactions were mild or moderate:

- Pain in the arm where the shot was given
- Irritability
- Crying
- Sleepiness


No reports of myocarditis†

Everyone ages 6 months and older should receive recommended COVID-19 vaccines to protect against severe illness and death



* Study of 21,915 children enrolled in v-safe and 667 reports received and processed by VAERS — United States, June 18–August 21, 2022
** 19 serious reports to VAERS after approximately 1 million children vaccinated. VAERS reports are classified as “serious” only if one of the following events are reported: hospitalization, prolongation of hospitalization, life-threatening illness, permanent disability, congenital anomaly or birth defect, or death.
† Myocarditis, or inflammation of the heart muscle, is a rare adverse event that has been associated with mRNA COVID-19 vaccines

bit.ly/mm7135a3
SEPTEMBER 2, 2022






APPENDIX

Ohio, Indiana

Ohio Confirmed New Cases / 1M (7 days average)




Indiana Confirmed New Cases / 1M (7 days average)






Wisconsin, Michigan

Wisconsin Confirmed New Cases / 1M (7 days average)




Michigan Confirmed New Cases / 1M (7 days average)






Illinois, Minnesota

Illinois Confirmed New Cases / 1M (7 days average)



Minnesota Confirmed New Cases / 1M (7 days average)




CDC COVID-19 Community Levels are defined by County Case Rates and Health Service Area (HSA) Hospitalizations

COVID-19 Community Levels – Use the Highest Level that Applies to Your Community


New COVID-19 Cases Per 100,000 people in the past 7 days	Indicators	Low	Medium	High
Fewer than 200	New COVID-19 admissions per 100,000 population (7-day total)	<10.0	10.0-19.9	≥20.0
	Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average)	<10.0%	10.0-14.9%	≥15.0%
200 or more	New COVID-19 admissions per 100,000 population (7-day total)	NA	<10.0	≥10.0
	Proportion of staffed inpatient beds occupied by COVID-19 patients (7-day average)	NA	<10.0%	≥10.0%

Health Service Areas




Adjusted new cases by on-set

New confirmed cases by onset actual and adjusted as of September 20, 2022 (-2 days)




Adjusted new cases by on-set, recent trends

New confirmed cases by onset actual and adjusted as of September 20, 2022 (-2 days)




Daily average deaths per million people by race and ethnicity have plateaued

Average daily deaths per million people by race



Average daily deaths per million people by ethnicity



- Deaths are lagging indicator of other metrics
- Currently, the American Indian/Alaska Native population has the highest death rate (2.17 deaths/million)

Note: Death information sourced from MDHHS and reflects date of death of confirmed and probable cases.
Source: MDHHS – Michigan Disease Surveillance System