

MI COVID RESPONSE DATA AND MODELING UPDATE

NOTE: All data as of August 16 unless otherwise noted

August 17, 2021

Executive Summary – All Indicators Show Increases

Michigan is now in [High Transmission](#)

Percent Positivity (7.7%) is increasing for seven weeks (up from 7.0% last week), and **Case Rate** (102.0 cases/million) have increased for over a month (up from 77.2 last week)

Michigan has 30th lowest number of cases (33rd last week), and 5th lowest case rate (9th last week) in the last 7 days

99% of positive tests available for sequencing in Michigan were **Delta variant** in the last 4 weeks

Percent of inpatient beds occupied by individuals with COVID (4.0%) has increased for four weeks (up from 2.9% last week).

Michigan has 9th lowest inpatient bed utilization (10th last week), and 10th lowest adult ICU bed utilization (10th last week)

Deaths (0.6 deaths/million) are increasing (0.5 deaths/million last week). There were 41 COVID deaths between Aug 3 and Aug 9.

Michigan has the T28th lowest number of deaths (21st last week), and T16th lowest death rate (T6th last week) in the last 7 days

7-day average **state testing rate** is steady at 2,068.0 tests/million/day. **Daily diagnostic tests (PCR)** is 20.5K per day, and the **weekly average for PCR and antigen tests** conducted in Michigan is 34.3K.

9.98 million **COVID-19 vaccine** doses reported to CDC, 4.95 million people have completed their vaccine series

Science Round Up

Models are projecting a continued increase in hospitalizations and deaths over the next four to six weeks, maybe longer

Children can be impacted by the spread of COVID-19

A larger proportion of those who become cases (98%), are hospitalized (95%), and died (95%) from COVID are unvaccinated

mRNA vaccine are 96% effective at preventing hospitalizations among elderly

Among individuals previously infected, vaccination provides additional protection to prevent reinfection


Layered mitigation, especially for those not vaccinated and those not yet eligible for vaccination can avoid unnecessary surge in cases and unintended school closures due to classroom outbreaks

Global and National Comparisons: US cases increasing

What we see today (data through 8/16):

- Globally, 207,366,191 cases and 4,365,541 deaths*
- Countries with the highest case count are U.S. (36,681,559), India (1,081,284), and Brazil (987,525)*
- Nearly all US jurisdictions have high community transmission
- Within the U.S., Rhode Island (14,876/100,000), North Dakota (14,858/100,000), & South Dakota (14,327/100,000) lead the nation in cumulative cases/capita
- Michigan has had 10,300 cases per 100,000 since March 1, 2020
- Michigan currently has identified 14,922 variants of concern (VOC)*
 - Cumulatively, the vast majority are B.1.1.7 (13,652 which is 91.5%)
 - Other VOCs include B.1.351 (0.6%), P.1 (2.2%) and B.1.617.2 (5.7%)
 - 571 VOC reported with specimen date in the 4 most recent weeks
 - Delta (B.1.617.2) 99%
 - Alpha (B.1.1.7) <1%
 - Gamma (P.1) <1%

* CDC removed Epsilon (B.1.427/B.1.429) from the lists of VOCs



National Comparison

Spread

Severity


Public Health
Response

Other
Indicators





Science
Round-up

State Comparisons: Missouri and Arkansas

Missouri Confirmed New Cases / 1M (7 days average)




Arkansas Confirmed New Cases / 1M (7 days average)








State Comparisons: Florida and Michigan

Florida Confirmed New Cases / 1M (7 days average)




Michigan Confirmed New Cases / 1M (7 days average)








State Comparisons: Nevada and Louisiana

Nevada Confirmed New Cases / 1M (7 days average)



Louisiana Confirmed New Cases / 1M (7 days average)



Key Messages: COVID-19 is Spreading Faster with Delta

Statewide positivity has increased to 7.7% (last week: 7.0%)

- One week percent change is up 11% (vs. up 21% last week)
- Increasing for seven weeks (over 6 times higher the Jun 26 low)
- Positivity is increasing in all MERC regions; five regions are >7% and one region > 10%

Case rate (102.0 cases/million) increasing for over a month (last week: 77.2 cases/million)

- One week increase of 15% (vs. 34% increase last week)
- Increasing for over a month (611% increase since Jun 26 low)
- Cases per million are increasing in all MERC regions
- Select variants in Michigan: 13,652 confirmed Alpha (B.1.1.7); 85 confirmed Beta (B.1.351); 329 confirmed Gamma (P.1); and 856 confirmed Delta (B.1.617.2)

Michigan is at High Transmission level

- More than half of the counties in Michigan are at high transmission level
- CDC would recommend all individuals, regardless of vaccination status, should mask indoors
- The U.S. is at high transmission level (252 cases/100,000 in last 7 days) with 54 states/territories in substantial or high transmission

Number of active outbreaks is up 36% from last week

- Eighty-six new outbreaks were identified in the past week
- Manufacturing/Construction reported the most new and ongoing outbreaks this week

National Comparison

Spread

Severity


Public Health
Response

Other
Indicators

Science
Round-up


Confirmed and probable case indicators

Table Date: 8/16/2021 (7 days from date table was produced: 8/9/2021)



	CDC Transmission Risk Level	Absolute Cases (per million)	CDC Case Trend	Average Percent Positivity	Positivity Trend	Tests (per million)	% IP Beds Occupied by COVID-19 Cases	% Occupied IP Beds Trend	Absolute Deaths (per million)	Death Trend
Detroit	High	100.4	elevated incidence growth	6.8	Increase - 4wk	2196.5	4.2	Increase - 4wk	0.4	Increase - 1wk
Grand Rapids	Substantial	89.3	elevated incidence growth	8.6	Increase - 6wk	2017.4	3.6	Increase - 4wk	0.8	<20 wkly deaths
Kalamazoo	High	126.7	elevated incidence plateau	10.7	Increase - 6wk	1728.1	5.0	Increase - 2wk	0.6	<20 wkly deaths
Saginaw	High	112.0	elevated incidence growth	9.7	Increase - 6wk	1557.4	2.8	Increase - 3wk	0.7	<20 wkly deaths
Lansing	High	99.2	elevated incidence growth	7.3	Decrease - 1wk	1773.0	4.1	Increase - 4wk	0.0	<20 wkly deaths
Traverse City	Substantial	97.7	elevated incidence plateau	5.8	Decrease - 1wk	1806.0	4.0	Increase - 3wk	2.6	<20 wkly deaths
Jackson	High	112.6	elevated incidence growth	9.3	Increase - 6wk	2022.1	6.9	Increase - 4wk	0.9	<20 wkly deaths
Upper Peninsula	High	97.8	elevated incidence growth	6.8	Increase - 3wk	1407.4	1.8	Decrease - 1wk	0.0	<20 wkly deaths
Michigan	High	102.0	elevated incidence growth	7.7	Increase - 7wk	2068.0	4.0	Increase - 4wk	0.6	Increase - 2wk

Cases



Positivity



National Comparison

Spread

Severity

Public Health Response

Other Indicators

Science Round-up

Overview of metrics for individuals <12 years


	Region	Population (<12 yrs)	Population (<18 yrs)	Cumulative Case Count (<12 yrs)	7-day Average Daily Case Count (<12 yrs)	7-day Average Daily Case Rate per Million (<12 yrs)	7-day Average Daily Pediatric Hospitalization Count (<18 yrs)	7-day Average Daily Pediatric Hospitalization Rate per Million (<18 yrs)	7-day Average Daily Death Count (<12 yrs)
1	Detroit	735529	1134247	29870	47.1	64.0	16.7	14.7	0
2	Grand Rapids	230120	350652	10154	12.9	56.1	5.6	16.0	0
3	Kalamazoo	140422	214801	5571	13.0	92.6	0.9	4.2	0
4	Saginaw	78759	122834	3367	5.7	72.4	0.0	0.0	0
5	Lansing	78140	119915	3310	5.6	71.7	3.3	27.5	0
6	Traverse City	53099	83462	1617	3.4	64.0	0.6	7.2	0
7	Jackson	41274	64091	1546	2.4	58.1	0.4	6.2	0
8	Upper Peninsula	34645	53875	1454	2.1	60.6	0.3	5.6	0
99	Michigan	1391988	2143877	56944	92.7	66.6	27.7	12.9	0

Note: Data as of 8/16; case data 8/9, hospitalization data 8/16. Hospitalization data is for pediatric patients (<18)



Michigan at High Transmission Level

[Dashboard](#) | [CDC](#) | [MI Start Map](#) for most recent data by reporting date



National Comparison

Spread

Severity



Public Health
Response

Other
Indicators

Science
Round-up

Cases per Capita Increasing Statewide

Daily new confirmed and probable cases per million (7-day rolling average) *by Onset Date*




- Case rates by onset date in Michigan are in substantial transmission
- Case rates in most counties are above 40 cases per million population

Note: Case information sourced from MDHHS and reflects date of onset of symptoms
Source: MDHHS – Michigan Disease Surveillance System



Case Rate Trends are Increasing for All Age Groups

Daily new confirmed and probable cases per million by age group (7-day rolling average)



- Case rate trends for all age groups are increasing
- Case rates for all age groups are between 60 and 145 cases per million (through 8/9)
- Case rate trends are highest for 20-29-year-olds followed by 30-39, 40-49, and 10-19

Note: Case information sourced from MDHHS and reflects date of onset of symptoms
Source: MDHHS – Michigan Disease Surveillance System

National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up

Number of Cases and Case Rates are Increasing for All Age Groups

Daily new confirmed and probable cases per million by age group (7-day rolling average)

Age Group	Average daily cases	Average Daily Case Rate	One Week % Change (Δ #)
0-9	70.3	61.0	19% (+11)
10-19	125.9	100.3	11% (+12)
20-29	201.0	145.7	9% (+16)
30-39	164.6	135.7	13% (+19)
40-49	146.4	124.2	18% (+22)
50-59	125.4	92.9	16% (+17)
60-69	107.4	84.2	28% (+23)
70-79	54.7	71.4	18% (+8)
80+	24.7	59.7	21% (+1-5)
Total [¶]	1022.7	102.0	15% (+133.6)

Note: Case information sourced from MDHHS and reflects date of onset of symptoms
Source: MDHHS – Michigan Disease Surveillance System

- Average daily number of cases (201) is highest for those aged 20-29
- Avg. daily case rate (145.7 cases/mil) is currently highest for 20-29
- Case rates for all age groups are between 60-145 cases per million
- Case rate trends are increasing for all age groups
- Case rates bottomed out on June 26, 2021

* Highest 7-day avg. following spring 2021 surge

[¶] Total may not reflect state due to missing age data

National Comparison

Spread

Severity


Public Health
Response

Other
Indicators


Science
Round-up

Racial and Ethnic Case Rates are Increasing

Daily new confirmed and probable cases per million (7 day rolling average) by race category



Daily new confirmed and probable cases per million (7 day rolling average) by ethnicity category



Updates since last week:

- Cases per million are increasing for all races and ethnicities
- **Hispanics and Whites have the highest case rates**
- In the past 30 days, 17% (↔) of race data and 20% (↔) ethnicity data was either missing or reported as unknown

Note: Case information sourced from MDHHS and reflects date of death of confirmed and probable cases.
Source: MDHHS – Michigan Disease Surveillance System

National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up

Identified COVID-19 Cases Caused by All Variants of Concern (VOC) in US and Michigan

Variants Circulating in United States, Jul 18 – Jul 31 (NOWCAST)

8/7/21

B.1.617.2

AY.3


USA				
WHO label	Lineage #	Type	%Total	95%PI
Alpha	B.1.1.7	VOC	0.9%	0.2-2.0%
Beta	B.1.351	VOC	0.0%	0.0-0.2%
Gamma	P.1	VOC	0.5%	0.0-1.2%
Delta	B.1.617.2	VOC	83.4%	79.7-87.1%
	AY.3	VOC	13.4%	10.2-16.9%
	AY.2	VOC	0.5%	0.0-1.2%
	AY.1	VOC	0.1%	0.0-0.2%
Eta	B.1.525	VOI	0.0%	0.0-0.2%
Iota	B.1.526	VOI	0.1%	0.0-0.5%
	B.1.621		0.6%	0.0-1.5%
	B.1.621.1		0.2%	0.0-0.7%
	B.1.628		0.2%	0.0-0.7%
	Other*		0.1%	0.0-0.5%
	A.2.5		0.0%	0.0-0.2%
	B.1.626		0.0%	0.0-0.2%
	B.1.429	VOI	0.0%	0.0-0.2%
	B.1.427	VOI	0.0%	0.0-0.2%

* Enumerated lineages are VOI/VOC or are circulating >1% in at least one HHS region during at least one two week period; remaining lineages are aggregated as "Other".

** These data include Nowcast estimates, which are modeled projections that may differ from weighted estimates generated at later dates

Sublineages of P.1 and B.1.351 (P.1.1, P.1.2, B.1.351.2, B.1.351.3) are aggregated with the parent lineage and included in parent lineage's proportion. AY.3.1 is aggregated with its parent lineage AY.3.

Variants of Concern in Michigan, Aug 16



Variant	MI Reported Cases [¶]	# of Counties	% Specimens in last 4 wks
B.1.1.7 (alpha)	13,652*	81	<1%
B.1.351 (beta)	85	24	<1%
P.1 (gamma)	329	35	<1%
B.1.617.2 (delta)	856 (↑506)	58 (↑8)	99%

* 534 cases within MDOC; [¶] 37 cases with county not yet determined

Data last updated Aug 16, 2021

Source: <https://covid.cdc.gov/covid-data-tracker/#variant-proportions> and MDSS

National Comparison

Spread

Severity


Public Health
Response

Other
Indicators

Science
Round-up

Number of Outbreaks Reported has Increased

Number of outbreak investigations by site type, week ending Aug 12



1. Based on a setting's level of control and the extent of time patrons/residents spend in the particular setting, different settings have differing levels of ability to ascertain whether a case derived from that setting

NOTE: Many factors, including the lack of ability to conduct effective contact tracing in certain settings, may result in significant underreporting of outbreaks. This chart does not provide a complete picture of outbreaks in Michigan and the absence of identified outbreaks in a particular setting in no way provides evidence that, in fact, that setting is not having outbreaks.

Source: LHD Weekly Sitreps

Key Messages: COVID-19 and Healthcare Capacity and COVID Severity

Hospitalizations and ICU utilization are increasing

- 2.1% of ED visits are for COVID-like illness (CLI) (up from 1.6% last week)
- Hospital admissions are increasing for all age groups under 80 years this week
- Hospitalizations up 23% since last week (vs. 52% increase week prior)
- All regions are showing increasing trends in hospitalization trends this week, except Region 7
 - Hospitalization for COVID-19 is highest in Regions 2N, 2S, 3 and 6
 - Fastest growth is in Regions 3, 6 and 8
- Volume of COVID-19 patients in intensive care has increased 15% since last week (vs. 57% increase last week)

Death rate is 0.6 daily deaths per million people

- Death rate has increased two weeks
- 41% increase since Jul 22 low
- 30-day proportion of deaths among those under 60 years of age is steady from the prior week

National Comparison

Spread

Severity


Public Health
Response

Other
Indicators

Science
Round-up

Michigan Trends in Emergency Department (ED) Visits for COVID-19-Like Illness (CLI) saw the largest increase in over 3 months

Percentage of Emergency Department visits with Diagnosed COVID-19 in Michigan, All Ages



- Trends for ED visits have increased to 2.1% since last week (up from 1.6% week prior)
- Trends vary by age groups with all age groups seeing an increase
- Over the past week, those 40-49 years have seen the highest number of avg. daily ED CLI visits, but those between 25 and 49 are all above the state average

Source: <https://covid.cdc.gov/covid-data-tracker/#ed-visits>

National Comparison

Spread


Severity

Public Health
Response

Other
Indicators

Science
Round-up

Average Hospital Admissions Are Increase for all Age Groups




Source: CHECC & EM Resource

- Trends for daily average hospital admissions have increased 28% since last week (vs. 68% increase prior week)
- This week, all age groups under 80 have experienced increases in daily hospital admissions
- Over the past week, those 60-69 years have seen the highest number of avg. daily hospital admissions (25 admissions)


Statewide Hospitalization Trends: Total COVID+ Census

Hospitalization Trends 3/1/2021 – 8/16/2021
Confirmed Positive & Persons Under Investigation (PUI)




COVID+ census in hospitals has increased 23% from last week (previous week was up 52%). The rate of growth in hospitalizations has slowed from last week.

Hospitalized COVID Positive Long Term Trend (beginning March 2020)



Statewide Hospitalization Trends: Regional COVID+ Census

Hospitalization Trends 3/1/2021 – 8/16/2021
Confirmed Positive by Region




All regions except Region 7 show increasing hospitalization trends this week. The fastest growth is in Regions 3, 6 and 8.

All regions except Region 8 are above 50/M population hospitalized with Regions 2S, 2N and 3 approaching 100/M.

Region	COVID+ Hospitalizations (% Δ from last week)	COVID+ Hospitalizations / MM
Region 1	65 (7%)	60/M
Region 2N	217 (23%)	98/M
Region 2S	214 (13%)	96/M
Region 3	112 (60%)	99/M
Region 5	66 (5%)	69/M
Region 6	102 (52%)	70/M
Region 7	32 (-9%)	64/M
Region 8	15 (200%)	48/M

Statewide Hospitalization Trends: ICU COVID+ Census

Hospitalization Trends 3/1/2021 – 8/16/2021
Confirmed Positive in ICUs




Overall, the census of COVID+ patients in ICUs has increased by 15% from last week.

6 regions have increasing ICU COVID+ hospital census from last week. Region 1 and 6 have 10%+ of ICU beds occupied with COVID+ patients.

Region	Adult COVID+ in ICU (% Δ from last week)	Adult ICU Occupancy	% of Adult ICU beds COVID+
Region 1	19 (19%)	81%	10%
Region 2N	47 (24%)	69%	8%
Region 2S	55 (10%)	78%	8%
Region 3	28 (100%)	82%	9%
Region 5	12 (-20%)	60%	6%
Region 6	25 (4%)	75%	11%
Region 7	6 (-50%)	60%	3%
Region 8	3 (50%)	57%	4%


Average and total new deaths, by age group

Daily confirmed and probable deaths per million by age group (7 day rolling average)



Total confirmed and probable deaths by age group (past 30 days, ending 8/9/2021)

- 26% of deaths below age sixty



- Overall trends for daily average deaths are increasing since last week
- Through 8/9, the 7-day avg. death rate is below 1.0 daily deaths per million people for those under the age of 70

Note: Death information sourced from MDHHS and reflects date of death of confirmed and probable cases.

Source: MDHHS – Michigan Disease Surveillance System

National Comparison

Spread

Severity


Public Health
Response

Other
Indicators


Science
Round-up

30-day rolling average daily deaths per million people by race and ethnicity

Average daily deaths per million people by race



Average daily deaths per million people by ethnicity



- Additional reviews of vital records death data were performed the weeks of 7/6 and 8/9 to search for race and ethnicity
- This review has resulted in an adjustment of deaths for American Indian and Alaskan Natives from previous weeks
- **Currently, American Indian/Alaskan Natives have the highest death rate**

Note: Death information sourced from MDHHS and reflects date of death of confirmed and probable cases.
Source: MDHHS – Michigan Disease Surveillance System

National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up

COVID-19 Vaccination

Administration (doses administered)

- 3,994 first doses administered each day (7 day rolling average*)
- Most administered frequently by pharmacies, local health departments, and hospitals

Coverage (people vaccinated)

- 65.4% (+0.6) of aged 18+ have had first dose of vaccine; 86.5% (+0.3) of aged 65+ have had first dose
- 4,955,984 people in Michigan have completed vaccination series (4,916,256 and 4,890,859 last 2 weeks)
- Initiation highest among Asian, Native Hawaiian or Pacific Islander and American Indian/Alaskan Native individuals (MI COVID Vaccine Dashboard 8/10/21)
- Less than 1% of Vaccinated Individuals Later Tested Positive for COVID-19 (Number of cases who are fully vaccinated (n= 12,121)

*https://covid.cdc.gov/covid-data-tracker/#vaccination-trends_vacctrends-onedose-daily

National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up


Doses Administered as of 8/16/2021

12,199,070 doses delivered to providers and 9,893,319 doses administered (CDC tracker)

- 44,903 doses administered week ending 8/14 week
- 3,994 first doses/day on average

Aug 8 – Aug 14 (inclusive), doses were most frequently administered by

- Pharmacies (30.6K)
- LHD (4.2K) and hospitals (3.2K)
- Family practice (2.1K), and FQHCs (1.7K), and Pediatricians (790)



National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up

Over 4.9 Million Michiganders fully vaccinated

4.96 million people in the state are fully vaccinated

82.2% of people aged 65 and older have completed the series


Race/Ethnicity for those 12 years and older:

- Initiation coverage highest among those of Non-Hispanic (NH) Asian, Native Hawaiian or Pacific Islander Race (53.1%), then NH American Indian (49.4%), NH White (46.2%), NH Black or African American Races (35.2%).
- Initiation is at 46.3% for those of Hispanic ethnicity
- Completion follows the same pattern
- 20.7% data missing or unknown

Vaccination Coverage in Michigan as of 8/17/21

Age Group	% At Least One Dose	% Fully Vaccinated	Number Fully Vaccinated
Total Population	54.3	49.6	4,955,984
≥ 12 years	63.1	57.7	4,955,844
≥ 18 years	65.4	60.1	4,714,549
≥ 65 years	86.5	82.2	1,450,355

Coverage by Race - State Level



National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up

Potential COVID-19 Vaccination Breakthrough Cases

Michigan part of CDC's nationwide investigation ([COVID-19 Breakthrough Case Investigations and Reporting | CDC](#))

Michigan Data (1/1/21 through 8/10/21):

- **12,121 cases met criteria based on a positive test 14 or more days after being fully vaccinated**
- **Less than 1% of people who were fully vaccinated met this case definition**
 - Includes 247 deaths (217 in persons ages 65 years or older)
 - 711 cases were hospitalized
- Vaccine breakthrough cases are expected. COVID-19 vaccines are effective and are a critical tool to bring the pandemic under control. However, no vaccines are 100% effective at preventing illness in vaccinated people. There will be a small percentage of fully vaccinated people who still get sick, are hospitalized, or die from COVID-19.
- More than 166 million people in the United States have been fully vaccinated as of August 9, 2021. Like with other vaccines, vaccine breakthrough cases will occur, even though the vaccines are working as expected. Asymptomatic infections among vaccinated people will also occur.
- There is some evidence that vaccination may make illness less severe for those who are vaccinated and still get sick.
- Current data suggest that COVID-19 vaccines authorized for use in the United States offer protection against most SARS-CoV-2 variants currently circulating in the United States. However, variants will cause some vaccine breakthrough cases.

National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up

Science Round Up

What is delta and what does it mean?

- Delta variant has led to high transmission throughout the U.S. with some states experiencing highest COVID-19 cases and hospitalizations to date
- In Michigan, delta has quickly become the predominant variant and cause for the current increase in cases and hospitalizations
- Models are projecting a continued increase in hospitalizations and deaths over the next four to six weeks, maybe longer
- With return to school year, lack of layered mitigation measures will likely mean increases in cases and severe outcomes among children (e.g., hospitalizations, MIS-C, and long-COVID)

Are vaccinations working?

- A larger proportion of those who become cases (98%), are hospitalized (95%), and died (95%) from COVID are unvaccinated
- mRNA vaccine are 96% effective at preventing hospitalizations among elderly
- Among individuals previously infected, vaccination provides additional protection to prevent reinfection

What can we do about case increases?

- Layered mitigation, especially for those not vaccinated and those not yet eligible for vaccination can avoid unnecessary surge in cases and unintended school closures due to classroom outbreaks
- Treatment when exposed: FDA revised EUA for REGEN-COV for post-exposure prophylaxis is a new tool for preventing severe COVID-19 outcomes

National Comparison

Spread

Severity

Public Health
Response

Other
Indicators


Science
Round-up

**What does Delta Variant mean
for Michigan**

Cumulative COVID-19 Case Rates: States with high Delta Comparison

New cases of Covid-19, reported to CDC, in AR, FL, MI, MO, NV, and LA

Seven-day moving average of new cases (per 100K), by number of days since .01 average daily cases (per 100K) first recorded.



- Average daily incidence per 100,000 cases in Michigan is currently lower than other states experiencing a surge in delta cases

Source: [CDC COVID Data Tracker – State Trend Comparison](#)

National Comparison

Spread

Severity


Public Health
Response

Other
Indicators

Science
Round-up

What if Scenarios: Hospitalizations if we follow Wave 2 or 3




If this wave continues to grow and follows growth patterns of prior Michigan waves, we would face growing hospitalizations through September with a peak in late September/early October.



Modeling scenarios for Michigan: COVID surge on the horizon

- If vaccination slowing and increased social contact rates continue, model simulations project a surge is likely, potentially similar size to spring
- If contact rates return to low levels and/or vaccinations increase to April uptake, the surge can be reduced/stopped

Model projections are scenarios rather than forecasts—actual contact patterns may not reflect the projected scenarios. Model calibrated to MDSS case data (through 7/16, as of 7/23), using mobility data (Unacast encounter rate), increased transmission probability in June for the Delta variant. Vaccination rates based on MCIR. Uncertainty: top 10% of 1000 parameter estimates.




How do these cases translate to hospitalizations?

- Projected hospital admissions, based on the model simulations and fraction of hospital admissions for different age groups over March-May
- These projections assume the same age distribution of cases and hospitalizations as the spring surge
- Projected hospital admissions ranges are based only on the best fit and median simulations (not the full uncertainty range)

Age group	Projected total hospital admissions Aug-Nov
0-17	204 – 428
18-19	110 – 229
20-29	959 – 2007
30-39	1217 – 2547
40-49	1463 – 3061
50-59	2256 – 4722
60-69	2550 – 5336
70-79	2041 – 4271
80+	1592 – 3331
Total	12,186 – 25,505

Delta will increase transmission in Children: SARS-CoV-2 can Negatively Impact Children Directly and Indirectly

- Children can get infected with SARS-CoV-2: proportion of kids getting sick with COVID-19 is increasing



- Children can transmit the virus to others and can be sources for outbreaks

Characteristics of COVID-19 Cases and Outbreaks at Child Care Facilities — District of Columbia, July–December 2020

Weekly / May 21, 2021 /

Christine Kim, PhD^{1,*}; S. Nesbitt, MD³ ([View author profile](#))

SARS-CoV-2 Transmission and Infection Among Attendees of an Overnight Camp — Georgia, June 2020

Weekly / August 7, 2020 / 69(31);1023–1025

On July 31, 2020, this report was posted online as an MMWR Early Release.

Christine M. Szablewski, DVM^{1,2}; Karen T. Chang, PhD^{2,3}; Marie M. Brown, MPH¹; Victoria T. Chu, MD^{2,3}; Anna R. Yousaf, MD^{2,3}; Ndubuisi Anyalechi, MD¹; Peter A. Aryee,

Sources: Case data: MDSS; [Kim C, et al. Characteristics of COVID-19 Cases and Outbreaks at Child Care Facilities — District of Columbia, July–December 2020. MMWR Morb Mortal Wkly Rep 2021;70](#); [Szablewski CM, et al. SARS-CoV-2 Transmission and Infection Among Attendees of an Overnight Camp — Georgia, June 2020. MMWR Morb Mortal Wkly Rep 2020;69](#)

National Comparison

Spread

Public Health
Response

Other
Indicators

Science
Round-up

SARS-CoV-2 can Negatively Impact Children Directly and Indirectly

- **Missed in person school negatively impacts children and can occur from statewide lockdowns or large uncontrolled outbreaks within schools**
 - Remote learning disproportionately affects minorities and lower income children

Association of Children's Mode of School Instruction with Child and Parent Experiences and Well-Being During the COVID-19 Pandemic — COVID Experiences Survey, United States, October 8–November 13, 2020

Weekly / March 19, 2021 / 70(11);369–376

Jorge V. Verlenden, PhD^{1,2}; Sanjana Pampati, MPH^{1,3}; Catherine N. Rasberry, PhD^{1,2}; Nicole Liddon, PhD¹; Marci Hertz, MS^{1,2}; Greta Kilmer, MS¹; Melissa Heim Viox, MPH⁴;

- **Children can experience severe outcomes from COVID-19 including MIS-C, Hospitalization, and Death**
 - A JAMA study reported MIS-C incidence was 5.1 persons per 1,000,000 person-months and 316 persons per 1,000,000 SARS-CoV-2 infections in persons younger than 21 years
 - Incidence was higher among Black, Hispanic or Latino, and Asian or Pacific Islander persons compared with White persons and in younger persons compared with older persons

Sources: [Verlenden JV, Pampati S, Rasberry CN, et al. Association of Children's Mode of School Instruction with Child and Parent Experiences and Well-Being During the COVID-19 Pandemic — COVID Experiences Survey, United States, October 8–November 13, 2020. MMWR Morb Mortal Wkly Rep 2021;70](#); [Payne AB, et al. Incidence of Multisystem Inflammatory Syndrome in Children Among US Persons Infected With SARS-CoV-2. JAMA Netw Open. 2021;4\(6\)](#)

National Comparison

Spread


Public Health
Response

Other
Indicators


Science
Round-up

SARS-CoV-2 can Negatively Impact Children Directly and Indirectly

- Children can experience severe health outcomes from COVID-19 including MIS-C and Hospitalization
 - Hospitalizations among children nationwide is higher than it's ever been***
 - Nearly half of children hospitalized have no reported underlying conditions†**



U.S. Pediatric Hospitalizations | Underlying Medical Conditions



Sources: * [CDC COVID Data Tracker > New Hospital Admissions](#); † [COVIDNET](#)

National Comparison

Spread


Public Health
Response

Other
Indicators

Science
Round-up

Multiple states are experiencing a sharp increase in pediatric hospitalizations

- States across the southern US have seen extremely rapid and concerning surges in pediatric hospitalization rates
 - Highest pediatric COVID hospitalization rates seen so far in the pandemic for many areas
- Potential for a similar surge in Michigan as cases and hospitalizations rise





Sick kids running out of room at Omaha-area hospitals as start of school, COVID loom

THE OKLAHOMAN 

RSV + COVID: "At Oklahoma Children's Hospital, available pediatric beds are scarce as RSV surges"



Baton Rouge: Kids sick with Covid are filling up children's hospitals in areas seeing spikes

daily advertiser

Louisiana remains COVID-19 capital of U.S. as hospitals struggle to find beds and staff

Tennessean.

'All of them': Tennessee health chief says children's hospitals will fill up as the delta variant surges: "on pace to completely fill children's hospitals across the state by the end of next week."

Arkansas Democrat  Gazette

After record admissions, Arkansas ICUs down to last 8 beds



Dallas County has no pediatric ICU beds left, county judge says


Quote from county judge Clay Jenkins: "That means if your child's in a car wreck, if your child has a congenital heart defect or something and needs an ICU bed, or more likely if they have Covid and need an ICU bed, we don't have one. Your child will wait for another child to die," Jenkins said. "Your child will just not get on the ventilator, your child will be CareFlighted to Temple or Oklahoma City or wherever we can find them a bed, but they won't be getting one here unless one clears."

SARS-CoV-2 can Negatively Impact Children Directly and Indirectly

Multisystem Inflammatory Syndrome in Children (MIS-C)

- Higher community transmissions is followed by higher incidence of MIS-C cases

Daily MIS-C Cases and COVID-19 Cases Reported to CDC (7-Day Moving Average)



Source: [MDHHS and MIS-C Data and Reporting](#)

National Comparison

Spread

Public Health
Response

Other
Indicators

Science
Round-up

SARS-CoV-2 can Negatively Impact Children Directly and Indirectly

Multisystem Inflammatory Syndrome in Children (MIS-C)

- Higher community transmissions is followed by higher incidence of MIS-C cases
 - Many of those who experience MIS-C in Michigan are admitted to intensive care, school age, and are Black/African American

Multisystem Inflammatory Syndrome in Children (MIS-C) Michigan Data Summary 7/29/2021


# Cases Confirmed and Reported to CDC*	160
MIS-C associated Deaths	5 or fewer
Cases admitted to ICU	113 (70.6%)
Onset Date Range	4/14/20 to 7/2/2021
Age Range	0-20 years

*Meets CDC Case definition
<https://emergency.cdc.gov/han/2020/han00432.asp>


- 70.6% of MIS-C cases admitted to ICU

DEMOGRAPHIC INFORMATION (N=160)					
Age Group	Count	%	Race	Count	%
0-4 yrs	42	26.3%	Black/African American	70	43.7%
5-10 yrs	65	40.6%	Caucasian	66	41.3%
>10 yrs	53	33.1%	All Others / Unknown	24	15.0%
Gender	Counts	%	Ethnicity	Count	%
Male	92	57.5%	Not Hispanic or Latino	114	71.3%
Female	68	42.5%	Hispanic or Latino	12	7.5%
Unknown	0	0.0%	Unknown	34	21.2%

- 44% of MIS-C cases are black or African American Children



Source: [MDHHS and MIS-C Data and Reporting](#)



SARS-CoV-2 can Negatively Impact Children Directly and Indirectly

- While many school-aged children fully recover from COVID-19, **1 in 20 can experience symptoms last longer than four weeks and 1 in 50 can experience symptoms for more than 8 weeks**

Illness duration and symptom profile in symptomatic UK school-aged children tested for SARS-CoV-2

Erika Molteni*, Carole H Sudre*, Liane S Canas, Sunil S Bhopal, Robert C Hughes, Michela Antonelli, Benjamin Murray, Kerstin Kläser, Eric Kerfoot,

- Children experience many **Indirect Impacts** when there is uncontrolled spread of SARS-CoV-2
 - Loss of loved ones/caregivers: more than 136,000 children in the US lost a primary or secondary care giver ([orphanhood-report.pdf \(cdc.gov\)](#))
 - Adverse outcome to mental and physical health
 - Interferences with developmental milestones

Sources: [Aerosol Dynamics Model for Estimating the Risk from Short-Range Airborne Transmission and Inhalation of Expiratory Droplets of SARS-CoV-2](#); [Southern Nevada Health District](#); [LA County](#); [Retraction Notice](#).

National Comparison

Spread

Public Health
Response

Other
Indicators

Science
Round-up

Are Vaccinations Working?

COVID-19 in Michigan: Cases by Vaccination Status, January 15 – July 28

Fully Vaccinated People (4,631,476)		
Cases	Hospitalization	Deaths
Percent of Cases In People Not Fully Vaccinated (388,584 / 398,302) 97.6%	Percent of Hospitalizations In People Not Fully Vaccinated (11,048 / 11,691) 94.5%	Percent of Deaths In People Not Fully Vaccinated (4,642 / 4,888) 95.0%
388,584 Total Cases Not Fully Vaccinated	11,048 Total Hospitalized Not Fully Vaccinated	4,642 Total Deaths Not Fully Vaccinated
Total Breakthrough Cases 9,718	Total Breakthrough Hospitalizations 643	Total Breakthrough Deaths 246
0.210% Percent of Fully Vaccinated People who Developed COVID-19 (9,718 / 4,631,476)	0.014% Percent of Fully Vaccinated People Who Were Hospitalized for COVID-19 (643 / 4,631,476)	0.005% Percent of Fully Vaccinated People Who Died of COVID-19 (246 / 4,631,476)
2.4% Percent of Cases Who Were Fully Vaccinated (9,718 / 398,302)	5.5% Percent of Hospitalizations Who Were Fully Vaccinated (643 / 11,691)	5.0% Percent of Deaths Who Were Fully Vaccinated (246 / 4,888)
Total Cases: 398,302	Total Hospitalizations: 11,691	Total Deaths: 4,888

Michigan Disease Surveillance System may underestimate the frequency of COVID-19 hospitalizations:

- Case investigation and follow-up is more difficult for individuals who get vaccinated (e.g., they are too ill to speak to investigators, don't answer their phone, or otherwise).
- These hospitalizations include individuals who are hospitalized for issues other than COVID19 (the same as breakthrough COVID-19).
- Individuals who get hospitalization will lag after infection and may occur after case investigation.

National Comparison

Spread

Severity


Public Health
Response

Other
Indicators


Science
Round-up

Potential COVID-19 Vaccination Breakthrough Cases

Breakthrough Case Rate vs. Unvaccinated Rate



Breakthrough Death Rate vs. Unvaccinated Death Rate



- Trends over time show that both the case rate and death rate among the vaccinated (aka breakthrough infections and deaths) are lower than the unvaccinated rate in Michigan
- The *proportion* of breakthrough cases and deaths among all cases and deaths has shown some increases as more people become fully vaccinated
 - However, the risk of infection and death remains significantly lower among the unvaccinated
 - This principle indicates that the absolute number or the proportion of absolute number of breakthrough alone is not sufficient to measure, especially as more people become vaccinated. Instead, proportion among appropriate denominator is preferred.

National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up

All three vaccines effective at preventing hospitalization


Effectiveness of COVID-19 Vaccines in Preventing Hospitalization Among Adults Aged ≥ 65 Years — COVID-NET, 13 States, February–April 2021

COVID-NET: data on laboratory-confirmed COVID-19–associated hospitalizations in 99 U.S. counties in 13 states including Michigan.

Among adults aged 65–74 years, effectiveness of full vaccination for preventing hospitalization was 96% for Pfizer-BioNTech, 96% for Moderna, and 84% for Janssen COVID-19 vaccines

Among adults aged ≥ 75 years, effectiveness of full vaccination for preventing hospitalization was 91% for Pfizer-BioNTech, 96% for Moderna, and 85% for Janssen COVID-19 vaccines.

FIGURE 2. Estimates of vaccine effectiveness in preventing COVID-19–associated hospitalization among patients aged ≥ 65 years for the COVID-NET catchment area, by vaccine product and age group using the screening method — COVID-NET, 13 states,* February 1–April 30, 2021[†]



Abbreviations: COVID-NET = Coronavirus Disease 2019–Associated Hospitalization Surveillance Network; Janssen = Janssen (Johnson & Johnson).

* COVID-NET data included in this analysis were from the following states: California, Colorado, Connecticut, Georgia, Maryland, Michigan, Minnesota, New Mexico, New York, Ohio, Oregon, Tennessee, and Utah.

[†] Confidence intervals indicated by error bars.

Moline HL, Whitaker M, Deng L, et al. Effectiveness of COVID-19 Vaccines in Preventing Hospitalization Among Adults Aged ≥ 65 Years — COVID-NET, 13 States, February–April 2021. MMWR Morb Mortal Wkly Rep. ePub: 6 August 2021. DOI: http://dx.doi.org/10.15585/mmwr.mm7032e3external_icon.

National Comparison

Spread

Severity


Public Health
Response

Other
Indicators


Science
Round-up

Understanding breakthrough cases: When more people are vaccinated, more cases will come from the vaccinated population—even if the vaccine is working

Scenario 1: 50% Vaccinated



Scenario 2: 90% Vaccinated



● Unvaccinated
 ● Unvaccinated & Infected
 ● Prevented Infection
 ● Vaccinated
 ● Vaccinated & Infected

Both Scenarios: Vaccine reduces disease by 80%, 2.5% infection level, 2000 total people

Among previously infected individuals: **being unvaccinated is associated with significantly higher odds of reinfection compared with being fully vaccinated**

Laboratory evidence suggests that antibody responses after COVID-19 vaccination provide better neutralization than natural infection, however few epidemiologic studies under real world conditions have been completed

Recent report from a case control study in Kentucky: Among those infected with SARS-CoV-2 in 2020, **being unvaccinated was associated with 2.34 times the odds of reinfection compared with being fully vaccinated**

Among individuals previously infected, full vaccination provides additional protection against reinfection

How many people in Michigan have been either vaccinated or previously infected?

Michiganders



Vaccinated	
Vaccinated and previously infected	Previous infection

- 28% of Michiganders have been previously infected with COVID based on seroprevalence (nucleocapsid - measures previous infection only)
- 54% of Michiganders (all ages) have received at least one dose; 49% fully vaccinated
- **If** people are equally likely to get vaccinated whether they have been previously infected or not, then:
 - ~67% would either received at least one dose or previously infected (estimate)
 - ~63% would either fully vaccinated or previously infected (estimate)
- Uncertainty around protection from previous infection

Is this enough to prevent another surge? Puerto Rico provides comparison point—

Puerto Rico: 74.8% of the population has either been previously infected or vaccinated—but they are currently at high transmission levels

- **75% of Puerto Ricans have either been vaccinated or previously infected** based on seroprevalence (spike protein – measures vaccination or previous infection)
- 70% of Puerto Ricans (all ages) have received at least one dose; **61% fully vaccinated**
- However, **Puerto Rico is seeing a rapid surge** similar to spring peak, reaching CDC **high transmission** levels



What Can We Do About Case Increases?


Layers of Defense Against COVID-19 in Schools

CDC recommended prevention strategies can be layered in different ways – the number and intensity of the layers can increase if community transmission increases

As community transmission increases, more holes appear in the defenses, meaning more layers of protection may be needed.



As the vaccination rate within a building or facility increases, fewer holes will appear in the defenses.



Low SARS-CoV-2 Transmission in Elementary Schools — Salt Lake County, Utah, December 3, 2020–January 31, 2021

Weekly / March 26, 2021 / 70(12);442–448

- Layered strategy: high adherence to masking + classroom cohorting and other measures—but classroom seats were a median of 3 ft apart
- “In a high community transmission setting, low school-associated transmission was observed with a 0.7% secondary attack rate.”

Pilot Investigation of SARS-CoV-2 Secondary Transmission in Kindergarten Through Grade 12 Schools Implementing Mitigation Strategies — St. Louis County and City of Springfield, Missouri, December 2020

Weekly / March 26, 2021 / 70(12);449–455

Layered prevention strategies including masking

Secondary transmission in only 2 of 102 close contacts tested


Clusters of SARS-CoV-2 Infection Among Elementary School Educators and Students in One School District — Georgia, December 2020–January 2021

Weekly / February 26, 2021 / 70(8);289–292

Five of the nine transmission clusters involved inadequate mask use by students

Modeling: Masks can substantially reduce transmission in school settings, even with delta variant

If 1 infectious child attends a class of 25 students, how long does it take for there to be a >50% chance of transmission occurring?




Estimates from the [COVID-19 Indoor Safety Guideline](#), based on [Bazant and Bush, A guideline to limit indoor airborne transmission of COVID-19, PNAS 2021](#). Simulations assume: delta strain, normal talking (not singing/etc.), with child age group for elementary and average between adult and child age groups for high school. Vaccine coverage was assumed to be 0% for elementary and 33% for high school, based on age-specific coverage rates as of 8/6/21. We assumed 95% mask fit/compliance for 'Complete Masking' and 75% for 'Imperfect Masking'.

Modeling study in NC: Masking and testing in K-12 schools can prevent 40-70% of new infections

New Infections among 500 Students after 1 semester

- Elementary School Setting (incoming protection = 30%)
- Middle School Setting (incoming protection = 40%)
- High School Setting (incoming protection = 50%)




Copyright - COVSIM Research Group

- Children under 12 are not yet eligible for vaccines and so are at higher risk
- **Without masks or testing up to 90% of susceptible students may become infected by the end of the semester** (if only 30% have protection due to previous infection or vaccination)
- **Masking reduces projected infections by 40-75% in for elementary, middle, and high schools**

Mask mandates implemented after delta surge saw slower rates of increases in jurisdictions compared to those who didn't require masking

- Delta variant contributed to surges in several U.S. states in July
- Several states prohibited local masks mandates
- To date, there have been no national universal mask mandates regardless of vaccinations, only recommendations
- Several local jurisdictions passed mask mandates for indoor settings, regardless of vaccination status, in response to spread of the delta variant
 - Three were: LA County (LAC), St. Louis (STL), and Southern Nevada Health District (NV) including Clark County (Las Vegas)
 - In contrast, Orleans Parish (Louisiana), Pulaski County (Little Rock, Arkansas), and Duval County (Jacksonville, Florida) had no such mandates
- **While other factors could have also contributed to lower case rates, jurisdictions with mask mandates have experienced lower spread of COVID-19 during the delta surge**



Sources: [St. Louis City COVID-19 dashboard](#); [LA County COVID-19 dashboard](#); CDC COVID Data Tracker; population sizes from [CDC Wonder Bridged Race Estimates](#)

National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up

MONOCLONAL ANTIBODY POST EXPOSURE PROPHYLAXIS: FDA revision of Emergency Use Authorization of REGEN-COV

Post-exposure prophylaxis using REGEN-COV in individuals 12 years and older who are


- **Are at high risk for progression to severe COVID-19**, and
- **Are not fully vaccinated** or are not expected to mount an adequate immune response, and
- **Have been exposed** to an individual infected with SARS-CoV-2 as close contact or because of COVID-19 infection in other individuals in same institutional setting (for example, nursing homes or prisons)

Use is in addition to prior authorization to treat non-hospitalized patients with mild to moderate COVID-19 in adult and pediatric patients, with positive SARS-CoV-2 viral testing, and who are at high risk for progression to severe COVID 19.

REGEN-COV is expected to be effective against circulating variants.

Post-exposure prophylaxis is not a substitute for vaccination against COVID-19.

Additional information on monoclonal antibody therapy is available at www.michigan.gov/covidtherapy.



National Comparison

Spread

Severity

Public Health
Response

Other
Indicators

Science
Round-up

CDC Guidance for Implementing COVID-19 Prevention Strategies in the Context of Varying Community Transmission Levels and Vaccination Coverage

Public health system needs to assess use of prevention strategies to avoid stressing health care capacity to provide adequate COVID-19 and non-COVID-19 care

CDC recommends five critical factors be considered to inform local decision making:

- 1) Level of SARS-CoV-2 community transmission
- 2) Health system capacity
- 3) COVID-19 vaccination coverage
- 4) Capacity for early detection of increases in COVID-19 cases
- 5) Populations at increased risk for severe outcomes from COVID-19

Proven effective strategies against transmission, beyond vaccination:

- 1) Using masks consistently and correctly
- 2) Maximizing ventilation
- 3) Maintaining physical distance and avoiding crowds
- 4) Staying home when sick
- 5) Handwashing
- 6) Regular cleaning of high-touch surfaces

Prevention strategies should be strengthened or added if transmission worsens.

Prevention strategies should only be relaxed after several weeks of continuous improvement in level of community transmission



Sources: [Guidance for Implementing COVID-19 Prevention Strategies in the Context of Varying Community Transmission Levels and Vaccination Coverage](#)

National Comparison

Spread

Severity

Public Health
Response



Other
Indicators

Science
Round-up

Appendix

Nearly all States and Territories are at High CDC Transmission Level

- 2 jurisdictions have substantial transmission (orange states); down 12 from 7 days ago
- 52 jurisdictions have high transmission (red states); up 12 from 7 days ago
- CDC recommends masking when indoor public spaces; regardless of vaccination status



National Comparison

Spread

Severity

Public Health
Response


Other
Indicators

Science
Round-up

Cumulative COVID-19 Case Rates: Midwest Comparison

Cumulative cases of Covid-19, reported to CDC, in IL, IN, MI, MN, OH, and WI

Cumulative cases (per 100K), by number of days since .3 total cases (per 100K) first recorded.



- Cumulative incidence per 100,000 cases in Michigan has been lower than other states in the Midwest following spring 2020 surge
- Michigan's mitigation policies helped control the spread of SARS-CoV-2 relative to other states in the Midwest, particular during surge in November and December
- The current trajectory in Michigan continues to be in the range of cumulative case rates of our Midwest neighbors

National Comparison

Spread

Severity


Public Health
Response

Other
Indicators


Science
Round-up

Most Michigan Counties at Substantial or High Transmission Levels (CDC)

This Week, 8/15



Last week, 8/8



Transmission Levels

Low

Moderate

Substantial

High

of counties

This week

Last week

1

4

10

20

34

42

38


17

Updates since last week:

- 1 of 83 counties met low transmission level this week, a 3 county decrease from last week
- 10 of 83 counties met moderate transmission classification, a 10 county decrease
- 34 of 83 counties met substantial transmission classification, an 8 county decrease
- 38 of 83 counties met high transmission classification, a 21 county increase from last week


Comparing CDC community transmission thresholds to MI levels

Case Rate*†




† CDC uses cases / 100,000 / week (conversion to MI metrics in paratheses)

Percent Positivity



Cases per Capita Increasing Within All MERC Regions




- Case rate trends for all regions are increasing
- Kalamazoo, Jackson, and Saginaw currently have the highest case rates

Identified COVID-19 Delta Variants by County

This week (Aug 16, 2021)*

Delta (B.1.617.2) Variant by County




B.1.617.2 Delta Variant Not Identified

Confirmed Delta (B.1.617.2) Variant Reported

Note: 18 cases in Wayne County attributed to Detroit City

Last week (Aug 9, 2021)

Delta (B.1.617.2) Variant by County
Aug 9



B.1.617.2 Delta Variant Not Identified

Confirmed Delta (B.1.617.2) Variant Reported

Note: 10 cases in Wayne County attributed to Detroit City

Note: The low number of specimens recently submitted for sequencing limits the ability to estimate the prevalence of variants in Michigan

* Delta found in wastewater surveillance samples

Data last updated Aug 16, 2021
Source: MDSS

National Comparison

Spread

Severity


Public Health Response

Other Indicators

Science Round-up



Comparing COVID hospitalization growth rates across states

Michigan Wave 2 (Fall/Winter) and Wave 3 (Spring) vs. Other states current summer 2021 waves



CDC model projections for Michigan: increasing hospitalizations and deaths


- *Ensemble model suggests increasing trends for hospitalizations and deaths*
- Uncertainty ranges from flat to increasing
- Case data projections also range from flat to increasing (not shown) and recent data has shown increases
- Individual models shown as grey lines, ensemble shown in red




Data Sources: [CDC mathematical model forecasting](#), [CovidComplete Data Center](#) model forecast evaluations

Understanding breakthrough cases: as more people are infected, there will be more cases among both vaccinated and unvaccinated people

Scenario 2: 2.5% of unvaccinated people infected



Scenario 3: 5% of unvaccinated people infected



● Unvaccinated ● Unvaccinated & Infected ● Prevented Infection ● Vaccinated ● Vaccinated & Infected

Both Scenarios: 90% Vaccinated, Vaccine reduces disease by 80%, 2000 total people



QUESTIONS?

[Michigan.gov/Coronavirus](https://michigan.gov/Coronavirus)