

the
YellowDog
Peridotite
& a Possible Buried
Igneous Complex of
Lower Keweenaw Age in the
Northern Peninsula of Michigan

Geological Survey
Michigan DNR
Report of
Investigation 24

Report of Investigation 24

**THE YELLOW DOG PERIDOTITE AND A
POSSIBLE BURIED IGNEOUS COMPLEX OF
LOWER KEWEENAWAN AGE IN THE NORTHERN
PENINSULA OF MICHIGAN**

by

John S. Klasner, David W. Snider, W. F. Cannon, and
John F. Slack

Lansing, Michigan
1979

PREFACE

This report is the result of a joint effort by the United States Geological Survey and the Geological Survey Division, Michigan Department of Natural Resources. Specific units involved in the project were the Eastern Mineral Resources Branch of the U.S. Geological Survey and the Geology and Minerals Research Unit, Michigan Geological Survey.

Both agencies became interested in a somewhat anomalous peridotite body outcropping in an otherwise flat sand plain in northern Marquette County, Michigan. Discussions between the two agencies culminated in an agreement to combine efforts in a thorough investigation of this curious rock body.

The report presents the multidisciplinary approach of geology, geochemistry and geophysics to the problem. The use of such an approach has proven to produce more scientifically complete results than would have been possible with a single discipline. In addition, the study has been extremely cost-efficient.

David W. Snider, Geologist
Geology & Minerals Research Unit
Geological Survey Division
Michigan Department of Natural Resources
Lansing, Michigan 48909

CONTENTS

PREFACE	2
ILLUSTRATIONS	2
<i>Figures</i>	2
<i>Tables</i>	2
ABSTRACT	2
INTRODUCTION	3
<i>Acknowledgments</i>	3
GENERAL GEOLOGY	4
PETROLOGY	4
<i>Texture, Mineralogy and Bulk Composition</i>	4
<i>Composition of Major Primary Silicate Minerals</i>	5

<i>Opaque Mineralogy</i>	6
<i>Trace Element Content</i>	8
PALEOMAGNETIC STUDIES	9
FIELD GEOPHYSICAL STUDIES	10
<i>Field Procedures and Data Reduction</i>	10
Gravity	10
Ground Magnetics	10
VLF-EM.....	10
<i>Interpretation of Geophysical Data</i>	11
Interpretation of Individual VLF-EM Anomalies	13
Regional Survey	13
GEOCHEMICAL SURVEY	16
SUMMARY AND CONCLUSIONS	16
REFERENCES	18

ILLUSTRATIONS

Figures

Figure 1. <i>Yellow Dog Plains general regional geology and location.</i>	3
Figure 2. <i>Location of analyzed samples. See Figure 3 for locations</i>	6
Figure 3. <i>Complete Bouguer gravity anomaly.</i>	11
Figure 4. <i>Ground magnetics of detailed study area.</i>	11
Figure 5. <i>Filtered VLF-EM anomalies</i>	14
Figure 6. <i>Dikes inferred from ground magnets.</i>	14
Figure 7. <i>Geology of the detailed study area inferred from geophysical surveys.</i>	15
Figure 8. <i>Gravity profile along A – A', showing how residual gravity anomalies were determined.</i>	15
Figure 9. <i>Gravity residual – 1, anomaly caused by deep body.</i>	15
Figure 10. <i>Gravity residual – 2, anomaly caused by near-surface dikes.</i>	15
Figure 11a. <i>Composite geophysical profiles and geologic model for cross section A – A'.</i>	17
Figure 11b. <i>Composite geophysical profiles and geologic model for cross section B – B'.</i>	17
Figure 12. <i>Composite geophysical profiles along regional lines 1 and 3.</i>	18
Figure 13. <i>Location of geochemical soil profiles.</i>	18
Figure 14. <i>Geochemical profiles in soil along profile A.</i>	18
Figure 15. <i>Geochemical profiles in soil along profile B.</i>	18

Tables

Table 1. <i>Chemical analyses of the Yellow Dog Peridotite expressed in weight percent.</i>	5
Table 2. <i>Electron microprobe analyses of olivines from the Yellow Dog Peridotite expressed in weight percent.</i>	6
Table 3. <i>Electron microprobe analyses of pyroxenes from the Yellow Dog Peridotite expressed in weight percent.</i>	8
Table 4. <i>Trace element content, of the Yellow Dog Peridotite expressed in parts per million (ppm).</i>	9
Table 5. <i>Comparison of selected trace element contents for average ultramafic rock and Yellow Dog Peridotite expressed in parts per million (ppm).</i>	9
Table 6. <i>Summary of data and interpretations of anomalies. See Figure 5 for locations.</i>	16

ABSTRACT

Partly serpentinized peridotite of early Keweenawan age crops out in two places along a 20-kilometer-long zone of

positive aeromagnetic anomalies in northern Marquette County, Michigan. Most of the area is mantled by Pleistocene drift with few bedrock exposures.

Petrographic and electron microprobe studies show that the peridotite was originally a plagioclase lherzolite containing 40 to 50 percent olivine (Fo_{80}) and approximately 10 to 15 percent each of enstatite ($\text{En}_{78}\text{Wo}_{04}\text{Fs}_{18}$) and diopsidic augite ($\text{En}_{47}\text{Wo}_{42}\text{Fs}_{11}$). The plagioclase varies from 5 to 10 percent, and according to Morris (1977) is labradorite, with a composition of An_{57-65} . Major oxide minerals (4 to 6 percent) are ilmenite and magnetite. Sulfides comprise 1 to 2 percent of the peridotite and are chiefly pyrrhotite, pentlandite, and chalcopyrite.




Figure 1. Yellow Dog Plains general regional geology and location.

Ground magnetic, gravity, and very low frequency electromagnetic (VLF-EM) surveys have refined the location and magnitude of anomalies previously known only from aeromagnetic studies. These surveys together with soil geochemical studies suggest that peridotite, and possibly other mafic rocks forming a differentiated igneous complex, may occur throughout a belt 20 kilometers long (east-west) and 1 to 2 kilometers wide (north-south).

Differentiated igneous complexes in many parts of the world are hosts for copper, nickel, chromium, or precious-metal deposits. The peridotite in the area of this study is anomalously rich in copper and sulfur compared to world-wide averages for peridotite. Positive electro-magnetic anomalies found near the peridotite outcrops may be caused by sulfide-rich zones in the igneous rocks and should be explored further for copper-nickel mineralization.

INTRODUCTION

The Yellow Dog Plains in northern Marquette County, Michigan, is a nearly featureless area covered by Pleistocene sand and gravel and is nearly devoid of bedrock exposures. The bedrock beneath the plains is mostly slate and other metasedimentary rock of the Michigamme Formation and has been shown as such on virtually all geologic compilations of the region. However, in Sections 11 and 12, T50N, R29W, there are two outcrops of massive, partly serpentinized peridotite. The outcrops lie on and form the highest magnetic peaks of a west-trending linear aeromagnetic anomaly which is about 20 kilometers long (see Figure 1). The anomaly lies parallel to another anomaly of comparable magnitude about 0.5 to 1 kilometer to the south. Both anomalies appear to cut the regional structural trends at low angles.

The two anomalies were first evaluated by Case and Gair (1965) and were inferred by them to be caused by magnetic sedimentary units in the Michigamme Formation. At that time, the existence of the peridotite outcrops was not generally known. Because of the close association of the peridotite outcrops with the observed aeromagnetic anomalies and because the anomalies appear to cut across structural trends in the Michigamme Formation, we believe that both anomalies are associated with peridotite.

The relatively fresh, undeformed nature of the peridotite suggests that it is younger than the Penokean orogeny which deformed and metamorphosed other rocks in the area about 1.9 billion years ago. Because komatiitic magmas are not common in rocks younger than about 2.5 billion years and because the texture and chemical composition of the peridotite suggest that it is a crystal cumulate, we estimate it is more likely that the peridotite is a differentiate from a less mafic magma rather than a product of a parental ultramafic magma.

- 1/ U. S. Geological Survey, Reston, Virginia
- 2/ Geological Survey Division, Michigan Department of Natural Resources, Lansing, Michigan

The present study was undertaken to determine if a relatively large differentiated igneous complex is beneath the Yellow Dog Plains and, if so, to determine its configuration and potential economic mineralization. An integrated program including ground magnetic, very low frequency electromagnetic (VLF-EM), gravity, paleomagnetic, rock chemistry and mineralogy, and soil chemistry studies was carried out in 1976 as a cooperative project between the U. S. Geological Survey, Eastern Mineral Resources Branch and the Geology and Minerals Research Unit of the Geological Survey Division, Michigan Department of Natural Resources. This report presents the results of that study and our inferences of the geology and mineral resource potential of the area of the Yellow Dog Peridotite and associated anomalies.

Acknowledgments.

R. Lilienthal, P. Geraci, and A. Grosz performed much of the geophysical field work, data reduction, and plotting. S. Guam assisted in interpreting and modeling the gravity

data. Kenneth Books performed the magnetic susceptibility and remanence studies. L. B. Wiggins assisted in the microprobe analyses. M. P. Foose reviewed the manuscript. Michigan Geological Survey Division personnel were instrumental in bringing the report to final form for publication. Final editing was accomplished by Beverly L. Champion; layout, graphics and word processing were coordinated by Steven E. Wilson; drafting by Darrell Hodge and Gregory A. Wilson; and manuscript typing by Lois J. DeClaire.

GENERAL GEOLOGY

The region surrounding the Yellow Dog Peridotite contains Precambrian rocks of three ages. A gneiss-greenstone complex older than 2.5 billion years (Precambrian W) forms the basement for roughly 2-billion-year-old (Precambrian X) rocks of the Marquette Range Supergroup. The Marquette Range Supergroup is preserved in grabens of Precambrian W rocks formed during the Penokean orogeny about 1.9 billion years ago (Cannon, 1973). Figure 1 shows the aerial distribution of Precambrian W, X and Y rocks near Yellow Dog Plains. The plains, formed of Pleistocene sand and gravel, are underlain mostly by the Michigamme Formation, part of the Marquette Range Supergroup, in a structural trough known informally as the Baraga Basin. They are flanked by Precambrian W gneiss and greenstone on the north, east, and south. Because of the extensive Pleistocene cover in the basin, little is known of the nature of the Michigamme Formation. Most exposures are along the southern flank of the basin where basal conglomerate, quartzite, and arkose as much as 20 meters thick underlie black slate and argillite. The Michigamme beneath the study area consists of fine-grained clastic rocks, largely black slate and argillite as shown by recent drilling (Trow, 1978 a,b,c). These rocks were deformed and metamorphosed to the greenschist facies about 1.9 billion years ago, during the Penokean orogeny.

The third age of Precambrian rocks is represented by west-trending diabase dikes of early Keweenawan age. The dikes were intruded and cooled during a period of reversed magnetic polarity about 1.1 billion years ago (Precambrian Y). Their negative remanent magnetization produces pronounced linear magnetic lows over the dikes and allows them to be identified and traced with certainty on aeromagnetic maps.

The Yellow Dog Peridotite is known from two exposures (see Figure 1). Because it is undeformed and only moderately serpentinized, it was judged from field evidence to be younger than the Penokean orogeny that deformed and metamorphosed the surrounding Precambrian W and X rocks into which the peridotite was emplaced. Because the peridotite causes prominent positive aeromagnetic anomalies, it was initially believed that it was not related to lower Keweenawan diabase dikes that typically produce magnetic lows. Subsequent paleomagnetic studies by Kenneth Books of the U.S. Geological Survey, however, confirm that when the soft components of remanent magnetization are removed the

peridotite does have a reversed remanent pole position very similar to that of other lower Keweenawan rocks in the region.

PETROLOGY

Texture, Mineralogy and Bulk Composition

The peridotite is reddish brown on weathered surfaces and greenish black on fresh surfaces. The weathered rind is generally confined to a few millimeters of the surface. The rock is massive and coarse grained, without visible layering, banding, or foliation. Poikilitic pyroxene is commonly a centimeter or more in length; olivine generally is 2 to 5 millimeters in diameter.

Microscopic examination indicates that the rock was originally a plagioclase lherzolite containing 40 to 50 percent olivine (now from 1/2 to 2/3 replaced by serpentine) and 20 to 30 percent pyroxene, with clinopyroxene generally slightly more abundant than orthopyroxene. Plagioclase comprises about 5 to 10 percent of the rock, and chlorite is present in amounts ranging from 5 to 15 percent. Opaque minerals, discussed in more detail below, make up 5 to 8 percent. A variety of other minerals including both brown and green amphibole, red-brown biotite, sericite, talc, carbonate, clinozoisite, and rare colorless spinel (?) together comprise 5 to 10 percent of the rock.

Olivine occurs as euhedral phenocrysts, now partly altered to serpentine. Adjacent euhedral phenocrysts commonly are in contact with each other and appear to be accumulations of crystals that have settled from the magma rather than crystals that have grown in place, since there is no indication that the shape of the olivine grains is controlled by contact with adjacent grains.

Olivine phenocrysts typically are surrounded by subhedral to anhedral pyroxene varying from about the same size as olivines to large grains poikilitically enclosing olivine phenocrysts. Some orthopyroxenes are euhedral, however, indicating crystal growth contemporaneous with olivine. Plagioclase generally occurs as subhedral and anhedral grains interstitial to pyroxene crystals. In a few places, it forms euhedral laths entirely within orthopyroxene, suggesting crystallization of at least some plagioclase prior to both pyroxene types. Green amphibole (actinolite) occurs as an alteration product of clinopyroxene and is clearly secondary. Brown hornblende forms rims on pyroxene grains and locally exhibits intercumulus textures suggesting a primary origin. Biotite occurs chiefly as subhedral laths which in places poikilitically enclose euhedral olivine. Most appears to have formed as an alteration of pyroxene, although the genesis of some euhedral intercumulus grains is not clear.

No cataclastic textures or other metamorphic fabrics are present on a microscopic scale. Minor deformation is displayed by shattered plagioclase grains and bent and kinked biotite flakes. The deformation may be due to volume expansion during the serpentinization of olivine

(i.e., Hostettler et al, 1966; Coleman and Keith, 1971). Commonly, the shattering of plagioclase is most intense where crystal terminations of serpentinized olivine grains impinge on plagioclase grains.

	no. 6	no. 35
SiO ₂	45.0	40.2
TiO ₂	0.83	0.92
Al ₂ O ₃	4.4	6.0
Fe ₂ O ₃	14.3	6.4
FeO (total Fe)		7.7
MnO	0.18	0.18
MgO	24.9	24.9
CaO	4.4	3.1
Na ₂ O	0.36	0.40
K ₂ O	0.11	0.20
P ₂ O ₅	0.10	0.046
F	na	0.03
LOI ¹	5.56	7.8
Moisture (at 105°C)	na	0.01
CO ₂	na	0.1
Total	100.15	97.99
na - not analyzed		
¹ LOI - loss on ignition at 1,000°C		
no.6 from the section 12 outcrop and was analyzed by the Rocky Mountain Geochemical Corporation		
no.35 from section 11 outcrop and was analyzed by Skyline Labs, Inc.		

Table 1. Chemical analyses of the Yellow Dog Peridotite expressed in weight percent.

The origin of the peridotite is important in assessing the potential for contained mineral deposits. Economic copper, nickel, and precious-metal ores are characteristically associated with layered mafic or ultramafic intrusions (i.e., Sudbury, Duluth), rather than with alpine-type intrusions. Utilizing criteria outlined by Thayer (1960), the Yellow Dog Peridotite displays textural and mineralogical features strongly suggestive of a stratiform mafic-ultramafic genesis. If it is a product of a parental ultramafic magma (a komatiite), it might contain nickel sulfide deposits. If, on the other hand, it is a differentiate from a less mafic magma, disseminated copper-nickel sulfides, chromite, or precious-metal ores could be present. The chemical composition of the peridotite (Table 1) suggests that the rock is not a komatiite and that the present composition more likely is a result of differentiation from basaltic magma. Specifically, the high TiO₂/MgO ratio, and the low Al₂O₃ content for a corresponding FeO/FeO+MgO ratio shown by the peridotite are more characteristic of a tholeiite than of a komatiite, according to data summarized by Naldrett and Cabri (1976) and Arndt et al. (1977). This, together with

the presence of cumulus olivine textures suggests that the bulk composition of the magma from which the peridotite crystallized was considerably less mafic than the present rock.

Composition of Major Primary Silicate Minerals

Chemical analyses of olivines and pyroxenes were obtained with an ARL-EMX-SM electron microprobe operated at an accelerating potential of 15 KV and a beam current of 0.05 mA. Crystals used were ADP, LiF, and RAP, with 20-second counting intervals on 2 to 3 micron-sized spots. Data reduction was performed by the method of Bence and Albee (1968). Two samples from the Section 12 outcrop (Figure 2) were studied. Compositions in Tables 2 and 3 represent means and standard deviations for 3 or more spots on several different grains for each slide.

Olivines (Table 2) are chrysolites, Fo₈₀₋₈₁, and of similar composition from both samples. No significant differences in major or minor oxide content between samples is apparent. Orthopyroxenes (Table 3) are enstatites, En₇₇₋₇₉, Wo₀₄ Fs₁₇₋₁₉, with only minor variations in composition between the two sample areas. The western sample (Sample 7) contains orthopyroxenes slightly higher in FeO and TiO₂ and lower in MgO than Sample 25. Clinopyroxenes are diopsidic augite, En_{41,53} Wo_{36,48} Fs₁₁, with identical FeO contents (6.5 weight percent) for both samples. Significant compositional differences occur for TiO₂, Al₂O₃, MgO and CaO (Table 3); Cr₂O₃ and Na₂O display minor variations between samples.

Compositions of individual silicate minerals may be of partial use in determining parental magma type. The Yellow Dog Peridotite contains olivines with 80 to 81 mole-percent forsterite, a common composition for olivines found within many types of ultramafic rocks. Peridotitic komatiites typically contain more magnesian olivines, at Fo₈₉₋₉₄ (Nesbitt, 1971; Pyke et al., 1973; Arndt et al., 1977). These compositions are not limited to komatiitic rocks, however, as cumulate parts of several layered mafic-ultramafic intrusions of tholeiitic origin (e.g., Stillwater) locally contain high-magnesian (>Fo₉₀) olivines. Other elements do not appear to be diagnostic indicators. Olivine compositions are thus of little use in distinguishing komatiitic from tholeiitic magma suites.

Chemical data for komatiitic pyroxenes are scarce, but one genetic discriminating parameter may be the alumina content of clinopyroxenes. Microprobe studies of ultramafic lavas from Ontario (Pyke et al., 1973; Arndt et al., 1977) and western Australia (Nesbitt, 1971) indicate unusually high contents of Al₂O₃ in clinopyroxene, generally from 5 to over 8 weight percent. Such alumina contents normally imply high formational pressure (Green, 1967), a condition clearly precluded by the subaerial environment of these lavas. High alumina contents of komatiitic clinopyroxenes are probably the result of disequilibrium crystallization during rapid quenching of ultramafic liquids under surface conditions. Nevertheless, most other ultramafic rocks formed at low pressures have

low Al₂O₃ in clinopyroxene (O'Hara, 1967), providing an empirical method to distinguish tholeiitic from komatiitic magmas crystallized under volcanic and subvolcanic conditions. Clinopyroxenes from the Yellow Dog Peridotite contain less than 3 weight percent alumina, and are compatible with bulk chemical data suggesting a tholeiitic rather than komatiitic affinity.




Figure 2. Location of analyzed samples. See Figure 3 for locations

Differences in chemistry of clinopyroxenes for the two sample areas suggest somewhat different petrologic histories for the dikes. Modal proportions of olivine, orthopyroxene, and clinopyroxene are similar for both samples, and so variations in clinopyroxene composition reflect differences in crystal-melt equilibria during crystallization and/or slight differences in original bulk chemistry of magma between the two sites. The occurrence of primary calcic plagioclase (An₅₇₋₆₅) coexisting with spinel (and hornblende?) suggests low pressure conditions along the univariant breakdown curve between the orthopyroxene - plagioclase - lherzolite facies and the spinel - lherzolite facies (O'Hara, 1967; Frost, 1976). This assemblage is stable within a much higher temperature-pressure range than local greenschist-facies rocks, thus reflecting original conditions of magma crystallization. Further work employing pyroxene geothermometers and geobarometers might define more closely the petrogenesis of the peridotite.

Nickel contents of the olivines and pyroxenes are contrary to predicted values based on experimentally determined distribution coefficients. In theory, olivines should contain much more nickel than coexisting pyroxenes.

Recent detailed microprobe studies of similar Yellow Dog samples by W. J. Morris (1977) suggest that the olivines contain about 0.4 weight percent nickel, or roughly three times that of coexisting ortho- and clinopyroxenes. Since

the values obtained in this investigation do not parallel the findings of W. J. Morris, we suspect an instrumental error in the anomalously low nickel values reported for our Yellow Dog olivines (Table 2).

Olivines		
Oxide	YDP-7	YDP-25
SiO ₂	38.88 ± .76	39.90 ± .41
TiO ₂	0.04 ± .04	0.01 ± .00
Al ₂ O ₃	0.05 ± .04	0.04 ± .02
FeO*	18.44 ± .93	17.81 ± .83
Cr ₂ O ₃	0.05 ± .04	0.03 ± .03
MnO	0.21 ± .08	0.20 ± .04
NiO	0.01 ± .02	0.00 ± .00
MgO	40.74 ± .79	42.21 ± .58
CaO	0.22 ± .09	0.26 ± .05
Na ₂ O	0.01 ± .01	0.02 ± .02
Total	98.65	100.48
Number of ions on the basis of 4 oxygens		
Si	1.010	1.010
Al	0.001	0.000
Ti	0.000	0.000
ΣFe	0.400	0.380
Cr	0.000	0.000
Mg	1.570	1.590
Ni	0.000	0.000
Mn	0.004	0.004
Ca	0.005	0.006
Na	0.000	0.000
Total	2.990	2.990
*Total iron as FeO		

Table 2. Electron microprobe analyses of olivines from the Yellow Dog Peridotite expressed in weight percent.

Opaque Mineralogy

Oxide and sulfide minerals typically make up 5 to 8 volume percent of the peridotite. Oxides are most abundant, and include ilmenite, magnetite, and minor to trace amounts of chromite and rutile. Ilmenite is the most common opaque mineral, occurring as rounded to irregular skeletal grains up to 2 millimeters in size. Most are interstitial to olivine and pyroxene grains, but some are poikilitically enclosed by large pyroxene crystals. Ilmenite grains are generally fractured and show textures suggesting partial resorption. Characteristically, they

occur isolated without associated oxides or sulfides. The majority of ilmenite grains are optically homogeneous, but some display fine, 1- to 3-micron-wide, oriented [0001] lamellae of titanomagnetite and chromite (?). One presumably secondary inclusion of hematite was found along the margins of one ilmenite grain.

Magnetite occurs as both primary and secondary grains of variable size and shape. Primary magnetite is found as interstitial grains between silicates and as angular to rounded inclusions up to 0.5 millimeters in orthopyroxene. Secondary magnetite, formed by serpentinization of olivine and pyroxene, occurs as thin (5 to 10 microns) stringers between relict olivine crystals, and as isolated grains or clusters of grains surrounding or within altered pyroxenes. Some magnetite grains are complexly intergrown with sulfide minerals.

Chromite is a minor oxide phase found as polygonal to subrounded grains 0.1 millimeter in diameter, generally intergrown with relict olivine. Trace amounts of rutile occur as irregular isolated grains 0.1 millimeter in length; one crystal displays color zoning in transmitted light with yellowish, orange, and dark red-brown bands.

Sulfide minerals typically comprise 1 to 2 percent of the peridotite. Major sulfides are pyrrhotite, pentlandite, and chalcopyrite, with minor pyrite and cubanite and trace amounts of mackinawite. Traces of secondary marcasite, bornite and covellite are present in a few sections. Sulfide minerals occur as composite grains up to 3 millimeters in diameter; the majority are less than 0.1 millimeter in size. Most are irregularly shaped, fractured, and interstitial to silicates; some form 0.1 millimeter rounded inclusions or globules within pyroxene crystals. Sulfides show preferential association with magnetite, chiefly as chalcopyrite containing laths of magnetite. Less commonly, chalcopyrite occurs as myrmekitic intergrowths within certain magnetite grains. Cubanite and pentlandite generally occur with pyrrhotite or chalcopyrite. Complex intergrowths are common, mainly fine lamellae of chalcopyrite in cubanite and leaf-like blades of cubanite in pyrrhotite. Mackinawite occurs as small inclusions in cubanite.

Orthopyroxenes		
Oxide	YDP-7	YDP-25
SiO ₂	54.26 ± .59	55.24 ± .63
TiO ₂	.57 ± .05	0.40 ± .10
Al ₂ O ₃	1.55 ± .25	1.84 ± .44
FeO*	12.17 ± .22	11.33 ± .48
Cr ₂ O ₃	0.32 ± .07	0.37 ± .08
MnO	0.25 ± .05	0.20 ± .04
NiO	0.11 ± .14	0.02 ± .03
MgO	27.84 ± .40	29.02 ± .31
CaO	1.99 ± .14	1.90 ± .17

Na ₂ O	0.05 ± .04	0.04 ± .02
Total	99.11	100.36
Number of ions on the basis of 6 oxygens		
Si	1.953	1.952
Al	0.065	0.076
Ti	0.015	0.010
ΣFe	0.366	0.334
Cr	0.008	0.010
Mg	1.493	1.528
Ni	0.003	0.000
Mn	0.007	0.005
Ca	0.076	0.071
Na	0.003	0.002
Total	3.989	3.988
Clinopyroxenes		
SiO ₂	51.90 ± .55	51.71 ± .34
TiO ₂	0.45 ± .12	1.45 ± .10
Al ₂ O ₃	1.85 ± .37	2.71 ± .16
FeO*	6.53 ± .58	6.52 ± .17
Cr ₂ O ₃	0.92 ± .07	0.56 ± .05
MnO	0.13 ± .04	0.16 ± .03
NiO	0.09 ± .09	0.04 ± .07
MgO	18.49 ± .21	16.29 ± .39
CaO	17.57 ± .92	19.25 ± .39
Na ₂ O	0.29 ± .02	0.43 ± .04
Total	98.22	99.12
Number of ions on the basis of 6 oxygens		
Si	1.934	1.914
Al	0.080	0.117
Ti	0.012	0.040
ΣFe	0.203	0.201
Cr	0.027	0.016
Mg	1.025	0.898
Ni	0.002	0.001
Mn	0.004	0.004
Ca	0.700	0.763
Na	0.020	0.030
Total	4.007	3.984
*Total iron as FeO		

Table 3. Electron microprobe analyses of pyroxenes from the Yellow Dog Peridotite expressed in weight percent.

Native nickel-iron alloy, probably awaruite, occurs as rare angular grains up to 15 microns in diameter both in areas of secondary silicates such as serpentine and talc, and as isolated inclusions in olivine or pyroxene crystals. It is most likely the result of a reducing environment generated during serpentinization, producing native metals from decomposed silicates and altered sulfides (Eckstrand, 1975; Botto and Morrison, 1976).

Textures of certain oxide and sulfide phases suggest liquid immiscibility with the crystallizing peridotite. Sulfide globules within pyroxenes are likely the result of sulfur saturation and subsequent sulfide liquid separation. Rounded inclusions of magnetite in pyroxenes and myrmekitic magnetite-chalcopyrite intergrowths suggest contemporaneous immiscibility of oxides and sulfides from the silicate melt.

Trace Element Content

Selected trace element contents from the Sections 11 and 12 outcrops are given in Table 4. Some elements show significant increases in concentration relative to world averages for peridotite and other ultramafic rocks (Table 5). Most important are large anomalies for copper and sulfur, as well as zinc and possibly silver. Zinc contents are approximately double those for average ultramafic rock. Varying limits of detection shown on Table 5 preclude an accurate assessment of average silver values, but the presence of 0.2 ppm silver in four samples (Table 4) is clearly anomalous. Cobalt and nickel contents fall within typical ultramafic ranges; chromium is depleted by a factor of about two. Semiquantitative spectrographic analyses of 10 samples for 50 other elements failed, within detection limits, to identify any unusual metal concentrations.

Copper contents of 37 samples of the Yellow Dog Peridotite range from 90 to 350 ppm and average 150 ppm, compared to worldwide means of less than 50 ppm reported by various authors (Table 5). The copper is not evenly distributed between the two outcrops, with the smaller Section 11 outcrop considerably enriched relative to the Section 12 outcrop. The two outcrops average 270 and 120 ppm, respectively. Copper values are clearly anomalous in relation to most other ultramafic rocks.

Sample	Ag	Cu	Co	Cr	Ni	Zn
1	na	160	na	na	1400	na
2	na	145	na	na	1400	na
3	na	335	na	na	1700	na
4	na	115	na	na	1200	na
5	na	340	na	na	1700	na
6	na	125	na	na	1300	na
7	<.2	90	50	3000	1300	<200

8	<.2	110	na	na	1450	na
9	<.2	190	na	na	1450	na
10	<.2	115	na	na	1300	na
11	<.2	90	na	na	1300	na
12	<.2	100	na	na	1250	na
13	<.2	95	na	na	1300	na
14	<.2	90	na	na	1350	na
15	<.2	105	na	na	1350	na
16	<.2	100	na	na	1250	na
17	<.2	115	na	na	1400	na
18	<.2	115	na	na	1150	na
19	<.2	110	na	na	1250	na
20	<.2	90	na	na	1250	na
21	<.2	100	na	na	1250	na
22	<.2	95	na	na	1200	na
23	<.2	105	na	na	1100	na
24	.2	120	na	na	1050	na
25	<.1	120	93	1300	1400	110
26	<.1	120	95	1300	1400	120
27	<.1	90	85	1300	1300	130
28	<.1	290	93	1400	1700	110
29	<.1	120	91	1300	1500	110
30	<.1	120	86	1400	1400	120
31	<.1	88	90	1400	1400	110
32	<.1	120	86	1300	1300	130
33	<.1	310	90	1300	1400	110
34	<.1	190	95	1300	1400	120
35	.2	175	na	na	1400	na
36	.2	340	na	na	1450	na
37	.2	355	50	2000	1250	<200

Samples 1 through 6 from unspecified location on Section 12 outcrop, and 7 through 24 and 35 through 37 from locations shown, on Figure 2. Analyzed by atomic absorption techniques by Rocky Mountain Geochemical Corp. except for Co, Cr, and Zn, which were spectrographic analyses by Skyline Labs Inc. Samples 25 through 34 are from locations shown on Figure 2, and analyses are by a semiquantitative spectrographic technique at U.S. Geological Survey analytical laboratories, Reston, Va.

na - not analyzed.

Table 4. Trace element content, of the Yellow Dog Peridotite expressed in parts per million (ppm).

Sulfur content is also distinctly higher in Yellow Dog Plains samples than in average ultramafic rock (Table 5). The most recent compilation of Wedepohl (1975) suggests 400 ppm (0.04 percent) as a mean sulfur value for ultramafic rock. Four Yellow Dog Peridotite samples yielded a range from 700 to 1740 ppm sulfur (0.070 to 0.174 weight percent), with an average of 1340 ppm (0.134 percent). Sulfur and copper contents may be used as effective discriminators of ore potential, utilizing the method of Cameron et al. (1971). The sulfur contents of over 1000 ultramafic rocks of the Canadian Shield were grouped by them into barren, minor ore, and ore categories, based on copper-nickel production figures. Barren ultramafics (N = 616 samples) display sulfur contents ranging from 0.011 to 0.327 weight percent, with a mean of 0.059 and standard deviation of 0.107. Minor ore samples (N = 91) vary from 0.029 to 0.879, with a mean and standard deviation of 0.177 and 0.559, respectively. Ultramafic rocks associated with 16 major copper-nickel ore deposits (N = 372) contain from 0.112 to 1.92 percent sulfur, with a mean of 0.582 and standard deviation of 1.37. Although there is much scatter in the data, sulfur contents of the Yellow Dog Peridotite are most similar to the minor ore category, as defined by Cameron et al. (1971). Nickel and cobalt contents of major ore suites are not significantly enriched relative to typical ultramafic rocks (Cameron et al., 1971, p. 308), a pattern followed by Yellow Dog Plains samples (Table 5). Notably, copper was found to be the second most important discriminator of ore potential, after sulfur. Anomalously high copper and sulfur contents determined for Yellow Dog Plains samples suggest the possibility of economic copper-nickel mineralization at depth.

PALEOMAGNETIC STUDIES

Seven hand specimens were collected from the two outcrops of the peridotite. Two cores from each sample were analyzed by Kenneth Books of the U.S. Geological Survey in the Reston magnetic laboratory. Measurements were made for susceptibility, direction and intensity of remanence. Susceptibility was converted to intensity of induced magnetism. Remanence data for natural remanent magnetization and for succeeding steps of alternating current cleaning up to 500 oersteds were

analyzed by computer and a Fisher statistical analysis run for each step. The optimum cleaning step (least scatter of direction) appeared to be at 200 oersteds but even at that step the samples showed considerably more scatter than other lower Keweenawan rocks previously analyzed from the region. The virtual geomagnetic pole data are: latitude -58.68; longitude 142.62 east; delta m = 38.58; and delta p = 34.29. The "optimum" cleaned direction of magnetization for the peridotite is not significantly different statistically from some other lower Keweenawan directions of magnetization and is very similar to directions obtained for diabase dikes nearby (DuBois, 1962). The peridotite undoubtedly can be assigned a lower Keweenawan age.

The relatively large scatter and differing polarizations of the natural remanent magnetization obtained from different samples indicate secondary components of magnetization. These components may be at least partially responsible for the net north-seeking down magnetization in the peridotite and could produce the positive magnetic anomaly over the peridotite. Although probably small, another contributor to a positive anomaly over the peridotite might be the moderate intensity of the induced magnetization (2.50×10^{-3} emu/cm²). This induced magnetization may be compared to the average intensity of remanent magnetization for the seven samples of 69.20×10^{-3} emu/cm².

It is possible that secondary magnetite produced during serpentinization of part of the olivine has a normal remanent direction and is at least partly responsible for the scatter of pole positions and the positive magnetic anomaly.

	Ag	Cu	Co	Cr	Ni	Zn	S*
Turekian & Wedepohl (1961)	0.06	10	150	1600	2000	50	300
Vinogradov (1962)	0.05	20	200	2000	2000	30	100
Goles (1967)	--	30	110	2400	1500	--	--
Fisher et al. (1969)**	--	41	101	3040	1581	--	--
Wedepohl (1975)	0.05	47	110	3090	1450	56	400
Yellow Dog Peridotite	0.2	150	87	1525	1350	117	1340
Sample size(n)	(31)	(37)	(12)	(12)	(37)	(10)	(4)

* Sulfur analyses provided by the Institute of Mineral Research, Michigan Technological University.

** Data recalculated specifically for 41 peridotites, excluding other ultramafic rock types.

-- = not given.

Table 5. Comparison of selected trace element contents for average ultramafic rock and Yellow Dog Peridotite expressed in

parts per million (ppm).

FIELD GEOPHYSICAL STUDIES

Although regional aeromagnetic data (Case and Gair, 1965) and regional gravity data (Klasner, 1977) are already available, further geophysical studies were conducted to determine more precisely the size and location of the peridotite bodies. These studies consisted of detailed surveys near the peridotite outcrops in Sections 11 and 12, T50N, R29W, plus more widely spaced surveys located along an east-west belt about 20 kilometers long and at a greater distance from the outcrops. Gravity (Figure 3), ground magnetic (Figure 4) and VLF-EM (Figure 5) techniques were used in the detailed survey area. Ground magnetic and VLF-EM profiles were measured along selected lines outside the detailed area (see Figures 1 and 6).

The combined use of all three geophysical techniques greatly restricts the spectrum of possible geologic bodies responsible for the measured anomalies and allows a closer approach to a unique solution than would be possible with any single technique. Complications of interpretation arise, however, from several factors. Most important are: 1) the complexities of the magnetic field caused by the interaction of the induced field and the irregularly oriented natural remanent field; 2) the possible variations in density of the peridotite due to variable degrees of serpentinization; 3) imprecise knowledge of the densities of all rock types in the area; 4) variations in thickness of Pleistocene drift; and 5) the imprecise understanding of the composition of the buried conductive bodies that produced a measurable VLF-EM response.

In spite of these difficulties, much useful information has been obtained on the geologic nature of the area, and geophysical models were prepared that seem consistent with the observed geological and geophysical data.

Field Procedures and Data Reduction

Gravity

Figure 3 shows the gravity coverage in the detailed study area and the location of two regional profiles. In the detailed study area, nine gravity profiles spaced 400 meters apart were measured in a north-south direction, roughly perpendicular to the trend of the aeromagnetic anomaly. Station spacing on the profiles was 152 meters. The two regional profiles extend across the southernmost aeromagnetic anomaly as well as the northern anomaly.

Gravity values were measured with a LaCoste and Romberg model G land gravimeter. Drift checks, taken at bench mark 1422 every three hours or less, had less than 0.1 milligal drift per hour. The gravity readings were tied to the Marquette gravity base.

Most elevations were determined by leveling between U.S. Geological Survey benchmarks. North of the base line (see Figure 3), surveys closed within 5 centimeters;

south of the baseline, where swamps made surveying difficult, elevation surveys closed within 18 centimeters or better. A few elevations shown on Figure 3 were determined by altimeter with an estimated accuracy based on repeated readings of L6 meters or less. All distances were measured with a steel tape and have an estimated accuracy of 98% or greater.

The gravity data were reduced using the 1967 international gravity formula, a 427-meter above sea level datum, and a Bouguer density of 1.7 g/cm^3 . The 1.7 g/cm^3 value is the density of the sand and silt that covers the bedrock and was determined from a density profile measured over a small topographic feature in the sand plains.

Ground Magnetics

Figure 4 shows the ground magnetic coverage in the detailed study area as well as several profiles that were extended southward into Sections 13 and 14. North of the base line, seventeen profiles were measured; south of the base line ten profiles were measured. Station spacing along the profiles was 30.3 meters. North of the base line, station spacing was measured with a steel tape and south of the base line distances were determined by pace.

Figure 6 shows the regional ground magnetic coverage. Outside of the detailed study area, twenty profiles were measured with spacing between profiles from 800 to 1,200 meters. Five profiles from Figure 4 are included in Figure 6 to maintain continuity of coverage. Station spacing for the regional profiles was 30.3 meters and was determined by pace.

All readings were taken with a Geometrix model G816 proton precession magnetometer in the backpack mode. Base checks for magnetic drift were made at bench mark 1422 every three hours or less. Data were discarded if the magnetic drift exceeded 20 gammas per hour.

Inasmuch as all data were tied to a common base, this base served as a reference for the magnitude of the total magnetic field in this area. It was assigned a value of 59,780 gammas, based on the average of 25 readings which had a total range of 60 gammas (59,766 to 59,826 gammas).

VLF-EM

VLF-EM coverage in the detailed study area consisted of 16 lines, shown on Figure 5. All readings were taken with a Geonics EM-16 unit facing north. The station spacing along the profiles was 30 meters, determined by pace. Transmitter locations used in the survey were NLK, Seattle, Washington (18.6 kHz) and NAA, Cutler, Maine (17.8 kHz).

The location of two lines of regional VLF-EM coverage is shown on Figure 1. Line 1 is the southern extension of the westernmost line in the detailed study area. It extends across the southernmost aeromagnetic anomaly. Line 3, located 9 kilometers west of the detailed study area, crosses both the northern and southern aeromagnetic

anomalies. Station spacing on both lines was 30 meters, determined by pace.

To make the VLF-EM data easier to interpret and to smooth noisy data, all of the profiles, including both the in-phase and quadrature components, were filtered by a method outlined by Fraser (1969). The filter is a difference operator and low-pass filter that smoothes the data and transforms zero crossovers into peaks and troughs, depending on the direction of the crossover. These filtered data are then contoured into positive and negative anomalies as shown on Figure 5.

Interpretation of Geophysical Data

Figure 5 shows that both positive and negative VLF-EM anomalies are present in the detailed study area. Positive in-phase anomalies, which ideally have a positive peak flanked by nearly symmetrical negative shoulders, generally indicate the presence of anomalous conductivity. Negative in-phase anomalies ideally have a pronounced trough with nearly symmetrical positive shoulders. Negative VLF-EM anomalies may be caused when: A) a resistive body is located in a field of more conductive material, such as a resistive dike outcropping in a swamp or area of conductive clay (Ronka, Patterson, written communication), or B) a dike with a high magnetic permeability and a low degree of conductivity causes a response in the VLF-EM receiver coils due to the natural magnetic field overriding the electromagnetically induced field (Grant and West, 1963).

It appears from this study that the negative in-phase VLF-EM anomalies are caused by conditions similar to the second situation (B) outlined above. Two of the negative anomalies (Bn and Cn) occur over the peridotite outcrops in areas of thick dry sand. And all of the negative VLF-EM anomalies have associated positive magnetic anomalies.

The positive VLF-EM anomalies (Figure 3), on the other hand, do not have associated positive magnetic anomalies. In fact there seems to be little, if any, associated magnetic expression.

It is significant that, with the exception of anomalies Cp and the northwest extension of Bp, all VLF-EM anomalies, both positive and negative, have associated positive residual gravity anomalies suggesting that the bodies which cause the electromagnetic anomalies are denser than the enclosing country rock. A cluster of mafic dikes, denser than the enclosing country rock as shown in Figure 7, could cause this unique combination of geophysical anomalies. The nature of these dikes is discussed in the following paragraphs




Figure 3. Complete Bouguer gravity anomaly.




Figure 4. Ground magnetics of detailed study area.

In order to interpret the gravity data, it is necessary to separate the Bouguer gravity anomaly map (Figure 3) into regional and residual components. This was done graphically by the cross profile method, and Figure 8 shows the different components into which the Bouguer anomaly was separated. R_1 represents the regional gravity gradient. No map has been prepared for this surface. R_2 is a regional surface, possibly caused by basement faulting or by variations in thickness of Pleistocene cover or both, as discussed below. R_3 is the observed Bouguer gravity anomaly. Residual gravity maps were prepared by subtracting R_1 from R_2 to form

gravity residual-1, as shown in Figure 9, and R_2 from R_3 to form gravity residual-2, as shown in Figure 10.

Theoretical two-dimensional gravity models (Talwani et al., 1939) were constructed along profiles AA' and BB' (Figures 11a and 11b) to try to explain the residual gravity-2 anomalies.

Preliminary geologic reports (Trow, 1978 a,b,c) for three holes drilled recently in the Baraga Basin near the area of study provide valuable data for construction of the gravity models. Approximate location of the three holes is shown on Figure 1. Holes DL-1 and DL-7 are located near the north edge of the Baraga Basin, whereas DL-3 is located in the south half of the basin. For the purpose of the gravity modeling, the most important points to be considered from the drilling are: a) depth of overburden varies from zero to over 260 feet, b) gamma-ray density logs indicate that the bulk density of middle Precambrian rocks ranges from 2.80 to 2.83, and c) in the north half of this part of the Baraga Basin, depth to basement is greater than 343 meters and in the south half it is greater than 1,000 meters.

Rock density measurements from chip samples of Yellow Dog Peridotite and rock density data from published work provide additional information for construction of the models. Density measurements on two separate samples of the peridotite indicate that the density may vary from at least 2.83 gm/cm³ to 2.93 gm/cm³. This range is much lower than the average density of fresh peridotite (3.23 gm/cm³) as reported in the Handbook of Physical Constants (Clark, 1966) and is probably related to the degree of serpentinization of the peridotite. Recent mapping in Archean terrain south of the Baraga Basin shows that basement rocks are predominantly syenite with 10 to 23 percent mafic gneiss. North of the basin, drill hole DL-7 encountered granodiorite in the basement, but this area has not been mapped in sufficient detail to determine how much, if any, mafic gneiss is intermixed with the granodiorite. According to Clark (1966), the average density for syenite is 2.76 gm/cm³ and for granodiorite it is 2.72 gm/cm³. Considering a density of 2.92 to 3.13 gm/cm³ for mafic gneiss (Klasner and Cannon, 1974; Klasner et al., 1978), basement rocks south of Baraga Basin could have an average density as high as 2.83 gm/cm³ and the rocks north of the basin could have an average density of 2.72 gm/cm³ or greater, depending on the percentage of mafic gneiss intermixed with granodiorite. At any rate, Precambrian W basement rocks appear to have about the same density as the overlying Precambrian X rocks.

In summary, then, the major factors to be considered in construction of the gravity models and interpretation of the gravity residual maps (Figures 9 and 10) are as follows:

A) Density ranges permissible for modeling are 1.7 gm/cm³ for Pleistocene drift, about 2.80 to 2.83 gm/cm³ for Precambrian X rocks, about 2.75 to 2.83 gm/cm³ for Precambrian W basement rocks, and 2.83 to 2.93 for peridotite.

B) The density contrast between drift and underlying bedrock is about -1.10 gm/cm³. This large contrast in

density makes variations in drift thickness an important factor in modeling. For example, a change in thickness of drift from 30.3 meters to 4.5.5 meters creates a 1.3+ milligal anomaly, which is an amount nearly as large as the variation in residual gravity shown on Figure 9.

C) Depths to Archean basement must be greater than the .545-meter and 1,000-meter drill hole depths in the north and south parts of the basin, respectively.

Because of the small density contrast between basement and Precambrian X rocks, and our imprecise knowledge of variations in drift thickness, no attempt was made to model the depth and configuration of the Baraga Basin. It is possible, however, that the south-sloping gradient in the gravity residual-1 (Figure 9) is caused by a down-to-the-south fault in the basement as suggested in previously constructed preliminary models (Klasner et al., 1977), or it may reflect the presence of a deeply buried mafic body of rock beneath the Baraga Basin.

The east-west trending, positive gravity anomaly (gravity residual-2) shown in Figure 10 can be explained, for the most part, by three east-west trending dikes as shown on Figures 11a and 11b. The combined gravitational effect of the dikes explains the 1,200-meter width of the anomaly. The position of the dikes is controlled by outcrops at Bn and Cn, the location of positive magnetic anomalies and negative electromagnetic anomalies (An), and the location of positive electromagnetic anomalies (Ap and Bp). Depth to the top of the dikes and dike width are not extremely critical parameters in the model. Therefore, these factors in the model should be considered rough approximations only.

Although the gravity residual-2 can be roughly explained by the dikes, it was necessary to include a layer of Pleistocene drift of varying thickness to create a closer match between observed and calculated gravity. This layer of very low density material, however, lowered the average amplitude of calculated gravity to a value well below that of the observed gravity residual-2. To counteract this, a large pod of dense mafic rock buried at ten to fifteen thousand feet below the ground surface is included in the model. Inasmuch as the mineralogy and chemical composition of the peridotite suggests that it formed by differentiation from a basaltic magma, the presence of such a mafic body at depth has geologic merit. However, more information is necessary to either confirm or refute this; therefore, this part of the model is intended primarily to illustrate that the gravity data permit the interpretation that a large mafic rock mass exists at depth. Other models could be constructed to explain the observed gravity data.

The important point to note in these two models (Figures 11a and 11b) is that the gravity data suggest excess mass beneath the magnetic and electromagnetic anomalies. The only alternate explanation that might be employed to explain such a unique combination of gravity, magnetic, and electromagnetic anomalies is abrupt high-frequency variations in drift thickness with accompanying magnetic and electromagnetic effects. In particular, this rationale might be used to explain anomalies Ap and Bp. Although

they have positive electromagnetic anomalies, they do not have accompanying magnetic anomalies. It is possible that ridges of graphitic slate stick up into the overlying drift and cause both the electromagnetic and gravity expression.

On the Bouguer gravity anomaly map of the Baraga-Dead River basins area (Klasner, 1977), a large, low amplitude, positive gravity anomaly is centered about two miles northeast of the detailed study area. This anomaly, which is at least one mile in diameter and about two milligals in amplitude, may represent a buried mafic body of rock that was once the magma chamber from which the Yellow Dog Peridotite differentiated. The anomaly appears geophysically continuous with the positive gravity anomalies over the detailed study area and may represent the northeastward continuation of the deeply buried mafic body postulated in the two-dimensional gravity models.

Interpretation of Individual VLF-EM Anomalies

The map of VLF-EM anomalies (Figure 3) shows four positive anomalies, Ap, Bp, Cp and Dp. Anomaly Ap, which has the highest amplitude, is probably offset by a small northwest-trending fault. Gravity model AA' (Figures 9, 11a) shows that it can be caused by a 30-meter-thick dike. Because it does not have an associated ground magnetic anomaly, the causative material may be something other than peridotite, and the anomalous conductivity indicated by the positive VLF-EM anomaly suggests that it may contain sulfide mineralization. Estimated depth to the top of the conductive body from VLF-EM data is 37 meters.

VLF-EM anomaly Bp is probably a compound anomaly. The high-intensity part of the anomaly (20 arbitrary filtered units) may be due, in part, to clay and electrolyte solutions in the underlying swamp. However, the EM profile along 1000W has a negative in-phase/quadrature relationship which may indicate a subsurface conductor other than the swamp. This possibility is supported by gravity model BB' which shows a 76-meter-thick dike at Bp near line 600W. There is only a slight magnetic anomaly over the dike, indicating small amounts of magnetite, but the positive EM anomaly indicates the possibility of sulfide mineralization. VLF-EM data on line 1000W suggests that the conductive body is 45 meters below the surface.

Both the positive VLF-EM anomaly Cp and the northwest extension of Bp trend north-northwest at a large angle to other electromagnetic anomalies in the area. The VLF-EM anomalies may be caused by electrolytic solutions or metallic mineralization in the shear zone of faults. Their rather low intensity may be due to the fact that the strike of the causative feature is at a high angle to the survey direction and not perpendicular to the nearly north-south transmitted electromagnetic field direction, so that a less-than-maximum coupling of the conductor with the transmitted field was achieved.

VLF-EM anomaly Dp has a maximum intensity of slightly more than ten arbitrary units. It falls within an area of a slight positive gravity anomaly but has no magnetic

expression. VLF-EM data indicate that the top of the causative body is about 30 meters below the surface. The nature of the causative body is not known.

Three negative anomalies occur in the filtered VLF-EM data, An, Bn, and Cn, shown on Figure 5. Anomaly An is continuous in an east-west direction for about 2,800 meters, the total extent of the detailed study area. Locally it has an amplitude of 20 units, which may be due to the causative body being very near the surface. The anomaly is very weak in the area of the northwest-trending fault zone (lines 00 and 200E). Anomaly An is coincident with a magnetic anomaly (compare Figure 5 with Figure 4 and see also Figures 11a and 11b). Gravity data suggest that the causative body is a vertical dike about 30 to 45 meters thick. As with all of the negative VLF-EM anomalies, the combined VLF-EM and magnetic data indicate that the causative bodies have high magnetic permeability, probably caused by magnetite.

Negative VLF-EM anomaly Bn has the highest absolute intensity of any of the EM anomalies. It overlies a peridotite outcrop but is much larger in area than the outcrop. It is coincident with an intense (1700 gammas above background) ground magnetic anomaly. Gravity model AA' (Figure 11a) shows that the causative feature may be a vertical dike about 90 meters thick. All geophysical data indicate that the dike ends abruptly near the west edge of the outcrop. It is probably truncated by a north west-trending fault.

VLF-EM anomaly Cn is also associated with a peridotite outcrop. The anomaly trends east-west and is much narrower and smaller in amplitude than Bn. Gravity model BB' (Figure 11b) indicates that the anomaly may be caused by a 30-meter thick vertical dike. This dike is also abruptly terminated at the northwest-trending fault zone.

A summary of data and interpretation of the anomalies shown on Figure 5 is listed on Table 6.

Regional Survey

Regional geophysical coverage outside of the area of the detailed survey was designed to accomplish two objectives: 1) to test the continuity of the inferred dikes in an east-west direction, and 2) to examine the VLF-EM response in the area of the aeromagnetic anomalies at three locations.




Figure 7. Geology of the detailed study area inferred from geophysical surveys.




Figure 8. Gravity profile along A – A', showing how residual gravity anomalies were determined.




Figure 9. Gravity residual – 1, anomaly caused by deep body.




Figure 10. Gravity residual – 2, anomaly caused by near-surface dikes.

VLF-EM			Gravity	Causative Feature	Orientation and Extent	Estimated Depth to Body	Remarks
Anomaly	Sign	Ground magnetics					
An	-	+ 59,900 gammas	positive anomaly, modelled 61 m thick	nearly vertical dike	generally N 75° W 2.8 km long	-	probable magnetite mineralization
Bn	-	+ 61,500 gammas	positive anomaly, modelled 120 m thick	nearly vertical dike	E-W about 800 m long	outcrops	see petrographic analyses
Cn	-	+ 60,000 gammas, complex body suggested	positive anomaly, modelled 30 m thick	nearly vertical dike	N 80° E 600 m long	outcrops	see petrographic analyses
Ap	+	no apparent expression	positive anomaly, modelled 30 m thick	nearly vertical dike	N 80° W 1 km long	37m, VLF-EM	possible sulfide mineralization
Bp	+	only minor magnetic expression	positive anomaly, modelled on line 600 W 76 m. thick	compound swamp, dike, and fault	N 45° W 900 m long	45m on line 1000 W, VLF-EM	compound anomaly possible sulfide mineralization
Cp	+	no apparent expression	zero to negative anomaly	fault zone	N 25° W about 800 m long	-	geophysics does not indicate sulfide mineralization
Dp	+	no apparent expression	slight positive anomaly, not modelled	unknown	irregular generally E-W about 400 m long	-	causative body not known but VLF-EM suggests conductivity

Table 6. Summary of data and interpretations of anomalies. See Figure 5 for locations.

GEOCHEMICAL SURVEY

Because the Yellow Dog Peridotite is much richer than the surrounding rocks in certain elements such as copper, nickel, cobalt, zinc, chromium, and magnesium, geochemical analysis of soils in areas of the positive aeromagnetic anomalies might give additional confirmation of the presence of buried peridotite. Near the peridotite outcrops and in most of Yellow Dog Plains, the Pleistocene drift cover is relatively thick and is mostly porous stratified sand and gravel. The probability that a geochemical anomaly from buried bedrock would be present in soil developed on the drift is slight and such areas were not tested. The westernmost extension of the magnetically anomalous area has thinner drift cover and the drift is mostly till. We determined that chances were good to detect anomalous bedrock compositions by soil analyses in that area.

Samples were collected along north-south traverses, shown in Figure 13, that crossed both the northern and

southern aeromagnetic anomalies. Samples were taken roughly 40 centimeters below the surface, generally below the organic-rich upper soil horizons. Most of the material sampled was reddish brown sandy soil. In the laboratory the samples were sieved and the minus 80-mesh fraction was analyzed for 65 elements by a semi-quantitative spectrometric technique at the U.S. Geological Survey analytical laboratories in Reston, Virginia.

The results show that weak geochemical anomalies occur at or near the magnetic anomalies on two of the three profiles (Profiles A and B, Figure 13). The third profile (C) had no anomalies but is mostly on thick sand deposits of the Yellow Dog Plains through which effective geochemical communication between soil and bedrock is unlikely. Figures 14 and 15 show the content of elements along Profiles A and B. On Profile A, a broad anomalous area with higher than average values for cobalt, copper, nickel, and zinc coincides with the northernmost aeromagnetic anomaly. A narrower, weaker anomaly in these elements is suggested near the southernmost aeromagnetic anomaly. Silver and chromium also appear to be slightly anomalous in the areas of the magnetic anomalies. Likewise, two separate anomalous areas occur on Profile B. A weak cobalt-copper-nickel anomaly is near the northernmost aeromagnetic anomaly. A more pronounced anomaly in silver, cobalt, copper, nickel, and zinc is about 0.4 kilometer south of the southernmost magnetic anomaly. There is a suggestion of higher-than-average chromium with both anomalies but the relationship is weak. The southernmost anomaly on B also has higher-than-average magnesium content, but the three remaining anomalies do not have high magnesium values. This may indicate that the anomalies are caused mostly by elements such as copper, cobalt, nickel, and zinc being released from sulfide minerals during weathering, whereas magnesium and chromium, although abundant in the peridotite or mafic rocks, are held in less readily-weathered silicate or oxide minerals.

Earlier work by the U.S. Geological Survey in the region immediately west of the area of this study disclosed weak soil anomalies that are clearly related to weakly anomalous zones in the Michigamme Formation. There, the anomalies in both bedrock and soil are expressed by all or some of the elements that are anomalous on profiles A and B, plus lead and molybdenum. The soil anomalies A and B do not have anomalous lead and molybdenum values and chromium appears more consistently high there than in anomalies related to the Michigamme Formation. The suite of anomalous elements in profiles A and B mirrors the element distribution of peridotite more closely than the element distribution in the Michigamme Formation.

SUMMARY AND CONCLUSIONS

Peridotite of early Keweenawan age is exposed at two outcrops in Sections 11 and 12, T50N, R29W, in an area generally mantled with Pleistocene outwash and till. Geophysical studies indicate that the peridotite is an east-west elongate body at least 1.5 kilometers long. Gravity,

ground magnetic, and VLF-EM surveys have disclosed several other anomalies nearby with the same trend as the peridotite. Some of these anomalies extend at least 15 kilometers west and 5 kilometers east of the peridotite outcrops and are coincident with a 20-kilometer-long aeromagnetically anomalous zone consisting of two parallel positive anomalies about 1 kilometer apart.

Eight to 12 kilometers west of the outcrops, geochemical soil anomalies of several elements, abundant in the peridotite, are coincident with the aeromagnetic positive anomalies and may indicate that peridotite extends that far west. Therefore, with decreasing degrees of certainty, peridotite and related rocks can be inferred in a belt from as little as 1.5 to as much as 20 kilometers long.

The chemical composition and texture of the peridotite from the outcrops lead the authors to believe that it is a differentiate from a less mafic magma rather than a komatiite-like ultramafic rock. Thus, it seems likely that a larger body of mafic rock from which the peridotite has formed by fractional crystallization and differentiation lies buried somewhere in the region. A broad, low frequency and low amplitude gravity anomaly near the peridotite outcrops may be caused by such a body at depth.




Figure 11a. Composite geophysical profiles and geologic model

for cross section A – A'.

Highly differentiated mafic rocks can be hosts for a variety of ore deposits, including copper, nickel, chromium and precious metals. The peridotite in the two outcrops contains an average of 150 ppm copper and some samples contain as much as 355 ppm. Sulfur contents range from 0.07 to 0.174 percent. The rock is considerably richer in both copper and sulfur than average peridotite, and in that respect resembles ore-bearing peridotites in other parts of the Canadian Shield. Positive EM anomalies such as Ap and Bp in areas of positive gravity anomalies may be caused by zones of sulfide concentrations in peridotite or mafic rocks, and need further study to determine the nature of the conductive body. Should the anomalies be shown to be caused by sulfide mineralization, the 20-kilometer-long magnetically anomalous zone would be a well-defined exploration target for extensions of that mineralization.

The Yellow Dog Peridotite is the only lower Keweenaw peridotite known in northern Michigan. Its tectonic significance is not yet known. Paleomagnetic data indicate that it is approximately the same age as the Duluth Complex in Minnesota. Prior to middle Keweenaw rifting it probably was much closer to the Duluth Complex than it is now. Perhaps there is a genetic relationship between the Yellow Dog and Duluth rocks. If the two are comagmatic, the Yellow Dog Peridotite may be a favorable host rock for copper-nickel mineralization similar to that found in parts of the Duluth Complex.




Figure 11b. Composite geophysical profiles and geologic model

for cross section B – B'.




Figure 12. Composite geophysical profiles along regional lines 1 and 3.




Figure 13. Location of geochemical soil profiles.

Figure 14. Geochemical profiles in soil along profile A.




Figure 15. Geochemical profiles in soil along profile B.

REFERENCES

- Arndt, N. T., A. J. Naldrett, and D. R. Pyke, 1977, KOMATIITIC AND IRON-RICH THOLEIITE LAVAS OF MUNRO TOWNSHIP, NORTHEAST ONTARIO, *Journal of Petrology*, v. 18, p. 319-369.
- Bence, A. E., and A. L. Albee, 1968, EMPIRICAL CORRECTION FACTORS FOR THE ELECTRON MICROANALYSIS OF SILICATES AND OXIDES *Journal of Geology*, v. 76, p. 382-403.
- Botto, R. L., and G. H. Morrison, 1976, JOSEPHINITE: A UNIQUE NICKEL-IRON, *American Journal of Science*, v. 276, p. 241-274.
- Cameron, E. M., G. Siddeley, and C. C. Durham, 1971, DISTRIBUTION OF ORE ELEMENTS IN ROCKS FOR EVALUATING ORE POTENTIAL; NICKEL, COPPER, COBALT, AND SULPHUR IN ULTRAMAFIC ROCKS OF THE CANADIAN SHIELD, *The Canadian Institute of Mining & Metallurgy Special Volume 11*, p. 298-313.
- Cannon, W. F., 1973, THE PENOKAN OROGENY IN NORTHERN MICHIGAN,

- Cannon, W. F., and J. S. Klasner, 1977,
GEOLOGIC MAP AND GEOPHYSICAL INTERPRETATION OF THE
WITCH LAKE QUADRANGLE, MICHIGAN,
United States Geological Survey
Miscellaneous Geologic Investigations Map I-987.
- Case, J. E., and J. E. Gair, 1965,
AEROMAGNETIC MAP OF PARTS OF MARQUETTE, DICKINSON,
BARAGA, ALGER AND SCHOOLCRAFT COUNTIES, MICHIGAN
AND ITS GEOLOGIC INTERPRETATION,
United States Geological Survey Geophysical
Investigations Map GP-467.
- Clark, S. P., Jr., 1966,
HANDBOOK OF PHYSICAL CONSTANTS,
Geological Society of America, Memoir 97, 587 p.
- Coleman, R. G., and T. E. Keith, 1971,
A CHEMICAL STUDY OF SERPENTINIZATION -BURRO
MOUNTAIN, CALIFORNIA,
Journal of Petrology, v. 12, p. 311-328.
- Dubois, P. M., 1962,
PALEOMAGNETISM AND CORRELATION OF KEWEENAWAN
ROCKS,
Geological Survey of Canada Bulletin 71, 75p.
- Eckstrand, O. R., 1975
THE DUMONT SERPENTINITE; A MODEL FOR THE CONTROL
OF NICKELIFEROUS OPAQUE MINERAL ASSEMBLAGES BY
ALTERATION REACTIONS IN ULTRAMAFIC ROCKS,
Economic Geology, v. 70, p. 183-201
- Fisher, D. E., O. Joensuu, and K. Bostrom,
ELEMENTAL ABUNDANCES IN ULTRAMAFIC ROCK AND THEIR
RELATION TO THE UPPER MANTLE,
Journal of Geophysical Research, v. 74, p. 3865-3873.
- Fraser, D. C., 1969,
CONTOURING VLF-EM DATA,
Geophysics, v. 34, p. 958-967.
- Frost, B. R., 1976,
LIMITS TO THE ASSEMBLAGE FORSTERITE-ANORTHITE AS
INFERRED FROM PERIDOTITE HORNFELSES, ICICLE CREEK,
WASHINGTON,
American Mineralogist, v. 61, p. 732-750.
- Goles, G. G., 1967,
TRACE ELEMENTS IN ULTRAMAFIC ROCKS, In; ULTRAMAFIC
AND RELATED ROCKS,
Wyllie, P. J., (Ed.), John Wiley and Sons,
New York: New York, p. 352-362.
- Grant, F. S., and F. G. West, 1965,
INTERPRETATION THEORY IN APPLIED GEOPHYSICS,
McGraw-Hill Book Co., New York: New York.
- Green, D. H., 1967,
HIGH-TEMPERATURE PERIDOTITE INTRUSIONS, In;
ULTRAMAFIC AND RELATED ROCKS,
Wyllie, P. J., (Ed.), John Wiley and Sons,
New York: New York, p. 212-222.
- Hostetler, P. B., et al., 1966,
BRUCITE IN ALPINE SERPENTINITES,
American Mineralogist, v. 51, p. 75-98.
- Klasner, J. S., 1977,
BOUGUER GRAVITY ANOMALY MAP, BARAGA-DEAD RIVER
BASINS AREA, NORTHERN MICHIGAN,
United States Geological Survey Open-File
Report no. 77-746.
- Klasner, J. S., and W. F. Cannon, 1974,
GEOLOGIC INTERPRETATION OF GRAVITY PROFILES IN THE
WESTERN MARQUETTE DISTRICT, NORTHERN MICHIGAN,
Geological Society of America Bulletin,
v. 85, p. 213-218.
- Klasner, J. S., et al., 1977,
THE YELLOW DOG PERIDOTITE AND A POSSIBLE BURIED
IGNEOUS COMPLEX OF LOWER KEWEENAWAN AGE IN THE
NORTHERN PENINSULA OF MICHIGAN,
United States Geological Survey Open-File
Report no. 77-93, 49 p.
- Klasner, J. S., et al., 1978,
BOUGUER GRAVITY ANOMALY MAP OF THE NORTHERN
MICHIGAN-LAKE SUPERIOR REGION,
United States Geological Survey Open-File
Report no. 78-211.
- Morris, W. J., 1977,
GEOCHEMISTRY AND ORIGIN OF THE YELLOW DOG PLAINS
PERIDOTITE, MARQUETTE COUNTY, MICHIGAN,
Master of Science Thesis, Michigan State University, East
Lansing, Michigan.
- Naldrett, A. J., and L. J. Cabri, 1976,
ULTRAMAFIC AND RELATED MAFIC ROCKS: THEIR
CLASSIFICATION AND GENESIS WITH SPECIAL REFERENCE TO
THE CONCENTRATION OF NICKEL SULFIDE AND PLATINUM-
GROUP ELEMENTS,
Economic Geology, v. 71, p. 1131-1158.
- Nesbitt, R. W., 1971,
SKELETAL CRYSTAL FORMS IN THE ULTRAMAFIC ROCKS OF
THE YILGARN BLOCK, WESTERN AUSTRALIA: EVIDENCE FOR
AN ARCHEAN ULTRAMAFIC LIQUID,
Geological Society of Australia Special Publication 3, p.
331-347.
- O'Hara, M. J., 1967,
MINERAL FACIES IN ULTRABASIC ROCKS, In; ULTRAMAFIC
AND RELATED ROCKS,
Wyllie, P. J., (Ed.), John Wiley and Sons,
New York: New York, p. 7-18.
- Pyke, D. R., A. J. Naldrett, and O. R. Eckstrand, 1973,
ARCHEAN ULTRAMAFIC FLOWS IN MUNRO TOWNSHIP,
ONTARIO,
Geological Society of America Bulletin,
v. 84, p. 955-978.
- Talwani, M., J. L. Worzel, and M. G. Landisman, 1959,
RAPID GRAVITY COMPUTATIONS FOR TWO-DIMENSIONAL
BODIES WITH APPLICATION TO THE MENDOCINO SUBMARINE
FRACTURE ZONE,
Journal of Geophysical Research, v. 64, p. 49-59.
- Thayer, T. P., 1960,
SOME CRITICAL DIFFERENCES BETWEEN ALPINE - TYPE AND

STRATIFORM PERIDOTITE - GABBRO COMPLEXES,
21st International Geological Congress, Copenhagen,
Reports, pt. 13, p. 247-259.

Trow, J., 1978a,
PRELIMINARY GEOLOGIC DATA -DIAMOND - DRILLING FOR
GEOLOGIC INFORMATION IN THE MIDDLE PRECAMBRIAN
BASINS IN THE WESTERN PORTION OF NORTHERN MICHIGAN,
Michigan Department of Natural Resources, Geological
Survey Open File Report no. DOE OFR 78-1, Prepared for
the United States Department of Energy, Project No. 20-
77-2132.

Trow, J., 1978b,
PRELIMINARY GEOLOGIC DATA — DIAMOND - DRILLING FOR
GEOLOGIC INFORMATION IN THE MIDDLE PRECAMBRIAN
BASINS IN THE WESTERN PORTION OF NORTHERN MICHIGAN,
Michigan Department of Natural Resources, Geological
Survey Open File Report no. DOE OFR 78-3, Prepared for
the United States Department of Energy, Project No. 20-
77-2132.

Trow, J., 1978c,
PRELIMINARY GEOLOGIC DATA -DIAMOND - DRILLING FOR
GEOLOGIC INFORMATION IN THE MIDDLE PRECAMBRIAN
BASINS IN THE WESTERN PORTION OF NORTHERN MICHIGAN,
Michigan Department of Natural Resources, Geological
Survey Open File Report no. DOE OFR 78-7, Prepared for
the United States Department of Energy, Project No. 20-
77-2132.

Turekian, K. K., and K. H. Wedepohl, 1961,
DISTRIBUTION OF THE ELEMENTS IN SOME MAJOR UNITS OF
THE EARTH'S CRUST,
Geological Society of America Bulletin,
v. 72, p. 175-192.

Vinogradov, A. P., 1962,
AVERAGE CONTENTS OF CHEMICAL ELEMENTS IN THE
PRINCIPAL TYPES OF IGNEOUS ROCKS OF THE EARTH'S
CRUST,
Geochemistry no. 7, p. 641-664.

Wedepohl, K. H., 1975,
THE CONTRIBUTIONS OF CHEMICAL DATA TO ASSUMPTIONS
ABOUT THE ORIGIN OF MAGMAS FROM THE MANTLE,
Fortschritte der Mineralogie, v. 52, p. 141-172.

STATE OF MICHIGAN
Governor
William G. Milliken

DEPARTMENT OF NATURAL RESOURCES
Director
Howard A. Tanner

GEOLOGICAL SURVEY DIVISION
Chief and State Geologist
Arthur E. Slaughter

NATURAL RESOURCES COMMISSION
Hillary F. Snell, CHAIRMAN, Grand Rapids
Jacob A. Hoefer, East Lansing
Carl T. Johnson, Cadillac
E. M. Laitala, Hancock
(Mrs.) Joan L. Wolfe, Belmont
Charles G. Younglove, Allen Park
John M. Robertson, Executive Assistant



Published by authority of State of Michigan Compiled Laws of 1970 section 321.6

Copies of this and all other Department of Natural Resources publication are available from:
Information Services Center
Michigan Department of Natural Resources
Box 30028
Lansing, Michigan 48909

On deposit in many libraries in Michigan and other selected locations