DEPARTMENT OF NATURAL RESOURCES AND ENVIRONMENT WATER BUREAU APRIL 2010

STAFF REPORT

A BIOLOGICAL SURVEY OF THE TAHQUAMENON, TWO HEARTED, MANISTIQUE, AND MILLECOQUINS RIVERS WATERSHEDS SCHOOLCRAFT, MACKINAC, CHIPPEWA, LUCE, ALGER, AND DELTA COUNTIES, MICHIGAN JUNE AND JULY 2009

INTRODUCTION

Biological, chemical, and physical habitat conditions of selected streams located in the Tahquamenon (HUC 04020202), Two Hearted (HUC 04020201), Manistique (HUC 04060106), and Millecoquins (HUC 04060107) Rivers watersheds (Figures 1 and 2) were assessed by staff from the Surface Water Assessment Section (SWAS).

The specific survey objective of these monitoring activities included the following:

- Evaluate the current biological and physical conditions at targeted and randomly selected stations in the Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds for attainment of Michigan Water Quality Standards (WQS).
- Support water quality-based effluent limit development for National Pollutant Discharge Elimination System (NPDES) permits.
- Identify potential nonpoint source problems and evaluate the effectiveness of specific nonpoint source projects.
- Evaluate general water quality trends within the watershed.
- Satisfy monitoring requests submitted by internal and external customers.

GENERAL WATERSHED HISTORY AND BACKGROUND INFORMATION

The Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds are part of the Northern Lakes and Forests ecoregion (Omernik and Gallant, 1988). Numerous water quality reports documented the biological and habitat conditions found within the watersheds since the 1990s (Goodwin, 2000; Holden, 2005; Madison and Lockwood, 2004; Michigan Department of Natural Resources (MDNR), 1988; Sayles, 1990; Taft, 1989; 1990; 1992a; 1992b; 1994a; 1994b; 1994c; 1998; 2000a; 2000b; and 2005; and Wolf, 2005).

Tahquamenon River Watershed

The Tahquamenon River, made famous in H.W. Longfellow's 1855 epic poem, "*The Song of Hiawatha*," flows approximately 87 miles eastward discharging to Whitefish Bay of Lake Superior. The Tahquamenon River originates in coarse glacial till or outwash materials, flowing down a relatively steep gradient (Waybrant and Zorn, 2008) before entering a large riparian wetland complex that extends for miles to Whitefish Bay.

The headwaters of the Tahquamenon River are considered quality trout water due to the relatively high stream gradient, excellent large woody structure, and an abundance of exposed cobble and gravel. Considerable habitat enhancement work conducted in recent years has protected eroding banks, exposed rock and gravel beds, and scoured the sand bedload to



Figure 1. Upper Peninsula Locations (West) 2009.



Figure 2. Upper Peninsula Locations (East) 2009.

produce deeper holding waters. The fish community reflects a traditional brook trout – mottled sculpin species complex, with large numbers of wild brook trout (Waybrant and Zorn, 2008).

Much of the Tahquamenon River is dark amber colored due to tannins that originate from wetland drainage and organic soils found extensively within the watershed. The lower watershed has light inorganic soils near the main branch, which are well drained and sandy. The sand throughout the region is easily eroded and deposited into rivers and streams (Holden, 2005). The lower river has the second largest waterfall found east of the Mississippi River called Tahquamenon Falls. The Upper Falls has a drop of nearly 50 feet and is more than 200 feet across. The Lower Falls, four miles downstream (Figure 3), is a series of 5 smaller falls (Exploring the North, Inc., 2009).

Two Hearted River Watershed

Two Hearted River originates northwest of Newberry and flows northeast through state forest lands before discharging to Lake Superior. This famous trout stream drains deep sand outwash plains, moraine ridges, and kettle depressions that contain peat bogs (MDNR, 2010 [Draft]). This river system is a snowmelt driven system, which makes the associated groundwater base flow rate very steady (Fongers, 2007). Also within this watershed, the Betsv River discharges to Whitefish Bay of Lake Superior and the Sucker River discharges



to Lake Superior near Grand Marais. Department of Natural Resources and Environment (DNRE) staff monitored this watershed in 1999 and 2004 (Taft, 2000b; Holden, 2005).

Manistique River Watershed

The Manistique River watershed lies within portions of 5 central Upper Peninsula counties (Alger, Delta, Luce, Mackinac, and Schoolcraft) and drains 1,461 square miles (MDEQ, 2004). This large river system was named for the Ojibway word "Monistique," which means "vermillion" (a vivid reddish orange) (New deal/WPA Art in Manistique, Michigan, 2009). The "Monistique River" name was misspelled when registered with the state and became the Manistique River.

The lower 1.7 miles of the Manistique River from the Paper Mill Dam downstream to Lake Michigan are listed as a federal Area of Concern (MDNR, 1987) due to historic problems with polychlorinated biphenyl-contaminated sediment and combined sewer overflows.

Manistique River watershed surveys by DNRE staff were conducted in August 1999 and August 2004. Qualitative biological surveys and habitat evaluations were completed at 10 locations in 1999 (Goodwin, 2000). In 1999, the macroinvertebrate communities rated acceptable and habitat rated from fair to good. Based on the results of the surveys, sampled reaches were determined to attain WQS. Eighteen monitoring sites throughout the entire Manistique River

watershed were evaluated in August 2004 (Wolf, 2005). All stations rated the macroinvertebrate communities either acceptable or excellent and were meeting WQS.

Millecoquins River Watershed

The Millecoquins River watershed has been surveyed by the DNRE three times since 1990 (Taft, 1990; 2000a; and 2005). The waters of the upper watershed originate from springs and lakes south of the city of Newberry and from wetland drainage from the Furlong Creek watershed. All of the flow enters Millecoquins Lake before entering the outlet channel to Lake Michigan through a series of sand dune ridges.

The federal Clean Water Act Section 319 nonpoint source implementation projects that included cattle exclusion, fencing, stream crossing upgrades, and erosion prevention projects were first implemented within the Doe Creek and Furlong Creek watersheds in the early 1990s. Numerous Best Management Practices and stream habitat improvement projects using federal Section 319 or the Clean Michigan Initiative bond monies have been implemented since 1990 by the Luce-West Mackinac County Conservation District. Several of the sites assessed during 2009 are discussed below.

METHODS

The habitat and macroinvertebrate community were qualitatively evaluated using the SWAS Procedure 51 (MDEQ, 1990; Creal et al., 1996) at 37 sites and water chemistry samples were collected at 9 sites (Tables 1, 2, and 5).

The macroinvertebrate communities were assessed and scored with metrics that rate the communities on a scale from excellent to poor. Scores can range from 9 to -9 for macroinvertebrates. Stations with a score greater than or equal to +5 are considered excellent. Stations with a score less than or equal to -5 are classified as poor. Stations with a score of -4 through +4 are classified as acceptable (moderately impaired). Habitat evaluations are based on 10 metrics, with a possible maximum total score of 200. Stations are classified as excellent with a habitat score >154, good with a score between 105 and154, marginal with a score between 56-104, and poor with a score <56.

Two site selection methods were used to assess monitoring locations within the Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds in 2009: (1) stratified random; and (2) targeted. A probabilistic monitoring approach, using stratified random site selection to address statewide and regional questions about water quality, was used to select several stations within the Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds (DNREa, in preparation). In addition to probabilistic monitoring, sites within the Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds were selected for targeted monitoring to fulfill specific monitoring requests, assess known or potential areas of concern, collect information and assess attainment of designated uses from areas where historic information was lacking, or to provide information for NPDES activities.

All of the samples from the rivers were collected, preserved (if necessary), stored at 4° Celsius, and transported to the DNRE's Environmental Laboratory for chemical analysis using standard protocols (MDNR, 1994). The targeted river samples were analyzed for nutrients and the point source discharges were analyzed for nutrients, metals, and conventional parameters.

Three stations were assessed using Michigan's Qualitative Biological and Habitat Survey

Protocols for Nonwadeable Rivers (DNREb, in preparation). These nonwadeable stations were located on both the East Branch Fox River and the Manistique River. Using this nonwadeable procedure, macroinvertebrate communities were scored with metrics that rate water bodies from excellent to poor. Macroinvertebrate ratings from 76-100 are considered excellent, 50-75 good, 25-49 fair, and 0-24 are considered poor.

RESULTS

Probabilistic Sites

Stations 1-25

The DNRE, Water Bureau, has recently begun to incorporate a stratified random sampling design component into the annual watershed assessments (DNREa, in preparation). The purpose of this probabilistic monitoring is to collect biological data for attainment status and temporal trend analysis. Probabilistic sampling allows the conclusions from a limited number of sampling stations to be extended to an entire watershed. The resulting data can be used to infer the condition of the state's waters at site-specific, watershed, or statewide scales.

The DNRE's Macroinvertebrate Community Status and Trend Monitoring Procedure (DNREa, in preparation) is used to estimate the number of river miles supporting the other indigenous aquatic life and wildlife designated use. The status and trend program utilizes river valley segments to provide the basic sampling unit. A river valley segment is defined as a stream reach that is relatively homogenous with respect to ecological segments of rivers and streams that share common geologic, flow, and temperature characteristics. The smallest river unit (i.e., valley segment) that can be interpreted from large-scale geologic maps, are in reality relatively large with each segment having its own unique identification number coupled with a known specific length. As such, valley segments provide a randomly selectable sampling unit that can be stratified by size and/or thermal regime.

To develop a statistically-based estimate of attainment status in the Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds, a total of 25 randomly selected sites were assessed using Procedure 51 (N = 22) and our draft nonwadable procedure (N = 3) (DNREb, in preparation). Sites were stratified based on size and water temperature. The strata included small cold, medium cold, small warm, medium warm, large warm, very large warm, and coastal rivers. Large cold and very large cold river segments are not represented in the Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds. Two sites located on the Manistique River (Stations 22 and 23) and one in the East Branch Fox River (Station 15) were sampled using the nonwadable protocol (DNREb, in preparation). The number of sites in each category, total watershed miles, and probabilistic results are summarized in Table 1. The probabilistic sites correspond to Stations 1-25 in Figures 1 and 2.

All of the 25 probabilistic sites in the 2009 Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds were supporting the "other indigenous aquatic life" component of the designated use specified in Rule 323.1100(1)(e) of the Michigan WQS. Based on the data and probabilistic monitoring methodology, we are 95 percent confident that at least 89 percent of the river miles in these watersheds are attaining the "other indigenous aquatic life" component of the designated use.

The overall mean Procedure 51 score for benthic macroinvertebrates at the 22 wadable probabilistic sites was 3.2. Three of 22 wadable sites rated excellent (+5 or greater) with the

remaining 19 stations rating acceptable. Only 2 of 22 sites had a negative, but still acceptable metric score. The lowest score (-2) occurred in the Shelldrake River (Station 20) that drains a vast wetland landscape.

The total number of taxa in the wadable probabilistic sites ranged from 16-39 (Tables 3A and 3B). The number of Ephemeroptera, Plecoptera, and Trichoptera (EPT) families ranged from 4-15. Mayflies and caddisflies composed 18-65 percent of the macroinvertebrate communities at the wadable probabilistic sites.

Three stations were assessed using Michigan's Qualitative Biological and Habitat Survey Protocols for Nonwadeable Rivers. All 3 sites surveyed in 2009 scored either good or excellent.

Targeted Sites

Station 26

A DNRE, Fisheries Division, status and trends site is located in the **upper Tahquamenon River** at a historic logging site called the Eagle's Nest location. A Procedure 51 biological survey was conducted at this station to compliment Fisheries Division data. The macroinvertebrate community rated excellent with a combined metric score of +8. Thirty-four taxa were found at Station 26, and approximately 60 percent of the individuals found at this station were EPT taxa. The habitat conditions rated excellent.

Station 27

The **Indian River off M-94** is a DNRE, Fisheries Division, status and trends site. This station was previously assessed by DNRE staff in 1999 and 2004 (Goodwin, 2000 and Wolf, 2005). In 1999, the macroinvertebrate community rated acceptable, 24 taxa were found, and approximately 50 percent of the individuals found at this station were EPT taxa. The habitat conditions rated good in 1999 (Goodwin, 2000). In 2004, the macroinvertebrate community rated excellent, 30 taxa were found, and approximately 50 percent of the individuals found at percent of the individuals found were EPT taxa. The habitat conditions rated good in 2004 (Wolf, 2005). The 2009 macroinvertebrate community rated excellent with a combined metric score of +7. Thirty-three taxa were found at Station 27, and approximately 50 percent of the individuals found at this station were EPT taxa. The habitat conditions rated good.

Station 28

Biological surveys on wadable sites in the **lower Manistique River** have not been conducted in the past; therefore, the boat launch location, upstream of the town, was assessed to provide additional watershed coverage. This launch site is upstream of the Manistique Area of Concern, but within the river segment where the limestone river bottom was modified by pine log sorting activities that ended in the early 1930s (Crowe, 1979). The macroinvertebrate community rated acceptable with a combined metric score of +1. Thirty-two taxa were found at Station 28 and approximately 9 percent of the individuals found at this station were EPT taxa. The habitat conditions rated good.

Station 29

A DNRE Water Chemistry Monitoring Program (WCMP) minimally impacted site is located on the **Fox River at the Fox River Campground**. This station was assessed to provide a link

between the WCMP data and macroinvertebrate community. The macroinvertebrate community rated acceptable with a combined metric score of +4. Thirty taxa were found at Station 29, and approximately 40 percent of the individuals found at this station were EPT taxa. The habitat conditions rated excellent.

The Fox River source water is primarily composed of filtered groundwater emanating from the Kingston sand plains. Water data taken in 2004 (Aiello, 2006) indicates this river system is very soft (hardness range of $39-60 \text{ CaCo}_3$ parts per million [ppm]) with low conductivity (85-114 micromhos per centimeter), elevated dissolved oxygen (10-12 ppm), and neutral pH (7.6-8.1 standard units).

Station 30

The Cut River, a small Lake Michigan coastal tributary, was sampled upstream of US-2 in

Mackinac County, southeast of Rexton (Figure 4). This river flows through large areas of forested wetland and steep sand dune ridges before entering Lake Michigan through a deep gorge. The DNRE has conducted only limited macroinvertebrate sampling in the Cut River headwaters. Our Procedure 51 macroinvertebrate survey indicated an acceptable community (+4) with excellent habitat conditions. Eighteen taxa were collected, with baetid mayflies, midges,



and blackflies dominating the community.

Station 31

The **Black River**, a Lake Michigan tributary, was sampled near the Black River State Forest Campground in Mackinac County, southwest of Garnet. This water body flows through large areas of swamp and sand ridges before entering Lake Michigan. The Procedure 51 macroinvertebrate survey indicated an excellent macroinvertebrate community (+6) with good habitat conditions. Thirty-two taxa were collected with lepidostomatid and limnephilid caddisflies, midges, and baetid mayflies dominating the community.

Station 32

Davenport Creek, a Lake Michigan coastal tributary, was sampled upstream of US-2 in Mackinac County, south of Rexton. This stream flows through large areas of swamp and sand dune ridges before entering Lake Michigan. We conducted Procedure 51 fish and

macroinvertebrate sampling at this specific site approximately 15 years ago (Taft, 1994c). The 2009 Procedure 51 macroinvertebrate survey indicated an acceptable community (+1) with good available habitat present upstream of the highway. Twenty-one taxa were collected with brachycentrid and limnephilid caddisflies, midges, and scuds dominating the community.

Station 33

Davenport Creek at Great Lake Gas Natural Gas Pipeline Crossing - This stream segment was last visited and photographed in 1994 (Figure 5). This site needed to be reassessed to determine the impacts of continued off-road vehicle use at the ford location. The images below (Figures 5 and 6) show that this site has revegetated with speckled alder (*Alnus incana*) and the off-road vehicle usage is much reduced due to fencing that has been erected by the natural gas pipeline company (Figure 6). Procedure 51 macroinvertebrate community and habitat scores were acceptable (+1) and good, respectively. This revegetation appears to have substantially reduced the sand load eroding from this site.



Station 34

Furlong Creek downstream Pleasant Avenue - Furlong Creek is a tributary to Millecoquins Lake and was the target of nonpoint source habitat improvement projects in the early 1990s. A macroinvertebrate survey was conducted at this site for the third time (Taft, 1990; 2000a; and 2005) to verify that the post-project water quality improvements are still being realized. A new properly engineered culvert (Figure 8) was installed to replace a cast-off steel form (Figure 7)





used during the construction of the Mackinac Bridge. This short tube was perched, and created a barrier to the upstream migration of fish as well as a large, 85-foot eddy below the road that eroded the stream bank. Perched culverts have been identified as having numerous negative environmental consequences (Stranahan, 2009). The 2009 Procedure 51 macroinvertebrate survey indicated an excellent community (+5) with good habitat conditions present. Thirty-two taxa were collected with greater than 60 percent of the macroinvertebrate community composed of mayflies and caddisflies.

Station 35

Doe Creek - Historically, this small tributary to Furlong Creek has been impacted by nutrients from local farmsteads, field runoff, and disturbed riparian cover from unrestricted animal access. It is currently listed in Michigan's Clean Water Act Section 305(b), 303(d) and 313 Integrated Report (LeSage and Smith, 2008) as having insufficient water quality information. This creek tends to lose its base flow in dry years so conducting macroinvertebrate assessments in mid-summer can be problematic. The estimated flow on June 10, 2009, was 17 cubic feet per second, which was due to recent rains. The 2009 macroinvertebrate community and habitat were assessed using Procedure 51 at the Raski Road crossing. The June 2009 macroinvertebrate survey indicated an acceptable community (0) with good available habitat conditions present downstream of M-117. Twenty-eight taxa were collected with large numbers of midges, hydroptilid caddisflies, baetid mayflies, and haliplid beetles present. These insects are typically found in abundance in small streams with variable flow conditions and wetland influences.

Stations 36-39

Water samples were collected (where flow conditions permitted) from Station 35 plus 4 additional sites (Stations 36-39) in June 2009 (Table 5). The total phosphorus concentrations were variable with the highest levels found in close proximity to working farms with associated livestock and pasture lands, as would be expected within an agricultural landscape. The total phosphorus concentrations have been relatively constant when compared with past sampling events (Taft, 1990; 2000a).

Station 40

Lower Millecoquins River at the Lower Rapids - This site is downstream of Millecoquins Lake and located within the Hiawatha Club property between US-2 and the Hiawatha Trail (Figure 2). The lower Millecoquins River has not been sampled for macroinvertebrates in the past by the DNRE. This site was selected and sampled for macroinvertebrates and water chemistry to assess WQS attainment. The Procedure 51 macroinvertebrate survey indicated an acceptable community (+4) with excellent available habitat conditions present at this gradient break upstream of the confluence with Lake Michigan. The broken limestone outcropping in the streambed provided optimal hard substrate conditions for colonization by crayfish, snails, mayflies, and caddisflies that require such habitat. Thirty taxa were collected with large numbers of algae grazing glossosomatid caddisflies and lymnaeid snails present.

Station 41

Crow River near Mouth - The State of Michigan acquired most of the property within the Crow River watershed along Lake Michigan within the last 15 years. This place, known locally as Simmons Woods was held by steel companies that needed a source of limestone close to the

surface for flux material. We sampled the Crow River at one station below Amadon Pond in 1999 (Taft, 2000a), which indicated that the stream was meeting WQS with a macroinvertebrate metric score of acceptable (+2). The 2009 Procedure 51 macroinvertebrate survey indicated an acceptable community (+2) with good available habitat conditions present upstream of the confluence with Lake Michigan (Figure 9). Thirty-two taxa were collected with hydropsychid and limnephilid caddisflies, midges, and baetid mayflies dominating the community. We have encountered fishermen that were catching brook trout in past surveys.



NPDES Program Support

To support water quality-based effluent limit development, water chemistry samples were collected at locations on the Tahquamenon River near the Newberry Wastewater Treatment Plant (WWTP). This river has been monitored several times in the past by DNRE staff (Goodwin, 2000; Holden, 2005; and Taft, 1989).

The Newberry WWTP is authorized by their NPDES permit to discharge treated sanitary wastewater to the Tahquamenon River. The Tahquamenon River in the vicinity of the Newberry WWTP was not wadable; therefore, Procedure 51 biological surveys were not conducted upstream and downstream of the WWTP. However, water chemistry grab samples were collected from the WWTP effluent and from the Tahquamenon River upstream of the WWTP. Nutrient concentrations in the WWTP effluent were higher (as expected) than background concentrations measured upstream of the WWTP. The Newberry WWTP is currently in compliance with their total phosphorus limitations and no nuisance plant conditions were observed during our monitoring. Metals such as cadmium, chromium, lead, selenium, silver, and mercury were not detected in either sample; however, it should be noted that mercury was

analyzed using USEPA Method 245.1 with a detection level of 0.2 micrograms per liter, which is significantly higher than the current WQS of 0.0013 micrograms per liter. Chemical data generated for the stations indicate that no toxic chemicals were detected at levels that exceed their respective Rule 57 water quality values.

Nonpoint Source Program Support

The DNRE conducted biological and channel morphology assessments on the Fox River in the vicinity of a bank stabilization project supported by federal Clean Water Act Section 319 funds (Taft, 1992b; and Suppnick, 1996). Biological and physical habitat assessments were also conducted by the DNRE on the Driggs River to collect baseline water quality data prior to the implementation of a bank stabilization project (Taft, 1994b).

As previously discussed, the Luce-West Mackinac Conservation District staff received \$214,000 Section 319 implementation funding for the Doe Creek and Furlong Creek watersheds to address nutrient levels, livestock access, and stream habitat issues in the early 1990s. In 1996, a four-mile section of Furlong Creek was placed on Michigan's Clean Water Act Section 303(d) list for an impaired macroinvertebrate community and excess nutrients due to unrestricted cattle access. The landowner, in collaboration with the DNRE and Michigan Department of Agriculture, developed and implemented a farm management plan, which was successful in recovering the stream reach. Subsequently, the DNRE removed this site from the Section 303(d) list in 2005. Furlong Creek is now recognized as a Section 319 nonpoint source success story, which can be found on the USEPA's Web site (USEPA, 2008).

WATER CHEMISTRY

Ambient water chemistry samples were taken at eight locations within the Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds, along with the primary outfall 001A of the Newberry WWTP (Table 5). The Newberry WWTP effluent total phosphorus concentration was 0.56 milligrams per liter (mg/l), which is well below the 1.0 mg/l monthly average effluent limit in their NPDES permit. A water hardness concentration of 229 mg/l calcium carbonate (CaCo₃) was measured at the Newberry WWTP outfall 001A. No chemicals were detected upstream of the Newberry WWTP or in the outfall 001A discharge that exceeded the Michigan WQS.

Flowing waters directly influenced by dairy and cattle farming activities on Doe Creek had higher phosphorus concentrations as compared to forested or pasture sites. The results appear to be very consistent with past sampling events as the watershed land use has changed little over the last 20 years (Taft, 1990 and 2000a).

CONCLUSION

All 25 probabilistic sites in the 2009 Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds were supporting the "other indigenous aquatic life" component of the designated use (DNREa and DNREb in preparation) specified in Rule 323.1100(1)(e) of the Michigan WQS. Based on the data and probabilistic monitoring methodology, we are 95 percent confident that at least 89 percent of the river miles in these watersheds are attaining the "other indigenous aquatic life" component of the designated use. This is encouraging given the extensive historic logging, limestone mining activities, and past nutrient enrichment pollution problems documented by DNRE staff (Taft, 1990; 1992a; 1992b; 1994a; and 2000a) in the Tahquamenon, Two Hearted, Manistique, and Millecoquins Rivers watersheds. Among 12 targeted sites assessed for benthic macroinvertebrates, 3 scored excellent and 9 rated acceptable (only 1 of the 9 had a metric score less than 0). The habitat condition at five of these targeted sites rated excellent and seven rated good. All targeted stations monitored were meeting Michigan WQS.

Field Work By: William Taft, Aquatic Biologist Matt Wesener, Aquatic Biologist Kay Edly, Aquatic Biologist Dawn Roush, Aquatic Biologist Surface Water Assessment Section Water Bureau

> Seth Wright Great Lakes Environmental Center

Report By: Kay Edly, Aquatic Biologist William Taft, Aquatic Biologist Surface Water Assessment Section Water Bureau

REFERENCES

- Aiello, C. 2006. Michigan Water Chemistry Monitoring: Great Lakes Tributaries 2004 Report. MI/DEQ/WB-06/045.
- Creal, W., S. Hanshue, S. Kosek, M. Oemke, and M. Walterhouse. 1996. Update of GLEAS Procedure 51 Metric Scoring and Interpretation. MDEQ Report #MI/DEQ/SWQ-96/068. Revised May 1998.
- Crowe, W.S. 1979. Lumberjack: Inside an Era in the Manistique, Mich. Region. Senger Publishers. Manistique, MI.
- DNREa. In preparation. Macroinvertebrate Community Status and Trend Monitoring Procedure.
- DNREb. In preparation. Qualitative Biological and Habitat Survey Protocols for Nonwadeable Rivers.
- Exploring the North, Inc. 1997/2009. *Tahquamenon Falls State Park* https://exploringthenorth.com/tahqua/tahqua.html. Viewed March 12, 2009.
- Fongers, D. 2007. Two Hearted River Watershed Hydrologic Study. Land and Water Management Division, MDEQ. January 18, 2007.
- Goodwin, K. 2000. A Biological Survey of the Tahquamenon River and Manistique River Watersheds: Luce, Chippewa, Mackinac, Schoolcraft, and Alger Counties, Michigan. August and September 1999. MDEQ Report #MI/DEQ/SWQ-00/107.
- Holden, S. 2005. A Biological Survey of the Tahquamenon River, Two Hearted River, and Selected Tributaries to Lake Superior Located in Alger, Chippewa, and Luce Counties. August 2004. MDEQ Report #MI/DEQ/WB-05/125.
- LeSage, S. and J. Smith. 2008. Water Quality and Pollution Control in Michigan: 2008 Sections 303(d) and 305(b) Integrated Report. MDEQ Report #MI/DEQ/WB-08/007.
- Madison, G. and R. Lockwood. 2004. Manistique River Assessment, MDNR, Fisheries Division.
- MDEQ. 2004. Manistique River Watershed System Streambank Stabilization fact sheet. (The link provided was broken and has been removed) Viewed January 5, 2010.
- MDEQ. 1990. SWAS Procedure 51 Qualitative Biological and Habitat Survey Protocols for Wadable Streams and Rivers, April 24, 1990. Revised June 1991, August 1996, January 1997, and May 2002.
- MDNR. 1987. Remedial Action Plan for the Manistique River Area of Concern. Michigan Department of Natural Resources, Surface Water Quality Division. Lansing, MI.
- MDNR. 1988. Fox River Natural River Plan. Revised March 12, 2002. Alger, Luce, and Schoolcraft Counties, Fisheries Division.

- MDNR. 1994. Quality Assurance for Water and Sediment Sampling. Environmental Protection Bureau, Lansing, Michigan. Publication No. 3730-0028.
- MDNR. 2010. Section 4 Eastern Upper Peninsula, Management Unit 31 Regional State Forest Plan (In Draft).

New Deal/WPA Art in Manistique, Michigan. (*The link provided was broken and has been removed*). Viewed on January 5, 2010.

- Omernik, J.M. and Gallant. 1988. Ecoregions of the Upper Midwest States. United States Environmental Protection Agency, Environmental Research Laboratory, EPA/600/3-88/037.
- Sayles, B. 1990. A Biological Survey of the Milakokia River in the Vicinity of Inland Lime and Stone Co., Gulliver, Schoolcraft County, Michigan, August 2, 1988. MDEQ Staff Report #MI/DNR/SWQ-90/007.
- Stranahan, S. 2009. Out of Sight but not Out of Mind: Why Culverts Wreak Havoc on Trout Streams. Trout Unlimited Magazine Fall. pp 36-39.
- Suppnick, J. 1996. Cross Section Survey of the Fox River: Schoolcraft County, Michigan. July 8, 1991 and July 16, 1995. MI/DEQ/SWQ-96/008.
- Taft, W. 1989. A Biological and Water Chemistry Survey of the Tahquamenon River in the Vicinity of the Newberry WWTP, Luce County, Michigan. August 22, 1989. MI/DNR/SWQ-89/163
- Taft, W. 1990. An Intensive Biological and Water Chemistry Survey of Doe and Furlong Creeks in the Vicinity of Engadine, Michigan, Mackinac County, Michigan, May-August 1989. MI/DNR/SWQ-90/022
- Taft, W. 1992a. A Biological Survey of Carlson Creek, Luce County, Michigan, September 17, 1991. MI/DNR/SWQ-92/207.
- Taft, W. 1992b. A Biological Survey of the Fox River in the Immediate Vicinity of a Bank Stabilization Project: Schoolcraft County, Michigan. July 8, 1991.
- Taft, W. 1994a. A Biological Survey of the Driggs River in the Immediate Vicinity of a Future Bank Stabilization Project: Schoolcraft County, Michigan. July 26, 1994. MI/DNR/SWQ-94/094.
- Taft, W. 1994b. A Biological Survey of the Upper Tahquamenon River, Luce County, Michigan, July 26, 1994. MI/DNR/SWQ-94/095.
- Taft, W. 1994c. A Biological Survey of Davenport Creek, Mackinac County, Michigan. MI/DNR/SWQ-94/096.
- Taft, W. 1998. A Biological and Water Chemistry Survey of the Tahquamenon River in the Vicinity of the Newberry WWTP, Luce County, Michigan. August 22, 1989. MI/DNR/SWQ-89/163.

- Taft, W. 2000a. A Biological Survey of Selected Coastal Lake Michigan Tributaries in Schoolcraft and Mackinac Counties, June 1999. MI/DEQ/SWQ-00/003
- Taft, W. 2000b. A Biological Survey of Selected Coastal Lake Superior Tributaries in Eastern Lake Superior Tributaries in Eastern Alger, Luce, and Chippewa Counties, June 1999 MI/DEQ/SWQ-00/004.
- Taft, W. 2005. A Biological Survey of Selected Coastal Lake Michigan Tributaries in Schoolcraft and Mackinac Counties. June 2004. MI/DEQ/SWQ-00/003. MI/DEQ/WB-05/100
- USEPA. 2008. Michigan: Furlong Creek. Section 319 Nonpoint Source Success Stories. (The link provided was broken and has been removed.) Viewed January 5, 2010.
- Waybrant, J.R. and T.G. Zorn. 2008. Tahquamenon River Assessment. Michigan Department of Natural Resources, Fisheries Special Report 45, Ann Arbor.
- Wolf, S. 2005. A Biological Survey of the Manistique River Watershed Luce, Mackinac, Alger, Schoolcraft, and Delta Counties, Michigan. August 2004. MI/DEQ/WB-05/106.

| Station | Waterbody | Location | County | Storet | Lat | Long | TRS | Sample Type |
|-----------------|--------------------------|--|-------------|--------|----------|----------|-----------------|-------------|
| Cold Small | | | | | | | | |
| 1 | Syphon Creek | County Road 442 | Luce | 480038 | 46.39071 | 85.74531 | 46N/12W/12 | М, Н |
| 2 | Star Creek | County Road 454 | Manistique | 020124 | 46.37216 | 86.39305 | 46N/17W/23 | М, Н |
| 3 | Grass Creek | Old RR Grade off Sunset Landing | Alger | 020159 | 46.52231 | 86.12086 | 48N/15W/25 | М, Н |
| 4 | E B Fox River | M-77 | Schoolcraft | 770090 | 46.40607 | 85.94795 | 46N/13W/05 | М, Н |
| 5 | Kilpecker Creek | Two Track off FF2438 | Delta | 210305 | 46.11395 | 86.49214 | 43N/18W/24 | M, H |
| 6 | E B Tahquamenon River | Arbutus Truck Trail | Chippewa | 170288 | 46.3177 | 84.9527 | 45N/05W/05 | M, H |
| 7 | Mead Creek | M-77 | Schoolcraft | 770089 | 46.17484 | 85.92751 | 44N/13W/27 | M, H |
| 8 | Big Murphy Creek | County Road 437 | Schoolcraft | 770159 | 46.07281 | 86.46690 | 43N/17W/32 | M, H |
| 9 | Clear Creek | M-77 | Schoolcraft | 770108 | 46.44928 | 85.93728 | 47N/13W/21 | M, H |
| 10 | Chenev Creek | upstream Tahquamenon River confluence | Chippewa | 170191 | 46.56306 | 85.08312 | 48N/06W/08 | М. Н |
| 11 | Little Fox River | Stanley Lake Road | Schoolcraft | 770157 | 46.47622 | 86.13987 | 47N/15W/11 | M. H |
| 12 | E B Two Hearted River | County Road 414 | Luce | 480028 | 46.64188 | 85.47916 | 49N/09W/18 | M, H |
| 13 | S B Stutts Creek | Clear Lake Road | Schoolcraft | 770158 | 46.22240 | 86.41703 | 44N/17W/10 | M, H |
| Cold Medium | | | | | | | | |
| 14 | Sucker River | Grand Marais Truck Trail | Alger | 020160 | 46.66203 | 85.86893 | 49N/13W/12 | М, Н |
| 15 | E B Fox River | M-28 | Luce | 480068 | 46.33551 | 85.81582 | 46N/12W/33 | M, H |
| Warm Small | | | | | | | | |
| 16 | Milakokia River | Betty Doe Lake Road (Huntspur Road) | Schoolcraft | 490065 | 46.02832 | 85.84404 | 42N/12W/18 | M, H |
| Warm Medium | | | | | | | | |
| 17 | Creighton River | Creighton Truck Trail | Schoolcraft | 770156 | 46.24108 | 86.24185 | 44N/16W/01 | M, H |
| 18 | W B Two Hearted River | County Road 418 | Luce | 480067 | 46.56948 | 85.68017 | 48N/11W/09 | M, H |
| 19 | W B Manistique River | Hickey Truck Trail | Schoolcraft | 770155 | 46.25291 | 86.26238 | 45N/16W/35 | M, H |
| 20 | Shelldrake (Betsy) River | Betsy River Truck Trail | Chippewa | 170287 | 46.6729 | 85.2339 | 49N/07W/06 | M, H |
| 21 | W B Manistique River | off Creighton Truck Trail | Schoolcraft | 770160 | 46.22857 | 86.23506 | 44N/15W/07 | M, H |
| Warm Large | | 05 (00 | | | 10.01005 | | 4=11/4 014/10 4 | |
| 22 | Manistique River | CR 498 | Schoolcraft | 770161 | 46.24695 | 85.92453 | 45N/13W/34 | M, H |
| Warm Very Large | | M : O I D I | <u> </u> | 770400 | 40.00400 | 00 10070 | | |
| 23 | Manistique River | Merwin Creek Road | Schoolcraft | 770162 | 46.03433 | 86.12276 | 42N/15W/13 | М, Н |
| Coastal | De avrie Oren l | Liberry the Tay 2 | N4 | 400405 | 40.4040 | 05 45 45 | 4001/0704//05 | |
| 24 | Paquin Creek | Hiawatha I rail | Mackinac | 490195 | 46.1813 | 85.1545 | 43N/07VV/35 | M, H |
| 25 | Hog Island Creek | Strickler Fruck Frail | iviackinac | 490200 | 46.1228 | 85.254 | 43N/08W/13 | М, Н |

Table 1. 2009 Tahquamenon, Two-Hearted, Manistique, and Millecoquins Rivers watersheds probabilistic sampling sites.

M- Benthic macroinvertebrate community assessment

H- Habitat assessment

| Station | Waterbody | Location | County | Storet | Lat | Long | TRS | Sample Type |
|---------|------------------------|-------------------------|-------------|--------|----------|----------|------------|-------------|
| 26 | Tahquamenon River | Near Eagles Nest | Luce | 480066 | 46.42482 | 85.79771 | 47N/12W/34 | M, H |
| 27 | Indian River | off M-94 | Schoolcraft | 770085 | 46.16791 | 86.41597 | 44N/17W/34 | М, Н |
| 28 | Manistique River | at Boat Launch | Schoolcraft | 770154 | 45.97088 | 86.24275 | 41N/16W/01 | М, Н |
| 29 | Fox River | Fox River Campground | Schoolcraft | 770082 | 46.39997 | 86.02787 | 46N/14W/11 | М, Н |
| 30 | Cut River | US-2 | Mackinac | 490002 | 46.0452 | 85.1242 | 42N/07W/12 | М, Н |
| 31 | Black River | State Forest Campground | Mackinac | 490198 | 46.11758 | 85.36597 | 43N/09W/13 | М, Н |
| 32 | Davenport Creek | US-2 | Mackinac | 490059 | 46.06772 | 85.26448 | 42N/08W/02 | М, Н |
| 33 | Davenport Creek | at gas pipeline | Mackinac | 490199 | 46.09928 | 85.2595 | 43N/08W/26 | М, Н |
| 34 | Furlong Creek | Pleasant Avenue | Mackinac | 490068 | 46.13899 | 85.59196 | 43N/10W/08 | М, Н |
| 35 | Doe Creek | Raski Road | Mackinac | 490104 | 46.14193 | 85.56093 | 43N/10W/09 | M, H, WC |
| 36 | Doe Creek | Kover | Mackinac | 490202 | 46.18031 | 85.59601 | 44N/10W/30 | WC |
| 37 | Doe Creek | Mile/Pleasant | Mackinac | 490070 | 46.16545 | 85.5923 | 44N/10W/31 | WC |
| 38 | Doe Creek | M-117 | Mackinac | 490053 | 46.14761 | 85.5714 | 44N/10W/05 | WC |
| 39 | Doe Creek | Krause Road | Mackinac | 490069 | 46.13668 | 85.55039 | 43N/10W09 | WC |
| 40 | Millecoquins River | Lower Rapids | Mackinac | 490196 | 46.11568 | 85.47358 | 43N/09W/18 | M, H, WC |
| 41 | Crow River | end of Summer Trail | Mackinac | 490197 | 46.03036 | 85.61533 | 42N/11W/13 | M, H, WC |
| 42 | Tahquamenon River | upstream Newberry WWTP | Luce | 480002 | 46.3713 | 85.50972 | 46N/10W/24 | WC |
| 43 | Newberry WWTP Effluent | Oufall 001A | Luce | 480039 | 46.3723 | 85.509 | 46N/10W/24 | WC |

| Table 2. | 2009 Tahquamenon | , Two-Hearted, | Manistique, | and Millecoquins | Rivers watersh | eds targeted sa | ampling sites. |
|----------|------------------|----------------|-------------|------------------|----------------|-----------------|----------------|
|----------|------------------|----------------|-------------|------------------|----------------|-----------------|----------------|

M - Benthic macroinvertebrate community assessment

H - Habitat assessment WC - Water Chemistry

| TAXA | Syphon Creek M-442 7/29/2009 STATION 1 | Star Creek Off Star Siding Road 6/24/2009 STATION 2 | Grass Creek Railroad Grade off Sunset Landing 7/28/2009 STATION 3 | East Branch Fox River M-77 crossing 6/9/2009 STATION 4 |
|--------------------------------------|---|--|--|---|
| ANNELIDA (segmented worms) | | | | |
| Hirudinea (leeches) | 1 | | 2 | 2 |
| APTHPOPODA | 49 | 2 | 2 | 2 |
| Crustacea | | | | |
| Decapoda (crayfish) | | | 3 | |
| Isopoda (sowbugs) | | | | 10 |
| Arachnoidea | | | | |
| Hydracarina | 3 | 1 | | 1 |
| Insecta | | | | |
| Baetiscidae | | 1 | | |
| Baetidae | 35 | 1 | 14 | 76 |
| Caenidae | 4 | 7 | 5 | |
| Ephemerellidae | 1 | | | 43 |
| Ephemeridae | 4 | | | |
| Heptageniidae | | | 2 | |
| Odonata Anicontero (droconflico) | | | | |
| Anisoptera (dragonines) Aeshnidae | | 3 | 3 | |
| Cordulegastridae | | 5 | 2 | |
| Gomphidae | | 9 | 1 | |
| Zygoptera (damselflies) | | | | |
| Calopterygidae | | 3 | 1 | |
| Plecoptera (stoneflies) | | | | |
| Nemouridae | 11 | 2 | | 1 |
| Pteronarcyidae | | 2 | | |
| Hemiptera (true bugs) | | 1 | | |
| Corixidae | | 4 | | |
| Gerridae | 1 | | 8 | |
| Megaloptera | | | | |
| Corydalidae (dobson flies) | | 3 | | |
| Sialidae (alder flies) | | | 2 | |
| Prochycontrideo | 5 | | 53 | 27 |
| Glossosomatidae | 5 | 3 | 55 | 1 |
| Helicopsychidae | | 10 | | - |
| Hydropsychidae | 7 | 3 | 2 | 1 |
| Hydroptilidae | | | 4 | |
| Lepidostomatidae | 5 | 7 | | |
| Leptoceridae | 24 | 4 | 0 | 10 |
| Limnephilidae | 24 | 24 | 9 | 18 |
| Philopotamidae | 3 | 1 | | 1 |
| Phryganeidae | 5 | | 2 | 1 |
| Polycentropodidae | | 4 | | |
| Psychomyiidae | | | 1 | |
| Rhyacophilidae | | | | 1 |
| Uenoidae | | 1 | | |
| Coleoptera (beetles) | 1 | | 2 | |
| Gyrinidae (adults) | 1 4 | | 3 | |
| Hydrophilidae (total) | - | | 4 | |
| Dryopidae | | 1 | | |
| Elmidae | | 5 | | |
| Diptera (flies) | | | | |
| Athericidae | _ | 1 | _ | |
| Ceratopogonidae | 5 | 15 | 3 | 20 |
| Dividae | 51 | 15 | 94 | 50 |
| Ptychopteridae | 32 | | 1 | |
| Simuliidae | 37 | | 5 | 96 |
| Tabanidae | | | 1 | |
| Tipulidae | 1 | | 2 | 2 |
| MOLLUSCA | | | | |
| Gastropoda (snails) | | 1 | | |
| Physidae | | 1 4 | 2 | 1 |
| Viviparidae | | 7 | 2 | 1 |
| Pelecypoda (bivalves) | | | - | |
| Sphaeriidae (clams) | 48 | 3 | 29 | |
| TOTAL INDIVIDUALS | 221 | 120 | 250 | 321 |
| I O I AL INDI VIDUALS | 321 | 129 | 239 | 341 |

Table 3A. Qualitative macroinvertebrate sampling results for the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites.

| METRIC | Syphon Creek M-442 7/29/2009 STATION 1 Value Score | | Star Creek Off Star Siding Road 6/24/2009 STATION 2 Value Score | | Grass Creek Railroad Grade off Sunset Landing 7/28/2009 STATION 3 Value Score | | | East Branch Fox River M-77 crossing 6/9/2009 STATION 4 Value Score | |
|------------------------------|--|---------|---|-----------|---|-------|---------|--|---------|
| TOTAL NUMBER OF TAXA | 24 | 0 | 29 |) 1 | | 28 | 1 | 16 | 0 |
| NUMBER OF MAYFLY TAXA | 4 | 0 | | -1 | | 3 | 0 | 2 | -1 |
| NUMBER OF CADDISFLY TAXA | 6 | 1 | ç |) 1 | | 6 | 1 | 6 | 1 |
| NUMBER OF STONEFLY TAXA | 1 | 0 | 2 | 2 1 | | 0 | -1 | 1 | 0 |
| PERCENT MAYFLY COMP. | 13.71 | 0 | 6.2 | 0 0 | | 8.11 | 0 | 37.07 | 1 |
| PERCENT CADDISFLY COMP. | 14.02 | 0 | 44.1 | 9 1 | | 27.41 | 0 | 18.38 | 0 |
| PERCENT DOMINANT TAXON | 15.26 | 1 | 18.6 | 0 0 | | 36.29 | -1 | 29.91 | -1 |
| PERCENT ISOPOD, SNAIL, LEECH | 0.31 | 1 | 3.8 | 8 1 | | 1.16 | 1 | 3.43 | 1 |
| PERCENT SURF. AIR BREATHERS | 11.84 | 0 | 3.1 | 0 1 | | 6.18 | 0 | 0.00 | 1 |
| TOTAL SCORE | | 3 | | 5 | | | 1 | | 2 |
| MACROINV. COMMUNITY RATING | | ACCEPT. | | EXCELLENT | | A | ACCEPT. | | ACCEPT. |

| Table 3B. Macroinvertebrate metric evaluation of the Tahquamenon | , Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites. |
|--|---|
|--|---|

| | Kilpecker Creek two-track off Federal Forest Rd 2438 7/27/2009 | East Branch Tahquamenon River Arbutus Truck Trail 6/11/2009 | Mead Creek M-77 crossing 6/9/2009 | Big Murphy Creek County Road 437 6/22/2009 |
|---------------------------------|--|---|---|--|
| TAXA | STATION 5 | STATION 6 | STATION 7 | STATION 8 |
| ANNELIDA (segmented worms) | | | | |
| Hirudinea (leeches) | 1 | | | |
| Oligochaeta (worms) | 1 | 3 | 11 | |
| ARTHROPODA | | | | |
| Crustacea | | | | |
| Amphipoda (scuds) | | | 26 | |
| Decapoda (crayfish) | | | 1 | 5 |
| Isopoda (sowbugs) | | | | 11 |
| Arachnoidea | 2 | | | |
| Hydracarina | 2 | 1 | | |
| Enhomoroptora (maxiflios) | | | | |
| Paotisgidae | | | | 1 |
| Baetidae | 14 | | 9 | 1 |
| Caenidae | 14 | | 1 | 7 |
| Ephemerellidae | 2 | 17 | 1 | 8 |
| Heptageniidae | 1 | | 1 | 4 |
| Odonata | | | | |
| Anisoptera (dragonflies) | | | | |
| Aeshnidae | | | 5 | 2 |
| Cordulegastridae | | | 2 | |
| Gomphidae | | | 3 | 1 |
| Zygoptera (damselflies) | | | | |
| Calopterygidae | | | 4 | |
| Coenagrionidae | | | 4 | |
| Plecoptera (stoneflies) | | 50 | | |
| Leuctridae | 6 | 60 | | 1 |
| Devlidee | 20 | | | 1 |
| Pteroparcyidae | | | | 1 |
| Hemintera (true bugs) | | | | 1 |
| Gerridae | | 1 | 3 | |
| Megaloptera | | - | - | |
| Corydalidae (dobson flies) | | | | 1 |
| Sialidae (alder flies) | | 3 | | |
| Trichoptera (caddisflies) | | | | |
| Brachycentridae | 24 | | | 33 |
| Helicopsychidae | | | 3 | |
| Hydropsychidae | | | | 56 |
| Lepidostomatidae | 5 | 61 | | |
| Leptoceridae | | _ | 1 | 1 |
| Limnephilidae | 15 | 2 | 40 | 2 |
| Molannidae Dhiles etcaside e | 1 | | | |
| Philopotamidae | 2 | | 1 | |
| Lenoidae | 1 | | 1 | |
| Coleoptera (beetles) | 1 | | | |
| Dytiscidae (total) | 1 | | | 1 |
| Gyrinidae (adults) | | | 7 | - |
| Hydrophilidae (total) | | 1 | | |
| Elmidae | | | 18 | 3 |
| Diptera (flies) | | | | |
| Athericidae | 2 | | | |
| Ceratopogonidae | 2 | 3 | 4 | |
| Chironomidae | 100 | 80 | 113 | 14 |
| Ptychopteridae | 1 | 2 | | |
| Simuliidae | 6 | 8 | 46 | 2 |
| Stratiomyidae | | | | 17 |
| Tabanidae | 1 | 1 | 2 | |
| Tipulidae | | 2 | 2 | 1 |
| MOLLUSCA Gestropode (speils) | | | | |
| Hydrobiidae | | | А | |
| Lymnaeidae | | | 4 | 5 |
| Physidae | 2 | | | 5 |
| Pelecypoda (bivalves) | 2 | | | |
| Sphaeriidae (clams) | 54 | 3 | 1 | 5 |
| Unionidae (mussels) | | - | 1 | - |
| · | | | | |
| TOTAL INDIVIDUALS | 270 | 248 | 312 | 183 |

| Table 3D con t. Macromyertebrate met | in evaluation of the ranguament | 1011, I | wo-meante | u, mainsi | ique, and wine | coquins Kiv | er probabilist | ic sampning | sites. | | |
|--------------------------------------|---------------------------------|---------|-------------------------------|-----------|------------------|-------------|----------------|------------------|-----------|----------|----|
| | Kilpecker Creek | | East Branch Tahquamenon River | | | Mead Creek | | Big Murphy Creek | | | |
| | two-track off Federal Fores | t Rd 2 | 438 | А | rbutus Truck Tra | ail | M-77 cro | ssing | County R | load 437 | |
| | 7/27/2009 | | | | 6/11/2009 | | 6/9/20 | 09 | 6/22/2009 | | |
| | STATION 5 | | | | STATION 6 | | STATIC | ON 7 | STAT | ION 8 | |
| METRIC | Value | | Score | Value | | Score | Value | Score | Value | Score | |
| TOTAL NUMBER OF TAXA | | 23 | 0 | | 16 | 0 | 26 | 0 | 24 | | 0 |
| NUMBER OF MAYFLY TAXA | | 3 | 0 | | 1 | -1 | 4 | 0 | 4 | | 0 |
| NUMBER OF CADDISFLY TAXA | | 6 | 1 | | 2 | -1 | 4 | 0 | 4 | | 0 |
| NUMBER OF STONEFLY TAXA | | 2 | 1 | | 1 | 0 | 0 | -1 | 3 | | 1 |
| PERCENT MAYFLY COMP. | 6 | 5.30 | 0 | | 6.85 | 0 | 3.85 | 0 | 10.93 | | 0 |
| PERCENT CADDISFLY COMP. | 17 | 7.78 | 0 | | 25.40 | 0 | 14.42 | 0 | 50.27 | | 1 |
| PERCENT DOMINANT TAXON | 37 | .04 | -1 | | 32.26 | -1 | 36.22 | -1 | 30.60 | | -1 |
| PERCENT ISOPOD, SNAIL, LEECH | 1 | .11 | 1 | | 0.00 | 1 | 1.28 | 1 | 8.74 | | 0 |
| PERCENT SURF. AIR BREATHERS | 0 |).74 | 1 | | 1.61 | 1 | 3.21 | 1 | 9.84 | | 0 |
| TOTAL SCORE | | | 3 | | | -1 | | 0 | | | 1 |
| MACROINV. COMMUNITY RATING | | А | CCEPT. | | 1 | ACCEPT. | | ACCEPT. | | ACCEPT. | |

Table 3B con't. Macroinvertebrate metric evaluation of the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites.

| | Clear Creek upstream M-77 6/9/2009 | Cheney Creek u/s Tahq. River confluence 6/10/2009 | Little Fox River Stanley Lake Road 6/23/2009 | East Branch Two Hearted River County Road 414 7/30/2009 |
|--------------------------------------|--|---|--|---|
| TAXA | STATION 9 | STATION 10 | STATION 11 | STATION 12 |
| ANNELIDA (segmented worms) | | | | |
| Hirudinea (leeches) | | | 1 | |
| Oligochaeta (worms) | 4 | 3 | 2 | 39 |
| ARTHROPODA | | | | |
| Crustacea | | | | 2 |
| Amphipoda (scuds) | 1 | | 1 | 2 |
| Isopoda (sowbugs) | | | 92 | |
| Arachnoidea | | | - | |
| Hydracarina | 1 | | | 10 |
| Insecta | | | | |
| Ephemeroptera (mayflies) | | | | |
| Baetiscidae | | 20 | 1 | 1 |
| Baetidae | 116 | 20 | 11 | 39 |
| Enhemerellidae | 5 | | 5 | 1 |
| Ephemeridae | 5 | | 2 | 1 |
| Heptageniidae | | 8 | 7 | 2 |
| Odonata | | | | |
| Anisoptera (dragonflies) | | | | |
| Aeshnidae | | 1 | 1 | 1 |
| Cordulegastridae | | | 3 | 1 |
| Gomphidae Zygoptera (damselflies) | | | 1 | |
| Calontervgidae | | | 5 | |
| Plecoptera (stoneflies) | | | 5 | |
| Leuctridae | 8 | 10 | | |
| Nemouridae | 18 | 3 | | |
| Perlidae | | | 2 | |
| Perlodidae | | 3 | | _ |
| Pteronarcyidae | | | | 9 |
| Corivideo | | | 1 | 1 |
| Gerridae | | 1 | 1 | 1 |
| Veliidae | | • | 1 | 5 |
| Megaloptera | | | | |
| Sialidae (alder flies) | | | 1 | |
| Trichoptera (caddisflies) | | | _ | |
| Brachycentridae | 61 | | 2 | 1 |
| Hencopsychidae | 4 | 1 | 1 39 | |
| Lepidostomatidae | - | 6 | 57 | 6 |
| Leptoceridae | | - | 7 | - |
| Limnephilidae | 52 | 22 | 7 | 14 |
| Molannidae | | | 1 | |
| Philopotamidae | 1 | 1 | _ | |
| Phryganeidae | | | 2 | |
| Venoidae | | | 1 | |
| Coleoptera (beetles) | | | - | |
| Dytiscidae (total) | 2 | 2 | | 1 |
| Gyrinidae (adults) | | | | 1 |
| Hydrophilidae (total) | | 1 | | 2 |
| Elmidae | | 1 | 2 | |
| Diptera (flies) | | | | |
| Athericidae | | 1 | 5 | 1 |
| Chironomidae | 10 | 100 | 27 | 2 64 |
| Simuliidae | 69 | 70 | 6 | 7 |
| Tabanidae | ~~ | 1 | 3 | 3 |
| Tipulidae | 1 | 1 | | |
| MOLLUSCA | | | | |
| Gastropoda (snails) | | | | |
| Hydrobiidae Physidae | | | 1 | 1 |
| r nysiuae Pelecypoda (biyalves) | | | 1 | 1 |
| Sphaeriidae (clams) | 16 | | 6 | 22 |
| | | | | |
| TOTAL INDIVIDUALS | 369 | 256 | 252 | 243 |

Table 3A con't. Qualitative macroinvertebrate sampling results for the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites.

| METRIC | Clear C upstream 6/9/20 STATIC Value | reek M-77 09 DN 9 Score | Cheney Cr u/s Tahq. River cr 6/10/200 STATION Value | eek onfluence 9 10 Score | Value | Little Fox Riv Stanley Lake R 6/23/2009 STATION 1 | er oad I Score | East Branch Two County R 7/30/2 STATI Value |) Hearted River toad 414 2009 ON 12 Score |
|------------------------------|--|-------------------------------------|---|--------------------------------------|-------|--|-------------------------|---|---|
| TOTAL NUMBER OF TAXA | 16 | 0 | 20 | 0 | | 34 | 1 | 27 | 0 |
| NUMBER OF MAYFLY TAXA | 2 | õ | 2 | -1 | | 5 | 1 | 6 | 1 |
| NUMBER OF CADDISFLY TAXA | 4 | 0 | 4 | 0 | | 9 | 1 | 3 | 0 |
| NUMBER OF STONEFLY TAXA | 2 | 1 | 3 | 1 | | 1 | 0 | 1 | 0 |
| PERCENT MAYFLY COMP. | 32.79 | 1 | 10.94 | 0 | | 10.32 | 0 | 21.4 | 0 0 |
| PERCENT CADDISFLY COMP. | 31.98 | 1 | 11.72 | 0 | | 25.40 | 0 | 8.6 | 4 0 |
| PERCENT DOMINANT TAXON | 31.44 | -1 | 39.06 | -1 | | 36.51 | -1 | 26.34 | 4 0 |
| PERCENT ISOPOD, SNAIL, LEECH | 0.00 | 1 | 0.00 | 1 | | 37.70 | -1 | 0.4 | 1 1 |
| PERCENT SURF. AIR BREATHERS | 0.54 | 1 | 1.56 | 1 | | 0.79 | 1 | 3.2 | 9 1 |
| TOTAL SCORE | | 4 | | 1 | | | 2 | | 3 |
| MACROINV. COMMUNITY RATING | | ACCEPT. | | ACCEPT. | | 1 | ACCEPT. | | ACCEPT. |

Table 3B con't. Macroinvertebrate metric evaluation of the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites.

| | South Branch Stutts Creek Clear Lake Road 6/25/2009 | Sucker River Grand Marais Truck Trail 7/29/2009 | Creighton River Creighton Truck Trail 6/24/2009 | Milakokia River Betty Doe Lake Rd 6/9/2009 |
|--------------------------------------|---|---|---|--|
| TAXA | STATION 13 | STATION 14 | STATION 16 | STATION 17 |
| PORIFERA (sponges) | | | | 1 |
| ANNELIDA (segmented worms) | | | | |
| Hirudinea (leeches) | 1 | 1 | | |
| Oligochaeta (worms) | 3 | 6 | 2 | 3 |
| ARTHROPODA | | | | |
| Crustacea | | | | 10 |
| Amphipoda (scuds) | 2 | 2 | 2 | 10 |
| Decapoda (crayfish) | | 2 | 3 | 1 |
| Hudrocerine | | | 2 | |
| Insecta | | | 5 | |
| Enhemeroptera (mayflies) | | | | |
| Baetiscidae | 2 | | 5 | |
| Baetidae | 16 | 35 | 10 | 23 |
| Caenidae | 1 | 3 | | 5 |
| Ephemerellidae | | 4 | 24 | |
| Ephemeridae | 15 | | 1 | |
| Heptageniidae | 9 | 7 | | 21 |
| Leptophlebiidae | 3 | | | 18 |
| Odonata | | | | |
| Anisoptera (dragonflies) | | | - | c |
| Aeshnidae | | 4 | 5 | 8 |
| Cordulegastridae | 1 | 2 | 1 | 20 |
| Gomphidae Zugoptore (domostifice) | 1 | 11 | 3 | 50 |
| Caloptera (danisennes) | | 2 | 2 | 1 |
| Plecontera (stoneflies) | | 2 | <u>2</u> | 1 |
| Leuctridae | | | 3 | |
| Nemouridae | | | 5 | 1 |
| Perlidae | | 2 | 4 | 4 |
| Perlodidae | | | 3 | |
| Pteronarcyidae | 2 | 24 | 3 | |
| Hemiptera (true bugs) | | | | |
| Corixidae | 4 | | 11 | |
| Gerridae | | 4 | 3 | 1 |
| Megaloptera | | | | |
| Corydalidae (dobson flies) | 2 | | 1 | 17 |
| Sialidae (alder flies) | 10 | | | |
| Development (caddisflies) | | 2 | | |
| Haliaanayahidaa | | 3 | | 2 |
| Hydronsychidae | 1 | 15 | 11 | 5 |
| Lepidostomatidae | 15 | 5 | 1 | 5 |
| Leptoceridae | 3 | 5 | 1 | |
| Limnephilidae | 52 | 1 | 20 | 25 |
| Molannidae | 2 | | 1 | 1 |
| Philopotamidae | | | 3 | |
| Polycentropodidae | 4 | | 2 | |
| Uenoidae | 2 | 1 | | 1 |
| Coleoptera (beetles) | | | | |
| Dytiscidae (total) | 1 | | 3 | |
| Haliplidae (adults) | 1 | | 1 | |
| Hydrophilidae (total) | 1 | - | 3 | 4 |
| Dryopidae | | 2 | 1 | c |
| Elmidae | 4 | 4 | | 8 |
| Diptera (Illes) | | 4 | 2 | |
| Ceretopogonideo | n | 4 | ے ۱ | 1 |
| Chironomidae | 12 | 5 72 | 1 | 1 6 |
| Ptychopteridae | 12 | 12 | 1 | 0 |
| Simuliidae | 2 | 13 | | 2 |
| Tabanidae | 4 | 2 | 2 | - |
| Tipulidae | | 10 | | |
| MOLLUSCA | | | | |
| Gastropoda (snails) | | | | |
| Ancylidae (limpets) | | | | 1 |
| Physidae | 2 | 2 | | |
| Pelecypoda (bivalves) | | | | |
| Sphaeriidae (clams) | 61 | 4 | 94 | 75 |
| TOTAL NIDBUDUALS | 242 | 240 | 224 | 276 |
| I UTAL INDIVIDUALS | 242 | 248 | 234 | 276 |

| Table 3A con't. Qualitative macroinvertebrate sampling results for the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River proba | bilistic |
|---|----------|
| sampling sites. | |

| Table 3B con't. Macroinvertebrate metric evaluation of the Tahquameno | n, Two-Hearted, Manistique, and Millecoquins River a probabilistic sampling sites. |
|---|--|
|---|--|

| | South Branch S Clear Lake 6/25/20 STATIO | South Branch Stutts Creek Sucker River Clear Lake Road Grand Marais Truck Trail 6/25/2009 7/29/2009 STATION 13 STATION 14 | | Creighton River Creighton Truck Trail 6/24/2009 STATION 16 | | Milakokia River Betty Doe Lake Rd 6/9/2009 STATION 17 | | |
|------------------------------|---|--|-------|---|-------|--|-------|---------|
| METRIC | Value | Score | Value | Score | Value | Score | Value | Score |
| TOTAL NUMBER OF TAXA | 33 | 1 | 29 | 1 | 34 | 1 | 27 | 0 |
| NUMBER OF MAYFLY TAXA | 6 | 1 | 4 | 0 | 4 | 0 | 4 | 0 |
| NUMBER OF CADDISFLY TAXA | 7 | 1 | 5 | 0 | 7 | 1 | 5 | 0 |
| NUMBER OF STONEFLY TAXA | 1 | 0 | 2 | 1 | 4 | 1 | 2 | 1 |
| PERCENT MAYFLY COMP. | 19.01 | 0 | 19.76 | 0 | 17.09 | 0 | 24.28 | 1 |
| PERCENT CADDISFLY COMP. | 32.64 | 1 | 10.08 | 0 | 16.67 | 0 | 12.68 | 0 |
| PERCENT DOMINANT TAXON | 25.21 | 0 | 29.03 | -1 | 40.17 | -1 | 27.17 | 0 |
| PERCENT ISOPOD, SNAIL, LEECH | 1.24 | 1 | 1.21 | 1 | 0.00 | 1 | 0.36 | 1 |
| PERCENT SURF. AIR BREATHERS | 3.31 | 1 | 1.61 | 1 | 8.97 | 0 | 1.81 | 1 |
| TOTAL SCORE | | 6 | | 3 | | 3 | | 4 |
| MACROINV. COMMUNITY RATING | | EXCELLENT | | ACCEPT. | | ACCEPT. | 1 | ACCEPT. |

| | West Branch Two Hearted River County Road 418 | West Branch Manistique River Hickey Truck Trail | Shelldrake River Betsy River Truck Trail | West Branch Manistique River off Crieghton Truck Trail |
|--|--|--|---|---|
| TAXA | 7/29/2009 STATION 18 | 7/27/2009 STATION 19 | 6/10/2009 STATION 20 | 6/24/2009 STATION 21 |
| mm | SIMIONIO | Similar () | 5111101120 | SIMION2I |
| ANNELIDA (segmented worms Hirudinea (leeches) | S) 1 | 1 | | |
| Oligochaeta (worms) | 13 | 1 | 4 | 23 |
| ARTHROPODA | 10 | - | | 23 |
| Crustacea | | | | |
| Amphipoda (scuds) | 16 | 15 | 1 | 2 |
| Decapoda (crayfish) | | 53 | | 1 |
| Arachnoidea | | | | |
| Hydracarina | 3 | | 1 | |
| Insecta | | | | |
| Ephemeroptera (mayflies) | | 2 | | |
| Baetiscidae | 11 | 12 | 12 | 4 |
| Caenidae | 11 | 12 | 43 | 1 |
| Enhemerellidae | 3 | 7 | 4 | 19 |
| Ephemeridae | 14 | 1 | 7 | 19 |
| Heptageniidae | 3 | 5 | | 1 |
| Leptophlebiidae | 1 | - | | - |
| Odonata | | | | |
| Anisoptera (dragonflies) | | | | |
| Aeshnidae | | 10 | 1 | 6 |
| Cordulegastridae | 1 | 2 | | 2 |
| Libellulidae | | | 1 | |
| Zygoptera (damselflies) | | | | |
| Calopterygidae | | 20 | 2 | 2 |
| Coenagrionidae | | | 1 | |
| Plecoptera (stoneflies) | | | | |
| Leuctridae | | 1 | | 4 |
| Perlidae | | 1 | | 1 |
| Perlodidae | | | | 3 |
| Pteronarcyidae | 2 | 2 | | 5 |
| Hemiptera (true bugs) | 10 | 2 | | |
| Conixidae | 10 | 2 | | |
| Valiidaa | 2 | 1 | 2 | 1 |
| Magaloptara | | 5 | 2 | 1 |
| Corvdalidae (dobson flies) | 1 | 3 | | |
| Sialidae (alder flies) | 1 | 5 | 1 | 3 |
| Trichoptera (caddisflies) | | | 1 | 5 |
| Brachycentridae | 18 | 8 | | |
| Helicopsychidae | 10 | 2 | | |
| Hydropsychidae | | 11 | | 21 |
| Hydroptilidae | 1 | | | |
| Lepidostomatidae | | | 2 | 1 |
| Leptoceridae | | 1 | | 5 |
| Limnephilidae | 5 | 10 | 14 | 9 |
| Molannidae | 2 | | | |
| Philopotamidae | | | | 1 |
| Coleoptera (beetles) | | | | |
| Dytiscidae (total) | 2 | 5 | | |
| Gyrinidae (adults) | | 1 | 2 | |
| Haliplidae (adults) | | 1 | 1 | |
| Hydrophilidae (total) | 1 | | 1 | |
| Dryopidae | | 6 | | 4 |
| Eimidae | 1 | 1 | | |
| Gyrinidae (larvae) | | 1 | | |
| Athericidae | А | | | 1 |
| Ceratonogonidae | 4 | 1 | n | i |
| Chironomidae | 28 | 20 | 2 30 | 41 |
| Dixidae | 20 | 1 | 50 | 41 |
| Ptychopteridae | | 1 | | 1 |
| Simuliidae | 14 | 6 | 163 | 4 |
| Tabanidae | 7 | 4 | - 30 | 2 |
| Tipulidae | | 3 | | 2 |
| MOLLUSCA | | - | | |
| Gastropoda (snails) | | | | |
| Ancylidae (limpets) | | 1 | | |
| Physidae | 12 | 7 | 3 | |
| Planorbidae | | | 1 | |
| Pelecypoda (bivalves) | | | | |
| Sphaeriidae (clams) | 86 | 9 | 45 | 71 |
| | | | | |
| TOTAL INDIVIDUALS | 266 | 246 | 325 | 241 |

Table 3A con't. Qualitative macroinvertebrate sampling results for the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites.

| able 5D con t. Macromycritebrate metric evaluation of the randuallenon, rwo-ricarted, Mainstique, and Minecolumis Kiver probabilistic sampling sites. | | | | | | | | | |
|---|--|----------|------------------------------|-------------|-------------------|---------|------------------------------|---------|--|
| | West Branch Two Hearted River County Road 418 | | West Branch Manistique River | | Shelldrake Rive | r | West Branch Manistique River | | |
| | | | Hickey ' | Truck Trail | Betsy River Truck | Frail | off Crieghton Truck Trail | | |
| | 7 | /29/2009 | 7/2 | 7/2009 | 6/10/2009 | | 6/24/200 | 9 | |
| | ST | ATION 18 | STAT | TON 19 | STATION 20 | | STATION | 21 | |
| METRIC | Value | Score | Value | Score | Value | Score | Value | Score | |
| TOTAL NUMBER OF TAXA | 29 | 1 | 39 | 1 | 22 | 0 | 29 | 1 | |
| NUMBER OF MAYFLY TAXA | 6 | 1 | 6 | 1 | 2 | -1 | 4 | 0 | |
| NUMBER OF CADDISFLY TAXA | 4 | 0 | 5 | 0 | 2 | -1 | 5 | 0 | |
| NUMBER OF STONEFLY TAXA | 1 | 0 | 3 | 1 | 0 | -1 | 4 | 1 | |
| PERCENT MAYFLY COMP. | 13.16 | 0 | 12.20 | 0 | 14.46 | 0 | 10.37 | 0 | |
| PERCENT CADDISFLY COMP. | 9.77 | 0 | 13.01 | 0 | 4.92 | 0 | 15.35 | 0 | |
| PERCENT DOMINANT TAXON | 32.33 | -1 | 21.54 | 0 | 50.15 | -1 | 29.46 | -1 | |
| PERCENT ISOPOD, SNAIL, LEECH | 4.89 | 0 | 3.66 | 1 | 1.23 | 1 | 0.00 | 1 | |
| PERCENT SURF. AIR BREATHERS | 5.64 | 0 | 6.10 | 0 | 1.85 | 1 | 0.83 | 1 | |
| TOTAL SCORE | | 1 | | 4 | | -2 | | 3 | |
| MACROINV. COMMUNITY RATING | | ACCEPT. | | ACCEPT. | | ACCEPT. | | ACCEPT. | |

Table 3B con't. Macroinvertebrate metric evaluation of the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites.

Table 3A con't. Qualitative macroinvertebrate sampling results for the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites.

| ТАХА | Paquin Creek Hiawatha Trail 6/7/2009 STATION 24 | Hog Island Creek Strickler Truck Trail 6/10/2009 STATION 25 |
|----------------------------|--|--|
| ANNELIDA (segmented worms) | | |
| Oligochaeta (worms) | | 2 |
| | | 2 |
| Crustacea | | |
| Amphipoda (scuds) | | 54 |
| Arachnoidea | | 54 |
| Hydracarina | 1 | |
| Insecta | 1 | |
| Ephemeroptera (mayflies) | | |
| Baetidae | 103 | 26 |
| Ephemerellidae | | |
| Heptageniidae | | 4 |
| Leptophlebiidae | 1 | |
| Odonata | | |
| Anisoptera (dragonflies) | | |
| Aeshnidae | 1 | |
| Cordulegastridae | 2 | |
| Libellulidae | | 1 |
| Plecoptera (stoneflies) | | |
| Leuctridae | 3 | |
| Nemouridae | 16 | 1 |
| Perlodidae | 1 | |
| Hemiptera (true bugs) | | |
| Gerridae | 1 | 1 |
| Trichoptera (caddisflies) | | |
| Brachycentridae | | 2 |
| Hydropsychidae | 3 | 2 |
| Lepidostomatidae | 2 | 13 |
| Limnephilidae | 42 | 50 |
| Philopotamidae | | 6 |
| Uenoidae | 11 | 3 |
| Lepidoptera (moths) | | |
| Noctuidae | 1 | |
| Coleoptera (beetles) | | |
| Dytiscidae (total) | 12 | 3 |
| Hydrophilidae (total) | | 2 |
| Diptera (flies) | | |
| Ceratopogonidae | 3 | |
| Chironomidae | 33 | 11 |
| Ptychopteridae | 1 | 1 |
| Simuliidae | 52 | 58 |
| Tabanidae | 3 | 1 |
| Tipulidae | 1 | 1 |
| MOLLUSCA | | |
| Gastropoda (snails) | | |
| Planorbidae | 1 | |
| TOTAL INDIVIDUALS | 294 | 245 |

29

Table 3B con't. Macroinvertebrate metric evaluation of the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites.

| | | Paquin Creek Hiawatha Trail 6/7/2009 STATION 24 | | Hog Island Creek Strickler Truck Trail 6/10/2009 STATION 25 | | |
|------------------------------|-------|--|---------|--|-------|---------|
| METRIC | Value | | Score | | Value | Score |
| TOTAL NUMBER OF TAXA | | 22 | | 0 | 21 | 0 |
| NUMBER OF MAYFLY TAXA | | 2 | | -1 | 3 | 0 |
| NUMBER OF CADDISFLY TAXA | | 4 | | 0 | 6 | 1 |
| NUMBER OF STONEFLY TAXA | | 3 | | 1 | 1 | 0 |
| PERCENT MAYFLY COMP. | | 35.37 | | 1 | 13.47 | 0 |
| PERCENT CADDISFLY COMP. | | 19.73 | | 0 | 31.02 | 1 |
| PERCENT DOMINANT TAXON | | 35.03 | | -1 | 23.67 | 0 |
| PERCENT ISOPOD, SNAIL, LEECH | | 0.34 | | 1 | 0.00 | 1 |
| PERCENT SURF. AIR BREATHERS | | 4.76 | | 1 | 2.86 | 1 |
| TOTAL SCORE | | | | 2 | | 4 |
| MACROINV. COMMUNITY RATING | | | ACCEPT. | | 1 | ACCEPT. |

| Table 3A con't. Qualitative macroinvertebrate sampling results for the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River | |
|---|--|
| targeted sampling sites. | |

| | Tahquamenon River near Eagles Nest 7/30/2009 | Indian River 2-track east off 94 7/27/2009 | Manistique River Manistique boat launch 7/28/2009 | Fox River Fox River campground 7/28/2009 |
|---|--|--|---|--|
| TAXA | STATION 26 | STATION 27 | STATION 28 | STATION 29 |
| ANNELIDA (segmented worms) | ~ | 20 | | 11 |
| Oligochaeta (worms) ARTHROPODA | 5 | 28 | | 11 |
| Crustacea | | | | |
| Amphipoda (scuds) | 19 | 2 | 80 | 1 |
| Isopoda (sowbugs) | | 5 | 2 | 36 |
| Arachnoidea | | | | |
| Hydracarina | | | 1 | |
| Ephemeroptera (mayflies) | | | | |
| Baetiscidae | 1 | 1 | | 1 |
| Baetidae | 16 | 10 | 6 | 26 23 |
| Ephemerellidae | 2 | 1 | 1 | 11 |
| Ephemeridae | | 1 | | 2 |
| Heptageniidae | 3 | 2 | 1 | 3 |
| Siphlonuridae | | 2 | 1 | |
| Tricorythidae | 59 | | | |
| Odonata Anisoptera (dragonflies) | | | | |
| Aeshnidae | 1 | 1 | 3 | |
| Cordulegastridae | _ | | | 1 |
| Gomphidae Macromiidae | 3 | 1 | 1 | 1 |
| Zygoptera (damselflies) | | | 1 | |
| Calopterygidae | | 2 | 3 | |
| Coenagrionidae Plecontera (stoneflies) | | | 2 | |
| Leuctridae | 1 | | | |
| Nemouridae | | _ | | 3 |
| Perlidae Perlodidae | 6 | 3 | 1 | 5 |
| Pteronarcyidae | 2 | 10 | | 13 |
| Hemiptera (true bugs) | _ | | | |
| Corixidae Gerridae | 2 | 1 | 1 | 1 2 |
| Mesoveliidae | 1 | 1 | 2 | 2 |
| Veliidae | | 1 | | |
| Megaloptera Corvdalidae (dobson flies) | | 1 | | |
| Trichoptera (caddisflies) | | | | |
| Brachycentridae | 28 | 50 | 4 | 4 |
| Glossosomatidae Helicopsychidae | 9 | | 3 | 13 |
| Hydropsychidae | 18 | 18 | 5 | |
| Lepidostomatidae | 5 | 1 | | |
| Leptoceridae | 2 | 4 | 1 | |
| Molannidae | 1 | | - | |
| Philopotamidae | 36 | 16 | 2 | |
| Uenoidae | 1 | | 2 | |
| Lepidoptera (moths) | | | | |
| Pyralidae | | | 1 | |
| Dytiscidae (total) | 1 | 1 | | 2 |
| Gyrinidae (adults) | | | | 1 |
| Haliplidae (adults) | 6 | 5 | 1 | 4 |
| Ptilodactylidae (larvae) | 0 | 5 | 4 | 4 |
| Diptera (flies) | | | | |
| Athericidae | 4 | 3 | | 11 |
| Chironomidae | 77 | 18 | 2 | 35 |
| Ephydridae | | | | 1 |
| Ptychopteridae Simuliidae | 3 | 2 | | 1 |
| Tabanidae | 1 | 1 | 1 | 14 |
| Tipulidae | 2 | | | 3 |
| MOLLUSCA Gastropoda (spails) | | | | |
| Ancylidae (limpets) | | 1 | 1 | |
| Hydrobiidae | | | 30 | |
| Lymnaeidae Physidae | 1 | 1 | 42 | |
| Viviparidae | 1 | 2 | 43 | |
| Pelecypoda (bivalves) | 2 | 4 | Ō | 22 |
| spnaernuae (ciams) | 3 | 61 | 9 | 32 |
| TOTAL INDIVIDUALS | 338 | 262 | 253 | 263 |

Table 3B con't. Macroinvertebrate metric evaluation of the Tahquamenon, Two-Hearted, Manistique, and Millecoquins River targeted sampling sites.

| | Tahquamenon River near Eagles Nest 7/30/2009 STATION 26 | | Indian River 2-track east off 94 7/27/2009 STATION 27 | | Manistique River Manistique boat launch 7/28/2009 STATION 28 | | Fox River Fox River campground 7/28/2009 STATION 29 | |
|------------------------------|--|-----------|--|----------|---|---------|--|---------|
| METRIC | Value | Score | Value | Score | Value | Score | Value | Score |
| TOTAL NUMBER OF TAXA | 34 | 1 | 33 | 1 | 32 | 1 | 30 | 1 |
| NUMBER OF MAYFLY TAXA | 6 | 1 | 7 | 1 | 5 | 1 | 6 | 1 |
| NUMBER OF CADDISFLY TAXA | 8 | 1 | 6 | 1 | 5 | 0 | 2 | -1 |
| NUMBER OF STONEFLY TAXA | 4 | 1 | 2 | 1 | 1 | 0 | 3 | 1 |
| PERCENT MAYFLY COMP. | 26.63 | 1 | 8.78 | 0 | 3.95 | 0 | 25.10 | 1 |
| PERCENT CADDISFLY COMP. | 29.59 | 1 | 35.50 | 1 | 4.35 | 0 | 6.46 | 0 |
| PERCENT DOMINANT TAXON | 22.78 | 0 | 23.28 | 0 | 31.62 | -1 | 13.69 | 1 |
| PERCENT ISOPOD, SNAIL, LEECH | 0.30 | 1 | 1.53 | 1 | 46.64 | -1 | 13.69 | -1 |
| PERCENT SURF. AIR BREATHERS | 1.18 | 1 | 1.15 | 1 | 1.58 | 1 | 2.66 | 1 |
| TOTAL SCORE | | 8 | | 7 | | 1 | | 4 |
| MACROINV. COMMUNITY RATING | | EXCELLENT | r i | EXCELLEN | Г | ACCEPT. | | ACCEPT. |

| Table 3A con't. Qualitative macroinvertebrate sampling results for the | Tahquamenon, Two-Hearted, Manistique, and Millecoquins River |
|--|--|
| targeted sampling sites. | |

| | Millecoquins River Lower Rapids 6/8/2009 | Furlong Creek Pleasant Ave 6/7/2009 | Davenport Creek US-2 6/7/2009 | Cut River US-2 6/7/2009 |
|----------------------------|--|---|-------------------------------------|-------------------------------|
| TAXA | STATION 30 | STATION 31 | STATION 32 | STATION 33 |
| ANNELIDA (segmented worms) | | | | |
| Oligochaeta (worms) | 6 | 4 | 2 | 4 |
| Crustacea | | | | |
| Amphipoda (scuds) | | 1 | 52 | |
| Decapoda (crayfish) | 1 | 4 | | |
| Isopoda (sowbugs) | | | 2 | 1 |
| Insecta | | | | |
| Ephemeroptera (mayflies) | 10 | | 10 | |
| Baetidae | 13 | 28 | 13 | 98 |
| Enhemerellidae | 26 | 5 | 17 | 4 |
| Heptageniidae | 25 | 69 | 17 | 4 |
| Isonychiidae | 6 | | | |
| Leptophlebiidae | | 44 | | |
| Odonata | | | | |
| Anisoptera (dragonflies) | | _ | | |
| Aeshnidae | 1 | 5 | | |
| Gomphidae | 1 | 5 | | |
| Zygoptera (damselflies) | 2 | | | |
| Calopterygidae | 2 | | | |
| Plecoptera (stoneflies) | | | | |
| Leuctridae | | | 1 | 3 |
| Nemouridae | | | 4 | |
| Perlidae | 2 | 5 | | |
| Periodidae | 2 | | | 4 |
| Gerridae | 2 | 1 | 1 | |
| Veliidae | 2 | 1 | 1 | 1 |
| Megaloptera | | | | - |
| Corydalidae (dobson flies) | 1 | 11 | | |
| Trichoptera (caddisflies) | | | | |
| Brachycentridae | | | 29 | |
| Glossosomatidae | 115 | 2 | | 3 |
| Helicopsychidae | 10 | 4 | 10 | 4 |
| Hydroptilidae | 12 | 9 | 17 | 4 |
| Lepidostomatidae | | , | 5 | 7 |
| Leptoceridae | 3 | 1 | | |
| Limnephilidae | 1 | 52 | 25 | 7 |
| Philopotamidae | | 2 | | 5 |
| Polycentropodidae | 1 | | | 2 |
| Knyacophilidae | | 1 | 1 | 1 |
| Coleoptera (beetles) | | 1 | | 15 |
| Dytiscidae (total) | | | 1 | |
| Hydrophilidae (total) | | | 1 | |
| Dryopidae | | 2 | 1 | |
| Elmidae | 25 | 29 | | |
| Diptera (flies) | , | | | |
| Athericidae | 6 | 1 | | |
| Chironomidae | 2 6 | 5 28 | 87 | 63 |
| Simuliidae | 1 | 12 | 10 | 85 |
| Tabanidae | | 1 | 1 | |
| Tipulidae | 1 | 2 | 2 | |
| MOLLUSCA | | | | |
| Gastropoda (snails) | | | | |
| Hydrobiidae | 117 | 1 | | |
| Lymnaeidae Physidae | 116 | 2 | | |
| Pelecypoda (biyalves) | | 2 | | |
| Sphaeriidae (clams) | 13 | 21 | | |
| Unionidae (mussels) | 1 | 1 | | |
| | | | | |
| TOTAL INDIVIDUALS | 411 | 359 | 275 | 309 |

| Table 3B con't. | Macroinvertebrate metric evaluation of the Tahquameno | n, Two-Hearted, Manistique | , and Millecoquins River targeted | |
|-----------------|---|----------------------------|-----------------------------------|--|
| sampling sites. | | | | |

| | Millecoquins River Lower Rapids 6/8/2009 STATION 30 | | Furlong Creek Pleasant Ave 6/7/2009 STATION 31 | | Davenport Creek US-2 6/7/2009 STATION 32 | | Cut River US-2 6/7/2009 STATION 33 | |
|------------------------------|--|---------|---|----------|---|---------|---|---------|
| METRIC | Value | Score | Value | Score | Value | Score | Value | Score |
| TOTAL NUMBER OF TAXA | 30 | 1 | 32 | 1 | 21 | 0 | 18 | 0 |
| NUMBER OF MAYFLY TAXA | 5 | 1 | 4 | 0 | 2 | -1 | 3 | 0 |
| NUMBER OF CADDISFLY TAXA | 6 | 1 | 8 | 1 | 5 | 0 | 8 | 1 |
| NUMBER OF STONEFLY TAXA | 2 | 1 | 1 | 0 | 2 | 1 | 2 | 1 |
| PERCENT MAYFLY COMP. | 17.52 | 0 | 40.67 | 1 | 10.91 | 0 | 34.30 | 1 |
| PERCENT CADDISFLY COMP. | 36.01 | 1 | 20.61 | 0 | 28.73 | 0 | 13.59 | 0 |
| PERCENT DOMINANT TAXON | 28.22 | -1 | 19.22 | 0 | 31.64 | -1 | 31.72 | -1 |
| PERCENT ISOPOD, SNAIL, LEECH | 28.22 | -1 | 0.84 | 1 | 0.73 | 1 | 0.32 | 1 |
| PERCENT SURF. AIR BREATHERS | 0.49 | 1 | 0.28 | 1 | 1.45 | 1 | 0.32 | 1 |
| TOTAL SCORE | | 4 | | 5 | | 1 | | 4 |
| MACROINV. COMMUNITY RATING | | ACCEPT. | | EXCELLEN | г. | ACCEPT. | | ACCEPT. |

| Table 3A con't. | Qualitative macroinvertebrate | sampling results for the | Tahquamenon, | Two-Hearted, | Manistique, ar | nd Millecoquins | River |
|-----------------|-------------------------------|--------------------------|--------------|--------------|----------------|-----------------|-------|
| targeted sampli | ng sites. | | | | | | |

| | Davenport Creek at gas pipeline 6/10/2009 | Doe Creek Raski Road 6/10/2009 | Black River State Forest Campground bridge 6/8/2009 | Crow River end of Summer Trail 6/8/2009 |
|------------------------------------|---|--------------------------------------|---|---|
| TAXA | STATION 34 | STATION 35 | STATION 36 | STATION 37 |
| ANNELIDA (segmented worms) | | | | |
| Hirudinea (leeches) | | 1 | 1 | |
| Oligochaeta (worms) | 1 | 2 | 3 | 1 |
| ARTHROPODA | | | | |
| Crustacea | 2 | 2 | 1 | 2 |
| Decapoda (cravfish) | 3 | 3 | 1 | 3 |
| Arachnoidea | | 1 | | |
| Hydracarina | 1 | | 1 | |
| Insecta | | | | |
| Ephemeroptera (mayflies) | | | | |
| Baetiscidae | | | | 6 |
| Baetidae | 55 | 21 | 39 | 35 |
| Caenidae | 20 | 5 | 11 | 1 |
| Ephemeridae | 23 | | 11 | 1 |
| Heptageniidae | | 3 | 1 | 8 |
| Isonychiidae | | | 6 | |
| Odonata | | | | |
| Anisoptera (dragonflies) | | | | |
| Aeshnidae | | 1 | | 8 |
| Gomphidae | | | | 8 |
| Lygoptera (damseiffies) | | | | 5 |
| Coepagrionidae | | 1 | | 5 |
| Plecoptera (stoneflies) | | 1 | | |
| Leuctridae | | | 12 | |
| Nemouridae | 4 | | 1 | |
| Perlidae | 3 | | | |
| Perlodidae | | | 9 | |
| Hemiptera (true bugs) | | | | 10 |
| Corrixidae | 1 | | 1 | 18 |
| Megaloptera | 1 | | 1 | |
| Corvdalidae (dobson flies) | | 3 | | 1 |
| Trichoptera (caddisflies) | | | | |
| Brachycentridae | 77 | | 24 | 1 |
| Helicopsychidae | | 1 | 1 | |
| Hydropsychidae | 12 | | 3 | 22 |
| Hydroptilidae | 16 | 32 | 5 | |
| Lepidostomatidae | 16 | | 40 | 1 |
| Limnenhilidae | 47 | 3 | 40 | 28 |
| Molannidae | | 5 | 10 | 1 |
| Philopotamidae | 1 | | | |
| Polycentropodidae | | | | 1 |
| Rhyacophilidae | | | 1 | |
| Coleoptera (beetles) | | 10 | | |
| Dytiscidae (total) | | 19 | I | |
| Halinlidae (adults) | 1 | 10 | | |
| Hydrophilidae (total) | 1 | 2 | 1 | |
| Dryopidae | | - 1 | - 1 | |
| Elmidae | | 10 | 1 | |
| Diptera (flies) | | | | |
| Athericidae | | _ | _ | 1 |
| Chironomidae | 1 | 9 | 3 | 1 |
| Divideo | 55 | 121 | 52 | 45 |
| Empididae | | 1 | | 1 |
| Ptychopteridae | | - | 1 | |
| Simuliidae | 14 | 6 | 17 | 31 |
| Tabanidae | | | 1 | 3 |
| Tipulidae | 1 | 1 | 2 | 1 |
| MOLLUSCA | | | | |
| Gastropoda (snails) Hydrobiidae | | 3 | | 4 |
| Lymnaeidae | | 5 | | + 2 |
| Physidae | | 1 | 1 | 4 |
| Planorbidae | | 2 | | 4 |
| Pelecypoda (bivalves) | | | | |
| Sphaeriidae (clams) | 4 | 1 | 16 | 6 |
| | | | | |
| TOTAL INDIVIDUALS | 324 | 265 | 298 | 257 |

| | Davenport Creek at gas pipeline 6/10/2009 STATION 34 | | Doe Creek Raski Road 6/10/2009 STATION 35 | | Black River State Forest Campground bridge 6/8/2009 STATION 36 | | | Crow River end of Summer Trail 6/8/2009 STATION 37 | |
|------------------------------|---|---------|--|---------|---|-------|-----------|---|---------|
| METRIC | Value | Score | Value | Score | Value | | Score | Value | Score |
| TOTAL NUMBER OF TAXA | 19 | 0 | 28 | 1 | | 32 | 1 | 32 | 1 |
| NUMBER OF MAYFLY TAXA | 2 | -1 | 3 | 0 | | 4 | 0 | 6 | 1 |
| NUMBER OF CADDISFLY TAXA | 5 | 0 | 3 | 0 | | 8 | 1 | 7 | 1 |
| NUMBER OF STONEFLY TAXA | 2 | 1 | 0 | -1 | | 3 | 1 | 0 | -1 |
| PERCENT MAYFLY COMP. | 25.93 | 1 | 10.94 | 0 | | 19.13 | 0 | 20.62 | 0 |
| PERCENT CADDISFLY COMP. | 47.22 | 1 | 13.58 | 0 | | 38.59 | 1 | 22.18 | 0 |
| PERCENT DOMINANT TAXON | 23.77 | 0 | 45.66 | -1 | | 17.45 | 0 | 17.51 | 0 |
| PERCENT ISOPOD, SNAIL, LEECH | 0.00 | 1 | 3.02 | 1 | | 0.67 | 1 | 5.45 | 0 |
| PERCENT SURF. AIR BREATHERS | 0.62 | 1 | 11.70 | 0 | | 1.34 | 1 | 7.00 | 0 |
| TOTAL SCORE | | 4 | | 0 | | | 6 | | 2 |
| MACROINV. COMMUNITY RATING | | ACCEPT. | | ACCEPT. | | E | EXCELLENT | Г 4 | ACCEPT. |

Table 4. Habitat evaluation for 2009 Tahquamenon, Two-Hearted, Manistique, and Millecoquins Rivers probabilistic sampling sites.

| | Syphon Creek M-442 GLIDE/POOL STATION 1 | Star Creek Off Star Siding Road GLIDE/POOL STATION 2 | Star Creek Grass Creek tar Siding Road Railroad Grade off Sunset Landin LIDE/POOL GLIDE/POOL TATION 2 STATION 3 | | Kilpecker Creek two-track off Federal Forest Rd 2438 GLIDE/POOL STATION 5 |
|--|--|---|---|---------------------------------|--|
| HABITAT METRIC | | | | | |
| Substrate and Instream Cover | | | | | |
| Epifaunal Substrate/ Avail Cover (20) | 10 | 10 | 10 | 12 | 14 |
| Embeddedness (20)* | | | | | |
| Velocity/Depth Regime (20)* | | | | | |
| Pool Substrate Characterization (20)** | 6 | 11 | 10 | 10 | 8 |
| Pool Variability (20)** | 1 | 8 | 5 | 13 | 11 |
| Channel Morphology | | | | | |
| Sediment Deposition (20) | 2 | 8 | 11 | 17 | 8 |
| Flow Status - Maint. Flow Volume (10) | 10 | 9 | 10 | 10 | 10 |
| Flow Status - Flashiness (10) | 10 | 8 | 10 | 10 | 10 |
| Channel Alteration (20) | 20 | 20 | 20 | 15 | 20 |
| Frequency of Riffles/Bends (20)* | | | | | |
| Channel Sinuosity (20)** | 15 | 14 | 15 | 12 | 18 |
| Riparian and Bank Structure | | | | | |
| Bank Stability (L) (10) | 10 | 9 | 10 | 10 | 10 |
| Bank Stability (R) (10) | 10 | 9 | 10 | 10 | 10 |
| Vegetative Protection (L) (10) | 10 | 8 | 10 | 10 | 9 |
| Vegetative Protection (R) (10) | 10 | 8 | 10 | 10 | 10 |
| Riparian Veg. Zone Width (L) (10) | 10 | 10 | 10 | 10 | 10 |
| Riparian Veg. Zone Width (R) (10) | 10 | 10 | 10 | 6 | 10 |
| TOTAL SCORE (200): | 134 | 142 | 151 | 155 | 158 |
| HABITAT RATING: | GOOD (SLIGHTLY IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) | EXCELLENT (NON- IMPAIRED) | EXCELLENT (NON- IMPAIRED) |

Note: Individual metrics may better describe conditions directly affecting the biological community while the Habitat Rating describe the general riverine environment at the site(s).

| Date: | 7/29/2009 | | 6/24/2009 | | 7/28/2009 | | 6/9/2009 | | 7/27/2009 | |
|-------------------------|--------------|----------|--------------------|----------|-----------------------|-----------|------------------|----------|-------------------------|----------------|
| Weather: | Sunny | | Sunny | | Sunny | | Cloudy | | Cloudy | r |
| Air Temperature: | 70 | Deg. F. | 75 | Deg. F. | 75 | Deg. F. | 49 | Deg. F. | | Deg. F. |
| Water Temperature: | 48 | Deg. F. | 62 | Deg. F. | 66 | Deg. F. | 47 | Deg. F. | 50 | Deg. F. |
| Ave. Stream Width: | 15 | Feet | 18 | Feet | 8 | Feet | 22.5 | Feet | 15 | Feet |
| Ave. Stream Depth: | 0.83 | Feet | 1.5 | Feet | 1.5 | Feet | 1.8 | Feet | 0.67 | Feet |
| Surface Velocity: | 1.67 | Ft./Sec. | 0.33 | Ft./Sec. | 1 | Ft./Sec. | 1.5 | Ft./Sec. | 1 | Ft./Sec. |
| Estimated Flow: | 20.7915 | CFS | 8.91 | CFS | 12 | CFS | 60.75 | CFS | 10.05 | CFS |
| Stream Modifications: | None | | None | | None | | Relocated | | None | : |
| Nuisance Plants (Y/N): | N | | N | | Ν | | N | | N | |
| Report Number: | | | | | | | | | | |
| STORET No.: | 480038 | | 20124 | | 20159 | | 770090 | | 210305 | |
| Stream Name: | Syphon Creek | | Star Creek | | Grass Creek | East | Branch Fox River | | Kilpecker Creek | |
| Road Crossing/Location: | M-442 | | Off Star Siding Ro | ad | Railroad Grade off Su | unset Lan | M-77 crossing | | two-track off Federal I | Forest Rd 2438 |
| County Code: | 48 | | 02 | | 02 | | 77 | | 21 | |
| TRS: | 46N12W12 | | 46N17W23 | | 48N15W25 | | 46N13W05 | | 43N18W24 | |
| Latitude (dd): | 46.39083 | | 46.37216 | | 46.52231 | | 46.40607 | | 46.11395 | |
| Longitude (dd): | -85.74472 | | -86.39325 | | -86.12086 | | -85.94795 | | -86.49214 | |
| Ecoregion: | NLAF | | NLAF | | NLAF | | NLAF | | NLAF | 1 |
| Stream Type: | Coldwater | | Coldwater | | Coldwater | | Coldwater | | Coldwater | |
| USGS Basin Code: | 4020202 | | 4060106 | | 4060106 | | 4060106 | | 4060106 | |

* Applies only to Riffle/Run stream Survey ** Applies only to Glide/Pool stream Surveys

Table 4 con't. Habitat evaluation for 2009 Tabquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites

| | East Branch Tahouamenon River | Mead Creek | Big Murphy Creek | Clear Creek | Chenev Creek |
|--|-------------------------------|---------------|------------------|---------------|----------------------------|
| | Arbutus Truck Trail | M-77 crossing | County Road 437 | upstream M-77 | u/s Tahq. River confluence |
| | GLIDE/POOL | GLIDE/POOL | GLIDE/POOL | GLIDE/POOL | GLIDE/POOL |
| | STATION 6 | STATION 7 | STATION 8 | STATION 9 | STATION 10 |
| HABITAT METRIC | | | | | |
| Substrate and Instream Cover | | | | | |
| Epifaunal Substrate/ Avail Cover (20) | 9 | 9 | 10 | 8 | 11 |
| Embeddedness (20)* | | | | | |
| Velocity/Depth Regime (20)* | | | | | |
| Pool Substrate Characterization (20)** | 12 | 12 | 8 | 11 | 8 |
| Pool Variability (20)** | 10 | 14 | 11 | 13 | 14 |
| Channel Morphology | | | | | |
| Sediment Deposition (20) | 20 | 14 | 6 | 20 | 12 |
| Flow Status - Maint. Flow Volume (10) | 9 | 8 | 9 | 10 | 9 |
| Flow Status - Flashiness (10) | 9 | 10 | 9 | 10 | 10 |
| Channel Alteration (20) | 20 | 20 | 20 | 20 | 20 |
| Frequency of Riffles/Bends (20)* | | | | | |
| Channel Sinuosity (20)** | 11 | 11 | 11 | 9 | 10 |
| Riparian and Bank Structure | | | | | |
| Bank Stability (L) (10) | 10 | 9 | 8 | 10 | 6 |
| Bank Stability (R) (10) | 10 | 9 | 8 | 10 | 6 |
| Vegetative Protection (L) (10) | 10 | 10 | 9 | 10 | 9 |
| Vegetative Protection (R) (10) | 10 | 10 | 9 | 10 | 9 |
| Riparian Veg. Zone Width (L) (10) | 10 | 10 | 10 | 10 | 9 |
| Riparian Veg. Zone Width (R) (10) | 10 | 10 | 10 | 10 | 9 |
| TOTAL SCORE (200): | 160 | 156 | 138 | 161 | 142 |
| HABITAT BATING: | EXCELLENT | EXCELLENT | GOOD | EXCELLENT | GOOD |
| habitat Kaling. | (NON- | (NON- | (SUIGHTI V | (NON- | (SUGHTI V |
| | IMPAIRED) | IMPAIRED) | IMPAIRED) | IMPAIRED) | IMPAIRED) |

Note: Individual metrics may better describe conditions directly affecting the biological community while the Habitat Rating describes the general riverine environment at the site(s).

| Date: | 6/11/2009 | | 6/9/2009 | | 6/22/2009 | | 6/9/2009 | | 6/10/2009 | , |
|-------------------------|-------------------------------|----------|---------------|----------|---------------|----------|---------------|----------|----------------|------------|
| Weather: | Sunny | | Cloudy | | Partly Cloudy | | Cloudy | | Sunny | , |
| Air Temperature: | 50 | Deg. F. | 52 | Deg. F. | 80 | Deg. F. | 48 | Deg. F. | | Deg. F. |
| Water Temperature: | 50 | Deg. F. | 53 | Deg. F. | 66 | Deg. F. | 44 | Deg. F. | 52 | Deg. F. |
| Ave. Stream Width: | 7.5 | Feet | 19 | Feet | 20 | Feet | 10 | Feet | 16 | Feet |
| Ave. Stream Depth: | 0.43 | Feet | 1.2 | Feet | 1.5 | Feet | 0.75 | Feet | 1.25 | Feet |
| Surface Velocity: | 0.8 | Ft./Sec. | 1.2 | Ft./Sec. | 1 | Ft./Sec. | 1.4 | Ft./Sec. | 0.77 | Ft./Sec. |
| Estimated Flow: | 2.58 | CFS | 27.36 | CFS | 30 | CFS | 10.5 | CFS | 15.4 | CFS |
| Stream Modifications: | None | | None | | None | | None | | None | • |
| Nuisance Plants (Y/N): | Ν | | N | | Ν | | N | | N | |
| Report Number: | | | | | | | | | | |
| STORET No.: | 170288 | | 770089 | | 770159 | | 770108 | | 170191 | |
| Stream Name: | East Branch Tahquamenon River | | Mead Creek | Big | Murphy Creek | | Clear Creek | | Cheney Creek | |
| Road Crossing/Location: | Arbutus Truck Trail | | M-77 crossing | | County Road 4 | 37 | upstream M-77 | 7 | u/s Tahq. Rive | r confluei |
| County Code: | 17 | | 77 | | 77 | | 77 | | 17 | , |
| TRS: | 45N05W05 | | 44N13W27 | | 43N17W32 | | 47N13W21 | | 48N06W08 | ; |
| Latitude (dd): | 46.3177 | | 46.17484 | | 46.07281 | | 46.44928 | | 46.56306 | |
| Longitude (dd): | -84.9527 | | -85.92751 | | -86.4669 | | -85.93728 | | -85.08312 | |
| Ecoregion: | NLAF | | NLAF | | NLAF | | NLAF | | NLAF | 7 |
| Stream Type: | Coldwater | | Warmwater | | Coldwater | | Coldwater | | Coldwater | |
| USGS Basin Code: | 4020202 | | 4060106 | | 4060106 | | 4060106 | | 4020202 | |

* Applies only to Riffle/Run stream Surveys ** Applies only to Glide/Pool stream Surveys

Table 4 con't. Habitat evaluation for 2009 Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites

| | Little Fox River | East Branch Two Hearted River | South Branch Stutts Creek | Sucker River | Creighton River |
|--|--------------------------------|---------------------------------|--------------------------------|--------------------------------|--------------------------------|
| | Stanley Lake Road | County Road 414 | Clear Lake Road | Grand Marais Truck Trail | Creighton Truck Trail |
| | GLIDE/POOL | GLIDE/POOL | GLIDE/POOL | GLIDE/POOL | GLIDE/POOL |
| | STATION 11 | STATION 12 | STATION 13 | STATION 14 | STATION 16 |
| HABITAT METRIC | | | | | |
| Substrate and Instream Cover | | | | | |
| Epifaunal Substrate/ Avail Cover (20) | 11 | 10 | 10 | 10 | 9 |
| Embeddedness (20)* | | | | | |
| Velocity/Depth Regime (20)* | | | | | |
| Pool Substrate Characterization (20)** | 12 | 11 | 11 | 7 | 8 |
| Pool Variability (20)** | 9 | 13 | 5 | 16 | 11 |
| Channel Morphology | | | | | |
| Sediment Deposition (20) | 8 | 10 | 10 | 6 | 7 |
| Flow Status - Maint. Flow Volume (10) | 10 | 10 | 8 | 10 | 9 |
| Flow Status - Flashiness (10) | 10 | 10 | 7 | 10 | 6 |
| Channel Alteration (20) | 19 | 20 | 20 | 20 | 20 |
| Frequency of Riffles/Bends (20)* | | | | | |
| Channel Sinuosity (20)** | 15 | 13 | 13 | 13 | 13 |
| Riparian and Bank Structure | | | | | |
| Bank Stability (L) (10) | 10 | 9 | 7 | 2 | 8 |
| Bank Stability (R) (10) | 10 | 9 | 7 | 2 | 6 |
| Vegetative Protection (L) (10) | 9 | 10 | 10 | 10 | 10 |
| Vegetative Protection (R) (10) | 9 | 10 | 10 | 10 | 10 |
| Riparian Veg. Zone Width (L) (10) | 10 | 10 | 8 | 10 | 10 |
| Riparian Veg. Zone Width (R) (10) | 10 | 10 | 10 | 10 | 10 |
| TOTAL SCORE (200): | 152 | 155 | 136 | 136 | 137 |
| HABITAT RATING: | GOOD (SLIGHTLY IMPAIRED) | EXCELLENT (NON- IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) |

Note: Individual metrics may better describe conditions directly affecting the biological community while the Habitat Rating describes the general riverine environment at the site(s).

| Date: | 6/23/2009 | | 7/30/2009 | | 6/25/2009 | | 7/29/2009 | | 6/24/2009 | |
|-------------------------|------------------|-------------|-------------------|----------|-----------------------|----------|--------------------|----------|-----------------|----------|
| Weather: | Sunny | | Partly Cloudy | | Cloudy | | Cloudy | | Sunny | |
| Air Temperature: | 80 | Deg. F. | 70 | Deg. F. | 75 | Deg. F. | 70 | Deg. F. | 85 | Deg. F. |
| Water Temperature: | 62 | Deg. F. | 54 | Deg. F. | | Deg. F. | 64 | Deg. F. | 62 | Deg. F. |
| Ave. Stream Width: | 12 | Feet | 85 | Feet | 22 | Feet | 30 | Feet | 32 | Feet |
| Ave. Stream Depth: | 1 | Feet | 1.5 | Feet | 1.25 | Feet | 0.83 | Feet | 2 | Feet |
| Surface Velocity: | 1.2 | Ft./Sec. | 1 | Ft./Sec. | 1 | Ft./Sec. | 1.67 | Ft./Sec. | 0.4 | Ft./Sec. |
| Estimated Flow: | 14.4 | CFS | 127.5 | CFS | 27.5 | CFS | 41.583 | CFS | 25.6 | CFS |
| Stream Modifications: | None | | None | | None | | None | | None | |
| Nuisance Plants (Y/N): | N | | N | | Ν | | N | | N | |
| Report Number: | | | | | | | | | | |
| STORET No.: | 770157 | | 480028 | | 770158 | | 20160 | | 770156 | |
| Stream Name: | Little Fox River | East Branch | Two Hearted River | Sout | h Branch Stutts Creek | | Sucker River | . (| Creighton River | |
| Road Crossing/Location: | Stanley Lake Roa | ıd | County Road 414 | | Clear Lake Road | Grand M | Iarais Truck Trail | Creigh | ton Truck Trail | |
| County Code: | 77 | | 48 | | 77 | | 02 | | 77 | |
| TRS: | 47N15W11 | | 49N09W18 | | 44N17W10 | | 49N13W12 | | 44N16W01 | |
| Latitude (dd): | 46.47622 | | 46.64188 | | 46.2224 | | 46.66203 | | 46.24108 | |
| Longitude (dd): | -86.13987 | | -85.47916 | | -86.41703 | | -85.86893 | | -86.24185 | |
| Ecoregion: | NLAF | 1 | NLAF | | NLAF | 1 | NLAF | | NLAF | 1 |
| Stream Type: | Coldwater | | Coldwater | | Coldwater | | Coldwater | | Warmwater | |
| USGS Basin Code: | 4060106 | | 4020201 | | 4060106 | | 4020201 | | 4060106 | |

* Applies only to Riffle/Run stream Surveys ** Applies only to Glide/Pool stream Surveys

Table 4 con't. Habitat evaluation for 2009 Tahquamenon, Two-Hearted, Manistique, and Millecoquins River probabilistic sampling sites.

| | Milakokia River Betty Doe Lake Rd GLIDE/POOL STATION 17 | lakokia River West Branch Two Hearted River West Branch Manistique River y Doe Lake Rd County Road 418 Hickey Truck Trail LIDE/POOL GLIDE/POOL GLIDE/POOL TATION 17 STATION 18 STATION 19 | | Shelldrake River Betsy River Truck Trail GLIDE/POOL STATION 20 | West Branch Manistique River off Crieghton Truck Trail GLIDE/POOL STATION 21 |
|--|--|---|--------------------------------|---|---|
| HABITAT METRIC | | | | | |
| Substrate and Instream Cover | | | | | |
| Epifaunal Substrate/ Avail Cover (20) | 18 | 12 | 7 | 12 | 8 |
| Embeddedness (20)* | | | | | |
| Velocity/Depth Regime (20)* | | | | | |
| Pool Substrate Characterization (20)** | 17 | 7 | 8 | 13 | 9 |
| Pool Variability (20)** | 15 | 3 | 13 | 12 | 11 |
| Channel Morphology | | | | | |
| Sediment Deposition (20) | 19 | 11 | 6 | 16 | 6 |
| Flow Status - Maint. Flow Volume (10) | 9 | 10 | 9 | 9 | 9 |
| Flow Status - Flashiness (10) | 10 | 10 | 7 | 9 | 5 |
| Channel Alteration (20) | 20 | 20 | 20 | 20 | 20 |
| Frequency of Riffles/Bends (20)* | | | | | |
| Channel Sinuosity (20)** | 10 | 17 | 14 | 13 | 15 |
| Riparian and Bank Structure | | | | | |
| Bank Stability (L) (10) | 10 | 1 | 4 | 9 | 4 |
| Bank Stability (R) (10) | 10 | 10 | 4 | 9 | 4 |
| Vegetative Protection (L) (10) | 10 | 8 | 6 | 10 | 10 |
| Vegetative Protection (R) (10) | 10 | 10 | 6 | 10 | 10 |
| Riparian Veg. Zone Width (L) (10) | 10 | 10 | 10 | 10 | 7 |
| Riparian Veg. Zone Width (R) (10) | 10 | 10 | 9 | 8 | 10 |
| TOTAL SCORE (200): | 178 | 139 | 123 | 160 | 128 |
| HABITAT RATING: | EXCELLENT (NON- IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) | EXCELLENT (NON- IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) |

Note: Individual metrics may better describe conditions directly affecting the biological community while the Habitat Rating describes the gener riverine environment at the site(s).

| Date: | 6/9/2009 | 7/29/2009 |) | 7/27/2009 | | 6/10/2009 | 6/24/2 | 009 |
|-------------------------|-----------------|-------------------------------|----------|------------------------------|----------|------------------|--------------------------|------------|
| Weather: | Cloudy | Cloudy | <i>,</i> | Partly Cloudy | | Sunny | Su | nny |
| Air Temperature: | 53 | Deg. F. | Deg. F. | | Deg. F. | 58 | Deg. F. | 80 Deg. F. |
| Water Temperature: | 52 | Deg. F. 56 | Deg. F. | 64 | Deg. F. | 57 | Deg. F. | 62 Deg. F. |
| Ave. Stream Width: | 19 | Feet 25 | Feet | 35 | Feet | 23 | Feet | 45 Feet |
| Ave. Stream Depth: | 0.64 | Feet 3 | Feet | 2.5 | Feet | 1.5 | Feet | 1 Feet |
| Surface Velocity: | 1.3 | Ft./Sec. 1.25 | Ft./Sec. | 0.71 | Ft./Sec. | 0.77 | Ft./Sec. | 1 Ft./Sec. |
| Estimated Flow: | 15.808 | CFS 93.75 | CFS | 62.125 | CFS | 26.565 | CFS | 45 CFS |
| Stream Modifications: | None | None | e | None | | None | N | one |
| Nuisance Plants (Y/N): | N | N | 1 | N | | N | | N |
| Report Number: | | | | | | | | |
| STORET No.: | 490065 | 480067 | | 770155 | | 170287 | 7701 | 60 |
| Stream Name: | Ailakokia River | West Branch Two Hearted River | r | West Branch Manistique River | 5 | Shelldrake River | Vest Branch Manistique R | iveı |
| Road Crossing/Location: | y Doe Lake Rd | County Road 418 | 3 | Hickey Truck Trail | Betsy R | iver Truck Trail | off Crieghton Truck | rail |
| County Code: | 49 | 48 | 3 | 77 | | 17 | - | 77 |
| TRS: | 42N12W18 | 48N11W09 |) | 45N16W35 | | 49N07W06 | 44N15V | V07 |
| Latitude (dd): | 46.02832 | 46.56948 | | 46.25291 | | 46.6729 | 46.228 | 57 |
| Longitude (dd): | -85.84404 | -85.68017 | | -86.26238 | | -85.2339 | -86.235 | 06 |
| Ecoregion: | NLAF | NLAF | 7 | NLAF | | NLAF | NI | AF |
| Stream Type: | Warmwater | Warmwater | r | Warmwater | | Warmwater | Warmw | ater |
| USGS Basin Code: | 4060107 | 4020201 | | 4060106 | | 4020201 | 40601 | 06 |

* Applies only to Riffle/Run stream Survey ** Applies only to Glide/Pool stream Survey!

| Table 4 con't. Habitat evaluation for 2009 | Tahquamenon, Two-Hearte | d. Manistique, and Millecoquir | s River probabilistic sampling sites |
|--|-------------------------|--------------------------------|--------------------------------------|
| | | | |

| | Paquin Creek Hiawatha Trail GLIDE/POOL STATION 24 | H Sti | log Island Cree rickler Truck Tr RIFFLE/RUN STATION 25 | ek rail |
|--|--|------------------------------------|---|---------------------------|
| HABITAT METRIC | | | | |
| Substrate and Instream Cover | | | | |
| Epifaunal Substrate/ Avail Cover (20) | 10 | | 18 | |
| Embeddedness (20)* | | | 15 | |
| Velocity/Depth Regime (20)* | | | 14 | |
| Pool Substrate Characterization (20)** | 8 | | | |
| Pool Variability (20)** | 8 | | | |
| Channel Morphology | | | | |
| Sediment Deposition (20) | 15 | | 18 | |
| Flow Status - Maint. Flow Volume (10) | 8 | | 7 | |
| Flow Status - Flashiness (10) | 8 | | 10 | |
| Channel Alteration (20) | 20 | | 20 | |
| Frequency of Riffles/Bends (20)* | | | 18 | |
| Channel Sinuosity (20)** | 11 | | | |
| Riparian and Bank Structure | | | | |
| Bank Stability (L) (10) | 7 | | 10 | |
| Bank Stability (R) (10) | 7 | | 10 | |
| Vegetative Protection (L) (10) | 10 | | 10 | |
| Vegetative Protection (R) (10) | 10 | | 10 | |
| Riparian Veg. Zone Width (L) (10) | 10 | | 10 | |
| Riparian Veg. Zone Width (R) (10) | 10 | | 10 | |
| TOTAL SCORE (200): | 142 | | 180 | |
| HABITAT RATING: | GOOD (SLIGHTLY IMPAIRED) | | EXCELLENT (NON- IMPAIRED) | |
| | Note: Individua biological com environment at | al metric munity v the site(| s may better de vhile the Habita s). | scribe con at Rating d |
| Date: | 6/7/2009 | | 6/10/2009 | |
| Weather: | Cloudy | | Sunny | |
| Air Temperature: | 52 | Deg. F. | | Deg. F. |
| Water Temperature: | 50 | Deg. F. | 49 | Deg. F. |
| Ave. Stream Width: | 12 | Feet | 8.3 | Feet |
| Ave. Stream Depth: | 0.44 | Feet | 0.41 | Feet |
| Surface Velocity: | 0.9 | Ft./Sec. | 0.4 | Ft./Sec. |
| Estimated Flow: | 4.752 | CFS | 1.3612 | CFS |

conditions directly affecting the g describes the general riverine

| () outlion | Cioudy | | Buility | |
|-------------------------|----------------|----------|---------------|----------|
| Air Temperature: | 52 | Deg. F. | | Deg. F. |
| Water Temperature: | 50 | Deg. F. | 49 | Deg. F. |
| Ave. Stream Width: | 12 | Feet | 8.3 | Feet |
| Ave. Stream Depth: | 0.44 | Feet | 0.41 | Feet |
| Surface Velocity: | 0.9 | Ft./Sec. | 0.4 | Ft./Sec. |
| Estimated Flow: | 4.752 | CFS | 1.3612 | CFS |
| Stream Modifications: | None | | None | |
| Nuisance Plants (Y/N): | N | | Ν | |
| Report Number: | | | | |
| | | | | |
| STORET No.: | 490195 | | 490200 | |
| Stream Name: | Paquin Creek | Hog | Island Creek | |
| Road Crossing/Location: | Hiawatha Trail | Strickle | r Truck Trail | |
| County Code: | 49 | | 49 | |
| TRS: | 43N07W35 | | 43N08W13 | |
| | | | | |
| Latitude (dd): | 46.0813 | | 46.1228 | |
| Longitude (dd): | -85.1545 | | -85.254 | |
| Ecoregion: | NLAF | i | NLAF | |
| Stream Type: | Coldwater | | Coldwater | |
| | | | | |
| USGS Basin Code: | 4060107 | | 4060107 | |
| | | | | |

* Applies only to Riffle/Run stream Surveys

** Applies only to Glide/Pool stream Surveys

| Table 4 con't. Habitat evaluation for 2009 Tahquamenon, Two-Hearted, Manistique, and Millecoquins River targeted sampling sites. Manistique, and Millecoquins River Manistique River Millecoquing River Tahquamenon River Indian River Manistique River Fay River Millecoquing River | | | | | | | | | | | | | |
|--|---|---|--|---|--|--|--|--|--|--|--|--|--|
| | Tahquamenon River near Eagles Nest RIFFLE/RUN STATION 26 | Indian River 2-track east off 94 GLIDE/POOL STATION 27 | Manistique River Manistique boat launch GLIDE/POOL STATION 28 | Fox River Fox River campground GLIDE/POOL STATION 29 | Millecoquins River Lower Rapids RIFFLE/RUN STATION 30 | | | | | | | | |
| HABITAT METRIC | | | | | | | | | | | | | |
| Substrate and Instream Cover | | | | | | | | | | | | | |
| Epifaunal Substrate/ Avail Cover (20) | 17 | 9 | 11 | 16 | 19 | | | | | | | | |
| Embeddedness (20)* | 14 | | | | 20 | | | | | | | | |
| Velocity/Depth Regime (20)* | 15 | | | | 15 | | | | | | | | |
| Pool Substrate Characterization (20)** | | 8 | 8 | 14 | | | | | | | | | |
| Pool Variability (20)** | | 13 | 2 | 15 | | | | | | | | | |
| Channel Morphology | | | | | | | | | | | | | |
| Sediment Deposition (20) | 8 | 7 | 15 | 11 | 20 | | | | | | | | |
| Flow Status - Maint. Flow Volume (10) | 10 | 9 | 9 | 10 | 9 | | | | | | | | |
| Flow Status - Flashiness (10) | 10 | 9 | 9 | 10 | 9 | | | | | | | | |
| Channel Alteration (20) | 20 | 19 | 13 | 20 | 20 | | | | | | | | |
| Frequency of Riffles/Bends (20)* | 16 | | | | 16 | | | | | | | | |
| Channel Sinuosity (20)** | | 13 | 5 | 13 | | | | | | | | | |
| Riparian and Bank Structure | | | | | | | | | | | | | |
| Bank Stability (L) (10) | 10 | 7 | 9 | 9 | 10 | | | | | | | | |
| Bank Stability (R) (10) | 9 | 9 | 9 | 9 | 10 | | | | | | | | |
| Vegetative Protection (L) (10) | 10 | 9 | 9 | 10 | 10 | | | | | | | | |
| Vegetative Protection (R) (10) | 10 | 10 | 9 | 10 | 10 | | | | | | | | |
| Riparian Veg. Zone Width (L) (10) | 10 | 10 | 4 | 10 | 10 | | | | | | | | |
| Riparian Veg. Zone Width (R) (10) | 9 | 10 | 7 | 9 | 10 | | | | | | | | |
| TOTAL SCORE (200): | 168 | 142 | 119 | 166 | 188 | | | | | | | | |
| HABITAT RATING: | EXCELLENT (NON- IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) | GOOD (SLIGHTLY IMPAIRED) | EXCELLENT (NON- IMPAIRED) | EXCELLENT (NON- IMPAIRED) | | | | | | | | |

Note: Individual metrics may better describe conditions directly affecting the biological community while the Habita Rating describes the general riverine environment at the site(s).

| Date: | 7/30/2009 | | 7/27/2009 | | 7/28/2009 | | 7/28/2009 | | 6/8/2009 | |
|-------------------------|-------------------|----------|------------------|----------|-------------------|----------|------------------|----------|------------------|----------|
| Weather: | Cloudy | | Cloudy | | Sunny | | Partly Cloudy | | Cloudy | |
| Air Temperature: | 70 | Deg. F. | 70 | Deg. F. | 70 | Deg. F. | 75 | Deg. F. | 57 | Deg. F. |
| Water Temperature: | 61 | Deg. F. | 60 | Deg. F. | 68 | Deg. F. | 61 | Deg. F. | 56 | Deg. F. |
| Ave. Stream Width: | 45 | Feet | 30 | Feet | 150 | Feet | 38 | Feet | 49 | Feet |
| Ave. Stream Depth: | 0.83 | Feet | 2.5 | Feet | 3 | Feet | 2 | Feet | 1.3 | Feet |
| Surface Velocity: | 1.67 | Ft./Sec. | 1 | Ft./Sec. | 0.71 | Ft./Sec. | 1.67 | Ft./Sec. | 1.4 | Ft./Sec. |
| Estimated Flow: | 62.3745 | CFS | 75 | CFS | 319.5 | CFS | 126.92 | CFS | 89.18 | CFS |
| Stream Modifications: | None | Ba | nk Stabilization | | None | Ba | nk Stabilization | | None | |
| Nuisance Plants (Y/N): | N | | N | | N | | Ν | | N | |
| Report Number: | | | | | | | | | | |
| STORET No.: | 480066 | | 770085 | | 770154 | | 770082 | | 490196 | |
| Stream Name: | Tahquamenon River | | Indian River | | Manistique River | | Fox River | Mi | llecoquins River | |
| Road Crossing/Location: | near Eagles Nest | | 2-track east off | 94 | Manistique boat l | aunch | Fox River cam | pground | Lower Rapids | |
| County Code: | 48 | | 77 | | 77 | | 77 | | 49 | |
| TRS: | 47N12W34 | | 44N17W34 | | 41N16W01 | | 46N14W11 | | 43N09W18 | |
| Latitude (dd): | 46.42482 | | 46.17226 | | 45.97088 | | 46.39997 | | 46.11568 | |
| Longitude (dd): | -85.79771 | | -86.4213 | | -86.24275 | | -86.02787 | | -85.47358 | |
| Ecoregion: | NLAF | | NLAF | | NLAF | | NLAF | | NLAF | ; |
| Stream Type: | Coldwater | | Coldwater | | Warmwater | | Coldwater | | Warmwater | ſ |
| USGS Basin Code: | 4020202 | | 4060106 | | 4060106 | | 4060106 | | 4060107 | |

* Applies only to Riffle/Run stream Survey: ** Applies only to Glide/Pool stream Survey:

| Table 4 con't. Habitat evaluation for 2009 | Tahquamenon, Two | o-Hearted, Manistique, | and Millecoquins Rive | er targeted sampling site | s. |
|--|------------------|------------------------|-----------------------|---------------------------|------------|
| | Furlong Creek | Davenport Creek | Cut River | Davenport Creek | Doe Creek |
| | Pleasant Ave | US-2 | US-2 | at gas pipeline | Raski Road |
| | RIFFLE/RUN | GLIDE/POOL | RIFFLE/RUN | GLIDE/POOL | GLIDE/POOL |
| | STATION 31 | STATION 32 | STATION 33 | STATION 34 | STATION 35 |
| HABITAT METRIC | | | | | |
| Substrate and Instream Cover | | | | | |
| Epifaunal Substrate/ Avail Cover (20) | 20 | 10 | 19 | 11 | 11 |
| Embeddedness (20)* | 20 | | 13 | | |
| Velocity/Depth Regime (20)* | 20 | | 10 | | |
| Pool Substrate Characterization (20)** | | 9 | | 7 | 9 |
| Pool Variability (20)** | | 13 | | 7 | 13 |
| Channel Morphology | | | | | |
| Sediment Deposition (20) | 19 | 8 | 16 | 16 | 16 |
| Flow Status - Maint. Flow Volume (10) | 8 | 9 | 8 | 10 | 7 |
| Flow Status - Flashiness (10) | 9 | 9 | 10 | 10 | 10 |
| Channel Alteration (20) | 19 | 19 | 20 | 15 | 20 |
| Frequency of Riffles/Bends (20)* | 18 | | 19 | | |
| Channel Sinuosity (20)** | | 12 | | 7 | 9 |
| Riparian and Bank Structure | | | | | |
| Bank Stability (L) (10) | 7 | 9 | 8 | 10 | 9 |
| Bank Stability (R) (10) | 8 | 9 | 8 | 10 | 9 |
| Vegetative Protection (L) (10) | 10 | 10 | 10 | 10 | 6 |
| Vegetative Protection (R) (10) | 10 | 10 | 10 | 10 | 6 |
| Riparian Veg. Zone Width (L) (10) | 5 | 10 | 10 | 10 | 4 |
| Riparian Veg. Zone Width (R) (10) | 7 | 10 | 10 | 10 | 4 |
| TOTAL SCORE (200): | 180 | 147 | 171 | 143 | 133 |
| | | | | | |
| HABITAT RATING | EXCELLENT | GOOD | EXCELLENT | GOOD | GOOD |
| | (NON- | (SLIGHTLY | (NON- | (SLIGHTLY | (SLIGHTLY |
| | IMPAIRED) | IMPAIRED) | IMPAIRED) | IMPAIRED) | IMPAIRED) |

Note: Individual metrics may better describe conditions directly affecting the biological community while the Habitat Rating describes the general riverine environment at the site(s).

| Date: | 6/7/2009 |) | 6/7/2009 | | 6/7/2009 | | 6/10/2009 | | 6/10/2009 | |
|-------------------------|---------------|---------|-----------------|----------|-----------|----------|-----------------|----------|---------------|----------|
| Weather: | Cloudy | , | Cloudy | | Rainy | | Partly Cloudy | | Partly Cloudy | |
| Air Temperature: | 54 | Deg. F | . 52 | Deg. F. | 49 | Deg. F. | | Deg. F. | | Deg. F. |
| Water Temperature: | 59 | Deg. F | . 50 | Deg. F. | 48 | Deg. F. | 46 | Deg. F. | | Deg. F. |
| Ave. Stream Width: | 16.5 | Feet | 18.5 | Feet | 15.6 | Feet | 14.5 | Feet | 16 | Feet |
| Ave. Stream Depth: | 0.38 | Feet | 0.77 | Feet | 0.44 | Feet | 0.72 | Feet | 0.76 | Feet |
| Surface Velocity: | 1.4 | Ft./Sec | 2.4 | Ft./Sec. | 2.5 | Ft./Sec. | 1.7 | Ft./Sec. | 1.4 | Ft./Sec. |
| Estimated Flow: | 8.778 | CFS | 34.188 | CFS | 17.16 | CFS | 17.748 | CFS | 17.024 | CFS |
| Stream Modifications: | t Improvement | | None | | None | C | anopy Removal | | None | |
| Nuisance Plants (Y/N): | Ň | | Ν | | Ν | | N | | Ν | |
| Report Number: | | | | | | | | | | |
| STORET No.: | 490068 | | 490059 | | 490002 | | 490199 | | 490104 | |
| Stream Name: | Furlong Creek | : 1 | Davenport Creek | | Cut River | D | avenport Creek | | Doe Creek | |
| Road Crossing/Location: | Pleasant Ave | | US-2 | | US-2 | | at gas pipeline | | Raski Road | |
| County Code: | 49 | | 49 | | 49 | | 49 | | 49 | |
| TRS: | 43N10W08 | | 42N08W02 | | 42N07W12 | | 43N08W26 | | 43N10W09 | |
| Latitude (dd): | 46.13899 | | 46.06772 | | 46.0452 | | 46.09928 | | 46.14193 | |
| Longitude (dd): | -85.59196 | | -85.26448 | | -85.1242 | | -85.2595 | | -85.56093 | |
| Ecoregion: | NLAI | 1 | NLAF | 1 | NLAF | | NLAF | | NLAF | 1 |
| Stream Type: | Warmwater | | Coldwater | | Coldwater | | Coldwater | | Warmwater | |
| USGS Basin Code: | 4060107 | | 4060107 | | 4060107 | | 4060107 | | 4060107 | |

* Applies only to Riffle/Run stream Surveys

** Applies only to Glide/Pool stream Surveys

| | Black River | Crow River |
|-----------------------------------|--------------------------------|---------------------|
| | State Forest Campground bridge | end of Summer Trail |
| | GLIDE/POOL | GLIDE/POOL |
| | STATION 36 | STATION 37 |
| HABITAT METRIC | | |
| Substrate and Instream Cover | | |
| Epifaunal Substrate/ Avail Cover | (20) 9 | 12 |
| Embeddedness (20)* | | |
| Velocity/Depth Regime (20)* | | |
| Pool Substrate Characterization (| 20)** 10 | 10 |
| Pool Variability (20)** | 12 | 13 |
| Channel Morphology | | |
| Sediment Deposition (20) | 9 | 11 |
| Flow Status - Maint. Flow Volum | ne (10) 7 | 9 |
| Flow Status - Flashiness (10) | 8 | 10 |
| Channel Alteration (20) | 20 | 20 |
| Frequency of Riffles/Bends (20)* | | |
| Channel Sinuosity (20)** | 17 | 17 |
| Riparian and Bank Structure | | |
| Bank Stability (L) (10) | 8 | 5 |
| Bank Stability (R) (10) | 8 | 5 |
| Vegetative Protection (L) (10) | 7 | 10 |
| Vegetative Protection (R) (10) | 7 | 10 |
| Riparian Veg. Zone Width (L) (1 | 0) 10 | 10 |
| Riparian Veg. Zone Width (R) (1 | 0) 10 | 10 |
| TOTAL SCORE (200): | 142 | 152 |

HABITAT RATING:

GOOD (SLIGHTLY IMPAIRED)

Note: Individual metrics may better describe conditions directly affecting the biological community while the Habitat Ratin describes the general riverine environment at the site(s)

| Date: | 6/8/2009 | | 6/8/2009 | |
|-------------------------|------------------|-----------------|---------------|----------|
| Weather: | Rainy | | Rainy | |
| Air Temperature: | 48 | Deg. F. | | Deg. F. |
| Water Temperature: | 48 | Deg. F. | 50 | Deg. F. |
| Ave. Stream Width: | 27 | Feet | 27.5 | Feet |
| Ave. Stream Depth: | 0.87 | Feet | 1.4 | Feet |
| Surface Velocity: | 1 | Ft./Sec. | 0.87 | Ft./Sec. |
| Estimated Flow: | 23.49 | CFS | 33.495 | CFS |
| Stream Modifications: | None | | None | |
| Nuisance Plants (Y/N): | N | | N | |
| Report Number: | | | | |
| STORET No.: | 490198 | | 490197 | |
| Stream Name: | Black River | | Crow River | |
| Road Crossing/Location: | State Forest Car | npground bridge | end of Summer | Trail |
| County Code: | 49 | | 49 | |
| TRS: | 43N09W13 | | 42N11W13 | |
| Latitude (dd): | 46.11758 | | 46.03036 | |
| Longitude (dd): | -85.36597 | | -85.61533 | |
| Ecoregion: | NLAF | | NLAF | |
| Stream Type: | Coldwater | | Coldwater | |
| USGS Basin Code: | 4060107 | | 4060107 | |

* Applies only to Riffle/Run stream Survey: ** Applies only to Glide/Pool stream Survey:

GOOD (SLIGHTLY IMPAIRED)

Table 5. Analytical results for 2009 Tahquamenon, Two-Hearted, Manistique, and Millecoquins Rivers watersheds sampling sites.

| Location Station STORET ID # Parameter | Units | Doe Creek Station 35 Raski Road 490104 | Doe Creek Station 36 Kovar Rd 490202 | Ν | Doe Creek Station 37 /iile/Pleasar 490070 | nt | Doe Creek Station 38 M-117 490053 | Doe Creek Station 39 Krause Road 490069 | Γ | Millecoquins River Station 40 Lower Rapids 490196 | En | Crow River Station 41 nd Simmon Tra 490197 | u, il | /s Newberry WWTP Station 42 M-123 crossing 480002 | Newberry WWTP Effluent Station 43 Outfall 001A 480039 |
|---|--------|---|---|---|--|----|--|--|---|--|----|---|----------|---|--|
| Ammonia | mg N/L | 0.016 | 0.021 | | 0.023 | | 0.015 | 0.012 | | 0.011 | | 0.01 | | 0.02 | 15 |
| Boron | ug/L | - | - | | - | | - | - | | - | | - | | - | - |
| Calcium | mg/L | - | - | | - | | - | - | | - | | - | | 20.3 | 64.1 |
| Chromium | ug/L | - | - | | - | | - | - | | - | | - | | ND (< 1) | ND (< 1) |
| Hardness - Calculated | mg/L | - | - | | - | | - | - | | - | | - | | - | 229 |
| Iron | ug/L | - | - | | - | | - | - | | - | | - | | - | - |
| Lithium | ug/L | - | - | | - | | - | - | | - | | - | | - | - |
| Magnesium | mg/L | - | - | | - | | - | - | | - | | - | | 5.3 | 16.8 |
| Mercury | ug/L | - | - | | - | | - | - | | - | | - | | ND (<0.2) | ND (<0.2) |
| Nitrate + Nitrite | mg N/L | 0.013 | 0.013 | | 0.086 | | 0.005 | 0.046 | | 0.005 | | 0.059 | | ND (< 0.01) | ND (<0.2) |
| Nitrogen - Kjeldahl | mg N/L | 0.87 | 0.81 | | 0.85 | | 0.87 | 0.66 | | 0.43 | | 0.31 | | 0.63 | 17 |
| Phosphorus | mg P/L | 0.06 | 0.03 | | 0.036 | | 0.064 | 0.034 | | 0.024 | | 0.01 | | 0.027 | 0.56 |
| Potassium | mg/L | - | - | | - | | - | - | | - | | - | | - | - |
| Selenium | ug/L | - | - | | - | | - | - | | - | | - | | ND (< 1) | ND (< 1) |
| Sodium | mg/L | - | - | | - | | - | - | | - | | - | | - | - |
| TOC | mg/L | 20 | 19 | | 22 | | 21 | 18 | | 9.3 | | 6.1 | | 18 | 13 |
| COD | mg/L | 46 | 46 | | 50 | | 49 | 38 | | 19 | | 12 | | 38 | 42 |
| Arsenic | ug/L | - | - | | - | | - | - | | - | | - | | 1.0 | ND (<1) |
| Barium | ug/L | - | - | | - | | - | - | | - | | - | | 27 | 39 |
| Cadmium | ug/L | - | - | | - | | - | - | | - | | - | | ND (0.2) | ND (<0.2) |
| Copper | ug/L | - | - | | - | | - | - | | - | | - | | ND (< 1) | 9.9 |
| Lead | ug/L | - | - | | - | | - | - | | - | | - | | ND (< 1) | ND (< 1) |
| Nickel | ug/L | - | - | | - | | - | - | | - | | - | | - | - |
| Silver | ug/L | - | - | | - | | - | - | | - | | - | | ND (<0.2) | ND (< 0.2) |
| Zinc | ug/L | - | - | | - | | - | - | | - | | - | | ND (< 10) | 28 |

ND - Value reported is less than level of detection. Metal values given are as total metals.