



MICHIGAN DEPARTMENT OF NATURAL RESOURCES  
Wildlife Division Report No. 3670  
March 2019

## 2017 MICHIGAN FURBEARER HARVEST SURVEY

Brian J. Frawley

### ABSTRACT

*A sample of furtakers was contacted after the 2017 hunting and trapping seasons to estimate the number of participants, days afield (effort), and furbearer harvests. In 2017, 22,981 people purchased a fur harvester license, which was 11% fewer than in 2016. About 11,397 license buyers either hunted or trapped furbearers in 2017. About 30% of the license buyers trapped (6,879 trappers), 32% hunted (7,331 hunters), and 12% (2,814) both trapped and hunted. The number of active furtakers in 2017 declined significantly by 11% from the number in 2016. The number of furtakers pursuing most species also declined between 2016 and 2017; however, the declines were significant only among hunters pursuing raccoon (-21%) and bobcat (-12%). Changes for hunting and trapping effort and harvest between 2016 and 2017 generally followed changes in the number of furtakers, although most of these declines were not significant. Hunters most commonly sought coyotes, raccoons, and bobcats, while trappers most frequently sought raccoons, coyotes, and muskrats. Trends in harvest can be affected by both changes in furtaker and furbearer numbers; thus, harvest per furtaker was examined for trends. The mean number of raccoon and opossum taken per furtaker has increased since the 1950s. The mean harvest of red fox by both hunters and trappers has declined since the mid-1980s. These trends suggest raccoon and opossum may have been increasing in abundance since their harvest has been monitored, while red fox numbers may have been declining. An estimated 171 trappers caught and released 237 bobcats that were caught in traps set for another species in 2017. In addition, these trappers registered an additional 19 incidental bobcats. Hunters and trappers combined spent an average of \$431 per year pursuing furbearers. Collectively, furtakers spent about \$4,910,000 hunting and trapping furbearers.*



A contribution of Federal Aid in Wildlife Restoration, Michigan Project W-147-R

#### Equal Rights for Natural Resource Users

The Michigan Department of Natural Resources provides equal opportunities for employment and access to Michigan's natural resources. Both State and Federal laws prohibit discrimination on the basis of race, color, national origin, religion, disability, age, sex, height, weight or marital status under the U.S. Civil Rights Acts of 1964 as amended, 1976 MI PA 453, 1976 MI PA 220, Title V of the Rehabilitation Act of 1973 as amended, and the 1990 Americans with Disabilities Act, as amended.

If you believe that you have been discriminated against in any program, activity, or facility, or if you desire additional information, please write:  
Human Resources, Michigan Department of Natural Resources, PO Box 30028, Lansing MI 48909-7528, or  
Michigan Department of Civil Rights, Cadillac Place, 3054 West Grand Blvd, Suite 3-600, Detroit, MI 48202, or  
Division of Federal Assistance, U.S. Fish & Wildlife Service, 4401 North Fairfax Drive, Mail Stop MBSP-4020, Arlington, VA 22203.

For information or assistance on this publication, contact Michigan Department of Natural Resources, Wildlife Division, P.O. Box 30444, MI 48909.  
This publication is available in alternative formats upon request.

## INTRODUCTION

The Department of Natural Resources (DNR) has the authority and responsibility to protect and manage wildlife resources in Michigan, while the Natural Resources Commission (NRC) has the authority to regulate the taking of game (Natural Resources and Environmental Protection Act, Public Act 451 of 1994). Harvest surveys are one of the management tools used by the DNR to accomplish its statutory responsibility. Estimating harvests and hunter participation are the primary objectives of these surveys. Information from harvest surveys, mandatory registration, and other indices are used to monitor furbearer populations and help establish harvest regulations.

The primary furbearing animals harvested for their pelts in Michigan during recent years have been badger (*Taxidea taxus*), beaver (*Castor canadensis*), bobcat (*Lynx rufus*), coyote (*Canis latrans*), fisher (*Martes pennanti*), gray fox (*Urocyon cinereoargenteus*), marten (*Martes americana*), mink (*Mustela vison*), muskrat (*Ondatra zibethica*), opossum (*Didelphis virginiana*), raccoon (*Procyon lotor*), red fox (*Vulpes vulpes*), river otter (*Lontra canadensis*), striped skunk (*Mephitis mephitis*), and weasels (*Mustela* spp.) (Frawley 2017). Coyote, opossum, weasels, and skunks could be taken year-round with a fur harvester license. The remaining furbearers could be harvested in 2017 during late fall through spring by a person possessing a fur harvester license (Table 1); however, nonresidents could not trap badger, bobcat, fisher, marten, or otter.

Landowners or their designees could take raccoons, coyotes, and skunks throughout the year on their property without a license if these animals were doing or about to do damage. Coyotes could also be taken by resident hunters possessing a base license. In addition, a mentored hunting program was started in 2012. Under this program, a mentored youth hunting license was created and could be purchased by youth hunters aged 9 and younger. The youth hunter had to participate with a mentor who was at least 21 years old. The mentored youth hunting license allowed the youth hunter to hunt small game, turkey, deer, trap furbearers, and fish for all species. Hunters taking furbearers on their own land without a license, or taking furbearers with either a base license or a mentored youth hunting license were not included in our sample. Thus, harvest estimates from this survey do not represent all possible forms of harvest, but only those taken by people with a fur harvester license.

## METHODS

Following the 2017 hunting and trapping seasons, a questionnaire (Appendix A) was sent to a random sample of people (4,996) who had purchased a fur harvester license (Table 2). This level of sampling should produce statewide estimates with a margin of error of less than 20% for the most commonly pursued species. All licensees had an equal chance of being included in the random sample. After the sample was selected, licensees were grouped into one of four strata on the basis of their residence. These strata included residents of the Upper Peninsula (UP), Northern Lower Peninsula (NLP), Southern Lower Peninsula (SLP), and nonresidents (Figure 1). People receiving the questionnaire were asked to report whether they pursued furbearers, number of days spent afield, and whether they harvested any furbearing animals.

Estimates were calculated using a stratified random sampling design (Cochran 1977). Using stratification, furtakers were placed into similar groups (strata) based on their county of residence. Residents of the UP, NLP, SLP, and nonresidents and licensees with unknown residency were grouped into separate strata (Figure 1). The overall sample consisted of 641 people from the UP stratum (N= 2,970), 1,078 people from the NLP stratum (N= 5,030), 3,195 from the SLP stratum (N= 14,520), and 86 people from the nonresident and unknown residency stratum (N=461). Estimates were derived for each group separately. The statewide estimate was then derived by combining group estimates so the influence of each group matched the proportion its members represented in the statewide population of furtakers. The primary reason for using a stratified sampling design was to produce more precise estimates. Improved precision means similar estimates should be obtained if this survey was repeated.

Estimates were subject to both sampling and nonsampling error. When a sample rather than the entire population has been surveyed, there is a chance that the sample estimates may differ from the true population values they represent. The difference, or sampling error, varies depending on the particular sample selected, and this variability was measured by the 95% confidence limit (CL). In theory, this CL can be added and subtracted from the estimate to calculate the 95% confidence interval. The confidence interval was a measure of the precision associated with the estimate and implies the true value would be within this interval 95 times out of 100.

Estimates also were affected by nonsampling error. The nonsampling error could occur for many reasons, including the failure to include a segment of the survey population, the inability to obtain data from all units in the sample, the inability or unwillingness of respondents to provide data, mistakes made by respondents, and errors made in the collection or processing of the data. It is very difficult to measure this error. Thus, estimates were not adjusted for nonsampling error. Furthermore, harvest estimates did not include animals taken legally outside the open season (e.g., nuisance animals).

Statistical tests are used routinely to determine the likelihood the differences among estimates are larger than expected by chance alone. The overlap of 95% confidence intervals was used to determine whether estimates differed. Non-overlapping 95% confidence intervals were equivalent to stating the difference between the means was larger than would be expected 95 out of 100 times ( $P < 0.05$ ), if the study had been repeated (Payton et al. 2003).

Estimates of events that occur infrequently are difficult to estimate precisely using common sampling designs (Cochran 1977). Relatively few furtakers harvest river otter, bobcat, badger, fisher, and marten; thus, some estimates associated with these species should be viewed cautiously. More precise harvest estimates were obtained for these species through tallying registration reports. All furtakers harvesting a river otter, bobcat, fisher, or marten were required to present these animals at a DNR office for registration. Prior to 2003, furtakers were also required to register badger; however, this requirement was eliminated in 2003. In this report, the marten harvest was determined only by registration.

During recent years, all licensed furtakers attempting to harvest bobcat, fisher, marten, and otter in Michigan were required to obtain a free harvest tag from the DNR. The list of furtakers obtaining these harvest tags formed a complete list of statewide trappers pursuing these species. Using these lists, the DNR was able to design separate harvest surveys that

provided more precise estimates (i.e., narrower confidence intervals) than previous surveys of all furtakers. Separate surveys were conducted to estimate furtaker participation, harvest, and effort for bobcat (Frawley 2019c), fisher and marten (Frawley 2019a), and otter (Frawley 2019b) seasons during recent years.

While the primary objectives of the fur harvesters' survey were estimating harvest, number of participants, and trapping and hunting effort, this survey also provided an opportunity to collect information about management issues. Questions were added to the questionnaire to determine furtakers satisfaction with furbearer numbers, animals harvested, and overall hunting or trapping experience. In addition, furtakers were asked to report how much they spent on things related to hunting and trapping furbearers (e.g., fuel, food, lodging, equipment, and ammunition) during 2017 seasons. Trappers also were asked whether they caught any bobcats incidentally in traps set for another species.

## **RESULTS AND DISCUSSION**

Questionnaires were mailed initially in late April 2017. Up to two follow-up questionnaires were sent to non-respondents. Questionnaires were undeliverable to 68 people, primarily because of changes in residence. Questionnaires were returned by 2,468 people, yielding a 50% adjusted response rate.

In 2017, 22,982 fur harvester licenses were purchased by 22,981 people (Figure 2, Table 2). The number of license buyers in 2017 was 11% fewer than in 2016. Most license buyers were men (97%), with an average age of 48 years (Figure 3). About 4% of the license buyers (843) were younger than 17 years of age. Furtakers less than 10 years of age using a mentored youth license were not included in the analyses.

Compared to 10 years ago, the number of people buying a fur harvester license in 2017 decreased by about 5% (24,296 people purchased a license in 2007). In addition, there were fewer license buyers for most age classes between 11 and 54 years of age in 2017, compared to 2007 (Figure 4). However, there were increased furtakers among the oldest age classes in 2017. The increased furtakers in the oldest age classes likely represented the rising share of older people in the population as the baby-boom generation aged and life expectancies have increased.

### **Mail Harvest Survey**

Overall, approximately 50% of license buyers either hunted or trapped furbearers during 2017 (Table 3). The number of active furtakers in 2017 declined significantly by 11% from the number of furtakers in 2016. About 30% of the license buyers trapped and 32% hunted furbearers during 2017. Trappers most often pursued raccoon, coyote, and muskrat (Table 4). Hunters most commonly sought coyote, raccoon, and bobcats. Coyotes and raccoons also ranked as the most frequently sought furbearers when trappers and hunters were combined.

The estimated number of hunters and trappers statewide has declined gradually since 2013, although current estimates are well above the lowest estimates reported during the mid-

1990s (Figure 5). Recent changes in furtaker numbers have paralleled declining fur prices (e.g., Dhuey 2018, Rees 2015, Conlee and Johnston 2018, Evelsizer 2018). Historically, the peaks in furtaker numbers corresponded closely to periods when pelt values peaked for many species such as muskrat, raccoon, and red fox (Evelsizer 2018, Conlee and Johnston 2018). Between 1999 and 2012, the number of people hunting furbearers was greater than the number of people trapping; however, the number of trappers and hunters was nearly equal the last six years (Figure 5).

The number of furtakers pursuing most furbearers declined between 2016 and 2017 (Table 4); however, the declines were significant only among hunters pursuing raccoon (-21%) and bobcat (-12%). Changes for hunting and trapping effort and harvest between 2016 and 2017 generally followed changes in the number of furtakers, although most of these declines were not significant (Table 4).

Harvest for most species in 2017 was near the low end of their historical ranges, except coyote, opossum, otter, and raccoon (Figures 6-8). Many factors influence harvest trends such as furtaker numbers, wildlife population size, harvest regulations, weather, habitat conditions, and fur prices; thus, any interpretations of trends should be viewed cautiously.

Trends in harvest per furtaker (Figures 9 and 10) were examined because this measure may eliminate some of the effects of changing furtaker and furbearer numbers over time, although many other factors may still complicate interpretations of these trends (Poole and Mowat 2001). The mean number of raccoon taken per trapper has generally increased since the early 1950s (Figures 9 and 10). The mean harvest of fox by both hunters and trappers has declined since the mid-1980s. These trends suggest raccoon may have been increasing in abundance since harvest has been monitored, while red fox numbers may have been declining.

These trends in furbearer numbers are not unique to Michigan. Increasing raccoon numbers and declining red fox numbers have been reported in many Midwestern states since the 1980s (e.g., Gehrt et al. 2002, Conlee and Johnston 2018, Evelsizer 2018). The decline in red fox numbers has been attributed largely to competition from increased coyote and bobcat (Sovada et al. 1995, Conlee and Johnston 2018, Evelsizer 2018). Gray fox numbers may also have been reduced because of the distemper virus associated with raccoons (Conlee and Johnston 2018).

The mean harvest of fisher and bobcat per trapper has declined during the last twenty years (Figure 9). Frawley (2019a) also reported increasing effort expended by trappers for each fisher registered during the last ten years. Both the declining mean harvest of fisher per trapper and the increasing effort per registered fisher suggest fisher numbers may have declined over the last twenty years. Using fisher trapper effort data with harvest at age information, researchers reported a 70% decline in fisher abundance in the Upper Peninsula (unpublished data; J.R. Skalski, School of Aquatic & Fishery Sciences, University of Washington, Seattle). The seasonal harvest limit for fisher was lowered from three to a combined bag limit of one fisher or one marten in 2011, and this reduction likely contributed to the decline in fisher taken per trapper in recent years (Frawley 2019a).

The mean number of bobcats taken per trapper declined from 2003 to 2017 (Figure 9). The seasonal harvest limit for bobcats was lowered from three to two bobcats in 2004, and the UP hunting and trapping season lengths were reduced in 2009, which likely contributed to the decline of bobcats taken per trapper since 2003 (Frawley 2019c).

## **Registration Data**

Compared to 2016, fewer otter (-6%) and fisher (-5%) were registered in 2017, but increased numbers of marten (20%) and bobcat (10%) were taken (Figure 11, Table 5). Registration totals excluded harvest by tribal members. In addition, registration totals only included animals that were registered and returned to the furtaker.

## **Incidental Capture of Bobcats**

An estimated 171 trappers caught a bobcat incidentally in a trap set for another species (Table 6). These trappers caught 237 incidental bobcats that were released alive from their traps. In addition, trappers caught an estimated 19 incidental bobcats that were registered. Because incidental bobcats could be captured more than once, the estimate of incidental bobcats caught by trappers probably does not represent unique bobcats.

## **Beaver Trapping Activity by Otter Trappers**

In order to trap otter, trappers were required to obtain a free otter harvest tag in addition to a fur harvester license. A separate survey was sent to these trappers obtaining an otter harvest tag to estimate their trapping activity (Frawley 2019b). Because otter trappers frequently sought beaver, these trappers also were asked to report information about their beaver trapping activity. However, these estimates associated with beaver trapping only represent the participation, effort, or harvest of trappers that obtained an otter harvest tag. In order to put these estimates into a broader perspective, it is important to know what proportion of beaver trapping activity was attributed to trappers having an otter harvest tag.

An estimated 1,939 furtakers sought beavers (Tables 4 and 7). About 59% of these trappers possessed an otter harvest tag (Table 7), and they were responsible for an estimated 75% of the beaver taken.

## **Furtaker Satisfaction**

Furtakers were asked to identify the furbearer species they primarily sought, and then report how satisfied they were the number of animals seen, number of animals taken, and their overall hunting or trapping experience for this primary species. At least 50% of furtakers were either very satisfied or somewhat satisfied with the number of raccoon, fox, coyote, fisher, muskrat, and beaver seen during 2017 (Table 8). Over 50% of furtakers seeking raccoon, muskrat, and beaver were satisfied with the number of animals taken; otherwise, less than 50% of furtakers were satisfied with the number of animals they harvested (Table 9). Over 50% of furtakers pursuing all species were either very satisfied or somewhat satisfied with their overall hunting or trapping experiences (Table 10). Fox and fisher were the species with the lowest levels of overall satisfaction.

## **Expenditures by Furtakers**

The average furtaker devoted  $28.9 \pm 1.9$  days hunting or trapping furbearers and spent an average of  $\$431 \pm \$44$  in 2017. Expenditures included the costs of fuel, food, lodging, equipment, and ammunition. Collectively, furtakers spent about  $\$4,910,000 (\pm \$502,000)$  on hunting and trapping furbearers in the 2017 seasons.

## **ACKNOWLEDGEMENTS**

I thank all the furtakers that provided information. Theresa Riebow completed data entry, and Marshall Strong prepared Figure 1. Adam Bump and Dwayne Etter reviewed a draft version of this report.

## **LITERATURE CITED**

- Cochran, W. G. 1977. Sampling techniques. John Wiley & Sons, New York. USA.
- Conlee, L. and S. Johnston. 2018. 2017 furbearer program annual report. Unpublished report. Missouri Department of Conservation, Jefferson City, USA.
- Dhuey, B. 2018. Wisconsin fur buyers report, 2017-2018. Unpublished report. Wisconsin Department of Natural Resources, Madison, USA.
- Evelsizer, V. 2018. Trends in Iowa wildlife populations and harvest – 2017-2018. Iowa Department of Natural Resources, Des Moines, Iowa, USA.
- Frawley, B. J. 2017. 2015 Michigan furbearer harvest survey. Wildlife Division Report 3637. Michigan Department of Natural Resources, Lansing, USA.
- Frawley, B. J. 2019a. 2017 marten and fisher harvest survey. Wildlife Division Report 3663. Michigan Department of Natural Resources, Lansing, USA.
- Frawley, B. J. 2019b. 2017 Michigan otter and beaver harvest survey. Wildlife Division Report 3668. Michigan Department of Natural Resources, Lansing, USA.
- Frawley, B. J. 2019c. 2017 bobcat hunter and trapper harvest in Michigan. Wildlife Division Report 3665. Michigan Department of Natural Resources, Lansing, USA.
- Gehrt, S. D., G. F. Huber, and J. A. Ellis. 2002. Long-term population trends of raccoons in Illinois. Wildlife Society Bulletin 30:457-463.
- Payton, M. E., M. H. Greenstone, and N. Schenker. 2003. Overlapping confidence intervals or standard error intervals: what do they mean in terms of statistical significance? Journal of Insect Science 3:34.

Poole, K. G. and G. Mowat. 2001. Alberta furbearer harvest data analysis. Alberta Sustainable Resource Development, Fish and Wildlife Division, Alberta Species at Risk Report No. 31. Edmonton, Alberta, Canada.

Rees, J. 2015. Wisconsin fur buyers report, 2014-2015. Unpublished report. Wisconsin Department of Natural Resources, Madison, USA.

Sovada, M. A., A. B. Sargeant, and J. W. Grier. 1995. Differential effects of coyotes and red foxes on duck nest success. *Journal of Wildlife Management* 59:19.






Figure 1. Stratum boundaries used for the analysis of the Michigan furbearer harvest survey.




Figure 2. Number of fur harvester licenses sold in Michigan, 1986-2017.




Figure 3. Ages of people that purchased a license to hunt or trap furbearers in Michigan for the 2017 hunting and trapping seasons (mean = 48 years).



Figure 4. Number of fur harvester license buyers in Michigan by age and sex during 2005 and 2017 hunting seasons. The number of people buying a license (all ages combined) was 24,296 in 2005 and 22,981 in 2017.




Figure 5. Estimated number of furtakers (trappers and hunters) in Michigan, 1957-2017. Estimates included only license buyers that actually trapped or hunted furbearers (any species). Data were not available for all years.




Figure 6. Estimated furbearer harvest by trappers and the number of trappers in Michigan estimated from mail harvest surveys, 1957-2017. Mail survey questionnaires were sent to a random sample of Trapping license buyers during 1957-1969. The sample also included Sportsman's license buyers in 1970-1972. During 1980-1983, the sample included Trapping and Senior Hunting license buyers. During 1986-2013, the sample was selected from people buying either Resident Fur Harvester, Senior Fur Harvester, Junior Fur Harvester, Military Fur Harvester, or Nonresident Fur Harvester licenses. The sample also included Senior Hunting license buyers during 1986-1988. During 1996-2013, samples also included people buying Resident Fur Harvester (trap only) and Junior Fur Harvester (trap only) licenses. Starting in 2014, license types were consolidated into a fur harvesters license type. Data were not available for all years.




Figure 6 (Continued). Estimated furbearer harvest by trappers and the number of trappers in Michigan estimated from mail harvest surveys, 1957-2017. Mail survey questionnaires were sent to a random sample of Trapping license buyers during 1957-1969. The sample also included Sportsman's license buyers in 1970-1972. During 1980-1983, the sample included Trapping and Senior Hunting license buyers. During 1986-2013, the sample was selected from people buying either Resident Fur Harvester, Senior Fur Harvester, Junior Fur Harvester, Military Fur Harvester, or Nonresident Fur Harvester licenses. The sample also included Senior Hunting License buyers during 1986-1988. During 1996-2013, samples also included people buying Resident Fur Harvester (trap only) and Junior Fur Harvester (trap only) licenses. Starting in 2014, license types were consolidated into a fur harvesters license type. Data were not available for all years.




Figure 7. Estimated furbearer harvest by trappers and the number of trappers in Michigan estimated from mail harvest surveys, 1980-2017. The mail survey was sent to a random sample of Trapping and Senior Hunting license buyers during 1980-1983. During 1986-2013, the sample was selected from people buying either Resident Fur Harvester, Senior Fur Harvester, Junior Fur Harvester, Military Fur Harvester, or Nonresident Fur Harvester licenses. The sample also included Senior Hunting license buyers during 1986-1988. During 1996-2013, samples also included people buying Resident Fur Harvester (trap only) and Junior Fur Harvester (trap only) licenses. Starting in 2014, license types were consolidated into a fur harvesters license type. Data were not available for all years.






Figure 7 (Continued). Estimated furbearer harvest by trappers and the number of trappers in Michigan estimated from mail harvest surveys, 1980-2017. The mail survey was sent to a random sample of Trapping and Senior Hunting license buyers during 1980-1983. During 1986-2013, the sample was selected from people buying either Resident Fur Harvester, Senior Fur Harvester, Junior Fur Harvester, Military Fur Harvester, or Nonresident Fur Harvester licenses. The sample also included Senior Hunting license buyers during 1986-1988. During 1996-2013, samples also included people buying Resident Fur Harvester (trap only) and Junior Fur Harvester (trap only) licenses. Starting in 2014, license types were consolidated into a fur harvesters license type. Data were not available for all years.




Figure 8. Estimated furbearer harvest by hunters and the number of hunters in Michigan estimated from mail harvest surveys, 1980-2017. The mail survey was sent to a random sample of people buying either small game licenses, Senior Hunting licenses, or Sportsman's licenses during 1980-1985. During 1986-2013, the sample was selected from people buying either Resident Fur Harvester, Senior Fur Harvester, Junior Fur Harvester, Military Fur Harvester, or Nonresident Fur Harvester licenses. The sample also included Senior Hunting license buyers during 1986-1988. During 1996-2013, samples also included people buying Resident Fur Harvester (trap only) and Junior Fur Harvester (trap only) licenses. Starting in 2014, license types were consolidated into a fur harvesters license type. Data were not available for all years.






Figure 8 (Continued). Estimated furbearer harvest by hunters and the number of hunters in Michigan estimated from mail harvest surveys, 1980-2017. The mail survey was sent to a random sample of people buying either small game licenses, Senior Hunting licenses, or Sportsman's licenses during 1980-1985. During 1986-2013, the sample was selected from people buying either Resident Fur Harvester, Senior Fur Harvester, Junior Fur Harvester, Military Fur Harvester, or Nonresident Fur Harvester licenses. The sample also included Senior Hunting license buyers during 1986-1988. During 1996-2013, samples also included people buying Resident Fur Harvester (trap only) and Junior Fur Harvester (trap only) licenses. Starting in 2014, license types were consolidated into a fur harvesters license type. Data were not available for all years.





Figure 9. Mean number of furbearers harvested annually per trapper in Michigan estimated from mail harvest surveys, 1954-2017. Data were not available for all years.





Figure 9 (continued). Mean number of furbearers harvested annually per trapper in Michigan estimated from mail harvest surveys, 1954-2017. Data were not available for all years.







Figure 9 (continued). Mean number of furbearers harvested annually per trapper in Michigan estimated from mail harvest surveys, 1954-2017. Data were not available for all years.





Figure 9 (continued). Mean number of furbearers harvested annually per trapper in Michigan estimated from mail harvest surveys, 1954-2017. Data were not available for all years.







Figure 10. Mean number of furbearers harvested annually per hunter in Michigan estimated from mail harvest surveys, 1954-2017. Data were not available for all years.




Figure 10 (continued). Mean number of furbearers harvested annually per hunter in Michigan estimated from mail harvest surveys, 1954-2017. Data were not available for all years.




Figure 11. Number of bobcat, otter, fisher, badger, and marten registered by furtakers in Michigan, 1985-2017. Badger and fisher seasons were established in 1989, and marten season started in 2000. Totals for 2013 were preliminary. Beginning in 2003, badger were no longer registered. Registration totals only included animals that were registered and returned to the furtaker and excluded harvest by tribal members.



Table 1. Trapping and hunting seasons when furbearing animals could be harvested in Michigan during 2017 seasons.<sup>a</sup>

Season, species, and area	Season dates
Trapping seasons <sup>b</sup>	
Muskrat and Mink	
UP	October 25 – March 1
NLP	November 1 – March 1
SLP	November 10 – March 1
Raccoon	
Statewide	October 15 – March 31
Fox and Coyote	
Statewide	October 15 – March 1
Bobcat <sup>c</sup>	
UP (units A and B)	December 1 – February 1
NLP (units C, D, E, and F)	December 10 – 20
Badger <sup>c</sup>	
UP and NLP	October 15 – November 14
SLP	November 1 – March 1
Fisher and Marten <sup>c</sup>	
UP	December 1 – 15
Beaver and Otter <sup>c,d</sup>	
UP	October 25 – April 15
NLP	November 1 – April 15
SLP	November 10 – March 31
Hunting seasons	
Bobcat <sup>c</sup>	
UP (units A B and C)	January 1 – March 1
NLP (Unit D)	January 1 – February 1
NLP (units E and F)	January 1 – 11
Fox	
Statewide	October 15 – March 1
Raccoon	
Statewide	October 1 – January 31
Coyote	
Statewide	Year-round

<sup>a</sup>No closed season for opossum, weasel, and skunk.

<sup>b</sup>Nonresidents may trap from November 15 through the regular season closing date, except nonresidents could not trap badger, bobcat, fisher, marten, or otter. In addition, the opening date for nonresident beaver trapping varied by area.

<sup>c</sup>No nonresident season existed for badger, bobcat, fisher, marten, and otter.

<sup>d</sup>Resident seasons only. Nonresident beaver season occurred during November 15-April 16 (UP), November 24-April 16 (NLP), and December 15 – March 31 (SLP).

Table 2. Number of fur harvester licenses sold and people receiving and returning harvest questionnaire, 2014-2017.

Item	Year			
	2014	2015	2016	2017
Licenses sold <sup>a</sup>	28,035	26,873	25,948	22,982
Individuals buying licenses <sup>a,b</sup>	28,029	26,865	25,938	22,981
Mentored youth license buyers <sup>c</sup>	12,915	11,929	11,342	11,691
Questionnaires mailed	4,200	4,200	4,200	5,000
Non-deliverable questionnaires	69	85	66	68
Questionnaires returned	2,329	2,267	2,269	2,468
Questionnaires returned (%) <sup>d</sup>	56	55	55	50

<sup>a</sup>License types included Fur Harvester, Senior Fur Harvester, and Lifetime Fur Harvester.

<sup>b</sup>A person was counted only once, regardless of how many licenses they purchased.

<sup>c</sup>The mentored youth hunting license was created in 2012 and was valid for hunting small game, waterfowl, turkey, and deer. These youth could also trap furbearers and fish all species. Although these license buyers were eligible to take furbearers, they were not included in survey sample.

<sup>d</sup>Response rate adjusted to exclude non-deliverable questionnaires.

Table 3. Estimated number of fur harvester license buyers who trapped or hunted furbearers in Michigan, 2015-2017.

Activity	2015		2016		2017		Change between 2016 and 2017 (%)
	Estimate	95% CL	Estimate	95% CL	Estimate	95% CL	
<b>Trapped</b>							
Number	9,077	500	7,525	462	6,879	391	-9
%	34	2	29	2	30	2	1
<b>Hunted</b>							
Number	9,392	505	8,266	473	7,331	399	-11*
%	35	2	32	2	32	2	0
<b>Trapped or hunted<sup>a</sup></b>							
Number	14,792	528	12,762	510	11,397	429	-11*
%	55	2	49	2	50	2	0
<b>Trapped only</b>							
Number	5,400	424	4,496	386	4,066	327	-10
%	20	2	17	1	18	1	0
<b>Hunted only</b>							
Number	5,715	434	5,237	408	4,518	340	-14
%	21	2	20	2	20	1	-1
<b>Trapped and hunted</b>							
Number	3,676	364	3,029	327	2,814	280	-7
%	14	1	12	1	12	1	1

<sup>a</sup>A person was counted only once, although they may have both trapped and hunted furbearers.

\*Non-overlapping 95% confidence intervals indicated estimates differed significantly between 2016 and 2017 (P<0.05).

Table 4. Estimated number of participants, harvest, and days afield during Michigan furbearer seasons, 2016 and 2017.

Species and season	Participants (No.)				Harvest (No.)				Days afield (No.)			
	Year		95% CL <sup>a</sup>	Change (%)	Year		95% CL <sup>a</sup>	Change (%)	Year		95% CL <sup>a</sup>	Change (%)
	2016	2017			2016	2017			2016	2017		
<b>Trapping</b>												
Mink	1,716	1,699	223	-1	4,663	5,176	1,460	11	42,098	46,728	8,545	11
Raccoon	4,281	3,927	321	-8	39,797	43,391	8,015	9	96,475	102,091	12,825	6
Opossum	1,895	1,782	228	-6	15,474	12,143	2,811	-22	52,806	46,325	8,907	-12
Skunk	1,227	922	168	-25	4,360	3,093	1,357	-29	32,954	28,291	7,637	-14
Weasel	416	362	106	-13	1,028	589	307	-43	6,819	9,468	4,126	39
Red fox	1,833	1,714	225	-7	3,624	3,472	869	-4	49,566	50,735	9,854	2
Gray fox	665	650	142	-2	353	350	176	-1	15,900	19,024	5,404	20
Coyote	3,541	3,219	297	-9	8,881	7,449	1,780	-16	96,711	87,204	12,469	-10
Bobcat <sup>b</sup>	1,129	1,185	98	5	161	229	50	42	14,570	13,196	1,566	-9
Beaver <sup>c</sup>	2,207	1,939	239	-12	13,531	12,172	3,127	-10	40,210	38,121	7,185	-5
Muskrat	2,837	2,643	272	-7	103,411	108,237	25,580	5	64,050	68,815	10,148	7
Otter <sup>c</sup>	979	942	60	-4	909	893	82	-2	17,425	16,003	1,705	-8
Fisher <sup>d</sup>	581	517	41	-11	203	149	23	-26*	5,011	4,585	450	-8
Badger	140	84	52	-40	105	38	35	-64	1,263	791	789	-37
<b>Hunting</b>												
Raccoon	2,784	2,197	250	-21*	34,127	34,544	11,601	1	50,177	39,891	8,039	-21
Red fox	2,261	1,782	230	-21	1,404	1,215	374	-13	26,276	28,852	7,003	10
Gray fox	994	734	150	-26	139	102	61	-27	10,643	13,027	4,819	22
Coyote	6,497	5,948	376	-8	9,112	13,501	4,031	48	79,906	84,221	11,252	5
Bobcat <sup>b</sup>	1,846	2,058	122	12*	260	298	54	14	15,136	16,248	1,874	7
<b>Trapping and hunting combined</b>												
Raccoon	6,029	5,247	357	-13	73,923	77,935	14,309	5	146,652	141,982	16,067	-3
Red fox	3,694	3,099	293	-16	5,028	4,687	964	-7	75,842	79,587	12,675	5
Gray fox	1,556	1,226	192	-21	492	452	188	-8	26,543	32,052	7,984	21
Coyote	8,588	7,713	405	-10	17,993	20,950	4,431	16	176,617	171,425	17,348	-3
Bobcat <sup>b</sup>	2,693	2,956	136	10*	422	527	74	25	29,706	29,444	2,476	-1

<sup>a</sup>95% CL for the 2017 estimate.

<sup>b</sup>Bobcat estimates from separate mail harvest survey (Frawley 2019c). See Table 5 for registration totals.

<sup>c</sup>Otter estimates from separate mail harvest survey (Frawley 2019b). See Table 5 for registration totals.

<sup>d</sup>Fisher estimates from separate mail harvest survey (Frawley 2019a). See Table 5 for registration totals.

\*Non-overlapping 95% confidence intervals indicated estimates differed significantly between 2016 and 2017 (P<0.05).

Table 5. Number of bobcat, otter, fisher, badger and marten registered by furtakers in Michigan, 1985-2017.<sup>a</sup>

Year	Species							
	Bobcat (by method of capture)				Otter	Fisher <sup>a</sup>	Badger <sup>b,c</sup>	Marten <sup>d</sup>
	Hunting	Trapping	Unknown	Total				
1985	193	100	14	307	791			
1986	268	390	11	669	1,431			
1987	315	277	5	597	1,030			
1988	327	170	0	497	731			
1989	178	91	0	269	900	94	28	
1990	265	85	0	350	654	125	52	
1991	292	79	0	371	877	68	35	
1992	276	104	0	380	896	139	63	
1993	285	163	0	448	1,252	425	90	
1994	373	422	0	795	1,552	417	124	
1995	311	137	1	449	1,143	210	75	
1996	463	420	0	883	1,438	471	109	
1997	347	771	0	1118	1,324	609	117	
1998	331	373	0	704	1,026	455	91	
1999	434	343	0	777	1,097	291	82	
2000	379	307	0	686	1,006	236	85	85
2001	465	727	0	1,192	1,204	381	174	97
2002	482	741	0	1,223	1,221	348	173	85
2003	340	621	0	961	1,496	442		149
2004	321	637	0	958	1,358	368		184
2005	309	508	0	817	1,526	322		164
2006	336	515	0	851	1,154	390		192
2007	336	299	0	635	663	280		316
2008	284	364	0	648	707	326		290
2009	331	270	0	601	997	216		247
2010	365	344	0	709	935	312		274
2011	290	367	0	657	1,360	205		187
2012	311	367	0	678	1,234	237		279
2013	217	308	0	525	849	280		284
2014	333	325	0	658	834	191		289
2015	286	297	0	583	856	237		280
2016	259	140	0	399	711	171		131
2017 <sup>e</sup>	251	186	0	437	665	162		157

<sup>a</sup>Registration totals included only animals legally harvested by furtakers during hunting and trapping seasons; excluded harvest by tribal members. Also, totals only included animals that were registered and returned to the furtaker (i.e., excluded accidental take).

<sup>b</sup>Badger and fisher seasons were established in 1989.

<sup>c</sup>Furtakers no longer were required to register badgers beginning in 2003.

<sup>d</sup>Marten season was established in 2000.

<sup>e</sup>Preliminary totals.

Table 6. Estimated number of trappers that caught an incidental bobcat and number of incidental bobcats caught and registered in Michigan, 2017.

Region <sup>a</sup>	Trappers		Incidental bobcats captured and released alive <sup>b</sup>		Incidental bobcats captured and registered <sup>b</sup>	
	No.	95% CL	No.	95% CL	No.	95% CL
Upper Peninsula	19	25	19	34	0	0
Northern Lower Peninsula	134	66	200	116	19	25
Southern Lower Peninsula	18	24	18	24	0	0
Unknown	0	0	0	0	0	0
Statewide	171	74	237	123	19	25

<sup>a</sup>See Figure 1 for region boundaries.

<sup>b</sup>Incidental bobcats caught in counties.

Table 7. Estimated number of beaver trappers, beaver harvested, and trapping effort (days afield), summarized by trappers with and without an otter harvest tag in Michigan, 2017.

Beaver trapper group	Trappers		Days afield		Harvest	
	No.	95% CL	No.	95% CL	No.	95% CL
Without an otter harvest tag	797	156	13,914	3,744	3,004	1,293
With an otter harvest tag	1,142	186	9,169	2,856	9,169	2,856
Combined	1,939	239	38,121	7,185	12,172	3,127

Table 8. Furtakers' level of satisfaction with the number of animal or animal sign seen during the 2017 hunting and trapping seasons, summarized by the primary species the furtaker targeted.<sup>a</sup>

Species	Satisfaction level							
	Very satisfied or somewhat satisfied		Neutral		Very dissatisfied or somewhat dissatisfied		No answer	
	%	95%	%	95%	%	95%	%	95%
		CL		CL		CL		CL
Raccoon	71	6	18	5	8	4	3	2
Fox	65	17	20	15	7	9	8	10
Coyote	64	5	25	4	9	3	3	2
Bobcat	51	13	25	11	24	11	0	0
Fisher	64	27	27	25	9	16	0	0
Mink	62	32	13	23	25	28	0	0
Muskrat	70	8	19	7	7	5	4	3
Beaver	79	9	12	7	7	6	1	2

<sup>a</sup>Furtakers were grouped in subgroups based on the primary species they targeted, and then satisfaction was summarized for each subgroup separately.

Table 9. Furtakers' level of satisfaction with the number of animal harvested during the 2017 hunting and trapping seasons, summarized by the primary species the furtaker targeted.<sup>a</sup>

Species	Satisfaction level							
	Very satisfied or somewhat satisfied		Neutral		Very dissatisfied or somewhat dissatisfied		No answer	
	%	95%	%	95%	%	95%	%	95%
		CL		CL		CL		CL
Raccoon	60	7	21	5	15	5	4	3
Fox	36	17	38	17	18	14	7	9
Coyote	30	5	34	5	31	5	4	2
Bobcat	16	9	39	13	39	13	6	6
Fisher	45	28	36	27	9	16	9	17
Mink	25	28	0	0	63	32	13	22
Muskrat	57	9	24	7	16	7	3	3
Beaver	67	10	16	8	15	8	1	2

<sup>a</sup>Furtakers were grouped in subgroups based on the primary species they targeted, and then satisfaction was summarized for each subgroup separately.

Table 10. Furtakers' level of satisfaction with their overall hunting or trapping experience during 2017, summarized by the primary species the furtaker targeted.<sup>a</sup>

Species	Satisfaction level							
	Very satisfied or somewhat satisfied		Neutral		Very dissatisfied or somewhat dissatisfied		No answer	
	%	95%	%	95%	%	95%	%	95%
		CL		CL		CL		CL
Raccoon	70	6	16	5	8	4	5	3
Fox	51	18	30	17	7	9	11	11
Coyote	59	5	27	4	10	3	4	2
Bobcat	61	13	14	9	22	11	4	5
Fisher	55	28	36	27	9	16	0	0
Mink	63	32	25	28	0	0	13	22
Muskrat	74	8	13	6	10	5	4	3
Beaver	76	9	14	8	9	6	1	2

<sup>a</sup>Furtakers were grouped in subgroups based on the primary species they targeted, and then satisfaction was summarized for each subgroup separately.

Appendix A. Questionnaire used to collect data for 2017 fur harvesters survey in Michigan.





MICHIGAN DEPARTMENT OF NATURAL RESOURCES, WILDLIFE DIVISION  
 PO BOX 30030 LANSING MI 48909-7530

**2017-2018 FUR HARVESTER'S REPORT**

This information is requested under authority of Part 435, 1994 PA 451, M.C.L. 324.43539.



*It is important you return this report even if you did not capture any furbearers.  
 If you did not hunt or trap, answer "No" to questions 1 and 2 and return this report.  
 Do not report hunting and trapping activity done as part of a nuisance control business.  
 Do not report incidental captures you were not allowed to keep.*

1. Did you attempt to hunt any of the animals listed in the tables below during 2017-2018 seasons?    1  Yes    2  No
2. Did you set a trap (including cable restraints or snares) for any of the animals listed in the tables below during 2017-2018 seasons?    1  Yes    2  No

- If you did not attempt to hunt or trap furbearers during 2017-2018 seasons, you can skip the remaining questions and return the questionnaire. Otherwise, please read the following instructions. In addition, questions continue on the back side of this page.
- Place an "X" in the box next to the name of each animal you attempted to hunt or trap. Record your hunting and trapping activity separately. If you attempted to harvest a species but were unsuccessful, you still should report that you sought this species and report the number of days you tried to capture this species.
- For each species you attempted to hunt or trap, list the three primary COUNTIES where you hunted or trapped. Also report the number of DAYS you hunted or trapped and the number of animals TAKEN in each county you listed. If you do not know the county name, give the name of the nearest village, town or city.

**3. Please report your HUNTING ACTIVITY in the table below:**

SPECIES HUNTED	1 <sup>ST</sup> COUNTY	NO. OF DAYS	NO. TAKEN	2 <sup>ND</sup> COUNTY	NO. OF DAYS	NO. TAKEN	3 <sup>RD</sup> COUNTY	NO. OF DAYS	NO. TAKEN
1 <input type="checkbox"/> RACCOON									
2 <input type="checkbox"/> RED FOX									
3 <input type="checkbox"/> GRAY FOX									
4 <input type="checkbox"/> COYOTE									
5 <input type="checkbox"/> BOBCAT									

**4. Please report your TRAPPING ACTIVITY in the table below (include snaring):**

SPECIES TRAPPED	1 <sup>ST</sup> COUNTY	NO. OF DAYS	NO. TAKEN	2 <sup>ND</sup> COUNTY	NO. OF DAYS	NO. TAKEN	3 <sup>RD</sup> COUNTY	NO. OF DAYS	NO. TAKEN
6 <input type="checkbox"/> RACCOON									
7 <input type="checkbox"/> RED FOX									
8 <input type="checkbox"/> GRAY FOX									
9 <input type="checkbox"/> COYOTE									
10 <input type="checkbox"/> BOBCAT									
11 <input type="checkbox"/> OPPOSUM									
12 <input type="checkbox"/> SKUNK									
13 <input type="checkbox"/> WEASEL									
14 <input type="checkbox"/> BADGER									
15 <input type="checkbox"/> FISHER									
16 <input type="checkbox"/> MINK									
17 <input type="checkbox"/> MUSKRAT									
18 <input type="checkbox"/> BEAVER									

Please continue on back

If you set a trap (including cable restraints or snares) during the 2017-2018 seasons, please continue with the following questions. If you did not trap, skip the remaining questions.

5. Did you incidentally catch any bobcat while trapping for other species that you have not already reported in Question #4.

1  Yes      2  No, Skip to question number 7.

6. If you answered yes in the previous question, please report the location and number of incidental bobcats you captured. Please do not report bobcat already reported in question #4.

COUNTY WHERE INCIDENTAL BOBCAT CAUGHT (List each county that you caught an incidental bobcat.)	NUMBER OF INCIDENTAL BOBCAT CAUGHT AND RELEASED (Count only incidental bobcats you released alive from your traps.)	NUMBER OF INCIDENTAL BOBCAT CAUGHT AND REGISTERED (Count incidental bobcats that were registered including catches that were not returned to you.)

7. What was the primary furbearer species you sought during the past year?

(Select one.)

- 1  Raccoon      2  Fox      3  Coyote      4  Bobcat  
 5  Fisher      6  Mink      7  Muskrat      8  Beaver  
 9  Other (please specify: \_\_\_\_\_)

8. During the fur harvesters trapping and hunting seasons, indicate how satisfied or dissatisfied you were with the following for the primary furbearer species you hunted or trapped.

	Very Satisfied	Somewhat Satisfied	Neutral	Somewhat Dissatisfied	Very Dissatisfied
a. Number of animals or sign (e.g., tracks) seen.	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
b. Number of animals harvested.	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
c. Your overall hunting or trapping experience.	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>

In the next two questions, you will be asked about the time and costs you spend hunting or trapping furbearers during the 2017 seasons.

9. How many total days did you hunt or trap furbearers during the 2017 seasons? Please exclude days spent scouting.

\_\_\_\_\_ Days

10. How much did you spend on things related to hunting or trapping furbearers during the 2017 seasons (for example, fuel, food, lodging, equipment, and ammunition)? Please report only costs incurred during the 2017 seasons.

\$ \_\_\_\_\_ Total cost

11. Do you have any comments or suggestions about furbearer management in Michigan?

\_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_

Please return questionnaire in the enclosed postage-paid envelope.  
 Thank you for your help.