

# BIANNUAL HIV SURVEILLANCE REPORT

## CALHOUN COUNTY

### January 2013

## Table of Contents: HIV Surveillance Statistics of Persons Diagnosed in Calhoun County

Front Matter	Page
Acronyms and Definitions	i
Prevalence Estimate Calculations	i
Risk Transmission Categories	i
<b>Section 1: Data on Prevalent Cases</b>	
Table 1. Demographic Information on Prevalent HIV Infection Cases	1
Table 2. Sex, Race, and Risk Among Prevalent HIV Infection Cases	2
Table 3. Prevalent HIV Infection Cases According to City of Residence at Diagnosis	3
<b>Section 2: New Diagnoses, Deaths, and Prevalence</b>	
Table 4. New Diagnoses, Deaths, and Prevalence of HIV Infection, by Year	4
Figure 1. New Diagnoses, Deaths, and Prevalence of HIV Infection, by Year	5
Table 5. Comparison of HIV Infection Prevalence to Michigan	5

HIV/STD/VH/TB Epidemiology Section  
Division of Communicable Disease  
Bureau of Disease Control, Prevention and Epidemiology  
Michigan Department of Community Health

Lansing - HIV Surveillance Office  
201 Townsend St., 5th Floor  
Lansing, MI 48913  
517-335-8165

MDCH-South Oakland Health  
Center  
27725 Greenfield Rd, Office 57A  
Southfield, MI 48076  
248-424-7910

**\*\*WEBSITE\*\***

<http://www.michigan.gov/hivstd>

**\*\*WEBSITE\*\***



## General HIV

### AIDS (Acquired Immune Deficiency Syndrome)

Now referred to as stage 3 HIV infection (see "HIV Infection Stages" below).

### HIV (Human Immunodeficiency Virus)

Diagnosis with HIV requires both a positive HIV screening and positive supplemental antibody test or detectable quantity on a virologic test. A standard case definition for HIV infection is used by all states for surveillance. Specific information is required in order to count a case of HIV infection, including a method to uniquely identify an individual. Each case is classified in a HIV infection stage (see below). Once a case reaches stage 3 (AIDS), the case is always considered stage 3 for surveillance purposes, even if his/her health improves (MMWR; December 5, 2008 / Vol. 57 / No. RR--10 / Pg. 1 - 12).

### HIV Infection Stages

*Stage 1:* A case does not have any of the conditions associated with severe HIV infection (called an AIDS-defining condition) and has  $\geq 500$  CD4 cells/ $\mu$ l.

*Stage 2:* A case has no AIDS-defining condition, but the level of CD4 cells has fallen to 200-499 cells/ $\mu$ l.

*Stage 3:* Diagnosis with any one of 26 AIDS-defining conditions which are indicative of a severe immune deficiency, or a laboratory test demonstrating severe immune deficiency: CD4 count  $<200$  cells/ $\mu$ l or CD4 percent  $<14\%$ . **Previously referred to as AIDS.**

*Stage unknown:* A case of HIV without information available on CD4 levels or AIDS-defining conditions.

## Prevalence Estimate Calculations

HIV prevalence estimates include all persons living in Calhoun County at diagnosis of HIV Infection (any stage) and incorporates an approximation of those not reported or not yet diagnosed. All estimates are rounded to the nearest ten, and the minimum estimate given is 10. Prevalence estimates were most recently revised in January 2013, so the estimates in this report may vary in comparison to reports published before the revision.

## Risk Transmission Categories - Definitions

### Blood Recipients

Hemophiliacs, blood transfusion recipients, and organ recipients who received blood products prior to 1985 & persons documented to have ever received an infected organ or unit of blood.

### Heterosexual Contact (HC):

*Heterosexual Contact w/ Female Risk (HCFR):* Males whose female sexual partners are known to be HIV-infected or at high risk for HIV. These partners meet one of the following criteria: IDU, hemophiliac, HIV infected transfusion recipient, or other HIV infected person of unknown risk.

*Heterosexual Contact w/ Male (HCM):* Females who have had sex with a male regardless of what is known about the male's HIV status or behaviors.

### Please note:

-Males whose only documented risk is sex with a female, and their female partner's risk and HIV status is unknown fall into the undetermined risk category.

### Injection Drug Users (IDU)

Persons who have a history of injection drug use.

### Men who have sex with men (MSM)

Males who have a history of sexual contact with other men.

### MSM/IDU

MSM who also have a history of injection drug use.

### Perinatal

HIV transmission from mother to child during birth or through breastfeeding.

### Undetermined

Males and females with no identified risk.

**TABLE 1. Demographic Information on Prevalent HIV Infection Cases Living in CALHOUN CO. at Diagnosis**

	REPORTED PREVALENCE <sup>†</sup>								CENSUS ESTIMATES 2011	
	EST PREV*	HIV Infection Non-Stage 3		HIV Infection Stage 3 (AIDS)		Total		Rate per 100,000 <sup>§</sup>	Num	Percent
	Num	Num	Percent	Num	Percent	Num	Percent			
<b>RACE/ ETHNICITY<sup>‡</sup></b>										
White	90	32	47%	37	52%	69	50%	63.8	108,073	80%
Black	70	26	38%	28	39%	54	39%	369.2	14,626	11%
Hispanic	10	<5	**	<5	**	<5	**	**	6,259	5%
Asian/PI	20	8	12%	<5	**	12	9%	537.6	2,232	2%
Am Indian/AN	0	0	0%	0	0%	0	0%	**	721	1%
Multi/Other/Unk	10	<5	**	<5	**	<5	**	N/A	3,579	3%
<b>SEX &amp; RACE</b>										
Male	130	47	69%	50	70%	97	70%	146.5	66,205	49%
White Male	60	25	37%	23	32%	48	35%	91.4	52,525	39%
Black Male	50	17	25%	22	31%	39	28%	535.3	7,285	5%
Hispanic Male	10	0	0%	<5	**	<5	**	**	3,211	2%
Other Male	10	5	7%	<5	**	9	6%	---	3,184	2%
Female	50	21	31%	21	30%	42	30%	60.6	69,285	51%
White Female	30	7	10%	14	20%	21	15%	37.8	55,548	41%
Black Female	20	9	13%	6	8%	15	11%	204.3	7,341	5%
Hispanic Female	10	<5	**	0	0%	<5	**	**	3,048	2%
Other Female	10	<5	**	<5	**	5	4%	---	3,348	2%
<b>RISK*</b>										
Male-Male Sex (MSM)	80	31	46%	27	38%	58	42%	--	--	--
Injection Drug Use (IDU)	20	<5	**	10	14%	13	9%	--	--	--
MSM/IDU	10	<5	**	<5	**	<5	**	--	--	--
Blood Products	10	0	0%	<5	**	<5	**	--	--	--
Heterosexual Contact (HC)	50	19	28%	20	28%	39	28%	--	--	--
HCFR (Males)	10	<5	**	6	8%	10	7%	--	--	--
HCM (Females)	40	15	22%	14	20%	29	21%	--	--	--
Perinatal	10	<5	**	<5	**	<5	**	--	--	--
Undetermined	30	12	18%	9	13%	21	15%	--	--	--
<b>AGE AT HIV DIAGNOSIS</b>										
0 - 12 years	10	<5	**	<5	**	<5	**	--	--	--
13 - 19 years	10	<5	**	7	10%	10	7%	--	--	--
20 - 24 years	30	13	19%	10	14%	23	17%	--	--	--
25 - 29 years	30	13	19%	11	15%	24	17%	--	--	--
30 - 39 years	50	16	24%	24	34%	40	29%	--	--	--
40 - 49 years	40	18	26%	10	14%	28	20%	--	--	--
50 - 59 years	10	<5	**	7	10%	10	7%	--	--	--
60 years and over	10	<5	**	<5	**	<5	**	--	--	--
Unspecified	0	0	0%	0	0%	0	0%	--	--	--
<b>TOTAL</b>	<b>180</b>	<b>68</b>	<b>100%</b>	<b>71</b>	<b>100%</b>	<b>139</b>	<b>100%</b>	<b>103</b>	<b>135,490</b>	<b>100%</b>

\*See front matter page i for descriptions of prevalence estimate calculations and risk category groupings. Risk categories used in Michigan are redefined as of January 2012. NOTE: Heterosexual contact for males includes only males whose sexual partners are known to be HIV infected or at high risk for HIV (HCFR). Heterosexual contact for females includes all females who have had sex with a male regardless of what is known about the male's HIV status or behaviors (HCM).

<sup>†</sup> Includes reports that contain patient name or are otherwise unduplicated. <5 and "\*\*" = 1, 2, 3, or 4 cases.

<sup>‡</sup> To calculate "1 out of x" statements, divide the census number by total reported prevalence. For example, for non-Hispanic whites: 108,073 / 69 = 1,566. Thus, 1 out of every 1,566 non-Hispanic white persons in CALHOUN CO. are living with HIV. Rates and "1 out of x" statements are not reliable for <10 cases. Thus, rates for <10 cases are shown as "----".

<sup>§</sup> Rates are not reported for risk categories and age at diagnosis because no reliable denominator data exist for these groups.

<sup>¶</sup> In this report, persons described as white, black, Asian/Pacific Islander (PI), or American Indian/Alaska Native (AN) are all non-Hispanic; persons described as Hispanic might be of any race.

**TABLE 2. Sex, Race, and Risk Among Prevalent HIV Infection Cases Living in CALHOUN CO. at Diagnosis**

	White		Black		Hispanic		Other or Unknown		All Male	
	Num	Percent	Num	Percent	Num	Percent	Num	Percent	Num	Percent
<b>MALE</b>										
Male-Male sex	39	81%	18	46%	0	0%	<5	**	58	60%
Injection Drug Use	0	0%	5	13%	<5	**	<5	**	7	7%
MSM/IDU	<5	**	<5	**	0	0%	0	0%	<5	**
Blood Products	<5	**	0	0%	0	0%	0	0%	<5	**
Heterosexual Contact (HCFR)	<5	**	6	15%	0	0%	<5	**	10	10%
Perinatal	0	0%	0	0%	0	0%	0	0%	0	0%
Undetermined	<5	**	8	21%	0	0%	6	67%	17	18%
<b>Male Subtotal</b>	<b>48</b>	<b>49%</b>	<b>39</b>	<b>40%</b>	<b>&lt;5</b>	<b>**</b>	<b>9</b>	<b>9%</b>	<b>97</b>	<b>100%</b>
<b>FEMALE</b>										
Injection Drug Use	<5	**	<5	**	0	0%	0	0%	6	14%
Blood Products	<5	**	0	0%	0	0%	0	0%	<5	**
Heterosexual Contact (HCM)	13	54%	13	87%	<5	**	<5	**	29	69%
Perinatal	<5	**	0	0%	0	0%	<5	**	<5	**
Undetermined	<5	**	0	0%	0	0%	<5	**	<5	**
<b>Female Subtotal</b>	<b>24</b>	<b>57%</b>	<b>15</b>	<b>36%</b>	<b>&lt;5</b>	<b>**</b>	<b>&lt;5</b>	<b>**</b>	<b>42</b>	<b>100%</b>
<b>ALL</b>										
Male-Male sex	39	54%	18	33%	0	0%	<5	**	58	42%
Injection Drug Use	<5	**	7	13%	<5	**	<5	**	13	9%
MSM/IDU	<5	**	<5	**	0	0%	0	0%	<5	**
Blood Products	5	7%	0	0%	0	0%	0	0%	<5	**
Heterosexual Contact (HC)	16	22%	19	35%	<5	**	<5	**	39	28%
HCFR (Males)	<5	**	6	11%	0	0%	<5	**	10	7%
HCM (Females)	13	18%	13	24%	<5	**	<5	**	29	21%
Perinatal	<5	**	0	0%	0	0%	<5	**	<5	**
Undetermined	5	7%	8	15%	0	0%	8	62%	21	15%
<b>RACE ALL</b>	<b>72</b>	<b>52%</b>	<b>54</b>	<b>39%</b>	<b>&lt;5</b>	<b>**</b>	<b>13</b>	<b>9%</b>	<b>139</b>	<b>100%</b>

**TABLE 4. New Diagnoses, Deaths, and Prevalence of HIV Infection in CALHOUN CO., by Year**

Year	<i>HIV Infection (all stages)</i>			<i>HIV Infection Stage 3 (AIDS)</i>		
	New HIV Diagnoses	Deaths	Prevalence	New Stage 3 Diagnoses	Deaths	Prevalence
1983	1	0	1	1	0	1
1984	1	0	2	1	0	2
1985	2	1	3	0	1	1
1986	9	0	12	3	0	4
1987	5	3	14	4	3	5
1988	8	3	19	5	3	7
1989	8	3	24	5	3	9
1990	6	3	27	3	3	9
1991	10	2	35	6	2	13
1992	7	10	32	8	10	11
1993	13	6	39	9	5	15
1994	11	5	45	7	5	17
1995	17	5	57	12	4	25
1996	10	2	65	8	2	31
1997	13	4	74	6	3	34
1998	5	6	73	1	5	30
1999	5	1	77	2	1	31
2000	10	7	80	3	6	28
2001	4	1	83	4	1	31
2002	6	3	86	7	3	35
2003	6	1	91	6	1	40
2004	4	7	88	5	7	38
2005	2	2	88	6	2	42
2006	6	0	94	3	0	45
2007	9	1	102	6	1	50
2008	3	1	104	1	1	50
2009	8	2	110	2	1	51
2010	7	1	116	8	1	58
2011	16	0	132	7	0	65
2012	8	1	<b>139</b>	7	1	<b>71</b>
<b>TOTAL</b>	<b>220</b>	<b>81</b>		<b>146</b>	<b>75</b>	


The prevalence of HIV in Michigan has steadily increased, since persons with HIV are living longer. This is largely due to improved anti-retroviral therapy.

The increase in HIV prevalence is also reflected in Figure 1 on page 4, which shows that the number of persons diagnosed, while stable for the last several years, is greater than the number of deaths each year. This directly contributes to the increase in prevalence. The current reported prevalence of HIV infection in CALHOUN CO. is 139. The prevalence of Stage 3 infection, which is a subset of the overall HIV infection prevalence, is 71.

As implied, the HIV infection section displays data on all persons with HIV, including those with Stage 3 infection, as well as those who have not progressed to Stage 3. Thus, persons represented in the Stage 3 section are also represented in the HIV infection section. The number of reported deaths includes deaths directly attributable to presence of HIV infection as well as deaths due to other causes.

NOTE: Reporting for recent years may not be complete. Data are not adjusted to account for reporting delays.

**FIGURE 1. New Diagnoses, Deaths, and Prevalence of HIV Infection in CALHOUN CO., by Year**



**TABLE 5. Comparison of HIV Infection Prevalence in CALHOUN CO. to State**

<i>Residence at Diagnosis</i>	<i>Estimated Prevalence</i>	<i>Reported Prevalence</i>				<i>Cumulative Stage 3</i>
		<b>HIV Infection Non-Stage 3</b>	<b>HIV Infection Stage 3 (AIDS)</b>	<b>Total</b>	<b>Rate</b>	
<b>Calhoun Co.</b>	180	68	71	139	103	146
<b>Michigan Total</b>	19,800	7,169	7,912	15,081	153	18,130