

GIST

Guideline Implementation Steps & Tools

Asthma Guideline Implementation Steps & Tools (GIST)

A provider education and practice redesign project by the Michigan Department of Community Health based on the 2007 NAEPP Asthma Guidelines. GIST makes it easier for primary care clinicians to use the asthma guidelines in their everyday care of patients with asthma. Visit GetAsthmaHelp.org/GIST for more information.

“Questions About Your Breathing”
or
“Asthma Control Test”

Given prior to seeing clinician

- Gets “Questions” form if being seen for respiratory complaints- history and symptom questionnaire
- “ACT” form for returning asthma patient, patient control assessment

“Asthma Diagnosis Tool”
or
“Asthma Patient Follow-Up Tool”

Patient with clinician, who uses:

- “Diagnosis” form if patient in for respiratory complaints- diagnoses/rules out asthma, finds severity level if asthma, step care started
- “Follow-Up” form for returning asthma patient- finds control level, steps up, down or maintains as needed

“Stepwise Approach to
Managing Asthma”
Asthma Action Plan, Ed & Meds

- Use Stepwise Approach to find medication needs, management approach
- Opportunity for patient ed about asthma triggers, meds, barriers, etc.
- Patient receives prescription(s) and Asthma Action Plan
- No specified AAP for GIST, many good ones available at GetAsthmaHelp.org

Evaluate,
improve system,
repeat

Monitor progress by tracking these or other outcomes:

- number of asthma patients in practice
- number (or %) of asthma patients with current asthma action plan
- number (or %) of asthma patients with control assessment at last visit
- number (or %) of asthma patients with appropriate prescription of ICS

ASTHMA DIAGNOSIS TOOL *Consider the diagnosis of asthma if patient states any of the following:*

- Family history of asthma, allergies or eczema
- Symptoms occur seasonally
- Symptoms when near chemicals, dusts, fumes at work
- Symptoms worsened by URI lasting longer than ten days, smoke, allergens or exercise

AND SPIROMETRY DEMONSTRATES OBSTRUCTION AND/OR REVERSIBILITY BY AN INCREASE IN FEV₁ OF 12% OR MORE AFTER BRONCHODILATOR.

Rule out co-morbid conditions. If in doubt, consult with an asthma specialist.

HIGHEST LEVEL OF CHECKED BOX = SEVERITY LEVEL / FOLLOW SEVERITY LEVEL DOWN TO FIND TREATMENT STEP → SEE TREATMENT STEPWISE APPROACH

	INTERMITTENT	MILD PERSISTENT	MODERATE PERSISTENT	SEVERE PERSISTENT
IMPAIRMENT	<p>SYMPTOMS:</p> <ul style="list-style-type: none"> <input type="checkbox"/> 2x/week or less <p>NIGHTTIME AWAKENINGS:</p> <ul style="list-style-type: none"> <input type="checkbox"/> 2x/month or less <p>INTERFERENCE W/NORMAL ACTIVITY:</p> <ul style="list-style-type: none"> <input type="checkbox"/> None <p>SHORT-ACTING B₂-AGONIST USE:</p> <ul style="list-style-type: none"> <input type="checkbox"/> 2 days/week or less <p>LUNG FUNCTION:</p> <ul style="list-style-type: none"> <input type="checkbox"/> FEV₁ more than 80% pred. 	<p>SYMPTOMS:</p> <ul style="list-style-type: none"> <input type="checkbox"/> More than 2x/week, not daily <p>NIGHTTIME AWAKENINGS:</p> <ul style="list-style-type: none"> <input type="checkbox"/> More than 2x/month <p>INTERFERENCE W/NORMAL ACTIVITY:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Minor limitation <p>SHORT-ACTING B₂-AGONIST USE:</p> <ul style="list-style-type: none"> <input type="checkbox"/> More than 2 days/week but not daily or more than 1x/day <p>LUNG FUNCTION:</p> <ul style="list-style-type: none"> <input type="checkbox"/> FEV₁ more than 80% pred. 	<p>SYMPTOMS:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Daily <p>NIGHTTIME AWAKENINGS:</p> <ul style="list-style-type: none"> <input type="checkbox"/> About 1x/week, not nightly <p>INTERFERENCE W/NORMAL ACTIVITY:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Some limitation <p>SHORT-ACTING B₂-AGONIST USE:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Daily <p>LUNG FUNCTION:</p> <ul style="list-style-type: none"> <input type="checkbox"/> FEV₁ 60-80% pred. 	<p>SYMPTOMS:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Throughout the day <p>NIGHTTIME AWAKENINGS:</p> <ul style="list-style-type: none"> <input type="checkbox"/> More than 1x/week, often nightly <p>INTERFERENCE W/NORMAL ACTIVITY:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Extremely limited <p>SHORT-ACTING B₂-AGONIST USE:</p> <ul style="list-style-type: none"> <input type="checkbox"/> Several times/day <p>LUNG FUNCTION:</p> <ul style="list-style-type: none"> <input type="checkbox"/> FEV₁ less than 60% pred.
RISK	<p>EXACERBATIONS REQUIRING ORAL STEROIDS:</p> <ul style="list-style-type: none"> <input type="checkbox"/> All ages: 0-1/year 	<p>EXACERBATIONS REQUIRING ORAL STEROIDS: consider severity and interval since last exacerbation</p> <ul style="list-style-type: none"> <input type="checkbox"/> Age 0-4: more than 2 in 6 months or more than 4 wheezing episodes/year lasting more than 1 day <input type="checkbox"/> All ages: more than 2/year <p>• Exacerbations of any severity may occur in patients in any severity category. • Frequency and severity may fluctuate over time.</p>		
TREATMENT STEP	<p>✓ All ages: STEP 1</p>	<p>✓ All ages: STEP 2</p>	<p>✓ All Ages: STEP 3; consider short course oral steroids option</p>	<p>✓ Age 0-4: STEP 3; short course oral steroids option</p> <p>✓ Age 5-11: STEP 3; STEP 4 short course oral steroids option</p> <p>✓ Age 12 & over: STEP 4 or 5; short course oral steroids option</p>
<p>TREATMENT FOR PERSISTENT ASTHMA: ✓ Daily inhaled steroids (see treatment stepwise approach) ✓ Assess response within 2-6 weeks</p>				
<p>FOR ALL PATIENTS WITH ASTHMA: <input type="checkbox"/> Rescue medication for all ages, all severity levels: Short-acting B₂-agonist PRN. Treatment intensity depends on symptom severity. <input type="checkbox"/> Provide written Asthma Action Plan <input type="checkbox"/> Identify & avoid triggers <input type="checkbox"/> Flu vaccine recommended annually, pneumococcal vaccine for adults <input type="checkbox"/> Review correct device technique each visit</p>				

Quick reference medication guides, asthma action plans and more: GetAsthmaHelp.org/GIST

Intermittent Asthma

Persistent Asthma: Daily Medication

Step up as indicated although address possible poor adherence to medication. Re-assess in 2 to 6 weeks.

Step down if well controlled and re-assess in 3 months. If very stable then assess control every 3 to 6 months.

All LABAs and combination agents containing LABAs have a black box warning.

Reference: National Heart, Lung, and Blood Institute's Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma 2007, NIH Publication 07-4051. This tool was adapted from the Colorado Clinical Guidelines Collaborative.

STEP 1 (all ages)

Short-acting beta-agonist (e.g., albuterol prn)

If used more than 2 days per week (other than for exercise) consider inadequate control and the need to step up treatment.

STEP 2

ALL AGES

Preferred: Low-dose inhaled steroid

Alternative: Leukotriene blocker **or** cromolyn

AGE 0-4 YRS

Consider referral (especially if diagnosis is in doubt)

STEP 3

AGE 12+ YRS

Preferred: Low-dose inhaled steroid + long-acting beta agonist **or** Medium-dose inhaled steroid

Alternative: Low-dose inhaled steroid + leukotriene blocker

AGE 5-11 YRS

Low-dose inhaled steroid + long-acting beta agonist **or** leukotriene blocker **or** Medium-dose inhaled steroid

AGE 0-4 YRS

Medium-dose inhaled steroid + referral

STEP 4

AGE 12+ YRS

Preferred: Medium-dose inhaled steroid + long-acting beta agonist

Alternative: Medium-dose inhaled steroid + leukotriene blocker

AGE 5-11 YRS

Same as 12+ yrs

AGE 0-4 YRS

Medium-dose inhaled steroid + either long-acting beta-agonist **or** leukotriene blocker

STEP 5

AGE 12+ YRS

High-dose inhaled steroid + long-acting beta agonist **—and—** Consider omalizumab if allergies

AGE 5-11 YRS

Preferred: High-dose inhaled steroid + long-acting beta agonist

Alternative: High-dose inhaled steroid + leukotriene blocker

AGE 0-4 YRS

High-dose inhaled steroid + either long-acting beta-agonist **or** leukotriene blocker

STEP 6

AGE 12+ YRS

High-dose inhaled steroid + long-acting beta agonist + oral steroid **—and—** Consider omalizumab if allergies

AGE 5-11 YRS

Preferred: High-dose inhaled steroid + long-acting beta agonist

Alternative: High-dose inhaled steroid + leukotriene blocker + **oral steroid**

AGE 0-4 YRS

High-dose inhaled steroid + either long-acting beta-agonist **or** leukotriene blocker + **oral steroid**

All ages Steps 4 through 6: Consult with asthma specialist

Consider immunotherapy if allergic asthma

RESCUE MEDICATION: Short-acting beta-agonist (e.g. albuterol) as needed for symptoms. Treatment intensity depends on symptom severity. Frequent or increasing use of rescue medication may indicate inadequate control and the need to step up treatment.

ASTHMA PATIENT FOLLOW-UP TOOL *Assess patient's asthma control and device technique.*

ACT™ Test Score _____ **Severity level at diagnosis:** Intermittent Mild Persistent Moderate Persistent Severe Persistent

HIGHEST LEVEL OF CHECKED BOX = CONTROL LEVEL / FOLLOW CONTROL LEVEL DOWN TO FIND TREATMENT STEP → SEE TREATMENT STEPWISE APPROACH

	WELL CONTROLLED	NOT WELL CONTROLLED	VERY POORLY CONTROLLED
IMPAIRMENT	<p>SYMPTOMS:</p> <input type="checkbox"/> 2 day/week or less, not more than once per day	<p>SYMPTOMS:</p> <input type="checkbox"/> More than 2 days/week or multiple times on 2 days/week or less	<p>SYMPTOMS:</p> <input type="checkbox"/> Throughout the day
	<p>NIGHTTIME AWAKENINGS:</p> <input type="checkbox"/> No more than once/month	<p>NIGHTTIME AWAKENINGS:</p> <input type="checkbox"/> Ages 0-4: More than once/month <input type="checkbox"/> Ages 5-11: 2 times/month or more <input type="checkbox"/> Age 12 & over: 1-3 times/week	<p>NIGHTTIME AWAKENINGS:</p> <input type="checkbox"/> Ages 0-4: More than once/week <input type="checkbox"/> Ages 5-11: 2 times/week or more <input type="checkbox"/> Age 12 & over: 4 times/week or more
	<p>INTERFERENCE W/NORMAL ACTIVITY:</p> <input type="checkbox"/> None	<p>INTERFERENCE W/NORMAL ACTIVITY:</p> <input type="checkbox"/> Some limitation	<p>INTERFERENCE W/NORMAL ACTIVITY:</p> <input type="checkbox"/> Extremely limited
	<p>SHORT-ACTING B₂-AGONIST USE:</p> <input type="checkbox"/> 2 days/week or less	<p>SHORT-ACTING B₂-AGONIST USE:</p> <input type="checkbox"/> More than 2 days/week	<p>SHORT-ACTING B₂-AGONIST USE:</p> <input type="checkbox"/> Several times/day
	<p>FEV₁ OR PEAK FLOW:</p> <input type="checkbox"/> Age 5 & over: More than 80% predicted/personal best	<p>FEV₁ OR PEAK FLOW:</p> <input type="checkbox"/> Age 5 & over: 60-80% pred./personal best	<p>FEV₁ OR PEAK FLOW:</p> <input type="checkbox"/> Age 5 & over: Less than 60% pred./personal best
	<p>FEV₁/FVC:</p> <input type="checkbox"/> Age 5 & over: more than 80%	<p>FEV₁/FVC:</p> <input type="checkbox"/> Age 5 & over: 75-80%	<p>FEV₁/FVC:</p> <input type="checkbox"/> Age 5 & over: less than 75%
	<p>ACT SCORE:</p> <input type="checkbox"/> 20 or more	<p>ACT SCORE:</p> <input type="checkbox"/> 16-19	<p>ACT SCORE:</p> <input type="checkbox"/> 15 or less
RISK	<p>EXACERBATIONS REQUIRING ORAL STEROIDS</p> <input type="checkbox"/> All ages: 0-1/year	<p>EXACERBATIONS REQUIRING ORAL STEROIDS</p> <input type="checkbox"/> Age 0-4: 2-3/year <input type="checkbox"/> Age 5 & over: 2/year or more; consider severity	<p>EXACERBATIONS REQUIRING ORAL STEROIDS</p> <input type="checkbox"/> Age 0-4: More than 3/year <input type="checkbox"/> Age 5 & over: 2/year or more; consider severity
TREATMENT STEP	<input type="checkbox"/> Maintain current step	<input checked="" type="checkbox"/> Check adherence & environmental control <input type="checkbox"/> Step up 1 step and assess response in 2-6 weeks	<input checked="" type="checkbox"/> Check adherence & environmental control <input type="checkbox"/> Consider short course of oral corticosteroids <input type="checkbox"/> Consider co-morbid conditions <input type="checkbox"/> Step up 1-2 steps and assess response in 2 weeks
	<input type="checkbox"/> Consider step down if well controlled for at least 3 months	<input type="checkbox"/> For side effects, consider alternative treatment options	
	<input type="checkbox"/> Rescue medication for all ages, all severity/control levels: Short-acting B ₂ -agonist PRN. Treatment intensity depends on symptom severity. <input type="checkbox"/> Provide written Asthma Action Plan; review/update <input type="checkbox"/> Spirometry annually for age 5 & over <input type="checkbox"/> Flu vaccine recommended annually, pneumooccal vaccine for adults <input type="checkbox"/> Consider referral to a specialist if not well controlled within 3-6 months using stepwise approach OR 2 or more ED visits or hospitalizations for asthma in a year.		

Today's Date: _____

Patient's Name: _____

FOR PATIENTS:

Take the Asthma Control Test™ (ACT) for people 12 yrs and older. Know your score. Share your results with your doctor.

Step 1 Write the number of each answer in the score box provided.

Step 2 Add the score boxes for your total.

Step 3 Take the test to the doctor to talk about your score.

1. In the past 4 weeks, how much of the time did your **asthma** keep you from getting as much done at work, school or at home?

All of the time	1	Most of the time	2	Some of the time	3	A little of the time	4	None of the time	5
-----------------	---	------------------	---	------------------	---	----------------------	---	------------------	---

2. During the past 4 weeks, how often have you had shortness of breath?

More than once a day	1	Once a day	2	3 to 6 times a week	3	Once or twice a week	4	Not at all	5
----------------------	---	------------	---	---------------------	---	----------------------	---	------------	---

3. During the past 4 weeks, how often did your **asthma** symptoms (wheezing, coughing, shortness of breath, chest tightness or pain) wake you up at night or earlier than usual in the morning?

4 or more nights a week	1	2 or 3 nights a week	2	Once a week	3	Once or twice	4	Not at all	5
-------------------------	---	----------------------	---	-------------	---	---------------	---	------------	---

4. During the past 4 weeks, how often have you used your rescue inhaler or nebulizer medication (such as albuterol)?

3 or more times per day	1	1 or 2 times per day	2	2 or 3 times per week	3	Once a week or less	4	Not at all	5
-------------------------	---	----------------------	---	-----------------------	---	---------------------	---	------------	---

5. How would you rate your **asthma** control during the past 4 weeks?

Not controlled at all	1	Poorly controlled	2	Somewhat controlled	3	Well controlled	4	Completely controlled	5
-----------------------	---	-------------------	---	---------------------	---	-----------------	---	-----------------------	---

SCORE

TOTAL

Copyright 2002, by QualityMetric Incorporated.
Asthma Control Test is a trademark of QualityMetric Incorporated.

**If your score is 19 or less, your asthma may not be controlled as well as it could be.
Talk to your doctor.**

FOR PHYSICIANS:

The ACT is:

- A simple, 5-question tool that is self-administered by the patient
- Clinically validated by specialist assessment and spirometry¹
- Recognized by the National Institutes of Health

QUESTIONS ABOUT YOUR BREATHING

Please answer the questions below for **ONLY THE PATIENT** seeing the doctor today,
you OR your child.

Name: _____

Date of Birth: _____

Today's Date: _____

1. Have you/has your child had shortness of breath, coughing, wheezing (whistling in the chest) during the day?

Yes No

2. Have you/has your child had breathing trouble at night or early in the morning Yes No

3. Has breathing trouble kept you/kept your child from school/work/normal activities? Yes No

4. Have you/has your child ever been to a doctor, urgent care, emergency room or a hospital for breathing trouble? Yes No

5. Do you/does your child get colds that settle in the chest, or coughing that lasts 10 days or more after a cold is gone?

Yes No

6. Have you/has your child ever needed steroid pills or syrup (prednisone, prednisolone, prelone) for breathing trouble?

Yes No

If yes, how many times has this happened? _____

7. Have you/has your child ever taken any other medicine (pills, inhalers, puffers, syrup) for breathing trouble? Yes No

If yes, please list: _____

8. Do you/does your child have a history of eczema, hay fever or other allergies, including foods? Yes No

If yes, please tell us about them: _____

9. At what age did you/did your child start having breathing trouble? _____

10. Do any blood relatives (parent, brother, sister, child) have:

Asthma Allergies

11. Do you or anyone in the family smoke? Yes No

12. Are you/is your child ever in smoky places? Yes No

13. Check any of the things that make your/your child's breathing worse, or tell us about others.

Breathing in chemicals, dusts, fumes at work

Colds or flu

Strong odors, like cleaners or perfumes

Animals

Weather

Dust

Exercise

Pollen and mold

Cigarette and other smoke

Medicines: _____

Other things: _____

Please list any medicine that you/that your child takes: _____

Thank you for your help! Please give this form to the doctor who sees you/your child today.