

TABLE 1. Demographic Information on Prevalent HIV/AIDS Cases Living in JACKSON CO. at Diagnosis

	EST PREV*	REPORTED PREVALENCE[†]					CENSUS 2006 ESTIMATES	
		HIV, not AIDS		AIDS		Total	Rate per 100,000[§]	Number Percent
	Number	Number Percent	Number Percent	Number Percent	Number Percent		Number Percent	
RACE/ ETHNICITY[¶]								
White	130	42 (72%)	48 (84%)	90 (78%)	63.3	142,265 (87%)		
Black	30	14 (24%)	8 (14%)	22 (19%)	172.8	12,730 (8%)		
Hispanic	10	<5 **	<5 **	<5 **	**	4,255 (3%)		
Asian/PI	0	0 (0%)	0 (0%)	0 (0%)	**	1,177 (1%)		
Am Indian/AN	0	0 (0%)	0 (0%)	0 (0%)	**	624 (<1%)		
Multi/Unk	0	0 (0%)	0 (0%)	0 (0%)	N/A	2,800 (2%)		
SEX & RACE								
Males	130	46 (79%)	50 (88%)	96 (83%)	114.7	83,715 (51%)		
White Males	100	31 (53%)	42 (74%)	73 (63%)	102.8	71,040 (43%)		
Black Males	30	13 (22%)	7 (12%)	20 (17%)	248.0	8,065 (5%)		
Hispanic Males	10	<5 **	<5 **	<5 **	**	2,273 (1%)		
Other Males	0	0 (0%)	0 (0%)	0 (0%)	**	2,337 (1%)		
Females	30	12 (21%)	7 (12%)	19 (17%)	23.7	80,136 (49%)		
White Females	20	11 (19%)	6 (11%)	17 (15%)	23.9	71,225 (43%)		
Black Females	10	<5 **	<5 **	<5 **	**	4,665 (3%)		
Hispanic Fmls	0	0 (0%)	0 (0%)	0 (0%)	**	1,982 (1%)		
Other Females	0	0 (0%)	0 (0%)	0 (0%)	**	2,264 (1%)		
RISK*								
Male-Male Sex	80	25 (43%)	34 (60%)	59 (51%)	N/A	N/A N/A		
Injection Drug Use	10	<5 **	<5 **	5 (4%)	N/A	N/A N/A		
MSM/IDU	10	<5 **	<5 **	<5 **	N/A	N/A N/A		
Blood Products	10	<5 **	<5 **	<5 **	N/A	N/A N/A		
Heterosexual	30	13 (22%)	9 (16%)	22 (19%)	N/A	N/A N/A		
HRH	20	11 (19%)	6 (11%)	17 (15%)	N/A	N/A N/A		
PH-Female	10	<5 **	<5 **	5 (4%)	N/A	N/A N/A		
Perinatal	10	<5 **	0 (0%)	<5 **	N/A	N/A N/A		
Undetermined	30	13 (22%)	9 (16%)	22 (19%)	N/A	N/A N/A		
PH-Male	20	8 (14%)	6 (11%)	14 (12%)	N/A	N/A N/A		
Unknown	10	5 (9%)	<5 **	8 (7%)	N/A	N/A N/A		
AGE AT HIV DIAGNOSIS								
0 - 12 years	10	<5 **	0 (0%)	<5 **	N/A	N/A N/A		
13 - 19 years	10	<5 **	<5 **	<5 **	N/A	N/A N/A		
20 - 24 years	30	12 (21%)	6 (11%)	18 (16%)	N/A	N/A N/A		
25 - 29 years	30	9 (16%)	13 (23%)	22 (19%)	N/A	N/A N/A		
30 - 39 years	60	20 (34%)	21 (37%)	41 (36%)	N/A	N/A N/A		
40 - 49 years	20	8 (14%)	7 (12%)	15 (13%)	N/A	N/A N/A		
50 - 59 years	20	<5 **	7 (12%)	11 (10%)	N/A	N/A N/A		
60 years and over	10	0 (0%)	<5 **	<5 **	N/A	N/A N/A		
Unspecified	0	0 (0%)	0 (0%)	0 (0%)	N/A	N/A N/A		
TOTAL	160	58 (100%)	57 (100%)	115 (100%)	70	163,851 (100%)		

*See page 4 for descriptions of prevalence estimate calculations and new risk category groupings. Risk categories used in Michigan are newly defined as of the July 2007 quarter.

[†] Includes reports that contain patient name or are otherwise unduplicated. <5 and ** = 1, 2, 3, or 4 cases.

[§] To calculate "1 out of x" statements, divide the census number by total reported prevalence. For example, for non-Hispanic whites: 142,265/90 = 1,581. Thus, 1 out of every 1,581 non-Hispanic white persons in JACKSON CO. are living with HIV.

[¶] In this report, persons described as white, black, Asian/Pacific Islander (PI), or American Indian/Alaska Native (AN) are all non-Hispanic; persons described as Hispanic might be of any race.

TABLE 2. Sex, Race, and Risk Among Prevalent HIV/AIDS Cases in JACKSON CO.

MALES	White		Black		Hispanic		Other or Unknown		Male Subtotal	
Male-Male sex	52	(71%)	5	(25%)	<5	**	0	(0%)	59	(61%)
Injecting Drug Use	<5	**	<5	**	0	(0%)	0	(0%)	5	(5%)
Male-Male Sex/IDU	<5	**	<5	**	0	(0%)	0	(0%)	<5	**
Blood Products	<5	**	<5	**	0	(0%)	0	(0%)	<5	**
Heterosexual*	<5	**	<5	**	0	(0%)	0	(0%)	5	(5%)
Perinatal	0	(0%)	0	(0%)	0	(0%)	0	(0%)	0	(0%)
Undetermined	11	(15%)	9	(45%)	<5	**	0	(0%)	21	(22%)
<i>PH-Male</i>	7	(10%)	7	(35%)	0	(0%)	0	(0%)	14	(15%)
<i>Unknown</i>	<5	**	<5	**	<5	**	0	(0%)	7	(7%)
Male Subtotal	73	(76%)	20	(21%)	<5	**	0	(0%)	96	(100%)
FEMALES	White		Black		Hispanic		Other or Unknown		Female Subtotal	
Injecting Drug Use	0	(0%)	0	(0%)	0	(0%)	0	(0%)	0	(0%)
Blood Products	0	(0%)	0	(0%)	0	(0%)	0	(0%)	0	(0%)
Heterosexual	15	(88%)	<5	**	0	(0%)	0	(0%)	17	(89%)
<i>HRH</i>	11	(65%)	<5	**	0	(0%)	0	(0%)	12	(63%)
<i>PH-Female</i>	<5	**	<5	**	0	(0%)	0	(0%)	5	(26%)
Perinatal	<5	**	0	(0%)	0	(0%)	0	(0%)	<5	**
Undetermined*	<5	**	0	(0%)	0	(0%)	0	(0%)	<5	**
Female Subtotal	17	(89%)	<5	**	0	(0%)	0	(0%)	19	(100%)
TOTAL	White		Black		Hispanic		Other or Unknown		Risk Total	
Male-Male sex	52	(58%)	5	(23%)	<5	**	0	(0%)	59	(51%)
Injecting Drug Use	<5	**	<5	**	0	(0%)	0	(0%)	5	(4%)
Male-Male Sex/IDU	<5	**	<5	**	0	(0%)	0	(0%)	<5	**
Blood Products	<5	**	<5	**	0	(0%)	0	(0%)	<5	**
Heterosexual	19	(21%)	<5	**	0	(0%)	0	(0%)	22	(19%)
<i>HRH</i>	15	(17%)	<5	**	0	(0%)	0	(0%)	17	(15%)
<i>PH-Female</i>	<5	**	<5	**	0	(0%)	0	(0%)	5	(4%)
Perinatal	<5	**	0	(0%)	0	(0%)	0	(0%)	<5	**
Undetermined	12	(13%)	9	(41%)	<5	**	0	(0%)	22	(19%)
<i>PH-Male</i>	7	(8%)	7	(32%)	0	(0%)	0	(0%)	14	(12%)
<i>Unknown</i>	5	(6%)	<5	**	<5	**	0	(0%)	8	(7%)
RACE TOTAL	90	(78%)	22	(19%)	<5	**	0	(0%)	115	(100%)

*In the male subset all cases in the heterosexual category are HRH because the PH-Female category is not applicable to males and, likewise, in the female subset, all cases in the undetermined category are of unknown risk because the PH-Male category is not applicable to females. See page 4 for additional details about the changes to Michigan's risk categories.

**TABLE 3. Prevalent HIV/AIDS Cases in JACKSON CO.
According to City of Residence at Diagnosis**

<i>CITY OF RESIDENCE AT DIAGNOSIS</i>	<i>EST PREV*</i>	<i>REPORTED PREVALENCE[†]</i>			
		<i>HIV, not AIDS</i>	<i>AIDS</i>	<i>Total</i>	
				<i>Number</i>	<i>Percent</i>
Brooklyn	10	<5	<5	<5	**
Cement City	10	<5	0	<5	**
Concord	10	<5	0	<5	**
Grass Lake	10	<5	<5	<5	**
Jackson	130	47	45	92	(80%)
Michigan Center	10	<5	<5	<5	**
Munith	10	<5	<5	<5	**
Parma	10	0	<5	<5	**
Pleasant Lake	10	<5	<5	<5	**
Rives Junction	10	0	<5	<5	**
Spring Arbor	10	0	<5	<5	**
Springport	10	<5	0	<5	**
UNKNOWN	10	<5	0	<5	**
TOTAL	160	58	57	115	(100%)

*See page 5 for descriptions of prevalence estimate calculations.

[†] Includes reports that contain patient name or are otherwise unduplicated. <5 and "**" = 1, 2, 3, or 4 cases.

TABLE 3. New Diagnoses, Deaths, and Prevalence of HIV/AIDS in JACKSON CO. by Year

Year	<i>HIV/AIDS</i>				<i>AIDS</i>		
	New HIV Diagnoses	Deaths	Prevalence		New AIDS Diagnoses	Deaths	Prevalence
1983	0	0	0		0	0	0
1984	0	0	0		0	0	0
1985	3	0	3		0	0	0
1986	6	2	7		3	2	1
1987	7	0	14		0	0	1
1988	4	1	17		2	1	2
1989	10	3	24		7	3	6
1990	13	1	36		7	1	12
1991	8	3	41		5	3	14
1992	10	4	47		7	3	18
1993	3	6	44		6	6	18
1994	9	7	46		8	7	19
1995	8	6	48		6	6	19
1996	12	9	51		8	8	19
1997	8	4	55		9	3	25
1998	6	3	58		7	3	29
1999	4	1	61		4	1	32
2000	6	2	65		3	1	34
2001	12	3	74		7	3	38
2002	9	1	82		4	1	41
2003	14	4	92		8	3	46
2004	6	1	97		1	0	47
2005	8	0	105		1	0	48
2006	4	1	108		4	0	52
2007	7	0	115		5	0	57
TOTAL	177	62			112	55	

The prevalence of HIV in Michigan has steadily increased, since persons with HIV are living longer. This is largely due to improved anti-retroviral therapy.

The increase in HIV prevalence is also reflected in Figure 1 on page 5, which shows that the number of persons diagnosed, while stable for the last several years, is greater than the number of deaths each year. This directly contributes to the increase in prevalence. The current reported prevalence of HIV/AIDS in JACKSON CO. is 115. The prevalence of AIDS, which is a subset of HIV/AIDS prevalence, is 57.

As implied, the HIV/AIDS section displays data on all persons with HIV, including those with AIDS, as well as those who have not been diagnosed with AIDS. Thus, persons represented in the AIDS section are also represented in the HIV/AIDS section. The number of reported deaths includes deaths directly attributable to presence of HIV/AIDS as well as deaths due to other causes.

NOTE: Reporting for recent years may not be complete. Data are not adjusted to account for reporting delays.




TABLE 4. Comparison of HIV/AIDS Prevalence in JACKSON CO. to State

<i>Residence at Diagnosis</i>	<i>Estimated Prevalence</i>	<i>Reported Prevalence</i>				<i>Cumulative AIDS</i>
		<i>HIV, not AIDS</i>	<i>AIDS</i>	<i>Total</i>	<i>Rate</i>	
Jackson Co.	160	58	57	115	70	112
Michigan Total	18,000	6,325	6,995	13,320	132	15,579

ADDITIONAL NOTES

Prevalence Estimate Calculations

HIV prevalence estimates include all persons living in JACKSON CO. at diagnosis of HIV or AIDS and incorporates an approximation of those not reported or not yet diagnosed. All estimates are rounded to the nearest ten, and the minimum estimate given is 10. Prevalence estimates were most recently revised in January 2008, so the estimates in this report may vary in comparison to reports published before the revision.

Risk Groups

In June 2007, the CSTE issued a position statement proposing the creation of a "presumed heterosexual" category. As a result, the following changes were made to Michigan's HIV risk categories.

Heterosexual risk is made up of two sub-categories, (1) HRH [High Risk Heterosexual]: Males and females whose sexual partners are known to be HIV-infected or at high risk for HIV. The partners meet one of the following criteria: a history of sexual contact with bi-sexual males (for females), IDU, hemophiliacs, HIV+ transfusion recipients or other HIV+ persons of unknown risk and (2) PH [Presumed Heterosexual]-female: Females who are not injection drug users and whose only reported risk is heterosexual contact, but their male partners' risk and HIV status are unknown.

Undetermined risk is made up of two sub-categories, (1) PH [Presumed Heterosexual]-Male: Males whose only reported risk is heterosexual contact, and their female partners' risk and HIV status are unknown and (2) Unknown: Males and Females with no identified risk.



Michigan Department of Community Health
HIV/STD/VH/TB Epidemiology Section
Division of Communicable Disease
Bureau of Epidemiology

Jennifer M. Granholm, Governor
Janet Olszewski, Director
<http://www.michigan.gov/hivstd>

Lansing Office 201 Townsend St., 5th Floor
Lansing, MI 48913
517-335-8165

Detroit Office 1151 Taylor St., Room 211B
Detroit, MI 48202
313-876-0353