

**TABLE 1: Characteristics of WASHTENAW CO.  
 Residents Living with HIV or AIDS as of July 1, 2007**

	Estimate of HIV Prevalence <sup>1</sup>	Reported Living with AIDS <sup>2</sup>		Reported Living with HIV not AIDS <sup>2</sup>	
		Number	Percent	Number	Percent
WASHTENAW CO. TOTAL	<b>600</b>	210	100%	219	100%
<b>SEX</b>					
Male	<b>500</b>	185	88%	175	80%
Female	<b>100</b>	25	12%	44	20%
<b>BEHAVIOR</b>					
Male-Male sex	<b>370</b>	126	60%	137	63%
Injecting Drug Use	<b>30</b>	13	6%	8	4%
Male-Male Sex/IDU	<b>30</b>	14	7%	10	5%
Blood Products	<b>10</b>	<5	*	0	0%
Heterosexual <sup>3</sup>	<b>100</b>	29	14%	44	20%
<i>HRH</i>	<b>80</b>	24	11%	32	15%
<i>PH-Female</i>	<b>20</b>	5	2%	12	5%
Perinatal	<b>10</b>	<5	*	<5	*
Undetermined <sup>4</sup>	<b>60</b>	23	11%	19	9%
<i>PH-Male</i>	<b>40</b>	17	8%	12	5%
<i>Unknown</i>	<b>20</b>	6	3%	7	3%
<b>AGE AT HIV DIAGNOSIS</b>					
0 -12 years	<b>10</b>	<5	*	<5	*
13 -19 years	<b>10</b>	<5	*	<5	*
20 -24 years	<b>70</b>	27	13%	26	12%
25 -29 years	<b>120</b>	38	18%	48	22%
30 -39 years	<b>230</b>	79	38%	86	39%
40 -49 years	<b>130</b>	48	23%	47	21%
50 -59 years	<b>30</b>	13	6%	7	3%
60 years and over	<b>10</b>	<5	*	<5	*
Unspecified	Not Applicable	0	0%	0	0%
<b>RACE / ETHNICITY</b>					
White, Non-Hisp.	<b>330</b>	115	55%	124	57%
Black, Non-Hisp.	<b>230</b>	84	40%	83	38%
Hispanic	<b>30</b>	9	4%	10	5%
Asian	<b>10</b>	<5	*	0	0%
American Indian	<b>10</b>	0	0%	0	0%
Unspecified/Other/Multi-race	Not Applicable	<5	*	<5	*

1. This estimate includes all persons living in WASHTENAW CO. at diagnosis of HIV or AIDS, including those not reported or not yet diagnosed. All estimates are rounded to the nearest ten, and the minimum estimate given is 10. The formula used to gain these estimates was updated October 2003, January 2004 and January 2007 so that numbers are only rounded at the end of the estimate calculation and so that cases with no identified risk are no longer part of the equation. This may result in minor variations when compared to past estimates.

2. Includes reports that contain patient name or are otherwise unduplicated. <5 and '\*' = 1, 2, 3, or 4 cases.

3. Heterosexual risk is made up of two sub-categories, (1) HRH [High Risk Heterosexual]: Males and females whose sexual partners are known to be HIV-infected or at high risk for HIV. The partners meet one of the following criteria: a history of sexual contact with bi-sexual males (for females), IDU, hemophiliacs, HIV+ transfusion recipients or other HIV+ persons of unknown risk and (2) PH [Presumed Heterosexual]-female: Females whose only reported risk is heterosexual contact, and their male partners' risk and HIV status are unknown.

4. Undetermined risk is made up of two sub-categories, (1) PH [Presumed Heterosexual]-Male: Males whose only reported risk is heterosexual contact, and their female partners' risk and HIV status are unknown and (2) Unknown: Males and Females with no identified risk.

**TABLE 2: Washtenaw County Residents Reported Living with HIV or AIDS: Sex by Race by Behavior**

<b>MALES:</b>	<b>White</b>		<b>Black</b>		<b>Hispanic</b>		<b>Other or Unknown</b>		<b>TOTAL</b>	
Male-Male Sex	178	82%	74	59%	10	67%	<5	*	263	73%
Injecting Drug Use	5	2%	12	10%	0	0%	0	0%	17	5%
Male-Male Sex/IDU	17	8%	7	6%	0	0%	0	0%	24	7%
Blood Recipient	<5	*	0	0%	0	0%	0	0%	<5	*
Heterosexual <sup>1</sup>	<5	*	9	7%	<5	*	0	0%	11	3%
Perinatal	0	0%	<5	*	<5	*	0	0%	<5	*
Undetermined <sup>2</sup>	14	6%	23	18%	<5	*	0	0%	40	11%
<i>PH-Male</i>	9	4%	17	13%	<5	*	0	0%	29	8%
<i>Unknown</i>	5	2%	6	5%	0	0%	0	0%	11	3%
<b>Male Subtotal</b>	<b>218</b>	<b>61%</b>	<b>126</b>	<b>35%</b>	<b>15</b>	<b>4%</b>	<b>&lt;5</b>	<b>*</b>	<b>360</b>	<b>100%</b>
<b>FEMALES:</b>	<b>White</b>		<b>Black</b>		<b>Hispanic</b>		<b>Other or Unknown</b>		<b>TOTAL</b>	
Injecting Drug Use	<5	*	<5	*	0	0%	0	0%	<5	*
Blood Recipient	<5	*	0	0%	0	0%	0	0%	<5	*
Heterosexual <sup>1</sup>	18	86%	37	90%	<5	*	<5	*	62	90%
<i>HRH</i>	15	71%	25	61%	<5	*	<5	*	45	65%
<i>PH-Female</i>	<5	*	12	29%	<5	*	<5	*	17	25%
Perinatal	0	0%	0	0%	0	0%	0	0%	0	0%
Undetermined <sup>2</sup>	0	0%	<5	*	0	0%	0	0%	<5	*
<b>Female Subtotal</b>	<b>21</b>	<b>30%</b>	<b>41</b>	<b>59%</b>	<b>&lt;5</b>	<b>*</b>	<b>&lt;5</b>	<b>*</b>	<b>69</b>	<b>100%</b>
<b>Total</b>	<b>White</b>		<b>Black</b>		<b>Hispanic</b>		<b>Other or Unknown</b>		<b>TOTAL</b>	
Male-Male Sex	178	74%	74	44%	10	53%	<5	*	263	61%
Injecting Drug Use	7	3%	14	8%	0	0%	0	0%	21	5%
Male-Male Sex/IDU	17	7%	7	4%	0	0%	0	0%	24	6%
Blood Recipient	<5	*	0	0%	0	0%	0	0%	<5	*
Heterosexual <sup>1</sup>	19	8%	46	28%	5	26%	<5	*	73	17%
<i>HRH</i>	16	7%	34	20%	<5	*	<5	*	56	13%
<i>PH-Female</i>	<5	*	12	7%	<5	*	<5	*	17	4%
Perinatal	0	0%	<5	*	<5	*	0	0%	<5	*
Undetermined <sup>2</sup>	14	6%	25	15%	<5	*	0	0%	42	10%
<i>PH-Male</i>	9	4%	17	10%	<5	*	0	0%	29	7%
<i>Unknown</i>	5	2%	8	5%	0	0%	0	0%	13	3%
<b>GRAND TOTAL</b>	<b>239</b>	<b>56%</b>	<b>167</b>	<b>39%</b>	<b>19</b>	<b>4%</b>	<b>&lt;5</b>	<b>*</b>	<b>429</b>	<b>100%</b>

1. Heterosexual risk is made up of two sub-categories, (1) HRH [High Risk Heterosexual]: Males and females whose sexual partners are known to be HIV-infected or at high risk for HIV. The partners meet one of the following criteria: a history of sexual contact with bi-sexual males (for females), IDU, hemophiliacs, HIV+ transfusion recipients or other HIV+ persons of unknown risk and (2) PH [Presumed Heterosexual]-female: Females whose only reported risk is heterosexual contact, and their male partners' risk and HIV status are unknown. \*\*In the male subset, all cases in the heterosexual category are HRH because the PH-Female category is not applicable to males.

2. Undetermined risk is made up of two sub-categories, (1) PH [Presumed Heterosexual]-Male: Males whose only reported risk is heterosexual contact, and their female partners' risk and HIV status are unknown and (2) Unknown: Males and Females with no identified risk. \*\*In the female subset, all cases in the undetermined category of unknown risk because the PH-Male category is not applicable to females.

NOTE: Includes reports that contain patient name or are otherwise unduplicated. <5 and "\*" = 1, 2, 3, or 4 cases.


**Table 3: Washtenaw County HIV/AIDS Prevalence  
 through July 1, 2007**

Year	HIV/AIDS Deaths/Year	HIV Incidence	AIDS Incidence	HIV/AIDS Prevalence	AIDS Prevalence
1981	0	0	0	0	0
1982	0	0	0	0	0
1983	0	1	0	1	0
1984	0	1	0	2	0
1985	3	22	7	21	4
1986	5	21	6	37	5
1987	3	30	13	64	15
1988	9	33	18	88	24
1989	9	37	20	116	35
1990	10	32	22	138	47
1991	14	44	31	168	64
1992	19	35	37	184	82
1993	19	30	34	195	97
1994	30	18	21	183	88
1995	24	29	33	188	97
1996	17	47	25	218	108
1997	10	23	17	231	115
1998	9	27	18	249	125
1999	8	27	17	268	135
2000	2	17	11	283	144
2001	8	24	14	299	151
2002	4	22	14	317	161
2003	1	28	12	344	172
2004	4	32	14	372	182
2005	3	27	21	396	201
2006	8	30	15	418	210
2007	0	11	0	429	210
<b>TOTAL</b>	<b>219</b>	<b>648</b>	<b>420</b>		

\*\*\*Reported HIV/AIDS prevalence (all persons currently living with an HIV or AIDS diagnosis) is 429 persons through July 1, 2007\*\*\*

\*\*\*Reported AIDS prevalence (all persons currently living with an AIDS diagnosis & note this is a subset of the HIV/AIDS prevalence listed above) is 210 persons through July 1, 2007\*\*\*

Prevalence of Washtenaw Co. residents living with HIV/AIDS and AIDS



Information based on the July 1, 2007 Michigan Quarterly HIV/AIDS Analysis

Residence at time of diagnosis	Estimate of HIV Prevalence	Reported prevalence rate	Persons living with AIDS	Persons living with HIV not AIDS
Washtenaw Co.	600	125	210	219
Michigan Total	17,000	129	6,805	6,177

**Note:** This estimate includes all persons living with HIV or AIDS, including those not yet diagnosed.

Rates are calculated per 100,000 population in 2000.

Data includes reports of HIV infection and AIDS that contain patient name or are otherwise unduplicated.