


Administering Influenza Vaccines


Intramuscular (IM) Administration of Trivalent Inactivated Influenza Vaccines (TIV): TIV (IM) and TIV High Dose

1. Use a needle long enough to reach deep into the muscle. Infants aged 6 through 11 months: 1"; 1 through 2 years: 1-1 $\frac{1}{4}$ "; children and adults 3 years and older: 1-1 $\frac{1}{2}$ ".
2. Choose the appropriate site (see chart below). With your left hand¹, bunch up the muscle.
3. With your right hand¹, insert the needle at a 90° angle to the skin with a quick thrust.
4. Push down on the plunger and inject the entire contents of the syringe. There is no need to aspirate.
5. Remove the needle and simultaneously apply pressure to the injection site with dry cotton ball or gauze. Hold in place for several seconds.
6. If there is any bleeding cover the injection site with a bandage.
7. Put the used syringe in a sharps container.



Intranasal Administration of Live Attenuated Influenza Vaccine (LAIV)

1. FluMist® (LAIV) is for intranasal administration only. Do not inject FluMist®.
2. Remove rubber tip protector. Do not remove dose-divider clip at the other end of the sprayer.
3. With the patient in an upright position (i.e., head not tilted back), place the tip inside the nostril to ensure LAIV is into the nose. The patient should breathe normally.
4. With a single motion, depress plunger as rapidly as possible until the dose-divider clip prevents you from going further.
5. Pinch and remove the dose-divider clip from the plunger.
6. Place the tip just inside the other nostril, and with a single motion depress plunger as rapidly as possible to deliver the remaining vaccine.
7. Dispose of the applicator in a sharps container.



TIV Dosage

Age	Dosage ²	Site ³	Needle Gauge
6-35 months	0.25 mL	Anterolateral thigh	22-25 g
3-8 years	0.5 mL	Anterolateral thigh or deltoid	22-25 g
9 years & older	0.5mL	Anterolateral thigh or deltoid	22-25 g

LAIV Dosage

Age	Dosage ²	Route	Site
Healthy, non-pregnant persons 2-49 years	0.2 mL Spray 0.1mL dose into each nostril as indicated by the dose-divider clip on the sprayer	Intranasal	Each nostril

¹ Use the opposite hand if you are left handed

² Children needing 2 doses of flu vaccine should receive the age-appropriate vaccine & dosage each time. Refer to: "Who needs 2 Doses of 2012-13 Seasonal Flu Vaccine" at www.michigan.gov/flu

³ Needle size and site of injection assessment must be made for each person on the basis of the size of the muscle and the thickness of adipose tissue at the injection site