

**Prevalence Estimates for Risk Factors and Health Indicators  
by Race-Ethnicity**

**State of Michigan**

**Selected Tables**

**Michigan Behavioral Risk Factor Survey**

**2020**

Lifecourse Epidemiology and Genomics Division  
Bureau of Epidemiology and Population Health  
Michigan Department of Health and Human Services

## Table of Contents

Introduction.....	2
Table 1: Fair or Poor General Health <sup>a</sup> by Race-Ethnicity .....	3
Table 2: Poor Physical Health on at Least 14 Days in the Past Month <sup>a</sup> by Race-Ethnicity .....	4
Table 3: Poor Mental Health on at Least 14 Days in the Past Month <sup>a</sup> by Race-Ethnicity .....	5
Table 4: Activity Limitation on at Least 14 Days in the Past Month <sup>a</sup> by Race-Ethnicity .....	6
Table 5: Disability <sup>a</sup> by Race-Ethnicity .....	7
Table 6: Obesity <sup>a</sup> by Race-Ethnicity.....	8
Table 7: No Health Care Coverage among Adults Aged 18 - 64 Years <sup>a</sup> by Race-Ethnicity .....	9
Table 8: No Personal Health Care Provider <sup>a</sup> by Race-Ethnicity .....	10
Table 9: No Health Care Access during Past 12 Months Due to Cost <sup>a</sup> by Race-Ethnicity .....	11
Table 10: No Leisure Time Physical Activity <sup>a</sup> by Race-Ethnicity .....	12
Table 11: Current Smoking <sup>a</sup> by Race-Ethnicity.....	13
Table 12: Current E-Cigarette Use <sup>a</sup> by Race-Ethnicity .....	14
Table 13: Current Hookah Use <sup>a</sup> by Race-Ethnicity .....	15
Table 14: Any Alcohol Consumption <sup>a</sup> by Race-Ethnicity .....	16
Table 15: Heavy Drinking <sup>a</sup> by Race-Ethnicity .....	17
Table 16: Binge Drinking <sup>a</sup> by Race-Ethnicity .....	18
Table 17: Drove Motor Vehicle After Drinking <sup>a</sup> by Race-Ethnicity .....	19
Table 18: Seatbelt Use <sup>a</sup> by Race-Ethnicity .....	20
Table 19: No Routine Checkup in Past Year <sup>a</sup> by Race-Ethnicity.....	21
Table 20: Mammography in the Past Two Years among Women Aged 40 and Older <sup>a</sup> by Race-Ethnicity .....	22
Table 21: Appropriate Timed Pap Test among Women Aged 18 and Older <sup>a</sup> by Race-Ethnicity.....	23
Table 22: Prostate-Specific Antigen (PSA) Test Within the Past Year among Men Aged 50 and Older <sup>a</sup> by Race-Ethnicity.....	24
Table 23: Appropriate Colorectal Cancer Screening among Adults Aged 50 and Older <sup>a</sup> by Race-Ethnicity .....	25
Table 24: No Dental Visit in the Past Year <sup>a</sup> by Race-Ethnicity .....	26
Table 25: Had Flu Vaccine in Past Year among Adults 65 Years and Older <sup>a</sup> by Race-Ethnicity .....	27
Table 26: Ever Had Pneumonia Vaccine among Adults 65 Years and Older <sup>a</sup> by Race-Ethnicity .....	28
Table 27: Ever Had an HIV Test among Adults 18 – 64 Years of Age <sup>a</sup> by Race-Ethnicity .....	29
Table 28: Lifetime Asthma Prevalence - Adult <sup>a</sup> by Race-Ethnicity .....	30
Table 29: Current Asthma Prevalence - Adult <sup>a</sup> by Race-Ethnicity .....	31
Table 30: Ever Told Chronic Obstructive Pulmonary Disease (COPD) <sup>a</sup> by Race-Ethnicity .....	32
Table 31: Ever Told Arthritis <sup>a</sup> by Race-Ethnicity .....	33
Table 32: Ever Told Heart Attack <sup>a</sup> by Race-Ethnicity .....	34
Table 33: Ever Told Angina or Coronary Heart Disease <sup>a</sup> by Race-Ethnicity .....	35
Table 34: Ever Told Stroke <sup>a</sup> by Race-Ethnicity .....	36
Table 35: Ever Told Any Cardiovascular Disease <sup>a</sup> by Race-Ethnicity .....	37
Table 36: Ever Told Skin Cancer <sup>a</sup> by Race-Ethnicity.....	38
Table 37: Ever Told Any Other Type of Cancer <sup>a</sup> by Race-Ethnicity .....	39
Table 38: Ever Told Cancer <sup>a</sup> by Race-Ethnicity.....	40
Table 39: Ever Told Diabetes <sup>a</sup> by Race-Ethnicity .....	41
Table 40: Ever Told Kidney Disease <sup>a</sup> by Race-Ethnicity .....	42
Table 41: Ever Told Depression <sup>a</sup> by Race-Ethnicity .....	43

## Introduction

The Michigan Behavioral Risk Factor Survey (MiBRFS) is an annual, statewide telephone survey of Michigan adults aged 18 years and older that is conducted through a collaborative effort among the Population Health Surveillance Branch (PHSB) of the Centers for Disease Control and Prevention (CDC), the Michigan State University Institute for Public Policy and Social Research (IPPSR), and the Michigan Department of Health and Human Services. Michigan BRFSS data contribute to the national [Behavioral Risk Factor Surveillance System](#) that is managed by the PHSB at the CDC.

In 2020, the total sample size of completed interviews for the core survey was 7,269 (land line = 2,205; cell phone = 5,064) which consisted of 5,872 White, non-Hispanics, 665 Black, non-Hispanics, 399 Other/Multiracial, non-Hispanics, and 198 Hispanics. Due to relatively small racial-ethnic subgroup sample sizes we were required to combine demographic groups in this analysis. Some of the estimates for these racial-ethnic subgroups are relatively imprecise (i.e., have large confidence intervals) and should be used with caution. Furthermore, only comparisons between estimates with non-overlapping 95% confidence intervals should be considered significantly different.

For the 2020 Michigan BRFSS, the sample of land line telephone numbers that were utilized for data collection was selected using a list-assisted, random-digit-dialed methodology with a disproportionate stratification based on phone bank density, and whether or not the phone numbers were directory listed. The sample of cell phone numbers used within the 2020 Michigan BRFSS data collection process was randomly selected from dedicated cellular telephone banks sorted on the basis of area code and exchange within the State of Michigan.

Iterative proportional fitting or raking was the weighting methodology used to make the 2020 Michigan BRFSS data representative of the current Michigan adult population. Estimates based on this methodology were weighted to adjust for the probabilities of selection and a raking adjustment factor that adjusted for the distribution of Michigan adults by telephone source (land line vs. cell phone), detailed race/ethnicity, education level, marital status, age by gender, gender by race/ethnicity, age by race/ethnicity, and renter/owner status.

State-specific, population-based prevalence estimates and asymmetric confidence intervals were calculated for indicators of health status, health risk behaviors, clinical preventive practices, and chronic conditions among the adult population in Michigan. Respondents who refused to answer a question were removed from the denominator during the analysis of such questions. Those who responded “Don’t Know/Not Sure” were also excluded from the denominator unless indicated otherwise. All analyses were performed in SAS-Callable SUDAAN®, a statistical computing program that was designed for complex sample surveys.

In an effort to provide the most reliable information possible, prevalence estimates are suppressed if they are based on a denominator of less than 50 respondents or have a relative standard error of greater than 30%.

Due to the BRFSS methodology changes that took place in 2011, the estimates provided within the following tables should only be compared to Michigan BRFSS estimates from 2011-2017 and not to Michigan BRFSS estimates from years prior to 2011.

If you have any questions regarding any of the estimates within the following tables, please contact Yan Tian at [TianY@michigan.gov](mailto:TianY@michigan.gov).

2020 Michigan BRFSS Estimates by Race-Ethnicity  
September 2021

**Table 1: Fair or Poor General Health<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	15.5	(14.4-16.7)	14.5	(13.3-15.8)	21.9	(18.4-26.0)	13.7	(9.6-19.1)	16.9	(11.5-24.0)
<b>Age</b>										
18 - 49	11.1	(9.7-12.7)	9.9	(8.4-11.7)	17.2	(12.8-22.8)	8.5	(4.7-14.8)	14.6	(8.9-22.9)
50 +	20.2	(18.5-21.9)	18.6	(16.9-20.5)	27.3	(21.9-33.5)	28.3	(18.8-40.3)	25.6	(14.1-41.8)
<b>Gender</b>										
Male	13.4	(11.9-15.1)	12.8	(11.2-14.6)	20.7	(15.7-26.9)	11.1	(6.7-17.7)	-- <sup>c</sup>	-- <sup>c</sup>
Female	17.6	(16.0-19.3)	16.2	(14.5-18.0)	23.0	(18.2-28.6)	17.1	(10.4-26.9)	23.6	(15.4-34.5)
<b>Education</b>										
High school graduate or less	22.1	(19.9-24.5)	20.8	(18.3-23.5)	25.0	(19.3-31.7)	30.5	(19.1-45.1)	25.0	(15.9-37.1)
Some college or more	11.4	(10.3-12.5)	10.8	(9.7-12.0)	19.2	(14.9-24.3)	6.9	(4.6-10.3)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	26.0	(23.4-28.8)	25.9	(22.8-29.2)	27.1	(21.5-33.6)	27.5	(17.1-41.0)	21.1	(11.9-34.5)
\$35,000 +	9.4	(8.3-10.6)	9.0	(7.9-10.3)	13.1	(8.8-19.1)	7.5	(4.5-12.2)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported that their health, in general, was either fair or poor.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 2: Poor Physical Health on at Least 14 Days in the Past Month<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	11.3	(10.4-12.4)	11.3	(10.3-12.5)	12.1	(9.2-15.9)	11.9	(8.1-17.0)	9.1	(5.2-15.3)
<b>Age</b>										
18 - 49	7.6	(6.4-9.0)	7.4	(6.1-9.0)	9.9	(6.2-15.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	15.3	(13.8-17.0)	14.8	(13.2-16.6)	14.8	(10.5-20.4)	27.2	(18.0-38.9)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	10.1	(8.7-11.7)	10.1	(8.5-11.8)	10.5	(6.6-16.5)	10.9	(6.4-18.1)	-- <sup>c</sup>	-- <sup>c</sup>
Female	12.6	(11.2-14.0)	12.6	(11.1-14.2)	13.5	(9.6-18.6)	13.1	(7.7-21.6)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	15.3	(13.3-17.5)	15.2	(13.0-17.7)	14.5	(9.6-21.3)	26.4	(15.6-41.0)	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	8.8	(7.9-9.9)	8.9	(7.9-10.1)	10.2	(7.2-14.1)	6.4	(4.0-10.3)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	19.1	(16.7-21.7)	19.3	(16.6-22.3)	16.8	(11.9-23.2)	32.4	(21.1-46.1)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	6.9	(6.0-7.9)	7.6	(6.6-8.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported 14 or more days of physical health that was not good, which includes physical illness and injury, during the past 30 days.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 3: Poor Mental Health on at Least 14 Days in the Past Month<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	15.8	(14.7-17.0)	15.8	(14.5-17.1)	16.4	(13.0-20.4)	16.5	(12.0-22.4)	14.9	(10.0-21.6)
<b>Age</b>										
18 - 49	19.8	(18.0-21.9)	20.8	(18.6-23.2)	19.6	(14.4-26.1)	15.2	(10.0-22.5)	15.6	(9.9-23.7)
50 +	12.0	(10.8-13.4)	11.6	(10.3-13.1)	12.8	(9.1-17.6)	20.6	(12.4-32.0)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	11.6	(10.1-13.2)	11.4	(9.9-13.2)	11.9	(7.7-17.9)	13.1	(7.9-20.9)	-- <sup>c</sup>	-- <sup>c</sup>
Female	19.9	(18.2-21.7)	20.0	(18.1-22.0)	20.2	(15.5-26.0)	21.1	(13.9-30.7)	17.6	(10.8-27.4)
<b>Education</b>										
High school graduate or less	18.6	(16.5-21.0)	18.6	(16.2-21.3)	14.4	(9.6-21.1)	29.9	(18.4-44.7)	22.4	(13.9-34.0)
Some college or more	14.1	(12.9-15.4)	14.1	(12.8-15.6)	17.9	(13.7-23.2)	11.5	(7.6-16.9)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	22.4	(19.8-25.2)	23.1	(20.0-26.5)	19.5	(14.3-26.0)	28.2	(17.1-42.7)	21.2	(12.2-34.2)
\$35,000 +	12.0	(10.7-13.4)	12.3	(10.9-13.9)	12.3	(8.1-18.4)	8.7	(5.0-14.7)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported 14 or more days of mental health that was not good, which includes stress, depression, and problems with emotions, during the past 30 days.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 4: Activity Limitation on at Least 14 Days in the Past Month<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	9.1	(8.3-10.1)	8.5	(7.6-9.5)	10.7	(8.1-14.0)	14.1	(9.8-20.1)	8.7	(5.0-14.7)
<b>Age</b>										
18 - 49	7.8	(6.6-9.2)	7.2	(5.9-8.7)	8.5	(5.5-12.9)	11.8	(6.8-19.5)	-- <sup>c</sup>	-- <sup>c</sup>
50 +	10.5	(9.3-11.8)	9.7	(8.6-11.1)	12.9	(8.8-18.4)	21.2	(13.2-32.3)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	7.2	(6.1-8.5)	6.4	(5.3-7.7)	10.4	(6.5-16.2)			-- <sup>c</sup>	-- <sup>c</sup>
Female	10.9	(9.7-12.3)	10.6	(9.2-12.1)	10.9	(7.8-15.1)	18.6	(11.8-28.2)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	11.9	(10.2-13.8)	11.1	(9.3-13.1)	11.2	(7.3-16.8)	31.4	(19.1-47.1)	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	7.4	(6.5-8.4)	7.0	(6.1-8.1)	10.3	(7.1-14.8)	7.9	(4.8-12.6)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	16.3	(14.1-18.8)	15.7	(13.2-18.6)	14.4	(10.2-20.0)	30.7	(18.9-45.6)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	5.3	(4.5-6.2)	5.3	(4.5-6.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported 14 or more days in the past 30 days in which either poor physical health or poor mental health kept respondents from doing their usual activities, such as self-care, work, and recreation.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 5: Disability<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	27.5	(26.2-28.9)	26.7	(25.2-28.2)	33.0	(28.5-37.8)	26.4	(20.7-33.1)	27.7	(20.6-36.1)
<b>Age</b>										
18 - 49	20.5	(18.5-22.6)	18.9	(16.7-21.2)	28.5	(22.5-35.3)	18.0	(12.0-26.2)	24.0	(16.5-33.6)
50 +	34.7	(32.7-36.6)	33.5	(31.5-35.5)	38.4	(32.0-45.3)	46.8	(35.5-58.4)	41.1	(26.2-57.8)
<b>Gender</b>										
Male	25.4	(23.4-27.5)	24.6	(22.4-26.8)	30.8	(24.0-38.5)	22.5	(15.6-31.4)	29.9	(19.6-42.6)
Female	29.5	(27.6-31.5)	28.7	(26.6-30.9)	34.9	(29.2-41.2)	31.5	(22.5-42.0)	26.1	(17.1-37.7)
<b>Education</b>										
High school graduate or less	39.0	(36.2-41.8)	37.2	(34.1-40.4)	41.8	(34.3-49.7)	56.9	(42.6-70.1)	41.5	(29.8-54.3)
Some college or more	20.2	(18.9-21.6)	20.3	(18.9-21.9)	24.8	(20.0-30.5)	14.8	(10.4-20.7)	13.9	(8.1-23.0)
<b>Household Income</b>										
< \$35,000	43.8	(40.7-47.0)	44.5	(40.9-48.2)	41.4	(34.4-48.7)	52.9	(40.3-65.2)	37.8	(25.3-52.1)
\$35,000 +	16.6	(15.2-18.1)	17.4	(15.9-19.1)	17.0	(11.8-23.9)	6.7	(3.9-11.4)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported being limited in any activities because of physical, mental, or emotional problems, or reported that they required use of special equipment (such as a cane, a wheelchair, a special bed, or a special telephone) due to a health problem.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.



**Table 6: Obesity<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	35.2	(33.7-36.7)	34.2	(32.7-35.9)	42.7	(38.0-47.7)	22.6	(17.4-28.7)	46.7	(37.6-56.0)
<b>Age</b>										
18 - 49	33.6	(31.3-35.9)	32.6	(30.0-35.2)	43.7	(36.8-50.8)	18.3	(12.8-25.5)	41.3	(30.8-52.8)
50 +	37.1	(35.2-39.0)	35.8	(33.8-37.8)	42.0	(35.4-48.8)	33.0	(22.7-45.3)	64.8	(49.2-77.8)
<b>Gender</b>										
Male	33.4	(31.4-35.6)	34.1	(31.9-36.4)	34.3	(27.5-41.9)	16.2	(11.2-22.9)	45.4	(33.3-58.0)
Female	37.1	(34.9-39.2)	34.4	(32.1-36.7)	50.3	(44.0-56.7)	31.1	(22.0-42.1)	47.8	(34.9-61.0)
<b>Education</b>										
High school graduate or less	36.9	(34.2-39.7)	35.7	(32.7-38.7)	38.5	(31.2-46.3)	36.6	(24.3-51.0)	49.2	(36.5-62.1)
Some college or more	34.1	(32.4-35.9)	33.4	(31.6-35.3)	46.3	(40.2-52.4)	16.7	(12.0-22.7)	44.2	(31.5-57.8)
<b>Household Income</b>										
< \$35,000	39.6	(36.6-42.7)	38.8	(35.3-42.5)	40.1	(33.4-47.2)	31.8	(21.4-44.4)	54.2	(40.0-67.8)
\$35,000 +	34.8	(32.9-36.8)	34.0	(32.0-36.0)	50.9	(43.2-58.6)	19.4	(13.3-27.3)	40.9	(27.7-55.6)

Note: BMI, body mass index, is defined as weight (in kilograms) divided by height (in meters) squared [weight in kg/(height in meters)<sup>2</sup>]. Weight and height were self-reported. Pregnant women were excluded.

<sup>a</sup> The proportion of respondents whose BMI was greater than or equal to 30.0.

<sup>b</sup> Michigan estimates include all racial groups.

**Table 7: No Health Care Coverage among Adults Aged 18 - 64 Years<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	8.4	(7.3-9.6)	7.5	(6.3-8.8)	9.7	(6.9-13.5)	9.1	(5.9-13.7)	14.3	(9.3-21.5)
<b>Age</b>										
18 - 49	9.1	(7.7-10.7)	8.2	(6.7-10.1)	10.6	(6.9-15.9)	9.1	(5.6-14.5)	14.9	(9.3-23.0)
50 - 64	6.5	(5.1-8.2)	5.9	(4.4-7.7)	8.2	(4.6-14.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	10.3	(8.7-12.3)	9.4	(7.6-11.7)	11.7	(7.1-18.6)	11.7	(7.2-18.5)	18.8	(10.4-31.8)
Female	6.3	(5.1-7.6)	5.5	(4.3-7.1)	7.9	(5.0-12.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	11.2	(9.1-13.7)	11.0	(8.6-14.1)	10.8	(6.3-18.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	6.4	(5.4-7.5)	5.4	(4.4-6.6)	8.0	(5.3-11.9)	8.8	(5.2-14.5)	14.4	(8.2-24.1)
<b>Household Income</b>										
< \$35,000	12.4	(10.0-15.2)	12.2	(9.2-15.8)	13.3	(8.8-19.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	5.7	(4.7-6.8)	5.1	(4.2-6.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	14.4	(7.8-25.1)

<sup>a</sup> Among those aged 18-64 years, the proportion who reported having no health care coverage, including health insurance, prepaid plans such as HMOs, or government plans, such as Medicare or Indian Health Services.

<sup>b</sup> Michigan estimates include all racial groups

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 8: No Personal Health Care Provider<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	14.5	(13.3-15.7)	13.2	(11.9-14.5)	14.2	(10.9-18.2)	23.2	(17.8-29.7)	23.9	(17.3-32.0)
<b>Age</b>										
18 - 49	23.0	(20.9-25.1)	22.1	(19.8-24.6)	20.4	(15.1-26.9)	29.0	(21.8-37.4)	29.7	(21.4-39.6)
50 +	5.5	(4.7-6.5)	5.2	(4.4-6.2)	7.5	(4.5-12.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	19.9	(18.0-22.0)	18.5	(16.5-20.8)	18.3	(12.8-25.4)	27.1	(19.5-36.3)	41.0	(28.9-54.4)
Female	9.1	(8.0-10.4)	8.1	(6.8-9.5)	10.6	(7.3-15.2)	18.1	(11.6-27.2)	12.8	(7.2-21.7)
<b>Education</b>										
High school graduate or less	17.7	(15.5-20.2)	16.7	(14.2-19.5)	17.1	(11.7-24.4)	17.9	(10.2-29.6)	31.7	(21.0-44.6)
Some college or more	12.2	(11.1-13.5)	11.0	(9.8-12.4)	11.7	(8.4-16.1)	25.5	(18.9-33.4)	16.2	(9.7-25.7)
<b>Household Income</b>										
< \$35,000	18.3	(15.9-21.0)	16.5	(13.8-19.6)	18.9	(13.6-25.6)	24.6	(15.4-37.0)	28.6	(17.5-43.1)
\$35,000 +	11.9	(10.6-13.4)	11.0	(9.6-12.5)	9.7	(5.8-15.6)	24.0	(16.5-33.5)	19.9	(12.0-31.3)

<sup>a</sup> The proportion who reported that they did not have anyone that they thought of as their personal doctor or health care provider.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 9: No Health Care Access during Past 12 Months Due to Cost<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	7.9	(7.1-8.8)	7.1	(6.2-8.1)	9.0	(6.9-11.8)	11.7	(8.5-16.0)	13.4	(8.7-20.1)
<b>Age</b>										
18 - 49	10.4	(9.1-11.9)	9.7	(8.1-11.5)	11.2	(7.9-15.7)	13.1	(9.1-18.5)	13.4	(8.1-21.4)
50 +	5.4	(4.5-6.3)	4.9	(4.1-5.9)	6.5	(4.1-10.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	7.3	(6.1-8.6)	6.4	(5.2-7.9)	8.7	(5.7-13.1)	12.0	(7.7-18.3)	-- <sup>c</sup>	-- <sup>c</sup>
Female	8.5	(7.4-9.8)	7.8	(6.6-9.2)	9.3	(6.5-13.2)	11.3	(6.9-17.8)	14.0	(7.9-23.4)
<b>Education</b>										
High school graduate or less	8.5	(7.0-10.2)	7.8	(6.1-9.8)	7.7	(4.8-12.2)	11.8	(6.5-20.4)	16.7	(9.4-27.9)
Some college or more	7.5	(6.6-8.5)	6.7	(5.8-7.8)	9.5	(6.7-13.4)	11.7	(7.9-17.0)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	14.0	(12.0-16.3)	13.6	(11.1-16.5)	12.6	(9.0-17.3)	20.2	(12.3-31.2)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	5.2	(4.3-6.2)	5.0	(4.1-6.1)	-- <sup>c</sup>	-- <sup>c</sup>	8.4	(4.9-13.9)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported that in the past 12 months, they could not see a doctor when they needed to due to the cost.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 10: No Leisure Time Physical Activity<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	20.8	(19.6-22.1)	19.1	(17.8-20.4)	30.9	(26.5-35.5)	18.4	(13.9-24.0)	21.2	(15.2-28.9)
<b>Age</b>										
18 - 49	17.2	(15.3-19.1)	14.7	(12.7-16.9)	30.3	(24.2-37.3)	15.8	(10.6-22.9)	18.4	(12.0-27.1)
50 +	24.0	(22.4-25.7)	22.6	(21.0-24.4)	31.5	(25.7-38.0)	24.0	(16.3-33.8)	32.2	(19.0-49.1)
<b>Gender</b>										
Male	19.0	(17.3-20.9)	17.4	(15.6-19.4)	30.0	(23.5-37.3)	16.0	(10.9-23.0)	20.9	(12.7-32.4)
Female	22.2	(20.5-24.0)	20.7	(18.9-22.5)	31.6	(26.1-37.8)	21.6	(14.4-31.1)	21.4	(13.5-32.2)
<b>Education</b>										
High school graduate or less	28.8	(26.3-31.3)	26.7	(24.1-29.5)	40.6	(33.3-48.3)	25.0	(15.2-38.3)	25.9	(16.6-38.0)
Some college or more	15.4	(14.2-16.7)	14.4	(13.1-15.7)	22.2	(17.7-27.5)	15.8	(11.3-21.7)	16.4	(9.8-26.2)
<b>Household Income</b>										
< \$35,000	29.0	(26.3-31.9)	26.4	(23.5-29.6)	38.3	(31.5-45.5)	27.3	(17.5-39.9)	21.2	(11.9-34.8)
\$35,000 +	15.2	(13.9-16.6)	14.8	(13.4-16.4)	20.2	(15.0-26.7)	11.0	(6.9-17.0)	18.3	(10.8-29.4)

<sup>a</sup> The proportion who reported not participating in any leisure time physical activities or exercises such as running, calisthenics, golf, gardening, or walking for exercise in the past month.

<sup>b</sup> Michigan estimates include all racial groups.

**Table 11: Current Smoking<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	18.4	(17.1-19.7)	17.9	(16.5-19.4)	19.5	(15.8-24.0)	18.1	(13.5-24.0)	20.1	(13.9-28.1)
<b>Age</b>										
18 - 49	20.5	(18.6-22.6)	21.0	(18.8-23.5)	19.3	(14.1-25.8)	16.6	(11.1-23.9)	21.8	(14.5-31.3)
50 +	16.1	(14.6-17.8)	15.3	(13.7-17.1)	20.0	(14.9-26.4)	22.4	(14.1-33.8)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	19.4	(17.5-21.4)	18.2	(16.1-20.4)	24.0	(17.8-31.7)	20.2	(13.5-29.0)	26.5	(16.7-39.5)
Female	17.2	(15.6-18.9)	17.6	(15.9-19.6)	15.6	(11.6-20.5)	15.5	(10.0-23.2)	15.4	(8.5-26.4)
<b>Education</b>										
High school graduate or less	27.2	(24.7-29.9)	27.1	(24.2-30.1)	25.0	(18.7-32.6)	39.8	(26.9-54.4)	26.6	(16.9-39.3)
Some college or more	12.6	(11.4-13.9)	12.5	(11.2-13.8)	14.8	(10.8-19.9)	10.0	(6.5-15.0)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	28.5	(25.7-31.5)	29.5	(26.1-33.1)	24.0	(18.2-31.0)	39.3	(27.4-52.5)	25.9	(15.4-40.2)
\$35,000 +	13.8	(12.4-15.3)	14.2	(12.7-15.9)	12.7	(8.4-18.9)	8.2	(4.8-13.5)	14.1	(7.7-24.4)

<sup>a</sup> The proportion who reported that they had ever smoked at least 100 cigarettes (5 packs) in their life and that they smoke cigarettes now, either every day or on some days.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

2020 Michigan BRFSS Estimates by Race-Ethnicity  
September 2021

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	6.4	(5.5-7.6)	6.7	(5.6-8.0)	6.1	(3.4-10.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	11.2	(9.4-13.5)	12.1	(9.9-14.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	1.9	(1.3-2.6)	2.0	(1.4-2.9)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	7.1	(5.6-8.9)	7.1	(5.5-9.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	6.0	(4.8-7.6)	6.4	(4.9-8.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	8.7	(6.6-11.2)	8.4	(6.2-11.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	5.3	(4.3-6.4)	5.8	(4.7-7.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	7.9	(5.7-10.7)	7.4	(5.2-10.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	5.7	(4.5-7.1)	6.4	(5.0-8.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> Among all adults, the proportion who reported that they currently use e-cigarettes or other electronic vaping products, either every day or on some days.  
<sup>b</sup> Michigan estimates include all racial groups.  
<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Due to significant BRFSS methodology changes that took place in 2011, the estimates within these tables should not be compared to Michigan BRFSS estimates from prior to 2011.

**Table 13: Current Hookah Use<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	2.2	(1.7-2.9)	1.9	(1.4-2.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	3.5	(2.5-4.8)	2.9	(1.9-4.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	1.1	(0.7-1.8)	1.0	(0.6-1.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	2.9	(2.1-4.2)	2.5	(1.7-3.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	1.6	(1.1-2.5)	1.3	(0.7-2.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	1.9	(1.1-3.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	2.5	(1.8-3.4)	2.4	(1.7-3.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	3.0	(1.8-4.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	2.0	(1.4-2.9)	1.8	(1.2-2.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> Among all adults, the proportion who reported smoking tobacco using a hookah, narghile, or water pipe on one or more days during the previous thirty days.  
<sup>b</sup> Michigan estimates include all racial groups.  
<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Due to significant BRFSS methodology changes that took place in 2011, the estimates within these tables should not be compared to Michigan BRFSS estimates from prior to 2011.



**Table 14: Any Alcohol Consumption<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	55.6	(54.1-57.2)	57.9	(56.2-59.6)	47.9	(43.0-52.8)	38.8	(32.5-45.5)	60.7	(52.0-68.8)
<b>Age</b>										
18 - 49	60.2	(57.7-62.6)	62.4	(59.6-65.2)	56.7	(49.4-63.7)	44.4	(36.2-53.1)	61.0	(50.7-70.4)
50 +	51.4	(49.5-53.4)	54.1	(52.0-56.1)	38.0	(31.7-44.6)	29.0	(20.0-39.9)	61.0	(44.9-75.0)
<b>Gender</b>										
Male	58.5	(56.2-60.8)	61.9	(59.4-64.4)	50.1	(42.4-57.8)	36.7	(28.8-45.4)	54.6	(41.7-66.9)
Female	52.9	(50.8-55.0)	54.0	(51.8-56.3)	46.0	(39.9-52.1)	41.6	(31.8-52.0)	65.0	(53.5-75.0)
<b>Education</b>										
High school graduate or less	44.7	(41.8-47.5)	45.2	(42.0-48.5)	41.3	(33.8-49.2)	29.6	(19.2-42.8)	56.6	(43.8-68.6)
Some college or more	62.6	(60.9-64.3)	65.4	(63.6-67.2)	53.9	(47.8-59.8)	42.4	(35.0-50.2)	64.7	(53.2-74.8)
<b>Household Income</b>										
< \$35,000	44.5	(41.4-47.6)	45.9	(42.3-49.6)	42.2	(35.3-49.4)	33.1	(23.3-44.6)	49.4	(35.8-63.1)
\$35,000 +	65.6	(63.7-67.5)	66.8	(64.7-68.7)	58.8	(51.0-66.1)	49.0	(39.4-58.6)	77.3	(66.7-85.3)

<sup>a</sup> The proportion who reported consuming at least one drink of any alcoholic beverage in the previous month.  
<sup>b</sup> Michigan estimates include all racial groups.

Due to significant BRFSS methodology changes that took place in 2011, the estimates within these tables should not be compared to Michigan BRFSS estimates from prior to 2011.

**Table 15: Heavy Drinking<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	6.8	(6.0-7.6)	7.4	(6.5-8.3)	4.7	(3.0-7.2)	3.2	(1.9-5.4)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	7.5	(6.3-8.9)	8.4	(6.9-10.1)	5.5	(3.1-9.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	6.1	(5.3-7.1)	6.6	(5.6-7.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	7.6	(6.4-8.9)	8.2	(6.9-9.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	6.0	(5.1-7.0)	6.5	(5.5-7.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	7.0	(5.7-8.6)	7.4	(5.9-9.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	6.6	(5.8-7.5)	7.3	(6.4-8.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	5.8	(4.5-7.4)	6.4	(4.8-8.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	8.2	(7.1-9.4)	8.8	(7.6-10.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported consuming on average more than two alcoholic beverages per day for men or more than one per day for women in the previous month.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 16: Binge Drinking<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	17.4	(16.3-18.7)	18.0	(16.7-19.4)	14.6	(11.3-18.7)	12.0	(8.5-16.7)	25.4	(17.6-35.2)
<b>Age</b>										
18 - 49	23.7	(21.7-25.9)	25.3	(23.0-27.8)	19.0	(14.2-25.1)	13.6	(9.3-19.6)	27.5	(18.1-39.4)
50 +	11.6	(10.3-13.0)	11.8	(10.5-13.3)	9.5	(5.6-15.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	22.3	(20.4-24.2)	23.8	(21.7-26.1)	16.4	(11.1-23.6)	13.6	(8.9-20.2)	23.9	(14.8-36.2)
Female	13.2	(11.7-14.8)	12.5	(11.0-14.2)	13.1	(9.4-18.0)	-- <sup>c</sup>	-- <sup>c</sup>	26.4	(15.6-41.1)
<b>Education</b>										
High school graduate or less	17.0	(14.9-19.3)	17.2	(14.8-19.8)	14.4	(9.4-21.3)	15.0	(8.3-25.6)	22.3	(13.5-34.4)
Some college or more	18.0	(16.6-19.5)	18.5	(17.1-20.1)	15.0	(10.9-20.3)	10.9	(7.0-16.4)	28.5	(16.8-44.2)
<b>Household Income</b>										
< \$35,000	15.4	(13.2-18.0)	15.4	(12.8-18.4)	16.0	(11.1-22.4)	12.9	(7.0-22.6)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	20.8	(19.1-22.5)	21.1	(19.4-22.9)	13.9	(9.1-20.5)	13.9	(8.8-21.4)	37.6	(24.5-52.8)

<sup>a</sup> The proportion who reported consuming five or more drinks per occasion at least once in the previous month.  
<sup>b</sup> Michigan estimates include all racial groups.  
<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Due to significant BRFSS methodology changes that took place in 2011, the estimates within these tables should not be compared to Michigan BRFSS estimates from prior to 2011.

**Table 17: Drove Motor Vehicle After Drinking<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	2.2	(1.7-2.8)	1.7	(1.3-2.3)	--b	--b	--b	--b	--b	--b
<b>Age</b>										
18 - 49	2.6	(1.8-3.7)	1.8	(1.1-2.9)	--b	--b	--b	--b	--b	--b
50 +	1.8	(1.2-2.6)	1.7	(1.1-2.4)	--b	--b	--b	--b	--b	--b
<b>Gender</b>										
Male	2.9	(2.1-4.0)	2.5	(1.8-3.6)	--b	--b	--b	--b	--b	--b
Female	1.5	(0.9-2.3)	--b	--b	--b	--b	--b	--b	--b	--b
<b>Education</b>										
High school graduate or less	2.7	(1.7-4.5)	--b	--b	--b	--b	--b	--b	--b	--b
Some college or more	2.0	(1.5-2.7)	1.8	(1.3-2.5)	--b	--b	--b	--b	--b	--b
<b>Household Income</b>										
< \$35,000	2.5	(1.4-4.5)	--b	--b	--b	--b	--b	--b	--b	--b
\$35,000 +	2.2	(1.6-3.0)	2.1	(1.5-3.0)	--b	--b	--b	--b	--b	--b

<sup>a</sup> Among adults who reporting drinking any alcohol within the past 30 days, the proportion who reported driving after having too much to drink at least once in the previous month.

<sup>b</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 18: Seatbelt Use<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	89.6	(88.5-90.6)	90.0	(88.7-91.1)	88.5	(85.0-91.3)	90.2	(84.8-93.8)	88.0	(80.8-92.7)
<b>Age</b>										
18 - 49	86.1	(84.1-87.9)	86.0	(83.7-88.1)	85.3	(79.3-89.8)	88.3	(81.6-92.8)	85.7	(76.7-91.6)
50 +	93.4	(92.3-94.3)	93.5	(92.4-94.4)	92.2	(88.4-94.9)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	86.9	(85.0-88.5)	86.6	(84.6-88.5)	86.6	(80.4-91.0)	89.0	(81.3-93.8)	-- <sup>c</sup>	-- <sup>c</sup>
Female	92.4	(91.1-93.6)	93.2	(91.7-94.4)	90.2	(86.0-93.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	86.8	(84.5-88.8)	87.1	(84.4-89.4)	85.8	(79.3-90.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	91.6	(90.4-92.6)	91.7	(90.5-92.8)	90.9	(87.0-93.7)	91.3	(85.0-95.1)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	88.0	(85.6-90.0)	88.3	(85.3-90.7)	88.0	(82.6-91.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	90.6	(89.3-91.8)	90.5	(89.0-91.8)	90.3	(84.4-94.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported always using a seatbelt when driving or riding in a car.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 19: No Routine Checkup in Past Year<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	23.4	(22.1-24.8)	23.4	(22.0-24.9)	18.0	(14.5-22.1)	31.4	(25.4-38.1)	27.2	(20.3-35.3)
<b>Age</b>										
18 - 49	31.3	(29.1-33.6)	33.0	(30.4-35.7)	20.9	(15.7-27.3)	34.4	(26.8-42.8)	30.5	(22.2-40.2)
50 +	15.4	(14.1-16.9)	15.2	(13.8-16.7)	15.1	(10.8-20.5)	23.5	(14.9-35.0)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	28.0	(25.9-30.2)	26.8	(24.5-29.2)	27.2	(20.9-34.5)	38.4	(29.9-47.7)	37.1	(25.7-50.2)
Female	18.9	(17.3-20.6)	20.2	(18.4-22.2)	10.1	(7.1-14.3)	22.0	(15.2-30.8)	20.4	(13.0-30.6)
<b>Education</b>										
High school graduate or less	24.1	(21.7-26.7)	25.2	(22.4-28.2)	16.6	(11.3-23.6)	23.5	(14.3-36.0)	31.5	(21.0-44.3)
Some college or more	22.8	(21.3-24.3)	22.3	(20.7-24.0)	19.1	(14.7-24.3)	34.4	(27.3-42.3)	22.8	(15.1-32.9)
<b>Household Income</b>										
< \$35,000	23.2	(20.5-26.0)	24.2	(21.1-27.7)	17.6	(12.6-24.0)	30.1	(19.7-43.0)	27.4	(16.8-41.4)
\$35,000 +	23.1	(21.5-24.8)	22.8	(21.0-24.7)	18.4	(13.0-25.3)	32.7	(24.5-42.2)	26.4	(17.2-38.3)

<sup>a</sup> The proportion who reported that they did not have a routine checkup in the past year.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 20: Mammography in the Past Two Years among Women Aged 40 and Older<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	72.7	(70.5-74.7)	72.1	(69.8-74.3)	76.1	(68.3-82.5)	66.7	(52.1-78.7)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	68.4	(64.3-72.3)	67.8	(63.4-72.0)	69.2	(55.3-80.4)	48.2	(23.8-73.4)	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	75.4	(73.0-77.7)	74.6	(72.0-77.1)	80.9	(71.8-87.5)	78.7	(67.1-87.0)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	67.1	(62.5-71.4)	67.7	(62.8-72.3)	66.8	(54.5-77.1)	48.4	(27.4-70.1)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	76.6	(73.9-79.2)	75.1	(72.1-77.9)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> Among women aged 40 years and older, the proportion who reported having a mammogram within the past two years.  
<sup>b</sup> Michigan estimates include all racial groups.  
<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 21: Appropriate Timed Pap Test among Women Aged 18 and Older<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	68.7	(66.4-70.9)	67.6	(65.0-70.1)	75.4	(68.7-81.0)	62.4	(50.2-73.1)	73.3	(61.0-82.9)
<b>Age</b>										
18 - 49	73.2	(69.8-76.3)	73.4	(69.5-77.0)	78.2	(69.3-85.1)	59.3	(44.3-72.7)	73.9	(60.1-84.2)
50 +	62.5	(59.3-65.6)	60.9	(57.5-64.2)	70.2	(59.4-79.2)	66.8	(49.1-80.8)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	65.4	(60.8-69.7)	63.6	(58.3-68.6)	75.2	(64.0-83.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	70.7	(68.1-73.2)	69.7	(66.8-72.5)	76.0	(67.6-82.7)	70.7	(59.0-80.1)	72.8	(56.6-84.6)
<b>Household Income</b>										
< \$35,000	63.2	(58.5-67.6)	61.6	(55.8-67.1)	69.2	(59.5-77.4)	48.3	(30.5-66.5)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	73.5	(70.6-76.3)	72.0	(68.7-75.0)	84.9	(74.0-91.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> Among women aged 18 years and older, the proportion who reported having a Pat test within the previous three years.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.



**Table 22: Prostate-Specific Antigen (PSA) Test Within the Past Year among Men Aged 50 and Older<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	32.9	(30.2-35.7)	34.4	(31.5-37.4)	28.0	(19.6-38.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	23.5	(19.3-28.4)	25.1	(20.3-30.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	38.3	(35.0-41.8)	39.6	(36.1-43.1)	37.7	(25.7-51.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	24.5	(19.5-30.4)	25.7	(20.1-32.3)	25.3	(13.7-42.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	35.4	(32.0-38.9)	36.2	(32.7-39.8)	28.2	(16.3-44.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> Among men aged 50 years and older, the proportion who reported having a PSA test within the past year.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 23: Appropriate Colorectal Cancer Screening among Adults Aged 50 and Older<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	75.6	(73.9-77.3)	76.0	(73.2-78.5)	73.7	(64.1-81.4)	66.6	(48.7-80.7)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	75.2	(72.6-77.7)	76.0	(73.2-78.5)	73.7	(64.1-81.4)	66.6	(48.7-80.7)	-- <sup>c</sup>	-- <sup>c</sup>
Female	76.2	(73.8-78.4)	76.7	(74.1-79.0)	75.9	(66.6-83.2)	63.8	(47.3-77.6)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	73.4	(70.1-76.6)	75.3	(71.8-78.5)	62.4	(50.7-72.8)	64.7	(43.1-81.6)	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	77.3	(75.3-79.1)	77.0	(75.0-79.0)	84.1	(77.8-88.8)	66.1	(51.8-77.9)	63.9	(41.2-81.8)
<b>Household Income</b>										
< \$35,000	70.4	(66.4-74.1)	71.6	(67.1-75.6)	66.1	(55.4-75.4)	77.8	(62.4-88.1)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	77.7	(75.6-79.7)	77.4	(75.1-79.5)	85.3	(76.5-91.2)	60.2	(42.9-75.3)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> Among those aged 50 years and older, the proportion who had either a fecal occult blood test within the past year, a sigmoidoscopy within the past five years, or a colonoscopy within the past ten years.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 24: No Dental Visit in the Past Year<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	30.8	(29.3-32.2)	28.7	(27.2-30.3)	37.2	(32.6-42.0)	35.8	(29.5-42.6)	39.1	(30.6-48.3)
<b>Age</b>										
18 - 49	33.2	(30.9-35.5)	31.5	(28.9-34.1)	37.4	(30.8-44.4)	36.3	(28.4-44.9)	38.7	(28.9-49.6)
50 +	28.2	(26.4-30.0)	26.2	(24.4-28.1)	37.0	(30.8-43.7)	35.1	(25.1-46.6)	41.2	(26.1-58.2)
<b>Gender</b>										
Male	32.9	(30.7-35.1)	31.0	(28.7-33.4)	39.9	(32.8-47.6)	40.2	(31.5-49.6)	36.2	(25.0-49.1)
Female	28.6	(26.7-30.6)	26.5	(24.5-28.6)	34.7	(29.1-40.9)	30.1	(21.8-40.0)	41.0	(29.4-53.6)
<b>Education</b>										
High school graduate or less	39.8	(37.1-42.6)	39.2	(36.1-42.3)	42.5	(35.1-50.4)	40.9	(28.4-54.6)	40.4	(28.9-53.1)
Some college or more	24.8	(23.2-26.5)	22.4	(20.8-24.0)	32.2	(26.9-38.1)	33.8	(26.7-41.8)	37.7	(25.8-51.4)
<b>Household Income</b>										
< \$35,000	43.5	(40.4-46.6)	44.5	(40.9-48.2)	43.6	(36.7-50.8)	43.1	(31.1-56.0)	33.0	(21.3-47.2)
\$35,000 +	23.1	(21.4-24.9)	21.5	(19.8-23.3)	25.6	(19.5-32.9)	31.6	(23.2-41.4)	38.6	(26.0-52.9)

<sup>a</sup> The proportion who reported that they had not visited a dentist or dental clinic for any reason in the previous year.

<sup>b</sup> Michigan estimates include all racial groups.

**Table 25: Had Flu Vaccine in Past Year among Adults 65 Years and Older<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	71.7	(69.4-73.9)	72.9	(70.4-75.2)	65.2	(55.6-73.6)	69.9	(51.7-83.5)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	71.9	(68.2-75.4)	73.1	(69.3-76.6)	59.8	(43.0-74.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	72.4	(69.4-75.3)	72.7	(69.4-75.7)	69.0	(58.3-78.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	70.1	(65.7-74.2)	71.3	(66.7-75.6)	56.1	(40.5-70.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	73.6	(71.0-76.1)	73.9	(71.1-76.5)	71.9	(60.6-81.1)	65.3	(49.0-78.7)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	68.5	(63.9-72.6)	66.8	(61.8-71.5)	73.0	(60.0-83.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	75.8	(72.6-78.7)	76.6	(73.3-79.5)	63.2	(47.2-76.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion of those aged 65 years and older who reported that they had a flu vaccine, either by an injection in the arm or sprayed in the nose during the past 12 months.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 26: Ever Had Pneumonia Vaccine among Adults 65 Years and Older<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	71.8	(69.4-74.2)	74.0	(71.5-76.4)	56.4	(46.6-65.7)	62.8	(41.1-80.4)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	66.7	(62.7-70.5)	69.1	(65.0-72.9)	46.6	(31.0-62.9)	55.2	(30.0-77.9)	-- <sup>c</sup>	-- <sup>c</sup>
Female	76.2	(73.1-79.1)	78.0	(74.8-80.9)	63.6	(51.7-74.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	70.1	(65.5-74.4)	73.1	(68.3-77.4)	47.5	(32.5-63.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	73.3	(70.5-75.9)	74.6	(71.8-77.2)	63.2	(50.9-74.0)	65.5	(48.8-79.2)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	71.2	(66.7-75.3)	73.0	(68.3-77.3)	60.6	(46.8-73.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	74.5	(71.1-77.6)	75.3	(71.8-78.4)	62.7	(46.4-76.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion of those aged 65 years and older who reported that they ever had a pneumococcal vaccine.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 27: Ever Had an HIV Test among Adults 18 – 64 Years of Age<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	43.2	(41.3-45.1)	38.2	(36.1-40.4)	65.9	(60.3-71.2)	39.3	(32.3-46.8)	51.9	(42.5-61.2)
<b>Age</b>										
18 - 49	45.5	(43.0-48.0)	41.1	(38.3-43.9)	68.1	(61.0-74.4)	39.9	(31.6-48.8)	52.3	(41.7-62.7)
50 - 64	38.6	(35.7-41.6)	33.7	(30.6-36.9)	62.0	(52.4-70.7)	40.1	(27.4-54.4)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	38.0	(35.4-40.8)	33.8	(30.9-36.9)	63.0	(54.0-71.1)	35.9	(27.2-45.7)	36.8	(25.0-50.5)
Female	48.0	(45.3-50.7)	42.8	(39.9-45.8)	68.7	(61.5-75.0)	44.1	(33.0-55.8)	62.8	(50.2-73.8)
<b>Education</b>										
High school graduate or less	44.6	(41.1-48.2)	37.4	(33.4-41.6)	65.9	(56.7-74.1)	51.8	(35.3-67.9)	58.0	(44.3-70.5)
Some college or more	41.8	(39.7-44.1)	38.7	(36.4-41.1)	65.5	(58.4-71.8)	35.0	(27.7-43.1)	46.0	(32.9-59.7)
<b>Household Income</b>										
< \$35,000	52.3	(48.3-56.3)	44.6	(39.8-49.6)	67.3	(58.9-74.7)	52.4	(38.4-66.0)	55.0	(40.2-69.0)
\$35,000 +	40.4	(38.1-42.8)	37.5	(35.0-40.0)	67.8	(58.9-75.6)	36.7	(27.7-46.7)	48.3	(34.3-62.7)

<sup>a</sup> The proportion of those aged 18-64 years who reported that they ever had been tested for HIV, apart from tests that were part of a blood donation.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 28: Lifetime Asthma Prevalence - Adult<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	15.4	(14.4-16.6)	15.0	(13.8-16.3)	17.3	(14.0-21.1)	17.4	(12.8-23.1)	14.3	(9.5-20.8)
<b>Age</b>										
18 - 49	16.9	(15.1-18.8)	17.4	(15.3-19.8)	16.0	(11.8-21.4)	15.5	(10.4-22.4)	14.6	(9.2-22.4)
50 +	13.9	(12.6-15.2)	12.8	(11.6-14.1)	18.6	(13.9-24.3)	24.0	(15.0-36.3)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	11.7	(10.3-13.4)	11.4	(9.8-13.3)	12.7	(8.7-18.4)	13.5	(8.6-20.5)	-- <sup>c</sup>	-- <sup>c</sup>
Female	18.8	(17.2-20.5)	18.3	(16.6-20.1)	21.2	(16.6-26.6)	22.5	(14.8-32.6)	16.1	(9.6-25.7)
<b>Education</b>										
High school graduate or less	17.0	(14.9-19.3)	15.0	(12.7-17.5)	22.4	(16.8-29.0)	30.1	(18.6-44.7)	18.3	(10.6-29.7)
Some college or more	14.4	(13.2-15.6)	15.0	(13.7-16.4)	13.0	(9.6-17.3)	12.2	(8.5-17.3)	10.1	(5.8-17.0)
<b>Household Income</b>										
< \$35,000	18.5	(16.2-21.0)	16.7	(14.2-19.6)	18.9	(14.1-24.8)	34.3	(23.1-47.6)	20.1	(11.0-33.8)
\$35,000 +	13.2	(11.9-14.5)	13.4	(12.0-14.9)	13.7	(9.2-19.9)	11.0	(6.9-17.0)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported that they were told by a doctor, nurse, or other health care professional that they had asthma.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 29: Current Asthma Prevalence - Adult<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	11.0	(10.1-12.0)	10.5	(9.5-11.6)	13.5	(10.7-17.0)	13.2	(9.2-18.5)	10.5	(6.5-16.4)
<b>Age</b>										
18 - 49	11.3	(9.8-13.0)	11.2	(9.5-13.3)	12.0	(8.5-16.8)	11.3	(7.0-17.8)	10.1	(5.8-17.1)
50 +	10.8	(9.7-12.1)	9.9	(8.8-11.0)	15.0	(10.8-20.4)	19.3	(11.3-31.0)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	7.3	(6.1-8.8)	6.9	(5.6-8.5)	10.0	(6.5-15.2)	8.4	(4.9-14.0)	-- <sup>c</sup>	-- <sup>c</sup>
Female	14.5	(13.1-16.0)	13.9	(12.5-15.6)	16.5	(12.5-21.5)	19.4	(12.2-29.5)	13.7	(7.8-22.8)
<b>Education</b>										
High school graduate or less	12.6	(10.7-14.7)	11.0	(9.0-13.3)	16.9	(12.1-23.1)	23.4	(13.2-37.9)	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	10.1	(9.1-11.2)	10.3	(9.2-11.4)	10.7	(7.7-14.6)	9.1	(6.0-13.4)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	14.0	(11.9-16.2)	12.8	(10.6-15.4)	14.7	(10.5-20.2)	24.0	(14.4-37.1)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	9.3	(8.3-10.5)	9.3	(8.2-10.6)	10.4	(6.6-16.0)	8.7	(5.2-14.3)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported that they still had asthma.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.



**Table 30: Ever Told Chronic Obstructive Pulmonary Disease (COPD)<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	8.3	(7.6-9.2)	8.9	(8.0-9.8)	6.7	(4.6-9.7)	7.0	(4.2-11.4)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	4.0	(3.1-5.1)	3.9	(2.9-5.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	13.0	(11.7-14.4)	13.3	(12.0-14.8)	12.0	(7.9-17.6)	13.0	(7.5-21.6)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	7.3	(6.2-8.5)	7.7	(6.5-9.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	9.5	(8.4-10.8)	10.0	(8.8-11.4)	7.4	(4.6-11.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	13.2	(11.5-15.1)	13.6	(11.7-15.7)	10.4	(6.5-16.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	5.4	(4.7-6.2)	6.1	(5.3-7.0)	3.5	(2.0-6.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	15.0	(12.9-17.3)	17.5	(15.0-20.3)	9.1	(5.6-14.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	4.8	(4.0-5.6)	5.2	(4.4-6.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported ever being told by a doctor that they had chronic obstructive pulmonary disease (COPD), emphysema or chronic bronchitis.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 31: Ever Told Arthritis<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	30.1	(28.8-31.4)	31.8	(30.3-33.3)	28.1	(24.2-32.4)	23.2	(18.0-29.3)	18.3	(12.8-25.6)
<b>Age</b>										
18 - 49	11.6	(10.2-13.2)	11.7	(10.1-13.5)	11.5	(8.0-16.2)	11.5	(7.0-18.3)	11.1	(6.3-18.8)
50 +	49.2	(47.3-51.1)	49.6	(47.6-51.6)	46.3	(40.0-52.8)	52.1	(40.7-63.3)	45.5	(30.3-61.6)
<b>Gender</b>										
Male	24.4	(22.6-26.2)	25.8	(23.9-27.9)	22.0	(16.7-28.3)	16.6	(11.2-23.9)	17.0	(9.7-28.3)
Female	35.6	(33.7-37.5)	37.4	(35.3-39.5)	33.5	(28.1-39.3)	31.7	(22.9-42.1)	19.2	(11.9-29.5)
<b>Education</b>										
High school graduate or less	34.5	(32.1-37.1)	36.9	(34.1-39.8)	26.6	(20.8-33.5)	35.9	(24.0-49.9)	24.9	(15.9-36.8)
Some college or more	27.4	(26.0-28.9)	28.7	(27.1-30.3)	29.4	(24.4-35.0)	18.1	(13.1-24.5)	11.6	(6.4-20.1)
<b>Household Income</b>										
< \$35,000	38.9	(36.0-41.8)	44.2	(40.7-47.8)	29.5	(23.8-35.9)	40.1	(28.5-52.9)	20.6	(11.9-33.4)
\$35,000 +	24.4	(22.8-26.0)	25.7	(24.0-27.4)	23.7	(17.9-30.6)	14.3	(9.0-21.9)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who had ever being told by a health care professional that they had some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 32: Ever Told Heart Attack<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	5.1	(4.5-5.8)	5.3	(4.6-6.0)	4.7	(2.9-7.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	1.2	(0.8-1.8)	1.3	(0.8-2.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	8.9	(7.8-10.1)	8.9	(7.8-10.1)	8.9	(5.4-14.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	6.6	(5.6-7.7)	7.4	(6.3-8.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	3.6	(2.9-4.4)	3.3	(2.6-4.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	7.0	(5.8-8.4)	7.0	(5.7-8.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	3.8	(3.2-4.4)	4.2	(3.6-5.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	6.9	(5.5-8.5)	8.3	(6.7-10.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	3.9	(3.2-4.7)	4.1	(3.4-5.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who had ever been told by a doctor that they had a heart attack or myocardial infarction.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 33: Ever Told Angina or Coronary Heart Disease<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	5.0	(4.4-5.7)	5.5	(4.8-6.3)	3.0	(1.8-4.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	1.2	(0.7-1.9)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	9.0	(7.9-10.2)	9.4	(8.3-10.8)	5.9	(3.6-9.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	6.2	(5.3-7.2)	7.0	(5.9-8.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	4.0	(3.2-4.9)	4.1	(3.3-5.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	5.7	(4.6-7.1)	6.2	(4.9-7.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	4.6	(4.0-5.3)	5.1	(4.4-5.9)	3.7	(2.1-6.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	6.6	(5.4-8.1)	7.8	(6.3-9.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	4.0	(3.4-4.7)	4.2	(3.5-5.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who had ever been told by a doctor that they had angina or coronary heart disease.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 34: Ever Told Stroke<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	3.5	(3.0-4.0)	3.6	(3.1-4.2)	3.7	(2.5-5.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	1.3	(0.8-1.9)	1.0	(0.6-1.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	5.7	(4.9-6.6)	5.7	(4.9-6.7)	6.1	(3.9-9.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	3.0	(2.4-3.7)	3.4	(2.7-4.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	4.0	(3.3-4.8)	3.8	(3.0-4.7)	5.2	(3.3-8.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	4.5	(3.6-5.6)	4.3	(3.4-5.5)	5.0	(2.9-8.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	2.8	(2.3-3.4)	3.1	(2.5-3.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	5.7	(4.6-7.2)	6.6	(5.2-8.4)	3.8	(2.1-6.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	2.2	(1.8-2.8)	2.3	(1.8-3.0)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who had ever been told by a doctor that they had a stroke.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 35: Ever Told Any Cardiovascular Disease<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	10.0	(9.1-10.9)	10.5	(9.5-11.5)	9.4	(7.0-12.6)	6.4	(3.8-10.6)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	2.7	(2.0-3.6)	2.7	(2.0-3.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	17.3	(15.8-18.8)	17.3	(15.8-19.0)	17.3	(12.6-23.1)	15.7	(9.4-24.9)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	11.4	(10.2-12.8)	12.8	(11.3-14.4)	8.1	(5.1-12.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	8.5	(7.4-9.8)	8.3	(7.1-9.6)	10.6	(7.2-15.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	12.9	(11.3-14.8)	13.3	(11.4-15.5)	11.7	(7.6-17.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	8.0	(7.2-8.9)	8.7	(7.8-9.7)	7.5	(5.0-11.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	13.3	(11.5-15.3)	15.7	(13.5-18.2)	8.5	(5.3-13.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	7.5	(6.6-8.5)	7.9	(6.9-9.0)	7.8	(4.6-12.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who had ever been told by a doctor that they had a heart attack, coronary heart disease or a stroke.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 36: Ever Told Skin Cancer<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	6.2	(5.6-6.8)	7.8	(7.1-8.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	1.0	(0.7-1.5)	1.2	(0.8-1.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	11.4	(10.4-12.5)	13.6	(12.3-14.9)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	5.8	(5.0-6.7)	7.3	(6.3-8.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	6.5	(5.8-7.4)	8.3	(7.3-9.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	5.0	(4.1-6.0)	6.4	(5.3-7.8)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	7.0	(6.3-7.7)	8.6	(7.7-9.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	5.2	(4.2-6.5)	7.4	(6.0-9.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	6.1	(5.4-6.9)	7.2	(6.4-8.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported ever being told by a doctor that they had skin cancer.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 37: Ever Told Any Other Type of Cancer<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	7.6	(6.9-8.3)	8.2	(7.4-9.0)	6.4	(4.5-9.0)	3.1	(1.8-5.1)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	2.3	(1.7-3.3)	2.2	(1.5-3.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	12.8	(11.6-14.1)	13.4	(12.1-14.8)	10.8	(7.3-15.6)	9.3	(5.3-15.8)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	6.2	(5.3-7.2)	6.6	(5.7-7.8)	6.2	(3.6-10.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	8.8	(7.7-10.0)	9.6	(8.4-11.0)	6.5	(4.1-10.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	8.4	(7.0-9.9)	9.3	(7.7-11.2)	5.6	(3.1-9.9)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	7.0	(6.2-7.8)	7.5	(6.6-8.4)	7.2	(4.6-10.9)	3.3	(1.8-6.0)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	7.9	(6.6-9.5)	9.6	(7.9-11.8)	4.9	(2.8-8.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	6.7	(5.8-7.6)	6.9	(6.0-7.9)	8.4	(4.8-14.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported ever being told by a doctor that they had a form of cancer other than skin cancer.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.



**Table 38: Ever Told Cancer<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	12.4	(11.6-13.3)	14.3	(13.3-15.4)	7.0	(5.0-9.7)	4.9	(3.0-7.8)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	3.3	(2.5-4.3)	3.4	(2.5-4.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	21.7	(20.2-23.2)	23.9	(22.3-25.6)	11.6	(8.1-16.4)	12.7	(8.0-19.7)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	10.6	(9.5-11.9)	12.2	(10.9-13.7)	6.2	(3.6-10.6)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	14.1	(12.8-15.4)	16.2	(14.7-17.8)	7.7	(5.1-11.6)	6.1	(3.4-10.8)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	12.2	(10.6-13.9)	14.2	(12.4-16.3)	5.9	(3.3-10.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	12.6	(11.6-13.6)	14.3	(13.2-15.6)	8.1	(5.4-12.0)	4.5	(2.7-7.2)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	12.4	(10.7-14.2)	15.8	(13.6-18.3)	5.9	(3.6-9.6)	4.5	(2.8-7.1)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	11.6	(10.5-12.7)	12.8	(11.6-14.0)	8.7	(5.1-14.5)	10.0	(5.8-16.7)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported ever being told by a doctor that they had skin cancer or any other type of cancer.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 39: Ever Told Diabetes<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	12.3	(11.4-13.3)	11.8	(10.8-12.8)	16.7	(13.5-20.5)	11.9	(8.2-17.0)	8.2	(4.9-13.6)
<b>Age</b>										
18 - 49	20.0	(18.5-21.6)	3.9	(3.0-5.1)	8.0	(4.7-13.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	12.3	(11.3-13.2)	18.7	(17.2-20.4)	25.9	(20.7-31.9)	25.7	(16.9-37.0)	24.1	(13.4-39.4)
<b>Gender</b>										
Male	11.4	(10.2-12.8)	12.4	(11.0-14.1)	19.8	(14.4-26.5)	11.5	(7.0-18.2)	-- <sup>c</sup>	-- <sup>c</sup>
Female	12.2	(11.3-13.2)	11.1	(9.8-12.6)	14.1	(10.7-18.5)	12.4	(6.9-21.3)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	10.6	(9.6-11.6)	14.2	(12.2-16.4)	18.5	(13.5-24.7)	19.0	(10.7-31.6)	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	11.8	(10.8-12.8)	10.3	(9.3-11.5)	14.9	(11.1-19.9)	9.0	(5.5-14.5)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	9.6	(8.5-10.7)	15.5	(13.3-18.0)	18.6	(13.7-24.6)	23.9	(14.5-36.8)	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	20.0	(18.5-21.6)	9.6	(8.5-10.8)	12.9	(8.7-18.9)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported that they were ever told by a doctor that they had diabetes. Adults who had been told they have prediabetes and women who had diabetes only during pregnancy were classified as not having been diagnosed.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 40: Ever Told Kidney Disease<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	3.2	(2.7-3.7)	3.4	(2.9-4.0)	3.4	(2.1-5.4)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Age</b>										
18 - 49	1.3	(0.8-2.1)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
50 +	5.2	(4.4-6.1)	5.4	(4.5-6.3)	4.9	(2.8-8.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	2.6	(2.0-3.3)	2.7	(2.1-3.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Female	3.8	(3.1-4.7)	4.0	(3.2-5.0)	4.2	(2.4-7.5)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Education</b>										
High school graduate or less	3.9	(3.0-5.0)	4.0	(3.0-5.3)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
Some college or more	2.8	(2.3-3.4)	3.0	(2.5-3.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
<b>Household Income</b>										
< \$35,000	4.2	(3.2-5.4)	5.1	(3.8-6.7)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>
\$35,000 +	2.5	(2.0-3.2)	2.5	(2.0-3.2)	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported ever being told by a doctor that they had kidney disease.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

**Table 41: Ever Told Depression<sup>a</sup> by Race-Ethnicity  
2020 Michigan BRFSS**

Demographic Characteristics	Michigan <sup>b</sup>		White, Non-Hispanic		Black, Non-Hispanic		Other/Multiracial, Non-Hispanic		Hispanic	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Total</b>	19.5	(18.3-20.8)	20.2	(18.8-21.6)	15.6	(12.6-19.3)	16.9	(12.2-23.0)	25.3	(17.7-34.8)
<b>Age</b>										
18 - 49	22.8	(20.9-24.9)	23.9	(21.7-26.3)	18.3	(13.7-24.0)	18.9	(12.8-27.1)	25.3	(16.6-36.5)
50 +	16.6	(15.2-18.2)	17.1	(15.5-18.8)	12.9	(9.2-17.7)	13.5	(8.2-21.2)	-- <sup>c</sup>	-- <sup>c</sup>
<b>Gender</b>										
Male	13.9	(12.4-15.6)	14.3	(12.6-16.3)	13.1	(9.1-18.5)	10.6	(6.3-17.3)	-- <sup>c</sup>	-- <sup>c</sup>
Female	25.0	(23.2-26.9)	25.7	(23.8-27.8)	17.8	(13.6-23.0)	25.2	(16.8-36.1)	32.9	(21.9-46.2)
<b>Education</b>										
High school graduate or less	22.4	(20.1-24.8)	23.1	(20.4-25.9)	17.3	(12.6-23.3)	29.3	(17.4-45.0)	22.1	(13.6-33.9)
Some college or more	17.8	(16.5-19.2)	18.4	(17.0-19.9)	13.6	(10.0-18.3)	12.0	(8.3-16.9)	28.6	(17.1-43.7)
<b>Household Income</b>										
< \$35,000	27.7	(25.0-30.5)	30.5	(27.2-34.0)	20.2	(15.4-26.2)	27.4	(17.1-40.9)	28.4	(17.9-42.0)
\$35,000 +	14.7	(13.3-16.1)	15.2	(13.8-16.7)	7.3	(4.3-12.1)	12.4	(6.9-21.5)	-- <sup>c</sup>	-- <sup>c</sup>

<sup>a</sup> The proportion who reported ever being told by a doctor that they had a depressive disorder including depression, major depression, dysthymia, or minor depression.

<sup>b</sup> Michigan estimates include all racial groups.

<sup>c</sup> Suppressed due to a denominator < 50 and/or a relative standard error > 30%.