

# The 2020 National HIV/AIDS Strategy (NHAS) Indicators Assessment of Michigan's Progress 2010 - 2016



data as of January 1, 2018

The indicators/objectives used to measure the National HIV/AIDS Strategy's (NHAS) goals were updated in 2015. The indicators now measure improvements between baseline, 2010, and 2020. The three main NHAS goals are the same: 1) Reduce new HIV infections, 2) Increase access to care and improve health outcomes for people living with HIV, and 3) Reduce HIV-related health disparities.

The indicators have been updated to reflect current need among persons over 12 years old living with HIV (PLWH).

## Progress Key:

**Met** - 2020 goal is met.

**On Track** - Most recent annual goal was met.

**Needs Improvement** - Most recent annual goal was not met, but numbers are stable or heading in the right direction.

**Worsening** - Most recent annual goal was not met, and numbers are heading in the wrong direction.

**N/A** - Indicator not relevant or no Michigan specific data available at this time.

## NHAS Goal 1 Indicators relevant to Michigan:

- 1) Increase the percentage of PLWH who know their status to 90%
- 2) Reduce the number of new diagnoses by 25%
- 3) Reduce the percentage of young gay and bisexual men who have engaged in HIV-risk behaviors by 10%

## Progress

Unknown  
Needs Improvement

Unknown

## NHAS Goal 2 Indicators relevant to Michigan:

- 4) Increase one month linkage to care rates to 85%
- 5) Increase care retention rates among PLWH to 90%
- 6) Increase the proportion of virally suppressed PLWH to 80%
- 7) Reduce homelessness among PLWH to no more than 5%
- 8) Reduce the death rate among PLWH by 33%

N/A

N/A

On Track

N/A

On Track

## NHAS Goal 3 Indicators relevant to Michigan:

- 9) Reduce new diagnosis rate disparities by 15% among:
  - gay and bisexual men
  - young black gay men
  - black females


Worsening  
Worsening  
Met in 2015

- 10) Increase to 80% the proportion of virally suppressed:
  - youth (13-24 year olds)
  - persons who inject drugs

On Track  
Nearly On Track

The HIV Care Continuum was developed by the CDC to assess gaps in care. It is presented here in order to provide context to many of the NHAS goals. Based on the care continuums and other trends observed, Michigan programs need to focus on linking and retaining PLWH in care. This, in turn will improve the maintenance of low viral load levels, improving the health of PLWH and reducing the risk of HIV transmission.

Michigan HIV Care Continuum, 2016



HIV Infected - Persons aware and unaware of their infection.

Diagnosed - Persons diagnosed with HIV.

In Care - PLWH with at least 1 CD4, viral load, or genotype lab test.

Virally Suppressed - PLWH with less than or equal to 200 copies of HIV virus per milliliter of blood ( $\leq 200$  copies/mL).

Maintained Undetectable - PLWH who maintained viral load levels  $\leq 200$  copies/mL for at least 4-8 months.

HIV and STD Surveillance & Epidemiology Section  
Division of Communicable Disease  
Bureau of Epidemiology and Population Health  
Michigan Department of Health and Human Services

MDHHS - South Oakland Health Center  
27725 Greenfield Rd, Office 57A  
Southfield, MI 48076  
248-424-7910


Lansing - HIV Surveillance Office  
333 S. Grand Ave., 3rd Floor  
Lansing, MI 48913  
517-335-8165

[www.michigan.gov/hivstd](http://www.michigan.gov/hivstd)

## Goal 1: Reducing New HIV Infections

### Indicator 1: By 2020, increase the proportion of PLWH who know their HIV status to 90%

In 2015, 87% of persons living with HIV in Michigan were aware of their HIV status. This figure appears to be improving slowly. Based on calculations provided by the CDC, persons 13-34 are less likely to be diagnosed compared to those over 35, and non-Hispanic white persons are more likely to be diagnosed than all other race/ethnicity groups.




### Indicator 2: By 2020, reduce the annual number of new HIV infections by 25%

To reach this goal, Michigan needs to reduce the annual number of new HIV infections from 744 (in 2010) to 581 by 2020. During 2016, there were 744 new diagnoses. In order to reach the 2020 goal, a reduction of 42 new cases per year is needed.

It is important to note that this objective does not account for HIV prevalence (higher prevalence rates increase the probability of new infections). Prevalence in Michigan has been climbing since the beginning of the epidemic, but the number of new diagnoses has remained stable in recent years<sup>1</sup> resulting in an 22% drop in the transmission rate.

This indicator, however, measures reported case counts, not transmission rate. Unfortunately, Michigan failed to meet the annual goal between 2011 and 2016. The increase of new diagnoses among black gay and bisexual men in their twenties is the primary reason new diagnoses are not decreasing in the state.

The number of new HIV diagnoses in Michigan has remained relatively stable since 2010



### Indicator 3: By 2020, reduce the percentage of young gay and bisexual men who have engaged in HIV-risk behaviors by at least 10%


A sample of Michigan high school students (grades 9 - 12) participate in the Youth Risk Behavioral Surveillance System (YRBSS) survey biannually. Responses from male participants who have had sex with a male or identify as gay or bisexual are included in this indicator calculation. Gender identity data were not collected pre-2013; therefore data are not available for the years 2010 - 2012.

Of the gay and bisexual male participants, those who reported one of the following are categorized as "engaging in HIV-risk behavior":

- 1) Multiple sex partners (3 or more) in the 3 months preceding the interview
- 2) Did not use a condom during last sexual encounter within 3 months
- 3) Ever injected an illegal drug

Large confidence intervals present in 2015 inhibit conclusive assessment of directionality.

23% of young gay/bi 2015 Michigan YRBSS participants engaged in HIV-risk behavior




## Goal 2: Increasing Access to Care and Improving Health Outcomes for People Living with HIV

**Indicator 4: By 2020, increase the percentage of newly diagnosed persons linked to HIV medical care within one month of diagnosis to at least 85%**

CD4, viral load (vl), and genotype lab tests are proxies for clinical care visits. The first of these lab tests collected is used to calculate the time between HIV diagnosis and linkage to care. The earlier persons are linked to care the more likely they are to be in care in the future and the better their prognosis as 85% are virally suppressed. This, in turn, reduces transmission risk to others.


This NHAS indicator measures the proportion of persons newly diagnosed with HIV who received one of these lab tests within one month ( $\leq 1$  mo) of diagnosis. Tests run on the day of diagnoses are counted. The **'NHAS 1 month linkage rate'** is represented by the light orange bars below.

However, persons "linked" within 8 days of diagnosis have future care rates even lower than those linked after 1 month (right). The **'true linkage rate'** is the proportion linked more than ( $>$ ) 8 days and  $\leq 1$  or  $\leq 3$  months. This is represented by the solid, orange lines below. These rates are not expected to reach 85%, however they are important to track.



**Programs should focus on increasing the 'true linkage rates'.**

'True linkage rates' are steadily increasing among persons diagnosed in Michigan



Linked to Care: At least 1 CD4, vl, or genotype test within given time frame

**Indicator 5: By 2020, increase the percentage of persons with diagnosed HIV infection who are retained in HIV medical care to at least 90%**

CD4, viral load (vl), and genotype lab tests are proxies for clinical care visits. Persons who received at least one of these lab tests within one calendar year are **'in care'**. Persons who received at least two of these tests are **'retained in care'**. Receiving HIV care is extremely important in improving individuals' prognoses and, in turn, reducing transmission risk to others by achieving viral suppression and maintaining a low viral load. 'Retention in care' is not necessary to achieve viral suppression. However care retention is essential for monitoring and maintaining a low viral load.


In 2016, the care rate among PLWH increased to 80% and the retention rate increased to 57%.

There are four groups with care rates substantially lower than the state average of 80%:

- 1) Males of Hispanic/Latino descent - 72% in care
- 2) Persons who have never progressed to stage 3 - 74% in care
- 3) Persons who inject drugs - 74% in care
- 4) Persons 25-29 years old - 76% in care

Respectively, these groups account for 4%, 47%, 11% and 9% of all PLWH in Michigan (persons may occupy more than one group). These same groups, as well as black persons, and 20-39 year olds have lower retention rates than the state average.

The rate of Michiganders living with HIV 'in care' and 'retained in care' increased during 2015 and 2016 for the first time since 2010



In Care: At least 1 CD4, vl, or genotype lab test within the given calendar year  
 Retained in Care: At least 2 CD4, vl, or genotype lab tests within the given calendar year

## Goal 2: Increasing Access to Care and Improving Health Outcomes for People Living with HIV (continued)


### Indicator 6: By 2020, increase the percentage of persons with diagnosed HIV infection who are virally suppressed to at least 80%

An individual is considered to be virally suppressed if he/she has less than or equal to 200 copies of HIV virus per milliliter of blood ( $\leq 200$  copies/mL). Consistent suppression of the virus in an individual is an indication of routine access to care and treatment adherence. Those who maintain low viral loads have the best long term prognosis, and transmission of HIV is extremely low among virally suppressed individuals. Those who monitor and maintain an undetectable viral load for over 6 months have effectively no risk of transmitting HIV<sup>2</sup>. The transmission rate among persons diagnosed but not retained in care is much higher: 5.3 transmissions per 100 PLWH not retained in care per year<sup>3</sup>.

The NHAS indicator measures the proportion of PLWH whose last viral load during a given year was suppressed (percent virally suppressed out of all PLWH in Michigan). This is referred to as the **'Virally Suppressed Proportion'**. For context the **'Viral Suppression Rate'** - the percent virally suppressed out of those who received a viral load (aka 'in care') - is also displayed below. The 'Viral Suppression Rate', indicates that once Michiganders with HIV are in care, the likelihood of achieving viral suppression is high; emphasizing the need to improve care rates. Neither of these measure, however, assess the maintenance of viral suppression over time.

Michigan is on track to increase the proportion virally suppressed to 80% by 2020.

'Proportion Virally Suppressed' and 'Viral Suppression Rate' continue to rise




*Virally Suppressed:  $\leq 200$  copies/mL at last lab during given year (of all PLWH)*  
*Viral Suppression Rate:  $\leq 200$  copies/mL at last lab during given year of PLWH who received a viral load*

### Indicator 7: By 2020, reduce the percentage of persons in HIV medical care who are homeless to no more than 5%

At this time, Michigan does not have an estimate of how many PLWH are homeless. Until an estimate can be calculated, the national rate is displayed - In 2014, 9% of PLWH were homeless (91% had stable housing).

Homelessness has increased from 7.7% of PLWH (nationally) in 2010.




### Indicator 8: By 2020, reduce the all-cause death rate among persons with diagnosed HIV infection by at least 33%.

Michigan conducts three death matches each year with the Social Security and National Death Indices, and Michigan Vital Records. Michigan's death data is very complete, however, a lag in death data reporting causes underestimates for recent years. Only years with complete data are displayed below.

The all-cause death rate among Michiganders living with HIV was 22.9 deaths per 1,000 PLWH during 2015. Michigan is nearly on track to reach the 2020 goal provided the death rate is reduced by 1.3 deaths per 1,000 per year.

The all-cause death rate among PLWH in Michigan remained stable 2013 - 2015



### Goal 3: Reducing HIV-Related Health Disparities

**Indicator 9: By 2020, reduce disparities in the rate of new diagnoses by at least 15% among gay/bisexual men, young gay/bisexual black men, and black women.**

It is well known that the gay and black communities shoulder a disproportionate burden of the HIV epidemic. This goal aims to reduce the new diagnosis rate disparity (ratio) observed between these groups and the overall population by at least 15%.

Ratio calculation:

$$\text{New Diagnosis Rate Ratio} = \frac{\text{Rate of new diagnoses among group of interest}}{\text{Rate of new diagnoses among Michiganders}}$$


where the rate of new diagnoses =  $\frac{\text{Number of new diagnoses in given group}}{\text{Total population of given group}}$

The total population of gay and bisexual men was estimated from Lieb *et al*, 2011<sup>3</sup>.




The graph to the right displays the rates of new diagnoses for each group of interest. Black women had a new diagnosis rate of 15.6 per 100,000 black women in Michigan during 2010. That rate was 2 times higher than the overall new diagnosis rate the same year. The disparity ratio (2 to 1) is to be reduced by 15%. The indicator is *not* looking to reduce the rate (15.6 per 100,000) by 15%.

Overall, diagnosis rate disparities among **gay/bisexual men** increased in 2014 and have remained higher than baseline. During 2015, there was a large increase in the diagnosis rate disparity among **young (13-24 years old) gay/bisexual black men** (Note: as testing initiatives did not focus on this group until after 2010, baseline was calculated in 2011). Barring any future increases, Michigan met the 2020 disparity reduction goal for **black females** in 2015.

Young, gay/bisexual, black men have an extremely high rate of new diagnoses per 100,000 persons



Diagnosis rate disparities: Comparing diagnosis rates between the three groups to the overall population (note scale difference among groups).



### Goal 3: Reducing HIV-Related Health Disparities (continued)

**Indicator 10: By 2020, increase the percentage of youth and persons who inject drugs (PWID) with diagnosed HIV infection who are virally suppressed to at least 80%**


See Goal 2, Indicator 6 for an overview of viral suppression.

The percent of virally suppressed PLWH in an HIV positive population is the 'virally suppressed proportion'. The percent of virally suppressed PLWH in an HIV positive population who received a viral load (aka 'in care') is the 'viral suppression rate'.

Youth (persons 13 - 24 years old) have achieved viral suppression significantly less often than their older peers regardless of other demographic factors<sup>4</sup>. They are one of the only groups with low 'viral suppression proportions' (58%) despite a relatively average proportion of the group receiving care (81%). The viral suppression proportion (58%) is the indicator of interest.

Michigan met the annual goal as the proportion of virally suppressed youth steadily improves. Additionally, the 'viral suppression rate' is improving rapidly. Efforts aimed to improve viral suppression among youth need two foci: 1) keep more youth in care - the rapidly improving 'viral suppression rate' indicates more youth in care should lead to more virally suppressed youth, and 2) reducing the viral load among youth already receiving care by ensuring treatment adherence.

The proportion of virally suppressed youth as well as their 'viral suppression rate' continue to increase




*Virally Suppressed: ≤200 copies/mL at last lab during given year (of all PLWH)*  
*Viral Suppression Rate: ≤200 copies/mL at last lab during given year of PLWH who received a viral load*

Persons who inject drugs (PWID) are less likely to be in care and, therefore, less likely to be virally suppressed compared to non-PWIDs.

Michigan was on track to meet the 2020 goal among PWID until 2014. The high viral suppression rate (normal in Michigan) among PWID indicates that efforts meant to increase the proportion of PWID in care should increase the proportion virally suppressed as well.

Improvements in the viral suppression proportion among persons who inject drugs has slowed in recent years, but the viral suppression rate continues to grow



*Virally Suppressed: ≤200 copies/mL at last lab during given year (of all PLWH)*  
*Viral Suppression Rate: ≤200 copies/mL at last lab during given year of PLWH who received a viral load*

#### References

- 1 MDHHS HIV/STD/TB/VH Epidemiology Section (2015). Annual Review of HIV Trends in Michigan (2010-2014). [www.michigan.gov/hivstd](http://www.michigan.gov/hivstd)
- 2 [www.PreventionAccess.org](http://www.PreventionAccess.org)
- 3 Skarbinski J, Rosenberg E, Paz-Bailey G, Hall I, Rose C, Viall A, et al. (2015) Human Immunodeficiency Virus Transmission at Each Step of the Care Continuum in the United States. *JAMA Intern Med*.doi:10.1001/jamainternmed.2014.8180
- 3 Lieb S, Fallon S, Friedman S, Thompson D, Gates G, Liberti T, et al. (2011) Statewide Estimation of Racial/Ethnic Populations of Men Who Have Sex with Men in the U.S. *ASPH Public Health Reports* vol 126 pg 60-
- 4 MDHHS HIV/STD/TB/VH Epidemiology Section (2014). HIV Infected Youth More Likely to be in Care but Less Likely to Achieve Viral Suppression. [www.michigan.gov/hivstd](http://www.michigan.gov/hivstd)