

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	149	87.2	0.0	4.7	0.0	0.0	8.1	0.0
0102	478	85.8	0.0	6.1	0.4	0.0	7.7	0.0
0103	503	85.1	0.0	7.2	0.0	0.0	7.8	0.0
0104	165	92.7	0.0	3.6	0.0	0.0	3.6	0.0
0105	88	*	*	*	*	*	*	*
0250	147	76.2	0.7	5.4	2.7	0.0	15.0	0.0
0251	135	83.7	0.7	2.2	3.7	0.0	9.6	0.0
0252	177	32.2	0.0	2.8	0.0	0.0	63.8	1.1
0253	157	68.8	1.3	3.8	10.2	0.0	14.6	1.3
0401	630	88.7	0.2	4.4	0.0	0.2	6.5	0.0
0402	433	82.4	0.9	3.0	0.2	0.0	13.4	0.0
0403	185	93.0	0.0	1.1	0.5	0.0	5.4	0.0
0404	147	87.8	0.7	5.4	0.0	0.0	6.1	0.0
0501	599	94.0	0.5	2.5	0.3	0.0	2.7	0.0
0503	129	88.4	0.8	2.3	0.8	0.0	7.8	0.0
0504	521	86.2	0.6	3.3	1.7	0.2	8.1	0.0
0505	311	84.6	0.3	6.4	0.3	0.0	8.4	0.0
0506	213	88.3	0.5	3.3	0.5	0.0	7.5	0.0
0507	184	90.8	1.1	1.6	0.0	0.0	6.5	0.0
0508	73	*	*	*	*	*	*	*
0650	1,353	74.5	2.1	8.3	3.0	0.6	11.3	0.2

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis ^(3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0651	831	90.3	0.0	4.7	0.4	0.0	4.7	0.0
0652	426	94.8	0.0	2.6	0.0	0.0	2.6	0.0
0653	660	94.4	0.2	2.6	0.0	0.0	2.7	0.2
0654	479	88.1	0.4	6.1	0.0	0.0	5.0	0.4
0655	411	88.3	0.0	3.4	0.0	0.0	8.0	0.2
0657	108	94.4	0.0	2.8	0.0	0.0	2.8	0.0
0701	605	87.4	2.0	2.5	1.0	0.8	6.3	0.0
0702	96	*	*	*	*	*	*	*
0703	219	87.7	0.5	3.2	0.0	0.0	8.2	0.5
0704	78	*	*	*	*	*	*	*
0705	121	91.7	1.7	1.7	0.8	0.0	4.1	0.0
0706	12	*	*	*	*	*	*	*
0850	995	76.9	3.3	10.7	0.0	0.9	8.2	0.0
0851	888	90.5	0.6	5.7	0.0	0.2	2.8	0.1
0901	2,001	65.6	1.6	17.2	0.0	0.2	15.2	0.1
0902	59	*	*	*	*	*	*	*
1051	1,444	76.7	0.6	11.9	0.8	0.3	9.6	0.1
1101	1,823	32.8	49.4	9.6	0.2	1.1	6.9	0.0
1102	880	72.4	9.4	7.8	0.5	0.5	9.4	0.0
1106	104	84.6	1.9	9.6	0.0	0.0	1.9	1.9
1250	1,205	81.3	0.4	12.9	0.0	0.2	5.2	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1251	877	88.5	0.3	6.3	0.0	0.1	4.6	0.2
1252	802	51.0	0.2	40.8	0.0	0.0	7.6	0.4
1253	1,173	70.5	1.4	12.8	0.2	0.0	15.1	0.0
1350	2,628	49.2	18.0	10.2	0.0	7.3	15.3	0.1
1351	76	*	*	*	*	*	*	*
1352	655	66.0	16.9	8.1	0.0	0.0	9.0	0.0
1401	298	85.6	1.0	5.0	0.0	0.0	8.4	0.0
1402	180	52.8	0.0	31.7	8.3	0.0	7.2	0.0
1403	38	*	*	*	*	*	*	*
1501	1,028	43.0	4.3	44.9	0.2	0.0	7.6	0.0
1503	1,187	51.9	6.7	28.0	0.4	0.1	12.9	0.1
1504	2,431	31.9	4.9	52.7	0.0	1.9	8.3	0.2
1505	522	73.6	3.1	13.2	0.0	0.2	10.0	0.0
1506	793	76.2	1.5	17.2	0.3	0.0	4.9	0.0
1507	751	80.7	0.9	8.3	0.3	0.8	9.1	0.0
1509	511	39.3	1.4	50.3	0.0	0.0	9.0	0.0
1511	874	81.2	1.4	8.6	0.0	0.1	8.7	0.0
1512	470	76.6	1.5	13.4	0.2	0.0	8.3	0.0
1514	466	73.4	3.0	16.5	0.0	0.0	6.9	0.2
1701	758	38.9	0.4	5.7	22.4	0.4	32.2	0.0
1950	730	87.8	0.0	10.0	0.0	0.0	2.2	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1952	174	92.5	0.0	5.2	0.0	0.0	2.3	0.0
1953	413	85.0	0.2	8.0	0.0	0.0	6.8	0.0
1954	576	76.2	0.0	19.1	0.0	0.0	4.7	0.0
1955	166	70.5	0.0	21.1	0.6	0.0	7.8	0.0
1957	590	76.9	0.5	14.9	0.0	0.0	7.6	0.0
1963	102	75.5	0.0	16.7	0.0	0.0	7.8	0.0
2150	777	75.8	0.1	4.4	1.5	0.0	18.1	0.0
2151	200	79.0	0.0	10.5	0.5	0.0	10.0	0.0
2152	37	*	*	*	*	*	*	*
2251	569	95.6	0.0	1.4	0.0	0.0	3.0	0.0
2252	264	95.5	0.0	0.8	0.0	0.0	3.4	0.4
2554	89	*	*	*	*	*	*	*
2555	8,586	42.5	37.8	7.3	0.0	0.2	12.1	0.0
2556	634	18.5	60.3	5.8	0.2	0.0	15.3	0.0
2557	13	*	*	*	*	*	*	*
2558	7	*	*	*	*	*	*	*
2559	2	*	*	*	*	*	*	*
2560	6	*	*	*	*	*	*	*
2561	3	*	*	*	*	*	*	*
2562	32	*	*	*	*	*	*	*
2563	53	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2564	22	*	*	*	*	*	*	*
3001	1,674	13.4	62.0	17.6	0.0	0.4	6.6	0.0
3002	466	14.2	46.1	29.0	0.0	0.0	10.7	0.0
3004	269	48.3	0.0	45.7	0.0	0.7	5.2	0.0
3005	149	87.2	1.3	6.7	0.0	1.3	3.4	0.0
3007	25	*	*	*	*	*	*	*
3201	617	88.2	0.3	7.3	0.0	0.0	4.2	0.0
3301	4,886	25.7	30.3	21.5	0.1	5.4	16.8	0.1
3304	137	44.5	15.3	16.8	0.0	12.4	10.9	0.0
3305	262	57.3	12.2	20.6	0.0	1.5	8.4	0.0
3306	403	79.2	1.2	8.2	0.0	0.5	10.9	0.0
3308	137	86.1	0.0	6.6	0.0	0.0	7.3	0.0
3312	604	19.9	21.5	36.8	0.2	1.2	20.4	0.2
3550	73	*	*	*	*	*	*	*
3551	15	*	*	*	*	*	*	*
3552	10	*	*	*	*	*	*	*
3701	1,225	80.2	0.0	13.0	0.0	0.0	6.8	0.0
3850	4,112	59.3	13.1	8.7	0.1	0.2	18.7	0.0
3901	1,868	29.2	52.9	7.0	0.1	0.3	10.5	0.0
3902	71	*	*	*	*	*	*	*
3903	35	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3950	2,890	42.8	23.6	15.8	0.1	0.7	15.7	1.4
3954	25	*	*	*	*	*	*	*
4106	2,992	6.6	36.8	49.0	0.3	0.4	6.9	0.0
4107	3,334	33.2	26.9	25.2	0.2	1.5	12.9	0.1
4109	1,181	71.8	3.0	13.0	0.0	0.3	11.9	0.0
4110	5	*	*	*	*	*	*	*
4112	689	0.4	0.9	98.4	0.0	0.0	0.3	0.0
4115	4,459	23.7	29.0	24.8	0.3	12.0	10.2	0.0
4351	739	78.8	0.8	10.4	0.0	0.1	9.6	0.3
4352	214	62.1	8.4	7.9	0.0	0.0	21.5	0.0
4353	662	53.8	0.9	41.7	0.0	0.0	3.6	0.0
4354	953	81.1	3.6	5.7	0.1	0.4	9.0	0.1
4355	395	76.5	0.5	9.4	1.5	0.0	12.2	0.0
4356	556	75.7	0.4	12.6	0.2	0.4	10.8	0.0
4358	36	*	*	*	*	*	*	*
4359	793	63.6	0.1	34.2	0.0	0.0	2.0	0.1
4360	298	84.6	0.0	10.7	0.0	0.0	4.7	0.0
4361	343	89.8	0.0	5.8	0.0	0.0	4.4	0.0
4362	389	86.6	0.5	4.4	0.0	0.3	8.2	0.0
4363	1,024	89.3	0.0	4.6	0.0	0.1	5.9	0.2
4401	1,285	81.2	1.0	10.2	0.1	0.5	6.9	0.2

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4501	1,198	86.4	0.5	8.1	0.0	0.0	4.9	0.1
4601	1,550	64.0	1.7	24.1	0.1	0.1	10.0	0.1
4701	1,543	86.8	1.6	4.5	0.1	0.5	6.5	0.0
5050	1,788	43.2	39.4	4.1	0.1	1.2	12.0	0.0
5051	3,011	46.3	32.7	5.1	0.0	2.5	13.3	0.1
5052	3,343	28.9	37.9	2.8	0.1	19.1	11.2	0.1
5055	1,977	91.9	2.1	0.6	0.0	5.2	0.4	0.0
5101	2,010	60.9	10.7	13.7	0.2	1.4	12.8	0.2
5102	60	*	*	*	*	*	*	*
5202	972	80.2	1.2	4.4	1.0	0.4	12.7	0.0
5206	32	*	*	*	*	*	*	*
5601	1,224	84.6	1.1	5.5	0.2	0.2	8.3	0.0
5801	2,003	69.3	6.4	8.7	0.0	0.1	15.0	0.4
5803	201	82.6	1.0	9.0	0.0	0.5	7.0	0.0
5804	44	*	*	*	*	*	*	*
5806	181	60.2	13.3	11.6	0.0	0.0	14.4	0.6
6101	68	*	*	*	*	*	*	*
6102	177	75.1	1.1	10.7	0.0	0.0	13.0	0.0
6103	4,209	43.3	29.5	12.9	0.2	0.2	13.8	0.0
6105	35	*	*	*	*	*	*	*
6106	23	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6301	3,864	32.8	28.6	24.8	0.0	1.7	12.0	0.1
6312	2,372	34.7	46.9	4.8	0.0	1.5	12.0	0.0
6316	1,489	60.7	16.1	9.3	0.2	3.0	10.6	0.1
6318	71	*	*	*	*	*	*	*
6320	315	50.5	39.4	4.4	0.0	0.3	5.4	0.0
6325	386	89.6	1.0	2.8	0.3	0.0	6.2	0.0
6335	1,510	72.3	12.3	4.0	0.0	6.5	4.9	0.0
6345	114	88.6	2.6	6.1	0.0	0.0	2.6	0.0
6360	1,233	23.8	48.9	18.4	0.0	1.3	7.5	0.1
7306	2,771	36.1	28.0	23.5	0.0	0.2	12.0	0.2
7401	2,539	69.1	5.4	10.1	0.1	0.1	15.0	0.0
7601	698	90.3	0.1	5.9	0.0	0.0	3.7	0.0
7850	1,374	80.8	0.6	9.8	0.0	0.1	8.6	0.1
7852	29	*	*	*	*	*	*	*
8101	2,107	2.3	90.6	1.6	0.1	0.2	5.1	0.0
8102	548	9.9	10.4	75.9	0.0	0.2	3.6	0.0
8105	652	32.2	58.7	2.1	0.5	0.6	5.8	0.0
8106	610	16.1	57.7	14.1	0.0	0.0	11.8	0.3
8150	35	*	*	*	*	*	*	*
8151	3,930	31.1	39.6	11.8	0.1	1.9	15.5	0.1
8152	45	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8153	13	*	*	*	*	*	*	*
8155	43	*	*	*	*	*	*	*
8157	10	*	*	*	*	*	*	*
8201	3,361	38.1	39.4	6.0	0.1	3.8	12.5	0.1
8202	5,569	86.0	10.0	1.9	0.0	0.3	1.8	0.0
8203	517	53.2	27.3	5.6	0.0	0.8	13.2	0.0
8250	752	15.4	70.2	4.4	0.0	0.7	9.2	0.1
8252	191	1.0	90.6	1.0	0.0	0.5	6.8	0.0
8255	2,673	42.2	31.7	12.2	0.1	2.8	10.8	0.1
8257	2,670	52.5	9.5	0.6	0.1	35.7	1.7	0.0
8263	807	56.4	13.9	17.6	0.1	1.9	9.9	0.2
8264	907	58.2	25.4	5.8	0.0	1.4	9.2	0.0
8267	880	47.2	16.1	26.5	0.1	0.7	9.2	0.2
8901	1,137	1.8	92.7	1.6	0.0	0.4	3.5	0.0
8902	1,694	4.1	90.0	2.2	0.0	0.1	3.5	0.1
8903	1,696	2.2	92.0	1.5	0.1	0.8	3.3	0.1
8904	2,623	2.4	91.7	1.9	0.1	1.0	3.0	0.0
8905	1,999	62.2	23.9	9.8	0.1	0.3	3.7	0.0
8909	863	2.4	90.6	1.7	0.0	0.1	5.1	0.0
9045	2,217	6.9	25.5	63.6	0.2	0.0	3.6	0.1
9050	1,308	11.9	69.6	15.1	0.0	0.0	3.3	0.1

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9051	1,894	5.2	6.8	85.7	0.0	0.0	2.2	0.1
9053	1,097	4.9	89.1	1.5	0.5	1.2	2.8	0.1
9055	317	1.3	97.5	0.9	0.0	0.0	0.3	0.0
9058	1,296	19.1	65.9	0.8	0.0	10.7	3.5	0.0
9066	1,467	0.8	95.2	1.6	0.0	0.2	2.2	0.0
9068	1,164	0.8	97.1	0.3	0.0	0.2	1.6	0.0
9069	3,275	0.6	98.0	0.4	0.0	0.1	0.9	0.0
9070	1,705	3.4	91.5	1.5	0.2	1.3	2.0	0.1
9072	1,581	1.6	92.8	1.1	0.0	0.4	4.0	0.1
9080	3,396	20.5	73.8	1.9	0.0	0.3	3.5	0.0
9082	122	0.0	96.7	1.6	0.0	0.0	1.6	0.0
9083	352	27.0	59.9	8.0	0.0	2.8	2.3	0.0
9084	371	3.0	93.0	1.9	0.0	0.3	1.9	0.0
9087	35	*	*	*	*	*	*	*
Michigan	193,486	46.6	28.7	13.2	0.3	2.1	9.0	0.1
National (Prior Year)	712,311	37.6	23.5	23.2	1.4	2.6	6.5	5.1

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

(4) Prenatal program participation

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months	12 - 23 Months	24 - 59 Months
		%	%	%
0101	149	29.5	20.8	49.7
0102	478	28.7	22.4	49.0
0103	503	27.8	20.3	51.9
0104	165	29.1	21.2	49.7
0105	88	*	*	*
0250	147	25.9	27.2	46.9
0251	135	29.6	21.5	48.9
0252	177	19.2	20.9	59.9
0253	157	33.8	21.0	45.2
0401	630	27.6	23.3	49.0
0402	433	31.9	22.6	45.5
0403	185	29.7	27.0	43.2
0404	147	32.0	18.4	49.7
0501	599	36.4	17.2	46.4
0503	129	31.8	26.4	41.9
0504	521	30.7	21.1	48.2
0505	311	37.6	21.5	40.8
0506	213	24.4	23.9	51.6
0507	184	17.9	24.5	57.6
0508	73	*	*	*
0650	1,353	29.3	23.8	46.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0651	831	24.9	25.3	49.8
0652	426	27.5	23.9	48.6
0653	660	23.9	20.9	55.2
0654	479	28.8	21.9	49.3
0655	411	29.0	23.8	47.2
0657	108	22.2	24.1	53.7
0701	605	26.1	24.5	49.4
0702	96	*	*	*
0703	219	27.9	27.4	44.7
0704	78	*	*	*
0705	121	24.8	18.2	57.0
0706	12	*	*	*
0850	995	28.7	21.5	49.7
0851	888	29.3	22.2	48.5
0901	2,001	24.8	27.8	47.4
0902	59	*	*	*
1051	1,444	30.8	22.4	46.8
1101	1,823	37.1	18.9	44.0
1102	880	36.9	18.1	45.0
1106	104	33.7	16.3	50.0
1250	1,205	30.2	26.1	43.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months	12 - 23 Months	24 - 59 Months
		%	%	%
1251	877	34.7	18.9	46.4
1252	802	25.9	22.3	51.7
1253	1,173	28.0	21.5	50.5
1350	2,628	29.9	22.7	47.5
1351	76	*	*	*
1352	655	24.6	22.4	53.0
1401	298	36.2	17.8	46.0
1402	180	28.3	18.3	53.3
1403	38	*	*	*
1501	1,028	26.2	23.1	50.8
1503	1,187	26.6	21.8	51.6
1504	2,431	26.7	24.7	48.6
1505	522	27.8	21.8	50.4
1506	793	28.0	22.3	49.7
1507	751	27.0	22.2	50.7
1509	511	22.9	24.3	52.8
1511	874	24.7	26.2	49.1
1512	470	24.5	25.7	49.8
1514	466	23.8	26.4	49.8
1701	758	27.7	22.3	50.0
1950	730	34.9	19.6	45.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1952	174	35.1	18.4	46.6
1953	413	27.8	22.3	49.9
1954	576	35.6	19.4	45.0
1955	166	28.3	16.9	54.8
1957	590	28.5	22.7	48.8
1963	102	26.5	22.5	51.0
2150	777	30.4	25.5	44.1
2151	200	32.0	26.5	41.5
2152	37	*	*	*
2251	569	26.5	22.5	51.0
2252	264	27.7	21.2	51.1
2554	89	*	*	*
2555	8,586	33.5	20.3	46.2
2556	634	23.7	21.5	54.9
2557	13	*	*	*
2558	7	*	*	*
2559	2	*	*	*
2560	6	*	*	*
2561	3	*	*	*
2562	32	*	*	*
2563	53	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months	12 - 23 Months	24 - 59 Months
		%	%	%
2564	22	*	*	*
3001	1,674	30.2	18.5	51.3
3002	466	27.7	19.1	53.2
3004	269	36.8	22.3	40.9
3005	149	40.9	12.8	46.3
3007	25	*	*	*
3201	617	31.8	22.9	45.4
3301	4,886	32.8	21.6	45.6
3304	137	24.8	19.7	55.5
3305	262	29.0	21.0	50.0
3306	403	28.8	22.6	48.6
3308	137	35.0	14.6	50.4
3312	604	25.2	23.3	51.5
3550	73	*	*	*
3551	15	*	*	*
3552	10	*	*	*
3701	1,225	27.8	24.1	48.2
3850	4,112	27.7	22.4	49.9
3901	1,868	30.8	22.2	47.0
3902	71	*	*	*
3903	35	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
3950	2,890	31.1	21.8	47.2
3954	25	*	*	*
4106	2,992	27.9	21.8	50.4
4107	3,334	32.8	21.4	45.7
4109	1,181	27.3	23.9	48.9
4110	5	*	*	*
4112	689	29.2	19.6	51.2
4115	4,459	32.2	20.8	47.0
4351	739	26.3	24.2	49.5
4352	214	26.2	25.7	48.1
4353	662	24.6	20.5	54.8
4354	953	28.8	24.8	46.5
4355	395	24.1	23.8	52.2
4356	556	27.2	21.8	51.1
4358	36	*	*	*
4359	793	24.8	24.8	50.3
4360	298	25.8	21.8	52.3
4361	343	29.7	21.9	48.4
4362	389	28.5	23.4	48.1
4363	1,024	27.9	23.3	48.7
4401	1,285	24.4	26.0	49.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
4501	1,198	28.3	21.3	50.4
4601	1,550	34.3	20.3	45.5
4701	1,543	29.1	22.2	48.7
5050	1,788	32.9	24.9	42.2
5051	3,011	35.4	23.3	41.3
5052	3,343	33.3	24.8	41.9
5055	1,977	21.0	22.4	56.6
5101	2,010	29.0	23.3	47.7
5102	60	*	*	*
5202	972	26.0	25.4	48.6
5206	32	*	*	*
5601	1,224	35.0	20.3	44.8
5801	2,003	33.0	22.2	44.8
5803	201	23.9	22.9	53.2
5804	44	*	*	*
5806	181	62.4	17.1	20.4
6101	68	*	*	*
6102	177	22.0	27.1	50.8
6103	4,209	30.2	23.3	46.4
6105	35	*	*	*
6106	23	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
6301	3,864	36.8	19.3	43.9
6312	2,372	31.9	19.2	48.9
6316	1,489	32.5	20.6	46.9
6318	71	*	*	*
6320	315	38.7	14.9	46.3
6325	386	37.0	17.1	45.9
6335	1,510	30.1	20.1	49.8
6345	114	32.5	24.6	43.0
6360	1,233	35.3	17.4	47.4
7306	2,771	33.7	20.1	46.2
7401	2,539	28.7	26.0	45.4
7601	698	31.2	21.8	47.0
7850	1,374	25.9	25.0	49.1
7852	29	*	*	*
8101	2,107	32.1	18.9	49.0
8102	548	28.5	20.3	51.3
8105	652	32.5	17.2	50.3
8106	610	30.2	23.0	46.9
8150	35	*	*	*
8151	3,930	30.3	21.7	48.0
8152	45	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
8153	13	*	*	*
8155	43	*	*	*
8157	10	*	*	*
8201	3,361	37.5	20.7	41.7
8202	5,569	27.5	22.2	50.3
8203	517	31.3	19.0	49.7
8250	752	40.2	18.0	41.9
8252	191	19.9	23.0	57.1
8255	2,673	37.8	19.8	42.5
8257	2,670	30.8	20.0	49.1
8263	807	33.7	20.8	45.5
8264	907	32.3	23.0	44.7
8267	880	33.4	21.8	44.8
8901	1,137	31.0	19.6	49.4
8902	1,694	33.5	19.7	46.8
8903	1,696	33.5	20.2	46.2
8904	2,623	33.7	20.9	45.4
8905	1,999	34.6	18.8	46.6
8909	863	32.0	20.9	47.2
9045	2,217	31.3	19.7	49.0
9050	1,308	29.3	20.3	50.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Age Distribution by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
9051	1,894	32.8	21.6	45.6
9053	1,097	46.9	16.0	37.1
9055	317	86.8	2.5	10.7
9058	1,296	29.7	20.9	49.4
9066	1,467	27.5	25.6	46.8
9068	1,164	33.5	19.5	47.0
9069	3,275	30.4	21.8	47.8
9070	1,705	36.2	18.6	45.2
9072	1,581	28.1	23.3	48.6
9080	3,396	27.8	24.0	48.3
9082	122	40.2	23.8	36.1
9083	352	99.7	0.3	0.0
9084	371	88.1	2.7	9.2
9087	35	*	*	*
Michigan	193,486	31.2	21.6	47.2
National (Prior Year)	712,311	35.1	21.8	43.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low		High	Short Stature		Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0101	38	*	*	59	*	*	*	32	*	*	34	*
0102	101	7.9 (19)	5.9 (46)	180	10.6 (114)	3.3 (31)	11.1 (90)	77	*	*	72	*
0103	104	6.7 (11)	13.5 (110)	191	6.3 (27)	3.7 (38)	10.5 (78)	87	*	*	93	*
0104	33	*	*	68	*	*	*	33	*	*	36	*
0105	31	*	*	44	*	*	*	13	*	*	14	*
0250	33	*	*	73	*	*	*	24	*	*	30	*
0251	25	*	*	59	*	*	*	25	*	*	26	*
0252	28	*	*	80	*	*	*	44	*	*	43	*
0253	41	*	*	80	*	*	*	20	*	*	18	*
0401	142	4.9 (2)	9.2 (96)	305	8.2 (72)	5.2 (81)	6.6 (5)	112	15.2 (51)	10.7 (15)	86	*
0402	99	*	*	200	3.0 (2)	5.5 (91)	15.0 (123)	76	*	*	67	*
0403	43	*	*	76	*	*	*	23	*	*	28	*
0404	41	*	*	59	*	*	*	23	*	*	23	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0501	132	13.6 (82)	7.6 (75)	338	6.5 (30)	12.4 (131)	10.9 (86)	122	25.4 (102)	19.7 (93)	85	*
0503	30	*	*	77	*	*	*	30	*	*	18	*
0504	114	8.8 (24)	5.3 (40)	313	8.3 (74)	5.8 (97)	8.3 (15)	118	16.1 (59)	12.7 (36)	92	*
0505	79	*	*	169	10.7 (116)	3.0 (25)	10.7 (83)	57	*	*	56	*
0506	40	*	*	118	8.5 (81)	6.8 (107)	10.2 (69)	44	*	*	45	*
0507	18	*	*	110	10.0 (105)	0.0 (1)	16.4 (132)	59	*	*	39	*
0508	7	*	*	38	*	*	*	19	*	*	11	*
0650	293	10.9 (56)	10.2 (103)	690	7.2 (50)	4.5 (61)	11.3 (93)	253	13.8 (29)	19.0 (90)	220	5.9 (12)
0651	162	4.9 (2)	7.4 (70)	417	8.6 (83)	5.3 (85)	9.8 (57)	172	20.3 (95)	14.5 (56)	154	6.5 (15)
0652	92	*	*	199	12.1 (127)	3.0 (25)	12.1 (105)	77	*	*	80	*
0653	133	10.5 (48)	8.3 (84)	325	11.1 (121)	3.7 (38)	11.4 (96)	134	18.7 (88)	14.2 (51)	115	10.4 (33)
0654	113	9.7 (34)	8.0 (80)	208	10.1 (107)	4.8 (74)	9.6 (49)	89	*	*	66	*
0655	84	*	*	175	3.4 (3)	4.0 (45)	13.1 (115)	70	*	*	63	*
0657	17	*	*	54	*	*	*	20	*	*	20	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0701	135	8.9 (25)	10.4 (107)	195	8.7 (85)	2.1 (10)	10.3 (72)	96	*	*	112	3.6 (5)
0702	28	*	*	38	*	*	*	19	*	*	17	*
0703	41	*	*	88	*	*	*	42	*	*	46	*
0704	12	*	*	23	*	*	*	10	*	*	14	*
0705	24	*	*	40	*	*	*	27	*	*	25	*
0706	1	*	*	5	*	*	*	2	*	*	2	*
0850	206	10.7 (53)	7.8 (77)	593	6.9 (41)	4.4 (58)	11.5 (97)	219	14.6 (41)	17.8 (85)	141	6.4 (14)
0851	194	11.9 (67)	6.7 (61)	578	6.1 (24)	4.5 (61)	8.7 (22)	214	14.5 (39)	16.8 (80)	176	4.5 (7)
0901	450	12.7 (75)	8.7 (86)	836	8.4 (77)	1.6 (5)	11.7 (102)	362	18.5 (87)	14.6 (58)	393	9.4 (27)
0902	9	*	*	36	*	*	*	19	*	*	15	*
1051	335	10.1 (42)	8.7 (86)	741	3.5 (4)	8.0 (121)	7.7 (8)	265	20.8 (98)	10.2 (10)	250	22.0 (77)
1101	448	10.0 (39)	3.8 (12)	1,201	7.6 (61)	5.3 (85)	9.2 (38)	485	16.9 (66)	11.3 (21)	565	9.4 (27)
1102	225	6.2 (9)	6.7 (61)	567	7.2 (50)	3.9 (43)	11.5 (97)	211	17.1 (67)	15.2 (63)	272	2.6 (4)
1106	26	*	*	66	*	*	*	26	*	*	33	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
1250	315	9.2 (27)	7.9 (79)	579	10.9 (119)	6.9 (109)	9.0 (31)	182	13.7 (27)	8.2 (2)	166	19.9 (72)
1251	201	10.4 (44)	8.0 (80)	331	7.3 (52)	0.6 (3)	10.3 (72)	148	14.9 (45)	16.2 (74)	142	14.8 (52)
1252	138	8.0 (21)	8.7 (86)	334	7.5 (58)	5.4 (88)	9.6 (49)	142	18.3 (86)	16.2 (74)	126	13.5 (43)
1253	227	4.8 (1)	7.5 (74)	498	10.4 (110)	6.0 (100)	12.2 (107)	194	16.5 (64)	13.4 (44)	163	13.5 (43)
1350	667	11.8 (65)	6.1 (51)	1,506	7.0 (43)	7.5 (115)	9.9 (60)	458	15.1 (50)	15.7 (69)	435	19.1 (65)
1351	13	*	*	59	*	*	*	25	*	*	25	*
1352	135	14.1 (87)	9.6 (97)	298	9.4 (99)	9.1 (128)	11.7 (102)	91	*	*	97	*
1401	56	*	*	119	8.4 (77)	7.6 (118)	10.9 (86)	36	*	*	47	*
1402	37	*	*	95	*	*	*	38	*	*	33	*
1403	3	*	*	11	*	*	*	5	*	*	4	*
1501	222	10.4 (44)	9.0 (92)	669	6.4 (28)	5.7 (93)	12.0 (104)	272	14.7 (43)	14.0 (50)	291	7.9 (21)
1503	264	7.2 (13)	10.2 (103)	807	4.5 (9)	4.2 (53)	11.3 (93)	329	13.7 (27)	16.4 (76)	373	15.0 (53)
1504	427	7.7 (18)	6.8 (65)	1,391	4.5 (9)	3.9 (43)	10.4 (76)	585	17.4 (71)	15.0 (62)	701	22.4 (78)
1505	111	10.8 (55)	9.9 (100)	273	5.1 (14)	4.4 (58)	8.8 (26)	129	14.0 (32)	12.4 (33)	127	29.1 (87)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A 2020 Pediatric Nutrition Surveillance ⁽¹⁾ Michigan Comparison of Growth and Anemia Indicators by Clinic Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)												
Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)							Anemia Low Hb/Hct ⁽⁸⁾	
	Low	High		Short Stature	Under- Weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
1506	164	8.5 (23)	6.7 (61)	578	6.9 (41)	2.6 (15)	12.5 (111)	226	17.3 (70)	19.5 (92)	247	9.7 (29)
1507	156	12.2 (70)	9.0 (92)	410	4.9 (12)	3.2 (28)	8.0 (12)	169	15.4 (54)	10.1 (8)	228	43.9 (96)
1509	102	12.7 (75)	9.8 (99)	321	8.7 (85)	2.8 (19)	10.9 (86)	148	18.9 (90)	14.9 (61)	169	7.1 (19)
1511	162	10.5 (48)	4.9 (30)	578	5.7 (18)	4.5 (61)	10.0 (65)	236	15.7 (56)	13.1 (40)	292	20.5 (73)
1512	91	*	*	328	5.8 (19)	5.2 (81)	10.1 (67)	134	11.9 (10)	15.7 (69)	153	19.6 (69)
1514	92	*	*	285	7.4 (53)	4.2 (53)	12.3 (109)	120	13.3 (21)	15.8 (71)	134	37.3 (95)
1701	153	7.2 (13)	14.4 (111)	320	10.0 (105)	5.3 (85)	18.1 (134)	155	14.8 (44)	19.4 (91)	83	*
1950	181	10.5 (48)	6.6 (58)	352	8.8 (89)	5.7 (93)	9.7 (53)	134	13.4 (24)	11.9 (26)	124	1.6 (3)
1952	47	*	*	98	*	*	*	29	*	*	32	*
1953	71	*	*	224	7.1 (47)	2.2 (11)	11.6 (100)	83	*	*	82	*
1954	136	5.9 (7)	7.4 (70)	313	7.0 (43)	5.1 (78)	11.5 (97)	109	12.8 (16)	16.5 (77)	108	0.9 (2)
1955	34	*	*	89	*	*	*	34	*	*	30	*
1957	109	12.8 (77)	11.9 (109)	311	10.6 (114)	5.8 (97)	15.1 (124)	122	17.2 (69)	18.0 (86)	109	0.0 (1)
1963	16	*	*	58	*	*	*	24	*	*	20	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
2150	191	7.3 (16)	7.3 (68)	404	15.1 (134)	1.0 (4)	15.6 (129)	143	17.5 (75)	21.0 (99)	154	13.0 (41)
2151	50	*	*	105	3.8 (5)	2.9 (21)	17.1 (133)	41	*	*	49	*
2152	9	*	*	17	*	*	*	9	*	*	8	*
2251	113	9.7 (34)	6.2 (53)	236	8.1 (69)	4.7 (72)	12.3 (109)	98	*	*	103	12.6 (39)
2252	54	*	*	99	*	*	*	42	*	*	49	*
2554	14	*	*	70	*	*	*	44	*	*	43	*
2555	2,107	13.0 (78)	4.4 (19)	3,164	9.1 (95)	2.9 (21)	11.1 (90)	1,253	15.2 (51)	12.0 (28)	1,351	19.1 (65)
2556	94	*	*	300	12.7 (130)	2.0 (9)	13.0 (114)	141	9.2 (3)	17.0 (82)	124	29.8 (88)
2557	1	*	*	5	*	*	*	2	*	*	4	*
2558	0	*	*	0	*	*	*	0	*	*	0	*
2559	0	*	*	0	*	*	*	0	*	*	0	*
2560	0	*	*	2	*	*	*	2	*	*	1	*
2561	1	*	*	2	*	*	*	1	*	*	0	*
2562	1	*	*	17	*	*	*	9	*	*	10	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
2563	8	*	*	22	*	*	*	6	*	*	14	*
2564	9	*	*	10	*	*	*	1	*	*	2	*
3001	359	13.1 (79)	3.3 (9)	739	5.8 (19)	7.6 (118)	8.5 (18)	213	17.4 (71)	12.2 (31)	197	7.1 (19)
3002	78	*	*	214	4.7 (11)	7.5 (115)	7.9 (10)	72	*	*	73	*
3004	60	*	*	138	5.1 (14)	8.0 (121)	5.8 (4)	33	*	*	46	*
3005	40	*	*	69	*	*	*	16	*	*	16	*
3007	3	*	*	13	*	*	*	4	*	*	2	*
3201	142	11.3 (59)	8.5 (85)	243	10.7 (116)	4.1 (49)	9.1 (36)	79	*	*	87	*
3301	1,197	11.5 (61)	5.0 (35)	2,304	8.0 (67)	7.7 (120)	9.8 (57)	702	17.8 (79)	17.1 (83)	682	10.1 (31)
3304	18	*	*	84	*	*	*	43	*	*	37	*
3305	51	*	*	146	6.2 (26)	4.1 (49)	8.2 (14)	57	*	*	45	*
3306	77	*	*	190	6.8 (38)	4.2 (53)	5.3 (3)	67	*	*	70	*
3308	40	*	*	69	*	*	*	25	*	*	21	*
3312	101	10.9 (56)	8.9 (91)	282	7.4 (53)	4.6 (70)	10.6 (79)	111	8.1 (2)	10.8 (16)	87	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
3550	21	*	*	20	*	*	*	5	*	*	8	*
3551	3	*	*	2	*	*	*	1	*	*	1	*
3552	2	*	*	7	*	*	*	3	*	*	3	*
3701	282	6.4 (10)	10.3 (105)	697	4.4 (8)	7.3 (113)	8.8 (26)	235	18.7 (88)	13.2 (42)	251	4.0 (6)
3850	886	9.7 (34)	8.8 (89)	2,103	5.2 (16)	13.7 (133)	8.9 (28)	665	17.7 (76)	16.5 (77)	579	17.1 (60)
3901	422	10.0 (39)	3.6 (11)	896	5.5 (17)	7.4 (114)	8.4 (17)	260	19.2 (92)	13.1 (40)	318	19.8 (71)
3902	20	*	*	40	*	*	*	10	*	*	11	*
3903	5	*	*	21	*	*	*	7	*	*	10	*
3950	609	11.5 (61)	7.6 (75)	1,480	6.6 (33)	5.7 (93)	8.7 (22)	472	16.3 (60)	14.2 (51)	565	25.8 (83)
3954	4	*	*	8	*	*	*	2	*	*	2	*
4106	604	15.2 (99)	4.5 (20)	1,345	11.7 (126)	2.8 (19)	15.1 (124)	616	18.0 (83)	18.0 (86)	597	10.7 (34)
4107	796	15.1 (98)	6.0 (48)	1,541	11.4 (125)	4.9 (75)	9.0 (31)	572	15.0 (47)	9.4 (6)	584	23.3 (80)
4109	223	8.1 (22)	4.9 (30)	614	11.1 (121)	3.3 (31)	9.9 (60)	247	17.8 (79)	10.1 (8)	234	4.7 (9)
4110	2	*	*	0	*	*	*	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
4112	151	7.9 (19)	6.6 (58)	561	6.8 (38)	4.5 (61)	13.5 (116)	233	13.3 (21)	20.6 (98)	137	8.0 (22)
4115	979	14.4 (90)	5.3 (40)	1,835	10.5 (112)	2.3 (12)	10.6 (79)	766	14.2 (36)	12.3 (32)	805	13.5 (43)
4351	162	9.3 (31)	6.2 (53)	403	10.4 (110)	4.0 (45)	9.9 (60)	161	13.0 (19)	12.4 (33)	140	9.3 (25)
4352	50	*	*	99	*	*	*	43	*	*	34	*
4353	135	8.9 (25)	7.4 (70)	265	8.7 (85)	4.5 (61)	14.7 (122)	111	14.4 (38)	21.6 (101)	98	*
4354	207	7.2 (13)	6.3 (55)	476	7.4 (53)	3.8 (41)	10.9 (86)	188	13.8 (29)	11.2 (20)	171	5.3 (10)
4355	77	*	*	157	6.4 (28)	7.0 (111)	9.6 (49)	71	*	*	71	*
4356	122	10.7 (53)	4.9 (30)	245	8.2 (72)	4.5 (61)	9.0 (31)	103	12.6 (13)	13.6 (46)	83	*
4358	1	*	*	29	*	*	*	17	*	*	16	*
4359	167	5.4 (5)	9.0 (92)	354	8.5 (81)	3.7 (38)	11.6 (100)	132	18.9 (90)	19.7 (93)	104	6.7 (17)
4360	54	*	*	107	6.5 (30)	1.9 (7)	9.3 (40)	51	*	*	57	*
4361	74	*	*	124	6.5 (30)	0.0 (1)	14.5 (121)	56	*	*	51	*
4362	90	*	*	167	9.0 (91)	1.8 (6)	10.2 (69)	74	*	*	83	*
4363	229	9.6 (33)	10.0 (102)	378	8.7 (85)	1.9 (7)	12.7 (113)	168	22.6 (101)	16.1 (73)	176	10.2 (32)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
4401	252	12.3 (72)	9.9 (100)	574	5.9 (22)	4.2 (53)	9.6 (49)	210	16.7 (65)	14.3 (54)	239	19.2 (67)
4501	249	9.2 (27)	7.2 (67)	704	7.8 (65)	6.1 (103)	10.4 (76)	282	17.7 (76)	15.6 (68)	225	12.4 (38)
4601	394	6.1 (8)	8.1 (82)	544	9.4 (99)	2.6 (15)	9.4 (43)	231	12.1 (11)	13.0 (38)	260	31.5 (90)
4701	312	12.2 (70)	6.1 (51)	530	8.1 (69)	2.6 (15)	8.5 (18)	260	13.5 (25)	11.5 (23)	283	4.6 (8)
5050	439	12.5 (73)	6.6 (58)	673	6.7 (37)	3.6 (36)	10.1 (67)	242	17.4 (71)	7.0 (1)	241	8.7 (24)
5051	808	11.9 (67)	4.2 (15)	1,499	10.3 (108)	6.9 (109)	9.1 (36)	442	15.6 (55)	12.0 (28)	370	13.5 (43)
5052	840	14.8 (94)	4.5 (20)	1,372	9.2 (96)	5.2 (81)	9.7 (53)	451	12.6 (13)	11.1 (17)	434	6.9 (18)
5055	350	7.4 (17)	5.1 (38)	661	4.2 (6)	3.2 (28)	15.6 (129)	385	15.3 (53)	18.2 (88)	289	16.3 (58)
5101	481	10.0 (39)	6.7 (61)	955	5.8 (19)	5.0 (77)	9.0 (31)	325	14.2 (36)	14.2 (51)	306	12.1 (37)
5102	11	*	*	41	*	*	*	17	*	*	20	*
5202	202	5.0 (4)	7.4 (70)	579	9.3 (97)	3.6 (36)	12.1 (105)	214	20.6 (97)	15.4 (66)	123	6.5 (15)
5206	6	*	*	21	*	*	*	6	*	*	3	*
5601	261	9.2 (27)	10.3 (105)	685	9.9 (104)	2.9 (21)	10.7 (83)	245	19.2 (92)	14.7 (60)	224	13.8 (48)
5801	478	9.8 (38)	7.3 (68)	850	9.6 (102)	4.2 (53)	9.5 (45)	292	17.8 (79)	14.4 (55)	311	17.4 (61)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
5803	28	*	*	93	*	*	*	42	*	*	43	*
5804	8	*	*	16	*	*	*	8	*	*	7	*
5806	73	*	*	67	*	*	*	10	*	*	16	*
6101	14	*	*	33	*	*	*	12	*	*	11	*
6102	30	*	*	98	*	*	*	46	*	*	36	*
6103	1,075	12.6 (74)	4.8 (28)	2,032	8.4 (77)	4.0 (45)	12.5 (111)	665	17.7 (76)	15.3 (65)	684	15.5 (54)
6105	5	*	*	18	*	*	*	9	*	*	7	*
6106	4	*	*	23	*	*	*	8	*	*	11	*
6301	990	13.3 (80)	5.8 (44)	1,710	9.8 (103)	3.2 (28)	9.8 (57)	645	17.1 (67)	13.0 (38)	615	23.6 (81)
6312	524	13.9 (85)	5.0 (35)	1,038	11.2 (123)	4.5 (61)	8.5 (18)	432	16.4 (62)	9.0 (4)	396	17.4 (61)
6316	315	11.4 (60)	7.0 (66)	702	7.7 (63)	5.1 (78)	10.3 (72)	282	14.5 (39)	12.1 (30)	266	12.0 (35)
6318	55	*	*	48	*	*	*	3	*	*	3	*
6320	88	*	*	160	7.5 (58)	8.8 (127)	9.4 (43)	52	*	*	42	*
6325	86	*	*	226	7.5 (58)	3.5 (35)	9.7 (53)	78	*	*	78	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
6335	265	10.6 (52)	4.5 (20)	822	7.8 (65)	5.4 (88)	12.2 (107)	300	14.0 (32)	12.7 (36)	266	27.8 (85)
6345	20	*	*	57	*	*	*	16	*	*	13	*
6360	268	13.8 (84)	4.9 (30)	610	10.7 (116)	8.5 (124)	9.3 (40)	185	14.1 (34)	13.5 (45)	184	29.9 (89)
7306	598	13.7 (83)	5.5 (42)	1,416	6.1 (24)	5.4 (88)	9.5 (45)	429	16.3 (60)	15.2 (63)	514	16.9 (59)
7401	617	10.4 (44)	6.0 (48)	1,375	7.1 (47)	6.8 (107)	9.5 (45)	415	17.8 (79)	15.4 (66)	490	12.0 (35)
7601	154	5.8 (6)	9.1 (95)	257	12.1 (127)	2.3 (12)	15.2 (126)	104	13.5 (25)	21.2 (100)	114	13.2 (42)
7850	295	10.5 (48)	8.8 (89)	621	6.6 (33)	5.8 (97)	8.9 (28)	232	21.6 (99)	15.9 (72)	258	9.3 (25)
7852	8	*	*	15	*	*	*	8	*	*	4	*
8101	458	18.3 (111)	1.5 (1)	1,108	11.2 (123)	10.3 (130)	9.7 (53)	372	13.2 (20)	11.3 (21)	449	34.5 (93)
8102	106	9.4 (32)	11.3 (108)	272	7.4 (53)	8.5 (124)	14.3 (120)	122	16.4 (62)	19.7 (93)	112	19.6 (69)
8105	50	*	*	335	4.2 (6)	3.3 (31)	15.2 (126)	144	16.0 (58)	14.6 (58)	146	8.2 (23)
8106	144	16.0 (104)	6.3 (55)	218	8.3 (74)	4.6 (70)	10.6 (79)	66	*	*	85	*
8150	1	*	*	15	*	*	*	10	*	*	4	*
8151	898	14.1 (87)	4.2 (15)	1,573	7.1 (47)	5.1 (78)	9.9 (60)	654	15.7 (56)	11.9 (26)	624	23.1 (79)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
8152	15	*	*	17	*	*	*	6	*	*	11	*
8153	3	*	*	3	*	*	*	0	*	*	1	*
8155	5	*	*	27	*	*	*	21	*	*	12	*
8157	1	*	*	2	*	*	*	0	*	*	0	*
8201	871	13.5 (81)	5.5 (42)	1,062	8.9 (90)	4.1 (49)	9.9 (60)	378	20.4 (96)	10.6 (12)	375	19.5 (68)
8202	1,151	10.4 (44)	5.0 (35)	1,805	6.6 (33)	4.5 (61)	9.3 (40)	873	12.7 (15)	10.4 (11)	896	15.8 (55)
8203	116	12.1 (69)	7.8 (77)	205	4.9 (12)	3.4 (34)	11.2 (92)	95	*	*	80	*
8250	202	14.4 (90)	4.5 (20)	248	9.3 (97)	6.0 (100)	8.1 (13)	92	*	*	85	*
8252	26	*	*	68	*	*	*	30	*	*	36	*
8255	621	11.8 (65)	5.8 (44)	1,183	8.1 (69)	4.0 (45)	10.7 (83)	412	15.0 (47)	16.7 (79)	406	16.0 (57)
8257	516	14.5 (92)	1.9 (2)	1,102	8.0 (67)	6.2 (104)	13.5 (116)	543	11.0 (7)	17.7 (84)	423	18.4 (64)
8263	186	9.7 (34)	6.5 (57)	383	9.4 (99)	2.9 (21)	10.2 (69)	162	21.6 (99)	11.7 (25)	130	6.2 (13)
8264	210	7.1 (12)	5.2 (39)	311	8.4 (77)	4.5 (61)	10.0 (65)	113	15.0 (47)	10.6 (12)	104	5.8 (11)
8267	185	10.3 (43)	9.7 (98)	292	11.0 (120)	4.1 (49)	11.3 (93)	110	10.0 (4)	14.5 (56)	132	13.6 (47)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
8901	209	14.8 (94)	2.9 (5)	621	9.0 (91)	9.7 (129)	8.5 (18)	224	13.8 (29)	12.5 (35)	253	21.7 (76)
8902	318	16.0 (104)	1.9 (2)	690	6.8 (38)	7.5 (115)	9.0 (31)	263	12.9 (18)	9.9 (7)	309	46.6 (97)
8903	388	15.2 (99)	4.9 (30)	748	10.3 (108)	7.2 (112)	10.3 (72)	273	11.7 (8)	13.2 (42)	298	26.5 (84)
8904	525	15.6 (102)	4.6 (24)	1,403	12.1 (127)	8.5 (124)	10.6 (79)	467	10.9 (6)	16.9 (81)	493	33.9 (91)
8905	393	11.7 (64)	5.9 (46)	820	7.6 (61)	6.6 (105)	13.7 (118)	336	14.9 (45)	18.2 (88)	363	20.7 (74)
8909	190	15.8 (103)	3.2 (8)	374	14.2 (133)	8.0 (121)	13.9 (119)	128	13.3 (21)	25.8 (102)	139	35.3 (94)
9045	466	10.9 (56)	8.2 (83)	915	8.3 (74)	4.9 (75)	15.4 (128)	365	18.1 (84)	20.5 (97)	405	14.6 (50)
9050	277	15.2 (99)	4.7 (26)	498	6.6 (33)	3.0 (25)	9.2 (38)	203	12.3 (12)	13.8 (49)	201	34.3 (92)
9051	468	11.5 (61)	6.0 (48)	759	7.0 (43)	2.6 (15)	15.7 (131)	303	18.2 (85)	20.1 (96)	304	28.3 (86)
9053	360	16.4 (106)	3.9 (13)	642	13.6 (131)	4.7 (72)	7.9 (10)	156	19.9 (94)	11.5 (23)	143	9.8 (30)
9055	273	9.2 (27)	4.8 (28)	238	13.9 (132)	2.5 (14)	6.7 (6)	1	*	*	2	*
9058	280	15.0 (97)	2.5 (4)	412	9.0 (91)	6.6 (105)	9.5 (45)	179	12.8 (16)	10.6 (12)	170	15.9 (56)
9066	355	17.5 (109)	4.2 (15)	439	10.5 (112)	5.7 (93)	8.9 (28)	175	8.0 (1)	13.7 (48)	166	17.5 (63)
9068	299	17.7 (110)	4.7 (26)	503	6.0 (23)	3.8 (41)	8.7 (22)	171	11.7 (8)	11.1 (17)	179	12.8 (40)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
9069	730	16.7 (107)	3.0 (6)	1,421	7.7 (63)	6.0 (100)	7.4 (7)	510	14.1 (34)	8.2 (2)	564	13.8 (48)
9070	376	14.9 (96)	3.5 (10)	810	9.0 (91)	4.4 (58)	7.8 (9)	288	14.6 (41)	9.0 (4)	268	14.6 (50)
9072	333	14.7 (93)	3.0 (6)	573	7.0 (43)	5.2 (81)	8.7 (22)	226	10.2 (5)	11.1 (17)	240	25.4 (82)
9080	714	14.0 (86)	4.6 (24)	1,157	7.4 (53)	5.6 (92)	8.3 (15)	449	17.4 (71)	13.6 (46)	464	21.6 (75)
9082	34	*	*	62	*	*	*	12	*	*	17	*
9083	350	17.4 (108)	4.0 (14)	297	2.0 (1)	13.5 (132)	3.0 (1)	0	*	*	0	*
9084	323	14.2 (89)	4.3 (18)	302	8.6 (83)	17.5 (134)	4.0 (2)	1	*	*	2	*
9087	0	*	*	18	*	*	*	14	*	*	7	*
Michigan	43,761	11.8	6.0	88,932	8.1	5.3	10.3	33,223	15.8	14.1	33,184	16.4
National (Prior Year)	159,266	11.0	6.0	649,753	7.7	2.9	11.2	283,555	16.1	13.8	433,852	14.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
0101	38	*	15	*	21	*	14	*	14	*	76	*	149	2.0 (78)
0102	101	60.4 (77)	54	*	72	*	36	*	51	*	233	81.1 (77)	476	3.8 (132)
0103	104	63.5 (67)	52	*	80	*	27	*	51	*	262	87.4 (46)	503	6.2 (158)
0104	34	*	14	*	21	*	9	*	12	*	82	*	165	4.8 (146)
0105	31	*	13	*	13	*	6	*	11	*	29	*	88	*
0250	33	*	22	*	35	*	13	*	16	*	69	*	147	1.4 (52)
0251	25	*	23	*	21	*	19	*	23	*	70	*	135	1.5 (57)
0252	28	*	15	*	31	*	5	*	15	*	106	74.5 (109)	177	2.3 (86)
0253	41	*	30	*	26	*	14	*	21	*	73	*	157	3.2 (114)
0401	142	70.4 (46)	88	*	107	6.5 (90)	57	*	72	*	325	88.6 (39)	628	2.9 (106)
0402	99	*	84	*	102	18.6 (14)	52	*	75	*	210	84.3 (62)	433	2.3 (86)
0403	43	*	28	*	33	*	22	*	28	*	88	*	185	8.1 (160)
0404	41	*	28	*	16	*	18	*	23	*	81	*	147	2.7 (100)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
0501	147	68.7 (51)	80	*	76	*	46	*	74	*	280	85.4 (57)	597	4.4 (143)
0503	31	*	20	*	24	*	13	*	18	*	57	*	128	1.6 (65)
0504	117	89.7 (1)	57	*	90	*	32	*	52	*	255	89.4 (30)	519	1.5 (57)
0505	79	*	38	*	38	*	19	*	37	*	134	88.8 (36)	311	3.2 (114)
0506	40	*	21	*	35	*	9	*	17	*	109	88.1 (42)	212	9.4 (161)
0507	18	*	20	*	26	*	10	*	19	*	109	90.8 (19)	183	1.1 (43)
0508	7	*	7	*	13	*	3	*	7	*	48	*	73	*
0650	293	71.7 (40)	218	36.7 (10)	251	13.5 (43)	120	22.5 (16)	174	13.2 (13)	662	88.8 (36)	1,352	2.4 (90)
0651	162	69.1 (49)	139	30.2 (31)	167	12.6 (49)	75	*	118	11.0 (21)	434	90.6 (22)	830	4.3 (142)
0652	92	*	58	*	75	*	39	*	51	*	212	90.6 (22)	426	3.8 (132)
0653	134	79.1 (17)	88	*	102	21.6 (8)	53	*	77	*	385	82.9 (69)	660	5.0 (150)
0654	114	74.6 (31)	77	*	76	*	55	*	67	*	245	90.2 (27)	478	5.2 (151)
0655	84	*	53	*	69	*	34	*	43	*	203	74.9 (107)	409	5.4 (153)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
0657	17	*	16	*	19	*	13	*	14	*	60	*	108	5.6 (156)
0701	136	85.3 (3)	91	*	99	*	61	*	73	*	315	90.8 (19)	605	0.5 (17)
0702	28	*	17	*	18	*	12	*	17	*	42	*	95	*
0703	41	*	47	*	49	*	28	*	38	*	100	78.0 (90)	217	4.1 (139)
0704	12	*	13	*	20	*	9	*	12	*	40	*	78	*
0705	24	*	20	*	17	*	17	*	17	*	74	*	121	2.5 (93)
0706	1	*	7	*	5	*	5	*	7	*	4	*	12	*
0850	211	82.5 (10)	125	27.2 (45)	151	15.9 (30)	77	*	110	5.5 (55)	497	78.3 (88)	984	1.1 (43)
0851	199	77.9 (20)	100	39.0 (5)	135	20.7 (9)	65	*	90	*	438	90.4 (25)	885	1.5 (57)
0901	455	65.1 (63)	233	28.3 (42)	383	14.4 (39)	129	24.0 (14)	218	7.8 (39)	986	89.0 (34)	2,001	1.9 (74)
0902	9	*	6	*	9	*	2	*	6	*	37	*	59	*
1051	339	81.7 (12)	210	46.2 (1)	245	27.3 (2)	133	33.8 (2)	180	15.0 (8)	694	90.8 (19)	1,440	1.4 (52)
1101	451	68.3 (54)	285	35.8 (12)	249	17.3 (23)	192	19.3 (25)	240	10.0 (26)	797	71.6 (115)	1,800	3.8 (132)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
1102	225	73.8 (35)	147	33.3 (16)	115	13.0 (45)	104	22.1 (17)	130	11.5 (18)	394	81.0 (78)	869	4.9 (147)
1106	26	*	12	*	13	*	8	*	10	*	53	*	103	1.0 (34)
1250	318	70.8 (44)	137	27.0 (46)	229	14.4 (39)	67	*	121	7.4 (43)	530	79.2 (85)	1,204	2.7 (100)
1251	203	77.3 (23)	91	*	102	15.7 (31)	50	*	84	*	411	83.2 (67)	877	2.7 (100)
1252	140	87.1 (2)	89	*	139	20.1 (11)	50	*	71	*	411	84.9 (60)	792	0.8 (27)
1253	231	84.8 (4)	131	35.9 (11)	186	15.6 (32)	73	*	101	16.8 (4)	611	87.4 (46)	1,161	1.6 (65)
1350	675	70.2 (47)	361	28.0 (43)	470	16.4 (26)	190	14.7 (36)	278	8.6 (34)	1,256	91.6 (17)	2,618	1.0 (34)
1351	13	*	2	*	13	*	5	*	2	*	39	*	76	*
1352	140	60.7 (75)	86	*	110	10.9 (65)	38	*	66	*	348	99.4 (1)	654	0.2 (8)
1401	56	*	65	*	48	*	40	*	60	*	138	75.4 (104)	298	5.4 (153)
1402	37	*	21	*	21	*	10	*	20	*	101	76.2 (99)	180	0.0 (1)
1403	3	*	2	*	4	*	1	*	2	*	28	*	38	*
1501	233	67.4 (58)	124	31.5 (23)	188	13.8 (42)	81	*	112	13.4 (11)	532	86.7 (49)	1,018	1.5 (57)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
1503	265	72.8 (37)	163	33.1 (19)	228	12.3 (52)	108	28.7 (5)	144	17.4 (3)	615	96.6 (7)	1,181	0.9 (29)
1504	585	79.0 (18)	329	28.9 (39)	471	12.7 (47)	173	23.1 (15)	261	9.6 (27)	1,178	88.9 (35)	2,417	0.6 (21)
1505	112	82.1 (11)	44	*	76	*	24	*	40	*	266	98.5 (3)	516	1.2 (45)
1506	166	71.1 (42)	106	21.7 (63)	140	15.0 (36)	70	*	99	*	404	94.8 (12)	787	0.4 (14)
1507	158	84.8 (4)	102	33.3 (16)	111	9.0 (72)	65	*	83	*	165	83.0 (68)	744	2.6 (97)
1509	103	60.2 (78)	67	*	101	19.8 (12)	48	*	64	*	272	92.3 (15)	507	1.6 (65)
1511	168	76.2 (25)	122	32.0 (21)	167	18.6 (14)	67	*	110	14.5 (10)	442	89.4 (30)	870	1.6 (65)
1512	93	*	56	*	88	*	38	*	47	*	241	95.9 (10)	470	0.0 (1)
1514	92	*	72	*	103	29.1 (1)	37	*	51	*	232	99.1 (2)	463	0.2 (8)
1701	153	64.1 (65)	94	*	114	25.4 (3)	53	*	87	*	396	91.9 (16)	757	2.6 (97)
1950	181	84.0 (7)	90	*	105	18.1 (17)	44	*	71	*	326	84.0 (63)	727	4.7 (145)
1952	47	*	19	*	20	*	11	*	17	*	80	*	171	0.0 (1)
1953	73	*	44	*	70	*	17	*	37	*	205	82.4 (72)	409	2.4 (90)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
1954	136	75.7 (29)	81	*	80	*	43	*	69	*	257	82.9 (69)	563	2.0 (78)
1955	34	*	20	*	20	*	12	*	16	*	90	*	165	2.4 (90)
1957	110	78.2 (19)	60	*	89	*	33	*	55	*	283	80.2 (82)	586	2.0 (78)
1963	16	*	11	*	11	*	2	*	10	*	51	*	98	*
2150	192	81.3 (14)	119	42.0 (3)	146	13.0 (45)	81	*	92	*	358	86.3 (52)	773	2.3 (86)
2151	49	*	34	*	34	*	17	*	28	*	85	*	198	1.5 (57)
2152	9	*	9	*	7	*	8	*	8	*	16	*	37	*
2251	114	69.3 (48)	88	*	96	*	55	*	76	*	309	89.6 (29)	569	0.5 (17)
2252	54	*	30	*	46	*	25	*	26	*	139	88.5 (40)	264	3.8 (132)
2554	14	*	9	*	9	*	8	*	9	*	61	*	88	*
2555	2,113	54.9 (89)	1,197	21.5 (64)	1,335	12.3 (52)	653	15.2 (35)	964	6.6 (48)	3,980	81.8 (73)	8,543	2.2 (83)
2556	94	*	76	*	106	12.3 (52)	28	*	54	*	348	83.3 (66)	633	2.1 (82)
2557	1	*	0	*	0	*	1	*	0	*	8	*	13	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
2558	0	*	0	*	0	*	0	*	0	*	7	*	7	*
2559	0	*	0	*	0	*	0	*	0	*	2	*	2	*
2560	0	*	0	*	0	*	0	*	0	*	6	*	6	*
2561	1	*	0	*	0	*	0	*	0	*	1	*	2	*
2562	1	*	5	*	6	*	1	*	4	*	21	*	32	*
2563	8	*	6	*	14	*	4	*	5	*	23	*	53	*
2564	9	*	3	*	3	*	2	*	3	*	5	*	22	*
3001	358	57.5 (85)	182	10.4 (82)	212	7.5 (85)	89	*	180	2.2 (71)	854	71.7 (114)	1,664	3.7 (127)
3002	78	*	48	*	64	*	22	*	48	*	250	77.2 (94)	464	3.7 (127)
3004	61	*	38	*	39	*	15	*	38	*	112	75.9 (101)	265	3.4 (120)
3005	40	*	19	*	14	*	15	*	19	*	67	*	148	3.4 (120)
3007	3	*	5	*	3	*	4	*	5	*	12	*	25	*
3201	142	76.1 (26)	76	*	94	*	49	*	67	*	282	80.9 (79)	613	3.1 (113)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
3301	1,210	71.1 (42)	617	29.7 (35)	807	17.5 (21)	340	17.6 (28)	522	10.3 (24)	2,227	91.4 (18)	4,835	0.9 (29)
3304	18	*	12	*	20	*	6	*	11	*	76	*	134	0.0 (1)
3305	52	*	36	*	37	*	20	*	34	*	134	94.8 (12)	262	0.4 (14)
3306	78	*	53	*	55	*	24	*	46	*	199	97.5 (5)	401	1.0 (34)
3308	40	*	8	*	10	*	4	*	7	*	68	*	136	0.0 (1)
3312	101	66.3 (61)	75	*	97	*	38	*	63	*	316	92.7 (14)	598	1.3 (48)
3550	22	*	7	*	5	*	9	*	7	*	36	*	72	*
3551	3	*	4	*	1	*	1	*	4	*	5	*	14	*
3552	2	*	1	*	1	*	0	*	0	*	4	*	9	*
3701	284	83.1 (8)	148	31.1 (26)	237	17.7 (18)	82	*	138	15.2 (7)	589	85.6 (54)	1,224	2.8 (103)
3850	894	67.8 (55)	787	35.5 (13)	764	12.2 (55)	456	24.3 (13)	649	12.6 (16)	2,145	75.3 (105)	4,107	2.5 (93)
3901	431	59.4 (80)	270	30.4 (28)	304	12.5 (50)	172	25.6 (8)	246	8.9 (33)	823	72.9 (113)	1,860	1.7 (71)
3902	20	*	13	*	15	*	8	*	12	*	20	*	71	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
3903	5	*	2	*	5	*	0	*	2	*	17	*	35	*
3950	627	65.9 (62)	400	29.8 (34)	443	15.3 (34)	230	24.8 (11)	352	8.5 (36)	1,328	76.9 (95)	2,881	1.5 (57)
3954	5	*	3	*	7	*	3	*	3	*	9	*	24	*
4106	616	74.2 (34)	362	23.2 (59)	492	12.0 (57)	191	13.1 (38)	314	6.4 (50)	1,512	86.7 (49)	2,972	0.6 (21)
4107	815	74.4 (32)	445	31.2 (25)	554	16.2 (27)	255	29.0 (4)	404	12.9 (15)	1,478	80.1 (83)	3,284	1.6 (65)
4109	228	81.6 (13)	145	26.9 (48)	190	13.2 (44)	105	24.8 (11)	138	9.4 (30)	588	85.5 (55)	1,173	0.6 (21)
4110	2	*	1	*	0	*	0	*	1	*	0	*	5	*
4112	152	23.0 (111)	125	13.6 (80)	131	7.6 (82)	82	*	124	1.6 (73)	363	98.1 (4)	689	1.2 (45)
4115	996	76.1 (26)	564	30.3 (30)	714	17.6 (19)	297	17.8 (27)	440	8.6 (34)	1,405	80.9 (79)	4,423	0.5 (17)
4351	162	82.7 (9)	80	*	142	22.5 (7)	41	*	71	*	373	96.8 (6)	739	1.5 (57)
4352	50	*	29	*	46	*	23	*	28	*	105	73.3 (110)	213	6.6 (159)
4353	138	77.5 (22)	70	*	102	17.6 (19)	47	*	58	*	374	87.7 (44)	659	3.0 (109)
4354	208	72.6 (38)	165	28.5 (41)	196	10.2 (67)	116	25.0 (10)	137	7.3 (45)	462	89.2 (32)	950	3.2 (114)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
4355	80	*	67	*	83	*	33	*	51	*	210	90.5 (24)	394	2.5 (93)
4356	122	71.3 (41)	73	*	102	16.7 (25)	50	*	63	*	290	81.4 (76)	555	5.4 (153)
4358	1	*	0	*	5	*	1	*	0	*	23	*	36	*
4359	169	72.2 (39)	102	29.4 (37)	154	17.5 (21)	45	*	90	*	405	88.1 (42)	792	1.0 (34)
4360	54	*	45	*	49	*	34	*	38	*	167	90.4 (25)	298	2.3 (86)
4361	75	*	49	*	57	*	32	*	43	*	173	80.3 (81)	342	2.6 (97)
4362	90	*	63	*	71	*	31	*	52	*	156	89.1 (33)	389	4.9 (147)
4363	232	77.6 (21)	187	42.8 (2)	192	15.6 (32)	123	31.7 (3)	163	16.0 (6)	515	88.7 (38)	1,023	5.3 (152)
4401	253	80.2 (16)	193	36.8 (9)	262	20.2 (10)	112	16.1 (32)	172	7.6 (42)	644	88.5 (40)	1,282	1.4 (52)
4501	249	74.3 (33)	159	34.0 (15)	208	13.9 (41)	97	*	137	5.1 (57)	629	85.9 (53)	1,197	2.2 (83)
4601	398	62.6 (70)	210	30.0 (33)	233	22.7 (6)	127	20.5 (20)	197	12.2 (17)	705	78.0 (90)	1,548	2.5 (93)
4701	312	80.8 (15)	176	30.1 (32)	225	16.0 (29)	114	21.9 (18)	158	10.1 (25)	771	83.7 (65)	1,542	1.4 (52)
5050	442	58.1 (82)	345	31.0 (27)	378	12.2 (55)	255	17.6 (28)	307	14.7 (9)	757	85.3 (58)	1,784	1.3 (48)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
5051	810	60.7 (75)	516	27.3 (44)	580	11.4 (60)	383	16.7 (30)	458	10.5 (23)	1,201	73.3 (110)	2,992	0.9 (29)
5052	845	57.8 (83)	569	28.6 (40)	613	11.7 (59)	437	12.1 (41)	505	7.3 (45)	1,350	75.2 (106)	3,330	0.9 (29)
5055	356	47.8 (99)	209	15.3 (76)	355	3.9 (97)	96	*	180	1.1 (75)	1,184	79.4 (84)	1,970	0.5 (17)
5101	481	68.6 (52)	248	27.0 (46)	360	14.7 (38)	147	20.4 (22)	222	11.3 (19)	942	87.3 (48)	2,003	1.0 (34)
5102	11	*	6	*	10	*	2	*	6	*	23	*	59	*
5202	203	84.7 (6)	101	31.7 (22)	163	17.2 (24)	52	*	69	*	496	95.4 (11)	956	1.0 (34)
5206	6	*	7	*	7	*	2	*	3	*	18	*	32	*
5601	262	76.3 (24)	153	37.9 (7)	173	23.1 (5)	102	35.3 (1)	147	17.7 (2)	560	90.0 (28)	1,196	1.5 (57)
5801	481	64.7 (64)	335	30.4 (28)	342	12.0 (57)	234	20.9 (19)	295	6.4 (50)	940	76.5 (97)	1,997	4.0 (137)
5803	28	*	33	*	34	*	19	*	31	*	110	75.5 (103)	200	1.0 (34)
5804	8	*	7	*	6	*	7	*	7	*	25	*	44	*
5806	73	*	67	*	49	*	41	*	54	*	38	*	181	4.4 (143)
6101	14	*	7	*	12	*	8	*	5	*	33	*	68	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
6102	30	*	27	*	38	*	19	*	21	*	94	*	177	2.8 (103)
6103	1,078	63.8 (66)	602	25.6 (51)	778	11.4 (60)	419	20.5 (20)	506	9.3 (31)	1,970	67.1 (120)	4,195	2.9 (106)
6105	5	*	4	*	4	*	3	*	3	*	21	*	35	*
6106	4	*	4	*	4	*	0	*	0	*	9	*	23	*
6301	1,015	67.0 (59)	605	26.0 (50)	582	9.6 (69)	484	15.7 (33)	546	8.1 (38)	1,718	74.7 (108)	3,833	1.0 (34)
6312	541	74.7 (30)	370	41.9 (4)	356	23.3 (4)	285	27.4 (6)	314	18.8 (1)	1,175	83.8 (64)	2,351	0.6 (21)
6316	325	76.0 (28)	229	34.1 (14)	244	12.7 (47)	153	20.3 (23)	201	13.4 (11)	716	81.7 (74)	1,468	0.3 (11)
6318	56	*	6	*	1	*	6	*	5	*	9	*	71	*
6320	90	*	39	*	25	*	38	*	34	*	150	78.0 (90)	315	0.0 (1)
6325	86	*	59	*	49	*	39	*	52	*	178	76.4 (98)	384	3.9 (136)
6335	265	70.6 (45)	213	19.7 (67)	198	11.1 (63)	146	10.3 (45)	197	4.1 (60)	774	86.6 (51)	1,499	0.2 (8)
6345	20	*	12	*	19	*	12	*	12	*	49	*	114	3.5 (123)
6360	272	66.9 (60)	179	31.3 (24)	139	15.1 (35)	115	26.1 (7)	160	11.3 (19)	591	71.1 (116)	1,228	1.7 (71)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
7306	608	61.2 (74)	407	22.6 (62)	409	8.3 (79)	355	11.5 (43)	353	6.5 (49)	1,247	77.8 (93)	2,707	3.0 (109)
7401	619	62.2 (73)	302	21.2 (66)	546	11.4 (60)	163	19.0 (26)	285	9.1 (32)	1,159	76.6 (96)	2,537	3.2 (114)
7601	156	46.8 (100)	73	*	119	18.5 (16)	30	*	68	*	331	84.6 (61)	695	1.7 (71)
7850	297	67.7 (56)	183	33.3 (16)	291	14.8 (37)	102	19.6 (24)	162	13.0 (14)	669	87.6 (45)	1,370	2.8 (103)
7852	8	*	4	*	4	*	1	*	3	*	17	*	29	*
8101	465	40.4 (107)	251	10.8 (81)	284	5.3 (92)	141	5.7 (55)	223	2.7 (69)	1,019	62.8 (124)	2,071	3.0 (109)
8102	110	43.6 (104)	73	*	95	*	46	*	64	*	276	64.1 (122)	540	1.9 (74)
8105	78	*	50	*	86	*	27	*	44	*	330	96.1 (8)	636	0.0 (1)
8106	146	56.8 (86)	65	*	111	4.5 (95)	59	*	65	*	304	41.1 (137)	610	1.6 (65)
8150	1	*	8	*	5	*	3	*	6	*	22	*	34	*
8151	906	73.5 (36)	615	38.5 (6)	667	19.3 (13)	469	25.4 (9)	525	16.2 (5)	1,850	96.0 (9)	3,900	0.7 (25)
8152	15	*	5	*	2	*	7	*	3	*	21	*	45	*
8153	3	*	1	*	1	*	0	*	1	*	9	*	13	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
8155	5	*	4	*	1	*	4	*	4	*	32	*	42	*
8157	1	*	3	*	2	*	2	*	3	*	5	*	10	*
8201	899	51.8 (95)	620	26.3 (49)	540	7.8 (80)	488	13.7 (37)	554	9.6 (27)	1,451	85.5 (55)	3,338	1.0 (34)
8202	1,207	44.3 (103)	771	22.7 (61)	908	7.6 (82)	573	6.8 (51)	653	3.2 (67)	2,967	79.1 (86)	5,560	1.2 (45)
8203	120	67.5 (57)	101	32.7 (20)	83	*	77	*	89	*	262	78.2 (89)	517	1.9 (74)
8250	206	53.4 (92)	155	25.2 (53)	110	6.4 (91)	111	5.4 (56)	133	8.3 (37)	325	76.0 (100)	746	1.3 (48)
8252	27	*	19	*	32	*	14	*	15	*	108	49.1 (134)	191	3.7 (127)
8255	640	55.8 (88)	574	24.0 (56)	460	8.9 (75)	418	11.7 (42)	487	7.0 (47)	1,182	75.7 (102)	2,668	1.3 (48)
8257	564	57.8 (83)	447	37.6 (8)	390	16.2 (27)	348	2.9 (61)	395	4.1 (60)	1,343	81.6 (75)	2,636	0.3 (11)
8263	192	56.8 (86)	153	29.4 (37)	134	9.0 (72)	118	15.3 (34)	135	9.6 (27)	385	73.2 (112)	806	0.4 (14)
8264	213	63.4 (68)	209	23.0 (60)	164	11.0 (64)	139	16.5 (31)	188	10.6 (22)	418	78.9 (87)	906	0.9 (29)
8267	190	54.7 (91)	154	18.8 (70)	150	8.7 (76)	132	9.8 (46)	140	6.4 (50)	415	85.3 (58)	875	0.8 (27)
8901	213	39.4 (108)	159	15.1 (77)	139	7.2 (88)	82	*	129	5.4 (56)	555	60.2 (126)	1,125	4.1 (139)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
8902	354	62.4 (71)	244	25.4 (52)	228	7.5 (85)	151	10.6 (44)	203	7.4 (43)	793	55.9 (128)	1,678	2.9 (106)
8903	390	50.3 (97)	231	17.7 (72)	271	6.6 (89)	151	6.0 (53)	206	3.9 (63)	784	46.7 (136)	1,669	4.2 (141)
8904	532	54.9 (89)	395	17.7 (72)	370	8.6 (77)	207	8.2 (50)	351	4.3 (59)	1,194	53.3 (131)	2,584	3.0 (109)
8905	452	62.4 (71)	296	25.0 (54)	259	8.5 (78)	164	9.1 (49)	244	6.1 (53)	934	66.1 (121)	1,970	2.0 (78)
8909	202	63.4 (68)	120	25.0 (54)	141	7.8 (80)	84	*	104	7.7 (40)	416	50.2 (133)	856	3.6 (124)
9045	471	46.5 (101)	264	18.9 (69)	268	9.0 (72)	171	6.4 (52)	234	3.4 (65)	1,129	69.9 (118)	2,207	1.4 (52)
9050	295	48.5 (98)	215	23.7 (58)	224	4.5 (95)	168	5.4 (56)	177	2.3 (70)	535	69.3 (119)	1,303	2.2 (83)
9051	471	68.4 (53)	324	21.3 (65)	318	9.7 (68)	218	4.1 (59)	282	3.9 (63)	906	61.8 (125)	1,892	0.7 (25)
9053	363	59.8 (79)	158	29.7 (35)	145	12.4 (51)	104	12.5 (40)	143	7.7 (40)	410	50.7 (132)	1,088	3.4 (120)
9055	273	42.5 (105)	9	*	4	*	11	*	8	*	10	*	317	3.2 (114)
9058	297	58.2 (81)	175	18.3 (71)	220	10.5 (66)	144	9.7 (47)	149	4.7 (58)	667	82.8 (71)	1,293	1.9 (74)
9066	359	52.9 (93)	253	19.0 (68)	305	7.5 (85)	130	3.8 (60)	191	2.1 (72)	696	47.8 (135)	1,456	3.6 (124)
9068	299	40.5 (106)	146	13.7 (79)	171	4.7 (94)	125	9.6 (48)	132	1.5 (74)	556	70.9 (117)	1,161	4.9 (147)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
9069	732	33.2 (109)	499	14.6 (78)	556	5.0 (93)	368	6.0 (53)	456	2.9 (68)	1,662	55.2 (130)	3,268	3.6 (124)
9070	376	52.9 (93)	277	23.8 (57)	252	9.1 (71)	224	12.9 (39)	252	4.0 (62)	797	57.3 (127)	1,692	4.0 (137)
9072	342	51.8 (95)	169	16.0 (75)	291	7.6 (82)	99	*	144	5.6 (54)	785	55.4 (129)	1,556	3.7 (127)
9080	763	45.6 (102)	445	17.5 (74)	645	9.3 (70)	311	4.8 (58)	383	3.4 (65)	1,681	63.8 (123)	3,384	3.7 (127)
9082	34	*	21	*	24	*	21	*	20	*	44	*	122	5.7 (157)
9083	350	68.9 (50)	7	*	1	*	10	*	6	*	0	*	352	0.3 (11)
9084	323	32.2 (110)	6	*	9	*	8	*	5	*	34	*	371	3.2 (114)
9087	0	*	1	*	1	*	0	*	1	*	28	*	35	*
Michigan	44,718	63.9	27,565	28.2	31,867	13.2	17,815	17.6	23,867	9.3	91,478	79.0	192,304	2.0
National (Prior Year)	168,373	71.4	88,368	31.4	91,841	23.5	60,230	17.4	71,191	8.1	285,831	82.0	655,593	3.7

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* National includes previous year's data for all participating States and ITOs

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0101	98	*	*	365	7.9 (112)	2.2 (39)	11.5 (115)	171	17.0 (98)	15.8 (119)	254	17.7 (107)
0102	347	7.8 (25)	6.3 (61)	1,248	6.2 (49)	2.7 (71)	10.7 (88)	559	18.8 (132)	14.5 (93)	814	18.9 (115)
0103	339	7.7 (22)	11.2 (145)	1,323	6.8 (66)	2.0 (28)	12.6 (141)	600	17.0 (98)	13.3 (73)	877	12.7 (67)
0104	120	7.5 (19)	6.7 (68)	492	8.9 (157)	2.8 (81)	10.4 (73)	226	18.6 (129)	13.7 (81)	328	11.0 (56)
0105	91	*	*	243	14.4 (190)	4.9 (169)	9.5 (46)	66	*	*	144	20.8 (130)
0250	122	7.4 (17)	11.5 (148)	431	3.2 (4)	4.9 (169)	6.3 (6)	160	20.0 (145)	10.6 (20)	274	10.2 (44)
0251	81	*	*	311	4.8 (12)	2.6 (65)	13.8 (171)	125	18.4 (126)	21.6 (159)	197	16.2 (92)
0252	107	5.6 (3)	14.0 (152)	484	5.0 (15)	4.1 (149)	9.9 (58)	244	15.6 (52)	13.1 (64)	336	4.5 (8)
0253	126	13.5 (125)	5.6 (43)	439	8.0 (121)	4.1 (149)	11.4 (109)	167	19.2 (142)	18.6 (153)	269	6.7 (15)
0401	419	7.6 (21)	10.7 (142)	1,631	8.6 (147)	2.3 (51)	11.6 (119)	698	18.9 (136)	15.5 (112)	1,055	18.5 (111)
0402	338	8.9 (51)	8.3 (105)	1,136	5.5 (21)	3.6 (132)	11.4 (109)	478	20.1 (146)	16.1 (122)	673	20.7 (128)
0403	138	14.5 (134)	7.2 (79)	443	9.7 (170)	3.4 (123)	11.7 (123)	181	12.2 (6)	18.8 (154)	255	17.3 (103)
0404	101	6.9 (12)	11.9 (149)	367	5.7 (31)	3.3 (116)	11.4 (109)	156	14.1 (21)	17.3 (140)	245	20.0 (123)
0501	382	9.9 (78)	6.8 (72)	1,677	5.7 (31)	6.6 (185)	12.6 (141)	738	19.0 (138)	16.9 (135)	941	15.5 (84)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0502	0	*	*	1	*	*	*	1	*	*	1	*
0503	97	*	*	363	6.1 (42)	7.2 (187)	8.5 (22)	150	16.7 (85)	10.7 (22)	196	8.2 (26)
0504	407	9.3 (60)	7.1 (78)	1,574	8.4 (139)	4.1 (149)	9.0 (32)	680	16.9 (93)	10.7 (22)	949	8.4 (29)
0505	244	10.2 (83)	4.9 (31)	854	8.3 (134)	4.7 (164)	10.8 (93)	318	20.1 (146)	17.6 (144)	492	8.5 (31)
0506	143	7.7 (22)	6.3 (61)	671	7.0 (82)	5.5 (173)	11.6 (119)	338	17.5 (111)	15.4 (109)	420	10.7 (51)
0507	75	*	*	569	7.9 (112)	1.9 (24)	13.5 (166)	289	14.9 (34)	18.0 (149)	389	6.7 (15)
0508	24	*	*	198	11.1 (181)	4.0 (146)	18.7 (189)	107	16.8 (91)	21.5 (158)	135	5.2 (12)
0509	3	*	*	39	*	*	*	22	*	*	28	*
0650	949	9.5 (65)	8.1 (101)	3,324	7.4 (96)	3.3 (116)	11.8 (127)	1,321	17.0 (98)	18.3 (151)	1,929	10.6 (47)
0651	535	6.0 (5)	6.7 (68)	2,051	6.8 (66)	3.9 (140)	9.0 (32)	906	20.5 (152)	13.6 (79)	1,279	9.7 (40)
0652	289	9.0 (53)	9.0 (125)	1,068	7.4 (96)	2.1 (34)	10.8 (93)	486	15.8 (56)	15.2 (104)	664	11.4 (59)
0653	380	9.5 (65)	8.7 (117)	1,645	8.3 (134)	2.4 (54)	11.9 (129)	778	17.0 (98)	15.4 (109)	1,085	10.8 (52)
0654	312	9.0 (53)	9.3 (128)	1,159	7.5 (102)	3.0 (92)	10.4 (73)	509	15.1 (39)	15.5 (112)	725	15.6 (86)
0655	283	4.2 (1)	6.4 (64)	1,048	5.6 (24)	3.1 (104)	9.1 (37)	444	14.0 (19)	13.1 (64)	640	13.1 (70)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
0657	57	*	*	248	6.9 (73)	6.0 (176)	8.1 (17)	108	21.3 (159)	9.3 (8)	167	16.2 (92)
0701	324	7.1 (15)	9.9 (135)	1,216	8.9 (157)	1.8 (20)	8.9 (30)	577	16.6 (82)	9.5 (9)	835	8.5 (31)
0702	68	*	*	226	7.5 (102)	1.8 (20)	12.4 (138)	114	24.6 (164)	12.3 (49)	129	7.0 (17)
0703	159	9.4 (61)	9.4 (129)	667	4.5 (10)	3.0 (92)	13.3 (162)	314	15.9 (60)	18.8 (154)	434	7.4 (19)
0704	43	*	*	157	5.1 (17)	3.2 (114)	12.1 (132)	69	*	*	107	5.6 (13)
0705	82	*	*	350	8.3 (134)	1.7 (16)	12.6 (141)	166	15.1 (39)	16.3 (128)	221	7.2 (18)
0706	53	*	*	111	11.7 (184)	2.7 (71)	9.0 (32)	24	*	*	50	*
0850	690	9.0 (53)	8.7 (117)	2,903	8.1 (126)	2.9 (87)	12.3 (135)	1,249	22.3 (161)	16.9 (135)	1,660	8.1 (24)
0851	622	9.5 (65)	9.5 (131)	2,564	6.6 (62)	2.8 (81)	9.4 (43)	1,090	17.2 (104)	12.4 (51)	1,479	7.8 (23)
0901	1,389	10.2 (83)	8.3 (105)	5,189	7.9 (112)	1.6 (15)	12.6 (141)	2,272	18.2 (123)	17.5 (141)	3,496	11.3 (58)
0902	37	*	*	203	7.4 (96)	2.0 (28)	7.9 (13)	101	19.8 (144)	10.9 (25)	136	11.0 (56)
1051	1,105	9.1 (58)	11.4 (147)	3,702	4.5 (10)	6.1 (179)	9.7 (52)	1,599	20.5 (152)	12.4 (51)	2,194	18.6 (112)
1052	8	*	*	4	*	*	*	0	*	*	0	*
1101	1,590	10.1 (80)	5.4 (38)	5,930	6.9 (73)	4.2 (153)	10.2 (68)	2,498	16.2 (68)	13.2 (66)	3,710	15.9 (88)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under- Weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
1102	710	7.5 (19)	6.1 (56)	2,610	6.9 (73)	3.1 (104)	11.4 (109)	1,100	17.6 (117)	15.8 (119)	1,636	10.3 (45)
1104	3	*	*	3	*	*	*	0	*	*	0	*
1106	70	*	*	278	6.8 (66)	4.0 (146)	9.4 (43)	116	20.7 (155)	11.2 (30)	189	9.0 (36)
1250	932	8.3 (36)	7.9 (92)	3,077	8.7 (150)	3.9 (140)	11.2 (102)	1,240	16.0 (62)	13.4 (74)	1,714	10.3 (45)
1251	660	8.2 (32)	7.3 (81)	2,389	8.5 (144)	1.5 (11)	12.3 (135)	1,024	16.3 (70)	15.6 (115)	1,477	16.5 (95)
1252	521	8.6 (41)	6.7 (68)	1,921	7.4 (96)	2.7 (71)	11.5 (115)	810	18.8 (132)	15.7 (116)	1,207	13.7 (75)
1253	685	7.0 (13)	8.3 (105)	2,504	9.1 (161)	2.4 (54)	13.3 (162)	1,090	16.3 (70)	15.7 (116)	1,552	10.6 (47)
1350	2,070	10.1 (80)	7.0 (77)	7,580	8.0 (121)	3.9 (140)	12.6 (141)	3,140	17.0 (98)	16.1 (122)	4,628	20.1 (124)
1351	97	*	*	443	9.3 (165)	2.7 (71)	15.6 (182)	199	20.1 (146)	16.1 (122)	294	17.0 (100)
1352	358	13.7 (132)	8.1 (101)	1,393	7.3 (92)	4.8 (168)	14.3 (175)	627	16.1 (66)	17.5 (141)	862	17.4 (104)
1401	212	8.0 (27)	7.5 (83)	732	7.9 (112)	3.4 (123)	11.2 (102)	298	18.1 (122)	14.8 (100)	430	14.0 (76)
1402	113	9.7 (72)	8.0 (97)	504	5.6 (24)	3.8 (138)	12.3 (135)	223	21.1 (158)	17.9 (148)	317	7.6 (22)
1403	18	*	*	125	4.0 (7)	0.8 (2)	17.6 (188)	66	*	*	83	*
1501	708	7.8 (25)	9.7 (133)	2,858	5.6 (24)	3.7 (136)	13.6 (167)	1,269	16.6 (82)	17.7 (146)	1,809	11.4 (59)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature		Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
1503	855	8.0 (27)	8.4 (111)	3,610	5.6 (24)	2.4 (54)	12.1 (132)	1,640	16.5 (78)	16.9 (135)	2,287	19.2 (117)
1504	1,674	7.0 (13)	6.9 (75)	6,524	4.9 (13)	3.0 (92)	11.2 (102)	2,881	16.7 (85)	14.6 (95)	4,236	36.0 (169)
1505	379	11.1 (97)	11.1 (144)	1,556	7.3 (92)	2.6 (65)	9.8 (55)	763	14.0 (19)	11.1 (27)	1,003	27.9 (156)
1506	522	8.6 (41)	8.4 (111)	2,299	6.1 (42)	2.0 (28)	14.4 (176)	1,047	15.8 (56)	16.4 (131)	1,428	18.7 (113)
1507	525	12.0 (109)	12.0 (150)	2,053	5.6 (24)	3.6 (132)	7.8 (12)	890	14.2 (25)	9.7 (11)	1,357	36.0 (169)
1508	30	*	*	152	7.2 (90)	0.7 (1)	25.0 (191)	64	*	*	102	40.2 (176)
1509	353	7.4 (17)	10.2 (137)	1,555	6.1 (42)	3.6 (132)	12.9 (152)	727	17.5 (111)	16.5 (134)	1,058	19.0 (116)
1511	598	11.0 (96)	7.5 (83)	2,449	5.4 (19)	3.1 (104)	10.1 (63)	1,031	15.4 (46)	13.5 (75)	1,547	22.4 (136)
1512	318	8.8 (48)	8.5 (116)	1,449	8.1 (126)	3.3 (116)	11.7 (123)	665	14.9 (34)	12.8 (56)	913	27.4 (155)
1513	0	*	*	34	*	*	*	26	*	*	26	*
1514	247	13.4 (123)	10.1 (136)	1,019	6.9 (73)	3.1 (104)	9.6 (50)	480	16.3 (70)	10.0 (13)	649	32.8 (166)
1701	538	6.3 (7)	12.1 (151)	2,051	7.3 (92)	4.0 (146)	15.2 (181)	937	16.4 (74)	17.5 (141)	1,159	21.9 (133)
1950	530	7.7 (22)	6.8 (72)	1,830	8.6 (147)	2.6 (65)	10.5 (80)	739	14.9 (34)	13.7 (81)	1,153	4.2 (7)
1952	121	6.6 (9)	8.3 (105)	435	10.3 (178)	1.1 (6)	10.8 (93)	168	21.4 (160)	11.3 (32)	266	3.8 (5)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
1953	267	8.6 (41)	6.0 (52)	1,092	9.8 (172)	2.8 (81)	11.4 (109)	454	13.9 (18)	13.2 (66)	694	4.0 (6)
1954	418	8.4 (37)	8.4 (111)	1,585	7.5 (102)	3.0 (92)	11.3 (107)	652	16.4 (74)	16.1 (122)	982	3.0 (4)
1955	110	10.0 (79)	5.5 (39)	465	6.7 (65)	3.4 (123)	14.4 (176)	207	18.8 (132)	22.7 (162)	305	2.6 (2)
1957	385	10.6 (91)	10.4 (139)	1,530	9.5 (168)	2.2 (39)	13.0 (154)	642	20.2 (150)	14.2 (89)	946	2.5 (1)
1963	70	*	*	277	6.1 (42)	2.2 (39)	13.0 (154)	127	17.3 (106)	22.0 (160)	178	2.8 (3)
1964	23	*	*	172	11.6 (182)	1.2 (7)	15.1 (180)	69	*	*	132	4.5 (8)
2150	576	9.5 (65)	7.5 (83)	2,114	10.0 (173)	1.7 (16)	11.2 (102)	925	17.3 (106)	13.0 (61)	1,334	10.0 (43)
2151	184	8.7 (46)	8.7 (117)	663	7.4 (96)	3.0 (92)	12.8 (149)	246	15.4 (46)	16.3 (128)	402	8.7 (33)
2152	9	*	*	17	*	*	*	9	*	*	8	*
2251	362	9.7 (72)	8.3 (105)	1,398	6.0 (38)	3.1 (104)	10.5 (80)	623	15.6 (52)	14.8 (100)	861	19.2 (117)
2252	162	11.7 (107)	10.5 (140)	618	6.3 (51)	3.2 (114)	13.6 (167)	290	21.0 (157)	19.3 (157)	407	10.8 (52)
2554	50	*	*	332	6.3 (51)	1.5 (11)	9.6 (50)	196	18.9 (136)	10.2 (15)	235	10.6 (47)
2555	6,635	12.6 (115)	5.6 (43)	21,319	9.0 (160)	2.7 (71)	11.0 (100)	8,840	15.8 (56)	11.9 (38)	13,265	21.5 (131)
2556	516	19.2 (152)	2.3 (1)	2,030	12.3 (186)	2.7 (71)	11.9 (129)	858	12.1 (4)	13.5 (75)	1,201	30.1 (160)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
2557	6	*	*	37	*	*	*	25	*	*	27	*
2558	3	*	*	37	*	*	*	30	*	*	26	*
2559	1	*	*	31	*	*	*	28	*	*	26	*
2560	2	*	*	46	*	*	*	35	*	*	35	*
2561	1	*	*	10	*	*	*	9	*	*	8	*
2562	11	*	*	127	6.3 (51)	3.1 (104)	15.0 (179)	81	*	*	94	*
2563	40	*	*	119	16.0 (191)	1.7 (16)	16.0 (184)	45	*	*	73	*
2564	15	*	*	20	*	*	*	4	*	*	6	*
2565	0	*	*	3	*	*	*	3	*	*	3	*
3001	1,063	13.5 (125)	3.7 (16)	3,725	6.1 (42)	4.6 (159)	10.3 (70)	1,541	16.0 (62)	12.1 (43)	2,279	11.9 (61)
3002	276	10.5 (89)	5.8 (47)	1,063	5.8 (36)	3.8 (138)	11.1 (101)	482	16.6 (82)	14.1 (87)	648	9.1 (37)
3004	171	13.5 (125)	5.8 (47)	558	4.3 (9)	4.7 (164)	13.3 (162)	227	18.5 (128)	23.3 (164)	331	4.5 (8)
3005	57	*	*	134	9.7 (170)	7.5 (188)	7.5 (9)	51	*	*	62	*
3007	10	*	*	57	*	*	*	26	*	*	44	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
3009	1	*	*	4	*	*	*	2	*	*	4	*
3201	388	10.3 (86)	11.3 (146)	1,501	9.1 (161)	3.1 (104)	13.0 (154)	636	15.3 (44)	13.5 (75)	994	8.7 (33)
3301	3,693	11.3 (99)	5.9 (50)	11,769	8.7 (150)	4.1 (149)	10.7 (88)	4,507	17.7 (119)	14.4 (91)	6,770	13.1 (70)
3304	91	*	*	474	7.0 (82)	4.6 (159)	6.1 (5)	234	15.0 (38)	6.0 (1)	302	15.2 (83)
3305	150	8.7 (46)	8.7 (117)	724	7.9 (112)	2.2 (39)	9.1 (37)	339	16.5 (78)	8.6 (5)	477	9.6 (39)
3306	248	4.8 (2)	8.9 (123)	1,063	8.5 (144)	2.1 (34)	9.5 (46)	472	16.7 (85)	10.6 (20)	701	9.8 (42)
3308	79	*	*	304	7.9 (112)	3.0 (92)	7.6 (11)	128	11.7 (2)	6.3 (2)	191	13.1 (70)
3309	14	*	*	74	*	*	*	37	*	*	55	*
3312	368	9.8 (74)	7.9 (92)	1,579	7.8 (108)	3.5 (128)	10.6 (85)	767	15.1 (39)	13.7 (81)	983	12.6 (66)
3314	2	*	*	2	*	*	*	0	*	*	0	*
3550	55	*	*	212	6.6 (62)	1.9 (24)	16.5 (185)	92	*	*	144	17.4 (104)
3551	20	*	*	63	*	*	*	28	*	*	34	*
3552	27	*	*	148	6.1 (42)	2.0 (28)	20.9 (190)	74	*	*	101	16.8 (99)
3701	892	8.4 (37)	8.1 (101)	3,549	5.6 (24)	4.6 (159)	10.7 (88)	1,522	19.1 (141)	13.6 (79)	2,293	8.2 (26)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low		High	Short Stature		Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
3850	2,728	10.7 (92)	7.6 (87)	10,151	5.7 (31)	8.0 (189)	10.0 (61)	4,260	18.2 (123)	15.8 (119)	5,788	18.8 (114)
3901	1,360	11.3 (99)	5.3 (37)	4,686	5.5 (21)	4.7 (164)	10.3 (70)	1,842	16.4 (74)	13.2 (66)	2,871	30.5 (161)
3902	187	10.2 (83)	7.5 (83)	951	6.0 (38)	3.0 (92)	13.1 (158)	464	17.5 (111)	17.2 (139)	601	27.0 (154)
3903	32	*	*	129	7.0 (82)	4.7 (164)	13.2 (160)	56	*	*	91	*
3904	6	*	*	45	*	*	*	23	*	*	33	*
3905	11	*	*	55	*	*	*	22	*	*	36	*
3950	1,945	11.4 (102)	8.4 (111)	7,011	6.3 (51)	4.2 (153)	10.8 (93)	2,873	17.2 (104)	13.9 (86)	4,430	32.4 (165)
3954	31	*	*	103	4.9 (13)	1.0 (5)	13.6 (167)	42	*	*	70	*
4103	8	*	*	52	*	*	*	24	*	*	11	*
4106	2,015	12.9 (119)	5.1 (34)	7,936	10.3 (178)	1.8 (20)	13.2 (160)	3,594	17.6 (117)	15.7 (116)	5,065	12.9 (69)
4107	2,508	13.7 (132)	6.6 (67)	9,064	10.0 (173)	2.7 (71)	10.4 (73)	3,689	15.5 (48)	12.1 (43)	5,681	17.9 (109)
4109	884	9.8 (74)	6.9 (75)	3,621	10.1 (176)	2.5 (62)	9.0 (32)	1,565	16.7 (85)	10.4 (17)	2,300	7.4 (19)
4110	8	*	*	1	*	*	*	0	*	*	0	*
4112	479	6.7 (10)	6.5 (66)	2,024	5.6 (24)	3.9 (140)	12.4 (138)	899	16.8 (91)	17.1 (138)	1,107	7.4 (19)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low		High	Short Stature		Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
4113	16	*	*	331	4.2 (8)	0.9 (3)	13.9 (173)	189	17.5 (111)	16.4 (131)	178	5.1 (11)
4115	3,140	13.6 (129)	6.0 (52)	11,147	9.2 (163)	2.4 (54)	11.4 (109)	4,825	14.9 (34)	13.2 (66)	7,182	16.6 (96)
4351	476	9.2 (59)	6.1 (56)	1,849	6.9 (73)	2.2 (39)	10.7 (88)	817	15.3 (44)	12.1 (43)	1,161	14.8 (79)
4352	138	9.4 (61)	8.0 (97)	577	6.8 (66)	2.9 (87)	8.7 (25)	266	16.9 (93)	11.3 (32)	376	14.9 (80)
4353	361	8.6 (41)	5.5 (39)	1,394	7.0 (82)	2.3 (51)	12.8 (149)	684	17.3 (106)	17.8 (147)	912	10.6 (47)
4354	667	8.8 (48)	7.9 (92)	2,390	8.5 (144)	2.4 (54)	12.1 (132)	1,023	18.6 (129)	15.2 (104)	1,489	9.7 (40)
4355	275	9.8 (74)	9.5 (131)	1,042	7.4 (96)	2.6 (65)	11.2 (102)	455	16.9 (93)	15.4 (109)	682	16.1 (91)
4356	384	9.6 (69)	6.3 (61)	1,464	6.9 (73)	2.7 (71)	9.8 (55)	679	15.5 (48)	12.2 (46)	935	13.3 (73)
4358	80	*	*	448	7.8 (108)	1.3 (10)	13.4 (165)	228	22.4 (162)	18.0 (149)	337	20.2 (125)
4359	498	6.4 (8)	9.4 (129)	1,761	7.3 (92)	1.8 (20)	12.7 (147)	811	20.5 (152)	17.6 (144)	1,091	14.5 (78)
4360	185	5.9 (4)	9.7 (133)	694	6.1 (42)	1.2 (7)	11.8 (127)	323	16.7 (85)	14.6 (95)	464	11.9 (61)
4361	217	8.8 (48)	10.6 (141)	774	7.0 (82)	2.2 (39)	13.0 (154)	338	15.7 (55)	18.3 (151)	546	19.6 (121)
4362	267	10.1 (80)	7.9 (92)	944	8.4 (139)	2.2 (39)	10.4 (73)	440	17.5 (111)	13.0 (61)	641	16.4 (94)
4363	716	8.2 (32)	8.8 (121)	2,554	8.0 (121)	1.5 (11)	11.7 (123)	1,160	18.8 (132)	12.8 (56)	1,705	16.0 (89)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- Weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
4401	901	9.8 (74)	8.4 (111)	3,544	6.3 (51)	2.3 (51)	9.3 (41)	1,534	16.7 (85)	14.5 (93)	2,249	14.9 (80)
4501	855	8.4 (37)	8.9 (123)	3,319	6.9 (73)	3.3 (116)	9.9 (58)	1,431	17.3 (106)	14.4 (91)	2,008	8.9 (35)
4601	1,194	6.7 (10)	7.7 (90)	4,018	7.9 (112)	2.5 (62)	12.5 (140)	1,711	18.4 (126)	16.2 (126)	2,415	26.7 (153)
4602	7	*	*	47	*	*	*	26	*	*	33	*
4701	993	12.4 (112)	7.9 (92)	3,557	8.7 (150)	2.2 (39)	8.7 (25)	1,577	16.9 (93)	12.0 (40)	2,363	8.3 (28)
5050	1,439	12.4 (112)	5.9 (50)	4,354	7.8 (108)	2.2 (39)	11.7 (123)	1,626	15.2 (43)	11.2 (30)	2,527	15.6 (86)
5051	2,398	11.2 (98)	5.2 (36)	7,248	8.1 (126)	3.1 (104)	10.1 (63)	2,607	14.7 (29)	12.5 (53)	3,935	15.0 (82)
5052	2,543	12.2 (111)	4.9 (31)	7,798	8.1 (126)	3.0 (92)	10.5 (80)	2,948	14.1 (21)	12.9 (59)	4,533	12.2 (64)
5055	1,028	8.1 (30)	4.1 (22)	4,541	3.6 (5)	2.9 (87)	12.8 (149)	2,354	14.8 (31)	15.3 (107)	3,188	22.5 (137)
5101	1,498	10.5 (89)	6.2 (58)	5,347	6.9 (73)	2.6 (65)	10.8 (93)	2,189	16.3 (70)	14.6 (95)	3,315	17.0 (100)
5102	23	*	*	76	*	*	*	32	*	*	43	*
5202	673	8.6 (41)	8.2 (104)	2,591	10.0 (173)	2.9 (87)	13.6 (167)	1,131	20.1 (146)	15.3 (107)	1,284	8.4 (29)
5206	6	*	*	21	*	*	*	6	*	*	3	*
5601	871	8.2 (32)	10.2 (137)	3,393	9.3 (165)	1.9 (24)	14.0 (174)	1,350	19.0 (138)	14.7 (98)	1,942	16.0 (89)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ^(3,4)

Clinic	Birth Weight ⁽⁵⁾			Height and Weight ^(6,7)						Anemia Low Hb/Hct ⁽⁸⁾		
	Low	High		Short Stature	Under- Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
5801	1,379	9.4 (61)	6.7 (68)	4,798	7.9 (112)	2.6 (65)	10.6 (85)	2,022	17.8 (121)	15.0 (103)	2,980	17.2 (102)
5803	112	7.1 (15)	10.7 (142)	590	5.4 (19)	1.7 (16)	10.8 (93)	287	12.5 (7)	13.2 (66)	406	8.1 (24)
5804	14	*	*	97	*	*	*	52	*	*	73	*
5806	244	11.5 (105)	4.5 (25)	421	11.6 (182)	6.2 (181)	7.4 (8)	69	*	*	148	17.6 (106)
6101	38	*	*	162	12.3 (186)	2.5 (62)	17.3 (187)	76	*	*	111	20.7 (128)
6102	113	8.0 (27)	6.2 (58)	549	7.1 (89)	2.4 (54)	11.5 (115)	268	19.0 (138)	12.3 (49)	361	14.4 (77)
6103	3,371	11.5 (105)	5.1 (34)	11,351	8.2 (131)	2.7 (71)	12.6 (141)	4,690	17.0 (98)	14.3 (90)	7,054	22.6 (138)
6105	19	*	*	116	12.9 (188)	0.9 (3)	6.9 (7)	61	*	*	86	*
6106	112	17.9 (149)	3.6 (14)	249	8.4 (139)	6.4 (184)	10.4 (73)	73	*	*	125	24.0 (147)
6301	3,281	13.3 (121)	6.0 (52)	11,127	9.4 (167)	2.0 (28)	10.4 (73)	4,412	16.2 (68)	13.0 (61)	6,744	20.5 (126)
6312	1,659	11.9 (108)	5.8 (47)	6,616	8.3 (134)	2.8 (81)	8.0 (15)	2,893	14.8 (31)	10.2 (15)	4,239	22.1 (134)
6316	1,068	10.7 (92)	7.2 (79)	3,926	6.3 (51)	3.0 (92)	10.0 (61)	1,642	15.5 (48)	12.8 (56)	2,432	16.6 (96)
6318	65	*	*	56	*	*	*	3	*	*	3	*
6320	167	11.4 (102)	4.8 (30)	548	6.4 (57)	5.1 (171)	8.6 (23)	244	14.8 (31)	7.8 (4)	319	19.7 (122)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
6325	216	6.0 (5)	8.8 (121)	817	5.5 (21)	2.1 (34)	10.3 (70)	370	16.5 (78)	11.1 (27)	494	23.1 (141)
6335	905	10.7 (92)	4.6 (26)	3,745	6.5 (59)	4.6 (159)	10.4 (73)	1,594	14.7 (29)	12.7 (54)	2,350	24.8 (149)
6340	23	*	*	133	5.3 (18)	3.0 (92)	11.3 (107)	66	*	*	90	*
6345	58	*	*	171	8.2 (131)	3.5 (128)	12.9 (152)	58	*	*	79	*
6360	749	13.4 (123)	5.5 (39)	2,946	8.0 (121)	4.2 (153)	10.8 (93)	1,336	15.9 (60)	14.1 (87)	1,808	29.5 (159)
7304	28	*	*	159	5.7 (31)	1.9 (24)	8.2 (20)	69	*	*	117	12.0 (63)
7306	2,113	12.0 (109)	5.6 (43)	7,143	5.7 (31)	4.4 (158)	10.5 (80)	2,687	17.4 (110)	13.7 (81)	4,256	19.2 (117)
7308	30	*	*	167	3.0 (2)	3.0 (92)	8.4 (21)	72	*	*	118	13.6 (74)
7310	11	*	*	48	*	*	*	18	*	*	30	*
7401	1,981	8.4 (37)	8.0 (97)	6,928	7.6 (107)	3.0 (92)	10.7 (88)	2,791	19.7 (143)	15.2 (104)	4,137	10.9 (55)
7601	519	8.1 (30)	8.3 (105)	1,822	11.9 (185)	1.2 (7)	15.8 (183)	797	20.7 (155)	22.6 (161)	1,179	16.6 (96)
7850	967	9.4 (61)	7.8 (91)	3,646	6.0 (38)	3.5 (128)	10.5 (80)	1,624	18.3 (125)	15.5 (112)	2,384	10.8 (52)
7852	20	*	*	98	*	*	*	64	*	*	71	*
8101	1,552	18.6 (150)	2.4 (2)	5,624	9.2 (163)	6.3 (182)	10.2 (68)	2,183	14.4 (27)	12.0 (40)	3,399	42.3 (178)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature	Under- Weight	Obese		≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
8102	328	10.4 (87)	9.1 (127)	1,596	6.0 (38)	5.6 (174)	17.0 (186)	839	16.9 (93)	22.9 (163)	1,092	31.0 (162)
8105	956	12.6 (115)	4.9 (31)	4,281	6.9 (73)	2.9 (87)	11.6 (119)	1,833	16.1 (66)	14.9 (102)	2,755	28.0 (157)
8106	420	15.0 (138)	5.5 (39)	1,344	8.8 (154)	3.1 (104)	12.0 (131)	587	16.0 (62)	16.2 (126)	850	22.1 (134)
8150	17	*	*	167	3.0 (2)	2.4 (54)	6.0 (4)	106	22.6 (163)	7.5 (3)	124	17.7 (107)
8151	2,706	12.8 (118)	4.7 (28)	9,231	6.8 (66)	3.4 (123)	8.7 (25)	3,867	16.4 (74)	12.2 (46)	5,937	23.3 (143)
8152	37	*	*	152	3.9 (6)	4.6 (159)	9.9 (58)	76	*	*	117	24.8 (149)
8153	10	*	*	47	*	*	*	21	*	*	36	*
8155	13	*	*	148	6.8 (66)	6.1 (179)	11.5 (115)	97	*	*	118	33.9 (168)
8157	5	*	*	47	*	*	*	32	*	*	36	*
8201	2,345	12.5 (114)	6.2 (58)	6,534	7.9 (112)	3.3 (116)	10.6 (85)	2,494	15.8 (56)	13.2 (66)	3,606	24.4 (148)
8202	3,288	9.0 (53)	5.6 (43)	12,370	5.0 (15)	3.3 (116)	8.1 (17)	5,727	13.8 (17)	9.6 (10)	8,278	23.0 (140)
8203	343	13.1 (120)	7.3 (81)	1,260	5.8 (36)	2.1 (34)	8.8 (28)	549	18.6 (129)	11.7 (36)	786	12.2 (64)
8250	611	12.6 (115)	4.6 (26)	1,740	8.8 (154)	3.1 (104)	9.1 (37)	630	14.3 (26)	12.9 (59)	911	19.5 (120)
8252	115	19.1 (151)	2.6 (4)	421	10.2 (177)	2.1 (34)	9.7 (52)	177	17.5 (111)	9.0 (7)	261	21.8 (132)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature		Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
8255	2,132	11.4 (102)	6.8 (72)	6,607	8.6 (147)	2.7 (71)	10.1 (63)	2,430	14.6 (28)	13.2 (66)	3,616	20.6 (127)
8257	1,577	13.6 (129)	2.9 (6)	6,165	7.5 (102)	3.9 (140)	11.6 (119)	2,651	12.0 (3)	13.8 (85)	3,945	23.8 (144)
8263	592	9.6 (69)	7.6 (87)	2,085	7.2 (90)	2.2 (39)	9.0 (32)	897	20.2 (150)	10.5 (19)	1,212	6.1 (14)
8264	611	8.2 (32)	8.0 (97)	2,023	7.8 (108)	2.2 (39)	10.1 (63)	839	15.1 (39)	11.1 (27)	1,198	9.2 (38)
8267	657	9.6 (69)	9.0 (125)	2,028	8.9 (157)	1.5 (11)	13.1 (158)	687	13.0 (10)	14.7 (98)	1,184	12.8 (68)
8901	776	16.8 (147)	2.7 (5)	3,257	8.4 (139)	6.0 (176)	9.5 (46)	1,325	14.1 (21)	11.5 (35)	2,069	38.0 (171)
8902	1,049	15.5 (140)	2.5 (3)	3,961	6.8 (66)	6.0 (176)	9.5 (46)	1,564	13.7 (15)	11.3 (32)	2,471	38.8 (173)
8903	1,205	17.2 (148)	3.3 (8)	3,996	8.2 (131)	5.9 (175)	9.8 (55)	1,621	13.7 (15)	11.9 (38)	2,441	39.0 (174)
8904	1,629	16.5 (144)	3.8 (19)	6,490	8.4 (139)	6.3 (182)	9.4 (43)	2,460	12.9 (9)	12.0 (40)	3,903	40.7 (177)
8905	1,264	11.3 (99)	6.0 (52)	4,981	6.5 (59)	5.1 (171)	12.7 (147)	2,161	16.5 (78)	16.3 (128)	3,189	32.3 (164)
8909	523	16.4 (143)	3.4 (11)	1,737	9.6 (169)	6.9 (186)	10.1 (63)	569	13.5 (11)	13.5 (75)	960	38.6 (172)
9045	1,600	10.4 (87)	6.4 (64)	5,747	7.0 (82)	2.8 (81)	13.8 (171)	2,494	16.0 (62)	16.4 (131)	3,773	18.0 (110)
9050	863	14.7 (135)	4.1 (22)	2,950	7.5 (102)	2.4 (54)	9.3 (41)	1,288	12.1 (4)	12.7 (54)	1,908	39.5 (175)
9051	1,375	8.9 (51)	7.6 (87)	4,597	6.5 (59)	2.0 (28)	14.8 (178)	1,950	17.7 (119)	19.0 (156)	2,626	26.1 (151)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
9053	827	16.0 (141)	3.3 (8)	2,148	8.8 (154)	3.6 (132)	8.6 (23)	825	15.6 (52)	10.9 (25)	1,058	23.9 (146)
9055	469	9.0 (53)	4.3 (24)	412	12.9 (188)	2.2 (39)	5.8 (3)	1	*	*	2	*
9058	875	14.9 (136)	3.5 (12)	3,081	6.6 (62)	4.2 (153)	8.1 (17)	1,445	13.5 (11)	10.1 (14)	1,987	22.7 (139)
9066	1,030	16.5 (144)	3.6 (14)	3,154	8.7 (150)	3.9 (140)	8.9 (30)	1,361	11.5 (1)	11.7 (36)	1,954	33.6 (167)
9068	887	15.4 (139)	3.7 (16)	2,907	6.4 (57)	3.4 (123)	9.2 (40)	1,235	13.6 (14)	12.2 (46)	1,789	15.5 (84)
9069	2,588	16.0 (141)	3.1 (7)	8,376	7.0 (82)	4.3 (157)	8.8 (28)	3,471	13.5 (11)	10.4 (17)	5,124	23.8 (144)
9070	1,338	13.3 (121)	3.7 (16)	4,277	8.1 (126)	3.5 (128)	7.9 (13)	1,712	12.7 (8)	8.6 (5)	2,517	31.1 (163)
9072	1,275	14.9 (136)	3.5 (12)	4,065	8.3 (134)	3.3 (116)	7.5 (9)	1,678	14.1 (21)	9.9 (12)	2,477	28.3 (158)
9080	2,298	13.6 (129)	3.8 (19)	7,528	6.2 (49)	3.7 (136)	8.0 (15)	3,259	15.5 (48)	10.7 (22)	4,571	23.2 (142)
9082	129	10.9 (95)	4.7 (28)	289	10.7 (180)	2.8 (81)	9.7 (52)	67	*	*	133	26.3 (152)
9083	1,407	16.6 (146)	3.3 (8)	1,233	2.8 (1)	14.0 (190)	3.0 (1)	2	*	*	3	*
9084	961	13.5 (125)	3.9 (21)	891	8.0 (121)	15.3 (191)	4.0 (2)	3	*	*	5	*
9085	1	*	*	1	*	*	*	0	*	*	0	*
9086	3	*	*	2	*	*	*	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)						Anemia Low Hb/Hct (8)		
	Low		High	Short Stature		Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
9087	1	*	*	33	*	*	*	23	*	*	22	*
Michigan	140,119	11.4	6.2	494,719	7.5	3.4	10.8	207,671	16.2	14.5	303,575	19.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age ≤ 2.3rd percentile, underweight is defined as weight-for-length ≤ 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as ≥ 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age ≥ 85th to < 95th percentile, and obesity is defined as ≥ 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
0101	98	*	81	*	79	*	57	*	70	*	230	90.4 (38)	465	2.8 (110)
0102	350	65.4 (105)	248	31.9 (77)	235	11.1 (95)	179	17.9 (69)	217	11.5 (52)	804	90.2 (41)	1,577	3.7 (132)
0103	340	67.1 (95)	277	30.7 (82)	254	9.8 (105)	172	19.8 (56)	244	7.8 (90)	858	87.2 (69)	1,654	6.6 (181)
0104	121	70.2 (87)	93	*	79	*	69	*	80	*	315	89.5 (52)	607	4.6 (155)
0105	91	*	52	*	74	*	40	*	43	*	90	*	292	4.5 (154)
0250	122	75.4 (57)	100	35.0 (55)	114	15.8 (45)	70	*	86	*	231	90.5 (37)	517	1.7 (63)
0251	81	*	73	*	61	*	52	*	62	*	185	83.8 (105)	393	3.3 (124)
0252	106	79.2 (28)	57	*	99	*	25	*	54	*	343	74.3 (150)	586	2.7 (103)
0253	126	73.0 (72)	94	*	89	*	53	*	78	*	245	83.3 (107)	527	2.7 (103)
0401	419	71.1 (81)	305	29.2 (90)	368	10.9 (97)	221	17.2 (73)	264	8.3 (86)	1,021	81.4 (120)	1,999	4.3 (148)
0402	339	77.6 (42)	255	40.8 (23)	285	15.1 (52)	165	28.5 (20)	216	20.8 (5)	676	84.0 (103)	1,403	4.0 (138)
0403	138	70.3 (86)	96	*	105	11.4 (90)	75	*	84	*	263	86.7 (75)	561	8.7 (191)
0404	101	72.3 (75)	88	*	81	*	52	*	71	*	236	92.4 (20)	458	3.5 (129)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
0501	448	73.2 (68)	305	40.0 (25)	271	17.3 (32)	202	31.7 (14)	264	21.2 (4)	976	86.5 (77)	1,972	4.6 (155)
0502	0	*	0	*	0	*	0	*	0	*	1	*	1	*
0503	101	85.1 (8)	78	*	71	*	53	*	65	*	191	80.6 (127)	417	3.8 (133)
0504	415	85.5 (7)	312	49.7 (4)	281	23.1 (8)	181	40.3 (2)	258	20.5 (7)	892	87.1 (70)	1,801	2.7 (103)
0505	257	86.4 (4)	166	49.4 (5)	140	17.1 (35)	95	*	136	20.6 (6)	441	85.5 (90)	1,013	2.9 (115)
0506	148	75.7 (55)	117	41.0 (21)	116	14.7 (57)	73	*	100	11.0 (57)	435	87.6 (67)	772	8.8 (192)
0507	85	*	85	*	104	22.1 (11)	60	*	74	*	377	89.9 (46)	654	2.1 (79)
0508	24	*	31	*	40	*	23	*	30	*	150	84.7 (96)	238	2.5 (94)
0509	3	*	4	*	6	*	1	*	2	*	29	*	41	*
0650	953	72.7 (73)	735	36.3 (48)	724	14.4 (63)	478	23.0 (40)	609	12.5 (41)	1,898	90.6 (35)	4,047	2.7 (103)
0651	536	71.5 (77)	443	34.3 (64)	443	12.6 (76)	283	21.2 (52)	373	11.8 (48)	1,296	91.1 (29)	2,498	5.1 (164)
0652	290	72.4 (74)	216	36.6 (44)	216	12.5 (77)	145	17.9 (69)	180	10.0 (65)	695	87.9 (64)	1,324	6.2 (177)
0653	383	81.5 (23)	329	41.6 (18)	335	19.1 (20)	184	31.5 (15)	274	14.2 (30)	1,141	87.1 (70)	2,011	5.8 (171)
0654	316	73.7 (67)	256	34.0 (65)	269	19.0 (22)	163	25.2 (31)	223	11.7 (49)	727	85.7 (87)	1,443	5.9 (173)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
0655	283	75.6 (56)	240	42.5 (15)	232	12.5 (77)	152	21.1 (53)	188	11.2 (54)	635	83.9 (104)	1,301	6.7 (183)
0657	57	*	54	*	54	*	27	*	40	*	167	94.6 (12)	310	8.1 (190)
0701	328	86.0 (5)	272	53.3 (1)	286	30.8 (1)	183	35.5 (7)	238	18.5 (13)	882	90.9 (30)	1,657	0.7 (15)
0702	68	*	50	*	44	*	34	*	43	*	149	92.6 (19)	286	2.4 (91)
0703	160	73.1 (70)	149	34.9 (56)	142	15.5 (48)	99	*	130	16.2 (21)	404	81.4 (120)	805	5.5 (168)
0704	43	*	37	*	45	*	16	*	28	*	108	89.8 (47)	213	2.3 (86)
0705	82	*	80	*	66	*	52	*	68	*	244	93.0 (17)	444	1.6 (58)
0706	53	*	35	*	29	*	24	*	31	*	34	*	133	0.8 (19)
0850	717	82.7 (13)	579	38.0 (32)	530	17.7 (28)	354	15.8 (81)	479	8.1 (89)	1,691	80.1 (131)	3,349	2.0 (75)
0851	635	78.7 (33)	491	39.5 (27)	429	18.2 (25)	304	28.6 (19)	424	12.5 (41)	1,463	91.5 (24)	2,924	1.8 (67)
0901	1,397	65.8 (103)	807	29.5 (87)	1,159	13.1 (72)	471	22.3 (45)	735	6.0 (102)	3,219	88.8 (58)	6,469	2.5 (94)
0902	37	*	34	*	37	*	13	*	29	*	131	83.2 (108)	230	2.6 (101)
1051	1,112	82.6 (14)	724	50.3 (3)	812	25.0 (4)	416	37.0 (4)	580	15.2 (24)	2,206	90.6 (35)	4,633	1.5 (51)
1052	8	*	0	*	0	*	1	*	0	*	0	*	8	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
1101	1,597	68.4 (93)	1,065	30.4 (84)	838	14.9 (54)	831	18.3 (67)	906	8.5 (84)	3,078	75.3 (147)	6,635	3.8 (133)
1102	711	74.1 (62)	496	34.5 (62)	353	12.2 (82)	369	22.2 (46)	424	13.0 (36)	1,396	80.6 (127)	2,955	4.8 (160)
1104	3	*	1	*	0	*	1	*	1	*	0	*	3	*
1106	70	*	52	*	49	*	37	*	44	*	153	82.4 (114)	317	2.8 (110)
1250	944	73.8 (64)	573	36.8 (40)	672	15.3 (49)	392	26.8 (27)	488	13.5 (34)	1,741	80.5 (129)	3,760	3.3 (124)
1251	663	78.6 (35)	474	35.2 (54)	429	16.8 (38)	309	24.9 (35)	404	14.9 (26)	1,434	84.5 (98)	2,976	5.2 (166)
1252	524	83.4 (10)	381	36.7 (43)	380	14.5 (62)	239	21.8 (48)	289	9.3 (72)	1,178	84.3 (100)	2,408	1.4 (41)
1253	695	82.2 (17)	501	37.7 (35)	483	16.4 (41)	316	25.0 (33)	386	12.2 (44)	1,601	89.5 (52)	3,196	2.2 (83)
1350	2,085	71.2 (80)	1,398	31.3 (78)	1,530	14.4 (63)	898	18.8 (62)	1,097	9.6 (68)	4,316	88.8 (58)	8,852	1.6 (58)
1351	97	*	71	*	72	*	58	*	64	*	242	97.9 (4)	471	0.8 (19)
1352	364	64.3 (111)	254	24.4 (112)	304	11.5 (88)	144	18.8 (62)	200	12.0 (46)	961	98.3 (1)	1,781	0.5 (12)
1401	212	81.1 (24)	164	43.9 (12)	142	14.8 (56)	100	32.0 (12)	137	16.8 (17)	433	84.1 (102)	930	5.6 (170)
1402	113	89.4 (1)	78	*	78	*	47	*	65	*	318	79.2 (135)	601	1.2 (33)
1403	18	*	15	*	17	*	11	*	12	*	101	78.2 (140)	155	11.6 (193)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
1501	730	64.4 (108)	524	34.5 (62)	563	12.4 (79)	359	19.8 (56)	447	11.2 (54)	1,676	91.5 (24)	3,242	1.3 (37)
1503	856	73.9 (63)	624	38.0 (32)	709	14.7 (57)	457	24.1 (37)	521	17.1 (16)	2,113	94.2 (13)	4,038	1.5 (51)
1504	1,850	77.8 (41)	1,006	34.6 (60)	1,392	15.2 (51)	519	22.5 (43)	780	11.5 (52)	3,706	91.4 (26)	7,635	0.7 (15)
1505	386	78.8 (30)	203	37.9 (34)	284	15.1 (52)	128	30.5 (17)	171	16.4 (20)	946	96.2 (7)	1,800	1.0 (28)
1506	526	69.0 (91)	390	34.6 (60)	400	11.5 (88)	282	24.1 (37)	325	9.2 (73)	1,319	96.6 (6)	2,538	0.3 (6)
1507	531	80.6 (25)	379	38.3 (30)	379	13.2 (69)	269	36.1 (5)	297	18.9 (11)	946	86.9 (72)	2,418	2.5 (94)
1508	30	*	21	*	33	*	5	*	12	*	64	*	149	0.0 (1)
1509	362	69.1 (90)	236	28.0 (93)	306	13.1 (72)	155	19.4 (60)	194	10.3 (61)	916	91.7 (23)	1,760	2.5 (94)
1511	606	78.2 (38)	470	34.9 (56)	507	14.6 (59)	308	25.6 (29)	401	13.0 (36)	1,325	90.3 (40)	2,776	1.8 (67)
1512	322	76.7 (46)	237	32.1 (75)	295	14.9 (54)	152	21.1 (53)	200	10.0 (65)	822	92.0 (22)	1,608	0.4 (11)
1513	0	*	2	*	1	*	0	*	2	*	25	*	33	*
1514	252	88.1 (2)	159	44.0 (11)	219	27.4 (3)	99	*	124	21.8 (3)	636	98.0 (3)	1,204	0.3 (6)
1701	539	65.7 (104)	371	40.4 (24)	398	21.4 (14)	243	38.3 (3)	333	26.7 (1)	1,300	91.4 (26)	2,539	7.6 (187)
1950	531	81.9 (18)	374	34.8 (58)	311	16.4 (41)	233	25.3 (30)	308	11.7 (49)	1,032	82.8 (110)	2,242	5.1 (164)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
1952	121	86.8 (3)	89	*	71	*	51	*	70	*	242	84.7 (96)	523	1.5 (51)
1953	269	81.8 (19)	201	41.3 (20)	202	21.8 (12)	137	27.0 (26)	175	13.1 (35)	634	82.6 (112)	1,290	2.9 (115)
1954	418	79.4 (26)	315	39.7 (26)	271	24.4 (5)	202	31.2 (16)	270	13.7 (33)	894	86.4 (78)	1,853	1.8 (67)
1955	111	82.9 (12)	81	*	80	*	53	*	72	*	286	83.2 (108)	548	2.7 (103)
1957	386	81.6 (21)	266	35.7 (53)	265	15.8 (45)	163	25.2 (31)	225	10.2 (63)	881	84.3 (100)	1,828	1.6 (58)
1963	70	*	54	*	35	*	25	*	44	*	168	86.9 (72)	321	0.3 (6)
1964	23	*	32	*	22	*	20	*	29	*	83	*	172	2.9 (115)
2150	581	78.8 (30)	399	36.6 (44)	461	13.0 (75)	300	22.7 (41)	331	10.9 (58)	1,253	88.3 (62)	2,526	3.1 (120)
2151	183	59.0 (122)	136	27.2 (101)	136	2.9 (137)	100	16.0 (80)	120	10.8 (59)	339	93.2 (15)	769	3.6 (131)
2152	9	*	9	*	7	*	8	*	8	*	16	*	37	*
2251	364	70.9 (83)	247	43.3 (13)	292	21.6 (13)	183	33.3 (9)	218	23.9 (2)	911	89.4 (54)	1,750	1.0 (28)
2252	162	74.7 (59)	104	33.7 (66)	129	17.8 (27)	94	*	90	*	426	88.7 (60)	795	5.8 (171)
2554	50	*	43	*	43	*	36	*	39	*	238	81.1 (124)	355	6.2 (177)
2555	6,659	55.5 (131)	4,693	24.6 (110)	4,018	10.9 (97)	3,152	12.1 (91)	3,766	7.1 (93)	12,609	82.7 (111)	27,088	2.3 (86)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
2556	519	54.1 (134)	426	22.3 (122)	446	7.8 (118)	261	9.2 (106)	336	4.2 (114)	1,177	78.9 (136)	2,414	4.2 (142)
2557	6	*	3	*	1	*	3	*	3	*	31	*	45	*
2558	3	*	1	*	1	*	1	*	1	*	40	*	45	*
2559	1	*	0	*	2	*	1	*	0	*	31	*	34	*
2560	2	*	2	*	3	*	2	*	2	*	42	*	50	*
2561	1	*	0	*	0	*	0	*	0	*	10	*	11	*
2562	12	*	20	*	21	*	6	*	14	*	97	*	142	2.8 (110)
2563	40	*	26	*	36	*	18	*	20	*	67	*	156	3.2 (122)
2564	15	*	3	*	3	*	3	*	3	*	8	*	32	*
2565	0	*	0	*	0	*	0	*	0	*	3	*	3	*
3001	1,064	58.4 (125)	734	17.8 (130)	660	6.5 (124)	521	7.7 (112)	680	4.0 (115)	2,359	71.0 (157)	4,737	3.4 (128)
3002	276	51.1 (144)	205	16.6 (132)	178	4.5 (132)	142	10.6 (96)	191	3.1 (124)	701	85.9 (84)	1,339	2.5 (94)
3004	172	73.8 (64)	121	33.1 (71)	97	*	70	*	109	7.3 (92)	336	82.4 (114)	704	1.7 (63)
3005	57	*	27	*	18	*	25	*	27	*	107	89.7 (49)	213	2.3 (86)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
3007	10	*	12	*	11	*	8	*	11	*	42	*	79	*
3009	1	*	1	*	0	*	2	*	0	*	4	*	8	*
3201	389	78.1 (40)	313	36.4 (47)	278	13.7 (67)	197	11.7 (93)	266	5.6 (104)	936	85.8 (86)	1,902	4.1 (140)
3301	3,733	74.6 (60)	2,367	39.0 (28)	2,268	17.9 (26)	1,552	18.4 (66)	1,889	11.6 (51)	6,516	89.2 (55)	14,318	1.4 (41)
3304	92	*	73	*	75	*	58	*	63	*	284	93.0 (17)	525	0.2 (4)
3305	155	82.6 (14)	121	43.0 (14)	114	21.1 (15)	84	*	106	14.2 (30)	452	93.1 (16)	850	0.8 (19)
3306	250	78.8 (30)	205	44.4 (9)	184	23.4 (6)	128	27.3 (24)	175	16.0 (22)	670	95.1 (10)	1,287	1.4 (41)
3308	79	*	57	*	46	*	30	*	48	*	184	95.1 (10)	372	0.8 (19)
3309	14	*	15	*	10	*	9	*	12	*	42	*	76	*
3312	367	66.5 (97)	276	29.3 (89)	264	10.2 (104)	184	13.0 (88)	235	8.9 (78)	1,045	90.0 (43)	1,912	1.8 (67)
3314	2	*	0	*	0	*	0	*	0	*	0	*	2	*
3550	57	*	52	*	30	*	33	*	42	*	119	85.7 (87)	246	2.8 (110)
3551	20	*	14	*	7	*	11	*	14	*	32	*	74	*
3552	27	*	20	*	21	*	12	*	17	*	78	*	145	5.5 (168)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
3701	898	83.0 (11)	673	34.8 (58)	785	17.5 (29)	414	28.0 (22)	587	14.0 (32)	2,071	86.6 (76)	4,130	2.3 (86)
3850	2,743	70.1 (88)	2,520	35.9 (51)	2,199	11.6 (87)	1,433	21.6 (50)	2,044	12.9 (38)	6,316	77.5 (142)	12,402	3.5 (129)
3901	1,389	66.3 (99)	977	30.9 (80)	844	12.3 (80)	669	19.6 (59)	843	6.8 (94)	2,535	74.6 (149)	5,742	2.1 (79)
3902	187	58.8 (123)	114	36.8 (40)	94	*	78	*	96	*	489	80.8 (125)	996	1.9 (73)
3903	32	*	20	*	17	*	13	*	19	*	75	*	149	1.3 (37)
3904	6	*	5	*	9	*	3	*	5	*	27	*	45	*
3905	11	*	11	*	10	*	6	*	7	*	24	*	56	*
3950	1,980	70.5 (85)	1,420	32.0 (76)	1,282	14.6 (59)	941	21.3 (51)	1,198	9.0 (77)	4,031	78.6 (137)	8,635	1.8 (67)
3954	32	*	18	*	23	*	15	*	17	*	55	*	122	1.6 (58)
4103	8	*	9	*	6	*	5	*	8	*	25	*	53	*
4106	2,047	76.3 (51)	1,761	25.0 (108)	1,564	11.3 (93)	1,101	10.0 (102)	1,490	6.2 (100)	4,909	86.0 (82)	9,679	0.8 (19)
4107	2,534	75.9 (52)	2,065	32.6 (72)	1,833	15.8 (45)	1,365	20.3 (55)	1,796	12.1 (45)	5,048	84.5 (98)	10,950	2.3 (86)
4109	893	75.9 (52)	723	36.1 (50)	642	16.8 (38)	533	21.8 (48)	631	11.9 (47)	2,104	82.2 (116)	4,230	1.2 (33)
4110	8	*	1	*	0	*	0	*	1	*	0	*	11	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
4112	480	31.7 (152)	416	19.2 (127)	396	10.4 (103)	295	6.1 (120)	400	3.8 (117)	1,155	98.2 (2)	2,192	1.5 (51)
4113	18	*	22	*	40	*	11	*	20	*	203	97.0 (5)	336	0.0 (1)
4115	3,175	76.9 (45)	2,477	33.5 (68)	2,190	19.3 (18)	1,533	17.7 (71)	1,946	9.5 (70)	5,962	83.7 (106)	13,941	0.9 (25)
4351	476	79.0 (29)	281	37.4 (37)	396	18.9 (23)	168	33.9 (8)	243	17.3 (15)	1,132	96.1 (8)	2,222	1.4 (41)
4352	138	73.2 (68)	101	38.6 (29)	134	13.4 (68)	69	*	90	*	369	75.1 (148)	706	7.1 (184)
4353	365	77.0 (44)	239	42.3 (16)	287	16.4 (41)	152	22.4 (44)	198	9.1 (75)	1,021	87.3 (68)	1,828	2.7 (103)
4354	670	71.5 (77)	499	30.9 (80)	551	13.8 (66)	364	24.5 (36)	428	10.3 (61)	1,416	90.9 (30)	2,922	3.3 (124)
4355	278	75.9 (52)	202	44.6 (8)	260	18.5 (24)	140	32.1 (11)	172	18.6 (12)	652	92.2 (21)	1,297	4.2 (142)
4356	385	73.8 (64)	266	37.6 (36)	320	16.3 (44)	201	28.4 (21)	233	15.0 (25)	950	86.9 (72)	1,809	3.9 (135)
4358	80	*	70	*	85	*	45	*	64	*	253	89.7 (49)	464	2.4 (91)
4359	500	76.4 (50)	268	29.9 (85)	419	16.7 (40)	144	27.1 (25)	242	8.7 (81)	1,153	88.9 (56)	2,234	1.4 (41)
4360	185	81.6 (21)	134	48.5 (6)	169	20.7 (16)	85	*	121	16.5 (19)	472	89.8 (47)	897	4.6 (155)
4361	218	81.7 (20)	160	44.4 (9)	194	22.7 (10)	104	35.6 (6)	136	19.9 (8)	521	81.4 (120)	1,039	3.3 (124)
4362	269	85.9 (6)	194	50.5 (2)	197	28.4 (2)	117	45.3 (1)	160	19.4 (9)	552	85.5 (90)	1,216	6.0 (175)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
4363	721	76.6 (47)	517	41.6 (18)	599	19.2 (19)	321	29.6 (18)	441	14.7 (28)	1,660	87.8 (65)	3,270	4.6 (155)
4401	906	78.3 (37)	745	37.0 (39)	799	17.4 (31)	501	18.8 (62)	654	8.6 (83)	2,129	90.0 (43)	4,304	0.9 (25)
4501	857	75.1 (58)	673	37.1 (38)	652	14.1 (65)	435	17.0 (75)	556	5.2 (109)	1,951	87.7 (66)	3,862	3.1 (120)
4601	1,201	66.3 (99)	814	32.4 (73)	767	17.3 (32)	600	25.0 (33)	724	12.8 (39)	2,363	81.6 (119)	5,068	3.2 (122)
4602	7	*	5	*	8	*	1	*	3	*	29	*	51	*
4701	996	79.4 (26)	773	32.2 (74)	710	17.5 (29)	552	22.1 (47)	685	12.6 (40)	2,253	88.9 (56)	4,639	1.4 (41)
5050	1,447	60.6 (116)	1,076	27.7 (98)	1,038	13.1 (72)	858	16.1 (78)	951	12.3 (43)	2,282	85.2 (92)	5,530	1.7 (63)
5051	2,406	64.8 (107)	1,616	28.0 (93)	1,604	11.8 (85)	1,286	16.1 (78)	1,422	9.6 (68)	3,546	79.8 (132)	8,848	1.5 (51)
5052	2,554	59.8 (118)	1,789	27.3 (100)	1,771	10.9 (97)	1,401	12.3 (90)	1,568	6.6 (97)	4,094	77.9 (141)	9,862	2.0 (75)
5055	1,076	45.2 (148)	754	14.5 (136)	944	3.8 (135)	463	3.7 (124)	652	1.2 (131)	3,495	76.4 (145)	5,931	0.9 (25)
5101	1,507	69.4 (89)	1,002	31.3 (78)	1,170	14.6 (59)	566	19.8 (56)	785	10.2 (63)	3,065	86.3 (79)	6,533	1.4 (41)
5102	22	*	16	*	16	*	10	*	15	*	39	*	94	*
5202	676	84.3 (9)	421	36.8 (40)	523	17.0 (37)	242	33.1 (10)	336	14.9 (26)	1,547	93.6 (14)	3,071	2.1 (79)
5206	6	*	7	*	7	*	2	*	3	*	18	*	32	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
5601	874	78.5 (36)	681	38.3 (30)	584	22.9 (9)	549	31.9 (13)	605	18.5 (13)	1,828	90.9 (30)	3,875	1.7 (63)
5801	1,384	63.3 (113)	1,039	27.9 (96)	1,023	11.0 (96)	726	18.0 (68)	911	6.8 (94)	2,903	85.0 (94)	6,042	3.9 (135)
5803	114	57.9 (127)	112	23.2 (117)	113	9.7 (106)	71	*	99	*	400	88.0 (63)	715	1.4 (41)
5804	14	*	17	*	18	*	14	*	17	*	83	*	131	6.1 (176)
5806	245	68.6 (92)	202	33.2 (69)	114	9.6 (107)	145	13.8 (84)	164	6.7 (96)	105	84.8 (95)	540	4.4 (151)
6101	38	*	30	*	40	*	24	*	23	*	102	75.5 (146)	202	7.4 (186)
6102	113	74.3 (61)	94	*	111	17.1 (35)	66	*	72	*	351	85.2 (92)	638	4.1 (140)
6103	3,389	64.2 (112)	2,208	27.9 (96)	2,369	9.2 (111)	1,514	17.7 (71)	1,788	8.2 (88)	6,482	73.1 (153)	13,837	4.3 (148)
6105	19	*	19	*	29	*	12	*	16	*	79	*	136	0.0 (1)
6106	112	71.4 (79)	56	*	44	*	38	*	43	*	78	*	262	7.6 (187)
6301	3,322	66.8 (96)	2,532	28.0 (93)	2,005	11.4 (90)	1,896	14.5 (83)	2,202	8.8 (79)	6,016	80.7 (126)	13,457	1.2 (33)
6312	1,691	77.6 (42)	1,441	42.1 (17)	1,207	19.1 (20)	1,042	28.0 (22)	1,212	19.1 (10)	3,993	82.6 (112)	7,989	0.7 (15)
6316	1,083	76.6 (47)	882	33.2 (69)	725	13.2 (69)	610	16.9 (76)	769	9.5 (70)	2,288	90.4 (38)	4,754	0.8 (19)
6318	66	*	6	*	1	*	6	*	5	*	9	*	81	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
6320	170	82.4 (16)	100	47.0 (7)	54	*	92	*	89	*	373	80.2 (130)	709	0.3 (6)
6325	217	76.5 (49)	157	36.3 (48)	107	23.4 (6)	110	26.4 (28)	143	16.8 (17)	502	85.9 (84)	984	2.6 (101)
6335	911	71.1 (81)	777	27.4 (99)	563	10.8 (101)	567	13.4 (85)	680	8.4 (85)	2,285	91.4 (26)	4,494	0.3 (6)
6340	23	*	16	*	10	*	13	*	13	*	77	*	134	0.7 (15)
6345	58	*	38	*	31	*	37	*	33	*	100	86.0 (82)	234	2.1 (79)
6360	760	66.2 (101)	526	29.5 (87)	380	13.2 (69)	397	18.6 (65)	463	10.4 (60)	1,884	73.6 (152)	3,602	2.4 (91)
7304	27	*	29	*	25	*	21	*	26	*	83	*	159	1.3 (37)
7306	2,122	61.5 (115)	1,538	25.2 (107)	1,301	8.5 (115)	1,191	13.0 (88)	1,287	7.8 (90)	3,685	82.1 (117)	8,438	2.5 (94)
7308	30	*	24	*	26	*	9	*	13	*	78	*	166	1.2 (33)
7310	11	*	5	*	6	*	3	*	3	*	16	*	47	*
7401	1,994	65.9 (102)	1,375	28.1 (92)	1,496	12.1 (83)	871	18.9 (61)	1,206	11.2 (54)	3,940	79.3 (133)	8,255	4.3 (148)
7601	522	52.9 (137)	373	30.6 (83)	375	12.3 (80)	229	22.7 (41)	324	15.7 (23)	1,146	90.9 (30)	2,335	2.8 (110)
7850	970	73.1 (70)	763	35.9 (51)	832	17.2 (34)	486	16.7 (77)	640	9.1 (75)	2,181	88.7 (60)	4,464	3.0 (119)
7852	20	*	9	*	14	*	5	*	8	*	68	*	112	5.4 (167)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
8101	1,602	45.3 (147)	1,185	15.9 (134)	967	6.6 (123)	889	6.7 (114)	1,051	3.3 (123)	3,096	63.5 (163)	6,726	4.4 (151)
8102	336	50.0 (145)	260	15.8 (135)	230	4.3 (133)	190	6.3 (117)	228	2.2 (129)	1,097	71.1 (156)	1,913	2.7 (103)
8105	990	60.0 (117)	726	22.9 (118)	610	7.2 (120)	604	11.9 (92)	642	9.2 (73)	2,269	85.6 (89)	4,689	1.5 (51)
8106	430	59.8 (118)	202	12.4 (137)	289	3.5 (136)	163	10.4 (97)	189	1.6 (130)	886	49.4 (174)	1,769	1.8 (67)
8150	17	*	22	*	22	*	17	*	20	*	133	95.5 (9)	191	1.6 (58)
8151	2,721	78.2 (38)	2,093	40.9 (22)	1,911	19.6 (17)	1,523	23.7 (39)	1,746	14.5 (29)	5,530	90.2 (41)	11,700	1.3 (37)
8152	37	*	21	*	19	*	19	*	18	*	100	90.0 (43)	182	2.2 (83)
8153	10	*	9	*	9	*	8	*	9	*	33	*	58	*
8155	13	*	14	*	14	*	10	*	13	*	118	90.7 (34)	167	0.6 (13)
8157	5	*	9	*	6	*	4	*	8	*	37	*	55	*
8201	2,387	58.4 (125)	1,759	26.4 (102)	1,309	7.8 (118)	1,420	13.3 (86)	1,561	8.8 (79)	3,800	86.3 (79)	8,842	1.5 (51)
8202	3,398	43.4 (149)	2,562	25.6 (106)	2,547	9.5 (108)	1,819	6.3 (117)	2,187	4.0 (115)	8,571	82.0 (118)	16,290	1.4 (41)
8203	348	78.7 (33)	315	33.7 (66)	249	12.0 (84)	227	13.2 (87)	276	8.7 (81)	779	79.3 (133)	1,592	4.8 (160)
8250	618	64.4 (108)	459	25.9 (104)	335	8.1 (116)	347	8.6 (108)	396	5.6 (104)	912	74.2 (151)	2,254	1.9 (73)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
8252	116	70.7 (84)	79	*	99	*	62	*	70	*	263	55.5 (169)	553	4.2 (142)
8255	2,159	57.9 (127)	1,811	23.6 (116)	1,263	9.5 (108)	1,363	11.0 (94)	1,583	6.5 (98)	3,483	78.4 (139)	8,212	2.0 (75)
8257	1,674	71.9 (76)	1,439	36.6 (44)	1,212	15.3 (49)	1,055	4.4 (123)	1,234	3.7 (119)	3,818	77.1 (143)	7,720	0.6 (13)
8263	600	53.8 (135)	492	25.8 (105)	382	11.3 (93)	383	17.2 (73)	449	10.0 (65)	1,214	76.5 (144)	2,545	1.0 (28)
8264	615	56.7 (129)	559	24.3 (114)	439	11.8 (85)	399	15.0 (82)	503	8.3 (86)	1,243	81.3 (123)	2,646	1.1 (32)
8267	664	54.8 (132)	566	20.7 (126)	476	9.0 (113)	426	9.2 (106)	514	6.2 (100)	1,085	89.7 (49)	2,635	1.0 (28)
8901	804	51.7 (142)	677	19.2 (127)	519	6.7 (122)	495	5.7 (122)	572	3.0 (126)	1,786	57.9 (165)	3,834	6.3 (179)
8902	1,146	64.4 (108)	933	24.5 (111)	745	7.0 (121)	673	10.3 (98)	784	5.5 (107)	2,255	57.2 (166)	5,039	4.2 (142)
8903	1,238	52.4 (141)	877	18.5 (129)	706	5.1 (128)	675	5.9 (121)	775	2.8 (127)	2,279	53.5 (170)	5,017	6.3 (179)
8904	1,722	59.1 (121)	1,540	21.6 (123)	1,119	8.0 (117)	1,055	8.2 (109)	1,329	4.4 (112)	3,454	59.5 (164)	7,805	4.8 (160)
8905	1,382	65.2 (106)	1,045	26.1 (103)	831	9.3 (110)	756	10.2 (100)	883	5.5 (107)	3,014	71.2 (155)	6,223	2.5 (94)
8909	556	62.8 (114)	442	24.9 (109)	340	5.3 (127)	331	10.3 (98)	381	6.0 (102)	929	56.6 (167)	2,241	4.2 (142)
9045	1,618	52.5 (140)	1,025	24.4 (112)	934	9.1 (112)	665	9.8 (103)	883	6.5 (98)	3,650	70.4 (158)	7,260	2.2 (83)
9050	892	51.2 (143)	613	22.7 (121)	560	4.8 (131)	499	6.2 (119)	507	3.6 (120)	1,661	72.7 (154)	3,838	4.2 (142)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance ⁽¹⁾

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾

Children Aged < 5 Years ⁽³⁾

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household ^(4,5)

Clinic	Breastfeeding ⁽⁶⁾				Exclusive Breastfeeding ⁽⁷⁾				≤ 2 Hours TV Viewing/Day ⁽⁸⁾		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
9051	1,389	67.9 (94)	1,095	28.9 (91)	966	10.8 (101)	705	7.8 (111)	898	4.9 (111)	2,876	65.1 (160)	5,907	1.4 (41)
9053	831	59.7 (120)	408	29.9 (85)	321	10.9 (97)	315	10.8 (95)	342	5.6 (104)	1,138	53.4 (171)	2,652	8.0 (189)
9055	469	43.1 (150)	19	*	4	*	22	*	15	*	10	*	513	2.9 (115)
9058	908	58.7 (124)	606	23.8 (115)	641	11.4 (90)	493	9.3 (105)	520	3.1 (124)	2,071	86.3 (79)	4,033	2.0 (75)
9066	1,041	53.3 (136)	637	21.2 (124)	773	5.0 (129)	372	6.5 (116)	492	3.5 (122)	1,975	51.2 (173)	4,223	4.9 (163)
9068	888	46.3 (146)	587	17.0 (131)	544	5.0 (129)	469	7.9 (110)	526	2.7 (128)	1,752	78.6 (137)	3,655	4.4 (151)
9069	2,593	33.9 (151)	1,684	16.3 (133)	1,625	4.1 (134)	1,204	6.7 (114)	1,521	3.6 (120)	5,042	65.0 (161)	10,516	4.6 (155)
9070	1,341	54.8 (132)	911	22.9 (118)	716	6.0 (125)	752	9.6 (104)	810	4.4 (112)	2,384	52.4 (172)	5,268	7.1 (184)
9072	1,288	52.6 (139)	767	21.1 (125)	847	5.5 (126)	486	10.1 (101)	660	5.2 (109)	2,383	56.1 (168)	5,154	5.9 (173)
9080	2,357	52.8 (138)	1,484	22.8 (120)	1,702	8.9 (114)	1,036	7.6 (113)	1,251	3.8 (117)	4,772	66.3 (159)	9,901	3.9 (135)
9082	129	56.6 (130)	71	*	57	*	67	*	62	*	109	64.2 (162)	351	6.6 (181)
9083	1,408	66.5 (97)	85	*	1	*	91	*	53	*	2	*	1,407	0.2 (4)
9084	961	30.3 (153)	47	*	10	*	72	*	35	*	36	*	1,012	4.0 (138)
9085	1	*	0	*	0	*	0	*	0	*	0	*	1	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)
9086	3	*	0	*	0	*	0	*	0	*	3	*
9087	1	*	4	*	3	*	2	*	4	*	39	*
Michigan	142,230	65.8	101,810	30.4	96,308	13.0	70,351	16.9	86,399	9.3	289,548	81.0
											607,733	2.6

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.