

This report shows modules and locations where the specified region either needs to focus additional prevention efforts or is excelling in infection prevention. **Table 1** presents a cumulative attributable difference (CAD) determined using the 2020 HHS target standardized infection ratios (SIRs) for each module, using the NHSN 2015 Baselines, which is modeled after the data included in the CDC National and State Annual HAI Report. Numbers with “Need to Prevent” next to them show how many infections the region needs to prevent quarterly in order to reach the 2020 HHS target SIR. Numbers with “Prevented” next to them show the number of infections prevented beyond what was expected for the region according to the 2020 HHS target SIR. Corresponding SIRs for each module and location type are provided as well.

NHSN Module¹	Hospital Type	Number of Facilities²	Location³	SIR⁴	Significant (Y/N)⁵	CAD⁶	Prevented or Need to Prevent
CAUTI	Acute	11	All	0.417	N	-3.201	Prevented
		9	ICU	0.000	N	-2.171	Prevented
		11	WARD+	0.596	N	-1.030	Prevented
CLABSI	Acute	11	All	0.766	N	0.102	Need to Prevent
		9	ICU	0.000	N	-2.171	Prevented
		11	WARD+	1.089	N	1.557	Need to Prevent
		----	NICU	----	----	----	----
CDI	Acute	11	Facility-wide	0.596	Y	-3.297	Prevented
MRSA Bac	Acute	11	Facility-wide	0.311	N	-0.606	Prevented
SSI COLO	Acute	11	----	1.400	N	2.999	Need to Prevent
SSI HYST	Acute	9	----	0.910	N	0.231	Need to Prevent
SSI HPRO	Acute	9	----	0.469	N	-0.493	Prevented
SSI KPRO	Acute	9	----	1.298	N	0.921	Need to Prevent

¹CAUTI, catheter-associated urinary tract infection; CLABSI, central line-associated bloodstream infection; CDI, *Clostridioides difficile* infection LabID; MRSA Bac, methicillin-resistant *Staphylococcus aureus* bloodstream infection LabID; SSI COLO, admission/readmission colon surgical site infection; SSI HYST, admission/readmission abdominal hysterectomy surgical site infection; SSI HPRO, admission/readmission hip arthroplasty surgical site infection; SSI KPRO, admission/readmission knee arthroplasty surgical site infection.

²Note: facilities in which an SIR could not be calculated with a CAD of 0 may be excluded from this table. Data for locations with less than 5 facilities reporting have been suppressed.

³All includes all units for which in-plan data are reported; ICU includes all critical care units; WARD+ includes all WARD, WARD_ONC, SCA, STEP, or OTHER units; NICU includes all neonatal critical care units; Facility-wide includes all inpatient units.

⁴SIR: Standardized Infection Ratio: Ratio of observed events compared to the number of predicted events, accounting for unit type or other variables. An SIR of 1 can be interpreted as having the same number of events as predicted. An SIR that is between 0 and 1 represents fewer events than predicted, while an SIR of greater than 1 represents more events than predicted. SIRs were calculated using the 2015 NHSN Baselines. The SIR is only calculated if the number of predicted infections is ≥ 1.

⁵Significant (Y/N). A Y indicates that, based on the p-value and 95% Confidence Interval (CI), the SIR is statistically significantly different than 1. An N indicates that, based on the p-value and 95% CI, the SIR is not statistically significantly different than 1 (expected).

⁶CAD=Cumulative Attributable Difference. The number of infections that the region either needs to prevent to meet the 2020 HHS target or has prevented beyond the 2020 HHS target. 2020 HHS HAI Target SIRs: CAUTI = 0.75, CLABSI = 0.50, CDI = 0.70, MRSA bacteremia = 0.50, SSI = 0.70.

In addition to the NHSN modules reported above, other module data are reported to the MDHHS SHARP Unit. However, there are no corresponding HHS Target SIRs for these HAIs and therefore CAD values are not calculated. SIR values using the NHSN 2015 Baseline for these HAIs are shown in **Table 2** below.

NHSN Module¹	Hospital Type	Number of Facilities²	Location³	SIR⁴	Significant (Y/N)⁵
Total VAE	Acute	10	All	1.478	N
		8	ICU	1.171	N
		5	WARD+	2.004	N
IVAC	Acute	10	All	1.499	N
		8	ICU	0.493	N
		5	WARD+	3.057	N
MBI-CLABSI	Acute	11	All	.	.
		9	ICU	.	.
		11	WARD+	.	.

¹Total VAE, total ventilator-associated events; IVAC Plus, infection-related ventilator-associated complication; MBI-CLABSI, mucosal barrier injury central line-associated bloodstream infection.
²Note: facilities for which the number of predicted infections are 0 have been excluded. Data for locations with less than 5 facilities reporting have been suppressed.
³All includes all units for which in-plan data are reported; ICU includes all critical care units; WARD+ includes all WARD, WARD_ONC, SCA, STEP, or OTHER units; NICU includes all neonatal critical care units; Facility-wide includes all inpatient units.
⁴SIR: Standardized Infection Ratio: Ratio of observed events compared to the number of predicted events, accounting for unit type or other variables. An SIR of 1 can be interpreted as having the same number of events as predicted. An SIR that is between 0 and 1 represents fewer events than predicted, while an SIR of greater than 1 represents more events than predicted. SIRs were calculated using the 2015 NHSN Baselines. The SIR is only calculated if the number of predicted infections is ≥ 1 .
⁵Significant (Y/N). A Y indicates that, based on the p-value and 95% Confidence Interval (CI), the SIR is statistically significantly different than 1. An N indicates that, based on the p-value and 95% CI, the SIR is not statistically significantly different than 1 (expected).

Standardized utilization ratios (SURs) is a risk-adjusted comparison of the amount of indwelling device use to the amount expected to be used for each facility using the NHSN 2015 Baselines. There are currently no national or state targets for SURs. The SUR data reported to the MDHHS SHARP unit is included in **Table 3** below.

Indwelling Device Type	Hospital Type	Number of Facilities¹	Location²	SUR³	Significant (Y/N)⁴
Central Lines	Acute	11	All	0.860	Y
		9	ICU	0.948	Y
		11	WARD+	0.948	Y
		----	NICU	----	----
Urinary Catheters	Acute	11	All	0.913	Y
		9	ICU	0.912	Y
		11	WARD+	0.912	Y
Ventilators	Acute	10	All	0.553	Y
		8	ICU	0.850	Y
		5	WARD+	0.850	Y

¹Note: facilities for which the number of predicted device days are 0 have been excluded. Data for locations with less than 5 facilities reporting have been suppressed.

²All includes all units for which in-plan data are reported; ICU includes all critical care units; WARD+ includes all WARD, WARD_ONC, SCA, STEP, or OTHER units; NICU includes all neonatal critical care units; Facility-wide includes all inpatient units.

³SUR: Standardized Utilization Ratio: Ratio of observed device days compared to the number of predicted device days, accounting for unit type or other variables. An SUR of 1 can be interpreted as having the same number of device days as predicted. An SUR that is between 0 and 1 represents fewer device days than predicted, while an SUR of greater than 1 represents more device days than predicted. SURs were calculated using the 2015 Baselines. The SUR is only calculated if the number of predicted infections is ≥ 1 .

⁴Significant (Y/N). A Y indicates that, based on the p-value and 95% Confidence Interval (CI), the SUR is statistically significantly different than 1. An N indicates that, based on the p-value and 95% CI, the SUR is not statistically significantly different than 1 (expected).

Please contact Elli Stier (RayE7@michigan.gov) with questions, comments, or suggestions. All aggregate reports are posted at www.michigan.gov/hai