

Moxy[®] 2E Herbicide

FIFRA Section 24(c) Special Local Need Label (SLN)

**FOR DISTRIBUTION AND USE ONLY
IN THE STATE OF MICHIGAN**

**FOR USE ONLY BY MICHIGAN CUT FLOWER GLADIOLA GROWERS
For control of emerged weeds in cut flower gladiola plants**

Active Ingredient:

Octanoic acid ester of bromoxynil[®]
(3, 5-dibromo-4-hydroxybenzotrile) ..33.4%
Inert Ingredients66.6%
TOTAL.....100.0%

**KEEP OUT OF REACH OF CHILDREN
WARNING**

EPA Reg. No. 9779-346

SLN: MI-070002

This label expires and must not be distributed or used in accordance with this SLN registration after March 15, 2027.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label and the Federal label for this product must be in possession of the user at the time of pesticide application. Read the label affixed to the container for Moxy[®] 2E Herbicide and this Section 24(c) label before applying. Do not use in a manner inconsistent with applicable use directions, restrictions, precautions and limitations provided on the label affixed to the container for Moxy[®] 2E Herbicide and this Section 24(c) label.

CROP/FLOWER	RATE OF APPLICATION	COMMENTS
Gladiola Plants	0.75 to 1 pint/acre (0.1875 to 0.25 lb ai/acre)	<p>Use lower rates on coarse, sandy soils and higher rates on silt and clays. (See product label for more specific information on soil types)</p> <p>Moxy[®] 2E Herbicide should be applied to actively growing weed seedlings from crop emergence to the 2-leaf growth stage of the cut flower gladiola plants. To reduce the potential for temporary leaf burn, application should be made when crops are not stressed by excessive high, cool or cold evening temperatures, abrasive winds or mechanical injury just as that caused by hail, sleet, or insect feeding.</p> <p>One application of Moxy[®] 2E Herbicide should be sufficient for control of broadleaf weeds; however, an additional application is permitted. The total cumulative rate should not exceed 2 pints per acre per season.</p> <p>RESTRICTIONS and PRECAUTIONS:</p> <ul style="list-style-type: none"> • The total cumulative rate should not exceed 2 pints per acre per season • Do not plant rotational crops until the following season • Do not incorporate • Post-emergence application after the 2-leaf growth stage may result in increased crop injury • Addition of a spray additive or mixing with a fertilizer may cause excessive crop leaf burn

24(c) registrant:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589

1/0517/7 rev2 DGW