



FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Stinger[®]

EPA Reg. No. 62719-73

EPA 24(c) Special Local Need Registration SLN No. MI-110001

Postemergence Broadleaf Weed Control in Swiss Chard For Distribution and Use Only Within the State of Michigan

This label expires and must not be distributed or used in accordance with the SLN registration after December 31, 2025

Special Conditions and Risks of Use

USE OF Stinger[®] herbicide (THE "PRODUCT") ON SWISS CHARD (THE "CROP") MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Dow AgroSciences makes the Product available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Supplemental Labeling, you acknowledge and accept that:

- (1) You assume all risk of Crop injury, Crop yield reduction and/or Crop loss;
- (2) Dow AgroSciences does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental Labeling;
- (3) Dow AgroSciences does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Dow AgroSciences disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to inability to sell the Crop, Crop injury, Crop yield reduction and/or Crop loss;
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Dow AgroSciences, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Dow AgroSciences and not specified by the U.S. EPA or the State of Michigan.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Stinger® herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this supplemental labeling, use of Stinger is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger.

Directions for Use

Refer to product label for Precautions and Restrictions, Mixing Instructions and Application Directions.

Target Broadleaf Weeds, Application Rate, and Use Restrictions:

Target Broadleaf Weeds	Stinger (pint/acre)	Use Restrictions
annual sowthistle perennial sowthistle (suppression) prickly lettuce sweet clover wild buckwheat	1/4 - 1/2 (0.094 - 0.187 lb ae clopyralid/acre)	<ul style="list-style-type: none">• Preharvest interval: Do not apply within 30 days of harvest.• Make 1 broadcast application per crop per year.• For use under this SLN label, do not apply this product through any type of irrigation system.

Broadcast Application Rates: Apply uniformly with ground equipment in a minimum of 10 to 40 gallons of water per acre. Apply when crop is in the cotyledon to 8-leaf stage of growth and the weeds are young and actively growing, emergence to the 5-leaf stage of growth. Application to wild buckwheat should be made at the 1 to 3 leaf stage of growth, before vining begins.

Tank Mixes: Stinger may be tank mixed with other herbicides labeled for use on Swiss chard. Refer to the product labels of these herbicides for lists of other weeds controlled, rates of application and use precautions. To minimize crop response, avoid mixing Stinger with crop oil concentrate adjuvants unless required by the tank mix partner.

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners

R043-134

Accepted: __/__/__

Replaces: R043-094