

**RESTRICTED USE PESTICIDE  
DUE TO ACUTE TOXICITY  
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS ONLY- NOT TO BE  
USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A  
CERTIFIED APPLICATOR.**

**syngenta.**

**INTRA**

**Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MICHIGAN**

**Gramoxone® SL 2.0  
For Row Middle Weed Control in Lettuce (Head and Leaf)**

**EPA Reg. No. 100-1431  
EPA SLN No. MI-120002**

**This label expires and must not be distributed or used in accordance with this SLN  
registration after June 8, 2026.**

**KEEP OUT OF REACH OF CHILDREN /  
MANTENER FUERA DEL ALCANCE DE LOS NIÑOS.**



**DANGER / POISON  
PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail. See additional  
precautionary statements and directions for use inside booklet.

**DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

## Specific Use Directions

Use Pattern	Maximum No. of Applications per Season	Rate per Acre	Application Timing	Minimum Total Spray Volume per Acre
Shielded Row Middle Application	1	1.5 pints	2-3 weeks after seeding	40 gallons

### Application Instructions

- Apply with conventional ground equipment directing spray between the rows and using shields to prevent spray contact with crop plants.
- Apply when weeds and grasses are succulent and weed growth is 1 to 6 inches high.
- Do not allow spray to contact crop as injury or excessive residues may result.
- Add a nonionic surfactant at 0.25% v/v or 32 fl oz/100 gal of spray mix.

### Restrictions

- Do not allow animals to graze in treated areas.
- Do not apply where Gramoxone SL 2.0 (or other paraquat-containing products) has been used as a preplant preemergent spray.
- Do not apply more than 1.5 pints/A per season.
- DO NOT apply this product through any type of irrigation system.

### Precautions

- Weeds emerging after application will not be controlled.
- Outer leaves should be stripped at the time of harvest.

---

©2021 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company.

24(c) Registrant:  
Syngenta Crop Protection, LLC  
P. O. Box 18300  
Greensboro, NC 27419-8300

Label Code: MI1431048DA0821