

Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0101	44	*	*	*	*	*	*	*
0102	196	93.9	0.5	3.1	0.0	0.0	2.0	0.5
0103	184	96.2	0.0	2.2	0.0	0.0	1.6	0.0
0104	49	*	*	*	*	*	*	*
0250	60	*	*	*	*	*	*	*
0251	33	*	*	*	*	*	*	*
0252	42	*	*	*	*	*	*	*
0253	60	*	*	*	*	*	*	*
0401	200	94.5	0.5	0.5	0.0	1.0	3.0	0.5
0402	149	89.9	0.7	2.0	0.0	1.3	6.0	0.0
0403	54	*	*	*	*	*	*	*
0404	55	*	*	*	*	*	*	*
0501	231	95.2	0.0	2.2	0.0	0.0	2.6	0.0
0503	70	*	*	*	*	*	*	*
0504	209	91.4	0.5	1.9	2.9	1.4	1.9	0.0
0505	159	83.6	0.0	1.3	0.0	0.0	15.1	0.0
0506	75	*	*	*	*	*	*	*
0650	419	88.3	3.6	3.6	1.0	0.5	2.6	0.5
0651	269	94.8	0.0	1.9	0.0	0.0	1.9	1.5
0652	127	91.3	0.0	2.4	0.0	0.0	5.5	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0653	142	97.2	0.0	1.4	0.0	0.0	1.4	0.0
0654	154	95.5	1.9	0.0	0.0	0.6	1.9	0.0
0655	182	91.8	0.0	1.6	0.0	0.0	4.9	1.6
0657	27	*	*	*	*	*	*	*
0701	166	89.8	0.6	1.8	0.0	5.4	1.8	0.6
0702	36	*	*	*	*	*	*	*
0703	98	*	*	*	*	*	*	*
0704	30	*	*	*	*	*	*	*
0705	58	*	*	*	*	*	*	*
0850	470	87.4	4.7	4.9	0.2	0.9	1.7	0.2
0851	315	97.5	0.0	1.6	0.0	0.0	0.6	0.3
0901	674	82.8	2.1	9.2	0.3	0.1	4.7	0.7
0902	28	*	*	*	*	*	*	*
1051	620	91.5	1.1	3.9	0.8	0.3	1.6	0.8
1101	785	38.3	53.5	4.1	0.1	1.5	1.4	1.0
1102	375	77.3	10.4	5.1	0.3	2.1	2.9	1.9
1106	47	*	*	*	*	*	*	*
1250	411	84.4	0.7	11.9	0.0	1.2	1.0	0.7
1251	348	95.1	0.3	1.7	0.0	0.3	1.1	1.4
1252	1	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1253	631	77.2	2.4	16.3	0.0	0.2	3.2	0.8
1350	957	63.1	19.1	7.6	0.1	4.4	5.0	0.6
1351	49	*	*	*	*	*	*	*
1352	174	66.1	23.6	5.7	0.0	0.6	4.0	0.0
1401	83	*	*	*	*	*	*	*
1402	55	*	*	*	*	*	*	*
1403	9	*	*	*	*	*	*	*
1404	10	*	*	*	*	*	*	*
1501	445	51.0	7.0	37.1	0.0	0.0	1.3	3.6
1503	465	62.8	7.3	21.9	0.6	1.1	1.7	4.5
1504	1,016	47.6	4.0	40.2	0.1	5.3	1.1	1.7
1505	325	86.5	1.8	7.7	0.3	0.6	1.2	1.8
1506	250	80.4	0.8	12.4	0.0	0.4	4.0	2.0
1507	270	88.1	1.5	4.1	0.0	1.5	4.1	0.7
1508	66	*	*	*	*	*	*	*
1509	239	57.7	1.7	35.6	0.0	0.0	1.3	3.8
1511	315	94.0	2.2	2.5	0.0	0.0	1.0	0.3
1512	138	87.7	0.0	5.1	0.7	0.0	5.1	1.4
1701	294	59.2	0.3	1.0	20.7	0.3	18.0	0.3
1950	312	92.0	0.0	4.2	0.0	1.0	1.9	1.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
1952	56	*	*	*	*	*	*	*
1953	106	89.6	0.9	6.6	0.0	0.0	0.9	1.9
1954	258	89.1	0.0	10.1	0.4	0.0	0.4	0.0
1955	58	*	*	*	*	*	*	*
1957	235	89.8	1.7	6.8	0.0	0.4	0.4	0.9
1959	22	*	*	*	*	*	*	*
2150	286	86.7	0.0	2.4	1.7	0.3	8.7	0.0
2151	108	88.9	0.0	2.8	0.9	0.0	2.8	4.6
2251	188	88.3	0.0	4.3	0.0	0.5	3.2	3.7
2252	87	*	*	*	*	*	*	*
2551	149	29.5	65.1	3.4	0.0	0.0	2.0	0.0
2552	8	*	*	*	*	*	*	*
2554	44	*	*	*	*	*	*	*
2555	3,036	53.5	38.5	4.2	0.1	0.4	2.7	0.6
3001	369	16.0	68.6	12.5	0.0	0.8	1.9	0.3
3002	92	*	*	*	*	*	*	*
3003	2	*	*	*	*	*	*	*
3004	48	*	*	*	*	*	*	*
3007	9	*	*	*	*	*	*	*
3009	10	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3010	1	*	*	*	*	*	*	*
3201	215	92.6	0.0	4.2	0.0	1.4	0.9	0.9
3301	1,715	45.7	29.3	14.7	0.1	5.2	4.3	0.9
3302	96	*	*	*	*	*	*	*
3304	69	*	*	*	*	*	*	*
3305	69	*	*	*	*	*	*	*
3306	98	*	*	*	*	*	*	*
3308	19	*	*	*	*	*	*	*
3309	18	*	*	*	*	*	*	*
3550	38	*	*	*	*	*	*	*
3551	13	*	*	*	*	*	*	*
3552	18	*	*	*	*	*	*	*
3701	487	90.8	0.0	7.0	0.0	0.8	1.4	0.0
3805	1,316	76.8	12.5	4.9	0.2	0.5	5.1	0.2
3950	743	68.8	18.3	6.9	0.1	1.1	3.9	0.9
3952	2	*	*	*	*	*	*	*
3954	13	*	*	*	*	*	*	*
4001	529	34.6	51.6	8.9	0.0	0.8	3.6	0.6
4002	443	56.2	22.3	12.9	0.2	0.5	7.7	0.2
4103	38	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
4106	764	20.9	43.7	29.1	0.4	0.3	4.2	1.4
4107	1,310	58.3	21.2	13.1	0.6	2.0	4.6	0.2
4109	397	92.7	0.0	3.0	0.0	0.8	3.5	0.0
4110	409	39.4	10.3	45.2	1.0	0.0	4.2	0.0
4112	236	0.4	1.7	95.3	0.0	0.8	0.4	1.3
4113	327	11.3	21.4	63.9	0.0	1.8	1.5	0.0
4120	789	59.9	8.6	25.2	0.4	1.8	3.5	0.5
4160	785	44.8	30.2	11.0	0.3	8.8	4.7	0.3
4351	226	92.9	0.0	2.7	0.0	0.4	3.1	0.9
4352	73	*	*	*	*	*	*	*
4353	190	68.4	0.0	27.4	0.5	0.0	3.2	0.5
4354	309	89.3	4.2	1.9	0.0	1.3	1.6	1.6
4355	119	85.7	0.0	7.6	0.0	0.8	5.9	0.0
4356	217	88.9	0.0	7.8	0.0	0.0	2.8	0.5
4358	66	*	*	*	*	*	*	*
4359	219	72.6	0.0	24.2	0.0	0.0	0.0	3.2
4360	73	*	*	*	*	*	*	*
4361	89	*	*	*	*	*	*	*
4362	113	95.6	0.0	1.8	0.0	0.0	2.7	0.0
4363	396	94.9	0.5	2.3	0.0	0.5	1.3	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4401	536	88.8	0.4	4.7	0.0	0.0	5.4	0.7
4501	384	94.8	0.8	3.1	0.0	0.0	0.8	0.5
4502	1	*	*	*	*	*	*	*
4601	710	81.8	1.1	13.4	0.0	0.0	2.1	1.5
4701	609	92.0	0.5	4.3	0.2	0.8	1.3	1.0
5050	688	66.9	26.9	1.3	0.0	1.9	2.6	0.4
5051	1,181	64.7	24.6	3.2	0.1	2.5	4.3	0.5
5052	1,119	61.4	25.5	2.1	0.0	8.4	2.3	0.4
5101	929	74.4	9.5	11.0	0.1	1.4	2.9	0.8
5202	276	91.7	0.4	0.7	1.1	1.1	4.7	0.4
5203	44	*	*	*	*	*	*	*
5204	37	*	*	*	*	*	*	*
5601	484	92.4	1.2	2.7	0.2	0.6	2.5	0.4
5801	805	85.6	5.7	5.1	0.2	0.2	2.2	0.9
5803	68	*	*	*	*	*	*	*
5804	18	*	*	*	*	*	*	*
6101	25	*	*	*	*	*	*	*
6102	73	*	*	*	*	*	*	*
6103	1,374	67.1	21.6	6.3	0.1	0.1	4.1	0.5
6104	288	38.5	47.6	7.6	0.0	0.0	5.9	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6105	20	*	*	*	*	*	*	*
6106	9	*	*	*	*	*	*	*
6301	2,017	48.0	27.8	18.5	0.1	2.3	2.7	0.6
6312	1,231	38.3	54.3	1.7	0.2	2.0	2.8	0.6
6316	722	71.7	18.4	5.1	0.1	2.2	1.8	0.6
6320	72	*	*	*	*	*	*	*
6325	51	*	*	*	*	*	*	*
6330	178	34.3	49.4	13.5	0.6	0.0	2.2	0.0
6335	212	83.0	9.0	1.9	0.9	3.8	1.4	0.0
6340	53	*	*	*	*	*	*	*
6345	5	*	*	*	*	*	*	*
6350	90	*	*	*	*	*	*	*
7304	74	*	*	*	*	*	*	*
7306	1,105	49.4	33.8	13.3	0.1	0.5	2.2	0.7
7308	66	*	*	*	*	*	*	*
7310	25	*	*	*	*	*	*	*
7401	1,063	85.7	4.0	4.9	0.1	0.3	4.4	0.7
7601	267	91.0	0.0	7.5	0.0	0.4	0.7	0.4
7850	462	92.6	0.2	4.5	0.0	0.0	2.2	0.4
7852	38	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
8101	333	12.9	82.6	2.1	0.0	0.6	1.8	0.0
8102	18	*	*	*	*	*	*	*
8105	86	*	*	*	*	*	*	*
8106	33	*	*	*	*	*	*	*
8150	16	*	*	*	*	*	*	*
8151	1,243	43.0	37.3	9.7	0.3	3.2	6.3	0.2
8152	169	40.8	45.6	5.3	0.0	0.0	8.3	0.0
8153	14	*	*	*	*	*	*	*
8154	126	34.1	19.8	14.3	0.0	31.0	0.8	0.0
8155	21	*	*	*	*	*	*	*
8156	6	*	*	*	*	*	*	*
8157	2	*	*	*	*	*	*	*
8201	1,212	59.6	25.0	4.2	0.2	1.8	8.8	0.4
8202	1,393	86.1	9.3	2.9	0.1	1.2	0.2	0.1
8203	167	66.5	25.1	2.4	0.0	0.6	3.6	1.8
8250	377	29.4	60.5	5.3	0.5	0.8	3.4	0.0
8252	39	*	*	*	*	*	*	*
8255	1,311	65.6	22.5	7.2	0.3	1.8	2.4	0.2
8257	190	50.0	19.5	1.1	0.0	28.4	1.1	0.0
8258	491	71.5	18.7	6.1	0.2	1.2	1.6	0.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8260	313	47.0	31.9	1.6	0.0	18.2	1.3	0.0
8901	472	1.5	94.9	0.6	0.0	1.3	1.7	0.0
8902	376	4.8	92.6	0.5	0.0	0.3	1.6	0.3
8903	529	1.9	96.4	0.8	0.0	0.2	0.8	0.0
8904	518	3.1	90.3	0.4	0.4	4.1	1.7	0.0
8905	1,124	40.7	31.2	23.2	0.0	2.2	2.5	0.1
9012	18	*	*	*	*	*	*	*
9013	49	*	*	*	*	*	*	*
9015	20	*	*	*	*	*	*	*
9016	80	*	*	*	*	*	*	*
9017	31	*	*	*	*	*	*	*
9018	53	*	*	*	*	*	*	*
9019	23	*	*	*	*	*	*	*
9021	44	*	*	*	*	*	*	*
9041	14	*	*	*	*	*	*	*
9045	578	10.4	37.0	50.5	0.2	0.7	1.2	0.0
9050	550	10.7	51.3	36.2	0.0	0.9	0.7	0.2
9051	693	10.1	13.7	75.0	0.1	0.1	0.9	0.0
9053	1,037	2.0	95.6	0.8	0.0	0.9	0.7	0.1
9058	293	57.3	36.9	0.7	0.0	4.1	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9066	717	1.7	97.6	0.0	0.1	0.3	0.3	0.0
9068	477	1.9	96.2	0.4	0.2	0.4	0.8	0.0
9069	878	1.9	95.9	0.3	0.1	0.0	1.6	0.1
9070	620	2.1	87.7	0.2	0.0	8.4	1.3	0.3
9072	588	3.6	93.9	0.3	0.2	1.4	0.5	0.2
9074	353	2.0	96.0	0.8	0.3	0.0	0.8	0.0
9080	1,074	20.5	76.2	2.0	0.0	0.1	1.2	0.0
Unknown Clinics	117	28.2	17.9	14.5	12.0	5.1	15.4	6.8
Michigan	68,268	57.0	27.3	10.0	0.4	1.7	2.9	0.6
Nation (Prior Year)	1,162,961	41.6	22.7	29.2	1.1	3.4	1.3	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	44	*	*	*	*	*	*	*
0102	196	0.0	4.6	8.7	71.4	12.8	2.6	0.0
0103	184	0.0	8.2	13.6	62.5	15.2	0.5	0.0
0104	49	*	*	*	*	*	*	*
0250	60	*	*	*	*	*	*	*
0251	33	*	*	*	*	*	*	*
0252	42	*	*	*	*	*	*	*
0253	60	*	*	*	*	*	*	*
0401	200	0.0	4.0	11.5	68.0	16.0	0.5	0.0
0402	149	0.0	2.0	14.1	56.4	26.8	0.7	0.0
0403	54	*	*	*	*	*	*	*
0404	55	*	*	*	*	*	*	*
0501	231	0.0	5.2	12.1	66.2	16.0	0.4	0.0
0503	70	*	*	*	*	*	*	*
0504	209	0.0	1.9	9.1	69.4	17.7	1.9	0.0
0505	159	0.0	5.7	11.3	63.5	17.6	1.9	0.0
0506	75	*	*	*	*	*	*	*
0650	419	0.0	5.7	12.4	65.9	15.3	0.7	0.0
0651	269	0.0	3.3	14.9	66.5	13.4	1.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0652	127	0.0	3.9	11.0	64.6	19.7	0.8	0.0
0653	142	1.4	7.7	9.2	63.4	16.2	0.7	1.4
0654	154	0.0	5.8	14.9	60.4	15.6	3.2	0.0
0655	182	0.5	4.9	11.5	65.9	15.9	1.1	0.0
0657	27	*	*	*	*	*	*	*
0701	166	0.0	1.8	9.0	63.3	25.3	0.6	0.0
0702	36	*	*	*	*	*	*	*
0703	98	*	*	*	*	*	*	*
0704	30	*	*	*	*	*	*	*
0705	58	*	*	*	*	*	*	*
0850	470	0.0	5.3	10.2	66.2	17.0	1.3	0.0
0851	315	0.0	2.9	11.7	62.2	21.6	1.6	0.0
0901	674	0.1	5.2	11.1	61.7	19.7	2.1	0.0
0902	28	*	*	*	*	*	*	*
1051	620	0.3	5.8	9.4	59.7	21.8	1.6	1.5
1101	785	0.1	7.5	13.8	60.6	16.9	0.9	0.1
1102	375	0.3	5.1	15.2	61.3	17.1	1.1	0.0
1106	47	*	*	*	*	*	*	*
1250	411	0.0	5.6	11.9	63.5	17.3	1.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1251	348	0.3	6.9	13.5	64.7	13.5	1.1	0.0
1252	1	*	*	*	*	*	*	*
1253	631	0.5	5.4	11.3	65.1	16.2	1.6	0.0
1350	957	0.2	5.4	12.9	65.4	14.9	0.9	0.2
1351	49	*	*	*	*	*	*	*
1352	174	0.0	6.9	19.0	62.1	11.5	0.6	0.0
1401	83	*	*	*	*	*	*	*
1402	55	*	*	*	*	*	*	*
1403	9	*	*	*	*	*	*	*
1404	10	*	*	*	*	*	*	*
1501	445	0.2	5.4	13.5	60.9	18.2	1.8	0.0
1503	465	0.2	5.8	13.1	61.9	17.8	0.9	0.2
1504	1,016	0.2	5.4	11.3	61.0	20.6	1.5	0.0
1505	325	0.3	4.9	10.2	61.5	21.8	1.2	0.0
1506	250	0.4	3.6	9.6	68.8	16.8	0.8	0.0
1507	270	0.4	1.9	8.1	71.5	17.0	1.1	0.0
1508	66	*	*	*	*	*	*	*
1509	239	0.8	9.2	9.6	60.3	18.4	1.7	0.0
1511	315	0.0	2.5	13.3	66.7	17.1	0.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1512	138	0.0	3.6	12.3	68.1	15.9	0.0	0.0
1701	294	0.0	4.1	11.6	67.7	14.6	2.0	0.0
1950	312	0.0	5.1	15.7	67.9	10.6	0.3	0.3
1952	56	*	*	*	*	*	*	*
1953	106	0.0	6.6	4.7	72.6	14.2	1.9	0.0
1954	258	0.0	5.0	11.2	67.1	15.5	0.8	0.4
1955	58	*	*	*	*	*	*	*
1957	235	0.0	2.6	10.6	65.5	19.6	1.3	0.4
1959	22	*	*	*	*	*	*	*
2150	286	0.0	4.2	10.8	66.4	18.2	0.3	0.0
2151	108	0.0	3.7	15.7	63.0	16.7	0.9	0.0
2251	188	0.0	5.9	12.2	64.4	16.0	1.6	0.0
2252	87	*	*	*	*	*	*	*
2551	149	0.7	8.7	12.1	65.1	12.1	1.3	0.0
2552	8	*	*	*	*	*	*	*
2554	44	*	*	*	*	*	*	*
2555	3,036	0.3	5.8	13.3	62.7	16.6	1.3	0.0
3001	369	1.6	6.0	13.8	62.1	14.1	2.2	0.3
3002	92	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3003	2	*	*	*	*	*	*	*
3004	48	*	*	*	*	*	*	*
3007	9	*	*	*	*	*	*	*
3009	10	*	*	*	*	*	*	*
3010	1	*	*	*	*	*	*	*
3201	215	0.9	0.9	8.4	67.0	22.3	0.5	0.0
3301	1,715	0.5	4.4	11.1	65.2	17.6	1.1	0.2
3302	96	*	*	*	*	*	*	*
3304	69	*	*	*	*	*	*	*
3305	69	*	*	*	*	*	*	*
3306	98	*	*	*	*	*	*	*
3308	19	*	*	*	*	*	*	*
3309	18	*	*	*	*	*	*	*
3550	38	*	*	*	*	*	*	*
3551	13	*	*	*	*	*	*	*
3552	18	*	*	*	*	*	*	*
3701	487	0.4	6.6	15.0	64.3	12.5	1.0	0.2
3805	1,316	0.3	4.9	12.7	66.0	15.2	0.5	0.4
3950	743	0.5	5.5	11.6	62.6	18.7	0.8	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3952	2	*	*	*	*	*	*	*
3954	13	*	*	*	*	*	*	*
4001	529	0.0	8.1	14.7	61.1	14.9	1.1	0.0
4002	443	0.2	7.0	11.1	63.0	17.6	1.1	0.0
4103	38	*	*	*	*	*	*	*
4106	764	0.5	6.2	15.4	57.3	18.8	1.4	0.3
4107	1,310	0.2	4.0	10.2	63.4	20.7	1.1	0.5
4109	397	0.3	5.0	12.1	64.7	17.1	0.8	0.0
4110	409	0.2	5.6	12.5	60.4	20.5	0.7	0.0
4112	236	0.0	5.1	3.8	56.8	30.9	3.0	0.4
4113	327	0.0	6.4	7.0	41.0	19.9	2.1	23.5
4120	789	0.3	4.4	12.3	60.6	20.8	1.6	0.0
4160	785	0.1	3.4	9.8	61.4	22.9	2.2	0.1
4351	226	0.4	4.9	17.7	58.0	18.1	0.9	0.0
4352	73	*	*	*	*	*	*	*
4353	190	0.0	5.3	15.8	60.0	16.8	2.1	0.0
4354	309	0.3	3.6	13.9	65.4	14.6	2.3	0.0
4355	119	0.0	5.0	9.2	65.5	20.2	0.0	0.0
4356	217	0.5	6.5	12.9	68.7	11.1	0.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4358	66	*	*	*	*	*	*	*
4359	219	0.0	7.8	16.0	57.5	17.8	0.9	0.0
4360	73	*	*	*	*	*	*	*
4361	89	*	*	*	*	*	*	*
4362	113	0.0	5.3	10.6	69.0	14.2	0.9	0.0
4363	396	0.3	4.0	13.4	66.2	14.6	1.5	0.0
4401	536	0.2	4.3	10.3	62.9	21.1	1.3	0.0
4501	384	0.0	4.7	12.5	61.2	20.6	1.0	0.0
4502	1	*	*	*	*	*	*	*
4601	710	0.0	6.2	12.1	62.8	17.0	1.8	0.0
4701	609	0.2	2.8	7.1	62.9	25.1	2.0	0.0
5050	688	0.3	2.9	8.9	57.3	28.9	1.7	0.0
5051	1,181	0.1	2.6	8.0	64.5	23.4	1.4	0.0
5052	1,119	0.1	3.8	9.9	59.9	24.0	2.3	0.0
5101	929	0.0	2.8	9.1	60.2	25.1	2.8	0.0
5202	276	0.0	4.7	8.3	72.8	13.8	0.4	0.0
5203	44	*	*	*	*	*	*	*
5204	37	*	*	*	*	*	*	*
5601	484	0.0	3.1	11.8	67.8	16.5	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5801	805	0.2	6.3	13.7	63.7	14.9	1.0	0.1
5803	68	*	*	*	*	*	*	*
5804	18	*	*	*	*	*	*	*
6101	25	*	*	*	*	*	*	*
6102	73	*	*	*	*	*	*	*
6103	1,374	0.4	5.0	11.1	66.5	16.0	0.9	0.0
6104	288	1.0	11.5	18.4	59.7	8.7	0.7	0.0
6105	20	*	*	*	*	*	*	*
6106	9	*	*	*	*	*	*	*
6301	2,017	0.2	3.7	13.0	60.1	21.4	1.6	0.0
6312	1,231	0.0	1.9	7.1	58.3	29.7	2.9	0.1
6316	722	0.1	2.6	7.9	63.4	23.7	2.2	0.0
6320	72	*	*	*	*	*	*	*
6325	51	*	*	*	*	*	*	*
6330	178	0.0	7.3	11.2	55.1	22.5	1.1	2.8
6335	212	0.0	1.9	6.1	59.0	30.2	2.8	0.0
6340	53	*	*	*	*	*	*	*
6345	5	*	*	*	*	*	*	*
6350	90	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
7304	74	*	*	*	*	*	*	*
7306	1,105	0.4	5.2	12.4	63.9	16.7	1.4	0.0
7308	66	*	*	*	*	*	*	*
7310	25	*	*	*	*	*	*	*
7401	1,063	0.0	5.2	13.5	64.5	15.9	0.8	0.0
7601	267	0.0	4.5	13.5	58.8	20.6	2.2	0.4
7850	462	0.0	5.4	11.3	66.0	16.5	0.9	0.0
7852	38	*	*	*	*	*	*	*
8101	333	0.3	5.4	12.9	57.1	21.0	3.3	0.0
8102	18	*	*	*	*	*	*	*
8105	86	*	*	*	*	*	*	*
8106	33	*	*	*	*	*	*	*
8150	16	*	*	*	*	*	*	*
8151	1,243	0.0	2.5	7.1	64.0	24.9	1.6	0.0
8152	169	0.6	16.0	31.4	52.1	0.0	0.0	0.0
8153	14	*	*	*	*	*	*	*
8154	126	0.0	2.4	6.3	47.6	39.7	1.6	2.4
8155	21	*	*	*	*	*	*	*
8156	6	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8157	2	*	*	*	*	*	*	*
8201	1,212	0.1	3.5	11.3	60.1	23.2	1.8	0.1
8202	1,393	0.0	1.8	4.8	59.7	27.0	2.9	3.7
8203	167	0.0	4.2	9.0	62.9	24.0	0.0	0.0
8250	377	0.5	5.6	13.0	56.5	23.3	1.1	0.0
8252	39	*	*	*	*	*	*	*
8255	1,311	0.3	4.7	10.0	64.1	19.9	0.9	0.1
8257	190	0.0	3.7	3.7	55.8	34.2	2.6	0.0
8258	491	0.2	4.1	9.2	64.6	21.4	0.4	0.2
8260	313	0.0	3.2	11.8	58.5	22.4	3.5	0.6
8901	472	0.6	10.2	16.3	55.9	16.1	0.8	0.0
8902	376	0.0	7.7	15.7	56.1	19.7	0.8	0.0
8903	529	0.6	9.1	13.8	59.9	14.6	2.1	0.0
8904	518	0.0	5.6	13.9	55.2	24.1	1.2	0.0
8905	1,124	0.2	4.0	10.6	59.5	24.1	1.3	0.3
9012	18	*	*	*	*	*	*	*
9013	49	*	*	*	*	*	*	*
9015	20	*	*	*	*	*	*	*
9016	80	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9017	31	*	*	*	*	*	*	*
9018	53	*	*	*	*	*	*	*
9019	23	*	*	*	*	*	*	*
9021	44	*	*	*	*	*	*	*
9041	14	*	*	*	*	*	*	*
9045	578	0.7	7.1	11.6	56.2	22.7	1.6	0.2
9050	550	0.0	7.3	11.1	59.6	20.5	1.5	0.0
9051	693	0.6	4.9	8.8	53.5	29.3	2.7	0.1
9053	1,037	0.4	4.8	14.7	57.6	20.5	2.0	0.0
9058	293	0.0	3.1	7.2	53.9	32.8	3.1	0.0
9066	717	0.7	6.4	16.3	56.1	19.4	1.1	0.0
9068	477	0.2	6.5	15.7	56.6	19.5	1.5	0.0
9069	878	0.2	5.2	14.6	56.8	21.1	2.1	0.0
9070	620	0.0	6.8	11.9	58.4	21.9	1.0	0.0
9072	588	0.0	6.8	14.3	58.0	19.4	1.5	0.0
9074	353	0.8	3.4	10.5	64.0	19.8	1.4	0.0
9080	1,074	0.3	4.3	12.0	59.5	20.9	3.1	0.0
Unknown Clinics	117	0.0	1.7	3.4	48.7	43.6	1.7	0.9
Michigan	68,268	0.2	4.9	11.5	61.7	19.9	1.5	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Nation (Prior Year)	1,162,961	0.3	5.3	11.5	59.4	21.7	1.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	40	* *	* *	36	* *	* *	4	* *	6	* *	40	* *	40	* *
0102	174	3.4 (27)	56.9 (80)	143	23.1 (102)	55.9 (90)	20	* *	33	* *	182	6.6 (60)	182	11.0 (72)
0103	172	3.5 (31)	54.1 (54)	158	16.5 (42)	53.8 (75)	21	* *	21	* *	171	4.7 (23)	171	9.9 (57)
0104	46	* *	* *	38	* *	* *	5	* *	3	* *	44	* *	45	* *
0250	56	* *	* *	52	* *	* *	2	* *	9	* *	53	* *	54	* *
0251	33	* *	* *	26	* *	* *	2	* *	4	* *	28	* *	28	* *
0252	40	* *	* *	33	* *	* *	0	* *	9	* *	36	* *	36	* *
0253	60	* *	* *	56	* *	* *	4	* *	2	* *	58	* *	58	* *
0401	197	8.1 (117)	54.8 (64)	178	13.5 (11)	53.9 (78)	20	* *	42	* *	187	4.8 (26)	187	9.6 (50)
0402	148	3.4 (27)	57.4 (86)	119	18.5 (74)	47.1 (24)	15	* *	24	* *	136	6.6 (60)	137	8.0 (21)
0403	52	* *	* *	43	* *	* *	4	* *	9	* *	50	* *	50	* *
0404	53	* *	* *	49	* *	* *	12	* *	8	* *	52	* *	52	* *
0501	162	3.7 (41)	58.0 (92)	145	20.7 (94)	42.1 (7)	20	* *	21	* *	220	5.9 (44)	220	6.8 (9)
0503	68	* *	* *	64	* *	* *	9	* *	5	* *	69	* *	69	* *
0504	168	2.4 (5)	51.2 (26)	110	20.0 (86)	39.1 (3)	27	* *	23	* *	199	5.0 (29)	199	7.0 (11)
0505	146	5.5 (92)	54.1 (54)	123	24.4 (105)	50.4 (51)	10	* *	12	* *	150	6.0 (47)	150	12.7 (94)
0506	47	* *	* *	40	* *	* *	5	* *	5	* *	71	* *	71	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0650	414	6.3 (108)	57.0 (82)	372	23.4 (104)	46.8 (22)	44	* *	54	* *	394	5.6 (38)	394	8.1 (24)
0651	261	7.3 (114)	54.4 (59)	232	19.8 (84)	51.7 (62)	28	* *	45	* *	250	3.6 (8)	251	5.2 (5)
0652	125	2.4 (5)	60.8 (110)	107	15.0 (25)	56.1 (91)	13	* *	23	* *	119	8.4 (86)	119	10.9 (71)
0653	122	6.6 (112)	45.9 (4)	111	18.0 (64)	52.3 (66)	15	* *	20	* *	131	5.3 (33)	131	9.2 (43)
0654	150	8.0 (116)	55.3 (68)	134	17.9 (63)	50.0 (46)	21	* *	20	* *	147	3.4 (5)	147	8.2 (26)
0655	167	4.2 (61)	53.9 (50)	146	13.7 (12)	53.4 (73)	20	* *	47	* *	165	6.7 (63)	165	13.3 (103)
0657	26	* *	* *	22	* *	* *	1	* *	6	* *	25	* *	25	* *
0701	160	2.5 (8)	47.5 (9)	141	18.4 (71)	47.5 (28)	18	* *	42	* *	153	7.8 (79)	153	13.1 (102)
0702	30	* *	* *	26	* *	* *	2	* *	7	* *	32	* *	32	* *
0703	94	* *	* *	80	* *	* *	16	* *	9	* *	80	* *	80	* *
0704	29	* *	* *	27	* *	* *	2	* *	2	* *	29	* *	29	* *
0705	53	* *	* *	50	* *	* *	6	* *	10	* *	57	* *	57	* *
0850	449	4.9 (77)	55.9 (72)	377	14.6 (23)	56.8 (97)	57	* *	85	* *	423	4.3 (17)	421	11.4 (77)
0851	306	3.3 (21)	58.8 (98)	274	16.1 (37)	53.6 (74)	31	* *	52	* *	301	9.6 (97)	302	9.3 (45)
0901	583	3.4 (27)	57.8 (90)	487	20.1 (88)	49.3 (39)	65	* *	84	* *	626	9.7 (98)	626	12.3 (90)
0902	25	* *	* *	22	* *	* *	6	* *	8	* *	28	* *	28	* *
1051	593	4.0 (54)	47.0 (8)	517	14.1 (15)	50.9 (54)	71	* *	55	* *	577	3.6 (8)	577	7.5 (17)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1101	700	5.4 (88)	54.7 (61)	605	19.5 (82)	53.2 (71)	145	52.4 (34)	85	* *	718	3.5 (6)	719	6.4 (7)
1102	318	3.8 (46)	51.6 (32)	263	13.3 (7)	60.8(109)	51	* *	71	* *	325	3.1 (2)	324	4.6(3)
1106	39	* *	* *	31	* *	* *	5	* *	8	* *	41	* *	41	* *
1250	393	5.6 (95)	54.7 (61)	339	25.7(106)	40.7 (5)	64	* *	70	* *	373	9.7 (98)	373	5.1(4)
1251	333	5.1 (83)	59.8 (104)	291	16.8 (47)	58.8(105)	41	* *	65	* *	315	3.5 (6)	315	8.9(40)
1252	0	* *	* *	0	* *	* *	0	* *	1	* *	1	* *	1	* *
1253	503	6.0 (103)	52.7 (38)	388	17.3 (53)	47.4 (26)	86	* *	112	37.5(31)	581	12.4(111)	581	8.8(34)
1350	911	4.5 (69)	53.5 (46)	766	16.4 (40)	50.0 (46)	137	36.5 (17)	110	25.5(15)	860	6.7 (63)	854	10.8(68)
1351	45	* *	* *	41	* *	* *	11	* *	12	* *	47	* *	47	* *
1352	167	3.6 (35)	59.3 (100)	150	18.7 (75)	54.0 (79)	20	* *	15	* *	167	9.0 (91)	167	16.2(117)
1401	82	* *	* *	69	* *	* *	12	* *	12	* *	74	* *	74	* *
1402	53	* *	* *	44	* *	* *	7	* *	8	* *	48	* *	48	* *
1403	9	* *	* *	7	* *	* *	1	* *	1	* *	9	* *	9	* *
1404	6	* *	* *	3	* *	* *	1	* *	1	* *	3	* *	3	* *
1501	382	4.2 (61)	62.6 (116)	248	38.3(109)	39.5 (4)	40	* *	47	* *	384	10.9(108)	384	12.2(88)
1503	354	5.6 (95)	55.4 (70)	221	20.4 (89)	48.0 (31)	40	* *	60	* *	397	7.1 (69)	398	7.3(13)
1504	743	5.0 (81)	53.2 (42)	530	19.1 (79)	47.4 (26)	130	43.1 (28)	165	33.3(25)	892	9.1 (93)	897	8.0(21)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1505	307	5.5 (92)	45.9 (4)	230	20.4 (89)	42.6 (10)	33	* *	65	* *	302	4.3 (17)	303	7.9(20)
1506	223	4.0 (54)	59.6 (101)	95	* *	* *	30	* *	35	* *	226	15.0(115)	226	9.3(45)
1507	239	2.9 (16)	50.6 (21)	187	19.3 (80)	48.7 (35)	28	* *	32	* *	253	8.3 (85)	252	13.9(107)
1508	45	* *	* *	34	* *	* *	6	* *	7	* *	40	* *	40	* *
1509	217	2.3 (3)	57.6 (89)	48	* *	* *	23	* *	23	* *	206	4.4 (19)	206	6.3(6)
1511	310	3.2 (20)	55.2 (67)	247	15.4 (26)	51.0 (55)	37	* *	52	* *	297	7.7 (76)	298	8.7(32)
1512	133	3.8 (46)	53.4 (43)	81	* *	* *	21	* *	21	* *	126	3.2 (3)	126	9.5(48)
1701	288	3.5 (31)	53.5 (46)	251	17.5 (57)	49.4 (40)	28	* *	50	* *	273	5.1 (31)	273	11.4(77)
1950	299	6.4 (109)	58.2 (95)	253	15.8 (32)	51.0 (55)	40	* *	75	* *	287	7.7 (76)	286	11.5(80)
1952	55	* *	* *	49	* *	* *	13	* *	12	* *	50	* *	50	* *
1953	100	4.0 (54)	61.0 (112)	85	* *	* *	19	* *	29	* *	98	* *	98	* *
1954	226	6.2 (107)	49.1 (13)	202	17.8 (61)	47.5 (28)	42	* *	52	* *	240	4.2 (15)	241	13.7(105)
1955	48	* *	* *	37	* *	* *	5	* *	11	* *	54	* *	54	* *
1957	218	5.5 (92)	57.8 (90)	192	10.9 (4)	57.8(102)	26	* *	73	* *	220	5.0 (29)	219	13.7(105)
1959	20	* *	* *	17	* *	* *	3	* *	3	* *	20	* *	20	* *
2150	269	3.0 (17)	50.6 (21)	234	21.4 (98)	46.6 (21)	23	* *	42	* *	257	10.5(105)	257	10.5(64)
2151	103	4.9 (77)	53.4 (43)	91	* *	* *	9	* *	19	* *	91	* *	91	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2251	172	4.1 (58)	50.6 (21)	153	9.8 (3)	60.1(107)	17	* *	22	* *	170	4.1 (12)	171	12.3(90)
2252	85	* *	* *	70	* *	* *	10	* *	16	* *	76	* *	76	* *
2551	143	4.9 (77)	55.9 (72)	118	19.5 (82)	58.5(104)	2	* *	65	* *	126	12.7(112)	125	10.4(63)
2552	8	* *	* *	7	* *	* *	4	* *	1	* *	8	* *	8	* *
2554	44	* *	* *	39	* *	* *	4	* *	4	* *	40	* *	40	* *
2555	2,920	5.3 (87)	52.9 (40)	2,650	17.2 (51)	51.4 (57)	631	38.8 (19)	1,028	29.4(22)	2,829	8.2 (83)	2,825	13.0(99)
3001	350	4.3 (63)	61.7 (114)	281	22.8(101)	44.5 (11)	43	* *	220	36.8(29)	338	5.6 (38)	338	8.0(21)
3002	86	* *	* *	78	* *	* *	11	* *	51	* *	84	* *	84	* *
3003	2	* *	* *	2	* *	* *	0	* *	0	* *	2	* *	2	* *
3004	44	* *	* *	42	* *	* *	4	* *	4	* *	43	* *	43	* *
3007	7	* *	* *	3	* *	* *	0	* *	2	* *	3	* *	3	* *
3009	9	* *	* *	4	* *	* *	0	* *	0	* *	4	* *	4	* *
3010	1	* *	* *	1	* *	* *	0	* *	0	* *	1	* *	1	* *
3201	200	3.5 (31)	53.5 (46)	180	20.6 (91)	48.9 (36)	21	* *	27	* *	192	9.4 (96)	193	9.8(54)
3301	1,570	4.5 (69)	54.3 (58)	1,283	13.3 (7)	56.4 (95)	263	29.7 (7)	285	22.8(10)	1,552	4.8 (26)	1,552	9.7(51)
3302	95	* *	* *	79	* *	* *	18	* *	14	* *	87	* *	87	* *
3304	63	* *	* *	46	* *	* *	7	* *	19	* *	61	* *	61	* *

- (1) Reporting period is January 1 through December 31.
 - (2) Excludes records with unknown data or errors.
 - (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
 - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
 - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
 - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3305	68	* *	* *	62	* *	* *	9	* *	15	* *	63	* *	63	* *
3306	93	* *	* *	80	* *	* *	10	* *	20	* *	91	* *	91	* *
3308	17	* *	* *	15	* *	* *	3	* *	3	* *	17	* *	17	* *
3309	17	* *	* *	17	* *	* *	3	* *	6	* *	18	* *	18	* *
3550	38	* *	* *	35	* *	* *	5	* *	7	* *	35	* *	35	* *
3551	13	* *	* *	12	* *	* *	5	* *	7	* *	12	* *	12	* *
3552	18	* *	* *	16	* *	* *	2	* *	7	* *	16	* *	16	* *
3701	363	5.2 (84)	50.4 (19)	287	17.4 (54)	50.5 (52)	45	* *	70	* *	445	6.5 (58)	445	8.8(34)
3805	1,184	5.0 (81)	51.4 (30)	947	16.5 (42)	51.6 (59)	153	32.7 (10)	122	25.4(14)	1,175	6.1 (51)	1,174	10.1(59)
3950	687	3.6 (35)	56.0 (75)	544	14.3 (19)	55.1 (85)	111	39.6 (21)	144	29.2(21)	658	13.7(114)	658	9.3(45)
3952	2	* *	* *	2	* *	* *	0	* *	1	* *	2	* *	2	* *
3954	12	* *	* *	11	* *	* *	1	* *	4	* *	12	* *	12	* *
4001	433	3.9 (50)	54.0 (52)	321	28.0(108)	42.4 (9)	47	* *	62	* *	463	10.4(104)	462	7.4(15)
4002	368	4.1 (58)	57.1 (84)	286	18.2 (68)	54.5 (82)	45	* *	78	* *	401	15.0(115)	401	8.7(32)
4103	37	* *	* *	33	* *	* *	8	* *	7	* *	36	* *	37	* *
4106	685	3.4 (27)	56.9 (80)	538	21.6 (99)	44.6 (12)	108	38.9 (20)	106	39.6(35)	680	4.7 (23)	680	8.5(28)
4107	1,087	3.7 (41)	46.6 (7)	777	15.6 (29)	48.6 (33)	192	37.0 (18)	232	35.3(28)	1,183	6.6 (60)	1,183	12.1(86)

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4109	346	7.8 (115)	49.1 (13)	171	7.6 (1)	56.7 (96)	55	* *	66	* *	380	6.1 (51)	380	9.2(43)
4110	363	3.3 (21)	62.5 (115)	282	17.4 (54)	51.4 (57)	25	* *	38	* *	376	6.4 (56)	376	8.8(34)
4112	233	2.1 (2)	69.5 (119)	221	72.9(110)	9.0 (1)	14	* *	17	* *	226	15.9(117)	227	8.8(34)
4113	296	3.0 (17)	60.8 (110)	224	18.8 (76)	46.9 (23)	20	* *	19	* *	292	9.9(102)	293	8.5(28)
4120	704	3.1 (19)	50.6 (21)	562	14.1 (15)	53.0 (70)	109	22.0 (3)	144	25.7(16)	732	4.6 (21)	732	9.8(54)
4160	507	5.9 (101)	49.1 (13)	318	14.5 (22)	53.8 (75)	113	29.2 (6)	161	26.1(19)	724	4.7 (23)	724	9.1(41)
4351	210	3.3 (21)	61.0 (112)	181	8.8 (2)	58.0(103)	21	* *	51	* *	207	3.9 (10)	208	9.1(41)
4352	70	* *	* *	58	* *	* *	7	* *	11	* *	67	* *	67	* *
4353	166	3.6 (35)	56.6 (78)	141	21.3 (96)	44.7 (13)	17	* *	29	* *	167	10.8(107)	167	15.0(111)
4354	294	6.1 (105)	54.4 (59)	262	15.6 (29)	57.6(101)	39	* *	52	* *	282	6.0 (47)	282	7.4(15)
4355	104	5.8 (100)	51.9 (34)	79	* *	* *	15	* *	18	* *	100	9.0 (91)	100	11.0(72)
4356	193	5.2 (84)	53.4 (43)	173	13.3 (7)	55.5 (89)	18	* *	44	* *	202	4.0 (11)	201	9.5(48)
4358	51	* *	* *	35	* *	* *	5	* *	8	* *	52	* *	52	* *
4359	209	4.3 (63)	57.4 (86)	194	23.2(103)	44.8 (14)	22	* *	29	* *	203	12.8(113)	204	11.8(83)
4360	71	* *	* *	60	* *	* *	4	* *	22	* *	68	* *	69	* *
4361	83	* *	* *	74	* *	* *	7	* *	23	* *	78	* *	78	* *
4362	103	8.7 (118)	51.5 (31)	89	* *	* *	16	* *	22	* *	102	4.9 (28)	102	9.8(54)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4363	379	2.6 (9)	56.5 (76)	338	18.9 (78)	55.3 (87)	33	* *	87	* *	365	7.4 (75)	365	12.9(98)
4401	527	6.5 (110)	50.5 (20)	455	16.0 (34)	48.6 (33)	79	* *	90	* *	498	10.2(103)	498	10.8(68)
4501	351	4.6 (72)	49.3 (16)	321	20.6 (91)	40.8 (6)	35	* *	53	* *	362	10.5(105)	362	11.0(72)
4502	1	* *	* *	1	* *	* *	0	* *	0	* *	1	* *	1	* *
4601	660	4.7 (73)	51.7 (33)	573	21.1 (95)	45.2 (16)	125	20.8 (2)	139	19.4 (5)	646	6.5 (58)	647	12.5(93)
4701	598	3.3 (21)	50.2 (18)	532	13.9 (14)	56.8 (97)	85	* *	170	13.5 (1)	552	7.8 (79)	553	12.8(96)
5050	673	5.6 (95)	52.5 (36)	443	15.6 (29)	54.0 (79)	127	40.2 (23)	251	22.3 (8)	463	9.3 (95)	465	11.4(77)
5051	1,136	4.3 (63)	54.1 (54)	775	13.7 (12)	55.4 (88)	210	40.0 (22)	336	20.8 (7)	828	8.2 (83)	827	10.6(66)
5052	1,083	4.9 (77)	54.8 (64)	687	16.7 (46)	48.9 (36)	205	35.6 (16)	359	19.2 (4)	738	7.7 (76)	741	10.5(64)
5101	883	3.9 (50)	51.2 (26)	736	15.5 (27)	49.6 (42)	152	25.7 (4)	231	24.7(12)	841	6.4 (56)	840	11.9(84)
5202	257	5.4 (88)	45.9 (4)	227	14.1 (15)	57.3(100)	39	* *	65	* *	254	5.9 (44)	254	14.2(108)
5203	41	* *	* *	33	* *	* *	5	* *	10	* *	39	* *	38	* *
5204	35	* *	* *	33	* *	* *	8	* *	7	* *	35	* *	35	* *
5601	466	4.7 (73)	54.1 (54)	412	14.1 (15)	56.3 (93)	64	* *	63	* *	452	7.3 (74)	452	15.3(114)
5801	740	5.4 (88)	51.2 (26)	629	18.1 (66)	52.3 (66)	111	35.1 (15)	122	23.0(11)	725	5.7 (42)	727	14.4(109)
5803	65	* *	* *	57	* *	* *	7	* *	14	* *	60	* *	60	* *
5804	14	* *	* *	13	* *	* *	2	* *	2	* *	17	* *	17	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)																
Clinic	Prepregnancy BMI ⁽⁴⁾				Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy		
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)		Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6101	20	* *	* *		19	* *	* *		2	* *	1	* *	23	* *	22	* *
6102	63	* *	* *		57	* *	* *		13	* *	6	* *	69	* *	69	* *
6103	1,221	4.4 (67)	58.1 (94)		1,087	19.4 (81)	52.4 (68)		150	28.7 (5)	103	18.4 (3)	1,276	8.7 (89)	1,270	11.7(81)
6104	262	1.1 (1)	53.1 (41)		228	21.9(100)	46.5 (20)		26	* *	13	* *	259	5.4 (34)	256	10.2(61)
6105	18	* *	* *		17	* *	* *		1	* *	4	* *	19	* *	19	* *
6106	9	* *	* *		6	* *	* *		0	* *	0	* *	7	* *	7	* *
6301	1,953	4.4 (67)	53.6 (49)		1,722	16.4 (40)	51.6 (59)		379	31.7 (9)	407	25.3(13)	1,822	8.0 (82)	1,823	11.7(81)
6312	1,192	2.6 (9)	57.1 (84)		1,047	14.6 (23)	53.8 (75)		240	33.3 (12)	357	29.7(23)	1,118	7.1 (69)	1,119	13.0(99)
6316	717	3.8 (46)	47.8 (10)		634	15.5 (27)	55.2 (86)		134	20.1 (1)	194	20.1 (6)	644	5.6 (38)	651	7.8(19)
6320	62	* *	* *		45	* *	* *		17	* *	24	* *	61	* *	61	* *
6325	50	* *	* *		42	* *	* *		8	* *	15	* *	48	* *	48	* *
6330	150	2.7 (12)	55.3 (68)		119	14.3 (19)	54.6 (83)		21	* *	30	* *	161	6.8 (67)	159	15.1(112)
6335	196	3.6 (35)	51.0 (25)		134	17.2 (51)	52.2 (65)		34	* *	38	* *	190	6.8 (67)	190	6.8(9)
6340	44	* *	* *		31	* *	* *		3	* *	10	* *	50	* *	50	* *
6345	3	* *	* *		3	* *	* *		0	* *	0	* *	4	* *	4	* *
6350	84	* *	* *		58	* *	* *		13	* *	18	* *	80	* *	80	* *
7304	49	* *	* *		42	* *	* *		6	* *	7	* *	72	* *	72	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7306	950	3.3 (21)	56.5 (76)	733	18.1 (66)	51.8 (63)	164	41.5 (25)	115	22.6 (9)	1,018	6.3 (54)	1,015	11.3(76)
7308	55	* *	* *	46	* *	* *	9	* *	9	* *	58	* *	58	* *
7310	18	* *	* *	17	* *	* *	6	* *	3	* *	25	* *	25	* *
7401	1,029	4.5 (69)	47.9 (11)	879	17.7 (60)	50.7 (53)	125	32.8 (11)	218	15.1 (2)	956	7.2 (71)	958	8.5(28)
7601	258	5.4 (88)	53.9 (50)	233	18.0 (64)	46.4 (19)	24	* *	46	* *	247	9.7 (98)	248	8.1(24)
7850	455	6.8 (113)	52.1 (35)	390	13.3 (7)	52.6 (69)	54	* *	63	* *	420	6.7 (63)	419	8.4(27)
7852	37	* *	* *	33	* *	* *	7	* *	8	* *	37	* *	37	* *
8101	296	2.7 (12)	58.8 (98)	259	17.0 (49)	50.2 (49)	74	* *	68	* *	321	7.2 (71)	322	12.1(86)
8102	18	* *	* *	15	* *	* *	3	* *	9	* *	17	* *	17	* *
8105	70	* *	* *	63	* *	* *	24	* *	49	* *	82	* *	82	* *
8106	29	* *	* *	22	* *	* *	12	* *	12	* *	32	* *	32	* *
8150	15	* *	* *	12	* *	* *	1	* *	5	* *	14	* *	14	* *
8151	1,170	3.7 (41)	54.7 (61)	945	16.1 (37)	49.7 (44)	206	35.0 (14)	316	30.7(24)	1,104	7.9 (81)	1,107	9.7(51)
8152	128	9.4 (119)	43.8 (3)	84	* *	* *	26	* *	25	* *	147	4.1 (12)	147	8.8(34)
8153	14	* *	* *	10	* *	* *	2	* *	8	* *	13	* *	13	* *
8154	125	5.6 (95)	32.0 (1)	105	16.2 (39)	25.7 (2)	23	* *	34	* *	111	0.9 (1)	111	0.9(1)
8155	19	* *	* *	17	* *	* *	2	* *	3	* *	21	* *	21	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾				Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight		Over-weight		< Ideal		> Ideal		3rd Trimester		Postpartum			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number			% (Rank)
8156	6	* *	* *	4	* *	* *	2	* *	1	* *	5	* *	5	* *
8157	2	* *	* *	2	* *	* *	1	* *	0	* *	2	* *	2	* *
8201	1,100	4.0 (54)	51.3 (29)	874	12.6 (6)	54.8 (84)	235	31.1 (8)	595	26.1 (19)	1,087	8.4 (86)	1,086	11.0 (72)
8202	1,195	4.7 (73)	48.3 (12)	952	17.6 (59)	49.5 (41)	232	43.5 (29)	252	37.3 (30)	1,286	5.5 (35)	1,285	2.9 (2)
8203	155	2.6 (9)	56.8 (79)	119	14.3 (19)	56.3 (93)	30	* *	45	* *	140	12.1 (110)	141	12.8 (96)
8250	366	5.2 (84)	52.7 (38)	146	19.9 (85)	60.3 (108)	88	* *	43	* *	158	4.4 (19)	158	10.1 (59)
8252	31	* *	* *	14	* *	* *	10	* *	7	* *	37	* *	37	* *
8255	1,199	6.1 (105)	52.5 (36)	668	16.0 (34)	51.6 (59)	233	33.5 (13)	430	26.0 (18)	1,158	5.8 (43)	1,158	10.3 (62)
8257	167	6.0 (103)	55.1 (66)	83	* *	* *	25	* *	47	* *	164	9.8 (101)	164	7.3 (13)
8258	450	3.3 (21)	49.8 (17)	372	12.1 (5)	61.6 (110)	102	47.1 (31)	158	33.5 (26)	435	6.0 (47)	434	10.6 (66)
8260	231	5.6 (95)	43.3 (2)	163	16.0 (34)	44.8 (14)	55	* *	54	* *	304	10.9 (108)	304	8.6 (31)
8901	434	3.9 (50)	59.7 (103)	372	18.8 (76)	53.2 (71)	80	* *	60	* *	435	5.5 (35)	435	12.2 (88)
8902	351	2.8 (14)	62.7 (117)	309	17.5 (57)	50.2 (49)	63	* *	56	* *	353	4.2 (15)	353	12.7 (94)
8903	485	3.9 (50)	59.8 (104)	427	21.3 (96)	48.0 (31)	92	* *	75	* *	492	4.1 (12)	492	12.0 (85)
8904	479	2.3 (3)	60.3 (109)	412	18.2 (68)	49.8 (45)	101	63.4 (36)	78	* *	480	5.2 (32)	483	9.7 (51)
8905	1,071	3.5 (31)	57.0 (82)	951	17.8 (61)	47.2 (25)	171	40.4 (24)	169	39.6 (35)	1,045	6.7 (63)	1,048	7.2 (12)
9012	16	* *	* *	9	* *	* *	0	* *	0	* *	16	* *	16	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
9013	46	* *	* *	38	* *	* *	2	* *	4	* *	42	* *	42	* *
9015	15	* *	* *	8	* *	* *	2	* *	1	* *	19	* *	19	* *
9016	74	* *	* *	61	* *	* *	7	* *	6	* *	73	* *	73	* *
9017	25	* *	* *	13	* *	* *	0	* *	1	* *	28	* *	28	* *
9018	48	* *	* *	41	* *	* *	0	* *	0	* *	47	* *	47	* *
9019	16	* *	* *	13	* *	* *	1	* *	2	* *	22	* *	22	* *
9021	38	* *	* *	15	* *	* *	0	* *	1	* *	37	* *	37	* *
9041	11	* *	* *	7	* *	* *	0	* *	1	* *	12	* *	12	* *
9045	521	3.6 (35)	58.7 (97)	431	16.9 (48)	46.2 (18)	93	* *	64	* *	545	8.6 (88)	546	8.8(34)
9050	528	2.8 (14)	60.2 (108)	454	18.3 (70)	48.9 (36)	93	* *	77	* *	508	6.3 (54)	508	10.0(58)
9051	627	2.4 (5)	60.0 (107)	451	20.0 (86)	42.1 (7)	61	* *	113	25.7(16)	633	9.2 (94)	633	7.6(18)
9053	1,017	3.7 (41)	58.0 (92)	894	17.4 (54)	56.9 (99)	221	48.4 (32)	214	45.3(37)	960	6.0 (47)	961	15.2(113)
9058	189	4.8 (76)	54.0 (52)	145	16.6 (44)	47.6 (30)	38	* *	40	* *	285	8.8 (90)	285	6.7(8)
9066	692	3.6 (35)	57.4 (86)	622	16.6 (44)	51.9 (64)	143	42.0 (26)	194	39.2(34)	682	5.6 (38)	682	13.5(104)
9068	463	6.5 (110)	59.6 (101)	407	26.3(107)	49.6 (42)	80	* *	85	* *	444	6.1 (51)	444	14.9(110)
9069	861	4.3 (63)	59.8 (104)	766	18.4 (71)	56.1 (91)	165	49.7 (33)	242	38.0(32)	820	7.2 (71)	820	16.1(116)
9070	594	5.9 (101)	55.9 (72)	484	18.4 (71)	45.9 (17)	123	42.3 (27)	121	38.8(33)	558	5.9 (44)	562	12.3(90)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9072	579	4.1 (58)	58.4 (96)	514	20.6 (91)	50.0 (46)	113	45.1 (30)	112	34.8 (27)	539	4.6 (21)	538	13.0 (99)
9074	353	3.7 (41)	66.3 (118)	312	17.0 (49)	59.0 (106)	68	* *	70	* *	330	3.3 (4)	330	15.5 (115)
9080	937	3.8 (46)	55.6 (71)	739	15.8 (32)	54.3 (81)	175	57.7 (35)	167	55.1 (38)	1,009	5.5 (35)	1,009	10.8 (68)
Unknown Clinics	79	* *	* *	59	* *	* *	10	* *	23	* *	73	* *	72	* *
Michigan	62,450	4.4	54.4	50,780	17.6	51.0	10,073	35.7	13,477	28.1	61,404	7.0	61,421	10.5
Nation (Prior Year)	1,106,370	4.5	53.4	835,927	21.5	48.0	164,201	33.9	346,981	28.1	419,956	5.2	467,675	6.3

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	40	* *	* *		41	* *	27	* *	33	* *
0102	185	89.2 (22)	5.4 (49)		194	47.4 (34)	168	25.6 (35)	167	87.4 (45)
0103	175	88.0 (29)	5.1 (45)		180	50.6 (18)	151	26.5 (27)	157	91.1 (14)
0104	45	* *	* *		48	* *	48	* *	45	* *
0250	54	* *	* *		56	* *	50	* *	50	* *
0251	28	* *	* *		33	* *	21	* *	32	* *
0252	36	* *	* *		41	* *	39	* *	36	* *
0253	59	* *	* *		59	* *	57	* *	57	* *
0401	188	94.7 (5)	3.2 (22)		199	48.2 (31)	195	23.6 (41)	178	93.3 (9)
0402	139	75.5 (87)	18.0 (95)		144	45.8 (42)	123	26.0 (30)	130	86.2 (52)
0403	49	* *	* *		52	* *	51	* *	41	* *
0404	52	* *	* *		54	* *	35	* *	47	* *
0501	222	79.3 (70)	15.3 (88)		222	50.9 (17)	202	25.7 (33)	200	83.5 (70)
0503	68	* *	* *		69	* *	68	* *	60	* *
0504	201	80.6 (65)	17.4 (94)		206	50.5 (20)	204	37.3 (6)	179	86.6 (51)
0505	150	72.0(104)	21.3 (105)		159	54.7 (5)	154	27.9 (18)	140	82.1 (81)
0506	73	* *	* *		73	* *	66	* *	64	* *
0650	392	77.8 (74)	19.1 (99)		409	49.1 (26)	395	25.8 (31)	369	90.5 (19)
0651	251	62.2(112)	26.7 (112)		259	47.9 (32)	176	29.5 (14)	219	88.6 (35)
0652	119	78.2 (73)	12.6 (82)		123	45.5 (44)	114	27.2 (23)	104	94.2 (5)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0653	129	74.4 (93)	24.8 (109)		140	52.1 (13)	127	15.0 (79)	123	85.4 (59)
0654	147	72.1(103)	19.7 (102)		149	51.7 (15)	145	22.1 (43)	129	77.5 (94)
0655	162	88.3 (26)	4.9 (41)		176	33.0 (88)	170	20.0 (55)	151	96.0 (2)
0657	25	* *	* *		27	* *	27	* *	24	* *
0701	154	75.3 (88)	18.2 (96)		164	42.1 (61)	156	35.9 (8)	130	89.2 (31)
0702	33	* *	* *		35	* *	35	* *	30	* *
0703	80	* *	* *		95	* *	90	* *	81	* *
0704	29	* *	* *		30	* *	29	* *	28	* *
0705	57	* *	* *		57	* *	58	* *	51	* *
0850	412	86.2 (37)	7.8 (61)		460	40.4 (68)	337	23.7 (40)	375	85.1 (62)
0851	292	91.8 (8)	4.1 (33)		304	44.4 (50)	223	28.7 (17)	256	84.8 (65)
0901	626	74.8 (90)	18.4 (97)		655	52.5 (10)	646	20.9 (49)	595	82.5 (80)
0902	28	* *	* *		26	* *	25	* *	24	* *
1051	572	61.9(113)	32.2 (115)		612	46.6 (39)	513	32.2 (10)	515	87.8 (39)
1101	710	76.9 (77)	6.5 (53)		769	33.4 (87)	641	9.7 (104)	637	88.9 (33)
1102	320	89.1 (23)	4.1 (33)		362	41.4 (63)	160	36.9 (7)	305	91.1 (14)
1106	41	* *	* *		46	* *	42	* *	38	* *
1250	308	75.0 (89)	14.0 (84)		406	47.0 (38)	384	15.9 (74)	357	72.3 (111)
1251	304	89.8 (17)	4.6 (37)		342	40.9 (66)	317	12.0 (93)	275	85.8 (56)
1252	1	* *	* *		1	* *	0	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1253	572	67.8(111)	24.7 (108)	622	42.4 (58)	591	11.0 (99)	535	79.1 (93)
1350	847	82.2 (59)	9.2 (70)	947	39.8 (71)	609	25.8 (31)	793	75.9 (98)
1351	46	* *	* *	48	* *	37	* *	36	* *
1352	166	61.4(114)	26.5 (111)	169	51.5 (16)	146	16.4 (73)	148	73.6 (106)
1401	73	* *	* *	81	* *	65	* *	72	* *
1402	45	* *	* *	55	* *	45	* *	44	* *
1403	9	* *	* *	8	* *	9	* *	9	* *
1404	5	* *	* *	10	* *	6	* *	6	* *
1501	357	85.2 (41)	7.0 (56)	438	45.7 (43)	256	21.9 (45)	362	64.1 (114)
1503	389	84.8 (44)	5.9 (51)	459	42.7 (56)	275	32.0 (11)	362	73.5 (107)
1504	872	88.9 (24)	4.8 (40)	996	42.4 (58)	768	19.3 (64)	828	82.6 (78)
1505	298	86.2 (37)	7.7 (59)	322	38.2 (77)	235	29.8 (13)	258	85.7 (57)
1506	222	83.3 (52)	11.3 (77)	244	43.0 (55)	224	21.9 (45)	197	83.2 (74)
1507	253	85.0 (42)	10.7 (75)	267	49.1 (26)	206	29.1 (15)	230	91.3 (13)
1508	35	* *	* *	66	* *	11	* *	36	* *
1509	194	80.9 (64)	11.9 (80)	233	50.6 (18)	158	17.7 (71)	195	76.4 (97)
1511	293	91.5 (10)	2.7 (18)	315	45.4 (45)	279	18.3 (67)	256	81.6 (82)
1512	125	94.4 (6)	4.0 (32)	136	33.8 (84)	117	11.1 (97)	109	79.8 (87)
1701	268	75.7 (84)	18.7 (98)	287	57.5 (3)	203	33.5 (9)	265	75.8 (100)
1950	287	88.2 (27)	7.0 (56)	304	44.1 (52)	268	14.2 (86)	260	86.9 (48)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1952	50	* *	* *	* *	54	* *	45	* *	42	* *
1953	96	* *	* *	* *	102	39.2 (73)	91	* *	91	* *
1954	240	80.4 (66)	9.2 (70)		252	45.2 (48)	222	26.1 (28)	224	89.7 (27)
1955	52	* *	* *	* *	57	* *	45	* *	46	* *
1957	218	90.8 (13)	4.6 (37)		229	39.3 (72)	197	20.8 (50)	173	83.2 (74)
1959	20	* *	* *	* *	22	* *	18	* *	22	* *
2150	256	84.4 (46)	9.8 (72)		280	52.5 (10)	273	25.6 (35)	250	85.6 (58)
2151	87	* *	* *	* *	106	44.3 (51)	71	* *	83	* *
2251	170	91.2 (11)	6.5 (53)		187	47.1 (35)	176	23.9 (39)	155	94.8 (4)
2252	76	* *	* *	* *	83	* *	67	* *	73	* *
2551	123	90.2 (16)	4.9 (41)		147	66.0 (1)	96	* *	147	72.8 (110)
2552	8	* *	* *	* *	8	* *	5	* *	8	* *
2554	41	* *	* *	* *	43	* *	29	* *	37	* *
2555	2,823	83.3 (52)	2.5 (16)		2,995	20.5 (113)	2,158	8.1 (106)	2,246	79.7 (88)
3001	338	74.6 (91)	10.9 (76)		361	50.1 (22)	333	6.0 (110)	318	74.8 (104)
3002	84	* *	* *	* *	87	* *	86	* *	83	* *
3003	2	* *	* *	* *	2	* *	2	* *	0	* *
3004	39	* *	* *	* *	47	* *	41	* *	41	* *
3007	4	* *	* *	* *	9	* *	6	* *	5	* *
3009	3	* *	* *	* *	9	* *	8	* *	7	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
3010	1	* *	* *	1	* *	1	* *	1	* *	
3201	195	79.0 (72)	15.9 (90)	211	48.8 (29)	201	20.4 (52)	180	90.0 (25)	
3301	1,503	81.4 (62)	8.6 (67)	1,691	35.1 (81)	1,517	15.9 (74)	1,414	85.9 (54)	
3302	87	* *	* *	96	* *	61	* *	92	* *	
3304	57	* *	* *	66	* *	59	* *	57	* *	
3305	62	* *	* *	67	* *	62	* *	56	* *	
3306	91	* *	* *	97	* *	94	* *	80	* *	
3308	17	* *	* *	19	* *	16	* *	17	* *	
3309	17	* *	* *	18	* *	16	* *	14	* *	
3550	35	* *	* *	37	* *	38	* *	37	* *	
3551	12	* *	* *	12	* *	13	* *	13	* *	
3552	16	* *	* *	16	* *	16	* *	15	* *	
3701	446	95.1 (4)	2.5 (16)	476	50.4 (21)	462	10.4 (102)	427	95.8 (3)	
3805	1,153	84.7 (45)	7.9 (63)	1,294	47.6 (33)	771	26.1 (28)	1,112	75.9 (98)	
3950	656	84.0 (50)	9.8 (72)	728	41.8 (62)	700	18.9 (65)	619	84.5 (67)	
3952	2	* *	* *	2	* *	1	* *	1	* *	
3954	12	* *	* *	13	* *	13	* *	9	* *	
4001	462	60.0(115)	27.7 (114)	515	52.6 (9)	490	11.0 (99)	459	59.9 (115)	
4002	400	74.0 (97)	16.8 (93)	436	45.4 (45)	409	14.4 (82)	378	74.9 (103)	
4103	36	* *	* *	38	* *	38	* *	34	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4106	659	73.0(102)	16.5 (91)	744	38.6 (74)	661	8.9 (105)	635	85.4 (59)
4107	1,179	79.3 (70)	14.2 (85)	1,290	38.5 (75)	1,092	21.4 (47)	1,111	88.1 (37)
4109	378	83.1 (55)	11.4 (78)	393	40.2 (70)	373	19.8 (58)	337	92.6 (10)
4110	370	59.5(116)	35.9 (116)	405	58.3 (2)	301	12.3 (91)	403	85.4 (59)
4112	222	73.4 (99)	0.9 (3)	226	52.7 (8)	202	95.5 (1)	214	16.4 (116)
4113	286	44.8(117)	43.0 (117)	323	52.9 (7)	270	6.7 (108)	309	83.2 (74)
4120	731	84.3 (47)	7.7 (59)	773	34.4 (82)	738	15.6 (76)	642	93.8 (7)
4160	724	89.4 (19)	5.2 (46)	773	32.2 (91)	724	15.2 (78)	647	93.5 (8)
4351	203	90.6 (14)	3.9 (29)	225	47.1 (35)	194	19.6 (62)	193	84.5 (67)
4352	67	* *	* *	72	* *	65	* *	63	* *
4353	164	75.6 (86)	16.5 (91)	186	52.2 (12)	167	12.6 (90)	159	80.5 (86)
4354	282	87.6 (33)	8.2 (65)	306	41.2 (65)	175	86.9 (3)	259	91.9 (12)
4355	101	73.3(100)	13.9 (83)	117	43.6 (54)	102	25.5 (38)	103	80.6 (85)
4356	201	69.7(107)	26.9 (113)	212	54.2 (6)	200	22.0 (44)	189	86.8 (50)
4358	47	* *	* *	63	* *	50	* *	49	* *
4359	195	89.7 (18)	3.6 (27)	216	40.3 (69)	197	30.5 (12)	179	87.7 (41)
4360	67	* *	* *	73	* *	53	* *	68	* *
4361	77	* *	* *	85	* *	80	* *	68	* *
4362	102	74.5 (92)	19.6 (101)	112	49.1 (26)	109	25.7 (33)	104	87.5 (44)
4363	361	70.1(106)	22.4 (106)	387	51.9 (14)	264	27.7 (19)	342	84.2 (69)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4401	494	74.3 (94)	19.2 (100)	529	41.4 (63)	455	20.0 (55)	442	83.3 (72)
4501	361	69.0(109)	23.0 (107)	374	48.4 (30)	369	27.1 (24)	336	89.6 (29)
4502	1	* *	* *	1	* *	1	* *	1	* *
4601	632	83.2 (54)	7.9 (63)	693	35.4 (80)	394	15.0 (79)	563	90.4 (20)
4701	555	84.9 (43)	8.8 (69)	598	29.1 (98)	523	27.5 (21)	439	90.4 (20)
5050	457	88.0 (29)	4.4 (36)	681	32.6 (90)	422	26.8 (25)	565	90.8 (16)
5051	825	88.7 (25)	3.2 (22)	1,168	32.9 (89)	744	20.4 (52)	961	88.4 (36)
5052	731	88.0 (29)	3.0 (21)	1,111	30.9 (95)	626	22.4 (42)	966	90.3 (23)
5101	834	89.3 (21)	5.3 (47)	911	34.1 (83)	534	27.7 (19)	725	86.2 (52)
5202	251	91.6 (9)	5.6 (50)	275	42.2 (60)	231	27.3 (22)	237	90.3 (23)
5203	37	* *	* *	44	* *	33	* *	39	* *
5204	35	* *	* *	37	* *	31	* *	31	* *
5601	455	81.3 (63)	15.2 (87)	480	46.3 (41)	462	26.6 (26)	420	96.2 (1)
5801	721	76.1 (82)	15.4 (89)	789	46.4 (40)	763	19.8 (58)	679	89.8 (26)
5803	59	* *	* *	68	* *	63	* *	51	* *
5804	17	* *	* *	18	* *	17	* *	14	* *
6101	22	* *	* *	25	* *	19	* *	21	* *
6102	68	* *	* *	73	* *	66	* *	61	* *
6103	1,233	84.2 (49)	7.0 (56)	1,361	49.5 (24)	1,011	20.7 (51)	1,134	75.0 (101)
6104	255	91.0 (12)	2.4 (14)	283	56.5 (4)	206	15.5 (77)	261	73.2 (108)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6105	16	* *	* *	* *	20	* *	16	* *	14	* *
6106	7	* *	* *	* *	9	* *	5	* *	8	* *
6301	1,806	84.3 (47)	6.6 (55)		2,000	31.8 (92)	1,400	14.3 (83)	1,620	83.3 (72)
6312	1,104	87.9 (32)	4.3 (35)		1,222	26.2(101)	788	19.9 (57)	902	87.8 (39)
6316	641	89.4 (19)	5.0 (43)		717	28.5 (99)	517	14.3 (83)	546	87.9 (38)
6320	61	* *	* *	* *	71	* *	16	* *	56	* *
6325	48	* *	* *	* *	51	* *	47	* *	37	* *
6330	162	76.5 (80)	8.6 (67)		173	29.5 (96)	149	12.8 (89)	138	87.7 (41)
6335	190	86.3 (36)	11.6 (79)		208	43.8 (53)	41	* *	181	89.5 (30)
6340	50	* *	* *	* *	53	* *	17	* *	38	* *
6345	4	* *	* *	* *	5	* *	2	* *	4	* *
6350	81	* *	* *	* *	86	* *	51	* *	73	* *
7304	71	* *	* *	* *	73	* *	71	* *	63	* *
7306	984	68.3(110)	21.1 (104)		1,082	45.4 (45)	1,032	11.5 (95)	957	77.0 (96)
7308	59	* *	* *	* *	65	* *	59	* *	55	* *
7310	24	* *	* *	* *	25	* *	25	* *	23	* *
7401	945	73.1(101)	20.1 (103)		1,049	49.9 (23)	663	21.0 (48)	937	79.7 (88)
7601	244	81.6 (61)	12.3 (81)		266	45.1 (49)	155	29.0 (16)	225	79.6 (90)
7850	415	69.2(108)	24.8 (109)		451	49.4 (25)	443	19.6 (62)	401	83.5 (70)
7852	37	* *	* *	* *	36	* *	33	* *	31	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8101	321	75.7 (84)	1.6 (7)		322	9.9(120)	265	13.6 (87)	191	86.9 (48)
8102	16	* *	* *		16	* *	10	* *	6	* *
8105	83	* *	* *		84	* *	31	* *	40	* *
8106	32	* *	* *		33	* *	30	* *	18	* *
8150	14	* *	* *		16	* *	14	* *	12	* *
8151	1,106	82.5 (57)	8.3 (66)		1,228	35.5 (79)	1,168	18.0 (68)	1,022	87.1 (47)
8152	145	86.2 (37)	2.8 (19)		167	38.3 (76)	155	5.2 (111)	149	82.6 (78)
8153	13	* *	* *		14	* *	12	* *	10	* *
8154	112	97.3 (2)	0.9 (3)		125	33.6 (85)	112	91.1 (2)	95	* *
8155	21	* *	* *		21	* *	20	* *	20	* *
8156	5	* *	* *		6	* *	6	* *	5	* *
8157	2	* *	* *		2	* *	2	* *	1	* *
8201	1,083	88.2 (27)	3.8 (28)		1,200	22.5(109)	761	18.5 (66)	909	89.1 (32)
8202	1,211	79.5 (69)	14.9 (86)		1,366	42.7 (56)	1,164	6.4 (109)	1,206	77.5 (94)
8203	139	94.2 (7)	0.7 (2)		165	24.8(103)	134	17.9 (69)	131	93.9 (6)
8250	158	86.7 (34)	1.9 (9)		376	25.5(102)	121	42.1 (5)	343	90.4 (20)
8252	33	* *	* *		38	* *	37	* *	31	* *
8255	1,150	86.5 (35)	3.9 (29)		1,291	21.1(111)	795	20.4 (52)	985	90.6 (18)
8257	166	97.0 (3)	1.2 (5)		185	37.3 (78)	155	0.6 (114)	150	80.7 (84)
8258	433	90.5 (15)	3.2 (22)		484	24.0(105)	434	17.5 (72)	411	92.0 (11)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8260	294	79.9 (68)	7.8 (61)	312	26.6(100)	169	17.8 (70)	233	82.8 (77)
8901	425	77.4 (75)	1.9 (9)	450	23.1(108)	193	9.8 (103)	363	84.6 (66)
8902	351	82.3 (58)	2.0 (12)	372	24.5(104)	235	11.5 (95)	291	84.9 (63)
8903	493	75.9 (83)	2.0 (12)	512	20.9(112)	330	2.7 (112)	409	87.3 (46)
8904	479	82.7 (56)	1.9 (9)	513	23.2(106)	273	11.7 (94)	405	84.9 (63)
8905	1,045	85.5 (40)	4.6 (37)	1,099	33.5 (86)	761	11.0 (99)	922	79.6 (90)
9012	17	* *	* *	18	* *	17	* *	16	* *
9013	42	* *	* *	48	* *	44	* *	40	* *
9015	19	* *	* *	19	* *	20	* *	15	* *
9016	74	* *	* *	80	* *	79	* *	70	* *
9017	28	* *	* *	30	* *	30	* *	25	* *
9018	48	* *	* *	52	* *	51	* *	40	* *
9019	22	* *	* *	23	* *	22	* *	20	* *
9021	37	* *	* *	43	* *	42	* *	39	* *
9041	12	* *	* *	14	* *	14	* *	14	* *
9045	543	71.8(105)	10.5 (74)	568	31.3 (94)	289	19.7 (61)	476	79.6 (90)
9050	509	74.1 (96)	5.3 (47)	543	31.5 (93)	252	59.9 (4)	449	73.7 (105)
9051	614	81.8 (60)	6.0 (52)	681	47.1 (35)	285	25.6 (35)	611	64.5 (113)
9053	961	76.7 (79)	3.9 (29)	1,024	18.5(115)	776	14.9 (81)	796	71.9 (112)
9058	284	98.2 (1)	0.4 (1)	288	40.6 (67)	266	0.8 (113)	250	81.2 (83)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9066	677	76.8 (78)	1.6 (7)	706	13.2(119)	455	13.4 (88)	543	87.7 (41)
9068	443	74.0 (97)	3.2 (22)	471	20.2(114)	81	* *	356	88.8 (34)
9069	819	80.0 (67)	2.8 (19)	865	17.1(116)	783	12.1 (92)	699	89.7 (27)
9070	558	77.4 (75)	3.2 (22)	607	23.2(106)	303	19.8 (58)	516	75.0 (101)
9072	539	76.3 (81)	1.5 (6)	574	15.0(118)	323	11.1 (97)	433	90.8 (16)
9074	331	74.3 (94)	2.4 (14)	351	16.2(117)	322	14.3 (83)	269	85.9 (54)
9080	1,002	83.6 (51)	5.0 (43)	1,064	29.3 (97)	902	7.4 (107)	862	72.9 (109)
Unknown Clinics	84	* *	* *	112	22.3(110)	65	* *	39	* *
Michigan	60,703	81.1	9.2	67,137	36.7	51,249	18.8	55,764	83.4
Nation (Prior Year)	429,770	83.1	4.1	1,074,863	36.0	380,154	19.2	418,448	85.0

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	35	* *	40	* *	16	* *	33	* *	40	* *
0102	183	51.9 (109)	186	32.3 (112)	82	* *	167	21.0 (80)	186	6.5 (83)
0103	173	50.3 (107)	174	30.5 (110)	71	* *	158	27.8 (108)	174	6.9 (86)
0104	48	* *	45	* *	21	* *	45	* *	45	* *
0250	53	* *	54	* *	27	* *	50	* *	54	* *
0251	32	* *	28	* *	15	* *	32	* *	28	* *
0252	39	* *	36	* *	18	* *	36	* *	36	* *
0253	58	* *	58	* *	21	* *	57	* *	59	* *
0401	196	53.6 (112)	186	30.6 (111)	91	* *	178	16.9 (56)	187	2.7 (13)
0402	143	53.8 (113)	133	33.1 (113)	62	* *	130	20.8 (79)	137	4.4 (43)
0403	52	* *	50	* *	18	* *	41	* *	50	* *
0404	54	* *	52	* *	20	* *	48	* *	52	* *
0501	226	45.6 (95)	220	29.1 (106)	87	* *	200	26.5 (106)	220	11.8 (109)
0503	69	* *	69	* *	28	* *	60	* *	69	* *
0504	204	40.2 (76)	199	24.6 (83)	71	* *	179	15.6 (45)	199	5.5 (62)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0505	158	46.8 (101)	150	25.3 (89)	65	* *	140	23.6 (93)	149	6.0 (70)
0506	69	* *	73	* *	31	* *	64	* *	73	* *
0650	404	46.0 (98)	393	24.9 (84)	169	38.5 (30)	369	23.3 (92)	394	6.1 (74)
0651	243	59.3 (116)	251	38.6 (117)	121	29.8 (50)	220	29.1 (110)	250	9.6 (105)
0652	115	43.5 (87)	119	27.7 (97)	43	* *	104	25.0 (99)	118	8.5 (99)
0653	133	52.6 (111)	131	34.4 (115)	61	* *	123	32.5 (113)	131	17.6 (116)
0654	147	38.8 (70)	147	25.2 (87)	50	* *	129	31.8 (112)	147	12.9 (113)
0655	170	59.4 (117)	165	35.8 (116)	82	* *	151	23.2 (91)	165	6.1 (74)
0657	27	* *	25	* *	10	* *	24	* *	25	* *
0701	155	36.1 (60)	153	19.6 (61)	45	* *	128	5.5 (4)	153	0.7 (2)
0702	35	* *	33	* *	13	* *	30	* *	33	* *
0703	91	* *	80	* *	45	* *	81	* *	80	* *
0704	29	* *	29	* *	12	* *	28	* *	29	* *
0705	58	* *	57	* *	23	* *	51	* *	57	* *
0850	440	35.0 (55)	423	19.1 (58)	121	39.7 (28)	378	16.1 (52)	423	6.9 (86)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0851	290	35.9 (59)	299	23.7 (78)	93	* *	258	14.0 (36)	302	3.6 (32)
0901	649	49.2 (105)	626	24.1 (81)	272	36.0 (36)	596	18.1 (62)	627	4.6 (47)
0902	26	* *	28	* *	14	* *	24	* *	28	* *
1051	590	39.8 (72)	573	17.8 (51)	195	44.1 (18)	514	11.3 (19)	575	5.6 (65)
1101	728	31.9 (43)	719	17.8 (51)	189	36.0 (36)	637	18.4 (65)	719	5.4 (58)
1102	347	46.4 (99)	325	23.1 (76)	129	35.7 (39)	305	21.6 (86)	325	7.4 (91)
1106	44	* *	41	* *	14	* *	38	* *	41	* *
1250	393	37.2 (63)	373	22.8 (74)	121	27.3 (54)	357	21.3 (83)	373	4.6 (47)
1251	318	38.7 (69)	316	26.3 (91)	98	* *	275	18.9 (68)	316	5.1 (55)
1252	0	* *	1	* *	0	* *	0	* *	1	* *
1253	593	37.9 (66)	583	22.5 (72)	177	28.2 (52)	535	15.1 (42)	584	4.1 (38)
1350	882	37.8 (65)	857	18.8 (55)	275	33.5 (44)	796	19.0 (70)	861	4.8 (50)
1351	45	* *	47	* *	16	* *	36	* *	47	* *
1352	162	40.1 (74)	166	27.1 (95)	58	* *	147	24.5 (97)	166	6.0 (70)
1401	79	* *	73	* *	34	* *	72	* *	74	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1402	49	* *	48	* *	15	* *	44	* *	49	* *
1403	9	* *	9	* *	6	* *	9	* *	9	* *
1404	7	* *	3	* *	0	* *	6	* *	3	* *
1501	394	28.9 (33)	380	16.3 (45)	94	* *	362	12.4 (22)	385	5.5 (62)
1503	410	33.2 (50)	397	21.4 (70)	111	27.9 (53)	365	17.8 (60)	396	8.3 (98)
1504	948	26.1 (30)	896	11.0 (33)	203	48.8 (10)	832	5.6 (5)	898	1.9 (7)
1505	306	29.1 (34)	303	11.2 (34)	77	* *	262	6.1 (7)	301	3.7 (33)
1506	227	36.6 (61)	223	18.8 (55)	66	* *	196	12.8 (24)	226	2.2 (9)
1507	250	37.2 (63)	249	18.1 (54)	85	* *	227	8.8 (15)	254	5.5 (62)
1508	40	* *	40	* *	0	* *	36	* *	40	* *
1509	208	33.7 (51)	205	20.0 (64)	59	* *	198	8.1 (11)	207	4.3 (42)
1511	301	43.2 (85)	297	28.6 (102)	111	29.7 (51)	258	17.8 (60)	298	6.0 (70)
1512	128	29.7 (36)	125	14.4 (41)	31	* *	109	6.4 (8)	127	3.1 (21)
1701	277	54.5 (114)	270	29.3 (108)	135	39.3 (29)	266	15.8 (48)	273	5.1 (55)
1950	290	42.8 (83)	289	29.4 (109)	105	30.5 (49)	260	21.2 (81)	289	6.2 (78)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1952	54	* *	49	* *	15	* *	42	* *	50	* *
1953	97	* *	98	* *	40	* *	91	* *	98	* *
1954	246	42.7 (82)	240	28.3 (100)	88	* *	225	21.3 (83)	240	7.5 (93)
1955	51	* *	54	* *	15	* *	46	* *	54	* *
1957	213	38.0 (67)	218	18.8 (55)	67	* *	172	12.2 (21)	219	4.1 (38)
1959	22	* *	20	* *	11	* *	21	* *	20	* *
2150	276	43.8 (90)	258	27.9 (98)	95	* *	250	19.2 (72)	258	2.7 (13)
2151	97	* *	90	* *	41	* *	83	* *	91	* *
2251	177	52.0 (110)	171	22.8 (74)	74	* *	155	14.8 (41)	172	3.5 (30)
2252	81	* *	76	* *	34	* *	73	* *	76	* *
2551	146	35.6 (58)	126	13.5 (37)	46	* *	148	18.2 (63)	126	3.2 (24)
2552	8	* *	8	* *	1	* *	8	* *	8	* *
2554	41	* *	40	* *	12	* *	37	* *	40	* *
2555	2,763	33.9 (53)	2,831	17.8 (51)	756	41.4 (25)	2,247	18.3 (64)	2,837	3.7 (33)
3001	341	43.4 (86)	339	19.8 (63)	125	40.0 (27)	318	24.8 (98)	339	5.6 (65)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3002	86	* *	84	* *	28	* *	83	* *	84	* *
3003	2	* *	2	* *	0	* *	0	* *	2	* *
3004	42	* *	43	* *	3	* *	41	* *	43	* *
3007	6	* *	3	* *	0	* *	5	* *	3	* *
3009	8	* *	4	* *	0	* *	7	* *	4	* *
3010	1	* *	1	* *	1	* *	1	* *	1	* *
3201	203	45.3 (93)	195	29.2 (107)	78	* *	180	26.1 (105)	196	6.1 (74)
3301	1,612	34.4 (54)	1,550	17.0 (47)	441	40.1 (26)	1,414	16.8 (54)	1,553	4.8 (50)
3302	96	* *	87	* *	31	* *	92	* *	87	* *
3304	64	* *	61	* *	1	* *	56	* *	61	* *
3305	66	* *	63	* *	19	* *	56	* *	63	* *
3306	96	* *	91	* *	31	* *	80	* *	91	* *
3308	18	* *	16	* *	6	* *	17	* *	17	* *
3309	16	* *	18	* *	5	* *	14	* *	18	* *
3550	38	* *	35	* *	22	* *	37	* *	35	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3551	13	* *	12	* *	6	* *	13	* *	12	* *
3552	16	* *	16	* *	12	* *	15	* *	16	* *
3701	470	35.5 (57)	447	26.2 (90)	136	18.4 (58)	427	25.5 (104)	448	5.4 (58)
3805	1,216	44.1 (91)	1,173	23.8 (79)	449	37.6 (31)	1,116	18.9 (68)	1,176	4.8 (50)
3950	702	39.2 (71)	658	17.5 (50)	225	41.8 (23)	620	16.0 (50)	658	2.7 (13)
3952	1	* *	2	* *	0	* *	1	* *	2	* *
3954	13	* *	12	* *	2	* *	9	* *	12	* *
4001	498	42.0 (81)	464	21.3 (69)	164	34.8 (41)	460	20.0 (75)	464	7.1 (90)
4002	425	44.5 (92)	401	23.9 (80)	149	36.9 (34)	376	33.5 (115)	401	12.5 (112)
4103	38	* *	37	* *	16	* *	34	* *	37	* *
4106	693	31.3 (42)	681	15.7 (44)	177	41.8 (23)	636	13.2 (28)	681	3.5 (30)
4107	1,253	32.2 (45)	1,185	16.6 (46)	321	37.1 (33)	1,109	8.7 (14)	1,181	2.2 (9)
4109	384	41.9 (80)	382	20.2 (65)	138	36.2 (35)	337	13.6 (30)	382	3.1 (21)
4110	408	32.4 (47)	377	20.2 (65)	121	24.8 (57)	407	14.7 (39)	377	2.4 (11)
4112	221	3.6 (1)	227	1.8 (1)	6	* *	214	8.9 (16)	228	6.1 (74)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4113	317	17.0 (12)	291	5.2 (6)	43	* *	308	5.2 (2)	293	0.7 (2)
4120	747	30.1 (37)	734	14.0 (38)	176	42.6 (20)	642	6.4 (8)	734	1.6 (6)
4160	752	20.9 (22)	723	7.7 (17)	127	51.2 (6)	649	6.9 (10)	722	1.5 (5)
4351	211	41.2 (79)	208	28.4 (101)	75	* *	193	21.2 (81)	208	8.2 (97)
4352	67	* *	67	* *	36	* *	63	* *	67	* *
4353	171	40.9 (78)	167	25.1 (86)	61	* *	160	18.8 (67)	167	6.0 (70)
4354	292	45.5 (94)	283	29.0 (105)	106	25.5 (55)	259	14.3 (38)	283	4.2 (41)
4355	112	46.4 (99)	100	27.0 (94)	40	* *	102	18.6 (66)	100	7.0 (88)
4356	205	40.5 (77)	202	22.3 (71)	75	* *	189	15.3 (43)	202	4.0 (37)
4358	51	* *	52	* *	1	* *	49	* *	52	* *
4359	199	30.7 (40)	204	10.8 (31)	52	* *	179	17.3 (58)	204	3.4 (29)
4360	72	* *	69	* *	18	* *	68	* *	69	* *
4361	80	* *	78	* *	39	* *	68	* *	78	* *
4362	110	57.3 (115)	102	33.3 (114)	54	* *	104	20.2 (77)	101	5.0 (54)
4363	373	47.5 (103)	364	26.9 (93)	148	32.4 (46)	342	24.0 (94)	365	6.3 (80)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4401	509	39.9 (73)	496	23.4 (77)	172	35.5 (40)	446	20.2 (77)	498	6.8 (85)
4501	368	49.5 (106)	362	24.3 (82)	158	42.4 (22)	336	24.4 (96)	362	4.1 (38)
4502	1	* *	1	* *	1	* *	1	* *	1	* *
4601	652	29.1 (34)	644	21.1 (68)	154	25.3 (56)	514	12.8 (24)	565	6.4 (82)
4701	582	43.1 (84)	552	20.8 (67)	195	52.8 (5)	444	13.1 (26)	554	2.5 (12)
5050	655	35.3 (56)	463	15.1 (42)	127	42.5 (21)	566	15.4 (44)	463	3.2 (24)
5051	1,128	31.9 (43)	829	10.7 (30)	203	45.8 (16)	954	14.0 (36)	831	3.2 (24)
5052	1,078	25.6 (29)	743	10.4 (28)	157	47.8 (14)	965	12.4 (22)	741	2.7 (13)
5101	877	33.0 (49)	839	15.4 (43)	237	47.7 (15)	726	15.6 (45)	840	4.5 (45)
5202	263	50.6 (108)	253	28.9 (104)	114	36.0 (36)	237	16.0 (50)	254	2.0 (8)
5203	41	* *	39	* *	20	* *	39	* *	39	* *
5204	35	* *	35	* *	16	* *	32	* *	34	* *
5601	463	43.6 (88)	454	24.9 (84)	176	33.0 (45)	420	17.6 (59)	452	4.9 (53)
5801	764	45.7 (96)	727	27.2 (96)	280	33.6 (43)	679	13.8 (34)	727	7.0 (88)
5803	63	* *	60	* *	11	* *	51	* *	60	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5804	17	* *	17	* *	4	* *	14	* *	17	* *
6101	24	* *	24	* *	8	* *	21	* *	24	* *
6102	69	* *	69	* *	23	* *	59	* *	69	* *
6103	1,272	38.6 (68)	1,270	22.5 (72)	423	34.0 (42)	1,142	15.7 (47)	1,276	6.5 (83)
6104	273	43.6 (88)	258	25.2 (87)	91	* *	260	36.5 (116)	255	11.8 (109)
6105	19	* *	19	* *	7	* *	15	* *	19	* *
6106	9	* *	7	* *	5	* *	9	* *	7	* *
6301	1,899	27.2 (32)	1,824	10.8 (31)	410	48.5 (11)	1,624	13.1 (26)	1,825	2.9 (19)
6312	1,176	19.5 (16)	1,117	6.5 (8)	169	54.4 (4)	903	8.2 (12)	1,114	3.2 (24)
6316	689	25.5 (28)	648	9.6 (25)	142	45.8 (16)	545	10.5 (18)	645	5.4 (58)
6320	70	* *	61	* *	4	* *	56	* *	61	* *
6325	50	* *	48	* *	17	* *	37	* *	48	* *
6330	166	30.1 (37)	164	17.1 (49)	41	* *	137	16.8 (54)	163	3.1 (21)
6335	204	9.3 (5)	190	4.7 (4)	13	* *	181	5.0 (1)	190	1.1 (4)
6340	48	* *	50	* *	4	* *	38	* *	51	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6345	5	* *	4	* *	1	* *	4	* *	4	* *
6350	84	* *	81	* *	15	* *	73	* *	81	* *
7304	71	* *	74	* *	29	* *	63	* *	74	* *
7306	1,046	37.1 (62)	1,018	14.0 (38)	327	49.2 (8)	949	21.6 (86)	1,005	7.8 (96)
7308	60	* *	57	* *	23	* *	55	* *	58	* *
7310	25	* *	25	* *	8	* *	23	* *	24	* *
7401	1,013	49.1 (104)	953	26.5 (92)	401	37.2 (32)	940	25.0 (99)	957	7.6 (95)
7601	251	45.8 (97)	248	28.6 (102)	98	* *	226	19.9 (74)	246	4.5 (45)
7850	440	47.3 (102)	416	27.9 (98)	175	30.9 (48)	400	29.3 (111)	422	6.2 (78)
7852	33	* *	37	* *	13	* *	31	* *	37	* *
8101	295	20.3 (19)	321	8.7 (22)	40	* *	188	22.3 (90)	322	7.5 (93)
8102	15	* *	17	* *	2	* *	6	* *	17	* *
8105	83	* *	83	* *	7	* *	40	* *	83	* *
8106	28	* *	30	* *	8	* *	18	* *	32	* *
8150	15	* *	14	* *	1	* *	12	* *	14	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8151	1,183	30.4 (39)	1,107	13.0 (36)	258	48.4 (12)	1,021	16.5 (53)	1,105	5.1 (55)
8152	157	40.1 (74)	147	19.7 (62)	46	* *	149	28.2 (109)	147	5.4 (58)
8153	14	* *	13	* *	4	* *	10	* *	13	* *
8154	122	4.9 (2)	111	4.5 (3)	5	* *	98	* *	111	0.0 (1)
8155	21	* *	21	* *	2	* *	20	* *	20	* *
8156	6	* *	5	* *	1	* *	5	* *	5	* *
8157	2	* *	2	* *	0	* *	1	* *	2	* *
8201	1,166	32.8 (48)	1,085	14.0 (38)	275	50.9 (7)	906	13.9 (35)	1,091	3.7 (33)
8202	1,357	14.1 (7)	1,285	5.8 (7)	143	49.0 (9)	1,206	12.1 (20)	1,286	3.3 (28)
8203	155	31.0 (41)	140	19.3 (60)	35	* *	133	15.8 (48)	141	5.7 (67)
8250	361	16.9 (11)	157	7.0 (11)	15	* *	343	13.7 (32)	157	5.7 (67)
8252	37	* *	37	* *	7	* *	31	* *	37	* *
8255	1,232	32.3 (46)	1,157	17.0 (47)	284	43.7 (19)	983	13.4 (29)	1,159	3.7 (33)
8257	181	5.5 (3)	165	4.8 (5)	6	* *	150	6.0 (6)	162	7.4 (91)
8258	466	33.7 (51)	435	19.1 (58)	119	31.9 (47)	412	17.2 (57)	436	4.6 (47)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8260	293	15.7 (9)	304	8.2 (18)	38	* *	233	8.2 (12)	304	3.0 (20)
8901	418	20.8 (21)	433	7.2 (15)	69	* *	366	25.1 (101)	433	9.5 (104)
8902	354	20.3 (19)	353	7.1 (12)	52	* *	290	21.7 (88)	352	8.5 (99)
8903	474	18.1 (14)	491	10.4 (28)	64	* *	409	25.2 (102)	492	9.6 (105)
8904	494	21.1 (23)	483	9.9 (27)	78	* *	408	21.3 (83)	484	9.3 (103)
8905	1,056	16.2 (10)	1,049	8.5 (21)	141	48.2 (13)	924	13.6 (30)	1,050	5.9 (69)
9012	17	* *	17	* *	4	* *	16	* *	17	* *
9013	44	* *	42	* *	6	* *	40	* *	42	* *
9015	20	* *	19	* *	2	* *	15	* *	19	* *
9016	79	* *	75	* *	21	* *	70	* *	75	* *
9017	31	* *	28	* *	6	* *	25	* *	28	* *
9018	51	* *	48	* *	13	* *	40	* *	48	* *
9019	22	* *	22	* *	8	* *	20	* *	22	* *
9021	42	* *	37	* *	12	* *	38	* *	37	* *
9041	14	* *	12	* *	6	* *	14	* *	12	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9045	539	17.4 (13)	547	7.1 (12)	76	* *	477	14.7 (39)	547	10.4 (107)
9050	504	19.6 (17)	509	8.4 (20)	79	* *	460	20.0 (75)	509	12.2 (111)
9051	663	10.3 (6)	636	6.8 (10)	56	* *	624	5.4 (3)	635	2.8 (17)
9053	948	27.1 (31)	962	9.6 (25)	198	60.1 (2)	797	32.5 (113)	962	18.7 (117)
9058	281	8.5 (4)	285	4.2 (2)	19	* *	250	10.0 (17)	285	2.8 (17)
9066	643	14.3 (8)	682	6.5 (8)	69	* *	543	19.0 (70)	680	9.1 (102)
9068	431	18.6 (15)	443	9.3 (24)	51	* *	356	25.3 (103)	444	13.7 (115)
9069	821	24.5 (27)	823	8.7 (22)	155	55.5 (3)	697	24.1 (95)	821	8.8 (101)
9070	573	19.9 (18)	558	7.2 (15)	94	* *	521	21.7 (88)	558	10.8 (108)
9072	532	22.9 (25)	538	11.9 (35)	84	* *	436	26.6 (107)	535	13.3 (114)
9074	334	24.3 (26)	331	8.2 (18)	55	* *	268	19.8 (73)	331	6.3 (80)
9080	1,008	21.8 (24)	1,010	7.1 (12)	175	61.1 (1)	863	13.7 (32)	1,012	4.4 (43)
Unknown Clinics	69	* *	73	* *	9	* *	38	* *	73	* *
Michigan	64,049	32.8	61,427	16.9	16,751	39.5	55,787	16.8	61,403	5.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
Nation (Prior Year)	814,631	23.0	715,034	12.1	137,883	38.1	660,570	16.3	562,126	15.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	Number	% (Rank)						
0101	37	* * * *	* * * *		36	* *	29	* *	38	* *
0102	169	3.6 (3)	10.1 (100)		176	7.4 (10)	156	1.9 (10)	173	70.5 (33)
0103	169	5.9 (14)	7.1 (51)		169	10.1 (42)	151	1.3 (5)	169	61.5 (70)
0104	40	* * * *	* * * *		41	* *	35	* *	41	* *
0250	52	* * * *	* * * *		50	* *	46	* *	52	* *
0251	27	* * * *	* * * *		28	* *	24	* *	27	* *
0252	31	* * * *	* * * *		34	* *	24	* *	31	* *
0253	55	* * * *	* * * *		54	* *	49	* *	54	* *
0401	176	5.7 (9)	14.2 (114)		180	7.2 (8)	162	2.5 (22)	176	53.4 (89)
0402	131	8.4 (72)	5.3 (20)		130	8.5 (23)	116	5.2 (99)	132	65.2 (53)
0403	43	* * * *	* * * *		45	* *	39	* *	43	* *
0404	48	* * * *	* * * *		51	* *	44	* *	48	* *
0501	195	13.3(112)	6.2 (33)		200	13.0 (89)	165	6.1(109)	199	79.4 (8)
0503	66	* * * *	* * * *		67	* *	59	* *	66	* *

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)	Number						
0504	175	8.0 (63)	10.3 (101)		192	12.5 (83)	153	2.6 (29)	188	80.9 (6)
0505	138	5.8 (11)	9.4 (91)		146	8.9 (27)	127	2.4 (19)	143	83.9 (4)
0506	60	* * *	* *		67	* *	59	* *	64	* *
0650	366	6.8 (33)	5.5 (23)		376	10.9 (50)	325	2.5 (22)	364	77.5 (11)
0651	220	8.2 (68)	7.7 (67)		228	13.2 (92)	187	2.7 (34)	219	61.6 (69)
0652	105	7.6 (54)	11.4 (106)		104	6.7 (5)	95	* *	105	62.9 (64)
0653	122	6.6 (26)	10.7 (103)		126	7.1 (7)	114	4.4 (89)	122	73.8 (18)
0654	131	6.9 (36)	7.6 (62)		134	8.2 (16)	119	3.4 (63)	130	73.8 (18)
0655	146	7.5 (52)	7.5 (58)		153	8.5 (23)	132	4.5 (91)	147	76.2 (13)
0657	21	* * *	* *		24	* *	19	* *	21	* *
0701	147	6.8 (33)	8.2 (77)		147	10.2 (44)	131	3.1 (52)	146	86.3 (1)
0702	30	* * *	* *		32	* *	27	* *	30	* *
0703	76	* * *	* *		77	* *	70	* *	76	* *
0704	27	* * *	* *		29	* *	24	* *	27	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0705	51	* * * *	*	53	* *	43	* *	51	* *
0850	397	7.1 (45)	9.8 (97)	409	9.5 (31)	356	2.5 (22)	401	75.1 (16)
0851	277	6.9 (36)	8.7 (83)	282	9.6 (33)	245	3.3 (60)	277	71.8 (27)
0901	595	7.7 (57)	9.7 (94)	602	12.5 (83)	516	2.5 (22)	594	60.1 (73)
0902	24	* * * *	*	25	* *	20	* *	24	* *
1051	526	7.6 (54)	10.6 (102)	549	10.2 (44)	469	3.0 (46)	525	81.1 (5)
1101	678	10.6 (90)	7.1 (51)	691	12.0 (73)	588	3.2 (57)	679	48.0 (98)
1102	307	8.1 (65)	9.8 (97)	309	8.4 (19)	274	4.0 (81)	307	58.0 (80)
1106	38	* * * *	*	39	* *	35	* *	38	* *
1250	348	6.9 (36)	8.9 (86)	359	10.9 (50)	306	2.3 (17)	348	58.9 (79)
1251	286	7.3 (49)	8.4 (81)	303	7.6 (11)	264	3.4 (63)	288	70.1 (35)
1252	1	* * * *	*	1	* *	1	* *	1	* *
1253	546	7.5 (52)	8.8 (85)	561	11.2 (59)	483	2.9 (39)	549	66.8 (50)
1350	820	8.3 (70)	8.3 (79)	848	11.1 (57)	732	3.1 (52)	819	59.1 (77)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
1351	44	* * * *		45	* *	37	* *	44	* *
1352	154	7.8 (59)	9.7 (94)	158	8.2 (16)	142	2.1 (14)	154	48.7 (97)
1401	69	* * * *		71	* *	59	* *	69	* *
1402	46	* * * *		48	* *	43	* *	46	* *
1403	7	* * * *		8	* *	7	* *	7	* *
1404	3	* * * *		3	* *	3	* *	3	* *
1501	345	5.8 (11)	7.8 (68)	362	11.0 (56)	306	2.6 (29)	349	72.2 (24)
1503	359	7.0 (42)	7.5 (58)	378	9.8 (37)	319	4.1 (82)	370	64.3 (58)
1504	830	6.1 (18)	6.9 (46)	869	10.9 (50)	733	2.0 (12)	864	73.1 (21)
1505	281	6.4 (22)	12.5 (111)	298	8.4 (19)	255	2.0 (12)	293	79.9 (7)
1506	210	6.7 (30)	8.1 (74)	217	12.9 (85)	182	1.1 (3)	210	75.2 (15)
1507	227	7.0 (42)	11.5 (107)	244	11.5 (65)	200	2.5 (22)	237	72.6 (23)
1508	37	* * * *		40	* *	34	* *	37	* *
1509	190	6.8 (33)	7.9 (69)	199	8.5 (23)	174	3.4 (63)	196	70.9 (32)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
1511	280	5.7 (9)	9.6 (92)	293	8.9 (27)	254	3.1 (52)	284	73.6 (20)
1512	122	6.6 (26)	12.3 (110)	122	13.9 (94)	103	1.9 (10)	124	62.1 (67)
1701	247	6.5 (24)	15.0 (115)	254	9.1 (30)	222	3.6 (70)	247	68.8 (45)
1950	259	6.6 (26)	6.2 (33)	266	7.9 (13)	236	3.0 (46)	259	71.8 (27)
1952	49	*	* * *	48	* *	41	* *	49	* *
1953	88	*	* * *	88	* *	79	* *	88	* *
1954	218	6.9 (36)	5.5 (23)	226	11.9 (70)	187	3.7 (73)	221	72.9 (22)
1955	49	*	* * *	52	* *	41	* *	50	* *
1957	202	7.9 (62)	7.4 (57)	205	10.2 (44)	179	2.8 (36)	202	84.7 (3)
1959	20	*	* * *	20	* *	18	* *	20	* *
2150	241	8.3 (70)	8.7 (83)	245	13.1 (90)	205	2.9 (39)	240	65.8 (52)
2151	83	*	* * *	88	* *	76	* *	83	* *
2251	158	4.4 (6)	8.9 (86)	163	8.0 (15)	145	1.4 (6)	158	69.6 (39)
2252	71	*	* * *	69	* *	59	* *	71	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
2551	119	5.0 (7)	6.7 (41)	124	7.3 (9)	108	4.6 (93)	120	45.0 (101)	
2552	8	* *	* *	8	* *	8	* *	8	* *	
2554	38	* *	* *	39	* *	35	* *	38	* *	
2555	2,745	10.8 (97)	5.0 (14)	2,789	14.4 (96)	2,326	3.9 (75)	2,740	53.0 (91)	
3001	310	11.0 (99)	5.2 (18)	319	16.9 (113)	257	4.3 (88)	310	55.2 (85)	
3002	78	* *	* *	79	* *	73	* *	78	* *	
3003	2	* *	* *	2	* *	2	* *	2	* *	
3004	40	* *	* *	41	* *	35	* *	40	* *	
3007	1	* *	* *	1	* *	1	* *	1	* *	
3009	2	* *	* *	2	* *	2	* *	2	* *	
3010	1	* *	* *	1	* *	1	* *	1	* *	
3201	188	9.0 (81)	12.2 (109)	190	12.1 (77)	165	3.6 (70)	187	55.6 (84)	
3301	1,469	7.8 (59)	7.1 (51)	1,512	10.5 (47)	1,308	3.0 (46)	1,477	68.9 (43)	
3302	85	* *	* *	86	* *	78	* *	85	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
3304	55	* * * *	* * * *	56	* *	51	* *	55	* *	
3305	63	* * * *	* * * *	62	* *	56	* *	63	* *	
3306	88	* * * *	* * * *	89	* *	81	* *	88	* *	
3308	15	* * * *	* * * *	16	* *	14	* *	15	* *	
3309	18	* * * *	* * * *	18	* *	15	* *	18	* *	
3550	31	* * * *	* * * *	32	* *	27	* *	31	* *	
3551	11	* * * *	* * * *	11	* *	10	* *	11	* *	
3552	14	* * * *	* * * *	13	* *	12	* *	14	* *	
3701	423	6.1 (18)	7.6 (62)	427	9.8 (37)	380	2.9 (39)	424	69.1 (42)	
3805	1,134	8.9 (77)	8.1 (74)	1,143	12.9 (85)	978	3.0 (46)	1,132	63.9 (59)	
3950	635	6.9 (36)	9.3 (89)	637	10.0 (39)	560	2.9 (39)	638	69.9 (37)	
3952	2	* * * *	* * * *	2	* *	2	* *	2	* *	
3954	11	* * * *	* * * *	12	* *	10	* *	12	* *	
4001	411	10.7 (94)	6.3 (37)	443	15.3 (102)	343	2.9 (39)	422	62.6 (65)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
4002	373	8.0 (63)	6.7 (41)	385	11.2 (59)	327	2.4 (19)	376	66.0 (51)
4103	35	* * *	* *	37	* *	31	* *	35	* *
4106	634	11.5(103)	7.3 (56)	647	14.5 (99)	528	4.9 (96)	634	63.4 (62)
4107	1,121	7.7 (57)	8.1 (74)	1,147	11.2 (59)	995	2.5 (22)	1,118	68.6 (46)
4109	360	8.1 (65)	8.6 (82)	368	10.9 (50)	321	2.2 (15)	360	70.0 (36)
4110	352	6.5 (24)	4.8 (12)	365	11.5 (65)	309	3.2 (57)	348	62.6 (65)
4112	208	3.8 (4)	8.2 (77)	210	9.5 (31)	186	2.2 (15)	208	28.4(114)
4113	273	2.9 (2)	12.5 (111)	284	14.1 (95)	234	0.9 (2)	272	68.0 (49)
4120	698	8.2 (68)	8.3 (79)	709	12.4 (80)	607	3.5 (68)	697	68.1 (48)
4160	687	9.5 (86)	9.0 (88)	703	11.9 (70)	596	4.2 (85)	693	69.6 (39)
4351	195	6.7 (30)	11.8 (108)	196	6.6 (3)	181	2.8 (36)	194	63.9 (59)
4352	62	* * *	* *	63	* *	58	* *	62	* *
4353	155	11.0 (99)	5.8 (29)	160	16.9 (113)	129	3.9 (75)	157	72.0 (25)
4354	263	4.2 (5)	11.0 (104)	269	10.8 (49)	233	1.7 (8)	263	71.9 (26)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)											
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)		Number	%	Number	%	Number	%	
	Number	% (Rank)	%	% (Rank)							Number
4355	97	* * * *	* * * *		96	* *		92	* *	97	* *
4356	183	5.5 (8)	6.0 (31)		188	11.2 (59)		162	2.5 (22)	183	71.0 (31)
4358	45	* * * *	* * * *		45	* *		39	* *	45	* *
4359	187	5.9 (14)	13.4 (113)		192	8.9 (27)		170	1.2 (4)	188	79.3 (9)
4360	66	* * * *	* * * *		68	* *		59	* *	66	* *
4361	73	* * * *	* * * *		75	* *		65	* *	73	* *
4362	96	* * * *	* * * *		99	* *		87	* *	96	* *
4363	342	5.8 (11)	9.9 (99)		342	8.5 (23)		306	2.9 (39)	342	74.0 (17)
4401	473	7.6 (54)	7.6 (62)		479	11.1 (57)		416	2.6 (29)	474	68.4 (47)
4501	340	9.1 (82)	7.9 (69)		345	11.9 (70)		298	3.7 (73)	339	54.6 (87)
4502	1	* * * *	* * * *		1	* *		1	* *	1	* *
4601	608	7.2 (46)	6.4 (38)		624	8.2 (16)		551	3.4 (63)	613	69.3 (41)
4701	516	6.2 (21)	11.2 (105)		536	9.7 (36)		464	1.5 (7)	516	71.7 (29)
5050	458	12.2(107)	6.8 (44)		457	16.0 (107)		380	4.2 (85)	459	65.1 (54)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
5051	810	8.9 (77)	6.8 (44)	811	12.9 (85)	694	4.2 (85)	812	64.8 (55)
5052	715	9.2 (85)	5.6 (27)	728	12.9 (85)	620	3.9 (75)	720	59.9 (75)
5101	805	7.8 (59)	5.1 (17)	818	12.0 (73)	697	3.3 (60)	815	70.3 (34)
5202	247	6.1 (18)	9.3 (89)	247	10.1 (42)	221	2.7 (34)	247	78.5 (10)
5203	34	* * *	* *	36	* *	28	* *	34	* *
5204	33	* * *	* *	34	* *	32	* *	33	* *
5601	431	7.4 (50)	9.7 (94)	443	7.0 (6)	398	3.0 (46)	431	75.6 (14)
5801	700	8.9 (77)	6.9 (46)	709	10.7 (48)	620	4.5 (91)	699	64.4 (57)
5803	58	* * *	* *	60	* *	53	* *	59	* *
5804	16	* * *	* *	17	* *	15	* *	16	* *
6101	24	* * *	* *	24	* *	22	* *	24	* *
6102	66	* * *	* *	67	* *	60	* *	65	* *
6103	1,206	8.6 (75)	6.7 (41)	1,244	12.2 (78)	1,058	3.0 (46)	1,208	54.8 (86)
6104	244	10.7 (94)	3.3 (5)	250	12.4 (80)	209	7.2(113)	245	44.9(102)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
	Number	% (Rank)	% (Rank)	Number						
6105	19	* * * *	* * * *	19	* *	16	* *	19	* *	
6106	7	* * * *	* * * *	7	* *	7	* *	7	* *	
6301	1,773	9.1 (82)	5.4 (22)	1,804	10.9 (50)	1,576	4.1 (82)	1,774	53.2 (90)	
6312	1,081	9.7 (88)	6.6 (40)	1,103	11.6 (68)	952	2.6 (29)	1,086	61.1 (71)	
6316	629	8.9 (77)	7.6 (62)	639	11.7 (69)	553	3.1 (52)	628	60.0 (74)	
6320	54	* * * *	* * * *	58	* *	49	* *	57	* *	
6325	47	* * * *	* * * *	48	* *	43	* *	48	* *	
6330	150	12.7(109)	8.0 (73)	155	16.1 (108)	122	4.1 (82)	154	50.6 (96)	
6335	167	6.0 (16)	9.6 (92)	177	6.2 (2)	154	2.6 (29)	172	56.4 (83)	
6340	43	* * * *	* * * *	49	* *	37	* *	45	* *	
6345	3	* * * *	* * * *	4	* *	2	* *	3	* *	
6350	70	* * * *	* * * *	74	* *	57	* *	73	* *	
7304	70	* * * *	* * * *	71	* *	62	* *	70	* *	
7306	959	10.9 (98)	7.2 (54)	992	11.5 (65)	841	5.2 (99)	962	50.9 (95)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)							
7308	53	* * * *	* *	56	* *	49	* *	51	* *	
7310	25	* * * *	* *	25	* *	21	* *	25	* *	
7401	909	6.4 (22)	5.9 (30)	924	10.0 (39)	819	3.3 (60)	908	59.9 (75)	
7601	228	6.6 (26)	7.0 (49)	241	12.0 (73)	202	3.5 (68)	228	64.5 (56)	
7850	403	8.4 (72)	5.5 (23)	407	11.3 (63)	356	2.8 (36)	402	68.9 (43)	
7852	34	* * * *	* *	34	* *	27	* *	35	* *	
8101	306	19.6(115)	4.2 (10)	308	22.4 (115)	232	9.1(114)	306	44.4(104)	
8102	9	* * * *	* *	11	* *	6	* *	9	* *	
8105	74	* * * *	* *	80	* *	64	* *	76	* *	
8106	27	* * * *	* *	31	* *	19	* *	27	* *	
8150	12	* * * *	* *	14	* *	10	* *	13	* *	
8151	1,057	8.5 (74)	7.6 (62)	1,079	10.9 (50)	934	3.6 (70)	1,058	69.7 (38)	
8152	141	10.6 (90)	5.0 (14)	143	8.4 (19)	127	3.9 (75)	144	63.2 (63)	
8153	12	* * * *	* *	12	* *	12	* *	12	* *	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)							
8154	106	1.9 (1)	7.5 (58)		107	3.7 (1)	101	0.0 (1)	106	77.4 (12)
8155	20	* *	* *		21	* *	20	* *	20	* *
8156	5	* *	* *		5	* *	5	* *	5	* *
8157	1	* *	* *		1	* *	1	* *	1	* *
8201	1,044	7.4 (50)	6.2 (33)		1,081	7.9 (13)	952	4.9 (96)	1,054	57.6 (82)
8202	1,174	7.0 (42)	7.0 (49)		1,231	9.6 (33)	1,055	2.9 (39)	1,214	63.5 (61)
8203	130	6.9 (36)	6.9 (46)		140	10.0 (39)	116	1.7 (8)	130	60.8 (72)
8250	147	9.5 (86)	6.1 (32)		154	14.9 (101)	124	2.4 (19)	147	85.0 (2)
8252	34	* *	* *		36	* *	27	* *	35	* *
8255	1,134	7.2 (46)	7.5 (58)		1,138	11.4 (64)	991	2.3 (17)	1,139	54.0 (88)
8257	120	6.7 (30)	5.0 (14)		158	7.6 (11)	108	5.6 (105)	144	61.8 (68)
8258	416	6.0 (16)	7.2 (54)		427	6.6 (3)	382	3.1 (52)	423	52.2 (93)
8260	287	9.1 (82)	3.1 (2)		299	8.4 (19)	262	4.6 (93)	292	71.6 (30)
8901	416	11.5 (103)	2.6 (1)		411	15.3 (102)	339	5.6 (105)	417	30.7 (113)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8902	339	11.5(103)	3.2	(3)	347	14.4 (96)	287	5.9(108)	340	42.1(107)
8903	471	14.6(114)	3.6	(6)	472	16.5 (109)	380	6.1(109)	471	25.5(115)
8904	465	12.3(108)	6.2	(33)	474	16.7 (112)	385	5.7(107)	463	44.5(103)
8905	978	8.1 (65)	5.3	(20)	1,001	12.3 (79)	844	3.9 (75)	983	38.6(112)
9012	15	*	*	* *	15	* *	13	* *	15	* *
9013	41	*	*	* *	41	* *	34	* *	42	* *
9015	19	*	*	* *	18	* *	14	* *	19	* *
9016	73	*	*	* *	73	* *	62	* *	73	* *
9017	25	*	*	* *	27	* *	23	* *	26	* *
9018	47	*	*	* *	47	* *	41	* *	47	* *
9019	21	*	*	* *	22	* *	16	* *	21	* *
9021	34	*	*	* *	35	* *	28	* *	34	* *
9041	11	*	*	* *	12	* *	6	* *	11	* *
9045	509	10.6 (90)	7.9	(69)	525	12.0 (73)	440	3.9 (75)	516	43.6(105)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9050	483	8.7 (76)	6.4 (38)	497	14.5 (99)	407	3.4 (63)	488	39.8(109)
9051	582	7.2 (46)	7.9 (69)	609	13.1 (90)	499	3.2 (57)	605	47.6 (99)
9053	925	14.3(113)	3.8 (8)	946	15.8 (105)	777	6.6(111)	925	45.4(100)
9058	255	10.6 (90)	5.5 (23)	271	9.6 (33)	229	4.4 (89)	268	59.0 (78)
9066	658	11.7(106)	5.6 (27)	670	12.4 (80)	570	5.4(101)	658	40.9(108)
9068	420	9.8 (89)	3.6 (6)	432	14.4 (96)	356	5.1 (98)	426	43.4(106)
9069	795	13.0(110)	4.8 (12)	811	15.3 (102)	667	5.5(103)	800	52.5 (92)
9070	531	10.7 (94)	3.2 (3)	549	16.6 (110)	438	4.6 (93)	535	39.4(111)
9072	520	13.1(111)	4.0 (9)	525	16.6 (110)	425	6.6(111)	523	39.6(110)
9074	327	11.3(102)	5.2 (18)	329	15.8 (105)	276	5.4(101)	329	57.8 (81)
9080	971	11.1(101)	4.4 (11)	994	13.7 (93)	838	5.5(103)	969	51.2 (94)
Unknown Clinics	56	* * *	* *	69	* *	51	* *	58	* *
Michigan	58,040	8.6	7.1	59,523	11.7	50,780	3.6	58,402	60.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Nation (Prior Year)	880,933	8.0	6.7	948,286	10.7	749,107	3.4	871,811	68.2

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0101	152	95.4	1.3	1.3	0.7	0.0	0.7	0.7
0102	546	93.6	0.7	2.9	0.4	0.0	2.2	0.2
0103	523	96.0	0.2	1.7	0.0	0.2	1.9	0.0
0104	153	96.1	0.0	1.3	0.0	0.7	2.0	0.0
0250	185	82.7	0.5	0.5	0.5	0.0	14.6	1.1
0251	145	86.9	0.0	0.7	4.8	0.7	6.9	0.0
0252	150	55.3	2.7	1.3	1.3	3.3	36.0	0.0
0253	195	77.9	0.0	0.5	9.2	0.5	11.8	0.0
0401	645	94.1	0.5	0.8	0.0	1.1	2.8	0.8
0402	453	87.2	0.7	1.3	0.0	1.3	9.3	0.2
0403	162	98.1	0.0	0.6	1.2	0.0	0.0	0.0
0404	175	94.3	0.0	1.1	0.0	0.0	4.6	0.0
0501	694	94.4	0.3	2.2	0.0	0.3	2.9	0.0
0502	1	*	*	*	*	*	*	*
0503	220	91.8	0.0	1.8	1.4	0.5	4.1	0.5
0504	671	90.3	0.4	1.0	2.5	1.2	4.5	0.0
0505	465	83.2	0.6	3.2	0.2	0.0	12.7	0.0
0506	219	98.2	0.0	0.0	0.5	0.0	1.4	0.0
0650	1,320	89.0	3.9	2.4	1.2	0.8	2.3	0.4
0651	799	94.7	0.1	2.0	0.1	0.1	1.8	1.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0652	373	93.8	0.0	1.9	0.0	0.0	2.9	1.3
0653	501	97.8	0.0	1.2	0.0	0.2	0.8	0.0
0654	467	93.4	1.5	1.1	0.9	0.2	2.8	0.2
0655	482	92.9	0.0	0.8	0.0	0.2	4.4	1.7
0657	83	*	*	*	*	*	*	*
0701	527	89.6	0.9	1.3	0.0	5.9	1.7	0.6
0702	104	96.2	0.0	0.0	1.0	0.0	1.9	1.0
0703	310	93.5	0.0	1.6	1.9	0.3	1.3	1.3
0704	81	*	*	*	*	*	*	*
0705	192	99.0	0.0	0.5	0.0	0.0	0.5	0.0
0850	1,441	88.3	4.2	4.8	0.1	1.0	1.2	0.4
0851	1,060	95.2	0.4	2.9	0.1	0.2	0.6	0.7
0901	2,133	82.9	2.2	9.6	0.2	0.2	4.3	0.7
0902	90	*	*	*	*	*	*	*
1051	1,818	90.8	0.9	3.7	0.7	0.8	2.2	0.9
1101	2,369	37.3	53.9	4.5	0.2	1.3	1.5	1.4
1102	1,155	78.6	10.7	5.5	0.3	1.1	2.4	1.3
1106	151	84.8	1.3	10.6	0.0	0.0	1.3	2.0
1250	1,289	85.3	0.5	10.7	0.1	1.2	1.7	0.5
1251	1,070	93.9	0.5	2.1	0.0	0.1	1.4	2.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
1252	284	64.4	0.7	33.5	0.0	0.4	0.7	0.4
1253	1,581	78.0	3.4	14.7	0.0	0.1	2.9	0.9
1350	2,989	63.8	19.3	8.3	0.2	3.5	3.9	0.9
1351	162	93.2	0.6	2.5	0.0	0.6	2.5	0.6
1352	490	65.9	24.3	5.5	0.0	0.4	3.3	0.6
1401	278	88.8	0.7	6.5	0.7	0.0	2.2	1.1
1402	189	55.6	0.5	22.8	10.1	1.1	9.0	1.1
1403	24	*	*	*	*	*	*	*
1404	33	*	*	*	*	*	*	*
1501	1,413	46.4	6.7	39.1	0.4	0.1	1.9	5.3
1503	1,556	59.8	6.4	26.3	0.6	1.0	1.3	4.6
1504	3,110	47.4	4.3	39.6	0.1	5.3	1.3	2.0
1505	1,030	87.3	2.0	6.8	0.1	0.8	1.6	1.5
1506	747	81.3	1.1	12.6	0.3	0.3	2.5	2.0
1507	799	91.1	0.8	3.5	0.1	0.8	2.5	1.3
1508	213	0.9	0.0	84.5	0.0	0.0	0.0	14.6
1509	807	53.4	2.2	38.3	0.0	0.1	1.0	5.0
1510	48	*	*	*	*	*	*	*
1511	873	94.5	1.9	1.7	0.0	0.0	1.4	0.5
1512	368	91.8	0.3	4.1	0.3	0.0	3.0	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
1701	904	62.6	0.2	1.5	19.2	1.1	14.9	0.3
1712	28	*	*	*	*	*	*	*
1950	970	92.6	0.1	4.5	0.0	0.5	2.0	0.3
1952	171	93.6	0.0	4.1	0.0	0.6	1.2	0.6
1953	351	90.0	0.9	5.1	0.3	0.6	2.6	0.6
1954	792	89.5	0.0	9.6	0.1	0.3	0.3	0.3
1955	159	87.4	0.6	10.1	0.0	0.6	1.3	0.0
1957	667	90.7	1.3	5.2	0.3	0.6	1.0	0.7
1959	125	87.2	1.6	7.2	0.0	2.4	1.6	0.0
2150	821	90.0	0.0	1.5	1.2	0.1	6.7	0.5
2151	335	90.7	0.0	2.4	0.6	0.3	3.0	3.0
2251	548	89.6	0.2	2.6	0.0	0.4	5.3	2.0
2252	274	96.0	0.4	1.8	0.0	0.0	1.5	0.4
2551	2,562	33.2	63.5	1.7	0.0	0.2	1.3	0.0
2552	8	*	*	*	*	*	*	*
2554	118	85.6	5.9	4.2	0.0	0.8	2.5	0.8
2555	7,067	61.7	31.6	3.7	0.1	0.4	2.1	0.5
3001	1,165	14.7	67.7	15.3	0.0	1.0	1.2	0.1
3002	287	22.6	51.2	24.0	0.0	0.3	1.7	0.0
3003	7	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
3004	178	3.9	0.0	95.5	0.0	0.0	0.0	0.6
3007	34	*	*	*	*	*	*	*
3009	25	*	*	*	*	*	*	*
3010	9	*	*	*	*	*	*	*
3201	697	94.4	0.1	2.9	0.0	0.9	0.9	0.9
3301	5,250	46.7	28.1	15.5	0.2	4.4	4.2	0.8
3302	342	27.2	42.7	19.9	0.3	3.5	5.6	0.9
3304	165	43.0	16.4	5.5	0.0	30.9	2.4	1.8
3305	174	74.7	9.8	10.3	0.0	3.4	1.1	0.6
3306	287	92.3	0.0	3.1	0.3	0.0	2.1	2.1
3308	70	*	*	*	*	*	*	*
3309	42	*	*	*	*	*	*	*
3550	107	9.3	0.0	0.9	61.7	0.0	28.0	0.0
3551	31	*	*	*	*	*	*	*
3552	52	*	*	*	*	*	*	*
3701	1,386	90.8	0.4	6.9	0.0	0.5	1.4	0.0
3805	4,088	76.7	12.7	5.4	0.1	0.6	4.2	0.2
3950	2,490	68.7	17.5	6.8	0.1	1.6	4.6	0.7
3952	14	*	*	*	*	*	*	*
3953	5	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
3954	56	*	*	*	*	*	*	*
4001	1,479	31.9	54.8	8.9	0.3	1.0	2.7	0.4
4002	1,266	53.8	23.9	14.2	0.3	0.8	6.6	0.3
4103	107	13.1	71.0	6.5	0.9	0.9	7.5	0.0
4106	1,767	20.1	44.4	29.9	0.2	0.4	3.5	1.5
4107	3,974	55.9	24.1	13.2	0.3	2.1	4.2	0.3
4109	1,342	92.2	0.4	4.1	0.1	0.6	2.5	0.2
4110	1,713	40.2	9.3	44.7	0.5	0.8	4.4	0.2
4112	749	0.4	0.9	97.3	0.0	0.3	0.3	0.8
4113	870	10.5	20.7	65.9	0.0	1.4	1.6	0.0
4120	2,451	60.7	8.6	24.9	0.2	2.1	3.0	0.4
4160	2,335	45.2	31.2	11.1	0.1	8.1	4.1	0.2
4351	721	92.0	0.3	3.6	0.1	0.4	2.4	1.2
4352	203	75.9	15.3	3.0	0.5	0.5	3.9	1.0
4353	634	69.9	0.2	26.2	0.2	0.0	2.7	0.9
4354	970	89.5	4.0	2.3	0.0	0.6	2.7	0.9
4355	420	79.8	0.0	7.6	0.2	0.2	10.5	1.7
4356	697	85.4	0.6	8.8	0.6	0.4	3.4	0.9
4357	4	*	*	*	*	*	*	*
4358	247	2.8	0.0	93.5	0.0	0.0	0.0	3.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
4359	647	70.5	0.2	26.1	0.2	0.2	0.6	2.3
4360	212	93.9	0.0	4.7	0.5	0.5	0.0	0.5
4361	300	92.7	1.3	0.7	0.0	0.3	4.7	0.3
4362	393	95.2	0.0	2.0	0.3	0.0	2.0	0.5
4363	1,153	93.7	0.4	2.8	0.1	0.5	2.0	0.5
4401	1,643	89.7	0.4	4.5	0.0	0.3	4.4	0.7
4501	1,153	94.2	0.6	3.6	0.0	0.0	1.4	0.3
4502	1	*	*	*	*	*	*	*
4591	2	*	*	*	*	*	*	*
4601	2,014	79.3	1.2	15.4	0.0	0.1	1.9	1.9
4701	1,737	92.1	0.7	3.9	0.1	1.2	1.2	0.9
5050	1,888	67.3	26.0	1.6	0.1	1.4	3.2	0.5
5051	3,390	64.5	25.5	3.5	0.1	1.8	4.0	0.6
5052	3,057	63.8	23.6	2.1	0.1	7.7	2.3	0.5
5101	2,844	72.9	8.9	12.2	0.1	1.6	3.6	0.7
5202	817	90.2	0.2	1.3	1.2	0.6	5.8	0.6
5203	179	84.4	2.8	1.1	1.7	0.6	8.4	1.1
5204	163	87.1	3.7	1.8	0.6	0.0	6.7	0.0
5601	1,465	93.4	1.2	2.0	0.3	0.8	1.8	0.5
5801	2,574	87.1	4.8	4.7	0.2	0.3	1.9	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
5803	188	88.3	0.0	3.7	0.0	0.0	2.7	5.3
5804	49	*	*	*	*	*	*	*
6101	74	*	*	*	*	*	*	*
6102	207	90.3	0.5	4.8	0.5	0.0	3.9	0.0
6103	4,523	63.1	24.5	7.4	0.1	0.3	4.0	0.5
6104	542	34.7	48.7	10.5	0.0	0.6	5.2	0.4
6105	57	*	*	*	*	*	*	*
6106	45	*	*	*	*	*	*	*
6301	6,013	49.0	27.0	18.2	0.1	2.7	2.6	0.5
6312	3,541	40.2	52.7	1.7	0.1	2.2	2.4	0.6
6316	1,997	73.5	15.7	6.2	0.2	2.4	1.5	0.6
6320	259	71.0	22.0	1.5	0.0	3.9	1.5	0.0
6325	171	91.2	2.3	4.1	0.0	0.6	1.8	0.0
6330	550	30.5	53.1	14.2	0.2	0.2	1.8	0.0
6335	679	80.7	12.7	2.1	0.3	3.4	0.9	0.0
6340	153	41.2	23.5	28.8	0.0	3.9	2.6	0.0
6345	34	*	*	*	*	*	*	*
6350	324	59.0	35.5	2.5	0.0	0.6	2.5	0.0
7304	261	92.3	0.8	5.4	0.0	0.8	0.8	0.0
7306	3,247	50.7	32.4	13.7	0.1	0.6	1.8	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
7308	223	96.4	0.0	2.2	0.0	0.0	1.3	0.0
7310	75	*	*	*	*	*	*	*
7401	3,019	86.1	4.4	4.8	0.0	0.4	3.7	0.7
7402	65	*	*	*	*	*	*	*
7403	33	*	*	*	*	*	*	*
7404	31	*	*	*	*	*	*	*
7601	874	92.6	0.0	5.5	0.1	0.2	0.7	0.9
7850	1,305	91.8	0.2	4.7	0.0	0.0	2.5	0.8
7851	83	*	*	*	*	*	*	*
7852	106	98.1	0.0	0.0	0.0	0.0	1.9	0.0
8101	342	12.9	82.7	2.0	0.0	0.6	1.8	0.0
8102	18	*	*	*	*	*	*	*
8105	86	*	*	*	*	*	*	*
8106	33	*	*	*	*	*	*	*
8150	58	*	*	*	*	*	*	*
8151	3,760	43.4	38.5	9.5	0.5	2.6	5.1	0.3
8152	525	39.4	48.0	3.2	0.2	0.8	8.2	0.2
8153	29	*	*	*	*	*	*	*
8154	359	33.1	20.1	11.1	0.0	35.1	0.6	0.0
8155	73	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
8156	23	*	*	*	*	*	*	*
8157	26	*	*	*	*	*	*	*
8201	3,378	60.2	27.0	4.5	0.2	2.2	5.6	0.2
8202	3,983	85.6	9.0	3.4	0.1	1.0	0.8	0.2
8203	491	68.4	25.1	3.1	0.0	0.2	2.4	0.8
8250	1,090	36.5	54.4	3.9	0.3	2.0	2.8	0.0
8252	45	*	*	*	*	*	*	*
8255	3,377	67.9	21.7	6.5	0.1	1.6	1.9	0.3
8257	616	47.6	23.5	0.5	0.0	27.1	1.3	0.0
8258	1,580	66.6	23.9	5.3	0.1	0.9	2.2	0.9
8260	764	48.8	31.3	1.3	0.1	16.8	1.7	0.0
8901	1,298	1.3	95.4	0.6	0.0	1.0	1.7	0.0
8902	912	3.8	93.9	0.7	0.0	0.1	1.4	0.1
8903	1,547	1.9	96.1	0.8	0.0	0.3	1.0	0.0
8904	1,337	3.7	87.1	0.7	0.2	6.2	2.0	0.1
8905	3,341	39.2	29.8	24.3	0.1	3.8	2.6	0.3
9004	4	*	*	*	*	*	*	*
9012	48	*	*	*	*	*	*	*
9013	118	1.7	90.7	0.0	0.0	0.8	6.8	0.0
9015	39	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
9016	183	0.0	96.7	0.5	0.0	0.0	2.7	0.0
9017	67	*	*	*	*	*	*	*
9018	141	2.8	95.7	0.7	0.0	0.0	0.7	0.0
9019	51	*	*	*	*	*	*	*
9021	117	1.7	94.0	1.7	0.0	0.9	1.7	0.0
9040	8	*	*	*	*	*	*	*
9041	234	3.4	94.9	0.9	0.0	0.0	0.9	0.0
9045	1,639	9.8	43.0	45.4	0.1	0.3	1.2	0.3
9050	2,086	12.7	54.9	30.7	0.0	0.8	0.6	0.2
9051	2,406	9.5	12.9	76.4	0.1	0.1	1.0	0.0
9053	3,115	2.0	95.5	0.9	0.0	0.9	0.6	0.1
9055	55	*	*	*	*	*	*	*
9058	846	63.9	29.7	0.5	0.1	5.2	0.6	0.0
9066	2,478	1.4	97.6	0.2	0.1	0.2	0.4	0.1
9068	1,613	1.7	96.7	0.6	0.1	0.4	0.4	0.2
9069	3,040	2.1	96.1	0.3	0.0	0.1	1.3	0.1
9070	2,271	1.8	88.2	0.2	0.1	8.1	1.5	0.1
9072	2,006	2.8	94.1	0.2	0.2	1.9	0.6	0.1
9074	1,585	2.5	95.1	0.6	0.1	0.3	1.4	0.1
9080	3,025	22.1	73.8	2.0	0.0	0.2	1.7	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Unknown Clinics	242	26.4	16.1	14.0	10.7	4.5	19.0	9.1
Michigan	206,241	56.9	27.4	10.3	0.3	1.7	2.7	0.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	152	0.0	3.3	9.9	59.9	25.7	0.7	0.7
0102	546	0.2	5.1	11.9	67.4	13.4	2.0	0.0
0103	523	0.0	5.4	13.6	64.4	15.5	1.1	0.0
0104	153	0.0	8.5	11.1	66.0	13.1	1.3	0.0
0250	185	0.0	5.4	11.9	69.7	11.9	1.1	0.0
0251	145	0.0	5.5	12.4	62.8	17.2	2.1	0.0
0252	150	0.0	2.0	11.3	61.3	24.7	0.7	0.0
0253	195	0.0	6.2	14.4	56.4	21.0	2.1	0.0
0401	645	0.3	4.5	14.9	65.0	15.0	0.3	0.0
0402	453	0.2	2.9	13.9	59.2	23.2	0.7	0.0
0403	162	0.0	6.2	9.3	63.6	19.8	1.2	0.0
0404	175	0.0	4.6	11.4	62.3	20.0	1.7	0.0
0501	694	0.1	5.2	12.0	65.6	16.0	1.0	0.1
0502	1	*	*	*	*	*	*	*
0503	220	0.5	2.7	15.0	61.4	19.5	0.9	0.0
0504	671	0.1	3.0	13.3	63.9	18.5	1.2	0.0
0505	465	0.0	3.7	11.4	66.0	17.4	1.5	0.0
0506	219	0.0	12.3	13.2	58.0	15.1	1.4	0.0
0650	1,320	0.1	5.8	12.3	64.6	16.1	0.8	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0651	799	0.1	4.6	14.5	67.1	12.1	1.4	0.1
0652	373	0.3	4.6	11.5	64.1	16.9	2.4	0.3
0653	501	0.8	5.2	12.8	65.1	14.2	1.4	0.6
0654	467	0.2	7.3	14.6	60.0	15.4	2.6	0.0
0655	482	0.2	5.0	12.7	65.4	16.0	0.8	0.0
0657	83	*	*	*	*	*	*	*
0701	527	0.0	2.8	8.5	64.9	21.8	1.9	0.0
0702	104	0.0	3.8	9.6	70.2	15.4	1.0	0.0
0703	310	0.0	4.2	12.6	64.8	16.8	1.6	0.0
0704	81	*	*	*	*	*	*	*
0705	192	0.5	4.7	7.8	56.8	27.1	3.1	0.0
0850	1,441	0.1	5.1	11.2	66.5	16.3	0.8	0.0
0851	1,060	0.1	4.7	13.8	63.4	16.7	1.3	0.0
0901	2,133	0.1	5.4	12.9	61.7	18.4	1.4	0.0
0902	90	*	*	*	*	*	*	*
1051	1,818	0.3	4.8	10.8	62.6	19.6	1.4	0.5
1101	2,369	0.3	8.7	14.5	59.5	15.7	1.4	0.1
1102	1,155	0.4	5.6	14.7	61.0	16.8	1.3	0.1
1106	151	0.7	5.3	11.3	60.3	21.2	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1250	1,289	0.2	6.0	11.2	66.6	14.9	1.1	0.1
1251	1,070	0.2	5.5	13.3	66.4	13.6	1.0	0.0
1252	284	0.4	4.2	11.6	61.3	20.8	1.8	0.0
1253	1,581	0.4	5.4	12.4	64.1	16.7	1.1	0.0
1350	2,989	0.3	6.3	14.1	63.3	14.9	0.9	0.2
1351	162	0.0	3.7	11.7	70.4	14.2	0.0	0.0
1352	490	0.0	8.2	13.9	65.5	11.4	1.0	0.0
1401	278	0.4	9.7	8.3	60.8	18.3	1.4	1.1
1402	189	0.0	3.2	10.6	57.7	24.9	3.7	0.0
1403	24	*	*	*	*	*	*	*
1404	33	*	*	*	*	*	*	*
1501	1,413	0.3	6.7	13.3	61.1	17.0	1.6	0.0
1503	1,556	0.1	6.8	12.9	59.7	19.2	1.0	0.4
1504	3,110	0.3	6.4	10.8	61.1	20.2	1.2	0.0
1505	1,030	0.1	4.0	11.0	60.3	23.6	1.1	0.0
1506	747	0.4	5.8	12.4	63.5	17.3	0.7	0.0
1507	799	0.1	3.1	10.5	66.0	19.3	1.0	0.0
1508	213	0.5	8.0	12.2	53.1	23.5	2.8	0.0
1509	807	0.4	8.2	13.1	56.8	18.7	2.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1510	48	*	*	*	*	*	*	*
1511	873	0.1	4.4	13.5	66.0	14.8	1.3	0.0
1512	368	0.0	3.8	12.5	62.8	19.8	0.8	0.3
1701	904	0.0	4.3	12.4	66.0	16.3	1.0	0.0
1712	28	*	*	*	*	*	*	*
1950	970	0.1	6.7	17.6	62.6	11.8	0.6	0.6
1952	171	0.0	4.7	11.1	66.1	16.4	1.8	0.0
1953	351	0.3	4.6	13.7	67.5	12.8	1.1	0.0
1954	792	0.1	4.2	13.3	67.6	14.0	0.8	0.1
1955	159	0.0	5.0	13.2	69.2	11.9	0.6	0.0
1957	667	0.0	3.6	11.4	63.9	19.5	1.5	0.1
1959	125	0.0	1.6	10.4	69.6	16.8	1.6	0.0
2150	821	0.0	3.7	11.2	66.3	17.9	1.0	0.0
2151	335	0.3	4.2	9.6	66.3	18.2	1.2	0.3
2251	548	0.4	4.6	11.9	64.8	17.5	0.9	0.0
2252	274	1.1	5.8	10.2	65.0	17.9	0.0	0.0
2551	2,562	0.5	9.5	18.5	57.3	12.9	1.2	0.1
2552	8	*	*	*	*	*	*	*
2554	118	1.7	7.6	17.8	59.3	11.9	1.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2555	7,067	0.2	5.7	13.1	62.4	17.4	1.0	0.1
3001	1,165	0.9	10.5	16.3	57.2	13.6	1.3	0.3
3002	287	0.7	9.8	16.4	56.4	16.7	0.0	0.0
3003	7	*	*	*	*	*	*	*
3004	178	0.0	3.4	9.0	56.2	29.8	1.7	0.0
3007	34	*	*	*	*	*	*	*
3009	25	*	*	*	*	*	*	*
3010	9	*	*	*	*	*	*	*
3201	697	0.3	1.9	10.8	63.8	21.4	1.9	0.0
3301	5,250	0.3	5.7	12.6	62.9	17.2	1.1	0.1
3302	342	0.3	4.7	12.0	63.5	18.4	1.2	0.0
3304	165	0.0	0.6	4.2	52.7	39.4	2.4	0.6
3305	174	0.0	2.9	9.2	61.5	22.4	3.4	0.6
3306	287	0.0	4.5	10.8	61.3	22.6	0.7	0.0
3308	70	*	*	*	*	*	*	*
3309	42	*	*	*	*	*	*	*
3550	107	0.0	4.7	15.9	66.4	12.1	0.9	0.0
3551	31	*	*	*	*	*	*	*
3552	52	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3701	1,386	0.2	5.7	13.5	67.0	12.3	1.2	0.1
3805	4,088	0.3	6.0	14.3	63.1	14.5	1.0	0.7
3950	2,490	0.3	5.5	12.0	63.5	17.9	0.8	0.1
3952	14	*	*	*	*	*	*	*
3953	5	*	*	*	*	*	*	*
3954	56	*	*	*	*	*	*	*
4001	1,479	0.4	10.1	14.6	58.2	14.9	1.8	0.0
4002	1,266	0.2	6.6	12.6	64.0	15.5	1.1	0.0
4103	107	0.0	10.3	14.0	62.6	13.1	0.0	0.0
4106	1,767	0.4	6.0	14.3	59.0	18.2	1.6	0.5
4107	3,974	0.3	6.1	12.2	60.9	18.4	1.1	1.0
4109	1,342	0.3	5.4	10.4	62.4	20.4	1.0	0.1
4110	1,713	0.4	5.9	13.6	56.5	22.5	1.1	0.1
4112	749	0.1	4.9	6.1	56.5	29.1	2.9	0.3
4113	870	0.2	6.3	9.8	47.8	20.3	1.4	14.1
4120	2,451	0.3	5.0	10.9	62.8	19.7	1.1	0.1
4160	2,335	0.2	3.8	10.9	61.5	22.0	1.5	0.1
4351	721	0.3	7.4	14.7	59.5	17.5	0.7	0.0
4352	203	0.0	6.9	15.8	60.6	15.8	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4353	634	0.3	5.7	14.7	63.4	14.5	1.4	0.0
4354	970	0.1	4.9	14.8	65.7	12.6	1.8	0.1
4355	420	0.2	5.0	11.0	65.7	17.9	0.2	0.0
4356	697	0.1	6.5	15.8	64.3	12.3	1.0	0.0
4357	4	*	*	*	*	*	*	*
4358	247	0.8	5.7	6.9	57.9	26.7	2.0	0.0
4359	647	0.0	5.6	13.6	60.4	19.5	0.9	0.0
4360	212	0.0	7.1	12.7	60.8	17.5	1.9	0.0
4361	300	1.3	8.0	16.7	59.0	13.7	1.3	0.0
4362	393	0.0	5.6	12.0	66.7	15.0	0.8	0.0
4363	1,153	0.1	4.3	12.3	67.3	15.1	0.9	0.0
4401	1,643	0.1	3.7	12.5	63.9	18.7	1.2	0.0
4501	1,153	0.2	4.6	11.9	64.0	18.6	0.7	0.1
4502	1	*	*	*	*	*	*	*
4591	2	*	*	*	*	*	*	*
4601	2,014	0.1	6.0	13.4	62.6	16.5	1.3	0.0
4701	1,737	0.1	3.2	10.1	62.8	22.5	1.4	0.0
5050	1,888	0.2	4.4	10.0	57.3	26.6	1.5	0.1
5051	3,390	0.1	3.3	9.4	63.3	22.5	1.3	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5052	3,057	0.1	3.7	9.5	59.2	25.5	2.0	0.0
5101	2,844	0.0	3.1	9.1	59.7	25.5	2.5	0.0
5202	817	0.0	3.3	9.3	70.3	16.6	0.4	0.1
5203	179	0.0	2.8	10.1	63.7	22.3	0.6	0.6
5204	163	0.0	2.5	9.2	73.0	14.7	0.6	0.0
5601	1,465	0.0	4.1	12.2	67.2	15.8	0.7	0.0
5801	2,574	0.2	6.9	14.0	62.0	16.0	0.9	0.0
5803	188	0.0	2.1	8.0	62.8	23.4	3.7	0.0
5804	49	*	*	*	*	*	*	*
6101	74	*	*	*	*	*	*	*
6102	207	0.5	5.3	14.0	63.3	15.9	1.0	0.0
6103	4,523	0.2	6.3	13.0	64.4	15.3	0.8	0.0
6104	542	0.7	11.4	17.7	60.1	9.0	0.9	0.0
6105	57	*	*	*	*	*	*	*
6106	45	*	*	*	*	*	*	*
6301	6,013	0.3	4.9	12.6	60.2	20.5	1.5	0.0
6312	3,541	0.1	2.4	7.8	57.8	29.1	2.7	0.1
6316	1,997	0.1	3.0	7.7	62.9	24.3	2.0	0.0
6320	259	0.0	2.3	7.7	55.2	32.0	2.3	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
6325	171	0.0	6.4	10.5	62.6	19.3	0.0	1.2
6330	550	0.4	7.8	17.1	52.9	19.8	1.1	0.9
6335	679	0.0	2.9	7.1	60.1	27.4	2.5	0.0
6340	153	0.0	5.9	11.8	52.3	24.8	5.2	0.0
6345	34	*	*	*	*	*	*	*
6350	324	0.6	5.9	12.0	56.2	22.2	1.9	1.2
7304	261	0.0	6.1	11.5	62.1	18.8	1.5	0.0
7306	3,247	0.3	6.1	14.3	60.6	17.5	1.2	0.1
7308	223	0.0	1.8	8.1	70.4	19.7	0.0	0.0
7310	75	*	*	*	*	*	*	*
7401	3,019	0.2	5.1	13.4	65.0	15.5	0.8	0.0
7402	65	*	*	*	*	*	*	*
7403	33	*	*	*	*	*	*	*
7404	31	*	*	*	*	*	*	*
7601	874	0.1	4.5	11.3	63.3	18.9	1.6	0.3
7850	1,305	0.2	5.3	12.8	64.4	16.2	1.2	0.0
7851	83	*	*	*	*	*	*	*
7852	106	0.0	0.9	11.3	65.1	21.7	0.9	0.0
8101	342	0.3	5.3	13.2	56.7	21.1	3.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8102	18	*	*	*	*	*	*	*
8105	86	*	*	*	*	*	*	*
8106	33	*	*	*	*	*	*	*
8150	58	*	*	*	*	*	*	*
8151	3,760	0.0	3.2	7.8	63.4	23.8	1.7	0.0
8152	525	0.8	15.4	32.8	50.9	0.0	0.0	0.2
8153	29	*	*	*	*	*	*	*
8154	359	0.0	1.1	4.2	50.1	40.9	1.7	1.9
8155	73	*	*	*	*	*	*	*
8156	23	*	*	*	*	*	*	*
8157	26	*	*	*	*	*	*	*
8201	3,378	0.1	4.0	12.0	59.4	22.8	1.6	0.1
8202	3,983	0.1	2.4	6.4	58.4	25.1	3.1	4.5
8203	491	0.0	4.3	12.2	60.7	20.8	2.0	0.0
8250	1,090	0.4	5.2	13.5	57.2	21.7	1.9	0.0
8252	45	*	*	*	*	*	*	*
8255	3,377	0.3	5.1	11.5	61.9	18.2	1.0	2.0
8257	616	0.2	3.6	6.3	61.7	26.0	2.3	0.0
8258	1,580	0.2	4.5	10.9	62.8	20.4	1.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8260	764	0.1	3.8	10.1	58.5	23.0	3.8	0.7
8901	1,298	0.8	9.9	17.0	55.2	16.3	0.8	0.0
8902	912	0.2	6.3	16.1	57.1	19.0	1.3	0.0
8903	1,547	1.0	10.5	15.8	55.7	15.6	1.4	0.0
8904	1,337	0.4	5.8	13.6	55.1	23.5	1.5	0.1
8905	3,341	0.2	4.9	10.9	57.0	24.9	1.8	0.2
9004	4	*	*	*	*	*	*	*
9012	48	*	*	*	*	*	*	*
9013	118	0.0	6.8	19.5	49.2	23.7	0.8	0.0
9015	39	*	*	*	*	*	*	*
9016	183	2.2	12.0	20.8	49.2	14.8	1.1	0.0
9017	67	*	*	*	*	*	*	*
9018	141	0.0	8.5	12.8	53.9	24.1	0.7	0.0
9019	51	*	*	*	*	*	*	*
9021	117	0.0	4.3	13.7	53.8	25.6	2.6	0.0
9040	8	*	*	*	*	*	*	*
9041	234	0.0	6.4	15.8	54.7	22.2	0.9	0.0
9045	1,639	0.9	11.7	13.2	52.7	19.7	1.5	0.4
9050	2,086	0.3	7.4	12.0	56.0	21.9	2.3	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9051	2,406	0.3	6.4	10.3	54.4	26.3	2.2	0.1
9053	3,115	0.3	6.6	15.1	56.5	19.8	1.7	0.1
9055	55	*	*	*	*	*	*	*
9058	846	0.1	3.3	7.2	55.3	30.7	3.3	0.0
9066	2,478	0.4	7.6	18.4	52.2	18.7	1.3	1.4
9068	1,613	0.4	6.7	14.8	57.3	19.0	1.8	0.0
9069	3,040	0.2	6.2	15.3	56.6	19.7	1.9	0.0
9070	2,271	0.3	7.5	14.5	56.5	19.9	1.2	0.1
9072	2,006	0.3	7.7	17.4	55.5	17.3	1.6	0.0
9074	1,585	0.6	5.9	13.4	60.6	18.2	1.2	0.0
9080	3,025	0.3	5.3	12.5	58.7	20.9	2.3	0.0
Unknown Clinics	242	0.0	2.5	3.7	53.3	38.4	1.7	0.4
Michigan	206,241	0.2	5.5	12.4	60.9	19.2	1.4	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)
0101	140	5.0 (127)	60.7 (157)	121	16.5 (32)	55.4(147)	10	* *	28	* *	117	10.3(159)	117	11.1(102)
0102	500	4.0 (69)	59.8 (153)	402	28.1(150)	50.0 (65)	54	* *	83	* *	465	7.3(112)	466	10.1(79)
0103	496	4.0 (69)	52.4 (50)	438	26.7(144)	45.2 (30)	46	* *	72	* *	437	6.9 (98)	437	11.2(103)
0104	145	5.5 (142)	56.6 (113)	126	16.7 (37)	57.1(151)	16	* *	19	* *	126	7.1(107)	127	10.2(81)
0250	176	1.1 (4)	64.2 (164)	149	30.2(151)	47.7 (41)	12	* *	35	* *	149	8.7(143)	150	14.0(154)
0251	145	4.1 (75)	45.5 (6)	119	26.9(145)	38.7 (7)	12	* *	15	* *	106	5.7 (59)	105	11.4(110)
0252	147	2.7 (20)	57.8 (129)	126	19.0 (80)	60.3(160)	7	* *	39	* *	130	4.6 (27)	130	13.8(151)
0253	191	7.3 (168)	53.9 (74)	167	24.0(133)	42.5 (17)	12	* *	19	* *	164	8.5(141)	163	14.1(156)
0401	631	5.7 (149)	53.1 (61)	568	16.5 (32)	53.5(124)	85	* *	130	14.6 (9)	532	6.6 (90)	532	11.7(117)
0402	443	3.4 (37)	53.7 (69)	380	18.4 (65)	49.5 (62)	47	* *	87	* *	395	6.6 (90)	396	7.6(27)
0403	158	7.6 (169)	51.3 (35)	134	22.4(123)	47.8 (44)	9	* *	24	* *	134	6.0 (76)	134	9.7(66)
0404	171	5.3 (135)	55.6 (101)	158	18.4 (65)	48.1 (46)	24	* *	30	* *	153	4.6 (27)	153	15.7(165)
0501	523	4.2 (79)	54.3 (78)	457	23.6(131)	43.1 (21)	45	* *	64	* *	591	5.2 (41)	591	7.3(20)
0502	1	* *	* *	0	* *	* *	0	* *	1	* *	1	* *	1	* *
0503	203	5.9 (155)	53.2 (63)	182	19.2 (84)	42.3 (16)	22	* *	18	* *	190	4.2 (17)	190	13.7(150)
0504	530	3.0 (23)	52.8 (57)	390	19.7 (90)	45.4 (32)	57	* *	66	* *	572	4.7 (29)	572	7.5(24)
0505	407	3.7 (52)	54.3 (78)	335	20.6(107)	51.0 (82)	45	* *	44	* *	407	6.4 (85)	408	12.3(129)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	174	3.4 (37)	50.6 (27)	145	25.5(140)	48.3 (51)	13	* *	10	* *	188	2.1 (4)	188	6.9(15)
0650	1,295	5.0 (127)	55.4 (98)	1,109	20.0 (96)	50.9 (80)	146	21.2 (7)	218	21.1(28)	1,209	4.0 (14)	1,209	8.4(37)
0651	740	6.4 (162)	54.6 (86)	649	19.3 (85)	50.4 (75)	80	* *	110	26.4(51)	726	4.3 (19)	727	8.5(41)
0652	362	3.6 (47)	61.3 (161)	315	22.5(125)	52.1(102)	34	* *	70	* *	343	7.9(128)	344	11.9(121)
0653	438	6.4 (162)	48.4 (14)	383	23.8(132)	53.5(124)	46	* *	75	* *	453	5.7 (59)	453	9.9(74)
0654	449	6.7 (165)	50.8 (31)	384	18.0 (61)	49.5 (62)	50	* *	63	* *	426	3.3 (9)	426	10.3(83)
0655	455	4.6 (104)	53.0 (60)	394	15.2 (12)	55.3(144)	45	* *	132	17.4(14)	432	10.0(155)	432	10.6(87)
0657	81	* *	* *	73	* *	* *	8	* *	19	* *	76	* *	76	* *
0701	496	3.0 (23)	42.3 (3)	434	19.8 (91)	46.1 (33)	47	* *	117	10.3 (2)	442	6.6 (90)	442	8.8(48)
0702	90	* *	* *	76	* *	* *	9	* *	16	* *	87	* *	88	* *
0703	304	3.3 (33)	58.2 (133)	266	20.3(101)	55.3(144)	39	* *	50	* *	246	10.2(157)	246	15.4(164)
0704	77	* *	* *	73	* *	* *	6	* *	10	* *	73	* *	73	* *
0705	175	2.3 (13)	48.0 (12)	156	21.2(117)	48.1 (46)	27	* *	32	* *	161	5.0 (36)	161	8.1(36)
0850	1,366	4.3 (86)	55.3 (96)	1,149	13.0 (2)	57.7(156)	173	20.8 (5)	256	19.9(22)	1,295	5.4 (45)	1,297	9.4(62)
0851	1,027	3.7 (52)	54.4 (81)	877	15.6 (17)	54.0(130)	128	21.9 (9)	157	18.5(18)	978	7.0(100)	980	8.8(48)
0901	1,888	4.7 (112)	55.1 (94)	1,600	20.9(111)	50.0 (65)	229	31.0 (25)	357	23.0(35)	1,864	8.7(143)	1,867	10.9(96)
0902	85	* *	* *	74	* *	* *	13	* *	26	* *	83	* *	83	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1051	1,721	4.2 (79)	50.6 (27)	1,475	17.8 (57)	52.1(102)	183	32.2 (29)	267	21.0(27)	1,556	4.4 (21)	1,556	9.1(55)
1101	2,140	4.8 (120)	55.7 (103)	1,839	20.2 (99)	51.8 (96)	377	52.8 (71)	407	42.8(91)	2,127	5.0 (36)	2,128	7.6(27)
1102	1,053	5.4 (139)	52.0 (44)	894	13.8 (4)	58.2(157)	151	27.2 (17)	255	20.8(25)	1,014	4.0 (14)	1,011	5.5(4)
1106	127	7.1 (167)	45.7 (7)	97	* *	* *	27	* *	42	* *	129	2.3 (5)	129	5.4(3)
1250	1,236	5.3 (135)	54.8 (90)	1,064	23.0(128)	45.3 (31)	164	21.3 (8)	177	24.3(42)	1,164	7.2(111)	1,164	5.8(6)
1251	1,019	4.5 (98)	58.2 (133)	867	17.5 (53)	54.4(135)	117	13.7 (1)	198	32.8(74)	954	3.5 (11)	954	7.2(18)
1252	230	2.6 (17)	57.4 (125)	162	19.8 (91)	49.4 (61)	26	* *	29	* *	251	13.1(167)	251	7.6(27)
1253	1,263	5.1 (131)	51.5 (36)	985	17.1 (46)	50.7 (78)	196	34.7 (32)	273	32.6(73)	1,438	9.4(150)	1,439	10.1(79)
1350	2,840	4.0 (69)	54.3 (78)	2,389	19.3 (85)	51.2 (88)	455	37.1 (39)	366	28.4(61)	2,654	5.5 (51)	2,650	10.0(78)
1351	152	3.3 (33)	54.6 (86)	134	14.2 (6)	52.2(105)	30	* *	27	* *	150	2.7 (7)	150	6.0(7)
1352	473	3.6 (47)	59.8 (153)	415	20.5(106)	53.0(121)	50	* *	43	* *	451	7.3(112)	451	13.1(142)
1401	269	3.0 (23)	62.1 (162)	212	25.9(142)	44.3 (25)	24	* *	29	* *	219	5.9 (72)	218	10.6(87)
1402	176	3.4 (37)	59.7 (151)	158	15.8 (20)	51.9 (98)	19	* *	33	* *	164	7.3(112)	164	9.8(71)
1403	24	* *	* *	18	* *	* *	1	* *	5	* *	19	* *	19	* *
1404	28	* *	* *	12	* *	* *	4	* *	3	* *	12	* *	12	* *
1501	1,129	3.8 (59)	58.3 (136)	678	41.7(159)	35.8 (4)	137	48.2 (67)	132	27.3(55)	1,189	8.9(147)	1,189	9.4(62)
1503	1,170	5.0 (127)	58.0 (130)	709	27.9(149)	42.5 (17)	138	34.8 (33)	194	34.5(77)	1,292	6.7 (94)	1,291	8.4(37)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1504	2,344	3.8 (59)	52.6 (54)	1,523	21.1(112)	47.5 (39)	384	38.8 (43)	508	31.3(70)	2,749	8.3(139)	2,757	8.6(43)
1505	984	4.7 (112)	47.8 (11)	810	21.1(112)	43.5 (22)	119	44.5 (61)	220	28.6(63)	950	3.3 (9)	952	8.5(41)
1506	607	4.6 (104)	54.0 (75)	303	31.4(154)	41.3 (15)	85	* *	112	17.0(13)	671	10.4(161)	672	11.2(103)
1507	708	3.7 (52)	50.7 (30)	559	18.6 (71)	52.6(114)	93	* *	117	28.2(58)	729	7.1(107)	729	13.3(144)
1508	131	0.8 (2)	67.9 (167)	92	* *	* *	19	* *	15	* *	119	0.8 (1)	119	7.6(27)
1509	712	3.4 (37)	57.4 (125)	267	32.2(155)	39.7 (9)	64	* *	82	* *	676	7.8(125)	675	7.3(20)
1510	45	* *	* *	23	* *	* *	3	* *	6	* *	42	* *	42	* *
1511	853	3.5 (45)	57.0 (120)	670	17.3 (49)	54.9(140)	115	45.2 (64)	143	25.2(46)	803	7.0(100)	804	11.3(108)
1512	351	3.7 (52)	55.3 (96)	216	19.9 (95)	49.5 (62)	48	* *	74	* *	337	3.6 (12)	340	9.7(66)
1701	878	4.3 (86)	53.3 (64)	734	19.1 (82)	51.2 (88)	79	* *	161	23.6(38)	737	5.7 (59)	737	9.9(74)
1712	26	* *	* *	24	* *	* *	1	* *	1	* *	21	* *	21	* *
1950	940	5.4 (139)	54.6 (86)	806	16.0 (24)	50.0 (65)	114	21.1 (6)	255	14.1 (6)	884	6.1 (79)	882	12.1(127)
1952	166	5.4 (139)	58.4 (137)	148	14.2 (6)	52.0(100)	29	* *	50	* *	158	5.7 (59)	158	13.3(144)
1953	340	4.1 (75)	56.8 (118)	297	17.2 (48)	54.5(136)	44	* *	108	16.7(12)	323	10.2(157)	322	8.7(45)
1954	657	5.5 (142)	52.2 (47)	539	22.6(126)	48.2 (49)	98	* *	202	20.8(25)	729	4.5 (24)	732	11.2(103)
1955	136	6.6 (164)	58.8 (141)	99	* *	* *	15	* *	43	* *	150	4.0 (14)	150	9.3(59)
1957	629	5.1 (131)	56.1 (110)	541	14.8 (8)	54.5(136)	77	* *	192	13.0 (4)	602	5.1 (39)	600	12.3(129)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1959	116	1.7 (7)	55.2 (95)	94	* *	* *	20	* *	17	* *	114	7.9(128)	114	11.4(110)
2150	789	3.4 (37)	50.8 (31)	706	26.9(145)	45.0 (28)	66	* *	157	11.5 (3)	685	10.1(156)	685	12.8(135)
2151	320	5.9(155)	49.7 (18)	280	18.6 (71)	51.4 (92)	26	* *	61	* *	258	7.0(100)	258	9.3(59)
2251	519	3.9 (64)	47.4 (9)	453	15.2 (12)	52.3(110)	40	* *	68	* *	441	4.8 (31)	442	11.8(118)
2252	264	4.2 (79)	52.3 (48)	217	35.9(157)	37.3 (5)	17	* *	38	* *	210	2.9 (8)	210	11.9(121)
2551	2,433	4.6 (104)	53.6 (68)	1,358	20.4(104)	51.1 (84)	459	53.6 (72)	429	39.4(86)	1,537	7.3(112)	1,538	11.9(121)
2552	8	* *	* *	7	* *	* *	4	* *	1	* *	8	* *	8	* *
2554	116	6.0 (160)	51.7 (41)	99	* *	* *	9	* *	15	* *	98	* *	99	* *
2555	6,685	5.5 (142)	53.3 (64)	5,278	17.6 (54)	51.2 (88)	1,353	39.3 (45)	2,104	29.2(65)	5,816	8.1(135)	5,805	12.8(135)
3001	1,111	3.8 (59)	59.5 (149)	909	24.9(138)	42.9 (20)	133	42.1 (54)	680	32.5(72)	996	4.3 (19)	998	9.6(64)
3002	270	3.3 (33)	70.0 (168)	199	27.6(147)	53.3(123)	32	* *	167	30.5(68)	242	5.4 (45)	242	7.0(17)
3003	6	* *	* *	6	* *	* *	1	* *	1	* *	6	* *	6	* *
3004	166	0.0 (1)	67.5 (165)	147	25.2(139)	38.8 (8)	6	* *	21	* *	140	11.4(165)	140	8.6(43)
3007	29	* *	* *	18	* *	* *	1	* *	6	* *	18	* *	18	* *
3009	19	* *	* *	9	* *	* *	0	* *	2	* *	9	* *	9	* *
3010	8	* *	* *	1	* *	* *	0	* *	0	* *	1	* *	1	* *
3201	661	4.4 (94)	54.5 (84)	580	20.7(109)	52.2(105)	75	* *	100	25.0(45)	592	8.8(145)	595	9.2(57)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3301	4,853	4.2 (79)	54.4 (81)	3,948	15.7 (19)	55.7(149)	848	28.7 (20)	914	23.6(38)	4,695	5.1 (39)	4,694	10.6(87)
3302	329	1.5 (6)	60.8 (159)	276	20.7(109)	48.6 (56)	56	* *	60	* *	311	4.5 (24)	312	11.5(115)
3304	154	5.8 (152)	40.9 (2)	120	20.0 (96)	40.0 (10)	20	* *	49	* *	144	5.6 (54)	145	6.2(10)
3305	163	1.2 (5)	46.6 (8)	144	13.2 (3)	52.8(118)	22	* *	46	* *	158	7.0(100)	158	10.8(93)
3306	276	4.0 (69)	57.2 (121)	244	12.7 (1)	57.4(152)	33	* *	71	* *	263	4.9 (34)	263	8.7(45)
3308	66	* *	* *	55	* *	* *	12	* *	20	* *	64	* *	64	* *
3309	41	* *	* *	41	* *	* *	8	* *	13	* *	42	* *	42	* *
3550	106	4.7 (112)	53.8 (71)	96	* *	* *	14	* *	18	* *	91	* *	91	* *
3551	30	* *	* *	27	* *	* *	7	* *	14	* *	27	* *	27	* *
3552	51	* *	* *	48	* *	* *	5	* *	14	* *	46	* *	46	* *
3701	1,179	4.9 (124)	52.5 (53)	930	19.5 (88)	51.9 (98)	155	23.9 (13)	187	13.9 (5)	1,262	5.8 (68)	1,262	7.6(27)
3805	3,817	4.5 (98)	50.9 (34)	3,216	17.0 (44)	52.3(110)	463	32.0 (28)	407	25.8(49)	3,697	5.5 (51)	3,696	9.6(64)
3950	2,357	4.4 (94)	56.0 (109)	1,932	16.0 (24)	54.2(132)	344	40.7 (52)	465	27.7(56)	2,230	9.7(153)	2,229	8.0(35)
3952	14	* *	* *	11	* *	* *	0	* *	2	* *	12	* *	12	* *
3953	5	* *	* *	4	* *	* *	0	* *	0	* *	4	* *	4	* *
3954	53	* *	* *	49	* *	* *	3	* *	10	* *	51	* *	51	* *
4001	1,258	4.3 (86)	54.5 (84)	999	32.4(156)	41.2 (14)	154	51.9 (70)	177	35.0(79)	1,307	7.5(117)	1,308	9.9(74)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
4002	1,105	4.1 (75)	55.8 (104)	915	21.1 (112)	52.8 (118)	131	39.7 (46)	197	34.5 (77)	1,154	10.3 (159)	1,154	7.5 (24)
4103	104	3.8 (59)	74.0 (170)	89	* *	* *	19	* *	14	* *	101	7.9 (128)	102	20.6 (168)
4106	1,598	3.4 (37)	58.1 (131)	1,250	20.6 (107)	48.1 (46)	229	41.5 (53)	266	35.7 (80)	1,586	5.0 (36)	1,586	9.0 (52)
4107	3,499	3.9 (64)	49.3 (16)	2,532	17.6 (54)	50.0 (65)	558	40.5 (50)	722	29.9 (67)	3,600	5.6 (54)	3,603	11.4 (110)
4109	1,096	4.3 (86)	52.9 (58)	556	24.3 (136)	47.7 (41)	174	20.7 (4)	249	14.1 (6)	1,250	6.5 (87)	1,251	9.1 (55)
4110	1,494	2.7 (20)	57.2 (121)	1,096	20.1 (98)	47.5 (39)	168	31.5 (27)	199	26.1 (50)	1,575	5.3 (42)	1,576	7.6 (27)
4112	719	2.1 (10)	67.7 (166)	653	69.5 (160)	11.0 (1)	52	* *	52	* *	705	13.3 (168)	706	5.5 (4)
4113	807	2.6 (17)	57.2 (121)	656	24.2 (135)	42.7 (19)	71	* *	53	* *	786	6.9 (98)	786	9.0 (52)
4120	2,177	4.5 (98)	51.5 (36)	1,510	16.2 (29)	51.8 (96)	340	30.0 (21)	505	24.2 (41)	2,271	4.8 (31)	2,272	11.2 (103)
4160	1,594	5.3 (135)	49.5 (17)	883	21.1 (112)	47.8 (44)	388	31.4 (26)	489	22.3 (32)	2,154	5.8 (68)	2,153	9.7 (66)
4351	686	4.4 (94)	55.8 (104)	605	16.5 (32)	53.7 (127)	61	* *	156	21.8 (31)	614	4.2 (17)	615	11.4 (110)
4352	195	2.1 (10)	60.5 (155)	171	22.8 (127)	51.5 (93)	21	* *	43	* *	170	5.9 (72)	170	8.8 (48)
4353	576	3.3 (33)	58.7 (140)	497	27.8 (148)	44.1 (23)	53	* *	101	26.7 (54)	533	8.6 (142)	532	12.0 (126)
4354	938	4.4 (94)	56.1 (110)	826	19.1 (82)	55.1 (142)	105	37.1 (39)	191	31.9 (71)	822	5.7 (59)	822	9.2 (57)
4355	376	4.5 (98)	50.8 (31)	317	16.7 (37)	57.4 (152)	46	* *	84	* *	329	7.9 (128)	328	11.3 (108)
4356	612	4.6 (104)	54.9 (92)	539	18.4 (65)	51.2 (88)	61	* *	106	19.8 (21)	571	4.4 (21)	571	10.3 (83)
4357	3	* *	* *	3	* *	* *	0	* *	0	* *	2	* *	2	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4358	195	1.0 (3)	73.8 (169)	150	38.0(158)	28.7 (2)	16	* *	36	* *	177	6.8 (96)	177	6.2(10)
4359	610	4.9 (124)	57.4 (125)	534	21.7(121)	46.6 (36)	61	* *	98	* *	531	9.4(150)	532	13.2(143)
4360	209	4.8 (120)	58.9 (143)	184	18.5 (68)	57.6(154)	12	* *	68	* *	182	7.7(120)	183	11.5(115)
4361	289	4.8 (120)	51.9 (43)	257	19.5 (88)	55.3(144)	30	* *	84	* *	248	5.6 (54)	248	6.0(7)
4362	372	4.0 (69)	50.5 (24)	323	16.1 (27)	59.8(159)	43	* *	97	* *	323	5.3 (42)	323	7.7(33)
4363	1,114	3.1 (26)	54.2 (76)	987	18.5 (68)	54.1(131)	89	* *	286	21.7(30)	989	6.1 (79)	988	13.0(139)
4401	1,603	6.1 (161)	52.1 (45)	1,385	16.6 (35)	51.7 (94)	185	24.3 (14)	334	20.7(23)	1,388	7.8(125)	1,389	12.4(131)
4501	1,085	4.3 (86)	54.4 (81)	973	20.3(101)	47.3 (38)	105	23.8 (11)	154	6.5 (1)	1,029	8.0(134)	1,030	9.7(66)
4502	1	* *	* *	1	* *	* *	0	* *	0	* *	1	* *	1	* *
4591	2	* *	* *	1	* *	* *	0	* *	0	* *	1	* *	1	* *
4601	1,791	4.7 (112)	53.8 (71)	1,549	23.0(128)	44.2 (24)	316	23.4 (10)	417	25.2(46)	1,807	7.6(118)	1,809	11.4(110)
4701	1,704	3.7 (52)	50.6 (27)	1,513	15.3 (14)	57.6(154)	269	18.2 (2)	553	14.6 (9)	1,562	7.7(120)	1,564	12.4(131)
5050	1,844	5.6 (147)	52.6 (54)	779	15.8 (20)	52.2(105)	380	37.9 (41)	396	23.0(35)	798	8.9(147)	801	10.5(85)
5051	3,258	4.2 (79)	53.7 (69)	1,527	14.1 (5)	55.4(147)	631	36.9 (38)	598	21.2(29)	1,590	7.7(120)	1,592	11.0(100)
5052	2,967	4.3 (86)	55.8 (104)	1,242	16.3 (31)	51.1 (84)	582	38.3 (42)	576	20.7(23)	1,311	7.9(128)	1,315	11.0(100)
5101	2,712	3.6 (47)	50.5 (24)	2,249	15.3 (14)	52.7(116)	470	30.6 (22)	1,068	25.7(48)	2,475	7.1(107)	2,478	10.9(96)
5202	770	3.1 (26)	49.2 (15)	676	16.0 (24)	50.9 (80)	102	25.5 (16)	194	14.4 (8)	680	7.1(107)	679	10.8(93)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5203	163	3.7 (52)	58.9 (143)	140	25.7(141)	49.3 (60)	13	* *	37	* *	136	8.8(145)	135	8.9(51)
5204	153	4.6 (104)	54.2 (76)	132	17.4 (51)	47.7 (41)	13	* *	41	* *	123	5.7 (59)	123	13.8(151)
5601	1,404	4.7 (112)	53.4 (67)	1,252	15.6 (17)	54.3(133)	159	18.9 (3)	224	28.1(57)	1,291	7.0(100)	1,290	12.6(134)
5801	2,405	5.7 (149)	52.3 (48)	2,040	20.2 (99)	52.9(120)	327	30.9 (24)	480	22.3(32)	2,285	6.1 (79)	2,287	14.6(160)
5803	170	5.9 (155)	55.9 (107)	149	22.1(122)	53.7(127)	21	* *	48	* *	162	9.9(154)	162	13.6(149)
5804	41	* *	* *	39	* *	* *	4	* *	10	* *	47	* *	47	* *
6101	65	* *	* *	57	* *	* *	11	* *	7	* *	66	* *	65	* *
6102	184	4.9 (124)	54.9 (92)	165	18.2 (62)	50.3 (72)	29	* *	20	* *	194	8.2(137)	196	10.7(91)
6103	4,051	3.9 (64)	56.6 (113)	3,528	18.9 (78)	51.0 (82)	463	35.6 (37)	297	23.9(40)	4,115	7.9(128)	4,104	12.8(135)
6104	480	3.1 (26)	52.9 (58)	400	18.8 (76)	48.3 (51)	57	* *	29	* *	487	4.5 (24)	485	11.8(118)
6105	53	* *	* *	49	* *	* *	4	* *	9	* *	54	* *	54	* *
6106	42	* *	* *	28	* *	* *	1	* *	3	* *	36	* *	36	* *
6301	5,842	4.7 (112)	53.3 (64)	5,163	16.6 (35)	50.8 (79)	1,041	33.8 (31)	1,529	24.5(44)	5,345	7.8(125)	5,351	12.4(131)
6312	3,446	3.2 (30)	56.9 (119)	3,044	15.8 (20)	53.2(122)	684	33.3 (30)	1,097	28.2(58)	3,153	6.5 (87)	3,160	13.0(139)
6316	1,977	4.6 (104)	50.1 (23)	1,758	16.2 (29)	53.8(129)	354	25.1 (15)	646	18.9(19)	1,766	5.4 (45)	1,776	7.9(34)
6320	240	5.8 (152)	48.3 (13)	180	16.7 (37)	51.1 (84)	49	* *	71	* *	216	5.6 (54)	216	6.5(13)
6325	165	1.8 (8)	52.1 (45)	134	14.9 (11)	52.2(105)	29	* *	54	* *	148	8.1(135)	148	10.8(93)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6330	496	2.2 (12)	56.7 (116)	375	17.9 (59)	50.4 (75)	76	* *	106	44.3 (92)	458	5.7 (59)	456	14.0 (154)
6335	639	3.6 (47)	49.9 (20)	503	16.1 (27)	50.3 (72)	103	35.0 (36)	129	23.3 (37)	596	6.0 (76)	596	6.7 (14)
6340	132	6.8 (166)	59.1 (145)	98	* *	* *	13	* *	33	* *	134	8.2 (137)	134	9.7 (66)
6345	31	* *	* *	27	* *	* *	6	* *	11	* *	29	* *	29	* *
6350	303	5.0 (127)	51.8 (42)	237	14.8 (8)	52.7 (116)	54	* *	81	* *	279	6.1 (79)	281	9.3 (59)
7304	193	4.7 (112)	55.4 (98)	146	18.5 (68)	44.5 (26)	24	* *	35	* *	227	6.2 (83)	227	8.4 (37)
7306	2,775	3.9 (64)	56.3 (112)	2,184	19.0 (80)	51.1 (84)	429	40.6 (51)	369	26.6 (53)	2,764	6.6 (90)	2,761	10.9 (96)
7308	198	2.5 (16)	58.1 (131)	176	17.0 (44)	48.3 (51)	27	* *	35	* *	195	10.8 (164)	195	13.3 (144)
7310	60	* *	* *	52	* *	* *	16	* *	7	* *	67	* *	67	* *
7401	2,902	4.6 (104)	49.9 (20)	2,491	16.7 (37)	51.7 (94)	397	28.0 (19)	900	19.6 (20)	2,609	9.0 (149)	2,607	10.2 (81)
7402	62	* *	* *	59	* *	* *	12	* *	25	* *	56	* *	56	* *
7403	31	* *	* *	29	* *	* *	5	* *	17	* *	25	* *	24	* *
7404	30	* *	* *	26	* *	* *	1	* *	20	* *	21	* *	21	* *
7601	848	5.9 (155)	52.4 (50)	738	17.1 (46)	48.4 (54)	84	* *	153	17.6 (15)	759	9.5 (152)	760	8.4 (37)
7850	1,272	5.7 (149)	52.7 (56)	1,101	15.8 (20)	52.4 (112)	151	23.8 (11)	198	16.2 (11)	1,174	5.6 (54)	1,172	6.2 (10)
7851	82	* *	* *	71	* *	* *	11	* *	8	* *	74	* *	74	* *
7852	104	4.8 (120)	53.8 (71)	95	* *	* *	10	* *	19	* *	101	5.9 (72)	101	6.9 (15)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8101	302	2.6 (17)	59.3 (146)	262	16.8 (41)	50.4 (75)	74	* *	73	* *	330	7.6(118)	331	12.1(127)
8102	18	* *	* *	15	* *	* *	3	* *	9	* *	17	* *	17	* *
8105	70	* *	* *	63	* *	* *	24	* *	49	* *	82	* *	82	* *
8106	29	* *	* *	22	* *	* *	12	* *	12	* *	32	* *	32	* *
8150	53	* *	* *	40	* *	* *	4	* *	19	* *	52	* *	52	* *
8151	3,539	4.0 (69)	54.8 (90)	2,837	19.8 (91)	48.8 (58)	627	39.2 (44)	928	28.2(58)	3,292	7.7(120)	3,299	9.8(71)
8152	432	7.9 (170)	44.9 (5)	286	22.4(123)	54.5(136)	69	* *	104	47.1(94)	476	3.6 (12)	476	13.9(153)
8153	26	* *	* *	19	* *	* *	5	* *	16	* *	25	* *	25	* *
8154	355	5.9 (155)	32.4 (1)	296	17.6 (54)	29.1 (3)	53	* *	106	17.9(16)	309	1.0 (2)	309	1.0(1)
8155	65	* *	* *	54	* *	* *	10	* *	21	* *	68	* *	68	* *
8156	21	* *	* *	18	* *	* *	5	* *	5	* *	19	* *	19	* *
8157	22	* *	* *	15	* *	* *	2	* *	6	* *	22	* *	22	* *
8201	3,175	4.3 (86)	51.5 (36)	1,978	14.8 (8)	55.9(150)	789	27.8 (18)	1,078	24.3(42)	2,250	7.7(120)	2,254	9.9(74)
8202	3,504	4.5 (98)	47.7 (10)	2,509	19.8 (91)	46.4 (35)	686	43.6 (58)	771	35.8(81)	3,388	5.8 (68)	3,393	3.2(2)
8203	464	3.7 (52)	55.4 (98)	290	15.5 (16)	54.5(136)	88	* *	120	18.3(17)	320	10.6(163)	321	11.8(118)
8250	1,027	4.6 (104)	51.6 (39)	490	18.2 (62)	52.2(105)	291	44.3 (59)	164	32.9(75)	539	5.9 (72)	540	10.9(96)
8252	36	* *	* *	16	* *	* *	10	* *	8	* *	43	* *	43	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8255	3,150	5.8 (152)	52.4 (50)	1,096	18.8 (76)	50.3 (72)	718	30.8 (23)	761	22.5(34)	1,810	6.2 (83)	1,811	9.8(71)
8257	552	5.6 (147)	51.6 (39)	224	25.9(142)	41.1 (13)	100	43.0 (56)	109	28.4(61)	399	7.3(112)	399	7.3(20)
8258	1,468	3.8 (59)	53.1 (61)	887	18.6 (71)	54.3(133)	350	34.9 (35)	331	29.0(64)	999	6.5 (87)	998	9.0(52)
8260	638	5.5 (142)	43.6 (4)	483	23.0(128)	40.4 (11)	156	48.7 (68)	148	33.1(76)	718	10.4(161)	718	6.0(7)
8901	1,182	4.2 (79)	59.3 (146)	987	21.3(118)	52.6(114)	198	55.6 (75)	156	45.5(93)	1,147	4.9 (34)	1,147	13.3(144)
8902	816	3.2 (30)	60.9 (160)	695	18.7 (75)	50.2 (69)	145	50.3 (69)	146	49.3(97)	813	5.7 (59)	815	13.0(139)
8903	1,396	3.6 (47)	59.5 (149)	1,147	21.1(112)	48.6 (56)	246	56.5 (76)	196	51.0(98)	1,379	4.8 (31)	1,382	12.9(138)
8904	1,205	3.2 (30)	60.7 (157)	1,016	20.4(104)	48.9 (59)	248	56.9 (77)	206	51.0(98)	1,200	6.8 (96)	1,204	11.9(121)
8905	3,042	3.4 (37)	57.7 (128)	2,560	21.3(118)	45.0 (28)	397	46.3 (65)	418	40.9(89)	2,935	6.0 (76)	2,943	7.5(24)
9004	4	* *	* *	3	* *	* *	2	* *	0	* *	3	* *	3	* *
9012	40	* *	* *	24	* *	* *	1	* *	0	* *	43	* *	43	* *
9013	108	1.9 (9)	55.6 (101)	88	* *	* *	11	* *	12	* *	106	4.7 (29)	106	17.0(167)
9015	30	* *	* *	18	* *	* *	4	* *	1	* *	37	* *	37	* *
9016	170	5.3 (135)	50.0 (22)	145	24.8(137)	46.2 (34)	20	* *	21	* *	171	7.0(100)	171	14.6(160)
9017	50	* *	* *	27	* *	* *	2	* *	2	* *	60	* *	60	* *
9018	127	2.4 (15)	55.9 (107)	106	17.9 (59)	59.4(158)	1	* *	2	* *	126	2.4 (6)	126	8.7(45)
9019	41	* *	* *	31	* *	* *	4	* *	3	* *	47	* *	47	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9021	93	* *	* *	46	* *	* *	6	* *	4	* *	103	1.9 (3)	103	10.7(91)
9040	8	* *	* *	6	* *	* *	0	* *	1	* *	7	* *	7	* *
9041	202	4.5 (98)	58.4 (137)	169	17.8 (57)	52.1(102)	10	* *	9	* *	193	6.7 (94)	193	11.9(121)
9045	1,496	4.3 (86)	56.7 (116)	1,179	18.3 (64)	48.5 (55)	237	45.1 (63)	240	40.4(88)	1,491	7.0(100)	1,494	10.5(85)
9050	2,000	2.8 (22)	59.7 (151)	1,729	20.3(101)	47.0 (37)	314	40.1 (48)	459	29.8(66)	1,873	6.4 (85)	1,873	11.2(103)
9051	2,258	2.3 (13)	57.3 (124)	1,708	21.4(120)	44.5 (26)	198	34.8 (33)	574	26.5(52)	2,129	8.3(139)	2,119	7.4(23)
9053	3,046	4.1 (75)	58.8 (141)	2,657	17.4 (51)	55.0(141)	632	44.8 (62)	971	38.2(84)	2,815	5.4 (45)	2,813	14.2(157)
9055	45	* *	* *	42	* *	* *	5	* *	1	* *	24	* *	25	* *
9058	687	4.7 (112)	50.5 (24)	549	30.4(152)	38.3 (6)	104	40.4 (49)	131	30.5(68)	787	11.6(166)	787	7.2(18)
9066	2,399	4.2 (79)	58.2 (133)	2,120	18.6 (71)	50.2 (69)	453	43.5 (57)	711	41.4(90)	2,270	5.5 (51)	2,274	14.9(163)
9068	1,580	5.1 (131)	58.5 (139)	1,422	24.0(133)	50.2 (69)	253	55.3 (74)	325	48.9(96)	1,493	5.4 (45)	1,493	14.5(159)
9069	2,961	3.4 (37)	60.5 (155)	2,625	17.3 (49)	53.5(124)	598	40.0 (47)	1,042	37.4(82)	2,802	5.4 (45)	2,807	14.6(160)
9070	2,176	5.5 (142)	54.7 (89)	1,835	19.3 (85)	48.2 (49)	360	44.4 (60)	471	38.6(85)	2,044	5.8 (68)	2,054	13.3(144)
9072	1,962	3.5 (45)	59.4 (148)	1,770	18.9 (78)	52.4(112)	378	42.3 (55)	648	38.0(83)	1,828	5.3 (42)	1,831	16.1(166)
9074	1,557	3.1 (26)	62.6 (163)	1,360	16.9 (43)	55.1(142)	315	47.6 (66)	490	39.6(87)	1,423	4.4 (21)	1,427	14.2(157)
9080	2,724	3.9 (64)	56.6 (113)	2,210	16.8 (41)	52.0(100)	515	54.0 (73)	560	48.2(95)	2,759	5.7 (59)	2,761	10.6(87)
Unknown Clinics	153	5.2 (134)	49.7 (18)	109	31.2(153)	40.4 (11)	17	* *	54	* *	136	16.9(169)	135	22.2(169)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Michigan	189,986	4.3	54.4	149,046	19.2	50.4	29,664	36.1	41,421	27.9	174,730	6.6	174,828	10.6

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
0101	132	86.4 (60)	7.6 (108)	146	51.4 (16)	82	* *	86	* *	
0102	505	89.5 (29)	5.1 (80)	532	47.9 (41)	361	23.5 (52)	361	88.1 (57)	
0103	495	88.3 (44)	6.3 (98)	511	51.7 (15)	328	24.4 (46)	336	88.4 (53)	
0104	141	81.6(106)	10.6 (132)	150	50.7 (19)	109	25.7 (33)	98	* *	
0250	164	75.0(153)	20.7 (166)	180	65.6 (2)	123	16.3 (105)	123	78.9 (121)	
0251	123	83.7 (85)	10.6 (132)	144	60.4 (7)	61	* *	88	* *	
0252	134	83.6 (89)	4.5 (73)	146	61.6 (4)	99	* *	101	79.2 (119)	
0253	179	91.6 (15)	1.1 (8)	187	62.0 (3)	132	15.9 (110)	134	78.4 (129)	
0401	596	93.3 (7)	1.7 (20)	637	47.1 (48)	449	23.8 (50)	425	92.0 (15)	
0402	427	84.8 (75)	10.8 (135)	445	47.4 (46)	288	22.9 (54)	284	87.3 (66)	
0403	148	92.6 (10)	1.4 (12)	156	53.8 (10)	106	30.2 (18)	93	* *	
0404	166	83.1 (95)	12.0 (143)	171	43.9 (66)	100	29.0 (22)	114	86.8 (71)	
0501	647	82.8 (99)	11.7 (138)	674	53.7 (11)	446	29.6 (20)	444	82.4 (110)	
0502	1	* *	* *	1	* *	0	* *	0	* *	
0503	212	81.1(107)	10.4 (131)	217	46.1 (55)	153	28.1 (25)	136	85.3 (85)	
0504	633	87.8 (49)	9.6 (125)	657	51.3 (17)	470	41.9 (7)	419	85.2 (87)	
0505	436	77.8(132)	16.3 (158)	463	54.2 (9)	345	28.4 (24)	322	78.6 (124)	
0506	204	58.3(168)	32.4 (169)	214	60.7 (5)	152	21.1 (69)	149	78.5 (128)	
0650	1,224	83.3 (93)	11.8 (142)	1,299	45.5 (58)	1,030	22.2 (59)	974	88.2 (56)	
0651	737	71.8(160)	18.3 (161)	780	48.5 (36)	522	22.4 (58)	603	84.2 (98)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
0652	348	82.8 (99)	10.1 (129)	366	47.3 (47)	277	24.2 (47)	277	90.3 (30)	
0653	460	82.4(101)	14.1 (153)	494	49.4 (30)	403	17.6 (96)	383	89.6 (40)	
0654	442	79.0(126)	13.8 (150)	456	53.5 (12)	370	20.8 (75)	352	81.8 (111)	
0655	432	90.3 (19)	2.5 (36)	470	33.6(119)	376	21.0 (71)	348	94.8 (3)	
0657	78	* *	* *	82	* *	70	* *	66	* *	
0701	478	78.7(128)	11.7 (138)	521	41.8 (84)	361	32.4 (10)	306	91.2 (24)	
0702	93	* *	* *	103	50.5 (25)	83	* *	72	* *	
0703	271	92.6 (10)	2.6 (38)	306	44.4 (64)	220	25.5 (39)	199	95.0 (2)	
0704	77	* *	* *	80	* *	57	* *	51	* *	
0705	179	80.4(113)	7.3 (105)	187	36.9(111)	141	34.0 (8)	127	89.8 (37)	
0850	1,292	87.5 (52)	5.4 (85)	1,420	41.0 (90)	1,074	19.9 (78)	1,119	84.7 (92)	
0851	970	90.0 (24)	4.4 (72)	1,041	42.8 (78)	828	19.9 (78)	854	83.8 (99)	
0901	1,967	81.9(103)	11.5 (137)	2,097	49.2 (32)	1,470	20.6 (77)	1,497	83.7 (102)	
0902	87	* *	* *	88	* *	63	* *	60	* *	
1051	1,627	69.8(164)	23.9 (167)	1,789	46.8 (50)	1,185	29.9 (19)	1,152	86.6 (72)	
1101	2,097	77.3(135)	4.9 (79)	2,333	34.8(113)	1,800	9.7 (142)	1,747	88.8 (48)	
1102	1,001	89.2 (33)	2.5 (36)	1,133	40.8 (92)	667	22.8 (55)	850	91.8 (17)	
1106	125	82.4(101)	3.2 (49)	148	29.1(137)	122	14.8 (115)	105	88.6 (50)	
1250	1,041	81.9(103)	8.5 (115)	1,267	48.7 (35)	1,003	18.5 (87)	1,028	71.7 (147)	
1251	940	90.9 (17)	3.2 (49)	1,051	43.5 (70)	860	14.7 (117)	807	82.8 (106)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1252	255	78.8(127)	10.2 (130)	281	50.5 (25)	224	23.2 (53)	227	71.8 (146)
1253	1,435	70.3(163)	20.3 (165)	1,556	41.2 (88)	1,334	15.6 (112)	1,214	77.1 (135)
1350	2,647	86.1 (65)	6.6 (101)	2,952	39.4 (99)	2,056	29.3 (21)	2,401	79.3 (118)
1351	152	86.2 (63)	7.9 (111)	160	37.5(109)	131	19.1 (83)	130	90.8 (26)
1352	452	73.9(157)	16.4 (159)	481	50.3 (27)	425	17.4 (99)	416	78.6 (124)
1401	242	77.3(135)	12.8 (146)	272	60.7 (5)	172	42.4 (6)	181	72.9 (145)
1402	166	68.7(165)	18.7 (163)	187	43.9 (66)	115	20.9 (72)	108	79.6 (116)
1403	22	* *	* *	23	* *	17	* *	16	* *
1404	17	* *	* *	33	* *	18	* *	18	* *
1501	1,147	84.7 (76)	5.4 (85)	1,390	43.2 (74)	836	18.3 (91)	1,034	68.1 (150)
1503	1,188	84.6 (79)	3.0 (48)	1,530	40.8 (92)	831	32.7 (9)	1,100	76.6 (137)
1504	2,725	90.1 (23)	4.0 (62)	3,031	41.3 (87)	2,173	17.1 (102)	2,333	82.6 (108)
1505	944	89.8 (26)	5.1 (80)	1,019	39.6 (98)	751	26.9 (28)	761	88.3 (54)
1506	668	86.5 (59)	6.9 (103)	737	43.3 (72)	603	15.4 (114)	557	84.4 (94)
1507	729	89.3 (31)	6.3 (98)	793	45.5 (58)	574	24.9 (42)	616	88.0 (58)
1508	117	88.9 (38)	1.7 (20)	212	29.7(134)	87	* *	117	69.2 (149)
1509	653	79.5(120)	10.7 (134)	791	47.0 (49)	507	15.8 (111)	560	76.1 (141)
1510	44	* *	* *	48	* *	36	* *	36	* *
1511	806	92.4 (12)	1.5 (15)	866	41.9 (83)	714	17.6 (96)	668	84.4 (94)
1512	336	93.5 (6)	2.1 (29)	363	34.4(115)	291	14.1 (120)	267	81.3 (113)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1701	806	78.2(131)	14.0 (151)	886	53.5 (12)	436	31.9 (11)	561	73.8 (144)
1712	24	* *	* *	28	* *	9	* *	11	* *
1950	891	90.5 (18)	4.3 (68)	957	43.7 (68)	750	14.7 (117)	719	89.8 (37)
1952	156	85.3 (73)	4.5 (73)	166	28.9(139)	122	17.2 (101)	112	90.2 (32)
1953	324	88.3 (44)	4.0 (62)	344	40.4 (96)	261	13.8 (121)	256	94.1 (5)
1954	746	87.5 (52)	4.8 (77)	775	45.3 (60)	578	22.8 (55)	599	85.8 (78)
1955	149	88.6 (39)	2.7 (41)	157	49.7 (28)	111	22.5 (57)	112	77.7 (133)
1957	607	89.1 (35)	4.0 (62)	653	40.9 (91)	497	22.1 (61)	441	83.0 (103)
1959	115	93.0 (8)	4.3 (68)	124	48.4 (38)	83	* *	88	* *
2150	755	89.1 (35)	5.7 (91)	805	47.6 (42)	540	26.1 (32)	506	89.1 (46)
2151	277	79.1(124)	12.6 (145)	329	42.9 (77)	166	27.7 (27)	188	85.6 (81)
2251	495	90.3 (19)	4.2 (66)	538	50.6 (20)	338	25.1 (41)	331	92.7 (11)
2252	245	89.0 (37)	1.2 (9)	267	58.1 (8)	134	47.0 (5)	152	78.3 (131)
2551	1,593	80.9(108)	4.7 (75)	2,542	34.1(116)	1,297	5.4 (151)	1,684	78.6 (124)
2552	8	* *	* *	8	* *	5	* *	8	* *
2554	104	88.5 (40)	5.8 (95)	116	46.6 (51)	64	* *	79	* *
2555	5,898	83.7 (85)	2.7 (41)	6,974	23.5(150)	4,373	8.6 (146)	4,511	80.8 (115)
3001	1,061	74.9(154)	8.2 (113)	1,138	48.5 (36)	772	7.5 (148)	775	77.2 (134)
3002	265	87.2 (55)	0.0 (1)	277	49.1 (33)	206	10.2 (138)	198	71.7 (147)
3003	6	* *	* *	7	* *	3	* *	1	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)											
Clinic	Medical Care				WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)		Number	% (Rank)		Number	% (Rank)	Number	% (Rank)
3004	143	75.5(150)	18.9 (164)		175	66.9 (1)		119	10.1 (139)	116	41.4 (154)
3007	21	* *	* *		32	* *		20	* *	19	* *
3009	6	* *	* *		24	* *		10	* *	9	* *
3010	2	* *	* *		9	* *		4	* *	4	* *
3201	649	85.7 (68)	7.7 (109)		687	43.2 (74)		504	22.2 (59)	460	90.2 (32)
3301	4,642	83.2 (94)	6.0 (96)		5,189	33.0(122)		4,305	16.3 (105)	4,279	86.2 (73)
3302	307	80.5(111)	7.8 (110)		340	26.8(142)		175	28.6 (23)	277	89.5 (43)
3304	141	83.7 (85)	9.9 (126)		161	30.4(128)		135	31.9 (11)	131	87.8 (61)
3305	157	88.5 (40)	5.7 (91)		172	32.6(123)		147	23.8 (50)	128	90.6 (29)
3306	263	89.7 (27)	5.3 (84)		283	37.5(109)		242	16.5 (104)	219	93.2 (9)
3308	65	* *	* *		70	* *		58	* *	54	* *
3309	41	* *	* *		42	* *		35	* *	33	* *
3550	100	91.0 (16)	1.0 (5)		104	43.3 (72)		79	* *	73	* *
3551	30	* *	* *		30	* *		23	* *	22	* *
3552	50	* *	* *		48	* *		34	* *	32	* *
3701	1,275	95.5 (2)	1.3 (10)		1,368	45.0 (62)		1,045	11.5 (133)	1,024	93.4 (7)
3805	3,649	86.3 (61)	5.5 (88)		4,006	45.6 (57)		2,608	20.9 (72)	3,408	75.5 (142)
3950	2,230	85.5 (70)	7.3 (105)		2,460	40.7 (95)		2,043	17.1 (102)	1,865	84.5 (93)
3952	13	* *	* *		14	* *		10	* *	10	* *
3953	4	* *	* *		5	* *		3	* *	3	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3954	53	* *	* *	* *	55	* *	45	* *	40	* *
4001	1,317	72.2(159)	14.1 (153)		1,443	49.1 (33)	1,196	12.3 (129)	1,134	63.4 (152)
4002	1,154	80.3(114)	9.9 (126)		1,243	43.6 (69)	1,026	16.1 (107)	963	76.3 (139)
4103	99	* *	* *	* *	106	34.0(117)	85	* *	79	* *
4106	1,558	75.5(150)	14.6 (155)		1,729	38.6(104)	1,437	9.5 (143)	1,360	86.0 (77)
4107	3,628	83.1 (95)	8.8 (118)		3,921	38.2(105)	2,862	21.8 (64)	3,001	89.2 (45)
4109	1,243	84.5 (80)	8.5 (115)		1,327	39.1(101)	1,054	21.5 (67)	974	90.0 (35)
4110	1,564	73.8(158)	18.5 (162)		1,694	47.5 (44)	1,145	12.8 (128)	1,406	85.8 (78)
4112	702	65.4(166)	0.3 (2)		726	44.9 (63)	559	68.3 (3)	631	27.4 (155)
4113	741	55.5(170)	30.4 (168)		860	50.6 (20)	642	4.5 (152)	724	87.8 (61)
4120	2,293	85.8 (67)	5.7 (91)		2,411	33.3(121)	1,994	16.0 (109)	1,812	92.9 (10)
4160	2,161	87.9 (46)	4.2 (66)		2,303	30.0(130)	1,939	18.4 (88)	1,778	91.8 (17)
4351	660	90.0 (24)	2.7 (41)		713	47.5 (44)	435	20.9 (72)	464	84.3 (96)
4352	182	83.5 (91)	6.0 (96)		202	42.1 (80)	133	14.3 (119)	130	86.2 (73)
4353	566	82.9 (98)	8.7 (117)		615	48.3 (40)	431	17.4 (99)	410	85.6 (81)
4354	897	92.4 (12)	3.3 (52)		951	42.4 (79)	482	77.0 (2)	626	89.5 (43)
4355	351	76.6(141)	11.4 (136)		414	40.8 (92)	247	26.7 (29)	253	87.7 (64)
4356	621	79.5(120)	15.9 (157)		682	50.6 (20)	472	23.9 (49)	455	87.9 (59)
4357	4	* *	* *	* *	3	* *	0	* *	0	* *
4358	181	58.0(169)	37.6 (170)		235	52.8 (14)	118	16.1 (107)	138	53.6 (153)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prenatal (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4359	577	88.4 (43)	2.1 (29)	634	43.1 (76)	424	25.7 (33)	384	86.2 (73)
4360	193	77.7(134)	14.0 (151)	210	47.6 (42)	111	30.6 (16)	148	83.8 (99)
4361	269	85.5 (70)	5.2 (82)	295	45.1 (61)	201	21.4 (68)	184	93.5 (6)
4362	348	83.6 (89)	9.5 (123)	386	46.6 (51)	252	31.7 (13)	249	92.4 (13)
4363	1,039	80.1(115)	13.6 (148)	1,128	49.3 (31)	614	31.1 (14)	786	85.0 (91)
4401	1,484	80.8(110)	11.7 (138)	1,615	36.0(112)	1,001	21.1 (69)	1,003	80.9 (114)
4501	1,092	79.2(123)	13.6 (148)	1,129	48.4 (38)	832	24.9 (42)	771	91.8 (17)
4502	1	* *	* *	1	* *	1	* *	1	* *
4591	1	* *	* *	2	* *	1	* *	2	* *
4601	1,824	86.7 (58)	3.9 (61)	1,984	31.1(126)	986	18.2 (92)	1,362	90.1 (34)
4701	1,585	87.1 (56)	6.8 (102)	1,713	29.7(134)	1,269	25.2 (40)	1,108	89.8 (37)
5050	813	89.2 (33)	3.6 (60)	1,874	29.6(136)	974	24.0 (48)	1,377	91.3 (23)
5051	1,635	87.9 (46)	3.4 (55)	3,368	33.8(118)	1,688	22.0 (62)	2,397	87.9 (59)
5052	1,332	86.3 (61)	3.4 (55)	3,041	31.2(125)	1,492	21.8 (64)	2,325	88.9 (47)
5101	2,527	90.3 (19)	2.8 (45)	2,809	28.6(140)	1,366	26.4 (30)	1,806	89.6 (40)
5202	734	92.2 (14)	3.5 (57)	810	42.1 (80)	520	27.9 (26)	508	90.9 (25)
5203	153	94.1 (4)	3.3 (52)	178	50.6 (20)	99	* *	101	85.1 (88)
5204	143	90.2 (22)	6.3 (98)	162	50.6 (20)	85	* *	83	* *
5601	1,381	85.6 (69)	9.3 (121)	1,439	41.6 (85)	1,054	25.7 (33)	970	95.1 (1)
5801	2,320	80.0(116)	9.4 (122)	2,542	42.1 (80)	1,985	19.8 (80)	1,809	90.7 (28)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5803	163	76.1(144)	11.7 (138)	187	29.9(132)	155	30.3 (17)	120	90.8 (26)
5804	47	* *	* *	49	* *	39	* *	31	* *
6101	64	* *	* *	72	* *	52	* *	59	* *
6102	194	89.7 (27)	4.1 (65)	202	38.1(106)	167	25.7 (33)	163	87.1 (67)
6103	4,103	87.9 (46)	4.3 (68)	4,463	49.6 (29)	2,916	18.9 (85)	3,266	77.1 (135)
6104	485	87.6 (51)	2.1 (29)	535	51.0 (18)	374	15.5 (113)	468	75.0 (143)
6105	48	* *	* *	56	* *	46	* *	42	* *
6106	34	* *	* *	45	* *	30	* *	42	* *
6301	5,439	84.5 (80)	4.8 (77)	5,959	27.7(141)	3,417	14.8 (115)	3,956	83.8 (99)
6312	3,196	86.2 (63)	3.5 (57)	3,513	22.2(155)	1,918	18.8 (86)	2,259	88.5 (52)
6316	1,812	89.3 (31)	3.5 (57)	1,983	25.6(145)	1,146	17.5 (98)	1,258	88.6 (50)
6320	231	86.1 (65)	9.5 (123)	257	30.0(130)	30	* *	162	92.6 (12)
6325	153	85.0 (74)	3.3 (52)	167	22.8(153)	109	24.8 (44)	104	93.3 (8)
6330	474	76.6(141)	5.7 (91)	539	29.1(137)	352	11.9 (130)	340	85.3 (85)
6335	621	87.1 (56)	8.2 (113)	672	37.8(107)	88	* *	460	88.3 (54)
6340	139	78.4(129)	12.9 (147)	151	41.1 (89)	32	* *	86	* *
6345	30	* *	* *	33	* *	5	* *	15	* *
6350	288	84.7 (76)	5.2 (82)	319	24.8(147)	118	26.3 (31)	207	94.2 (4)
7304	237	83.5 (91)	8.9 (119)	257	41.6 (85)	180	13.3 (125)	161	82.6 (108)
7306	2,915	76.2(143)	12.2 (144)	3,191	43.4 (71)	2,351	13.1 (127)	2,195	77.9 (132)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7308	203	87.7 (50)	7.9 (111)	221	38.9(103)	160	25.6 (38)	145	92.4 (13)
7310	71	* *	* *	75	* *	53	* *	50	* *
7401	2,641	77.2(137)	15.2 (156)	2,992	46.2 (54)	1,742	20.7 (76)	2,191	78.9 (121)
7402	60	* *	* *	65	* *	26	* *	27	* *
7403	29	* *	* *	33	* *	7	* *	10	* *
7404	27	* *	* *	31	* *	12	* *	14	* *
7601	774	85.4 (72)	7.5 (107)	864	44.3 (65)	447	25.7 (33)	575	78.4 (129)
7850	1,175	76.1(144)	16.4 (159)	1,286	46.1 (55)	1,057	18.4 (88)	952	83.0 (103)
7851	75	* *	* *	82	* *	65	* *	65	* *
7852	100	80.0(116)	10.0 (128)	104	39.4 (99)	78	* *	72	* *
8101	330	76.1(144)	1.5 (15)	330	9.7(172)	270	13.7 (123)	191	86.9 (69)
8102	16	* *	* *	16	* *	10	* *	6	* *
8105	83	* *	* *	84	* *	31	* *	40	* *
8106	32	* *	* *	33	* *	30	* *	18	* *
8150	52	* *	* *	58	* *	39	* *	34	* *
8151	3,365	84.4 (84)	5.4 (85)	3,711	33.5(120)	2,941	17.9 (95)	2,669	87.7 (64)
8152	470	84.5 (80)	2.1 (29)	520	37.7(108)	398	6.8 (150)	380	86.1 (76)
8153	27	* *	* *	29	* *	19	* *	18	* *
8154	317	96.5 (1)	1.6 (17)	357	30.3(129)	263	86.7 (1)	232	23.3 (156)
8155	68	* *	* *	73	* *	50	* *	56	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
Clinic	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8156	20	* *	* *	* *	22	* *	18	* *	15	* *
8157	22	* *	* *	* *	26	* *	18	* *	21	* *
8201	2,301	88.5 (40)	2.7 (41)		3,355	22.0 (157)	1,970	19.1 (83)	2,377	90.3 (30)
8202	3,347	83.7 (85)	9.1 (120)		3,914	39.1 (101)	2,380	11.8 (131)	2,822	78.8 (123)
8203	327	93.0 (8)	0.9 (3)		488	22.1 (156)	318	18.2 (92)	356	91.6 (21)
8250	542	84.5 (80)	1.3 (10)		1,078	21.1 (161)	635	24.7 (45)	922	92.0 (15)
8252	39	* *	* *	* *	44	* *	42	* *	31	* *
8255	1,839	87.3 (54)	2.9 (47)		3,339	22.0 (157)	2,007	19.8 (80)	2,474	91.7 (20)
8257	415	93.7 (5)	1.4 (12)		604	31.3 (124)	407	1.0 (154)	443	87.8 (61)
8258	1,038	89.4 (30)	2.6 (38)		1,565	24.5 (149)	1,102	18.4 (88)	1,092	91.6 (21)
8260	711	80.9 (108)	5.5 (88)		759	25.7 (144)	475	13.3 (125)	516	85.1 (88)
8901	1,184	74.9 (154)	1.9 (26)		1,254	23.5 (150)	528	10.6 (136)	841	84.3 (96)
8902	844	79.6 (118)	1.4 (12)		899	22.9 (152)	439	9.8 (141)	595	82.9 (105)
8903	1,434	75.5 (150)	1.8 (25)		1,498	21.2 (160)	759	6.9 (149)	1,014	86.9 (69)
8904	1,252	81.9 (103)	2.1 (29)		1,314	22.7 (154)	655	9.5 (143)	888	85.1 (88)
8905	3,050	84.7 (76)	3.2 (49)		3,262	34.6 (114)	1,585	10.0 (140)	2,271	78.6 (124)
9004	3	* *	* *	* *	4	* *	0	* *	3	* *
9012	45	* *	* *	* *	46	* *	39	* *	37	* *
9013	106	70.8 (161)	2.8 (45)		116	16.4 (167)	92	* *	85	* *
9015	37	* *	* *	* *	38	* *	37	* *	30	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prenatal (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9016	173	64.7(167)	1.7 (20)	183	19.7(164)	162	10.5 (137)	140	85.7 (80)
9017	60	* *	* *	66	* *	60	* *	50	* *
9018	129	78.3(130)	1.6 (17)	140	20.7(162)	121	10.7 (135)	100	90.0 (35)
9019	48	* *	* *	51	* *	39	* *	41	* *
9021	103	79.6(118)	1.0 (5)	116	20.7(162)	99	* *	96	* *
9040	7	* *	* *	8	* *	1	* *	4	* *
9041	220	80.5(111)	0.9 (3)	227	24.7(148)	80	* *	86	* *
9045	1,501	74.8(156)	7.1 (104)	1,609	30.6(127)	1,007	11.6 (132)	1,150	81.5 (112)
9050	1,899	77.0(139)	2.6 (38)	2,044	29.9(132)	865	55.3 (4)	1,425	76.6 (137)
9051	2,124	83.0 (97)	4.7 (75)	2,355	46.4 (53)	834	30.7 (15)	1,605	65.8 (151)
9053	2,854	77.2(137)	2.2 (34)	3,075	17.2(166)	1,738	18.0 (94)	2,025	79.5 (117)
9055	52	* *	* *	55	* *	3	* *	5	* *
9058	802	94.6 (3)	1.0 (5)	832	40.3 (97)	663	2.0 (153)	631	82.7 (107)
9066	2,298	76.0(147)	1.7 (20)	2,442	13.7(171)	1,054	13.4 (124)	1,564	85.5 (83)
9068	1,517	75.7(148)	2.4 (35)	1,580	18.4(165)	537	22.0 (62)	1,021	87.1 (67)
9069	2,815	79.1(124)	2.0 (27)	2,987	14.2(170)	2,050	13.8 (121)	2,055	89.6 (40)
9070	2,057	77.8(132)	1.6 (17)	2,232	21.4(159)	928	21.7 (66)	1,519	76.3 (139)
9072	1,862	75.7(148)	1.7 (20)	1,959	14.6(169)	1,096	8.7 (145)	1,322	88.8 (48)
9074	1,450	76.8(140)	2.0 (27)	1,558	15.7(168)	1,045	11.4 (134)	1,036	85.4 (84)
9080	2,797	79.4(122)	4.3 (68)	2,985	26.6(143)	2,200	8.2 (147)	2,069	79.0 (120)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Unknown Clinics	160	70.6(162)	5.6 (90)	234	25.2(146)	131	19.8 (80)	89	* *
Michigan	177,865	82.9	6.4	203,107	35.6	131,724	18.8	144,705	83.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	128	51.6 (148)	136	36.8 (166)	46	* *	85	* *	117	6.8 (115)
0102	495	54.1 (156)	508	33.7 (157)	212	26.9 (102)	360	25.3 (132)	470	7.7 (128)
0103	460	53.3 (154)	496	34.9 (162)	187	29.4 (90)	336	27.1 (140)	441	7.3 (123)
0104	142	55.6 (161)	142	35.2 (165)	64	* *	98	* *	128	15.6 (163)
0250	172	61.6 (167)	163	35.0 (163)	81	* *	123	22.8 (121)	150	6.0 (96)
0251	134	51.5 (147)	125	31.2 (150)	52	* *	87	* *	108	8.3 (136)
0252	139	43.9 (108)	135	29.6 (143)	55	* *	101	25.7 (135)	130	5.4 (82)
0253	175	52.0 (150)	181	34.3 (160)	80	* *	132	26.5 (139)	164	11.0 (153)
0401	593	58.3 (164)	596	33.1 (154)	276	34.4 (64)	425	19.3 (91)	537	5.4 (82)
0402	413	53.5 (155)	421	32.5 (153)	184	26.1 (104)	282	22.0 (115)	400	4.8 (63)
0403	146	50.7 (144)	151	21.9 (89)	56	* *	93	* *	135	5.9 (93)
0404	157	49.7 (142)	166	33.7 (157)	65	* *	115	47.8 (156)	153	24.2 (169)
0501	638	47.3 (131)	657	26.5 (116)	238	31.1 (85)	443	26.2 (137)	589	12.1 (158)
0502	0	* *	1	* *	0	* *	0	* *	1	* *
0503	205	46.3 (119)	214	22.9 (96)	78	* *	136	11.8 (29)	192	4.2 (53)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0504	636	47.2 (130)	640	27.8 (132)	245	40.0 (40)	416	16.3 (65)	576	4.9 (66)
0505	435	47.6 (133)	442	26.9 (120)	172	32.6 (76)	321	24.9 (129)	410	6.1 (98)
0506	202	53.0 (153)	207	33.3 (155)	85	* *	150	36.7 (155)	192	17.7 (167)
0650	1,196	46.8 (125)	1,225	26.9 (120)	496	33.3 (71)	974	22.3 (118)	1,224	8.2 (134)
0651	703	57.2 (163)	742	39.4 (168)	344	23.5 (109)	600	32.0 (150)	738	11.4 (156)
0652	315	47.9 (136)	347	27.1 (125)	134	32.1 (79)	276	29.3 (146)	343	11.4 (156)
0653	457	52.1 (151)	470	33.4 (156)	210	27.1 (99)	383	30.8 (148)	468	16.9 (166)
0654	434	46.5 (122)	441	29.7 (144)	181	26.5 (103)	352	33.5 (152)	440	13.6 (160)
0655	430	58.8 (165)	439	39.4 (168)	211	23.7 (108)	347	24.5 (128)	437	6.6 (112)
0657	78	* *	78	* *	32	* *	66	* *	78	* *
0701	470	36.6 (75)	488	19.7 (72)	130	36.9 (49)	304	5.6 (8)	444	0.5 (6)
0702	98	* *	93	* *	40	* *	70	* *	89	* *
0703	282	55.0 (159)	274	23.7 (99)	112	42.9 (30)	198	16.7 (67)	249	4.0 (45)
0704	72	* *	79	* *	31	* *	51	* *	73	* *
0705	179	51.4 (146)	180	30.0 (148)	75	* *	127	11.8 (29)	160	0.6 (7)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	1,280	39.5 (86)	1,308	20.5 (79)	411	39.4 (44)	1,121	17.7 (79)	1,301	6.1 (98)
0851	956	42.4 (97)	989	23.8 (101)	361	33.0 (74)	853	16.8 (69)	985	5.8 (92)
0901	1,989	50.0 (143)	1,977	27.0 (122)	830	34.2 (65)	1,497	20.6 (104)	1,880	5.3 (80)
0902	78	* *	87	* *	38	* *	60	* *	83	* *
1051	1,634	45.0 (112)	1,664	21.8 (88)	580	41.0 (36)	1,151	12.3 (32)	1,567	4.5 (58)
1101	2,020	34.5 (65)	2,147	18.9 (68)	558	34.8 (62)	1,748	20.3 (100)	2,146	6.4 (108)
1102	1,030	47.6 (133)	1,020	26.2 (114)	396	31.6 (84)	850	24.4 (127)	1,017	7.6 (126)
1106	137	46.7 (124)	130	23.8 (101)	45	* *	106	20.8 (105)	130	3.8 (37)
1250	1,168	38.9 (85)	1,184	24.2 (103)	392	27.3 (97)	1,026	19.9 (96)	1,183	4.9 (66)
1251	924	39.9 (90)	972	27.4 (128)	317	23.0 (110)	807	17.5 (75)	963	5.2 (78)
1252	261	34.9 (66)	258	20.5 (79)	75	* *	226	15.0 (49)	258	2.3 (18)
1253	1,383	41.9 (96)	1,457	25.5 (112)	470	28.3 (93)	1,215	17.5 (75)	1,456	5.1 (74)
1350	2,640	40.8 (92)	2,670	22.5 (93)	889	29.1 (91)	2,397	18.2 (83)	2,669	4.5 (58)
1351	148	45.3 (113)	153	24.2 (103)	58	* *	130	12.3 (32)	153	3.9 (39)
1352	455	46.8 (125)	452	30.8 (149)	189	27.0 (100)	415	27.2 (141)	451	6.9 (117)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1401	260	45.8 (115)	239	23.0 (97)	87	* *	181	22.1 (116)	221	7.7 (128)
1402	162	33.3 (57)	178	19.1 (70)	45	* *	108	11.1 (25)	164	7.3 (123)
1403	24	* *	22	* *	11	* *	16	* *	19	* *
1404	28	* *	12	* *	0	* *	18	* *	12	* *
1501	1,182	28.2 (42)	1,192	16.1 (57)	280	33.9 (68)	1,043	13.3 (38)	1,194	5.4 (82)
1503	1,291	34.2 (62)	1,316	20.7 (81)	373	32.4 (77)	1,101	15.6 (55)	1,306	5.7 (90)
1504	2,752	27.9 (40)	2,779	12.8 (43)	679	45.5 (24)	2,339	7.1 (12)	2,765	1.8 (13)
1505	921	28.1 (41)	957	13.1 (45)	230	40.0 (40)	760	6.7 (10)	950	2.8 (21)
1506	662	38.8 (84)	677	21.1 (82)	218	31.7 (83)	556	17.4 (73)	677	4.4 (57)
1507	704	39.5 (86)	729	20.0 (77)	256	41.4 (33)	612	13.1 (36)	729	6.0 (96)
1508	141	0.7 (1)	118	2.5 (4)	1	* *	117	1.7 (1)	116	0.0 (1)
1509	671	30.6 (50)	684	19.4 (71)	181	28.7 (92)	568	10.6 (22)	681	3.4 (30)
1510	47	* *	42	* *	21	* *	34	* *	43	* *
1511	794	46.5 (122)	810	29.8 (146)	322	28.0 (94)	670	15.5 (54)	811	5.4 (82)
1512	323	36.5 (73)	339	18.9 (68)	101	35.6 (55)	267	7.9 (15)	339	4.1 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1701	812	51.8 (149)	826	29.8 (146)	332	33.1 (73)	565	17.7 (79)	742	5.5 (87)
1712	27	* *	24	* *	14	* *	11	* *	21	* *
1950	869	43.3 (105)	894	26.8 (119)	324	30.2 (88)	716	21.2 (106)	893	6.3 (106)
1952	152	41.4 (93)	156	28.8 (139)	52	* *	111	11.7 (26)	155	3.9 (39)
1953	308	48.4 (137)	325	27.7 (131)	133	38.3 (47)	255	15.3 (53)	324	4.9 (66)
1954	709	43.0 (103)	742	27.4 (128)	268	27.6 (96)	598	16.1 (62)	738	8.0 (132)
1955	146	43.2 (104)	151	26.5 (116)	54	* *	111	11.7 (26)	151	6.6 (112)
1957	579	41.8 (95)	607	22.1 (90)	202	38.6 (46)	439	15.7 (57)	603	3.6 (32)
1959	115	47.0 (129)	116	17.2 (61)	46	* *	85	* *	113	2.7 (20)
2150	745	46.0 (118)	761	26.5 (116)	263	33.5 (70)	505	17.4 (73)	689	3.3 (27)
2151	286	54.9 (157)	294	32.0 (151)	113	31.9 (81)	187	19.3 (91)	260	5.0 (71)
2251	488	52.7 (152)	501	29.1 (140)	199	35.7 (53)	332	13.9 (43)	445	5.2 (78)
2252	253	49.0 (138)	246	33.7 (157)	90	* *	151	22.5 (120)	212	4.2 (53)
2551	2,423	36.3 (71)	1,604	15.8 (55)	526	39.4 (44)	1,673	24.3 (125)	1,538	5.4 (82)
2552	8	* *	8	* *	1	* *	8	* *	8	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2554	108	43.5 (107)	103	27.2 (126)	35	* *	78	* *	97	* *
2555	6,200	36.2 (70)	5,920	18.3 (65)	1,531	41.4 (33)	4,513	20.3 (100)	5,828	4.5 (58)
3001	1,050	38.3 (80)	1,071	18.7 (67)	327	35.8 (52)	771	26.3 (138)	1,006	7.0 (119)
3002	267	31.1 (52)	265	21.5 (86)	66	* *	198	17.7 (79)	248	6.5 (110)
3003	3	* *	6	* *	0	* *	1	* *	6	* *
3004	161	6.2 (7)	154	3.2 (5)	9	* *	116	6.9 (11)	140	0.0 (1)
3007	25	* *	19	* *	1	* *	19	* *	18	* *
3009	12	* *	9	* *	0	* *	9	* *	9	* *
3010	4	* *	1	* *	1	* *	4	* *	1	* *
3201	621	45.9 (116)	647	27.0 (122)	239	31.8 (82)	460	21.7 (112)	605	5.1 (74)
3301	4,728	35.6 (69)	4,734	17.1 (60)	1,366	40.5 (39)	4,270	17.6 (78)	4,715	5.0 (71)
3302	305	37.4 (77)	316	19.9 (75)	89	* *	277	18.4 (85)	312	2.2 (16)
3304	144	3.5 (4)	145	1.4 (2)	3	* *	129	3.9 (4)	145	0.7 (8)
3305	156	33.3 (57)	159	12.6 (42)	39	* *	128	9.4 (19)	158	1.9 (15)
3306	250	37.6 (78)	264	21.2 (84)	77	* *	219	14.6 (47)	263	6.1 (98)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3308	63	* *	64	* *	21	* *	54	* *	65	* *
3309	36	* *	42	* *	13	* *	33	* *	42	* *
3550	97	* *	100	47.0 (171)	48	* *	73	* *	91	* *
3551	29	* *	30	* *	14	* *	22	* *	27	* *
3552	42	* *	49	* *	25	* *	32	* *	46	* *
3701	1,273	36.4 (72)	1,280	25.9 (113)	392	18.9 (111)	1,023	28.8 (144)	1,277	5.6 (89)
3805	3,682	45.3 (113)	3,687	26.3 (115)	1,443	32.3 (78)	3,409	21.3 (108)	3,685	6.1 (98)
3950	2,280	36.6 (75)	2,249	19.7 (72)	711	35.7 (53)	1,867	16.1 (62)	2,248	3.7 (35)
3952	13	* *	13	* *	4	* *	10	* *	13	* *
3953	5	* *	4	* *	3	* *	3	* *	4	* *
3954	53	* *	53	* *	11	* *	40	* *	52	* *
4001	1,359	38.7 (82)	1,324	21.1 (82)	430	30.9 (86)	1,132	18.9 (89)	1,320	8.3 (136)
4002	1,149	42.7 (99)	1,164	24.3 (106)	409	34.5 (63)	962	32.0 (150)	1,159	13.4 (159)
4103	91	* *	102	27.5 (130)	31	* *	78	* *	102	5.9 (93)
4106	1,566	31.5 (53)	1,594	14.8 (49)	392	40.8 (37)	1,359	13.2 (37)	1,587	4.1 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4107	3,639	31.6 (54)	3,653	15.9 (56)	966	40.6 (38)	2,992	10.4 (20)	3,633	3.2 (25)
4109	1,207	39.8 (89)	1,268	21.6 (87)	418	35.6 (55)	973	17.2 (70)	1,264	4.0 (45)
4110	1,603	28.6 (44)	1,598	17.3 (62)	411	27.7 (95)	1,412	13.5 (39)	1,589	3.2 (25)
4112	719	2.9 (3)	708	1.4 (2)	16	* *	628	8.9 (17)	708	5.1 (74)
4113	839	16.8 (18)	785	6.6 (12)	115	49.6 (10)	722	5.4 (6)	783	1.4 (10)
4120	2,178	29.4 (47)	2,301	14.6 (48)	538	39.6 (43)	1,810	8.3 (16)	2,298	1.8 (13)
4160	2,126	22.6 (30)	2,174	10.3 (34)	395	47.3 (21)	1,776	7.2 (14)	2,164	1.6 (12)
4351	645	43.9 (108)	664	29.2 (142)	234	25.2 (106)	465	21.9 (114)	622	8.2 (134)
4352	177	55.9 (162)	185	38.4 (167)	83	* *	130	36.2 (153)	169	5.3 (80)
4353	563	38.7 (82)	575	21.2 (84)	174	36.8 (50)	411	18.5 (86)	539	6.9 (117)
4354	888	47.5 (132)	900	32.3 (152)	332	24.4 (107)	626	17.3 (71)	828	5.9 (93)
4355	361	49.6 (141)	367	28.1 (134)	140	36.4 (51)	251	18.3 (84)	331	6.6 (112)
4356	629	42.8 (100)	637	24.2 (103)	214	34.1 (66)	455	18.7 (87)	576	4.9 (66)
4357	3	* *	4	* *	0	* *	0	* *	2	* *
4358	190	2.6 (2)	189	0.0 (1)	4	* *	137	2.2 (2)	181	0.0 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4359	559	33.6 (59)	589	18.5 (66)	141	33.3 (71)	383	15.7 (57)	537	4.1 (49)
4360	197	38.6 (81)	198	23.7 (99)	57	* *	147	28.6 (143)	181	8.3 (136)
4361	277	59.6 (166)	274	39.4 (168)	129	25.6 (105)	183	24.0 (124)	251	4.0 (45)
4362	359	54.9 (157)	359	34.8 (161)	155	30.3 (87)	248	20.2 (99)	326	7.7 (128)
4363	1,046	49.3 (139)	1,066	27.9 (133)	405	32.1 (79)	772	21.5 (109)	997	6.4 (108)
4401	1,454	44.6 (111)	1,506	25.3 (111)	522	35.1 (59)	1,013	19.3 (91)	1,401	7.4 (125)
4501	1,049	47.7 (135)	1,100	25.1 (110)	418	39.7 (42)	771	22.2 (117)	1,029	4.8 (63)
4502	1	* *	1	* *	1	* *	1	* *	1	* *
4591	2	* *	1	* *	0	* *	2	* *	1	* *
4601	1,753	32.5 (56)	1,836	22.2 (91)	477	27.0 (100)	1,270	12.2 (31)	1,630	5.5 (87)
4701	1,536	46.9 (127)	1,587	24.8 (109)	587	44.5 (27)	1,113	14.6 (47)	1,581	4.6 (61)
5050	1,764	38.2 (79)	823	12.9 (44)	204	48.5 (12)	1,369	15.6 (55)	799	3.8 (37)
5051	3,151	34.9 (66)	1,648	11.2 (39)	376	48.1 (14)	2,367	14.4 (46)	1,596	4.1 (49)
5052	2,886	28.9 (46)	1,352	10.0 (31)	267	49.1 (11)	2,317	14.0 (45)	1,322	3.4 (30)
5101	2,447	34.2 (62)	2,555	16.1 (57)	669	48.0 (15)	1,809	13.7 (42)	2,493	4.3 (56)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5202	746	51.3 (145)	747	28.2 (136)	301	34.9 (60)	509	16.5 (66)	682	2.5 (19)
5203	159	55.3 (160)	160	35.0 (163)	64	* *	101	21.8 (113)	137	5.1 (74)
5204	144	44.4 (110)	147	22.4 (92)	48	* *	83	* *	122	0.0 (1)
5601	1,301	46.3 (119)	1,380	24.6 (107)	493	34.1 (66)	968	18.7 (87)	1,292	5.0 (71)
5801	2,322	46.9 (127)	2,332	27.0 (122)	895	33.7 (69)	1,807	17.5 (75)	2,309	7.0 (119)
5803	156	25.0 (35)	162	14.8 (49)	26	* *	120	11.7 (26)	163	3.1 (24)
5804	39	* *	47	* *	14	* *	31	* *	47	* *
6101	66	* *	68	* *	21	* *	59	* *	67	* *
6102	182	43.4 (106)	197	28.4 (137)	74	* *	160	15.0 (49)	194	5.7 (90)
6103	4,040	39.5 (86)	4,175	22.8 (95)	1,400	34.9 (60)	3,278	16.0 (60)	4,154	7.1 (121)
6104	504	42.5 (98)	488	24.6 (107)	172	27.3 (97)	469	31.6 (149)	485	10.1 (147)
6105	52	* *	54	* *	21	* *	43	* *	54	* *
6106	42	* *	36	* *	15	* *	43	* *	36	* *
6301	5,261	28.3 (43)	5,512	12.4 (41)	1,196	44.5 (27)	3,959	12.6 (34)	5,377	3.9 (39)
6312	3,098	18.0 (21)	3,238	7.0 (14)	435	55.2 (6)	2,250	9.1 (18)	3,168	3.9 (39)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6316	1,751	28.7 (45)	1,828	12.2 (40)	392	46.2 (23)	1,260	10.7 (23)	1,783	3.6 (32)
6320	224	16.1 (16)	232	8.2 (20)	23	* *	162	4.3 (5)	216	0.9 (9)
6325	154	42.9 (102)	158	29.7 (144)	52	* *	104	21.2 (106)	148	6.8 (115)
6330	493	29.6 (48)	482	15.1 (52)	105	35.2 (58)	339	16.2 (64)	460	4.8 (63)
6335	613	14.7 (12)	629	8.9 (23)	73	* *	460	5.4 (6)	601	2.2 (16)
6340	133	19.5 (24)	141	9.2 (28)	17	* *	86	* *	136	1.5 (11)
6345	28	* *	31	* *	9	* *	15	* *	29	* *
6350	275	33.8 (60)	292	18.2 (64)	69	* *	207	13.5 (39)	281	3.9 (39)
7304	232	41.4 (93)	243	23.0 (97)	70	* *	161	25.5 (134)	232	15.1 (162)
7306	2,918	35.5 (68)	2,997	15.4 (54)	855	47.6 (18)	2,185	20.4 (102)	2,780	7.8 (131)
7308	205	40.5 (91)	206	19.9 (75)	72	* *	145	20.0 (97)	196	9.7 (144)
7310	66	* *	72	* *	28	* *	50	* *	67	* *
7401	2,814	49.4 (140)	2,694	27.3 (127)	1,129	37.6 (48)	2,178	22.9 (122)	2,623	8.0 (132)
7402	63	* *	61	* *	33	* *	27	* *	56	* *
7403	33	* *	29	* *	14	* *	10	* *	25	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7404	31	* *	28	* *	8	* *	14	* *	22	* *
7601	782	46.3 (119)	798	28.1 (134)	307	32.9 (75)	576	20.1 (98)	764	6.2 (103)
7850	1,188	45.9 (116)	1,180	29.1 (140)	453	29.6 (89)	951	25.0 (130)	1,185	6.1 (98)
7851	82	* *	74	* *	40	* *	65	* *	74	* *
7852	94	* *	102	22.5 (93)	36	* *	72	* *	102	4.9 (66)
8101	302	20.5 (26)	330	9.1 (26)	40	* *	188	22.3 (118)	331	7.6 (126)
8102	15	* *	17	* *	2	* *	6	* *	17	* *
8105	83	* *	83	* *	7	* *	40	* *	83	* *
8106	28	* *	30	* *	8	* *	18	* *	32	* *
8150	49	* *	52	* *	8	* *	34	* *	51	* *
8151	3,393	31.0 (51)	3,367	15.2 (53)	807	41.6 (31)	2,667	18.0 (82)	3,327	7.2 (122)
8152	486	42.8 (100)	476	20.4 (78)	170	47.6 (18)	381	30.7 (147)	476	8.8 (140)
8153	27	* *	26	* *	8	* *	18	* *	26	* *
8154	325	5.2 (5)	316	3.2 (5)	13	* *	235	2.6 (3)	315	0.3 (5)
8155	68	* *	67	* *	4	* *	56	* *	67	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8156	21	* *	20	* *	7	* *	15	* *	19	* *
8157	25	* *	22	* *	2	* *	22	* *	22	* *
8201	3,057	32.3 (55)	2,308	13.3 (47)	517	47.6 (18)	2,373	12.7 (35)	2,268	3.3 (27)
8202	3,597	13.9 (10)	3,482	6.1 (11)	339	47.8 (16)	2,823	11.0 (24)	3,417	3.3 (27)
8203	433	34.4 (64)	326	17.5 (63)	80	* *	360	15.0 (49)	323	4.0 (45)
8250	974	24.6 (34)	542	10.7 (37)	76	* *	917	17.3 (71)	539	4.6 (61)
8252	43	* *	43	* *	7	* *	31	* *	43	* *
8255	3,140	36.5 (73)	1,847	14.9 (51)	428	47.0 (22)	2,466	16.0 (60)	1,815	3.7 (35)
8257	548	5.5 (6)	413	4.8 (8)	12	* *	444	6.5 (9)	385	6.5 (110)
8258	1,427	34.1 (61)	1,044	19.7 (72)	260	35.4 (57)	1,089	16.7 (67)	1,002	4.2 (53)
8260	706	16.1 (16)	731	8.5 (22)	89	* *	515	10.5 (21)	719	3.9 (39)
8901	1,106	23.3 (32)	1,202	10.1 (32)	190	56.8 (1)	845	25.7 (135)	1,151	11.1 (155)
8902	788	15.7 (14)	849	5.8 (9)	84	* *	594	19.4 (94)	813	9.0 (141)
8903	1,332	18.7 (22)	1,436	9.7 (30)	185	43.8 (29)	1,013	23.5 (123)	1,384	9.8 (145)
8904	1,184	18.8 (23)	1,259	9.0 (24)	168	54.2 (7)	886	21.6 (110)	1,208	9.5 (143)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8905	2,882	16.0 (15)	3,093	9.1 (26)	361	41.6 (31)	2,282	13.6 (41)	2,951	6.2 (103)
9004	3	* *	3	* *	0	* *	3	* *	3	* *
9012	41	* *	45	* *	4	* *	37	* *	43	* *
9013	92	* *	106	17.0 (59)	15	* *	85	* *	106	10.4 (150)
9015	38	* *	37	* *	4	* *	30	* *	37	* *
9016	166	25.3 (36)	174	13.2 (46)	29	* *	140	36.4 (154)	174	16.1 (165)
9017	65	* *	60	* *	11	* *	50	* *	60	* *
9018	122	26.2 (39)	129	10.9 (38)	23	* *	100	25.0 (130)	127	6.3 (106)
9019	45	* *	48	* *	14	* *	41	* *	47	* *
9021	106	25.5 (37)	103	5.8 (9)	20	* *	95	* *	103	2.9 (23)
9040	4	* *	7	* *	1	* *	4	* *	7	* *
9041	213	20.7 (27)	222	9.0 (24)	36	* *	83	* *	198	9.1 (142)
9045	1,462	15.2 (13)	1,530	7.8 (17)	183	50.8 (9)	1,148	15.1 (52)	1,499	8.7 (139)
9050	1,812	17.1 (19)	1,930	7.7 (16)	254	56.3 (4)	1,438	19.1 (90)	1,889	10.0 (146)
9051	2,183	12.5 (9)	2,206	6.9 (13)	220	41.4 (33)	1,637	7.1 (12)	2,160	2.8 (21)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 - 2011 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9053	2,620	26.1 (38)	2,870	10.5 (36)	523	51.8 (8)	2,028	28.9 (145)	2,821	16.0 (164)
9055	35	* *	54	* *	2	* *	5	* *	24	* *
9058	787	7.0 (8)	806	3.8 (7)	38	* *	631	13.9 (43)	795	3.6 (32)
9066	2,075	14.1 (11)	2,308	7.1 (15)	219	48.4 (13)	1,567	20.4 (102)	2,273	10.7 (152)
9068	1,287	23.0 (31)	1,517	10.4 (35)	224	47.8 (16)	1,021	27.2 (141)	1,491	14.2 (161)
9069	2,599	21.5 (28)	2,826	8.1 (19)	455	55.4 (5)	2,049	25.4 (133)	2,808	10.3 (149)
9070	1,946	17.9 (20)	2,083	8.0 (18)	277	45.1 (25)	1,521	19.7 (95)	2,054	10.1 (147)
9072	1,648	19.9 (25)	1,861	10.2 (33)	240	44.6 (26)	1,321	21.6 (110)	1,831	11.0 (153)
9074	1,340	24.2 (33)	1,452	9.2 (28)	254	56.7 (3)	1,029	24.3 (125)	1,436	10.4 (150)
9080	2,660	22.1 (29)	2,825	8.2 (20)	465	56.8 (1)	2,067	15.9 (59)	2,785	6.2 (103)
Unknown Clinics	128	29.7 (49)	140	28.6 (138)	21	* *	81	* *	140	21.4 (168)
Michigan	183,796	33.7	180,268	17.8	47,745	37.6	144,508	17.3	175,660	6.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
0101	124	7.3 (68)	11.3 (155)	126	14.3 (148)	102	2.0 (20)	127	66.1 (68)
0102	450	5.3 (18)	11.6 (159)	480	7.3 (7)	408	2.0 (20)	462	65.4 (77)
0103	450	6.2 (37)	10.7 (148)	475	9.5 (43)	408	1.5 (9)	454	61.2(102)
0104	133	5.3 (18)	9.0 (119)	136	8.8 (26)	120	3.3 (88)	134	64.2 (89)
0250	155	10.3(139)	11.0 (151)	156	12.2 (120)	133	5.3(142)	154	65.6 (74)
0251	114	7.0 (65)	7.0 (65)	122	8.2 (17)	104	3.8(115)	114	57.9(116)
0252	123	8.1(103)	11.4 (158)	124	12.9 (132)	104	1.9 (17)	123	69.9 (43)
0253	162	8.6(117)	11.1 (153)	165	7.3 (7)	147	6.8(159)	160	56.9(120)
0401	557	8.1(103)	12.0 (162)	576	8.0 (16)	509	3.7(110)	557	49.2(142)
0402	399	4.5 (7)	8.8 (113)	410	7.6 (13)	366	2.2 (27)	400	69.0 (47)
0403	140	8.6(117)	7.1 (67)	140	7.1 (5)	127	5.5(144)	140	62.9 (97)
0404	152	7.2 (66)	9.2 (121)	159	9.4 (39)	140	3.6(103)	152	46.7(146)
0501	554	8.7(122)	8.5 (102)	603	10.0 (54)	494	4.3(124)	579	72.5 (30)
0502	1	*	* * *	1	* *	1	* *	1	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0503	197	8.1(103)	6.1 (41)	206	12.1 (116)	172	4.1(120)	197	83.2 (3)
0504	554	7.4 (78)	10.1 (139)	607	9.7 (46)	495	2.4 (34)	589	79.6 (6)
0505	391	6.4 (44)	8.7 (108)	419	8.4 (20)	359	3.3 (88)	405	77.0 (16)
0506	172	5.2 (16)	5.8 (32)	183	7.1 (5)	161	3.1 (78)	175	76.6 (17)
0650	1,133	6.5 (48)	6.4 (48)	1,176	12.6 (128)	984	2.7 (52)	1,127	70.9 (38)
0651	677	7.8 (92)	6.6 (54)	701	13.7 (143)	578	3.6(103)	673	57.5(119)
0652	306	8.2(108)	10.8 (149)	321	10.0 (54)	272	4.4(127)	308	59.4(108)
0653	437	6.9 (62)	9.6 (130)	447	8.3 (19)	400	3.8(115)	436	67.9 (53)
0654	398	6.0 (30)	10.3 (142)	409	8.8 (26)	358	2.5 (37)	396	70.2 (42)
0655	394	7.4 (78)	7.9 (86)	418	8.6 (23)	356	3.1 (78)	395	74.2 (23)
0657	70	*	* * *	72	* *	66	* *	70	* *
0701	455	6.4 (44)	9.7 (133)	472	10.8 (76)	404	2.7 (52)	454	80.6 (5)
0702	91	*	* * *	96	* *	81	* *	88	* *
0703	258	4.3 (5)	6.6 (54)	262	10.3 (63)	231	1.3 (8)	257	66.9 (61)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0704	71	* * * *			77	* *	62	* *	71	* *
0705	164	4.9 (13)	8.5 (102)		170	8.8 (26)	148	2.7 (52)	163	76.1 (19)
0850	1,253	6.1 (33)	7.7 (81)		1,282	10.5 (70)	1,116	2.4 (34)	1,256	71.9 (33)
0851	924	6.2 (37)	8.9 (117)		956	10.5 (70)	822	2.7 (52)	927	66.1 (68)
0901	1,872	7.5 (82)	9.8 (135)		1,914	11.1 (85)	1,653	3.0 (70)	1,867	58.0 (114)
0902	79	* * * *			82	* *	71	* *	78	* *
1051	1,524	6.6 (52)	10.0 (137)		1,595	11.2 (93)	1,345	2.5 (37)	1,526	82.2 (4)
1101	2,025	9.9 (136)	6.4 (48)		2,087	12.1 (116)	1,769	3.6 (103)	2,020	47.6 (144)
1102	960	6.6 (52)	9.1 (120)		975	9.4 (39)	856	2.8 (62)	959	52.6 (129)
1106	124	5.6 (24)	11.3 (155)		126	10.3 (63)	111	0.9 (5)	124	72.6 (28)
1250	1,101	6.2 (37)	8.3 (97)		1,132	9.8 (48)	976	2.3 (30)	1,102	59.9 (106)
1251	879	6.5 (48)	7.8 (82)		927	8.2 (17)	805	3.1 (78)	887	67.8 (54)
1252	240	4.6 (8)	7.1 (67)		250	9.6 (45)	216	2.8 (62)	242	62.8 (98)
1253	1,348	7.3 (68)	8.4 (100)		1,394	11.0 (81)	1,187	2.7 (52)	1,357	65.1 (81)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
1350	2,555	7.7 (89)	7.9 (86)	2,633	9.5 (43)	2,308	3.8(115)	2,559	53.7(127)
1351	148	5.4 (22)	12.2 (163)	151	11.9 (112)	130	1.5 (9)	148	72.3 (32)
1352	424	6.8 (60)	8.7 (108)	435	11.0 (81)	379	1.6 (11)	424	50.5(136)
1401	219	8.7(122)	8.7 (108)	231	10.4 (68)	195	5.1(140)	219	69.9 (43)
1402	169	5.9 (27)	12.4 (164)	171	8.8 (26)	154	2.6 (45)	169	84.6 (2)
1403	15	* * *	* *	19	* *	14	* *	15	* *
1404	11	* * *	* *	12	* *	10	* *	12	* *
1501	1,106	5.8 (26)	8.8 (113)	1,168	10.1 (58)	984	2.3 (30)	1,131	69.9 (43)
1503	1,197	6.6 (52)	8.5 (102)	1,285	11.1 (85)	1,053	3.3 (88)	1,238	66.4 (66)
1504	2,490	6.7 (55)	6.9 (61)	2,669	10.9 (80)	2,189	2.6 (45)	2,630	71.1 (36)
1505	897	6.4 (44)	11.1 (153)	932	9.4 (39)	811	2.0 (20)	913	77.3 (13)
1506	630	6.7 (55)	9.5 (128)	660	12.3 (123)	553	2.0 (20)	633	71.6 (34)
1507	666	6.3 (42)	9.5 (128)	715	11.6 (106)	590	2.5 (37)	695	71.1 (36)
1508	109	1.8 (1)	5.5 (26)	134	7.5 (11)	100	1.0 (6)	118	74.6 (22)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
1509	640	7.8 (92)	8.4 (100)	672	11.2 (93)	563	3.7 (110)	652	65.5 (76)
1510	43	* *	* *	44	* *	38	* *	44	* *
1511	762	5.9 (27)	8.5 (102)	791	9.2 (34)	687	2.9 (66)	774	70.8 (39)
1512	322	7.8 (92)	9.3 (125)	331	12.7 (130)	279	3.6 (103)	324	63.0 (96)
1701	746	5.5 (23)	11.9 (160)	789	7.4 (9)	680	2.6 (45)	745	67.7 (55)
1712	25	* *	* *	25	* *	24	* *	25	* *
1950	830	6.7 (55)	5.9 (36)	853	10.0 (54)	741	2.7 (52)	827	68.2 (52)
1952	154	5.2 (16)	7.1 (67)	153	9.8 (48)	134	0.7 (2)	153	65.4 (77)
1953	301	5.3 (18)	7.0 (65)	305	9.2 (34)	269	3.3 (88)	302	66.6 (64)
1954	647	7.3 (68)	7.1 (67)	710	11.1 (85)	565	3.5 (97)	686	66.5 (65)
1955	130	7.7 (89)	8.5 (102)	147	10.2 (60)	115	3.5 (97)	139	64.0 (90)
1957	561	7.8 (92)	9.3 (125)	585	13.0 (134)	483	2.9 (66)	566	74.9 (21)
1959	109	2.8 (2)	11.9 (160)	112	9.8 (48)	98	* *	111	77.5 (12)
2150	710	6.5 (48)	9.6 (130)	731	11.1 (85)	624	2.6 (45)	707	64.5 (87)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
2151	271	5.9 (27)	9.2 (121)	286	9.1 (33)	246	2.4 (34)	270	58.9(111)
2251	456	5.3 (18)	8.3 (97)	473	8.5 (21)	417	1.9 (17)	456	64.9 (83)
2252	227	6.2 (37)	4.8 (20)	234	11.1 (85)	198	2.5 (37)	227	59.9(106)
2551	1,529	9.7(133)	3.9 (12)	1,580	14.0 (146)	1,309	3.6(103)	1,533	47.2(145)
2552	8	* * *	* *	8	* *	8	* *	8	* *
2554	98	* * *	* *	101	5.0 (1)	91	* *	98	* *
2555	5,705	9.9(136)	5.6 (28)	5,833	13.4 (137)	4,884	3.6(103)	5,702	55.0(124)
3001	961	12.7(154)	4.2 (14)	1,012	14.1 (147)	824	6.4(155)	962	50.4(137)
3002	239	12.1(149)	5.4 (25)	247	11.3 (95)	209	7.7(161)	239	43.9(152)
3003	6	* * *	* *	6	* *	6	* *	6	* *
3004	141	5.0 (15)	0.7 (2)	146	13.0 (134)	122	2.5 (37)	141	76.6 (17)
3007	16	* * *	* *	17	* *	15	* *	15	* *
3009	9	* * *	* *	10	* *	9	* *	9	* *
3010	1	* * *	* *	2	* *	1	* *	1	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
3201	597	8.0(100)	10.9 (150)	623	11.4 (101)	526	3.2 (86)	599	53.6(128)
3301	4,507	7.3 (68)	7.6 (78)	4,656	10.5 (70)	4,012	2.6 (45)	4,539	65.3 (79)
3302	302	8.6(117)	6.6 (54)	313	14.4 (149)	259	2.3 (30)	304	58.9(111)
3304	132	7.6 (85)	8.3 (97)	136	7.4 (9)	123	3.3 (88)	133	94.7 (1)
3305	155	3.9 (4)	11.0 (151)	156	9.0 (32)	141	0.0 (1)	156	78.2 (10)
3306	251	6.0 (30)	15.1 (165)	257	9.3 (36)	226	2.2 (27)	252	67.5 (57)
3308	63	* * *	* *	65	* *	59	* *	63	* *
3309	41	* * *	* *	41	* *	36	* *	41	* *
3550	91	* * *	* *	92	* *	82	* *	91	* *
3551	29	* * *	* *	29	* *	26	* *	29	* *
3552	47	* * *	* *	45	* *	42	* *	47	* *
3701	1,213	5.6 (24)	6.8 (58)	1,242	11.1 (85)	1,078	2.5 (37)	1,214	64.6 (85)
3805	3,565	8.5(112)	7.9 (86)	3,598	11.1 (85)	3,121	3.4 (95)	3,562	60.5(103)
3950	2,164	7.4 (78)	8.8 (113)	2,198	10.7 (74)	1,908	2.8 (62)	2,169	64.3 (88)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾	Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
3952	13	* * * *	*	13	* *	12	* *	12	* *
3953	4	* * * *	*	4	* *	4	* *	4	* *
3954	49	* * * *	*	50	* *	45	* *	50	* *
4001	1,167	11.7(146)	4.7 (18)	1,261	15.5 (156)	971	4.3(124)	1,200	59.1(110)
4002	1,075	8.9(125)	6.0 (39)	1,116	12.3 (123)	933	3.1 (78)	1,084	63.8 (91)
4103	94	* * * *	*	98	* *	81	* *	94	* *
4106	1,494	10.6(142)	5.8 (32)	1,528	13.9 (144)	1,266	4.7(133)	1,492	63.4 (94)
4107	3,451	8.5(112)	7.8 (82)	3,541	11.5 (105)	3,046	3.1 (78)	3,448	66.0 (71)
4109	1,197	7.9 (98)	9.7 (133)	1,224	12.3 (123)	1,045	2.5 (37)	1,194	64.7 (84)
4110	1,483	6.7 (55)	7.1 (67)	1,531	10.6 (73)	1,319	3.0 (70)	1,487	62.8 (98)
4112	666	4.4 (6)	9.2 (121)	672	10.7 (74)	588	2.0 (20)	668	49.6(141)
4113	739	4.9 (13)	8.7 (108)	761	12.9 (132)	645	1.2 (7)	735	66.7 (63)
4120	2,173	7.3 (68)	8.9 (117)	2,221	11.3 (95)	1,915	2.2 (27)	2,178	65.6 (74)
4160	2,045	8.6(117)	8.2 (94)	2,115	10.8 (76)	1,807	4.5(128)	2,057	66.1 (68)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4351	617	7.3 (68)	11.3 (155)	636	7.5 (11)	565	4.2(123)	617	69.0 (47)
4352	174	9.2(128)	6.9 (61)	180	5.6 (3)	163	6.1(150)	174	58.0(114)
4353	527	7.8 (92)	8.2 (94)	546	13.0 (134)	455	3.3 (88)	531	65.2 (80)
4354	839	4.8 (12)	9.8 (135)	857	10.3 (63)	745	1.7 (14)	840	66.8 (62)
4355	341	7.9 (98)	10.3 (142)	356	9.8 (48)	307	3.6(103)	340	67.6 (56)
4356	588	6.1 (33)	8.2 (94)	604	11.3 (95)	519	3.1 (78)	586	67.4 (58)
4357	3	*	* * *	3	* *	3	* *	3	* *
4358	165	7.3 (68)	6.1 (41)	170	11.8 (110)	141	2.1 (26)	167	77.8 (11)
4359	538	4.6 (8)	10.0 (137)	563	8.5 (21)	492	1.6 (11)	539	75.3 (20)
4360	183	6.0 (30)	8.7 (108)	188	10.1 (58)	164	3.0 (70)	184	77.2 (15)
4361	252	8.3(109)	5.2 (24)	264	8.7 (24)	230	3.5 (97)	253	70.4 (40)
4362	332	6.9 (62)	9.6 (130)	345	9.9 (52)	299	3.0 (70)	334	79.6 (6)
4363	995	6.8 (60)	10.2 (140)	1,005	9.3 (36)	886	2.9 (66)	993	73.8 (25)
4401	1,413	8.1(103)	7.3 (74)	1,445	11.7 (108)	1,235	2.8 (62)	1,409	66.0 (71)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
4501	1,031	6.7 (55)	7.8 (82)	1,051	11.1 (85)	910	3.0 (70)	1,029	55.9(122)
4502	1	*	* * *	1	* *	1	* *	1	* *
4591	1	*	* * *	1	* *	1	* *	1	* *
4601	1,664	7.5 (82)	6.5 (51)	1,804	9.3 (36)	1,493	3.5 (97)	1,748	67.3 (59)
4701	1,497	6.1 (33)	10.4 (147)	1,539	8.8 (26)	1,355	2.0 (20)	1,495	68.5 (51)
5050	807	11.4(145)	5.6 (28)	808	15.1 (154)	677	4.7(133)	807	78.4 (9)
5051	1,603	8.5(112)	6.7 (57)	1,628	12.1 (116)	1,395	4.1(120)	1,604	77.3 (13)
5052	1,308	8.8(124)	5.7 (30)	1,336	11.8 (110)	1,148	3.1 (78)	1,316	74.1 (24)
5101	2,442	7.4 (78)	6.3 (44)	2,517	11.0 (81)	2,151	2.7 (52)	2,471	67.1 (60)
5202	720	7.2 (66)	8.8 (113)	729	11.9 (112)	630	3.0 (70)	720	71.4 (35)
5203	147	7.5 (82)	10.2 (140)	156	13.5 (138)	126	4.0(118)	148	60.1(105)
5204	141	8.5(112)	7.8 (82)	143	7.7 (15)	130	4.6(129)	141	73.8 (25)
5601	1,314	7.6 (85)	9.3 (125)	1,354	8.8 (26)	1,191	2.9 (66)	1,311	72.5 (30)
5801	2,220	8.5(112)	8.0 (91)	2,277	11.6 (106)	1,951	3.5 (97)	2,216	60.2(104)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	% ^(Rank)	Number	% ^(Rank)	Number	% ^(Rank)
	Number	% ^(Rank)	% ^(Rank)	Number						
5803	156	3.8 (3)	10.3 (142)	160	10.0 (54)	140	0.7 (2)	157	72.6 (28)	
5804	45	* *	* *	46	* *	40	* *	45	* *	
6101	63	* *	* *	65	* *	56	* *	64	* *	
6102	187	6.4 (44)	5.9 (36)	185	10.8 (76)	163	2.5 (37)	186	68.8 (49)	
6103	3,918	8.4(110)	6.9 (61)	4,038	12.2 (120)	3,407	3.5 (97)	3,926	51.4(135)	
6104	456	12.1(149)	3.7 (9)	474	13.5 (138)	389	6.9(160)	456	43.9(152)	
6105	53	* *	* *	52	* *	44	* *	53	* *	
6106	35	* *	* *	35	* *	32	* *	35	* *	
6301	5,309	8.9(125)	6.5 (51)	5,444	11.4 (101)	4,681	3.7(110)	5,323	52.6(129)	
6312	3,115	9.4(130)	6.5 (51)	3,198	12.4 (127)	2,720	2.6 (45)	3,127	62.2(100)	
6316	1,760	7.3 (68)	7.9 (86)	1,795	10.3 (63)	1,576	2.7 (52)	1,759	58.5(113)	
6320	212	8.0(100)	8.0 (91)	229	12.2 (120)	186	1.6 (11)	219	54.8(125)	
6325	150	9.3(129)	6.0 (39)	154	13.6 (140)	125	0.8 (4)	151	63.6 (92)	
6330	448	12.3(152)	6.3 (44)	468	15.6 (157)	370	4.3(124)	456	43.9(152)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
6335	568	6.2 (37)	8.6 (107)	604	7.6 (13)	520	3.3 (88)	586	49.7(140)
6340	128	6.3 (42)	6.3 (44)	136	11.0 (81)	114	1.8 (16)	132	52.3(131)
6345	28	* * *	* *	30	* *	23	* *	28	* *
6350	264	9.1(127)	7.6 (78)	284	12.3 (123)	228	4.8(138)	273	45.8(148)
7304	223	7.6 (85)	10.3 (142)	233	10.3 (63)	198	3.0 (70)	224	54.5(126)
7306	2,802	9.6(132)	7.5 (76)	2,907	11.3 (95)	2,466	4.7(133)	2,795	48.9(143)
7308	195	4.6 (8)	10.3 (142)	201	11.9 (112)	171	2.3 (30)	192	57.8(117)
7310	70	* * *	* *	72	* *	65	* *	71	* *
7401	2,579	7.8 (92)	6.8 (58)	2,638	11.4 (101)	2,283	2.7 (52)	2,581	56.7(121)
7402	61	* * *	* *	61	* *	55	* *	61	* *
7403	29	* * *	* *	29	* *	25	* *	29	* *
7404	27	* * *	* *	27	* *	24	* *	27	* *
7601	757	6.1 (33)	5.5 (26)	785	11.7 (108)	664	3.2 (86)	754	61.5(101)
7850	1,121	7.3 (68)	6.3 (44)	1,153	11.3 (95)	990	2.6 (45)	1,123	63.4 (94)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
Clinic	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	%	% (Rank)						
7851	72	* * * *	* * * *		74	* *	66	* *	72	* *
7852	95	* * * *	* * * *		97	* *	81	* *	96	* *
8101	313	19.5(165)	4.2 (14)		315	22.5 (168)	236	8.9(162)	313	44.7(150)
8102	9	* * * *	* * * *		11	* *	6	* *	9	* *
8105	74	* * * *	* * * *		80	* *	64	* *	76	* *
8106	27	* * * *	* * * *		31	* *	19	* *	27	* *
8150	45	* * * *	* * * *		51	* *	39	* *	46	* *
8151	3,188	8.6(117)	7.2 (72)		3,272	11.4 (101)	2,805	3.7(110)	3,193	69.6 (46)
8152	460	10.4(140)	6.1 (41)		471	10.2 (60)	408	4.7(133)	468	59.4(108)
8153	23	* * * *	* * * *		23	* *	21	* *	23	* *
8154	305	4.6 (8)	9.2 (121)		309	5.5 (2)	287	1.7 (14)	305	79.3 (8)
8155	62	* * * *	* * * *		65	* *	60	* *	62	* *
8156	19	* * * *	* * * *		19	* *	17	* *	20	* *
8157	19	* * * *	* * * *		20	* *	18	* *	19	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
8201	2,193	8.1(103)	7.4 (75)	2,298	9.7 (46)	1,964	4.0(118)	2,209	64.6 (85)
8202	3,090	6.5 (48)	6.9 (61)	3,352	10.2 (60)	2,763	3.0 (70)	3,263	65.1 (81)
8203	302	7.3 (68)	7.9 (86)	326	12.6 (128)	263	1.9 (17)	301	65.8 (73)
8250	517	9.7(133)	5.8 (32)	530	10.8 (76)	455	4.6(129)	518	73.7 (27)
8252	40	* * *	* *	42	* *	33	* *	41	* *
8255	1,768	7.7 (89)	7.6 (78)	1,809	11.3 (95)	1,546	2.7 (52)	1,776	66.4 (66)
8257	323	8.4(110)	5.0 (22)	394	9.9 (52)	290	4.8(138)	367	68.7 (50)
8258	957	7.6 (85)	7.5 (76)	1,027	8.7 (24)	859	3.4 (95)	981	55.9(122)
8260	694	9.7(133)	3.7 (9)	713	9.4 (39)	626	5.8(148)	703	70.3 (41)
8901	1,133	13.0(157)	3.5 (6)	1,154	16.4 (163)	919	6.4(155)	1,134	30.1(165)
8902	816	10.9(143)	3.7 (9)	833	15.6 (157)	682	4.7(133)	818	38.5(158)
8903	1,355	13.1(159)	3.2 (3)	1,379	15.6 (157)	1,111	6.3(154)	1,356	26.7(166)
8904	1,196	12.9(156)	5.7 (30)	1,225	14.9 (153)	1,007	6.2(151)	1,194	43.4(155)
8905	2,893	8.0(100)	6.4 (48)	2,980	12.0 (115)	2,501	4.1(120)	2,902	38.0(159)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
9004	3	* * * *		3	* *	2	* *	3	* *
9012	40	* * * *		42	* *	32	* *	41	* *
9013	102	16.7(163)	5.9 (36)	102	14.7 (152)	86	* *	103	46.6(147)
9015	36	* * * *		36	* *	27	* *	36	* *
9016	169	17.2(164)	0.6 (1)	171	18.1 (167)	138	11.6(163)	169	44.4(151)
9017	57	* * * *		59	* *	49	* *	58	* *
9018	125	12.8(155)	4.8 (20)	127	15.7 (161)	105	5.7(146)	126	50.0(139)
9019	46	* * * *		48	* *	39	* *	46	* *
9021	97	* * * *		101	13.9 (144)	82	* *	97	* *
9040	6	* * * *		7	* *	6	* *	6	* *
9041	202	11.9(147)	3.5 (6)	207	12.1 (116)	174	5.7(146)	199	36.7(160)
9045	1,424	10.4(140)	7.2 (72)	1,470	14.4 (149)	1,196	4.6(129)	1,439	45.5(149)
9050	1,826	9.9(136)	6.8 (58)	1,874	15.2 (155)	1,527	4.6(129)	1,836	34.9(163)
9051	1,994	6.9 (62)	8.0 (91)	2,118	12.7 (130)	1,715	3.1 (78)	2,093	51.6(133)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 - 2011 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9053	2,728	13.5(160)	4.2 (14)	2,824	16.0 (162)	2,272	6.4(155)	2,730	42.3(156)
9055	52	* * *	* *	54	* *	48	* *	52	* *
9058	723	9.5(131)	5.8 (32)	757	10.4 (68)	645	3.7(110)	747	63.5 (93)
9066	2,196	12.3(152)	4.3 (17)	2,278	13.6 (140)	1,880	6.2(151)	2,203	34.0(164)
9068	1,456	13.0(157)	3.6 (8)	1,477	17.9 (166)	1,171	5.5(144)	1,459	39.0(157)
9069	2,703	11.9(147)	5.0 (22)	2,777	14.5 (151)	2,281	5.3(142)	2,739	50.4(137)
9070	1,957	12.2(151)	3.4 (5)	2,049	15.6 (157)	1,627	6.2(151)	1,983	36.3(162)
9072	1,753	14.0(161)	3.9 (12)	1,812	17.5 (165)	1,417	6.6(158)	1,756	36.7(160)
9074	1,407	14.3(162)	3.2 (3)	1,429	17.4 (164)	1,148	6.0(149)	1,409	51.5(134)
9080	2,671	11.3(144)	4.7 (18)	2,744	13.6 (140)	2,293	5.2(141)	2,674	51.9(132)
Unknown Clinics	99	* * *	* *	140	5.7 (4)	89	* *	102	57.8(117)
Michigan	169,183	8.5	7.1	175,048	11.7	147,955	3.6	170,443	59.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

