

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 1

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	34	*	*	*	*	*	*	*
0102	147	85.0	2.0	4.1	0.0	0.7	8.2	0.0
0103	147	89.1	0.0	6.8	0.7	0.0	3.4	0.0
0104	41	*	*	*	*	*	*	*
0105	2	*	*	*	*	*	*	*
0250	47	*	*	*	*	*	*	*
0251	32	*	*	*	*	*	*	*
0252	40	*	*	*	*	*	*	*
0253	43	*	*	*	*	*	*	*
0401	161	93.2	0.6	1.2	0.6	0.6	3.1	0.6
0402	140	90.0	1.4	2.1	0.0	0.0	6.4	0.0
0403	53	*	*	*	*	*	*	*
0404	21	*	*	*	*	*	*	*
0501	170	97.1	0.0	1.2	0.0	0.0	1.8	0.0
0503	34	*	*	*	*	*	*	*
0504	156	88.5	1.3	2.6	1.3	0.0	6.4	0.0
0505	90	*	*	*	*	*	*	*
0506	55	*	*	*	*	*	*	*
0507	25	*	*	*	*	*	*	*
0508	10	*	*	*	*	*	*	*
0509	1	*	*	*	*	*	*	*
0650	360	80.0	2.2	6.9	3.9	1.4	5.3	0.3
0651	213	94.4	0.5	1.9	0.9	0.9	1.4	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 2

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0652	112	94.6	0.0	1.8	0.0	0.0	3.6	0.0
0653	117	96.6	0.0	1.7	0.0	0.0	1.7	0.0
0654	118	88.1	0.8	3.4	0.0	0.0	7.6	0.0
0655	119	91.6	0.0	0.8	0.0	0.0	5.9	1.7
0657	27	*	*	*	*	*	*	*
0701	120	90.0	0.8	1.7	0.8	3.3	2.5	0.8
0702	23	*	*	*	*	*	*	*
0703	66	*	*	*	*	*	*	*
0704	21	*	*	*	*	*	*	*
0705	42	*	*	*	*	*	*	*
0706	7	*	*	*	*	*	*	*
0850	267	84.3	4.9	6.4	0.0	2.6	1.5	0.4
0851	228	92.5	0.0	5.3	0.0	0.4	0.9	0.9
0901	519	79.2	1.7	12.5	0.0	0.2	5.6	0.8
0902	18	*	*	*	*	*	*	*
1051	403	84.6	0.7	5.0	1.2	1.2	6.5	0.7
1101	661	35.2	50.2	8.6	0.3	0.8	4.2	0.6
1102	295	73.2	10.8	7.8	1.4	1.0	5.4	0.3
1104	1	*	*	*	*	*	*	*
1106	29	*	*	*	*	*	*	*
1250	344	88.1	0.6	8.4	0.0	0.0	1.7	1.2
1251	252	92.9	0.4	3.2	0.0	0.4	2.0	1.2
1252	203	63.5	0.0	30.5	0.0	0.5	4.4	1.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 3

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1253	268	79.1	1.1	8.6	0.0	0.0	10.4	0.7
1350	758	60.6	20.8	6.3	0.3	5.4	6.2	0.4
1351	52	*	*	*	*	*	*	*
1352	120	70.8	20.8	3.3	0.0	0.0	4.2	0.8
1401	89	*	*	*	*	*	*	*
1402	36	*	*	*	*	*	*	*
1403	9	*	*	*	*	*	*	*
1501	281	54.8	2.5	36.3	0.0	0.0	5.3	1.1
1503	341	61.6	7.6	22.6	0.3	0.3	6.7	0.9
1504	709	39.8	6.3	44.9	0.3	3.5	4.2	1.0
1505	154	84.4	3.2	7.8	0.0	0.6	2.6	1.3
1506	207	78.3	1.0	17.9	0.0	1.0	1.9	0.0
1507	193	81.3	1.0	9.3	0.5	2.6	5.2	0.0
1508	16	*	*	*	*	*	*	*
1509	154	52.6	1.3	37.0	0.0	0.0	4.5	4.5
1511	226	89.4	2.2	5.3	0.0	0.0	2.7	0.4
1512	124	83.9	0.8	8.1	0.8	0.0	6.5	0.0
1514	106	83.0	2.8	12.3	0.0	0.9	0.0	0.9
1701	223	49.8	0.4	4.0	26.0	0.9	17.5	0.9
1950	198	91.9	0.5	4.5	0.0	0.0	2.5	0.5
1952	34	*	*	*	*	*	*	*
1953	114	88.6	0.0	9.6	0.0	0.0	1.8	0.0
1954	176	86.9	0.0	11.9	0.0	0.0	1.1	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 4

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1955	38	*	*	*	*	*	*	*
1957	147	81.6	0.7	12.2	0.0	0.0	4.8	0.7
1963	24	*	*	*	*	*	*	*
1964	16	*	*	*	*	*	*	*
2150	219	84.0	0.0	2.7	0.5	0.0	12.8	0.0
2151	86	*	*	*	*	*	*	*
2251	133	87.2	0.8	3.8	0.0	0.0	5.3	3.0
2252	66	*	*	*	*	*	*	*
2554	18	*	*	*	*	*	*	*
2555	2,619	49.2	37.2	5.2	0.0	0.2	7.9	0.3
2556	138	26.8	61.6	6.5	0.0	0.0	5.1	0.0
2557	1	*	*	*	*	*	*	*
2558	3	*	*	*	*	*	*	*
2560	2	*	*	*	*	*	*	*
2561	2	*	*	*	*	*	*	*
2562	11	*	*	*	*	*	*	*
2563	15	*	*	*	*	*	*	*
2564	7	*	*	*	*	*	*	*
3001	382	17.3	68.1	10.5	0.0	1.0	2.9	0.3
3002	116	20.7	55.2	18.1	0.0	0.0	6.0	0.0
3004	84	*	*	*	*	*	*	*
3005	19	*	*	*	*	*	*	*
3007	5	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 5

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3201	150	93.3	0.0	4.0	0.0	0.7	1.3	0.7
3301	1,467	37.8	32.5	14.6	0.1	5.6	9.0	0.3
3304	42	*	*	*	*	*	*	*
3305	65	*	*	*	*	*	*	*
3306	86	*	*	*	*	*	*	*
3308	22	*	*	*	*	*	*	*
3309	8	*	*	*	*	*	*	*
3312	158	38.0	25.9	24.7	0.0	0.6	10.8	0.0
3550	11	*	*	*	*	*	*	*
3551	7	*	*	*	*	*	*	*
3552	19	*	*	*	*	*	*	*
3701	355	86.2	0.0	9.3	0.0	0.3	3.9	0.3
3850	1,002	72.9	13.4	5.3	0.2	0.2	7.7	0.4
3901	562	33.3	53.0	5.9	0.2	0.5	6.9	0.2
3902	82	*	*	*	*	*	*	*
3903	18	*	*	*	*	*	*	*
3904	4	*	*	*	*	*	*	*
3905	9	*	*	*	*	*	*	*
3950	816	55.4	20.8	12.5	0.1	1.5	9.2	0.5
3954	10	*	*	*	*	*	*	*
4103	1	*	*	*	*	*	*	*
4106	867	12.8	39.7	42.0	0.2	0.7	3.9	0.7
4107	998	43.0	28.2	19.6	0.6	1.6	6.7	0.3

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 6

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4109	308	86.0	1.0	9.4	0.0	0.6	2.9	0.0
4110	5	*	*	*	*	*	*	*
4112	183	1.1	1.6	95.1	0.0	0.0	0.0	2.2
4115	1,253	31.4	31.4	21.4	0.5	9.8	5.2	0.2
4351	169	84.6	0.6	8.9	0.0	0.6	5.3	0.0
4352	56	*	*	*	*	*	*	*
4353	129	65.9	1.6	31.0	0.0	0.0	1.6	0.0
4354	250	86.4	5.2	2.8	0.0	0.0	5.2	0.4
4355	118	78.8	2.5	6.8	2.5	0.8	7.6	0.8
4356	144	78.5	1.4	13.2	0.0	0.7	6.3	0.0
4358	37	*	*	*	*	*	*	*
4359	174	70.7	0.0	27.6	0.6	0.0	1.1	0.0
4360	69	*	*	*	*	*	*	*
4361	92	*	*	*	*	*	*	*
4362	100	94.0	1.0	1.0	0.0	1.0	3.0	0.0
4363	282	91.1	0.4	2.8	0.0	0.7	4.3	0.7
4401	343	91.3	0.3	3.8	0.0	0.6	3.2	0.9
4501	302	93.0	0.0	4.6	0.0	0.0	2.3	0.0
4601	433	73.2	1.8	17.3	0.0	0.0	7.2	0.5
4602	1	*	*	*	*	*	*	*
4701	367	92.4	0.5	4.1	0.0	0.3	1.6	1.1
5050	548	50.7	38.3	2.2	0.0	2.0	6.8	0.0
5051	902	52.7	33.1	4.3	0.1	3.7	5.5	0.6

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 7

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
5052	984	37.1	38.4	2.4	0.0	15.9	5.6	0.5
5055	393	91.9	1.8	0.3	0.0	6.1	0.0	0.0
5101	543	69.8	11.4	12.0	0.2	2.2	4.4	0.0
5102	13	*	*	*	*	*	*	*
5202	266	90.6	0.4	1.9	0.8	0.0	6.4	0.0
5601	335	93.1	1.2	1.8	0.0	0.6	3.3	0.0
5801	604	81.6	6.0	5.8	0.2	0.5	5.5	0.5
5803	48	*	*	*	*	*	*	*
5804	8	*	*	*	*	*	*	*
5806	20	*	*	*	*	*	*	*
6101	11	*	*	*	*	*	*	*
6102	46	*	*	*	*	*	*	*
6103	1,215	53.5	27.7	9.7	0.4	0.5	7.8	0.3
6105	5	*	*	*	*	*	*	*
6106	86	*	*	*	*	*	*	*
6301	1,227	40.9	30.9	21.0	0.1	1.9	5.1	0.2
6312	620	35.8	52.9	3.4	0.0	1.1	6.3	0.5
6316	452	67.7	19.5	6.6	0.0	3.1	3.1	0.0
6318	14	*	*	*	*	*	*	*
6320	53	*	*	*	*	*	*	*
6325	105	89.5	3.8	4.8	0.0	0.0	1.9	0.0
6335	387	72.1	15.0	3.4	0.0	8.0	1.6	0.0
6340	20	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 8

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6345	37	*	*	*	*	*	*	*
6360	306	22.5	54.2	19.6	0.0	0.3	3.3	0.0
7304	10	*	*	*	*	*	*	*
7306	812	47.5	28.7	17.0	0.0	1.1	5.5	0.1
7308	10	*	*	*	*	*	*	*
7310	2	*	*	*	*	*	*	*
7401	765	81.8	6.3	6.4	0.4	0.4	4.4	0.3
7601	212	93.4	0.0	5.2	0.0	0.0	1.4	0.0
7850	407	89.9	0.5	4.9	0.0	0.0	4.4	0.2
7852	6	*	*	*	*	*	*	*
8101	557	3.6	93.2	0.5	0.0	0.0	2.7	0.0
8102	147	16.3	12.2	66.7	0.0	0.7	4.1	0.0
8105	543	30.8	63.5	2.4	0.0	0.9	2.4	0.0
8106	175	25.7	60.0	8.0	0.0	0.0	5.7	0.6
8150	12	*	*	*	*	*	*	*
8151	1,082	36.5	40.1	10.5	0.1	1.8	10.5	0.4
8152	13	*	*	*	*	*	*	*
8153	3	*	*	*	*	*	*	*
8155	4	*	*	*	*	*	*	*
8157	5	*	*	*	*	*	*	*
8201	814	45.6	38.2	4.9	0.4	3.4	6.9	0.6
8202	1,269	83.3	12.8	1.6	0.0	0.2	1.3	0.8
8203	136	58.8	30.9	2.9	0.0	1.5	4.4	1.5

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 9

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8250	244	17.2	74.2	2.5	0.0	1.6	3.7	0.8
8252	54	*	*	*	*	*	*	*
8255	878	51.6	33.3	8.8	0.2	2.5	3.3	0.3
8257	681	49.3	14.0	0.9	0.3	33.2	2.1	0.3
8263	245	66.9	13.5	13.5	0.4	2.4	3.3	0.0
8264	264	64.0	25.8	3.4	0.0	1.9	4.9	0.0
8267	248	52.4	15.3	27.4	0.0	0.0	4.4	0.4
8901	347	0.9	94.8	0.9	0.0	1.2	2.0	0.3
8902	502	2.8	92.8	2.0	0.0	0.0	2.4	0.0
8903	523	2.1	93.3	0.8	0.0	0.6	2.9	0.4
8904	812	3.0	93.5	1.1	0.1	0.5	1.8	0.0
8905	517	60.9	27.5	8.1	0.0	0.6	2.9	0.0
8909	236	3.0	93.6	0.8	0.0	0.4	2.1	0.0
9045	621	11.0	26.2	58.6	0.2	0.0	3.7	0.3
9050	329	14.0	68.1	14.3	0.0	0.0	3.6	0.0
9051	551	9.4	8.0	79.5	0.0	0.4	2.4	0.4
9053	341	3.5	92.1	0.9	0.3	1.2	1.8	0.3
9055	9	*	*	*	*	*	*	*
9058	347	15.3	74.9	1.7	0.0	6.9	1.2	0.0
9066	366	1.6	96.2	0.8	0.0	0.0	1.4	0.0
9068	299	2.0	97.3	0.3	0.0	0.0	0.3	0.0
9069	889	0.9	97.2	0.3	0.0	0.2	1.2	0.1
9070	540	3.7	90.6	0.7	0.0	2.8	2.0	0.2

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Page : 10

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9072	554	3.1	94.8	0.5	0.0	0.5	0.9	0.2
9080	951	18.2	78.1	1.8	0.1	0.1	1.5	0.2
9082	45	*	*	*	*	*	*	*
9083	67	*	*	*	*	*	*	*
9084	18	*	*	*	*	*	*	*
9085	4	*	*	*	*	*	*	*
9086	12	*	*	*	*	*	*	*
9087	1	*	*	*	*	*	*	*
Michigan	54,099	51.1	30.6	10.6	0.4	2.1	4.8	0.4
National (Prior Year)	181,949	53.8	23.0	14.9	1.5	2.8	3.7	0.4

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Page : 11

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	34	*	*	*	*	*	*	*
0102	147	0.0	2.7	6.1	68.0	21.8	1.4	0.0
0103	147	0.0	2.0	10.9	66.7	19.0	1.4	0.0
0104	41	*	*	*	*	*	*	*
0105	2	*	*	*	*	*	*	*
0250	47	*	*	*	*	*	*	*
0251	32	*	*	*	*	*	*	*
0252	40	*	*	*	*	*	*	*
0253	43	*	*	*	*	*	*	*
0401	161	0.0	3.1	7.5	67.1	20.5	1.9	0.0
0402	140	0.0	2.9	9.3	58.6	27.1	2.1	0.0
0403	53	*	*	*	*	*	*	*
0404	21	*	*	*	*	*	*	*
0501	170	0.0	1.2	8.2	62.4	28.2	0.0	0.0
0503	34	*	*	*	*	*	*	*
0504	156	0.0	1.3	7.1	65.4	23.7	2.6	0.0
0505	90	*	*	*	*	*	*	*
0506	55	*	*	*	*	*	*	*
0507	25	*	*	*	*	*	*	*
0508	10	*	*	*	*	*	*	*
0509	1	*	*	*	*	*	*	*
0650	360	0.3	2.2	7.5	69.2	20.0	0.8	0.0
0651	213	0.0	2.8	6.6	71.4	17.8	0.9	0.5
0652	112	0.0	1.8	3.6	73.2	18.8	2.7	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Page : 12

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0653	117	0.0	2.6	3.4	68.4	24.8	0.9	0.0
0654	118	0.8	5.1	8.5	66.1	18.6	0.8	0.0
0655	119	0.0	2.5	10.1	63.9	22.7	0.0	0.8
0657	27	*	*	*	*	*	*	*
0701	120	0.0	2.5	5.8	50.8	36.7	4.2	0.0
0702	23	*	*	*	*	*	*	*
0703	66	*	*	*	*	*	*	*
0704	21	*	*	*	*	*	*	*
0705	42	*	*	*	*	*	*	*
0706	7	*	*	*	*	*	*	*
0850	267	0.4	4.5	7.9	57.7	27.3	2.2	0.0
0851	228	0.0	3.1	7.5	61.8	25.4	1.3	0.9
0901	519	0.2	3.1	7.9	62.2	24.9	1.7	0.0
0902	18	*	*	*	*	*	*	*
1051	403	0.0	2.0	4.7	62.3	28.0	2.7	0.2
1101	661	0.2	3.2	10.0	59.9	22.4	1.5	2.9
1102	295	0.0	1.4	11.2	61.0	25.4	1.0	0.0
1104	1	*	*	*	*	*	*	*
1106	29	*	*	*	*	*	*	*
1250	344	0.0	1.2	11.0	60.2	24.1	3.5	0.0
1251	252	0.0	2.0	8.3	69.4	18.3	1.6	0.4
1252	203	0.0	3.4	10.8	59.1	24.1	2.5	0.0
1253	268	0.0	2.2	9.3	60.8	25.7	1.9	0.0
1350	758	0.4	2.6	8.0	64.8	23.0	1.1	0.1

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Page : 13

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1351	52	*	*	*	*	*	*	*
1352	120	0.8	3.3	8.3	64.2	22.5	0.0	0.8
1401	89	*	*	*	*	*	*	*
1402	36	*	*	*	*	*	*	*
1403	9	*	*	*	*	*	*	*
1501	281	0.0	2.1	12.1	60.1	23.1	1.8	0.7
1503	341	0.3	4.4	13.8	57.8	22.3	1.5	0.0
1504	709	0.0	3.5	7.8	59.5	28.1	1.1	0.0
1505	154	0.0	2.6	6.5	61.7	27.9	1.3	0.0
1506	207	0.0	4.8	6.8	58.5	27.5	2.4	0.0
1507	193	0.0	1.0	4.1	65.8	28.0	1.0	0.0
1508	16	*	*	*	*	*	*	*
1509	154	0.0	1.9	6.5	63.6	26.6	1.3	0.0
1511	226	0.0	3.5	8.4	64.6	21.2	2.2	0.0
1512	124	0.0	2.4	7.3	58.1	30.6	1.6	0.0
1514	106	0.0	3.8	5.7	55.7	33.0	1.9	0.0
1701	223	0.0	2.7	7.2	62.8	25.6	1.8	0.0
1950	198	0.0	2.5	7.1	73.2	15.7	1.5	0.0
1952	34	*	*	*	*	*	*	*
1953	114	0.9	6.1	6.1	65.8	19.3	1.8	0.0
1954	176	0.6	2.3	9.1	63.6	22.7	1.7	0.0
1955	38	*	*	*	*	*	*	*
1957	147	0.0	2.0	5.4	61.2	29.9	1.4	0.0
1963	24	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS
Table 4A
2019 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic

Run Date: 6/26/2020

Page : 14

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1964	16	*	*	*	*	*	*	*
2150	219	0.0	1.8	9.6	65.8	21.5	1.4	0.0
2151	86	*	*	*	*	*	*	*
2251	133	0.8	0.8	9.0	61.7	26.3	1.5	0.0
2252	66	*	*	*	*	*	*	*
2554	18	*	*	*	*	*	*	*
2555	2,619	0.2	2.7	7.6	64.8	23.1	1.5	0.1
2556	138	0.0	3.6	8.7	70.3	15.9	1.4	0.0
2557	1	*	*	*	*	*	*	*
2558	3	*	*	*	*	*	*	*
2560	2	*	*	*	*	*	*	*
2561	2	*	*	*	*	*	*	*
2562	11	*	*	*	*	*	*	*
2563	15	*	*	*	*	*	*	*
2564	7	*	*	*	*	*	*	*
3001	382	0.3	3.7	8.1	68.1	18.6	1.3	0.0
3002	116	0.0	4.3	10.3	65.5	19.0	0.9	0.0
3004	84	*	*	*	*	*	*	*
3005	19	*	*	*	*	*	*	*
3007	5	*	*	*	*	*	*	*
3201	150	0.0	1.3	4.0	67.3	25.3	2.0	0.0
3301	1,467	0.1	2.3	8.5	62.4	24.7	1.9	0.1
3304	42	*	*	*	*	*	*	*
3305	65	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS
Table 4A
2019 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic

Run Date: 6/26/2020

Page : 15

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3306	86	*	*	*	*	*	*	*
3308	22	*	*	*	*	*	*	*
3309	8	*	*	*	*	*	*	*
3312	158	0.0	1.9	10.8	57.6	28.5	0.6	0.6
3550	11	*	*	*	*	*	*	*
3551	7	*	*	*	*	*	*	*
3552	19	*	*	*	*	*	*	*
3701	355	0.0	1.7	9.9	62.0	25.4	1.1	0.0
3850	1,002	0.0	2.8	9.6	64.1	22.8	0.8	0.0
3901	562	0.2	3.4	11.6	59.3	23.8	1.8	0.0
3902	82	*	*	*	*	*	*	*
3903	18	*	*	*	*	*	*	*
3904	4	*	*	*	*	*	*	*
3905	9	*	*	*	*	*	*	*
3950	816	0.0	3.6	7.6	60.2	26.1	2.6	0.0
3954	10	*	*	*	*	*	*	*
4103	1	*	*	*	*	*	*	*
4106	867	0.0	3.9	8.7	58.6	25.8	2.7	0.3
4107	998	0.2	2.8	7.6	59.4	28.8	0.9	0.3
4109	308	0.3	2.6	5.5	64.9	25.0	1.6	0.0
4110	5	*	*	*	*	*	*	*
4112	183	0.0	3.8	10.4	50.3	31.7	3.3	0.5
4115	1,253	0.0	2.3	6.5	56.5	31.9	2.6	0.2
4351	169	0.0	2.4	10.7	59.8	26.0	1.2	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Page : 16

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4352	56	*	*	*	*	*	*	*
4353	129	0.0	6.2	10.1	65.1	17.8	0.8	0.0
4354	250	0.4	0.8	6.8	69.6	21.2	0.8	0.4
4355	118	0.0	2.5	13.6	58.5	22.0	3.4	0.0
4356	144	0.0	2.8	9.0	64.6	22.9	0.7	0.0
4358	37	*	*	*	*	*	*	*
4359	174	0.6	1.7	9.2	60.9	24.1	3.4	0.0
4360	69	*	*	*	*	*	*	*
4361	92	*	*	*	*	*	*	*
4362	100	0.0	8.0	10.0	62.0	20.0	0.0	0.0
4363	282	0.0	2.5	8.5	63.5	23.4	1.8	0.4
4401	343	0.0	3.2	5.5	66.5	23.9	0.9	0.0
4501	302	0.3	3.0	6.0	67.5	20.9	2.3	0.0
4601	433	0.0	1.8	9.2	65.4	22.4	1.2	0.0
4602	1	*	*	*	*	*	*	*
4701	367	0.0	2.5	4.1	61.9	29.2	2.5	0.0
5050	548	0.0	2.0	5.5	57.5	33.0	2.0	0.0
5051	902	0.0	0.9	5.0	62.2	29.4	2.5	0.0
5052	984	0.1	1.3	6.1	62.7	27.4	2.3	0.0
5055	393	0.0	0.0	1.0	53.9	41.0	3.8	0.3
5101	543	0.0	0.7	5.0	64.3	27.1	2.9	0.0
5102	13	*	*	*	*	*	*	*
5202	266	0.0	0.8	9.4	60.9	27.1	1.9	0.0
5601	335	0.0	1.5	4.5	66.6	25.4	2.1	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Page : 17

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5801	604	0.0	3.3	10.3	64.7	21.0	0.7	0.0
5803	48	*	*	*	*	*	*	*
5804	8	*	*	*	*	*	*	*
5806	20	*	*	*	*	*	*	*
6101	11	*	*	*	*	*	*	*
6102	46	*	*	*	*	*	*	*
6103	1,215	0.2	3.4	10.7	62.1	22.3	1.0	0.2
6105	5	*	*	*	*	*	*	*
6106	86	*	*	*	*	*	*	*
6301	1,227	0.2	2.0	8.1	61.0	26.5	2.1	0.0
6312	620	0.0	0.6	4.5	52.1	39.2	3.4	0.2
6316	452	0.0	0.4	2.4	55.5	39.4	2.2	0.0
6318	14	*	*	*	*	*	*	*
6320	53	*	*	*	*	*	*	*
6325	105	0.0	1.0	2.9	59.0	29.5	1.9	5.7
6335	387	0.0	0.5	3.1	57.4	35.1	3.9	0.0
6340	20	*	*	*	*	*	*	*
6345	37	*	*	*	*	*	*	*
6360	306	0.0	1.0	9.2	57.5	28.4	2.3	1.6
7304	10	*	*	*	*	*	*	*
7306	812	0.1	3.0	8.3	63.3	24.3	1.0	0.1
7308	10	*	*	*	*	*	*	*
7310	2	*	*	*	*	*	*	*
7401	765	0.0	3.3	8.0	63.7	23.1	2.0	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Page : 18

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
7601	212	0.0	3.3	12.7	68.4	15.6	0.0	0.0
7850	407	0.0	4.2	9.6	66.8	17.7	1.7	0.0
7852	6	*	*	*	*	*	*	*
8101	557	0.5	3.8	9.2	61.2	23.3	2.0	0.0
8102	147	0.0	1.4	3.4	68.7	24.5	2.0	0.0
8105	543	0.2	0.7	5.2	63.2	28.4	2.2	0.2
8106	175	0.0	2.9	5.1	64.0	26.3	1.7	0.0
8150	12	*	*	*	*	*	*	*
8151	1,082	0.1	1.8	7.2	59.9	29.1	1.8	0.1
8152	13	*	*	*	*	*	*	*
8153	3	*	*	*	*	*	*	*
8155	4	*	*	*	*	*	*	*
8157	5	*	*	*	*	*	*	*
8201	814	0.0	1.2	6.1	60.7	29.4	2.6	0.0
8202	1,269	0.0	1.2	4.6	53.9	35.1	4.3	0.9
8203	136	0.7	0.0	5.1	55.1	36.8	2.2	0.0
8250	244	0.0	1.2	8.6	61.9	27.0	1.2	0.0
8252	54	*	*	*	*	*	*	*
8255	878	0.0	1.0	6.2	67.4	23.2	2.1	0.1
8257	681	0.3	0.4	3.5	58.0	32.5	5.1	0.1
8263	245	0.0	1.2	5.3	66.9	24.9	1.6	0.0
8264	264	0.0	1.1	5.3	62.5	29.2	1.9	0.0
8267	248	0.4	2.4	7.3	64.1	23.8	2.0	0.0
8901	347	0.0	2.6	8.1	66.6	21.0	1.4	0.3

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS
Table 4A
2019 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic

Run Date: 6/26/2020

Page : 19

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8902	502	0.2	1.4	5.6	64.9	25.7	2.0	0.2
8903	523	0.2	3.4	7.8	63.3	23.1	2.1	0.0
8904	812	0.1	2.3	6.7	64.9	24.1	1.8	0.0
8905	517	0.0	1.4	3.3	55.5	36.0	3.9	0.0
8909	236	0.0	5.1	6.4	64.0	21.6	2.5	0.4
9045	621	0.2	4.0	8.1	55.6	29.1	3.1	0.0
9050	329	0.0	3.0	9.4	64.7	20.7	2.1	0.0
9051	551	0.0	2.9	6.9	53.4	30.3	6.5	0.0
9053	341	0.0	1.8	7.0	63.0	25.8	2.3	0.0
9055	9	*	*	*	*	*	*	*
9058	347	0.0	2.9	7.2	57.6	28.5	3.7	0.0
9066	366	0.0	3.0	6.0	59.6	28.4	3.0	0.0
9068	299	0.3	3.3	7.0	63.9	22.7	2.3	0.3
9069	889	0.0	2.9	6.3	63.1	25.2	2.4	0.1
9070	540	0.2	1.9	10.6	63.1	22.0	2.2	0.0
9072	554	0.2	2.3	10.1	62.5	24.2	0.7	0.0
9080	951	0.0	1.1	6.4	62.1	28.0	2.3	0.1
9082	45	*	*	*	*	*	*	*
9083	67	*	*	*	*	*	*	*
9084	18	*	*	*	*	*	*	*
9085	4	*	*	*	*	*	*	*
9086	12	*	*	*	*	*	*	*
9087	1	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS
Table 4A
2019 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic

Run Date: 6/26/2020

Page : 20

Michigan	54,102	0.1	2.4	7.4	61.8	26.1	2.0	0.2
National (Prior Year)	181,959	0.1	2.9	8.6	61.8	24.6	1.8	0.1

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 21

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	16	*	*	5	*	*	3	*	8	*	29	*	29	*
0102	87	*	*	34	*	*	22	*	26	*	133	4.5 (8)	133	10.5 (39)
0103	124	7.3 (105)	58.9 (29)	77	*	*	22	*	32	*	139	5.8 (32)	139	8.6 (21)
0104	34	*	*	16	*	*	2	*	9	*	36	*	36	*
0105	1	*	*	0	*	*	0	*	0	*	2	*	2	*
0250	47	*	*	33	*	*	6	*	6	*	40	*	40	*
0251	22	*	*	2	*	*	6	*	8	*	28	*	28	*
0252	39	*	*	36	*	*	4	*	11	*	36	*	36	*
0253	26	*	*	13	*	*	3	*	8	*	39	*	39	*
0401	151	5.3 (88)	65.6 (100)	104	23.1 (74)	50.0 (25)	20	*	27	*	148	4.7 (10)	148	18.2 (110)
0402	134	8.2 (107)	54.5 (7)	30	*	*	19	*	34	*	128	7.0 (49)	128	11.7 (55)
0403	49	*	*	19	*	*	6	*	13	*	48	*	48	*
0404	21	*	*	15	*	*	2	*	3	*	19	*	19	*
0501	131	4.6 (72)	55.7 (11)	28	*	*	26	*	22	*	166	7.2 (54)	166	10.8 (41)
0503	32	*	*	16	*	*	7	*	4	*	34	*	34	*
0504	141	4.3 (67)	64.5 (95)	56	*	*	24	*	20	*	147	4.8 (11)	147	18.4 (113)
0505	75	*	*	31	*	*	8	*	13	*	86	*	86	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 22

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	30	*	*	6	*	*	8	*	4	*	54	*	53	*
0507	14	*	*	2	*	*	2	*	6	*	25	*	25	*
0508	6	*	*	3	*	*	1	*	2	*	10	*	10	*
0509	1	*	*	1	*	*	0	*	0	*	1	*	1	*
0650	291	4.5 (71)	63.6 (87)	220	15.9 (17)	51.4 (34)	48	*	72	*	326	8.6 (87)	327	12.2 (59)
0651	199	6.0 (95)	53.3 (4)	120	10.0 (2)	52.5 (45)	22	*	35	*	199	8.0 (76)	199	10.1 (35)
0652	87	*	*	60	*	*	17	*	26	*	111	9.9 (104)	111	18.9 (115)
0653	107	1.9 (5)	58.9 (29)	67	*	*	17	*	18	*	107	9.3 (97)	107	15.9 (99)
0654	109	0.0 (1)	62.4 (67)	70	*	*	17	*	27	*	106	7.5 (63)	107	15.9 (99)
0655	107	4.7 (75)	57.0 (14)	83	*	*	16	*	16	*	104	6.7 (45)	105	20.0 (116)
0657	21	*	*	15	*	*	3	*	7	*	27	*	27	*
0701	87	*	*	34	*	*	12	*	43	*	106	6.6 (43)	106	6.6 (7)
0702	20	*	*	18	*	*	3	*	5	*	22	*	22	*
0703	57	*	*	21	*	*	8	*	10	*	59	*	59	*
0704	21	*	*	17	*	*	4	*	7	*	19	*	19	*
0705	27	*	*	12	*	*	9	*	11	*	37	*	37	*
0706	5	*	*	3	*	*	0	*	3	*	7	*	7	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 23

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	186	4.8 (80)	62.9 (77)	118	22.9 (72)	48.3 (17)	40	*	57	*	239	11.7 (112)	240	14.6 (86)
0851	121	6.6 (102)	51.2 (2)	34	*	*	37	*	53	*	205	5.9 (33)	205	11.2 (50)
0901	430	3.0 (16)	61.9 (61)	280	16.4 (23)	53.2 (49)	84	*	76	*	478	5.0 (15)	478	9.0 (23)
0902	17	*	*	11	*	*	3	*	1	*	16	*	16	*
1051	354	3.7 (47)	53.4 (5)	286	13.3 (6)	54.5 (57)	55	*	67	*	374	11.0 (111)	374	7.5 (13)
1101	557	4.1 (58)	62.7 (73)	424	22.9 (72)	47.4 (11)	81	*	75	*	608	8.2 (80)	608	17.6 (106)
1102	255	4.7 (75)	58.8 (24)	168	18.5 (41)	51.8 (37)	46	*	63	*	259	7.3 (58)	259	14.3 (79)
1104	0	*	*	0	*	*	0	*	0	*	1	*	1	*
1106	27	*	*	18	*	*	7	*	5	*	25	*	25	*
1250	264	4.9 (81)	61.0 (54)	125	19.2 (48)	52.8 (47)	66	*	65	*	317	7.6 (65)	317	10.1 (35)
1251	227	6.6 (102)	59.5 (36)	171	19.3 (50)	54.4 (55)	35	*	62	*	232	8.6 (87)	232	6.5 (6)
1252	180	6.1 (98)	68.9 (107)	56	*	*	27	*	37	*	185	5.9 (33)	185	9.7 (31)
1253	239	4.2 (61)	58.2 (21)	113	23.9 (75)	54.0 (54)	39	*	42	*	239	7.9 (72)	239	9.2 (25)
1350	615	3.9 (51)	62.9 (77)	336	20.2 (61)	52.1 (41)	95	*	162	27.2 (7)	679	9.9 (104)	678	10.9 (44)
1351	44	*	*	38	*	*	8	*	19	*	48	*	48	*
1352	68	*	*	25	*	*	16	*	29	*	112	4.5 (8)	112	2.7 (2)
1401	82	*	*	61	*	*	14	*	13	*	84	*	84	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 24

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1402	33	*	*	26	*	*	7	*	6	*	35	*	35	*
1403	9	*	*	9	*	*	0	*	3	*	9	*	9	*
1501	253	5.5 (92)	61.7 (60)	188	48.4 (80)	26.1 (2)	31	*	44	*	254	7.9 (72)	254	7.9 (17)
1503	303	5.3 (88)	58.7 (23)	228	18.4 (40)	50.0 (25)	45	*	36	*	311	8.7 (91)	313	14.7 (88)
1504	515	4.7 (75)	59.6 (38)	334	15.9 (17)	49.7 (24)	103	47.6 (13)	247	36.0 (27)	642	9.0 (94)	643	7.0 (8)
1505	110	2.7 (13)	60.0 (41)	50	*	*	26	*	49	*	141	13.5 (116)	141	12.1 (58)
1506	182	6.0 (95)	63.2 (83)	53	*	*	29	*	34	*	187	8.0 (76)	187	12.8 (66)
1507	164	6.1 (98)	60.4 (49)	81	*	*	17	*	30	*	181	9.4 (99)	181	9.4 (28)
1508	10	*	*	6	*	*	1	*	2	*	14	*	14	*
1509	120	4.2 (61)	62.5 (69)	58	*	*	20	*	45	*	122	12.3 (113)	122	7.4 (12)
1511	178	7.9 (106)	60.1 (44)	83	*	*	40	*	72	*	204	6.4 (38)	204	9.8 (33)
1512	90	*	*	26	*	*	26	*	44	*	117	4.3 (6)	117	5.1 (4)
1514	73	*	*	49	*	*	19	*	33	*	97	*	97	*
1701	194	5.2 (85)	55.7 (11)	132	12.9 (4)	56.1 (66)	27	*	48	*	197	5.1 (17)	197	12.7 (64)
1950	188	5.3 (88)	59.0 (32)	152	16.4 (23)	48.0 (14)	28	*	55	*	188	6.4 (38)	188	13.3 (70)
1952	31	*	*	28	*	*	4	*	15	*	34	*	34	*
1953	104	9.6 (108)	54.8 (9)	85	*	*	20	*	47	*	109	8.3 (84)	109	18.3 (111)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 25

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1954	171	2.3 (7)	64.3 (93)	146	19.2 (48)	52.1 (41)	24	*	34	*	165	12.7 (114)	165	14.5 (83)
1955	37	*	*	28	*	*	5	*	8	*	33	*	33	*
1957	137	3.6 (43)	59.1 (34)	111	16.2 (21)	52.3 (44)	20	*	44	*	138	9.4 (99)	138	15.2 (93)
1963	22	*	*	18	*	*	6	*	4	*	23	*	24	*
1964	15	*	*	9	*	*	3	*	5	*	14	*	14	*
2150	203	3.4 (33)	60.1 (44)	152	14.5 (12)	56.6 (69)	34	*	56	*	200	9.5 (101)	200	18.5 (114)
2151	81	*	*	65	*	*	10	*	20	*	72	*	73	*
2251	96	*	*	34	*	*	24	*	34	*	118	3.4 (2)	118	14.4 (82)
2252	55	*	*	28	*	*	7	*	13	*	54	*	54	*
2554	16	*	*	7	*	*	1	*	3	*	16	*	16	*
2555	2,030	5.4 (91)	59.1 (34)	1,032	18.3 (37)	52.7 (46)	494	48.0 (16)	485	30.1 (13)	2,415	7.1 (52)	2,416	15.1 (91)
2556	108	4.6 (72)	58.3 (22)	82	*	*	15	*	18	*	131	7.6 (65)	131	12.2 (59)
2557	1	*	*	0	*	*	1	*	0	*	1	*	1	*
2558	3	*	*	2	*	*	1	*	1	*	3	*	3	*
2560	1	*	*	1	*	*	0	*	0	*	1	*	1	*
2561	1	*	*	1	*	*	1	*	0	*	2	*	2	*
2562	10	*	*	3	*	*	3	*	2	*	10	*	10	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 26

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2563	10	*	*	2	*	*	2	*	5	*	15	*	15	*
2564	3	*	*	1	*	*	2	*	1	*	7	*	7	*
3001	361	3.9 (51)	65.7 (101)	275	20.7 (64)	50.5 (29)	69	*	239	31.0 (14)	356	5.9 (33)	356	13.8 (73)
3002	111	1.8 (3)	78.4 (108)	88	*	*	13	*	78	*	108	7.4 (61)	108	13.9 (75)
3004	78	*	*	63	*	*	12	*	18	*	75	*	75	*
3005	16	*	*	7	*	*	7	*	7	*	18	*	18	*
3007	5	*	*	3	*	*	0	*	4	*	5	*	5	*
3201	129	6.2 (100)	63.6 (87)	94	*	*	20	*	35	*	137	5.1 (17)	138	8.0 (18)
3301	1,229	3.7 (47)	63.4 (85)	954	18.1 (36)	52.2 (43)	264	47.7 (14)	284	32.4 (17)	1,369	7.5 (63)	1,369	11.2 (50)
3304	34	*	*	27	*	*	5	*	8	*	39	*	39	*
3305	55	*	*	40	*	*	10	*	18	*	55	*	55	*
3306	68	*	*	56	*	*	16	*	21	*	77	*	77	*
3308	21	*	*	17	*	*	5	*	9	*	19	*	19	*
3309	7	*	*	4	*	*	1	*	0	*	7	*	7	*
3312	120	4.2 (61)	60.0 (41)	101	16.8 (27)	53.5 (51)	32	*	29	*	152	7.2 (54)	153	7.2 (10)
3550	9	*	*	0	*	*	5	*	6	*	9	*	9	*
3551	6	*	*	1	*	*	1	*	3	*	7	*	7	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 27

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3552	14	*	*	4	*	*	4	*	7	*	18	*	18	*
3701	277	3.2 (24)	64.3 (93)	186	15.6 (15)	56.5 (68)	45	*	92	*	324	10.8 (110)	323	16.4 (102)
3850	719	4.2 (61)	62.0 (63)	425	20.0 (59)	54.4 (55)	144	38.9 (3)	154	33.8 (23)	913	7.1 (52)	915	9.3 (27)
3901	448	3.6 (43)	57.4 (16)	198	16.7 (26)	57.6 (71)	60	*	124	27.4 (8)	503	8.2 (80)	503	12.7 (64)
3902	64	*	*	27	*	*	8	*	20	*	70	*	70	*
3903	15	*	*	7	*	*	2	*	10	*	17	*	17	*
3904	3	*	*	1	*	*	2	*	0	*	4	*	4	*
3905	7	*	*	5	*	*	0	*	2	*	9	*	9	*
3950	674	3.4 (33)	63.1 (82)	382	18.3 (37)	51.8 (37)	119	39.5 (4)	192	38.0 (31)	745	10.1 (106)	745	10.2 (37)
3954	9	*	*	5	*	*	2	*	2	*	9	*	9	*
4103	0	*	*	0	*	*	0	*	0	*	1	*	1	*
4106	640	3.3 (28)	62.3 (66)	325	25.5 (78)	42.8 (6)	163	38.7 (2)	318	29.6 (12)	794	4.9 (13)	795	11.9 (57)
4107	796	3.6 (43)	59.7 (39)	349	14.3 (10)	48.7 (19)	184	51.6 (21)	409	34.7 (24)	921	7.2 (54)	923	14.5 (83)
4109	148	4.1 (58)	59.5 (36)	75	*	*	56	*	83	*	290	7.9 (72)	290	13.8 (73)
4110	5	*	*	5	*	*	0	*	0	*	5	*	5	*
4112	176	2.3 (7)	60.2 (47)	153	76.5 (81)	5.2 (1)	28	*	25	*	168	19.6 (117)	169	13.0 (67)
4115	728	6.0 (95)	57.0 (14)	147	19.7 (55)	46.9 (8)	250	41.2 (6)	420	33.3 (20)	1,168	8.0 (76)	1,169	13.9 (75)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 28

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4351	163	3.1 (21)	65.0 (97)	145	17.9 (34)	50.3 (28)	22	*	37	*	161	9.3 (97)	161	13.7 (72)
4352	50	*	*	35	*	*	7	*	10	*	50	*	51	*
4353	102	2.9 (15)	67.6 (103)	65	*	*	22	*	20	*	113	5.3 (21)	112	14.3 (79)
4354	209	3.3 (28)	61.2 (56)	120	12.5 (3)	55.8 (63)	27	*	56	*	236	8.1 (79)	235	14.0 (77)
4355	83	*	*	30	*	*	20	*	36	*	111	9.0 (94)	111	16.2 (101)
4356	124	3.2 (24)	58.9 (29)	78	*	*	32	*	33	*	135	10.4 (109)	135	11.1 (48)
4358	33	*	*	24	*	*	8	*	6	*	34	*	34	*
4359	151	3.3 (28)	60.3 (48)	114	19.3 (50)	48.2 (16)	31	*	38	*	165	10.3 (107)	165	14.5 (83)
4360	66	*	*	54	*	*	8	*	16	*	60	*	60	*
4361	81	*	*	63	*	*	11	*	26	*	85	*	85	*
4362	85	*	*	70	*	*	19	*	30	*	95	*	95	*
4363	248	5.2 (85)	57.7 (18)	132	19.7 (55)	47.0 (9)	39	*	51	*	253	6.7 (45)	253	13.0 (67)
4401	338	3.0 (16)	56.2 (13)	306	18.0 (35)	53.3 (50)	61	*	91	*	320	4.4 (7)	320	7.5 (13)
4501	223	3.1 (21)	61.4 (58)	168	16.1 (20)	58.9 (75)	39	*	64	*	287	9.8 (102)	287	15.3 (95)
4601	334	3.0 (16)	61.4 (58)	52	*	*	91	*	119	42.9 (32)	393	8.7 (91)	393	11.7 (55)
4602	1	*	*	0	*	*	0	*	0	*	1	*	1	*
4701	349	5.2 (85)	49.6 (1)	221	21.7 (68)	48.4 (18)	53	*	129	13.2 (1)	329	13.4 (115)	329	15.8 (97)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 29

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5050	523	3.3 (28)	65.0 (97)	457	20.4 (63)	51.4 (34)	121	46.3 (10)	151	27.8 (9)	502	6.4 (38)	502	11.2 (50)
5051	832	4.7 (75)	59.0 (32)	711	20.3 (62)	49.6 (23)	195	45.1 (8)	297	28.6 (10)	834	6.7 (45)	834	10.0 (34)
5052	907	3.5 (38)	61.2 (56)	762	19.3 (50)	51.2 (33)	196	36.2 (1)	253	26.9 (6)	913	8.7 (91)	914	9.5 (29)
5055	165	1.2 (2)	58.8 (24)	51	*	*	43	*	81	*	380	7.6 (65)	380	3.2 (3)
5101	458	3.5 (38)	57.6 (17)	281	17.1 (29)	57.7 (72)	90	*	170	20.6 (4)	504	8.5 (86)	504	12.5 (61)
5102	13	*	*	8	*	*	1	*	5	*	12	*	12	*
5202	236	3.4 (33)	60.6 (51)	173	16.2 (21)	53.8 (52)	38	*	59	*	230	4.8 (11)	232	16.4 (102)
5601	319	3.4 (33)	63.0 (79)	21	*	*	49	*	127	14.2 (2)	316	7.9 (72)	316	13.3 (70)
5801	505	3.0 (16)	64.0 (91)	320	24.4 (76)	47.2 (10)	115	50.4 (19)	125	32.0 (16)	556	7.0 (49)	556	15.8 (97)
5803	44	*	*	25	*	*	3	*	15	*	44	*	44	*
5804	7	*	*	7	*	*	0	*	4	*	7	*	7	*
5806	10	*	*	6	*	*	0	*	1	*	17	*	18	*
6101	11	*	*	7	*	*	1	*	1	*	9	*	9	*
6102	43	*	*	33	*	*	9	*	11	*	44	*	44	*
6103	1,085	4.4 (70)	63.7 (89)	696	19.7 (55)	51.7 (36)	157	47.1 (12)	160	20.0 (3)	1,109	8.4 (85)	1,110	18.0 (108)
6105	5	*	*	4	*	*	0	*	0	*	4	*	4	*
6106	82	*	*	58	*	*	14	*	8	*	79	*	79	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 30

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6301	1,102	3.5 (38)	58.0 (19)	753	18.9 (45)	55.1 (59)	213	45.5 (9)	250	31.2 (15)	1,119	6.5 (41)	1,120	11.3 (53)
6312	570	4.7 (75)	58.8 (24)	396	17.4 (30)	50.5 (29)	112	52.7 (22)	172	33.7 (22)	576	5.2 (20)	576	8.3 (19)
6316	416	3.8 (49)	60.1 (44)	302	15.9 (17)	56.6 (69)	98	*	103	35.0 (26)	409	5.6 (29)	409	9.5 (29)
6318	11	*	*	8	*	*	6	*	3	*	12	*	12	*
6320	39	*	*	27	*	*	8	*	19	*	49	*	49	*
6325	84	*	*	42	*	*	20	*	39	*	92	*	92	*
6335	265	4.9 (81)	54.7 (8)	100	15.0 (14)	50.0 (25)	54	*	83	*	370	8.6 (87)	370	7.6 (16)
6340	18	*	*	6	*	*	1	*	4	*	14	*	14	*
6345	31	*	*	18	*	*	10	*	15	*	32	*	32	*
6360	240	4.2 (61)	58.8 (24)	128	16.4 (23)	53.1 (48)	47	*	67	*	276	7.6 (65)	276	15.2 (93)
7304	10	*	*	8	*	*	1	*	3	*	10	*	10	*
7306	651	3.5 (38)	58.8 (24)	502	18.3 (37)	51.8 (37)	145	48.3 (17)	149	28.9 (11)	754	7.7 (69)	755	11.0 (45)
7308	9	*	*	7	*	*	0	*	0	*	8	*	8	*
7310	1	*	*	0	*	*	0	*	2	*	2	*	2	*
7401	719	6.3 (101)	55.6 (10)	632	17.4 (30)	55.5 (61)	108	40.7 (5)	129	37.2 (30)	708	8.2 (80)	709	15.7 (96)
7601	33	*	*	18	*	*	39	*	36	*	193	7.3 (58)	193	8.3 (19)
7850	392	3.8 (49)	63.0 (79)	328	18.6 (44)	47.6 (12)	64	*	88	*	369	9.8 (102)	370	14.3 (79)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 31

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7852	6	*	*	4	*	*	0	*	1	*	6	*	6	*
8101	395	5.1 (84)	68.6 (105)	212	19.3 (50)	55.7 (62)	99	*	87	*	523	5.5 (27)	524	18.3 (111)
8102	117	4.3 (67)	62.4 (67)	56	*	*	26	*	14	*	139	7.2 (54)	139	10.8 (41)
8105	501	3.2 (24)	62.7 (73)	278	17.6 (32)	50.7 (31)	104	67.3 (26)	111	45.0 (33)	491	6.3 (36)	490	11.0 (45)
8106	80	*	*	48	*	*	36	*	40	*	164	4.9 (13)	163	10.4 (38)
8150	11	*	*	6	*	*	4	*	6	*	11	*	11	*
8151	948	3.9 (51)	60.8 (52)	637	21.4 (66)	48.7 (19)	200	47.0 (11)	312	33.0 (19)	960	7.0 (49)	962	12.6 (63)
8152	11	*	*	8	*	*	2	*	4	*	10	*	10	*
8153	3	*	*	1	*	*	0	*	2	*	2	*	2	*
8155	4	*	*	3	*	*	1	*	3	*	4	*	4	*
8157	5	*	*	3	*	*	1	*	3	*	4	*	4	*
8201	581	4.1 (58)	62.7 (73)	384	18.5 (41)	56.0 (64)	151	49.7 (18)	272	32.7 (18)	727	6.7 (45)	728	10.6 (40)
8202	739	4.3 (67)	52.1 (3)	297	25.3 (77)	40.1 (4)	232	47.8 (15)	267	34.8 (25)	1,206	5.4 (24)	1,205	2.4 (1)
8203	119	3.4 (33)	63.9 (90)	103	9.7 (1)	61.2 (79)	29	*	36	*	122	7.4 (61)	122	21.3 (117)
8250	221	5.9 (94)	62.0 (63)	173	18.5 (41)	61.3 (80)	64	*	76	*	219	8.2 (80)	219	17.8 (107)
8252	46	*	*	30	*	*	21	*	15	*	54	*	54	*
8255	729	4.0 (55)	60.5 (50)	508	14.2 (8)	55.1 (59)	189	50.8 (20)	292	36.3 (28)	804	7.8 (71)	804	10.8 (41)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 32

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8257	481	4.0 (55)	54.3 (6)	320	28.4 (79)	33.4 (3)	121	55.4 (23)	187	36.4 (29)	643	10.3 (107)	643	5.1 (4)
8263	226	5.8 (93)	61.9 (61)	185	17.8 (33)	58.9 (75)	49	*	101	22.8 (5)	228	5.3 (21)	228	7.5 (13)
8264	184	3.3 (28)	60.9 (53)	108	13.0 (5)	62.0 (81)	57	*	87	*	239	3.3 (1)	239	7.1 (9)
8267	144	4.2 (61)	62.5 (69)	110	20.9 (65)	43.6 (7)	60	*	80	*	220	4.1 (4)	221	9.0 (23)
8901	301	2.3 (7)	62.8 (76)	209	20.1 (60)	47.8 (13)	68	*	57	*	318	4.1 (4)	319	14.7 (88)
8902	432	4.6 (72)	63.2 (83)	250	19.6 (54)	56.0 (64)	88	*	93	*	461	5.4 (24)	461	13.2 (69)
8903	472	2.3 (7)	66.1 (102)	292	13.4 (7)	60.6 (78)	89	*	71	*	488	5.7 (31)	487	16.4 (102)
8904	670	3.9 (51)	63.4 (85)	385	15.8 (16)	58.4 (73)	135	67.4 (27)	128	53.1 (34)	752	5.5 (27)	753	15.1 (91)
8905	424	2.6 (12)	63.0 (79)	217	18.9 (45)	53.9 (53)	89	*	95	*	487	8.6 (87)	487	11.5 (54)
8909	159	2.5 (11)	59.7 (39)	79	*	*	39	*	38	*	220	9.1 (96)	221	18.1 (109)
9045	437	1.8 (3)	62.2 (65)	312	21.8 (69)	48.1 (15)	105	43.8 (7)	89	*	567	6.5 (41)	567	8.8 (22)
9050	245	4.9 (81)	58.0 (19)	81	*	*	71	*	46	*	316	6.6 (43)	316	9.2 (25)
9051	448	3.6 (43)	68.5 (104)	303	19.1 (47)	41.9 (5)	80	*	59	*	521	7.3 (58)	520	7.3 (11)
9053	313	3.2 (24)	64.2 (92)	226	14.2 (8)	56.2 (67)	53	*	47	*	323	5.6 (29)	323	14.6 (86)
9055	9	*	*	6	*	*	0	*	0	*	9	*	9	*
9058	219	6.8 (104)	62.6 (72)	147	17.0 (28)	49.0 (22)	74	*	56	*	334	5.1 (17)	334	14.1 (78)
9066	231	3.0 (16)	64.5 (95)	119	14.3 (10)	54.6 (58)	78	*	56	*	342	5.0 (15)	342	17.5 (105)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Page : 33

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9068	284	2.1 (6)	68.7 (106)	173	21.4 (66)	60.1 (77)	47	*	33	*	280	3.6 (3)	280	12.5 (61)
9069	766	3.5 (38)	61.0 (54)	606	21.9 (70)	48.8 (21)	165	57.6 (24)	99	*	856	5.3 (21)	854	9.7 (31)
9070	478	4.0 (55)	60.0 (41)	302	21.9 (70)	51.0 (32)	87	*	75	*	500	5.4 (24)	500	11.0 (45)
9072	434	2.8 (14)	65.2 (99)	244	19.7 (55)	52.0 (40)	96	*	81	*	519	7.7 (69)	520	14.8 (90)
9080	701	3.1 (21)	62.5 (69)	393	14.8 (13)	58.8 (74)	185	60.5 (25)	152	33.6 (21)	886	6.3 (36)	886	11.1 (48)
9082	44	*	*	25	*	*	10	*	7	*	42	*	42	*
9083	64	*	*	61	*	*	0	*	0	*	65	*	65	*
9084	12	*	*	7	*	*	2	*	0	*	17	*	17	*
9085	4	*	*	2	*	*	0	*	1	*	2	*	2	*
9086	12	*	*	5	*	*	0	*	4	*	10	*	10	*
9087	1	*	*	1	*	*	0	*	1	*	1	*	1	*
Michigan	43,733	4.2	60.6	27,372	19.1	51.5	9,098	45.0	12,203	30.7	49,829	7.3	49,855	11.9
National (Prior Year)	159,852	4.1	60.3	114,229	22.5	48.3	44,569	18.2	57,476	18.5	100,642	10.0	99,796	9.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 34

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	30	*	*	33	*	30	*	26	*
0102	132	84.1 (65)	10.6 (87)	140	28.6 (69)	130	20.0 (55)	119	95.0 (5)
0103	137	78.8 (89)	10.9 (88)	146	36.3 (28)	129	17.8 (63)	115	94.8 (8)
0104	36	*	*	41	*	33	*	30	*
0105	2	*	*	1	*	0	*	1	*
0250	41	*	*	47	*	37	*	43	*
0251	28	*	*	32	*	31	*	26	*
0252	36	*	*	40	*	33	*	25	*
0253	39	*	*	39	*	38	*	35	*
0401	145	80.0 (86)	13.1 (94)	155	43.9 (6)	145	22.8 (41)	128	89.8 (40)
0402	132	81.1 (81)	12.9 (93)	137	35.8 (30)	126	20.6 (50)	112	89.3 (46)
0403	48	*	*	52	*	43	*	36	*
0404	19	*	*	20	*	17	*	15	*
0501	166	91.0 (25)	3.6 (38)	168	35.1 (35)	148	25.0 (36)	128	92.2 (29)
0503	34	*	*	33	*	31	*	29	*
0504	149	88.6 (41)	5.4 (52)	154	30.5 (61)	149	29.5 (21)	128	85.9 (72)
0505	86	*	*	89	*	82	*	75	*
0506	54	*	*	53	*	49	*	47	*
0507	24	*	*	24	*	20	*	14	*
0508	10	*	*	10	*	9	*	7	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 35

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0509	1	*	*	1	*	1	*	1	*
0650	333	85.3 (54)	9.6 (82)	355	34.1 (45)	313	24.9 (37)	279	92.8 (22)
0651	196	75.5 (102)	16.8 (106)	210	46.7 (2)	179	25.7 (32)	174	85.6 (73)
0652	111	86.5 (47)	9.9 (85)	112	34.8 (38)	99	*	79	*
0653	108	75.9 (100)	19.4 (111)	116	37.1 (26)	109	32.1 (18)	98	*
0654	106	74.5 (105)	17.9 (109)	118	34.7 (39)	104	17.3 (69)	88	*
0655	101	85.1 (57)	3.0 (27)	115	43.5 (7)	103	27.2 (26)	98	*
0657	27	*	*	27	*	23	*	22	*
0701	107	81.3 (79)	14.0 (99)	120	39.2 (19)	107	38.3 (10)	91	*
0702	21	*	*	22	*	22	*	18	*
0703	58	*	*	63	*	56	*	51	*
0704	19	*	*	21	*	20	*	17	*
0705	38	*	*	42	*	40	*	39	*
0706	7	*	*	7	*	6	*	4	*
0850	241	86.7 (46)	4.1 (42)	266	27.8 (72)	241	24.9 (37)	192	92.2 (29)
0851	203	89.7 (33)	7.4 (71)	226	33.2 (49)	212	25.9 (31)	178	95.5 (4)
0901	477	76.1 (97)	18.9 (110)	515	39.6 (17)	473	15.4 (84)	427	89.2 (47)
0902	16	*	*	18	*	18	*	18	*
1051	373	82.3 (73)	11.0 (90)	399	27.6 (73)	350	22.0 (43)	280	87.1 (64)
1101	594	69.5 (115)	14.1 (100)	652	29.1 (67)	573	14.8 (87)	487	86.4 (70)
1102	251	74.9 (104)	13.5 (96)	290	35.2 (33)	259	16.2 (78)	221	92.3 (27)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 36

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
1104	1	*	*	1	*	1	*	0 *
1106	25	*	*	29	*	29	*	23 *
1250	279	82.8 (71)	9.0 (76)	340	32.9 (53)	296	17.2 (70)	251 81.7 (98)
1251	229	80.3 (84)	11.8 (91)	251	31.9 (56)	216	19.0 (57)	189 84.1 (83)
1252	178	66.9 (116)	19.7 (112)	200	40.0 (15)	188	17.6 (66)	167 88.6 (52)
1253	230	74.3 (107)	15.7 (103)	264	35.2 (33)	236	21.2 (46)	205 87.3 (60)
1350	664	88.0 (44)	6.5 (62)	743	36.2 (29)	510	21.2 (46)	561 82.4 (94)
1351	48	*	*	52	*	47	*	36 *
1352	110	92.7 (9)	6.4 (61)	118	33.1 (50)	100	4.0 (109)	92 *
1401	84	*	*	89	*	74	*	71 *
1402	35	*	*	36	*	29	*	28 *
1403	9	*	*	9	*	7	*	6 *
1501	257	81.7 (76)	13.6 (97)	278	41.4 (12)	183	18.0 (61)	225 83.1 (91)
1503	311	81.0 (82)	10.9 (88)	337	44.8 (3)	278	33.8 (13)	277 80.5 (101)
1504	633	90.5 (28)	2.7 (23)	697	30.7 (59)	606	29.9 (20)	516 82.4 (94)
1505	138	92.0 (17)	5.8 (58)	154	26.0 (79)	118	28.0 (23)	99 *
1506	184	71.7 (113)	21.7 (117)	205	34.6 (42)	176	2.8 (112)	168 94.6 (9)
1507	178	91.6 (18)	3.9 (41)	189	42.3 (10)	146	25.3 (34)	147 89.1 (48)
1508	14	*	*	15	*	15	*	15 *
1509	125	94.4 (4)	1.6 (13)	151	33.1 (50)	118	18.6 (59)	116 90.5 (35)
1511	204	90.7 (27)	2.5 (20)	223	35.0 (37)	182	15.9 (81)	166 83.7 (88)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 37

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1512	116	92.2 (14)	5.2 (50)	123	25.2 (80)	96	*	84	*
1514	96	*	*	104	29.8 (65)	80	*	70	*
1701	196	85.2 (55)	9.2 (78)	217	39.2 (19)	180	42.8 (4)	168	84.5 (79)
1950	187	71.7 (113)	20.9 (114)	197	35.5 (31)	176	25.6 (33)	155	93.5 (18)
1952	34	*	*	34	*	29	*	25	*
1953	109	76.1 (97)	17.4 (108)	113	38.9 (21)	102	20.6 (50)	94	*
1954	164	76.2 (96)	16.5 (104)	172	37.8 (24)	152	28.3 (22)	136	95.6 (3)
1955	33	*	*	37	*	33	*	30	*
1957	138	85.5 (51)	9.4 (80)	144	32.6 (54)	123	26.8 (28)	104	89.4 (45)
1963	23	*	*	24	*	22	*	20	*
1964	14	*	*	16	*	16	*	16	*
2150	199	92.5 (11)	1.5 (11)	217	42.9 (8)	198	24.2 (39)	175	92.6 (26)
2151	71	*	*	85	*	64	*	60	*
2251	117	89.7 (33)	6.8 (66)	131	35.1 (35)	120	40.8 (7)	108	96.3 (1)
2252	56	*	*	66	*	51	*	51	*
2554	17	*	*	18	*	15	*	14	*
2555	2,414	86.3 (48)	4.2 (43)	2,572	24.7 (81)	1,996	3.9 (110)	1,879	86.2 (71)
2556	124	77.4 (92)	6.5 (62)	135	14.8 (118)	96	*	64	*
2557	1	*	*	1	*	1	*	1	*
2558	3	*	*	3	*	3	*	3	*
2560	1	*	*	2	*	2	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 38

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
2561	2	*	*	2	*	2	*	2 *
2562	10	*	*	10	*	9	*	9 *
2563	15	*	*	14	*	11	*	6 *
2564	7	*	*	7	*	6	*	5 *
3001	352	81.5 (77)	6.8 (66)	380	36.6 (27)	327	11.9 (95)	309 81.6 (99)
3002	107	94.4 (4)	0.9 (3)	116	44.8 (3)	107	11.2 (97)	102 85.3 (74)
3004	76	*	*	84	*	76	*	71 *
3005	18	*	*	19	*	16	*	10 *
3007	4	*	*	4	*	4	*	4 *
3201	137	87.6 (45)	8.0 (73)	149	32.2 (55)	126	32.5 (17)	112 87.5 (57)
3301	1,357	84.8 (61)	5.7 (56)	1,454	29.2 (66)	1,300	8.0 (107)	1,087 90.0 (38)
3304	39	*	*	42	*	41	*	35 *
3305	56	*	*	65	*	62	*	49 *
3306	76	*	*	84	*	76	*	61 *
3308	19	*	*	22	*	17	*	16 *
3309	7	*	*	8	*	8	*	8 *
3312	151	88.7 (40)	2.0 (15)	157	23.6 (86)	131	10.7 (100)	113 92.0 (31)
3550	9	*	*	11	*	11	*	10 *
3551	7	*	*	7	*	7	*	6 *
3552	18	*	*	19	*	17	*	16 *
3701	318	95.3 (2)	2.8 (25)	353	33.4 (48)	312	21.5 (45)	261 91.6 (32)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 39

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3850	908	84.1 (65)	8.5 (74)	989	38.7 (22)	878	17.1 (71)	798	85.3 (74)
3901	499	85.0 (58)	7.6 (72)	558	44.4 (5)	512	14.5 (89)	467	79.7 (102)
3902	70	*	*	82	*	79	*	69	*
3903	18	*	*	18	*	17	*	12	*
3904	4	*	*	4	*	4	*	4	*
3905	9	*	*	9	*	9	*	9	*
3950	740	84.3 (64)	9.1 (77)	807	34.7 (39)	722	16.5 (75)	620	84.4 (81)
3954	9	*	*	10	*	9	*	7	*
4103	1	*	*	1	*	1	*	1	*
4106	775	85.8 (50)	4.9 (49)	860	22.7 (90)	727	7.0 (108)	661	87.4 (59)
4107	924	85.2 (55)	5.3 (51)	983	21.8 (93)	817	26.3 (30)	717	91.5 (33)
4109	287	77.7 (91)	14.3 (101)	306	31.0 (58)	283	15.9 (81)	222	95.0 (5)
4110	5	*	*	5	*	5	*	0	*
4112	168	47.6 (117)	0.0 (1)	178	27.0 (75)	168	95.2 (1)	156	92.9 (20)
4115	1,159	90.3 (29)	4.5 (48)	1,238	22.5 (92)	1,093	16.1 (79)	920	90.4 (36)
4351	161	88.8 (39)	3.7 (40)	169	40.2 (14)	65	*	136	93.4 (19)
4352	52	*	*	56	*	51	*	48	*
4353	111	73.0 (111)	21.6 (116)	129	34.1 (45)	106	23.6 (40)	96	*
4354	235	88.1 (42)	6.0 (59)	247	41.7 (11)	184	17.4 (68)	196	88.8 (51)
4355	112	84.8 (61)	13.4 (95)	116	30.2 (62)	102	77.5 (2)	93	*
4356	133	77.4 (92)	16.5 (104)	142	33.8 (47)	124	22.6 (42)	117	87.2 (62)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 40

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
4358	35	*	*	36	*	35	*	31 *
4359	161	74.5 (105)	19.9 (113)	172	33.1 (50)	147	27.2 (26)	139 92.8 (22)
4360	60	*	*	68	*	63	*	55 *
4361	83	*	*	92	*	83	*	75 *
4362	95	*	*	98	*	85	*	73 *
4363	252	73.4 (110)	15.5 (102)	273	39.6 (17)	240	27.9 (25)	223 82.1 (96)
4401	319	82.8 (71)	10.3 (86)	337	30.0 (64)	309	14.6 (88)	257 94.9 (7)
4501	288	81.9 (75)	9.7 (84)	295	31.5 (57)	257	26.8 (28)	218 94.0 (13)
4601	386	92.2 (14)	3.6 (38)	428	23.4 (87)	372	18.3 (60)	321 89.1 (48)
4602	1	*	*	1	*	1	*	1 *
4701	328	91.2 (22)	3.0 (27)	359	21.2 (95)	267	41.6 (5)	237 90.3 (37)
5050	501	91.2 (22)	3.4 (33)	534	17.2 (109)	355	18.0 (61)	355 94.6 (9)
5051	831	92.5 (11)	2.0 (15)	882	17.0 (110)	661	14.2 (90)	582 93.6 (17)
5052	906	90.3 (29)	2.4 (19)	972	20.5 (99)	633	10.9 (99)	659 93.9 (14)
5055	377	85.4 (53)	13.8 (98)	386	50.3 (1)	338	8.9 (104)	348 84.8 (76)
5101	499	90.0 (32)	3.2 (29)	534	20.4 (100)	350	28.0 (23)	382 93.7 (16)
5102	12	*	*	11	*	10	*	4 *
5202	228	91.2 (22)	4.4 (46)	264	34.5 (43)	201	40.3 (8)	217 89.9 (39)
5601	316	94.3 (7)	1.3 (7)	333	21.0 (98)	299	33.4 (15)	241 94.2 (11)
5801	554	83.2 (69)	7.0 (69)	597	30.7 (59)	542	20.1 (54)	473 89.6 (43)
5803	43	*	*	47	*	39	*	29 *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 41

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
5804	7	*	*	8	*	7	*	3 *
5806	18	*	*	20	*	16	*	11 *
6101	9	*	*	11	*	10	*	9 *
6102	43	*	*	46	*	41	*	37 *
6103	1,094	88.1 (42)	5.5 (53)	1,184	39.7 (16)	1,006	15.6 (83)	920 81.6 (99)
6105	3	*	*	5	*	4	*	4 *
6106	75	*	*	85	*	81	*	78 *
6301	1,097	84.9 (59)	8.8 (75)	1,221	27.9 (71)	954	11.6 (96)	919 84.5 (79)
6312	556	89.2 (36)	5.6 (54)	615	24.1 (83)	477	16.4 (76)	437 83.5 (89)
6316	408	91.4 (19)	2.9 (26)	449	23.2 (89)	374	19.0 (57)	317 92.7 (24)
6318	12	*	*	14	*	13	*	13 *
6320	49	*	*	51	*	32	*	27 *
6325	91	*	*	103	21.4 (94)	91	*	64 *
6335	368	89.1 (37)	9.2 (78)	382	38.7 (22)	124	33.1 (16)	305 93.8 (15)
6340	14	*	*	20	*	18	*	20 *
6345	31	*	*	37	*	33	*	23 *
6360	273	82.1 (74)	9.5 (81)	304	28.0 (70)	231	17.7 (64)	220 87.7 (54)
7304	10	*	*	10	*	10	*	10 *
7306	743	83.0 (70)	6.5 (62)	808	30.2 (62)	702	16.1 (79)	608 86.5 (67)
7308	8	*	*	10	*	10	*	10 *
7310	2	*	*	2	*	2	*	2 *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 42

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7401	702	72.2 (112)	21.1 (115)	755	37.4 (25)	629	20.8 (48)	609	84.2 (82)
7601	191	89.5 (35)	7.3 (70)	211	40.3 (13)	196	36.2 (12)	183	92.3 (27)
7850	366	76.5 (95)	16.9 (107)	404	34.7 (39)	361	20.8 (48)	318	86.8 (66)
7852	6	*	*	6	*	6	*	6	*
8101	524	75.6 (101)	3.2 (29)	544	18.2 (106)	272	15.1 (85)	378	86.5 (67)
8102	135	83.7 (68)	4.4 (46)	144	35.4 (32)	83	*	117	82.1 (96)
8105	493	92.1 (16)	1.2 (5)	532	23.3 (88)	476	13.9 (92)	371	92.7 (24)
8106	157	81.5 (77)	5.7 (56)	171	19.9 (102)	124	3.2 (111)	110	87.3 (60)
8150	11	*	*	12	*	12	*	12	*
8151	963	90.3 (29)	3.2 (29)	1,055	26.6 (77)	939	16.4 (76)	808	91.0 (34)
8152	10	*	*	13	*	12	*	12	*
8153	2	*	*	3	*	3	*	2	*
8155	4	*	*	4	*	4	*	4	*
8157	4	*	*	5	*	4	*	4	*
8201	721	92.8 (8)	1.2 (5)	800	17.5 (108)	610	36.4 (11)	522	84.1 (83)
8202	1,189	94.4 (4)	3.4 (33)	1,252	28.9 (68)	784	10.2 (101)	976	89.1 (48)
8203	122	91.0 (25)	1.6 (13)	135	15.6 (116)	114	9.6 (102)	79	*
8250	216	86.1 (49)	1.4 (8)	242	12.8 (119)	199	20.6 (50)	161	87.6 (56)
8252	54	*	*	54	*	50	*	44	*
8255	795	91.4 (19)	1.0 (4)	860	16.9 (111)	480	30.4 (19)	574	92.9 (20)
8257	643	96.7 (1)	1.4 (8)	671	27.3 (74)	587	33.7 (14)	521	84.1 (83)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 43

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8263	228	94.7 (3)	0.4 (2)	242	16.1 (114)	100	41.0 (6)	156	96.2 (2)
8264	237	92.4 (13)	3.4 (33)	257	16.7 (112)	211	17.5 (67)	171	94.2 (11)
8267	217	92.6 (10)	1.4 (8)	247	21.1 (97)	217	56.7 (3)	191	89.5 (44)
8901	315	80.0 (86)	3.5 (36)	341	23.8 (84)	226	13.7 (93)	261	83.5 (89)
8902	461	85.5 (51)	2.2 (18)	496	21.2 (95)	334	17.7 (64)	364	82.7 (93)
8903	480	80.2 (85)	3.3 (32)	509	23.8 (84)	350	21.7 (44)	358	74.3 (104)
8904	746	84.5 (63)	1.5 (11)	786	18.3 (105)	496	16.9 (72)	547	86.5 (67)
8905	483	91.3 (21)	2.1 (17)	504	26.4 (78)	347	25.1 (35)	368	84.0 (86)
8909	221	89.1 (37)	2.7 (23)	234	22.6 (91)	136	20.6 (50)	174	89.7 (41)
9045	570	74.0 (108)	11.8 (91)	617	34.5 (43)	168	39.9 (9)	505	84.6 (78)
9050	310	76.1 (97)	6.8 (66)	323	20.4 (100)	264	19.7 (56)	230	84.8 (76)
9051	502	78.7 (90)	9.6 (82)	542	42.4 (9)	490	16.7 (73)	452	78.3 (103)
9053	324	84.9 (59)	2.5 (20)	337	16.6 (113)	277	11.2 (97)	204	89.7 (41)
9055	9	*	*	9	*	2	*	0	*
9058	334	84.1 (65)	6.3 (60)	347	24.5 (82)	232	8.6 (106)	253	87.7 (54)
9066	343	75.2 (103)	6.7 (65)	362	15.2 (117)	274	12.0 (94)	234	87.2 (62)
9068	280	78.9 (88)	4.3 (44)	293	27.0 (75)	234	15.0 (86)	219	83.1 (91)
9069	856	81.2 (80)	2.6 (22)	879	15.8 (115)	494	9.3 (103)	550	87.5 (57)
9070	498	73.9 (109)	5.6 (54)	534	18.9 (104)	315	14.0 (91)	357	87.1 (64)
9072	513	77.0 (94)	4.3 (44)	544	17.6 (107)	425	8.9 (104)	381	84.0 (86)
9080	879	80.4 (83)	3.5 (36)	939	19.2 (103)	647	16.7 (73)	680	87.9 (53)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Page : 44

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9082	42	*	*	43	*	40	*	34	*
9083	66	*	*	66	*	47	*	3	*
9084	17	*	*	18	*	10	*	8	*
9085	2	*	*	4	*	4	*	4	*
9086	10	*	*	12	*	11	*	12	*
9087	1	*	*	1	*	1	*	0	*
Michigan	49,415	84.9	6.6	53,324	28.5	42,172	19.1	39,761	39761.0
National (Prior Year)	125,577	72.2	5.9	159,369	19.0	98,317	21.7	96,069	88.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Page : 45

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	30	*	29	*	6	*	26	*	29	*
0102	130	43.1 (108)	133	29.3 (109)	45	*	119	16.8 (90)	133	3.0 (68)
0103	130	43.1 (108)	138	31.2 (114)	48	*	115	18.3 (95)	137	5.8 (106)
0104	34	*	36	*	13	*	30	*	36	*
0105	1	*	2	*	0	*	0	*	2	*
0250	45	*	41	*	19	*	43	*	41	*
0251	31	*	28	*	12	*	26	*	28	*
0252	34	*	36	*	5	*	25	*	36	*
0253	38	*	39	*	16	*	35	*	39	*
0401	146	51.4 (115)	148	34.5 (117)	64	*	128	12.5 (65)	148	3.4 (76)
0402	127	50.4 (113)	128	27.3 (104)	50	*	112	14.3 (77)	128	4.7 (94)
0403	43	*	48	*	11	*	36	*	48	*
0404	17	*	19	*	3	*	15	*	19	*
0501	150	38.7 (94)	166	30.1 (111)	48	*	128	20.3 (101)	166	2.4 (53)
0503	31	*	34	*	12	*	29	*	34	*
0504	149	42.3 (107)	147	25.2 (98)	53	*	128	13.3 (70)	147	4.8 (95)
0505	82	*	86	*	33	*	75	*	84	*
0506	49	*	54	*	18	*	47	*	54	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Page : 46

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0507	20	*	25	*	3	*	14	*	25	*
0508	9	*	10	*	2	*	7	*	10	*
0509	1	*	1	*	1	*	1	*	1	*
0650	326	38.7 (94)	330	24.5 (93)	101	34.7 (18)	279	14.3 (77)	331	2.7 (63)
0651	189	50.8 (114)	199	27.1 (103)	81	*	174	14.4 (79)	199	3.5 (77)
0652	99	*	111	30.6 (112)	35	*	79	*	111	10.8 (117)
0653	110	34.5 (77)	108	18.5 (74)	30	*	98	*	108	2.8 (64)
0654	104	40.4 (101)	107	28.0 (106)	28	*	88	*	107	3.7 (80)
0655	105	52.4 (116)	105	32.4 (116)	46	*	98	*	105	4.8 (95)
0657	23	*	27	*	11	*	22	*	27	*
0701	111	31.5 (72)	106	22.6 (88)	28	*	91	*	106	0.9 (13)
0702	22	*	22	*	6	*	18	*	22	*
0703	57	*	59	*	24	*	51	*	59	*
0704	20	*	19	*	7	*	17	*	19	*
0705	41	*	37	*	14	*	39	*	37	*
0706	6	*	7	*	0	*	4	*	7	*
0850	242	38.8 (98)	240	24.2 (92)	66	*	192	6.8 (23)	240	2.9 (67)
0851	212	33.5 (75)	205	17.1 (67)	57	*	178	9.0 (49)	205	5.4 (103)
0901	477	32.5 (73)	480	22.7 (89)	132	21.2 (25)	426	15.0 (81)	479	4.8 (95)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Page : 47

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0902	18	*	16	*	5	*	18	*	16	*
1051	350	35.1 (81)	374	20.1 (77)	101	35.6 (14)	280	8.9 (48)	374	2.1 (41)
1101	571	27.1 (61)	609	13.5 (56)	125	40.0 (9)	487	12.5 (65)	609	4.3 (87)
1102	260	41.9 (105)	259	22.0 (83)	82	*	221	15.8 (88)	259	6.2 (107)
1104	1	*	1	*	0	*	0	*	1	*
1106	29	*	25	*	8	*	23	*	25	*
1250	297	33.3 (74)	318	17.9 (71)	75	*	250	11.6 (60)	318	2.2 (47)
1251	216	44.0 (110)	232	31.0 (113)	77	*	189	17.5 (93)	232	3.0 (68)
1252	187	36.9 (89)	184	18.5 (74)	54	*	166	15.7 (87)	183	3.3 (74)
1253	236	41.9 (105)	238	26.5 (102)	72	*	204	10.8 (55)	237	4.2 (86)
1350	664	27.1 (61)	676	18.3 (72)	141	24.8 (23)	570	10.4 (53)	678	2.4 (53)
1351	48	*	48	*	10	*	36	*	48	*
1352	101	34.7 (79)	110	21.8 (82)	31	*	92	*	112	1.8 (37)
1401	82	*	84	*	27	*	71	*	84	*
1402	32	*	35	*	9	*	29	*	35	*
1403	8	*	9	*	3	*	6	*	9	*
1501	251	21.5 (35)	255	12.2 (51)	43	*	225	6.2 (18)	255	1.6 (30)
1503	314	23.2 (44)	313	10.2 (36)	64	*	277	8.3 (38)	313	1.3 (23)
1504	620	18.1 (24)	643	11.0 (43)	100	39.0 (11)	516	2.9 (6)	643	0.6 (7)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic

Page : 48

Comparison of Smoking Indicators <small>(2,3)</small>										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes <small>(4)</small>	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1505	130	22.3 (40)	141	10.6 (39)	22	*	99	*	140	2.1 (41)
1506	187	22.5 (42)	186	13.4 (55)	34	*	167	1.8 (3)	187	0.0 (1)
1507	168	31.0 (71)	181	13.8 (57)	46	*	146	14.4 (79)	181	2.8 (64)
1508	15	*	14	*	1	*	15	*	14	*
1509	127	26.0 (58)	123	12.2 (51)	27	*	115	12.2 (63)	123	0.8 (12)
1511	199	38.7 (94)	204	22.1 (84)	61	*	167	7.2 (27)	204	2.5 (56)
1512	108	25.9 (57)	117	17.1 (67)	26	*	85	*	117	0.0 (1)
1514	93	*	97	*	8	*	69	*	96	*
1701	190	44.7 (111)	195	31.8 (115)	70	*	167	15.0 (81)	197	6.6 (113)
1950	177	41.8 (104)	188	27.7 (105)	60	*	155	14.2 (76)	189	5.3 (102)
1952	29	*	34	*	10	*	25	*	34	*
1953	104	41.3 (103)	110	25.5 (99)	37	*	94	*	110	4.5 (92)
1954	153	35.3 (82)	165	20.0 (76)	45	*	136	8.8 (47)	165	2.4 (53)
1955	33	*	33	*	10	*	30	*	33	*
1957	131	29.8 (68)	136	17.6 (69)	31	*	104	7.7 (29)	138	2.2 (47)
1963	23	*	23	*	8	*	20	*	24	*
1964	16	*	14	*	5	*	16	*	14	*
2150	198	46.5 (112)	200	25.0 (97)	77	*	175	13.7 (73)	200	5.5 (105)
2151	76	*	73	*	22	*	60	*	72	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic

Page : 49

Comparison of Smoking Indicators ^(2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes ⁽⁴⁾	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2251	120	34.2 (76)	118	26.3 (100)	35	*	108	8.3 (38)	117	1.7 (32)
2252	54	*	54	*	20	*	51	*	54	*
2554	15	*	16	*	3	*	14	*	16	*
2555	2,252	26.6 (59)	2,429	15.2 (63)	466	35.6 (14)	1,875	9.0 (49)	2,421	1.7 (32)
2556	111	23.4 (47)	132	14.4 (59)	18	*	66	*	132	1.5 (28)
2557	1	*	1	*	1	*	1	*	1	*
2558	3	*	3	*	0	*	3	*	3	*
2560	2	*	1	*	0	*	2	*	1	*
2561	2	*	2	*	1	*	2	*	2	*
2562	9	*	10	*	0	*	9	*	10	*
2563	11	*	15	*	2	*	6	*	15	*
2564	6	*	7	*	1	*	5	*	7	*
3001	328	35.4 (83)	356	14.9 (62)	103	45.6 (6)	309	11.3 (59)	355	2.5 (56)
3002	107	24.3 (53)	108	11.1 (46)	24	*	102	9.8 (52)	108	0.9 (13)
3004	76	*	75	*	16	*	71	*	75	*
3005	16	*	18	*	7	*	10	*	18	*
3007	4	*	5	*	0	*	4	*	5	*
3201	136	39.7 (100)	138	22.5 (87)	43	*	112	18.8 (97)	138	5.1 (101)
3301	1,303	18.3 (25)	1,369	10.4 (37)	199	37.7 (13)	1,079	8.3 (38)	1,362	3.8 (82)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Page : 50

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3304	41	*	39	*	3	*	34	*	39	*
3305	62	*	55	*	6	*	49	*	55	*
3306	76	*	77	*	11	*	61	*	76	*
3308	18	*	19	*	4	*	16	*	19	*
3309	8	*	7	*	1	*	8	*	7	*
3312	131	19.8 (31)	153	11.1 (46)	23	*	113	4.4 (10)	153	0.7 (10)
3550	11	*	9	*	3	*	10	*	9	*
3551	7	*	7	*	2	*	6	*	7	*
3552	17	*	18	*	12	*	16	*	18	*
3701	312	36.2 (85)	324	22.2 (85)	95	*	261	11.1 (58)	324	2.5 (56)
3850	897	38.7 (94)	914	26.3 (100)	288	30.2 (19)	796	13.1 (69)	915	2.8 (64)
3901	516	24.8 (54)	500	14.0 (58)	103	30.1 (20)	467	6.9 (24)	500	1.8 (37)
3902	78	*	69	*	17	*	69	*	70	*
3903	17	*	18	*	4	*	12	*	18	*
3904	4	*	4	*	2	*	4	*	4	*
3905	9	*	9	*	2	*	9	*	8	*
3950	726	23.7 (50)	746	15.4 (65)	136	25.0 (22)	619	8.4 (43)	744	1.2 (21)
3954	9	*	9	*	3	*	7	*	9	*
4103	1	*	1	*	0	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Page : 51

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4106	745	15.0 (11)	796	8.3 (29)	93	*	657	0.8 (1)	792	0.9 (13)
4107	890	22.8 (43)	927	11.0 (43)	148	44.6 (7)	719	7.8 (32)	925	3.1 (72)
4109	284	28.5 (65)	291	14.8 (61)	63	*	221	3.2 (7)	290	1.4 (24)
4110	5	*	5	*	0	*	0	*	5	*
4112	169	4.1 (3)	169	1.8 (3)	7	*	127	6.3 (20)	158	2.5 (56)
4115	1,105	15.4 (15)	1,168	7.2 (23)	131	39.7 (10)	912	4.5 (11)	1,166	0.9 (13)
4351	148	34.5 (77)	161	24.8 (96)	47	*	136	15.4 (86)	161	3.1 (72)
4352	51	*	51	*	28	*	48	*	51	*
4353	106	23.6 (48)	113	12.4 (53)	20	*	96	*	113	0.9 (13)
4354	217	36.4 (86)	236	24.6 (94)	72	*	196	16.3 (89)	236	3.0 (68)
4355	106	36.8 (88)	111	28.8 (107)	34	*	93	*	111	5.4 (103)
4356	126	38.1 (92)	135	21.5 (81)	44	*	117	13.7 (73)	135	4.4 (90)
4358	35	*	34	*	7	*	31	*	34	*
4359	161	28.0 (63)	165	10.9 (41)	39	*	139	4.3 (9)	165	0.6 (7)
4360	64	*	60	*	17	*	55	*	60	*
4361	83	*	85	*	26	*	75	*	85	*
4362	85	*	95	*	32	*	73	*	95	*
4363	245	35.9 (84)	253	29.2 (108)	71	*	223	13.5 (72)	253	4.3 (87)
4401	309	26.9 (60)	320	15.3 (64)	57	*	257	6.2 (18)	320	0.9 (13)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Page : 52

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4501	258	39.1 (99)	288	23.6 (90)	84	*	218	12.8 (68)	288	1.4 (24)
4601	373	34.9 (80)	393	15.8 (66)	93	*	317	8.5 (44)	394	4.3 (87)
4602	1	*	1	*	0	*	1	*	1	*
4701	317	28.4 (64)	326	18.4 (73)	65	*	237	5.9 (14)	328	2.1 (41)
5050	462	25.5 (55)	500	11.4 (48)	75	*	355	11.0 (57)	502	1.2 (21)
5051	796	21.1 (33)	834	10.6 (39)	122	46.7 (5)	582	8.1 (35)	831	2.3 (50)
5052	842	19.2 (30)	916	10.4 (37)	122	47.5 (3)	659	8.3 (38)	916	1.7 (32)
5055	378	2.9 (2)	380	0.8 (1)	10	*	348	5.5 (13)	380	1.1 (19)
5101	476	21.2 (34)	502	8.4 (30)	81	*	382	6.0 (17)	504	2.0 (40)
5102	10	*	12	*	2	*	4	*	12	*
5202	242	40.5 (102)	233	30.0 (110)	82	*	218	8.3 (38)	232	3.0 (68)
5601	300	29.0 (66)	316	20.3 (79)	69	*	241	7.1 (25)	315	3.5 (77)
5801	543	37.4 (90)	555	21.1 (80)	170	38.2 (12)	473	12.1 (62)	554	2.3 (50)
5803	39	*	44	*	7	*	29	*	43	*
5804	7	*	7	*	0	*	3	*	7	*
5806	17	*	18	*	3	*	11	*	18	*
6101	11	*	9	*	3	*	9	*	9	*
6102	41	*	44	*	13	*	37	*	43	*
6103	1,048	30.1 (69)	1,106	20.1 (77)	270	23.0 (24)	921	12.3 (64)	1,110	4.1 (83)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic

Page : 53

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
6105	5	*	4	*	0	*	4	*	4	*
6106	82	*	78	*	28	*	77	*	78	*
6301	1,079	15.0 (11)	1,120	8.9 (32)	135	35.6 (14)	917	5.2 (12)	1,120	1.7 (32)
6312	559	5.4 (4)	577	1.9 (4)	18	*	436	2.1 (4)	576	0.3 (3)
6316	414	15.5 (16)	409	6.1 (17)	51	*	315	3.5 (8)	409	0.7 (10)
6318	14	*	12	*	4	*	13	*	12	*
6320	49	*	48	*	3	*	27	*	48	*
6325	94	*	92	*	14	*	64	*	92	*
6335	368	6.3 (5)	370	3.8 (7)	15	*	302	1.7 (2)	370	1.4 (24)
6340	20	*	14	*	1	*	20	*	14	*
6345	35	*	31	*	7	*	23	*	31	*
6360	267	15.0 (11)	277	7.2 (23)	29	*	220	15.0 (81)	276	2.5 (56)
7304	10	*	10	*	3	*	10	*	9	*
7306	705	36.5 (87)	760	13.0 (54)	215	57.2 (1)	607	10.4 (53)	761	1.1 (19)
7308	10	*	8	*	1	*	10	*	8	*
7310	2	*	2	*	0	*	2	*	2	*
7401	691	38.4 (93)	711	22.4 (86)	216	35.2 (17)	607	18.9 (98)	707	7.2 (115)
7601	199	30.7 (70)	193	17.6 (69)	48	*	183	12.0 (61)	193	2.6 (62)
7850	368	38.0 (91)	370	24.1 (91)	118	28.8 (21)	318	15.1 (84)	369	3.5 (77)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Page : 54

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
7852	6	*	6	*	3	*	6	*		
8101	452	15.0 (11)	523	7.5 (25)	53	*	375	9.6 (51)	525	6.3 (109)
8102	132	22.0 (38)	137	11.7 (50)	25	*	119	5.9 (14)	135	4.4 (90)
8105	486	12.8 (8)	492	4.1 (9)	41	*	370	8.6 (45)	492	0.6 (7)
8106	138	23.2 (44)	164	11.0 (43)	23	*	110	7.3 (28)	164	1.8 (37)
8150	12	*	11	*	0	*	12	*	11	*
8151	951	23.3 (46)	964	11.5 (49)	177	46.9 (4)	806	7.7 (29)	964	1.7 (32)
8152	12	*	10	*	1	*	12	*	10	*
8153	3	*	2	*	0	*	2	*	2	*
8155	4	*	4	*	0	*	4	*	4	*
8157	4	*	4	*	1	*	4	*	4	*
8201	702	17.0 (20)	727	10.9 (41)	76	*	501	7.8 (32)	728	1.5 (28)
8202	1,132	6.8 (6)	1,206	1.9 (4)	68	*	976	6.3 (20)	1,206	2.1 (41)
8203	117	29.1 (67)	122	24.6 (94)	20	*	78	*	122	3.3 (74)
8250	202	23.8 (51)	218	9.2 (34)	39	*	161	8.7 (46)	218	1.4 (24)
8252	51	*	54	*	9	*	44	*	54	*
8255	743	25.7 (56)	804	14.4 (59)	138	41.3 (8)	570	8.2 (36)	803	1.6 (30)
8257	624	1.4 (1)	643	0.9 (2)	6	*	524	2.1 (4)	644	0.3 (3)
8263	208	23.6 (48)	228	10.1 (35)	39	*	156	7.1 (25)	228	0.4 (5)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Page : 55

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8264	234	21.8 (37)	239	5.9 (15)	41	*	171	8.2 (36)	239	0.4 (5)
8267	225	19.1 (29)	220	8.2 (28)	25	*	185	5.9 (14)	220	2.3 (50)
8901	287	24.0 (52)	318	8.5 (31)	61	*	259	19.3 (99)	318	5.0 (99)
8902	432	16.7 (19)	461	3.5 (6)	61	*	365	15.1 (84)	458	4.8 (95)
8903	424	17.5 (22)	488	9.0 (33)	64	*	359	20.9 (102)	487	7.2 (115)
8904	676	17.6 (23)	752	6.6 (21)	94	*	545	17.8 (94)	752	6.6 (113)
8905	450	15.8 (18)	487	6.6 (21)	50	*	366	6.3 (20)	485	4.1 (83)
8909	207	18.4 (26)	221	5.9 (15)	28	*	175	12.6 (67)	219	4.1 (83)
9045	563	11.0 (7)	571	4.2 (10)	48	*	505	7.9 (34)	571	2.5 (56)
9050	263	22.1 (39)	315	6.3 (19)	55	*	229	17.0 (91)	315	2.2 (47)
9051	495	13.1 (9)	521	4.0 (8)	57	*	452	7.7 (29)	520	2.1 (41)
9053	279	17.2 (21)	324	5.6 (12)	43	*	201	20.9 (102)	324	4.6 (93)
9055	2	*	9	*	0	*	0	*	9	*
9058	299	15.7 (17)	332	5.7 (13)	37	*	250	10.8 (55)	332	2.1 (41)
9066	288	20.8 (32)	341	5.0 (11)	43	*	234	22.2 (104)	340	6.2 (107)
9068	247	21.5 (35)	281	5.7 (13)	43	*	219	13.7 (73)	281	5.0 (99)
9069	708	13.6 (10)	858	6.1 (17)	78	*	548	18.4 (96)	858	6.4 (110)
9070	441	18.4 (26)	501	6.4 (20)	64	*	355	19.7 (100)	498	6.4 (110)
9072	445	18.7 (28)	520	7.5 (25)	72	*	381	17.1 (92)	519	6.4 (110)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Page : 56

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
9080	807	22.4 (41)	886	8.0 (27)	151	53.6 (2)	681	13.4 (71)	886	3.7 (80)
9082	42	*	42	*	5	*	34	*	42	*
9083	49	*	65	*	0	*	2	*	65	*
9084	17	*	17	*	1	*	8	*	17	*
9085	4	*	2	*	0	*	4	*	2	*
9086	12	*	10	*	1	*	12	*	10	*
9087	1	*	1	*	0	*	0	*	1	*
Michigan	47,367	24.7	49,881	13.5	9,363	38.4	39,647	10.1	49,826	2.9
National (Prior Year)	155,621	27.7	167,457	14.5	26,902	35.4	120,434	11.5	146,763	7.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 57

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	28	*	*	28	*	24	*	28	*
0102	121	6.6 (18)	5.8 (44)	126	8.7 (12)	110	4.5 (61)	122	60.7 (87)
0103	129	6.2 (13)	7.0 (66)	132	4.5 (1)	122	4.9 (72)	129	75.2 (51)
0104	32	*	*	36	*	29	*	32	*
0105	1	*	*	0	*	0	*	1	*
0250	40	*	*	40	*	35	*	40	*
0251	28	*	*	28	*	26	*	28	*
0252	34	*	*	36	*	33	*	34	*
0253	39	*	*	35	*	32	*	39	*
0401	133	12.8 (94)	12.8 (114)	135	9.6 (19)	117	6.8 (92)	132	75.8 (43)
0402	120	8.3 (44)	6.7 (59)	121	10.7 (36)	104	2.9 (25)	120	75.8 (43)
0403	45	*	*	45	*	39	*	45	*
0404	19	*	*	19	*	18	*	19	*
0501	142	7.7 (33)	7.7 (79)	163	9.2 (15)	127	3.9 (42)	157	73.2 (59)
0503	31	*	*	33	*	27	*	33	*
0504	141	6.4 (15)	6.4 (51)	144	13.9 (87)	121	0.0 (1)	142	87.3 (4)
0505	72	*	*	82	*	63	*	78	*
0506	48	*	*	50	*	41	*	48	*
0507	22	*	*	23	*	19	*	24	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 58

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	10	*	*	10	*	10	*	10	*
0509	1	*	*	1	*	1	*	1	*
0650	307	8.5 (46)	7.8 (81)	322	12.1 (61)	272	3.7 (39)	307	78.8 (33)
0651	191	4.7 (4)	3.1 (9)	193	10.4 (31)	168	1.8 (7)	191	79.6 (30)
0652	101	13.9 (103)	6.9 (62)	107	15.0 (98)	85	*	102	73.5 (58)
0653	101	5.9 (8)	6.9 (62)	107	15.0 (98)	87	*	101	87.1 (5)
0654	103	10.7 (75)	3.9 (21)	106	6.6 (3)	97	*	105	77.1 (40)
0655	102	2.9 (1)	4.9 (29)	99	*	88	*	102	75.5 (48)
0657	24	*	*	25	*	21	*	24	*
0701	97	*	*	100	6.0 (2)	91	*	97	*
0702	22	*	*	22	*	21	*	22	*
0703	59	*	*	56	*	50	*	59	*
0704	18	*	*	19	*	18	*	18	*
0705	33	*	*	35	*	30	*	33	*
0706	6	*	*	6	*	4	*	6	*
0850	222	8.1 (39)	9.0 (97)	238	11.3 (45)	197	5.1 (77)	230	80.4 (25)
0851	197	8.1 (39)	10.7 (105)	201	10.9 (39)	174	4.0 (47)	197	81.7 (19)
0901	452	8.6 (48)	6.6 (58)	465	10.1 (25)	408	4.4 (59)	453	62.5 (82)
0902	16	*	*	16	*	16	*	16	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 59

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1051	353	6.2 (13)	13.6 (115)	364	8.8 (13)	318	1.9 (8)	353	82.4 (17)
1101	578	8.5 (46)	5.4 (36)	593	11.1 (44)	506	4.9 (72)	580	71.2 (62)
1102	238	6.7 (21)	4.2 (22)	248	8.1 (8)	217	2.3 (15)	238	71.4 (61)
1104	1	*	*	1	*	1	*	1	*
1106	24	*	*	25	*	24	*	24	*
1250	299	7.4 (29)	8.7 (92)	312	10.9 (39)	266	1.9 (8)	302	75.5 (48)
1251	215	6.0 (9)	10.7 (105)	225	7.6 (5)	197	2.0 (11)	215	80.9 (24)
1252	176	9.1 (51)	7.4 (76)	176	13.1 (76)	150	4.7 (67)	176	84.7 (11)
1253	223	7.2 (26)	5.8 (44)	233	8.6 (9)	202	1.5 (4)	227	81.5 (20)
1350	649	9.2 (52)	8.0 (83)	656	9.5 (17)	578	4.8 (69)	650	73.8 (57)
1351	46	*	*	47	*	42	*	46	*
1352	108	10.2 (65)	5.6 (40)	110	11.8 (56)	95	*	109	70.6 (63)
1401	82	*	*	84	*	74	*	82	*
1402	33	*	*	33	*	30	*	33	*
1403	9	*	*	9	*	9	*	9	*
1501	236	5.5 (6)	8.9 (96)	246	10.2 (28)	208	1.9 (8)	242	64.5 (78)
1503	294	7.1 (25)	6.8 (60)	303	9.9 (23)	263	3.0 (29)	294	74.1 (55)
1504	605	6.8 (22)	7.1 (70)	621	10.1 (25)	541	2.8 (23)	614	76.9 (41)
1505	130	9.2 (52)	12.3 (112)	139	10.8 (37)	116	4.3 (56)	133	86.5 (7)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 60

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1506	178	7.9 (35)	11.2 (108)	179	8.9 (14)	160	5.0 (76)	178	75.8 (43)
1507	167	6.6 (18)	10.2 (103)	173	11.0 (42)	148	1.4 (3)	168	82.7 (14)
1508	13	*	*	13	*	12	*	13	*
1509	111	3.6 (2)	16.2 (116)	115	10.4 (31)	98	*	115	75.7 (47)
1511	198	9.6 (60)	8.6 (91)	200	13.0 (74)	170	2.4 (17)	198	82.3 (18)
1512	110	7.3 (27)	10.0 (102)	111	15.3 (103)	94	*	111	81.1 (22)
1514	91	*	*	95	*	83	*	94	*
1701	179	9.5 (58)	11.7 (111)	182	11.5 (49)	154	2.6 (19)	179	61.5 (85)
1950	175	7.4 (29)	5.1 (31)	180	13.3 (80)	153	2.6 (19)	175	85.7 (9)
1952	32	*	*	33	*	26	*	32	*
1953	101	7.9 (35)	6.9 (62)	105	10.5 (33)	90	*	101	87.1 (5)
1954	155	5.8 (7)	11.0 (107)	158	9.5 (17)	136	2.9 (25)	156	87.8 (2)
1955	30	*	*	30	*	27	*	30	*
1957	134	8.2 (41)	12.7 (113)	134	9.7 (20)	118	3.4 (36)	134	85.8 (8)
1963	21	*	*	22	*	20	*	21	*
1964	13	*	*	13	*	13	*	13	*
2150	191	9.4 (55)	7.3 (73)	192	10.9 (39)	169	4.1 (51)	191	80.1 (28)
2151	69	*	*	70	*	61	*	69	*
2251	114	7.9 (35)	10.5 (104)	116	8.6 (9)	104	4.8 (69)	114	78.1 (35)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 61

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2252	47	*	*	51	*	40	*	47	*
2554	15	*	*	16	*	15	*	15	*
2555	2,329	11.4 (82)	6.2 (47)	2,367	13.6 (84)	2,001	5.6 (86)	2,329	55.2 (100)
2556	127	12.6 (92)	3.1 (9)	130	20.0 (115)	100	8.0 (101)	130	53.8 (104)
2557	1	*	*	1	*	1	*	1	*
2558	2	*	*	3	*	1	*	3	*
2560	1	*	*	1	*	1	*	1	*
2561	2	*	*	2	*	1	*	2	*
2562	10	*	*	10	*	8	*	10	*
2563	15	*	*	14	*	13	*	15	*
2564	7	*	*	7	*	7	*	7	*
3001	333	12.9 (95)	3.0 (7)	343	15.5 (104)	283	3.9 (42)	333	61.0 (86)
3002	105	10.5 (71)	1.9 (1)	105	10.5 (33)	94	*	105	56.2 (99)
3004	73	*	*	73	*	67	*	73	*
3005	18	*	*	18	*	16	*	18	*
3007	5	*	*	4	*	2	*	5	*
3201	127	11.8 (86)	7.1 (70)	132	12.1 (61)	112	3.6 (38)	127	78.7 (34)
3301	1,273	9.7 (61)	6.4 (51)	1,329	11.7 (54)	1,123	4.2 (55)	1,287	77.9 (36)
3304	38	*	*	39	*	35	*	39	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 62

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3305	50	*	*	54	*	46	*	51	*
3306	69	*	*	74	*	66	*	69	*
3308	18	*	*	18	*	16	*	18	*
3309	7	*	*	7	*	7	*	7	*
3312	147	12.9 (95)	8.8 (94)	150	17.3 (109)	122	7.4 (97)	147	69.4 (64)
3550	9	*	*	9	*	8	*	9	*
3551	6	*	*	7	*	6	*	6	*
3552	18	*	*	18	*	15	*	18	*
3701	312	8.7 (49)	9.0 (97)	316	11.7 (54)	274	2.9 (25)	312	82.7 (14)
3850	871	10.2 (65)	7.3 (73)	885	11.5 (49)	768	3.9 (42)	870	71.7 (60)
3901	468	14.3 (105)	5.3 (34)	489	14.3 (91)	404	5.9 (87)	476	66.4 (75)
3902	67	*	*	70	*	61	*	67	*
3903	16	*	*	17	*	15	*	16	*
3904	4	*	*	4	*	3	*	4	*
3905	9	*	*	9	*	7	*	9	*
3950	703	9.7 (61)	8.5 (88)	721	12.2 (65)	613	2.8 (23)	709	74.9 (52)
3954	7	*	*	8	*	7	*	7	*
4103	1	*	*	1	*	0	*	1	*
4106	751	12.6 (92)	4.3 (23)	775	15.2 (102)	634	4.9 (72)	755	77.5 (38)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 63

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4107	878	13.2 (100)	5.8 (44)	903	16.7 (108)	728	5.4 (83)	879	80.3 (27)
4109	276	9.4 (55)	7.6 (78)	284	13.4 (82)	238	2.9 (25)	277	75.8 (43)
4110	5	*	*	5	*	0	*	5	*
4112	156	6.4 (15)	6.4 (51)	155	9.7 (20)	137	2.2 (12)	155	27.7 (116)
4115	1,108	12.5 (90)	6.4 (51)	1,132	14.8 (95)	936	4.3 (56)	1,112	77.2 (39)
4351	153	7.8 (34)	6.5 (56)	158	13.9 (87)	132	3.0 (29)	154	82.5 (16)
4352	47	*	*	48	*	37	*	47	*
4353	109	6.4 (15)	6.4 (51)	111	7.2 (4)	101	4.0 (47)	109	74.3 (54)
4354	228	6.1 (12)	8.8 (94)	228	11.4 (46)	200	1.5 (4)	229	68.6 (67)
4355	100	13.0 (97)	3.0 (7)	104	17.3 (109)	81	*	100	85.0 (10)
4356	126	10.3 (67)	9.5 (101)	131	11.5 (49)	110	4.5 (61)	126	74.6 (53)
4358	33	*	*	32	*	29	*	33	*
4359	156	4.5 (3)	9.0 (97)	158	10.8 (37)	138	2.2 (12)	156	79.5 (31)
4360	50	*	*	54	*	46	*	50	*
4361	74	*	*	82	*	69	*	74	*
4362	88	*	*	90	*	76	*	89	*
4363	233	6.0 (9)	7.7 (79)	235	9.8 (22)	202	3.0 (29)	233	77.7 (37)
4401	306	6.9 (23)	8.2 (85)	309	10.0 (24)	271	3.0 (29)	306	80.4 (25)
4501	271	8.9 (50)	8.5 (88)	272	13.6 (84)	229	2.2 (12)	270	81.5 (20)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 64

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4601	369	7.0 (24)	6.5 (56)	389	12.1 (61)	321	1.6 (6)	368	73.9 (56)
4602	1	*	*	1	*	1	*	1	*
4701	322	10.6 (73)	6.8 (60)	322	12.7 (71)	279	3.9 (42)	322	84.5 (12)
5050	491	12.4 (89)	5.5 (39)	489	14.3 (91)	414	5.3 (80)	491	64.4 (79)
5051	806	10.3 (67)	5.6 (40)	808	12.6 (69)	689	4.5 (61)	809	66.1 (76)
5052	889	10.6 (73)	5.7 (42)	901	14.2 (89)	757	4.4 (59)	891	62.1 (83)
5055	332	6.6 (18)	5.4 (36)	358	7.8 (6)	304	2.3 (15)	355	46.8 (111)
5101	486	8.2 (41)	7.0 (66)	490	12.2 (65)	419	3.1 (33)	489	67.7 (69)
5102	12	*	*	11	*	10	*	11	*
5202	225	9.3 (54)	11.6 (110)	231	12.1 (61)	198	4.5 (61)	226	87.6 (3)
5601	303	7.3 (27)	11.2 (108)	313	9.3 (16)	274	2.6 (19)	303	82.8 (13)
5801	528	10.8 (76)	6.3 (50)	543	11.6 (53)	462	5.2 (78)	529	66.7 (73)
5803	40	*	*	39	*	34	*	40	*
5804	6	*	*	6	*	5	*	6	*
5806	18	*	*	18	*	16	*	18	*
6101	9	*	*	9	*	7	*	9	*
6102	44	*	*	44	*	41	*	44	*
6103	1,053	10.4 (70)	5.0 (30)	1,073	14.2 (89)	884	4.1 (51)	1,055	68.4 (68)
6105	4	*	*	4	*	4	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 65

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6106	78	*	*	78	*	66	*	78	*
6301	1,082	12.5 (90)	7.4 (76)	1,107	12.8 (72)	939	4.7 (67)	1,088	66.9 (72)
6312	556	12.2 (88)	7.0 (66)	567	11.8 (56)	488	4.1 (51)	560	79.8 (29)
6316	391	11.3 (80)	7.9 (82)	400	11.0 (42)	347	3.5 (37)	394	79.4 (32)
6318	10	*	*	12	*	9	*	11	*
6320	44	*	*	46	*	38	*	45	*
6325	89	*	*	90	*	81	*	89	*
6335	353	7.9 (35)	7.1 (70)	357	10.1 (25)	314	4.8 (69)	356	68.8 (66)
6340	13	*	*	14	*	10	*	13	*
6345	31	*	*	32	*	29	*	31	*
6360	265	12.1 (87)	4.5 (25)	273	15.0 (98)	227	6.2 (89)	267	67.0 (70)
7304	10	*	*	10	*	10	*	10	*
7306	736	10.9 (77)	5.3 (34)	747	13.5 (83)	635	3.9 (42)	736	64.4 (79)
7308	8	*	*	8	*	6	*	8	*
7310	2	*	*	2	*	2	*	2	*
7401	682	6.0 (9)	9.1 (100)	697	10.2 (28)	608	3.3 (35)	685	69.1 (65)
7601	186	7.5 (31)	4.8 (27)	191	13.6 (84)	160	0.0 (1)	186	56.5 (97)
7850	354	8.2 (41)	6.2 (47)	358	12.3 (67)	310	3.2 (34)	354	75.4 (50)
7852	6	*	*	6	*	5	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 66

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8101	486	19.3 (116)	2.1 (2)	510	21.0 (116)	374	8.0 (101)	504	43.8 (113)
8102	129	10.9 (77)	8.5 (88)	134	14.9 (96)	106	6.6 (91)	132	53.8 (104)
8105	480	11.3 (80)	4.8 (27)	484	13.0 (74)	412	4.1 (51)	480	58.5 (91)
8106	153	16.3 (114)	3.3 (15)	157	19.1 (114)	120	9.2 (105)	154	60.4 (88)
8150	10	*	*	10	*	10	*	10	*
8151	916	11.2 (79)	5.2 (33)	928	12.6 (69)	796	5.3 (80)	915	81.0 (23)
8152	10	*	*	10	*	10	*	10	*
8153	2	*	*	2	*	1	*	2	*
8155	4	*	*	4	*	3	*	4	*
8157	4	*	*	4	*	4	*	4	*
8201	696	11.4 (82)	6.2 (47)	712	11.4 (46)	611	4.9 (72)	704	66.6 (74)
8202	1,123	8.3 (44)	5.7 (42)	1,150	8.6 (9)	1,016	4.0 (47)	1,148	43.3 (114)
8203	116	9.5 (58)	6.9 (62)	118	10.2 (28)	104	3.8 (41)	116	87.9 (1)
8250	214	13.1 (99)	4.7 (26)	218	10.6 (35)	191	7.9 (99)	215	67.0 (70)
8252	53	*	*	54	*	46	*	53	*
8255	773	10.3 (67)	7.0 (66)	788	11.9 (59)	673	4.6 (65)	775	58.3 (93)
8257	581	13.6 (102)	3.1 (9)	625	13.1 (76)	494	7.3 (95)	598	76.6 (42)
8263	220	10.0 (63)	8.2 (85)	224	12.9 (73)	191	2.6 (19)	220	54.5 (102)
8264	226	4.9 (5)	8.4 (87)	229	7.9 (7)	202	2.5 (18)	227	55.1 (101)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 67

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8267	213	9.4 (55)	8.0 (83)	220	11.8 (56)	189	3.7 (39)	214	52.3 (107)
8901	305	15.4 (110)	3.3 (15)	312	15.7 (106)	254	7.5 (98)	306	44.8 (112)
8902	399	14.0 (104)	2.8 (4)	451	17.3 (109)	335	5.4 (83)	429	60.1 (89)
8903	451	17.5 (115)	2.9 (5)	466	18.0 (112)	361	8.9 (104)	460	50.0 (109)
8904	674	15.7 (111)	3.3 (15)	721	18.9 (113)	534	7.3 (95)	709	54.2 (103)
8905	427	10.1 (64)	7.3 (73)	464	11.4 (46)	369	4.3 (56)	459	64.7 (77)
8909	191	16.2 (112)	3.1 (9)	215	14.9 (96)	159	6.9 (93)	202	62.9 (81)
9045	533	10.5 (71)	5.4 (36)	556	13.1 (76)	461	4.6 (65)	536	53.2 (106)
9050	292	13.0 (97)	3.8 (19)	301	13.3 (80)	252	7.9 (99)	296	57.8 (94)
9051	493	7.5 (31)	8.7 (92)	495	12.5 (68)	424	4.0 (47)	497	62.0 (84)
9053	311	15.1 (108)	3.2 (13)	318	14.5 (94)	263	7.2 (94)	312	59.9 (90)
9055	9	*	*	9	*	9	*	9	*
9058	317	15.1 (108)	3.2 (13)	331	11.5 (49)	280	9.3 (106)	324	57.7 (95)
9066	315	16.2 (112)	2.5 (3)	338	15.1 (101)	268	8.2 (103)	316	58.5 (91)
9068	272	13.2 (100)	5.1 (31)	273	16.1 (107)	224	5.4 (83)	272	47.1 (110)
9069	802	14.5 (106)	3.4 (18)	834	14.4 (93)	682	6.0 (88)	803	35.7 (115)
9070	486	11.5 (84)	2.9 (5)	496	11.9 (59)	424	5.2 (78)	487	57.7 (95)
9072	494	15.0 (107)	4.3 (23)	505	15.6 (105)	411	6.3 (90)	494	52.0 (108)
9080	859	11.5 (84)	3.8 (19)	867	13.1 (76)	738	5.3 (80)	863	56.3 (98)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Page : 68

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9082	40	*	*	40	*	33	*	39	*
9083	62	*	*	65	*	50	*	62	*
9084	14	*	*	17	*	12	*	14	*
9085	2	*	*	2	*	2	*	2	*
9086	10	*	*	10	*	10	*	10	*
9087	1	*	*	1	*	1	*	1	*
Michigan	47,135	10.4	6.4	48,449	12.5	40,866	4.5	47,531	67.8
National (Prior Year)	121,176	9.3	6.7	121,542	11.8	103,169	3.9	119,843	56.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 69

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	116	92.2	0.9	3.4	0.0	0.0	3.4	0.0
0102	505	89.1	0.8	3.0	0.4	0.2	6.3	0.2
0103	442	90.5	0.5	4.5	0.5	0.0	3.6	0.5
0104	154	92.2	0.0	3.9	0.0	0.6	2.6	0.6
0105	9	*	*	*	*	*	*	*
0250	158	85.4	0.0	2.5	0.6	0.0	11.4	0.0
0251	100	86.0	0.0	1.0	8.0	0.0	5.0	0.0
0252	116	54.3	0.9	0.9	0.9	0.0	43.1	0.0
0253	154	79.9	0.6	0.0	3.9	0.0	15.6	0.0
0401	497	93.6	0.6	2.2	0.2	0.4	2.6	0.4
0402	409	86.3	0.7	2.7	0.0	0.5	9.8	0.0
0403	153	96.1	0.0	2.6	0.0	0.0	0.7	0.7
0404	122	92.6	0.8	1.6	0.0	0.0	4.9	0.0
0501	525	95.0	0.4	2.1	0.0	0.8	1.7	0.0
0503	114	89.5	0.0	5.3	0.0	0.9	4.4	0.0
0504	498	87.8	1.0	2.2	1.8	0.4	6.8	0.0
0505	323	83.6	0.9	4.3	0.6	0.3	10.2	0.0
0506	193	94.3	0.5	0.5	0.5	0.0	4.1	0.0
0507	81	*	*	*	*	*	*	*
0508	35	*	*	*	*	*	*	*
0509	6	*	*	*	*	*	*	*
0650	1,144	80.2	3.6	5.6	3.0	1.9	5.6	0.1
0651	624	92.9	0.3	3.4	0.3	0.5	2.1	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 70

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0652	325	95.1	0.0	2.2	0.0	0.3	2.5	0.0
0653	409	95.6	0.2	1.7	0.0	0.0	2.2	0.2
0654	385	92.2	1.3	1.8	0.5	0.0	3.9	0.3
0655	365	91.2	0.3	2.2	0.0	0.0	5.2	1.1
0657	69	*	*	*	*	*	*	*
0701	408	87.3	1.7	2.2	0.2	4.9	2.7	1.0
0702	67	*	*	*	*	*	*	*
0703	225	90.7	1.3	1.8	0.0	0.4	3.6	2.2
0704	54	*	*	*	*	*	*	*
0705	136	94.1	0.7	0.0	0.0	0.0	4.4	0.7
0706	12	*	*	*	*	*	*	*
0850	873	84.8	5.3	5.7	0.1	1.1	2.5	0.5
0851	745	93.7	0.4	4.4	0.0	0.4	0.8	0.3
0901	1,660	79.7	2.2	10.8	0.0	0.4	6.1	0.8
0902	43	*	*	*	*	*	*	*
1051	1,342	84.2	1.1	6.6	1.3	1.1	5.0	0.7
1101	1,921	36.1	50.5	7.9	0.3	0.8	3.4	1.0
1102	844	73.7	10.7	7.0	0.8	0.8	6.2	0.8
1104	1	*	*	*	*	*	*	*
1106	79	*	*	*	*	*	*	*
1250	1,061	87.2	0.5	9.5	0.0	0.3	2.0	0.6
1251	816	93.3	0.4	2.3	0.0	0.4	2.7	1.0
1252	604	61.6	0.3	33.6	0.0	0.3	3.5	0.7
1253	830	78.6	3.9	9.9	0.0	0.1	6.3	1.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 71

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1350	2,435	59.1	20.1	7.0	0.2	6.3	6.7	0.5
1351	135	86.7	3.7	5.9	0.0	0.0	3.0	0.7
1352	376	66.5	23.9	5.6	0.0	0.0	3.5	0.5
1401	247	90.7	0.8	3.2	1.2	0.4	2.0	1.6
1402	132	59.1	0.8	22.7	7.6	1.5	7.6	0.8
1403	25	*	*	*	*	*	*	*
1501	815	51.4	3.6	36.2	0.0	0.2	4.9	3.7
1503	1,097	61.4	7.1	22.9	0.3	0.5	6.8	0.9
1504	2,183	40.6	5.5	44.7	0.1	3.9	4.2	1.1
1505	433	84.3	3.9	6.7	0.0	0.7	3.0	1.4
1506	602	77.1	0.8	17.3	0.3	0.5	2.8	1.2
1507	625	85.1	1.0	7.0	0.2	1.4	5.0	0.3
1508	86	*	*	*	*	*	*	*
1509	450	50.9	2.4	39.3	0.2	0.0	3.1	4.0
1511	741	88.7	2.0	4.3	0.0	1.1	3.2	0.7
1512	367	87.7	0.3	5.2	0.8	0.5	5.2	0.3
1513	54	*	*	*	*	*	*	*
1514	235	84.7	1.3	8.9	0.0	0.9	3.0	1.3
1701	706	54.7	0.4	2.7	20.8	0.4	20.3	0.6
1950	604	92.9	0.2	3.8	0.3	0.2	2.2	0.5
1952	124	94.4	0.0	3.2	0.0	0.0	2.4	0.0
1953	331	90.9	0.3	6.3	0.0	0.0	2.1	0.3
1954	574	86.2	0.2	12.4	0.0	0.2	1.0	0.0
1955	100	86.0	1.0	11.0	0.0	0.0	1.0	1.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 72

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1957	486	82.1	0.4	14.4	0.0	0.0	2.9	0.2
1963	35	*	*	*	*	*	*	*
1964	48	*	*	*	*	*	*	*
2150	681	86.2	0.1	2.2	0.3	0.7	10.3	0.1
2151	247	76.9	0.8	9.7	0.4	0.0	8.5	3.6
2251	421	91.4	0.5	2.6	0.5	0.0	4.0	1.0
2252	203	90.6	0.5	1.0	0.0	0.0	4.9	3.0
2554	70	*	*	*	*	*	*	*
2555	8,009	47.8	39.4	4.9	0.1	0.3	7.0	0.5
2556	334	30.8	56.0	6.0	0.0	0.3	6.6	0.3
2557	10	*	*	*	*	*	*	*
2558	7	*	*	*	*	*	*	*
2559	9	*	*	*	*	*	*	*
2560	4	*	*	*	*	*	*	*
2561	2	*	*	*	*	*	*	*
2562	13	*	*	*	*	*	*	*
2563	20	*	*	*	*	*	*	*
2564	7	*	*	*	*	*	*	*
3001	1,145	16.0	67.0	12.6	0.1	0.9	3.2	0.3
3002	296	21.3	50.3	23.0	0.0	0.0	5.4	0.0
3004	193	38.3	0.0	59.1	0.5	0.0	2.1	0.0
3005	19	*	*	*	*	*	*	*
3007	26	*	*	*	*	*	*	*
3009	6	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 73

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3201	460	94.1	0.0	3.5	0.2	0.2	1.1	0.9
3301	4,371	38.5	30.8	15.4	0.2	5.9	8.8	0.3
3304	107	41.1	15.0	11.2	0.0	29.0	3.7	0.0
3305	162	67.3	10.5	11.7	0.0	5.6	4.3	0.6
3306	257	82.9	3.5	7.0	0.4	0.8	4.3	1.2
3308	74	*	*	*	*	*	*	*
3309	37	*	*	*	*	*	*	*
3312	429	36.8	28.0	24.9	0.5	1.2	8.2	0.5
3550	65	*	*	*	*	*	*	*
3551	29	*	*	*	*	*	*	*
3552	52	*	*	*	*	*	*	*
3701	1,092	89.3	0.1	7.3	0.0	0.2	3.0	0.1
3850	3,160	71.8	14.7	5.5	0.1	0.4	7.2	0.3
3901	1,142	33.2	52.9	6.1	0.3	0.8	5.9	0.9
3902	350	49.4	15.1	17.4	0.0	0.9	17.1	0.0
3903	28	*	*	*	*	*	*	*
3904	7	*	*	*	*	*	*	*
3905	11	*	*	*	*	*	*	*
3950	2,114	57.8	21.9	9.7	0.1	1.4	8.5	0.5
3954	12	*	*	*	*	*	*	*
4001	512	30.5	55.3	8.0	0.0	1.4	4.7	0.2
4002	327	48.9	25.1	18.0	0.3	0.6	7.0	0.0
4003	2	*	*	*	*	*	*	*
4004	4	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 74

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4103	42	*	*	*	*	*	*	*
4106	2,566	14.8	42.3	36.6	0.2	0.7	4.5	0.8
4107	3,237	45.1	26.7	18.4	0.4	1.9	7.3	0.3
4109	1,039	84.6	1.4	8.7	0.1	1.3	3.9	0.0
4110	5	*	*	*	*	*	*	*
4112	568	1.8	1.1	95.4	0.0	0.7	0.0	1.1
4113	343	4.7	18.4	74.3	0.0	0.6	2.0	0.0
4115	3,610	35.4	27.5	19.5	0.4	11.4	5.3	0.5
4351	582	88.5	0.2	6.2	0.0	0.7	4.3	0.2
4352	167	69.5	13.8	6.6	0.0	0.0	9.6	0.6
4353	421	67.7	0.7	29.2	0.2	0.0	1.7	0.5
4354	774	87.7	3.9	3.1	0.3	0.3	4.4	0.4
4355	357	79.0	1.4	8.4	2.2	0.6	8.1	0.3
4356	467	84.2	1.3	10.7	0.0	0.2	3.6	0.0
4358	141	39.0	0.0	56.7	0.0	0.0	2.8	1.4
4359	485	69.7	0.2	27.0	0.2	0.6	1.2	1.0
4360	230	94.3	0.4	4.8	0.0	0.0	0.4	0.0
4361	268	95.5	0.0	1.5	0.4	0.0	1.9	0.7
4362	304	92.1	0.3	2.3	0.3	1.0	3.6	0.3
4363	907	92.3	0.1	3.2	0.0	0.2	3.6	0.6
4401	1,077	89.2	0.4	5.1	0.0	0.5	4.4	0.5
4501	990	91.9	0.5	4.5	0.0	0.2	2.7	0.1
4601	1,396	75.1	1.8	16.3	0.0	0.1	5.9	0.7
4602	23	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 75

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4701	1,116	91.3	0.4	3.6	0.2	1.3	2.1	1.2
5050	1,740	51.6	38.3	2.7	0.0	1.6	5.6	0.2
5051	2,729	52.5	34.5	4.3	0.1	2.9	5.2	0.4
5052	2,959	40.0	36.9	2.0	0.1	15.1	5.1	0.5
5055	1,193	93.7	1.3	0.2	0.1	3.8	0.3	0.6
5101	1,833	69.2	11.2	13.2	0.1	1.6	4.8	0.0
5102	14	*	*	*	*	*	*	*
5202	795	86.9	0.5	3.3	0.9	0.4	7.5	0.5
5203	4	*	*	*	*	*	*	*
5204	4	*	*	*	*	*	*	*
5601	1,078	93.4	0.8	1.8	0.1	0.9	2.8	0.2
5801	1,772	81.7	6.2	6.2	0.1	0.5	4.7	0.7
5803	137	88.3	0.0	10.2	0.0	0.0	1.5	0.0
5804	17	*	*	*	*	*	*	*
5806	50	*	*	*	*	*	*	*
6101	35	*	*	*	*	*	*	*
6102	122	86.1	0.8	3.3	0.0	0.0	9.0	0.8
6103	4,048	53.9	29.0	8.9	0.3	0.5	7.0	0.3
6105	22	*	*	*	*	*	*	*
6106	103	35.9	51.5	7.8	0.0	0.0	4.9	0.0
6301	4,171	41.5	30.6	20.4	0.1	2.2	4.7	0.4
6312	2,184	33.6	56.5	3.4	0.0	1.6	4.4	0.5
6316	1,283	66.9	19.6	7.0	0.2	3.3	2.8	0.2
6318	14	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 76

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6320	129	62.0	24.8	6.2	0.0	0.8	5.4	0.8
6325	192	89.6	3.6	4.2	0.0	0.0	2.6	0.0
6330	118	24.6	55.1	17.8	0.0	0.8	1.7	0.0
6335	1,059	75.9	14.3	2.5	0.1	5.9	1.2	0.1
6340	50	*	*	*	*	*	*	*
6345	60	*	*	*	*	*	*	*
6360	641	22.8	53.2	16.7	0.0	0.8	6.6	0.0
7304	92	*	*	*	*	*	*	*
7306	2,555	44.7	32.8	16.3	0.0	0.6	5.0	0.6
7308	85	*	*	*	*	*	*	*
7310	22	*	*	*	*	*	*	*
7401	2,336	82.4	5.6	5.9	0.1	0.4	5.1	0.5
7601	650	93.4	0.2	4.8	0.0	0.0	1.5	0.2
7850	1,172	88.8	0.3	5.8	0.1	0.1	4.5	0.3
7852	21	*	*	*	*	*	*	*
8101	1,913	5.3	91.4	0.5	0.1	0.2	2.4	0.2
8102	426	13.1	10.1	72.8	0.0	0.2	3.3	0.5
8105	1,657	31.7	61.3	3.0	0.1	0.7	2.8	0.2
8106	486	23.9	62.3	8.0	0.4	0.0	5.1	0.2
8150	33	*	*	*	*	*	*	*
8151	3,273	36.6	39.6	10.1	0.2	2.9	10.0	0.6
8152	54	*	*	*	*	*	*	*
8153	7	*	*	*	*	*	*	*
8155	20	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 77

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8157	9	*	*	*	*	*	*	*
8201	2,628	46.7	38.4	4.8	0.3	3.4	6.0	0.5
8202	4,023	85.7	10.4	1.8	0.0	0.4	1.1	0.6
8203	445	58.0	29.7	4.0	0.2	1.8	5.2	1.1
8250	814	20.1	68.6	4.1	0.1	1.1	5.7	0.4
8252	160	2.5	92.5	0.0	0.0	1.9	3.1	0.0
8255	2,629	53.6	31.5	7.8	0.2	2.7	3.8	0.5
8257	2,115	49.4	14.8	0.9	0.1	32.7	1.7	0.4
8263	727	68.6	14.4	10.3	0.3	2.2	3.9	0.3
8264	701	64.6	25.7	3.9	0.1	1.6	3.7	0.4
8266	9	*	*	*	*	*	*	*
8267	887	56.3	16.9	21.0	0.1	0.1	4.8	0.6
8901	1,143	1.6	94.0	0.7	0.1	1.2	2.2	0.3
8902	1,503	3.7	92.0	1.7	0.0	0.1	2.4	0.1
8903	1,445	2.3	92.7	1.0	0.0	0.9	2.8	0.3
8904	2,472	2.3	93.0	1.1	0.2	0.7	2.5	0.1
8905	1,609	59.0	29.4	8.5	0.0	0.7	2.2	0.2
8908	18	*	*	*	*	*	*	*
8909	641	2.2	94.4	0.8	0.0	0.5	1.9	0.3
9016	7	*	*	*	*	*	*	*
9018	3	*	*	*	*	*	*	*
9045	1,885	10.3	26.5	58.7	0.1	0.3	3.7	0.4
9050	968	12.4	70.9	14.3	0.0	0.2	2.3	0.0
9051	1,753	9.2	7.9	80.8	0.0	0.3	1.6	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 78

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9053	1,011	3.5	90.1	1.1	0.1	3.2	1.8	0.3
9055	29	*	*	*	*	*	*	*
9058	1,062	20.3	67.8	1.4	0.0	8.9	1.5	0.1
9066	1,140	1.7	96.8	0.5	0.0	0.1	0.9	0.1
9068	962	2.2	96.5	0.5	0.0	0.1	0.5	0.2
9069	2,883	0.8	97.5	0.3	0.0	0.2	0.9	0.1
9070	1,401	3.0	92.1	0.6	0.0	2.4	1.6	0.1
9072	1,809	2.3	95.2	0.4	0.0	0.4	1.4	0.2
9080	2,885	20.0	76.2	1.8	0.2	0.1	1.6	0.2
9081	16	*	*	*	*	*	*	*
9082	165	2.4	95.8	1.2	0.0	0.0	0.6	0.0
9083	123	30.1	59.3	7.3	0.0	1.6	1.6	0.0
9084	43	*	*	*	*	*	*	*
9085	53	*	*	*	*	*	*	*
9086	48	*	*	*	*	*	*	*
9087	1	*	*	*	*	*	*	*
Michigan	167,191	51.6	30.3	10.5	0.4	2.2	4.6	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 79

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	116	0.0	1.7	7.8	59.5	31.0	0.0	0.0
0102	505	0.0	4.0	8.9	65.3	21.0	0.8	0.0
0103	442	0.0	2.5	8.8	66.3	21.0	1.4	0.0
0104	154	0.0	4.5	9.1	71.4	14.9	0.0	0.0
0105	9	*	*	*	*	*	*	*
0250	158	0.0	4.4	5.7	71.5	18.4	0.0	0.0
0251	100	0.0	2.0	7.0	60.0	29.0	2.0	0.0
0252	116	0.0	3.4	5.2	62.9	26.7	1.7	0.0
0253	154	0.0	4.5	7.8	65.6	22.1	0.0	0.0
0401	497	0.0	3.6	9.5	64.8	20.5	1.6	0.0
0402	409	0.2	1.7	7.6	66.0	23.5	1.0	0.0
0403	153	0.0	2.6	4.6	73.2	19.0	0.7	0.0
0404	122	0.0	2.5	8.2	63.1	23.0	3.3	0.0
0501	525	0.0	2.3	9.0	63.8	24.6	0.4	0.0
0503	114	0.0	0.9	5.3	56.1	35.1	2.6	0.0
0504	498	0.0	0.8	8.8	64.3	24.7	1.4	0.0
0505	323	0.0	1.5	8.7	63.8	25.1	0.9	0.0
0506	193	0.0	2.6	10.9	64.2	21.2	1.0	0.0
0507	81	*	*	*	*	*	*	*
0508	35	*	*	*	*	*	*	*
0509	6	*	*	*	*	*	*	*
0650	1,144	0.1	2.7	8.0	65.5	22.6	0.9	0.2
0651	624	0.3	2.7	9.5	67.8	17.9	1.6	0.2
0652	325	0.0	2.2	6.5	68.3	20.6	2.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 80

Michigan

Comparison of Age Distribution by Clinic Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0653	409	0.0	2.7	8.1	68.0	20.8	0.5	0.0
0654	385	0.3	5.2	10.9	60.8	22.1	0.8	0.0
0655	365	0.0	3.0	11.2	63.0	22.2	0.3	0.3
0657	69	*	*	*	*	*	*	*
0701	408	0.0	2.5	6.6	57.6	29.7	3.4	0.2
0702	67	*	*	*	*	*	*	*
0703	225	0.0	1.3	9.3	63.1	25.3	0.9	0.0
0704	54	*	*	*	*	*	*	*
0705	136	0.0	2.9	9.6	56.6	27.2	3.7	0.0
0706	12	*	*	*	*	*	*	*
0850	873	0.1	2.2	9.4	60.6	26.0	1.3	0.5
0851	745	0.0	3.1	9.3	64.6	21.5	1.3	0.3
0901	1,660	0.1	2.6	8.8	64.5	22.3	1.7	0.0
0902	43	*	*	*	*	*	*	*
1051	1,342	0.0	2.6	6.5	62.1	26.2	2.3	0.3
1101	1,921	0.1	3.9	9.3	62.2	21.8	1.4	1.4
1102	844	0.1	3.3	10.8	62.3	21.8	1.5	0.1
1104	1	*	*	*	*	*	*	*
1106	79	*	*	*	*	*	*	*
1250	1,061	0.0	2.0	9.3	61.6	23.9	3.0	0.1
1251	816	0.1	2.7	10.2	67.4	18.5	1.0	0.1
1252	604	0.0	3.0	12.6	61.6	20.7	2.2	0.0
1253	830	0.0	3.3	8.7	66.1	20.7	1.2	0.0
1350	2,435	0.2	3.1	9.1	63.5	22.8	1.3	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 81

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1351	135	0.0	3.0	6.7	58.5	31.9	0.0	0.0
1352	376	0.3	4.0	7.7	67.8	18.6	1.3	0.3
1401	247	0.0	3.6	7.7	60.7	25.9	1.6	0.4
1402	132	0.0	0.8	6.8	59.8	29.5	1.5	1.5
1403	25	*	*	*	*	*	*	*
1501	815	0.0	2.9	11.8	60.5	22.5	2.0	0.4
1503	1,097	0.3	4.1	10.1	60.8	23.1	1.6	0.0
1504	2,183	0.2	3.6	8.0	60.7	25.8	1.8	0.0
1505	433	0.0	1.6	6.7	62.4	28.2	0.9	0.2
1506	602	0.0	3.5	8.3	60.8	26.1	1.3	0.0
1507	625	0.0	1.1	5.3	66.6	25.4	1.6	0.0
1508	86	*	*	*	*	*	*	*
1509	450	0.0	4.9	7.3	61.3	23.6	2.9	0.0
1511	741	0.3	3.2	11.1	62.2	21.7	1.5	0.0
1512	367	0.0	2.2	6.8	62.1	28.1	0.8	0.0
1513	54	*	*	*	*	*	*	*
1514	235	0.0	2.6	5.1	51.5	39.1	1.7	0.0
1701	706	0.1	1.4	7.4	62.3	27.2	1.6	0.0
1950	604	0.0	3.5	11.1	68.4	15.9	1.2	0.0
1952	124	0.8	1.6	8.9	63.7	22.6	2.4	0.0
1953	331	0.6	2.7	7.9	70.1	17.5	1.2	0.0
1954	574	0.2	2.6	8.2	63.8	23.5	1.7	0.0
1955	100	0.0	2.0	6.0	68.0	23.0	1.0	0.0
1957	486	0.0	3.1	8.4	63.6	23.0	1.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 82

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1963	35	*	*	*	*	*	*	*
1964	48	*	*	*	*	*	*	*
2150	681	0.0	2.1	9.3	63.6	24.2	0.9	0.0
2151	247	0.0	5.7	10.1	62.3	20.2	1.6	0.0
2251	421	0.2	1.9	9.3	60.1	26.6	1.7	0.2
2252	203	0.0	2.5	9.4	59.1	27.6	1.5	0.0
2554	70	*	*	*	*	*	*	*
2555	8,009	0.2	2.7	8.4	64.9	22.4	1.4	0.1
2556	334	0.6	2.4	8.4	68.0	19.8	0.9	0.0
2557	10	*	*	*	*	*	*	*
2558	7	*	*	*	*	*	*	*
2559	9	*	*	*	*	*	*	*
2560	4	*	*	*	*	*	*	*
2561	2	*	*	*	*	*	*	*
2562	13	*	*	*	*	*	*	*
2563	20	*	*	*	*	*	*	*
2564	7	*	*	*	*	*	*	*
3001	1,145	0.5	4.5	9.1	65.7	18.8	1.4	0.1
3002	296	0.0	4.4	11.5	63.5	19.6	1.0	0.0
3004	193	0.0	2.1	6.7	63.2	25.9	2.1	0.0
3005	19	*	*	*	*	*	*	*
3007	26	*	*	*	*	*	*	*
3009	6	*	*	*	*	*	*	*
3201	460	0.0	2.0	7.4	64.6	25.0	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 83

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3301	4,371	0.2	2.8	8.0	62.5	24.4	2.0	0.1
3304	107	0.0	0.0	0.0	52.3	47.7	0.0	0.0
3305	162	0.0	1.9	7.4	46.9	42.0	1.9	0.0
3306	257	0.0	10.5	6.6	54.9	27.2	0.4	0.4
3308	74	*	*	*	*	*	*	*
3309	37	*	*	*	*	*	*	*
3312	429	0.0	2.3	7.2	60.6	28.2	1.4	0.2
3550	65	*	*	*	*	*	*	*
3551	29	*	*	*	*	*	*	*
3552	52	*	*	*	*	*	*	*
3701	1,092	0.0	2.7	9.2	66.7	20.5	0.9	0.0
3850	3,160	0.1	3.3	9.2	66.3	20.1	0.8	0.2
3901	1,142	0.1	3.5	10.9	62.1	21.7	1.7	0.0
3902	350	0.0	3.1	9.1	61.1	24.0	2.6	0.0
3903	28	*	*	*	*	*	*	*
3904	7	*	*	*	*	*	*	*
3905	11	*	*	*	*	*	*	*
3950	2,114	0.0	3.2	8.2	62.1	24.3	2.1	0.0
3954	12	*	*	*	*	*	*	*
4001	512	0.4	4.5	10.5	61.7	20.3	2.5	0.0
4002	327	0.0	4.6	7.6	60.6	26.0	1.2	0.0
4003	2	*	*	*	*	*	*	*
4004	4	*	*	*	*	*	*	*
4103	42	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 84

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4106	2,566	0.3	3.9	9.7	59.0	24.7	2.1	0.2
4107	3,237	0.1	2.9	8.0	61.3	26.0	1.5	0.2
4109	1,039	0.2	2.6	6.1	64.5	24.7	1.8	0.1
4110	5	*	*	*	*	*	*	*
4112	568	0.0	5.3	8.3	51.9	30.6	3.7	0.2
4113	343	0.0	2.3	7.9	47.8	32.9	3.5	5.5
4115	3,610	0.1	2.3	6.5	58.8	29.8	2.4	0.1
4351	582	0.0	2.4	11.2	66.2	19.6	0.7	0.0
4352	167	0.0	1.8	10.8	67.1	18.6	1.8	0.0
4353	421	0.0	4.5	8.8	66.5	18.8	1.4	0.0
4354	774	0.1	2.5	9.2	68.5	18.7	0.9	0.1
4355	357	0.0	4.2	13.4	57.7	21.6	2.8	0.3
4356	467	0.2	3.9	6.2	68.7	20.6	0.4	0.0
4358	141	0.7	2.8	5.0	63.8	25.5	2.1	0.0
4359	485	0.2	2.9	7.6	61.6	24.5	3.1	0.0
4360	230	0.0	1.7	10.9	69.1	16.1	2.2	0.0
4361	268	0.0	1.5	8.2	64.2	25.0	1.1	0.0
4362	304	0.0	4.3	7.6	67.4	20.1	0.7	0.0
4363	907	0.0	2.8	7.6	66.4	22.2	1.0	0.1
4401	1,077	0.1	3.0	7.5	65.6	22.7	1.0	0.1
4501	990	0.2	2.6	8.1	65.4	21.8	1.8	0.1
4601	1,396	0.0	2.4	9.7	65.0	22.0	0.8	0.0
4602	23	*	*	*	*	*	*	*
4701	1,116	0.0	2.0	6.2	60.3	29.7	1.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 85

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5050	1,740	0.0	1.9	5.5	59.3	31.7	1.7	0.0
5051	2,729	0.0	0.8	4.4	65.0	27.8	2.0	0.0
5052	2,959	0.1	2.2	6.0	62.8	26.6	2.3	0.0
5055	1,193	0.0	0.2	0.8	54.3	40.5	4.1	0.1
5101	1,833	0.1	0.8	5.3	62.5	28.7	2.6	0.0
5102	14	*	*	*	*	*	*	*
5202	795	0.1	2.4	7.4	60.0	28.9	1.1	0.0
5203	4	*	*	*	*	*	*	*
5204	4	*	*	*	*	*	*	*
5601	1,078	0.0	1.7	7.1	66.0	23.7	1.5	0.1
5801	1,772	0.0	3.3	10.5	63.7	21.4	1.1	0.1
5803	137	0.0	0.7	2.2	59.1	35.8	2.2	0.0
5804	17	*	*	*	*	*	*	*
5806	50	*	*	*	*	*	*	*
6101	35	*	*	*	*	*	*	*
6102	122	0.0	9.0	8.2	56.6	24.6	1.6	0.0
6103	4,048	0.3	3.6	9.6	64.7	20.5	1.3	0.1
6105	22	*	*	*	*	*	*	*
6106	103	0.0	4.9	8.7	67.0	18.4	1.0	0.0
6301	4,171	0.1	2.7	8.3	61.3	25.4	2.1	0.0
6312	2,184	0.0	1.1	4.8	56.7	34.5	2.9	0.0
6316	1,283	0.0	0.8	3.7	57.3	35.0	3.2	0.0
6318	14	*	*	*	*	*	*	*
6320	129	0.0	0.0	0.8	56.6	41.1	1.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 86

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
6325	192	0.0	1.6	3.1	63.5	27.6	1.0	3.1
6330	118	0.0	1.7	10.2	61.9	26.3	0.0	0.0
6335	1,059	0.0	0.9	4.0	54.0	37.9	3.2	0.0
6340	50	*	*	*	*	*	*	*
6345	60	*	*	*	*	*	*	*
6360	641	0.2	1.1	8.3	61.8	25.6	2.2	0.9
7304	92	*	*	*	*	*	*	*
7306	2,555	0.2	2.7	8.4	65.1	22.1	1.5	0.0
7308	85	*	*	*	*	*	*	*
7310	22	*	*	*	*	*	*	*
7401	2,336	0.1	3.3	8.3	64.1	22.9	1.3	0.0
7601	650	0.0	3.8	9.4	63.4	22.8	0.6	0.0
7850	1,172	0.1	3.2	10.1	68.6	16.7	1.3	0.0
7852	21	*	*	*	*	*	*	*
8101	1,913	0.4	2.9	9.2	63.5	21.6	2.4	0.0
8102	426	0.0	1.9	3.8	62.7	29.6	1.9	0.2
8105	1,657	0.1	1.8	4.9	61.6	29.5	2.0	0.1
8106	486	0.0	3.1	5.1	64.4	24.7	2.7	0.0
8150	33	*	*	*	*	*	*	*
8151	3,273	0.2	2.4	6.5	60.3	28.0	2.4	0.1
8152	54	*	*	*	*	*	*	*
8153	7	*	*	*	*	*	*	*
8155	20	*	*	*	*	*	*	*
8157	9	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 87

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8201	2,628	0.1	2.5	6.1	62.2	27.0	2.1	0.0
8202	4,023	0.0	1.1	3.8	55.7	34.1	4.3	1.0
8203	445	0.4	1.3	7.0	57.8	31.2	2.2	0.0
8250	814	0.2	1.7	8.2	60.0	28.6	1.2	0.0
8252	160	0.0	1.9	5.0	71.9	20.0	1.3	0.0
8255	2,629	0.0	1.7	6.2	66.3	24.3	1.4	0.0
8257	2,115	0.1	0.9	3.9	58.5	31.7	4.7	0.1
8263	727	0.0	0.8	5.2	65.3	27.2	1.4	0.0
8264	701	0.0	2.6	5.3	63.6	26.0	2.6	0.0
8266	9	*	*	*	*	*	*	*
8267	887	0.1	3.4	6.5	62.2	26.2	1.6	0.0
8901	1,143	0.1	2.8	8.1	65.3	21.7	1.9	0.1
8902	1,503	0.3	1.8	7.7	64.4	23.6	2.2	0.1
8903	1,445	0.1	3.0	8.0	63.7	23.5	1.6	0.1
8904	2,472	0.1	2.9	7.2	65.8	22.1	1.9	0.0
8905	1,609	0.0	1.3	4.4	57.7	33.3	3.3	0.0
8908	18	*	*	*	*	*	*	*
8909	641	0.3	4.5	8.1	60.8	24.3	1.7	0.2
9016	7	*	*	*	*	*	*	*
9018	3	*	*	*	*	*	*	*
9045	1,885	0.1	3.6	8.0	56.9	28.6	2.6	0.1
9050	968	0.3	2.9	8.0	61.9	24.4	2.6	0.0
9051	1,753	0.1	3.1	7.5	52.0	31.0	6.3	0.0
9053	1,011	0.0	2.3	6.6	66.6	22.3	2.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 88

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9055	29	*	*	*	*	*	*	*
9058	1,062	0.2	2.1	5.1	59.6	29.3	3.7	0.1
9066	1,140	0.1	3.1	7.0	63.3	24.3	2.2	0.0
9068	962	0.1	3.0	8.9	62.5	23.5	1.9	0.1
9069	2,883	0.1	2.9	6.4	64.7	23.9	2.0	0.0
9070	1,401	0.2	2.6	9.6	63.1	22.6	1.9	0.0
9072	1,809	0.2	2.5	9.6	65.2	21.2	1.3	0.0
9080	2,885	0.0	1.8	6.3	62.2	27.4	2.2	0.0
9081	16	*	*	*	*	*	*	*
9082	165	0.0	6.1	7.9	49.7	36.4	0.0	0.0
9083	123	0.0	0.0	4.1	64.2	26.8	4.1	0.8
9084	43	*	*	*	*	*	*	*
9085	53	*	*	*	*	*	*	*
9086	48	*	*	*	*	*	*	*
9087	1	*	*	*	*	*	*	*
Michigan	167,191	0.1	2.6	7.6	62.5	25.2	1.9	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Maternal Health Indicators by Clinic Combined 3 Years

Page : 89

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	68	*	*	15	*	*	15	*	27	*	106	3.8 (3)	106	10.4 (56)
0102	362	5.0 (114)	60.8 (95)	161	15.5 (20)	52.2 (58)	77	*	101	35.6 (51)	466	5.2 (15)	466	8.4 (23)
0103	369	4.3 (79)	59.9 (75)	237	19.8 (96)	48.5 (21)	59	*	99	*	412	8.0 (110)	412	10.7 (63)
0104	126	3.2 (26)	54.8 (20)	66	*	*	16	*	29	*	140	7.1 (78)	140	8.6 (27)
0105	8	*	*	6	*	*	1	*	0	*	9	*	9	*
0250	150	4.0 (62)	64.0 (129)	88	*	*	18	*	33	*	142	7.0 (73)	143	9.8 (39)
0251	73	*	*	17	*	*	10	*	27	*	94	*	93	*
0252	115	2.6 (12)	62.6 (116)	101	26.7 (129)	56.4 (115)	11	*	29	*	106	9.4 (141)	106	14.2 (128)
0253	95	*	*	25	*	*	13	*	35	*	143	8.4 (122)	143	18.9 (157)
0401	465	3.7 (46)	59.6 (71)	349	19.5 (90)	55.0 (103)	64	*	109	29.4 (32)	457	4.4 (6)	457	10.9 (68)
0402	383	7.0 (146)	55.9 (31)	176	16.5 (32)	53.4 (76)	68	*	71	*	383	8.1 (112)	383	8.9 (31)
0403	145	7.6 (147)	51.7 (4)	85	*	*	22	*	36	*	143	4.9 (10)	143	8.4 (23)
0404	116	0.9 (1)	62.9 (118)	90	*	*	16	*	23	*	117	6.8 (66)	117	11.1 (78)
0501	320	4.4 (88)	56.9 (36)	95	*	*	78	*	84	*	495	5.9 (37)	495	8.3 (21)
0503	103	7.8 (149)	51.5 (2)	58	*	*	15	*	16	*	109	7.3 (82)	109	11.0 (73)
0504	439	4.1 (67)	55.4 (22)	195	18.5 (72)	50.3 (35)	85	*	71	*	475	6.1 (41)	475	12.4 (98)
0505	266	3.4 (31)	56.8 (35)	145	19.3 (87)	53.8 (83)	47	*	47	*	302	6.6 (53)	303	8.9 (31)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 90

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	125	8.0 (150)	58.4 (56)	34	*	*	21	*	20	*	183	8.7 (130)	183	10.4 (56)
0507	47	*	*	20	*	*	6	*	20	*	81	*	81	*
0508	22	*	*	8	*	*	4	*	6	*	34	*	34	*
0509	6	*	*	5	*	*	0	*	1	*	6	*	6	*
0650	948	4.2 (73)	61.4 (99)	639	17.2 (43)	54.1 (88)	173	32.9 (10)	234	26.1 (19)	1,064	7.4 (89)	1,065	11.0 (73)
0651	591	5.2 (123)	59.2 (67)	445	17.3 (45)	51.7 (50)	86	*	90	*	579	6.2 (45)	579	8.1 (18)
0652	225	3.6 (40)	55.6 (27)	144	19.4 (88)	49.3 (28)	55	*	51	*	317	6.6 (53)	317	13.2 (111)
0653	362	2.8 (15)	58.8 (62)	243	11.5 (2)	54.7 (98)	52	*	65	*	386	5.7 (28)	387	13.2 (111)
0654	346	2.3 (9)	65.6 (137)	227	22.0 (117)	52.4 (64)	48	*	87	*	347	7.8 (103)	347	12.1 (91)
0655	332	5.1 (121)	59.0 (64)	197	17.8 (55)	57.4 (125)	44	*	71	*	334	7.2 (81)	337	15.7 (145)
0657	47	*	*	25	*	*	8	*	14	*	68	*	68	*
0701	269	4.8 (106)	44.2 (1)	113	15.0 (15)	44.2 (6)	58	*	127	14.2 (3)	374	5.3 (18)	375	9.3 (33)
0702	61	*	*	39	*	*	5	*	12	*	58	*	58	*
0703	170	4.1 (67)	60.0 (78)	75	*	*	31	*	44	*	194	8.8 (134)	194	10.3 (50)
0704	48	*	*	32	*	*	9	*	14	*	46	*	46	*
0705	71	*	*	26	*	*	23	*	35	*	123	8.9 (135)	123	9.8 (39)
0706	8	*	*	5	*	*	0	*	4	*	12	*	12	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 91

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	593	4.2 (73)	60.0 (78)	389	18.8 (78)	52.7 (68)	147	42.2 (26)	177	16.9 (7)	798	10.3 (148)	801	15.9 (147)
0851	422	5.9 (139)	53.8 (15)	128	15.6 (24)	54.7 (98)	102	36.3 (13)	163	16.6 (6)	675	7.0 (73)	676	12.0 (90)
0901	1,377	4.1 (67)	60.8 (95)	1,004	19.2 (86)	52.2 (58)	252	32.9 (10)	229	18.3 (9)	1,551	6.7 (57)	1,551	10.1 (48)
0902	39	*	*	28	*	*	3	*	5	*	40	*	40	*
1051	1,214	4.4 (88)	51.6 (3)	995	15.9 (26)	53.5 (80)	189	56.1 (63)	194	37.6 (60)	1,267	8.0 (110)	1,269	9.9 (42)
1101	1,625	3.8 (50)	61.8 (106)	1,224	20.2 (101)	52.5 (66)	304	51.3 (51)	261	35.6 (51)	1,768	6.3 (47)	1,767	14.4 (133)
1102	703	3.1 (24)	59.3 (68)	494	15.4 (18)	56.7 (119)	115	37.4 (16)	172	26.7 (23)	751	6.1 (41)	751	9.5 (36)
1104	0	*	*	0	*	*	0	*	0	*	1	*	1	*
1106	70	*	*	50	*	*	12	*	22	*	68	*	69	*
1250	793	5.5 (131)	61.2 (97)	379	17.9 (59)	51.7 (50)	176	14.8 (1)	177	26.6 (22)	973	8.4 (122)	973	7.9 (16)
1251	741	5.8 (138)	58.0 (50)	531	19.6 (92)	51.6 (47)	104	37.5 (17)	169	27.2 (24)	754	5.7 (28)	754	6.8 (12)
1252	484	3.9 (58)	63.4 (123)	210	13.8 (9)	61.0 (131)	81	*	105	30.5 (38)	560	5.7 (28)	560	13.9 (126)
1253	659	4.4 (88)	58.1 (51)	348	18.4 (70)	52.0 (55)	118	31.4 (7)	157	27.4 (25)	736	7.5 (93)	738	12.2 (92)
1350	2,029	3.1 (24)	61.6 (102)	1,145	19.9 (97)	54.1 (88)	328	47.3 (41)	462	34.0 (45)	2,206	7.9 (107)	2,202	10.0 (44)
1351	113	0.9 (1)	66.4 (143)	100	20.0 (98)	57.0 (121)	16	*	40	*	129	10.9 (151)	129	10.9 (68)
1352	209	6.2 (144)	53.6 (14)	101	21.8 (114)	54.5 (93)	46	*	81	*	351	5.7 (28)	351	4.3 (5)
1401	226	4.9 (108)	53.5 (13)	167	18.0 (60)	55.7 (109)	41	*	35	*	229	7.4 (89)	229	10.0 (44)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Maternal Health Indicators by Clinic Combined 3 Years

Page : 92

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1402	121	2.5 (10)	66.1 (142)	93	*	*	22	*	20	*	123	8.1 (112)	123	6.5 (9)
1403	24	*	*	19	*	*	1	*	6	*	24	*	24	*
1501	727	4.3 (79)	63.4 (123)	512	49.0 (130)	23.6 (2)	84	*	121	24.0 (17)	712	7.0 (73)	712	8.7 (29)
1503	951	3.6 (40)	60.7 (91)	605	16.9 (39)	53.9 (84)	123	43.1 (27)	137	29.9 (34)	998	7.0 (73)	1,000	12.9 (104)
1504	1,750	4.7 (100)	59.8 (73)	1,192	18.0 (60)	47.4 (12)	323	55.4 (60)	591	42.1 (76)	1,995	9.2 (139)	1,998	8.5 (26)
1505	332	3.3 (30)	56.9 (36)	193	15.5 (20)	51.8 (52)	63	*	129	38.8 (66)	404	8.2 (117)	404	12.6 (102)
1506	491	4.7 (100)	61.5 (100)	157	17.2 (43)	52.9 (71)	90	*	106	29.2 (31)	549	7.3 (82)	549	9.7 (37)
1507	489	4.5 (94)	55.2 (21)	237	13.5 (8)	51.1 (40)	65	*	116	39.7 (69)	573	8.2 (117)	573	10.5 (61)
1508	41	*	*	15	*	*	4	*	7	*	67	*	67	*
1509	346	3.8 (50)	65.6 (137)	189	20.1 (100)	51.9 (54)	53	*	99	*	366	10.1 (145)	367	9.3 (33)
1511	585	5.1 (121)	60.5 (86)	279	21.9 (115)	48.0 (15)	115	40.9 (22)	193	32.1 (40)	681	6.9 (70)	684	13.3 (117)
1512	254	4.3 (79)	55.5 (26)	77	*	*	68	*	96	*	343	5.8 (32)	344	10.5 (61)
1513	46	*	*	31	*	*	5	*	12	*	49	*	49	*
1514	168	6.0 (141)	55.4 (22)	123	11.4 (1)	50.4 (36)	36	*	89	*	221	5.9 (37)	221	10.0 (44)
1701	519	4.0 (62)	59.9 (75)	254	16.5 (32)	56.3 (114)	89	*	122	27.9 (27)	632	7.3 (82)	631	11.9 (89)
1950	569	4.2 (73)	61.9 (109)	454	19.6 (92)	52.0 (55)	78	*	149	18.1 (8)	564	8.5 (126)	566	14.7 (138)
1952	115	6.1 (143)	63.5 (126)	99	*	*	21	*	43	*	120	6.7 (57)	120	11.7 (85)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 93

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1953	304	6.6 (145)	53.9 (16)	255	23.9 (124)	49.0 (25)	49	*	106	18.9 (10)	314	9.2 (139)	314	16.6 (152)
1954	516	5.0 (114)	59.5 (70)	374	17.1 (41)	54.8 (101)	80	*	107	5.6 (1)	534	10.5 (149)	534	14.0 (127)
1955	93	*	*	63	*	*	15	*	20	*	90	*	90	*
1957	445	4.7 (100)	61.6 (102)	344	15.4 (18)	54.1 (88)	66	*	132	14.4 (4)	453	8.2 (117)	453	14.3 (129)
1963	32	*	*	28	*	*	7	*	9	*	33	*	34	*
1964	45	*	*	34	*	*	9	*	13	*	43	*	43	*
2150	483	2.7 (14)	55.7 (29)	283	15.5 (20)	53.7 (82)	86	*	152	26.3 (21)	607	11.2 (153)	607	13.5 (118)
2151	167	3.6 (40)	52.1 (6)	119	13.4 (7)	54.6 (97)	36	*	60	*	207	5.3 (18)	208	11.5 (84)
2251	325	5.2 (123)	52.9 (12)	122	11.5 (2)	48.4 (20)	56	*	87	*	378	4.2 (5)	378	13.2 (111)
2252	144	7.6 (147)	52.8 (10)	44	*	*	26	*	36	*	168	8.9 (135)	168	13.1 (108)
2554	60	*	*	41	*	*	9	*	14	*	65	*	65	*
2555	6,345	5.4 (128)	57.7 (48)	3,103	18.2 (66)	52.3 (62)	1,458	50.5 (49)	1,327	33.8 (44)	7,452	7.6 (97)	7,452	15.5 (144)
2556	283	3.5 (35)	56.5 (34)	231	21.6 (112)	51.1 (40)	35	*	55	*	322	7.8 (103)	322	16.8 (155)
2557	4	*	*	0	*	*	6	*	1	*	8	*	8	*
2558	6	*	*	4	*	*	3	*	1	*	7	*	7	*
2559	7	*	*	4	*	*	3	*	0	*	9	*	9	*
2560	3	*	*	1	*	*	0	*	0	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 94

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2561	1	*	*	1	*	*	1	*	0	*	2	*	2	*
2562	11	*	*	3	*	*	5	*	3	*	11	*	11	*
2563	15	*	*	4	*	*	2	*	6	*	20	*	20	*
2564	3	*	*	1	*	*	2	*	1	*	7	*	7	*
3001	1,070	3.5 (35)	63.6 (127)	807	23.0 (120)	45.8 (8)	208	51.0 (50)	724	42.3 (78)	1,066	5.4 (20)	1,065	10.9 (68)
3002	284	1.1 (4)	81.0 (151)	223	20.6 (105)	51.1 (40)	45	*	206	44.2 (81)	278	4.7 (8)	278	13.7 (123)
3004	179	2.2 (8)	65.9 (140)	152	24.3 (126)	38.2 (4)	26	*	36	*	174	5.2 (15)	174	8.6 (27)
3005	16	*	*	7	*	*	7	*	7	*	18	*	18	*
3007	22	*	*	12	*	*	1	*	9	*	19	*	19	*
3009	4	*	*	0	*	*	0	*	1	*	1	*	1	*
3201	377	5.0 (114)	64.7 (134)	289	16.6 (35)	51.2 (44)	57	*	113	22.1 (15)	430	7.9 (107)	431	12.5 (101)
3301	3,707	4.2 (73)	60.5 (86)	2,961	19.1 (84)	52.0 (55)	759	47.0 (38)	700	34.6 (47)	4,100	6.8 (66)	4,101	10.3 (50)
3304	84	*	*	69	*	*	11	*	18	*	102	17.6 (156)	102	5.9 (6)
3305	135	5.9 (139)	60.0 (78)	107	20.6 (105)	54.2 (91)	21	*	30	*	149	10.1 (145)	149	3.4 (3)
3306	220	5.5 (131)	52.7 (9)	186	12.4 (4)	57.0 (121)	47	*	67	*	243	3.7 (2)	243	7.8 (14)
3308	67	*	*	54	*	*	6	*	25	*	68	*	68	*
3309	35	*	*	24	*	*	6	*	9	*	34	*	34	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 95

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3312	354	5.4 (128)	64.1 (130)	294	17.7 (52)	52.7 (68)	89	*	68	*	408	8.1 (112)	409	8.1 (18)
3550	57	*	*	28	*	*	12	*	21	*	60	*	60	*
3551	23	*	*	10	*	*	2	*	15	*	27	*	27	*
3552	40	*	*	14	*	*	6	*	12	*	46	*	46	*
3701	726	4.4 (88)	61.8 (106)	483	14.9 (12)	55.5 (107)	160	24.4 (2)	284	14.1 (2)	1,002	8.2 (117)	1,000	15.8 (146)
3850	2,334	5.7 (136)	60.2 (81)	1,346	18.6 (74)	53.4 (76)	476	46.4 (36)	542	41.5 (73)	2,913	6.1 (41)	2,916	10.7 (63)
3901	895	4.6 (98)	57.4 (44)	424	19.6 (92)	51.2 (44)	146	61.6 (66)	257	40.1 (70)	1,042	6.7 (57)	1,042	10.9 (68)
3902	295	3.4 (31)	63.7 (128)	151	13.2 (5)	55.0 (103)	43	*	74	*	321	9.7 (142)	322	7.8 (14)
3903	24	*	*	15	*	*	2	*	16	*	27	*	27	*
3904	6	*	*	1	*	*	3	*	1	*	7	*	7	*
3905	9	*	*	7	*	*	0	*	2	*	11	*	11	*
3950	1,697	3.4 (31)	59.7 (72)	992	17.0 (40)	55.2 (105)	300	44.7 (29)	545	36.5 (56)	1,919	9.1 (138)	1,919	10.0 (44)
3954	10	*	*	6	*	*	2	*	3	*	11	*	11	*
4001	427	4.2 (73)	60.4 (85)	173	18.5 (72)	49.1 (27)	54	*	72	*	448	7.1 (78)	449	14.3 (129)
4002	284	4.9 (108)	63.4 (123)	159	15.7 (25)	59.7 (130)	37	*	48	*	301	7.6 (97)	301	8.3 (21)
4003	2	*	*	1	*	*	1	*	0	*	2	*	2	*
4004	4	*	*	3	*	*	0	*	0	*	4	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 96

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4103	19	*	*	5	*	*	5	*	11	*	34	*	34	*
4106	1,895	3.5 (35)	62.0 (110)	985	21.9 (115)	46.3 (10)	452	50.2 (48)	773	38.8 (66)	2,349	5.8 (32)	2,350	11.0 (73)
4107	2,641	4.3 (79)	57.3 (42)	1,325	15.9 (26)	50.2 (33)	584	47.6 (42)	1,132	37.7 (62)	3,004	6.7 (57)	3,006	13.1 (108)
4109	464	5.6 (133)	52.8 (10)	193	18.1 (64)	53.4 (76)	154	30.5 (5)	265	18.9 (10)	966	7.6 (97)	971	14.3 (129)
4110	5	*	*	5	*	*	0	*	0	*	5	*	5	*
4112	559	3.0 (20)	62.8 (117)	504	75.2 (131)	6.5 (1)	77	*	73	*	535	17.6 (156)	536	11.4 (82)
4113	300	2.0 (5)	68.3 (149)	64	*	*	52	*	90	*	330	10.6 (150)	329	10.3 (50)
4115	2,184	5.0 (114)	55.4 (22)	664	19.0 (83)	48.3 (19)	633	45.2 (30)	1,249	38.0 (64)	3,366	7.6 (97)	3,368	11.8 (87)
4351	559	3.0 (20)	63.3 (121)	445	22.5 (118)	49.0 (25)	81	*	117	28.2 (28)	552	7.8 (103)	552	11.2 (80)
4352	149	8.7 (151)	61.7 (104)	102	17.6 (48)	52.9 (71)	20	*	34	*	152	7.9 (107)	153	12.4 (98)
4353	350	2.6 (12)	64.3 (131)	225	17.8 (55)	51.1 (40)	63	*	81	*	380	6.6 (53)	379	13.7 (123)
4354	641	4.4 (88)	56.9 (36)	376	17.3 (45)	54.0 (86)	109	28.4 (4)	178	20.2 (12)	725	5.5 (22)	724	13.0 (106)
4355	285	3.5 (35)	58.2 (53)	87	*	*	53	*	98	*	335	8.1 (112)	335	11.3 (81)
4356	369	4.9 (108)	54.7 (19)	158	21.5 (111)	46.2 (9)	84	*	95	*	434	6.7 (57)	434	11.8 (87)
4358	119	2.5 (10)	66.4 (143)	92	*	*	18	*	20	*	123	7.3 (82)	123	11.4 (82)
4359	419	3.8 (50)	62.1 (112)	237	18.6 (74)	49.4 (29)	67	*	98	*	457	11.2 (153)	458	12.9 (104)
4360	219	5.0 (114)	57.5 (45)	176	17.6 (48)	53.4 (76)	27	*	46	*	214	1.9 (1)	214	11.7 (85)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 97

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4361	235	4.7 (100)	58.3 (54)	187	15.5 (20)	59.4 (129)	31	*	66	*	252	6.7 (57)	252	10.3 (50)
4362	266	6.0 (141)	56.0 (32)	209	18.2 (66)	55.5 (107)	57	*	84	*	287	8.7 (130)	287	9.8 (39)
4363	846	3.8 (50)	58.7 (61)	611	20.6 (105)	50.4 (36)	121	39.7 (21)	190	30.0 (36)	836	6.1 (41)	836	12.3 (95)
4401	1,059	4.5 (94)	56.2 (33)	931	15.9 (26)	54.7 (98)	185	38.9 (18)	290	32.4 (41)	1,012	4.1 (4)	1,012	6.1 (8)
4501	738	5.3 (127)	56.9 (36)	570	15.3 (17)	56.8 (120)	124	25.0 (3)	185	21.6 (14)	948	8.4 (122)	948	12.4 (98)
4601	1,111	3.7 (46)	57.5 (45)	262	21.4 (110)	47.7 (14)	284	46.8 (37)	337	38.9 (68)	1,273	7.5 (93)	1,274	13.0 (106)
4602	15	*	*	8	*	*	1	*	7	*	23	*	23	*
4701	1,035	5.0 (114)	52.5 (8)	684	17.4 (47)	52.5 (66)	179	34.1 (12)	378	15.1 (5)	1,018	12.0 (155)	1,018	14.7 (138)
5050	1,668	3.9 (58)	61.8 (106)	1,446	18.1 (64)	53.9 (84)	376	48.1 (44)	435	32.9 (43)	1,599	7.6 (97)	1,597	12.3 (95)
5051	2,579	4.7 (100)	58.4 (56)	2,223	18.4 (70)	52.8 (70)	545	47.0 (38)	785	35.4 (49)	2,533	7.7 (102)	2,533	10.7 (63)
5052	2,779	4.5 (94)	60.7 (91)	2,350	20.3 (102)	49.4 (29)	596	44.1 (28)	694	34.1 (46)	2,751	8.4 (122)	2,752	10.4 (56)
5055	628	2.1 (6)	54.6 (18)	255	17.6 (48)	53.3 (74)	111	41.4 (23)	294	37.4 (59)	1,143	8.7 (130)	1,144	4.2 (4)
5101	1,645	4.4 (88)	58.4 (56)	1,208	15.9 (26)	52.2 (58)	331	41.7 (24)	459	30.3 (37)	1,704	9.7 (142)	1,704	12.3 (95)
5102	14	*	*	9	*	*	1	*	5	*	13	*	13	*
5202	697	3.6 (40)	55.4 (22)	510	19.4 (88)	47.5 (13)	104	36.5 (14)	198	23.7 (16)	704	5.1 (11)	710	14.4 (133)
5203	3	*	*	2	*	*	0	*	0	*	3	*	3	*
5204	4	*	*	2	*	*	1	*	0	*	4	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 98

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5601	934	2.8 (15)	60.2 (81)	189	21.2 (109)	54.5 (93)	157	31.8 (9)	400	25.3 (18)	1,023	6.0 (40)	1,024	12.7 (103)
5801	1,334	3.8 (50)	59.9 (75)	765	22.6 (119)	48.8 (24)	312	52.9 (53)	354	35.6 (51)	1,618	7.0 (73)	1,618	13.6 (120)
5803	126	4.8 (106)	57.1 (41)	85	*	*	17	*	44	*	128	10.9 (151)	128	14.8 (140)
5804	9	*	*	8	*	*	0	*	5	*	15	*	15	*
5806	32	*	*	25	*	*	1	*	12	*	47	*	48	*
6101	33	*	*	25	*	*	1	*	9	*	32	*	32	*
6102	115	5.2 (123)	55.7 (29)	86	*	*	16	*	22	*	118	8.5 (126)	118	14.4 (133)
6103	3,727	4.0 (62)	61.5 (100)	2,650	20.3 (102)	50.2 (33)	537	45.4 (32)	461	27.8 (26)	3,719	8.1 (112)	3,725	14.3 (129)
6105	22	*	*	17	*	*	0	*	4	*	20	*	21	*
6106	99	*	*	70	*	*	16	*	12	*	96	*	96	*
6301	3,816	3.8 (50)	58.1 (51)	2,729	19.5 (90)	51.6 (47)	782	48.1 (44)	855	35.4 (49)	3,863	5.6 (25)	3,865	10.7 (63)
6312	2,022	3.7 (46)	60.5 (86)	1,501	16.5 (32)	52.2 (58)	440	50.0 (47)	597	40.2 (71)	2,047	4.5 (7)	2,047	8.0 (17)
6316	1,215	4.0 (62)	57.8 (49)	981	14.9 (12)	57.1 (123)	257	45.5 (33)	332	34.6 (47)	1,181	6.8 (66)	1,181	10.4 (56)
6318	11	*	*	8	*	*	6	*	3	*	12	*	12	*
6320	95	*	*	46	*	*	18	*	42	*	119	10.1 (145)	119	6.7 (11)
6325	141	5.7 (136)	58.9 (63)	61	*	*	29	*	73	*	177	5.1 (11)	177	16.4 (151)
6330	63	*	*	17	*	*	14	*	15	*	107	6.5 (51)	108	14.8 (140)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 99

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6335	691	3.9 (58)	51.7 (4)	223	24.2 (125)	44.4 (7)	148	45.9 (34)	210	30.5 (38)	1,006	8.5 (126)	1,006	8.4 (23)
6340	41	*	*	14	*	*	2	*	8	*	43	*	42	*
6345	50	*	*	31	*	*	13	*	24	*	55	*	55	*
6360	471	4.9 (108)	58.6 (59)	185	18.9 (79)	49.7 (32)	99	*	136	41.9 (74)	586	6.7 (57)	585	13.2 (111)
7304	75	*	*	56	*	*	14	*	20	*	87	*	87	*
7306	2,056	4.1 (67)	60.7 (91)	1,535	20.0 (98)	50.4 (36)	419	54.4 (56)	439	36.4 (55)	2,381	6.7 (57)	2,381	10.8 (67)
7308	78	*	*	60	*	*	11	*	9	*	75	*	75	*
7310	6	*	*	3	*	*	2	*	5	*	19	*	19	*
7401	2,212	5.6 (133)	57.0 (40)	1,898	16.8 (37)	56.2 (112)	324	49.7 (46)	458	37.6 (60)	2,138	6.9 (70)	2,145	13.7 (123)
7601	183	4.9 (108)	59.0 (64)	72	*	*	105	31.4 (7)	122	21.3 (13)	594	6.6 (53)	594	9.9 (42)
7850	1,124	4.7 (100)	60.6 (89)	932	19.1 (84)	48.2 (18)	175	38.9 (18)	236	35.6 (51)	1,081	8.6 (129)	1,082	12.2 (92)
7852	20	*	*	16	*	*	3	*	4	*	21	*	21	*
8101	1,524	4.0 (62)	65.0 (136)	969	18.9 (79)	52.3 (62)	325	67.1 (72)	256	53.9 (88)	1,790	6.5 (51)	1,792	16.3 (149)
8102	324	3.4 (31)	67.0 (146)	164	17.7 (52)	53.0 (73)	65	*	41	*	392	8.2 (117)	392	12.2 (92)
8105	1,337	4.3 (79)	63.0 (119)	850	18.9 (79)	53.5 (80)	321	60.7 (65)	360	44.7 (82)	1,529	8.7 (130)	1,529	13.5 (118)
8106	241	3.7 (46)	55.6 (27)	154	18.2 (66)	56.5 (117)	92	*	95	*	457	5.5 (22)	456	9.4 (35)
8150	26	*	*	11	*	*	5	*	10	*	28	*	28	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 100

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8151	2,957	3.8 (50)	59.8 (73)	1,950	20.4 (104)	49.6 (31)	607	54.2 (55)	886	37.9 (63)	2,938	7.4 (89)	2,939	13.2 (111)
8152	50	*	*	31	*	*	13	*	17	*	49	*	49	*
8153	7	*	*	5	*	*	0	*	3	*	6	*	6	*
8155	18	*	*	12	*	*	3	*	7	*	19	*	19	*
8157	9	*	*	4	*	*	2	*	4	*	7	*	7	*
8201	1,987	5.0 (114)	59.1 (66)	1,166	17.6 (48)	54.0 (86)	519	47.2 (40)	776	29.9 (34)	2,400	7.3 (82)	2,401	10.3 (50)
8202	2,530	4.9 (108)	52.2 (7)	1,224	23.0 (120)	42.5 (5)	728	53.2 (54)	912	43.2 (79)	3,835	6.7 (57)	3,832	3.3 (2)
8203	392	2.8 (15)	62.0 (110)	337	14.2 (10)	56.1 (111)	88	*	135	32.6 (42)	414	7.5 (93)	414	16.7 (154)
8250	708	3.8 (50)	60.6 (89)	555	17.7 (52)	56.6 (118)	199	45.2 (30)	241	46.9 (85)	742	8.9 (135)	742	16.3 (149)
8252	140	4.3 (79)	67.9 (148)	101	14.9 (12)	56.4 (115)	41	*	38	*	154	7.1 (78)	154	11.0 (73)
8255	2,171	4.6 (98)	58.3 (54)	1,443	13.2 (5)	55.9 (110)	575	47.7 (43)	829	37.0 (58)	2,424	7.3 (82)	2,425	10.3 (50)
8257	1,399	4.1 (67)	54.0 (17)	747	26.4 (128)	36.8 (3)	353	55.5 (61)	594	43.6 (80)	1,994	9.7 (142)	1,994	5.9 (6)
8263	648	3.9 (58)	60.2 (81)	503	18.3 (69)	54.5 (93)	147	30.6 (6)	278	28.8 (30)	678	7.8 (103)	678	11.1 (78)
8264	476	3.2 (26)	58.6 (59)	302	16.2 (31)	55.3 (106)	149	38.9 (18)	230	26.1 (19)	635	6.8 (66)	635	8.8 (30)
8266	4	*	*	2	*	*	0	*	2	*	9	*	9	*
8267	558	5.6 (133)	57.3 (42)	362	17.1 (41)	53.3 (74)	209	42.1 (25)	239	29.7 (33)	795	6.2 (45)	797	8.2 (20)
8901	1,013	3.2 (26)	63.3 (121)	687	18.0 (60)	52.4 (64)	192	67.2 (73)	146	63.0 (93)	1,056	5.8 (32)	1,058	14.4 (133)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 101

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8902	1,294	5.2 (123)	61.2 (97)	818	18.6 (74)	54.8 (101)	247	63.2 (67)	249	55.8 (89)	1,393	5.1 (11)	1,393	13.2 (111)
8903	1,286	3.0 (20)	64.4 (132)	810	17.8 (55)	56.2 (112)	256	74.6 (76)	202	60.9 (92)	1,351	5.4 (20)	1,349	16.8 (155)
8904	2,091	3.6 (40)	62.2 (113)	1,381	18.0 (60)	54.5 (93)	455	68.4 (74)	372	57.0 (90)	2,302	5.6 (25)	2,304	15.0 (142)
8905	1,267	2.9 (18)	59.4 (69)	730	17.8 (55)	51.0 (39)	263	55.9 (62)	286	42.0 (75)	1,508	7.4 (89)	1,508	11.0 (73)
8908	16	*	*	9	*	*	1	*	1	*	18	*	18	*
8909	509	4.3 (79)	64.4 (132)	315	16.8 (37)	58.1 (128)	123	64.2 (69)	90	*	603	7.3 (82)	605	15.2 (143)
9016	6	*	*	3	*	*	0	*	2	*	7	*	7	*
9018	3	*	*	3	*	*	0	*	0	*	3	*	3	*
9045	1,413	2.1 (6)	64.8 (135)	1,108	23.4 (123)	48.1 (16)	313	46.0 (35)	234	36.8 (57)	1,749	6.4 (49)	1,750	7.3 (13)
9050	788	4.3 (79)	60.7 (91)	427	24.6 (127)	48.7 (23)	173	72.8 (75)	119	57.1 (91)	928	6.3 (47)	927	10.2 (49)
9051	1,362	2.9 (18)	65.9 (140)	867	16.0 (30)	46.7 (11)	185	37.3 (15)	208	28.4 (29)	1,631	7.5 (93)	1,630	6.6 (10)
9053	937	3.5 (35)	63.0 (119)	741	14.6 (11)	57.5 (126)	164	64.6 (70)	122	48.4 (86)	962	5.2 (15)	962	13.6 (120)
9055	27	*	*	23	*	*	3	*	2	*	29	*	29	*
9058	709	5.4 (128)	62.5 (115)	505	18.6 (74)	48.1 (16)	195	52.8 (52)	181	38.7 (65)	1,016	6.9 (70)	1,017	13.6 (120)
9066	858	3.0 (20)	66.9 (145)	547	15.2 (16)	57.2 (124)	225	54.7 (57)	161	52.8 (87)	1,079	5.6 (25)	1,078	16.0 (148)
9068	933	3.2 (26)	67.2 (147)	701	23.1 (122)	54.2 (91)	163	55.2 (59)	88	*	905	5.1 (11)	905	9.7 (37)
9069	2,578	3.6 (40)	60.2 (81)	2,128	19.7 (95)	51.6 (47)	529	55.0 (58)	285	40.4 (72)	2,757	5.8 (32)	2,755	10.4 (56)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 102

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9070	1,113	4.1 (67)	57.5 (45)	739	21.1 (108)	51.8 (52)	239	66.9 (71)	174	46.6 (84)	1,322	5.5 (22)	1,323	13.1 (108)
9072	1,525	4.2 (73)	62.3 (114)	1,029	21.6 (112)	51.3 (46)	345	56.2 (64)	230	45.2 (83)	1,718	6.4 (49)	1,719	14.4 (133)
9080	2,282	4.3 (79)	61.7 (104)	1,516	16.7 (36)	57.7 (127)	577	63.6 (68)	485	42.1 (76)	2,740	5.9 (37)	2,740	10.9 (68)
9081	11	*	*	7	*	*	3	*	2	*	12	*	12	*
9082	157	4.5 (94)	65.6 (137)	88	*	*	32	*	20	*	150	4.7 (8)	151	16.6 (152)
9083	117	0.9 (1)	71.8 (150)	111	18.9 (79)	48.6 (22)	3	*	1	*	120	5.8 (32)	120	2.5 (1)
9084	35	*	*	30	*	*	2	*	1	*	42	*	42	*
9085	52	*	*	19	*	*	22	*	5	*	47	*	47	*
9086	46	*	*	25	*	*	15	*	8	*	40	*	40	*
9087	1	*	*	1	*	*	0	*	1	*	1	*	1	*
Michigan	136,140	4.2	59.6	87,782	19.1	51.5	27,604	47.9	35,578	35.0	154,877	7.1	154,947	11.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 103

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	108	85.2 (63)	9.3 (100)	115	29.6 (101)	106	24.5 (54)	89	*
0102	458	85.2 (63)	9.2 (97)	497	34.6 (73)	451	18.6 (87)	410	91.5 (38)
0103	410	78.3 (122)	13.4 (123)	435	37.0 (49)	387	21.2 (72)	358	91.9 (35)
0104	140	81.4 (105)	9.3 (100)	154	37.0 (49)	136	22.1 (63)	123	90.2 (55)
0105	9	*	*	8	*	5	*	6	*
0250	144	73.6 (145)	20.1 (150)	156	46.2 (13)	112	28.6 (29)	130	91.5 (38)
0251	94	*	*	100	47.0 (11)	88	*	79	*
0252	106	73.6 (145)	21.7 (153)	115	36.5 (55)	104	28.8 (27)	88	*
0253	144	80.6 (107)	12.5 (119)	147	55.1 (3)	141	20.6 (78)	134	90.3 (51)
0401	453	83.9 (83)	9.9 (107)	486	37.0 (49)	453	26.3 (41)	397	92.7 (26)
0402	387	83.5 (86)	9.3 (100)	404	35.4 (68)	360	23.6 (58)	322	88.8 (67)
0403	142	84.5 (73)	4.2 (43)	150	32.0 (88)	130	28.5 (31)	119	91.6 (37)
0404	117	84.6 (70)	8.5 (93)	120	44.2 (17)	111	25.2 (47)	96	*
0501	498	94.0 (3)	1.4 (8)	520	33.8 (80)	467	27.0 (37)	399	89.0 (64)
0503	110	88.2 (45)	7.3 (82)	111	33.3 (83)	104	28.8 (27)	90	*
0504	475	90.1 (25)	5.7 (65)	494	29.1 (104)	461	32.1 (20)	388	87.1 (87)
0505	304	84.2 (78)	8.2 (90)	320	36.3 (58)	298	26.8 (38)	268	89.2 (61)
0506	184	83.2 (87)	9.2 (97)	187	48.1 (10)	169	20.7 (77)	156	83.3 (120)
0507	80	*	*	80	*	72	*	50	*
0508	33	*	*	34	*	32	*	24	*
0509	6	*	*	6	*	4	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 104

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	1,071	85.0 (66)	8.8 (95)	1,131	36.3 (58)	976	27.7 (33)	911	93.2 (20)
0651	574	76.5 (130)	15.5 (136)	616	44.3 (16)	540	18.5 (88)	521	88.3 (73)
0652	315	81.6 (103)	14.0 (128)	322	35.7 (66)	286	28.0 (32)	248	85.5 (103)
0653	387	74.9 (139)	19.4 (149)	407	43.5 (19)	379	29.3 (25)	342	88.3 (73)
0654	348	74.7 (142)	19.0 (148)	384	42.2 (20)	350	20.6 (78)	322	78.9 (138)
0655	330	84.5 (73)	5.5 (61)	356	44.9 (14)	325	24.6 (51)	300	90.3 (51)
0657	68	*	*	67	*	59	*	56	*
0701	378	80.2 (108)	14.6 (131)	393	35.6 (67)	364	44.5 (9)	318	92.1 (33)
0702	57	*	*	66	*	64	*	55	*
0703	190	90.5 (18)	2.1 (11)	219	35.2 (69)	196	28.6 (29)	171	95.9 (1)
0704	45	*	*	54	*	50	*	44	*
0705	124	87.9 (49)	9.7 (104)	136	33.8 (80)	126	66.7 (2)	113	90.3 (51)
0706	12	*	*	12	*	9	*	6	*
0850	798	88.3 (44)	3.0 (28)	868	26.6 (113)	775	22.5 (61)	643	93.3 (19)
0851	676	89.9 (30)	6.7 (75)	735	34.3 (75)	678	23.7 (57)	579	94.6 (5)
0901	1,539	74.0 (143)	20.5 (151)	1,647	41.8 (22)	1,511	15.2 (119)	1,372	84.3 (115)
0902	40	*	*	43	*	39	*	36	*
1051	1,266	77.8 (124)	14.6 (131)	1,324	29.9 (100)	1,178	23.6 (58)	932	87.6 (81)
1101	1,732	72.3 (150)	12.2 (117)	1,902	28.0 (109)	1,628	13.1 (129)	1,416	85.9 (97)
1102	736	73.1 (147)	15.6 (137)	828	36.0 (65)	706	17.0 (102)	635	94.3 (8)
1104	1	*	*	1	*	1	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 105

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1106	68	*	*	78	*	69	*	57	*
1250	833	82.2 (97)	9.7 (104)	1,043	36.2 (61)	912	21.8 (65)	834	80.3 (132)
1251	749	79.3 (115)	11.3 (113)	806	36.7 (54)	706	18.1 (92)	623	84.1 (118)
1252	538	70.3 (151)	17.7 (142)	593	41.1 (26)	554	16.4 (109)	494	89.1 (63)
1253	712	73.9 (144)	16.9 (140)	814	34.2 (77)	719	18.2 (90)	637	88.7 (68)
1350	2,143	88.9 (38)	5.1 (58)	2,394	36.5 (55)	1,591	19.3 (84)	1,844	79.6 (135)
1351	129	87.6 (53)	6.2 (71)	135	29.6 (101)	116	25.0 (48)	94	*
1352	349	90.0 (27)	4.6 (51)	372	37.9 (46)	325	2.8 (153)	299	89.3 (59)
1401	228	81.6 (103)	12.3 (118)	245	31.8 (89)	166	32.5 (19)	186	87.6 (81)
1402	122	75.4 (137)	13.9 (127)	130	30.8 (97)	96	*	87	*
1403	24	*	*	23	*	15	*	17	*
1501	712	83.7 (84)	11.4 (114)	800	39.6 (30)	523	16.4 (109)	619	82.1 (124)
1503	982	84.7 (69)	8.4 (91)	1,075	43.6 (18)	899	40.9 (11)	892	77.6 (140)
1504	1,933	90.9 (16)	2.9 (26)	2,157	31.8 (89)	1,578	26.2 (42)	1,538	74.7 (141)
1505	393	90.8 (17)	5.3 (60)	428	27.1 (110)	314	27.4 (34)	278	89.2 (61)
1506	543	72.7 (149)	22.3 (156)	595	38.0 (44)	518	6.4 (149)	480	93.1 (21)
1507	564	90.2 (23)	5.0 (57)	614	38.3 (41)	460	26.1 (43)	463	90.5 (49)
1508	63	*	*	84	*	66	*	67	*
1509	365	90.1 (25)	4.4 (47)	442	41.2 (24)	334	24.6 (51)	356	84.8 (109)
1511	673	88.7 (39)	4.8 (55)	728	36.1 (62)	535	19.3 (84)	553	85.7 (99)
1512	341	90.0 (27)	6.7 (75)	365	26.6 (113)	254	31.9 (23)	263	90.5 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 106

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1513	49	*	*	53	*	45	*	45	*
1514	218	92.7 (6)	4.6 (51)	232	25.9 (119)	173	26.6 (39)	147	92.5 (29)
1701	636	85.8 (60)	10.4 (109)	680	46.3 (12)	590	49.0 (8)	566	84.5 (113)
1950	557	78.5 (119)	15.3 (135)	598	38.8 (38)	520	21.5 (68)	468	91.2 (43)
1952	120	81.7 (102)	13.3 (122)	120	34.2 (77)	106	21.7 (67)	93	*
1953	313	82.7 (92)	12.8 (121)	322	36.3 (58)	293	21.8 (65)	250	94.4 (6)
1954	531	80.2 (108)	13.4 (123)	564	36.9 (52)	509	27.3 (35)	456	93.4 (17)
1955	91	*	*	97	*	85	*	75	*
1957	454	86.1 (59)	6.8 (77)	476	31.7 (92)	403	24.8 (50)	350	88.9 (66)
1963	33	*	*	35	*	28	*	21	*
1964	42	*	*	48	*	45	*	40	*
2150	608	91.0 (14)	2.6 (20)	671	44.4 (15)	617	24.3 (55)	559	92.5 (29)
2151	204	84.3 (75)	12.7 (120)	241	36.1 (62)	189	24.3 (55)	178	88.2 (75)
2251	380	89.5 (31)	7.4 (83)	414	39.4 (33)	385	33.0 (18)	342	94.2 (9)
2252	175	78.9 (116)	18.3 (146)	203	52.7 (4)	172	57.6 (3)	177	86.4 (92)
2554	67	*	*	69	*	58	*	54	*
2555	7,412	86.4 (56)	4.0 (42)	7,855	24.3 (124)	6,219	8.0 (146)	5,756	88.4 (71)
2556	312	77.9 (123)	4.2 (43)	323	10.2 (159)	243	14.0 (125)	114	84.2 (117)
2557	7	*	*	10	*	8	*	9	*
2558	6	*	*	7	*	7	*	7	*
2559	9	*	*	9	*	8	*	9	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 107

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2560	3	*	*	4	*	4	*	4	*
2561	2	*	*	2	*	2	*	2	*
2562	11	*	*	12	*	11	*	11	*
2563	20	*	*	17	*	11	*	6	*
2564	7	*	*	7	*	6	*	5	*
3001	1,062	76.3 (132)	8.1 (88)	1,139	34.7 (72)	995	9.0 (141)	926	80.5 (131)
3002	277	89.2 (35)	0.7 (3)	296	38.5 (40)	271	8.5 (143)	258	84.5 (113)
3004	172	93.6 (5)	2.3 (14)	192	52.1 (6)	179	19.0 (86)	170	71.2 (144)
3005	18	*	*	19	*	16	*	10	*
3007	21	*	*	25	*	22	*	21	*
3009	1	*	*	6	*	4	*	4	*
3201	426	83.1 (88)	11.7 (116)	457	38.9 (37)	387	36.7 (13)	347	87.0 (90)
3301	4,055	84.6 (70)	7.4 (83)	4,326	28.9 (105)	3,863	7.6 (147)	3,271	90.2 (55)
3304	102	84.3 (75)	10.8 (110)	104	36.5 (55)	103	13.6 (127)	82	*
3305	150	88.0 (48)	8.0 (86)	162	31.5 (93)	152	7.2 (148)	120	94.2 (9)
3306	239	84.1 (80)	8.4 (91)	254	31.1 (94)	233	5.6 (151)	187	95.2 (3)
3308	68	*	*	72	*	59	*	46	*
3309	34	*	*	36	*	33	*	30	*
3312	403	85.6 (62)	4.7 (54)	426	26.5 (116)	361	11.1 (138)	324	89.8 (58)
3550	60	*	*	64	*	59	*	57	*
3551	27	*	*	28	*	24	*	21	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 108

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3552	46	*	*	52	*	45	*	44	*
3701	993	91.6 (12)	4.3 (46)	1,083	34.3 (75)	975	20.2 (81)	826	93.7 (12)
3850	2,890	85.7 (61)	7.4 (83)	3,121	39.3 (34)	2,748	15.4 (118)	2,559	86.0 (94)
3901	1,033	82.9 (90)	6.4 (72)	1,124	35.0 (70)	966	12.4 (133)	823	78.7 (139)
3902	318	82.4 (95)	9.7 (104)	345	41.2 (24)	316	14.9 (121)	272	80.9 (128)
3903	27	*	*	28	*	26	*	12	*
3904	7	*	*	7	*	7	*	7	*
3905	11	*	*	11	*	10	*	9	*
3950	1,896	82.1 (99)	10.1 (108)	2,086	33.9 (79)	1,866	17.7 (94)	1,592	85.1 (108)
3954	11	*	*	12	*	11	*	7	*
4001	445	74.8 (141)	16.4 (139)	504	60.9 (1)	506	15.2 (119)	509	72.3 (143)
4002	302	76.2 (134)	17.9 (143)	327	57.8 (2)	320	20.0 (82)	327	83.2 (122)
4003	2	*	*	2	*	2	*	2	*
4004	4	*	*	4	*	4	*	4	*
4103	34	*	*	40	*	36	*	38	*
4106	2,315	84.8 (68)	5.8 (66)	2,536	27.1 (110)	2,210	8.1 (145)	2,013	87.3 (84)
4107	2,999	84.1 (80)	6.9 (78)	3,191	25.2 (120)	2,359	21.3 (69)	2,375	91.9 (35)
4109	963	80.8 (106)	13.5 (125)	1,030	30.9 (96)	934	17.7 (94)	750	93.6 (13)
4110	5	*	*	5	*	5	*	0	*
4112	535	44.1 (157)	0.0 (1)	556	24.8 (122)	534	93.3 (1)	500	92.8 (24)
4113	325	89.5 (31)	0.0 (1)	341	28.4 (106)	274	17.2 (99)	324	88.0 (77)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 109

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4115	3,339	88.5 (42)	4.9 (56)	3,567	25.2 (120)	3,073	16.2 (113)	2,669	91.0 (45)
4351	552	90.2 (23)	3.4 (36)	580	40.0 (29)	212	52.4 (5)	482	93.4 (17)
4352	154	69.5 (152)	18.8 (147)	166	50.6 (8)	149	12.1 (136)	141	80.9 (128)
4353	369	76.2 (134)	17.9 (143)	417	38.1 (43)	363	21.2 (72)	326	79.4 (137)
4354	722	87.8 (52)	5.8 (66)	759	38.3 (41)	556	23.6 (58)	596	90.3 (51)
4355	335	89.3 (34)	8.1 (88)	352	31.8 (89)	315	54.0 (4)	282	93.6 (13)
4356	432	82.2 (97)	11.6 (115)	463	36.9 (52)	411	29.0 (26)	383	91.4 (40)
4358	119	67.2 (156)	21.0 (152)	134	32.8 (85)	114	25.4 (46)	101	73.3 (142)
4359	447	74.9 (139)	17.9 (143)	482	39.2 (35)	406	26.1 (43)	380	90.8 (46)
4360	214	69.2 (154)	22.0 (154)	226	41.6 (23)	168	32.1 (20)	180	85.6 (100)
4361	251	86.9 (55)	8.0 (86)	266	42.1 (21)	242	26.4 (40)	218	90.8 (46)
4362	287	75.6 (136)	17.4 (141)	299	39.5 (31)	267	22.1 (63)	243	85.6 (100)
4363	833	76.7 (128)	15.0 (133)	881	40.7 (27)	674	31.5 (24)	707	84.6 (111)
4401	1,010	83.0 (89)	11.1 (112)	1,057	30.7 (98)	978	16.0 (115)	817	93.5 (15)
4501	946	77.8 (124)	14.1 (129)	972	36.1 (62)	867	27.3 (35)	738	92.8 (24)
4601	1,265	88.1 (46)	5.5 (61)	1,372	26.2 (118)	1,215	21.3 (69)	1,039	87.1 (87)
4602	23	*	*	23	*	15	*	13	*
4701	1,016	91.1 (13)	3.2 (33)	1,097	21.8 (133)	816	38.5 (12)	724	92.1 (33)
5050	1,585	90.5 (18)	3.2 (33)	1,699	19.0 (145)	1,132	18.2 (90)	1,189	93.1 (21)
5051	2,513	91.0 (14)	3.0 (28)	2,678	19.6 (141)	1,993	16.8 (107)	1,860	93.1 (21)
5052	2,729	87.9 (49)	3.4 (36)	2,914	21.2 (137)	1,972	14.2 (124)	2,100	92.7 (26)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 110

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5055	1,117	83.7 (84)	14.2 (130)	1,170	52.6 (5)	729	13.0 (130)	1,018	81.6 (125)
5101	1,683	90.0 (27)	4.2 (43)	1,809	23.8 (126)	1,344	24.6 (51)	1,316	93.5 (15)
5102	13	*	*	12	*	11	*	4	*
5202	700	88.7 (39)	6.1 (70)	786	37.4 (47)	586	36.0 (16)	626	91.4 (40)
5203	3	*	*	4	*	4	*	4	*
5204	4	*	*	4	*	3	*	4	*
5601	1,021	87.6 (53)	8.6 (94)	1,063	29.4 (103)	967	32.0 (22)	810	95.9 (1)
5801	1,612	82.3 (96)	9.2 (97)	1,748	32.3 (86)	1,597	18.3 (89)	1,382	87.3 (84)
5803	126	88.1 (46)	5.6 (63)	132	26.5 (116)	118	14.4 (123)	82	*
5804	15	*	*	17	*	15	*	10	*
5806	48	*	*	49	*	31	*	15	*
6101	32	*	*	33	*	30	*	26	*
6102	117	84.6 (70)	5.1 (58)	119	34.5 (74)	104	25.0 (48)	91	*
6103	3,665	86.2 (58)	6.5 (73)	3,938	39.1 (36)	3,446	17.2 (99)	3,150	79.8 (133)
6105	20	*	*	22	*	19	*	16	*
6106	92	*	*	102	40.2 (28)	92	*	80	*
6301	3,822	85.2 (63)	6.5 (73)	4,140	26.6 (113)	3,340	11.1 (138)	3,105	87.0 (90)
6312	1,997	88.5 (42)	4.4 (47)	2,152	21.8 (133)	1,685	15.5 (117)	1,551	87.1 (87)
6316	1,174	89.1 (36)	3.1 (30)	1,265	24.7 (123)	1,059	21.1 (74)	894	92.6 (28)
6318	12	*	*	14	*	13	*	13	*
6320	119	95.0 (2)	2.5 (16)	124	22.6 (128)	60	*	80	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 111

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6325	175	92.6 (7)	4.6 (51)	188	21.3 (136)	148	16.9 (106)	115	86.1 (93)
6330	107	78.5 (119)	15.0 (133)	118	48.3 (9)	92	*	118	81.4 (126)
6335	1,003	88.7 (39)	9.3 (100)	1,042	39.5 (31)	269	26.0 (45)	835	91.3 (42)
6340	43	*	*	50	*	32	*	43	*
6345	53	*	*	60	*	47	*	38	*
6360	581	85.0 (66)	6.9 (78)	629	23.8 (126)	381	21.0 (75)	448	85.3 (106)
7304	85	*	*	91	*	82	*	75	*
7306	2,343	78.5 (119)	10.8 (110)	2,529	33.0 (84)	2,127	15.8 (116)	1,906	86.0 (94)
7308	74	*	*	83	*	78	*	72	*
7310	19	*	*	22	*	21	*	19	*
7401	2,138	69.2 (154)	22.6 (157)	2,314	38.0 (44)	1,977	18.0 (93)	1,880	85.9 (97)
7601	579	84.3 (75)	8.8 (95)	643	38.7 (39)	504	41.9 (10)	519	87.3 (84)
7850	1,064	76.6 (129)	16.1 (138)	1,163	37.4 (47)	1,056	16.4 (109)	920	87.9 (79)
7852	21	*	*	21	*	20	*	19	*
8101	1,784	77.6 (126)	2.6 (20)	1,875	18.3 (146)	1,127	16.2 (113)	1,273	88.1 (76)
8102	382	82.5 (94)	4.5 (49)	403	35.0 (70)	251	20.3 (80)	330	81.2 (127)
8105	1,531	86.4 (56)	3.1 (30)	1,633	22.5 (129)	1,460	16.6 (108)	1,174	92.5 (29)
8106	444	76.8 (127)	5.9 (68)	475	19.6 (141)	347	6.1 (150)	326	84.7 (110)
8150	28	*	*	31	*	29	*	28	*
8151	2,936	90.4 (20)	2.8 (24)	3,224	30.0 (99)	2,919	17.0 (102)	2,508	91.1 (44)
8152	49	*	*	52	*	50	*	49	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 112

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8153	6	*	*	7	*	5	*	3	*
8155	19	*	*	19	*	18	*	16	*
8157	7	*	*	9	*	8	*	8	*
8201	2,390	91.8 (10)	1.6 (9)	2,590	18.3 (146)	1,903	34.4 (17)	1,758	89.0 (64)
8202	3,779	90.3 (21)	7.0 (81)	3,966	31.0 (95)	2,733	12.3 (134)	3,106	84.6 (111)
8203	414	92.5 (8)	1.0 (4)	444	17.3 (151)	332	16.3 (112)	291	95.2 (3)
8250	736	82.9 (90)	2.4 (15)	803	15.3 (157)	630	22.4 (62)	548	87.8 (80)
8252	153	76.5 (130)	2.6 (20)	157	16.6 (154)	128	13.3 (128)	114	88.6 (69)
8255	2,404	90.3 (21)	1.3 (7)	2,579	17.3 (151)	1,575	36.7 (13)	1,762	92.5 (29)
8257	1,980	96.1 (1)	1.1 (6)	2,079	28.3 (107)	1,880	19.6 (83)	1,621	84.3 (115)
8263	677	93.8 (4)	1.0 (4)	718	15.2 (158)	390	36.4 (15)	479	94.4 (6)
8264	630	92.4 (9)	2.7 (23)	693	19.2 (144)	577	17.0 (102)	490	94.1 (11)
8266	8	*	*	9	*	9	*	9	*
8267	790	89.0 (37)	2.9 (26)	881	19.5 (143)	767	49.4 (6)	645	88.4 (71)
8901	1,046	79.6 (112)	3.5 (39)	1,117	24.0 (125)	728	12.6 (132)	850	82.6 (123)
8902	1,379	84.2 (78)	2.2 (12)	1,479	21.8 (133)	977	17.0 (102)	1,065	85.4 (104)
8903	1,339	78.6 (118)	3.7 (40)	1,413	22.4 (130)	960	17.1 (101)	1,028	79.8 (133)
8904	2,280	82.7 (92)	2.5 (16)	2,408	18.2 (148)	1,394	17.4 (97)	1,696	87.6 (81)
8905	1,498	89.5 (31)	3.4 (36)	1,567	28.2 (108)	1,101	21.3 (69)	1,172	85.4 (104)
8908	18	*	*	18	*	15	*	18	*
8909	599	84.1 (80)	2.2 (12)	631	19.8 (140)	443	17.4 (97)	460	89.3 (59)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 113

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9016	7	*	*	7	*	7	*	7	*
9018	3	*	*	3	*	3	*	3	*
9045	1,752	73.1 (147)	13.6 (126)	1,852	33.6 (82)	394	49.2 (7)	1,480	85.3 (106)
9050	909	76.3 (132)	6.9 (78)	946	22.4 (130)	755	20.9 (76)	687	83.4 (119)
9051	1,561	69.5 (152)	22.0 (154)	1,705	51.6 (7)	1,518	14.5 (122)	1,511	70.5 (145)
9053	963	78.8 (117)	6.0 (69)	1,006	20.3 (139)	804	11.1 (138)	668	80.8 (130)
9055	29	*	*	28	*	16	*	8	*
9058	1,015	87.9 (49)	4.5 (49)	1,052	22.4 (130)	767	5.6 (151)	771	88.5 (70)
9066	997	80.0 (110)	3.3 (35)	1,127	16.4 (155)	742	12.7 (131)	752	88.0 (77)
9068	907	79.5 (114)	2.5 (16)	946	26.7 (112)	746	13.8 (126)	716	85.6 (100)
9069	2,752	79.7 (111)	2.8 (24)	2,853	16.4 (155)	1,668	8.9 (142)	1,843	89.9 (57)
9070	1,310	75.1 (138)	5.6 (63)	1,380	17.8 (149)	836	17.6 (96)	922	86.0 (94)
9072	1,697	79.6 (112)	3.7 (40)	1,786	17.4 (150)	1,427	8.2 (144)	1,261	83.3 (120)
9080	2,727	82.1 (99)	3.1 (30)	2,842	20.8 (138)	1,971	11.8 (137)	2,074	90.6 (48)
9081	12	*	*	16	*	15	*	16	*
9082	151	82.1 (99)	2.0 (10)	162	17.3 (151)	147	12.2 (135)	112	79.5 (136)
9083	121	91.7 (11)	2.5 (16)	122	1.6 (160)	88	*	11	*
9084	42	*	*	43	*	18	*	8	*
9085	45	*	*	52	*	48	*	52	*
9086	40	*	*	48	*	47	*	48	*
9087	1	*	*	1	*	1	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 114

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Michigan	153,335	84.1	7.0	164,632	29.5	129,731	19.1	124,345	87.4
----------	---------	------	-----	---------	------	---------	------	---------	------

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic Combined 3 Years

Page : 115

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	107	46.7 (141)	106	29.2 (139)	37	*	89	*	106	3.8 (108)
0102	452	49.6 (150)	466	34.8 (154)	189	24.3 (96)	410	22.2 (137)	466	4.5 (126)
0103	399	45.1 (138)	414	32.1 (149)	155	24.5 (95)	358	20.7 (133)	413	5.8 (145)
0104	138	52.2 (154)	140	35.0 (155)	60	*	123	16.3 (111)	140	3.6 (105)
0105	6	*	9	*	1	*	5	*	9	*
0250	137	48.9 (148)	144	29.2 (139)	59	*	130	18.5 (127)	144	0.7 (10)
0251	89	*	94	*	43	*	79	*	94	*
0252	106	38.7 (113)	106	28.3 (136)	31	*	88	*	106	3.8 (108)
0253	141	49.6 (150)	143	30.8 (145)	66	*	134	18.7 (128)	143	2.8 (78)
0401	457	49.2 (149)	456	29.6 (141)	180	32.8 (73)	397	17.9 (122)	456	4.2 (119)
0402	362	52.2 (154)	382	28.5 (137)	155	38.7 (50)	322	16.8 (117)	383	3.7 (107)
0403	130	46.2 (139)	144	30.6 (144)	54	*	119	26.1 (144)	144	9.0 (155)
0404	111	43.2 (133)	117	23.1 (115)	40	*	96	*	117	4.3 (124)
0501	471	42.3 (129)	495	30.3 (143)	154	25.3 (93)	399	19.3 (129)	495	5.7 (143)
0503	105	48.6 (147)	110	32.7 (152)	42	*	90	*	110	10.0 (156)
0504	461	37.3 (106)	475	20.4 (103)	147	37.4 (53)	388	11.3 (74)	475	3.4 (99)
0505	298	39.9 (121)	302	22.2 (110)	107	40.2 (44)	268	11.6 (77)	301	6.0 (147)
0506	171	46.8 (142)	184	34.2 (153)	74	*	156	26.3 (145)	184	8.2 (153)
0507	72	*	81	*	18	*	50	*	81	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 116

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0508	32	*	34	*	6	*	24	*	34	*
0509	4	*	6	*	3	*	3	*	6	*
0650	1,045	39.5 (119)	1,066	23.3 (116)	339	39.5 (47)	911	14.3 (100)	1,069	3.0 (88)
0651	566	48.4 (146)	578	31.0 (146)	235	26.0 (91)	521	17.9 (122)	578	4.5 (126)
0652	286	48.3 (145)	317	28.1 (134)	119	36.1 (58)	248	23.0 (141)	317	7.9 (151)
0653	382	41.1 (125)	388	25.3 (125)	132	26.5 (86)	341	18.2 (125)	388	4.6 (128)
0654	352	44.3 (136)	347	28.2 (135)	126	26.2 (88)	322	22.7 (139)	347	4.6 (128)
0655	330	48.2 (144)	338	31.1 (148)	136	34.6 (65)	300	20.3 (131)	338	5.9 (146)
0657	58	*	68	*	24	*	56	*	68	*
0701	371	29.1 (75)	375	19.2 (91)	93	*	317	6.3 (25)	375	1.1 (17)
0702	64	*	58	*	19	*	55	*	58	*
0703	196	50.5 (153)	193	38.9 (157)	72	*	171	14.6 (101)	194	4.1 (115)
0704	50	*	46	*	15	*	44	*	46	*
0705	129	37.2 (105)	123	26.0 (127)	40	*	113	12.4 (83)	123	3.3 (97)
0706	9	*	12	*	2	*	6	*	12	*
0850	780	39.9 (121)	802	22.1 (107)	236	38.6 (51)	643	7.8 (43)	802	2.5 (66)
0851	686	34.7 (94)	678	19.6 (95)	191	39.3 (48)	579	9.2 (53)	678	2.8 (78)
0901	1,515	38.9 (114)	1,554	24.1 (120)	531	30.3 (81)	1,371	13.7 (94)	1,554	3.8 (108)
0902	39	*	40	*	11	*	36	*	40	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic Combined 3 Years

Page : 117

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1051	1,179	35.4 (100)	1,270	21.3 (106)	339	34.8 (63)	932	9.7 (61)	1,270	3.1 (91)
1101	1,648	26.9 (68)	1,772	13.3 (62)	355	40.8 (42)	1,416	11.6 (77)	1,772	3.6 (105)
1102	732	35.5 (101)	751	20.0 (100)	199	35.7 (60)	635	15.3 (105)	751	4.1 (115)
1104	1	*	1	*	0	*	0	*	1	*
1106	70	*	69	*	15	*	57	*	69	*
1250	957	33.2 (90)	975	19.4 (93)	259	33.6 (69)	833	12.5 (85)	975	3.1 (91)
1251	709	44.3 (136)	754	29.8 (142)	253	26.5 (86)	623	16.9 (118)	754	4.9 (131)
1252	555	35.0 (98)	559	18.6 (88)	152	34.9 (62)	493	15.0 (103)	558	3.4 (99)
1253	721	39.1 (115)	737	22.1 (107)	213	36.6 (57)	636	11.8 (80)	737	3.5 (103)
1350	2,119	30.0 (78)	2,196	18.5 (86)	537	26.1 (90)	1,862	11.5 (76)	2,206	2.6 (70)
1351	120	31.7 (85)	129	16.3 (80)	28	*	94	*	129	1.6 (38)
1352	328	37.5 (107)	349	18.9 (90)	108	25.0 (94)	299	1.7 (3)	351	1.1 (17)
1401	219	43.8 (134)	229	31.0 (146)	79	*	187	13.4 (92)	229	3.9 (111)
1402	115	28.7 (74)	123	11.4 (56)	25	*	88	*	123	0.8 (11)
1403	22	*	23	*	10	*	17	*	24	*
1501	695	21.4 (41)	714	11.1 (50)	124	24.2 (97)	619	5.2 (18)	711	1.7 (42)
1503	984	25.3 (62)	1,001	12.6 (61)	216	28.7 (82)	893	7.2 (32)	1,001	1.9 (49)
1504	1,878	20.1 (33)	1,997	10.1 (42)	337	45.4 (29)	1,553	4.6 (15)	1,998	1.3 (28)
1505	372	24.5 (57)	402	11.2 (52)	78	*	281	3.9 (12)	403	2.2 (58)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 118

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1506	542	24.0 (53)	548	15.1 (73)	111	16.2 (101)	479	2.9 (8)	549	0.9 (14)
1507	543	30.9 (82)	572	15.2 (74)	148	43.9 (34)	465	10.5 (68)	574	2.6 (70)
1508	70	*	67	*	3	*	67	*	67	*
1509	386	24.1 (54)	367	13.6 (65)	80	*	356	9.6 (58)	368	2.2 (58)
1511	641	34.9 (96)	683	19.8 (98)	191	30.4 (80)	555	11.0 (69)	684	2.2 (58)
1512	316	30.1 (79)	342	19.6 (95)	90	*	265	4.2 (14)	344	1.5 (34)
1513	47	*	49	*	13	*	45	*	49	*
1514	210	16.7 (15)	220	8.2 (32)	26	*	145	1.4 (2)	218	0.0 (1)
1701	634	43.1 (131)	629	32.3 (151)	231	17.3 (100)	569	23.6 (142)	629	14.5 (157)
1950	533	42.8 (130)	566	25.1 (124)	189	35.4 (61)	469	16.6 (115)	567	5.1 (134)
1952	109	42.2 (128)	120	25.8 (126)	38	*	93	*	120	2.5 (66)
1953	297	40.1 (123)	315	22.2 (110)	97	*	250	12.4 (83)	315	3.2 (93)
1954	518	40.2 (124)	534	23.4 (117)	174	37.4 (53)	456	9.9 (63)	534	1.5 (34)
1955	86	*	91	*	30	*	75	*	91	*
1957	432	33.8 (92)	452	19.5 (94)	121	46.3 (26)	351	8.3 (47)	455	1.5 (34)
1963	29	*	33	*	8	*	21	*	34	*
1964	46	*	43	*	13	*	40	*	43	*
2150	620	43.9 (135)	608	24.2 (121)	218	34.4 (66)	559	12.5 (85)	608	2.8 (78)
2151	216	48.1 (143)	208	27.9 (133)	75	*	178	16.9 (118)	207	2.9 (83)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 119

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2251	394	39.1 (115)	378	26.5 (129)	122	25.4 (92)	342	7.3 (33)	379	1.6 (38)
2252	185	46.5 (140)	168	36.3 (156)	71	*	176	14.8 (102)	168	5.4 (139)
2554	58	*	65	*	14	*	54	*	65	*
2555	6,835	28.5 (73)	7,488	15.8 (77)	1,624	36.1 (58)	5,758	9.3 (55)	7,485	2.2 (58)
2556	259	19.3 (30)	323	14.9 (72)	28	*	115	13.0 (89)	322	3.4 (99)
2557	9	*	8	*	1	*	9	*	8	*
2558	7	*	7	*	0	*	7	*	7	*
2559	9	*	9	*	2	*	9	*	9	*
2560	4	*	3	*	1	*	4	*	3	*
2561	2	*	2	*	1	*	2	*	2	*
2562	11	*	11	*	0	*	11	*	11	*
2563	12	*	20	*	2	*	6	*	20	*
2564	6	*	7	*	1	*	5	*	7	*
3001	997	34.7 (94)	1,066	15.8 (77)	295	45.8 (28)	926	9.8 (62)	1,065	2.9 (83)
3002	272	25.0 (61)	278	11.2 (52)	64	*	258	7.4 (35)	278	1.1 (17)
3004	182	20.9 (37)	174	10.3 (45)	31	*	170	7.1 (31)	174	0.6 (6)
3005	16	*	18	*	7	*	10	*	18	*
3007	22	*	19	*	2	*	21	*	19	*
3009	4	*	1	*	0	*	4	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 120

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3201	428	35.3 (99)	431	20.4 (103)	127	40.2 (44)	349	16.3 (111)	431	3.9 (111)
3301	3,885	17.6 (21)	4,099	11.2 (52)	556	34.0 (67)	3,264	7.4 (35)	4,094	4.1 (115)
3304	103	2.9 (2)	102	0.0 (1)	3	*	81	*	102	0.0 (1)
3305	152	18.4 (24)	149	9.4 (37)	19	*	120	3.3 (9)	149	1.3 (28)
3306	233	21.5 (42)	243	10.7 (48)	39	*	187	5.3 (20)	242	2.5 (66)
3308	60	*	68	*	11	*	46	*	68	*
3309	33	*	34	*	9	*	30	*	34	*
3312	362	20.2 (34)	409	15.4 (75)	65	*	324	5.6 (22)	409	2.7 (73)
3550	59	*	60	*	28	*	57	*	60	*
3551	24	*	27	*	10	*	21	*	27	*
3552	45	*	44	*	27	*	44	*	46	*
3701	976	38.2 (108)	1,005	22.1 (107)	312	31.4 (79)	826	11.7 (79)	1,005	2.8 (78)
3850	2,845	39.2 (117)	2,916	26.2 (128)	948	28.5 (83)	2,557	13.4 (92)	2,916	2.7 (73)
3901	974	25.4 (63)	1,039	14.5 (67)	190	33.2 (70)	823	7.7 (41)	1,039	1.4 (33)
3902	316	22.2 (46)	321	14.6 (68)	59	*	272	7.4 (35)	322	1.9 (49)
3903	26	*	28	*	4	*	12	*	28	*
3904	7	*	7	*	2	*	7	*	7	*
3905	10	*	11	*	2	*	9	*	10	*
3950	1,871	26.8 (67)	1,919	15.8 (77)	393	31.6 (78)	1,591	9.6 (58)	1,919	1.5 (34)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 121

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3954	11	*	11	*	3	*	7	*	11	*
4001	510	34.9 (96)	448	18.8 (89)	152	32.9 (72)	510	12.5 (85)	449	2.9 (83)
4002	327	34.3 (93)	301	16.6 (81)	103	38.8 (49)	327	11.0 (69)	300	1.7 (42)
4003	2	*	2	*	1	*	2	*	2	*
4004	4	*	4	*	2	*	4	*	4	*
4103	38	*	34	*	7	*	38	*	34	*
4106	2,242	20.3 (35)	2,353	10.0 (40)	367	40.3 (43)	2,008	2.8 (7)	2,349	1.3 (28)
4107	2,875	21.7 (45)	3,014	11.2 (52)	463	42.5 (36)	2,377	9.3 (55)	3,011	3.4 (99)
4109	936	29.6 (76)	974	14.7 (70)	220	42.3 (37)	748	6.1 (24)	973	1.2 (24)
4110	5	*	5	*	0	*	0	*	5	*
4112	534	3.4 (4)	537	1.5 (4)	14	*	470	7.0 (29)	526	3.0 (88)
4113	326	6.1 (6)	330	3.0 (9)	21	*	324	0.3 (1)	331	0.0 (1)
4115	3,188	17.2 (19)	3,364	7.7 (31)	432	44.2 (31)	2,664	4.6 (15)	3,366	1.2 (24)
4351	536	41.2 (126)	552	24.6 (122)	192	32.3 (74)	481	16.4 (113)	552	2.5 (66)
4352	150	60.0 (156)	152	32.2 (150)	79	*	141	24.8 (143)	152	4.6 (128)
4353	363	30.6 (80)	380	19.2 (91)	94	*	326	7.4 (35)	380	1.1 (17)
4354	682	39.6 (120)	725	24.8 (123)	228	32.0 (75)	596	16.1 (110)	725	3.9 (111)
4355	323	36.8 (103)	336	22.9 (114)	103	26.2 (88)	282	11.0 (69)	336	4.2 (119)
4356	416	33.7 (91)	436	19.7 (97)	120	36.7 (56)	383	9.9 (63)	435	3.2 (93)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 122

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4358	115	17.4 (20)	123	11.4 (56)	18	*	101	5.0 (17)	124	0.8 (11)
4359	435	28.3 (71)	458	13.5 (64)	104	44.2 (31)	380	7.9 (45)	458	0.9 (14)
4360	201	32.3 (87)	215	18.1 (84)	58	*	180	16.7 (116)	215	4.2 (119)
4361	246	43.1 (131)	254	27.6 (131)	88	*	218	15.6 (107)	254	1.2 (24)
4362	271	49.8 (152)	287	28.9 (138)	121	39.7 (46)	243	22.6 (138)	287	5.2 (137)
4363	794	37.0 (104)	836	26.6 (130)	241	19.9 (99)	707	14.0 (97)	836	4.3 (124)
4401	978	31.4 (83)	1,011	14.2 (66)	232	44.0 (33)	817	3.5 (11)	1,010	0.6 (6)
4501	873	38.4 (110)	949	22.4 (112)	279	34.8 (63)	738	14.1 (98)	950	2.7 (73)
4601	1,224	38.2 (108)	1,277	17.4 (83)	355	47.3 (23)	1,035	9.6 (58)	1,279	3.9 (111)
4602	15	*	23	*	7	*	13	*	23	*
4701	981	32.5 (88)	1,013	18.5 (86)	230	41.7 (39)	727	8.0 (46)	1,018	2.7 (73)
5050	1,485	25.6 (65)	1,597	12.1 (59)	279	47.0 (24)	1,190	9.2 (53)	1,600	1.8 (47)
5051	2,413	22.5 (48)	2,533	10.1 (42)	421	48.5 (22)	1,862	7.3 (33)	2,531	2.0 (53)
5052	2,601	21.3 (40)	2,753	11.4 (56)	418	41.1 (40)	2,103	8.6 (50)	2,754	1.7 (42)
5055	1,127	2.9 (2)	1,146	0.8 (2)	28	*	1,039	4.1 (13)	1,145	1.1 (17)
5101	1,617	23.4 (51)	1,700	10.4 (46)	306	46.7 (25)	1,320	5.2 (18)	1,703	1.9 (49)
5102	11	*	13	*	2	*	4	*	13	*
5202	722	41.8 (127)	710	27.7 (132)	255	33.7 (68)	630	10.0 (65)	709	2.1 (56)
5203	4	*	3	*	3	*	4	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 123

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
5204	4	*	4	*	0	*	4	*	4	*
5601	968	29.8 (77)	1,024	19.8 (98)	237	28.3 (84)	809	8.5 (48)	1,023	2.2 (58)
5801	1,601	38.4 (110)	1,619	23.7 (119)	488	33.2 (70)	1,382	13.0 (89)	1,619	3.0 (88)
5803	118	28.0 (70)	128	14.8 (71)	22	*	82	*	127	1.6 (38)
5804	15	*	15	*	3	*	10	*	15	*
5806	32	*	48	*	5	*	15	*	48	*
6101	31	*	32	*	11	*	26	*	32	*
6102	104	32.7 (89)	118	16.9 (82)	33	*	91	*	117	5.1 (134)
6103	3,554	31.5 (84)	3,714	20.9 (105)	962	27.1 (85)	3,153	13.1 (91)	3,728	5.3 (138)
6105	20	*	21	*	1	*	16	*	21	*
6106	94	*	95	*	28	*	79	*	95	*
6301	3,714	17.8 (22)	3,868	9.6 (38)	516	38.2 (52)	3,108	5.9 (23)	3,868	1.3 (28)
6312	1,972	6.0 (5)	2,048	2.2 (7)	83	*	1,552	2.3 (5)	2,045	0.6 (6)
6316	1,196	16.8 (17)	1,182	7.5 (29)	134	43.3 (35)	893	3.4 (10)	1,182	0.8 (11)
6318	14	*	12	*	4	*	13	*	12	*
6320	118	6.8 (8)	118	2.5 (8)	7	*	80	*	118	0.0 (1)
6325	169	24.3 (55)	177	18.1 (84)	26	*	115	13.9 (96)	177	1.1 (17)
6330	118	22.9 (50)	108	15.7 (76)	25	*	118	8.5 (48)	108	1.9 (49)
6335	1,001	6.8 (8)	1,008	3.8 (11)	48	*	833	1.9 (4)	1,008	0.9 (14)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 124

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
6340	47	*	43	*	6	*	43	*	*	
6345	57	*	54	*	12	*	38	*	54	*
6360	540	18.1 (23)	586	9.6 (38)	78	*	448	11.8 (80)	586	2.2 (58)
7304	84	*	86	*	35	*	75	*	86	*
7306	2,202	36.0 (102)	2,388	13.3 (62)	665	52.0 (17)	1,906	12.7 (88)	2,390	2.1 (56)
7308	78	*	75	*	19	*	72	*	75	*
7310	22	*	19	*	6	*	19	*	19	*
7401	2,099	39.2 (117)	2,150	23.6 (118)	684	31.9 (77)	1,878	17.8 (121)	2,146	5.4 (139)
7601	580	31.7 (85)	591	20.1 (101)	163	22.1 (98)	529	11.2 (73)	593	2.7 (73)
7850	1,060	38.5 (112)	1,081	22.8 (113)	353	32.0 (75)	919	16.4 (113)	1,081	3.3 (97)
7852	20	*	21	*	7	*	19	*	21	*
8101	1,557	21.0 (39)	1,793	8.2 (32)	256	56.3 (14)	1,266	13.7 (94)	1,794	5.4 (139)
8102	362	18.8 (28)	390	8.5 (34)	58	*	332	5.4 (21)	388	2.8 (78)
8105	1,479	15.6 (13)	1,531	6.3 (20)	157	51.0 (19)	1,173	10.1 (67)	1,531	2.3 (65)
8106	397	20.7 (36)	456	10.1 (42)	69	*	325	7.7 (41)	456	1.8 (47)
8150	29	*	28	*	2	*	28	*	28	*
8151	2,935	23.8 (52)	2,949	11.1 (50)	530	45.3 (30)	2,506	7.5 (39)	2,947	1.7 (42)
8152	50	*	48	*	11	*	49	*	49	*
8153	5	*	6	*	0	*	3	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 125

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8155	18	*	19	*	2	*	16	*	19	*
8157	8	*	7	*	1	*	8	*	7	*
8201	2,306	18.6 (27)	2,399	11.0 (49)	307	37.1 (55)	1,736	10.0 (65)	2,402	2.0 (53)
8202	3,656	6.1 (6)	3,837	1.8 (6)	191	63.9 (4)	3,120	6.3 (25)	3,834	2.2 (58)
8203	396	30.6 (80)	413	20.1 (101)	89	*	290	11.4 (75)	414	1.7 (42)
8250	689	26.9 (68)	741	10.0 (40)	132	58.3 (11)	548	11.1 (72)	741	2.6 (70)
8252	136	24.3 (55)	154	3.2 (10)	29	*	114	15.8 (109)	154	3.2 (93)
8255	2,241	28.3 (71)	2,424	14.6 (68)	477	41.9 (38)	1,760	9.0 (52)	2,424	1.6 (38)
8257	1,948	2.1 (1)	1,994	1.3 (3)	32	*	1,628	2.3 (5)	1,994	0.6 (6)
8263	629	24.8 (60)	677	10.6 (47)	123	50.4 (20)	479	6.9 (28)	678	1.2 (24)
8264	628	24.5 (57)	634	8.8 (36)	118	53.4 (16)	491	9.4 (57)	635	1.3 (28)
8266	9	*	9	*	1	*	9	*	9	*
8267	796	25.6 (65)	796	12.1 (59)	141	41.1 (40)	637	7.8 (43)	796	1.1 (17)
8901	940	24.6 (59)	1,056	7.5 (29)	203	58.6 (9)	848	18.3 (126)	1,056	5.0 (132)
8902	1,274	16.7 (15)	1,391	4.5 (14)	178	65.7 (2)	1,067	14.2 (99)	1,389	6.0 (147)
8903	1,205	22.7 (49)	1,353	8.5 (34)	233	54.9 (15)	1,030	20.7 (133)	1,351	7.4 (150)
8904	2,064	21.6 (44)	2,301	6.7 (24)	353	69.1 (1)	1,693	17.4 (120)	2,302	7.0 (149)
8905	1,414	15.8 (14)	1,507	7.0 (27)	174	51.7 (18)	1,170	6.3 (25)	1,506	2.9 (83)
8908	18	*	18	*	6	*	18	*	18	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 126

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8909	563	19.9 (31)	605	4.5 (14)	95	*	460	15.4 (106)	603	5.6 (142)
9016	7	*	7	*	2	*	7	*	7	*
9018	3	*	3	*	1	*	3	*	3	*
9045	1,681	11.7 (11)	1,759	4.4 (13)	162	50.0 (21)	1,484	8.6 (50)	1,759	2.9 (83)
9050	791	25.5 (64)	928	5.2 (17)	190	65.3 (3)	688	17.9 (122)	927	3.5 (103)
9051	1,618	10.2 (10)	1,631	3.9 (12)	137	46.0 (27)	1,514	7.0 (29)	1,631	2.0 (53)
9053	850	18.5 (26)	963	5.6 (19)	135	60.7 (6)	666	15.6 (107)	964	5.0 (132)
9055	17	*	29	*	2	*	8	*	29	*
9058	901	12.2 (12)	1,014	4.9 (16)	84	*	768	7.6 (40)	1,014	3.2 (93)
9066	909	20.9 (37)	1,078	6.5 (22)	156	60.3 (8)	754	19.5 (130)	1,077	4.1 (115)
9068	805	22.4 (47)	909	6.8 (25)	153	61.4 (5)	716	15.2 (104)	909	4.2 (119)
9069	2,343	16.9 (18)	2,762	6.5 (22)	315	60.6 (7)	1,841	20.6 (132)	2,761	8.1 (152)
9070	1,160	19.1 (29)	1,325	6.8 (25)	173	58.4 (10)	919	22.7 (139)	1,319	5.7 (143)
9072	1,475	21.5 (42)	1,721	6.4 (21)	269	57.2 (12)	1,261	21.8 (136)	1,720	5.1 (134)
9080	2,442	19.9 (31)	2,738	7.0 (27)	407	56.8 (13)	2,074	11.8 (80)	2,740	4.2 (119)
9081	16	*	12	*	0	*	16	*	12	*
9082	152	18.4 (24)	151	5.3 (18)	23	*	112	21.4 (135)	151	8.6 (154)
9083	92	*	120	1.7 (5)	0	*	10	*	118	0.0 (1)
9084	34	*	42	*	1	*	8	*	42	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 127

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
9085	52	*	47	*	11	*	52	*	47	*
9086	48	*	40	*	1	*	48	*	40	*
9087	1	*	1	*	0	*	0	*	1	*
Michigan	146,996	26.0	155,033	13.8	31,092	38.9	124,354	10.6	155,026	3.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 128

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	99	*	*	101	10.9 (62)	89	*	99	*
0102	441	7.0 (31)	7.9 (75)	451	8.6 (19)	403	4.5 (95)	442	64.9 (113)
0103	393	7.4 (40)	9.4 (120)	396	7.3 (8)	359	3.6 (66)	393	68.2 (102)
0104	133	5.3 (8)	6.8 (53)	137	6.6 (3)	125	4.8 (104)	133	74.4 (78)
0105	8	*	*	7	*	7	*	8	*
0250	138	6.5 (18)	14.5 (154)	140	10.0 (41)	122	0.8 (1)	138	75.4 (74)
0251	90	*	*	93	*	85	*	90	*
0252	102	3.9 (3)	12.7 (151)	104	5.8 (1)	96	*	102	81.4 (29)
0253	132	12.9 (133)	9.8 (125)	129	10.9 (62)	113	8.0 (145)	132	69.7 (96)
0401	425	10.1 (100)	12.0 (147)	434	9.2 (25)	383	4.7 (100)	423	71.9 (90)
0402	356	9.3 (83)	7.9 (75)	364	10.2 (46)	315	4.4 (93)	356	78.7 (54)
0403	137	11.7 (124)	6.6 (50)	137	12.4 (105)	118	5.9 (126)	137	73.7 (82)
0404	115	2.6 (1)	12.2 (150)	115	9.6 (30)	103	1.9 (7)	115	69.6 (97)
0501	382	8.6 (70)	9.2 (118)	478	8.8 (20)	344	3.5 (59)	451	78.0 (56)
0503	100	6.0 (14)	12.0 (147)	105	7.6 (9)	92	*	102	83.3 (17)
0504	448	7.4 (40)	8.5 (101)	463	9.5 (28)	404	3.2 (48)	455	83.1 (20)
0505	263	8.4 (69)	8.0 (82)	293	8.2 (14)	242	2.9 (29)	285	86.3 (6)
0506	163	8.0 (56)	9.8 (125)	171	9.4 (26)	144	5.6 (123)	169	80.5 (36)
0507	65	*	*	74	*	57	*	75	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 129

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	31	*	*	32	*	28	*	32	*
0509	6	*	*	6	*	6	*	6	*
0650	987	8.3 (66)	7.9 (75)	1,034	9.7 (33)	896	4.8 (104)	988	77.7 (62)
0651	546	5.9 (13)	7.9 (75)	558	9.5 (28)	491	2.9 (29)	546	72.7 (87)
0652	288	11.5 (120)	8.0 (82)	299	11.7 (83)	251	5.2 (117)	289	70.9 (91)
0653	357	8.1 (60)	9.0 (109)	378	11.4 (76)	317	3.5 (59)	358	84.4 (11)
0654	331	6.6 (21)	9.4 (120)	339	9.4 (26)	301	4.0 (77)	333	80.2 (38)
0655	323	6.5 (18)	7.4 (62)	323	9.9 (36)	287	3.1 (42)	324	74.4 (78)
0657	60	*	*	61	*	54	*	60	*
0701	339	4.4 (5)	12.1 (149)	351	7.1 (6)	303	2.6 (18)	341	87.4 (5)
0702	54	*	*	57	*	49	*	54	*
0703	190	7.4 (40)	8.4 (97)	187	11.8 (85)	163	2.5 (17)	190	72.6 (89)
0704	44	*	*	46	*	44	*	44	*
0705	111	7.2 (34)	9.9 (128)	117	7.7 (10)	102	4.9 (109)	110	83.6 (13)
0706	10	*	*	10	*	7	*	10	*
0850	760	7.9 (52)	8.7 (104)	798	10.5 (56)	683	2.8 (24)	779	83.3 (17)
0851	648	8.3 (66)	10.3 (133)	665	9.9 (36)	580	4.1 (85)	655	80.2 (38)
0901	1,468	7.5 (45)	8.6 (102)	1,503	9.9 (36)	1,325	3.4 (57)	1,469	66.8 (106)
0902	39	*	*	40	*	39	*	39	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 130

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1051	1,203	7.7 (47)	11.6 (145)	1,234	8.8 (20)	1,089	2.8 (24)	1,203	85.0 (9)
1101	1,679	9.2 (81)	6.4 (44)	1,737	10.8 (60)	1,490	4.8 (104)	1,684	69.8 (95)
1102	693	6.3 (15)	7.5 (67)	722	8.2 (14)	633	3.5 (59)	693	74.6 (77)
1104	1	*	*	1	*	1	*	1	*
1106	67	*	*	68	*	61	*	67	*
1250	921	6.5 (18)	8.0 (82)	943	10.8 (60)	813	2.1 (14)	930	79.1 (52)
1251	693	5.6 (11)	8.8 (106)	729	9.7 (33)	624	2.1 (14)	694	79.4 (48)
1252	518	7.7 (47)	6.8 (53)	529	11.9 (89)	450	2.9 (29)	520	83.5 (14)
1253	699	8.7 (73)	7.9 (75)	716	10.1 (43)	624	2.6 (18)	705	84.0 (12)
1350	2,092	9.4 (85)	7.4 (62)	2,136	10.9 (62)	1,843	3.8 (73)	2,096	74.0 (81)
1351	123	5.7 (12)	8.1 (85)	124	8.1 (13)	113	3.5 (59)	123	82.9 (22)
1352	339	10.0 (98)	8.6 (102)	345	11.9 (89)	300	5.0 (111)	340	65.0 (111)
1401	218	7.3 (37)	10.6 (136)	220	10.9 (62)	193	1.6 (5)	218	85.3 (8)
1402	114	5.3 (8)	7.9 (75)	119	10.1 (43)	101	1.0 (2)	114	88.6 (3)
1403	23	*	*	22	*	22	*	23	*
1501	663	6.8 (28)	10.1 (131)	681	10.3 (49)	583	2.7 (23)	673	64.6 (114)
1503	929	7.4 (40)	8.2 (89)	948	12.1 (94)	804	2.9 (29)	931	77.9 (58)
1504	1,892	7.7 (47)	8.2 (89)	1,952	11.1 (68)	1,677	2.8 (24)	1,916	77.2 (65)
1505	381	10.8 (112)	12.9 (153)	395	10.4 (52)	341	5.3 (118)	386	82.1 (27)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 131

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1506	518	6.6 (21)	10.8 (137)	529	10.2 (46)	461	3.3 (53)	520	73.3 (85)
1507	524	9.4 (85)	12.8 (152)	549	12.0 (91)	457	2.6 (18)	531	79.5 (46)
1508	65	*	*	64	*	55	*	65	*
1509	344	6.4 (17)	9.9 (128)	349	11.2 (71)	302	3.0 (36)	350	70.9 (91)
1511	652	9.2 (81)	9.5 (122)	659	12.1 (94)	566	2.8 (24)	656	80.2 (38)
1512	327	10.1 (100)	9.8 (125)	335	15.2 (142)	278	4.0 (77)	328	82.3 (26)
1513	44	*	*	47	*	41	*	46	*
1514	211	11.4 (119)	9.0 (109)	216	13.9 (125)	181	3.9 (75)	215	91.2 (1)
1701	591	8.1 (60)	11.2 (141)	589	11.2 (71)	505	2.6 (18)	593	68.6 (100)
1950	539	8.0 (56)	6.5 (47)	548	11.5 (81)	473	3.2 (48)	540	83.5 (14)
1952	114	4.4 (5)	9.6 (124)	113	8.8 (20)	102	2.0 (12)	114	83.3 (17)
1953	293	6.8 (28)	8.2 (89)	299	10.4 (52)	260	1.9 (7)	293	84.6 (10)
1954	507	7.3 (37)	9.9 (128)	516	10.9 (62)	446	3.1 (42)	508	82.5 (24)
1955	84	*	*	86	*	77	*	84	*
1957	437	8.9 (77)	10.8 (137)	437	10.3 (49)	385	3.6 (66)	438	85.4 (7)
1963	31	*	*	32	*	26	*	31	*
1964	39	*	*	40	*	37	*	39	*
2150	568	7.6 (46)	8.8 (106)	580	10.3 (49)	508	3.0 (36)	568	79.2 (49)
2151	201	10.0 (98)	9.0 (109)	200	12.5 (109)	173	4.0 (77)	201	59.7 (123)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 132

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2251	363	7.7 (47)	8.3 (94)	370	8.4 (18)	329	3.3 (53)	363	75.8 (72)
2252	150	9.3 (83)	9.3 (119)	158	11.4 (76)	132	5.3 (118)	150	80.7 (34)
2554	63	*	*	65	*	57	*	63	*
2555	7,229	11.9 (125)	5.6 (31)	7,309	13.6 (120)	6,177	5.7 (124)	7,243	57.3 (132)
2556	312	16.3 (151)	2.6 (1)	313	19.2 (152)	244	9.4 (149)	315	53.3 (137)
2557	7	*	*	8	*	6	*	8	*
2558	6	*	*	7	*	5	*	7	*
2559	9	*	*	9	*	7	*	9	*
2560	3	*	*	3	*	3	*	3	*
2561	2	*	*	2	*	1	*	2	*
2562	11	*	*	11	*	9	*	11	*
2563	19	*	*	17	*	15	*	19	*
2564	7	*	*	7	*	7	*	7	*
3001	1,000	13.1 (137)	4.1 (19)	1,020	14.2 (130)	857	4.8 (104)	1,000	59.2 (124)
3002	269	11.2 (117)	4.5 (21)	274	12.0 (91)	238	5.0 (111)	269	50.6 (145)
3004	164	7.3 (37)	9.1 (116)	164	10.4 (52)	147	4.1 (85)	164	80.5 (36)
3005	18	*	*	18	*	16	*	18	*
3007	18	*	*	17	*	14	*	18	*
3009	1	*	*	1	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 133

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3201	404	8.7 (73)	11.4 (143)	415	11.3 (75)	359	3.1 (42)	405	77.5 (64)
3301	3,854	10.1 (100)	6.4 (44)	3,995	12.4 (105)	3,370	4.6 (97)	3,882	78.7 (54)
3304	95	*	*	98	*	86	*	98	*
3305	137	8.0 (56)	9.5 (122)	146	8.2 (14)	127	4.7 (100)	139	87.8 (4)
3306	223	6.7 (25)	11.7 (146)	232	7.8 (11)	205	2.0 (12)	223	82.5 (24)
3308	63	*	*	64	*	59	*	63	*
3309	34	*	*	33	*	29	*	34	*
3312	388	11.6 (122)	7.7 (71)	403	13.6 (120)	337	6.2 (130)	391	67.3 (105)
3550	56	*	*	59	*	51	*	55	*
3551	24	*	*	27	*	23	*	24	*
3552	45	*	*	46	*	37	*	45	*
3701	965	9.0 (80)	6.8 (53)	980	12.3 (103)	843	2.8 (24)	967	81.9 (28)
3850	2,759	9.5 (89)	7.4 (62)	2,822	12.5 (109)	2,405	3.6 (66)	2,759	73.4 (84)
3901	966	12.3 (128)	5.9 (36)	1,003	13.7 (123)	830	5.1 (114)	981	69.0 (99)
3902	301	8.6 (70)	11.3 (142)	304	11.8 (85)	262	5.0 (111)	302	68.2 (102)
3903	25	*	*	27	*	24	*	25	*
3904	6	*	*	6	*	5	*	6	*
3905	11	*	*	11	*	9	*	11	*
3950	1,799	9.9 (95)	9.1 (116)	1,856	12.4 (105)	1,566	3.1 (42)	1,813	75.3 (75)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 134

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3954	9	*	*	10	*	9	*	9	*
4001	424	10.1 (100)	5.2 (26)	431	13.5 (117)	362	4.7 (100)	425	72.7 (87)
4002	288	6.6 (21)	8.3 (94)	297	11.4 (76)	255	1.6 (5)	289	68.5 (101)
4003	2	*	*	2	*	2	*	2	*
4004	3	*	*	4	*	3	*	3	*
4103	32	*	*	32	*	24	*	32	*
4106	2,212	12.7 (131)	4.7 (22)	2,286	13.9 (125)	1,895	6.0 (128)	2,231	76.5 (71)
4107	2,845	12.0 (127)	7.7 (71)	2,942	14.8 (135)	2,417	4.8 (104)	2,851	77.7 (62)
4109	928	9.4 (85)	8.1 (85)	943	13.9 (125)	796	2.6 (18)	932	79.1 (52)
4110	5	*	*	5	*	0	*	5	*
4112	516	5.4 (10)	6.4 (44)	512	10.4 (52)	455	1.5 (4)	515	26.8 (155)
4113	316	8.2 (64)	7.9 (75)	327	12.5 (109)	275	4.0 (77)	317	81.1 (31)
4115	3,198	10.7 (109)	7.1 (58)	3,276	13.5 (117)	2,749	3.6 (66)	3,207	80.0 (43)
4351	516	7.4 (40)	8.7 (104)	534	12.4 (105)	453	2.9 (29)	516	77.1 (66)
4352	142	9.9 (95)	4.9 (25)	145	19.3 (153)	113	6.2 (130)	142	76.8 (70)
4353	365	7.7 (47)	6.6 (50)	366	7.1 (6)	335	4.5 (95)	364	76.9 (68)
4354	691	7.2 (34)	7.7 (71)	696	11.5 (81)	605	3.0 (36)	692	74.4 (78)
4355	313	9.9 (95)	10.5 (135)	322	11.2 (71)	273	4.0 (77)	312	81.1 (31)
4356	417	9.6 (91)	7.2 (59)	427	13.8 (124)	357	3.9 (75)	417	77.0 (67)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 135

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4358	117	3.4 (2)	6.0 (40)	115	6.1 (2)	107	1.9 (7)	118	78.0 (56)
4359	428	6.3 (15)	8.2 (89)	433	11.1 (68)	379	2.9 (29)	428	76.9 (68)
4360	183	4.4 (5)	10.9 (139)	201	7.0 (5)	169	3.0 (36)	183	79.8 (45)
4361	227	7.9 (52)	11.5 (144)	245	10.6 (58)	201	3.0 (36)	227	81.1 (31)
4362	266	10.2 (104)	7.5 (67)	272	11.0 (67)	234	5.1 (114)	267	89.5 (2)
4363	781	7.0 (31)	8.1 (85)	792	10.2 (46)	682	3.4 (57)	782	80.2 (38)
4401	965	7.9 (52)	8.4 (97)	973	10.5 (56)	850	3.2 (48)	966	79.5 (46)
4501	900	8.3 (66)	10.4 (134)	914	11.4 (76)	786	3.1 (42)	899	75.6 (73)
4601	1,204	6.6 (21)	8.2 (89)	1,246	10.0 (41)	1,072	3.0 (36)	1,204	73.1 (86)
4602	21	*	*	22	*	21	*	21	*
4701	991	9.6 (91)	9.0 (109)	1,003	12.0 (91)	869	3.6 (66)	991	79.2 (49)
5050	1,566	11.1 (116)	5.6 (31)	1,557	14.8 (135)	1,310	4.2 (88)	1,567	60.9 (120)
5051	2,457	10.7 (109)	5.9 (36)	2,472	12.6 (113)	2,112	4.0 (77)	2,462	65.6 (108)
5052	2,679	10.8 (112)	4.7 (22)	2,701	13.4 (116)	2,292	4.9 (109)	2,685	61.5 (119)
5055	949	8.1 (60)	3.6 (14)	1,073	8.3 (17)	863	4.1 (85)	1,054	43.2 (153)
5101	1,639	8.0 (56)	6.8 (53)	1,667	11.7 (83)	1,434	3.3 (53)	1,646	70.7 (93)
5102	13	*	*	12	*	11	*	12	*
5202	686	10.2 (104)	9.0 (109)	704	11.8 (85)	604	5.5 (122)	689	83.5 (14)
5203	3	*	*	3	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 136

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5204	4	*	*	4	*	3	*	4	*
5601	974	6.7 (25)	10.2 (132)	999	9.7 (33)	875	2.3 (16)	975	77.8 (59)
5801	1,538	8.7 (73)	6.3 (42)	1,578	11.1 (68)	1,353	3.5 (59)	1,523	65.2 (110)
5803	118	6.8 (28)	11.0 (140)	119	10.1 (43)	104	1.9 (7)	117	58.1 (129)
5804	14	*	*	14	*	12	*	14	*
5806	47	*	*	47	*	40	*	47	*
6101	32	*	*	30	*	27	*	32	*
6102	116	8.6 (70)	7.8 (74)	113	8.8 (20)	103	2.9 (29)	116	82.8 (23)
6103	3,531	10.7 (109)	5.2 (26)	3,580	14.8 (135)	2,931	4.3 (89)	3,536	65.3 (109)
6105	21	*	*	21	*	18	*	21	*
6106	95	*	*	95	*	76	*	95	*
6301	3,701	12.3 (128)	5.9 (36)	3,792	13.3 (115)	3,195	5.3 (118)	3,722	66.6 (107)
6312	1,969	11.2 (117)	5.8 (35)	2,003	12.3 (103)	1,713	4.0 (77)	1,981	77.8 (59)
6316	1,142	10.6 (107)	7.5 (67)	1,158	12.2 (101)	994	3.5 (59)	1,147	79.2 (49)
6318	10	*	*	12	*	9	*	11	*
6320	109	11.9 (125)	8.3 (94)	113	8.0 (12)	99	*	111	80.2 (38)
6325	166	4.2 (4)	8.4 (97)	173	6.9 (4)	155	1.9 (7)	169	79.9 (44)
6330	99	*	*	105	21.0 (156)	78	*	101	51.5 (142)
6335	958	8.9 (77)	5.9 (36)	970	9.6 (30)	858	5.1 (114)	968	69.5 (98)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 137

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6340	39	*	*	41	*	32	*	40	*
6345	52	*	*	53	*	46	*	52	*
6360	557	12.9 (133)	5.2 (26)	569	15.1 (141)	466	6.7 (134)	565	63.0 (116)
7304	83	*	*	83	*	77	*	83	*
7306	2,314	11.5 (120)	5.7 (34)	2,337	12.2 (101)	2,018	4.6 (97)	2,316	61.8 (117)
7308	75	*	*	74	*	68	*	75	*
7310	18	*	*	19	*	16	*	18	*
7401	2,046	7.2 (34)	8.1 (85)	2,087	10.7 (59)	1,819	3.2 (48)	2,048	70.1 (94)
7601	574	7.1 (33)	7.3 (60)	585	12.1 (94)	499	1.0 (2)	574	58.5 (127)
7850	1,033	7.9 (52)	7.4 (62)	1,047	9.6 (30)	930	3.3 (53)	1,033	74.9 (76)
7852	21	*	*	21	*	20	*	21	*
8101	1,632	16.9 (153)	3.2 (9)	1,746	20.0 (154)	1,281	6.9 (137)	1,700	46.2 (150)
8102	356	9.6 (91)	9.0 (109)	363	17.6 (148)	275	4.4 (93)	360	47.8 (149)
8105	1,477	10.6 (107)	5.6 (31)	1,501	12.1 (94)	1,284	4.0 (77)	1,481	63.3 (115)
8106	416	14.7 (143)	5.5 (30)	441	14.3 (133)	347	7.5 (143)	425	60.2 (122)
8150	24	*	*	24	*	23	*	24	*
8151	2,810	10.8 (112)	5.4 (29)	2,866	13.6 (120)	2,427	4.7 (100)	2,746	80.6 (35)
8152	49	*	*	47	*	42	*	48	*
8153	6	*	*	6	*	4	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 138

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8155	18	*	*	18	*	17	*	18	*
8157	6	*	*	6	*	6	*	6	*
8201	2,316	10.9 (115)	6.5 (47)	2,357	12.1 (94)	2,020	4.6 (97)	2,326	61.8 (117)
8202	3,498	8.1 (60)	6.2 (41)	3,688	8.9 (24)	3,160	4.3 (89)	3,663	46.2 (150)
8203	401	10.5 (106)	7.0 (57)	408	11.8 (85)	354	3.1 (42)	401	83.0 (21)
8250	713	11.6 (122)	4.8 (24)	728	12.5 (109)	618	6.1 (129)	716	68.2 (102)
8252	152	17.1 (154)	6.6 (50)	151	14.6 (134)	128	9.4 (149)	152	59.2 (124)
8255	2,336	9.6 (91)	7.4 (62)	2,373	11.4 (76)	2,047	4.3 (89)	2,343	59.0 (126)
8257	1,773	12.5 (130)	3.7 (16)	1,927	12.1 (94)	1,529	6.9 (137)	1,855	77.8 (59)
8263	659	8.8 (76)	9.0 (109)	670	9.9 (36)	592	3.2 (48)	658	58.4 (128)
8264	595	8.2 (64)	8.4 (97)	616	9.9 (36)	529	3.6 (66)	598	55.5 (134)
8266	7	*	*	9	*	6	*	9	*
8267	774	8.9 (77)	7.6 (70)	788	11.2 (71)	689	3.5 (59)	775	55.5 (134)
8901	997	14.7 (143)	3.0 (6)	1,027	16.7 (147)	816	6.9 (137)	1,013	50.6 (145)
8902	1,248	14.0 (139)	2.8 (3)	1,359	16.6 (146)	1,036	5.3 (118)	1,304	57.4 (131)
8903	1,244	16.2 (150)	2.7 (2)	1,303	17.7 (149)	1,004	8.3 (147)	1,271	48.4 (148)
8904	2,082	14.0 (139)	3.0 (6)	2,221	17.7 (149)	1,683	6.4 (133)	2,168	54.8 (136)
8905	1,326	9.4 (85)	7.3 (60)	1,438	12.6 (113)	1,135	4.3 (89)	1,414	60.5 (121)
8908	16	*	*	18	*	13	*	16	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 139

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8909	552	16.5 (152)	2.9 (5)	587	17.9 (151)	442	7.0 (140)	565	57.5 (130)
9016	7	*	*	7	*	6	*	7	*
9018	3	*	*	3	*	3	*	3	*
9045	1,643	9.5 (89)	6.5 (47)	1,695	13.5 (117)	1,403	3.8 (73)	1,661	52.8 (138)
9050	864	13.0 (136)	3.8 (17)	888	15.3 (144)	722	6.8 (135)	876	51.8 (141)
9051	1,468	6.7 (25)	8.8 (106)	1,499	12.1 (94)	1,261	3.6 (66)	1,490	73.6 (83)
9053	909	12.9 (133)	4.1 (19)	944	14.1 (129)	781	6.3 (132)	911	52.0 (140)
9055	28	*	*	28	*	25	*	28	*
9058	945	14.8 (145)	3.5 (12)	991	13.9 (125)	811	8.0 (145)	971	55.8 (133)
9066	1,009	15.2 (146)	3.0 (6)	1,066	15.0 (139)	853	7.7 (144)	1,015	51.3 (143)
9068	866	14.4 (142)	3.9 (18)	885	16.2 (145)	720	7.4 (142)	867	45.6 (152)
9069	2,613	14.1 (141)	3.4 (10)	2,698	15.0 (139)	2,207	5.9 (126)	2,615	36.7 (154)
9070	1,276	13.5 (138)	3.4 (10)	1,306	14.2 (130)	1,079	6.8 (135)	1,279	51.1 (144)
9072	1,617	15.3 (147)	3.5 (12)	1,688	15.2 (142)	1,362	7.0 (140)	1,619	49.4 (147)
9080	2,632	12.7 (131)	3.6 (14)	2,672	14.8 (135)	2,216	5.8 (125)	2,644	52.4 (139)
9081	11	*	*	11	*	10	*	11	*
9082	144	15.3 (147)	2.8 (3)	148	14.2 (130)	120	9.2 (148)	143	65.0 (111)
9083	112	16.1 (149)	6.3 (42)	120	20.0 (154)	90	*	113	81.4 (29)
9084	37	*	*	42	*	32	*	37	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/26/2020

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Page : 140

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9085	47	*	*	46	*	39	*	47	*
9086	40	*	*	40	*	36	*	40	*
9087	1	*	*	1	*	1	*	1	*
Michigan	146,255	10.3	6.6	150,442	12.5	126,827	4.4	147,505	57.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.