

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 1

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
0101	40	*	*	*	*	*	*	*
0102	151	91.4	0.7	0.7	0.7	0.0	6.6	0.0
0103	121	94.2	0.0	0.8	0.0	0.0	4.1	0.8
0104	39	*	*	*	*	*	*	*
0105	11	*	*	*	*	*	*	*
0250	39	*	*	*	*	*	*	*
0251	29	*	*	*	*	*	*	*
0252	30	*	*	*	*	*	*	*
0253	46	*	*	*	*	*	*	*
0401	153	96.1	0.7	1.3	0.0	0.0	2.0	0.0
0402	118	87.3	0.8	4.2	0.8	0.0	6.8	0.0
0403	53	*	*	*	*	*	*	*
0404	42	*	*	*	*	*	*	*
0501	159	97.5	0.0	1.3	0.0	0.6	0.6	0.0
0503	34	*	*	*	*	*	*	*
0504	135	88.1	2.2	2.2	1.5	0.0	5.9	0.0
0505	89	*	*	*	*	*	*	*
0506	48	*	*	*	*	*	*	*
0507	20	*	*	*	*	*	*	*
0508	10	*	*	*	*	*	*	*
0650	343	79.6	4.4	4.1	5.0	0.9	5.8	0.3
0651	182	95.6	0.0	2.2	0.0	0.0	2.2	0.0
0652	96	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 2

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0653	152	94.7	0.0	3.3	0.0	0.0	2.0	0.0
0654	130	87.7	3.8	3.1	0.0	0.0	5.4	0.0
0655	101	97.0	0.0	1.0	0.0	1.0	1.0	0.0
0657	19	*	*	*	*	*	*	*
0701	159	89.9	3.1	2.5	0.0	1.9	1.9	0.6
0702	29	*	*	*	*	*	*	*
0703	49	*	*	*	*	*	*	*
0704	20	*	*	*	*	*	*	*
0705	31	*	*	*	*	*	*	*
0706	3	*	*	*	*	*	*	*
0850	236	81.4	4.7	11.9	0.0	1.3	0.4	0.4
0851	216	93.5	0.0	4.6	0.0	0.9	0.9	0.0
0901	504	79.4	2.0	10.9	0.0	0.2	7.1	0.4
0902	12	*	*	*	*	*	*	*
1051	374	85.3	0.5	7.8	1.9	0.5	3.5	0.5
1101	525	34.9	51.0	8.4	0.0	1.3	4.0	0.4
1102	279	74.2	11.5	7.9	0.4	0.7	5.4	0.0
1106	29	*	*	*	*	*	*	*
1250	345	85.8	0.3	10.1	0.0	0.9	1.4	1.4
1251	216	95.4	0.0	2.3	0.5	0.0	1.9	0.0
1252	167	62.3	0.0	31.7	0.0	0.0	6.0	0.0
1253	252	77.0	1.6	12.3	0.0	0.0	8.7	0.4
1350	763	54.5	20.1	9.4	0.8	7.1	7.9	0.3

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 3

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
1351	27	*	*	*	*	*	*	*
1352	144	73.6	16.7	3.5	0.0	0.0	6.3	0.0
1401	64	*	*	*	*	*	*	*
1402	40	*	*	*	*	*	*	*
1403	5	*	*	*	*	*	*	*
1501	262	47.3	5.0	43.5	0.0	0.0	2.3	1.9
1503	286	61.9	7.7	21.0	0.7	0.0	8.0	0.7
1504	663	34.7	5.3	49.9	0.2	4.5	4.7	0.8
1505	109	75.2	4.6	15.6	0.0	0.0	3.7	0.9
1506	191	84.3	1.0	11.5	0.0	0.5	1.6	1.0
1507	171	80.7	1.2	9.9	0.0	0.0	6.4	1.8
1508	6	*	*	*	*	*	*	*
1509	122	46.7	0.0	47.5	0.0	0.0	4.9	0.8
1511	188	89.9	2.7	4.8	0.5	0.5	1.6	0.0
1512	111	83.8	0.9	8.1	0.0	0.9	6.3	0.0
1514	117	83.8	3.4	6.0	0.9	0.9	5.1	0.0
1701	191	55.5	1.0	1.0	18.8	0.5	23.0	0.0
1950	213	89.7	0.0	6.6	0.0	0.5	3.3	0.0
1952	52	*	*	*	*	*	*	*
1953	82	*	*	*	*	*	*	*
1954	152	86.2	0.0	11.8	0.7	0.0	1.3	0.0
1955	34	*	*	*	*	*	*	*
1957	115	81.7	2.6	11.3	0.0	0.0	3.5	0.9

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 4

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
1963	16	*	*	*	*	*	*	*
1964	1	*	*	*	*	*	*	*
2150	212	85.8	0.0	2.4	1.4	0.0	9.9	0.5
2151	60	*	*	*	*	*	*	*
2152	4	*	*	*	*	*	*	*
2251	129	95.3	0.0	2.3	0.0	0.0	2.3	0.0
2252	58	*	*	*	*	*	*	*
2554	11	*	*	*	*	*	*	*
2555	2,396	47.0	40.1	6.0	0.1	0.2	6.4	0.3
2556	88	*	*	*	*	*	*	*
2557	1	*	*	*	*	*	*	*
2558	1	*	*	*	*	*	*	*
2560	2	*	*	*	*	*	*	*
2561	1	*	*	*	*	*	*	*
2562	3	*	*	*	*	*	*	*
2563	8	*	*	*	*	*	*	*
2564	15	*	*	*	*	*	*	*
3001	383	19.6	61.4	14.9	0.0	0.8	3.4	0.0
3002	92	*	*	*	*	*	*	*
3004	71	*	*	*	*	*	*	*
3005	47	*	*	*	*	*	*	*
3007	4	*	*	*	*	*	*	*
3201	145	91.0	0.7	6.2	0.0	0.0	2.1	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 5

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
3203	1	*	*	*	*	*	*	*
3301	1,386	36.1	31.3	16.9	0.1	6.0	9.2	0.4
3304	27	*	*	*	*	*	*	*
3305	50	*	*	*	*	*	*	*
3306	90	*	*	*	*	*	*	*
3308	45	*	*	*	*	*	*	*
3312	113	27.4	26.5	36.3	1.8	1.8	6.2	0.0
3550	26	*	*	*	*	*	*	*
3551	4	*	*	*	*	*	*	*
3552	13	*	*	*	*	*	*	*
3701	308	88.6	0.3	8.1	0.0	0.0	2.9	0.0
3850	983	70.0	14.6	5.6	0.2	0.5	9.0	0.1
3901	510	33.1	52.7	7.3	0.0	1.0	5.7	0.2
3902	33	*	*	*	*	*	*	*
3903	5	*	*	*	*	*	*	*
3950	719	52.0	24.1	13.2	0.3	1.3	8.6	0.6
3954	5	*	*	*	*	*	*	*
4106	738	11.2	44.6	38.2	0.4	0.7	4.9	0.0
4107	982	40.2	28.1	22.2	0.3	2.1	6.8	0.1
4109	245	78.8	2.9	11.8	0.0	0.4	6.1	0.0
4110	2	*	*	*	*	*	*	*
4112	173	1.2	0.6	97.1	0.6	0.6	0.0	0.0
4115	1,151	31.6	30.9	20.5	0.2	12.3	4.2	0.3

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 6

### Michigan

#### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
4351	178	90.4	1.1	5.1	0.0	0.0	2.8	0.6
4352	53	*	*	*	*	*	*	*
4353	147	58.5	1.4	35.4	0.7	0.0	4.1	0.0
4354	231	84.8	4.8	2.2	0.4	1.3	6.1	0.4
4355	97	*	*	*	*	*	*	*
4356	137	83.9	1.5	8.0	0.0	1.5	5.1	0.0
4358	23	*	*	*	*	*	*	*
4359	182	65.9	0.5	31.3	0.5	0.0	0.0	1.6
4360	67	*	*	*	*	*	*	*
4361	85	*	*	*	*	*	*	*
4362	95	*	*	*	*	*	*	*
4363	258	91.1	0.0	2.7	0.0	0.0	5.8	0.4
4401	272	90.8	0.7	5.1	0.0	0.4	2.9	0.0
4501	277	93.1	0.4	5.1	0.0	0.4	1.1	0.0
4601	440	73.0	2.3	20.5	0.0	0.2	3.9	0.2
4701	320	92.2	1.6	3.1	0.0	0.6	2.2	0.3
5050	504	50.6	39.1	3.0	0.0	1.4	5.4	0.6
5051	929	51.3	36.6	4.4	0.0	2.7	4.8	0.1
5052	939	35.6	40.6	2.6	0.3	17.0	3.4	0.4
5055	404	85.4	5.4	0.2	0.0	8.4	0.5	0.0
5101	541	68.8	12.0	11.1	0.4	2.0	5.7	0.0
5102	12	*	*	*	*	*	*	*
5202	231	84.8	1.3	3.9	1.7	1.3	6.9	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 7

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
5206	5	*	*	*	*	*	*	*
5601	300	89.3	1.3	3.7	0.0	2.3	3.3	0.0
5801	611	81.3	7.2	5.4	0.2	0.3	4.4	1.1
5803	34	*	*	*	*	*	*	*
5804	8	*	*	*	*	*	*	*
5806	20	*	*	*	*	*	*	*
6101	10	*	*	*	*	*	*	*
6102	32	*	*	*	*	*	*	*
6103	1,127	49.8	32.4	8.9	0.4	0.5	7.7	0.4
6105	4	*	*	*	*	*	*	*
6106	31	*	*	*	*	*	*	*
6301	1,122	38.7	31.2	22.6	0.1	1.4	5.8	0.2
6312	589	35.5	53.8	3.2	0.0	2.4	5.1	0.0
6316	361	65.7	19.4	7.2	0.3	3.3	3.9	0.3
6318	22	*	*	*	*	*	*	*
6320	113	36.3	52.2	4.4	0.9	0.9	5.3	0.0
6325	103	94.2	0.0	1.0	0.0	0.0	3.9	1.0
6335	311	71.7	17.7	1.9	0.0	6.4	2.3	0.0
6340	1	*	*	*	*	*	*	*
6345	30	*	*	*	*	*	*	*
6360	334	29.9	51.2	12.9	0.0	1.5	4.5	0.0
7306	730	44.5	32.1	15.1	0.0	0.7	7.0	0.7
7401	684	80.0	6.9	5.6	0.0	0.6	6.9	0.1

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 8

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
7601	205	92.7	0.0	5.4	0.5	0.5	0.5	0.5
7850	331	87.0	0.3	9.7	0.0	0.0	3.0	0.0
7852	9	*	*	*	*	*	*	*
8101	557	4.5	91.6	0.4	0.0	0.4	3.2	0.0
8102	140	10.7	13.6	69.3	0.0	1.4	5.0	0.0
8105	234	31.6	63.7	2.6	0.4	0.4	0.9	0.4
8106	181	24.9	55.2	11.6	0.6	1.1	6.1	0.6
8150	1	*	*	*	*	*	*	*
8151	986	37.7	41.5	10.2	0.1	2.7	7.4	0.3
8152	17	*	*	*	*	*	*	*
8153	4	*	*	*	*	*	*	*
8155	5	*	*	*	*	*	*	*
8157	1	*	*	*	*	*	*	*
8201	956	45.2	38.3	4.9	0.2	3.9	7.0	0.5
8202	1,371	82.0	14.2	1.8	0.0	0.5	1.1	0.4
8203	140	62.1	27.9	4.3	0.7	1.4	3.6	0.0
8250	262	23.7	65.6	5.3	0.0	0.4	5.0	0.0
8252	58	*	*	*	*	*	*	*
8255	779	54.0	31.2	9.1	0.1	2.4	3.1	0.0
8257	686	53.5	12.2	0.4	0.3	32.1	1.5	0.0
8263	239	66.1	16.7	11.3	0.0	1.7	4.2	0.0
8264	281	63.7	28.1	4.3	0.0	1.4	2.5	0.0
8267	207	53.6	19.8	23.7	0.0	0.0	2.9	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 9

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
8901	308	3.9	90.6	1.0	0.0	0.3	4.2	0.0
8902	479	6.3	89.6	2.5	0.0	0.0	1.3	0.4
8903	515	3.9	90.5	2.5	0.0	0.6	2.3	0.2
8904	820	3.7	90.6	2.0	0.1	0.9	2.4	0.4
8905	499	63.7	25.7	8.0	0.0	0.2	2.4	0.0
8909	274	5.5	90.9	1.1	0.0	0.0	2.6	0.0
9045	498	10.2	31.3	56.0	0.2	0.4	1.6	0.2
9050	334	14.1	67.7	14.7	0.0	0.0	3.6	0.0
9051	497	8.7	5.2	83.9	0.0	0.0	2.0	0.2
9053	380	7.4	87.9	1.6	0.3	1.1	1.8	0.0
9055	3	*	*	*	*	*	*	*
9058	328	17.4	70.4	1.8	0.0	8.5	1.5	0.3
9066	440	2.0	94.3	0.5	0.0	0.5	2.7	0.0
9068	330	2.1	96.1	0.3	0.0	0.6	0.9	0.0
9069	740	1.2	97.6	0.8	0.0	0.1	0.3	0.0
9070	417	4.6	92.1	1.4	0.0	1.0	1.0	0.0
9072	327	2.8	94.2	0.6	0.3	0.6	1.5	0.0
9080	831	19.3	76.4	2.2	0.1	0.4	1.7	0.0
9082	40	*	*	*	*	*	*	*
9083	1	*	*	*	*	*	*	*
9084	5	*	*	*	*	*	*	*
9087	4	*	*	*	*	*	*	*
<b>Michigan</b>	<b>50,034</b>	<b>50.8</b>	<b>30.8</b>	<b>10.9</b>	<b>0.4</b>	<b>2.3</b>	<b>4.6</b>	<b>0.3</b>

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Page : 10

### Michigan

#### Comparison of Racial and Ethnic Distribution by Clinic

National (Prior Year)	146,574	54.6	25.3	12.2	1.2	2.5	3.9	0.3
-----------------------	---------	------	------	------	-----	-----	-----	-----

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Page : 11

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	40	*	*	*	*	*	*	*
0102	151	0.0	4.0	7.9	62.3	23.8	2.0	0.0
0103	121	0.0	4.1	5.0	66.9	23.1	0.8	0.0
0104	39	*	*	*	*	*	*	*
0105	11	*	*	*	*	*	*	*
0250	39	*	*	*	*	*	*	*
0251	29	*	*	*	*	*	*	*
0252	30	*	*	*	*	*	*	*
0253	46	*	*	*	*	*	*	*
0401	153	0.0	2.0	10.5	63.4	23.5	0.7	0.0
0402	118	0.0	0.8	13.6	59.3	23.7	2.5	0.0
0403	53	*	*	*	*	*	*	*
0404	42	*	*	*	*	*	*	*
0501	159	0.6	1.3	5.0	69.8	20.8	2.5	0.0
0503	34	*	*	*	*	*	*	*
0504	135	0.0	3.0	3.0	62.2	28.9	3.0	0.0
0505	89	*	*	*	*	*	*	*
0506	48	*	*	*	*	*	*	*
0507	20	*	*	*	*	*	*	*
0508	10	*	*	*	*	*	*	*
0650	343	0.0	3.2	7.9	65.0	23.0	0.9	0.0
0651	182	0.5	6.0	6.0	62.6	23.6	1.1	0.0
0652	96	*	*	*	*	*	*	*
0653	152	0.0	1.3	6.6	63.2	27.0	2.0	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Page : 12

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0654	130	0.0	3.1	9.2	58.5	27.7	0.8	0.8
0655	101	0.0	1.0	13.9	57.4	26.7	1.0	0.0
0657	19	*	*	*	*	*	*	*
0701	159	0.0	1.9	5.0	56.6	35.2	1.3	0.0
0702	29	*	*	*	*	*	*	*
0703	49	*	*	*	*	*	*	*
0704	20	*	*	*	*	*	*	*
0705	31	*	*	*	*	*	*	*
0706	3	*	*	*	*	*	*	*
0850	236	0.0	2.1	10.6	61.4	23.3	1.7	0.8
0851	216	0.0	1.9	7.4	67.1	21.3	2.3	0.0
0901	504	0.0	2.4	8.5	63.1	24.6	1.4	0.0
0902	12	*	*	*	*	*	*	*
1051	374	0.3	2.4	8.0	57.0	29.1	2.4	0.8
1101	525	0.0	5.7	7.8	60.4	21.3	1.7	3.0
1102	279	0.0	1.8	11.8	60.2	23.3	2.2	0.7
1106	29	*	*	*	*	*	*	*
1250	345	0.0	2.6	10.4	62.6	22.6	1.7	0.0
1251	216	0.0	2.8	10.2	67.1	18.5	0.5	0.9
1252	167	0.6	1.8	9.0	62.9	24.6	1.2	0.0
1253	252	0.0	1.2	8.7	70.6	19.0	0.4	0.0
1350	763	0.0	3.7	6.6	64.6	23.2	1.6	0.4
1351	27	*	*	*	*	*	*	*
1352	144	0.0	2.1	6.9	64.6	22.9	2.8	0.7

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Page : 13

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1401	64	*	*	*	*	*	*	*
1402	40	*	*	*	*	*	*	*
1403	5	*	*	*	*	*	*	*
1501	262	0.0	0.8	8.0	61.1	26.7	1.9	1.5
1503	286	0.3	4.2	10.5	60.5	24.1	0.3	0.0
1504	663	0.3	2.1	6.9	63.0	24.9	2.4	0.3
1505	109	0.0	0.9	3.7	75.2	17.4	1.8	0.9
1506	191	0.0	1.0	6.3	60.2	31.4	1.0	0.0
1507	171	0.0	1.8	5.8	63.2	25.7	3.5	0.0
1508	6	*	*	*	*	*	*	*
1509	122	0.0	1.6	7.4	62.3	24.6	3.3	0.8
1511	188	0.0	2.7	11.2	59.0	26.1	1.1	0.0
1512	111	0.0	2.7	5.4	58.6	28.8	4.5	0.0
1514	117	0.0	0.0	3.4	65.0	29.1	1.7	0.9
1701	191	0.0	1.6	8.4	64.9	23.0	2.1	0.0
1950	213	0.0	2.3	14.6	61.5	20.2	0.9	0.5
1952	52	*	*	*	*	*	*	*
1953	82	*	*	*	*	*	*	*
1954	152	0.7	3.3	5.9	64.5	23.7	1.3	0.7
1955	34	*	*	*	*	*	*	*
1957	115	0.0	1.7	5.2	54.8	37.4	0.9	0.0
1963	16	*	*	*	*	*	*	*
1964	1	*	*	*	*	*	*	*
2150	212	0.0	2.4	6.6	56.6	34.4	0.0	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Page : 14

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2151	60	*	*	*	*	*	*	*
2152	4	*	*	*	*	*	*	*
2251	129	0.0	3.1	6.2	59.7	30.2	0.8	0.0
2252	58	*	*	*	*	*	*	*
2554	11	*	*	*	*	*	*	*
2555	2,396	0.3	2.6	7.9	63.9	23.8	1.5	0.0
2556	88	*	*	*	*	*	*	*
2557	1	*	*	*	*	*	*	*
2558	1	*	*	*	*	*	*	*
2560	2	*	*	*	*	*	*	*
2561	1	*	*	*	*	*	*	*
2562	3	*	*	*	*	*	*	*
2563	8	*	*	*	*	*	*	*
2564	15	*	*	*	*	*	*	*
3001	383	0.0	2.6	8.6	64.5	20.9	2.1	1.3
3002	92	*	*	*	*	*	*	*
3004	71	*	*	*	*	*	*	*
3005	47	*	*	*	*	*	*	*
3007	4	*	*	*	*	*	*	*
3201	145	0.0	0.7	8.3	57.2	31.7	2.1	0.0
3203	1	*	*	*	*	*	*	*
3301	1,386	0.1	3.3	8.4	61.1	25.3	1.7	0.1
3304	27	*	*	*	*	*	*	*
3305	50	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**WIC PNSS & PedNSS**  
**Table 4A**  
**2020 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Run Date: 4/21/2021

Page : 15

<b>Comparison of Age Distribution (2)</b>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
3306	90	*	*	*	*	*	*	*
3308	45	*	*	*	*	*	*	*
3312	113	0.0	0.9	7.1	67.3	21.2	3.5	0.0
3550	26	*	*	*	*	*	*	*
3551	4	*	*	*	*	*	*	*
3552	13	*	*	*	*	*	*	*
3701	308	0.0	3.2	9.7	66.6	19.8	0.6	0.0
3850	983	0.0	2.8	7.5	63.5	24.6	1.0	0.5
3901	510	0.2	3.7	7.6	58.6	26.9	2.2	0.8
3902	33	*	*	*	*	*	*	*
3903	5	*	*	*	*	*	*	*
3950	719	0.1	3.3	7.1	62.7	24.6	1.7	0.4
3954	5	*	*	*	*	*	*	*
4106	738	0.0	2.6	9.5	58.5	26.0	2.3	1.1
4107	982	0.2	2.4	8.1	58.6	29.9	0.7	0.0
4109	245	0.0	2.0	7.8	64.5	24.9	0.8	0.0
4110	2	*	*	*	*	*	*	*
4112	173	0.6	4.0	8.7	58.4	26.6	1.2	0.6
4115	1,151	0.1	1.3	6.1	57.4	32.8	1.7	0.5
4351	178	0.0	1.1	9.0	68.0	20.8	1.1	0.0
4352	53	*	*	*	*	*	*	*
4353	147	0.0	3.4	6.8	66.0	19.0	4.8	0.0
4354	231	0.0	2.2	5.2	66.2	25.1	1.3	0.0
4355	97	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Page : 16

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4356	137	0.0	3.6	7.3	67.2	20.4	1.5	0.0
4358	23	*	*	*	*	*	*	*
4359	182	0.0	1.6	5.5	61.5	27.5	3.3	0.5
4360	67	*	*	*	*	*	*	*
4361	85	*	*	*	*	*	*	*
4362	95	*	*	*	*	*	*	*
4363	258	0.0	2.3	10.5	67.4	18.6	1.2	0.0
4401	272	0.0	2.9	5.5	62.9	27.6	0.7	0.4
4501	277	0.0	2.9	6.1	63.5	24.5	2.5	0.4
4601	440	0.2	1.8	7.7	60.0	28.9	1.1	0.2
4701	320	0.0	0.9	5.3	61.6	30.6	1.6	0.0
5050	504	0.0	1.2	4.6	58.5	33.1	2.6	0.0
5051	929	0.1	1.5	4.6	57.3	34.0	2.5	0.0
5052	939	0.0	1.5	3.9	58.3	33.8	2.4	0.1
5055	404	0.0	0.0	1.0	51.2	43.8	4.0	0.0
5101	541	0.0	1.5	4.6	59.1	31.4	3.3	0.0
5102	12	*	*	*	*	*	*	*
5202	231	0.4	0.4	3.9	58.9	32.5	3.9	0.0
5206	5	*	*	*	*	*	*	*
5601	300	0.0	2.0	5.3	62.7	29.0	1.0	0.0
5801	611	0.0	2.5	7.9	61.4	26.0	2.1	0.2
5803	34	*	*	*	*	*	*	*
5804	8	*	*	*	*	*	*	*
5806	20	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Page : 17

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
6101	10	*	*	*	*	*	*	*
6102	32	*	*	*	*	*	*	*
6103	1,127	0.3	3.5	9.2	61.8	23.2	1.7	0.4
6105	4	*	*	*	*	*	*	*
6106	31	*	*	*	*	*	*	*
6301	1,122	0.0	3.0	6.2	61.3	27.0	2.4	0.0
6312	589	0.0	0.5	2.5	54.0	40.6	2.4	0.0
6316	361	0.0	0.3	2.5	60.1	34.6	2.5	0.0
6318	22	*	*	*	*	*	*	*
6320	113	0.0	2.7	0.9	58.4	35.4	2.7	0.0
6325	103	0.0	1.0	7.8	62.1	27.2	1.0	1.0
6335	311	0.0	0.0	1.9	53.1	40.2	4.8	0.0
6340	1	*	*	*	*	*	*	*
6345	30	*	*	*	*	*	*	*
6360	334	0.0	2.7	8.1	57.2	27.5	3.0	1.5
7306	730	0.1	3.2	8.8	61.9	24.9	1.1	0.0
7401	684	0.3	2.3	7.6	65.9	22.2	1.5	0.1
7601	205	0.0	1.5	6.3	68.8	22.4	1.0	0.0
7850	331	0.0	2.4	6.9	68.0	22.1	0.6	0.0
7852	9	*	*	*	*	*	*	*
8101	557	0.0	2.2	8.4	59.8	27.3	2.0	0.4
8102	140	0.0	3.6	5.0	57.1	31.4	2.9	0.0
8105	234	0.0	2.6	5.1	53.0	37.6	1.7	0.0
8106	181	0.0	1.7	5.5	66.3	23.8	2.8	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Page : 18

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8150	1	*	*	*	*	*	*	*
8151	986	0.0	1.2	5.5	56.9	33.9	2.3	0.2
8152	17	*	*	*	*	*	*	*
8153	4	*	*	*	*	*	*	*
8155	5	*	*	*	*	*	*	*
8157	1	*	*	*	*	*	*	*
8201	956	0.1	1.8	4.6	58.8	32.4	2.1	0.2
8202	1,371	0.0	1.0	3.3	55.4	34.8	4.7	0.8
8203	140	0.0	0.7	5.0	55.0	37.9	1.4	0.0
8250	262	0.0	2.7	5.3	60.3	27.9	3.8	0.0
8252	58	*	*	*	*	*	*	*
8255	779	0.0	1.7	7.3	62.0	27.1	1.9	0.0
8257	686	0.0	0.4	3.4	59.8	31.2	5.1	0.1
8263	239	0.0	0.0	5.0	64.9	28.0	2.1	0.0
8264	281	0.0	1.4	2.1	62.3	32.0	2.1	0.0
8267	207	0.0	4.3	5.3	60.9	26.6	2.9	0.0
8901	308	0.6	3.9	8.4	57.8	26.0	2.9	0.3
8902	479	0.2	1.3	5.4	60.3	31.5	1.3	0.0
8903	515	0.0	4.1	9.1	61.9	22.7	1.9	0.2
8904	820	0.4	2.9	6.6	62.1	26.2	1.8	0.0
8905	499	0.0	1.4	3.8	53.5	36.3	4.8	0.2
8909	274	0.0	1.5	8.0	57.7	29.9	2.6	0.4
9045	498	0.2	2.8	6.8	54.0	31.9	4.2	0.0
9050	334	0.0	1.8	6.6	56.0	34.4	1.2	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**WIC PNSS & PedNSS**  
**Table 4A**  
**2020 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Run Date: 4/21/2021

Page : 19

<b>Comparison of Age Distribution (2)</b>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
9051	497	0.2	2.0	8.7	53.5	30.4	5.2	0.0
9053	380	0.0	3.2	6.8	56.8	30.5	2.1	0.5
9055	3	*	*	*	*	*	*	*
9058	328	0.0	0.9	6.7	56.7	31.7	3.7	0.3
9066	440	0.5	2.7	6.4	63.0	25.0	2.3	0.2
9068	330	0.3	1.8	5.2	60.6	29.1	2.7	0.3
9069	740	0.1	3.9	6.1	54.9	32.2	2.7	0.1
9070	417	0.2	2.6	5.5	61.6	26.4	3.6	0.0
9072	327	0.6	2.8	10.4	57.8	26.6	1.5	0.3
9080	831	0.0	1.8	5.8	57.8	32.5	2.2	0.0
9082	40	*	*	*	*	*	*	*
9083	1	*	*	*	*	*	*	*
9084	5	*	*	*	*	*	*	*
9087	4	*	*	*	*	*	*	*
<b>Michigan</b>	<b>50,036</b>	<b>0.1</b>	<b>2.2</b>	<b>6.8</b>	<b>60.6</b>	<b>27.9</b>	<b>2.1</b>	<b>0.3</b>
<b>National (Prior Year)</b>	<b>146,580</b>	<b>0.1</b>	<b>2.9</b>	<b>8.8</b>	<b>61.5</b>	<b>24.9</b>	<b>1.8</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Page : 20

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	28	*	*	12	*	*	7	*	1	*	36	*	36	*
0102	92	*	*	24	*	*	12	*	1	*	134	7.5 (42)	134	12.7 (61)
0103	104	5.8 (89)	66.3 (89)	38	*	*	5	*	3	*	106	4.7 (6)	106	13.2 (67)
0104	27	*	*	7	*	*	2	*	1	*	37	*	38	*
0105	7	*	*	1	*	*	0	*	0	*	11	*	11	*
0250	37	*	*	26	*	*	1	*	1	*	32	*	32	*
0251	20	*	*	11	*	*	0	*	0	*	26	*	26	*
0252	30	*	*	27	*	*	4	*	1	*	28	*	28	*
0253	45	*	*	39	*	*	0	*	0	*	42	*	42	*
0401	138	6.5 (94)	58.7 (26)	64	*	*	11	*	2	*	132	8.3 (56)	133	18.0 (101)
0402	103	1.9 (8)	60.2 (44)	30	*	*	3	*	0	*	106	9.4 (74)	106	8.5 (12)
0403	49	*	*	32	*	*	4	*	1	*	48	*	48	*
0404	41	*	*	31	*	*	3	*	0	*	40	*	40	*
0501	101	4.0 (47)	57.4 (15)	31	*	*	10	*	3	*	148	8.8 (62)	148	12.8 (62)
0503	31	*	*	10	*	*	3	*	0	*	31	*	31	*
0504	120	5.0 (77)	59.2 (31)	60	*	*	3	*	0	*	125	6.4 (24)	125	15.2 (85)
0505	63	*	*	15	*	*	7	*	2	*	82	*	82	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Page : 21

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	38	*	*	5	*	*	4	*	1	*	43	*	43	*
0507	16	*	*	4	*	*	0	*	1	*	20	*	20	*
0508	8	*	*	4	*	*	0	*	0	*	10	*	10	*
0650	298	4.7 (70)	60.4 (48)	179	15.6 (20)	48.6 (13)	20	*	11	*	308	8.8 (62)	308	9.4 (24)
0651	165	3.0 (24)	60.0 (41)	101	18.8 (42)	49.5 (18)	10	*	7	*	168	7.7 (45)	168	10.7 (35)
0652	73	*	*	57	*	*	6	*	2	*	91	*	92	*
0653	146	2.7 (19)	65.8 (86)	115	7.8 (1)	61.7 (65)	6	*	4	*	142	10.6 (90)	143	17.5 (98)
0654	111	3.6 (36)	61.3 (52)	73	*	*	10	*	1	*	123	5.7 (15)	123	10.6 (33)
0655	99	*	*	79	*	*	7	*	0	*	86	*	86	*
0657	12	*	*	7	*	*	2	*	0	*	18	*	18	*
0701	141	1.4 (4)	56.0 (8)	104	20.2 (51)	42.3 (3)	7	*	1	*	140	8.6 (60)	140	11.4 (43)
0702	29	*	*	23	*	*	0	*	3	*	28	*	28	*
0703	45	*	*	39	*	*	4	*	0	*	47	*	47	*
0704	20	*	*	15	*	*	0	*	0	*	18	*	18	*
0705	28	*	*	19	*	*	3	*	0	*	28	*	28	*
0706	3	*	*	2	*	*	0	*	0	*	3	*	3	*
0850	140	7.9 (100)	56.4 (10)	62	*	*	11	*	6	*	218	11.9 (99)	218	17.0 (92)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Page : 22

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0851	83	*	*	15	*	*	18	*	3	*	198	14.1 (107)	198	14.6 (80)
0901	420	3.1 (25)	63.3 (70)	215	16.7 (27)	52.6 (34)	28	*	3	*	458	5.7 (15)	458	11.1 (39)
0902	9	*	*	3	*	*	0	*	1	*	9	*	9	*
1051	313	5.1 (80)	51.8 (2)	240	15.0 (12)	48.3 (12)	21	*	2	*	343	5.2 (10)	345	9.3 (22)
1101	455	2.9 (22)	64.4 (78)	326	19.0 (43)	57.4 (61)	46	*	33	*	467	9.2 (71)	467	20.1 (109)
1102	251	4.8 (74)	58.6 (24)	155	13.5 (2)	58.1 (63)	24	*	20	*	231	7.8 (46)	231	16.9 (90)
1106	23	*	*	19	*	*	4	*	2	*	23	*	23	*
1250	172	4.1 (50)	65.1 (83)	72	*	*	14	*	3	*	314	10.5 (87)	314	7.6 (9)
1251	179	5.0 (77)	66.5 (90)	134	14.9 (10)	50.0 (19)	4	*	6	*	204	5.4 (12)	204	4.4 (5)
1252	133	4.5 (63)	65.4 (84)	38	*	*	8	*	1	*	149	7.4 (40)	149	8.7 (14)
1253	222	3.6 (36)	58.6 (24)	93	*	*	8	*	7	*	230	9.1 (69)	229	10.9 (36)
1350	574	5.1 (80)	56.3 (9)	339	20.6 (53)	50.4 (23)	34	*	9	*	677	9.3 (73)	678	11.2 (40)
1351	20	*	*	9	*	*	3	*	0	*	22	*	22	*
1352	69	*	*	31	*	*	3	*	1	*	130	3.1 (2)	130	3.8 (3)
1401	52	*	*	28	*	*	4	*	1	*	59	*	59	*
1402	37	*	*	20	*	*	0	*	0	*	36	*	36	*
1403	3	*	*	2	*	*	0	*	0	*	5	*	5	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Page : 23

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1501	212	1.9 (8)	69.3 (96)	92	*	*	22	*	14	*	235	11.9 (99)	235	8.9 (17)
1503	240	4.2 (55)	58.8 (29)	165	15.2 (14)	52.7 (35)	18	*	11	*	270	7.8 (46)	271	13.3 (71)
1504	285	7.0 (98)	54.7 (6)	129	14.7 (8)	57.4 (61)	47	*	41	*	591	9.1 (69)	592	8.3 (11)
1505	65	*	*	13	*	*	4	*	6	*	96	*	96	*
1506	146	4.1 (50)	63.0 (64)	19	*	*	11	*	10	*	169	12.4 (102)	170	11.8 (46)
1507	108	2.8 (20)	63.9 (75)	19	*	*	11	*	7	*	156	9.0 (67)	156	5.8 (7)
1508	4	*	*	3	*	*	0	*	1	*	4	*	4	*
1509	88	*	*	62	*	*	7	*	13	*	107	5.6 (14)	107	10.3 (31)
1511	157	4.5 (63)	59.9 (39)	83	*	*	16	*	12	*	178	12.9 (106)	178	12.9 (63)
1512	83	*	*	18	*	*	7	*	8	*	101	4.0 (5)	101	13.9 (77)
1514	46	*	*	23	*	*	9	*	16	*	113	6.2 (21)	113	18.6 (104)
1701	172	2.9 (22)	62.8 (62)	124	15.3 (15)	50.8 (25)	4	*	4	*	170	8.2 (54)	172	9.9 (25)
1950	192	2.1 (14)	62.0 (58)	98	*	*	10	*	3	*	195	12.8 (105)	195	11.8 (46)
1952	51	*	*	28	*	*	6	*	2	*	50	*	50	*
1953	75	*	*	44	*	*	6	*	3	*	74	*	74	*
1954	140	1.4 (4)	68.6 (93)	93	*	*	9	*	3	*	138	18.8 (111)	138	17.4 (96)
1955	34	*	*	21	*	*	1	*	1	*	32	*	32	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 24

### Michigan

### Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1957	102	4.9 (76)	72.5 (100)	57	*	*	7	*	0	*	104	19.2 (112)	104	17.3 (94)
1963	14	*	*	9	*	*	0	*	0	*	14	*	15	*
1964	1	*	*	1	*	*	0	*	0	*	1	*	1	*
2150	198	2.5 (18)	63.6 (71)	150	19.3 (45)	52.7 (35)	11	*	3	*	191	12.0 (101)	191	12.0 (50)
2151	53	*	*	44	*	*	1	*	0	*	48	*	48	*
2152	4	*	*	2	*	*	0	*	0	*	3	*	4	*
2251	96	*	*	36	*	*	9	*	4	*	114	0.9 (1)	115	14.8 (82)
2252	39	*	*	10	*	*	4	*	1	*	49	*	49	*
2554	10	*	*	7	*	*	2	*	0	*	11	*	11	*
2555	1,640	4.3 (58)	60.1 (42)	723	16.2 (23)	52.3 (32)	144	42.4 (1)	35	*	2,157	8.0 (50)	2,161	15.5 (87)
2556	80	*	*	35	*	*	3	*	4	*	81	*	81	*
2557	1	*	*	1	*	*	0	*	0	*	1	*	1	*
2558	1	*	*	0	*	*	0	*	0	*	0	*	0	*
2560	1	*	*	0	*	*	0	*	0	*	1	*	1	*
2561	1	*	*	1	*	*	1	*	0	*	1	*	1	*
2562	0	*	*	0	*	*	0	*	0	*	3	*	3	*
2563	3	*	*	1	*	*	0	*	1	*	6	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 25

### Michigan

### Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2564	8	*	*	1	*	*	1	*	1	*	11	*	11	*
3001	348	3.7 (41)	63.2 (67)	165	21.8 (60)	43.0 (5)	17	*	21	*	364	7.4 (40)	364	10.2 (29)
3002	82	*	*	33	*	*	4	*	6	*	84	*	84	*
3004	42	*	*	14	*	*	1	*	1	*	64	*	64	*
3005	44	*	*	19	*	*	4	*	2	*	43	*	43	*
3007	3	*	*	0	*	*	0	*	0	*	3	*	3	*
3201	127	4.7 (70)	69.3 (96)	92	*	*	10	*	3	*	141	9.9 (78)	142	12.0 (50)
3203	1	*	*	0	*	*	0	*	0	*	1	*	1	*
3301	1,108	3.7 (41)	63.2 (67)	886	20.4 (52)	54.3 (47)	87	*	17	*	1,306	6.8 (27)	1,306	13.2 (67)
3304	20	*	*	15	*	*	0	*	1	*	24	*	24	*
3305	37	*	*	30	*	*	4	*	0	*	48	*	48	*
3306	73	*	*	62	*	*	3	*	0	*	86	*	86	*
3308	40	*	*	34	*	*	3	*	0	*	42	*	42	*
3312	92	*	*	67	*	*	4	*	0	*	102	12.7 (104)	102	17.6 (99)
3550	22	*	*	6	*	*	1	*	0	*	22	*	22	*
3551	2	*	*	0	*	*	0	*	0	*	3	*	3	*
3552	13	*	*	4	*	*	1	*	0	*	11	*	11	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 26

### Michigan

### Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3701	231	6.1 (92)	60.2 (44)	167	17.4 (32)	53.3 (40)	10	*	4	*	284	9.5 (75)	284	16.9 (90)
3850	682	4.5 (63)	61.6 (54)	463	14.9 (10)	55.3 (55)	36	*	9	*	914	8.4 (58)	916	11.0 (37)
3901	391	2.0 (12)	59.3 (32)	165	17.0 (30)	54.5 (49)	24	*	5	*	444	9.9 (78)	444	10.1 (26)
3902	26	*	*	12	*	*	3	*	0	*	22	*	22	*
3903	5	*	*	2	*	*	0	*	1	*	5	*	5	*
3950	579	4.5 (63)	60.1 (42)	319	15.4 (17)	51.1 (27)	29	*	11	*	615	9.9 (78)	615	11.2 (40)
3954	4	*	*	4	*	*	0	*	0	*	5	*	5	*
4106	409	3.7 (41)	61.9 (56)	169	20.7 (54)	44.4 (8)	52	*	23	*	681	7.0 (32)	681	12.3 (59)
4107	697	4.3 (58)	58.4 (23)	273	16.1 (21)	53.8 (42)	52	*	22	*	878	7.1 (35)	879	15.1 (84)
4109	136	2.2 (16)	57.4 (15)	95	*	*	14	*	5	*	224	7.1 (35)	224	17.4 (96)
4110	1	*	*	1	*	*	0	*	0	*	2	*	2	*
4112	172	1.7 (6)	68.0 (92)	101	62.4 (67)	17.8 (1)	13	*	2	*	158	17.1 (110)	158	13.9 (77)
4115	574	4.5 (63)	59.6 (36)	135	22.2 (61)	45.9 (9)	68	*	25	*	1,040	10.0 (81)	1,041	13.2 (67)
4351	166	4.2 (55)	64.5 (80)	106	21.7 (59)	42.5 (4)	9	*	5	*	162	11.7 (98)	162	19.8 (108)
4352	52	*	*	40	*	*	3	*	0	*	48	*	48	*
4353	116	0.9 (2)	69.0 (95)	87	*	*	5	*	5	*	128	12.5 (103)	128	19.5 (106)
4354	163	4.3 (58)	68.7 (94)	56	*	*	19	*	5	*	213	5.2 (10)	213	19.7 (107)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Page : 27

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4355	46	*	*	21	*	*	4	*	3	*	90	*	91	*
4356	127	0.0 (1)	57.5 (18)	102	14.7 (8)	61.8 (66)	4	*	3	*	127	8.7 (61)	127	17.3 (94)
4358	15	*	*	14	*	*	0	*	0	*	22	*	22	*
4359	153	5.2 (83)	62.7 (61)	117	19.7 (47)	48.7 (14)	4	*	9	*	168	10.7 (92)	168	12.5 (60)
4360	66	*	*	44	*	*	1	*	1	*	57	*	57	*
4361	72	*	*	33	*	*	2	*	1	*	77	*	77	*
4362	74	*	*	51	*	*	2	*	5	*	90	*	90	*
4363	218	4.1 (50)	57.3 (14)	149	18.1 (38)	50.3 (21)	15	*	5	*	242	10.3 (86)	242	9.1 (20)
4401	263	4.2 (55)	57.4 (15)	223	17.9 (36)	54.7 (50)	12	*	5	*	247	4.9 (7)	247	11.7 (45)
4501	220	4.1 (50)	63.2 (67)	170	25.3 (66)	51.8 (31)	16	*	4	*	255	11.0 (94)	256	12.9 (63)
4601	279	4.3 (58)	59.9 (39)	173	14.5 (6)	50.3 (21)	27	*	8	*	398	10.1 (83)	399	14.3 (79)
4701	313	3.2 (27)	57.2 (13)	240	15.0 (12)	52.9 (37)	17	*	8	*	296	10.5 (87)	296	18.2 (103)
5050	483	4.6 (68)	62.1 (59)	412	15.5 (19)	57.3 (60)	26	*	6	*	465	8.8 (62)	465	11.6 (44)
5051	848	3.2 (27)	61.9 (56)	690	16.4 (25)	55.2 (53)	52	*	17	*	849	7.9 (49)	850	9.1 (20)
5052	870	4.0 (47)	59.4 (33)	728	21.0 (56)	50.1 (20)	43	*	15	*	881	10.8 (93)	881	11.0 (37)
5055	148	4.7 (70)	52.0 (3)	34	*	*	16	*	6	*	392	14.3 (109)	392	3.3 (2)
5101	427	5.6 (87)	56.7 (12)	244	13.9 (3)	56.1 (57)	37	*	15	*	496	11.1 (96)	496	13.3 (71)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Page : 28

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5102	11	*	*	7	*	*	3	*	0	*	12	*	12	*
5202	219	5.0 (77)	59.8 (37)	154	17.5 (33)	51.3 (29)	15	*	9	*	205	8.8 (62)	207	12.1 (55)
5206	4	*	*	4	*	*	0	*	0	*	5	*	5	*
5601	227	3.1 (25)	63.0 (64)	16	*	*	15	*	4	*	268	11.6 (97)	268	9.0 (18)
5801	533	5.4 (85)	60.6 (50)	393	15.3 (15)	53.9 (43)	35	*	12	*	548	8.8 (62)	549	18.0 (101)
5803	32	*	*	25	*	*	3	*	2	*	30	*	30	*
5804	8	*	*	6	*	*	1	*	0	*	7	*	7	*
5806	11	*	*	7	*	*	0	*	0	*	19	*	19	*
6101	10	*	*	9	*	*	0	*	0	*	9	*	9	*
6102	31	*	*	26	*	*	0	*	0	*	31	*	31	*
6103	999	3.3 (32)	66.0 (88)	767	17.7 (34)	50.8 (25)	39	*	8	*	1,036	10.0 (81)	1,032	19.2 (105)
6105	3	*	*	3	*	*	0	*	0	*	4	*	4	*
6106	28	*	*	19	*	*	0	*	2	*	26	*	26	*
6301	897	3.2 (27)	61.3 (52)	492	16.7 (27)	51.2 (28)	71	*	20	*	1,001	6.0 (18)	1,000	12.2 (57)
6312	514	3.3 (32)	59.5 (35)	312	18.3 (39)	52.9 (37)	39	*	11	*	525	3.6 (3)	525	10.3 (31)
6316	311	5.1 (80)	55.0 (7)	184	20.1 (49)	53.3 (40)	22	*	5	*	324	7.1 (35)	324	9.3 (22)
6318	7	*	*	0	*	*	2	*	0	*	21	*	21	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Page : 29

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6320	75	*	*	37	*	*	7	*	1	*	100	9.0 (67)	100	12.0 (50)
6325	75	*	*	38	*	*	7	*	3	*	85	*	85	*
6335	153	6.5 (94)	53.6 (4)	50	*	*	20	*	6	*	296	14.2 (108)	296	10.1 (26)
6340	1	*	*	0	*	*	0	*	0	*	0	*	0	*
6345	19	*	*	14	*	*	0	*	0	*	27	*	27	*
6360	205	3.9 (46)	53.7 (5)	80	*	*	13	*	2	*	300	7.3 (39)	300	13.0 (65)
7306	560	3.2 (27)	60.5 (49)	332	14.5 (6)	53.0 (39)	47	*	12	*	662	8.3 (56)	663	10.6 (33)
7401	651	3.5 (34)	58.7 (26)	546	18.7 (41)	54.4 (48)	43	*	9	*	616	7.1 (35)	617	13.0 (65)
7601	32	*	*	10	*	*	9	*	5	*	171	7.6 (43)	171	8.8 (16)
7850	298	4.4 (62)	64.8 (82)	240	17.9 (36)	51.3 (29)	12	*	5	*	292	9.2 (71)	293	13.3 (71)
7852	8	*	*	7	*	*	0	*	0	*	8	*	8	*
8101	483	2.3 (17)	64.4 (78)	358	22.3 (62)	47.8 (11)	40	*	10	*	521	8.1 (52)	521	17.1 (93)
8102	79	*	*	38	*	*	8	*	3	*	129	10.1 (83)	130	8.5 (12)
8105	186	4.8 (74)	71.5 (98)	85	*	*	31	*	3	*	189	6.9 (30)	189	4.2 (4)
8106	80	*	*	42	*	*	11	*	2	*	168	5.4 (12)	168	16.7 (89)
8150	1	*	*	0	*	*	0	*	0	*	1	*	1	*
8151	756	3.8 (45)	60.2 (44)	478	19.9 (48)	49.2 (17)	41	*	18	*	907	9.5 (75)	907	13.3 (71)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Page : 30

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8152	14	*	*	8	*	*	2	*	0	*	15	*	15	*
8153	4	*	*	1	*	*	1	*	0	*	3	*	3	*
8155	3	*	*	1	*	*	0	*	1	*	4	*	4	*
8157	1	*	*	1	*	*	0	*	0	*	1	*	1	*
8201	582	4.6 (68)	57.7 (19)	357	14.0 (4)	52.4 (33)	39	*	19	*	863	5.8 (17)	861	8.7 (14)
8202	697	5.9 (90)	50.4 (1)	317	22.7 (63)	43.5 (6)	74	*	23	*	1,315	3.6 (3)	1,314	2.1 (1)
8203	115	5.2 (83)	64.3 (77)	76	*	*	12	*	3	*	123	8.1 (52)	123	17.9 (100)
8250	182	6.6 (97)	57.7 (19)	105	19.0 (43)	55.2 (53)	17	*	6	*	226	8.4 (58)	225	12.0 (50)
8252	33	*	*	9	*	*	3	*	2	*	51	*	51	*
8255	592	5.4 (85)	61.0 (51)	403	14.4 (5)	55.3 (55)	64	*	17	*	678	5.0 (9)	678	14.6 (80)
8257	248	3.6 (36)	58.1 (21)	91	*	*	23	*	11	*	647	8.0 (50)	648	4.9 (6)
8263	189	3.7 (41)	58.2 (22)	117	15.4 (17)	65.0 (67)	16	*	3	*	199	10.6 (90)	198	10.1 (26)
8264	171	4.7 (70)	63.7 (72)	113	21.2 (58)	50.4 (23)	19	*	5	*	239	6.7 (26)	239	12.1 (55)
8267	115	6.1 (92)	61.7 (55)	84	*	*	12	*	4	*	192	6.3 (22)	190	7.4 (8)
8901	259	1.9 (8)	62.9 (63)	164	18.3 (39)	54.9 (51)	21	*	6	*	288	4.9 (7)	288	12.2 (57)
8902	249	2.8 (20)	59.0 (30)	99	*	*	31	*	5	*	437	6.9 (30)	437	11.9 (49)
8903	362	1.7 (6)	65.5 (85)	173	17.3 (31)	59.0 (64)	22	*	5	*	470	6.8 (27)	469	21.5 (112)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Page : 31

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8904	613	3.6 (36)	63.8 (74)	350	17.7 (34)	54.9 (51)	63	*	7	*	744	7.8 (46)	746	13.3 (71)
8905	280	3.6 (36)	56.4 (10)	110	23.6 (64)	43.6 (7)	31	*	7	*	457	10.5 (87)	459	10.2 (29)
8909	182	7.1 (99)	63.7 (72)	77	*	*	14	*	1	*	255	8.2 (54)	255	13.7 (76)
9045	313	1.3 (3)	71.6 (99)	157	21.0 (56)	39.5 (2)	26	*	6	*	463	10.2 (85)	463	7.8 (10)
9050	261	5.7 (88)	59.4 (33)	60	*	*	17	*	4	*	312	6.1 (19)	312	11.2 (40)
9051	308	1.9 (8)	65.9 (87)	140	20.7 (54)	46.4 (10)	23	*	2	*	465	9.7 (77)	465	9.0 (18)
9053	342	4.1 (50)	66.7 (91)	225	16.9 (29)	54.2 (45)	19	*	4	*	355	7.0 (32)	355	13.2 (67)
9055	3	*	*	0	*	*	0	*	0	*	3	*	3	*
9058	169	5.9 (90)	59.8 (37)	84	*	*	22	*	2	*	301	11.0 (94)	301	15.0 (83)
9066	288	2.1 (14)	58.7 (26)	127	25.2 (65)	48.8 (15)	25	*	5	*	416	7.0 (32)	419	20.5 (110)
9068	304	2.0 (12)	64.1 (76)	209	20.1 (49)	54.1 (44)	17	*	3	*	312	6.4 (24)	313	16.0 (88)
9069	532	3.2 (27)	63.0 (64)	279	16.1 (21)	49.1 (16)	55	*	4	*	710	6.3 (22)	710	12.0 (50)
9070	372	4.0 (47)	60.2 (44)	227	16.3 (24)	54.2 (45)	36	*	3	*	380	6.1 (19)	382	15.2 (85)
9072	245	6.5 (94)	62.4 (60)	118	19.5 (46)	56.8 (59)	25	*	4	*	294	6.8 (27)	294	20.7 (111)
9080	490	3.5 (34)	64.5 (80)	231	16.5 (26)	56.3 (58)	60	*	8	*	787	7.6 (43)	787	11.8 (46)
9082	34	*	*	14	*	*	1	*	0	*	37	*	37	*
9083	0	*	*	0	*	*	0	*	0	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 32

### Michigan

### Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9084	4	*	*	3	*	*	1	*	1	*	4	*	4	*
9087	4	*	*	2	*	*	0	*	0	*	3	*	3	*
Michigan	36,925	4.0	60.9	21,575	18.2	52.2	2,834	39.8	956	30.9	45,597	8.3	45,629	12.4
National (Prior Year)	128,555	4.0	61.6	89,471	22.3	49.7	34,531	19.0	48,409	17.0	94,843	9.2	94,594	9.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 33

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
0101	36	*	*	40	*	35	*	31 *
0102	135	83.0 (68)	8.1 (76)	145	38.6 (20)	139	21.6 (54)	128 84.4 (89)
0103	105	79.0 (86)	14.3 (93)	121	28.9 (62)	110	20.9 (60)	97 *
0104	38	*	*	39	*	36	*	33 *
0105	11	*	*	11	*	11	*	11 *
0250	31	*	*	38	*	27	*	32 *
0251	25	*	*	28	*	23	*	21 *
0252	28	*	*	30	*	28	*	25 *
0253	42	*	*	46	*	41	*	38 *
0401	131	80.9 (77)	16.8 (102)	152	42.8 (10)	138	21.7 (53)	127 95.3 (4)
0402	108	71.3 (105)	15.7 (99)	115	34.8 (38)	102	29.4 (18)	92 *
0403	49	*	*	52	*	46	*	39 *
0404	40	*	*	42	*	37	*	30 *
0501	149	95.3 (9)	1.3 (8)	159	33.3 (47)	134	33.6 (15)	116 91.4 (40)
0503	31	*	*	32	*	25	*	23 *
0504	125	84.8 (56)	6.4 (61)	134	27.6 (66)	121	33.1 (16)	96 *
0505	84	*	*	87	*	83	*	73 *
0506	43	*	*	48	*	45	*	41 *
0507	20	*	*	20	*	16	*	14 *
0508	10	*	*	10	*	9	*	8 *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 34

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	309	84.8 (56)	8.4 (78)	342	34.8 (38)	298	27.9 (26)	275	89.5 (53)
0651	171	83.0 (68)	8.8 (82)	181	45.3 (5)	150	25.3 (37)	153	88.9 (56)
0652	91	*	*	96	*	85	*	73	*
0653	145	65.5 (108)	28.3 (111)	151	47.0 (2)	141	34.8 (14)	122	87.7 (67)
0654	123	80.5 (80)	14.6 (95)	127	45.7 (4)	119	28.6 (22)	106	81.1 (98)
0655	86	*	*	98	*	86	*	73	*
0657	18	*	*	19	*	18	*	15	*
0701	138	77.5 (89)	15.2 (97)	156	32.7 (53)	134	36.6 (11)	119	92.4 (28)
0702	28	*	*	28	*	24	*	20	*
0703	45	*	*	49	*	42	*	32	*
0704	18	*	*	20	*	20	*	19	*
0705	28	*	*	30	*	29	*	30	*
0706	3	*	*	3	*	3	*	3	*
0850	213	85.0 (55)	4.2 (38)	234	31.6 (54)	211	27.5 (30)	174	93.1 (24)
0851	195	91.8 (21)	3.6 (27)	214	27.1 (70)	190	24.2 (40)	150	96.7 (3)
0901	463	73.0 (101)	19.9 (105)	501	41.5 (13)	461	13.7 (84)	415	84.3 (90)
0902	9	*	*	12	*	12	*	8	*
1051	346	76.9 (93)	15.3 (98)	371	27.0 (72)	325	26.5 (33)	255	89.0 (55)
1101	452	72.1 (104)	9.5 (85)	515	29.5 (61)	447	15.7 (77)	371	85.7 (82)
1102	226	77.0 (92)	14.6 (95)	271	36.9 (30)	249	25.3 (37)	209	95.2 (6)
1106	22	*	*	29	*	26	*	21	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Page : 35

Michigan

## Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1250	260	79.2 (83)	11.2 (88)	336	31.0 (57)	290	23.1 (45)	265	84.9 (86)
1251	202	83.2 (66)	9.4 (84)	214	37.9 (23)	187	16.6 (72)	164	90.9 (44)
1252	144	65.3 (109)	18.8 (104)	165	39.4 (18)	154	26.6 (32)	136	90.4 (48)
1253	219	64.4 (110)	22.8 (110)	249	34.9 (37)	220	16.4 (74)	194	90.7 (45)
1350	659	93.6 (12)	3.6 (27)	755	33.6 (44)	520	28.1 (24)	543	81.0 (100)
1351	22	*	*	25	*	26	*	26	*
1352	130	93.1 (15)	4.6 (43)	140	32.9 (51)	132	2.3 (105)	108	88.9 (56)
1401	59	*	*	63	*	58	*	49	*
1402	35	*	*	39	*	31	*	27	*
1403	5	*	*	5	*	5	*	3	*
1501	231	79.7 (82)	12.6 (91)	259	33.2 (50)	168	17.3 (70)	200	84.0 (92)
1503	262	86.6 (46)	8.0 (75)	282	44.0 (7)	243	21.0 (59)	219	80.8 (101)
1504	591	90.7 (29)	3.7 (32)	653	31.4 (55)	577	27.7 (27)	483	86.5 (76)
1505	95	*	*	107	36.4 (33)	88	*	108	89.8 (51)
1506	170	73.5 (100)	21.2 (106)	187	41.2 (14)	170	7.6 (99)	148	95.3 (4)
1507	150	90.7 (29)	6.0 (60)	169	33.7 (43)	147	23.8 (43)	123	91.1 (42)
1508	4	*	*	6	*	5	*	6	*
1509	108	91.7 (24)	4.6 (43)	122	41.8 (11)	105	19.0 (65)	94	*
1511	177	95.5 (8)	1.1 (7)	186	37.6 (27)	170	15.9 (76)	149	91.9 (35)
1512	101	96.0 (6)	2.0 (10)	111	25.2 (84)	91	*	69	*
1514	113	95.6 (7)	3.5 (26)	113	15.9 (113)	71	*	40	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 36

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1701	172	96.5 (5)	2.9 (20)	186	46.8 (3)	172	34.9 (13)	157	82.2 (96)
1950	193	83.9 (62)	9.8 (87)	209	34.4 (40)	176	26.1 (34)	157	94.3 (14)
1952	50	*	*	51	*	44	*	38	*
1953	75	*	*	80	*	73	*	64	*
1954	138	86.2 (49)	7.2 (73)	150	40.7 (15)	129	25.6 (35)	125	93.6 (20)
1955	32	*	*	33	*	29	*	25	*
1957	103	86.4 (48)	3.9 (34)	115	29.6 (60)	78	*	87	*
1963	16	*	*	16	*	10	*	9	*
1964	1	*	*	1	*	1	*	1	*
2150	190	91.6 (25)	2.1 (11)	211	35.5 (35)	192	28.1 (24)	169	94.1 (15)
2151	44	*	*	59	*	41	*	31	*
2152	4	*	*	3	*	2	*	0	*
2251	117	91.5 (26)	5.1 (47)	129	33.3 (47)	119	22.7 (48)	99	*
2252	51	*	*	57	*	50	*	48	*
2554	11	*	*	11	*	7	*	6	*
2555	2,136	87.3 (43)	3.6 (27)	2,366	28.0 (64)	1,738	5.9 (104)	1,718	87.8 (66)
2556	79	*	*	87	*	47	*	62	*
2557	1	*	*	1	*	0	*	0	*
2558	0	*	*	1	*	0	*	1	*
2560	1	*	*	2	*	2	*	2	*
2561	1	*	*	1	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 37

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
2562	3	*	*	3	*	1	*	3 *
2563	6	*	*	8	*	3	*	3 *
2564	11	*	*	15	*	13	*	12 *
3001	361	79.2 (83)	8.3 (77)	379	33.8 (42)	333	12.9 (87)	304 84.2 (91)
3002	82	*	*	92	*	86	*	83 *
3004	64	*	*	69	*	65	*	51 *
3005	44	*	*	47	*	46	*	40 *
3007	3	*	*	4	*	3	*	2 *
3201	142	78.2 (88)	5.6 (55)	144	27.1 (70)	119	23.5 (44)	100 92.0 (33)
3203	1	*	*	1	*	1	*	1 *
3301	1,298	87.6 (41)	6.8 (69)	1,373	27.5 (67)	1,212	6.4 (101)	1,040 88.8 (59)
3304	23	*	*	27	*	26	*	19 *
3305	47	*	*	50	*	44	*	35 *
3306	86	*	*	90	*	75	*	65 *
3308	42	*	*	44	*	37	*	24 *
3312	101	86.1 (50)	8.9 (83)	113	41.6 (12)	100	6.0 (103)	90 *
3550	23	*	*	25	*	23	*	22 *
3551	3	*	*	4	*	4	*	4 *
3552	11	*	*	13	*	13	*	13 *
3701	286	89.9 (32)	6.6 (67)	307	32.9 (51)	272	21.3 (56)	219 95.0 (8)
3850	907	82.8 (71)	9.7 (86)	970	37.9 (23)	822	20.2 (63)	772 86.7 (74)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A 2020 Pregnancy Nutrition Surveillance (1) Michigan

Page : 38

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3901	442	85.3 (54)	5.2 (48)	500	38.2 (22)	444	14.9 (78)	399	82.7 (95)
3902	22	*	*	33	*	31	*	32	*
3903	5	*	*	5	*	5	*	4	*
3950	614	84.7 (60)	6.4 (61)	715	31.3 (56)	614	18.6 (66)	511	87.9 (64)
3954	5	*	*	5	*	5	*	5	*
4106	675	86.8 (45)	4.3 (41)	731	25.9 (79)	601	9.0 (96)	539	83.7 (93)
4107	873	86.1 (50)	5.4 (52)	973	22.3 (94)	828	27.2 (31)	682	89.9 (50)
4109	221	79.2 (83)	13.6 (92)	243	33.3 (47)	216	20.4 (61)	182	94.5 (11)
4110	2	*	*	2	*	2	*	0	*
4112	160	43.1 (111)	0.0 (1)	171	24.0 (89)	155	63.2 (1)	155	92.9 (25)
4115	1,009	88.9 (36)	5.9 (59)	1,141	23.5 (90)	920	16.4 (74)	842	92.2 (32)
4351	161	89.4 (34)	5.6 (55)	174	44.3 (6)	53	*	146	93.2 (22)
4352	48	*	*	52	*	46	*	40	*
4353	124	72.6 (102)	21.8 (108)	143	40.6 (16)	137	25.5 (36)	112	82.1 (97)
4354	212	92.0 (20)	5.7 (58)	230	37.8 (25)	191	14.7 (81)	188	92.6 (27)
4355	92	*	*	96	*	83	*	72	*
4356	125	78.4 (87)	11.2 (88)	133	43.6 (8)	120	24.2 (40)	104	93.3 (21)
4358	22	*	*	23	*	23	*	23	*
4359	162	67.9 (107)	21.6 (107)	181	35.4 (36)	146	21.2 (58)	130	91.5 (39)
4360	57	*	*	64	*	61	*	53	*
4361	77	*	*	85	*	81	*	70	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Page : 39

Michigan

## Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4362	90	*	*	92	*	85	*	76	*
4363	238	74.8 (99)	18.1 (103)	252	38.5 (21)	221	27.6 (28)	193	90.2 (49)
4401	248	76.2 (98)	14.5 (94)	266	26.7 (74)	235	21.3 (56)	198	93.9 (16)
4501	259	76.8 (94)	16.6 (101)	276	39.1 (19)	253	28.5 (23)	215	94.9 (9)
4601	392	91.8 (21)	3.3 (23)	439	30.5 (58)	380	21.8 (52)	327	92.0 (33)
4701	295	85.8 (52)	7.8 (74)	316	26.3 (76)	254	39.8 (8)	219	93.2 (22)
5050	463	93.3 (13)	1.5 (9)	497	20.9 (99)	346	10.1 (92)	349	95.1 (7)
5051	845	92.8 (16)	2.1 (11)	913	23.4 (92)	680	11.0 (90)	631	94.5 (11)
5052	873	91.0 (27)	3.6 (27)	922	24.9 (85)	632	10.0 (94)	674	93.8 (18)
5055	391	83.1 (67)	16.1 (100)	397	48.6 (1)	352	7.1 (100)	330	89.4 (54)
5101	493	90.7 (29)	4.9 (46)	531	26.0 (78)	218	37.2 (10)	379	93.9 (16)
5102	12	*	*	12	*	10	*	10	*
5202	201	89.1 (35)	6.5 (64)	229	36.7 (32)	166	38.6 (9)	179	86.6 (75)
5206	5	*	*	4	*	1	*	0	*
5601	268	91.0 (27)	2.6 (16)	298	33.6 (44)	273	28.9 (21)	228	94.7 (10)
5801	546	83.5 (63)	8.4 (78)	602	30.1 (59)	537	22.9 (46)	442	88.9 (56)
5803	30	*	*	34	*	34	*	28	*
5804	7	*	*	8	*	7	*	4	*
5806	18	*	*	20	*	16	*	14	*
6101	9	*	*	9	*	10	*	9	*
6102	30	*	*	32	*	26	*	26	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 40

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6103	1,011	87.1 (44)	5.3 (50)	1,103	37.1 (29)	815	22.5 (49)	835	86.1 (79)
6105	4	*	*	4	*	4	*	2	*
6106	26	*	*	31	*	29	*	31	*
6301	995	86.6 (46)	6.5 (64)	1,114	27.4 (68)	874	12.5 (88)	808	88.2 (62)
6312	523	92.4 (18)	3.4 (24)	587	24.4 (87)	434	24.2 (40)	415	91.1 (42)
6316	327	92.4 (18)	3.4 (24)	360	26.9 (73)	289	22.1 (50)	251	94.4 (13)
6318	21	*	*	22	*	14	*	7	*
6320	95	*	*	111	19.8 (105)	58	*	67	*
6325	84	*	*	102	26.5 (75)	81	*	64	*
6335	294	89.5 (33)	8.5 (80)	308	36.0 (34)	82	*	241	91.7 (36)
6340	0	*	*	1	*	1	*	1	*
6345	27	*	*	30	*	24	*	19	*
6360	288	87.5 (42)	5.2 (48)	327	25.4 (82)	198	29.3 (19)	237	90.7 (45)
7306	654	82.9 (70)	7.0 (71)	725	33.5 (46)	631	16.6 (72)	550	87.3 (70)
7401	616	70.5 (106)	21.8 (108)	681	39.6 (17)	579	22.1 (50)	548	86.1 (79)
7601	169	88.2 (39)	6.5 (64)	204	43.6 (8)	188	48.9 (3)	168	91.7 (36)
7850	289	84.8 (56)	11.8 (90)	328	37.2 (28)	225	24.4 (39)	254	83.5 (94)
7852	8	*	*	9	*	6	*	7	*
8101	510	80.4 (81)	2.7 (17)	535	21.9 (96)	298	14.8 (79)	359	85.2 (84)
8102	131	77.1 (90)	6.9 (70)	137	26.3 (76)	84	*	95	*
8105	191	98.4 (1)	1.0 (6)	233	36.9 (30)	231	18.6 (66)	233	92.3 (30)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 41

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8106	168	76.8 (94)	4.2 (38)	180	21.1 (98)	138	18.1 (68)	113	87.6 (69)
8150	1	*	*	1	*	1	*	1	*
8151	902	91.8 (21)	4.2 (38)	969	28.4 (63)	843	13.3 (86)	690	93.8 (18)
8152	15	*	*	17	*	8	*	17	*
8153	3	*	*	3	*	3	*	3	*
8155	4	*	*	5	*	4	*	5	*
8157	1	*	*	1	*	1	*	1	*
8201	856	92.8 (16)	2.2 (13)	947	17.5 (110)	597	45.4 (6)	562	92.3 (30)
8202	1,292	95.2 (10)	2.9 (20)	1,353	27.2 (69)	796	6.4 (101)	1,029	89.8 (51)
8203	123	94.3 (11)	4.1 (37)	138	22.5 (93)	101	42.6 (7)	109	97.2 (1)
8250	225	88.9 (36)	2.7 (17)	257	16.7 (112)	176	29.0 (20)	187	85.0 (85)
8252	49	*	*	55	*	48	*	44	*
8255	669	93.3 (13)	0.6 (3)	773	19.9 (104)	466	47.6 (5)	539	91.7 (36)
8257	637	97.8 (3)	0.8 (4)	667	27.9 (65)	467	10.1 (92)	511	88.1 (63)
8263	199	98.0 (2)	0.5 (2)	239	19.7 (106)	116	48.3 (4)	173	97.1 (2)
8264	239	97.1 (4)	0.8 (4)	281	25.3 (83)	202	35.1 (12)	195	92.8 (26)
8267	187	88.2 (39)	4.3 (41)	206	20.4 (102)	156	61.5 (2)	149	91.3 (41)
8901	288	83.3 (65)	3.8 (33)	307	18.2 (108)	141	19.1 (64)	201	81.1 (98)
8902	435	84.6 (61)	2.8 (19)	474	20.5 (101)	241	20.3 (62)	322	88.5 (60)
8903	459	76.5 (97)	3.9 (34)	502	23.5 (90)	342	17.8 (69)	341	80.1 (102)
8904	746	80.8 (79)	3.6 (27)	807	20.0 (103)	457	16.8 (71)	545	86.2 (78)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 42

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8905	452	88.9 (36)	2.9 (20)	490	24.5 (86)	363	32.0 (17)	361	87.0 (71)
8909	255	83.5 (63)	2.4 (15)	268	17.2 (111)	182	13.7 (84)	165	87.9 (64)
9045	462	85.5 (53)	7.1 (72)	492	34.1 (41)	254	27.6 (28)	378	87.0 (71)
9050	302	81.5 (76)	5.3 (50)	332	24.4 (87)	263	22.8 (47)	241	85.9 (81)
9051	455	84.8 (56)	6.4 (61)	481	37.8 (25)	441	13.8 (83)	402	90.5 (47)
9053	354	82.5 (72)	2.3 (14)	375	18.7 (107)	288	11.5 (89)	198	92.4 (28)
9055	3	*	*	2	*	2	*	1	*
9058	301	82.1 (74)	5.6 (55)	325	25.5 (80)	240	8.3 (98)	215	86.5 (76)
9066	409	72.6 (102)	5.4 (52)	435	14.7 (114)	289	8.7 (97)	210	84.8 (87)
9068	311	82.3 (73)	5.5 (54)	325	21.5 (97)	255	14.5 (82)	231	88.3 (61)
9069	712	81.9 (75)	3.9 (34)	733	18.0 (109)	397	9.6 (95)	433	86.8 (73)
9070	387	76.7 (96)	6.7 (68)	410	22.2 (95)	277	14.8 (79)	264	84.8 (87)
9072	293	77.1 (90)	8.5 (80)	325	25.5 (80)	315	10.8 (91)	326	87.7 (67)
9080	779	80.9 (77)	4.7 (45)	819	20.9 (99)	560	21.4 (55)	585	85.6 (83)
9082	36	*	*	40	*	35	*	25	*
9083	1	*	*	1	*	0	*	0	*
9084	4	*	*	5	*	5	*	5	*
9087	3	*	*	4	*	4	*	4	*
<b>Michigan</b>	<b>45,167</b>	<b>85.6</b>	<b>6.4</b>	<b>49,385</b>	<b>29.3</b>	<b>38,062</b>	<b>20.3</b>	<b>36,374</b>	<b>36374.0</b>
<b>National (Prior Year)</b>	<b>100,055</b>	<b>77.9</b>	<b>5.0</b>	<b>134,286</b>	<b>21.1</b>	<b>82,984</b>	<b>20.5</b>	<b>79,295</b>	<b>89.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

## 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic

Page : 43

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	35	*	36	*	13	*	31	*	36	*
0102	139	38.8 (99)	135	31.1 (111)	44	*	128	18.8 (98)	135	3.0 (80)
0103	110	36.4 (92)	106	27.4 (108)	28	*	97	*	106	6.6 (110)
0104	36	*	38	*	7	*	33	*	38	*
0105	11	*	11	*	5	*	11	*	11	*
0250	34	*	31	*	11	*	32	*	31	*
0251	23	*	26	*	8	*	21	*	26	*
0252	28	*	28	*	8	*	25	*	28	*
0253	41	*	42	*	13	*	38	*	42	*
0401	139	42.4 (107)	133	25.6 (107)	45	*	127	12.6 (77)	132	1.5 (52)
0402	102	35.3 (83)	105	19.0 (84)	31	*	92	*	106	4.7 (104)
0403	46	*	47	*	17	*	39	*	48	*
0404	38	*	40	*	9	*	30	*	40	*
0501	135	35.6 (85)	149	24.2 (103)	41	*	116	19.8 (100)	149	3.4 (91)
0503	25	*	31	*	7	*	23	*	31	*
0504	124	40.3 (103)	125	17.6 (76)	36	*	96	*	125	0.8 (27)
0505	83	*	82	*	25	*	72	*	82	*
0506	45	*	43	*	13	*	41	*	43	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Page : 44

Michigan

## Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0507	16	*	20	*	6	*	14	*	20	*
0508	9	*	10	*	3	*	8	*	10	*
0650	318	32.4 (78)	309	18.4 (82)	85	*	275	8.7 (50)	308	2.6 (74)
0651	165	44.8 (109)	169	33.1 (112)	64	*	153	21.6 (102)	169	1.2 (42)
0652	86	*	92	*	31	*	73	*	92	*
0653	143	38.5 (98)	144	23.6 (100)	43	*	122	18.9 (99)	144	3.5 (94)
0654	119	34.5 (81)	123	18.7 (83)	37	*	106	16.0 (93)	123	4.1 (99)
0655	86	*	86	*	23	*	73	*	86	*
0657	18	*	18	*	10	*	15	*	18	*
0701	134	37.3 (95)	140	23.6 (100)	40	*	119	4.2 (14)	140	0.7 (23)
0702	24	*	28	*	5	*	20	*	28	*
0703	43	*	47	*	14	*	32	*	47	*
0704	20	*	18	*	5	*	19	*	18	*
0705	30	*	28	*	12	*	30	*	28	*
0706	3	*	3	*	1	*	3	*	3	*
0850	210	40.0 (101)	218	18.3 (80)	71	*	174	6.9 (32)	218	1.8 (57)
0851	191	38.2 (96)	198	19.7 (87)	52	*	149	10.7 (64)	198	0.5 (17)
0901	462	40.3 (103)	463	22.2 (98)	160	30.0 (14)	415	12.8 (78)	463	4.1 (99)
0902	12	*	9	*	2	*	8	*	9	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Page : 45

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1051	325	36.3 (90)	348	17.0 (75)	89	*	255	9.8 (59)	348	3.2 (88)
1101	448	27.2 (63)	467	13.3 (61)	89	*	371	14.6 (88)	467	3.0 (80)
1102	250	37.2 (94)	230	18.3 (80)	59	*	209	12.0 (70)	231	2.2 (66)
1106	26	*	23	*	3	*	21	*	23	*
1250	312	27.2 (63)	315	13.3 (61)	68	*	264	9.1 (55)	315	0.3 (9)
1251	189	42.9 (108)	204	27.5 (109)	67	*	164	14.0 (84)	203	1.5 (52)
1252	154	26.0 (61)	150	14.0 (67)	32	*	136	8.8 (52)	150	2.7 (77)
1253	219	36.1 (89)	230	17.8 (77)	68	*	194	8.2 (45)	228	2.2 (66)
1350	648	27.6 (66)	675	13.9 (66)	132	34.8 (10)	550	7.6 (39)	676	1.3 (49)
1351	26	*	21	*	9	*	26	*	22	*
1352	130	24.6 (58)	129	12.4 (57)	30	*	107	2.8 (6)	130	0.0 (1)
1401	59	*	59	*	21	*	49	*	59	*
1402	34	*	36	*	10	*	27	*	36	*
1403	5	*	5	*	1	*	3	*	5	*
1501	226	17.7 (35)	234	7.7 (40)	30	*	195	4.1 (12)	235	0.4 (13)
1503	251	24.7 (59)	271	14.0 (67)	56	*	219	4.6 (18)	271	0.4 (13)
1504	586	15.4 (25)	593	7.4 (37)	63	*	484	2.5 (5)	594	0.3 (9)
1505	108	21.3 (47)	96	*	20	*	103	4.9 (21)	96	*
1506	170	24.1 (57)	170	15.3 (72)	31	*	148	7.4 (38)	170	1.8 (57)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Page : 46

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1507	150	30.0 (71)	156	9.0 (44)	32	*	120	10.0 (60)	155	3.9 (97)
1508	6	*	4	*	0	*	6	*	4	*
1509	108	19.4 (41)	107	13.1 (59)	17	*	94	*	107	1.9 (61)
1511	172	38.4 (97)	178	19.1 (85)	53	*	149	4.7 (19)	178	2.2 (66)
1512	93	*	100	7.0 (33)	11	*	69	*	100	1.0 (38)
1514	98	*	113	7.1 (34)	5	*	37	*	110	0.9 (33)
1701	180	33.9 (80)	172	23.8 (102)	52	*	155	12.3 (73)	172	3.5 (94)
1950	190	41.6 (105)	194	24.2 (103)	59	*	157	21.0 (101)	194	5.2 (106)
1952	43	*	50	*	8	*	38	*	50	*
1953	74	*	74	*	19	*	64	*	74	*
1954	137	31.4 (74)	137	21.2 (94)	37	*	126	14.3 (86)	138	0.7 (23)
1955	30	*	32	*	9	*	25	*	32	*
1957	94	*	104	14.4 (70)	27	*	87	*	104	0.0 (1)
1963	12	*	16	*	0	*	9	*	16	*
1964	1	*	1	*	0	*	1	*	1	*
2150	192	45.3 (110)	191	25.1 (106)	66	*	169	12.4 (74)	191	1.6 (54)
2151	45	*	48	*	9	*	31	*	48	*
2152	2	*	4	*	0	*	0	*	4	*
2251	120	35.8 (88)	115	24.3 (105)	34	*	99	*	115	0.9 (33)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Page : 47

Michigan

## Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2252	52	*	49	*	17	*	47	*	48	*
2554	8	*	11	*	5	*	6	*	11	*
2555	2,046	22.5 (51)	2,162	12.9 (58)	356	37.4 (8)	1,713	8.2 (45)	2,159	2.5 (73)
2556	70	*	83	*	14	*	62	*	84	*
2557	0	*	1	*	0	*	0	*	1	*
2558	1	*	0	*	0	*	1	*	0	*
2560	2	*	1	*	0	*	2	*	1	*
2561	1	*	1	*	1	*	1	*	1	*
2562	3	*	3	*	1	*	3	*	3	*
2563	4	*	6	*	0	*	3	*	6	*
2564	14	*	11	*	4	*	12	*	11	*
3001	334	31.4 (74)	363	11.6 (54)	91	*	304	11.8 (69)	364	4.1 (99)
3002	86	*	84	*	17	*	82	*	83	*
3004	65	*	63	*	6	*	51	*	64	*
3005	46	*	43	*	12	*	40	*	43	*
3007	3	*	3	*	0	*	2	*	3	*
3201	128	35.2 (82)	142	21.1 (93)	39	*	100	13.0 (81)	142	4.9 (105)
3203	1	*	1	*	0	*	1	*	1	*
3301	1,223	17.2 (30)	1,307	10.8 (51)	165	32.7 (12)	1,034	4.4 (15)	1,306	1.7 (55)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

## 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic

Page : 48

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3304	26	*	24	*	1	*	19	*	24	*
3305	46	*	48	*	2	*	35	*	47	*
3306	77	*	86	*	13	*	66	*	85	*
3308	38	*	43	*	8	*	24	*	43	*
3312	100	25.0 (60)	102	10.8 (51)	21	*	90	*	101	1.0 (38)
3550	23	*	23	*	11	*	22	*	23	*
3551	4	*	3	*	2	*	4	*	3	*
3552	13	*	11	*	8	*	13	*	11	*
3701	271	28.0 (67)	283	20.1 (88)	61	*	219	5.0 (22)	286	2.4 (70)
3850	861	35.3 (83)	917	21.8 (96)	265	30.2 (13)	772	8.2 (45)	917	1.2 (42)
3901	447	23.0 (52)	443	10.6 (50)	86	*	396	7.1 (35)	440	1.8 (57)
3902	32	*	22	*	4	*	32	*	22	*
3903	5	*	5	*	0	*	4	*	5	*
3950	619	23.4 (55)	613	11.1 (53)	100	37.0 (9)	509	6.1 (29)	612	0.8 (27)
3954	5	*	5	*	1	*	5	*	5	*
4106	616	16.2 (27)	681	6.2 (26)	76	*	535	3.2 (9)	681	0.1 (7)
4107	856	21.6 (48)	879	10.5 (48)	124	50.8 (3)	681	8.1 (44)	877	1.1 (41)
4109	217	26.7 (62)	224	13.4 (63)	46	*	182	6.0 (28)	223	0.0 (1)
4110	2	*	2	*	0	*	0	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

## 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic

Page : 49

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4112	162	4.3 (4)	161	0.0 (1)	6	*	117	12.0 (70)	155	1.9 (61)
4115	1,027	11.8 (14)	1,040	4.5 (13)	94	*	830	2.0 (4)	1,033	0.4 (13)
4351	164	27.4 (65)	162	15.4 (74)	38	*	146	9.6 (57)	162	1.9 (61)
4352	46	*	48	*	24	*	40	*	48	*
4353	136	30.1 (72)	128	14.1 (69)	28	*	111	9.0 (53)	128	3.1 (84)
4354	205	31.7 (76)	212	22.2 (98)	54	*	188	12.8 (78)	212	2.4 (70)
4355	84	*	91	*	32	*	72	*	91	*
4356	121	42.1 (106)	127	29.9 (110)	41	*	104	12.5 (76)	127	2.4 (70)
4358	23	*	22	*	1	*	23	*	22	*
4359	162	19.1 (40)	167	8.4 (42)	29	*	130	9.2 (56)	168	1.2 (42)
4360	61	*	57	*	14	*	53	*	56	*
4361	81	*	77	*	24	*	69	*	76	*
4362	84	*	90	*	33	*	75	*	90	*
4363	222	28.4 (68)	243	20.2 (89)	51	*	193	16.1 (95)	242	5.4 (107)
4401	236	35.6 (85)	247	17.8 (77)	59	*	198	6.6 (31)	247	2.0 (64)
4501	253	35.6 (85)	259	20.8 (91)	73	*	215	11.6 (68)	259	1.2 (42)
4601	385	30.6 (73)	399	13.8 (65)	90	*	327	12.2 (72)	399	3.0 (80)
4701	290	31.7 (76)	295	21.4 (95)	69	*	221	5.9 (27)	295	0.7 (23)
5050	457	21.2 (46)	465	8.4 (42)	77	*	349	10.3 (63)	465	0.9 (33)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Page : 50

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
5051	819	16.2 (27)	849	6.6 (30)	97	*	631	3.6 (10)	848	1.4 (50)
5052	812	17.4 (32)	881	7.4 (37)	108	45.4 (6)	670	5.2 (24)	881	0.9 (33)
5055	383	1.8 (2)	392	1.3 (5)	5	*	330	4.5 (17)	392	1.0 (38)
5101	480	16.5 (29)	495	10.1 (47)	65	*	378	6.9 (32)	497	0.4 (13)
5102	12	*	12	*	2	*	10	*	12	*
5202	215	33.5 (79)	212	20.8 (91)	66	*	179	10.1 (62)	212	1.4 (50)
5206	2	*	5	*	0	*	0	*	5	*
5601	273	29.7 (70)	268	15.3 (72)	64	*	228	7.0 (34)	265	0.8 (27)
5801	539	36.7 (93)	549	18.0 (79)	149	43.6 (7)	442	12.9 (80)	549	5.8 (109)
5803	34	*	30	*	8	*	28	*	30	*
5804	7	*	7	*	1	*	4	*	7	*
5806	17	*	19	*	8	*	14	*	19	*
6101	10	*	9	*	3	*	9	*	9	*
6102	27	*	31	*	6	*	26	*	31	*
6103	955	29.6 (69)	1,034	19.1 (85)	225	28.4 (15)	838	10.7 (64)	1,038	3.3 (89)
6105	4	*	4	*	0	*	2	*	4	*
6106	31	*	25	*	8	*	31	*	26	*
6301	986	10.9 (11)	1,001	6.7 (31)	84	*	806	6.2 (30)	1,001	0.7 (23)
6312	530	5.7 (5)	525	1.1 (3)	18	*	414	3.1 (8)	524	0.2 (8)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

## 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic

Page : 51

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
6316	333	14.4 (20)	326	7.1 (34)	36	*	252	4.4 (15)	326	0.3 (9)
6318	14	*	21	*	1	*	7	*	21	*
6320	102	8.8 (7)	100	2.0 (6)	5	*	67	*	99	*
6325	97	*	85	*	17	*	63	*	84	*
6335	295	7.5 (6)	296	3.7 (9)	16	*	241	1.7 (3)	296	0.3 (9)
6340	1	*	0	*	0	*	1	*	0	*
6345	29	*	27	*	3	*	19	*	27	*
6360	292	15.1 (23)	298	7.4 (37)	36	*	235	7.2 (36)	300	0.0 (1)
7306	633	36.3 (90)	664	13.6 (64)	181	54.7 (2)	550	11.5 (67)	663	3.3 (89)
7401	614	39.7 (100)	619	21.8 (96)	197	33.5 (11)	548	14.1 (85)	618	3.4 (91)
7601	189	19.6 (43)	171	15.2 (71)	27	*	166	7.8 (40)	171	0.6 (21)
7850	301	40.2 (102)	294	20.7 (90)	93	*	252	13.1 (82)	293	7.8 (111)
7852	7	*	8	*	2	*	7	*	8	*
8101	430	14.9 (22)	521	5.4 (21)	51	*	357	9.0 (53)	520	2.9 (78)
8102	112	10.7 (9)	130	4.6 (14)	9	*	93	*	131	0.8 (27)
8105	233	9.9 (8)	190	3.2 (7)	18	*	232	0.9 (2)	189	0.5 (17)
8106	147	23.1 (53)	169	11.8 (55)	27	*	113	8.0 (43)	169	0.6 (21)
8150	1	*	1	*	0	*	1	*	1	*
8151	845	17.9 (36)	902	6.9 (32)	121	47.9 (5)	687	5.4 (25)	906	0.8 (27)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Page : 52

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8152	17	*	15	*	6	*	17	*	15	*
8153	3	*	3	*	1	*	3	*	3	*
8155	5	*	4	*	1	*	5	*	4	*
8157	1	*	1	*	0	*	1	*	1	*
8201	793	17.2 (30)	858	7.1 (34)	102	50.0 (4)	557	5.7 (26)	863	1.2 (42)
8202	1,226	3.8 (3)	1,315	1.2 (4)	38	*	1,028	5.0 (22)	1,316	1.2 (42)
8203	124	21.8 (49)	122	13.1 (59)	17	*	109	7.3 (37)	123	0.8 (27)
8250	222	19.4 (41)	224	5.4 (21)	25	*	187	11.2 (66)	226	0.9 (33)
8252	48	*	51	*	8	*	44	*	51	*
8255	676	20.9 (45)	674	12.2 (56)	92	*	542	7.9 (41)	676	1.8 (57)
8257	621	0.8 (1)	650	0.9 (2)	3	*	510	0.4 (1)	648	1.2 (42)
8263	219	23.3 (54)	199	8.0 (41)	30	*	172	4.1 (12)	198	0.5 (17)
8264	261	13.4 (19)	239	5.0 (18)	26	*	194	3.6 (10)	239	0.0 (1)
8267	182	12.1 (16)	191	9.4 (46)	14	*	150	4.7 (19)	192	0.5 (17)
8901	241	14.5 (21)	289	9.3 (45)	30	*	201	10.0 (60)	287	3.8 (96)
8902	383	13.3 (18)	438	4.6 (14)	40	*	323	8.4 (48)	438	3.4 (91)
8903	414	15.5 (26)	468	4.9 (17)	43	*	344	16.0 (93)	469	2.6 (74)
8904	675	17.6 (33)	752	6.4 (28)	88	*	547	12.4 (74)	752	4.4 (103)
8905	449	12.0 (15)	459	3.7 (9)	45	*	356	8.7 (50)	458	2.2 (66)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Page : 53

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8909	218	18.8 (39)	256	5.1 (20)	32	*	165	17.6 (97)	256	3.9 (97)
9045	436	11.0 (12)	465	4.7 (16)	41	*	378	7.9 (41)	463	1.7 (55)
9050	273	19.8 (44)	312	6.4 (28)	54	*	242	16.1 (95)	311	2.9 (78)
9051	447	11.2 (13)	465	3.2 (7)	43	*	402	3.0 (7)	465	0.0 (1)
9053	306	10.8 (10)	355	3.9 (11)	27	*	197	13.2 (83)	355	2.0 (64)
9055	2	*	3	*	0	*	1	*	3	*
9058	272	12.9 (17)	302	6.3 (27)	27	*	209	8.6 (49)	302	3.0 (80)
9066	295	17.6 (33)	419	5.7 (23)	33	*	210	14.8 (89)	418	3.1 (84)
9068	269	21.9 (50)	313	10.5 (48)	48	*	232	15.5 (91)	313	4.2 (102)
9069	602	15.1 (23)	712	6.0 (24)	63	*	430	15.6 (92)	710	5.4 (107)
9070	346	17.9 (36)	386	6.0 (24)	49	*	257	14.4 (87)	383	2.6 (74)
9072	326	23.6 (56)	294	4.1 (12)	68	*	327	15.3 (90)	294	3.1 (84)
9080	694	18.2 (38)	786	5.0 (18)	104	59.6 (1)	584	9.6 (57)	786	3.1 (84)
9082	36	*	37	*	2	*	26	*	37	*
9083	0	*	1	*	0	*	0	*	1	*
9084	5	*	4	*	1	*	5	*	4	*
9087	4	*	3	*	0	*	4	*	3	*
Michigan	43,650	22.6	45,656	11.5	7,693	39.4	36,252	8.6	45,631	2.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Page : 54

### Michigan

#### Comparison of Smoking Indicators by Clinic

National (Prior Year)	126,132	26.9	133,382	14.1	21,263	35.7	97,728	11.9	118,446	6.6
-----------------------	---------	------	---------	------	--------	------	--------	------	---------	-----

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 55

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	35	*	*	36	*	35	*	35	*
0102	124	11.3 (71)	8.1 (76)	125	8.8 (14)	110	8.2 (92)	124	62.1 (72)
0103	101	6.9 (16)	11.9 (105)	104	7.7 (7)	93	*	101	63.4 (69)
0104	33	*	*	36	*	31	*	33	*
0105	9	*	*	10	*	8	*	9	*
0250	31	*	*	30	*	26	*	31	*
0251	24	*	*	24	*	20	*	24	*
0252	28	*	*	28	*	27	*	28	*
0253	42	*	*	42	*	36	*	42	*
0401	128	2.3 (1)	7.8 (71)	129	7.8 (8)	118	0.0 (1)	128	73.4 (41)
0402	102	2.9 (2)	9.8 (96)	102	8.8 (14)	91	*	102	78.4 (24)
0403	44	*	*	43	*	39	*	44	*
0404	39	*	*	39	*	32	*	39	*
0501	136	14.0 (92)	9.6 (92)	149	12.8 (64)	117	6.8 (85)	147	70.7 (51)
0503	30	*	*	29	*	26	*	31	*
0504	114	7.9 (25)	5.3 (33)	119	10.1 (29)	101	3.0 (22)	118	89.8 (1)
0505	78	*	*	77	*	66	*	78	*
0506	39	*	*	39	*	36	*	39	*
0507	17	*	*	19	*	15	*	17	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 56

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	10	*	*	10	*	9	*	10	*
0650	297	9.8 (50)	10.8 (104)	303	12.5 (60)	260	5.4 (68)	298	71.1 (48)
0651	159	4.4 (5)	8.2 (77)	165	6.7 (4)	149	2.7 (14)	158	71.5 (46)
0652	90	*	*	90	*	82	*	90	*
0653	131	7.6 (23)	9.9 (98)	134	9.7 (25)	117	2.6 (13)	132	82.6 (11)
0654	115	10.4 (58)	7.0 (64)	117	10.3 (32)	102	6.9 (86)	116	73.3 (42)
0655	81	*	*	81	*	71	*	81	*
0657	18	*	*	18	*	14	*	18	*
0701	131	9.2 (46)	13.0 (106)	137	8.0 (9)	119	5.9 (75)	132	84.8 (8)
0702	27	*	*	27	*	25	*	27	*
0703	43	*	*	45	*	38	*	43	*
0704	18	*	*	18	*	15	*	18	*
0705	27	*	*	27	*	23	*	27	*
0706	3	*	*	3	*	3	*	3	*
0850	207	10.1 (56)	7.2 (67)	215	11.2 (44)	183	3.3 (28)	211	85.8 (6)
0851	182	10.4 (58)	5.5 (35)	194	10.8 (42)	162	4.9 (60)	186	80.1 (18)
0901	440	10.5 (64)	9.3 (88)	454	13.4 (73)	381	4.2 (51)	445	64.7 (65)
0902	8	*	*	8	*	8	*	8	*
1051	322	8.1 (29)	7.1 (66)	331	9.1 (18)	291	3.4 (30)	324	81.5 (15)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 57

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1101	447	8.7 (38)	4.3 (17)	454	10.6 (38)	393	5.6 (72)	448	68.8 (55)
1102	216	5.1 (9)	6.0 (46)	219	5.0 (1)	199	3.5 (32)	216	70.8 (50)
1106	23	*	*	23	*	19	*	23	*
1250	301	9.0 (44)	8.0 (74)	306	8.5 (10)	268	4.1 (47)	304	72.4 (44)
1251	197	8.6 (37)	8.6 (83)	199	10.1 (29)	176	4.0 (44)	198	77.8 (25)
1252	140	7.1 (17)	9.3 (88)	141	8.5 (10)	126	3.2 (25)	142	87.3 (3)
1253	216	4.6 (6)	7.9 (72)	223	9.9 (27)	193	1.0 (3)	218	86.2 (5)
1350	656	10.4 (58)	6.6 (55)	662	10.7 (39)	582	4.6 (56)	659	71.8 (45)
1351	20	*	*	19	*	18	*	20	*
1352	120	13.3 (85)	10.0 (99)	125	10.4 (33)	106	6.6 (84)	125	61.6 (73)
1401	52	*	*	56	*	46	*	52	*
1402	35	*	*	35	*	28	*	35	*
1403	4	*	*	5	*	3	*	4	*
1501	213	8.5 (36)	10.3 (100)	225	9.8 (26)	189	3.7 (38)	224	70.5 (52)
1503	253	6.3 (12)	8.3 (79)	256	12.1 (57)	219	4.1 (47)	254	73.6 (40)
1504	424	7.5 (22)	6.1 (48)	579	11.6 (51)	366	2.7 (14)	577	80.1 (18)
1505	87	*	*	91	*	75	*	89	*
1506	162	7.4 (20)	4.9 (26)	163	13.5 (75)	140	2.1 (9)	162	74.1 (37)
1507	143	8.4 (34)	9.8 (96)	149	11.4 (47)	127	5.5 (70)	143	86.7 (4)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 58

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1508	4	*	*	4	*	3	*	4	*
1509	98	*	*	103	15.5 (95)	83	*	99	*
1511	163	8.0 (27)	5.5 (35)	171	10.5 (36)	144	4.2 (51)	168	79.2 (22)
1512	97	*	*	100	12.0 (54)	86	*	97	*
1514	106	17.9 (108)	8.5 (82)	109	22.0 (111)	82	*	109	85.3 (7)
1701	160	8.8 (39)	13.8 (108)	160	13.8 (80)	134	3.7 (38)	160	66.3 (61)
1950	182	10.4 (58)	6.6 (55)	188	12.2 (58)	158	3.8 (41)	183	82.5 (12)
1952	46	*	*	47	*	43	*	46	*
1953	74	*	*	72	*	64	*	74	*
1954	136	6.6 (14)	6.6 (55)	135	10.4 (33)	120	1.7 (6)	136	76.5 (31)
1955	29	*	*	29	*	25	*	29	*
1957	100	14.0 (92)	13.0 (106)	103	14.6 (86)	86	*	101	79.2 (22)
1963	14	*	*	15	*	14	*	14	*
1964	1	*	*	1	*	1	*	1	*
2150	175	5.1 (9)	9.1 (87)	187	9.6 (24)	158	1.9 (7)	176	82.4 (13)
2151	44	*	*	46	*	41	*	44	*
2152	4	*	*	3	*	3	*	4	*
2251	113	9.7 (49)	6.2 (51)	114	15.8 (96)	95	*	114	71.1 (48)
2252	47	*	*	46	*	42	*	47	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 59

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2554	9	*	*	10	*	8	*	9	*
2555	2,083	11.2 (70)	4.4 (19)	2,117	12.5 (60)	1,808	5.4 (68)	2,087	55.8 (86)
2556	78	*	*	82	*	60	*	78	*
2557	1	*	*	1	*	0	*	1	*
2558	0	*	*	0	*	0	*	0	*
2560	1	*	*	1	*	1	*	1	*
2561	1	*	*	1	*	1	*	1	*
2562	3	*	*	3	*	2	*	3	*
2563	6	*	*	6	*	6	*	6	*
2564	11	*	*	11	*	11	*	11	*
3001	348	13.5 (87)	4.0 (12)	353	13.3 (72)	300	6.0 (77)	348	58.3 (79)
3002	81	*	*	82	*	73	*	81	*
3004	57	*	*	60	*	47	*	59	*
3005	40	*	*	40	*	35	*	40	*
3007	3	*	*	3	*	3	*	3	*
3201	135	10.4 (58)	10.4 (101)	141	13.5 (75)	119	5.0 (61)	135	77.0 (27)
3203	0	*	*	0	*	0	*	0	*
3301	1,235	11.5 (73)	5.2 (32)	1,271	13.5 (75)	1,067	4.6 (56)	1,241	72.9 (43)
3304	22	*	*	22	*	19	*	22	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 60

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3305	47	*	*	48	*	42	*	48	*
3306	83	*	*	85	*	74	*	83	*
3308	40	*	*	40	*	34	*	39	*
3312	97	*	*	100	15.0 (90)	82	*	98	*
3550	22	*	*	22	*	18	*	22	*
3551	3	*	*	3	*	3	*	3	*
3552	10	*	*	11	*	9	*	10	*
3701	277	7.2 (18)	9.7 (94)	281	8.9 (16)	252	3.6 (33)	276	83.3 (9)
3850	872	8.8 (39)	9.4 (90)	890	10.8 (42)	781	4.7 (58)	875	69.3 (54)
3901	419	8.8 (39)	4.8 (24)	434	12.0 (54)	364	4.1 (47)	426	60.3 (75)
3902	22	*	*	22	*	18	*	22	*
3903	5	*	*	5	*	4	*	5	*
3950	569	10.2 (57)	6.9 (61)	602	10.0 (28)	508	2.8 (19)	581	67.1 (59)
3954	5	*	*	5	*	5	*	5	*
4106	644	13.2 (84)	4.7 (22)	663	14.8 (88)	550	5.8 (74)	652	74.2 (36)
4107	827	13.5 (87)	5.7 (39)	865	16.5 (100)	688	5.5 (70)	842	76.2 (32)
4109	214	9.8 (50)	6.5 (53)	221	15.4 (94)	182	2.7 (14)	214	79.9 (20)
4110	2	*	*	2	*	0	*	2	*
4112	147	8.2 (30)	7.5 (69)	150	12.7 (63)	128	3.9 (42)	148	16.9 (108)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 61

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4115	978	13.3 (85)	5.4 (34)	1,022	14.5 (85)	834	5.3 (66)	991	77.6 (26)
4351	155	8.4 (34)	6.5 (53)	159	11.9 (53)	137	3.6 (33)	155	81.9 (14)
4352	46	*	*	48	*	41	*	46	*
4353	120	10.0 (53)	5.8 (41)	120	17.5 (105)	96	*	122	79.5 (21)
4354	205	7.8 (24)	6.8 (60)	207	8.7 (12)	187	3.2 (25)	206	73.8 (38)
4355	81	*	*	83	*	69	*	83	*
4356	114	12.3 (78)	7.0 (64)	117	13.7 (78)	96	*	114	73.7 (39)
4358	20	*	*	21	*	20	*	20	*
4359	154	3.9 (3)	8.4 (80)	156	7.1 (6)	142	1.4 (4)	155	74.8 (34)
4360	45	*	*	49	*	39	*	45	*
4361	61	*	*	77	*	54	*	62	*
4362	87	*	*	86	*	78	*	87	*
4363	227	7.9 (25)	9.7 (94)	227	12.8 (64)	194	2.1 (9)	228	80.7 (16)
4401	239	10.5 (64)	10.5 (102)	238	13.9 (81)	202	5.0 (61)	239	80.3 (17)
4501	248	8.9 (43)	6.9 (61)	251	10.4 (33)	222	4.5 (55)	248	75.0 (33)
4601	385	4.9 (7)	8.8 (85)	393	5.9 (2)	362	2.8 (19)	385	65.5 (62)
4701	283	9.2 (46)	6.0 (46)	287	12.5 (60)	247	2.0 (8)	282	83.0 (10)
5050	449	12.0 (75)	6.2 (51)	453	15.2 (93)	377	4.0 (44)	450	58.2 (80)
5051	819	10.4 (58)	4.3 (17)	829	11.5 (48)	714	4.2 (51)	821	61.5 (74)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 62

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5052	852	13.7 (89)	4.9 (26)	858	14.8 (88)	713	6.0 (77)	856	57.9 (82)
5055	350	7.4 (20)	4.9 (26)	360	6.1 (3)	330	3.9 (42)	355	48.7 (98)
5101	482	8.3 (32)	7.5 (69)	485	9.3 (20)	432	3.2 (25)	483	69.8 (53)
5102	12	*	*	12	*	11	*	12	*
5202	203	4.9 (7)	7.9 (72)	202	6.9 (5)	187	2.7 (14)	203	87.7 (2)
5206	5	*	*	4	*	4	*	5	*
5601	261	6.5 (13)	9.6 (92)	264	9.1 (18)	235	2.1 (9)	261	77.0 (27)
5801	527	6.6 (14)	7.2 (67)	534	11.2 (44)	466	1.5 (5)	527	65.5 (62)
5803	30	*	*	30	*	26	*	30	*
5804	6	*	*	7	*	5	*	6	*
5806	17	*	*	18	*	15	*	17	*
6101	9	*	*	8	*	6	*	9	*
6102	31	*	*	31	*	30	*	31	*
6103	993	11.0 (67)	5.1 (30)	994	12.8 (64)	849	5.9 (75)	996	64.1 (66)
6105	4	*	*	4	*	4	*	4	*
6106	25	*	*	26	*	22	*	25	*
6301	945	12.8 (83)	5.8 (41)	986	13.7 (78)	805	5.1 (64)	963	68.4 (57)
6312	485	11.1 (68)	5.6 (37)	517	12.2 (58)	425	3.3 (28)	494	76.9 (29)
6316	298	11.1 (68)	6.7 (58)	319	15.0 (90)	252	3.6 (33)	304	76.6 (30)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 63

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6318	18	*	*	20	*	17	*	19	*
6320	90	*	*	95	*	73	*	92	*
6325	83	*	*	85	*	76	*	83	*
6335	281	10.0 (53)	3.9 (10)	280	13.9 (81)	241	4.1 (47)	281	71.5 (46)
6340	0	*	*	0	*	0	*	0	*
6345	25	*	*	27	*	23	*	25	*
6360	290	12.4 (80)	5.9 (44)	293	15.0 (90)	244	6.1 (81)	292	68.5 (56)
7306	625	12.2 (76)	6.1 (48)	652	13.0 (67)	545	4.4 (54)	633	62.9 (70)
7401	594	9.9 (52)	6.7 (58)	604	11.6 (51)	526	4.0 (44)	594	64.0 (67)
7601	166	5.4 (11)	8.4 (80)	170	9.4 (22)	150	2.7 (14)	168	49.4 (96)
7850	279	8.2 (30)	9.0 (86)	287	11.5 (48)	246	2.8 (19)	280	68.2 (58)
7852	8	*	*	8	*	7	*	8	*
8101	485	16.5 (104)	2.5 (3)	489	19.0 (110)	374	8.3 (93)	490	42.9 (105)
8102	124	7.3 (19)	10.5 (102)	128	10.2 (31)	110	3.6 (33)	126	46.8 (102)
8105	148	4.1 (4)	4.1 (13)	189	9.0 (17)	138	0.7 (2)	181	54.1 (90)
8106	160	14.4 (96)	6.9 (61)	167	17.4 (103)	134	3.0 (22)	163	58.9 (77)
8150	1	*	*	1	*	1	*	1	*
8151	864	12.4 (80)	4.4 (19)	881	16.2 (98)	715	3.6 (33)	873	74.7 (35)
8152	14	*	*	15	*	14	*	14	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2020 Pregnancy Nutrition Surveillance (1)

Page : 64

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8153	3	*	*	2	*	2	*	3	*
8155	4	*	*	4	*	3	*	4	*
8157	1	*	*	1	*	0	*	1	*
8201	807	12.4 (80)	5.6 (37)	845	13.0 (67)	696	6.0 (77)	827	54.2 (89)
8202	1,193	9.1 (45)	4.8 (24)	1,267	9.3 (20)	1,076	5.0 (61)	1,249	46.1 (103)
8203	115	12.2 (76)	8.7 (84)	119	16.8 (101)	93	*	119	65.5 (62)
8250	217	16.6 (105)	5.1 (30)	220	13.2 (69)	184	8.7 (97)	221	50.7 (93)
8252	47	*	*	49	*	41	*	49	*
8255	645	11.8 (74)	5.7 (39)	670	10.7 (39)	574	5.7 (73)	662	56.2 (84)
8257	561	13.7 (89)	2.1 (2)	626	10.5 (36)	491	8.4 (94)	606	58.9 (77)
8263	187	8.0 (27)	8.0 (74)	195	8.7 (12)	171	4.7 (58)	190	57.9 (82)
8264	226	8.8 (39)	5.8 (41)	232	13.4 (73)	197	3.0 (22)	228	62.7 (71)
8267	180	8.3 (32)	9.4 (90)	187	10.7 (39)	161	2.5 (12)	184	52.7 (92)
8901	270	17.0 (107)	1.9 (1)	285	18.9 (109)	219	9.1 (99)	268	43.3 (104)
8902	379	14.0 (92)	2.6 (5)	428	17.1 (102)	320	5.3 (66)	416	55.8 (86)
8903	426	14.6 (98)	4.2 (15)	450	18.2 (108)	341	8.5 (95)	428	50.2 (95)
8904	716	15.2 (100)	4.5 (21)	730	17.7 (106)	579	7.1 (88)	719	48.7 (98)
8905	391	9.5 (48)	6.1 (48)	443	14.4 (84)	328	3.7 (38)	444	63.5 (68)
8909	236	11.4 (72)	3.8 (8)	243	11.5 (48)	205	6.3 (82)	243	54.3 (88)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic

Page : 65

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9045	441	10.0 (53)	8.2 (77)	455	13.2 (69)	380	3.4 (30)	443	47.2 (101)
9050	275	14.9 (99)	4.7 (22)	306	16.3 (99)	227	7.0 (87)	292	50.7 (93)
9051	440	10.5 (64)	5.9 (44)	437	14.6 (86)	364	5.2 (65)	443	67.0 (60)
9053	336	13.7 (89)	3.9 (10)	346	11.3 (46)	294	8.5 (95)	338	59.2 (76)
9055	2	*	*	2	*	1	*	2	*
9058	282	14.2 (95)	2.5 (3)	295	9.5 (23)	253	8.7 (97)	296	58.1 (81)
9066	390	16.9 (106)	3.8 (8)	409	17.4 (103)	322	7.5 (90)	396	49.2 (97)
9068	301	15.6 (102)	5.0 (29)	303	13.2 (69)	258	9.3 (100)	301	40.5 (106)
9069	669	15.8 (103)	3.0 (6)	699	16.0 (97)	558	7.5 (90)	671	35.0 (107)
9070	367	12.3 (78)	4.1 (13)	376	12.0 (54)	317	6.0 (77)	368	53.5 (91)
9072	275	15.3 (101)	3.3 (7)	292	17.8 (107)	231	6.5 (83)	281	55.9 (85)
9080	710	14.5 (97)	4.2 (15)	765	14.2 (83)	603	7.3 (89)	756	47.6 (100)
9082	35	*	*	37	*	30	*	35	*
9083	1	*	*	1	*	1	*	1	*
9084	4	*	*	4	*	4	*	4	*
9087	3	*	*	3	*	3	*	3	*
<b>Michigan</b>	<b>42,825</b>	<b>10.8</b>	<b>6.1</b>	<b>44,372</b>	<b>12.6</b>	<b>37,125</b>	<b>4.8</b>	<b>43,607</b>	<b>64.8</b>
<b>National (Prior Year)</b>	<b>99,198</b>	<b>9.7</b>	<b>6.5</b>	<b>100,365</b>	<b>12.4</b>	<b>84,851</b>	<b>4.1</b>	<b>99,674</b>	<b>52.7</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 66

### Michigan

#### Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
0101	117	91.5	0.9	2.6	0.0	0.0	5.1	0.0
0102	490	89.6	0.8	2.2	0.6	0.2	6.3	0.2
0103	407	91.2	0.0	3.7	0.5	0.0	3.9	0.7
0104	139	93.5	0.0	3.6	0.0	0.0	2.9	0.0
0105	20	*	*	*	*	*	*	*
0250	146	83.6	0.7	2.1	0.0	0.0	13.7	0.0
0251	89	*	*	*	*	*	*	*
0252	103	48.5	1.0	1.9	1.0	0.0	47.6	0.0
0253	136	78.7	0.0	0.0	4.4	0.0	16.9	0.0
0401	462	95.0	0.4	1.5	0.2	0.2	2.4	0.2
0402	386	88.6	1.0	2.8	0.3	0.0	7.3	0.0
0403	157	95.5	0.6	1.9	0.6	0.0	0.6	0.6
0404	112	89.3	1.8	0.9	0.9	0.0	7.1	0.0
0501	492	95.9	0.4	1.8	0.0	0.4	1.4	0.0
0503	109	89.0	0.0	4.6	0.0	0.9	5.5	0.0
0504	457	88.4	1.1	2.6	2.0	0.2	5.7	0.0
0505	297	84.5	0.7	4.7	0.7	0.0	9.4	0.0
0506	173	93.1	0.6	1.2	0.0	0.6	4.6	0.0
0507	76	*	*	*	*	*	*	*
0508	29	*	*	*	*	*	*	*
0509	3	*	*	*	*	*	*	*
0650	1,091	80.3	3.7	5.6	3.8	0.9	5.5	0.2
0651	593	94.6	0.3	2.5	0.3	0.3	1.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 67

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0652	325	95.1	0.0	1.8	0.0	0.0	3.1	0.0
0653	416	95.7	0.2	2.2	0.0	0.0	1.7	0.2
0654	375	89.9	2.1	2.4	0.3	0.0	5.3	0.0
0655	346	93.1	0.0	1.4	0.0	0.3	4.3	0.9
0657	63	*	*	*	*	*	*	*
0701	427	87.6	2.1	2.8	0.2	3.7	2.3	1.2
0702	74	*	*	*	*	*	*	*
0703	188	89.9	1.6	1.1	0.0	0.5	4.8	2.1
0704	56	*	*	*	*	*	*	*
0705	115	91.3	1.7	0.0	0.0	0.0	6.1	0.9
0706	13	*	*	*	*	*	*	*
0850	799	83.7	5.0	7.6	0.1	1.3	1.8	0.5
0851	717	93.0	0.1	4.9	0.0	0.6	1.1	0.3
0901	1,564	79.6	1.5	11.3	0.0	0.1	6.8	0.6
0902	43	*	*	*	*	*	*	*
1051	1,213	83.8	0.8	7.2	1.6	1.1	4.8	0.8
1101	1,811	35.2	50.5	8.4	0.2	1.1	3.7	0.9
1102	852	72.9	10.8	7.5	0.8	0.9	6.3	0.7
1104	1	*	*	*	*	*	*	*
1106	79	*	*	*	*	*	*	*
1250	1,069	86.6	0.5	9.4	0.0	0.5	2.1	1.0
1251	734	94.0	0.4	2.6	0.1	0.3	1.9	0.7
1252	570	61.2	0.4	32.8	0.0	0.2	4.7	0.7
1253	794	78.1	2.0	10.5	0.0	0.1	8.3	1.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 68

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1350	2,370	58.9	19.7	7.5	0.3	6.1	7.2	0.4
1351	119	84.0	3.4	7.6	0.0	0.0	4.2	0.8
1352	388	70.4	20.1	4.6	0.0	0.0	4.4	0.5
1401	230	91.3	0.9	3.5	1.3	0.0	1.7	1.3
1402	118	57.6	0.0	23.7	10.2	0.8	7.6	0.0
1403	24	*	*	*	*	*	*	*
1501	809	50.7	3.7	38.1	0.0	0.1	4.7	2.7
1503	973	61.7	7.6	21.8	0.4	0.3	7.3	0.9
1504	2,083	38.3	5.5	46.3	0.1	3.9	4.9	1.0
1505	398	81.7	4.3	8.5	0.0	0.5	3.8	1.3
1506	585	80.2	0.9	15.0	0.3	0.7	2.2	0.7
1507	577	83.0	0.9	8.3	0.2	1.0	5.9	0.7
1508	52	*	*	*	*	*	*	*
1509	423	50.6	1.4	41.1	0.2	0.0	3.5	3.1
1511	679	89.0	2.5	4.9	0.1	0.4	2.7	0.4
1512	355	85.1	0.6	7.3	0.3	0.6	6.2	0.0
1513	21	*	*	*	*	*	*	*
1514	297	84.2	2.4	7.7	0.3	0.7	3.7	1.0
1701	657	54.3	0.6	2.7	20.4	0.6	20.5	0.6
1950	609	91.6	0.2	4.8	0.2	0.2	3.0	0.2
1952	133	95.5	0.0	1.5	0.0	0.0	2.3	0.8
1953	310	91.9	0.3	6.5	0.0	0.0	1.3	0.0
1954	521	86.8	0.0	11.7	0.2	0.0	1.3	0.0
1955	106	85.8	0.9	10.4	0.0	0.0	1.9	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 69

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1957	425	81.4	0.9	13.6	0.0	0.0	3.5	0.5
1963	51	*	*	*	*	*	*	*
1964	36	*	*	*	*	*	*	*
2150	651	85.9	0.0	2.0	0.6	0.3	11.1	0.2
2151	227	78.4	0.9	9.7	0.4	0.0	6.6	4.0
2152	4	*	*	*	*	*	*	*
2251	406	92.1	0.5	2.5	0.0	0.0	3.9	1.0
2252	188	89.9	0.5	0.5	0.5	0.0	5.9	2.7
2554	52	*	*	*	*	*	*	*
2555	7,582	47.7	39.2	5.3	0.1	0.3	7.1	0.4
2556	345	29.6	58.3	5.2	0.3	0.3	6.4	0.0
2557	9	*	*	*	*	*	*	*
2558	6	*	*	*	*	*	*	*
2559	3	*	*	*	*	*	*	*
2560	6	*	*	*	*	*	*	*
2561	3	*	*	*	*	*	*	*
2562	16	*	*	*	*	*	*	*
2563	28	*	*	*	*	*	*	*
2564	22	*	*	*	*	*	*	*
3001	1,150	18.2	65.0	12.9	0.0	0.7	3.2	0.1
3002	299	22.4	50.2	22.7	0.0	0.0	4.7	0.0
3004	215	47.9	0.0	49.3	0.5	0.5	1.9	0.0
3005	66	*	*	*	*	*	*	*
3007	16	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 70

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
3009	3	*	*	*	*	*	*	*
3201	431	93.3	0.2	4.2	0.2	0.2	1.6	0.2
3203	1	*	*	*	*	*	*	*
3301	4,278	37.6	31.3	16.0	0.1	5.6	9.0	0.4
3304	103	49.5	9.7	9.7	0.0	24.3	6.8	0.0
3305	164	67.7	9.1	13.4	0.0	4.3	5.5	0.0
3306	264	84.8	3.0	6.4	0.4	0.8	3.8	0.8
3308	94	*	*	*	*	*	*	*
3309	20	*	*	*	*	*	*	*
3312	394	33.0	27.9	29.4	0.5	1.0	8.1	0.0
3550	61	*	*	*	*	*	*	*
3551	21	*	*	*	*	*	*	*
3552	46	*	*	*	*	*	*	*
3701	1,021	88.0	0.2	8.1	0.0	0.1	3.5	0.1
3850	3,066	71.7	14.1	5.3	0.1	0.4	8.1	0.3
3901	1,526	33.3	53.1	6.5	0.1	0.7	6.0	0.4
3902	318	47.8	17.0	15.7	0.0	0.6	18.9	0.0
3903	31	*	*	*	*	*	*	*
3904	7	*	*	*	*	*	*	*
3905	11	*	*	*	*	*	*	*
3950	2,180	55.0	22.1	11.5	0.2	1.4	9.4	0.4
3954	17	*	*	*	*	*	*	*
4001	92	*	*	*	*	*	*	*
4002	58	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 71

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4103	16	*	*	*	*	*	*	*
4106	2,494	13.4	42.5	38.1	0.3	0.7	4.5	0.5
4107	3,044	42.0	27.8	20.2	0.4	1.9	7.4	0.3
4109	936	82.8	1.9	9.5	0.1	1.0	4.7	0.0
4110	7	*	*	*	*	*	*	*
4112	545	1.7	1.1	95.6	0.2	0.7	0.0	0.7
4113	116	4.3	16.4	75.0	0.0	0.9	3.4	0.0
4115	3,573	33.0	29.7	20.6	0.3	11.3	4.7	0.4
4351	547	88.1	0.5	6.0	0.0	0.5	4.6	0.2
4352	157	73.9	11.5	4.5	0.0	0.0	9.6	0.6
4353	421	63.7	1.2	32.5	0.2	0.0	2.4	0.0
4354	757	86.1	4.1	2.9	0.4	0.7	5.3	0.5
4355	326	78.8	1.5	7.7	2.8	0.3	8.6	0.3
4356	426	81.9	1.6	11.0	0.0	0.7	4.7	0.0
4358	114	41.2	0.0	57.0	0.0	0.0	0.9	0.9
4359	509	68.6	0.4	28.5	0.4	0.4	0.8	1.0
4360	222	89.6	0.5	7.7	0.0	0.0	2.3	0.0
4361	253	93.3	0.4	3.6	0.0	0.0	2.4	0.4
4362	303	90.4	0.3	3.6	0.0	1.0	4.3	0.3
4363	835	91.4	0.1	2.9	0.0	0.2	4.9	0.5
4401	992	89.6	0.6	5.1	0.0	0.5	3.8	0.3
4501	936	92.0	0.1	5.1	0.0	0.1	2.6	0.1
4601	1,350	73.9	1.9	18.1	0.0	0.1	5.5	0.5
4602	10	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 72

### Michigan

#### Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4701	1,066	92.5	0.8	3.6	0.0	0.8	1.9	0.6
5050	1,631	51.2	38.4	2.6	0.0	1.5	6.1	0.2
5051	2,713	52.6	34.4	4.5	0.1	2.9	5.2	0.3
5052	2,947	37.7	38.6	2.4	0.2	15.8	4.7	0.5
5055	1,189	89.9	3.1	0.2	0.1	5.9	0.4	0.4
5101	1,702	69.0	11.9	11.9	0.2	1.8	5.3	0.0
5102	26	*	*	*	*	*	*	*
5202	791	86.9	0.6	3.3	1.4	0.4	7.3	0.1
5206	5	*	*	*	*	*	*	*
5601	971	92.0	1.1	2.4	0.0	1.4	3.1	0.0
5801	1,792	81.3	6.1	6.3	0.1	0.4	5.1	0.7
5803	126	86.5	0.8	9.5	0.0	0.8	2.4	0.0
5804	17	*	*	*	*	*	*	*
5806	59	*	*	*	*	*	*	*
6101	32	*	*	*	*	*	*	*
6102	117	82.1	1.7	5.1	0.0	0.9	10.3	0.0
6103	3,728	52.6	29.6	9.2	0.3	0.5	7.5	0.3
6105	18	*	*	*	*	*	*	*
6106	134	35.8	50.7	6.7	0.0	0.7	6.0	0.0
6301	3,762	40.7	31.0	20.8	0.1	1.8	5.3	0.2
6312	1,911	34.2	55.2	3.2	0.0	1.5	5.4	0.5
6316	1,211	66.8	18.6	7.3	0.2	3.4	3.6	0.2
6318	36	*	*	*	*	*	*	*
6320	207	45.9	42.0	4.8	0.5	0.5	6.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 73

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6325	262	91.2	1.9	3.1	0.0	0.0	3.4	0.4
6330	4	*	*	*	*	*	*	*
6335	1,050	73.8	15.1	2.4	0.0	7.0	1.5	0.1
6340	37	*	*	*	*	*	*	*
6345	79	*	*	*	*	*	*	*
6360	865	25.7	51.2	16.1	0.0	0.9	6.1	0.0
7304	39	*	*	*	*	*	*	*
7306	2,378	44.8	31.7	16.5	0.0	0.7	5.7	0.5
7308	46	*	*	*	*	*	*	*
7310	12	*	*	*	*	*	*	*
7401	2,221	81.4	6.2	6.0	0.1	0.4	5.6	0.4
7601	625	92.8	0.0	5.4	0.2	0.2	1.1	0.3
7850	1,119	89.0	0.4	6.3	0.1	0.1	4.0	0.1
7852	20	*	*	*	*	*	*	*
8101	1,774	4.8	91.7	0.3	0.1	0.2	2.9	0.1
8102	411	14.1	11.7	69.1	0.0	0.7	4.1	0.2
8105	1,288	32.0	61.6	2.6	0.2	0.6	2.7	0.2
8106	514	24.3	58.6	10.3	0.4	0.4	5.6	0.4
8150	23	*	*	*	*	*	*	*
8151	3,103	36.7	40.4	10.4	0.2	2.3	9.3	0.6
8152	53	*	*	*	*	*	*	*
8153	9	*	*	*	*	*	*	*
8155	14	*	*	*	*	*	*	*
8157	7	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 74

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
8201	2,663	45.5	38.9	4.6	0.2	3.7	6.5	0.5
8202	4,053	83.5	12.5	1.8	0.0	0.3	1.3	0.5
8203	404	59.7	29.2	4.0	0.5	2.0	4.0	0.7
8250	791	20.1	68.9	4.6	0.1	0.9	5.1	0.4
8252	161	3.7	92.5	0.0	0.0	0.6	3.1	0.0
8255	2,532	53.0	32.3	8.4	0.2	2.3	3.5	0.3
8257	2,128	49.7	13.8	0.8	0.2	33.5	1.9	0.2
8263	705	67.8	14.6	10.9	0.3	2.1	4.1	0.1
8264	730	64.7	26.2	4.1	0.1	1.2	3.6	0.1
8267	778	54.5	17.0	23.3	0.0	0.1	4.8	0.4
8901	1,049	2.0	93.1	0.6	0.1	1.0	3.0	0.2
8902	1,493	4.6	91.1	1.8	0.0	0.0	2.2	0.3
8903	1,549	2.7	92.0	1.4	0.0	0.8	2.9	0.2
8904	2,452	2.9	91.9	1.4	0.2	0.8	2.5	0.2
8905	1,547	60.3	28.6	8.1	0.0	0.5	2.5	0.1
8909	726	3.7	92.8	0.8	0.0	0.3	2.2	0.1
9045	1,785	10.6	27.4	57.5	0.2	0.3	3.6	0.3
9050	965	13.0	69.3	14.8	0.0	0.0	2.9	0.0
9051	1,611	9.2	7.2	81.4	0.0	0.1	1.7	0.2
9053	938	5.0	89.0	1.3	0.2	2.3	2.0	0.1
9055	19	*	*	*	*	*	*	*
9058	1,020	18.2	70.3	1.8	0.0	7.8	1.8	0.1
9066	1,155	1.7	95.5	0.7	0.0	0.2	1.8	0.1
9068	968	2.0	96.5	0.4	0.0	0.3	0.7	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 75

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9069	2,616	0.9	97.5	0.5	0.0	0.2	0.9	0.1
9070	1,433	3.8	91.8	1.0	0.0	1.7	1.6	0.1
9072	1,466	2.5	95.1	0.5	0.1	0.5	1.2	0.1
9080	2,673	19.3	76.4	1.9	0.2	0.2	1.8	0.1
9082	150	3.3	94.7	1.3	0.0	0.0	0.7	0.0
9083	104	26.9	61.5	8.7	0.0	1.9	1.0	0.0
9084	35	*	*	*	*	*	*	*
9085	50	*	*	*	*	*	*	*
9086	48	*	*	*	*	*	*	*
9087	5	*	*	*	*	*	*	*
Michigan	159,551	51.2	30.4	10.7	0.4	2.2	4.8	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 76

### Michigan

#### Comparison of Age Distribution by Clinic

#### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	117	0.0	0.9	6.8	64.1	25.6	2.6	0.0
0102	490	0.0	3.7	8.2	64.5	22.4	1.2	0.0
0103	407	0.0	3.2	8.1	66.3	21.1	1.2	0.0
0104	139	0.0	5.8	7.2	70.5	15.8	0.7	0.0
0105	20	*	*	*	*	*	*	*
0250	146	0.0	6.2	6.8	63.0	23.3	0.7	0.0
0251	89	*	*	*	*	*	*	*
0252	103	0.0	2.9	4.9	53.4	37.9	1.0	0.0
0253	136	0.0	3.7	8.8	65.4	22.1	0.0	0.0
0401	462	0.0	3.7	9.1	64.5	21.2	1.5	0.0
0402	386	0.3	1.8	10.4	60.6	25.4	1.6	0.0
0403	157	0.0	3.2	5.7	67.5	22.9	0.6	0.0
0404	112	0.0	0.0	8.9	67.9	23.2	0.0	0.0
0501	492	0.2	1.6	7.7	65.7	23.8	1.0	0.0
0503	109	0.0	0.9	9.2	55.0	32.1	2.8	0.0
0504	457	0.0	1.3	5.7	63.9	26.9	2.2	0.0
0505	297	0.0	0.7	7.7	71.4	19.9	0.3	0.0
0506	173	0.0	2.9	11.0	65.3	20.8	0.0	0.0
0507	76	*	*	*	*	*	*	*
0508	29	*	*	*	*	*	*	*
0509	3	*	*	*	*	*	*	*
0650	1,091	0.1	2.8	8.2	65.3	22.6	0.9	0.1
0651	593	0.3	3.7	7.8	66.6	20.1	1.3	0.2
0652	325	0.0	2.5	4.6	71.1	19.4	2.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 77

### Michigan

#### Comparison of Age Distribution by Clinic

#### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0653	416	0.0	1.9	7.2	67.3	22.6	1.0	0.0
0654	375	0.3	4.8	10.4	61.6	21.9	0.8	0.3
0655	346	0.0	2.3	11.6	61.3	24.0	0.6	0.3
0657	63	*	*	*	*	*	*	*
0701	427	0.0	2.3	5.2	56.9	32.1	3.3	0.2
0702	74	*	*	*	*	*	*	*
0703	188	0.0	0.5	8.0	64.4	26.6	0.5	0.0
0704	56	*	*	*	*	*	*	*
0705	115	0.0	2.6	9.6	56.5	28.7	2.6	0.0
0706	13	*	*	*	*	*	*	*
0850	799	0.1	2.4	10.4	60.1	25.0	1.5	0.5
0851	717	0.0	2.2	8.4	64.2	23.2	1.8	0.3
0901	1,564	0.1	2.6	9.1	63.1	23.5	1.7	0.0
0902	43	*	*	*	*	*	*	*
1051	1,213	0.1	2.5	6.7	60.7	27.2	2.6	0.3
1101	1,811	0.1	4.0	9.2	61.6	21.4	1.5	2.2
1102	852	0.0	2.6	11.5	60.6	23.6	1.5	0.2
1104	1	*	*	*	*	*	*	*
1106	79	*	*	*	*	*	*	*
1250	1,069	0.0	2.1	10.1	61.8	23.4	2.6	0.0
1251	734	0.1	2.6	10.2	66.8	18.9	1.0	0.4
1252	570	0.2	2.1	10.4	63.3	21.8	2.3	0.0
1253	794	0.0	3.0	8.1	67.0	20.9	1.0	0.0
1350	2,370	0.1	3.1	8.4	63.8	23.0	1.3	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 78

### Michigan

#### Comparison of Age Distribution by Clinic

#### Combined 3 Years

Comparison of Age Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1351	119	0.0	2.5	6.7	53.8	35.3	1.7	0.0
1352	388	0.3	3.6	7.5	65.7	20.9	1.5	0.5
1401	230	0.0	3.5	7.0	67.4	20.9	1.3	0.0
1402	118	0.0	0.0	3.4	56.8	36.4	3.4	0.0
1403	24	*	*	*	*	*	*	*
1501	809	0.0	2.2	10.4	59.7	24.8	2.0	0.9
1503	973	0.4	3.4	11.0	60.7	23.0	1.4	0.0
1504	2,083	0.2	2.9	8.0	61.0	25.7	2.1	0.1
1505	398	0.0	1.8	5.5	65.6	25.4	1.3	0.5
1506	585	0.0	2.9	6.8	59.8	29.1	1.4	0.0
1507	577	0.0	1.0	5.0	66.2	25.6	2.1	0.0
1508	52	*	*	*	*	*	*	*
1509	423	0.0	4.0	6.6	62.9	24.1	2.1	0.2
1511	679	0.0	3.4	10.8	61.1	23.3	1.5	0.0
1512	355	0.0	2.5	6.5	59.4	29.3	2.3	0.0
1513	21	*	*	*	*	*	*	*
1514	297	0.0	1.7	4.4	56.9	35.0	1.7	0.3
1701	657	0.2	1.8	7.8	62.7	25.9	1.7	0.0
1950	609	0.0	3.4	12.0	66.3	16.9	1.1	0.2
1952	133	0.0	0.8	9.8	65.4	22.6	1.5	0.0
1953	310	0.3	3.5	7.4	69.0	18.1	1.6	0.0
1954	521	0.4	2.5	8.4	64.5	22.5	1.5	0.2
1955	106	0.0	0.9	9.4	64.2	24.5	0.9	0.0
1957	425	0.0	2.1	6.8	60.9	28.7	1.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 79

### Michigan

#### Comparison of Age Distribution by Clinic

#### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1963	51	*	*	*	*	*	*	*
1964	36	*	*	*	*	*	*	*
2150	651	0.0	2.3	8.1	61.4	27.5	0.6	0.0
2151	227	0.0	4.4	7.0	67.0	19.8	1.8	0.0
2152	4	*	*	*	*	*	*	*
2251	406	0.2	1.7	8.6	61.1	27.1	1.2	0.0
2252	188	0.0	3.7	8.0	63.3	23.9	1.1	0.0
2554	52	*	*	*	*	*	*	*
2555	7,582	0.2	2.6	7.8	64.4	23.5	1.5	0.1
2556	345	0.3	2.3	8.1	68.4	19.7	1.2	0.0
2557	9	*	*	*	*	*	*	*
2558	6	*	*	*	*	*	*	*
2559	3	*	*	*	*	*	*	*
2560	6	*	*	*	*	*	*	*
2561	3	*	*	*	*	*	*	*
2562	16	*	*	*	*	*	*	*
2563	28	*	*	*	*	*	*	*
2564	22	*	*	*	*	*	*	*
3001	1,150	0.3	3.7	8.6	65.7	19.7	1.6	0.4
3002	299	0.0	5.7	9.0	64.2	19.7	1.3	0.0
3004	215	0.0	1.4	5.1	60.9	30.2	2.3	0.0
3005	66	*	*	*	*	*	*	*
3007	16	*	*	*	*	*	*	*
3009	3	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 80

### Michigan

#### Comparison of Age Distribution by Clinic

#### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3201	431	0.0	1.4	6.5	62.2	28.1	1.9	0.0
3203	1	*	*	*	*	*	*	*
3301	4,278	0.2	2.9	8.3	62.1	24.6	1.9	0.1
3304	103	0.0	0.0	1.0	52.4	45.6	1.0	0.0
3305	164	0.0	1.8	9.1	50.0	37.2	1.8	0.0
3306	264	0.4	8.0	8.3	53.0	28.4	1.1	0.8
3308	94	*	*	*	*	*	*	*
3309	20	*	*	*	*	*	*	*
3312	394	0.0	2.0	8.6	61.9	25.6	1.5	0.3
3550	61	*	*	*	*	*	*	*
3551	21	*	*	*	*	*	*	*
3552	46	*	*	*	*	*	*	*
3701	1,021	0.0	2.7	8.9	66.1	21.4	0.9	0.0
3850	3,066	0.0	3.1	8.7	65.0	22.1	0.8	0.3
3901	1,526	0.1	3.7	10.0	60.6	23.6	1.8	0.3
3902	318	0.0	3.8	10.7	58.5	24.8	2.2	0.0
3903	31	*	*	*	*	*	*	*
3904	7	*	*	*	*	*	*	*
3905	11	*	*	*	*	*	*	*
3950	2,180	0.1	3.2	7.4	62.5	24.5	2.1	0.2
3954	17	*	*	*	*	*	*	*
4001	92	*	*	*	*	*	*	*
4002	58	*	*	*	*	*	*	*
4103	16	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 81

### Michigan

#### Comparison of Age Distribution by Clinic

#### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4106	2,494	0.1	3.3	9.5	58.9	25.5	2.2	0.5
4107	3,044	0.2	2.6	8.5	59.4	28.0	1.1	0.1
4109	936	0.1	2.2	6.3	64.7	25.1	1.4	0.1
4110	7	*	*	*	*	*	*	*
4112	545	0.2	3.7	8.4	54.3	30.6	2.4	0.4
4113	116	0.0	0.9	8.6	41.4	34.5	7.8	6.9
4115	3,573	0.1	2.0	6.3	57.6	31.7	2.1	0.3
4351	547	0.0	1.8	9.1	67.3	20.8	0.9	0.0
4352	157	0.0	1.3	7.0	69.4	21.0	1.3	0.0
4353	421	0.0	4.5	8.1	65.6	19.2	2.6	0.0
4354	757	0.1	1.7	8.1	67.8	21.1	1.1	0.1
4355	326	0.0	2.1	11.0	57.1	25.5	4.0	0.3
4356	426	0.2	3.8	7.0	66.9	21.1	0.9	0.0
4358	114	0.9	1.8	7.9	60.5	26.3	2.6	0.0
4359	509	0.2	2.4	7.3	62.3	25.0	2.8	0.2
4360	222	0.0	0.5	11.3	68.0	18.5	1.8	0.0
4361	253	0.0	2.0	8.3	64.4	23.7	1.2	0.4
4362	303	0.0	3.0	8.9	68.3	19.8	0.0	0.0
4363	835	0.0	3.0	8.0	65.9	21.7	1.3	0.1
4401	992	0.0	3.4	6.5	63.6	25.1	1.2	0.2
4501	936	0.1	2.9	7.3	65.2	22.6	1.8	0.1
4601	1,350	0.1	2.1	9.1	63.3	24.4	1.0	0.1
4602	10	*	*	*	*	*	*	*
4701	1,066	0.0	1.7	5.6	61.5	29.3	1.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 82

### Michigan

#### Comparison of Age Distribution by Clinic

#### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5050	1,631	0.0	1.9	4.9	57.8	33.0	2.4	0.0
5051	2,713	0.0	1.0	4.8	61.4	30.6	2.2	0.0
5052	2,947	0.0	1.7	5.5	60.8	29.5	2.5	0.0
5055	1,189	0.0	0.1	1.1	52.1	42.5	4.1	0.1
5101	1,702	0.1	1.0	4.8	61.7	29.6	2.9	0.0
5102	26	*	*	*	*	*	*	*
5202	791	0.3	1.1	6.8	60.6	29.3	1.9	0.0
5206	5	*	*	*	*	*	*	*
5601	971	0.0	1.9	5.5	65.7	25.5	1.4	0.0
5801	1,792	0.0	2.8	9.3	63.1	23.2	1.6	0.1
5803	126	0.0	0.0	1.6	52.4	42.9	3.2	0.0
5804	17	*	*	*	*	*	*	*
5806	59	*	*	*	*	*	*	*
6101	32	*	*	*	*	*	*	*
6102	117	0.0	6.8	8.5	59.8	22.2	2.6	0.0
6103	3,728	0.3	3.5	9.7	63.5	21.6	1.3	0.2
6105	18	*	*	*	*	*	*	*
6106	134	0.0	3.7	9.0	68.7	17.2	0.7	0.7
6301	3,762	0.1	2.6	8.1	61.2	25.7	2.3	0.0
6312	1,911	0.0	0.7	3.9	55.1	37.3	2.9	0.1
6316	1,211	0.0	0.5	2.8	58.7	35.2	2.8	0.0
6318	36	*	*	*	*	*	*	*
6320	207	0.0	1.4	1.0	58.9	36.7	1.9	0.0
6325	262	0.0	1.1	5.3	62.2	27.5	1.1	2.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 83

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
6330	4	*	*	*	*	*	*	*
6335	1,050	0.0	0.6	2.9	54.8	38.4	3.4	0.0
6340	37	*	*	*	*	*	*	*
6345	79	*	*	*	*	*	*	*
6360	865	0.1	1.7	8.2	58.5	27.5	2.7	1.3
7304	39	*	*	*	*	*	*	*
7306	2,378	0.1	2.9	9.0	63.2	23.4	1.3	0.0
7308	46	*	*	*	*	*	*	*
7310	12	*	*	*	*	*	*	*
7401	2,221	0.1	3.0	7.9	64.7	22.6	1.5	0.1
7601	625	0.0	2.2	9.0	66.1	21.8	1.0	0.0
7850	1,119	0.1	3.1	8.5	68.5	18.5	1.3	0.0
7852	20	*	*	*	*	*	*	*
8101	1,774	0.3	2.6	9.2	62.1	23.7	2.0	0.1
8102	411	0.0	2.2	4.1	62.5	29.4	1.7	0.0
8105	1,288	0.1	1.5	4.8	60.7	31.1	1.7	0.1
8106	514	0.0	1.9	5.3	66.3	23.5	2.9	0.0
8150	23	*	*	*	*	*	*	*
8151	3,103	0.1	1.8	6.1	59.0	30.5	2.4	0.2
8152	53	*	*	*	*	*	*	*
8153	9	*	*	*	*	*	*	*
8155	14	*	*	*	*	*	*	*
8157	7	*	*	*	*	*	*	*
8201	2,663	0.0	2.0	5.3	60.3	29.9	2.3	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 84

### Michigan

#### Comparison of Age Distribution by Clinic

#### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8202	4,053	0.0	1.2	3.8	55.3	34.5	4.4	0.7
8203	404	0.2	0.7	5.7	56.7	34.9	1.7	0.0
8250	791	0.3	2.0	7.7	59.4	28.3	2.3	0.0
8252	161	0.0	1.2	6.8	73.3	18.0	0.6	0.0
8255	2,532	0.0	1.6	6.8	64.8	25.0	1.7	0.0
8257	2,128	0.1	0.6	3.6	58.8	31.9	4.9	0.1
8263	705	0.0	0.6	5.2	66.1	26.5	1.6	0.0
8264	730	0.0	2.3	3.2	63.0	28.9	2.6	0.0
8267	778	0.1	3.7	6.8	61.2	26.5	1.7	0.0
8901	1,049	0.3	3.0	8.3	64.2	22.1	2.0	0.2
8902	1,493	0.1	1.7	6.4	63.5	26.4	1.9	0.1
8903	1,549	0.1	3.4	8.8	62.2	23.8	1.7	0.1
8904	2,452	0.2	2.6	6.7	63.7	24.9	1.9	0.0
8905	1,547	0.0	1.4	3.8	56.0	34.8	3.9	0.1
8909	726	0.1	3.6	8.0	59.8	26.0	2.2	0.3
9045	1,785	0.2	3.4	7.4	56.1	29.6	3.2	0.1
9050	965	0.1	2.6	7.5	60.0	28.0	1.9	0.0
9051	1,611	0.1	2.7	7.9	52.5	30.5	6.4	0.0
9053	938	0.0	2.3	7.5	60.1	27.6	2.2	0.2
9055	19	*	*	*	*	*	*	*
9058	1,020	0.2	2.0	6.2	59.1	28.7	3.6	0.2
9066	1,155	0.3	2.7	6.4	63.2	24.8	2.5	0.1
9068	968	0.2	2.8	7.1	61.5	26.0	2.2	0.2
9069	2,616	0.0	3.4	6.3	60.9	26.7	2.5	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 85

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9070	1,433	0.1	2.2	8.9	62.3	23.9	2.4	0.0
9072	1,466	0.3	2.5	9.8	63.4	23.1	1.0	0.1
9080	2,673	0.0	1.8	6.2	60.4	29.4	2.2	0.0
9082	150	0.0	5.3	9.3	51.3	32.7	1.3	0.0
9083	104	0.0	0.0	4.8	60.6	28.8	4.8	1.0
9084	35	*	*	*	*	*	*	*
9085	50	*	*	*	*	*	*	*
9086	48	*	*	*	*	*	*	*
9087	5	*	*	*	*	*	*	*
<b>Michigan</b>	<b>159,551</b>	<b>0.1</b>	<b>2.4</b>	<b>7.3</b>	<b>61.7</b>	<b>26.3</b>	<b>2.0</b>	<b>0.2</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

## 2018 - 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Maternal Health Indicators by Clinic Combined 3 Years

Page : 86

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	67	*	*	21	*	*	18	*	21	*	107	6.5 (43)	107	10.3 (48)
0102	314	5.7 (134)	60.8 (79)	123	14.6 (12)	51.2 (47)	66	*	68	*	444	5.0 (9)	444	10.8 (62)
0103	349	5.2 (125)	61.3 (89)	187	19.8 (89)	50.8 (43)	47	*	63	*	370	6.2 (37)	370	9.7 (33)
0104	108	2.8 (18)	54.6 (17)	52	*	*	12	*	23	*	129	3.1 (4)	130	6.9 (9)
0105	15	*	*	7	*	*	1	*	0	*	20	*	20	*
0250	139	3.6 (39)	62.6 (107)	93	*	*	14	*	24	*	124	8.9 (114)	125	11.2 (68)
0251	59	*	*	19	*	*	7	*	16	*	82	*	81	*
0252	102	3.9 (54)	65.7 (134)	90	*	*	12	*	20	*	94	*	94	*
0253	96	*	*	59	*	*	7	*	22	*	126	11.1 (145)	126	17.5 (147)
0401	428	4.9 (112)	61.0 (84)	273	20.1 (95)	52.7 (64)	50	*	64	*	417	5.8 (22)	418	14.6 (122)
0402	358	5.9 (137)	57.5 (34)	120	12.5 (4)	55.8 (103)	45	*	48	*	355	9.0 (119)	355	10.1 (43)
0403	146	6.2 (144)	57.5 (34)	84	*	*	17	*	25	*	144	5.6 (16)	144	10.4 (50)
0404	107	3.7 (42)	61.7 (93)	82	*	*	10	*	14	*	106	8.5 (106)	106	10.4 (50)
0501	315	4.8 (109)	54.9 (19)	86	*	*	60	*	55	*	463	7.8 (81)	463	9.9 (39)
0503	101	5.9 (137)	52.5 (8)	50	*	*	15	*	10	*	104	6.7 (52)	104	9.6 (31)
0504	405	4.0 (65)	59.0 (54)	183	14.2 (10)	50.3 (34)	57	*	47	*	433	6.5 (43)	433	14.5 (119)
0505	229	3.5 (36)	59.8 (68)	90	*	*	36	*	31	*	275	5.1 (10)	276	12.0 (78)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 87

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	115	6.1 (141)	61.7 (93)	23	*	*	22	*	9	*	161	8.1 (93)	161	9.3 (25)
0507	46	*	*	14	*	*	3	*	15	*	76	*	76	*
0508	19	*	*	8	*	*	1	*	6	*	28	*	28	*
0509	3	*	*	2	*	*	0	*	1	*	3	*	3	*
0650	894	4.1 (69)	61.6 (91)	581	17.4 (46)	50.3 (34)	132	28.8 (6)	167	22.2 (10)	999	7.5 (71)	1,000	11.0 (65)
0651	553	4.7 (106)	57.7 (38)	374	16.8 (36)	50.0 (27)	61	*	68	*	551	6.4 (40)	551	9.4 (27)
0652	234	3.8 (49)	58.1 (41)	155	19.4 (83)	53.5 (76)	42	*	41	*	314	8.3 (101)	315	14.3 (116)
0653	378	2.9 (23)	61.1 (85)	276	9.4 (1)	58.3 (120)	46	*	42	*	390	7.9 (87)	391	15.6 (135)
0654	333	2.1 (8)	62.8 (111)	219	24.7 (118)	50.7 (41)	44	*	61	*	345	7.5 (71)	345	13.0 (95)
0655	321	3.7 (42)	61.7 (93)	235	19.1 (77)	49.8 (24)	37	*	48	*	308	8.1 (93)	310	17.1 (145)
0657	46	*	*	26	*	*	6	*	12	*	62	*	62	*
0701	331	4.2 (78)	50.2 (1)	183	18.6 (71)	41.0 (5)	51	*	96	*	382	6.5 (43)	383	10.2 (47)
0702	70	*	*	53	*	*	4	*	11	*	68	*	68	*
0703	164	3.0 (24)	58.5 (46)	86	*	*	21	*	29	*	168	10.7 (143)	168	10.7 (58)
0704	55	*	*	43	*	*	8	*	9	*	50	*	50	*
0705	67	*	*	37	*	*	20	*	20	*	105	7.6 (76)	105	9.5 (29)
0706	9	*	*	5	*	*	0	*	3	*	13	*	13	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 88

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	538	4.6 (101)	60.8 (79)	314	19.1 (77)	52.5 (61)	98	*	114	15.8 (4)	726	10.3 (139)	728	15.5 (133)
0851	358	6.7 (146)	52.8 (11)	94	*	*	90	*	116	16.4 (6)	653	9.2 (126)	653	12.3 (81)
0901	1,315	3.4 (32)	61.9 (98)	856	17.2 (42)	52.6 (62)	201	29.9 (7)	146	15.8 (4)	1,444	6.2 (37)	1,444	10.1 (43)
0902	39	*	*	25	*	*	3	*	4	*	38	*	38	*
1051	1,064	4.1 (69)	52.3 (5)	854	15.3 (17)	52.3 (60)	133	54.1 (51)	128	32.8 (32)	1,125	8.6 (109)	1,128	9.6 (31)
1101	1,544	3.6 (39)	63.1 (117)	1,153	20.0 (94)	53.1 (70)	243	47.3 (34)	194	28.4 (23)	1,647	7.6 (76)	1,646	16.3 (141)
1102	729	3.8 (49)	59.3 (58)	480	15.6 (22)	55.4 (99)	105	37.1 (11)	129	17.8 (7)	735	7.1 (62)	735	13.5 (105)
1104	0	*	*	0	*	*	0	*	0	*	1	*	1	*
1106	69	*	*	52	*	*	12	*	15	*	66	*	67	*
1250	718	5.7 (134)	62.5 (105)	358	17.9 (55)	53.4 (75)	142	16.2 (1)	121	25.6 (18)	977	8.5 (106)	977	8.3 (19)
1251	658	5.6 (133)	60.9 (82)	492	16.5 (30)	53.7 (79)	78	*	117	29.1 (24)	679	6.0 (29)	679	6.2 (7)
1252	472	4.4 (91)	64.8 (127)	176	15.3 (17)	57.4 (115)	67	*	71	*	522	5.9 (25)	522	11.3 (69)
1253	673	4.5 (97)	58.1 (41)	320	21.9 (109)	49.4 (21)	90	*	97	*	713	7.4 (67)	712	10.7 (58)
1350	1,902	4.0 (65)	60.1 (70)	1,101	20.9 (104)	52.8 (66)	254	44.9 (25)	327	29.4 (25)	2,118	8.9 (114)	2,119	10.6 (54)
1351	99	*	*	77	*	*	14	*	30	*	110	9.1 (121)	110	10.0 (41)
1352	213	6.1 (141)	56.3 (27)	98	*	*	37	*	55	*	361	3.9 (5)	361	4.2 (4)
1401	201	5.0 (116)	51.7 (3)	139	13.7 (7)	54.7 (90)	29	*	29	*	216	6.0 (29)	216	12.5 (86)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 89

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1402	108	2.8 (18)	72.2 (147)	75	*	*	10	*	14	*	110	13.6 (151)	110	5.5 (5)
1403	22	*	*	17	*	*	0	*	4	*	24	*	24	*
1501	699	4.0 (65)	66.0 (138)	456	46.9 (120)	26.8 (2)	80	*	100	25.0 (16)	721	9.2 (126)	721	8.6 (24)
1503	839	4.1 (69)	60.2 (71)	561	16.9 (38)	52.8 (66)	105	41.0 (16)	84	*	894	8.4 (105)	897	14.5 (119)
1504	1,389	4.8 (109)	59.5 (63)	886	16.4 (28)	50.0 (27)	253	47.4 (35)	486	38.7 (53)	1,886	8.9 (114)	1,890	8.3 (19)
1505	281	3.9 (54)	57.3 (33)	136	13.2 (6)	50.7 (41)	52	*	95	*	367	8.2 (97)	367	12.3 (81)
1506	480	4.6 (101)	62.9 (115)	123	15.4 (19)	56.1 (106)	67	*	76	*	529	9.1 (121)	530	11.1 (66)
1507	438	4.3 (87)	57.8 (40)	177	11.3 (3)	57.1 (114)	52	*	86	*	529	8.3 (101)	529	7.9 (14)
1508	27	*	*	13	*	*	3	*	6	*	41	*	41	*
1509	323	5.3 (128)	63.5 (119)	189	18.0 (58)	50.3 (34)	45	*	94	*	352	9.1 (121)	353	8.2 (18)
1511	542	5.4 (130)	59.6 (64)	266	19.2 (81)	49.2 (19)	97	*	146	30.1 (27)	632	8.7 (111)	633	12.3 (81)
1512	247	4.5 (97)	60.7 (78)	68	*	*	57	*	82	*	330	4.8 (8)	331	9.7 (33)
1513	19	*	*	7	*	*	0	*	4	*	17	*	17	*
1514	166	6.0 (140)	51.2 (2)	110	10.0 (2)	52.7 (64)	37	*	81	*	281	6.4 (40)	281	13.2 (97)
1701	541	4.1 (69)	60.3 (72)	348	16.1 (25)	54.3 (84)	69	*	90	*	587	7.2 (64)	588	11.1 (66)
1950	566	4.6 (101)	61.1 (85)	393	17.6 (52)	51.7 (51)	66	*	109	21.1 (9)	570	10.2 (138)	571	13.7 (109)
1952	124	3.2 (27)	66.1 (139)	91	*	*	15	*	32	*	129	7.0 (59)	129	12.4 (84)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 90

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1953	285	6.7 (146)	55.4 (22)	214	21.5 (107)	50.0 (27)	41	*	80	*	287	8.0 (89)	287	18.5 (151)
1954	485	3.9 (54)	63.5 (119)	371	17.5 (49)	55.0 (94)	62	*	77	*	482	14.1 (152)	482	15.4 (131)
1955	102	0.0 (1)	65.7 (134)	72	*	*	10	*	17	*	97	*	97	*
1957	387	4.4 (91)	66.1 (139)	280	16.4 (28)	51.1 (45)	43	*	89	*	398	11.6 (147)	398	14.6 (122)
1963	46	*	*	37	*	*	7	*	9	*	47	*	49	*
1964	34	*	*	26	*	*	6	*	9	*	31	*	31	*
2150	534	2.4 (11)	59.2 (56)	374	16.3 (27)	54.0 (81)	71	*	111	22.5 (11)	587	10.6 (141)	587	13.1 (96)
2151	184	4.9 (112)	53.8 (15)	141	14.2 (10)	53.2 (73)	28	*	42	*	187	5.9 (25)	188	14.4 (117)
2152	4	*	*	2	*	*	0	*	0	*	3	*	4	*
2251	311	5.5 (132)	52.4 (6)	119	16.0 (24)	50.4 (39)	50	*	59	*	367	2.7 (3)	368	14.7 (125)
2252	139	8.6 (150)	53.2 (13)	45	*	*	18	*	26	*	156	7.7 (79)	156	13.5 (105)
2554	45	*	*	27	*	*	7	*	7	*	48	*	48	*
2555	5,677	5.0 (116)	58.5 (46)	2,620	17.4 (46)	52.1 (57)	1,134	50.3 (43)	911	32.4 (30)	6,953	7.7 (79)	6,962	15.5 (133)
2556	292	3.8 (49)	58.9 (52)	204	24.0 (117)	49.0 (18)	33	*	43	*	327	7.0 (59)	327	18.0 (149)
2557	5	*	*	1	*	*	4	*	1	*	8	*	8	*
2558	6	*	*	4	*	*	1	*	1	*	5	*	5	*
2559	3	*	*	2	*	*	1	*	0	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 91

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2560	4	*	*	1	*	*	0	*	0	*	4	*	4	*
2561	2	*	*	2	*	*	2	*	0	*	3	*	3	*
2562	11	*	*	3	*	*	5	*	3	*	14	*	14	*
2563	18	*	*	5	*	*	2	*	7	*	26	*	26	*
2564	11	*	*	2	*	*	3	*	2	*	18	*	18	*
3001	1,068	3.7 (42)	63.9 (125)	707	21.8 (108)	46.5 (13)	158	46.8 (33)	502	39.0 (55)	1,075	6.1 (33)	1,074	11.5 (71)
3002	281	1.4 (4)	79.0 (150)	191	20.4 (99)	52.9 (68)	30	*	148	37.8 (49)	280	6.1 (33)	280	13.9 (110)
3004	178	2.2 (9)	66.3 (141)	128	21.1 (105)	37.5 (4)	20	*	29	*	193	7.8 (81)	193	10.9 (63)
3005	60	*	*	26	*	*	11	*	9	*	61	*	61	*
3007	14	*	*	5	*	*	1	*	6	*	13	*	13	*
3009	2	*	*	0	*	*	0	*	1	*	1	*	1	*
3201	362	5.0 (116)	66.9 (143)	273	15.4 (19)	55.3 (98)	47	*	75	*	409	7.6 (76)	411	12.7 (89)
3203	1	*	*	0	*	*	0	*	0	*	1	*	1	*
3301	3,531	4.2 (78)	62.0 (99)	2,810	19.6 (86)	53.1 (70)	578	46.7 (32)	473	32.6 (31)	4,002	6.9 (55)	4,003	11.6 (74)
3304	79	*	*	62	*	*	8	*	15	*	95	*	95	*
3305	134	7.5 (148)	57.5 (34)	106	17.9 (55)	54.7 (90)	20	*	26	*	151	10.6 (141)	151	7.3 (10)
3306	217	3.7 (42)	53.5 (14)	183	14.8 (13)	55.7 (101)	35	*	45	*	250	2.4 (2)	250	8.4 (23)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 92

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3308	83	*	*	67	*	*	8	*	17	*	86	*	86	*
3309	19	*	*	13	*	*	2	*	4	*	18	*	18	*
3312	322	4.3 (87)	63.0 (116)	258	20.2 (96)	51.2 (47)	58	*	42	*	368	7.9 (87)	369	10.0 (41)
3550	50	*	*	9	*	*	9	*	12	*	53	*	53	*
3551	14	*	*	3	*	*	2	*	7	*	18	*	18	*
3552	36	*	*	8	*	*	6	*	10	*	40	*	40	*
3701	714	4.6 (101)	61.1 (85)	482	16.2 (26)	55.2 (96)	120	25.8 (4)	198	15.7 (3)	935	9.1 (121)	933	16.8 (144)
3850	2,219	5.0 (116)	61.2 (88)	1,343	16.6 (31)	54.1 (83)	351	41.6 (18)	357	38.1 (50)	2,831	6.9 (55)	2,836	10.6 (54)
3901	1,190	3.4 (32)	58.2 (44)	518	17.8 (53)	53.5 (76)	154	57.8 (55)	222	36.0 (43)	1,368	7.8 (81)	1,368	11.3 (69)
3902	268	4.1 (69)	65.3 (132)	121	14.0 (8)	57.9 (118)	43	*	56	*	279	10.4 (140)	280	6.8 (8)
3903	27	*	*	16	*	*	2	*	16	*	30	*	30	*
3904	6	*	*	1	*	*	3	*	1	*	7	*	7	*
3905	9	*	*	7	*	*	0	*	2	*	11	*	11	*
3950	1,786	3.9 (54)	60.3 (72)	1,032	16.6 (31)	53.6 (78)	240	43.8 (21)	384	38.3 (51)	1,950	9.2 (126)	1,950	10.3 (48)
3954	14	*	*	10	*	*	2	*	3	*	16	*	16	*
4001	74	*	*	26	*	*	0	*	12	*	79	*	79	*
4002	50	*	*	27	*	*	0	*	10	*	48	*	48	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 93

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4103	7	*	*	3	*	*	2	*	4	*	13	*	13	*
4106	1,705	3.2 (27)	62.0 (99)	802	22.6 (112)	44.8 (8)	360	45.6 (28)	621	36.2 (44)	2,295	6.4 (40)	2,295	11.7 (76)
4107	2,392	4.1 (69)	58.5 (46)	1,113	14.9 (14)	51.1 (45)	407	48.2 (36)	826	38.7 (53)	2,800	6.6 (49)	2,802	14.5 (119)
4109	438	3.9 (54)	55.3 (21)	228	19.3 (82)	55.7 (101)	120	24.2 (3)	177	14.7 (1)	860	7.8 (81)	864	15.3 (130)
4110	6	*	*	6	*	*	0	*	0	*	7	*	7	*
4112	537	2.0 (6)	64.6 (126)	425	73.4 (121)	8.2 (1)	63	*	51	*	505	16.2 (153)	506	12.5 (86)
4113	102	1.0 (2)	73.5 (149)	25	*	*	15	*	32	*	112	12.5 (149)	112	8.0 (15)
4115	1,987	4.9 (112)	57.0 (32)	451	19.1 (77)	46.3 (11)	503	45.7 (29)	899	37.7 (48)	3,297	8.3 (101)	3,300	12.9 (92)
4351	518	3.3 (29)	62.5 (105)	389	21.9 (109)	45.8 (9)	67	*	80	*	511	9.2 (126)	511	14.1 (112)
4352	148	8.1 (149)	62.8 (111)	110	18.2 (62)	48.2 (15)	16	*	20	*	145	9.0 (119)	146	15.1 (129)
4353	344	2.0 (6)	65.7 (134)	234	17.9 (55)	51.7 (51)	50	*	57	*	376	8.0 (89)	375	14.4 (117)
4354	591	4.2 (78)	60.6 (77)	307	19.9 (92)	52.1 (57)	84	*	127	18.9 (8)	699	5.9 (25)	698	14.6 (122)
4355	217	2.3 (10)	60.4 (74)	71	*	*	37	*	71	*	309	7.4 (67)	310	12.9 (92)
4356	358	2.5 (13)	55.6 (23)	211	17.1 (41)	54.5 (87)	65	*	61	*	394	7.4 (67)	394	12.4 (84)
4358	95	*	*	75	*	*	12	*	12	*	103	12.6 (150)	103	9.7 (33)
4359	433	4.4 (91)	62.4 (103)	291	20.3 (97)	48.5 (17)	53	*	81	*	477	10.9 (144)	478	13.2 (97)
4360	214	4.7 (106)	61.7 (93)	164	22.6 (112)	50.6 (40)	19	*	33	*	198	1.0 (1)	198	14.1 (112)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 94

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4361	214	2.8 (18)	55.6 (23)	148	20.3 (97)	56.1 (106)	23	*	42	*	235	8.9 (114)	235	9.8 (37)
4362	254	5.1 (123)	58.3 (45)	193	18.7 (73)	54.9 (93)	46	*	59	*	284	9.9 (134)	284	9.5 (29)
4363	746	4.2 (78)	59.4 (61)	485	18.4 (67)	49.9 (26)	91	*	127	28.3 (22)	765	8.0 (89)	765	11.8 (77)
4401	974	4.3 (87)	55.7 (25)	855	16.7 (35)	54.4 (86)	134	38.8 (13)	213	31.9 (29)	922	4.1 (7)	922	7.3 (10)
4501	715	5.0 (116)	59.7 (67)	558	18.3 (65)	56.5 (110)	109	21.1 (2)	132	22.7 (12)	884	9.4 (130)	885	13.6 (107)
4601	1,028	3.9 (54)	58.1 (41)	338	17.8 (53)	47.9 (14)	213	49.8 (40)	248	41.5 (58)	1,224	8.6 (109)	1,226	13.4 (103)
4602	7	*	*	3	*	*	0	*	2	*	10	*	10	*
4701	1,021	4.4 (91)	52.7 (9)	714	17.2 (42)	51.7 (51)	131	33.6 (10)	262	15.3 (2)	983	11.7 (148)	982	16.4 (142)
5050	1,567	4.3 (87)	62.6 (107)	1,361	17.4 (46)	54.7 (90)	267	49.8 (40)	313	32.9 (33)	1,507	8.0 (89)	1,505	12.2 (80)
5051	2,513	4.2 (78)	60.0 (69)	2,134	18.7 (73)	52.2 (59)	434	46.3 (31)	570	34.2 (37)	2,510	7.5 (71)	2,511	9.9 (39)
5052	2,747	4.1 (69)	61.4 (90)	2,314	20.6 (101)	49.4 (21)	427	42.4 (19)	514	33.1 (35)	2,748	9.4 (130)	2,749	10.5 (52)
5055	584	2.7 (16)	56.7 (29)	200	18.5 (69)	54.5 (87)	96	*	197	35.5 (41)	1,144	11.3 (146)	1,143	3.7 (3)
5101	1,462	4.4 (91)	57.7 (38)	987	15.0 (15)	54.3 (84)	245	37.6 (12)	322	27.6 (20)	1,569	10.1 (136)	1,570	13.2 (97)
5102	25	*	*	16	*	*	4	*	5	*	25	*	25	*
5202	713	3.9 (54)	56.8 (30)	523	18.9 (76)	49.3 (20)	91	*	148	23.0 (14)	702	5.7 (20)	707	16.1 (138)
5206	4	*	*	4	*	*	0	*	0	*	5	*	5	*
5601	828	3.1 (26)	61.8 (97)	87	*	*	110	30.0 (8)	299	22.7 (12)	904	8.3 (101)	905	12.7 (89)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 95

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5801	1,496	4.2 (78)	60.9 (82)	984	20.5 (100)	49.8 (24)	243	52.3 (47)	245	35.9 (42)	1,620	7.8 (81)	1,621	15.4 (131)
5803	116	5.2 (125)	65.5 (133)	81	*	*	13	*	35	*	116	7.8 (81)	116	19.8 (152)
5804	16	*	*	13	*	*	1	*	4	*	14	*	14	*
5806	36	*	*	26	*	*	1	*	7	*	55	*	56	*
6101	30	*	*	24	*	*	1	*	6	*	29	*	29	*
6102	111	4.5 (97)	58.6 (50)	83	*	*	12	*	18	*	114	8.8 (113)	114	14.9 (128)
6103	3,366	3.9 (54)	63.5 (119)	2,390	19.5 (84)	50.3 (34)	363	44.1 (22)	331	26.6 (19)	3,414	8.5 (106)	3,414	16.6 (143)
6105	17	*	*	14	*	*	0	*	4	*	16	*	17	*
6106	127	3.9 (54)	52.8 (11)	89	*	*	16	*	14	*	122	7.4 (67)	122	22.1 (153)
6301	3,289	3.8 (49)	59.2 (56)	2,125	18.2 (62)	51.8 (55)	554	46.0 (30)	559	34.5 (38)	3,429	5.7 (20)	3,430	10.6 (54)
6312	1,739	3.7 (42)	59.6 (64)	1,184	16.6 (31)	51.4 (49)	291	51.5 (45)	367	42.5 (61)	1,760	4.0 (6)	1,760	8.3 (19)
6316	1,110	4.1 (69)	56.5 (28)	811	16.6 (31)	56.8 (112)	196	44.9 (25)	223	34.5 (38)	1,101	6.8 (54)	1,101	9.8 (37)
6318	18	*	*	8	*	*	8	*	3	*	33	*	33	*
6320	148	3.4 (32)	52.7 (9)	75	*	*	22	*	33	*	188	10.1 (136)	188	10.1 (43)
6325	198	6.1 (141)	55.1 (20)	92	*	*	33	*	62	*	230	5.2 (12)	230	14.8 (127)
6330	1	*	*	1	*	*	0	*	0	*	4	*	4	*
6335	645	4.7 (106)	51.9 (4)	227	23.3 (115)	42.3 (7)	119	41.2 (17)	161	32.9 (33)	1,000	9.9 (134)	1,000	8.3 (19)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 96

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6340	32	*	*	11	*	*	1	*	5	*	30	*	29	*
6345	59	*	*	39	*	*	10	*	20	*	71	*	71	*
6360	609	4.4 (91)	57.6 (37)	244	18.0 (58)	52.0 (56)	100	54.0 (50)	120	43.3 (65)	784	7.3 (66)	784	12.9 (92)
7304	32	*	*	19	*	*	2	*	9	*	37	*	37	*
7306	1,887	4.0 (65)	59.3 (58)	1,307	18.4 (67)	50.9 (44)	325	53.2 (48)	323	34.1 (36)	2,203	7.2 (64)	2,205	10.6 (54)
7308	41	*	*	28	*	*	4	*	1	*	37	*	37	*
7310	3	*	*	1	*	*	0	*	2	*	9	*	9	*
7401	2,104	5.4 (130)	56.8 (30)	1,801	18.1 (60)	55.8 (103)	242	44.2 (23)	318	36.5 (45)	2,026	6.9 (55)	2,030	13.9 (110)
7601	108	6.5 (145)	54.6 (17)	51	*	*	82	*	83	*	553	6.9 (55)	553	8.1 (17)
7850	1,057	4.2 (78)	62.7 (109)	871	19.7 (88)	48.3 (16)	129	39.5 (14)	178	35.4 (40)	1,018	9.1 (121)	1,020	13.6 (107)
7852	18	*	*	15	*	*	0	*	3	*	19	*	19	*
8101	1,439	3.4 (32)	65.7 (134)	977	20.6 (101)	50.3 (34)	240	63.3 (62)	193	54.9 (71)	1,665	6.7 (52)	1,668	15.8 (137)
8102	290	2.4 (11)	68.3 (146)	156	19.9 (92)	55.1 (95)	51	*	35	*	380	9.7 (133)	381	10.5 (52)
8105	1,068	3.8 (49)	65.1 (130)	588	18.7 (73)	51.7 (51)	233	59.7 (58)	236	43.2 (63)	1,152	8.2 (97)	1,151	11.5 (71)
8106	232	3.9 (54)	53.9 (16)	134	17.2 (42)	59.7 (121)	75	*	65	*	479	5.6 (16)	478	11.5 (71)
8150	19	*	*	8	*	*	4	*	7	*	20	*	20	*
8151	2,668	3.7 (42)	60.8 (79)	1,765	19.8 (89)	49.5 (23)	436	50.2 (42)	627	37.6 (47)	2,800	8.1 (93)	2,802	13.3 (100)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 97

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8152	48	*	*	33	*	*	9	*	14	*	47	*	47	*
8153	9	*	*	4	*	*	1	*	2	*	7	*	7	*
8155	11	*	*	7	*	*	1	*	6	*	12	*	12	*
8157	7	*	*	4	*	*	2	*	3	*	5	*	5	*
8201	1,834	5.0 (116)	58.8 (51)	1,140	17.3 (45)	52.9 (68)	364	48.9 (37)	545	30.8 (28)	2,416	6.5 (43)	2,415	10.1 (43)
8202	2,434	4.9 (112)	52.4 (6)	1,114	23.3 (115)	41.7 (6)	582	52.2 (46)	656	42.2 (60)	3,867	5.4 (13)	3,865	2.6 (2)
8203	346	3.5 (36)	63.3 (118)	279	12.5 (4)	55.2 (96)	66	*	80	*	367	7.1 (62)	367	17.4 (146)
8250	655	5.2 (125)	59.4 (61)	481	19.1 (77)	56.1 (106)	141	42.6 (20)	173	48.6 (69)	704	8.9 (114)	703	14.2 (114)
8252	122	5.7 (134)	64.8 (127)	68	*	*	30	*	28	*	149	6.0 (29)	149	9.4 (27)
8255	2,039	5.0 (116)	59.1 (55)	1,386	14.1 (9)	55.9 (105)	435	49.0 (38)	584	37.0 (46)	2,287	7.0 (59)	2,287	12.1 (79)
8257	1,230	4.1 (69)	56.2 (26)	646	25.4 (119)	37.2 (3)	267	53.6 (49)	411	42.1 (59)	2,001	9.6 (132)	2,002	5.5 (5)
8263	597	4.2 (78)	58.5 (46)	433	18.5 (69)	57.0 (113)	115	26.1 (5)	168	25.0 (16)	633	8.2 (97)	632	9.3 (25)
8264	489	3.7 (42)	62.2 (102)	318	17.0 (40)	53.1 (70)	118	40.7 (15)	161	24.2 (15)	648	6.6 (49)	648	9.7 (33)
8267	450	5.3 (128)	59.3 (58)	313	15.7 (23)	53.7 (79)	146	44.5 (24)	177	29.4 (25)	702	5.8 (22)	701	8.0 (15)
8901	929	2.7 (16)	63.7 (123)	612	18.1 (60)	53.3 (74)	150	64.7 (63)	98	*	971	5.1 (10)	972	13.3 (100)
8902	1,113	4.6 (101)	60.5 (75)	591	19.5 (84)	54.0 (81)	201	58.7 (56)	178	48.3 (68)	1,370	5.4 (13)	1,370	12.6 (88)
8903	1,296	3.0 (24)	64.9 (129)	759	16.9 (38)	57.7 (116)	216	70.8 (65)	144	62.5 (73)	1,428	5.6 (16)	1,426	18.4 (150)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 98

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8904	1,968	3.5 (36)	62.8 (111)	1,217	18.3 (65)	54.6 (89)	353	62.3 (60)	262	56.1 (72)	2,260	6.2 (37)	2,263	14.2 (114)
8905	1,142	2.8 (18)	59.6 (64)	621	18.2 (62)	50.1 (30)	203	49.3 (39)	206	43.2 (63)	1,440	8.7 (111)	1,442	10.9 (63)
8909	526	4.8 (109)	62.7 (109)	275	17.5 (49)	57.8 (117)	96	*	70	*	677	7.5 (71)	679	15.6 (135)
9045	1,252	1.7 (5)	67.2 (144)	857	22.8 (114)	46.3 (11)	249	45.4 (27)	182	40.7 (57)	1,644	7.5 (71)	1,645	7.5 (12)
9050	759	5.1 (123)	60.5 (75)	257	21.4 (106)	50.2 (32)	135	71.1 (66)	89	*	918	5.8 (22)	918	10.7 (58)
9051	1,193	2.6 (14)	67.8 (145)	732	17.5 (49)	46.0 (10)	148	33.1 (9)	134	27.6 (20)	1,504	8.2 (97)	1,503	7.8 (13)
9053	845	3.3 (29)	65.2 (131)	579	15.2 (16)	56.1 (106)	108	63.0 (61)	73	*	880	6.1 (33)	880	13.3 (100)
9055	19	*	*	12	*	*	2	*	2	*	19	*	19	*
9058	597	5.9 (137)	61.6 (91)	377	18.6 (71)	50.1 (30)	152	54.6 (52)	130	38.5 (52)	968	8.1 (93)	968	14.7 (125)
9066	773	2.8 (18)	63.5 (119)	387	16.8 (36)	56.6 (111)	160	51.3 (44)	119	49.6 (70)	1,085	6.1 (33)	1,088	17.9 (148)
9068	916	2.6 (14)	66.5 (142)	626	21.9 (109)	55.4 (99)	124	58.9 (57)	69	*	912	5.4 (13)	913	12.7 (89)
9069	2,188	3.3 (29)	62.0 (99)	1,620	19.6 (86)	50.2 (32)	394	56.6 (53)	200	42.5 (61)	2,504	6.0 (29)	2,503	10.7 (58)
9070	1,201	4.2 (78)	58.9 (52)	757	20.7 (103)	52.6 (62)	208	64.9 (64)	147	47.6 (67)	1,333	5.6 (16)	1,335	13.4 (103)
9072	1,162	4.5 (97)	62.4 (103)	655	19.8 (89)	51.6 (50)	238	56.7 (54)	163	45.4 (66)	1,376	6.5 (43)	1,377	16.2 (139)
9080	1,951	3.9 (54)	62.8 (111)	1,134	15.5 (21)	57.9 (118)	431	60.6 (59)	343	39.4 (56)	2,526	6.5 (43)	2,526	11.6 (74)
9082	138	3.6 (39)	63.8 (124)	68	*	*	21	*	15	*	136	6.6 (49)	136	16.2 (139)
9083	100	1.0 (2)	73.0 (148)	95	*	*	3	*	1	*	101	5.9 (25)	101	2.0 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 99

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9084	27	*	*	21	*	*	3	*	1	*	33	*	33	*
9085	49	*	*	17	*	*	22	*	5	*	44	*	44	*
9086	46	*	*	25	*	*	15	*	8	*	40	*	40	*
9087	5	*	*	3	*	*	0	*	1	*	4	*	4	*
Michigan	125,954	4.1	60.3	78,065	18.8	51.5	21,022	46.2	25,209	34.2	146,794	7.5	146,883	11.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 100

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	108	88.0 (57)	6.5 (84)	116	22.4 (130)	105	27.6 (37)	89	*
0102	440	85.5 (71)	8.6 (101)	476	33.4 (70)	440	20.0 (84)	403	89.8 (62)
0103	365	79.7 (113)	13.2 (124)	402	36.3 (51)	360	21.1 (78)	329	90.3 (56)
0104	130	82.3 (95)	9.2 (105)	139	38.8 (28)	124	19.4 (87)	110	91.8 (40)
0105	20	*	*	19	*	16	*	17	*
0250	125	77.6 (128)	17.6 (145)	144	50.7 (1)	105	28.6 (31)	124	90.3 (56)
0251	81	*	*	88	*	76	*	68	*
0252	94	*	*	103	30.1 (95)	89	*	76	*
0253	126	77.8 (126)	10.3 (110)	132	47.0 (6)	123	16.3 (106)	115	89.6 (64)
0401	411	82.5 (94)	12.4 (121)	454	40.7 (21)	419	22.9 (65)	372	92.5 (30)
0402	361	79.5 (115)	11.9 (119)	379	34.3 (64)	336	26.2 (46)	305	90.5 (53)
0403	144	85.4 (73)	2.1 (12)	154	28.6 (102)	130	23.1 (63)	113	93.8 (16)
0404	106	93.4 (7)	2.8 (26)	110	40.0 (23)	97	*	83	*
0501	465	93.5 (6)	1.9 (9)	489	33.5 (69)	424	30.7 (25)	360	90.3 (56)
0503	105	88.6 (55)	5.7 (73)	105	30.5 (91)	92	*	80	*
0504	434	89.2 (45)	5.3 (66)	453	27.4 (107)	419	33.2 (17)	344	89.0 (69)
0505	279	86.0 (69)	7.9 (94)	293	36.2 (52)	278	27.3 (40)	245	89.0 (69)
0506	162	84.6 (79)	6.2 (79)	167	38.3 (33)	156	21.2 (77)	145	79.3 (139)
0507	75	*	*	75	*	65	*	48	*
0508	28	*	*	29	*	26	*	22	*
0509	3	*	*	3	*	2	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 101

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	1,006	85.3 (75)	8.3 (97)	1,080	35.2 (59)	936	27.1 (41)	868	91.4 (44)
0651	550	77.5 (131)	13.8 (127)	586	44.2 (10)	499	22.8 (66)	493	87.4 (86)
0652	312	82.7 (93)	10.6 (111)	324	33.0 (73)	287	28.6 (31)	245	83.7 (122)
0653	393	72.3 (145)	22.4 (153)	414	42.8 (13)	384	31.3 (23)	340	88.2 (79)
0654	345	76.8 (135)	17.4 (143)	371	41.0 (17)	337	23.1 (63)	300	80.0 (137)
0655	304	87.2 (59)	4.6 (51)	337	42.4 (15)	302	25.5 (49)	276	90.9 (48)
0657	62	*	*	63	*	54	*	50	*
0701	384	79.4 (116)	14.1 (131)	418	33.7 (68)	373	38.3 (14)	324	92.3 (35)
0702	67	*	*	72	*	67	*	55	*
0703	164	92.7 (12)	1.8 (8)	184	32.6 (77)	162	30.2 (27)	138	98.6 (1)
0704	50	*	*	56	*	53	*	48	*
0705	106	88.7 (53)	7.5 (91)	114	31.6 (84)	107	59.8 (3)	100	89.0 (69)
0706	13	*	*	13	*	11	*	9	*
0850	722	87.1 (60)	3.7 (39)	794	30.5 (91)	717	23.7 (59)	593	92.4 (32)
0851	649	89.1 (50)	7.1 (90)	708	32.1 (80)	657	23.9 (55)	549	95.4 (5)
0901	1,444	74.3 (142)	19.8 (146)	1,554	40.9 (18)	1,434	14.4 (122)	1,298	85.4 (113)
0902	38	*	*	43	*	43	*	38	*
1051	1,130	79.3 (117)	13.8 (127)	1,201	28.1 (105)	1,065	23.8 (56)	844	87.3 (88)
1101	1,604	71.6 (146)	11.3 (115)	1,785	28.7 (100)	1,549	14.5 (121)	1,314	86.5 (97)
1102	716	74.7 (141)	13.8 (127)	833	34.9 (61)	738	20.5 (81)	629	94.4 (9)
1104	1	*	*	1	*	1	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 102

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
1106	66	*	*	79	*	74	*	60 *
1250	837	81.0 (103)	9.9 (107)	1,050	33.0 (73)	922	22.0 (72)	819 83.3 (123)
1251	673	80.4 (108)	11.3 (115)	727	35.1 (60)	629	17.0 (100)	551 86.0 (107)
1252	505	69.3 (150)	17.0 (142)	562	39.0 (27)	524	18.9 (90)	466 90.1 (60)
1253	684	71.5 (147)	17.4 (143)	780	33.8 (67)	702	18.8 (91)	617 88.8 (74)
1350	2,068	90.2 (30)	4.9 (56)	2,336	35.3 (57)	1,595	22.4 (69)	1,752 80.9 (132)
1351	110	83.6 (88)	10.0 (108)	117	31.6 (84)	105	26.7 (42)	87 *
1352	359	91.9 (18)	5.3 (66)	382	35.6 (55)	346	2.6 (149)	304 89.5 (65)
1401	215	83.7 (86)	10.7 (112)	229	31.4 (86)	180	32.8 (18)	178 88.8 (74)
1402	109	78.9 (120)	14.7 (137)	117	36.8 (46)	95	*	79 *
1403	24	*	*	22	*	16	*	15 *
1501	719	83.6 (88)	10.7 (112)	799	36.9 (44)	512	19.3 (88)	622 83.9 (121)
1503	877	83.5 (90)	8.9 (104)	958	42.8 (13)	796	32.0 (20)	777 80.2 (136)
1504	1,856	90.8 (23)	3.0 (29)	2,053	31.3 (88)	1,678	28.1 (34)	1,472 80.4 (134)
1505	361	89.2 (45)	6.4 (83)	393	28.5 (104)	302	25.8 (47)	293 89.8 (62)
1506	525	73.3 (143)	21.3 (150)	577	37.4 (40)	506	6.5 (148)	461 94.6 (8)
1507	518	89.8 (37)	5.4 (69)	570	37.4 (40)	447	24.4 (54)	428 90.0 (61)
1508	40	*	*	50	*	42	*	44 *
1509	355	90.4 (26)	4.2 (48)	417	38.1 (35)	339	19.2 (89)	326 87.1 (92)
1511	631	92.4 (13)	2.5 (19)	669	34.7 (62)	548	16.8 (102)	510 87.3 (88)
1512	331	92.1 (17)	4.8 (54)	354	24.3 (118)	272	31.3 (23)	237 92.4 (32)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 103

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1513	18	*	*	21	*	21	*	21	*
1514	281	92.9 (11)	5.0 (59)	291	22.7 (127)	210	32.4 (19)	151	94.0 (13)
1701	590	89.2 (45)	6.9 (88)	636	44.2 (10)	559	41.5 (9)	523	83.2 (125)
1950	564	78.2 (125)	15.4 (139)	601	36.4 (50)	517	23.8 (56)	462	92.9 (26)
1952	129	79.1 (118)	16.3 (140)	131	32.8 (75)	115	25.2 (51)	99	*
1953	287	81.9 (96)	14.3 (134)	305	37.0 (43)	276	22.5 (67)	245	94.7 (7)
1954	481	80.0 (110)	12.9 (123)	513	38.0 (36)	459	26.4 (45)	417	94.0 (13)
1955	98	*	*	103	40.8 (20)	88	*	78	*
1957	397	86.1 (67)	6.3 (81)	418	31.3 (88)	335	30.4 (26)	305	88.9 (73)
1963	49	*	*	51	*	38	*	30	*
1964	31	*	*	36	*	33	*	30	*
2150	586	92.2 (16)	1.5 (7)	643	40.9 (18)	591	25.5 (49)	525	93.0 (25)
2151	180	83.9 (84)	12.2 (120)	223	33.2 (71)	164	27.4 (39)	147	90.5 (53)
2152	4	*	*	3	*	2	*	0	*
2251	372	89.8 (37)	6.7 (86)	402	36.8 (46)	373	29.2 (30)	329	93.6 (18)
2252	164	83.5 (90)	12.8 (122)	187	46.5 (8)	157	54.1 (5)	156	87.8 (82)
2554	50	*	*	51	*	42	*	37	*
2555	6,922	86.4 (65)	4.1 (46)	7,459	25.4 (114)	5,864	7.1 (145)	5,453	87.8 (82)
2556	316	78.8 (121)	5.4 (69)	337	17.8 (150)	232	14.2 (124)	162	83.3 (123)
2557	7	*	*	9	*	6	*	7	*
2558	4	*	*	6	*	5	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 104

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2559	3	*	*	3	*	3	*	3	*
2560	4	*	*	6	*	6	*	6	*
2561	3	*	*	3	*	3	*	3	*
2562	14	*	*	15	*	12	*	14	*
2563	26	*	*	25	*	14	*	9	*
2564	18	*	*	22	*	19	*	17	*
3001	1,067	77.6 (128)	7.7 (92)	1,142	33.9 (66)	988	11.5 (132)	915	81.2 (131)
3002	277	90.3 (28)	1.1 (5)	299	42.1 (16)	274	9.5 (137)	261	85.8 (108)
3004	193	90.2 (30)	4.7 (52)	213	50.7 (1)	199	20.1 (83)	175	74.9 (143)
3005	62	*	*	66	*	62	*	50	*
3007	12	*	*	15	*	13	*	12	*
3009	0	*	*	3	*	2	*	2	*
3201	409	81.9 (96)	8.8 (102)	429	32.2 (79)	364	31.6 (21)	311	89.4 (67)
3203	1	*	*	1	*	1	*	1	*
3301	3,968	86.8 (62)	6.1 (76)	4,238	29.0 (97)	3,769	6.8 (146)	3,194	90.6 (51)
3304	94	*	*	102	38.2 (34)	99	*	77	*
3305	151	89.4 (41)	7.9 (94)	164	32.3 (78)	153	8.5 (143)	121	90.9 (48)
3306	247	84.6 (79)	8.5 (100)	262	31.7 (83)	228	8.8 (142)	192	95.8 (4)
3308	86	*	*	92	*	77	*	56	*
3309	18	*	*	19	*	18	*	16	*
3312	364	86.5 (64)	4.9 (56)	393	30.5 (91)	336	9.2 (138)	301	90.4 (55)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 105

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3550	54	*	*	59	*	54	*	51	*
3551	18	*	*	20	*	20	*	18	*
3552	40	*	*	46	*	42	*	41	*
3701	930	93.0 (10)	3.9 (43)	1,016	32.7 (76)	906	21.3 (76)	754	94.4 (9)
3850	2,812	85.1 (76)	7.9 (94)	3,029	39.1 (26)	2,654	17.2 (99)	2,449	85.8 (108)
3901	1,360	83.8 (85)	6.5 (84)	1,505	38.7 (30)	1,347	13.3 (125)	1,203	79.7 (138)
3902	276	81.2 (102)	10.9 (114)	317	49.5 (4)	311	15.8 (112)	300	82.3 (128)
3903	31	*	*	31	*	29	*	16	*
3904	7	*	*	7	*	7	*	7	*
3905	11	*	*	11	*	10	*	9	*
3950	1,935	84.1 (83)	8.4 (99)	2,156	33.2 (71)	1,908	17.9 (94)	1,610	86.1 (105)
3954	16	*	*	17	*	16	*	12	*
4001	77	*	*	91	*	91	*	92	*
4002	48	*	*	58	*	55	*	58	*
4103	13	*	*	14	*	16	*	16	*
4106	2,265	86.4 (65)	4.9 (56)	2,469	25.7 (112)	2,100	8.1 (144)	1,900	86.5 (97)
4107	2,793	84.4 (82)	6.3 (81)	3,009	24.0 (120)	2,376	23.8 (56)	2,189	91.4 (44)
4109	855	80.0 (110)	13.8 (127)	928	31.8 (82)	845	17.6 (95)	684	93.6 (18)
4110	7	*	*	7	*	7	*	0	*
4112	508	46.9 (153)	0.0 (1)	534	25.8 (111)	501	83.6 (1)	476	92.0 (38)
4113	112	90.2 (30)	0.0 (1)	116	37.1 (42)	116	12.9 (126)	116	91.4 (44)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 106

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4115	3,253	89.0 (52)	5.0 (59)	3,538	23.9 (121)	3,036	16.1 (107)	2,625	91.7 (41)
4351	510	89.2 (45)	4.1 (46)	542	40.0 (23)	180	52.2 (6)	453	92.9 (26)
4352	147	64.6 (152)	21.1 (148)	156	47.4 (5)	139	12.9 (126)	130	81.5 (130)
4353	364	73.1 (144)	21.2 (149)	415	38.8 (28)	372	21.8 (73)	324	79.0 (140)
4354	696	89.9 (35)	5.2 (64)	743	38.4 (32)	592	20.3 (82)	586	90.6 (51)
4355	311	87.8 (58)	10.0 (108)	323	34.7 (62)	283	74.6 (2)	251	92.8 (28)
4356	390	80.8 (105)	11.5 (118)	419	36.0 (53)	370	28.1 (34)	336	92.3 (35)
4358	101	65.3 (151)	21.8 (152)	109	32.1 (80)	102	28.4 (33)	93	*
4359	465	71.0 (148)	20.6 (147)	506	36.6 (49)	428	23.4 (61)	387	91.5 (43)
4360	199	75.9 (138)	16.6 (141)	218	43.6 (12)	181	29.8 (29)	179	87.2 (91)
4361	234	91.0 (22)	4.3 (49)	253	37.9 (37)	233	27.5 (38)	206	90.8 (50)
4362	284	79.9 (112)	14.1 (131)	295	36.9 (44)	263	22.1 (70)	237	84.8 (119)
4363	762	77.2 (132)	14.4 (136)	814	39.6 (25)	686	29.9 (28)	650	86.2 (101)
4401	922	80.2 (109)	13.2 (124)	972	29.9 (96)	887	16.6 (105)	738	93.9 (15)
4501	888	78.8 (121)	13.6 (126)	922	35.6 (55)	819	26.7 (42)	699	93.8 (16)
4601	1,211	89.3 (42)	5.0 (59)	1,333	27.0 (109)	1,160	22.1 (70)	992	88.6 (77)
4602	10	*	*	10	*	8	*	8	*
4701	979	89.3 (42)	4.7 (52)	1,046	23.0 (124)	787	38.8 (12)	703	92.2 (37)
5050	1,496	91.4 (20)	2.7 (22)	1,597	18.5 (143)	1,093	15.1 (117)	1,103	94.3 (11)
5051	2,495	91.9 (18)	2.6 (21)	2,662	20.4 (135)	1,987	14.3 (123)	1,795	93.4 (21)
5052	2,731	89.9 (35)	3.1 (30)	2,898	22.5 (129)	1,979	11.8 (130)	2,065	93.4 (21)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 107

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5055	1,130	84.8 (78)	14.1 (131)	1,169	50.2 (3)	872	9.7 (136)	995	84.9 (118)
5101	1,558	90.4 (26)	4.0 (45)	1,676	24.2 (119)	1,041	26.7 (42)	1,204	93.4 (21)
5102	25	*	*	24	*	21	*	14	*
5202	693	89.2 (45)	5.9 (74)	782	35.3 (57)	585	36.1 (16)	611	90.3 (56)
5206	5	*	*	4	*	1	*	0	*
5601	907	90.0 (34)	5.2 (64)	963	29.0 (97)	868	31.5 (22)	725	94.9 (6)
5801	1,617	81.4 (100)	9.6 (106)	1,768	31.1 (90)	1,597	20.7 (79)	1,357	88.0 (81)
5803	115	88.7 (53)	6.1 (76)	122	27.9 (106)	112	16.1 (107)	82	*
5804	14	*	*	17	*	15	*	8	*
5806	55	*	*	59	*	43	*	28	*
6101	29	*	*	30	*	29	*	24	*
6102	112	89.3 (42)	3.6 (38)	117	35.9 (54)	100	25.0 (52)	91	*
6103	3,355	87.1 (60)	6.0 (75)	3,637	38.7 (30)	2,998	19.5 (86)	2,826	82.1 (129)
6105	16	*	*	18	*	16	*	12	*
6106	118	90.7 (25)	3.4 (35)	133	46.6 (7)	121	15.7 (113)	111	82.9 (126)
6301	3,384	86.1 (67)	6.9 (88)	3,733	27.4 (107)	2,967	11.1 (134)	2,786	86.4 (99)
6312	1,712	89.1 (50)	4.4 (50)	1,890	23.9 (121)	1,452	17.6 (95)	1,358	86.2 (101)
6316	1,100	90.1 (33)	3.1 (30)	1,201	24.7 (116)	992	21.8 (73)	851	93.3 (24)
6318	33	*	*	36	*	27	*	20	*
6320	183	95.1 (3)	2.7 (22)	200	18.0 (148)	103	38.8 (12)	118	94.1 (12)
6325	227	93.4 (7)	3.5 (36)	257	22.6 (128)	211	15.2 (116)	158	91.1 (47)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 108

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6330	4	*	*	4	*	2	*	4	*
6335	994	89.6 (39)	8.8 (102)	1,034	37.7 (39)	260	28.1 (34)	806	92.6 (29)
6340	30	*	*	37	*	26	*	31	*
6345	69	*	*	79	*	64	*	48	*
6360	767	85.9 (70)	6.8 (87)	849	25.2 (115)	533	24.8 (53)	618	87.7 (84)
7304	36	*	*	38	*	34	*	33	*
7306	2,173	80.8 (105)	8.3 (97)	2,355	31.4 (86)	2,038	16.7 (103)	1,758	86.9 (95)
7308	36	*	*	45	*	44	*	43	*
7310	10	*	*	12	*	11	*	11	*
7401	2,023	70.2 (149)	21.7 (151)	2,202	37.9 (37)	1,847	20.0 (84)	1,770	85.8 (108)
7601	542	86.7 (63)	7.7 (92)	621	40.4 (22)	535	44.9 (7)	512	89.5 (65)
7850	1,009	79.1 (118)	15.1 (138)	1,111	36.7 (48)	934	18.8 (91)	875	87.1 (92)
7852	19	*	*	20	*	17	*	17	*
8101	1,657	78.3 (123)	2.7 (22)	1,726	19.2 (141)	967	15.9 (110)	1,152	86.2 (101)
8102	374	80.7 (107)	5.1 (63)	397	30.5 (91)	227	20.7 (79)	300	80.7 (133)
8105	1,156	89.5 (40)	2.2 (13)	1,272	24.6 (117)	1,152	16.1 (107)	954	92.5 (30)
8106	469	77.0 (134)	5.3 (66)	507	20.5 (134)	384	8.9 (141)	336	86.6 (96)
8150	20	*	*	21	*	21	*	20	*
8151	2,797	90.8 (23)	3.3 (32)	3,053	28.6 (102)	2,708	15.5 (115)	2,288	91.7 (41)
8152	47	*	*	53	*	41	*	49	*
8153	7	*	*	8	*	7	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 109

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8155	12	*	*	14	*	12	*	12	*
8157	5	*	*	7	*	6	*	6	*
8201	2,399	92.4 (13)	1.9 (9)	2,628	18.1 (147)	1,834	36.9 (15)	1,668	88.2 (79)
8202	3,811	94.0 (4)	3.9 (43)	3,999	28.8 (99)	2,509	10.4 (135)	3,097	87.4 (86)
8203	367	92.4 (13)	2.5 (19)	401	18.0 (148)	324	22.5 (67)	271	97.0 (2)
8250	699	84.5 (81)	2.7 (22)	779	15.5 (154)	611	23.6 (60)	546	87.0 (94)
8252	147	74.8 (140)	5.4 (69)	157	17.8 (150)	131	11.5 (132)	113	92.0 (38)
8255	2,262	91.4 (20)	0.9 (4)	2,493	18.2 (145)	1,505	40.2 (10)	1,694	92.4 (32)
8257	1,982	96.7 (1)	1.3 (6)	2,088	28.7 (100)	1,746	17.0 (100)	1,610	85.2 (116)
8263	632	95.7 (2)	0.6 (3)	697	17.4 (152)	317	43.2 (8)	476	96.2 (3)
8264	645	94.0 (4)	1.9 (9)	723	21.0 (133)	560	23.4 (61)	489	93.5 (20)
8267	693	88.2 (56)	3.5 (36)	774	19.6 (138)	651	55.8 (4)	565	88.5 (78)
8901	964	81.7 (99)	3.3 (32)	1,032	21.7 (131)	646	14.7 (119)	756	82.7 (127)
8902	1,365	84.9 (77)	2.3 (16)	1,473	21.2 (132)	913	17.4 (97)	1,045	86.2 (101)
8903	1,405	78.3 (123)	3.3 (32)	1,515	22.9 (125)	1,059	18.2 (93)	1,068	78.7 (141)
8904	2,244	83.1 (92)	2.4 (18)	2,398	18.2 (145)	1,381	17.4 (97)	1,631	87.3 (88)
8905	1,429	90.3 (28)	2.3 (16)	1,514	26.4 (110)	1,074	25.7 (48)	1,104	86.1 (105)
8909	675	85.5 (71)	2.2 (13)	715	19.3 (139)	484	16.7 (103)	482	89.0 (69)
9045	1,644	76.7 (136)	11.3 (115)	1,763	34.1 (65)	540	39.6 (11)	1,406	85.4 (113)
9050	897	77.7 (127)	6.1 (76)	949	22.9 (125)	766	21.8 (73)	679	84.1 (120)
9051	1,455	77.2 (132)	14.3 (134)	1,571	45.6 (9)	1,439	14.8 (118)	1,336	78.6 (142)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 110

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9053	879	81.9 (96)	3.8 (40)	929	20.0 (136)	730	12.3 (129)	568	86.4 (99)
9055	19	*	*	18	*	10	*	6	*
9058	967	85.4 (73)	5.4 (69)	1,012	23.2 (123)	717	6.6 (147)	708	87.7 (84)
9066	1,046	75.5 (139)	5.0 (59)	1,145	15.5 (154)	729	11.7 (131)	665	85.3 (115)
9068	911	79.7 (113)	3.8 (40)	955	25.5 (113)	761	14.6 (120)	710	85.8 (108)
9069	2,509	80.9 (104)	2.9 (27)	2,588	16.5 (153)	1,455	9.2 (138)	1,608	89.1 (68)
9070	1,333	76.2 (137)	6.2 (79)	1,412	19.3 (139)	875	15.7 (113)	932	85.8 (108)
9072	1,360	77.6 (128)	4.8 (54)	1,447	19.8 (137)	1,213	9.2 (138)	1,131	85.0 (117)
9080	2,510	81.4 (100)	3.8 (40)	2,633	19.1 (142)	1,785	15.9 (110)	1,875	88.7 (76)
9082	135	83.7 (86)	2.2 (13)	147	18.4 (144)	134	12.7 (128)	102	80.4 (134)
9083	102	93.1 (9)	2.9 (27)	103	1.9 (156)	75	*	10	*
9084	33	*	*	35	*	19	*	13	*
9085	42	*	*	49	*	46	*	49	*
9086	40	*	*	48	*	47	*	48	*
9087	4	*	*	5	*	5	*	4	*
<b>Michigan</b>	<b>145,457</b>	<b>84.9</b>	<b>6.6</b>	<b>157,300</b>	<b>29.1</b>	<b>123,217</b>	<b>19.5</b>	<b>117,076</b>	<b>88.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 7A

### 2018 - 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic Combined 3 Years

Page : 111

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	105	46.7 (145)	107	30.8 (145)	38	*	89	*	107	1.9 (59)
0102	441	46.3 (142)	445	33.9 (150)	169	20.1 (90)	403	19.9 (137)	445	2.7 (85)
0103	363	40.5 (127)	369	29.0 (140)	121	22.3 (88)	329	20.4 (139)	368	6.3 (146)
0104	126	42.1 (130)	130	29.2 (141)	44	*	110	14.5 (101)	130	3.8 (110)
0105	17	*	20	*	6	*	16	*	20	*
0250	133	44.4 (136)	125	26.4 (133)	47	*	124	15.3 (111)	125	0.8 (11)
0251	77	*	82	*	31	*	68	*	82	*
0252	91	*	94	*	23	*	76	*	94	*
0253	123	46.3 (142)	126	27.8 (135)	50	*	115	19.1 (133)	126	2.4 (78)
0401	423	47.8 (147)	418	28.7 (139)	163	34.4 (61)	372	12.4 (86)	417	2.4 (78)
0402	338	45.3 (138)	353	24.9 (123)	127	36.2 (56)	305	13.1 (93)	355	4.2 (122)
0403	130	46.2 (140)	144	29.9 (143)	49	*	113	18.6 (129)	145	6.9 (147)
0404	98	*	106	25.5 (127)	29	*	83	*	106	2.8 (88)
0501	428	38.8 (119)	464	26.3 (132)	131	31.3 (75)	360	19.4 (134)	464	5.0 (133)
0503	93	*	105	24.8 (121)	33	*	80	*	105	7.6 (149)
0504	422	39.6 (124)	433	20.3 (98)	134	37.3 (48)	344	8.4 (55)	433	2.3 (73)
0505	278	38.8 (119)	276	21.0 (101)	97	*	244	10.7 (76)	274	4.7 (128)
0506	157	43.3 (133)	162	32.1 (148)	60	*	145	24.1 (143)	162	6.2 (144)
0507	65	*	76	*	18	*	48	*	76	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 112

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0508	26	*	28	*	8	*	22	*	28	*
0509	2	*	3	*	2	*	2	*	3	*
0650	997	36.5 (104)	1,003	21.5 (107)	297	38.0 (45)	868	12.4 (86)	1,004	3.1 (93)
0651	534	48.5 (148)	552	31.7 (147)	222	27.5 (84)	493	17.8 (125)	552	3.3 (101)
0652	288	44.8 (137)	315	28.3 (138)	109	33.0 (67)	245	22.4 (142)	315	8.6 (151)
0653	389	38.6 (118)	393	23.4 (116)	122	28.7 (80)	340	15.0 (107)	393	3.1 (93)
0654	337	40.4 (126)	345	25.5 (127)	109	29.4 (77)	300	20.3 (138)	345	4.6 (127)
0655	306	45.4 (139)	310	31.6 (146)	110	34.5 (60)	276	18.5 (128)	310	5.5 (139)
0657	54	*	62	*	26	*	50	*	62	*
0701	379	31.4 (84)	383	21.4 (106)	99	*	323	5.3 (20)	383	0.8 (11)
0702	67	*	68	*	18	*	55	*	68	*
0703	163	46.6 (144)	168	35.1 (152)	62	*	138	15.9 (117)	168	3.0 (92)
0704	53	*	50	*	16	*	48	*	50	*
0705	111	40.5 (127)	105	33.3 (149)	37	*	100	15.0 (107)	105	3.8 (110)
0706	11	*	13	*	3	*	9	*	13	*
0850	717	39.9 (125)	728	20.5 (100)	223	39.9 (39)	593	7.9 (46)	728	2.1 (63)
0851	658	36.9 (108)	653	18.7 (90)	188	44.1 (32)	548	8.9 (61)	653	2.5 (81)
0901	1,438	38.2 (114)	1,452	23.4 (116)	485	28.7 (80)	1,297	13.3 (94)	1,451	4.0 (117)
0902	43	*	38	*	11	*	38	*	38	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 113

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1051	1,066	35.0 (98)	1,132	19.6 (95)	304	37.2 (49)	844	10.0 (70)	1,132	2.8 (88)
1101	1,553	26.4 (71)	1,651	13.0 (62)	319	42.3 (37)	1,314	12.9 (89)	1,651	3.8 (110)
1102	743	37.6 (112)	734	19.9 (96)	200	39.0 (43)	629	14.8 (106)	735	3.9 (113)
1104	1	*	1	*	0	*	0	*	1	*
1106	74	*	67	*	14	*	60	*	67	*
1250	954	30.7 (80)	979	17.6 (86)	231	32.9 (68)	817	11.3 (78)	979	1.6 (44)
1251	633	43.4 (134)	679	29.2 (141)	224	29.5 (76)	551	17.1 (122)	678	4.0 (117)
1252	524	32.1 (87)	522	17.8 (88)	137	37.2 (49)	465	12.9 (89)	521	3.1 (93)
1253	702	38.2 (114)	712	21.1 (103)	209	36.8 (52)	616	10.6 (75)	709	3.1 (93)
1350	2,050	29.1 (76)	2,111	17.4 (84)	474	27.0 (86)	1,768	9.7 (65)	2,119	2.0 (62)
1351	107	34.6 (95)	109	17.4 (84)	29	*	87	*	110	0.9 (19)
1352	345	30.7 (80)	358	15.9 (81)	96	*	303	2.0 (6)	361	0.8 (11)
1401	208	42.8 (131)	216	27.8 (135)	76	*	179	14.0 (96)	216	4.2 (122)
1402	104	33.7 (91)	110	11.8 (60)	25	*	80	*	110	0.9 (19)
1403	21	*	24	*	8	*	15	*	24	*
1501	701	19.5 (35)	721	10.4 (51)	114	27.2 (85)	617	4.5 (17)	721	1.2 (29)
1503	870	25.1 (66)	898	13.5 (65)	192	31.8 (72)	777	5.7 (25)	898	1.6 (44)
1504	1,807	18.2 (30)	1,891	9.5 (43)	271	46.9 (24)	1,478	3.1 (11)	1,891	0.6 (8)
1505	353	23.2 (57)	365	11.5 (59)	72	*	289	3.8 (12)	365	3.6 (107)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 114

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1506	525	23.8 (59)	529	15.3 (77)	103	13.6 (94)	460	3.9 (13)	530	0.9 (19)
1507	502	31.9 (86)	527	13.1 (63)	136	45.6 (28)	424	9.9 (68)	527	2.7 (85)
1508	44	*	41	*	1	*	44	*	41	*
1509	365	24.7 (64)	354	13.8 (68)	76	*	326	8.6 (56)	354	1.7 (55)
1511	596	35.1 (99)	632	19.3 (93)	169	34.3 (63)	512	8.2 (50)	633	2.1 (63)
1512	301	25.2 (67)	329	14.6 (71)	68	*	239	2.9 (10)	330	0.9 (19)
1513	21	*	17	*	9	*	21	*	17	*
1514	261	14.2 (14)	281	7.8 (34)	19	*	146	0.7 (3)	277	0.4 (4)
1701	593	40.5 (127)	586	30.2 (144)	203	16.3 (93)	523	18.4 (126)	587	10.1 (153)
1950	536	43.1 (132)	570	27.0 (134)	187	31.6 (73)	462	18.4 (126)	571	5.3 (138)
1952	115	33.9 (93)	129	21.7 (108)	34	*	99	*	129	2.3 (73)
1953	279	39.4 (122)	288	21.9 (110)	87	*	245	14.3 (99)	288	3.5 (104)
1954	469	36.0 (103)	481	21.0 (101)	142	35.9 (57)	418	9.8 (66)	482	1.5 (38)
1955	90	*	98	*	28	*	78	*	98	*
1957	368	32.9 (89)	396	17.7 (87)	103	43.7 (34)	305	8.2 (50)	399	1.3 (31)
1963	41	*	49	*	8	*	30	*	50	*
1964	34	*	31	*	10	*	30	*	31	*
2150	591	46.2 (140)	588	24.8 (121)	218	37.2 (49)	525	13.5 (95)	588	2.7 (85)
2151	191	43.5 (135)	188	25.0 (125)	53	*	147	15.6 (115)	187	3.2 (97)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 115

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2152	2	*	4	*	0	*	0	*	4	*
2251	376	37.0 (109)	368	26.1 (130)	118	21.2 (89)	329	7.9 (46)	368	1.4 (35)
2252	166	47.0 (146)	156	37.2 (153)	62	*	155	16.1 (119)	155	5.8 (141)
2554	43	*	48	*	14	*	37	*	48	*
2555	6,498	26.2 (70)	6,987	14.8 (74)	1,361	36.4 (54)	5,448	8.6 (56)	6,977	2.1 (63)
2556	271	20.3 (41)	330	14.8 (74)	38	*	163	9.8 (66)	330	2.1 (63)
2557	7	*	8	*	1	*	7	*	8	*
2558	6	*	5	*	0	*	6	*	5	*
2559	3	*	3	*	1	*	3	*	3	*
2560	6	*	4	*	1	*	6	*	4	*
2561	3	*	3	*	2	*	3	*	3	*
2562	14	*	14	*	1	*	14	*	14	*
2563	16	*	26	*	2	*	9	*	26	*
2564	20	*	18	*	5	*	17	*	18	*
3001	991	33.7 (91)	1,074	14.9 (76)	290	47.6 (22)	915	11.7 (82)	1,074	3.4 (103)
3002	274	24.8 (65)	280	11.4 (58)	62	*	260	9.6 (64)	279	2.2 (70)
3004	200	21.5 (49)	192	9.9 (47)	31	*	175	8.6 (56)	193	1.6 (44)
3005	62	*	61	*	19	*	50	*	61	*
3007	13	*	13	*	1	*	12	*	13	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 7A

### 2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 116

#### Michigan

#### Comparison of Smoking Indicators by Clinic

#### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3009	2	*	1	*	0	*	2	*	1	*
3201	392	36.5 (104)	411	22.4 (112)	121	36.4 (54)	312	15.4 (112)	411	4.9 (131)
3203	1	*	1	*	0	*	1	*	1	*
3301	3,789	17.6 (27)	4,003	10.9 (56)	534	33.3 (66)	3,180	6.4 (30)	3,996	3.2 (97)
3304	99	*	95	*	4	*	76	*	95	*
3305	155	13.5 (11)	151	6.6 (25)	12	*	121	2.5 (8)	150	0.7 (9)
3306	230	21.7 (50)	250	9.2 (42)	43	*	193	7.3 (37)	248	1.2 (29)
3308	79	*	87	*	14	*	56	*	87	*
3309	18	*	18	*	6	*	16	*	18	*
3312	336	20.5 (43)	369	13.8 (68)	60	*	301	4.3 (16)	368	1.6 (44)
3550	54	*	54	*	26	*	51	*	54	*
3551	20	*	18	*	8	*	18	*	18	*
3552	42	*	40	*	24	*	41	*	40	*
3701	905	34.5 (94)	935	21.1 (103)	259	29.0 (78)	754	8.6 (56)	938	2.3 (73)
3850	2,740	37.4 (111)	2,838	24.7 (120)	874	28.8 (79)	2,447	10.7 (76)	2,839	2.1 (63)
3901	1,359	24.4 (61)	1,364	12.9 (61)	268	35.1 (58)	1,200	7.3 (37)	1,361	1.6 (44)
3902	312	22.8 (56)	279	14.7 (72)	62	*	300	7.3 (37)	280	1.4 (35)
3903	29	*	31	*	4	*	16	*	31	*
3904	7	*	7	*	2	*	7	*	7	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 117

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3905	10	*	11	*	2	*	9	*	10	*
3950	1,917	24.2 (60)	1,947	13.7 (67)	349	31.5 (74)	1,607	7.7 (42)	1,946	1.1 (25)
3954	16	*	16	*	4	*	12	*	16	*
4001	92	*	79	*	24	*	92	*	79	*
4002	58	*	48	*	13	*	58	*	48	*
4103	16	*	13	*	3	*	16	*	13	*
4106	2,138	18.0 (29)	2,298	8.5 (38)	316	39.6 (41)	1,891	2.5 (8)	2,294	0.8 (11)
4107	2,678	21.9 (52)	2,807	10.7 (55)	416	46.2 (26)	2,190	8.2 (50)	2,803	2.3 (73)
4109	846	28.6 (74)	866	14.0 (70)	189	45.0 (29)	682	5.3 (20)	864	0.9 (19)
4110	7	*	7	*	0	*	0	*	7	*
4112	508	3.3 (3)	510	1.0 (1)	16	*	408	8.3 (53)	493	2.6 (82)
4113	116	3.4 (4)	112	3.6 (8)	4	*	116	0.0 (1)	112	0.0 (1)
4115	3,168	15.0 (19)	3,296	6.4 (24)	373	42.9 (36)	2,606	4.2 (15)	3,289	1.1 (25)
4351	501	34.9 (97)	511	20.2 (97)	153	34.6 (59)	453	14.1 (98)	511	2.2 (70)
4352	140	61.4 (150)	146	34.2 (151)	79	*	130	22.3 (141)	146	4.1 (120)
4353	371	28.8 (75)	376	15.7 (80)	81	*	323	7.7 (42)	376	1.6 (44)
4354	661	35.9 (101)	698	23.6 (118)	203	28.6 (82)	586	15.0 (107)	698	3.2 (97)
4355	290	37.2 (110)	310	25.5 (127)	98	*	251	14.3 (99)	310	3.9 (113)
4356	375	36.5 (104)	396	24.5 (119)	117	25.6 (87)	336	11.6 (80)	396	3.5 (104)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 118

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4358	102	14.7 (17)	102	9.8 (46)	13	*	93	*	103	1.9 (59)
4359	461	24.5 (62)	477	10.5 (52)	96	*	387	7.8 (45)	478	0.8 (11)
4360	202	29.7 (77)	199	17.1 (83)	48	*	179	14.5 (101)	198	3.5 (104)
4361	234	38.5 (117)	237	26.2 (131)	78	*	205	14.6 (105)	236	0.4 (4)
4362	266	48.5 (148)	284	28.2 (137)	112	38.4 (44)	236	21.6 (140)	284	4.2 (122)
4363	733	33.4 (90)	766	24.9 (123)	200	19.0 (92)	650	14.0 (96)	765	4.4 (125)
4401	888	31.3 (83)	921	14.7 (72)	201	43.3 (35)	738	4.9 (19)	920	1.1 (25)
4501	824	36.5 (104)	889	22.0 (111)	250	34.4 (61)	699	12.6 (88)	890	1.8 (57)
4601	1,170	34.8 (96)	1,229	15.6 (79)	308	46.1 (27)	988	9.9 (68)	1,230	3.7 (108)
4602	8	*	10	*	5	*	8	*	10	*
4701	945	31.6 (85)	979	19.5 (94)	223	36.8 (52)	706	6.2 (29)	981	2.1 (63)
5050	1,418	24.6 (63)	1,504	11.2 (57)	254	44.5 (31)	1,104	10.3 (73)	1,507	1.6 (44)
5051	2,383	20.1 (39)	2,510	8.8 (40)	357	47.3 (23)	1,796	6.6 (32)	2,506	1.9 (59)
5052	2,552	19.6 (37)	2,750	10.1 (50)	376	44.1 (32)	2,061	7.5 (40)	2,751	1.5 (38)
5055	1,124	2.9 (2)	1,144	1.1 (2)	24	*	1,007	4.6 (18)	1,144	1.0 (24)
5101	1,490	20.7 (45)	1,567	9.6 (44)	247	44.9 (30)	1,204	6.4 (30)	1,570	1.5 (38)
5102	23	*	25	*	4	*	14	*	25	*
5202	722	38.2 (114)	712	25.4 (126)	234	32.9 (68)	613	9.3 (63)	710	2.1 (63)
5206	2	*	5	*	0	*	0	*	5	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 119

### Michigan

#### Comparison of Smoking Indicators by Clinic

#### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
5601	869	29.9 (78)	905	19.2 (92)	210	31.9 (71)	725	7.6 (41)	901	2.3 (73)
5801	1,602	37.8 (113)	1,622	21.3 (105)	472	37.5 (47)	1,357	12.9 (89)	1,622	3.3 (101)
5803	112	30.4 (79)	116	15.5 (78)	22	*	82	*	115	2.6 (82)
5804	15	*	14	*	1	*	8	*	14	*
5806	45	*	56	*	13	*	28	*	56	*
6101	30	*	29	*	9	*	24	*	29	*
6102	101	32.7 (88)	114	22.8 (115)	31	*	91	*	113	6.2 (144)
6103	3,220	31.1 (82)	3,409	20.3 (98)	840	28.6 (82)	2,831	12.2 (85)	3,422	4.5 (126)
6105	17	*	17	*	0	*	12	*	17	*
6106	125	35.2 (100)	120	21.7 (108)	36	*	110	14.5 (101)	121	9.1 (152)
6301	3,334	14.6 (16)	3,431	7.8 (34)	386	39.1 (42)	2,783	5.4 (23)	3,431	1.1 (25)
6312	1,708	5.6 (5)	1,761	2.0 (6)	63	*	1,356	2.4 (7)	1,759	0.5 (6)
6316	1,123	16.7 (24)	1,104	7.0 (29)	139	40.3 (38)	851	4.0 (14)	1,104	0.7 (9)
6318	28	*	33	*	5	*	20	*	33	*
6320	187	8.0 (8)	187	1.6 (4)	9	*	118	0.0 (1)	186	0.0 (1)
6325	236	25.4 (68)	230	18.7 (90)	36	*	157	12.1 (84)	229	1.3 (31)
6330	4	*	4	*	2	*	4	*	4	*
6335	994	6.5 (7)	1,000	3.6 (8)	47	*	804	1.5 (4)	1,000	0.8 (11)
6340	34	*	30	*	4	*	31	*	30	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 120

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
6345	75	*	70	*	12	*	48	*	70	*
6360	748	16.6 (23)	783	8.4 (37)	100	34.0 (64)	616	10.2 (72)	784	1.5 (38)
7304	34	*	36	*	16	*	33	*	36	*
7306	2,045	35.9 (101)	2,211	13.3 (64)	597	53.8 (15)	1,757	11.6 (80)	2,212	2.2 (70)
7308	44	*	37	*	8	*	43	*	37	*
7310	12	*	9	*	2	*	11	*	9	*
7401	1,983	39.4 (122)	2,037	22.7 (114)	638	33.7 (65)	1,768	17.6 (124)	2,032	5.5 (139)
7601	569	26.5 (72)	552	16.3 (82)	120	20.0 (91)	512	9.0 (62)	552	1.3 (31)
7850	1,019	38.9 (121)	1,022	22.5 (113)	329	32.8 (70)	873	15.7 (116)	1,019	4.7 (128)
7852	18	*	19	*	8	*	17	*	19	*
8101	1,416	16.9 (25)	1,668	6.8 (27)	182	51.1 (19)	1,146	11.3 (78)	1,668	4.8 (130)
8102	340	17.1 (26)	379	7.7 (33)	49	*	300	5.3 (20)	378	2.9 (91)
8105	1,170	15.0 (19)	1,154	6.0 (20)	119	56.3 (12)	952	7.1 (35)	1,153	1.6 (44)
8106	415	20.2 (40)	480	10.6 (53)	68	*	336	7.7 (42)	479	1.5 (38)
8150	21	*	20	*	1	*	20	*	20	*
8151	2,726	21.8 (51)	2,806	9.9 (47)	457	46.6 (25)	2,284	6.7 (33)	2,809	1.4 (35)
8152	50	*	46	*	12	*	49	*	47	*
8153	7	*	7	*	1	*	6	*	7	*
8155	13	*	12	*	2	*	12	*	12	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

## 2018 - 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic Combined 3 Years

Page : 121

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8157	6	*	5	*	1	*	6	*	5	*
8201	2,267	17.6 (27)	2,411	9.9 (47)	286	37.8 (46)	1,641	7.9 (46)	2,418	1.6 (44)
8202	3,654	5.7 (6)	3,868	1.6 (4)	176	61.9 (4)	3,107	5.8 (27)	3,868	1.8 (57)
8203	357	27.2 (73)	366	18.0 (89)	62	*	270	10.0 (70)	367	1.6 (44)
8250	667	22.6 (54)	701	8.3 (36)	103	53.4 (16)	546	10.3 (73)	703	1.7 (55)
8252	134	20.9 (47)	149	4.7 (13)	24	*	113	15.0 (107)	149	4.0 (117)
8255	2,165	25.4 (68)	2,284	13.6 (66)	389	39.8 (40)	1,696	8.6 (56)	2,285	1.6 (44)
8257	1,944	1.6 (1)	2,004	1.2 (3)	23	*	1,614	1.5 (4)	2,003	0.8 (11)
8263	622	23.5 (58)	632	9.7 (45)	104	56.7 (11)	475	5.7 (25)	632	0.8 (11)
8264	656	19.1 (31)	648	6.8 (27)	93	*	488	7.0 (34)	648	0.5 (6)
8267	693	20.5 (43)	701	10.6 (53)	95	*	558	5.9 (28)	702	1.3 (31)
8901	854	22.1 (53)	972	8.5 (38)	168	54.2 (14)	753	16.9 (121)	970	5.2 (135)
8902	1,237	16.4 (22)	1,370	4.9 (14)	167	62.3 (3)	1,048	12.9 (89)	1,368	5.2 (135)
8903	1,279	19.6 (37)	1,428	7.6 (32)	196	51.0 (20)	1,072	19.5 (135)	1,427	5.9 (142)
8904	2,019	19.4 (34)	2,267	7.0 (29)	298	64.4 (2)	1,631	15.5 (114)	2,267	6.0 (143)
8905	1,358	13.9 (13)	1,441	5.6 (18)	143	55.2 (13)	1,097	7.1 (35)	1,439	2.6 (82)
8909	612	19.3 (33)	680	5.1 (16)	96	*	482	16.0 (118)	678	5.0 (133)
9045	1,592	11.4 (10)	1,655	4.3 (12)	147	49.7 (21)	1,410	8.3 (53)	1,653	2.4 (78)
9050	780	22.6 (54)	918	5.8 (19)	166	60.8 (5)	679	16.5 (120)	917	3.2 (97)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 122

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9051	1,463	11.3 (9)	1,503	3.9 (11)	142	51.4 (18)	1,337	5.6 (24)	1,503	1.5 (38)
9053	778	15.7 (21)	880	4.9 (14)	106	58.5 (8)	564	15.4 (112)	881	3.7 (108)
9055	10	*	19	*	1	*	6	*	19	*
9058	859	13.5 (11)	966	6.0 (20)	89	*	699	8.0 (49)	967	2.8 (88)
9066	856	20.8 (46)	1,086	6.6 (25)	132	65.2 (1)	667	18.7 (130)	1,085	4.1 (120)
9068	808	21.0 (48)	915	8.9 (41)	143	53.1 (17)	711	14.5 (101)	915	3.9 (113)
9069	2,103	14.9 (18)	2,511	6.3 (22)	236	57.2 (9)	1,603	18.7 (130)	2,509	7.1 (148)
9070	1,186	19.2 (32)	1,341	7.1 (31)	176	60.2 (7)	923	19.6 (136)	1,334	5.2 (135)
9072	1,258	20.3 (41)	1,378	5.4 (17)	223	60.5 (6)	1,132	18.7 (130)	1,377	4.9 (131)
9080	2,238	19.5 (35)	2,524	6.3 (22)	362	56.9 (10)	1,874	11.8 (83)	2,525	3.9 (113)
9082	139	14.4 (15)	136	3.7 (10)	15	*	103	17.5 (123)	136	8.1 (150)
9083	77	*	101	2.0 (6)	0	*	9	*	101	0.0 (1)
9084	29	*	33	*	2	*	13	*	33	*
9085	49	*	44	*	10	*	49	*	44	*
9086	48	*	40	*	1	*	48	*	40	*
9087	5	*	4	*	0	*	4	*	4	*
Michigan	139,618	24.5	146,978	13.0	27,387	38.9	116,894	9.8	146,919	2.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2018 - 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 123

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	102	6.9 (22)	3.9 (17)	104	4.8 (1)	97	*	102	67.6 (94)
0102	417	8.4 (57)	7.4 (68)	425	9.2 (19)	375	5.1 (109)	418	64.6 (110)
0103	348	7.2 (28)	10.9 (137)	353	6.8 (4)	320	3.4 (49)	348	67.5 (95)
0104	118	5.1 (3)	8.5 (105)	125	8.8 (13)	109	3.7 (60)	118	73.7 (74)
0105	17	*	*	17	*	15	*	17	*
0250	124	5.6 (9)	11.3 (143)	122	11.5 (70)	108	0.9 (1)	124	76.6 (57)
0251	79	*	*	80	*	71	*	79	*
0252	92	*	*	94	*	88	*	92	*
0253	122	11.5 (114)	5.7 (37)	119	12.6 (100)	103	6.8 (127)	122	73.0 (79)
0401	394	7.1 (25)	10.2 (131)	399	8.5 (12)	358	2.8 (25)	393	73.8 (72)
0402	331	7.3 (30)	8.5 (105)	336	9.8 (27)	292	3.1 (36)	331	78.9 (41)
0403	135	11.9 (119)	8.9 (117)	134	12.7 (102)	116	6.0 (119)	135	74.8 (67)
0404	105	6.7 (19)	12.4 (149)	104	12.5 (99)	91	*	105	74.3 (69)
0501	392	10.2 (94)	7.9 (84)	457	9.4 (22)	350	4.9 (102)	445	75.7 (61)
0503	98	*	*	101	8.9 (16)	88	*	101	85.1 (12)
0504	406	8.1 (50)	7.6 (75)	420	10.7 (44)	360	3.1 (36)	414	86.2 (7)
0505	244	10.2 (94)	5.3 (31)	264	10.2 (35)	218	4.1 (76)	259	85.7 (9)
0506	144	6.9 (22)	8.3 (95)	148	10.1 (32)	126	4.0 (72)	149	76.5 (58)
0507	62	*	*	71	*	54	*	70	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 124

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	27	*	*	27	*	25	*	27	*
0509	3	*	*	3	*	3	*	3	*
0650	943	8.6 (65)	9.1 (121)	977	10.7 (44)	845	4.6 (93)	945	75.0 (66)
0651	525	5.3 (4)	7.4 (68)	535	8.8 (13)	475	2.7 (20)	524	74.0 (70)
0652	293	10.6 (99)	8.2 (93)	303	11.2 (60)	259	4.6 (93)	294	72.4 (84)
0653	362	7.2 (28)	9.4 (125)	378	11.4 (66)	320	2.2 (10)	363	83.7 (15)
0654	327	9.2 (77)	8.0 (86)	334	8.4 (11)	298	6.4 (123)	330	75.5 (63)
0655	298	5.4 (6)	7.0 (64)	296	9.1 (17)	266	2.3 (12)	298	73.5 (76)
0657	57	*	*	58	*	47	*	57	*
0701	352	6.5 (17)	11.9 (148)	367	7.1 (5)	320	4.4 (86)	355	87.0 (4)
0702	67	*	*	67	*	60	*	67	*
0703	162	8.0 (48)	11.1 (140)	161	10.6 (42)	142	2.8 (25)	162	74.7 (68)
0704	49	*	*	50	*	46	*	49	*
0705	96	*	*	99	*	86	*	95	*
0706	11	*	*	11	*	9	*	11	*
0850	685	7.7 (44)	8.9 (117)	721	10.4 (40)	617	3.2 (39)	704	83.5 (16)
0851	618	8.4 (57)	9.2 (123)	641	9.8 (27)	557	4.5 (90)	624	80.3 (31)
0901	1,374	8.7 (71)	8.4 (101)	1,417	10.9 (51)	1,227	3.6 (56)	1,380	66.7 (100)
0902	36	*	*	37	*	36	*	36	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 125

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1051	1,070	7.6 (40)	11.0 (138)	1,097	9.1 (17)	965	2.5 (17)	1,072	83.2 (19)
1101	1,566	8.6 (65)	5.7 (37)	1,608	10.3 (38)	1,389	4.9 (102)	1,569	68.8 (91)
1102	679	6.3 (15)	6.2 (47)	701	6.7 (3)	624	3.4 (49)	679	72.9 (81)
1104	1	*	*	1	*	1	*	1	*
1106	66	*	*	67	*	60	*	66	*
1250	924	7.9 (47)	7.4 (68)	950	10.6 (42)	815	2.7 (20)	934	75.6 (62)
1251	637	6.8 (20)	8.5 (105)	662	9.4 (22)	573	3.0 (32)	638	79.5 (35)
1252	486	7.6 (40)	7.0 (64)	494	10.9 (51)	428	3.0 (32)	489	84.0 (14)
1253	669	6.6 (18)	7.6 (75)	692	9.2 (19)	604	2.2 (10)	675	84.4 (13)
1350	2,035	9.1 (74)	7.5 (74)	2,063	10.1 (32)	1,810	4.2 (80)	2,041	72.8 (82)
1351	104	3.8 (1)	8.7 (115)	104	9.6 (25)	93	*	104	86.5 (6)
1352	344	11.3 (110)	7.8 (80)	352	11.4 (66)	303	5.3 (114)	350	64.6 (110)
1401	204	7.4 (33)	7.4 (68)	212	12.7 (102)	179	1.1 (2)	204	83.3 (18)
1402	105	9.5 (83)	7.6 (75)	107	12.1 (90)	91	*	105	92.4 (1)
1403	23	*	*	23	*	21	*	23	*
1501	667	7.5 (39)	10.5 (133)	695	10.2 (35)	589	3.2 (39)	687	66.4 (101)
1503	837	6.8 (20)	7.4 (68)	854	11.8 (81)	729	3.2 (39)	838	75.4 (65)
1504	1,649	7.3 (30)	7.3 (67)	1,845	10.9 (51)	1,458	2.8 (25)	1,818	78.7 (42)
1505	342	9.1 (74)	11.4 (144)	358	10.9 (51)	306	4.9 (102)	348	82.8 (23)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 126

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1506	499	7.4 (33)	8.0 (86)	507	11.2 (60)	439	3.2 (39)	500	73.0 (79)
1507	485	9.5 (83)	11.8 (147)	507	13.0 (106)	422	3.3 (45)	487	82.5 (25)
1508	40	*	*	39	*	34	*	40	*
1509	330	6.1 (14)	10.0 (130)	337	11.9 (85)	287	2.1 (8)	336	70.5 (88)
1511	601	9.5 (83)	7.7 (79)	615	12.2 (93)	521	3.8 (65)	607	80.1 (32)
1512	314	8.0 (48)	8.0 (86)	323	15.5 (136)	266	1.9 (6)	315	78.4 (45)
1513	14	*	*	16	*	14	*	15	*
1514	267	13.5 (133)	8.6 (111)	274	15.3 (132)	225	3.6 (56)	273	87.9 (2)
1701	546	7.3 (30)	11.7 (146)	549	11.5 (70)	468	2.4 (15)	547	67.5 (95)
1950	537	8.2 (53)	6.9 (61)	550	11.6 (76)	471	3.0 (32)	539	82.0 (26)
1952	120	5.0 (2)	8.3 (95)	122	8.2 (9)	110	1.8 (5)	120	86.7 (5)
1953	270	7.4 (33)	6.7 (56)	277	11.6 (76)	238	1.7 (4)	270	85.9 (8)
1954	461	7.4 (33)	8.7 (115)	468	11.5 (70)	402	2.7 (20)	462	81.4 (29)
1955	89	*	*	91	*	79	*	89	*
1957	387	9.8 (89)	11.1 (140)	389	12.1 (90)	336	3.3 (45)	388	83.2 (19)
1963	45	*	*	47	*	40	*	45	*
1964	30	*	*	30	*	29	*	30	*
2150	547	7.1 (25)	9.1 (121)	565	9.9 (29)	491	3.1 (36)	548	81.8 (27)
2151	179	7.8 (46)	8.9 (117)	181	9.9 (29)	160	3.8 (65)	179	57.0 (127)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 127

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2152	4	*	*	3	*	3	*	4	*
2251	357	8.1 (50)	8.4 (101)	362	11.0 (57)	315	3.5 (52)	358	73.7 (74)
2252	140	9.3 (79)	10.7 (135)	145	9.7 (26)	125	4.0 (72)	140	78.6 (43)
2554	45	*	*	47	*	42	*	45	*
2555	6,739	11.7 (117)	5.4 (33)	6,830	13.3 (115)	5,793	5.6 (116)	6,746	56.6 (130)
2556	317	16.7 (148)	3.5 (9)	323	20.4 (150)	246	8.9 (140)	320	52.2 (140)
2557	7	*	*	8	*	6	*	8	*
2558	4	*	*	5	*	3	*	5	*
2559	3	*	*	3	*	2	*	3	*
2560	4	*	*	4	*	4	*	4	*
2561	3	*	*	3	*	2	*	3	*
2562	14	*	*	14	*	11	*	14	*
2563	25	*	*	23	*	21	*	25	*
2564	18	*	*	18	*	18	*	18	*
3001	1,015	13.0 (127)	4.0 (18)	1,034	14.0 (119)	872	4.8 (98)	1,015	59.0 (121)
3002	272	11.4 (111)	4.4 (22)	275	12.0 (87)	240	5.0 (105)	272	52.9 (137)
3004	181	8.8 (73)	8.3 (95)	185	13.0 (106)	158	4.4 (86)	183	73.8 (72)
3005	58	*	*	58	*	51	*	58	*
3007	12	*	*	11	*	9	*	12	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 128

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3009	1	*	*	1	*	1	*	1	*
3201	383	9.7 (88)	11.2 (142)	398	11.3 (63)	343	4.1 (76)	383	78.6 (43)
3203	0	*	*	0	*	0	*	0	*
3301	3,762	10.6 (99)	6.1 (46)	3,896	12.9 (105)	3,276	4.6 (93)	3,788	76.4 (59)
3304	89	*	*	91	*	79	*	91	*
3305	140	8.6 (65)	6.4 (53)	149	9.4 (22)	128	3.9 (71)	143	87.4 (3)
3306	232	5.6 (9)	10.8 (136)	241	8.3 (10)	212	1.9 (6)	232	79.7 (33)
3308	80	*	*	80	*	71	*	79	*
3309	18	*	*	17	*	15	*	18	*
3312	353	11.9 (119)	8.5 (105)	363	16.0 (139)	297	5.7 (117)	354	67.8 (93)
3550	52	*	*	52	*	45	*	51	*
3551	15	*	*	18	*	15	*	15	*
3552	38	*	*	40	*	32	*	38	*
3701	905	8.4 (57)	8.0 (86)	918	10.9 (51)	805	2.9 (30)	905	83.4 (17)
3850	2,688	9.6 (87)	8.1 (92)	2,748	11.5 (70)	2,377	4.2 (80)	2,691	71.9 (87)
3901	1,279	11.2 (108)	5.8 (40)	1,327	13.3 (115)	1,104	4.7 (97)	1,300	65.8 (102)
3902	263	9.1 (74)	11.0 (138)	267	12.4 (96)	230	5.2 (110)	264	65.5 (104)
3903	28	*	*	30	*	26	*	28	*
3904	6	*	*	6	*	5	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 129

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3905	11	*	*	11	*	9	*	11	*
3950	1,822	10.1 (93)	8.6 (111)	1,893	11.7 (79)	1,597	2.8 (25)	1,844	72.7 (83)
3954	14	*	*	15	*	14	*	14	*
4001	76	*	*	77	*	68	*	76	*
4002	46	*	*	48	*	40	*	46	*
4103	12	*	*	11	*	8	*	12	*
4106	2,156	13.0 (127)	4.7 (24)	2,233	14.3 (123)	1,842	5.8 (118)	2,178	77.3 (56)
4107	2,648	12.7 (126)	6.9 (61)	2,747	15.9 (138)	2,219	5.0 (105)	2,665	77.6 (53)
4109	833	8.6 (65)	7.8 (80)	850	13.1 (110)	723	2.9 (30)	834	78.1 (47)
4110	7	*	*	7	*	0	*	7	*
4112	478	5.9 (12)	6.7 (56)	477	10.7 (44)	420	2.1 (8)	478	22.6 (150)
4113	105	9.5 (83)	9.5 (126)	111	16.2 (141)	88	*	106	83.0 (22)
4115	3,128	12.1 (121)	6.3 (51)	3,219	14.0 (119)	2,676	4.5 (90)	3,145	78.3 (46)
4351	484	7.6 (40)	6.2 (47)	498	12.0 (87)	426	3.3 (45)	485	79.4 (36)
4352	135	11.1 (106)	8.9 (117)	139	15.8 (137)	113	5.3 (114)	135	73.3 (78)
4353	360	8.3 (54)	6.7 (56)	360	10.8 (48)	316	3.8 (65)	361	77.8 (52)
4354	663	7.4 (33)	7.8 (80)	671	11.8 (81)	579	2.6 (18)	665	74.0 (70)
4355	285	9.8 (89)	9.8 (129)	292	12.3 (95)	247	4.0 (72)	287	77.4 (55)
4356	368	10.9 (103)	6.8 (60)	380	14.7 (128)	310	4.2 (80)	368	75.5 (63)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 130

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4358	97	*	*	97	*	90	*	97	*
4359	448	5.4 (6)	8.5 (105)	454	10.8 (48)	398	2.3 (12)	449	78.0 (48)
4360	166	5.4 (6)	11.4 (144)	183	7.1 (5)	151	2.6 (18)	166	80.7 (30)
4361	200	8.5 (62)	10.5 (133)	233	12.0 (87)	176	4.0 (72)	201	83.1 (21)
4362	267	9.4 (82)	7.9 (84)	268	10.1 (32)	234	4.3 (83)	268	85.4 (11)
4363	717	7.4 (33)	8.6 (111)	723	11.3 (63)	620	2.7 (20)	718	79.4 (36)
4401	882	8.4 (57)	8.6 (111)	887	11.3 (63)	768	3.4 (49)	882	79.7 (33)
4501	839	8.5 (62)	9.2 (123)	855	11.2 (60)	736	3.3 (45)	838	76.1 (60)
4601	1,169	5.7 (11)	8.2 (93)	1,206	8.8 (13)	1,057	2.7 (20)	1,170	69.0 (90)
4602	9	*	*	10	*	9	*	9	*
4701	951	10.0 (92)	8.3 (95)	960	12.4 (96)	831	3.5 (52)	950	81.8 (27)
5050	1,466	11.7 (117)	5.8 (40)	1,464	15.4 (135)	1,222	4.4 (86)	1,468	61.0 (115)
5051	2,430	10.4 (98)	5.5 (34)	2,446	12.1 (90)	2,100	4.1 (76)	2,436	65.0 (107)
5052	2,666	11.5 (114)	5.0 (26)	2,691	13.7 (118)	2,267	5.2 (110)	2,676	60.9 (117)
5055	1,008	7.1 (25)	4.2 (20)	1,060	7.4 (7)	932	3.6 (56)	1,045	44.8 (146)
5101	1,512	8.6 (65)	6.7 (56)	1,532	11.4 (66)	1,323	3.6 (56)	1,519	70.0 (89)
5102	25	*	*	24	*	22	*	24	*
5202	686	8.6 (65)	8.5 (105)	696	10.9 (51)	609	4.1 (76)	687	85.7 (9)
5206	5	*	*	4	*	4	*	5	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 131

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5601	867	6.9 (22)	9.7 (128)	891	9.9 (29)	776	2.3 (12)	867	79.0 (39)
5801	1,551	8.3 (54)	6.6 (54)	1,583	11.5 (70)	1,361	3.0 (32)	1,552	65.8 (102)
5803	108	9.3 (79)	10.2 (131)	109	11.9 (85)	93	*	108	62.0 (114)
5804	12	*	*	13	*	10	*	12	*
5806	53	*	*	55	*	47	*	53	*
6101	29	*	*	27	*	22	*	29	*
6102	113	5.3 (4)	6.2 (47)	113	6.2 (2)	106	2.8 (25)	113	77.9 (51)
6103	3,252	10.9 (103)	5.2 (28)	3,285	14.3 (123)	2,724	4.8 (98)	3,258	65.5 (104)
6105	17	*	*	17	*	16	*	17	*
6106	120	13.3 (131)	4.2 (20)	121	18.2 (147)	98	*	120	65.0 (107)
6301	3,274	12.2 (122)	6.3 (51)	3,367	13.1 (110)	2,826	5.2 (110)	3,304	67.1 (98)
6312	1,672	11.1 (106)	6.0 (44)	1,724	11.5 (70)	1,467	3.8 (65)	1,689	77.6 (53)
6316	1,050	10.2 (94)	7.2 (66)	1,081	12.6 (100)	913	3.2 (39)	1,059	78.0 (48)
6318	28	*	*	32	*	26	*	30	*
6320	170	11.2 (108)	5.3 (31)	178	14.0 (119)	146	4.8 (98)	173	82.7 (24)
6325	224	6.3 (15)	8.0 (86)	227	7.5 (8)	207	2.4 (15)	223	78.0 (48)
6330	4	*	*	4	*	4	*	4	*
6335	950	9.2 (77)	5.2 (28)	955	10.7 (44)	840	4.8 (98)	954	72.1 (86)
6340	29	*	*	30	*	24	*	29	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 132

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6345	68	*	*	71	*	61	*	68	*
6360	751	12.3 (123)	5.9 (43)	764	14.5 (127)	635	6.1 (121)	758	67.0 (99)
7304	36	*	*	36	*	35	*	36	*
7306	2,127	11.0 (105)	5.7 (37)	2,165	12.2 (93)	1,858	4.3 (83)	2,136	62.7 (112)
7308	37	*	*	37	*	33	*	37	*
7310	9	*	*	9	*	9	*	9	*
7401	1,949	7.7 (44)	8.3 (95)	1,988	10.8 (48)	1,732	3.5 (52)	1,952	68.4 (92)
7601	536	6.0 (13)	7.8 (80)	549	11.7 (79)	472	1.1 (2)	538	53.7 (133)
7850	971	8.1 (50)	7.6 (75)	988	10.5 (41)	867	3.2 (39)	972	73.4 (77)
7852	19	*	*	19	*	17	*	19	*
8101	1,534	16.9 (149)	3.0 (3)	1,606	19.6 (149)	1,198	7.6 (135)	1,580	44.7 (147)
8102	356	8.4 (57)	9.6 (127)	365	14.8 (129)	293	3.8 (65)	362	50.6 (143)
8105	1,088	10.6 (99)	4.9 (25)	1,137	11.8 (81)	954	3.7 (60)	1,121	59.8 (120)
8106	447	15.0 (139)	5.8 (40)	468	16.2 (141)	370	6.5 (126)	455	60.4 (118)
8150	17	*	*	16	*	16	*	17	*
8151	2,680	11.6 (116)	5.0 (26)	2,732	14.2 (122)	2,291	4.5 (90)	2,665	79.0 (39)
8152	46	*	*	47	*	44	*	46	*
8153	7	*	*	6	*	4	*	7	*
8155	12	*	*	12	*	10	*	12	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

### 2018 - 2020 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 133

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8157	5	*	*	5	*	4	*	5	*
8201	2,308	11.4 (111)	6.2 (47)	2,370	12.8 (104)	1,998	5.0 (105)	2,336	60.1 (119)
8202	3,583	8.3 (54)	5.6 (35)	3,725	9.2 (19)	3,234	4.3 (83)	3,692	45.5 (145)
8203	350	11.4 (111)	8.3 (95)	357	13.2 (113)	301	3.7 (60)	354	79.4 (36)
8250	678	13.1 (129)	5.2 (28)	693	11.4 (66)	594	7.2 (130)	684	62.3 (113)
8252	144	13.9 (135)	5.6 (35)	146	11.6 (76)	124	7.3 (132)	146	61.0 (115)
8255	2,195	10.6 (99)	6.9 (61)	2,247	11.0 (57)	1,936	5.0 (105)	2,216	57.6 (125)
8257	1,775	12.6 (125)	3.1 (4)	1,936	11.8 (81)	1,539	7.3 (132)	1,860	72.2 (85)
8263	606	8.7 (71)	8.4 (101)	622	10.3 (38)	543	3.7 (60)	609	56.7 (129)
8264	608	7.6 (40)	7.4 (68)	626	10.2 (35)	541	3.7 (60)	612	57.7 (124)
8267	678	9.3 (79)	8.4 (101)	695	11.1 (59)	606	3.5 (52)	683	54.3 (131)
8901	917	15.4 (143)	2.3 (1)	949	17.3 (145)	747	8.0 (138)	929	49.6 (144)
8902	1,194	14.0 (136)	2.5 (2)	1,341	16.9 (144)	998	5.2 (110)	1,286	58.4 (122)
8903	1,307	16.4 (146)	3.1 (4)	1,377	18.4 (148)	1,044	8.7 (139)	1,332	51.1 (142)
8904	2,071	15.0 (139)	3.7 (12)	2,194	18.0 (146)	1,666	6.9 (129)	2,150	52.8 (138)
8905	1,270	10.2 (94)	6.6 (54)	1,384	13.2 (113)	1,082	4.4 (86)	1,372	64.7 (109)
8909	615	15.0 (139)	3.4 (8)	655	15.0 (130)	511	6.8 (127)	639	57.0 (127)
9045	1,551	9.8 (89)	6.0 (44)	1,606	13.6 (117)	1,327	3.8 (65)	1,564	53.1 (135)
9050	839	13.8 (134)	4.1 (19)	882	15.2 (131)	703	7.5 (134)	864	51.3 (141)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/21/2021

## Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Page : 134

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9051	1,395	8.5 (62)	8.0 (86)	1,404	13.1 (110)	1,183	4.6 (93)	1,412	67.5 (95)
9053	838	13.4 (132)	3.7 (12)	863	13.0 (106)	722	7.2 (130)	842	58.1 (123)
9055	18	*	*	18	*	17	*	18	*
9058	902	15.5 (145)	3.2 (6)	948	12.4 (96)	787	9.3 (141)	933	57.6 (125)
9066	1,012	16.4 (146)	3.8 (14)	1,073	16.7 (143)	842	7.7 (136)	1,023	53.0 (136)
9068	880	14.8 (137)	4.4 (22)	893	15.3 (132)	740	7.8 (137)	880	44.7 (147)
9069	2,362	15.0 (139)	3.2 (6)	2,454	15.3 (132)	1,986	6.4 (123)	2,366	35.3 (149)
9070	1,289	12.5 (124)	3.6 (10)	1,320	13.0 (106)	1,107	6.0 (119)	1,291	53.8 (132)
9072	1,295	14.8 (137)	3.6 (10)	1,352	16.0 (139)	1,088	6.3 (122)	1,302	53.4 (134)
9080	2,387	13.1 (129)	3.8 (14)	2,464	14.4 (126)	2,019	6.4 (123)	2,441	52.4 (139)
9082	130	15.4 (143)	3.8 (14)	133	14.3 (123)	108	10.2 (142)	129	65.1 (106)
9083	96	*	*	101	21.8 (151)	75	*	96	*
9084	28	*	*	33	*	26	*	28	*
9085	44	*	*	43	*	36	*	44	*
9086	40	*	*	40	*	36	*	40	*
9087	4	*	*	4	*	4	*	4	*
<b>Michigan</b>	<b>138,554</b>	<b>10.4</b>	<b>6.4</b>	<b>142,789</b>	<b>12.5</b>	<b>120,167</b>	<b>4.6</b>	<b>140,109</b>	<b>57.4</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.