

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2021 Pregnancy Nutrition Surveillance (1)

Page : 1

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	35	*	*	*	*	*	*	*
0102	135	91.9	0.7	2.2	0.7	0.0	4.4	0.0
0103	124	90.3	0.8	4.8	0.0	0.0	4.0	0.0
0104	42	*	*	*	*	*	*	*
0105	3	*	*	*	*	*	*	*
0250	39	*	*	*	*	*	*	*
0251	31	*	*	*	*	*	*	*
0252	28	*	*	*	*	*	*	*
0253	36	*	*	*	*	*	*	*
0401	142	93.0	0.7	2.1	0.7	0.0	3.5	0.0
0402	125	87.2	0.0	2.4	0.0	0.0	10.4	0.0
0403	43	*	*	*	*	*	*	*
0404	38	*	*	*	*	*	*	*
0501	137	92.7	0.0	2.2	0.0	0.0	5.1	0.0
0503	22	*	*	*	*	*	*	*
0504	119	83.2	0.8	3.4	0.8	1.7	10.1	0.0
0505	69	*	*	*	*	*	*	*
0506	60	*	*	*	*	*	*	*
0507	34	*	*	*	*	*	*	*
0508	5	*	*	*	*	*	*	*
0650	355	76.1	4.2	8.2	2.3	0.3	9.0	0.0
0651	183	92.9	0.5	3.3	0.0	0.5	2.7	0.0
0652	115	93.9	0.0	2.6	0.0	0.0	3.5	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2021 Pregnancy Nutrition Surveillance (1)

Page : 2

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0653	131	93.9	0.0	2.3	0.0	1.5	2.3	0.0
0654	138	94.2	0.0	3.6	0.0	0.0	2.2	0.0
0655	104	93.3	0.0	1.9	0.0	0.0	3.8	1.0
0657	19	*	*	*	*	*	*	*
0701	184	89.1	0.5	2.2	1.6	2.7	3.8	0.0
0702	17	*	*	*	*	*	*	*
0703	50	*	*	*	*	*	*	*
0704	14	*	*	*	*	*	*	*
0705	14	*	*	*	*	*	*	*
0850	217	88.0	3.2	7.8	0.0	0.5	0.5	0.0
0851	186	91.4	0.5	5.4	0.5	0.0	1.6	0.5
0901	501	79.8	2.2	11.6	0.0	0.2	6.2	0.0
0902	11	*	*	*	*	*	*	*
1051	398	84.4	1.3	6.0	3.5	0.8	3.5	0.5
1052	1	*	*	*	*	*	*	*
1101	488	33.0	52.5	8.0	0.2	2.0	3.5	0.8
1102	263	77.2	10.6	7.2	0.0	0.4	3.0	1.5
1106	16	*	*	*	*	*	*	*
1250	326	83.1	1.2	12.6	0.0	0.3	1.8	0.9
1251	206	93.2	0.5	2.4	0.0	0.0	1.9	1.9
1252	165	57.0	0.6	38.8	0.0	0.0	3.0	0.6
1253	215	78.1	0.9	9.8	0.0	0.5	10.7	0.0
1350	831	59.0	21.5	7.9	0.2	5.3	5.7	0.4

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2021 Pregnancy Nutrition Surveillance (1)

Page : 3

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1351	1	*	*	*	*	*	*	*
1352	127	69.3	17.3	6.3	0.0	0.0	4.7	2.4
1401	73	*	*	*	*	*	*	*
1402	37	*	*	*	*	*	*	*
1403	5	*	*	*	*	*	*	*
1501	242	53.3	5.0	36.0	0.4	0.8	2.5	2.1
1503	267	58.1	7.1	21.7	0.0	0.4	9.4	3.4
1504	626	39.9	5.6	43.5	0.2	4.5	5.8	0.6
1505	22	*	*	*	*	*	*	*
1506	185	75.7	2.7	17.3	0.0	1.1	2.7	0.5
1507	192	84.9	3.6	7.3	0.0	1.6	2.6	0.0
1509	143	47.6	0.7	46.2	0.0	1.4	1.4	2.8
1511	207	85.5	2.9	7.2	0.5	0.5	2.4	1.0
1512	110	89.1	0.0	1.8	0.0	1.8	7.3	0.0
1514	210	69.5	4.8	16.2	0.5	1.9	6.7	0.5
1701	180	57.2	0.6	3.9	17.2	1.1	20.0	0.0
1950	201	90.5	1.0	7.5	0.0	0.0	0.5	0.5
1952	44	*	*	*	*	*	*	*
1953	88	*	*	*	*	*	*	*
1954	118	83.1	0.8	11.9	0.0	0.0	2.5	1.7
1955	35	*	*	*	*	*	*	*
1957	132	84.1	0.8	9.1	0.0	0.0	6.1	0.0
1963	23	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2021 Pregnancy Nutrition Surveillance (1)

Page : 4

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2150	200	82.0	1.5	2.5	1.5	1.5	11.0	0.0
2151	77	*	*	*	*	*	*	*
2152	13	*	*	*	*	*	*	*
2251	141	95.0	0.0	1.4	0.0	0.0	2.1	1.4
2252	47	*	*	*	*	*	*	*
2554	19	*	*	*	*	*	*	*
2555	2,371	48.8	37.9	6.2	0.0	0.4	6.3	0.3
2556	61	*	*	*	*	*	*	*
2560	1	*	*	*	*	*	*	*
2562	2	*	*	*	*	*	*	*
2563	3	*	*	*	*	*	*	*
3001	391	26.1	52.2	13.6	0.3	0.3	7.4	0.3
3002	54	*	*	*	*	*	*	*
3004	95	*	*	*	*	*	*	*
3005	50	*	*	*	*	*	*	*
3007	4	*	*	*	*	*	*	*
3201	124	95.2	0.0	4.8	0.0	0.0	0.0	0.0
3301	1,467	40.0	31.2	15.3	0.1	4.9	8.3	0.1
3304	28	*	*	*	*	*	*	*
3305	41	*	*	*	*	*	*	*
3306	66	*	*	*	*	*	*	*
3308	31	*	*	*	*	*	*	*
3312	137	38.0	27.0	23.4	0.0	1.5	10.2	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2021 Pregnancy Nutrition Surveillance (1)

Page : 5

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3550	26	*	*	*	*	*	*	*
3551	3	*	*	*	*	*	*	*
3552	1	*	*	*	*	*	*	*
3701	292	87.0	0.3	8.9	0.0	0.0	3.8	0.0
3850	960	71.7	15.2	6.5	0.1	0.9	5.6	0.0
3901	563	39.4	48.3	5.7	0.0	0.7	5.3	0.5
3902	7	*	*	*	*	*	*	*
3903	2	*	*	*	*	*	*	*
3950	662	51.5	24.9	13.0	0.5	0.8	8.5	0.9
4106	705	12.8	38.3	44.0	0.3	0.9	3.8	0.0
4107	945	40.2	29.7	20.8	0.3	2.2	6.3	0.3
4109	207	87.4	1.4	8.2	0.0	0.0	2.9	0.0
4112	198	1.0	1.5	96.0	0.0	0.0	0.0	1.5
4115	1,130	28.2	32.0	23.0	0.3	11.4	4.9	0.2
4351	159	85.5	0.0	6.3	0.6	1.3	6.3	0.0
4352	57	*	*	*	*	*	*	*
4353	161	60.2	0.0	36.0	0.6	0.0	3.1	0.0
4354	257	90.3	1.6	3.9	0.4	0.0	3.5	0.4
4355	88	*	*	*	*	*	*	*
4356	119	73.9	2.5	15.1	0.8	0.8	5.0	1.7
4359	205	60.0	0.5	35.1	0.5	0.5	2.9	0.5
4360	71	*	*	*	*	*	*	*
4361	94	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2021 Pregnancy Nutrition Surveillance (1)

Page : 6

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4362	92	*	*	*	*	*	*	*
4363	214	89.7	1.4	2.3	0.0	0.5	6.1	0.0
4401	255	88.2	1.2	5.9	0.0	0.8	3.9	0.0
4501	251	93.6	0.0	4.4	0.0	0.0	2.0	0.0
4601	389	69.9	3.6	19.8	0.0	0.3	5.7	0.8
4701	340	89.1	1.5	4.1	0.0	0.9	3.2	1.2
5050	496	45.0	47.2	2.4	0.0	1.0	4.2	0.2
5051	883	50.1	38.8	3.5	0.1	2.3	4.5	0.7
5052	848	30.0	44.0	3.3	0.2	18.6	3.8	0.1
5055	438	76.7	11.0	1.6	0.2	9.8	0.7	0.0
5101	518	65.6	15.6	12.7	0.2	1.9	3.9	0.0
5102	3	*	*	*	*	*	*	*
5202	222	87.4	2.3	2.7	2.3	0.0	5.0	0.5
5206	4	*	*	*	*	*	*	*
5601	330	86.4	2.1	6.4	0.0	2.4	2.4	0.3
5801	596	80.7	7.0	6.7	0.0	0.2	5.0	0.3
5803	11	*	*	*	*	*	*	*
5804	8	*	*	*	*	*	*	*
5806	9	*	*	*	*	*	*	*
6101	19	*	*	*	*	*	*	*
6102	27	*	*	*	*	*	*	*
6103	1,097	58.1	23.9	9.8	0.3	0.4	7.1	0.5
6105	2	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2021 Pregnancy Nutrition Surveillance (1)

Page : 7

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6301	1,067	40.3	29.4	22.9	0.2	2.4	4.6	0.2
6312	590	35.9	54.9	4.4	0.2	1.7	2.7	0.2
6316	354	57.6	27.7	7.3	0.0	4.0	3.1	0.3
6320	186	38.2	51.6	2.7	0.0	2.7	4.8	0.0
6325	106	88.7	1.9	7.5	0.0	0.0	1.9	0.0
6335	298	68.1	21.1	3.4	0.0	4.7	2.7	0.0
6345	39	*	*	*	*	*	*	*
6360	385	34.3	44.7	15.3	0.0	1.6	3.6	0.5
7306	729	46.1	33.3	15.1	0.0	0.1	5.1	0.3
7401	590	81.4	5.1	7.5	0.0	0.5	5.4	0.2
7601	173	94.8	0.0	2.3	0.0	0.6	1.2	1.2
7850	331	87.3	0.6	5.7	0.0	0.3	5.7	0.3
7852	3	*	*	*	*	*	*	*
8101	579	7.1	89.1	1.2	0.0	0.3	1.9	0.3
8102	196	22.4	16.8	58.2	0.0	1.0	1.5	0.0
8105	1	*	*	*	*	*	*	*
8106	196	27.0	53.6	13.8	0.0	0.5	5.1	0.0
8150	5	*	*	*	*	*	*	*
8151	1,079	36.8	40.8	9.9	0.1	3.0	9.0	0.5
8152	2	*	*	*	*	*	*	*
8155	3	*	*	*	*	*	*	*
8201	922	42.5	43.2	4.7	0.2	3.8	5.1	0.5
8202	1,245	87.2	10.0	1.5	0.0	0.3	0.8	0.1

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2021 Pregnancy Nutrition Surveillance (1)

Page : 8

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8203	132	51.5	35.6	7.6	0.0	0.0	5.3	0.0
8250	159	20.8	70.4	1.9	0.6	1.9	4.4	0.0
8252	57	*	*	*	*	*	*	*
8255	688	55.4	28.2	10.8	0.4	2.3	2.8	0.1
8257	572	54.0	10.1	0.7	0.2	33.6	1.2	0.2
8263	155	63.9	14.8	15.5	0.0	1.3	4.5	0.0
8264	239	55.2	30.5	6.3	0.0	2.1	5.9	0.0
8267	238	56.7	18.1	21.4	0.4	0.8	2.1	0.4
8901	279	12.2	82.8	1.8	0.0	1.8	1.1	0.4
8902	425	9.6	85.4	1.9	0.0	0.2	2.8	0.0
8903	470	7.7	84.0	1.9	0.2	1.3	4.9	0.0
8904	678	3.5	91.4	1.6	0.1	1.0	1.8	0.4
8905	753	59.6	28.4	7.3	0.3	1.9	2.3	0.3
8909	424	5.4	90.3	2.8	0.0	0.0	0.9	0.5
9045	454	11.0	23.3	63.4	0.0	0.4	1.5	0.2
9050	300	20.7	60.0	16.7	0.0	0.0	2.7	0.0
9051	471	8.1	4.5	85.4	0.0	0.2	1.3	0.6
9053	481	5.8	86.7	2.7	0.0	2.1	2.5	0.2
9058	470	16.2	72.1	1.3	0.0	8.3	2.1	0.0
9066	321	2.8	94.1	0.6	0.0	0.3	2.2	0.0
9068	277	2.9	94.2	0.0	0.0	0.4	1.8	0.7
9069	600	1.5	96.8	1.5	0.0	0.0	0.2	0.0
9070	562	4.4	90.6	1.8	0.4	1.6	1.2	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2021 Pregnancy Nutrition Surveillance (1)

Page : 9

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9072	59	*	*	*	*	*	*	*
9080	818	17.8	77.4	2.1	0.1	0.1	2.3	0.1
9082	46	*	*	*	*	*	*	*
9083	4	*	*	*	*	*	*	*
9087	8	*	*	*	*	*	*	*
Michigan	48,337	51.3	30.0	11.3	0.3	2.4	4.5	0.3
National (Prior Year)	99,423	47.9	24.1	19.8	1.6	2.8	3.6	0.2

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2021 Pregnancy Nutrition Surveillance (1)

Page : 10

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	35	*	*	*	*	*	*	*
0102	135	0.0	2.2	8.1	61.5	26.7	1.5	0.0
0103	124	0.0	3.2	10.5	61.3	24.2	0.8	0.0
0104	42	*	*	*	*	*	*	*
0105	3	*	*	*	*	*	*	*
0250	39	*	*	*	*	*	*	*
0251	31	*	*	*	*	*	*	*
0252	28	*	*	*	*	*	*	*
0253	36	*	*	*	*	*	*	*
0401	142	0.0	2.1	7.7	58.5	30.3	1.4	0.0
0402	125	0.8	2.4	8.0	64.0	24.0	0.8	0.0
0403	43	*	*	*	*	*	*	*
0404	38	*	*	*	*	*	*	*
0501	137	0.0	1.5	13.1	59.9	25.5	0.0	0.0
0503	22	*	*	*	*	*	*	*
0504	119	0.0	2.5	6.7	58.0	29.4	3.4	0.0
0505	69	*	*	*	*	*	*	*
0506	60	*	*	*	*	*	*	*
0507	34	*	*	*	*	*	*	*
0508	5	*	*	*	*	*	*	*
0650	355	0.0	2.0	7.9	64.5	23.4	2.0	0.3
0651	183	0.5	2.7	12.6	60.1	23.0	1.1	0.0
0652	115	0.0	0.9	12.2	61.7	24.3	0.9	0.0
0653	131	0.0	2.3	3.8	64.9	27.5	0.8	0.8

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2021 Pregnancy Nutrition Surveillance (1)

Page : 11

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0654	138	0.0	3.6	9.4	61.6	21.0	2.9	1.4
0655	104	0.0	5.8	4.8	53.8	32.7	2.9	0.0
0657	19	*	*	*	*	*	*	*
0701	184	0.0	1.6	8.7	56.5	31.0	2.2	0.0
0702	17	*	*	*	*	*	*	*
0703	50	*	*	*	*	*	*	*
0704	14	*	*	*	*	*	*	*
0705	14	*	*	*	*	*	*	*
0850	217	0.0	2.3	6.5	59.0	30.0	1.8	0.5
0851	186	0.0	2.7	5.4	63.4	26.9	0.0	1.6
0901	501	0.0	2.0	5.8	62.7	28.3	1.2	0.0
0902	11	*	*	*	*	*	*	*
1051	398	0.0	0.8	6.3	61.6	29.9	1.5	0.0
1052	1	*	*	*	*	*	*	*
1101	488	0.4	2.9	10.5	61.1	22.5	1.8	0.8
1102	263	0.0	3.4	6.1	62.7	25.9	0.8	1.1
1106	16	*	*	*	*	*	*	*
1250	326	0.0	1.8	7.4	62.6	23.9	4.0	0.3
1251	206	1.5	1.9	5.8	63.1	24.3	2.4	1.0
1252	165	0.0	3.6	8.5	63.6	20.0	4.2	0.0
1253	215	0.0	4.2	5.6	66.0	22.8	1.4	0.0
1350	831	0.0	2.4	7.8	63.3	24.2	1.4	0.8
1351	1	*	*	*	*	*	*	*
1352	127	0.0	0.0	7.9	70.9	20.5	0.8	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2021 Pregnancy Nutrition Surveillance (1)

Page : 12

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1401	73	*	*	*	*	*	*	*
1402	37	*	*	*	*	*	*	*
1403	5	*	*	*	*	*	*	*
1501	242	0.0	2.5	7.9	66.5	21.1	1.7	0.4
1503	267	0.4	2.6	10.1	58.4	25.1	1.9	1.5
1504	626	0.0	2.1	6.1	59.9	30.0	1.9	0.0
1505	22	*	*	*	*	*	*	*
1506	185	0.0	1.1	4.3	70.3	21.6	2.2	0.5
1507	192	0.0	1.0	5.7	57.3	32.8	2.6	0.5
1509	143	0.0	2.1	6.3	56.6	31.5	3.5	0.0
1511	207	0.0	2.4	6.3	57.0	32.9	1.0	0.5
1512	110	0.0	2.7	5.5	59.1	32.7	0.0	0.0
1514	210	0.5	0.0	4.8	66.7	23.8	3.8	0.5
1701	180	0.0	2.2	6.7	61.1	28.9	1.1	0.0
1950	201	0.0	4.0	8.0	57.7	28.4	1.5	0.5
1952	44	*	*	*	*	*	*	*
1953	88	*	*	*	*	*	*	*
1954	118	0.0	0.8	7.6	68.6	22.0	0.8	0.0
1955	35	*	*	*	*	*	*	*
1957	132	0.0	0.8	5.3	56.1	36.4	1.5	0.0
1963	23	*	*	*	*	*	*	*
2150	200	0.5	0.0	6.5	64.5	27.5	1.0	0.0
2151	77	*	*	*	*	*	*	*
2152	13	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2021 Pregnancy Nutrition Surveillance (1)

Page : 13

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2251	141	0.0	0.7	5.7	65.2	27.7	0.7	0.0
2252	47	*	*	*	*	*	*	*
2554	19	*	*	*	*	*	*	*
2555	2,371	0.2	2.0	7.2	63.1	25.6	1.8	0.2
2556	61	*	*	*	*	*	*	*
2560	1	*	*	*	*	*	*	*
2562	2	*	*	*	*	*	*	*
2563	3	*	*	*	*	*	*	*
3001	391	0.0	2.6	9.7	63.9	22.3	0.8	0.8
3002	54	*	*	*	*	*	*	*
3004	95	*	*	*	*	*	*	*
3005	50	*	*	*	*	*	*	*
3007	4	*	*	*	*	*	*	*
3201	124	0.0	0.8	1.6	61.3	35.5	0.8	0.0
3301	1,467	0.1	2.4	6.8	61.2	27.1	2.0	0.3
3304	28	*	*	*	*	*	*	*
3305	41	*	*	*	*	*	*	*
3306	66	*	*	*	*	*	*	*
3308	31	*	*	*	*	*	*	*
3312	137	0.0	2.2	8.0	62.0	27.0	0.7	0.0
3550	26	*	*	*	*	*	*	*
3551	3	*	*	*	*	*	*	*
3552	1	*	*	*	*	*	*	*
3701	292	0.3	1.7	6.2	65.8	23.6	1.7	0.7

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2021 Pregnancy Nutrition Surveillance (1)

Page : 14

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3850	960	0.1	2.9	7.4	62.5	24.8	1.6	0.7
3901	563	0.2	3.4	7.6	57.2	29.7	1.6	0.4
3902	7	*	*	*	*	*	*	*
3903	2	*	*	*	*	*	*	*
3950	662	0.0	2.6	7.3	61.6	26.1	1.8	0.6
4106	705	0.0	2.6	5.7	59.4	28.2	3.0	1.1
4107	945	0.0	2.9	6.8	61.0	26.1	2.3	1.0
4109	207	0.0	1.9	4.8	58.9	32.9	1.4	0.0
4112	198	0.5	6.6	10.1	51.0	27.8	4.0	0.0
4115	1,130	0.1	1.3	5.7	58.1	31.6	2.7	0.5
4351	159	0.0	0.6	5.0	71.7	19.5	2.5	0.6
4352	57	*	*	*	*	*	*	*
4353	161	0.0	0.6	7.5	67.1	21.7	2.5	0.6
4354	257	0.0	1.6	7.4	65.4	23.3	1.9	0.4
4355	88	*	*	*	*	*	*	*
4356	119	0.0	3.4	6.7	64.7	22.7	0.8	1.7
4359	205	0.0	3.9	7.3	58.5	26.3	2.9	1.0
4360	71	*	*	*	*	*	*	*
4361	94	*	*	*	*	*	*	*
4362	92	*	*	*	*	*	*	*
4363	214	0.0	2.8	7.5	63.6	25.7	0.5	0.0
4401	255	0.0	2.4	4.7	62.0	27.8	2.7	0.4
4501	251	0.0	2.4	6.4	63.3	25.9	1.6	0.4
4601	389	0.0	0.5	7.2	65.3	25.7	0.8	0.5

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2021 Pregnancy Nutrition Surveillance (1)

Page : 15

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4701	340	0.0	1.5	6.2	55.9	33.5	2.4	0.6
5050	496	0.0	0.8	3.8	58.3	35.1	2.0	0.0
5051	883	0.0	0.6	4.4	61.8	30.6	2.6	0.0
5052	848	0.0	1.5	6.0	56.1	34.2	2.1	0.0
5055	438	0.0	0.7	2.1	47.3	45.4	4.3	0.2
5101	518	0.2	0.6	2.9	59.1	35.7	1.5	0.0
5102	3	*	*	*	*	*	*	*
5202	222	0.0	3.2	3.2	64.9	27.0	1.8	0.0
5206	4	*	*	*	*	*	*	*
5601	330	0.0	1.2	7.9	63.9	25.5	1.5	0.0
5801	596	0.0	1.7	6.4	64.4	26.2	1.2	0.2
5803	11	*	*	*	*	*	*	*
5804	8	*	*	*	*	*	*	*
5806	9	*	*	*	*	*	*	*
6101	19	*	*	*	*	*	*	*
6102	27	*	*	*	*	*	*	*
6103	1,097	0.1	2.5	10.0	60.7	24.9	1.1	0.7
6105	2	*	*	*	*	*	*	*
6301	1,067	0.0	2.2	6.0	58.5	31.5	1.8	0.0
6312	590	0.0	0.5	3.4	53.7	39.3	2.9	0.2
6316	354	0.0	0.8	3.1	53.1	37.9	5.1	0.0
6320	186	0.0	0.0	2.7	64.0	29.0	3.8	0.5
6325	106	0.0	1.9	2.8	60.4	34.9	0.0	0.0
6335	298	0.0	0.0	2.7	54.0	39.6	3.7	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS
Table 4A
2021 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic

Run Date: 5/29/2024

Page : 16

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
6345	39	*	*	*	*	*	*	*
6360	385	0.0	2.9	9.4	60.0	26.0	1.0	0.8
7306	729	0.0	2.9	7.3	63.6	24.6	1.1	0.5
7401	590	0.0	1.7	7.8	63.4	23.9	2.2	1.0
7601	173	0.0	3.5	6.4	61.3	28.9	0.0	0.0
7850	331	0.0	3.3	6.6	65.3	24.2	0.6	0.0
7852	3	*	*	*	*	*	*	*
8101	579	0.2	2.6	5.9	55.6	32.5	2.6	0.7
8102	196	0.0	2.6	4.6	61.2	29.6	2.0	0.0
8105	1	*	*	*	*	*	*	*
8106	196	0.0	2.6	3.6	68.4	25.0	0.5	0.0
8150	5	*	*	*	*	*	*	*
8151	1,079	0.1	1.3	5.9	56.6	32.9	2.7	0.5
8152	2	*	*	*	*	*	*	*
8155	3	*	*	*	*	*	*	*
8201	922	0.0	1.3	6.1	58.2	32.9	1.5	0.0
8202	1,245	0.0	1.1	3.4	52.3	37.2	5.5	0.6
8203	132	0.0	0.8	6.8	59.1	28.0	5.3	0.0
8250	159	0.0	0.6	4.4	52.2	39.6	2.5	0.6
8252	57	*	*	*	*	*	*	*
8255	688	0.1	1.0	6.0	59.7	31.7	1.5	0.0
8257	572	0.0	0.7	3.1	51.9	39.5	4.7	0.0
8263	155	0.0	3.2	4.5	52.3	37.4	2.6	0.0
8264	239	0.0	0.0	2.1	59.0	35.6	3.3	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2021 Pregnancy Nutrition Surveillance (1)

Page : 17

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8267	238	0.0	0.8	6.3	64.7	26.1	2.1	0.0
8901	279	0.4	2.5	6.5	57.7	29.4	3.2	0.4
8902	425	0.0	1.4	3.1	58.6	33.2	3.3	0.5
8903	470	0.0	2.8	7.2	56.0	31.3	2.1	0.6
8904	678	0.0	1.8	6.8	56.6	32.4	1.9	0.4
8905	753	0.0	0.8	3.3	56.4	35.5	3.7	0.3
8909	424	0.0	1.4	7.3	62.0	26.4	2.8	0.0
9045	454	0.2	2.9	7.7	58.6	26.2	4.0	0.4
9050	300	0.3	2.7	6.0	59.3	29.0	2.0	0.7
9051	471	0.0	1.5	6.2	56.7	31.6	4.0	0.0
9053	481	0.0	2.9	6.2	55.7	32.4	2.3	0.4
9058	470	0.0	1.9	6.0	58.5	31.1	2.1	0.4
9066	321	0.3	2.5	6.5	54.5	33.6	2.2	0.3
9068	277	0.0	5.4	7.9	57.8	24.9	2.9	1.1
9069	600	0.2	3.5	5.8	54.2	34.0	1.8	0.5
9070	562	0.0	3.6	8.4	55.2	31.7	1.1	0.2
9072	59	*	*	*	*	*	*	*
9080	818	0.0	2.2	6.2	55.4	32.9	2.7	0.6
9082	46	*	*	*	*	*	*	*
9083	4	*	*	*	*	*	*	*
9087	8	*	*	*	*	*	*	*
Michigan	48,337	0.1	2.0	6.3	59.7	29.4	2.2	0.4
National (Prior Year)	99,433	0.1	2.4	7.1	59.5	28.4	2.3	0.2

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 18

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	26	*	*	17	*	*	0	*	2	*	32	*	32	*
0102	99	*	*	60	*	*	0	*	1	*	122	8.2 (41)	122	10.7 (21)
0103	102	2.0 (7)	68.6 (95)	54	*	*	0	*	4	*	112	4.5 (2)	113	16.8 (89)
0104	28	*	*	11	*	*	0	*	1	*	39	*	39	*
0105	3	*	*	2	*	*	0	*	0	*	3	*	3	*
0250	33	*	*	27	*	*	0	*	0	*	37	*	37	*
0251	28	*	*	13	*	*	0	*	0	*	27	*	27	*
0252	26	*	*	19	*	*	0	*	0	*	25	*	25	*
0253	34	*	*	22	*	*	0	*	0	*	33	*	33	*
0401	127	4.7 (74)	65.4 (78)	73	*	*	0	*	0	*	134	11.9 (98)	134	19.4 (106)
0402	102	6.9 (98)	58.8 (17)	28	*	*	0	*	1	*	114	11.4 (91)	114	14.9 (73)
0403	40	*	*	17	*	*	0	*	1	*	39	*	39	*
0404	34	*	*	19	*	*	0	*	0	*	36	*	36	*
0501	73	*	*	22	*	*	11	*	16	*	125	8.8 (51)	125	11.2 (25)
0503	10	*	*	4	*	*	1	*	3	*	21	*	21	*
0504	99	*	*	36	*	*	11	*	19	*	112	8.0 (35)	112	13.4 (51)
0505	39	*	*	10	*	*	9	*	7	*	62	*	62	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 19

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	37	*	*	13	*	*	1	*	12	*	56	*	56	*
0507	16	*	*	5	*	*	1	*	7	*	34	*	34	*
0508	2	*	*	2	*	*	0	*	1	*	5	*	5	*
0650	303	6.9 (98)	60.1 (22)	181	13.3 (5)	53.6 (31)	3	*	6	*	334	11.1 (89)	335	13.7 (57)
0651	144	4.2 (64)	66.0 (84)	72	*	*	0	*	0	*	168	5.4 (5)	170	10.6 (20)
0652	106	1.9 (5)	62.3 (49)	85	*	*	0	*	1	*	113	7.1 (23)	113	15.9 (80)
0653	118	5.1 (84)	61.0 (32)	93	*	*	2	*	9	*	122	8.2 (41)	123	11.4 (28)
0654	77	*	*	34	*	*	0	*	1	*	126	7.1 (23)	127	11.8 (33)
0655	94	*	*	57	*	*	1	*	1	*	99	*	100	29.0 (112)
0657	11	*	*	2	*	*	0	*	0	*	19	*	19	*
0701	162	3.1 (28)	58.0 (15)	120	16.7 (32)	45.0 (5)	1	*	1	*	165	7.3 (28)	165	17.0 (91)
0702	17	*	*	14	*	*	0	*	0	*	16	*	16	*
0703	43	*	*	32	*	*	2	*	1	*	42	*	42	*
0704	12	*	*	4	*	*	1	*	0	*	10	*	11	*
0705	11	*	*	11	*	*	1	*	0	*	14	*	14	*
0850	124	6.5 (95)	62.9 (58)	46	*	*	3	*	1	*	201	9.0 (52)	201	17.4 (96)
0851	124	6.5 (95)	57.3 (11)	61	*	*	1	*	3	*	174	9.8 (66)	177	14.1 (63)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 20

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0901	398	3.3 (37)	61.8 (44)	254	18.5 (44)	58.7 (58)	0	*	2	*	470	10.0 (73)	470	13.4 (51)
0902	9	*	*	6	*	*	0	*	0	*	10	*	10	*
1051	321	5.9 (93)	55.5 (6)	243	19.8 (50)	52.3 (25)	3	*	1	*	366	9.6 (64)	367	9.5 (12)
1052	1	*	*	0	*	*	0	*	0	*	1	*	1	*
1101	410	4.1 (59)	61.5 (35)	291	17.5 (36)	55.0 (39)	24	*	19	*	451	9.1 (53)	452	19.9 (108)
1102	228	6.1 (94)	60.1 (22)	164	15.2 (17)	58.5 (57)	8	*	10	*	227	9.3 (58)	228	17.1 (92)
1106	13	*	*	9	*	*	0	*	0	*	12	*	12	*
1250	97	*	*	23	*	*	2	*	3	*	295	11.5 (94)	296	11.8 (33)
1251	160	4.4 (69)	61.9 (45)	102	15.7 (25)	49.0 (10)	1	*	9	*	184	6.5 (18)	185	7.6 (6)
1252	121	4.1 (59)	62.8 (56)	44	*	*	1	*	5	*	144	13.9 (106)	145	11.7 (31)
1253	185	8.1 (102)	60.5 (26)	56	*	*	2	*	5	*	187	12.3 (100)	187	13.4 (51)
1350	699	2.9 (22)	63.7 (66)	496	16.5 (31)	52.0 (23)	0	*	3	*	787	6.6 (19)	787	10.0 (15)
1351	0	*	*	0	*	*	0	*	0	*	1	*	1	*
1352	76	*	*	50	*	*	0	*	0	*	117	6.0 (14)	117	6.8 (5)
1401	51	*	*	22	*	*	2	*	5	*	66	*	66	*
1402	29	*	*	15	*	*	0	*	2	*	32	*	32	*
1403	4	*	*	2	*	*	0	*	0	*	4	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 21

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1501	196	3.1 (28)	67.9 (90)	76	*	*	22	*	23	*	218	14.2 (107)	217	12.9 (48)
1503	221	4.1 (59)	65.6 (81)	161	19.9 (52)	47.8 (8)	14	*	19	*	244	9.8 (66)	244	13.9 (58)
1504	134	5.2 (86)	53.0 (4)	65	*	*	32	*	79	*	562	9.8 (66)	562	10.3 (19)
1505	9	*	*	0	*	*	0	*	0	*	19	*	19	*
1506	143	4.9 (79)	67.1 (88)	20	*	*	8	*	24	*	172	15.1 (108)	172	12.2 (39)
1507	126	5.6 (91)	47.6 (1)	24	*	*	7	*	29	*	174	9.8 (66)	174	9.2 (11)
1509	104	3.8 (52)	71.2 (101)	43	*	*	6	*	19	*	126	9.5 (62)	127	7.9 (7)
1511	161	2.5 (15)	61.5 (35)	91	*	*	13	*	29	*	194	10.8 (84)	195	16.4 (84)
1512	83	*	*	33	*	*	7	*	30	*	104	3.8 (1)	104	16.3 (82)
1514	100	2.0 (7)	56.0 (7)	32	*	*	7	*	46	*	201	10.9 (85)	201	18.4 (103)
1701	154	3.2 (34)	59.7 (19)	106	13.2 (4)	56.6 (48)	0	*	1	*	165	12.7 (101)	165	7.9 (7)
1950	180	3.3 (37)	63.3 (62)	128	15.6 (24)	55.5 (42)	2	*	0	*	188	11.7 (96)	188	16.5 (86)
1952	36	*	*	30	*	*	2	*	1	*	42	*	42	*
1953	73	*	*	53	*	*	0	*	1	*	79	*	79	*
1954	106	3.8 (52)	70.8 (99)	77	*	*	0	*	0	*	113	13.3 (103)	113	14.2 (65)
1955	31	*	*	23	*	*	0	*	1	*	33	*	33	*
1957	121	4.1 (59)	71.1 (100)	78	*	*	0	*	0	*	125	19.2 (111)	125	14.4 (68)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 22

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1963	17	*	*	10	*	*	0	*	0	*	21	*	21	*
2150	192	3.1 (28)	60.4 (25)	142	20.4 (56)	45.1 (6)	12	*	31	*	183	8.7 (50)	183	14.8 (72)
2151	74	*	*	54	*	*	3	*	4	*	66	*	66	*
2152	13	*	*	10	*	*	1	*	0	*	12	*	12	*
2251	83	*	*	40	*	*	0	*	1	*	131	4.6 (3)	131	17.6 (97)
2252	26	*	*	5	*	*	0	*	0	*	43	*	43	*
2554	12	*	*	2	*	*	2	*	1	*	18	*	18	*
2555	1,612	5.2 (86)	59.7 (19)	797	19.9 (52)	51.9 (22)	33	*	47	*	2,186	9.1 (53)	2,191	17.3 (94)
2556	39	*	*	19	*	*	0	*	0	*	58	*	58	*
2560	1	*	*	0	*	*	0	*	0	*	1	*	1	*
2562	2	*	*	0	*	*	0	*	0	*	2	*	2	*
2563	3	*	*	3	*	*	1	*	0	*	3	*	3	*
3001	244	4.5 (72)	69.7 (97)	65	*	*	0	*	0	*	369	8.1 (37)	369	14.6 (70)
3002	34	*	*	8	*	*	0	*	0	*	51	*	51	*
3004	57	*	*	12	*	*	0	*	0	*	89	*	89	*
3005	35	*	*	4	*	*	0	*	0	*	49	*	49	*
3007	2	*	*	0	*	*	0	*	0	*	4	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 23

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3201	113	5.3 (89)	65.5 (80)	89	*	*	4	*	14	*	120	10.0 (73)	120	16.7 (88)
3301	1,215	3.6 (48)	61.5 (35)	884	19.6 (49)	56.3 (46)	10	*	2	*	1,400	8.1 (37)	1,401	12.3 (41)
3304	22	*	*	19	*	*	0	*	0	*	27	*	27	*
3305	34	*	*	25	*	*	0	*	0	*	38	*	38	*
3306	56	*	*	47	*	*	0	*	0	*	64	*	64	*
3308	30	*	*	22	*	*	0	*	0	*	26	*	26	*
3312	121	2.5 (15)	65.3 (77)	94	*	*	0	*	0	*	136	11.0 (86)	135	14.1 (63)
3550	21	*	*	10	*	*	0	*	0	*	25	*	25	*
3551	2	*	*	1	*	*	0	*	0	*	3	*	3	*
3552	1	*	*	1	*	*	0	*	0	*	1	*	1	*
3701	197	3.6 (48)	68.5 (94)	149	20.1 (54)	51.7 (19)	17	*	20	*	269	9.7 (65)	272	22.1 (111)
3850	630	3.3 (37)	57.0 (9)	421	15.4 (19)	51.8 (20)	15	*	36	*	901	8.2 (41)	902	12.4 (44)
3901	417	3.1 (28)	65.7 (82)	183	16.9 (33)	57.4 (54)	0	*	4	*	501	7.2 (26)	502	13.9 (58)
3902	6	*	*	2	*	*	0	*	0	*	7	*	7	*
3903	1	*	*	1	*	*	0	*	0	*	2	*	2	*
3950	492	3.0 (26)	66.1 (86)	257	12.5 (2)	58.0 (56)	0	*	2	*	616	10.6 (81)	617	11.7 (31)
4106	305	3.0 (26)	63.6 (64)	98	*	*	57	*	124	21.8 (1)	664	7.5 (29)	666	16.1 (81)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 24

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4107	615	3.6 (48)	61.0 (32)	258	17.4 (35)	53.9 (35)	62	*	195	23.1 (2)	865	7.9 (33)	867	19.8 (107)
4109	137	2.9 (22)	59.9 (21)	62	*	*	9	*	54	*	195	8.2 (41)	195	14.9 (73)
4112	188	0.5 (1)	68.6 (95)	153	48.4 (65)	26.8 (1)	1	*	4	*	187	17.6 (110)	187	18.7 (104)
4115	477	3.4 (41)	61.2 (34)	116	12.9 (3)	57.8 (55)	84	*	253	26.5 (3)	1,046	8.4 (48)	1,047	15.8 (79)
4351	140	2.9 (22)	60.7 (30)	53	*	*	3	*	1	*	146	11.0 (86)	146	17.1 (92)
4352	46	*	*	37	*	*	0	*	0	*	54	*	54	*
4353	122	4.9 (79)	69.7 (97)	93	*	*	5	*	6	*	147	13.6 (105)	147	12.2 (39)
4354	149	4.0 (57)	62.4 (52)	38	*	*	0	*	3	*	242	9.1 (53)	242	17.8 (98)
4355	35	*	*	16	*	*	0	*	0	*	82	*	82	*
4356	109	2.8 (20)	66.1 (86)	78	*	*	1	*	2	*	104	16.3 (109)	104	18.3 (101)
4359	190	3.2 (34)	67.9 (90)	136	25.7 (63)	44.1 (4)	0	*	2	*	190	8.4 (48)	190	18.9 (105)
4360	63	*	*	46	*	*	2	*	0	*	67	*	67	*
4361	84	*	*	48	*	*	0	*	1	*	87	*	87	*
4362	73	*	*	51	*	*	1	*	0	*	86	*	87	*
4363	175	7.4 (100)	57.7 (13)	130	23.1 (61)	50.8 (15)	0	*	0	*	207	12.1 (99)	207	16.9 (90)
4401	248	4.4 (69)	56.5 (8)	208	16.3 (29)	47.6 (7)	0	*	0	*	234	5.6 (6)	235	10.2 (18)
4501	188	3.7 (51)	66.0 (84)	133	15.8 (26)	49.6 (11)	6	*	5	*	245	11.0 (86)	245	13.5 (55)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 25

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4601	326	2.1 (12)	61.7 (40)	234	15.4 (19)	50.4 (14)	0	*	0	*	345	9.9 (71)	346	14.2 (65)
4701	310	4.8 (76)	57.7 (13)	200	20.5 (57)	54.0 (36)	1	*	0	*	308	8.1 (37)	309	20.1 (109)
5050	466	3.4 (41)	67.4 (89)	371	18.3 (43)	52.0 (23)	0	*	0	*	472	9.3 (58)	473	16.3 (82)
5051	833	3.1 (28)	62.3 (49)	679	17.8 (39)	56.8 (49)	0	*	0	*	810	9.8 (66)	810	10.1 (17)
5052	804	2.9 (22)	62.4 (52)	618	18.6 (46)	48.7 (9)	0	*	0	*	798	11.4 (91)	798	12.3 (41)
5055	250	1.2 (2)	63.6 (64)	124	18.5 (44)	51.6 (18)	0	*	0	*	422	10.2 (75)	422	5.2 (3)
5101	372	4.3 (66)	58.3 (16)	285	13.7 (6)	56.5 (47)	13	*	2	*	480	10.2 (75)	480	12.3 (41)
5102	2	*	*	1	*	*	0	*	0	*	2	*	2	*
5202	190	1.6 (3)	63.2 (61)	129	15.5 (22)	66.7 (65)	2	*	1	*	205	8.3 (45)	207	11.1 (23)
5206	1	*	*	0	*	*	0	*	0	*	4	*	4	*
5601	222	2.7 (17)	64.0 (70)	102	9.8 (1)	60.8 (61)	5	*	4	*	315	9.5 (62)	314	12.1 (37)
5801	557	4.3 (66)	64.8 (72)	442	15.8 (26)	56.1 (44)	0	*	0	*	552	8.3 (45)	552	18.1 (100)
5803	11	*	*	11	*	*	0	*	0	*	11	*	11	*
5804	8	*	*	5	*	*	0	*	0	*	8	*	8	*
5806	6	*	*	3	*	*	0	*	0	*	8	*	8	*
6101	18	*	*	12	*	*	0	*	0	*	17	*	18	*
6102	24	*	*	18	*	*	0	*	0	*	24	*	24	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 26

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6103	999	3.5 (45)	61.7 (40)	803	17.8 (39)	51.8 (20)	0	*	4	*	1,023	10.2 (75)	1,029	16.5 (86)
6105	1	*	*	0	*	*	0	*	0	*	2	*	2	*
6301	874	3.4 (41)	62.5 (54)	542	20.8 (59)	43.7 (3)	1	*	0	*	978	5.8 (9)	978	11.1 (23)
6312	494	4.7 (74)	59.5 (18)	299	14.0 (9)	57.2 (53)	0	*	0	*	536	6.9 (21)	536	9.0 (10)
6316	307	4.9 (79)	60.9 (31)	192	16.1 (28)	56.8 (49)	0	*	0	*	331	5.7 (7)	331	11.2 (25)
6320	125	4.8 (76)	64.8 (72)	64	*	*	1	*	0	*	163	13.5 (104)	163	17.8 (98)
6325	69	*	*	34	*	*	0	*	0	*	94	*	94	*
6335	196	7.7 (101)	57.1 (10)	99	*	*	1	*	0	*	282	11.3 (90)	282	14.5 (69)
6345	28	*	*	15	*	*	0	*	0	*	36	*	36	*
6360	260	3.5 (45)	61.9 (45)	117	15.4 (19)	54.7 (37)	0	*	0	*	346	6.6 (19)	346	13.9 (58)
7306	540	2.8 (20)	62.0 (48)	335	14.6 (14)	53.7 (34)	0	*	0	*	693	9.4 (60)	692	11.6 (29)
7401	555	4.1 (59)	63.1 (59)	437	14.4 (13)	55.6 (43)	4	*	13	*	533	8.1 (37)	534	15.4 (77)
7601	44	*	*	13	*	*	1	*	2	*	159	6.3 (15)	159	12.6 (46)
7850	289	5.2 (86)	63.7 (66)	236	23.3 (62)	50.8 (15)	1	*	1	*	305	5.9 (12)	305	6.6 (4)
7852	3	*	*	2	*	*	0	*	0	*	3	*	3	*
8101	523	4.2 (64)	65.2 (75)	349	17.5 (36)	56.2 (45)	0	*	0	*	551	8.3 (45)	551	18.3 (101)
8102	147	2.0 (7)	72.1 (102)	80	*	*	0	*	0	*	187	6.4 (16)	187	13.4 (51)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 27

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8105	1	*	*	0	*	*	0	*	0	*	0	*	0	*
8106	108	6.5 (95)	50.9 (3)	48	*	*	0	*	0	*	185	5.9 (12)	185	17.3 (94)
8150	4	*	*	2	*	*	0	*	0	*	5	*	5	*
8151	859	3.5 (45)	61.9 (45)	542	17.3 (34)	51.5 (17)	34	*	51	*	979	10.6 (81)	979	15.1 (75)
8152	1	*	*	1	*	*	0	*	1	*	2	*	2	*
8155	2	*	*	1	*	*	0	*	0	*	3	*	3	*
8201	662	2.7 (17)	60.6 (29)	430	14.0 (9)	60.2 (60)	0	*	4	*	813	11.4 (91)	815	11.8 (33)
8202	584	3.9 (55)	53.4 (5)	346	15.3 (18)	52.9 (29)	0	*	0	*	1,215	7.2 (26)	1,215	3.6 (2)
8203	113	2.7 (17)	60.2 (24)	74	*	*	0	*	1	*	114	10.5 (80)	113	9.7 (14)
8250	123	3.3 (37)	65.9 (83)	71	*	*	0	*	0	*	137	5.8 (9)	137	9.5 (12)
8252	40	*	*	16	*	*	0	*	0	*	48	*	48	*
8255	583	4.5 (72)	60.5 (26)	389	20.1 (54)	53.5 (30)	0	*	0	*	609	11.8 (97)	609	11.8 (33)
8257	217	5.5 (90)	49.8 (2)	120	29.2 (64)	36.7 (2)	0	*	0	*	550	8.0 (35)	550	3.3 (1)
8263	133	3.8 (52)	61.7 (40)	81	*	*	0	*	0	*	132	10.6 (81)	132	12.1 (37)
8264	190	2.1 (12)	68.4 (93)	128	14.1 (11)	61.7 (63)	0	*	0	*	216	7.9 (33)	215	13.5 (55)
8267	191	5.8 (92)	62.8 (56)	127	18.9 (47)	55.1 (41)	0	*	0	*	212	7.5 (29)	212	10.8 (22)
8901	229	4.4 (69)	64.2 (71)	145	19.3 (48)	50.3 (13)	0	*	0	*	262	7.6 (31)	263	15.6 (78)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 28

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8902	276	5.1 (84)	60.5 (26)	151	13.9 (8)	53.6 (31)	0	*	0	*	396	9.1 (53)	396	11.6 (29)
8903	299	2.3 (14)	63.9 (69)	205	13.7 (6)	60.0 (59)	0	*	0	*	434	7.1 (23)	437	15.1 (75)
8904	554	3.1 (28)	65.2 (75)	293	16.4 (30)	53.6 (31)	0	*	0	*	639	9.1 (53)	640	16.4 (84)
8905	507	3.9 (55)	61.5 (35)	282	18.1 (42)	52.5 (27)	0	*	0	*	719	12.8 (102)	719	14.0 (62)
8909	328	3.4 (41)	63.1 (59)	177	17.5 (36)	57.1 (52)	0	*	0	*	384	9.9 (71)	384	12.8 (47)
9045	321	1.9 (5)	68.2 (92)	190	14.7 (15)	50.0 (12)	0	*	0	*	438	9.4 (60)	438	8.0 (9)
9050	257	4.3 (66)	65.4 (78)	64	*	*	0	*	0	*	282	7.8 (32)	282	13.1 (50)
9051	148	2.0 (7)	57.4 (12)	53	*	*	0	*	0	*	442	11.5 (94)	442	10.0 (15)
9053	423	4.0 (57)	65.0 (74)	289	14.9 (16)	61.2 (62)	1	*	0	*	451	6.4 (16)	452	13.9 (58)
9058	309	4.9 (79)	63.4 (63)	208	17.8 (39)	54.8 (38)	0	*	0	*	448	10.3 (78)	449	14.7 (71)
9066	250	4.8 (76)	61.6 (39)	161	20.5 (57)	52.8 (28)	0	*	0	*	308	5.2 (4)	309	20.1 (109)
9068	256	2.0 (7)	62.5 (54)	204	22.1 (60)	56.9 (51)	0	*	0	*	258	5.8 (9)	259	11.2 (25)
9069	432	3.2 (34)	62.3 (49)	252	19.8 (50)	52.4 (26)	0	*	1	*	570	7.0 (22)	570	13.0 (49)
9070	492	4.9 (79)	63.8 (68)	334	14.1 (11)	61.7 (63)	0	*	1	*	529	5.7 (7)	529	14.2 (65)
9072	53	*	*	16	*	*	0	*	0	*	47	*	47	*
9080	515	1.7 (4)	61.7 (40)	291	15.5 (22)	55.0 (39)	0	*	1	*	777	10.4 (79)	775	12.5 (45)
9082	37	*	*	24	*	*	0	*	0	*	45	*	45	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2021 Pregnancy Nutrition Surveillance (1)

Page : 29

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Pregpregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9083	2	*	*	0	*	*	0	*	0	*	4	*	4	*
9087	8	*	*	5	*	*	0	*	0	*	7	*	7	*
Michigan	35,616	3.8	62.1	21,801	17.6	53.3	608	32.1	1,368	23.8	44,841	9.1	44,895	13.5
National (Prior Year)	76,624	4.0	60.8	54,177	21.0	48.0	13,936	20.1	4,844	20.1	52,072	11.4	51,357	13.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 30

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
0101	31	*	*	34	*	33	*	20 *
0102	112	90.2 (63)	1.8 (56)	131	35.1 (20)	125	27.2 (40)	93 *
0103	104	90.4 (60)	2.9 (81)	122	34.4 (25)	117	16.2 (84)	91 *
0104	38	*	*	41	*	40	*	32 *
0105	3	*	*	3	*	3	*	3 *
0250	31	*	*	38	*	36	*	30 *
0251	26	*	*	31	*	29	*	26 *
0252	25	*	*	26	*	24	*	19 *
0253	33	*	*	36	*	36	*	35 *
0401	127	92.1 (44)	1.6 (46)	137	31.4 (39)	135	26.7 (44)	106 96.2 (14)
0402	99	*	*	121	37.2 (15)	121	28.9 (30)	100 96.0 (15)
0403	39	*	*	43	*	42	*	33 *
0404	36	*	*	37	*	37	*	30 *
0501	123	96.7 (7)	1.6 (46)	130	30.8 (44)	128	23.4 (60)	101 94.1 (35)
0503	21	*	*	22	*	22	*	16 *
0504	107	95.3 (17)	0.9 (23)	117	29.9 (48)	116	41.4 (7)	89 *
0505	64	*	*	68	*	67	*	52 *
0506	53	*	*	58	*	55	*	34 *
0507	33	*	*	34	*	34	*	22 *
0508	4	*	*	5	*	5	*	1 *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 31

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	309	90.0 (65)	3.6 (87)	351	35.3 (19)	346	34.7 (20)	260	93.1 (43)
0651	157	89.8 (66)	3.2 (85)	181	39.2 (8)	169	28.4 (35)	148	89.2 (80)
0652	106	90.6 (59)	2.8 (80)	113	37.2 (15)	110	20.0 (71)	96	*
0653	111	79.3 (104)	7.2 (104)	128	35.9 (17)	124	25.0 (53)	93	*
0654	107	85.0 (92)	7.5 (105)	134	47.0 (1)	130	19.2 (74)	104	90.4 (73)
0655	93	*	*	102	43.1 (3)	94	*	86	*
0657	19	*	*	19	*	18	*	17	*
0701	153	90.2 (63)	1.3 (41)	180	31.7 (37)	178	39.3 (12)	138	97.8 (2)
0702	16	*	*	17	*	17	*	9	*
0703	42	*	*	50	*	45	*	29	*
0704	11	*	*	13	*	13	*	9	*
0705	13	*	*	14	*	13	*	9	*
0850	197	90.4 (60)	2.0 (62)	214	22.0 (88)	199	35.2 (17)	146	92.5 (50)
0851	167	92.2 (43)	1.2 (37)	184	27.7 (60)	180	36.1 (16)	121	98.3 (1)
0901	422	86.3 (88)	3.8 (90)	495	37.4 (13)	483	25.1 (52)	377	94.2 (32)
0902	10	*	*	11	*	10	*	8	*
1051	330	89.7 (69)	4.8 (100)	386	31.6 (38)	385	28.8 (32)	278	91.7 (57)
1052	1	*	*	1	*	1	*	1	*
1101	421	80.3 (103)	2.6 (76)	473	23.3 (79)	443	19.4 (73)	350	90.6 (68)
1102	205	84.9 (93)	3.9 (91)	261	31.0 (43)	244	23.0 (61)	172	94.8 (27)
1106	11	*	*	14	*	14	*	9	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 32

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1250	276	85.5 (91)	5.8 (102)	315	31.4 (39)	277	41.5 (6)	229	92.1 (53)
1251	177	93.8 (27)	0.6 (12)	202	27.2 (61)	189	39.7 (10)	148	97.3 (5)
1252	125	68.8 (107)	14.4 (108)	163	30.7 (45)	157	19.1 (76)	120	86.7 (90)
1253	140	75.7 (106)	11.4 (107)	207	39.6 (7)	206	22.3 (64)	163	87.7 (86)
1350	752	96.5 (9)	0.8 (19)	806	29.5 (51)	774	31.7 (23)	591	92.7 (47)
1351	1	*	*	1	*	1	*	1	*
1352	114	100.0 (1)	0.0 (1)	125	32.0 (34)	113	3.5 (107)	89	*
1401	58	*	*	70	*	70	*	56	*
1402	28	*	*	34	*	31	*	25	*
1403	4	*	*	4	*	5	*	3	*
1501	177	86.4 (87)	4.0 (93)	239	37.7 (11)	219	13.2 (90)	183	82.5 (96)
1503	236	94.1 (23)	1.7 (52)	263	37.3 (14)	245	19.2 (74)	198	76.3 (99)
1504	542	95.8 (13)	0.7 (15)	613	29.9 (48)	574	25.8 (49)	419	85.4 (93)
1505	18	*	*	22	*	13	*	22	*
1506	138	88.4 (76)	4.3 (97)	182	38.5 (9)	177	9.6 (101)	143	90.9 (65)
1507	167	96.4 (10)	1.2 (37)	187	29.4 (52)	175	25.7 (50)	124	97.6 (3)
1509	122	91.8 (49)	1.6 (46)	139	33.8 (27)	128	24.2 (57)	109	95.4 (21)
1511	182	93.4 (31)	1.6 (46)	203	34.5 (22)	199	24.6 (54)	154	91.6 (61)
1512	96	*	*	108	25.9 (70)	103	35.0 (19)	63	*
1514	187	93.0 (38)	1.1 (31)	201	18.9 (102)	163	28.8 (32)	114	96.5 (10)
1701	157	96.8 (6)	1.3 (41)	175	37.7 (11)	171	43.9 (5)	142	93.7 (36)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 33

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1950	180	92.8 (39)	2.2 (67)	194	32.5 (32)	163	27.6 (38)	151	96.7 (7)
1952	38	*	*	44	*	37	*	30	*
1953	73	*	*	87	*	74	*	64	*
1954	104	93.3 (33)	1.0 (30)	115	30.4 (46)	89	*	90	*
1955	32	*	*	34	*	28	*	24	*
1957	119	93.3 (33)	1.7 (52)	127	29.1 (53)	97	*	88	*
1963	21	*	*	22	*	16	*	16	*
2150	182	95.6 (14)	0.0 (1)	196	28.6 (56)	189	25.4 (51)	145	96.6 (8)
2151	62	*	*	75	*	65	*	57	*
2152	12	*	*	13	*	12	*	12	*
2251	126	93.7 (28)	4.0 (93)	140	35.0 (21)	134	33.6 (21)	108	94.4 (30)
2252	38	*	*	46	*	44	*	37	*
2554	19	*	*	19	*	16	*	18	*
2555	2,080	89.8 (66)	1.2 (37)	2,330	23.6 (77)	1,874	16.6 (82)	1,671	93.4 (41)
2556	54	*	*	61	*	49	*	52	*
2560	1	*	*	1	*	1	*	1	*
2562	2	*	*	2	*	2	*	2	*
2563	3	*	*	3	*	3	*	3	*
3001	350	88.0 (77)	3.1 (82)	385	34.5 (22)	374	13.9 (89)	312	88.5 (82)
3002	45	*	*	53	*	48	*	38	*
3004	85	*	*	91	*	90	*	68	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 34

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3005	45	*	*	48	*	49	*	43	*
3007	4	*	*	4	*	4	*	3	*
3201	110	89.1 (73)	1.8 (56)	122	23.0 (80)	118	44.1 (4)	81	*
3301	1,267	91.8 (49)	1.7 (52)	1,449	26.1 (67)	1,314	9.8 (100)	1,040	90.5 (69)
3304	26	*	*	28	*	28	*	17	*
3305	34	*	*	40	*	37	*	23	*
3306	55	*	*	65	*	58	*	45	*
3308	23	*	*	30	*	28	*	16	*
3312	118	90.7 (58)	3.4 (86)	132	31.8 (36)	125	10.4 (99)	101	92.1 (53)
3550	24	*	*	25	*	24	*	19	*
3551	3	*	*	3	*	3	*	3	*
3552	1	*	*	1	*	1	*	1	*
3701	265	93.6 (29)	1.9 (60)	284	24.6 (72)	275	20.0 (71)	191	96.3 (12)
3850	815	90.9 (57)	3.6 (87)	944	38.5 (9)	918	24.4 (56)	719	90.5 (69)
3901	467	92.7 (40)	0.6 (12)	547	32.5 (32)	527	16.5 (83)	413	89.3 (79)
3902	7	*	*	7	*	7	*	7	*
3903	2	*	*	2	*	2	*	1	*
3950	578	91.2 (53)	2.6 (76)	643	31.9 (35)	613	21.9 (65)	458	90.8 (66)
4106	617	87.2 (83)	2.4 (71)	693	22.9 (82)	620	10.6 (98)	510	88.4 (84)
4107	839	91.3 (52)	2.3 (68)	930	18.9 (102)	867	26.0 (47)	631	93.0 (45)
4109	174	93.1 (37)	1.1 (31)	200	30.0 (47)	194	27.8 (36)	132	95.5 (20)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 35

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4112	185	45.4 (108)	0.0 (1)	193	26.4 (66)	104	51.9 (1)	177	92.1 (53)
4115	953	93.3 (33)	2.4 (71)	1,106	18.9 (102)	812	17.5 (80)	755	95.1 (24)
4351	124	90.3 (62)	0.8 (19)	157	40.8 (5)	141	12.8 (91)	123	87.0 (89)
4352	50	*	*	54	*	54	*	45	*
4353	121	79.3 (104)	9.9 (106)	154	28.6 (56)	142	26.8 (42)	108	85.2 (95)
4354	222	96.4 (10)	1.4 (44)	252	34.1 (26)	242	11.6 (95)	187	90.4 (73)
4355	79	*	*	85	*	86	*	64	*
4356	93	*	*	119	31.1 (42)	111	29.7 (28)	91	*
4359	152	80.9 (102)	6.6 (103)	201	42.8 (4)	197	17.3 (81)	167	88.6 (81)
4360	63	*	*	68	*	69	*	56	*
4361	84	*	*	92	*	87	*	66	*
4362	82	*	*	90	*	81	*	66	*
4363	179	86.6 (84)	3.9 (91)	211	32.7 (30)	200	26.0 (47)	156	89.7 (77)
4401	214	92.1 (44)	4.2 (96)	251	32.7 (30)	241	30.3 (27)	178	94.9 (25)
4501	225	88.0 (77)	4.0 (93)	248	28.2 (59)	238	29.0 (29)	171	92.4 (51)
4601	338	93.2 (36)	0.0 (1)	379	24.8 (71)	371	23.7 (59)	273	92.7 (47)
4701	287	87.5 (80)	4.9 (101)	332	28.9 (55)	318	38.4 (13)	236	95.3 (23)
5050	465	94.6 (20)	0.2 (6)	484	21.1 (93)	414	14.0 (88)	318	97.5 (4)
5051	792	94.1 (23)	0.4 (9)	856	20.9 (94)	682	12.8 (91)	562	95.4 (21)
5052	777	94.1 (23)	0.8 (19)	826	22.8 (83)	674	10.7 (97)	609	94.4 (30)
5055	383	95.6 (14)	3.7 (89)	427	40.5 (6)	335	4.2 (106)	344	93.3 (42)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 36

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5101	451	92.7 (40)	2.0 (62)	513	26.5 (64)	249	40.6 (8)	369	95.7 (17)
5102	2	*	*	3	*	1	*	3	*
5202	191	89.0 (75)	4.7 (98)	218	35.8 (18)	171	22.8 (63)	183	95.6 (19)
5206	3	*	*	3	*	3	*	2	*
5601	296	91.9 (47)	2.7 (79)	323	24.1 (74)	313	24.6 (54)	218	96.3 (12)
5801	539	92.4 (42)	0.9 (23)	585	23.4 (78)	567	28.9 (30)	397	94.2 (32)
5803	11	*	*	11	*	11	*	9	*
5804	7	*	*	8	*	8	*	8	*
5806	8	*	*	9	*	8	*	8	*
6101	14	*	*	18	*	16	*	13	*
6102	19	*	*	26	*	21	*	22	*
6103	958	89.5 (71)	2.3 (68)	1,066	33.5 (29)	889	26.7 (44)	795	91.7 (57)
6105	2	*	*	2	*	2	*	1	*
6301	939	91.8 (49)	1.8 (56)	1,046	23.0 (80)	943	12.5 (94)	723	91.7 (57)
6312	521	94.4 (21)	1.3 (41)	571	24.3 (73)	504	23.0 (61)	398	89.7 (77)
6316	324	92.0 (46)	1.9 (60)	345	21.4 (92)	292	27.7 (37)	237	93.7 (36)
6320	159	95.6 (14)	3.1 (82)	183	22.4 (85)	131	31.3 (24)	138	94.2 (32)
6325	92	*	*	102	26.5 (64)	99	*	71	*
6335	267	96.6 (8)	1.1 (31)	293	34.5 (22)	150	30.7 (26)	220	92.7 (47)
6345	35	*	*	38	*	30	*	25	*
6360	326	94.2 (22)	0.9 (23)	378	28.6 (56)	313	26.8 (42)	290	93.1 (43)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 37

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7306	655	89.8 (66)	1.7 (52)	720	31.4 (39)	689	28.7 (34)	549	92.0 (56)
7401	443	86.5 (85)	4.7 (98)	582	44.2 (2)	561	26.6 (46)	474	87.1 (87)
7601	159	96.9 (5)	0.6 (12)	168	23.8 (75)	160	48.1 (2)	121	95.9 (16)
7850	296	97.0 (3)	0.7 (15)	326	21.8 (90)	84	*	210	87.1 (87)
7852	3	*	*	3	*	0	*	2	*
8101	531	82.1 (101)	2.6 (76)	561	20.5 (97)	499	20.6 (70)	380	90.3 (76)
8102	175	89.1 (73)	1.1 (31)	191	17.3 (108)	167	18.0 (79)	136	85.3 (94)
8105	0	*	*	1	*	0	*	1	*
8106	178	83.7 (95)	1.1 (31)	190	11.1 (113)	177	20.9 (68)	110	96.4 (11)
8150	4	*	*	5	*	5	*	4	*
8151	929	93.4 (31)	2.4 (71)	1,046	33.7 (28)	1,024	15.1 (86)	792	94.9 (25)
8152	1	*	*	2	*	1	*	1	*
8155	3	*	*	3	*	3	*	3	*
8201	805	94.9 (18)	0.7 (15)	906	22.0 (88)	634	39.4 (11)	616	91.7 (57)
8202	1,177	97.4 (2)	0.5 (10)	1,218	29.9 (48)	851	8.2 (103)	934	92.9 (46)
8203	110	93.6 (29)	0.9 (23)	129	22.5 (84)	96	*	99	*
8250	136	91.9 (47)	0.7 (15)	157	22.3 (87)	126	32.5 (22)	110	93.6 (38)
8252	45	*	*	55	*	51	*	41	*
8255	593	93.9 (26)	0.5 (10)	673	18.1 (105)	495	37.4 (14)	441	90.5 (69)
8257	540	97.0 (3)	0.0 (1)	558	26.9 (62)	371	5.4 (105)	434	78.1 (98)
8263	125	96.0 (12)	2.4 (71)	154	26.6 (63)	116	44.8 (3)	140	97.1 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 38

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8264	212	94.8 (19)	0.9 (23)	231	15.2 (109)	173	39.9 (9)	141	95.7 (17)
8267	210	89.5 (71)	2.4 (71)	233	14.6 (111)	179	35.2 (17)	156	93.6 (38)
8901	259	86.1 (89)	2.3 (68)	273	17.9 (106)	158	27.2 (40)	164	80.5 (97)
8902	387	91.2 (53)	0.3 (8)	416	20.7 (95)	282	31.2 (25)	294	94.6 (29)
8903	424	82.8 (100)	0.9 (23)	458	19.7 (99)	396	18.4 (77)	321	88.5 (82)
8904	611	84.0 (94)	1.8 (56)	657	19.3 (100)	531	21.5 (66)	430	91.4 (62)
8905	671	91.2 (53)	1.6 (46)	737	22.4 (85)	625	36.2 (15)	465	90.5 (69)
8909	374	87.4 (81)	1.1 (31)	414	21.7 (91)	365	18.4 (77)	301	90.7 (67)
9045	422	91.0 (56)	1.4 (44)	449	23.8 (75)	233	24.0 (58)	329	93.6 (38)
9050	245	86.5 (85)	0.8 (19)	293	29.0 (54)	262	27.5 (39)	214	86.0 (92)
9051	437	89.7 (69)	0.2 (6)	459	26.1 (67)	429	5.8 (104)	349	96.6 (8)
9053	439	87.9 (79)	0.9 (23)	468	12.2 (112)	389	15.9 (85)	264	94.7 (28)
9058	430	85.8 (90)	2.1 (64)	459	15.0 (110)	415	8.9 (102)	240	88.3 (85)
9066	292	82.9 (99)	3.1 (82)	319	26.0 (69)	289	12.8 (91)	274	92.3 (52)
9068	243	83.5 (97)	2.1 (64)	272	19.9 (98)	252	14.7 (87)	184	86.4 (91)
9069	567	83.6 (96)	1.2 (37)	591	19.0 (101)	366	11.5 (96)	354	91.2 (64)
9070	513	83.0 (98)	2.1 (64)	554	17.5 (107)	461	21.3 (67)	343	91.3 (63)
9072	46	*	*	59	*	59	*	59	*
9080	733	87.4 (81)	1.6 (46)	796	20.6 (96)	562	20.8 (69)	531	90.4 (73)
9082	44	*	*	46	*	41	*	28	*
9083	4	*	*	3	*	3	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2021 Pregnancy Nutrition Surveillance (1)

Page : 39

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9087	7	*	*	8	*	8	*	8	*
Michigan	42,342	90.6	2.0	47,286	27.0	40,570	22.9	33,883	33883.0
National (Prior Year)	77,410	69.2	7.1	83,553	19.4	60,693	22.5	61,433	89.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 40

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	21	*	32	*	4	*	20	*	32	*
0102	95	*	122	27.9 (111)	34	*	93	*	123	3.3 (97)
0103	91	*	112	20.5 (99)	24	*	91	*	113	6.2 (111)
0104	34	*	39	*	9	*	32	*	39	*
0105	3	*	3	*	1	*	3	*	3	*
0250	31	*	37	*	11	*	30	*	35	*
0251	26	*	27	*	8	*	26	*	27	*
0252	21	*	25	*	4	*	18	*	25	*
0253	35	*	32	*	15	*	35	*	33	*
0401	110	41.8 (99)	134	23.1 (104)	38	*	106	11.3 (89)	134	1.5 (45)
0402	102	38.2 (95)	114	20.2 (98)	35	*	100	6.0 (51)	114	1.8 (57)
0403	34	*	38	*	12	*	33	*	39	*
0404	31	*	35	*	7	*	30	*	36	*
0501	105	37.1 (93)	125	24.8 (108)	33	*	101	18.8 (99)	125	3.2 (93)
0503	16	*	21	*	3	*	16	*	21	*
0504	94	*	112	12.5 (69)	20	*	89	*	111	1.8 (57)
0505	54	*	62	*	10	*	52	*	62	*
0506	39	*	56	*	13	*	34	*	56	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic

Page : 41

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0507	24	*	34	*	4	*	22	*	34	*
0508	1	*	5	*	0	*	1	*	5	*
0650	270	23.0 (59)	335	13.4 (75)	55	*	259	6.6 (57)	334	0.6 (12)
0651	155	36.8 (92)	169	26.0 (109)	47	*	148	10.8 (85)	168	2.4 (76)
0652	97	*	113	28.3 (112)	39	*	96	*	113	8.8 (112)
0653	97	*	122	21.3 (100)	25	*	93	*	123	4.9 (109)
0654	110	28.2 (73)	126	18.3 (93)	26	*	104	15.4 (98)	127	4.7 (108)
0655	86	*	100	23.0 (103)	28	*	86	*	100	3.0 (89)
0657	17	*	19	*	2	*	17	*	19	*
0701	142	30.3 (81)	165	17.6 (90)	38	*	138	6.5 (55)	165	2.4 (76)
0702	11	*	16	*	4	*	9	*	16	*
0703	29	*	44	*	9	*	29	*	44	*
0704	9	*	11	*	2	*	9	*	11	*
0705	10	*	14	*	3	*	9	*	14	*
0850	160	38.1 (94)	202	16.3 (85)	46	*	146	4.8 (40)	202	3.0 (89)
0851	130	40.8 (96)	176	15.3 (84)	48	*	121	5.8 (46)	178	2.2 (67)
0901	393	34.1 (87)	470	17.9 (91)	121	36.4 (7)	377	8.5 (72)	475	2.7 (82)
0902	8	*	9	*	3	*	8	*	9	*
1051	306	35.9 (91)	361	11.9 (67)	96	*	278	4.3 (37)	367	1.9 (60)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 42

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)									
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)
1052	1	*	1	*	0	*	1	*	
1101	358	20.9 (51)	450	8.9 (53)	68	*	350	8.9 (76)	452 4.6 (106)
1102	184	32.1 (82)	227	16.3 (85)	48	*	172	12.2 (92)	227 0.9 (22)
1106	10	*	12	*	2	*	9	*	12 *
1250	243	21.4 (52)	295	13.6 (76)	39	*	227	10.1 (80)	297 1.3 (37)
1251	151	33.8 (86)	185	23.2 (105)	45	*	148	8.8 (74)	185 2.2 (67)
1252	123	22.0 (57)	144	11.8 (66)	22	*	120	7.5 (63)	145 4.1 (101)
1253	167	29.9 (79)	188	13.8 (77)	43	*	163	6.7 (58)	185 2.2 (67)
1350	617	25.9 (65)	761	9.9 (56)	145	35.2 (8)	589	2.7 (15)	783 0.9 (22)
1351	1	*	1	*	0	*	1	*	1 *
1352	89	*	116	4.3 (22)	14	*	88	*	117 0.0 (1)
1401	57	*	65	*	22	*	56	*	66 *
1402	26	*	32	*	4	*	25	*	32 *
1403	3	*	4	*	1	*	3	*	4 *
1501	188	23.4 (60)	209	6.2 (37)	41	*	183	1.6 (9)	218 1.4 (41)
1503	205	17.6 (44)	244	8.2 (50)	35	*	198	4.0 (32)	244 0.4 (6)
1504	442	15.6 (36)	563	6.6 (39)	60	*	420	1.7 (10)	564 0.9 (22)
1505	22	*	19	*	6	*	22	*	19 *
1506	147	18.4 (47)	168	8.9 (53)	25	*	143	4.2 (35)	172 1.7 (53)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 43

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1507	134	35.1 (89)	173	10.4 (57)	40	*	123	8.1 (69)	174	2.9 (87)
1509	111	21.6 (56)	126	14.3 (79)	23	*	109	4.6 (39)	127	2.4 (76)
1511	157	29.3 (76)	191	12.6 (70)	44	*	154	4.5 (38)	196	2.0 (62)
1512	71	*	103	7.8 (46)	11	*	63	*	103	1.9 (60)
1514	135	9.6 (15)	201	6.0 (36)	12	*	111	0.0 (1)	198	1.5 (45)
1701	145	27.6 (71)	164	19.5 (97)	36	*	141	7.8 (65)	165	3.0 (89)
1950	151	41.7 (98)	182	22.5 (102)	59	*	151	14.6 (97)	188	1.6 (51)
1952	32	*	42	*	8	*	30	*	42	*
1953	66	*	78	*	19	*	64	*	79	*
1954	92	*	112	18.8 (95)	32	*	90	*	112	0.9 (22)
1955	26	*	32	*	11	*	24	*	33	*
1957	90	*	125	15.2 (83)	28	*	88	*	126	2.4 (76)
1963	16	*	21	*	3	*	16	*	22	*
2150	156	35.3 (90)	183	22.4 (101)	46	*	145	10.3 (81)	183	1.6 (51)
2151	62	*	66	*	14	*	57	*	66	*
2152	12	*	12	*	8	*	12	*	12	*
2251	112	28.6 (74)	130	13.8 (77)	29	*	108	2.8 (16)	130	0.8 (19)
2252	36	*	43	*	23	*	37	*	43	*
2554	18	*	18	*	7	*	17	*	17	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 44

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2555	1,756	21.5 (54)	2,166	11.2 (62)	319	41.1 (5)	1,670	8.5 (72)	2,184	2.3 (73)
2556	52	*	57	*	15	*	52	*	58	*
2560	1	*	1	*	0	*	1	*	1	*
2562	2	*	2	*	0	*	2	*	2	*
2563	3	*	3	*	1	*	3	*	3	*
3001	318	30.2 (80)	367	11.2 (62)	94	*	312	11.2 (87)	369	4.6 (106)
3002	38	*	51	*	15	*	38	*	51	*
3004	74	*	89	*	17	*	68	*	89	*
3005	43	*	49	*	14	*	43	*	49	*
3007	2	*	4	*	2	*	3	*	4	*
3201	89	*	120	24.2 (107)	34	*	81	*	120	3.3 (97)
3301	1,102	15.8 (37)	1,398	7.4 (43)	157	47.1 (4)	1,038	3.3 (23)	1,399	2.2 (67)
3304	18	*	27	*	2	*	17	*	27	*
3305	25	*	39	*	3	*	23	*	39	*
3306	45	*	63	*	9	*	45	*	63	*
3308	21	*	26	*	1	*	16	*	26	*
3312	107	14.0 (29)	136	5.1 (31)	15	*	101	2.0 (13)	136	1.5 (45)
3550	19	*	25	*	10	*	19	*	25	*
3551	3	*	3	*	3	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 45

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3552	1	*	1	*	1	*	1	*		
3701	209	33.5 (85)	270	18.1 (92)	60	*	191	6.8 (60)	272	0.7 (17)
3850	751	29.4 (77)	903	18.3 (93)	204	30.4 (9)	719	6.0 (51)	902	2.1 (65)
3901	433	21.5 (54)	501	9.4 (55)	76	*	412	4.9 (41)	501	0.6 (12)
3902	6	*	6	*	4	*	7	*	7	*
3903	1	*	2	*	0	*	1	*	2	*
3950	477	21.4 (52)	615	11.7 (65)	96	*	458	4.1 (33)	617	1.8 (57)
4106	521	11.7 (21)	666	6.9 (41)	54	*	509	1.4 (7)	665	0.6 (12)
4107	681	15.1 (34)	869	7.8 (46)	84	*	627	5.1 (44)	868	1.4 (41)
4109	140	25.0 (63)	195	14.4 (80)	29	*	131	3.8 (26)	194	2.6 (80)
4112	177	1.7 (2)	187	2.1 (7)	3	*	138	9.4 (77)	172	2.3 (73)
4115	809	7.0 (8)	1,039	3.3 (12)	59	*	748	0.9 (5)	1,045	0.7 (17)
4351	127	34.6 (88)	145	24.1 (106)	37	*	123	3.3 (23)	146	2.1 (65)
4352	43	*	55	*	21	*	45	*	54	*
4353	109	19.3 (48)	147	12.9 (72)	21	*	108	6.5 (55)	147	2.0 (62)
4354	195	32.8 (84)	233	18.9 (96)	57	*	188	11.2 (87)	241	1.7 (53)
4355	69	*	82	*	18	*	64	*	83	*
4356	92	*	104	10.6 (59)	24	*	91	*	103	3.9 (100)
4359	171	17.5 (42)	190	11.6 (64)	28	*	168	4.2 (35)	190	0.5 (9)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 46

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4360	56	*	67	*	14	*	56	*	67	*
4361	69	*	87	*	19	*	66	*	87	*
4362	70	*	86	*	20	*	66	*	87	*
4363	165	26.1 (67)	207	16.4 (88)	38	*	155	10.3 (81)	207	1.4 (41)
4401	192	28.1 (72)	235	17.0 (89)	46	*	178	7.9 (67)	234	0.9 (22)
4501	181	32.6 (83)	245	16.3 (85)	58	*	171	14.0 (95)	245	2.9 (87)
4601	293	26.3 (70)	348	12.9 (72)	62	*	273	7.0 (62)	348	5.2 (110)
4701	250	26.0 (66)	307	11.1 (61)	57	*	236	3.8 (26)	307	1.0 (32)
5050	338	17.5 (42)	472	8.3 (51)	54	*	318	3.8 (26)	473	0.8 (19)
5051	608	12.8 (24)	810	5.7 (35)	65	*	561	3.9 (29)	807	1.5 (45)
5052	639	11.7 (21)	799	4.6 (27)	59	*	609	3.9 (29)	797	0.8 (19)
5055	358	3.1 (4)	422	0.9 (3)	9	*	344	4.1 (33)	422	0.2 (4)
5101	387	14.0 (29)	479	4.6 (27)	50	*	367	2.2 (14)	480	1.5 (45)
5102	3	*	2	*	0	*	3	*	2	*
5202	184	51.6 (100)	199	26.6 (110)	91	*	180	8.3 (71)	210	1.0 (32)
5206	3	*	4	*	1	*	2	*	4	*
5601	233	26.2 (68)	312	12.8 (71)	53	*	218	2.8 (16)	313	1.0 (32)
5801	430	25.8 (64)	555	13.0 (74)	97	*	397	8.8 (74)	552	4.2 (102)
5803	9	*	11	*	1	*	9	*	11	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 47

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
5804	8	*	8	*	2	*	8	*	8	*
5806	8	*	8	*	3	*	8	*	8	*
6101	15	*	18	*	7	*	13	*	18	*
6102	23	*	23	*	7	*	22	*	24	*
6103	831	28.6 (74)	1,024	14.4 (80)	211	39.8 (6)	797	6.8 (60)	1,026	1.1 (35)
6105	1	*	2	*	0	*	1	*	2	*
6301	777	9.7 (16)	975	4.3 (22)	69	*	724	3.0 (20)	977	1.4 (41)
6312	430	3.3 (5)	535	1.3 (4)	11	*	397	1.5 (8)	535	0.4 (6)
6316	255	8.6 (10)	331	4.2 (21)	20	*	237	1.7 (10)	329	0.6 (12)
6320	150	10.0 (17)	164	4.3 (22)	13	*	138	0.0 (1)	164	0.6 (12)
6325	77	*	93	*	15	*	71	*	93	*
6335	238	8.8 (11)	282	3.9 (20)	18	*	220	1.8 (12)	282	0.4 (6)
6345	27	*	36	*	6	*	25	*	36	*
6360	304	14.5 (31)	345	5.2 (32)	35	*	289	5.2 (45)	343	0.9 (22)
7306	561	29.6 (78)	663	8.1 (49)	159	53.5 (2)	548	10.9 (86)	694	2.6 (80)
7401	490	41.0 (97)	528	14.4 (80)	171	51.5 (3)	474	12.4 (93)	534	4.3 (103)
7601	127	13.4 (25)	159	10.7 (60)	15	*	121	5.8 (46)	159	1.3 (37)
7850	221	26.2 (68)	307	10.4 (57)	50	*	205	7.8 (65)	302	1.7 (53)
7852	2	*	3	*	1	*	2	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 48

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8101	401	15.0 (33)	551	3.8 (17)	54	*	380	11.6 (91)	551	2.2 (67)
8102	145	12.4 (23)	187	4.8 (30)	15	*	134	6.7 (58)	188	3.2 (93)
8105	1	*	0	*	0	*	1	*	0	*
8106	120	24.2 (62)	185	7.6 (44)	19	*	110	8.2 (70)	185	2.7 (82)
8150	4	*	5	*	2	*	4	*	5	*
8151	843	16.8 (39)	977	6.6 (39)	116	57.8 (1)	792	3.2 (21)	975	0.9 (22)
8152	1	*	2	*	0	*	1	*	2	*
8155	3	*	3	*	1	*	3	*	3	*
8201	673	13.5 (27)	815	6.4 (38)	76	*	615	2.8 (16)	817	1.3 (37)
8202	979	2.6 (3)	1,215	0.6 (2)	23	*	933	2.9 (19)	1,214	0.5 (9)
8203	102	23.5 (61)	113	8.0 (48)	17	*	99	*	114	0.0 (1)
8250	115	18.3 (46)	137	4.4 (25)	17	*	110	0.9 (5)	137	0.0 (1)
8252	47	*	49	*	5	*	41	*	48	*
8255	470	19.6 (49)	608	8.4 (52)	75	*	441	6.1 (53)	609	1.3 (37)
8257	467	0.2 (1)	551	0.5 (1)	2	*	438	0.5 (3)	550	0.9 (22)
8263	147	20.4 (50)	133	12.0 (68)	23	*	139	5.8 (46)	133	3.0 (89)
8264	151	17.2 (41)	216	3.7 (16)	21	*	141	3.5 (25)	215	0.9 (22)
8267	168	22.0 (57)	211	7.6 (44)	28	*	156	6.4 (54)	212	2.8 (84)
8901	178	16.9 (40)	266	5.6 (34)	27	*	166	9.6 (78)	265	3.4 (99)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 49

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8902	311	10.0 (17)	398	3.0 (11)	27	*	294	5.8 (46)	399	2.8 (84)
8903	328	15.2 (35)	436	5.3 (33)	43	*	319	11.3 (89)	436	2.8 (84)
8904	458	16.6 (38)	644	3.6 (15)	64	*	428	9.6 (78)	643	4.5 (104)
8905	493	9.1 (12)	718	3.3 (12)	43	*	460	3.9 (29)	718	1.7 (53)
8909	320	10.3 (19)	386	3.4 (14)	31	*	301	7.6 (64)	383	2.3 (73)
9045	345	5.5 (7)	439	1.6 (6)	16	*	328	4.9 (41)	437	0.5 (9)
9050	223	13.9 (28)	281	4.6 (27)	27	*	214	14.5 (96)	278	3.2 (93)
9051	360	5.3 (6)	443	1.4 (5)	19	*	350	0.6 (4)	443	0.2 (4)
9053	298	9.1 (12)	453	2.6 (8)	23	*	264	4.9 (41)	453	2.0 (62)
9058	258	9.3 (14)	449	3.8 (17)	21	*	239	5.9 (50)	448	0.9 (22)
9066	292	18.2 (45)	311	7.1 (42)	43	*	273	13.2 (94)	311	3.2 (93)
9068	190	14.7 (32)	259	2.7 (9)	22	*	184	10.3 (81)	259	1.2 (36)
9069	385	7.5 (9)	576	3.8 (17)	28	*	354	10.7 (84)	576	4.5 (104)
9070	365	13.4 (25)	530	2.8 (10)	43	*	341	7.9 (67)	527	1.5 (45)
9072	59	*	47	*	14	*	59	*	47	*
9080	546	10.8 (20)	777	4.5 (26)	54	*	530	3.2 (21)	778	2.2 (67)
9082	32	*	45	*	4	*	28	*	45	*
9083	3	*	4	*	0	*	2	*	4	*
9087	8	*	7	*	1	*	8	*	7	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2021 Pregnancy Nutrition Surveillance (1)

Page : 50

Michigan

Comparison of Smoking Indicators by Clinic

Michigan	35,714	19.7	44,742	9.2	6,188	42.4	33,791	6.1	44,858	1.9
National (Prior Year)	83,471	24.3	86,947	11.2	14,888	35.8	70,642	7.8	84,906	3.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 51

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	27	*	*	29	*	25	*	27	*
0102	110	11.8 (72)	11.8 (106)	117	13.7 (73)	95	*	109	72.5 (56)
0103	103	10.7 (60)	3.9 (12)	105	7.6 (5)	93	*	103	71.8 (59)
0104	34	*	*	38	*	30	*	34	*
0105	3	*	*	3	*	3	*	3	*
0250	34	*	*	34	*	29	*	34	*
0251	25	*	*	27	*	25	*	25	*
0252	20	*	*	24	*	18	*	20	*
0253	29	*	*	32	*	23	*	29	*
0401	127	9.4 (41)	9.4 (92)	131	13.0 (59)	107	3.7 (33)	126	66.7 (76)
0402	109	13.8 (89)	4.6 (21)	108	12.0 (43)	93	*	109	82.6 (12)
0403	37	*	*	39	*	34	*	37	*
0404	34	*	*	34	*	28	*	34	*
0501	106	4.7 (3)	11.3 (104)	117	9.4 (13)	93	*	116	81.0 (16)
0503	20	*	*	21	*	19	*	20	*
0504	104	10.6 (57)	7.7 (76)	108	9.3 (11)	94	*	104	90.4 (1)
0505	50	*	*	61	*	44	*	59	*
0506	48	*	*	53	*	40	*	50	*
0507	27	*	*	32	*	23	*	31	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 52

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	4	*	*	5	*	3	*	5	*
0650	315	10.8 (62)	8.9 (89)	330	9.4 (13)	287	5.9 (70)	315	74.9 (49)
0651	152	10.5 (54)	6.6 (56)	168	13.7 (73)	132	6.8 (83)	152	66.4 (77)
0652	104	9.6 (45)	5.8 (39)	109	10.1 (18)	91	*	104	67.3 (73)
0653	111	9.0 (37)	6.3 (48)	113	14.2 (80)	95	*	111	73.0 (55)
0654	118	11.9 (74)	6.8 (59)	120	13.3 (64)	99	*	118	74.6 (50)
0655	91	*	*	94	*	80	*	91	*
0657	17	*	*	19	*	14	*	17	*
0701	153	5.9 (12)	8.5 (87)	164	6.7 (4)	142	3.5 (30)	152	89.5 (2)
0702	16	*	*	16	*	16	*	16	*
0703	40	*	*	43	*	37	*	42	*
0704	10	*	*	10	*	7	*	10	*
0705	14	*	*	14	*	13	*	14	*
0850	187	12.8 (81)	8.0 (81)	197	13.2 (63)	161	5.0 (54)	189	85.7 (6)
0851	161	8.7 (33)	8.1 (83)	173	12.7 (54)	141	2.8 (18)	163	87.1 (5)
0901	435	9.7 (48)	8.7 (88)	463	10.2 (21)	393	4.1 (38)	436	69.0 (67)
0902	10	*	*	10	*	9	*	10	*
1051	337	8.6 (32)	13.6 (110)	353	10.5 (26)	300	4.0 (37)	339	89.1 (4)
1052	1	*	*	1	*	0	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 53

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1101	425	12.0 (78)	5.4 (33)	435	12.2 (45)	367	5.4 (61)	426	65.5 (80)
1102	211	7.6 (21)	7.6 (75)	224	8.0 (7)	192	4.7 (50)	212	75.5 (46)
1106	10	*	*	12	*	8	*	10	*
1250	284	9.5 (43)	8.1 (83)	291	10.7 (28)	249	6.8 (83)	285	77.5 (33)
1251	173	10.4 (53)	6.4 (52)	182	15.9 (93)	145	4.1 (38)	173	77.5 (33)
1252	135	5.9 (12)	11.9 (107)	140	11.4 (37)	120	0.8 (3)	135	77.8 (29)
1253	169	4.1 (2)	12.4 (108)	180	10.0 (17)	151	2.6 (15)	169	84.6 (7)
1350	731	9.8 (49)	6.6 (56)	758	10.4 (23)	648	5.4 (61)	733	75.4 (47)
1351	1	*	*	1	*	1	*	1	*
1352	111	5.4 (7)	7.2 (67)	115	6.1 (2)	104	2.9 (20)	112	71.4 (61)
1401	61	*	*	61	*	55	*	61	*
1402	28	*	*	31	*	27	*	28	*
1403	4	*	*	4	*	3	*	4	*
1501	198	9.6 (45)	6.1 (44)	214	11.7 (41)	175	2.9 (20)	207	75.8 (44)
1503	228	8.3 (28)	10.1 (97)	237	10.5 (26)	205	3.9 (35)	228	76.3 (38)
1504	464	10.1 (50)	9.3 (91)	550	12.9 (56)	400	4.5 (46)	539	78.8 (25)
1505	19	*	*	19	*	17	*	19	*
1506	156	7.1 (17)	8.3 (85)	166	13.3 (64)	135	0.7 (2)	160	70.0 (64)
1507	149	8.1 (27)	13.4 (109)	168	10.7 (28)	131	2.3 (14)	159	78.6 (26)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 54

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1509	114	5.3 (6)	7.9 (79)	120	12.5 (50)	99	*	113	67.3 (73)
1511	183	8.7 (33)	7.7 (76)	191	15.7 (91)	154	1.3 (4)	188	80.3 (19)
1512	98	*	*	102	16.7 (97)	81	*	98	*
1514	175	12.6 (80)	6.3 (48)	197	18.3 (106)	145	2.1 (10)	182	84.1 (9)
1701	147	5.4 (7)	10.2 (99)	159	6.3 (3)	136	2.2 (11)	146	78.1 (28)
1950	178	15.2 (97)	7.9 (79)	182	15.4 (88)	147	3.4 (27)	178	80.3 (19)
1952	39	*	*	42	*	34	*	39	*
1953	73	*	*	79	*	58	*	76	*
1954	103	7.8 (24)	4.9 (24)	108	9.3 (11)	93	*	103	81.6 (15)
1955	32	*	*	33	*	29	*	32	*
1957	120	19.2 (110)	7.5 (72)	120	20.8 (110)	93	*	121	89.3 (3)
1963	18	*	*	18	*	15	*	18	*
2150	168	4.8 (4)	10.1 (97)	180	8.3 (8)	152	1.3 (4)	168	76.2 (39)
2151	65	*	*	65	*	60	*	65	*
2152	11	*	*	12	*	8	*	11	*
2251	126	5.6 (10)	7.1 (64)	129	11.6 (40)	112	2.7 (16)	126	77.0 (37)
2252	40	*	*	42	*	38	*	40	*
2554	18	*	*	18	*	15	*	18	*
2555	2,064	11.7 (70)	4.9 (24)	2,160	13.8 (77)	1,775	5.5 (64)	2,066	60.1 (94)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 55

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2556	56	*	*	58	*	45	*	57	*
2560	1	*	*	1	*	1	*	1	*
2562	2	*	*	2	*	2	*	2	*
2563	3	*	*	3	*	3	*	3	*
3001	343	18.1 (108)	4.1 (14)	354	17.5 (104)	283	6.7 (82)	342	54.1 (100)
3002	47	*	*	48	*	38	*	47	*
3004	76	*	*	80	*	58	*	75	*
3005	47	*	*	47	*	45	*	48	*
3007	4	*	*	4	*	2	*	4	*
3201	111	2.7 (1)	9.9 (96)	118	5.9 (1)	105	0.0 (1)	110	77.3 (35)
3301	1,283	11.0 (65)	5.6 (36)	1,355	13.7 (73)	1,109	4.7 (50)	1,295	77.8 (29)
3304	26	*	*	27	*	22	*	26	*
3305	37	*	*	38	*	33	*	37	*
3306	62	*	*	63	*	58	*	61	*
3308	23	*	*	26	*	21	*	23	*
3312	129	9.3 (40)	6.2 (46)	133	12.8 (55)	111	1.8 (7)	130	73.8 (53)
3550	22	*	*	22	*	18	*	22	*
3551	2	*	*	3	*	2	*	3	*
3552	1	*	*	1	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 56

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3701	255	7.1 (17)	10.6 (101)	266	10.9 (31)	223	2.2 (11)	256	84.4 (8)
3850	839	10.6 (57)	7.0 (63)	873	11.9 (42)	740	4.7 (50)	841	71.6 (60)
3901	456	11.8 (72)	3.7 (10)	482	12.0 (43)	393	5.1 (58)	461	73.1 (54)
3902	4	*	*	7	*	4	*	4	*
3903	2	*	*	2	*	2	*	2	*
3950	552	11.1 (66)	9.8 (95)	595	13.1 (62)	476	5.0 (54)	553	75.8 (44)
4106	607	13.0 (83)	5.3 (32)	649	13.7 (73)	518	6.8 (83)	614	76.2 (39)
4107	803	13.1 (84)	6.5 (53)	856	15.1 (85)	688	5.8 (68)	812	76.2 (39)
4109	185	11.4 (69)	6.5 (53)	189	19.6 (109)	148	4.1 (38)	184	80.4 (18)
4112	179	5.0 (5)	3.4 (7)	176	10.8 (30)	155	3.2 (25)	179	26.8 (110)
4115	973	13.4 (87)	6.3 (48)	1,031	15.4 (88)	823	5.0 (54)	988	77.7 (31)
4351	131	9.2 (39)	10.7 (102)	142	13.4 (67)	115	6.1 (73)	132	81.8 (13)
4352	50	*	*	51	*	37	*	50	*
4353	138	8.7 (33)	9.4 (92)	136	11.0 (33)	119	4.2 (42)	140	80.0 (21)
4354	228	10.5 (54)	10.5 (100)	238	10.1 (18)	204	4.4 (45)	228	78.9 (23)
4355	67	*	*	76	*	55	*	67	*
4356	101	6.9 (16)	6.9 (62)	103	12.6 (51)	88	*	101	68.3 (70)
4359	175	12.0 (78)	4.6 (21)	176	13.6 (72)	147	5.4 (61)	175	76.0 (43)
4360	57	*	*	66	*	53	*	57	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 57

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4361	67	*	*	80	*	56	*	67	*
4362	81	*	*	84	*	70	*	81	*
4363	194	10.3 (52)	11.3 (104)	202	14.4 (83)	165	1.8 (7)	194	84.0 (11)
4401	224	6.3 (14)	4.9 (24)	228	9.6 (15)	199	4.5 (46)	224	78.6 (26)
4501	223	5.4 (7)	9.0 (90)	238	14.3 (81)	193	1.6 (6)	223	77.6 (32)
4601	327	9.5 (43)	6.1 (44)	346	11.0 (33)	290	4.1 (38)	327	74.0 (52)
4701	291	11.7 (70)	7.2 (67)	302	12.3 (46)	255	5.5 (64)	290	84.1 (9)
5050	460	10.2 (51)	4.3 (18)	465	18.3 (106)	373	3.5 (30)	458	62.0 (86)
5051	765	10.8 (62)	7.1 (64)	786	13.9 (78)	652	4.6 (48)	767	66.2 (78)
5052	749	13.4 (87)	4.1 (14)	775	14.3 (81)	634	6.2 (75)	750	64.5 (82)
5055	381	10.5 (54)	6.0 (43)	396	10.1 (18)	343	5.2 (59)	387	62.3 (85)
5101	465	8.8 (36)	5.2 (31)	470	13.0 (59)	403	2.7 (16)	467	74.5 (51)
5102	2	*	*	2	*	1	*	2	*
5202	200	8.0 (26)	11.0 (103)	204	8.3 (8)	181	3.9 (35)	201	75.1 (48)
5206	4	*	*	3	*	3	*	4	*
5601	298	7.7 (23)	9.4 (92)	307	10.4 (23)	268	2.2 (11)	298	77.2 (36)
5801	519	9.6 (45)	7.5 (72)	539	10.4 (23)	464	4.7 (50)	520	68.8 (68)
5803	11	*	*	11	*	10	*	11	*
5804	8	*	*	8	*	8	*	8	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 58

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5806	8	*	*	8	*	6	*	8	*
6101	15	*	*	17	*	13	*	16	*
6102	24	*	*	24	*	19	*	24	*
6103	959	10.6 (57)	7.4 (71)	988	12.3 (46)	829	4.2 (42)	962	61.5 (88)
6105	2	*	*	2	*	1	*	2	*
6301	893	13.8 (89)	6.8 (59)	949	12.6 (51)	771	6.5 (79)	906	69.9 (65)
6312	489	11.9 (74)	5.9 (41)	517	15.1 (85)	413	3.1 (24)	497	78.9 (23)
6316	299	8.4 (30)	5.0 (29)	318	12.3 (46)	264	3.4 (27)	303	80.9 (17)
6320	155	8.4 (30)	7.1 (64)	162	11.1 (35)	138	2.9 (20)	155	79.4 (22)
6325	90	*	*	91	*	78	*	89	*
6335	265	11.3 (68)	5.7 (38)	275	12.4 (49)	230	5.2 (59)	269	76.2 (39)
6345	36	*	*	36	*	31	*	36	*
6360	325	15.1 (96)	4.9 (24)	341	13.5 (70)	275	8.0 (87)	328	71.0 (63)
7306	660	13.3 (86)	6.2 (46)	678	13.4 (67)	570	5.6 (66)	656	68.8 (68)
7401	491	9.4 (41)	7.5 (72)	511	12.9 (56)	426	3.8 (34)	492	65.2 (81)
7601	144	9.0 (37)	4.9 (24)	156	10.9 (31)	124	5.6 (66)	149	58.4 (96)
7850	291	7.6 (21)	6.5 (53)	304	9.9 (16)	261	3.4 (27)	292	67.1 (75)
7852	2	*	*	2	*	2	*	2	*
8101	501	16.6 (99)	2.2 (3)	519	21.2 (111)	388	9.3 (91)	499	47.1 (106)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 59

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8102	172	6.4 (15)	5.8 (39)	182	11.5 (38)	150	3.3 (26)	173	60.7 (90)
8105	0	*	*	0	*	0	*	0	*
8106	173	13.9 (91)	3.5 (8)	181	16.6 (96)	143	6.3 (76)	175	53.7 (101)
8150	5	*	*	5	*	5	*	5	*
8151	895	12.8 (81)	6.3 (48)	945	13.5 (70)	771	6.1 (73)	903	81.8 (13)
8152	2	*	*	2	*	2	*	2	*
8155	2	*	*	3	*	2	*	2	*
8201	764	10.7 (60)	7.3 (69)	808	13.0 (59)	665	5.0 (54)	767	68.3 (70)
8202	1,097	7.2 (19)	5.0 (29)	1,173	9.0 (10)	993	3.5 (30)	1,148	61.7 (87)
8203	108	7.4 (20)	8.3 (85)	111	12.6 (51)	94	*	108	72.2 (57)
8250	126	7.9 (25)	5.6 (36)	133	17.3 (102)	105	1.9 (9)	126	60.3 (92)
8252	42	*	*	44	*	37	*	43	*
8255	571	11.9 (74)	6.8 (59)	596	11.2 (36)	500	6.0 (72)	575	65.9 (79)
8257	498	16.1 (98)	3.0 (6)	531	10.2 (21)	443	12.0 (97)	511	71.4 (61)
8263	125	11.2 (67)	8.0 (81)	131	16.8 (98)	105	2.9 (20)	124	62.9 (83)
8264	201	11.9 (74)	5.5 (35)	208	11.5 (38)	175	4.6 (48)	201	72.1 (58)
8267	194	5.7 (11)	6.7 (58)	208	7.7 (6)	177	2.8 (18)	196	60.2 (93)
8901	245	18.0 (107)	4.1 (14)	255	16.9 (100)	201	10.0 (95)	244	52.9 (103)
8902	369	17.3 (105)	3.5 (8)	388	17.5 (104)	301	8.6 (88)	374	67.9 (72)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 60

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8903	397	13.9 (91)	2.5 (4)	418	17.2 (101)	324	5.9 (70)	397	53.1 (102)
8904	594	17.0 (102)	2.0 (1)	612	17.3 (102)	478	6.5 (79)	600	57.3 (99)
8905	658	13.1 (84)	5.9 (41)	693	15.7 (91)	547	6.4 (77)	671	69.6 (66)
8909	360	17.2 (104)	4.7 (23)	374	16.8 (98)	300	9.7 (93)	361	59.0 (95)
9045	418	10.8 (62)	7.7 (76)	433	12.9 (56)	364	5.8 (68)	423	41.4 (108)
9050	259	14.3 (93)	5.4 (33)	271	14.0 (79)	220	6.4 (77)	259	60.6 (91)
9051	422	8.3 (28)	7.3 (69)	427	13.3 (64)	361	4.2 (42)	427	60.9 (89)
9053	415	16.6 (99)	4.1 (14)	442	15.4 (88)	348	8.6 (88)	415	62.7 (84)
9058	406	14.3 (93)	2.5 (4)	440	13.4 (67)	346	7.5 (86)	425	58.1 (97)
9066	285	17.5 (106)	3.9 (12)	304	16.1 (94)	244	9.8 (94)	286	50.3 (105)
9068	244	18.9 (109)	2.0 (1)	255	16.5 (95)	203	10.8 (96)	244	45.1 (107)
9069	535	17.0 (102)	4.5 (20)	563	18.5 (108)	436	9.4 (92)	532	37.4 (109)
9070	496	16.9 (101)	4.4 (19)	520	15.2 (87)	419	8.6 (88)	495	57.8 (98)
9072	43	*	*	47	*	36	*	43	*
9080	734	15.0 (95)	3.8 (11)	763	14.4 (83)	621	6.6 (81)	750	52.1 (104)
9082	43	*	*	45	*	42	*	43	*
9083	4	*	*	3	*	3	*	4	*
9087	6	*	*	7	*	4	*	6	*
Michigan	41,563	11.5	6.3	43,647	13.2	35,837	5.2	41,966	69.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2021 Pregnancy Nutrition Surveillance (1)

Page : 61

Michigan

Comparison of Infant Health Indicators by Clinic

National (Prior Year)	80,061	10.1	6.0	77,312	11.9	65,997	4.4	80,887	57.7
------------------------------	---------------	-------------	------------	---------------	-------------	---------------	------------	---------------	-------------

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 62

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	109	91.7	0.9	1.8	0.0	0.0	4.6	0.9
0102	433	89.4	1.2	2.3	0.5	0.2	6.5	0.0
0103	392	91.1	0.3	4.3	0.3	0.0	3.8	0.3
0104	122	95.1	0.0	3.3	0.0	0.0	1.6	0.0
0105	16	*	*	*	*	*	*	*
0250	125	83.2	0.8	2.4	2.4	0.0	11.2	0.0
0251	92	*	*	*	*	*	*	*
0252	98	*	*	*	*	*	*	*
0253	125	73.6	0.0	0.0	9.6	0.0	16.8	0.0
0401	456	94.1	0.7	1.5	0.4	0.2	2.9	0.2
0402	383	88.3	0.8	2.9	0.3	0.0	7.8	0.0
0403	149	96.0	0.7	1.3	0.7	0.0	0.7	0.7
0404	101	93.1	1.0	0.0	1.0	0.0	5.0	0.0
0501	466	95.9	0.0	1.5	0.0	0.2	2.4	0.0
0503	90	*	*	*	*	*	*	*
0504	410	86.8	1.5	2.7	1.2	0.5	7.3	0.0
0505	248	87.5	0.0	4.8	0.4	0.0	7.3	0.0
0506	163	92.6	0.0	1.2	0.6	0.6	4.9	0.0
0507	79	*	*	*	*	*	*	*
0508	25	*	*	*	*	*	*	*
0509	1	*	*	*	*	*	*	*
0650	1,058	78.5	3.6	6.4	3.7	0.9	6.7	0.2
0651	578	94.3	0.3	2.4	0.3	0.5	2.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 63

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0652	323	94.4	0.0	1.9	0.0	0.0	3.7	0.0
0653	400	95.0	0.0	2.5	0.0	0.5	2.0	0.0
0654	386	90.2	1.6	3.4	0.0	0.0	4.9	0.0
0655	324	93.8	0.0	1.2	0.0	0.3	3.7	0.9
0657	65	*	*	*	*	*	*	*
0701	463	89.6	1.5	2.2	0.9	2.6	2.8	0.4
0702	69	*	*	*	*	*	*	*
0703	165	84.8	2.4	1.8	0.0	1.2	7.3	2.4
0704	55	*	*	*	*	*	*	*
0705	87	*	*	*	*	*	*	*
0706	10	*	*	*	*	*	*	*
0850	720	84.4	4.3	8.6	0.0	1.5	0.8	0.3
0851	630	92.5	0.2	5.1	0.2	0.5	1.1	0.5
0901	1,524	79.5	2.0	11.7	0.0	0.2	6.3	0.4
0902	41	*	*	*	*	*	*	*
1051	1,175	84.8	0.9	6.2	2.2	0.9	4.5	0.6
1052	1	*	*	*	*	*	*	*
1101	1,674	34.5	51.1	8.4	0.2	1.3	3.9	0.6
1102	837	74.8	11.0	7.6	0.6	0.7	4.7	0.6
1104	1	*	*	*	*	*	*	*
1106	74	*	*	*	*	*	*	*
1250	1,015	85.7	0.7	10.3	0.0	0.4	1.7	1.2
1251	674	93.8	0.3	2.7	0.1	0.1	1.9	1.0
1252	535	61.1	0.2	33.5	0.0	0.2	4.5	0.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 64

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1253	735	78.1	1.2	10.2	0.0	0.1	9.9	0.4
1350	2,352	58.0	20.8	7.9	0.4	5.9	6.5	0.3
1351	80	*	*	*	*	*	*	*
1352	391	71.4	18.2	4.3	0.0	0.0	5.1	1.0
1401	226	89.8	0.9	3.5	1.3	0.0	3.5	0.9
1402	113	57.5	0.0	22.1	9.7	0.0	10.6	0.0
1403	19	*	*	*	*	*	*	*
1501	785	51.8	4.1	38.6	0.1	0.3	3.4	1.7
1503	894	60.6	7.5	21.8	0.3	0.2	7.9	1.6
1504	1,998	38.1	5.8	46.1	0.2	4.2	4.9	0.8
1505	285	80.7	4.6	10.5	0.0	0.4	2.8	1.1
1506	583	79.4	1.5	15.6	0.0	0.9	2.1	0.5
1507	556	82.4	2.0	8.8	0.2	1.4	4.7	0.5
1508	22	*	*	*	*	*	*	*
1509	419	49.2	0.7	43.2	0.0	0.5	3.6	2.9
1511	621	88.2	2.6	5.8	0.3	0.3	2.3	0.5
1512	345	85.5	0.6	6.1	0.3	0.9	6.7	0.0
1514	433	76.7	3.9	12.5	0.5	1.4	4.6	0.5
1701	594	53.9	0.7	3.0	21.0	0.8	20.0	0.3
1950	612	90.7	0.5	6.2	0.0	0.2	2.1	0.3
1952	130	93.8	0.8	3.1	0.0	0.0	1.5	0.8
1953	284	91.5	0.0	7.0	0.0	0.0	1.4	0.0
1954	446	85.7	0.2	11.9	0.2	0.0	1.6	0.4
1955	107	82.2	0.9	12.1	0.0	0.0	2.8	1.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 65

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1957	394	82.5	1.3	10.9	0.0	0.0	4.8	0.5
1963	63	*	*	*	*	*	*	*
1964	17	*	*	*	*	*	*	*
2150	631	84.0	0.5	2.5	1.1	0.5	11.3	0.2
2151	223	77.6	1.3	11.7	0.0	0.0	7.2	2.2
2152	17	*	*	*	*	*	*	*
2251	403	92.6	0.2	2.5	0.0	0.0	3.2	1.5
2252	171	88.9	0.0	1.8	0.6	0.0	5.8	2.9
2554	48	*	*	*	*	*	*	*
2555	7,386	48.4	38.4	5.8	0.0	0.3	6.9	0.3
2556	287	25.8	62.0	4.5	0.3	0.0	7.3	0.0
2557	2	*	*	*	*	*	*	*
2558	4	*	*	*	*	*	*	*
2560	5	*	*	*	*	*	*	*
2561	3	*	*	*	*	*	*	*
2562	16	*	*	*	*	*	*	*
2563	26	*	*	*	*	*	*	*
2564	22	*	*	*	*	*	*	*
3001	1,156	21.0	60.5	13.0	0.1	0.7	4.6	0.2
3002	262	22.5	50.8	21.0	0.0	0.0	5.7	0.0
3004	250	61.2	0.4	36.0	0.0	0.4	2.0	0.0
3005	116	88.8	0.0	6.0	0.0	1.7	2.6	0.9
3007	13	*	*	*	*	*	*	*
3201	419	93.1	0.2	5.0	0.0	0.2	1.2	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 66

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3203	1	*	*	*	*	*	*	*
3301	4,320	38.0	31.7	15.6	0.1	5.5	8.8	0.3
3304	97	*	*	*	*	*	*	*
3305	156	60.9	11.5	12.8	0.0	7.1	7.7	0.0
3306	242	86.4	2.9	6.2	0.4	1.2	2.5	0.4
3308	98	*	*	*	*	*	*	*
3309	8	*	*	*	*	*	*	*
3312	408	35.0	26.5	27.5	0.5	1.2	9.3	0.0
3550	63	*	*	*	*	*	*	*
3551	14	*	*	*	*	*	*	*
3552	33	*	*	*	*	*	*	*
3701	955	87.2	0.2	8.8	0.0	0.1	3.6	0.1
3850	2,945	71.5	14.4	5.8	0.2	0.5	7.4	0.2
3901	1,635	35.4	51.3	6.2	0.1	0.7	6.0	0.3
3902	122	52.5	29.5	7.4	0.0	1.6	9.0	0.0
3903	25	*	*	*	*	*	*	*
3904	4	*	*	*	*	*	*	*
3905	9	*	*	*	*	*	*	*
3950	2,197	53.1	23.1	12.9	0.3	1.2	8.8	0.6
3954	15	*	*	*	*	*	*	*
4103	1	*	*	*	*	*	*	*
4106	2,310	12.3	40.8	41.4	0.3	0.7	4.2	0.3
4107	2,925	41.2	28.6	20.9	0.4	2.0	6.6	0.2
4109	760	84.1	1.7	9.9	0.0	0.4	3.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 67

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4110	7	*	*	*	*	*	*	*
4112	554	1.1	1.3	96.0	0.2	0.2	0.0	1.3
4115	3,534	30.5	31.5	21.6	0.3	11.1	4.8	0.2
4351	506	87.0	0.6	6.7	0.2	0.6	4.7	0.2
4352	166	74.1	11.4	3.0	0.0	0.0	10.8	0.6
4353	437	61.3	0.9	34.3	0.5	0.0	3.0	0.0
4354	738	87.3	3.8	3.0	0.3	0.4	4.9	0.4
4355	303	81.2	1.3	6.3	2.6	0.3	7.9	0.3
4356	400	79.0	1.8	12.0	0.3	1.0	5.5	0.5
4358	60	*	*	*	*	*	*	*
4359	561	65.2	0.4	31.6	0.5	0.2	1.4	0.7
4360	207	87.4	0.0	7.7	0.0	0.0	4.8	0.0
4361	271	91.5	0.7	4.8	0.0	0.0	2.6	0.4
4362	287	91.6	0.3	3.1	0.0	0.7	4.2	0.0
4363	754	90.7	0.5	2.7	0.0	0.4	5.3	0.4
4401	870	90.2	0.7	4.8	0.0	0.6	3.3	0.3
4501	830	93.3	0.1	4.7	0.0	0.1	1.8	0.0
4601	1,262	72.1	2.5	19.2	0.0	0.2	5.5	0.5
4602	1	*	*	*	*	*	*	*
4701	1,027	91.2	1.2	3.8	0.0	0.6	2.3	0.9
5050	1,548	48.8	41.4	2.5	0.0	1.5	5.5	0.3
5051	2,714	51.4	36.2	4.1	0.1	2.9	5.0	0.4
5052	2,771	34.4	40.9	2.7	0.2	17.1	4.3	0.4
5055	1,235	84.4	6.2	0.7	0.1	8.2	0.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 68

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
5101	1,602	68.1	13.0	11.9	0.2	2.1	4.7	0.0
5102	28	*	*	*	*	*	*	*
5202	719	87.8	1.3	2.8	1.5	0.4	6.1	0.1
5206	9	*	*	*	*	*	*	*
5601	965	89.6	1.6	3.9	0.0	1.8	3.0	0.1
5801	1,811	81.2	6.7	6.0	0.1	0.3	5.0	0.7
5803	93	*	*	*	*	*	*	*
5804	24	*	*	*	*	*	*	*
5806	49	*	*	*	*	*	*	*
6101	40	*	*	*	*	*	*	*
6102	105	81.0	1.0	9.5	0.0	1.0	7.6	0.0
6103	3,439	53.7	28.0	9.5	0.3	0.5	7.6	0.4
6105	11	*	*	*	*	*	*	*
6106	117	35.0	49.6	7.7	0.0	0.9	6.8	0.0
6301	3,416	40.0	30.5	22.1	0.1	1.9	5.2	0.2
6312	1,799	35.7	53.9	3.7	0.1	1.7	4.7	0.2
6316	1,167	64.0	21.9	7.0	0.1	3.4	3.3	0.2
6318	36	*	*	*	*	*	*	*
6320	352	39.2	49.4	4.0	0.3	1.7	5.4	0.0
6325	314	90.8	1.9	4.5	0.0	0.0	2.5	0.3
6335	996	70.8	17.7	2.9	0.0	6.5	2.1	0.0
6340	21	*	*	*	*	*	*	*
6345	106	90.6	1.9	2.8	0.0	1.9	2.8	0.0
6360	1,025	29.4	49.7	15.8	0.0	1.2	3.8	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 69

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
7304	10	*	*	*	*	*	*	*
7306	2,271	46.1	31.3	15.8	0.0	0.7	5.9	0.4
7308	10	*	*	*	*	*	*	*
7310	2	*	*	*	*	*	*	*
7401	2,039	81.1	6.1	6.4	0.1	0.5	5.5	0.2
7601	590	93.6	0.0	4.4	0.2	0.3	1.0	0.5
7850	1,069	88.2	0.5	6.6	0.0	0.1	4.4	0.2
7852	18	*	*	*	*	*	*	*
8101	1,693	5.1	91.3	0.7	0.0	0.2	2.6	0.1
8102	483	17.2	14.5	64.0	0.0	1.0	3.3	0.0
8105	778	31.1	63.5	2.4	0.1	0.8	1.9	0.1
8106	552	25.9	56.2	11.2	0.2	0.5	5.6	0.4
8150	18	*	*	*	*	*	*	*
8151	3,147	37.0	40.8	10.2	0.1	2.5	9.0	0.4
8152	32	*	*	*	*	*	*	*
8153	7	*	*	*	*	*	*	*
8155	12	*	*	*	*	*	*	*
8157	6	*	*	*	*	*	*	*
8201	2,692	44.4	39.9	4.8	0.3	3.7	6.3	0.6
8202	3,885	84.1	12.4	1.6	0.0	0.4	1.1	0.4
8203	408	57.6	31.4	4.9	0.2	1.0	4.4	0.5
8250	665	20.6	69.9	3.5	0.2	1.2	4.4	0.3
8252	169	7.7	87.0	3.0	0.0	0.6	1.8	0.0
8255	2,345	53.5	31.1	9.5	0.3	2.4	3.1	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 70

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8257	1,939	52.2	12.2	0.7	0.3	32.9	1.6	0.2
8263	639	65.9	15.0	13.1	0.2	1.9	3.9	0.0
8264	784	61.2	28.1	4.6	0.0	1.8	4.3	0.0
8267	693	54.3	17.6	24.2	0.1	0.3	3.2	0.3
8901	934	5.2	89.8	1.2	0.0	1.1	2.5	0.2
8902	1,406	6.0	89.5	2.1	0.0	0.1	2.1	0.1
8903	1,508	4.4	89.5	1.7	0.1	0.8	3.3	0.2
8904	2,310	3.4	91.9	1.6	0.1	0.8	2.0	0.3
8905	1,769	61.2	27.4	7.7	0.1	1.0	2.5	0.1
8909	934	4.8	91.3	1.8	0.0	0.1	1.7	0.2
9045	1,573	10.7	27.0	59.2	0.1	0.3	2.4	0.3
9050	963	16.1	65.4	15.2	0.0	0.0	3.3	0.0
9051	1,519	8.8	6.0	82.8	0.0	0.2	1.9	0.4
9053	1,202	5.7	88.6	1.8	0.2	1.5	2.1	0.2
9055	12	*	*	*	*	*	*	*
9058	1,145	16.2	72.5	1.6	0.0	7.9	1.7	0.1
9066	1,127	2.1	94.9	0.6	0.0	0.3	2.1	0.0
9068	906	2.3	95.9	0.2	0.0	0.3	1.0	0.2
9069	2,229	1.2	97.2	0.8	0.0	0.1	0.6	0.0
9070	1,519	4.2	91.0	1.3	0.1	1.8	1.4	0.1
9072	940	3.0	94.1	0.6	0.1	0.6	1.4	0.1
9080	2,600	18.4	77.3	2.0	0.1	0.2	1.8	0.1
9082	131	0.8	97.7	0.8	0.0	0.0	0.8	0.0
9083	72	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 71

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9084	23	*	*	*	*	*	*	*
9085	4	*	*	*	*	*	*	*
9086	12	*	*	*	*	*	*	*
9087	13	*	*	*	*	*	*	*
Michigan	152,475	51.1	30.5	10.9	0.4	2.3	4.6	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 72

Michigan

Comparison of Age Distribution by Clinic Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	109	0.0	0.9	2.8	66.1	27.5	2.8	0.0
0102	433	0.0	3.0	7.4	64.0	24.0	1.6	0.0
0103	392	0.0	3.1	8.9	65.1	21.9	1.0	0.0
0104	122	0.0	6.6	7.4	64.8	20.5	0.8	0.0
0105	16	*	*	*	*	*	*	*
0250	125	0.0	5.6	4.8	61.6	27.2	0.8	0.0
0251	92	*	*	*	*	*	*	*
0252	98	*	*	*	*	*	*	*
0253	125	0.0	3.2	9.6	68.8	16.8	1.6	0.0
0401	456	0.0	2.4	8.6	63.2	24.6	1.3	0.0
0402	383	0.3	2.1	10.2	60.6	25.1	1.8	0.0
0403	149	0.0	2.7	7.4	62.4	26.2	1.3	0.0
0404	101	0.0	1.0	4.0	60.4	33.7	1.0	0.0
0501	466	0.2	1.3	8.6	64.2	24.9	0.9	0.0
0503	90	*	*	*	*	*	*	*
0504	410	0.0	2.2	5.6	62.2	27.1	2.9	0.0
0505	248	0.0	1.6	8.1	69.4	20.6	0.0	0.4
0506	163	0.0	2.5	7.4	67.5	22.7	0.0	0.0
0507	79	*	*	*	*	*	*	*
0508	25	*	*	*	*	*	*	*
0509	1	*	*	*	*	*	*	*
0650	1,058	0.1	2.5	7.8	66.3	22.1	1.2	0.1
0651	578	0.3	3.8	8.3	65.1	21.3	1.0	0.2
0652	323	0.0	2.2	7.1	69.7	19.8	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 73

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0653	400	0.0	2.0	4.8	65.3	26.5	1.3	0.3
0654	386	0.3	3.9	9.1	61.9	22.5	1.6	0.8
0655	324	0.0	3.1	9.6	58.6	27.2	1.2	0.3
0657	65	*	*	*	*	*	*	*
0701	463	0.0	1.9	6.7	55.1	33.9	2.4	0.0
0702	69	*	*	*	*	*	*	*
0703	165	0.0	0.0	9.1	55.8	34.5	0.6	0.0
0704	55	*	*	*	*	*	*	*
0705	87	*	*	*	*	*	*	*
0706	10	*	*	*	*	*	*	*
0850	720	0.1	3.1	8.3	59.3	26.8	1.9	0.4
0851	630	0.0	2.5	6.8	64.1	24.4	1.3	0.8
0901	1,524	0.1	2.5	7.4	62.7	25.9	1.4	0.0
0902	41	*	*	*	*	*	*	*
1051	1,175	0.1	1.7	6.3	60.3	29.0	2.2	0.3
1052	1	*	*	*	*	*	*	*
1101	1,674	0.2	3.9	9.4	60.4	22.1	1.7	2.3
1102	837	0.0	2.2	9.8	61.3	24.9	1.3	0.6
1104	1	*	*	*	*	*	*	*
1106	74	*	*	*	*	*	*	*
1250	1,015	0.0	1.9	9.7	61.8	23.5	3.1	0.1
1251	674	0.4	2.2	8.2	66.8	20.2	1.5	0.7
1252	535	0.2	3.0	9.5	61.7	23.0	2.6	0.0
1253	735	0.0	2.4	8.0	65.7	22.6	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 74

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1350	2,352	0.1	2.9	7.5	64.2	23.5	1.4	0.5
1351	80	*	*	*	*	*	*	*
1352	391	0.3	1.8	7.7	66.5	22.0	1.3	0.5
1401	226	0.0	3.5	5.3	64.2	26.5	0.4	0.0
1402	113	0.0	0.9	2.7	52.2	40.7	2.7	0.9
1403	19	*	*	*	*	*	*	*
1501	785	0.0	1.8	9.4	62.4	23.7	1.8	0.9
1503	894	0.3	3.8	11.6	58.8	23.7	1.2	0.4
1504	1,998	0.1	2.6	7.0	60.8	27.6	1.8	0.1
1505	285	0.0	1.8	5.6	65.6	24.9	1.8	0.4
1506	583	0.0	2.4	5.8	62.8	26.9	1.9	0.2
1507	556	0.0	1.3	5.2	62.1	29.0	2.3	0.2
1508	22	*	*	*	*	*	*	*
1509	419	0.0	1.9	6.7	60.9	27.7	2.6	0.2
1511	621	0.0	2.9	8.5	60.4	26.6	1.4	0.2
1512	345	0.0	2.6	6.1	58.6	30.7	2.0	0.0
1514	433	0.2	0.9	4.6	63.5	27.5	2.8	0.5
1701	594	0.0	2.2	7.4	63.0	25.8	1.7	0.0
1950	612	0.0	2.9	10.0	64.1	21.4	1.3	0.3
1952	130	0.0	0.8	6.2	67.7	23.1	2.3	0.0
1953	284	0.4	4.9	6.7	64.8	21.1	2.1	0.0
1954	446	0.4	2.2	7.6	65.2	22.9	1.3	0.2
1955	107	0.0	1.9	13.1	61.7	22.4	0.9	0.0
1957	394	0.0	1.5	5.3	57.6	34.3	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 75

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1963	63	*	*	*	*	*	*	*
1964	17	*	*	*	*	*	*	*
2150	631	0.2	1.4	7.6	62.3	27.7	0.8	0.0
2151	223	0.0	4.0	7.6	65.5	22.0	0.9	0.0
2152	17	*	*	*	*	*	*	*
2251	403	0.2	1.5	6.9	62.3	28.0	1.0	0.0
2252	171	0.0	4.7	7.0	59.6	27.5	1.2	0.0
2554	48	*	*	*	*	*	*	*
2555	7,386	0.2	2.4	7.6	64.0	24.1	1.6	0.1
2556	287	0.0	2.1	7.3	70.7	18.5	1.4	0.0
2557	2	*	*	*	*	*	*	*
2558	4	*	*	*	*	*	*	*
2560	5	*	*	*	*	*	*	*
2561	3	*	*	*	*	*	*	*
2562	16	*	*	*	*	*	*	*
2563	26	*	*	*	*	*	*	*
2564	22	*	*	*	*	*	*	*
3001	1,156	0.1	2.9	8.8	65.5	20.6	1.4	0.7
3002	262	0.0	4.6	6.5	66.0	21.4	1.5	0.0
3004	250	0.0	1.6	7.2	58.0	31.2	2.0	0.0
3005	116	0.0	0.9	6.0	67.2	22.4	3.4	0.0
3007	13	*	*	*	*	*	*	*
3201	419	0.0	1.0	4.8	62.1	30.5	1.7	0.0
3203	1	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 76

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3301	4,320	0.1	2.7	7.9	61.6	25.7	1.9	0.2
3304	97	*	*	*	*	*	*	*
3305	156	0.0	0.0	8.3	57.7	31.4	2.6	0.0
3306	242	0.4	4.5	7.4	52.1	32.6	2.1	0.8
3308	98	*	*	*	*	*	*	*
3309	8	*	*	*	*	*	*	*
3312	408	0.0	1.7	8.8	61.8	26.0	1.5	0.2
3550	63	*	*	*	*	*	*	*
3551	14	*	*	*	*	*	*	*
3552	33	*	*	*	*	*	*	*
3701	955	0.1	2.2	8.7	64.6	23.0	1.2	0.2
3850	2,945	0.0	2.9	8.2	63.4	24.0	1.1	0.4
3901	1,635	0.2	3.5	9.0	58.3	26.8	1.8	0.4
3902	122	0.0	3.3	10.7	59.8	23.8	2.5	0.0
3903	25	*	*	*	*	*	*	*
3904	4	*	*	*	*	*	*	*
3905	9	*	*	*	*	*	*	*
3950	2,197	0.0	3.2	7.3	61.4	25.6	2.0	0.3
3954	15	*	*	*	*	*	*	*
4103	1	*	*	*	*	*	*	*
4106	2,310	0.0	3.1	8.0	58.8	26.6	2.6	0.8
4107	2,925	0.1	2.7	7.5	59.6	28.3	1.3	0.4
4109	760	0.1	2.2	6.1	63.2	27.1	1.3	0.0
4110	7	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 77

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4112	554	0.4	4.9	9.7	53.1	28.7	2.9	0.4
4115	3,534	0.1	1.7	6.1	57.3	32.1	2.4	0.4
4351	506	0.0	1.4	8.3	66.4	22.1	1.6	0.2
4352	166	0.0	2.4	5.4	66.3	24.7	1.2	0.0
4353	437	0.0	3.2	8.0	66.1	19.7	2.7	0.2
4354	738	0.1	1.5	6.5	67.1	23.2	1.4	0.3
4355	303	0.3	1.7	8.3	55.4	29.4	4.3	0.7
4356	400	0.0	3.3	7.8	65.5	22.0	1.0	0.5
4358	60	*	*	*	*	*	*	*
4359	561	0.2	2.5	7.3	60.2	26.0	3.2	0.5
4360	207	0.0	0.0	10.6	64.7	23.2	1.4	0.0
4361	271	0.0	1.5	8.1	63.5	25.1	1.5	0.4
4362	287	0.0	3.5	10.1	61.7	24.7	0.0	0.0
4363	754	0.0	2.5	8.9	64.9	22.4	1.2	0.1
4401	870	0.0	2.9	5.3	64.0	26.2	1.4	0.2
4501	830	0.1	2.8	6.1	64.9	23.6	2.2	0.2
4601	1,262	0.1	1.4	8.1	63.5	25.7	1.0	0.2
4602	1	*	*	*	*	*	*	*
4701	1,027	0.0	1.7	5.2	59.8	31.1	2.1	0.2
5050	1,548	0.0	1.4	4.7	58.1	33.7	2.2	0.0
5051	2,714	0.0	1.0	4.7	60.4	31.4	2.5	0.0
5052	2,771	0.0	1.4	5.3	59.2	31.6	2.3	0.0
5055	1,235	0.0	0.2	1.4	50.7	43.5	4.0	0.2
5101	1,602	0.1	0.9	4.2	60.9	31.3	2.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 78

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5102	28	*	*	*	*	*	*	*
5202	719	0.1	1.4	5.7	61.5	28.8	2.5	0.0
5206	9	*	*	*	*	*	*	*
5601	965	0.0	1.6	5.9	64.5	26.5	1.6	0.0
5801	1,811	0.0	2.5	8.2	63.5	24.4	1.3	0.1
5803	93	*	*	*	*	*	*	*
5804	24	*	*	*	*	*	*	*
5806	49	*	*	*	*	*	*	*
6101	40	*	*	*	*	*	*	*
6102	105	0.0	6.7	9.5	58.1	22.9	2.9	0.0
6103	3,439	0.2	3.1	10.0	61.6	23.4	1.3	0.4
6105	11	*	*	*	*	*	*	*
6106	117	0.0	4.3	8.5	70.1	16.2	0.0	0.9
6301	3,416	0.1	2.4	6.9	60.3	28.2	2.1	0.0
6312	1,799	0.0	0.6	3.5	53.3	39.7	2.9	0.1
6316	1,167	0.0	0.5	2.7	56.2	37.4	3.2	0.0
6318	36	*	*	*	*	*	*	*
6320	352	0.0	0.9	1.7	61.4	32.7	3.1	0.3
6325	314	0.0	1.3	4.5	60.5	30.6	1.0	2.2
6335	996	0.0	0.2	2.6	55.0	38.1	4.1	0.0
6340	21	*	*	*	*	*	*	*
6345	106	0.0	0.0	2.8	61.3	34.9	0.9	0.0
6360	1,025	0.0	2.2	8.9	58.3	27.2	2.0	1.3
7304	10	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 79

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
7306	2,271	0.1	3.0	8.1	63.0	24.6	1.1	0.2
7308	10	*	*	*	*	*	*	*
7310	2	*	*	*	*	*	*	*
7401	2,039	0.1	2.5	7.8	64.3	23.1	1.9	0.3
7601	590	0.0	2.7	8.6	66.4	21.9	0.3	0.0
7850	1,069	0.0	3.4	7.9	66.7	21.0	1.0	0.0
7852	18	*	*	*	*	*	*	*
8101	1,693	0.2	2.8	7.8	58.8	27.8	2.2	0.4
8102	483	0.0	2.5	4.3	62.3	28.6	2.3	0.0
8105	778	0.1	1.3	5.1	60.2	31.1	2.1	0.1
8106	552	0.0	2.4	4.7	66.3	25.0	1.6	0.0
8150	18	*	*	*	*	*	*	*
8151	3,147	0.1	1.4	6.2	57.8	31.9	2.3	0.3
8152	32	*	*	*	*	*	*	*
8153	7	*	*	*	*	*	*	*
8155	12	*	*	*	*	*	*	*
8157	6	*	*	*	*	*	*	*
8201	2,692	0.0	1.4	5.6	59.2	31.6	2.0	0.1
8202	3,885	0.0	1.1	3.8	53.9	35.6	4.8	0.7
8203	408	0.2	0.5	5.6	56.4	34.3	2.9	0.0
8250	665	0.0	1.7	6.3	58.9	30.4	2.6	0.2
8252	169	0.0	1.2	6.5	67.5	24.3	0.6	0.0
8255	2,345	0.0	1.2	6.5	63.4	27.0	1.8	0.0
8257	1,939	0.1	0.5	3.4	56.8	34.1	5.0	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 80

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8263	639	0.0	1.3	5.0	62.6	29.1	2.0	0.0
8264	784	0.0	0.9	3.2	61.4	32.1	2.4	0.0
8267	693	0.1	2.5	6.3	63.3	25.4	2.3	0.0
8901	934	0.3	3.0	7.7	61.0	25.2	2.5	0.3
8902	1,406	0.1	1.4	4.8	61.5	29.9	2.1	0.2
8903	1,508	0.1	3.4	8.1	60.5	25.5	2.1	0.3
8904	2,310	0.2	2.4	6.7	61.5	27.3	1.9	0.1
8905	1,769	0.0	1.1	3.4	55.3	35.8	4.1	0.2
8909	934	0.0	2.4	7.3	61.2	26.2	2.7	0.2
9045	1,573	0.2	3.3	7.6	55.9	29.2	3.7	0.1
9050	963	0.1	2.5	7.4	60.0	28.0	1.8	0.2
9051	1,519	0.1	2.2	7.2	54.4	30.7	5.3	0.0
9053	1,202	0.0	2.7	6.7	58.2	30.0	2.2	0.3
9055	12	*	*	*	*	*	*	*
9058	1,145	0.0	1.9	6.6	57.7	30.5	3.1	0.3
9066	1,127	0.3	2.8	6.3	59.4	28.6	2.5	0.2
9068	906	0.2	3.4	6.6	60.8	25.7	2.6	0.6
9069	2,229	0.1	3.4	6.1	58.0	29.9	2.3	0.2
9070	1,519	0.1	2.7	8.4	59.8	26.8	2.2	0.1
9072	940	0.3	2.3	10.5	60.3	25.2	1.2	0.1
9080	2,600	0.0	1.7	6.2	58.6	31.0	2.4	0.2
9082	131	0.0	4.6	6.1	58.0	29.0	2.3	0.0
9083	72	*	*	*	*	*	*	*
9084	23	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 4A**2019 - 2021 Pregnancy Nutrition Surveillance (1)**

Page : 81

Michigan**Comparison of Age Distribution by Clinic****Combined 3 Years**

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9085	4	*	*	*	*	*	*	*
9086	12	*	*	*	*	*	*	*
9087	13	*	*	*	*	*	*	*
Michigan	152,475	0.1	2.2	6.9	60.8	27.7	2.1	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 82

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	70	*	*	34	*	*	10	*	11	*	97	*	97	*
0102	278	5.0 (104)	61.9 (80)	118	16.1 (33)	51.7 (49)	34	*	28	*	389	6.7 (28)	389	11.3 (46)
0103	330	5.2 (117)	64.2 (115)	169	14.8 (15)	55.0 (87)	27	*	39	*	357	5.0 (6)	358	12.6 (71)
0104	89	*	*	34	*	*	4	*	11	*	112	6.3 (22)	113	12.4 (67)
0105	11	*	*	3	*	*	0	*	0	*	16	*	16	*
0250	117	5.1 (111)	57.3 (24)	86	*	*	7	*	7	*	109	13.8 (139)	109	11.9 (57)
0251	70	*	*	26	*	*	6	*	8	*	81	*	81	*
0252	95	*	*	82	*	*	8	*	12	*	89	*	89	*
0253	105	0.0 (1)	65.7 (124)	74	*	*	3	*	8	*	114	8.8 (82)	114	17.5 (132)
0401	416	5.5 (125)	63.2 (100)	241	18.3 (71)	54.4 (81)	31	*	29	*	414	8.2 (67)	415	18.6 (138)
0402	339	5.9 (132)	57.5 (26)	88	*	*	22	*	35	*	348	9.2 (89)	348	11.8 (55)
0403	138	5.1 (111)	58.0 (31)	68	*	*	10	*	15	*	135	11.1 (130)	135	8.9 (14)
0404	96	*	*	65	*	*	5	*	3	*	95	*	95	*
0501	305	4.6 (96)	55.7 (18)	81	*	*	47	*	41	*	439	8.2 (67)	439	11.6 (51)
0503	73	*	*	30	*	*	11	*	7	*	86	*	86	*
0504	360	3.6 (43)	62.5 (90)	152	13.8 (9)	53.9 (76)	38	*	39	*	384	6.3 (22)	384	15.9 (119)
0505	177	4.5 (90)	62.1 (83)	56	*	*	24	*	22	*	230	5.7 (14)	230	15.7 (115)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 83

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	105	2.9 (23)	57.1 (22)	24	*	*	13	*	17	*	153	9.2 (89)	152	10.5 (32)
0507	46	*	*	11	*	*	3	*	14	*	79	*	79	*
0508	16	*	*	9	*	*	1	*	3	*	25	*	25	*
0509	1	*	*	1	*	*	0	*	0	*	1	*	1	*
0650	892	5.4 (123)	61.3 (70)	580	15.0 (17)	51.2 (41)	71	*	89	*	968	9.5 (99)	970	11.9 (57)
0651	508	4.5 (90)	59.1 (39)	293	14.0 (11)	53.9 (76)	32	*	42	*	535	7.1 (35)	537	10.4 (30)
0652	266	2.6 (15)	63.5 (105)	202	20.3 (102)	53.0 (63)	23	*	29	*	315	9.8 (106)	316	17.1 (129)
0653	371	3.2 (31)	62.3 (85)	275	13.1 (6)	56.7 (100)	25	*	31	*	371	9.4 (95)	373	15.0 (105)
0654	297	1.7 (5)	63.3 (102)	177	23.2 (112)	50.3 (27)	27	*	29	*	355	6.8 (30)	357	12.6 (71)
0655	300	4.0 (70)	60.3 (48)	219	21.0 (106)	51.6 (45)	24	*	17	*	289	10.0 (113)	291	22.0 (143)
0657	44	*	*	24	*	*	5	*	7	*	64	*	64	*
0701	390	2.6 (15)	55.6 (17)	258	18.2 (66)	43.4 (5)	20	*	45	*	411	7.5 (47)	411	12.4 (67)
0702	66	*	*	55	*	*	3	*	8	*	66	*	66	*
0703	145	2.8 (19)	57.9 (30)	92	*	*	14	*	11	*	148	14.2 (140)	148	14.2 (97)
0704	53	*	*	36	*	*	5	*	7	*	47	*	48	*
0705	66	*	*	42	*	*	13	*	11	*	79	*	79	*
0706	8	*	*	5	*	*	0	*	3	*	10	*	10	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 84

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	450	6.2 (133)	60.9 (60)	226	19.9 (97)	47.8 (13)	54	*	64	*	658	10.9 (127)	659	16.2 (122)
0851	328	6.4 (135)	54.9 (14)	110	16.4 (38)	53.6 (71)	56	*	59	*	577	9.9 (109)	580	13.3 (80)
0901	1,248	3.1 (27)	62.3 (85)	749	17.2 (53)	54.9 (85)	112	24.1 (1)	81	*	1,406	6.9 (32)	1,406	11.2 (43)
0902	35	*	*	20	*	*	3	*	2	*	35	*	35	*
1051	988	4.9 (103)	53.5 (9)	769	15.9 (28)	51.9 (52)	79	*	70	*	1,083	8.7 (80)	1,086	8.7 (12)
1052	1	*	*	0	*	*	0	*	0	*	1	*	1	*
1101	1,422	3.7 (50)	62.9 (97)	1,041	20.2 (100)	52.6 (57)	151	39.7 (10)	127	26.8 (10)	1,526	8.8 (82)	1,527	19.1 (139)
1102	734	5.2 (117)	59.1 (39)	487	15.8 (25)	56.1 (94)	78	*	93	*	717	8.1 (63)	718	16.0 (121)
1104	0	*	*	0	*	*	0	*	0	*	1	*	1	*
1106	63	*	*	46	*	*	11	*	7	*	60	*	60	*
1250	533	4.5 (90)	61.5 (76)	220	21.4 (109)	50.5 (30)	82	*	71	*	926	9.8 (106)	927	9.8 (21)
1251	566	5.5 (125)	62.4 (88)	407	17.0 (48)	51.6 (45)	40	*	77	*	620	6.9 (32)	621	6.1 (5)
1252	434	5.1 (111)	66.1 (130)	138	18.8 (76)	56.5 (97)	36	*	43	*	478	8.8 (82)	479	10.0 (24)
1253	646	5.1 (111)	59.0 (37)	262	23.7 (114)	51.5 (43)	49	*	54	*	656	9.6 (103)	655	11.0 (39)
1350	1,888	3.9 (69)	61.2 (69)	1,171	18.8 (76)	51.6 (45)	129	38.8 (7)	174	27.0 (11)	2,143	8.5 (74)	2,143	10.7 (34)
1351	64	*	*	47	*	*	11	*	19	*	71	*	71	*
1352	213	5.2 (117)	62.0 (82)	106	18.9 (79)	63.2 (118)	19	*	30	*	359	4.5 (3)	359	4.5 (3)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 85

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1401	185	4.3 (79)	49.2 (1)	111	11.7 (3)	55.9 (91)	20	*	19	*	209	6.7 (28)	209	11.5 (49)
1402	99	*	*	61	*	*	7	*	8	*	103	11.7 (134)	103	6.8 (6)
1403	16	*	*	13	*	*	0	*	3	*	18	*	18	*
1501	661	3.6 (43)	66.0 (129)	356	42.1 (119)	32.0 (2)	75	*	81	*	707	11.2 (131)	706	9.8 (21)
1503	764	4.6 (96)	60.7 (56)	554	17.9 (60)	50.2 (25)	77	*	66	*	825	8.7 (80)	828	14.0 (92)
1504	934	5.5 (125)	57.2 (23)	528	15.5 (22)	51.1 (40)	182	41.8 (11)	367	33.5 (29)	1,795	9.3 (93)	1,797	8.5 (11)
1505	184	3.8 (57)	59.2 (42)	63	*	*	30	*	55	*	256	9.4 (95)	256	10.2 (27)
1506	471	5.1 (111)	64.3 (116)	92	*	*	48	*	68	*	528	11.7 (134)	529	12.3 (60)
1507	398	5.0 (104)	57.3 (24)	124	10.5 (2)	57.3 (108)	35	*	66	*	511	9.4 (95)	511	8.2 (8)
1508	14	*	*	9	*	*	1	*	3	*	18	*	18	*
1509	312	4.5 (90)	64.1 (112)	163	17.2 (53)	53.4 (68)	33	*	77	*	355	9.3 (93)	356	8.4 (10)
1511	496	5.0 (104)	60.5 (53)	257	20.6 (104)	50.2 (25)	69	*	113	24.8 (8)	576	9.9 (109)	577	13.0 (77)
1512	256	5.1 (111)	61.7 (79)	77	*	*	40	*	82	*	322	4.0 (2)	322	11.5 (49)
1514	219	5.0 (104)	51.6 (2)	104	13.5 (8)	52.9 (61)	35	*	95	*	411	8.0 (57)	411	15.8 (118)
1701	520	3.8 (57)	59.2 (42)	362	13.8 (9)	54.4 (81)	31	*	53	*	532	8.5 (74)	534	10.3 (28)
1950	560	3.6 (43)	61.4 (72)	378	16.4 (38)	52.1 (53)	40	*	58	*	571	10.3 (120)	571	13.8 (89)
1952	118	2.5 (14)	71.2 (140)	86	*	*	12	*	18	*	126	7.1 (35)	126	16.7 (126)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Maternal Health Indicators by Clinic Combined 3 Years

Page : 86

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1953	252	7.5 (137)	56.7 (20)	182	19.8 (94)	50.5 (30)	26	*	51	*	262	8.0 (57)	262	19.8 (141)
1954	417	2.4 (13)	67.4 (135)	316	22.5 (111)	50.3 (27)	33	*	37	*	416	14.9 (141)	416	15.4 (111)
1955	102	2.0 (9)	70.6 (139)	72	*	*	6	*	10	*	98	*	98	*
1957	360	4.2 (75)	66.9 (133)	246	15.4 (20)	48.8 (15)	27	*	44	*	367	15.5 (142)	367	15.5 (113)
1963	53	*	*	37	*	*	6	*	4	*	58	*	60	*
1964	16	*	*	10	*	*	3	*	5	*	15	*	15	*
2150	593	3.0 (26)	61.4 (72)	444	18.0 (61)	51.6 (45)	57	*	90	*	574	10.1 (117)	574	15.2 (107)
2151	208	5.3 (121)	53.4 (8)	163	16.0 (30)	50.9 (37)	14	*	24	*	186	5.4 (10)	187	13.9 (90)
2152	17	*	*	12	*	*	1	*	0	*	15	*	16	*
2251	275	6.5 (136)	54.5 (12)	110	18.2 (66)	49.1 (16)	33	*	39	*	363	3.0 (1)	364	15.7 (115)
2252	120	9.2 (140)	53.3 (7)	43	*	*	11	*	14	*	146	9.6 (103)	146	13.0 (77)
2554	38	*	*	16	*	*	5	*	4	*	45	*	45	*
2555	5,282	5.0 (104)	59.6 (45)	2,552	18.2 (66)	52.4 (56)	671	45.9 (20)	567	30.9 (20)	6,758	8.0 (57)	6,768	15.9 (119)
2556	227	4.4 (86)	59.9 (47)	136	27.2 (117)	44.9 (7)	18	*	22	*	270	8.5 (74)	270	14.1 (95)
2557	2	*	*	1	*	*	1	*	0	*	2	*	2	*
2558	4	*	*	2	*	*	1	*	1	*	3	*	3	*
2560	3	*	*	1	*	*	0	*	0	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 87

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2561	2	*	*	2	*	*	2	*	0	*	3	*	3	*
2562	12	*	*	3	*	*	3	*	2	*	15	*	15	*
2563	16	*	*	6	*	*	3	*	6	*	24	*	24	*
2564	11	*	*	2	*	*	3	*	2	*	18	*	18	*
3001	953	4.0 (70)	65.8 (126)	505	20.4 (103)	47.7 (12)	86	*	260	31.5 (21)	1,089	7.2 (38)	1,089	12.9 (76)
3002	227	1.3 (3)	77.1 (141)	129	19.4 (87)	50.4 (29)	17	*	84	*	243	7.4 (43)	243	14.8 (102)
3004	177	1.7 (5)	67.8 (137)	89	*	*	13	*	19	*	228	9.2 (89)	228	12.3 (60)
3005	95	*	*	30	*	*	11	*	9	*	110	10.9 (127)	110	10.9 (36)
3007	10	*	*	3	*	*	0	*	4	*	12	*	12	*
3201	369	5.4 (123)	66.1 (130)	275	17.1 (51)	57.1 (105)	34	*	52	*	398	8.3 (69)	400	12.0 (59)
3203	1	*	*	0	*	*	0	*	0	*	1	*	1	*
3301	3,552	3.7 (50)	62.7 (94)	2,724	19.3 (84)	54.2 (80)	361	44.6 (15)	303	32.7 (25)	4,075	7.5 (47)	4,076	12.3 (60)
3304	76	*	*	61	*	*	5	*	9	*	90	*	90	*
3305	126	7.9 (138)	56.3 (19)	95	*	*	14	*	18	*	141	12.1 (137)	141	7.1 (7)
3306	197	4.1 (73)	60.9 (60)	165	17.0 (48)	58.2 (114)	19	*	21	*	227	5.3 (7)	227	10.1 (26)
3308	91	*	*	73	*	*	8	*	9	*	87	*	87	*
3309	7	*	*	4	*	*	1	*	0	*	7	*	7	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 88

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3312	333	3.6 (43)	62.2 (84)	262	19.1 (83)	50.8 (35)	36	*	29	*	390	10.0 (113)	390	12.3 (60)
3550	52	*	*	16	*	*	6	*	6	*	56	*	56	*
3551	10	*	*	2	*	*	1	*	3	*	13	*	13	*
3552	28	*	*	9	*	*	5	*	7	*	30	*	30	*
3701	705	4.3 (79)	64.1 (112)	502	17.5 (57)	54.0 (79)	72	*	116	8.6 (1)	877	10.0 (113)	879	18.3 (137)
3850	2,031	4.0 (70)	60.3 (48)	1,309	16.7 (43)	53.9 (76)	195	39.0 (8)	199	30.2 (17)	2,728	7.9 (56)	2,733	10.9 (36)
3901	1,256	2.9 (23)	60.7 (56)	546	16.8 (46)	56.6 (98)	84	*	133	26.3 (9)	1,448	8.4 (70)	1,449	12.4 (67)
3902	96	*	*	41	*	*	11	*	20	*	99	*	99	*
3903	21	*	*	10	*	*	2	*	11	*	24	*	24	*
3904	3	*	*	1	*	*	2	*	0	*	4	*	4	*
3905	7	*	*	5	*	*	0	*	2	*	9	*	9	*
3950	1,745	3.7 (50)	62.9 (97)	958	15.8 (25)	53.2 (65)	148	41.9 (12)	205	37.1 (34)	1,976	10.2 (118)	1,977	11.0 (39)
3954	13	*	*	9	*	*	2	*	2	*	14	*	14	*
4103	0	*	*	0	*	*	0	*	0	*	1	*	1	*
4106	1,354	3.3 (33)	62.5 (90)	592	23.8 (115)	42.9 (4)	272	38.2 (6)	465	27.1 (12)	2,139	6.4 (24)	2,142	13.4 (83)
4107	2,108	3.8 (57)	59.6 (45)	880	15.8 (25)	51.8 (51)	298	52.0 (30)	626	30.8 (19)	2,664	7.4 (43)	2,669	16.4 (123)
4109	421	3.1 (27)	58.9 (36)	232	15.9 (28)	57.3 (108)	79	*	142	18.3 (4)	709	7.8 (52)	709	15.2 (107)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 89

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4110	6	*	*	6	*	*	0	*	0	*	7	*	7	*
4112	536	1.5 (4)	65.7 (124)	407	62.4 (120)	16.5 (1)	42	*	31	*	513	18.1 (143)	514	15.4 (111)
4115	1,779	4.8 (101)	59.0 (37)	398	18.6 (73)	49.7 (20)	402	38.1 (5)	698	30.7 (18)	3,254	8.8 (82)	3,257	14.3 (98)
4351	469	3.4 (36)	63.5 (105)	304	21.1 (107)	46.7 (10)	34	*	43	*	469	10.7 (123)	469	16.8 (128)
4352	148	10.1 (141)	61.5 (76)	112	19.6 (90)	50.9 (37)	10	*	10	*	152	11.8 (136)	153	15.7 (115)
4353	340	2.9 (23)	68.8 (138)	245	18.0 (61)	53.5 (69)	32	*	31	*	388	10.8 (126)	387	15.2 (107)
4354	521	3.8 (57)	63.9 (109)	214	18.7 (74)	50.5 (30)	46	*	64	*	691	7.5 (47)	690	17.1 (129)
4355	164	4.3 (79)	55.5 (16)	67	*	*	24	*	39	*	283	9.9 (109)	284	12.7 (73)
4356	360	1.9 (8)	60.6 (54)	258	16.3 (36)	54.7 (83)	37	*	38	*	366	11.5 (133)	366	15.3 (110)
4358	48	*	*	38	*	*	8	*	6	*	56	*	56	*
4359	494	3.8 (57)	64.0 (111)	367	21.8 (110)	46.9 (11)	35	*	49	*	523	9.8 (106)	523	15.5 (113)
4360	195	4.1 (73)	62.6 (93)	144	24.3 (116)	49.3 (17)	11	*	17	*	184	4.9 (4)	184	13.6 (86)
4361	237	0.8 (2)	58.6 (33)	144	23.6 (113)	50.7 (34)	13	*	28	*	249	10.0 (113)	249	11.2 (43)
4362	232	4.3 (79)	57.8 (29)	172	13.4 (7)	63.4 (119)	22	*	35	*	271	10.3 (120)	272	12.5 (70)
4363	641	5.5 (125)	57.6 (28)	411	20.2 (100)	49.4 (18)	54	*	56	*	702	9.5 (99)	702	12.8 (75)
4401	849	3.8 (57)	56.7 (20)	737	17.5 (57)	52.1 (53)	73	*	96	*	801	4.9 (4)	802	9.6 (19)
4501	631	3.6 (43)	63.4 (104)	471	19.3 (84)	53.7 (72)	61	*	73	*	787	10.5 (122)	788	14.0 (92)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 90

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4601	939	3.1 (27)	61.0 (64)	459	15.3 (18)	50.1 (23)	118	52.5 (31)	127	40.9 (37)	1,136	9.5 (99)	1,138	13.4 (83)
4602	1	*	*	0	*	*	0	*	0	*	1	*	1	*
4701	972	4.4 (86)	54.6 (13)	661	18.9 (79)	51.7 (49)	71	*	137	12.4 (2)	933	10.7 (123)	934	18.0 (136)
5050	1,472	3.7 (50)	64.8 (119)	1,240	18.1 (64)	53.5 (69)	147	46.9 (23)	157	28.7 (15)	1,439	8.1 (63)	1,440	13.0 (77)
5051	2,513	3.7 (50)	61.1 (67)	2,080	18.2 (66)	53.8 (74)	247	43.3 (13)	314	27.7 (14)	2,493	8.1 (63)	2,494	9.7 (20)
5052	2,581	3.5 (41)	61.0 (64)	2,108	19.7 (93)	50.1 (23)	239	37.2 (4)	268	27.6 (13)	2,592	10.2 (118)	2,593	10.9 (36)
5055	563	2.1 (11)	59.1 (39)	209	18.7 (74)	56.0 (92)	59	*	87	*	1,194	10.7 (123)	1,194	3.9 (2)
5101	1,257	4.5 (90)	57.5 (26)	810	14.9 (16)	56.8 (101)	140	34.3 (3)	187	20.9 (5)	1,480	9.9 (109)	1,480	12.7 (73)
5102	26	*	*	16	*	*	4	*	5	*	26	*	26	*
5202	645	3.4 (36)	61.1 (67)	456	16.4 (38)	56.6 (98)	55	*	69	*	640	7.2 (38)	646	13.3 (80)
5206	5	*	*	4	*	*	0	*	0	*	9	*	9	*
5601	768	3.1 (27)	63.3 (102)	139	9.4 (1)	56.1 (94)	69	*	135	15.6 (3)	899	9.6 (103)	898	11.6 (51)
5801	1,595	4.3 (79)	63.1 (99)	1,155	18.0 (61)	52.9 (61)	150	48.7 (25)	137	32.1 (24)	1,656	8.0 (57)	1,657	17.3 (131)
5803	87	*	*	61	*	*	6	*	17	*	85	*	85	*
5804	23	*	*	18	*	*	1	*	4	*	22	*	22	*
5806	27	*	*	16	*	*	0	*	1	*	44	*	45	*
6101	39	*	*	28	*	*	1	*	1	*	35	*	36	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Maternal Health Indicators by Clinic Combined 3 Years

Page : 91

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6102	98	*	*	77	*	*	9	*	11	*	99	*	99	*
6103	3,083	3.8 (57)	63.8 (108)	2,266	18.4 (72)	51.5 (43)	196	46.9 (23)	172	20.9 (5)	3,168	9.5 (99)	3,171	17.9 (134)
6105	9	*	*	7	*	*	0	*	0	*	10	*	10	*
6106	110	3.6 (43)	53.6 (10)	77	*	*	14	*	10	*	105	8.6 (78)	105	21.0 (142)
6301	2,873	3.4 (36)	60.4 (51)	1,787	18.9 (79)	50.6 (33)	285	43.5 (14)	270	31.5 (21)	3,098	6.1 (19)	3,098	11.6 (51)
6312	1,578	4.2 (75)	59.3 (44)	1,007	16.7 (43)	53.2 (65)	151	51.0 (29)	183	33.9 (30)	1,637	5.3 (7)	1,637	9.2 (17)
6316	1,034	4.5 (90)	58.8 (34)	678	17.1 (51)	55.8 (89)	120	45.8 (18)	108	33.3 (27)	1,064	6.1 (19)	1,064	10.0 (24)
6318	18	*	*	8	*	*	8	*	3	*	33	*	33	*
6320	239	3.8 (57)	60.7 (56)	128	14.1 (12)	57.0 (104)	16	*	20	*	312	12.5 (138)	312	15.1 (106)
6325	228	5.7 (130)	52.6 (4)	114	12.3 (5)	64.9 (120)	27	*	42	*	271	7.4 (43)	271	16.6 (125)
6335	614	6.2 (133)	55.2 (15)	249	14.5 (13)	48.6 (14)	75	*	89	*	948	11.2 (131)	948	10.4 (30)
6340	19	*	*	6	*	*	1	*	4	*	14	*	14	*
6345	78	*	*	47	*	*	10	*	15	*	95	*	95	*
6360	705	3.8 (57)	58.4 (32)	325	16.9 (47)	53.8 (74)	60	*	69	*	922	7.2 (38)	922	14.0 (92)
7304	10	*	*	8	*	*	1	*	3	*	10	*	10	*
7306	1,751	3.2 (31)	60.4 (51)	1,169	16.2 (34)	52.7 (59)	192	45.8 (18)	161	29.8 (16)	2,109	8.4 (70)	2,110	11.0 (39)
7308	9	*	*	7	*	*	0	*	0	*	8	*	8	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 92

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7310	1	*	*	0	*	*	0	*	2	*	2	*	2	*
7401	1,925	4.7 (100)	58.8 (34)	1,615	17.0 (48)	55.2 (88)	155	39.4 (9)	151	38.4 (35)	1,857	7.8 (52)	1,860	14.7 (101)
7601	109	8.3 (139)	53.2 (6)	41	*	*	49	*	43	*	523	7.1 (35)	523	9.8 (21)
7850	979	4.4 (86)	63.7 (107)	804	19.8 (94)	49.6 (19)	77	*	94	*	966	8.4 (70)	968	11.6 (51)
7852	17	*	*	13	*	*	0	*	1	*	17	*	17	*
8101	1,401	3.8 (57)	65.9 (128)	919	19.8 (94)	52.8 (60)	139	61.9 (38)	97	*	1,595	7.3 (41)	1,596	17.9 (134)
8102	343	2.6 (15)	67.6 (136)	174	17.8 (59)	58.0 (113)	34	*	17	*	455	7.7 (50)	456	11.2 (43)
8105	688	3.6 (43)	65.0 (120)	363	16.3 (36)	51.0 (39)	135	64.4 (39)	114	45.6 (39)	680	6.5 (26)	679	9.1 (15)
8106	268	5.2 (117)	52.6 (4)	138	16.7 (43)	57.2 (107)	47	*	42	*	517	5.4 (10)	516	14.9 (103)
8150	16	*	*	8	*	*	4	*	6	*	17	*	17	*
8151	2,563	3.7 (50)	61.0 (64)	1,657	19.6 (90)	49.7 (20)	275	46.2 (21)	381	32.8 (26)	2,846	9.0 (88)	2,848	13.7 (88)
8152	26	*	*	17	*	*	4	*	5	*	27	*	27	*
8153	7	*	*	2	*	*	1	*	2	*	5	*	5	*
8155	9	*	*	5	*	*	1	*	4	*	11	*	11	*
8157	6	*	*	4	*	*	1	*	3	*	5	*	5	*
8201	1,825	3.8 (57)	60.3 (48)	1,171	15.5 (22)	56.4 (96)	190	48.9 (26)	295	31.9 (23)	2,403	8.0 (57)	2,404	10.3 (28)
8202	2,020	4.8 (101)	51.9 (3)	960	20.8 (105)	45.8 (8)	306	46.7 (22)	290	33.4 (28)	3,736	5.4 (10)	3,734	2.7 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 93

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8203	347	3.7 (50)	62.8 (95)	253	11.9 (4)	54.9 (85)	41	*	40	*	359	8.6 (78)	358	16.5 (124)
8250	526	5.5 (125)	61.4 (72)	349	18.1 (64)	57.9 (112)	81	*	82	*	582	7.7 (50)	581	13.6 (86)
8252	119	5.0 (104)	63.9 (109)	55	*	*	24	*	17	*	153	7.8 (52)	153	10.5 (32)
8255	1,904	4.6 (96)	60.7 (56)	1,300	16.0 (30)	54.7 (83)	253	49.0 (27)	309	36.2 (32)	2,091	8.1 (63)	2,091	12.3 (60)
8257	946	4.2 (75)	54.2 (11)	531	28.1 (118)	35.6 (3)	144	52.8 (32)	198	36.9 (33)	1,840	8.8 (82)	1,841	4.5 (3)
8263	548	4.6 (96)	60.6 (54)	383	15.7 (24)	59.3 (116)	65	*	104	24.0 (7)	559	8.4 (70)	558	9.5 (18)
8264	545	3.3 (33)	64.4 (118)	349	16.0 (30)	58.2 (114)	76	*	92	*	694	5.9 (17)	693	10.8 (35)
8267	450	5.3 (121)	62.4 (88)	321	17.4 (55)	53.3 (67)	72	*	84	*	624	5.9 (17)	623	9.1 (15)
8901	789	2.8 (19)	63.2 (100)	518	19.3 (84)	50.8 (35)	89	*	63	*	868	5.4 (10)	870	14.1 (95)
8902	957	4.3 (79)	61.3 (70)	500	18.8 (76)	53.0 (63)	119	55.5 (34)	98	*	1,294	7.0 (34)	1,294	12.3 (60)
8903	1,133	2.1 (11)	65.3 (122)	670	14.5 (13)	60.0 (117)	111	65.8 (41)	76	*	1,392	6.5 (26)	1,393	17.7 (133)
8904	1,837	3.5 (41)	64.1 (112)	1,028	16.6 (41)	55.8 (89)	198	61.6 (37)	135	54.8 (40)	2,135	7.4 (43)	2,139	14.9 (103)
8905	1,211	3.4 (36)	60.9 (60)	609	19.4 (87)	51.4 (42)	120	50.0 (28)	102	44.1 (38)	1,663	10.9 (127)	1,665	12.3 (60)
8909	669	4.2 (75)	62.5 (90)	333	16.2 (34)	57.4 (110)	53	*	39	*	859	9.2 (89)	860	14.4 (99)
9045	1,071	1.7 (5)	66.8 (132)	659	19.6 (90)	46.6 (9)	131	45.0 (17)	95	*	1,468	8.5 (74)	1,468	8.2 (8)
9050	763	5.0 (104)	60.9 (60)	205	16.6 (41)	57.1 (105)	88	*	50	*	910	6.8 (30)	910	11.1 (42)
9051	904	2.8 (19)	65.8 (126)	496	19.4 (87)	43.5 (6)	103	29.1 (2)	61	*	1,428	9.4 (95)	1,427	8.7 (12)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 94

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9053	1,078	3.8 (57)	65.3 (122)	740	15.3 (18)	57.6 (111)	73	*	51	*	1,129	6.4 (24)	1,130	13.9 (90)
9055	12	*	*	6	*	*	0	*	0	*	12	*	12	*
9058	697	5.7 (130)	62.3 (85)	439	18.2 (66)	52.6 (57)	96	*	58	*	1,083	8.9 (87)	1,084	14.6 (100)
9066	769	3.3 (33)	61.4 (72)	407	20.1 (98)	52.1 (53)	103	44.7 (16)	61	*	1,066	5.8 (16)	1,070	19.4 (140)
9068	844	2.0 (9)	65.2 (121)	586	21.2 (108)	56.8 (101)	64	*	36	*	850	5.3 (7)	852	13.4 (83)
9069	1,730	3.4 (36)	61.9 (80)	1,137	20.1 (98)	49.7 (20)	220	57.3 (35)	104	39.4 (36)	2,136	6.1 (19)	2,134	11.3 (46)
9070	1,342	4.3 (79)	61.5 (76)	863	17.4 (55)	56.0 (92)	123	65.0 (40)	79	*	1,409	5.7 (14)	1,411	13.3 (80)
9072	732	4.4 (86)	64.3 (116)	378	19.0 (82)	53.7 (72)	121	55.4 (33)	85	*	860	7.8 (52)	861	16.7 (126)
9080	1,706	2.8 (19)	62.8 (95)	915	15.4 (20)	56.9 (103)	245	57.6 (36)	161	34.2 (31)	2,450	8.0 (57)	2,448	11.8 (55)
9082	115	2.6 (15)	67.0 (134)	63	*	*	11	*	7	*	124	7.3 (41)	124	11.3 (46)
9083	66	*	*	61	*	*	0	*	0	*	70	*	70	*
9084	16	*	*	10	*	*	3	*	1	*	21	*	21	*
9085	4	*	*	2	*	*	0	*	1	*	2	*	2	*
9086	12	*	*	5	*	*	0	*	4	*	10	*	10	*
9087	13	*	*	8	*	*	0	*	1	*	11	*	11	*
Michigan	116,274	4.0	61.2	70,748	18.3	52.3	12,540	43.2	14,527	30.1	140,267	8.2	140,379	12.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 95

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	97	*	*	107	23.4 (115)	98	*	77	*
0102	379	85.5 (82)	7.1 (97)	416	34.1 (56)	394	22.8 (74)	340	90.0 (73)
0103	346	82.4 (109)	9.5 (110)	389	33.4 (60)	356	18.3 (105)	303	92.1 (45)
0104	112	85.7 (81)	7.1 (97)	121	40.5 (16)	109	22.9 (72)	95	*
0105	16	*	*	15	*	14	*	15	*
0250	103	81.6 (117)	13.6 (133)	123	52.0 (2)	100	34.0 (17)	105	93.3 (30)
0251	79	*	*	91	*	83	*	73	*
0252	89	*	*	96	*	85	*	69	*
0253	114	83.3 (97)	7.0 (96)	121	48.8 (4)	115	21.7 (80)	108	91.7 (51)
0401	403	84.1 (92)	10.7 (120)	444	39.6 (20)	418	23.7 (67)	361	93.6 (27)
0402	339	79.1 (125)	11.8 (128)	373	35.9 (40)	349	26.1 (53)	304	91.8 (47)
0403	136	85.3 (84)	1.5 (9)	147	27.2 (99)	131	23.7 (67)	108	97.2 (2)
0404	95	*	*	99	*	91	*	75	*
0501	438	94.1 (8)	2.3 (20)	457	33.3 (61)	410	27.3 (42)	345	92.5 (40)
0503	86	*	*	87	*	78	*	68	*
0504	381	89.2 (51)	4.5 (67)	405	29.4 (87)	386	34.2 (14)	313	90.7 (64)
0505	234	88.0 (59)	6.0 (87)	244	37.3 (31)	232	27.2 (44)	200	85.5 (120)
0506	150	82.7 (104)	6.0 (87)	159	31.4 (75)	149	24.2 (65)	122	83.6 (131)
0507	77	*	*	78	*	70	*	50	*
0508	24	*	*	25	*	23	*	16	*
0509	1	*	*	1	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 96

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	951	86.6 (74)	7.3 (99)	1,048	34.7 (50)	957	29.4 (31)	814	91.8 (47)
0651	524	82.3 (110)	10.1 (115)	572	43.9 (7)	498	26.5 (48)	475	87.8 (99)
0652	308	87.3 (67)	6.2 (92)	321	34.9 (49)	294	25.5 (58)	248	84.7 (125)
0653	364	72.8 (138)	19.2 (142)	395	40.5 (16)	374	30.7 (27)	313	87.9 (96)
0654	336	80.1 (122)	13.4 (132)	379	42.7 (9)	353	21.8 (79)	298	84.6 (127)
0655	280	89.3 (50)	3.2 (40)	315	41.0 (15)	283	26.9 (47)	257	91.4 (53)
0657	64	*	*	65	*	59	*	54	*
0701	398	83.4 (96)	9.5 (110)	456	34.0 (57)	419	38.2 (11)	348	95.4 (9)
0702	65	*	*	67	*	63	*	47	*
0703	145	91.0 (39)	2.8 (34)	162	31.5 (72)	143	30.8 (26)	112	99.1 (1)
0704	48	*	*	54	*	53	*	45	*
0705	79	*	*	86	*	82	*	78	*
0706	10	*	*	10	*	9	*	7	*
0850	651	87.3 (67)	3.5 (45)	714	27.3 (98)	651	28.9 (32)	512	92.6 (36)
0851	565	91.2 (35)	4.2 (60)	624	29.5 (86)	582	28.5 (36)	449	96.7 (5)
0901	1,362	78.2 (127)	14.5 (134)	1,511	39.5 (21)	1,417	18.1 (106)	1,219	89.1 (80)
0902	35	*	*	41	*	40	*	34	*
1051	1,049	82.8 (103)	10.5 (118)	1,156	28.7 (88)	1,060	25.8 (56)	813	89.3 (79)
1052	1	*	*	1	*	1	*	1	*
1101	1,467	73.4 (137)	9.4 (107)	1,640	27.6 (93)	1,463	16.5 (111)	1,208	87.4 (104)
1102	682	78.6 (126)	11.0 (124)	822	34.4 (52)	752	21.4 (81)	602	94.0 (21)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 97

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1104	1	*	*	1	*	1	*	0	*
1106	58	*	*	72	*	69	*	53	*
1250	815	82.6 (107)	8.6 (106)	991	31.8 (70)	863	27.0 (46)	745	86.0 (116)
1251	608	85.2 (86)	7.7 (103)	667	32.4 (69)	592	24.8 (63)	501	90.2 (71)
1252	447	66.9 (141)	17.9 (141)	528	36.9 (33)	499	20.8 (88)	423	88.7 (86)
1253	589	71.0 (139)	17.3 (139)	720	36.4 (38)	662	19.9 (96)	562	88.6 (87)
1350	2,075	92.9 (18)	3.5 (45)	2,304	33.0 (65)	1,804	27.7 (41)	1,695	85.5 (120)
1351	71	*	*	78	*	74	*	63	*
1352	354	95.2 (5)	3.7 (53)	383	32.6 (68)	345	3.2 (143)	289	89.6 (77)
1401	201	87.1 (69)	8.5 (105)	222	33.3 (61)	202	33.2 (20)	176	90.9 (60)
1402	98	*	*	109	39.4 (22)	91	*	80	*
1403	18	*	*	18	*	17	*	12	*
1501	665	82.3 (110)	10.7 (120)	776	37.5 (28)	570	16.0 (112)	608	83.2 (132)
1503	809	86.7 (72)	7.3 (99)	882	42.3 (10)	766	25.1 (61)	694	79.4 (139)
1504	1,766	92.2 (26)	2.4 (25)	1,963	30.7 (78)	1,757	27.8 (40)	1,418	84.7 (125)
1505	251	88.8 (52)	8.0 (104)	283	33.6 (58)	219	25.6 (57)	229	92.1 (45)
1506	492	77.0 (131)	16.7 (137)	574	38.0 (26)	523	6.7 (141)	459	93.7 (26)
1507	495	92.9 (18)	3.6 (49)	545	35.2 (47)	468	25.0 (62)	394	92.4 (43)
1508	18	*	*	21	*	20	*	21	*
1509	355	92.7 (22)	2.5 (27)	412	35.9 (40)	351	20.8 (88)	319	91.8 (47)
1511	563	93.1 (13)	1.8 (14)	612	35.6 (45)	551	19.1 (101)	469	88.9 (85)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 98

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1512	313	94.9 (6)	2.9 (37)	342	25.4 (104)	290	33.4 (18)	216	92.6 (36)
1514	396	92.9 (18)	3.5 (45)	418	20.8 (124)	314	30.9 (24)	224	95.1 (10)
1701	525	92.4 (24)	4.8 (73)	578	41.2 (14)	523	40.5 (7)	467	86.5 (112)
1950	560	82.7 (104)	11.1 (125)	600	34.2 (55)	515	26.4 (52)	463	94.8 (13)
1952	122	86.1 (78)	9.8 (113)	129	28.7 (88)	110	27.3 (42)	93	*
1953	257	83.3 (97)	10.5 (118)	280	33.6 (58)	249	26.5 (48)	222	96.8 (3)
1954	406	84.0 (94)	9.4 (107)	437	36.8 (35)	370	28.6 (35)	351	94.3 (18)
1955	97	*	*	104	41.3 (13)	90	*	79	*
1957	360	88.3 (55)	5.3 (81)	386	30.6 (79)	298	34.2 (14)	279	92.5 (40)
1963	60	*	*	62	*	48	*	45	*
1964	15	*	*	17	*	17	*	17	*
2150	571	93.2 (12)	1.2 (5)	624	35.9 (40)	579	25.9 (54)	489	94.3 (18)
2151	177	85.9 (79)	10.7 (120)	219	30.6 (79)	170	31.2 (23)	148	93.9 (24)
2152	16	*	*	16	*	14	*	12	*
2251	360	91.7 (31)	5.3 (81)	400	34.5 (51)	373	32.4 (21)	315	93.3 (30)
2252	145	86.9 (70)	9.7 (112)	169	40.2 (18)	145	55.9 (3)	136	89.0 (82)
2554	47	*	*	48	*	38	*	38	*
2555	6,630	87.7 (62)	3.1 (39)	7,268	25.4 (104)	5,608	8.8 (136)	5,268	89.0 (82)
2556	257	81.3 (119)	4.7 (72)	283	21.2 (121)	192	10.4 (133)	178	86.5 (112)
2557	2	*	*	2	*	1	*	1	*
2558	3	*	*	4	*	3	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 99

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2560	3	*	*	5	*	5	*	5	*
2561	3	*	*	3	*	3	*	3	*
2562	15	*	*	15	*	12	*	14	*
2563	24	*	*	25	*	17	*	12	*
2564	18	*	*	22	*	19	*	17	*
3001	1,063	82.9 (101)	6.1 (89)	1,144	35.0 (48)	1,034	13.0 (122)	925	84.8 (124)
3002	234	90.6 (42)	1.3 (7)	261	44.1 (6)	241	12.4 (124)	223	87.4 (104)
3004	225	91.1 (37)	3.6 (49)	244	43.0 (8)	231	22.5 (76)	190	83.2 (132)
3005	107	89.7 (48)	2.8 (34)	114	33.3 (61)	111	19.8 (97)	93	*
3007	11	*	*	12	*	11	*	9	*
3201	389	84.6 (90)	5.4 (83)	415	27.7 (92)	363	33.3 (19)	293	90.1 (72)
3203	1	*	*	1	*	1	*	1	*
3301	3,922	88.0 (59)	4.8 (73)	4,276	27.6 (93)	3,826	8.1 (140)	3,167	89.8 (74)
3304	88	*	*	97	*	95	*	71	*
3305	137	91.2 (35)	6.6 (94)	155	29.7 (83)	143	10.5 (130)	107	89.7 (75)
3306	217	90.8 (40)	4.6 (69)	239	30.1 (82)	209	10.5 (130)	171	93.6 (27)
3308	84	*	*	96	*	82	*	56	*
3309	7	*	*	8	*	8	*	8	*
3312	370	88.6 (53)	4.3 (61)	402	31.3 (76)	356	9.3 (135)	304	91.1 (59)
3550	56	*	*	61	*	58	*	51	*
3551	13	*	*	14	*	14	*	13	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 100

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
3552	30	*	*	33	*	31	*	30 *
3701	869	93.0 (15)	3.8 (54)	944	30.6 (79)	859	21.0 (87)	671 94.0 (21)
3850	2,630	85.8 (80)	7.4 (102)	2,903	38.4 (24)	2,618	20.6 (90)	2,289 87.4 (104)
3901	1,408	87.6 (64)	4.5 (67)	1,605	38.4 (24)	1,483	15.3 (114)	1,279 83.7 (129)
3902	99	*	*	122	50.0 (3)	117	20.5 (93)	108 88.0 (92)
3903	25	*	*	25	*	24	*	17 *
3904	4	*	*	4	*	4	*	4 *
3905	9	*	*	9	*	9	*	9 *
3950	1,932	86.5 (76)	6.3 (93)	2,165	32.7 (66)	1,949	18.8 (102)	1,589 87.4 (104)
3954	14	*	*	15	*	14	*	12 *
4103	1	*	*	1	*	1	*	1 *
4106	2,067	86.6 (74)	4.0 (56)	2,284	23.8 (113)	1,948	8.8 (136)	1,710 86.5 (112)
4107	2,636	87.4 (66)	4.4 (64)	2,886	21.0 (122)	2,512	26.5 (48)	2,030 91.4 (53)
4109	682	82.1 (112)	10.7 (120)	749	31.5 (72)	693	20.6 (90)	536 95.0 (12)
4110	7	*	*	7	*	7	*	0 *
4112	513	45.4 (142)	0.0 (1)	542	25.8 (103)	427	73.1 (2)	488 92.6 (36)
4115	3,121	90.7 (41)	4.3 (61)	3,485	21.7 (120)	2,825	16.6 (110)	2,517 92.4 (43)
4351	446	89.5 (49)	3.6 (49)	500	41.8 (12)	259	30.9 (24)	405 91.4 (53)
4352	150	69.3 (140)	14.7 (135)	162	42.0 (11)	151	20.5 (93)	133 82.7 (134)
4353	356	75.0 (134)	17.7 (140)	426	34.3 (53)	385	25.5 (58)	316 82.3 (136)
4354	669	92.1 (27)	4.3 (61)	729	37.9 (27)	617	14.3 (119)	571 90.5 (68)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 101

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4355	283	84.8 (89)	12.0 (129)	297	34.3 (53)	271	80.1 (1)	229	91.3 (56)
4356	351	82.1 (112)	11.1 (125)	394	36.3 (39)	355	25.4 (60)	312	90.7 (64)
4358	57	*	*	59	*	58	*	54	*
4359	475	74.3 (136)	16.2 (136)	554	37.4 (30)	490	21.4 (81)	436	90.8 (61)
4360	180	85.0 (88)	9.4 (107)	200	39.0 (23)	193	28.5 (36)	164	89.0 (82)
4361	244	90.2 (44)	4.1 (58)	269	35.7 (44)	251	28.3 (38)	211	90.5 (68)
4362	267	82.0 (115)	12.4 (130)	280	37.5 (28)	251	25.9 (54)	215	87.0 (110)
4363	669	77.4 (130)	13.3 (131)	736	37.2 (32)	661	27.2 (44)	572	86.9 (111)
4401	781	83.2 (99)	10.0 (114)	854	29.7 (83)	785	21.4 (81)	633	94.6 (14)
4501	772	82.0 (115)	10.4 (117)	819	33.1 (64)	748	28.1 (39)	604	93.9 (24)
4601	1,116	92.4 (24)	2.4 (25)	1,246	26.3 (100)	1,123	21.3 (84)	921	91.2 (58)
4602	1	*	*	1	*	1	*	1	*
4701	910	88.2 (57)	5.2 (78)	1,007	25.3 (107)	839	39.8 (9)	692	92.9 (34)
5050	1,429	93.0 (15)	1.7 (11)	1,515	19.7 (134)	1,115	14.1 (120)	1,022	95.7 (7)
5051	2,468	93.1 (13)	1.5 (9)	2,651	20.5 (128)	2,023	12.7 (123)	1,775	94.5 (15)
5052	2,556	91.7 (31)	2.3 (20)	2,720	22.7 (118)	1,939	10.5 (130)	1,942	94.0 (21)
5055	1,151	88.0 (59)	11.2 (127)	1,210	46.3 (5)	1,025	6.7 (141)	1,022	89.1 (80)
5101	1,443	91.1 (37)	3.4 (44)	1,578	24.3 (110)	817	34.3 (13)	1,130	94.4 (16)
5102	26	*	*	26	*	21	*	17	*
5202	620	89.8 (47)	5.2 (78)	711	35.6 (45)	538	34.2 (14)	579	90.7 (64)
5206	8	*	*	7	*	4	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 102

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5601	880	92.5 (23)	2.2 (19)	954	26.0 (102)	885	28.9 (32)	687	95.1 (10)
5801	1,639	86.3 (77)	5.5 (84)	1,784	28.1 (91)	1,646	24.1 (66)	1,312	90.8 (61)
5803	84	*	*	92	*	84	*	66	*
5804	21	*	*	24	*	22	*	15	*
5806	44	*	*	49	*	40	*	33	*
6101	32	*	*	38	*	36	*	31	*
6102	92	*	*	104	32.7 (66)	88	*	85	*
6103	3,063	88.2 (57)	4.4 (64)	3,353	36.9 (33)	2,710	21.3 (84)	2,550	86.2 (115)
6105	9	*	*	11	*	10	*	7	*
6106	101	90.1 (46)	4.0 (56)	116	53.4 (1)	110	14.5 (118)	109	82.6 (135)
6301	3,031	87.6 (64)	5.9 (86)	3,381	26.2 (101)	2,771	12.2 (126)	2,450	87.9 (96)
6312	1,600	91.9 (28)	3.5 (45)	1,773	24.3 (110)	1,415	21.1 (86)	1,250	88.0 (92)
6316	1,059	91.9 (28)	2.7 (32)	1,154	23.8 (113)	955	22.6 (75)	805	93.5 (29)
6318	33	*	*	36	*	27	*	20	*
6320	303	95.4 (4)	2.6 (29)	345	19.7 (134)	221	35.3 (12)	232	94.4 (16)
6325	267	93.6 (10)	2.6 (29)	307	24.8 (108)	271	22.1 (78)	199	95.5 (8)
6335	929	91.4 (33)	6.7 (95)	983	36.6 (37)	356	29.5 (29)	766	92.8 (35)
6340	14	*	*	21	*	19	*	21	*
6345	93	*	*	105	12.4 (148)	87	*	67	*
6360	887	88.3 (55)	5.0 (75)	1,009	27.4 (96)	742	24.7 (64)	747	90.8 (61)
7304	10	*	*	10	*	10	*	10	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 103

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7306	2,052	85.1 (87)	5.1 (77)	2,253	31.6 (71)	2,022	20.6 (90)	1,707	88.5 (88)
7308	8	*	*	10	*	10	*	10	*
7310	2	*	*	2	*	2	*	2	*
7401	1,761	75.2 (133)	17.2 (138)	2,018	40.1 (19)	1,769	23.1 (70)	1,631	85.7 (119)
7601	519	91.3 (34)	5.0 (75)	583	36.7 (36)	544	44.1 (6)	472	93.0 (33)
7850	951	85.4 (83)	10.3 (116)	1,058	31.5 (72)	670	23.1 (70)	782	85.8 (118)
7852	17	*	*	18	*	12	*	15	*
8101	1,565	79.4 (124)	2.9 (37)	1,640	20.2 (129)	1,069	17.6 (107)	1,117	87.4 (104)
8102	441	83.9 (95)	3.9 (55)	472	25.4 (104)	334	20.1 (95)	348	83.9 (128)
8105	684	93.9 (9)	1.2 (5)	766	27.5 (95)	707	15.4 (113)	605	92.6 (36)
8106	503	80.7 (121)	3.6 (49)	541	17.2 (144)	439	15.0 (115)	333	90.4 (70)
8150	16	*	*	18	*	18	*	17	*
8151	2,794	91.8 (30)	3.3 (42)	3,070	29.6 (85)	2,806	15.0 (115)	2,290	93.2 (32)
8152	26	*	*	32	*	21	*	30	*
8153	5	*	*	6	*	6	*	5	*
8155	11	*	*	12	*	11	*	12	*
8157	5	*	*	6	*	5	*	5	*
8201	2,382	93.5 (11)	1.4 (8)	2,653	19.0 (139)	1,841	40.4 (8)	1,700	89.6 (77)
8202	3,658	95.6 (3)	2.3 (20)	3,823	28.6 (90)	2,431	8.3 (139)	2,939	90.6 (67)
8203	355	93.0 (15)	2.3 (20)	402	20.1 (132)	311	28.9 (32)	287	96.2 (6)
8250	577	88.6 (53)	1.7 (11)	656	16.6 (145)	501	26.5 (48)	458	88.0 (92)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 104

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8252	148	75.0 (134)	6.1 (89)	164	20.7 (126)	149	11.4 (128)	129	91.5 (52)
8255	2,057	92.8 (21)	0.7 (2)	2,306	18.3 (141)	1,441	38.4 (10)	1,554	91.8 (47)
8257	1,820	97.2 (1)	0.8 (3)	1,896	27.4 (96)	1,425	18.6 (103)	1,466	83.7 (129)
8263	552	96.2 (2)	0.9 (4)	635	20.0 (133)	332	44.9 (5)	469	96.8 (3)
8264	688	94.8 (7)	1.7 (11)	769	19.4 (136)	586	30.2 (28)	507	94.1 (20)
8267	614	90.2 (44)	2.6 (29)	686	18.7 (140)	552	51.1 (4)	496	91.3 (56)
8901	862	82.9 (101)	3.2 (40)	921	20.2 (129)	525	19.2 (100)	626	81.9 (137)
8902	1,283	86.9 (70)	1.8 (14)	1,386	20.8 (124)	857	22.9 (72)	980	88.2 (91)
8903	1,363	79.8 (123)	2.8 (34)	1,469	22.4 (119)	1,088	19.3 (99)	1,020	80.7 (138)
8904	2,103	83.0 (100)	2.3 (20)	2,250	19.2 (138)	1,484	18.5 (104)	1,522	87.8 (99)
8905	1,606	90.6 (42)	2.1 (18)	1,731	24.1 (112)	1,335	32.1 (22)	1,194	87.4 (104)
8909	850	86.7 (72)	1.9 (17)	916	20.6 (127)	683	17.6 (107)	640	89.7 (75)
9045	1,454	82.6 (107)	7.3 (99)	1,558	31.3 (76)	655	29.5 (29)	1,212	87.8 (99)
9050	857	81.0 (120)	4.6 (69)	948	24.5 (109)	789	23.3 (69)	685	85.5 (120)
9051	1,394	84.1 (92)	5.6 (85)	1,482	35.9 (40)	1,360	12.4 (124)	1,203	87.7 (102)
9053	1,117	85.3 (84)	1.8 (14)	1,180	15.5 (146)	954	13.2 (121)	666	92.5 (40)
9055	12	*	*	11	*	4	*	1	*
9058	1,065	84.2 (91)	4.4 (64)	1,131	21.0 (122)	887	8.7 (138)	708	87.6 (103)
9066	1,044	76.3 (132)	5.2 (78)	1,116	18.1 (142)	852	11.2 (129)	718	88.4 (89)
9068	834	81.5 (118)	4.1 (58)	890	22.8 (117)	741	14.7 (117)	634	86.0 (116)
9069	2,135	82.1 (112)	2.7 (32)	2,203	17.4 (143)	1,257	10.0 (134)	1,337	88.3 (90)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 105

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9070	1,398	78.0 (128)	4.6 (69)	1,498	19.3 (137)	1,053	17.4 (109)	964	88.0 (92)
9072	852	77.7 (129)	6.1 (89)	928	23.1 (116)	799	11.8 (127)	766	85.4 (123)
9080	2,391	82.7 (104)	3.3 (42)	2,554	20.2 (129)	1,769	19.5 (98)	1,796	87.9 (96)
9082	122	87.7 (62)	2.5 (27)	129	13.2 (147)	116	22.4 (77)	87	*
9083	71	*	*	70	*	50	*	5	*
9084	21	*	*	23	*	15	*	13	*
9085	2	*	*	4	*	4	*	4	*
9086	10	*	*	12	*	11	*	12	*
9087	11	*	*	13	*	13	*	12	*
Michigan	136,924	86.9	5.1	149,995	28.3	120,804	20.8	110,018	89.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 106

Michigan

Comparison of Smoking Indicators by Clinic Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	86	*	97	*	23	*	77	*	97	*
0102	364	40.4 (126)	390	29.5 (140)	123	19.5 (82)	340	16.8 (130)	391	3.1 (95)
0103	331	36.6 (114)	356	26.7 (132)	100	23.0 (80)	303	19.1 (134)	356	6.2 (138)
0104	104	35.6 (104)	113	24.8 (125)	29	*	95	*	113	2.7 (83)
0105	15	*	16	*	6	*	14	*	16	*
0250	110	44.5 (138)	109	23.9 (121)	41	*	105	14.3 (115)	107	0.9 (17)
0251	80	*	81	*	28	*	73	*	81	*
0252	83	*	89	*	17	*	68	*	89	*
0253	114	45.6 (140)	113	24.8 (125)	44	*	108	16.7 (129)	114	4.4 (127)
0401	395	45.6 (140)	415	28.0 (136)	147	28.6 (75)	361	12.2 (95)	414	2.2 (68)
0402	331	42.0 (131)	347	22.5 (119)	116	38.8 (46)	304	10.9 (82)	348	3.7 (108)
0403	123	42.3 (133)	133	27.8 (135)	40	*	108	18.5 (133)	135	5.2 (134)
0404	86	*	94	*	19	*	75	*	95	*
0501	390	37.2 (118)	440	26.6 (131)	122	26.2 (78)	345	19.7 (136)	440	3.0 (91)
0503	72	*	86	*	22	*	68	*	86	*
0504	367	37.1 (117)	384	19.0 (94)	109	35.8 (54)	313	8.6 (64)	383	2.6 (79)
0505	219	36.1 (109)	230	21.7 (113)	68	*	199	8.5 (61)	228	5.3 (136)
0506	133	39.8 (125)	153	26.8 (133)	44	*	122	22.1 (138)	153	6.5 (139)
0507	60	*	79	*	13	*	50	*	79	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 107

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0508	19	*	25	*	5	*	16	*	25	*
0509	1	*	1	*	1	*	1	*	1	*
0650	914	31.8 (87)	974	18.8 (93)	241	33.2 (61)	813	10.0 (76)	973	2.0 (64)
0651	509	44.6 (139)	537	28.7 (137)	192	26.6 (77)	475	15.6 (121)	536	2.4 (76)
0652	282	42.2 (132)	316	29.1 (139)	105	33.3 (60)	248	20.6 (137)	316	9.5 (142)
0653	350	34.9 (103)	374	21.4 (110)	98	*	313	14.7 (116)	375	3.7 (108)
0654	333	34.2 (98)	356	21.3 (108)	91	*	298	16.1 (124)	357	4.2 (122)
0655	277	44.0 (136)	291	28.9 (138)	97	*	257	17.1 (131)	291	3.4 (102)
0657	58	*	64	*	23	*	54	*	64	*
0701	387	33.1 (94)	411	20.9 (105)	106	34.9 (56)	348	5.5 (32)	411	1.5 (44)
0702	57	*	66	*	15	*	47	*	66	*
0703	129	44.2 (137)	150	30.7 (141)	47	*	112	19.6 (135)	150	4.0 (115)
0704	49	*	48	*	14	*	45	*	48	*
0705	81	*	79	*	29	*	78	*	79	*
0706	9	*	10	*	1	*	7	*	10	*
0850	612	39.1 (123)	660	19.8 (97)	183	41.5 (36)	512	6.3 (43)	660	2.6 (79)
0851	533	37.0 (116)	579	17.4 (88)	157	49.0 (20)	448	8.7 (66)	581	2.8 (86)
0901	1,332	35.7 (105)	1,413	20.9 (105)	413	29.1 (72)	1,218	12.2 (95)	1,417	3.9 (114)
0902	38	*	34	*	10	*	34	*	34	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic Combined 3 Years

Page : 108

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1051	981	35.8 (106)	1,083	16.3 (84)	286	44.1 (27)	813	7.6 (53)	1,089	2.4 (76)
1052	1	*	1	*	0	*	1	*	1	*
1101	1,377	25.6 (70)	1,526	12.1 (64)	282	42.9 (31)	1,208	12.1 (94)	1,528	4.0 (115)
1102	694	37.6 (121)	716	19.0 (94)	189	42.9 (31)	602	13.5 (106)	717	3.2 (98)
1104	1	*	1	*	0	*	0	*	1	*
1106	65	*	60	*	13	*	53	*	60	*
1250	852	27.7 (76)	928	15.0 (79)	182	33.5 (59)	741	10.3 (81)	930	1.3 (33)
1251	556	40.8 (127)	621	27.5 (134)	189	29.1 (72)	501	13.8 (112)	620	2.3 (71)
1252	464	29.3 (80)	478	15.1 (80)	108	37.0 (50)	422	11.1 (86)	478	3.3 (100)
1253	622	36.7 (115)	656	19.8 (97)	183	39.9 (40)	561	8.7 (66)	650	2.9 (89)
1350	1,929	26.9 (74)	2,112	13.9 (73)	418	31.6 (67)	1,709	6.8 (46)	2,137	1.5 (44)
1351	75	*	70	*	19	*	63	*	71	*
1352	320	25.6 (70)	355	12.7 (68)	75	*	287	2.1 (7)	359	0.6 (5)
1401	198	40.9 (128)	208	22.1 (115)	70	*	176	12.5 (101)	209	4.3 (125)
1402	92	*	103	12.6 (66)	23	*	81	*	103	1.0 (20)
1403	16	*	18	*	5	*	12	*	18	*
1501	665	20.8 (48)	698	8.9 (47)	114	34.2 (58)	603	4.1 (14)	708	1.1 (24)
1503	770	22.2 (55)	828	10.9 (57)	155	32.3 (66)	694	5.9 (40)	828	0.7 (11)
1504	1,648	16.4 (29)	1,799	8.4 (45)	223	45.7 (25)	1,420	2.4 (9)	1,801	0.6 (5)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 109

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1505	260	22.7 (60)	256	10.2 (54)	48	*	224	3.1 (11)	255	3.5 (103)
1506	504	21.8 (53)	524	12.6 (66)	90	*	458	4.4 (17)	529	1.1 (24)
1507	452	31.9 (88)	510	11.2 (59)	118	42.4 (34)	389	11.1 (86)	510	3.1 (95)
1508	21	*	18	*	1	*	21	*	18	*
1509	346	22.5 (58)	356	13.2 (71)	67	*	318	7.5 (51)	357	1.7 (53)
1511	528	35.8 (106)	573	18.0 (91)	158	39.2 (44)	470	5.5 (32)	578	2.2 (68)
1512	272	21.7 (52)	320	10.9 (57)	48	*	217	4.6 (18)	320	0.9 (17)
1514	326	12.6 (16)	411	7.1 (36)	25	*	217	0.0 (1)	404	1.0 (20)
1701	515	36.1 (109)	531	25.4 (129)	158	17.1 (83)	463	11.9 (93)	534	4.5 (129)
1950	518	41.7 (130)	564	24.8 (125)	178	33.1 (62)	463	16.6 (127)	571	4.0 (115)
1952	104	27.9 (77)	126	19.8 (97)	26	*	93	*	126	3.2 (98)
1953	244	37.3 (120)	262	20.2 (102)	75	*	222	12.2 (95)	263	3.0 (91)
1954	382	33.8 (96)	414	20.0 (101)	114	35.1 (55)	352	11.4 (89)	415	1.4 (41)
1955	89	*	97	*	30	*	79	*	98	*
1957	315	30.2 (83)	365	15.9 (82)	86	*	279	5.0 (22)	368	1.6 (49)
1963	51	*	60	*	11	*	45	*	62	*
1964	17	*	15	*	5	*	17	*	15	*
2150	546	42.9 (134)	574	24.2 (123)	189	37.6 (49)	489	12.3 (100)	574	3.0 (91)
2151	183	36.1 (109)	187	21.4 (110)	45	*	148	12.8 (104)	186	2.7 (83)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 110

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2152	14	*	16	*	8	*	12	*	16	*
2251	352	33.0 (93)	363	21.2 (107)	98	*	315	5.4 (29)	362	1.1 (24)
2252	142	49.3 (142)	146	37.7 (143)	60	*	135	16.3 (126)	145	4.8 (131)
2554	41	*	45	*	15	*	37	*	44	*
2555	6,054	23.7 (65)	6,757	13.2 (71)	1,141	37.7 (48)	5,258	8.6 (64)	6,764	2.2 (68)
2556	233	24.5 (66)	272	14.7 (77)	47	*	180	8.3 (58)	274	1.5 (44)
2557	1	*	2	*	1	*	1	*	2	*
2558	4	*	3	*	0	*	4	*	3	*
2560	5	*	3	*	0	*	5	*	3	*
2561	3	*	3	*	2	*	3	*	3	*
2562	14	*	15	*	1	*	14	*	15	*
2563	18	*	24	*	3	*	12	*	24	*
2564	20	*	18	*	5	*	17	*	18	*
3001	980	32.3 (90)	1,086	12.5 (65)	288	49.3 (19)	925	11.5 (91)	1,088	3.8 (111)
3002	231	26.4 (73)	243	11.9 (62)	56	*	222	11.3 (88)	242	2.1 (66)
3004	215	23.3 (63)	227	7.9 (41)	39	*	190	14.2 (114)	228	3.1 (95)
3005	105	32.4 (91)	110	14.5 (76)	33	*	93	*	110	3.6 (105)
3007	9	*	12	*	2	*	9	*	12	*
3201	353	38.8 (122)	400	22.5 (119)	116	31.0 (69)	293	14.7 (116)	400	4.5 (129)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 111

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3203	1	*	1	*	0	*	1	*	1	*
3301	3,628	17.2 (33)	4,074	9.5 (50)	521	39.0 (45)	3,151	5.4 (29)	4,067	2.6 (79)
3304	85	*	90	*	6	*	70	*	90	*
3305	133	14.3 (21)	142	6.3 (28)	11	*	107	1.9 (6)	141	0.0 (1)
3306	198	19.2 (43)	226	10.2 (54)	33	*	172	5.2 (25)	224	1.8 (56)
3308	77	*	88	*	13	*	56	*	88	*
3309	8	*	7	*	1	*	8	*	7	*
3312	338	19.5 (44)	391	9.0 (48)	59	*	304	3.6 (13)	390	1.0 (20)
3550	53	*	57	*	24	*	51	*	57	*
3551	14	*	13	*	7	*	13	*	13	*
3552	31	*	30	*	21	*	30	*	30	*
3701	792	32.7 (92)	877	20.3 (103)	216	27.3 (76)	671	7.9 (55)	882	1.9 (61)
3850	2,509	34.8 (102)	2,734	22.1 (115)	757	30.3 (70)	2,287	9.2 (72)	2,734	2.0 (64)
3901	1,396	23.2 (62)	1,444	11.4 (60)	265	36.6 (51)	1,275	6.3 (43)	1,441	1.4 (41)
3902	116	25.0 (69)	97	*	25	*	108	8.3 (58)	99	*
3903	23	*	25	*	4	*	17	*	25	*
3904	4	*	4	*	2	*	4	*	4	*
3905	9	*	9	*	2	*	9	*	8	*
3950	1,822	23.0 (61)	1,974	12.9 (69)	332	32.8 (63)	1,586	6.4 (45)	1,973	1.3 (33)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 112

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3954	14	*	14	*	4	*	12	*	14	*
4103	1	*	1	*	0	*	1	*	1	*
4106	1,882	14.5 (22)	2,143	7.2 (37)	223	39.9 (40)	1,701	1.7 (4)	2,138	0.6 (5)
4107	2,427	20.2 (47)	2,675	9.8 (51)	356	47.8 (22)	2,027	7.1 (49)	2,670	1.9 (61)
4109	641	27.1 (75)	710	14.2 (74)	138	40.6 (37)	534	4.3 (16)	707	1.3 (33)
4110	7	*	7	*	0	*	0	*	7	*
4112	508	3.3 (3)	517	1.4 (4)	16	*	382	9.2 (72)	485	2.3 (71)
4115	2,941	11.8 (10)	3,247	5.1 (18)	284	42.6 (33)	2,490	2.6 (10)	3,244	0.6 (5)
4351	439	31.9 (88)	468	21.4 (110)	122	28.7 (74)	405	9.6 (75)	469	2.3 (71)
4352	140	55.0 (143)	154	35.1 (142)	73	*	133	23.3 (139)	153	7.2 (140)
4353	351	24.8 (67)	388	13.1 (70)	69	*	315	7.0 (48)	388	2.1 (66)
4354	617	33.7 (95)	681	21.9 (114)	183	25.1 (79)	572	13.5 (106)	689	2.3 (71)
4355	259	36.3 (113)	284	24.3 (124)	84	*	229	12.2 (95)	285	4.2 (122)
4356	339	37.2 (118)	366	21.3 (108)	109	31.2 (68)	312	13.5 (106)	365	3.6 (105)
4358	58	*	56	*	8	*	54	*	56	*
4359	494	21.5 (51)	522	10.3 (56)	96	*	437	5.7 (36)	523	0.8 (13)
4360	181	30.9 (85)	184	19.0 (94)	45	*	164	11.0 (85)	183	1.6 (49)
4361	233	34.3 (99)	249	25.3 (128)	69	*	210	10.0 (76)	248	1.2 (30)
4362	239	43.5 (135)	271	24.0 (122)	85	*	214	18.2 (132)	272	4.4 (127)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 113

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4363	632	30.7 (84)	703	22.3 (117)	160	20.6 (81)	571	13.5 (106)	702	3.8 (111)
4401	737	30.0 (82)	802	16.6 (86)	162	43.8 (29)	633	6.8 (46)	801	1.2 (30)
4501	692	36.1 (109)	792	20.5 (104)	215	38.1 (47)	604	12.7 (103)	792	1.8 (56)
4601	1,051	30.9 (85)	1,140	14.2 (74)	245	45.3 (26)	917	9.4 (74)	1,141	4.1 (118)
4602	1	*	1	*	0	*	1	*	1	*
4701	857	28.8 (79)	928	16.9 (87)	191	36.6 (51)	694	5.2 (25)	930	1.3 (33)
5050	1,257	21.8 (53)	1,437	9.4 (49)	206	42.2 (35)	1,022	8.5 (61)	1,440	1.0 (20)
5051	2,223	17.0 (32)	2,493	7.6 (38)	284	44.0 (28)	1,774	5.2 (25)	2,486	1.7 (53)
5052	2,293	16.5 (30)	2,596	7.6 (38)	289	46.7 (23)	1,938	5.9 (40)	2,594	1.2 (30)
5055	1,119	2.6 (2)	1,194	1.0 (2)	24	*	1,022	4.7 (19)	1,194	0.8 (13)
5101	1,343	17.4 (35)	1,476	7.7 (40)	196	50.5 (17)	1,127	5.1 (23)	1,481	1.3 (33)
5102	25	*	26	*	4	*	17	*	26	*
5202	641	41.3 (129)	644	25.9 (130)	239	32.6 (64)	577	8.8 (69)	654	1.8 (56)
5206	5	*	9	*	1	*	2	*	9	*
5601	806	28.4 (78)	896	16.2 (83)	186	36.6 (51)	687	5.7 (36)	893	1.8 (56)
5801	1,512	33.9 (97)	1,659	17.4 (88)	416	40.6 (37)	1,312	11.4 (89)	1,655	4.1 (118)
5803	82	*	85	*	16	*	66	*	84	*
5804	22	*	22	*	3	*	15	*	22	*
5806	42	*	45	*	14	*	33	*	45	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 114

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
6101	36	*	36	*	13	*	31	*	36	*
6102	91	*	98	*	26	*	85	*	98	*
6103	2,834	29.5 (81)	3,164	17.9 (90)	706	29.7 (71)	2,556	10.1 (78)	3,174	2.8 (86)
6105	10	*	10	*	0	*	7	*	10	*
6106	113	34.5 (101)	103	22.3 (117)	36	*	108	14.8 (118)	104	9.6 (143)
6301	2,842	12.1 (13)	3,096	6.8 (34)	288	34.4 (57)	2,447	4.9 (21)	3,098	1.3 (33)
6312	1,519	4.9 (5)	1,637	1.5 (5)	47	*	1,247	2.2 (8)	1,635	0.3 (2)
6316	1,002	13.4 (19)	1,066	5.8 (23)	107	43.0 (30)	804	3.2 (12)	1,064	0.6 (5)
6318	28	*	33	*	5	*	20	*	33	*
6320	301	9.6 (7)	312	3.2 (8)	21	*	232	0.0 (1)	311	0.3 (2)
6325	268	25.7 (72)	270	16.3 (84)	46	*	198	8.1 (57)	269	1.5 (44)
6335	901	7.3 (6)	948	3.8 (11)	49	*	763	1.7 (4)	948	0.7 (11)
6340	21	*	14	*	1	*	21	*	14	*
6345	91	*	94	*	16	*	67	*	94	*
6360	863	14.8 (23)	920	6.5 (30)	100	40.0 (39)	744	8.7 (66)	919	1.1 (24)
7304	10	*	10	*	3	*	10	*	9	*
7306	1,899	34.4 (100)	2,087	11.6 (61)	555	55.3 (10)	1,705	10.9 (82)	2,118	2.3 (71)
7308	10	*	8	*	1	*	10	*	8	*
7310	2	*	2	*	0	*	2	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 115

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
7401	1,795	39.6 (124)	1,858	19.9 (100)	584	39.4 (42)	1,629	15.4 (119)	1,859	5.1 (133)
7601	515	22.3 (56)	523	14.7 (77)	90	*	470	8.9 (70)	523	1.5 (44)
7850	890	35.8 (106)	971	18.7 (92)	261	32.6 (64)	775	12.5 (101)	964	4.3 (125)
7852	15	*	17	*	6	*	15	*	17	*
8101	1,283	15.0 (24)	1,595	5.5 (20)	158	52.5 (14)	1,112	10.1 (78)	1,596	3.8 (111)
8102	389	15.2 (26)	454	6.8 (34)	49	*	346	5.8 (39)	454	2.9 (89)
8105	720	11.8 (10)	682	3.8 (11)	59	*	603	5.6 (34)	681	0.6 (5)
8106	405	23.5 (64)	518	10.0 (53)	69	*	333	7.8 (54)	518	1.7 (53)
8150	17	*	17	*	2	*	17	*	17	*
8151	2,639	19.5 (44)	2,843	8.3 (44)	414	50.2 (18)	2,285	5.4 (29)	2,845	1.1 (24)
8152	30	*	27	*	7	*	30	*	27	*
8153	6	*	5	*	1	*	5	*	5	*
8155	12	*	11	*	2	*	12	*	11	*
8157	5	*	5	*	1	*	5	*	5	*
8201	2,168	16.0 (27)	2,400	8.0 (43)	254	48.0 (21)	1,673	5.3 (28)	2,408	1.3 (33)
8202	3,337	4.4 (4)	3,736	1.2 (3)	129	61.2 (5)	2,937	4.7 (19)	3,736	1.3 (33)
8203	343	24.8 (67)	357	15.4 (81)	54	*	286	8.4 (60)	359	1.4 (41)
8250	539	20.8 (48)	579	6.6 (32)	81	*	458	7.9 (55)	581	0.9 (17)
8252	146	18.5 (39)	154	5.8 (23)	22	*	129	10.9 (82)	153	2.6 (79)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 116

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8255	1,889	22.4 (57)	2,086	11.9 (62)	305	39.3 (43)	1,553	7.5 (51)	2,088	1.6 (49)
8257	1,712	0.9 (1)	1,844	0.8 (1)	11	*	1,472	1.0 (3)	1,842	0.8 (13)
8263	574	22.6 (59)	560	9.8 (51)	92	*	467	5.6 (34)	559	1.1 (24)
8264	646	17.3 (34)	694	4.9 (16)	88	*	506	5.1 (23)	693	0.4 (4)
8267	575	17.7 (37)	622	8.4 (45)	67	*	491	5.7 (36)	624	1.9 (61)
8901	706	19.0 (42)	873	7.9 (41)	118	50.8 (16)	626	13.7 (111)	870	4.1 (118)
8902	1,126	13.7 (20)	1,297	3.7 (10)	128	64.1 (3)	982	10.1 (78)	1,295	3.7 (108)
8903	1,166	16.1 (28)	1,392	6.5 (30)	150	46.0 (24)	1,022	16.2 (125)	1,392	4.2 (122)
8904	1,809	17.4 (35)	2,148	5.6 (22)	246	64.2 (2)	1,520	13.6 (110)	2,147	5.2 (134)
8905	1,392	12.2 (14)	1,664	4.4 (14)	138	55.1 (11)	1,182	6.1 (42)	1,661	2.5 (78)
8909	745	15.0 (24)	863	4.5 (15)	91	*	641	11.5 (91)	858	3.3 (100)
9045	1,344	9.6 (7)	1,475	3.6 (9)	105	51.4 (15)	1,211	7.1 (49)	1,471	1.6 (49)
9050	759	18.8 (40)	908	5.8 (23)	136	60.3 (6)	685	15.9 (122)	904	2.8 (86)
9051	1,302	10.3 (9)	1,429	2.9 (7)	119	58.8 (8)	1,204	4.1 (14)	1,428	0.8 (13)
9053	883	12.2 (14)	1,132	3.9 (13)	93	*	662	12.2 (95)	1,132	2.7 (83)
9055	4	*	12	*	0	*	1	*	12	*
9058	829	12.8 (18)	1,083	5.1 (18)	85	*	698	8.5 (61)	1,082	1.8 (56)
9066	875	18.9 (41)	1,071	5.9 (26)	119	59.7 (7)	717	16.6 (127)	1,069	4.1 (118)
9068	706	19.8 (46)	853	6.6 (32)	113	54.9 (13)	635	13.4 (105)	853	3.5 (103)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 117

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
9069	1,695	12.7 (17)	2,146	5.5 (20)	169	55.0 (12)	1,332	15.5 (120)	2,144	5.6 (137)
9070	1,152	16.7 (31)	1,417	4.9 (16)	156	64.7 (1)	953	14.1 (113)	1,408	3.6 (105)
9072	830	21.2 (50)	861	6.3 (28)	154	62.3 (4)	767	15.9 (122)	860	5.0 (132)
9080	2,047	17.9 (38)	2,449	5.9 (26)	309	56.6 (9)	1,795	9.1 (71)	2,450	3.0 (91)
9082	110	11.8 (10)	124	2.4 (6)	11	*	88	*	124	7.3 (141)
9083	52	*	70	*	0	*	4	*	70	*
9084	22	*	21	*	2	*	13	*	21	*
9085	4	*	2	*	0	*	4	*	2	*
9086	12	*	10	*	1	*	12	*	10	*
9087	13	*	11	*	1	*	12	*	11	*
Michigan	126,731	22.5	140,279	11.5	23,244	39.8	109,690	8.4	140,315	2.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾ Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 118

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	90	*	*	93	*	84	*	90	*
0102	355	9.9 (77)	8.5 (111)	368	10.3 (29)	315	5.7 (109)	355	64.8 (101)
0103	333	7.8 (33)	7.5 (79)	341	6.5 (1)	308	6.5 (118)	333	70.6 (82)
0104	99	*	*	110	10.9 (43)	90	*	99	*
0105	13	*	*	13	*	11	*	13	*
0250	105	9.5 (66)	9.5 (124)	104	13.5 (104)	90	*	105	81.9 (26)
0251	77	*	*	79	*	71	*	77	*
0252	82	*	*	88	*	78	*	82	*
0253	110	7.3 (20)	8.2 (104)	109	15.6 (126)	91	*	110	77.3 (51)
0401	388	8.2 (44)	10.1 (132)	395	10.1 (22)	342	3.5 (48)	386	72.0 (75)
0402	331	8.5 (55)	6.9 (61)	331	10.6 (37)	288	4.2 (68)	331	78.9 (38)
0403	126	10.3 (84)	8.7 (113)	127	9.4 (12)	112	6.3 (114)	126	79.4 (34)
0404	92	*	*	92	*	78	*	92	*
0501	384	9.1 (61)	9.4 (122)	429	10.5 (34)	337	4.2 (68)	420	74.5 (64)
0503	81	*	*	83	*	72	*	84	*
0504	359	8.1 (43)	6.4 (49)	371	11.3 (53)	316	2.2 (6)	364	89.0 (1)
0505	200	13.0 (113)	6.5 (52)	220	13.6 (106)	173	5.8 (110)	215	83.7 (15)
0506	135	10.4 (87)	9.6 (127)	142	11.3 (53)	117	6.0 (112)	137	80.3 (31)
0507	66	*	*	74	*	57	*	72	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 119

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	24	*	*	25	*	22	*	25	*
0509	1	*	*	1	*	1	*	1	*
0650	919	9.7 (72)	9.1 (118)	955	11.3 (53)	819	5.0 (93)	920	75.0 (60)
0651	502	6.4 (7)	5.8 (40)	526	10.3 (29)	449	3.6 (53)	501	73.1 (70)
0652	295	10.8 (95)	7.1 (69)	306	11.4 (60)	258	5.0 (93)	296	72.3 (74)
0653	343	7.6 (26)	7.9 (87)	354	12.7 (87)	299	2.3 (7)	344	80.8 (27)
0654	336	11.0 (96)	6.0 (44)	343	10.2 (26)	298	7.4 (126)	339	74.9 (62)
0655	274	4.7 (2)	7.3 (74)	274	10.2 (26)	239	1.3 (2)	274	74.5 (64)
0657	59	*	*	62	*	49	*	59	*
0701	381	7.1 (17)	11.5 (138)	401	7.0 (3)	352	4.8 (89)	381	87.7 (4)
0702	65	*	*	65	*	62	*	65	*
0703	142	7.0 (14)	7.7 (83)	144	10.4 (31)	125	1.6 (3)	144	75.0 (60)
0704	46	*	*	47	*	40	*	46	*
0705	74	*	*	76	*	66	*	74	*
0706	9	*	*	9	*	7	*	9	*
0850	616	10.2 (82)	8.1 (99)	650	11.8 (70)	541	4.4 (75)	630	83.8 (14)
0851	540	9.1 (61)	8.1 (99)	568	11.4 (60)	477	4.0 (63)	546	82.8 (21)
0901	1,327	9.6 (67)	8.2 (104)	1,382	11.2 (51)	1,182	4.2 (68)	1,334	65.4 (97)
0902	34	*	*	34	*	33	*	34	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 120

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1051	1,012	7.6 (26)	11.6 (139)	1,048	9.4 (12)	909	3.1 (33)	1,016	84.4 (11)
1052	1	*	*	1	*	0	*	1	*
1101	1,450	9.6 (67)	5.0 (27)	1,482	11.3 (53)	1,266	5.3 (100)	1,454	68.8 (87)
1102	665	6.5 (9)	5.9 (43)	691	7.1 (4)	608	3.5 (48)	666	72.5 (72)
1104	1	*	*	1	*	1	*	1	*
1106	57	*	*	60	*	51	*	57	*
1250	884	8.6 (56)	8.3 (106)	909	10.0 (21)	783	4.2 (68)	891	75.1 (59)
1251	585	8.2 (44)	8.7 (113)	606	10.9 (43)	518	3.3 (38)	586	78.8 (39)
1252	451	7.5 (22)	9.3 (121)	457	11.2 (51)	396	3.0 (26)	453	83.4 (18)
1253	608	5.4 (3)	8.4 (109)	636	9.4 (12)	546	1.6 (3)	614	84.0 (12)
1350	2,036	9.8 (76)	7.0 (64)	2,076	10.2 (26)	1,808	5.0 (93)	2,042	73.8 (66)
1351	67	*	*	67	*	61	*	67	*
1352	339	9.7 (72)	7.7 (83)	350	9.4 (12)	305	4.6 (84)	346	67.6 (92)
1401	195	5.6 (4)	9.2 (120)	201	10.4 (31)	175	2.3 (7)	195	85.1 (8)
1402	96	*	*	99	*	85	*	96	*
1403	17	*	*	18	*	15	*	17	*
1501	647	7.7 (30)	8.5 (111)	685	10.5 (34)	572	2.8 (17)	673	70.0 (85)
1503	775	7.2 (18)	8.3 (106)	796	10.8 (42)	687	3.6 (53)	776	74.6 (63)
1504	1,493	8.0 (38)	7.5 (79)	1,750	11.5 (65)	1,307	3.3 (38)	1,730	78.6 (40)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 121

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1505	236	6.8 (13)	11.9 (140)	249	12.4 (80)	208	2.9 (22)	241	83.0 (20)
1506	496	7.5 (22)	8.3 (106)	508	11.8 (70)	435	2.8 (17)	500	73.4 (68)
1507	459	7.6 (26)	11.1 (137)	490	11.0 (47)	406	3.0 (26)	470	82.6 (23)
1508	17	*	*	17	*	15	*	17	*
1509	323	5.9 (5)	10.5 (135)	338	12.7 (87)	280	1.1 (1)	327	68.8 (87)
1511	544	8.8 (59)	7.4 (76)	562	13.2 (99)	468	2.6 (15)	554	80.7 (29)
1512	305	9.2 (63)	7.5 (79)	313	14.7 (118)	261	2.3 (7)	306	78.1 (45)
1514	372	12.6 (111)	8.1 (99)	401	17.0 (134)	310	2.6 (15)	385	85.7 (6)
1701	486	8.0 (38)	11.9 (140)	501	10.6 (37)	424	2.8 (17)	485	68.0 (91)
1950	535	11.0 (96)	6.5 (52)	550	13.6 (106)	458	3.3 (38)	536	82.8 (21)
1952	117	7.7 (30)	7.7 (83)	122	11.5 (65)	103	2.9 (22)	117	88.0 (3)
1953	248	10.5 (90)	7.3 (74)	256	13.3 (101)	212	3.3 (38)	251	88.4 (2)
1954	394	6.6 (10)	7.9 (87)	401	9.7 (18)	349	1.7 (5)	395	82.3 (24)
1955	91	*	*	92	*	81	*	91	*
1957	354	13.6 (121)	11.0 (136)	357	14.8 (119)	297	4.0 (63)	356	85.1 (8)
1963	53	*	*	55	*	49	*	53	*
1964	14	*	*	14	*	14	*	14	*
2150	534	6.6 (10)	8.8 (115)	559	9.7 (18)	479	2.5 (12)	535	79.6 (33)
2151	178	7.9 (37)	7.9 (87)	181	8.8 (8)	162	4.9 (91)	178	58.4 (115)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 122

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2152	15	*	*	15	*	11	*	15	*
2251	353	7.6 (26)	7.9 (87)	359	12.0 (74)	311	3.5 (48)	354	75.4 (57)
2252	134	8.2 (44)	9.7 (128)	139	10.1 (22)	120	3.3 (38)	134	76.9 (52)
2554	42	*	*	44	*	38	*	42	*
2555	6,476	11.5 (102)	5.2 (31)	6,644	13.3 (101)	5,584	5.5 (105)	6,482	56.9 (121)
2556	261	14.6 (127)	3.4 (9)	270	20.7 (142)	205	7.3 (124)	265	49.8 (135)
2557	2	*	*	2	*	1	*	2	*
2558	2	*	*	3	*	1	*	3	*
2560	3	*	*	3	*	3	*	3	*
2561	3	*	*	3	*	2	*	3	*
2562	15	*	*	15	*	12	*	15	*
2563	24	*	*	23	*	22	*	24	*
2564	18	*	*	18	*	18	*	18	*
3001	1,024	14.8 (128)	3.7 (14)	1,050	15.4 (124)	866	5.5 (105)	1,023	57.8 (117)
3002	233	9.9 (77)	3.4 (9)	235	12.3 (79)	205	5.4 (102)	233	55.4 (124)
3004	206	11.2 (98)	6.8 (60)	213	15.5 (125)	172	4.1 (67)	207	71.5 (77)
3005	105	3.8 (1)	9.5 (124)	105	8.6 (7)	96	*	106	70.8 (79)
3007	12	*	*	11	*	7	*	12	*
3201	373	8.6 (56)	9.1 (118)	391	10.7 (40)	336	3.0 (26)	372	77.7 (47)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 123

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3203	0	*	*	0	*	0	*	0	*
3301	3,791	10.7 (93)	5.7 (38)	3,955	13.0 (91)	3,299	4.5 (82)	3,823	76.3 (55)
3304	86	*	*	88	*	76	*	87	*
3305	134	7.5 (22)	7.5 (79)	140	10.7 (40)	121	3.3 (38)	136	85.3 (7)
3306	214	6.1 (6)	7.0 (64)	222	7.2 (5)	198	2.5 (12)	213	77.5 (49)
3308	81	*	*	84	*	71	*	80	*
3309	7	*	*	7	*	7	*	7	*
3312	373	11.3 (100)	8.0 (95)	383	15.1 (122)	315	4.4 (75)	375	70.7 (81)
3550	53	*	*	53	*	44	*	53	*
3551	11	*	*	13	*	11	*	12	*
3552	29	*	*	30	*	25	*	29	*
3701	844	7.7 (30)	9.7 (128)	863	10.5 (34)	749	2.9 (22)	844	83.4 (18)
3850	2,582	9.9 (77)	7.9 (87)	2,648	11.4 (60)	2,289	4.5 (82)	2,586	70.8 (79)
3901	1,343	11.8 (105)	4.6 (22)	1,405	12.8 (90)	1,161	5.1 (97)	1,363	66.8 (94)
3902	93	*	*	99	*	83	*	93	*
3903	23	*	*	24	*	21	*	23	*
3904	4	*	*	4	*	3	*	4	*
3905	9	*	*	9	*	7	*	9	*
3950	1,824	10.3 (84)	8.4 (109)	1,918	11.8 (70)	1,597	3.4 (46)	1,843	72.7 (71)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 124

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3954	12	*	*	13	*	12	*	12	*
4103	1	*	*	1	*	0	*	1	*
4106	2,002	12.9 (112)	4.7 (25)	2,087	14.6 (116)	1,702	5.8 (110)	2,021	76.1 (56)
4107	2,508	13.3 (116)	6.0 (44)	2,624	16.1 (131)	2,104	5.6 (108)	2,533	77.7 (47)
4109	675	10.1 (81)	7.0 (64)	694	15.7 (128)	568	3.2 (36)	675	78.4 (43)
4110	7	*	*	7	*	0	*	7	*
4112	482	6.4 (7)	5.6 (36)	481	11.0 (47)	420	3.1 (33)	482	24.1 (141)
4115	3,059	13.0 (113)	6.0 (44)	3,185	14.9 (121)	2,593	4.8 (89)	3,091	77.5 (49)
4351	439	8.4 (51)	7.7 (83)	459	13.1 (93)	384	4.2 (68)	441	82.1 (25)
4352	143	11.9 (107)	9.8 (130)	147	17.0 (134)	115	5.2 (99)	143	73.4 (68)
4353	367	8.4 (51)	7.4 (76)	367	12.0 (74)	316	3.8 (57)	371	78.2 (44)
4354	661	8.2 (44)	8.8 (115)	673	10.1 (22)	591	3.0 (26)	663	73.8 (66)
4355	248	13.3 (116)	6.9 (61)	263	15.6 (126)	205	4.4 (75)	250	76.8 (53)
4356	341	10.0 (80)	7.9 (87)	351	12.5 (82)	294	4.4 (75)	341	72.4 (73)
4358	53	*	*	53	*	49	*	53	*
4359	485	7.0 (14)	7.2 (71)	490	10.6 (37)	427	3.0 (26)	486	76.7 (54)
4360	152	6.6 (10)	7.2 (71)	169	6.5 (1)	138	2.9 (22)	152	84.9 (10)
4361	202	9.4 (65)	9.9 (131)	239	11.7 (68)	179	3.9 (61)	203	83.7 (15)
4362	256	10.2 (82)	9.4 (122)	260	10.4 (31)	224	4.9 (91)	257	86.0 (5)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 125

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4363	654	8.0 (38)	9.5 (124)	664	12.2 (77)	561	2.3 (7)	655	80.6 (30)
4401	769	7.8 (33)	7.9 (87)	775	11.1 (49)	672	4.0 (63)	769	79.8 (32)
4501	742	7.8 (33)	8.1 (99)	761	12.7 (87)	644	2.8 (17)	741	78.1 (45)
4601	1,081	7.0 (14)	7.2 (71)	1,128	9.6 (16)	973	2.8 (17)	1,080	70.9 (78)
4602	1	*	*	1	*	1	*	1	*
4701	896	10.5 (90)	6.7 (57)	911	12.5 (82)	781	3.8 (57)	894	83.9 (13)
5050	1,400	11.6 (104)	5.4 (34)	1,407	15.9 (129)	1,164	4.3 (74)	1,399	61.6 (108)
5051	2,390	10.5 (90)	5.6 (36)	2,423	12.6 (85)	2,055	4.4 (75)	2,397	64.6 (103)
5052	2,490	12.5 (110)	5.0 (27)	2,534	14.4 (113)	2,104	5.5 (105)	2,497	61.4 (109)
5055	1,063	8.3 (49)	5.5 (35)	1,114	8.1 (6)	977	3.9 (61)	1,097	52.9 (129)
5101	1,433	8.4 (51)	6.6 (55)	1,445	11.5 (65)	1,254	3.0 (26)	1,439	70.6 (82)
5102	26	*	*	25	*	22	*	25	*
5202	628	7.5 (22)	10.2 (134)	637	9.3 (11)	566	3.7 (55)	630	83.7 (15)
5206	9	*	*	7	*	7	*	9	*
5601	862	7.2 (18)	10.1 (132)	884	9.6 (16)	777	2.3 (7)	862	79.1 (36)
5801	1,574	9.0 (60)	7.0 (64)	1,616	11.1 (49)	1,392	3.8 (57)	1,576	67.0 (93)
5803	81	*	*	80	*	70	*	81	*
5804	20	*	*	21	*	18	*	20	*
5806	43	*	*	44	*	37	*	43	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 126

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6101	33	*	*	34	*	26	*	34	*
6102	99	*	*	99	*	90	*	99	*
6103	3,005	10.7 (93)	5.8 (40)	3,055	13.1 (93)	2,562	4.7 (87)	3,013	64.8 (101)
6105	10	*	*	10	*	9	*	10	*
6106	103	9.7 (72)	4.9 (26)	104	14.4 (113)	88	*	103	65.0 (99)
6301	2,920	13.0 (113)	6.7 (57)	3,042	13.1 (93)	2,515	5.4 (102)	2,957	68.3 (90)
6312	1,530	11.8 (105)	6.2 (48)	1,601	13.0 (91)	1,326	3.5 (48)	1,551	78.6 (40)
6316	988	10.3 (84)	6.7 (57)	1,037	12.6 (85)	863	3.5 (48)	1,001	79.0 (37)
6318	28	*	*	32	*	26	*	30	*
6320	289	10.4 (87)	6.9 (61)	303	13.9 (109)	249	4.0 (63)	292	80.8 (27)
6325	262	9.2 (63)	8.8 (115)	266	9.8 (20)	235	3.8 (57)	261	78.5 (42)
6335	899	9.6 (67)	5.7 (38)	912	12.0 (74)	785	4.7 (87)	906	71.9 (76)
6340	13	*	*	14	*	10	*	13	*
6345	92	*	*	95	*	83	*	92	*
6360	880	13.3 (116)	5.1 (30)	907	14.4 (113)	746	6.8 (120)	887	69.0 (86)
7304	10	*	*	10	*	10	*	10	*
7306	2,021	12.1 (108)	5.8 (40)	2,077	13.3 (101)	1,750	4.6 (84)	2,025	65.3 (98)
7308	8	*	*	8	*	6	*	8	*
7310	2	*	*	2	*	2	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 127

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7401	1,767	8.3 (49)	7.9 (87)	1,812	11.4 (60)	1,560	3.7 (55)	1,771	66.3 (96)
7601	496	7.3 (20)	6.0 (44)	517	11.4 (60)	434	2.5 (12)	503	54.7 (126)
7850	924	8.0 (38)	7.1 (69)	949	11.3 (53)	817	3.2 (36)	926	70.6 (82)
7852	16	*	*	16	*	14	*	16	*
8101	1,472	17.5 (141)	2.2 (2)	1,518	20.4 (141)	1,136	8.5 (133)	1,493	44.6 (138)
8102	425	8.0 (38)	8.0 (95)	444	12.2 (77)	366	4.4 (75)	431	54.5 (127)
8105	628	9.6 (67)	4.6 (22)	673	11.9 (73)	550	3.3 (38)	661	57.3 (120)
8106	486	14.8 (128)	4.5 (21)	505	17.6 (138)	397	6.0 (112)	492	57.5 (119)
8150	16	*	*	16	*	16	*	16	*
8151	2,675	12.1 (108)	5.3 (33)	2,754	14.1 (111)	2,282	5.0 (93)	2,691	79.2 (35)
8152	26	*	*	27	*	26	*	26	*
8153	5	*	*	4	*	3	*	5	*
8155	10	*	*	11	*	8	*	10	*
8157	5	*	*	5	*	4	*	5	*
8201	2,267	11.5 (102)	6.4 (49)	2,365	12.5 (82)	1,972	5.3 (100)	2,298	62.7 (107)
8202	3,413	8.2 (44)	5.2 (31)	3,590	9.0 (9)	3,085	4.2 (68)	3,545	50.2 (134)
8203	339	9.7 (72)	8.0 (95)	348	13.2 (99)	291	3.1 (33)	343	75.2 (58)
8250	557	13.3 (116)	5.0 (27)	571	13.1 (93)	480	6.9 (121)	562	59.1 (113)
8252	142	14.8 (128)	3.5 (12)	147	10.9 (43)	124	7.3 (124)	145	62.8 (106)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 128

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8255	1,989	11.3 (100)	6.5 (52)	2,054	11.3 (53)	1,747	5.4 (102)	2,012	59.8 (112)
8257	1,640	14.4 (124)	2.7 (3)	1,782	11.3 (53)	1,428	9.1 (136)	1,715	68.8 (87)
8263	532	9.6 (67)	8.1 (99)	550	12.4 (80)	467	3.4 (46)	534	57.7 (118)
8264	653	8.4 (51)	6.6 (55)	669	10.9 (43)	574	3.3 (38)	656	63.0 (105)
8267	587	7.8 (33)	8.0 (95)	615	10.1 (22)	527	3.0 (26)	594	55.1 (125)
8901	820	16.7 (139)	3.0 (5)	852	17.1 (136)	674	8.8 (135)	818	46.7 (137)
8902	1,147	15.1 (132)	3.0 (5)	1,267	17.3 (137)	956	6.4 (117)	1,219	61.0 (110)
8903	1,274	15.4 (135)	3.2 (7)	1,334	17.8 (139)	1,026	7.8 (128)	1,285	51.1 (133)
8904	1,984	15.9 (138)	3.3 (8)	2,063	18.0 (140)	1,591	7.0 (122)	2,028	53.2 (128)
8905	1,476	11.2 (98)	6.4 (49)	1,600	14.1 (111)	1,244	5.1 (97)	1,574	66.5 (95)
8909	787	15.2 (133)	4.1 (18)	832	14.8 (119)	664	8.0 (129)	806	58.6 (114)
9045	1,392	10.4 (87)	7.0 (64)	1,444	13.1 (93)	1,205	4.6 (84)	1,402	47.7 (136)
9050	826	14.0 (123)	4.6 (22)	878	14.6 (116)	699	7.2 (123)	847	56.2 (123)
9051	1,355	8.7 (58)	7.4 (76)	1,359	13.5 (104)	1,149	4.4 (75)	1,367	63.3 (104)
9053	1,062	15.3 (134)	3.8 (15)	1,106	13.8 (108)	905	8.2 (130)	1,065	60.8 (111)
9055	11	*	*	11	*	10	*	11	*
9058	1,005	14.5 (126)	2.7 (3)	1,066	11.7 (68)	879	8.4 (131)	1,045	58.0 (116)
9066	990	16.9 (140)	3.4 (9)	1,051	16.3 (132)	834	8.4 (131)	998	52.5 (131)
9068	817	15.8 (137)	4.2 (20)	831	15.2 (123)	685	8.5 (133)	817	44.1 (139)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2019 - 2021 Pregnancy Nutrition Surveillance (1)

Page : 129

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9069	2,006	15.6 (136)	3.5 (12)	2,096	16.0 (130)	1,676	7.4 (126)	2,006	35.9 (140)
9070	1,349	13.7 (122)	3.8 (15)	1,392	13.1 (93)	1,160	6.6 (119)	1,350	56.6 (122)
9072	812	15.0 (131)	4.1 (18)	844	16.5 (133)	678	6.3 (114)	818	52.9 (129)
9080	2,303	13.5 (120)	4.0 (17)	2,395	13.9 (109)	1,962	6.3 (114)	2,369	52.2 (132)
9082	118	14.4 (124)	1.7 (1)	122	9.0 (9)	105	9.5 (137)	117	65.0 (99)
9083	67	*	*	69	*	54	*	67	*
9084	18	*	*	21	*	16	*	18	*
9085	2	*	*	2	*	2	*	2	*
9086	10	*	*	10	*	10	*	10	*
9087	10	*	*	11	*	8	*	10	*
Michigan	131,523	10.9	6.3	136,468	12.7	113,828	4.8	133,104	56.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.