

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2023 Pregnancy Nutrition Surveillance (1)

Page : 1

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	52	*	*	*	*	*	*	*
0102	118	92.4	0.0	2.5	0.0	0.0	4.2	0.8
0103	134	91.0	0.7	2.2	0.0	0.0	5.2	0.7
0104	43	*	*	*	*	*	*	*
0105	1	*	*	*	*	*	*	*
0250	40	*	*	*	*	*	*	*
0251	28	*	*	*	*	*	*	*
0252	36	*	*	*	*	*	*	*
0253	47	*	*	*	*	*	*	*
0401	131	95.4	0.8	2.3	0.0	0.0	1.5	0.0
0402	92	*	*	*	*	*	*	*
0403	37	*	*	*	*	*	*	*
0404	33	*	*	*	*	*	*	*
0501	131	86.3	1.5	5.3	0.8	1.5	4.6	0.0
0503	18	*	*	*	*	*	*	*
0504	96	*	*	*	*	*	*	*
0505	55	*	*	*	*	*	*	*
0506	52	*	*	*	*	*	*	*
0507	21	*	*	*	*	*	*	*
0508	4	*	*	*	*	*	*	*
0650	345	78.6	4.1	6.7	2.6	0.3	7.5	0.3
0651	162	90.7	0.6	4.9	0.6	0.6	2.5	0.0
0652	83	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2023 Pregnancy Nutrition Surveillance (1)

Page : 2

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0653	123	91.9	0.8	4.1	0.0	0.0	2.4	0.8
0654	137	92.0	1.5	1.5	0.0	0.7	3.6	0.7
0655	108	96.3	0.0	1.9	0.0	0.0	1.9	0.0
0657	7	*	*	*	*	*	*	*
0701	137	89.1	2.2	0.0	1.5	4.4	2.9	0.0
0702	33	*	*	*	*	*	*	*
0703	63	*	*	*	*	*	*	*
0704	21	*	*	*	*	*	*	*
0705	28	*	*	*	*	*	*	*
0706	1	*	*	*	*	*	*	*
0850	221	81.0	5.0	8.1	0.0	2.3	3.6	0.0
0851	216	89.4	0.5	6.0	0.5	1.9	1.9	0.0
0901	434	78.3	3.2	9.7	0.0	0.7	7.6	0.5
0902	3	*	*	*	*	*	*	*
1051	353	78.8	1.4	10.5	2.3	1.7	5.4	0.0
1101	455	37.1	46.2	9.9	0.4	3.1	2.4	0.9
1102	217	65.0	12.0	11.1	0.9	0.9	8.3	1.8
1106	21	*	*	*	*	*	*	*
1250	340	74.4	7.6	16.2	0.3	0.0	1.5	0.0
1251	237	92.0	0.4	3.8	0.0	0.4	3.4	0.0
1252	164	62.8	0.0	32.3	0.0	0.0	4.9	0.0
1253	206	79.1	1.0	11.7	0.0	0.0	7.3	1.0
1350	738	58.5	17.3	10.8	0.4	5.4	6.9	0.5

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2023 Pregnancy Nutrition Surveillance (1)

Page : 3

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1352	96	*	*	*	*	*	*	*
1401	75	*	*	*	*	*	*	*
1402	26	*	*	*	*	*	*	*
1403	2	*	*	*	*	*	*	*
1501	263	43.0	7.2	43.7	0.0	0.0	3.0	3.0
1503	258	57.8	7.8	22.9	0.4	0.8	7.0	3.5
1504	604	35.6	5.3	50.5	0.0	4.8	3.1	0.7
1506	173	74.6	0.0	18.5	0.0	0.6	5.8	0.6
1507	168	80.4	1.2	13.1	0.6	1.2	1.8	1.8
1509	109	51.4	1.8	39.4	0.0	1.8	3.7	1.8
1511	189	85.2	1.6	9.0	0.0	0.5	3.7	0.0
1512	119	79.8	2.5	10.1	0.8	1.7	5.0	0.0
1514	212	76.4	4.2	12.7	0.0	1.4	5.2	0.0
1701	161	60.9	1.2	3.1	16.8	1.9	16.1	0.0
1950	172	93.6	0.0	5.2	0.0	0.0	1.2	0.0
1952	44	*	*	*	*	*	*	*
1953	104	82.7	0.0	14.4	0.0	0.0	2.9	0.0
1954	117	83.8	0.0	13.7	0.0	0.0	2.6	0.0
1955	45	*	*	*	*	*	*	*
1957	116	81.9	4.3	12.9	0.0	0.9	0.0	0.0
1963	30	*	*	*	*	*	*	*
2150	183	82.5	0.0	3.8	2.7	0.0	10.9	0.0
2151	77	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2023 Pregnancy Nutrition Surveillance (1)

Page : 4

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2152	9	*	*	*	*	*	*	*
2251	114	91.2	0.0	2.6	0.0	0.0	5.3	0.9
2252	59	*	*	*	*	*	*	*
2554	11	*	*	*	*	*	*	*
2555	2,313	48.6	37.4	6.8	0.0	0.3	6.6	0.4
2556	21	*	*	*	*	*	*	*
2563	1	*	*	*	*	*	*	*
2564	1	*	*	*	*	*	*	*
3001	361	27.7	45.2	12.5	0.6	1.1	12.7	0.3
3002	199	37.7	37.2	18.1	0.0	0.0	7.0	0.0
3004	112	78.6	0.0	17.0	0.0	0.9	2.7	0.9
3005	54	*	*	*	*	*	*	*
3007	2	*	*	*	*	*	*	*
3201	148	91.2	0.7	5.4	0.0	0.0	0.7	2.0
3301	1,409	37.3	30.6	16.6	0.3	6.2	8.6	0.4
3304	24	*	*	*	*	*	*	*
3305	51	*	*	*	*	*	*	*
3306	62	*	*	*	*	*	*	*
3308	25	*	*	*	*	*	*	*
3312	153	43.8	23.5	17.6	1.3	3.3	10.5	0.0
3314	1	*	*	*	*	*	*	*
3550	10	*	*	*	*	*	*	*
3701	269	88.5	0.4	8.6	0.0	0.4	2.2	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2023 Pregnancy Nutrition Surveillance (1)

Page : 5

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3850	849	70.8	13.3	7.7	0.2	0.7	7.1	0.2
3901	449	34.7	48.1	7.8	0.2	1.6	6.9	0.7
3950	699	50.5	23.0	14.9	0.0	0.4	10.3	0.9
4106	752	11.7	34.0	48.1	0.1	1.2	4.3	0.5
4107	882	40.8	28.9	21.3	0.3	1.9	6.7	0.0
4109	241	82.6	2.1	9.5	0.4	1.7	2.9	0.8
4112	201	0.5	2.0	97.5	0.0	0.0	0.0	0.0
4115	1,110	31.6	31.2	21.6	0.2	9.6	5.6	0.2
4351	170	86.5	0.6	6.5	0.0	1.2	5.3	0.0
4352	58	*	*	*	*	*	*	*
4353	164	60.4	1.2	36.0	0.0	0.0	2.4	0.0
4354	251	87.3	2.0	5.6	0.0	0.4	4.0	0.8
4355	97	*	*	*	*	*	*	*
4356	128	74.2	1.6	14.1	0.0	0.8	8.6	0.8
4359	169	56.8	1.2	40.2	0.6	0.0	0.0	1.2
4360	78	*	*	*	*	*	*	*
4361	74	*	*	*	*	*	*	*
4362	83	*	*	*	*	*	*	*
4363	219	92.2	0.5	5.5	0.0	0.5	1.4	0.0
4401	244	88.5	0.4	7.0	0.0	0.0	3.7	0.4
4501	248	95.2	0.4	3.2	0.0	0.4	0.4	0.4
4601	391	74.2	2.6	18.4	0.0	0.5	3.8	0.5
4701	321	87.2	0.9	7.2	0.3	1.9	1.6	0.9

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2023 Pregnancy Nutrition Surveillance (1)

Page : 6

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
5050	547	45.5	42.6	3.5	0.0	1.6	6.4	0.4
5051	822	48.9	36.6	4.3	0.1	5.1	4.5	0.5
5052	892	26.7	39.2	3.0	0.1	25.9	4.6	0.4
5055	445	76.2	8.8	0.9	0.4	13.0	0.7	0.0
5101	489	58.5	12.9	19.6	0.0	2.0	6.5	0.4
5202	199	87.4	1.0	3.0	2.5	0.0	5.5	0.5
5601	305	87.5	2.3	6.6	0.0	1.3	2.3	0.0
5801	533	78.8	5.4	9.4	0.0	0.0	4.9	1.5
5803	3	*	*	*	*	*	*	*
5804	1	*	*	*	*	*	*	*
5806	32	*	*	*	*	*	*	*
6101	18	*	*	*	*	*	*	*
6102	14	*	*	*	*	*	*	*
6103	1,040	53.9	27.6	10.3	0.5	1.0	6.4	0.3
6105	1	*	*	*	*	*	*	*
6301	1,128	36.8	29.7	25.7	0.3	2.7	4.3	0.4
6312	600	31.8	56.0	6.5	0.2	2.0	3.3	0.2
6316	388	60.8	22.2	9.3	0.0	2.8	3.9	1.0
6318	10	*	*	*	*	*	*	*
6320	185	40.5	46.5	3.8	0.0	3.8	5.4	0.0
6325	110	90.9	1.8	2.7	0.0	0.0	4.5	0.0
6335	308	58.1	24.4	4.9	0.0	9.7	1.9	1.0
6345	60	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2023 Pregnancy Nutrition Surveillance (1)

Page : 7

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6360	262	29.0	43.1	19.1	0.0	4.6	4.2	0.0
7306	539	47.7	28.8	18.7	0.0	0.6	4.1	0.2
7401	558	79.9	4.7	8.4	0.4	0.5	5.7	0.4
7601	152	91.4	0.0	7.2	0.0	0.0	0.7	0.7
7850	346	91.6	0.3	4.0	0.0	0.0	3.2	0.9
7852	2	*	*	*	*	*	*	*
8101	535	6.5	90.1	1.9	0.0	0.2	1.3	0.0
8102	99	*	*	*	*	*	*	*
8106	157	23.6	52.2	14.6	0.0	0.0	8.9	0.6
8150	1	*	*	*	*	*	*	*
8151	1,044	37.0	33.6	17.5	0.1	4.2	7.1	0.5
8155	2	*	*	*	*	*	*	*
8201	768	41.0	42.2	6.5	0.1	4.3	5.2	0.7
8202	1,307	91.5	5.3	1.5	0.0	0.4	1.0	0.4
8203	123	52.0	29.3	8.9	0.0	0.0	9.8	0.0
8250	137	25.5	64.2	2.9	0.0	0.0	7.3	0.0
8252	28	*	*	*	*	*	*	*
8255	617	49.6	31.9	10.4	0.6	3.1	4.1	0.3
8257	512	53.5	7.0	1.0	0.0	36.7	1.8	0.0
8263	101	66.3	11.9	16.8	0.0	0.0	5.0	0.0
8264	286	54.9	32.5	4.9	0.3	3.5	3.8	0.0
8267	280	55.0	17.1	22.1	0.0	0.7	4.6	0.4
8901	232	6.0	85.3	2.6	0.0	2.6	3.4	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2023 Pregnancy Nutrition Surveillance (1)

Page : 8

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8902	384	7.3	88.8	1.0	0.0	0.8	1.8	0.3
8903	399	5.8	87.7	1.3	0.0	3.0	2.3	0.0
8904	584	6.0	88.0	2.4	0.0	1.0	2.4	0.2
8905	510	69.0	18.8	7.5	0.2	1.6	2.9	0.0
8909	348	7.2	87.6	2.6	0.0	0.0	2.6	0.0
9045	437	11.7	21.3	63.8	0.0	0.5	1.1	1.6
9050	262	18.3	59.5	18.3	0.0	0.4	3.4	0.0
9051	631	6.3	4.8	88.4	0.0	0.3	0.0	0.2
9053	591	10.2	80.0	3.0	0.0	3.7	2.9	0.2
9055	3	*	*	*	*	*	*	*
9058	460	13.9	77.0	0.9	0.0	6.3	1.7	0.2
9066	145	3.4	91.0	2.1	0.0	0.0	3.4	0.0
9068	298	3.7	92.3	1.3	0.0	0.3	2.0	0.3
9069	539	0.7	95.9	2.2	0.0	0.2	0.9	0.0
9070	666	6.6	88.7	1.7	0.0	1.8	0.9	0.3
9080	939	19.6	73.8	3.1	0.2	0.1	3.0	0.2
9082	38	*	*	*	*	*	*	*
9083	7	*	*	*	*	*	*	*
Michigan	46,160	50.6	28.4	12.8	0.3	2.8	4.6	0.4
National (Prior Year)	56,992	49.9	25.9	14.8	1.9	2.9	4.3	0.3

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2023 Pregnancy Nutrition Surveillance (1)

Page : 9

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	52	*	*	*	*	*	*	*
0102	118	0.0	0.0	7.6	61.0	26.3	4.2	0.8
0103	134	0.7	1.5	2.2	63.4	29.9	0.7	1.5
0104	43	*	*	*	*	*	*	*
0105	1	*	*	*	*	*	*	*
0250	40	*	*	*	*	*	*	*
0251	28	*	*	*	*	*	*	*
0252	36	*	*	*	*	*	*	*
0253	47	*	*	*	*	*	*	*
0401	131	0.0	0.8	3.1	61.8	32.8	1.5	0.0
0402	92	*	*	*	*	*	*	*
0403	37	*	*	*	*	*	*	*
0404	33	*	*	*	*	*	*	*
0501	131	0.0	1.5	9.9	59.5	24.4	3.1	1.5
0503	18	*	*	*	*	*	*	*
0504	96	*	*	*	*	*	*	*
0505	55	*	*	*	*	*	*	*
0506	52	*	*	*	*	*	*	*
0507	21	*	*	*	*	*	*	*
0508	4	*	*	*	*	*	*	*
0650	345	0.0	1.2	8.1	62.3	24.9	2.3	1.2
0651	162	0.0	1.2	6.8	61.7	26.5	1.2	2.5
0652	83	*	*	*	*	*	*	*
0653	123	0.0	1.6	3.3	61.8	28.5	0.8	4.1

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2023 Pregnancy Nutrition Surveillance (1)

Page : 10

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0654	137	0.0	2.9	5.8	69.3	16.1	1.5	4.4
0655	108	0.0	0.9	6.5	58.3	32.4	0.0	1.9
0657	7	*	*	*	*	*	*	*
0701	137	0.0	0.7	5.8	62.8	27.7	2.9	0.0
0702	33	*	*	*	*	*	*	*
0703	63	*	*	*	*	*	*	*
0704	21	*	*	*	*	*	*	*
0705	28	*	*	*	*	*	*	*
0706	1	*	*	*	*	*	*	*
0850	221	0.0	0.9	6.3	55.2	35.3	0.9	1.4
0851	216	0.0	2.3	7.4	61.6	25.5	1.4	1.9
0901	434	0.0	2.8	9.2	58.3	27.9	0.9	0.9
0902	3	*	*	*	*	*	*	*
1051	353	0.0	0.8	5.4	61.2	29.2	2.0	1.4
1101	455	0.2	2.2	8.4	61.3	22.4	2.6	2.9
1102	217	0.5	2.3	7.8	61.8	26.7	0.5	0.5
1106	21	*	*	*	*	*	*	*
1250	340	0.0	2.1	5.6	62.4	24.4	4.1	1.5
1251	237	0.4	1.7	6.3	69.6	17.3	2.1	2.5
1252	164	0.0	2.4	4.9	62.8	25.0	1.8	3.0
1253	206	0.0	2.9	5.8	59.7	25.7	1.0	4.9
1350	738	0.0	1.4	6.9	60.0	26.4	2.6	2.7
1352	96	*	*	*	*	*	*	*
1401	75	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS
Table 4A
2023 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic

Run Date: 6/28/2024

Page : 11

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1402	26	*	*	*	*	*	*	*
1403	2	*	*	*	*	*	*	*
1501	263	0.4	2.3	3.4	57.0	29.7	4.9	2.3
1503	258	0.0	1.6	6.6	60.5	25.2	3.1	3.1
1504	604	0.2	1.7	5.8	56.1	33.6	2.2	0.5
1506	173	0.0	0.6	8.1	61.3	24.3	1.7	4.0
1507	168	0.0	0.6	6.5	53.6	36.9	1.2	1.2
1509	109	0.0	0.9	9.2	56.9	30.3	0.9	1.8
1511	189	0.0	3.2	6.9	52.4	35.4	0.5	1.6
1512	119	0.0	0.0	2.5	67.2	28.6	0.8	0.8
1514	212	0.0	0.9	3.3	54.7	37.3	2.8	0.9
1701	161	0.0	1.2	3.1	64.6	27.3	0.6	3.1
1950	172	0.0	1.7	7.6	56.4	30.8	1.2	2.3
1952	44	*	*	*	*	*	*	*
1953	104	0.0	0.0	8.7	58.7	28.8	2.9	1.0
1954	117	0.0	2.6	6.0	59.0	31.6	0.9	0.0
1955	45	*	*	*	*	*	*	*
1957	116	0.0	0.0	1.7	64.7	31.0	0.9	1.7
1963	30	*	*	*	*	*	*	*
2150	183	0.0	1.1	4.9	62.3	28.4	2.7	0.5
2151	77	*	*	*	*	*	*	*
2152	9	*	*	*	*	*	*	*
2251	114	0.0	0.0	4.4	60.5	34.2	0.9	0.0
2252	59	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS
Table 4A
2023 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic

Run Date: 6/28/2024

Page : 12

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2554	11	*	*	*	*	*	*	*
2555	2,313	0.0	1.8	6.7	60.6	28.5	1.5	0.8
2556	21	*	*	*	*	*	*	*
2563	1	*	*	*	*	*	*	*
2564	1	*	*	*	*	*	*	*
3001	361	0.0	0.6	4.2	66.2	24.4	0.8	3.9
3002	199	0.0	1.5	7.0	53.8	30.2	3.0	4.5
3004	112	0.0	1.8	8.0	42.9	42.9	2.7	1.8
3005	54	*	*	*	*	*	*	*
3007	2	*	*	*	*	*	*	*
3201	148	0.0	0.7	2.0	58.8	35.1	1.4	2.0
3301	1,409	0.1	2.0	4.7	56.6	32.1	2.3	2.2
3304	24	*	*	*	*	*	*	*
3305	51	*	*	*	*	*	*	*
3306	62	*	*	*	*	*	*	*
3308	25	*	*	*	*	*	*	*
3312	153	0.0	2.0	7.2	53.6	30.7	3.9	2.6
3314	1	*	*	*	*	*	*	*
3550	10	*	*	*	*	*	*	*
3701	269	0.0	1.9	7.8	59.1	27.5	1.9	1.9
3850	849	0.0	2.4	6.4	60.0	26.0	0.8	4.5
3901	449	0.0	2.4	8.0	59.0	25.8	2.7	2.0
3950	699	0.1	1.9	4.9	60.1	28.8	1.7	2.6
4106	752	0.1	2.3	6.4	57.7	28.9	2.9	1.7

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2023 Pregnancy Nutrition Surveillance (1)

Page : 13

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4107	882	0.2	1.7	5.6	56.6	32.0	2.2	1.8
4109	241	0.0	1.7	2.5	55.6	36.5	2.9	0.8
4112	201	0.0	3.5	8.0	54.2	27.9	4.5	2.0
4115	1,110	0.1	1.1	4.0	53.4	37.2	3.4	0.8
4351	170	0.0	0.6	6.5	65.3	20.6	2.9	4.1
4352	58	*	*	*	*	*	*	*
4353	164	0.0	0.6	6.7	53.0	34.1	4.9	0.6
4354	251	0.0	2.4	6.0	63.3	24.3	2.4	1.6
4355	97	*	*	*	*	*	*	*
4356	128	0.0	0.8	8.6	58.6	28.1	0.8	3.1
4359	169	0.6	1.2	4.7	62.7	26.6	3.6	0.6
4360	78	*	*	*	*	*	*	*
4361	74	*	*	*	*	*	*	*
4362	83	*	*	*	*	*	*	*
4363	219	0.0	1.4	5.9	64.4	24.7	1.4	2.3
4401	244	0.0	1.2	4.9	62.3	29.1	0.8	1.6
4501	248	0.0	1.2	6.0	67.7	21.0	1.2	2.8
4601	391	0.0	1.5	6.9	62.9	26.3	1.8	0.5
4701	321	0.0	1.2	4.7	55.5	35.8	2.5	0.3
5050	547	0.0	1.1	2.9	57.0	36.2	2.4	0.4
5051	822	0.1	0.7	3.4	54.9	38.3	1.7	0.9
5052	892	0.0	0.7	3.5	56.4	35.8	2.1	1.6
5055	445	0.0	0.0	1.8	44.5	50.6	3.1	0.0
5101	489	0.0	1.2	2.9	55.8	36.6	3.1	0.4

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2023 Pregnancy Nutrition Surveillance (1)

Page : 14

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5202	199	0.0	1.0	6.5	60.3	30.2	2.0	0.0
5601	305	0.0	0.3	5.6	60.3	32.1	1.0	0.7
5801	533	0.0	0.9	6.9	63.4	26.3	1.5	0.9
5803	3	*	*	*	*	*	*	*
5804	1	*	*	*	*	*	*	*
5806	32	*	*	*	*	*	*	*
6101	18	*	*	*	*	*	*	*
6102	14	*	*	*	*	*	*	*
6103	1,040	0.3	2.2	5.4	62.0	25.3	1.7	3.1
6105	1	*	*	*	*	*	*	*
6301	1,128	0.2	1.8	6.2	56.6	32.5	1.8	1.0
6312	600	0.0	0.8	4.8	46.3	43.2	4.3	0.5
6316	388	0.0	0.3	2.6	50.5	41.2	5.4	0.0
6318	10	*	*	*	*	*	*	*
6320	185	0.0	0.5	4.3	48.1	41.1	4.9	1.1
6325	110	0.0	0.9	0.9	64.5	30.0	0.0	3.6
6335	308	0.0	1.0	3.9	48.1	44.5	1.9	0.6
6345	60	*	*	*	*	*	*	*
6360	262	0.0	0.8	1.9	58.8	34.0	2.7	1.9
7306	539	0.4	2.6	5.6	54.5	33.6	2.8	0.6
7401	558	0.0	2.0	5.2	61.5	28.1	1.4	1.8
7601	152	0.0	1.3	7.9	63.8	22.4	2.6	2.0
7850	346	0.0	1.7	6.4	64.2	25.4	1.4	0.9
7852	2	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2023 Pregnancy Nutrition Surveillance (1)

Page : 15

Michigan

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8101	535	0.0	2.1	7.5	51.6	34.0	3.4	1.5
8102	99	*	*	*	*	*	*	*
8106	157	0.0	0.6	6.4	53.5	36.3	3.2	0.0
8150	1	*	*	*	*	*	*	*
8151	1,044	0.0	1.4	4.8	51.4	37.3	4.2	0.9
8155	2	*	*	*	*	*	*	*
8201	768	0.0	1.0	5.2	53.4	36.3	3.0	1.0
8202	1,307	0.0	0.8	3.0	49.0	41.5	5.2	0.4
8203	123	0.0	1.6	5.7	61.0	27.6	3.3	0.8
8250	137	0.0	5.1	5.8	53.3	32.8	2.9	0.0
8252	28	*	*	*	*	*	*	*
8255	617	0.0	0.8	3.6	58.7	34.0	2.8	0.2
8257	512	0.0	0.4	1.6	53.7	38.7	5.7	0.0
8263	101	0.0	1.0	8.9	59.4	28.7	2.0	0.0
8264	286	0.0	0.3	3.8	61.2	31.5	2.8	0.3
8267	280	0.0	0.7	4.3	53.6	38.2	2.5	0.7
8901	232	0.0	2.2	5.6	49.6	36.6	3.4	2.6
8902	384	0.0	0.8	4.9	45.8	43.8	3.9	0.8
8903	399	0.0	2.3	4.8	52.6	35.8	2.0	2.5
8904	584	0.9	1.5	6.7	56.0	32.2	1.7	1.0
8905	510	0.0	0.6	2.9	55.7	38.0	2.4	0.4
8909	348	0.0	1.7	6.9	53.7	33.6	2.9	1.1
9045	437	0.0	2.3	5.5	53.5	34.6	3.0	1.1
9050	262	0.0	2.3	5.7	51.9	35.9	3.1	1.1

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS
Table 4A
2023 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic

Run Date: 6/28/2024

Page : 16

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9051	631	0.0	2.1	5.9	52.8	33.8	3.6	1.9
9053	591	0.0	1.7	3.9	58.5	31.6	2.7	1.5
9055	3	*	*	*	*	*	*	*
9058	460	0.0	0.7	5.0	51.5	39.8	2.6	0.4
9066	145	0.0	1.4	4.8	45.5	45.5	2.8	0.0
9068	298	0.0	2.7	5.7	53.0	33.6	1.7	3.4
9069	539	0.2	2.2	3.2	54.0	37.1	3.2	0.2
9070	666	0.0	2.0	7.1	53.8	33.3	1.8	2.1
9080	939	0.0	1.4	6.5	52.3	35.9	3.1	0.9
9082	38	*	*	*	*	*	*	*
9083	7	*	*	*	*	*	*	*
Michigan	46,161	0.1	1.5	5.3	56.9	32.3	2.5	1.4
National (Prior Year)	57,001	0.1	1.8	6.0	58.2	30.9	2.3	0.6

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 17

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	41	*	*	20	*	*	5	*	12	*	48	*	48	*
0102	88	*	*	34	*	*	12	*	25	*	105	9.5 (58)	106	13.2 (33)
0103	104	4.8 (66)	67.3 (96)	50	*	*	17	*	29	*	129	14.7 (101)	129	13.2 (33)
0104	34	*	*	11	*	*	3	*	15	*	41	*	41	*
0105	1	*	*	0	*	*	0	*	0	*	1	*	1	*
0250	32	*	*	12	*	*	3	*	12	*	35	*	35	*
0251	18	*	*	7	*	*	2	*	13	*	25	*	25	*
0252	33	*	*	26	*	*	0	*	6	*	32	*	32	*
0253	47	*	*	36	*	*	8	*	8	*	45	*	45	*
0401	120	5.0 (74)	61.7 (50)	36	*	*	19	*	28	*	120	10.8 (79)	121	20.7 (95)
0402	69	*	*	8	*	*	15	*	21	*	87	*	87	*
0403	32	*	*	19	*	*	5	*	4	*	33	*	33	*
0404	29	*	*	5	*	*	5	*	8	*	28	*	28	*
0501	102	3.9 (35)	57.8 (19)	16	*	*	20	*	32	*	126	7.1 (22)	126	12.7 (26)
0503	15	*	*	2	*	*	1	*	4	*	16	*	16	*
0504	90	*	*	39	*	*	16	*	21	*	90	*	90	*
0505	44	*	*	16	*	*	8	*	20	*	54	*	54	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 18

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	35	*	*	5	*	*	11	*	12	*	50	*	50	*
0507	11	*	*	3	*	*	2	*	3	*	20	*	20	*
0508	1	*	*	0	*	*	0	*	0	*	4	*	4	*
0650	273	4.0 (37)	65.2 (82)	119	11.8 (6)	59.7 (59)	45	*	93	*	317	12.3 (91)	317	18.0 (83)
0651	123	8.1 (101)	56.9 (16)	25	*	*	20	*	44	*	148	12.8 (94)	148	17.6 (80)
0652	65	*	*	52	*	*	16	*	12	*	78	*	78	*
0653	106	6.6 (96)	60.4 (33)	68	*	*	14	*	21	*	114	14.9 (103)	115	17.4 (78)
0654	106	6.6 (96)	59.4 (27)	75	*	*	11	*	42	*	129	12.4 (92)	129	24.0 (105)
0655	102	6.9 (99)	52.9 (5)	15	*	*	13	*	10	*	99	*	99	*
0657	6	*	*	5	*	*	0	*	3	*	7	*	7	*
0701	115	2.6 (8)	53.0 (6)	62	*	*	8	*	33	*	119	8.4 (35)	117	18.8 (87)
0702	27	*	*	2	*	*	2	*	4	*	29	*	29	*
0703	48	*	*	19	*	*	6	*	12	*	54	*	54	*
0704	19	*	*	11	*	*	1	*	3	*	16	*	16	*
0705	16	*	*	6	*	*	3	*	8	*	21	*	21	*
0706	1	*	*	1	*	*	0	*	0	*	1	*	1	*
0850	156	3.8 (31)	66.0 (89)	33	*	*	8	*	38	*	200	8.5 (37)	200	15.5 (61)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 19

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0851	147	5.4 (81)	60.5 (36)	42	*	*	12	*	45	*	202	10.4 (72)	201	16.9 (72)
0901	316	3.8 (31)	62.3 (56)	174	16.7 (31)	58.0 (52)	23	*	66	*	406	7.9 (29)	407	17.0 (73)
0902	3	*	*	3	*	*	0	*	1	*	3	*	3	*
1051	278	5.4 (81)	56.8 (14)	177	15.3 (19)	54.8 (36)	37	*	64	*	311	6.8 (17)	314	13.4 (39)
1101	366	5.2 (78)	66.4 (92)	226	18.6 (43)	52.2 (19)	46	*	91	*	397	8.8 (42)	398	19.6 (91)
1102	183	5.5 (86)	58.5 (21)	121	18.2 (42)	54.5 (33)	25	*	43	*	193	8.3 (33)	193	15.0 (52)
1106	18	*	*	14	*	*	0	*	7	*	17	*	19	*
1250	149	5.4 (81)	62.4 (59)	55	*	*	49	*	71	*	318	9.1 (45)	318	9.4 (6)
1251	219	3.7 (29)	61.6 (47)	177	15.3 (19)	57.6 (50)	28	*	39	*	215	16.3 (106)	216	11.1 (13)
1252	127	7.1 (100)	61.4 (44)	61	*	*	23	*	39	*	150	8.7 (40)	150	12.0 (19)
1253	181	5.5 (86)	56.9 (16)	96	*	*	35	*	54	*	190	6.3 (10)	189	14.3 (46)
1350	668	5.4 (81)	59.0 (23)	487	17.9 (41)	55.9 (42)	52	*	70	*	668	9.7 (62)	668	12.9 (29)
1352	85	*	*	54	*	*	8	*	11	*	87	*	87	*
1401	65	*	*	49	*	*	4	*	14	*	67	*	67	*
1402	13	*	*	7	*	*	4	*	1	*	23	*	23	*
1403	2	*	*	2	*	*	0	*	0	*	2	*	2	*
1501	201	4.0 (37)	64.7 (78)	87	*	*	37	*	43	*	234	13.2 (97)	235	15.3 (59)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 20

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1503	217	6.0 (92)	63.1 (63)	158	19.6 (49)	57.6 (50)	40	*	41	*	235	11.1 (82)	234	13.2 (33)
1504	231	4.8 (66)	51.9 (3)	66	*	*	79	*	120	19.2 (6)	549	10.2 (68)	550	8.7 (4)
1506	142	2.8 (13)	56.3 (12)	26	*	*	37	*	46	*	156	9.6 (60)	156	17.9 (82)
1507	115	8.7 (102)	57.4 (18)	19	*	*	19	*	43	*	159	15.1 (104)	159	11.9 (18)
1509	72	*	*	31	*	*	16	*	26	*	100	7.0 (20)	100	7.0 (3)
1511	154	3.2 (18)	65.6 (85)	98	*	*	27	*	44	*	178	14.0 (98)	178	12.9 (29)
1512	74	*	*	40	*	*	12	*	50	*	112	11.6 (88)	112	8.9 (5)
1514	115	2.6 (8)	54.8 (9)	33	*	*	22	*	86	*	204	8.3 (33)	204	15.7 (64)
1701	141	2.1 (6)	63.1 (63)	100	10.0 (2)	57.0 (47)	17	*	47	*	144	11.8 (89)	144	11.1 (13)
1950	156	2.6 (8)	64.7 (78)	105	14.3 (12)	61.0 (60)	10	*	30	*	162	9.3 (53)	162	16.7 (71)
1952	40	*	*	26	*	*	2	*	7	*	42	*	42	*
1953	90	*	*	66	*	*	4	*	28	*	97	*	97	*
1954	110	1.8 (2)	68.2 (99)	61	*	*	7	*	26	*	107	7.5 (25)	107	10.3 (8)
1955	39	*	*	26	*	*	0	*	10	*	42	*	42	*
1957	102	2.9 (14)	69.6 (101)	66	*	*	7	*	24	*	105	14.3 (99)	106	9.4 (6)
1963	28	*	*	18	*	*	0	*	8	*	27	*	27	*
2150	171	1.2 (1)	62.0 (52)	123	23.6 (62)	51.2 (17)	25	*	44	*	172	10.5 (73)	173	15.0 (52)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 21

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2151	75	*	*	62	*	*	12	*	23	*	71	*	71	*
2152	8	*	*	4	*	*	0	*	1	*	6	*	6	*
2251	100	4.0 (37)	73.0 (102)	70	*	*	0	*	6	*	100	4.0 (1)	100	14.0 (45)
2252	34	*	*	11	*	*	0	*	0	*	54	*	54	*
2554	5	*	*	0	*	*	1	*	1	*	9	*	9	*
2555	1,747	5.2 (78)	60.4 (33)	743	19.4 (48)	52.9 (22)	82	*	347	27.7 (17)	2,080	9.7 (62)	2,082	17.1 (75)
2556	16	*	*	5	*	*	0	*	2	*	20	*	20	*
2563	0	*	*	0	*	*	0	*	0	*	1	*	1	*
2564	1	*	*	0	*	*	0	*	0	*	1	*	1	*
3001	252	4.8 (66)	67.9 (98)	154	20.1 (52)	46.8 (7)	11	*	44	*	334	9.3 (53)	333	19.2 (89)
3002	159	6.3 (94)	56.6 (13)	88	*	*	11	*	30	*	188	5.9 (7)	188	12.8 (28)
3004	66	*	*	34	*	*	3	*	8	*	109	4.6 (2)	109	15.6 (62)
3005	32	*	*	17	*	*	1	*	6	*	49	*	49	*
3007	2	*	*	1	*	*	0	*	0	*	2	*	2	*
3201	122	6.6 (96)	64.8 (80)	54	*	*	20	*	39	*	142	9.2 (49)	142	11.3 (15)
3301	1,158	3.6 (25)	61.0 (41)	716	16.6 (30)	57.5 (49)	82	*	116	14.7 (1)	1,331	7.1 (22)	1,334	13.3 (37)
3304	23	*	*	12	*	*	1	*	0	*	24	*	24	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 22

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3305	44	*	*	29	*	*	3	*	1	*	46	*	46	*
3306	54	*	*	30	*	*	2	*	4	*	60	*	60	*
3308	23	*	*	15	*	*	4	*	2	*	21	*	22	*
3312	134	3.7 (29)	58.2 (20)	77	*	*	6	*	9	*	139	8.6 (38)	138	20.3 (93)
3314	1	*	*	0	*	*	0	*	0	*	1	*	1	*
3550	8	*	*	6	*	*	1	*	2	*	8	*	8	*
3701	230	3.0 (17)	61.7 (50)	175	18.9 (45)	50.3 (12)	43	*	77	*	256	10.2 (68)	256	16.0 (65)
3850	718	5.8 (91)	60.7 (37)	483	14.5 (14)	58.4 (55)	35	*	94	*	783	10.5 (73)	785	14.8 (49)
3901	359	3.6 (25)	59.3 (24)	178	11.8 (6)	65.7 (65)	18	*	67	*	409	7.3 (24)	408	15.4 (60)
3950	544	4.8 (66)	59.7 (30)	290	14.5 (14)	58.3 (53)	36	*	98	*	629	10.2 (68)	631	13.6 (41)
4106	275	4.4 (49)	60.4 (33)	65	*	*	109	33.9 (2)	220	15.0 (2)	704	9.1 (45)	706	20.1 (92)
4107	561	3.9 (35)	60.1 (32)	180	17.8 (39)	51.1 (16)	158	34.2 (3)	292	20.5 (9)	810	7.7 (26)	813	22.1 (98)
4109	131	4.6 (59)	59.5 (28)	71	*	*	30	*	89	*	230	9.6 (60)	229	18.8 (87)
4112	201	2.0 (4)	67.7 (97)	165	55.8 (65)	23.0 (1)	3	*	16	*	186	24.7 (107)	186	26.9 (106)
4115	567	5.5 (86)	61.6 (47)	188	17.6 (36)	46.3 (6)	155	34.8 (4)	390	25.6 (14)	1,057	9.1 (45)	1,058	17.0 (73)
4351	148	6.1 (93)	59.5 (28)	93	*	*	4	*	20	*	158	14.6 (100)	158	28.5 (107)
4352	53	*	*	41	*	*	1	*	7	*	50	*	50	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 23

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4353	122	4.9 (71)	66.4 (92)	88	*	*	10	*	26	*	151	11.9 (90)	151	13.2 (33)
4354	192	6.3 (94)	63.5 (67)	43	*	*	6	*	31	*	231	14.7 (101)	231	23.4 (103)
4355	90	*	*	57	*	*	5	*	19	*	88	*	88	*
4356	117	3.4 (22)	65.0 (81)	92	*	*	6	*	19	*	118	9.3 (53)	119	17.6 (80)
4359	150	4.0 (37)	66.0 (89)	109	24.8 (63)	47.7 (10)	5	*	26	*	156	7.7 (26)	156	23.1 (101)
4360	72	*	*	54	*	*	7	*	19	*	75	*	75	*
4361	69	*	*	56	*	*	3	*	11	*	72	*	72	*
4362	78	*	*	62	*	*	4	*	19	*	78	*	78	*
4363	201	4.5 (53)	55.7 (11)	138	16.7 (31)	55.8 (41)	6	*	43	*	204	9.8 (64)	204	11.8 (17)
4401	241	5.4 (81)	61.4 (44)	203	13.8 (10)	54.2 (29)	23	*	43	*	228	7.0 (20)	229	13.5 (40)
4501	196	4.6 (59)	63.3 (65)	106	20.8 (55)	54.7 (35)	25	*	34	*	232	9.9 (66)	232	12.1 (21)
4601	316	5.1 (76)	58.5 (21)	188	14.9 (16)	55.9 (42)	46	*	112	25.0 (13)	343	11.1 (82)	343	17.2 (77)
4701	257	4.7 (63)	56.8 (14)	147	20.4 (54)	54.4 (32)	24	*	78	*	304	11.5 (84)	305	22.3 (99)
5050	529	4.2 (44)	69.4 (100)	436	15.1 (17)	54.8 (36)	25	*	217	32.3 (21)	500	9.2 (49)	500	18.6 (86)
5051	758	4.4 (49)	62.4 (59)	552	15.9 (26)	55.3 (39)	57	*	341	28.7 (19)	755	9.3 (53)	754	16.3 (67)
5052	837	2.9 (14)	61.1 (42)	682	22.1 (59)	46.8 (7)	79	*	272	24.3 (11)	835	12.9 (95)	833	14.8 (49)
5055	411	2.4 (7)	62.3 (56)	347	22.5 (60)	43.8 (4)	20	*	61	*	429	15.6 (105)	429	10.7 (10)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 24

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5101	406	4.4 (49)	62.3 (56)	298	11.4 (3)	57.4 (48)	50	*	233	25.8 (16)	448	5.8 (6)	449	12.7 (26)
5202	162	5.6 (90)	52.5 (4)	99	*	*	2	*	35	*	180	6.7 (15)	182	11.0 (12)
5601	242	2.9 (14)	65.3 (84)	106	8.5 (1)	63.2 (63)	30	*	89	*	284	8.8 (42)	283	14.8 (49)
5801	502	5.2 (78)	65.7 (87)	389	21.1 (57)	50.9 (14)	38	*	113	16.8 (3)	475	8.8 (42)	477	22.4 (100)
5803	3	*	*	1	*	*	0	*	0	*	3	*	3	*
5804	1	*	*	1	*	*	0	*	0	*	1	*	1	*
5806	29	*	*	19	*	*	0	*	7	*	30	*	30	*
6101	16	*	*	13	*	*	1	*	1	*	18	*	18	*
6102	11	*	*	9	*	*	1	*	2	*	12	*	12	*
6103	896	4.6 (59)	63.6 (70)	640	17.7 (38)	54.2 (29)	35	*	42	*	950	10.0 (67)	949	20.8 (96)
6105	1	*	*	1	*	*	0	*	0	*	1	*	1	*
6301	976	4.5 (53)	61.2 (43)	672	16.1 (27)	53.1 (24)	88	*	202	25.7 (15)	1,001	6.6 (14)	1,001	12.4 (24)
6312	535	4.5 (53)	59.3 (24)	357	19.0 (46)	51.0 (15)	59	*	150	37.3 (22)	560	9.5 (58)	560	12.5 (25)
6316	347	4.9 (71)	54.5 (7)	246	15.4 (22)	56.1 (44)	24	*	69	*	357	9.8 (64)	357	10.9 (11)
6318	8	*	*	5	*	*	1	*	0	*	9	*	9	*
6320	148	2.7 (12)	61.5 (46)	65	*	*	10	*	34	*	165	11.5 (84)	165	19.4 (90)
6325	82	*	*	43	*	*	6	*	25	*	95	*	95	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 25

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)		(5)	(6)										
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6335	243	4.5 (53)	54.7 (8)	105	11.4 (3)	49.5 (11)	29	*	66	*	276	12.7 (93)	278	13.7 (43)
6345	48	*	*	24	*	*	3	*	10	*	51	*	51	*
6360	206	4.9 (71)	55.3 (10)	109	15.6 (24)	46.8 (7)	21	*	51	*	242	8.7 (40)	241	16.6 (70)
7306	470	4.0 (37)	66.2 (91)	372	19.6 (49)	54.6 (34)	60	*	70	*	511	11.5 (84)	511	15.1 (55)
7401	528	3.8 (31)	63.6 (70)	405	15.3 (19)	56.8 (46)	44	*	148	29.7 (20)	495	10.5 (73)	497	13.9 (44)
7601	95	*	*	10	*	*	11	*	42	*	138	9.4 (57)	138	12.3 (23)
7850	290	5.5 (86)	62.4 (59)	227	21.1 (57)	52.9 (22)	21	*	57	*	323	8.4 (35)	325	11.7 (16)
7852	2	*	*	1	*	*	0	*	0	*	1	*	1	*
8101	448	3.8 (31)	64.3 (76)	276	11.6 (5)	58.3 (53)	23	*	67	*	491	8.6 (38)	491	16.5 (68)
8102	84	*	*	46	*	*	2	*	8	*	91	*	91	*
8106	70	*	*	28	*	*	8	*	27	*	147	8.2 (31)	147	23.8 (104)
8150	1	*	*	0	*	*	0	*	1	*	1	*	1	*
8151	808	4.3 (46)	62.5 (62)	523	20.3 (53)	53.2 (26)	138	32.6 (1)	269	17.8 (4)	942	8.2 (31)	942	15.2 (56)
8155	2	*	*	0	*	*	0	*	0	*	2	*	2	*
8201	629	5.1 (76)	60.9 (39)	320	16.9 (35)	53.8 (27)	79	*	288	23.6 (10)	665	11.0 (81)	668	18.0 (83)
8202	307	4.6 (59)	48.2 (1)	110	20.9 (56)	44.5 (5)	74	*	175	18.3 (5)	1,261	6.1 (9)	1,261	3.4 (1)
8203	115	4.3 (46)	63.5 (67)	90	*	*	7	*	52	*	109	9.2 (49)	108	18.5 (85)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 26

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8250	111	3.6 (25)	64.0 (73)	58	*	*	10	*	50	*	124	8.1 (30)	124	14.5 (48)
8252	10	*	*	6	*	*	0	*	2	*	26	*	26	*
8255	552	4.0 (37)	65.2 (82)	294	14.3 (12)	58.5 (56)	69	*	234	24.8 (12)	549	10.6 (77)	551	16.5 (68)
8257	367	4.4 (49)	51.5 (2)	256	23.4 (61)	40.6 (3)	51	*	103	19.4 (7)	475	9.1 (45)	478	3.6 (2)
8263	81	*	*	33	*	*	0	*	27	*	77	*	77	*
8264	253	4.7 (63)	64.4 (77)	188	13.3 (9)	61.7 (62)	28	*	129	19.4 (7)	256	10.2 (68)	256	12.1 (21)
8267	253	4.7 (63)	65.6 (85)	136	19.1 (47)	52.2 (19)	45	*	127	28.3 (18)	258	7.8 (28)	258	20.9 (97)
8901	200	2.0 (4)	67.0 (94)	114	15.8 (25)	55.3 (39)	4	*	19	*	220	10.9 (80)	220	20.5 (94)
8902	311	4.8 (66)	62.1 (53)	155	16.8 (34)	52.3 (21)	17	*	56	*	347	4.9 (4)	347	15.6 (62)
8903	309	3.2 (18)	63.8 (72)	193	13.0 (8)	63.7 (64)	12	*	30	*	374	5.9 (7)	374	13.6 (41)
8904	474	3.4 (22)	63.3 (65)	266	16.2 (28)	53.8 (27)	23	*	67	*	539	6.5 (13)	539	17.4 (78)
8905	423	2.6 (8)	59.3 (24)	210	16.7 (31)	54.3 (31)	15	*	66	*	468	13.0 (96)	468	15.0 (52)
8909	296	4.1 (43)	62.2 (54)	190	14.2 (11)	61.1 (61)	14	*	43	*	318	6.9 (19)	318	16.0 (65)
9045	334	4.5 (53)	67.1 (95)	213	18.8 (44)	53.1 (24)	12	*	37	*	400	10.5 (73)	399	12.0 (19)
9050	219	5.0 (74)	63.5 (67)	69	*	*	10	*	23	*	239	6.3 (10)	238	13.0 (32)
9051	305	4.3 (46)	65.9 (88)	138	26.8 (64)	39.1 (2)	27	*	95	*	590	11.5 (84)	590	12.9 (29)
9053	540	3.3 (21)	59.8 (31)	447	15.2 (18)	55.0 (38)	23	*	41	*	567	6.3 (10)	567	10.4 (9)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2023 Pregnancy Nutrition Surveillance (1)

Page : 27

Michigan

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9055	2	*	*	1	*	*	0	*	0	*	2	*	2	*
9058	330	4.5 (53)	60.9 (39)	247	17.8 (39)	51.8 (18)	12	*	46	*	433	9.2 (49)	434	17.1 (75)
9066	125	3.2 (18)	61.6 (47)	84	*	*	0	*	7	*	135	6.7 (15)	134	23.1 (101)
9068	284	1.8 (2)	64.1 (75)	244	16.4 (29)	59.0 (58)	11	*	20	*	269	5.6 (5)	270	15.2 (56)
9069	466	3.6 (25)	62.2 (54)	345	19.7 (51)	50.4 (13)	28	*	54	*	502	6.8 (17)	502	13.3 (37)
9070	608	3.5 (24)	64.0 (73)	470	15.5 (23)	58.7 (57)	25	*	64	*	616	4.7 (3)	618	15.2 (56)
9080	743	4.2 (44)	60.7 (37)	358	17.6 (36)	56.4 (45)	37	*	98	*	876	10.6 (77)	876	14.3 (46)
9082	32	*	*	13	*	*	0	*	4	*	37	*	37	*
9083	7	*	*	5	*	*	0	*	2	*	6	*	6	*
Michigan	35,984	4.3	61.6	21,505	17.5	54.1	3,531	34.3	9,106	23.2	42,426	9.3	42,465	15.0
National (Prior Year)	43,167	4.0	61.9	28,282	19.8	49.9	3,540	26.3	4,123	24.5	45,204	10.0	45,068	15.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2023 Pregnancy Nutrition Surveillance (1)

Page : 28

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
0101	47	*	*	50	*	52	*	35 *
0102	95	*	*	116	29.3 (57)	115	27.8 (37)	90 *
0103	122	93.4 (43)	0.0 (1)	131	36.6 (25)	131	23.7 (59)	105 94.3 (17)
0104	39	*	*	43	*	43	*	33 *
0105	1	*	*	1	*	1	*	1 *
0250	32	*	*	36	*	40	*	32 *
0251	25	*	*	28	*	28	*	20 *
0252	32	*	*	36	*	35	*	28 *
0253	42	*	*	47	*	47	*	41 *
0401	116	94.8 (27)	0.0 (1)	124	20.2 (98)	130	31.5 (24)	81 *
0402	75	*	*	92	*	92	*	79 *
0403	32	*	*	36	*	34	*	29 *
0404	29	*	*	32	*	32	*	22 *
0501	120	97.5 (8)	0.0 (1)	130	32.3 (43)	119	31.1 (25)	102 88.2 (73)
0503	16	*	*	18	*	18	*	13 *
0504	88	*	*	96	*	94	*	79 *
0505	50	*	*	54	*	54	*	36 *
0506	45	*	*	52	*	52	*	42 *
0507	20	*	*	20	*	21	*	13 *
0508	4	*	*	4	*	4	*	3 *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2023 Pregnancy Nutrition Surveillance (1)

Page : 29

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	302	93.7 (40)	0.3 (39)	337	32.0 (47)	332	29.5 (29)	264	91.3 (39)
0651	140	90.7 (71)	1.4 (92)	158	36.7 (22)	132	24.2 (55)	120	86.7 (84)
0652	73	*	*	82	*	83	*	68	*
0653	100	94.0 (36)	2.0 (101)	122	38.5 (17)	120	35.8 (14)	100	88.0 (75)
0654	103	85.4 (93)	0.0 (1)	136	44.1 (5)	135	15.6 (90)	112	93.8 (23)
0655	92	*	*	107	41.1 (9)	107	47.7 (4)	91	*
0657	5	*	*	7	*	7	*	7	*
0701	103	95.1 (22)	0.0 (1)	136	33.8 (35)	134	44.0 (6)	102	89.2 (62)
0702	25	*	*	31	*	32	*	27	*
0703	54	*	*	62	*	60	*	45	*
0704	16	*	*	21	*	20	*	19	*
0705	21	*	*	28	*	26	*	19	*
0706	1	*	*	1	*	1	*	0	*
0850	198	90.9 (68)	1.5 (94)	219	27.9 (67)	218	28.4 (34)	149	94.6 (15)
0851	199	97.5 (8)	0.0 (1)	215	30.7 (51)	216	38.0 (10)	153	92.8 (25)
0901	364	91.8 (62)	1.1 (86)	430	41.2 (8)	428	24.8 (51)	339	88.5 (68)
0902	3	*	*	3	*	3	*	1	*
1051	286	91.3 (65)	1.4 (92)	345	29.3 (57)	344	26.2 (45)	263	87.5 (80)
1101	328	81.4 (101)	1.8 (100)	450	25.8 (80)	331	24.5 (52)	297	90.2 (54)
1102	157	89.8 (74)	0.6 (64)	213	37.1 (21)	164	23.2 (63)	150	92.7 (29)
1106	19	*	*	21	*	18	*	14	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2023 Pregnancy Nutrition Surveillance (1)

Page : 30

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1250	290	92.1 (59)	0.3 (39)	334	29.3 (57)	338	44.1 (5)	245	93.1 (24)
1251	209	94.7 (28)	0.0 (1)	230	30.0 (55)	233	28.8 (32)	173	90.2 (54)
1252	109	88.1 (86)	0.9 (78)	161	37.3 (20)	160	30.6 (26)	119	89.1 (66)
1253	154	90.9 (68)	0.0 (1)	199	36.7 (22)	204	20.1 (81)	155	87.1 (82)
1350	653	95.1 (22)	0.5 (52)	726	36.4 (26)	620	29.8 (28)	564	92.7 (29)
1352	86	*	*	94	*	92	*	74	*
1401	64	*	*	75	*	74	*	55	*
1402	23	*	*	26	*	25	*	20	*
1403	2	*	*	2	*	2	*	1	*
1501	187	94.7 (28)	0.5 (52)	259	34.4 (32)	248	41.1 (7)	194	87.1 (82)
1503	220	97.3 (10)	0.0 (1)	253	31.6 (48)	239	23.8 (58)	186	80.6 (96)
1504	532	97.9 (7)	0.2 (35)	591	26.7 (74)	576	20.7 (77)	408	91.9 (34)
1506	134	95.5 (18)	0.0 (1)	169	26.6 (75)	169	16.6 (87)	124	91.9 (34)
1507	156	98.1 (5)	0.0 (1)	163	36.2 (27)	156	37.2 (12)	113	97.3 (1)
1509	89	*	*	108	32.4 (42)	104	13.5 (95)	78	*
1511	173	89.0 (82)	0.0 (1)	186	32.3 (43)	182	35.2 (16)	141	92.2 (32)
1512	104	93.3 (44)	1.0 (84)	117	23.1 (86)	115	32.2 (23)	72	*
1514	195	94.9 (25)	0.5 (52)	208	27.9 (67)	207	33.3 (19)	135	89.6 (57)
1701	143	98.6 (3)	0.7 (69)	159	40.9 (10)	159	62.3 (2)	120	94.2 (19)
1950	152	95.4 (20)	0.0 (1)	171	33.9 (34)	161	58.4 (3)	132	96.2 (4)
1952	39	*	*	41	*	42	*	30	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2023 Pregnancy Nutrition Surveillance (1)

Page : 31

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
1953	92	*	*	102	20.6 (97)	96	*	76 *
1954	102	98.0 (6)	0.0 (1)	116	44.0 (6)	105	68.6 (1)	98 *
1955	42	*	*	44	*	43	*	25 *
1957	100	93.0 (52)	0.0 (1)	116	34.5 (31)	90	*	85 *
1963	27	*	*	30	*	29	*	19 *
2150	169	94.1 (35)	0.0 (1)	181	34.8 (30)	180	33.3 (19)	136 94.1 (20)
2151	66	*	*	75	*	49	*	57 *
2152	6	*	*	9	*	9	*	8 *
2251	94	*	*	109	36.7 (22)	111	39.6 (8)	92 *
2252	48	*	*	59	*	53	*	51 *
2554	9	*	*	11	*	8	*	10 *
2555	1,902	89.7 (75)	0.4 (46)	2,273	28.9 (61)	1,967	22.1 (68)	1,703 91.3 (39)
2556	16	*	*	21	*	17	*	17 *
2563	0	*	*	1	*	1	*	1 *
2564	1	*	*	1	*	1	*	1 *
3001	323	93.2 (48)	0.3 (39)	355	42.5 (7)	356	12.9 (96)	294 87.8 (76)
3002	182	95.1 (22)	0.0 (1)	196	32.1 (46)	197	14.2 (94)	160 90.0 (56)
3004	102	93.1 (50)	0.0 (1)	109	40.4 (13)	111	23.4 (61)	83 *
3005	47	*	*	52	*	54	*	43 *
3007	2	*	*	2	*	2	*	2 *
3201	117	89.7 (75)	0.9 (78)	145	25.5 (81)	145	37.9 (11)	110 91.8 (36)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2023 Pregnancy Nutrition Surveillance (1)

Page : 32

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3301	1,109	89.1 (81)	0.5 (52)	1,381	30.1 (54)	1,385	14.6 (93)	1,051	89.2 (62)
3304	17	*	*	23	*	23	*	20	*
3305	37	*	*	51	*	49	*	43	*
3306	56	*	*	60	*	62	*	52	*
3308	13	*	*	25	*	24	*	17	*
3312	116	93.1 (50)	0.0 (1)	151	40.4 (13)	151	17.2 (85)	121	88.4 (71)
3314	1	*	*	1	*	1	*	1	*
3550	8	*	*	10	*	10	*	8	*
3701	243	95.5 (18)	0.0 (1)	265	27.9 (67)	266	20.3 (79)	193	95.9 (6)
3850	717	92.9 (53)	0.7 (69)	845	38.5 (17)	839	20.9 (76)	653	89.6 (57)
3901	363	91.5 (63)	0.6 (64)	446	32.5 (41)	437	16.9 (86)	329	83.3 (93)
3950	595	90.9 (68)	0.5 (52)	690	30.0 (55)	676	23.4 (61)	502	89.2 (62)
4106	619	86.3 (92)	1.5 (94)	733	28.1 (63)	713	11.4 (100)	556	83.6 (91)
4107	758	93.3 (44)	0.5 (52)	869	24.7 (83)	806	36.8 (13)	652	92.5 (31)
4109	219	97.3 (10)	0.5 (52)	236	23.3 (85)	233	27.5 (39)	153	92.8 (25)
4112	187	77.0 (103)	0.0 (1)	199	20.1 (99)	3	*	173	86.7 (84)
4115	888	94.6 (31)	0.5 (52)	1,097	28.0 (65)	719	21.8 (70)	833	91.5 (38)
4351	133	88.7 (83)	1.5 (94)	168	51.2 (2)	146	21.9 (69)	140	84.3 (89)
4352	45	*	*	56	*	56	*	45	*
4353	115	92.2 (55)	0.9 (78)	161	37.9 (19)	160	30.6 (26)	126	91.3 (39)
4354	215	95.3 (21)	0.5 (52)	245	56.3 (1)	237	25.7 (49)	203	87.2 (81)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2023 Pregnancy Nutrition Surveillance (1)

Page : 33

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
4355	79	*	*	93	*	96	*	73 *
4356	95	*	*	122	44.3 (4)	124	21.8 (70)	98 *
4359	103	85.4 (93)	3.9 (103)	167	46.7 (3)	166	21.1 (75)	133 82.7 (94)
4360	65	*	*	76	*	78	*	60 *
4361	66	*	*	70	*	74	*	58 *
4362	68	*	*	83	*	80	*	69 *
4363	189	94.7 (28)	0.0 (1)	211	32.7 (38)	215	26.0 (48)	162 92.0 (33)
4401	192	92.2 (55)	0.0 (1)	242	32.6 (40)	242	29.3 (30)	193 94.3 (17)
4501	199	94.5 (32)	0.0 (1)	243	40.7 (11)	245	16.3 (88)	201 90.5 (50)
4601	314	92.0 (60)	0.3 (39)	387	31.0 (50)	384	20.6 (78)	290 91.7 (37)
4701	281	92.2 (55)	1.1 (86)	315	31.1 (49)	315	34.3 (17)	230 90.9 (44)
5050	485	94.0 (36)	0.8 (73)	530	20.9 (93)	521	15.5 (91)	360 95.3 (10)
5051	701	91.0 (67)	1.6 (99)	800	26.4 (77)	792	11.7 (99)	538 95.4 (9)
5052	802	91.1 (66)	0.4 (46)	869	20.8 (95)	834	12.1 (98)	595 93.9 (22)
5055	379	99.7 (1)	0.0 (1)	441	39.9 (16)	56	*	339 96.8 (2)
5101	428	93.7 (40)	0.2 (35)	468	20.9 (93)	309	34.0 (18)	335 95.2 (12)
5202	165	94.5 (32)	1.2 (88)	191	33.0 (37)	170	23.5 (60)	143 95.8 (7)
5601	263	95.8 (15)	0.8 (73)	303	25.1 (82)	302	26.5 (44)	217 91.2 (43)
5801	451	92.2 (55)	0.7 (69)	524	30.2 (53)	368	32.6 (21)	376 92.8 (25)
5803	2	*	*	3	*	2	*	3 *
5804	1	*	*	1	*	1	*	1 *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2023 Pregnancy Nutrition Surveillance (1)

Page : 34

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
5806	29	*	*	31	*	23	*	12 *
6101	15	*	*	18	*	18	*	14 *
6102	13	*	*	14	*	14	*	11 *
6103	892	90.7 (71)	0.4 (46)	1,025	35.8 (29)	1,006	21.2 (74)	792 94.7 (14)
6105	1	*	*	1	*	1	*	1 *
6301	920	94.9 (25)	0.4 (46)	1,115	29.0 (60)	984	25.5 (50)	808 90.7 (45)
6312	510	92.9 (53)	0.2 (35)	590	28.1 (63)	528	19.9 (82)	409 89.5 (59)
6316	324	95.7 (16)	1.2 (88)	375	28.0 (65)	334	28.4 (34)	256 90.6 (47)
6318	7	*	*	9	*	9	*	1 *
6320	158	96.8 (12)	0.0 (1)	185	30.3 (52)	181	29.3 (30)	149 95.3 (10)
6325	92	*	*	108	26.9 (71)	108	32.4 (22)	74 *
6335	272	96.0 (14)	0.4 (46)	304	28.6 (62)	300	28.3 (36)	220 94.5 (16)
6345	49	*	*	59	*	57	*	45 *
6360	231	94.4 (34)	0.9 (78)	257	21.4 (91)	249	24.1 (56)	181 95.6 (8)
7306	467	89.7 (75)	0.2 (35)	532	34.4 (32)	529	26.7 (42)	402 90.5 (50)
7401	390	89.5 (78)	1.5 (94)	550	40.5 (12)	552	24.1 (56)	431 90.7 (45)
7601	132	98.5 (4)	0.0 (1)	152	33.6 (36)	151	15.2 (92)	111 89.2 (62)
7850	304	96.1 (13)	1.0 (84)	340	40.0 (15)	138	39.1 (9)	255 82.7 (94)
7852	1	*	*	2	*	2	*	2 *
8101	465	83.9 (95)	0.9 (78)	524	20.8 (95)	428	22.4 (66)	353 88.1 (74)
8102	90	*	*	94	*	77	*	80 *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2023 Pregnancy Nutrition Surveillance (1)

Page : 35

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8106	132	83.3 (98)	1.5 (94)	154	19.5 (101)	156	22.4 (66)	103	96.1 (5)
8150	1	*	*	1	*	1	*	1	*
8151	888	93.2 (48)	0.5 (52)	1,023	27.5 (70)	1,014	20.2 (80)	755	89.3 (61)
8155	2	*	*	2	*	2	*	2	*
8201	645	93.3 (44)	0.6 (64)	758	15.8 (109)	610	26.2 (45)	492	87.6 (77)
8202	1,183	98.8 (2)	0.1 (34)	1,280	32.2 (45)	1,277	4.5 (103)	1,039	96.7 (3)
8203	105	89.5 (78)	0.0 (1)	123	26.8 (73)	116	26.7 (42)	81	*
8250	120	93.3 (44)	0.0 (1)	137	18.2 (106)	115	21.7 (72)	97	*
8252	22	*	*	28	*	27	*	25	*
8255	540	91.9 (61)	0.6 (64)	608	15.1 (110)	567	27.2 (40)	363	89.5 (59)
8257	472	95.6 (17)	0.0 (1)	507	24.5 (84)	387	3.6 (105)	409	69.9 (97)
8263	72	*	*	101	32.7 (38)	94	*	101	86.1 (87)
8264	250	94.0 (36)	0.0 (1)	284	18.3 (104)	260	28.8 (32)	188	83.5 (92)
8267	254	93.7 (40)	1.2 (88)	276	10.5 (111)	253	24.5 (52)	170	87.6 (77)
8901	215	80.5 (102)	0.9 (78)	228	21.5 (90)	204	11.3 (101)	158	88.6 (67)
8902	333	87.1 (89)	0.3 (39)	371	26.1 (79)	299	22.7 (65)	263	85.6 (88)
8903	360	82.8 (99)	0.3 (39)	390	20.0 (100)	270	24.4 (54)	270	84.1 (90)
8904	514	81.9 (100)	0.4 (46)	570	22.8 (87)	483	19.9 (82)	397	88.4 (71)
8905	447	94.0 (36)	0.0 (1)	501	26.9 (71)	438	35.4 (15)	358	90.5 (50)
8909	296	89.5 (78)	0.3 (39)	340	21.2 (92)	241	26.1 (47)	234	88.5 (68)
9045	377	91.5 (63)	0.8 (73)	431	26.5 (76)	163	12.3 (97)	327	95.1 (13)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2023 Pregnancy Nutrition Surveillance (1)

Page : 36

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9050	206	86.4 (91)	0.5 (52)	260	26.2 (78)	245	22.9 (64)	192	88.5 (68)
9051	555	90.6 (73)	0.5 (52)	616	22.4 (88)	434	0.7 (106)	488	92.8 (25)
9053	565	86.9 (90)	1.2 (88)	583	16.5 (108)	475	21.3 (73)	323	91.3 (39)
9055	2	*	*	3	*	3	*	2	*
9058	419	88.1 (86)	0.7 (69)	452	19.5 (101)	409	9.3 (102)	267	90.6 (47)
9066	128	88.3 (84)	0.0 (1)	144	36.1 (28)	115	27.0 (41)	145	87.6 (77)
9068	267	83.9 (95)	3.4 (102)	293	18.8 (103)	287	16.0 (89)	199	86.4 (86)
9069	499	87.8 (88)	0.6 (64)	532	17.5 (107)	422	4.3 (104)	357	94.1 (20)
9070	595	83.9 (95)	0.8 (73)	656	18.3 (104)	606	19.1 (84)	426	90.6 (47)
9080	836	88.2 (85)	0.8 (73)	935	22.0 (89)	689	27.7 (38)	668	90.4 (53)
9082	35	*	*	37	*	23	*	18	*
9083	6	*	*	7	*	7	*	7	*
Michigan	39,293	91.8	0.6	45,366	28.9	40,573	23.3	33,331	33331.0
National (Prior Year)	47,224	86.4	1.2	51,965	24.2	46,223	23.1	38,918	91.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 37

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	34	*	42	*	12	*	35	*	48	*
0102	89	*	96	*	27	*	90	*	107	8.4 (73)
0103	105	34.3 (94)	125	11.2 (83)	33	*	105	27.6 (96)	129	17.1 (104)
0104	33	*	41	*	11	*	33	*	41	*
0105	1	*	0	*	0	*	1	*	1	*
0250	27	*	34	*	14	*	32	*	35	*
0251	20	*	23	*	10	*	20	*	27	*
0252	28	*	31	*	6	*	28	*	33	*
0253	41	*	43	*	18	*	41	*	45	*
0401	78	*	115	13.9 (96)	22	*	81	*	121	6.6 (57)
0402	79	*	81	*	17	*	79	*	87	*
0403	29	*	32	*	12	*	29	*	33	*
0404	22	*	28	*	4	*	22	*	30	*
0501	101	19.8 (69)	124	6.5 (58)	19	*	102	17.6 (84)	126	11.1 (92)
0503	13	*	15	*	3	*	13	*	16	*
0504	77	*	87	*	19	*	78	*	90	*
0505	36	*	52	*	8	*	36	*	54	*
0506	42	*	48	*	17	*	42	*	51	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 38

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0507	13	*	19	*	1	*	13	*	20	*
0508	3	*	4	*	0	*	3	*	4	*
0650	255	21.6 (73)	309	11.7 (86)	59	*	266	13.5 (70)	317	10.7 (89)
0651	119	36.1 (95)	143	19.6 (102)	39	*	120	31.7 (97)	149	14.8 (99)
0652	67	*	75	*	16	*	68	*	77	*
0653	99	*	104	12.5 (91)	36	*	100	27.0 (95)	118	16.9 (103)
0654	112	25.0 (80)	120	20.0 (103)	27	*	112	25.0 (94)	129	27.1 (107)
0655	91	*	95	*	25	*	91	*	99	*
0657	7	*	7	*	0	*	7	*	7	*
0701	101	28.7 (88)	120	16.7 (98)	28	*	102	6.9 (26)	123	2.4 (19)
0702	27	*	29	*	9	*	27	*	29	*
0703	44	*	52	*	16	*	45	*	54	*
0704	19	*	15	*	5	*	19	*	16	*
0705	19	*	21	*	5	*	19	*	21	*
0706	0	*	1	*	0	*	0	*	1	*
0850	149	15.4 (44)	203	9.4 (77)	19	*	149	12.1 (66)	203	9.9 (81)
0851	149	12.8 (31)	201	6.0 (54)	21	*	152	11.8 (61)	202	5.4 (49)
0901	339	25.1 (81)	404	12.6 (92)	81	*	339	17.7 (85)	405	9.1 (79)
0902	1	*	3	*	0	*	1	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 39

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1051	261	26.1 (83)	304	12.2 (90)	62	*	263	17.5 (83)	319	10.7 (89)
1101	289	19.0 (61)	396	6.3 (55)	52	*	297	19.5 (90)	396	10.1 (84)
1102	146	24.7 (79)	191	13.6 (95)	36	*	151	21.9 (93)	193	10.4 (87)
1106	14	*	18	*	6	*	14	*	19	*
1250	244	15.6 (46)	309	7.1 (66)	35	*	245	14.7 (76)	318	10.7 (89)
1251	169	31.4 (91)	202	11.4 (84)	51	*	173	19.1 (88)	216	12.5 (97)
1252	118	16.1 (48)	147	6.8 (63)	16	*	119	10.1 (46)	150	10.0 (82)
1253	151	21.2 (72)	185	11.9 (87)	31	*	154	13.6 (71)	190	10.0 (82)
1350	561	18.9 (59)	662	5.9 (52)	100	57.0 (1)	564	7.3 (30)	666	4.8 (45)
1352	74	*	84	*	18	*	74	*	86	*
1401	55	*	63	*	20	*	55	*	67	*
1402	19	*	23	*	3	*	20	*	23	*
1403	1	*	2	*	0	*	1	*	2	*
1501	194	20.1 (70)	227	6.6 (59)	38	*	194	11.3 (57)	237	8.4 (73)
1503	185	21.1 (71)	235	8.9 (75)	37	*	186	10.8 (52)	237	2.1 (13)
1504	407	8.8 (18)	547	2.4 (21)	34	*	408	3.4 (5)	554	2.2 (14)
1506	124	19.4 (66)	148	9.5 (78)	23	*	124	7.3 (30)	158	10.1 (84)
1507	109	26.6 (84)	148	3.4 (32)	30	*	113	11.5 (59)	160	11.9 (95)
1509	78	*	98	*	14	*	78	*	100	7.0 (64)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 40

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1511	141	22.7 (75)	175	9.1 (76)	29	*	142	18.3 (87)	178	18.0 (105)
1512	69	*	111	4.5 (38)	12	*	72	*	112	4.5 (40)
1514	135	26.7 (85)	201	5.0 (44)	33	*	135	4.4 (11)	204	1.5 (7)
1701	116	14.7 (43)	144	18.8 (101)	17	*	120	14.2 (75)	148	7.4 (66)
1950	131	23.7 (77)	155	7.1 (66)	30	*	132	17.4 (82)	162	11.7 (94)
1952	30	*	40	*	4	*	30	*	42	*
1953	76	*	91	*	18	*	76	*	96	*
1954	98	*	96	*	34	*	98	*	107	16.8 (102)
1955	25	*	41	*	8	*	25	*	42	*
1957	85	*	105	8.6 (73)	14	*	85	*	106	8.5 (75)
1963	19	*	27	*	9	*	18	*	28	*
2150	135	28.9 (89)	169	17.2 (100)	37	*	135	16.3 (78)	173	10.4 (87)
2151	57	*	69	*	7	*	57	*	71	*
2152	8	*	6	*	2	*	8	*	6	*
2251	92	*	99	*	23	*	92	*	101	3.0 (26)
2252	49	*	53	*	14	*	51	*	54	*
2554	8	*	9	*	4	*	10	*	9	*
2555	1,701	17.8 (54)	2,025	6.7 (60)	268	44.4 (5)	1,705	11.8 (61)	2,071	6.7 (59)
2556	17	*	20	*	0	*	17	*	20	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 41

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2563	1	*	1	*	0	*	1	*	1	*
2564	1	*	1	*	0	*	1	*	1	*
3001	293	19.1 (63)	335	5.1 (45)	49	*	294	10.9 (54)	336	10.1 (84)
3002	159	14.5 (40)	188	5.3 (48)	21	*	159	18.2 (86)	188	9.0 (78)
3004	83	*	109	2.8 (26)	13	*	83	*	109	1.8 (9)
3005	42	*	49	*	7	*	43	*	49	*
3007	2	*	2	*	0	*	2	*	2	*
3201	110	23.6 (76)	142	8.5 (72)	25	*	110	17.3 (81)	141	3.5 (30)
3301	1,049	17.3 (50)	1,336	6.4 (56)	173	44.5 (4)	1,051	10.6 (50)	1,337	2.6 (20)
3304	20	*	24	*	2	*	20	*	24	*
3305	43	*	46	*	5	*	43	*	46	*
3306	52	*	60	*	8	*	52	*	60	*
3308	17	*	22	*	3	*	17	*	22	*
3312	122	11.5 (26)	138	4.3 (37)	13	*	121	9.9 (44)	139	1.4 (6)
3314	1	*	1	*	0	*	1	*	1	*
3550	8	*	8	*	3	*	8	*	8	*
3701	191	24.1 (78)	237	11.4 (84)	45	*	193	16.6 (79)	256	6.6 (57)
3850	653	19.4 (66)	788	10.7 (81)	115	32.2 (7)	653	13.6 (71)	784	8.0 (68)
3901	323	13.9 (36)	403	6.7 (60)	48	*	329	9.4 (42)	411	5.1 (46)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 42

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3950	499	18.0 (55)	615	6.7 (60)	80	*	501	10.6 (50)	631	6.8 (60)
4106	550	9.1 (19)	704	4.8 (43)	46	*	556	6.8 (24)	707	2.3 (17)
4107	638	18.2 (57)	795	4.5 (38)	121	57.0 (1)	646	6.8 (24)	807	4.7 (42)
4109	146	11.0 (25)	227	7.5 (70)	22	*	153	6.5 (22)	229	3.9 (36)
4112	176	8.5 (16)	185	1.6 (10)	10	*	175	12.0 (64)	186	3.2 (28)
4115	825	8.0 (14)	1,053	3.6 (33)	71	*	831	4.5 (12)	1,055	1.3 (5)
4351	140	32.1 (93)	150	16.7 (98)	42	*	138	19.6 (91)	149	15.4 (100)
4352	42	*	47	*	15	*	44	*	50	*
4353	126	19.0 (61)	150	13.3 (94)	21	*	126	7.9 (34)	150	6.0 (53)
4354	203	28.6 (87)	210	11.0 (82)	55	*	202	16.8 (80)	232	11.6 (93)
4355	73	*	82	*	26	*	73	*	89	*
4356	97	*	116	14.7 (97)	28	*	97	*	119	20.2 (106)
4359	133	10.5 (24)	156	4.5 (38)	13	*	133	2.3 (3)	153	3.3 (29)
4360	60	*	74	*	11	*	60	*	75	*
4361	58	*	69	*	21	*	58	*	72	*
4362	69	*	71	*	25	*	69	*	78	*
4363	162	19.1 (63)	201	9.5 (78)	28	*	162	19.1 (88)	205	12.7 (98)
4401	186	19.4 (66)	223	12.1 (88)	37	*	193	8.3 (36)	229	4.4 (38)
4501	200	31.5 (92)	231	12.6 (92)	63	*	201	20.9 (92)	232	12.1 (96)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 43

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4601	289	18.3 (58)	341	8.2 (71)	47	*	290	9.3 (39)	344	5.2 (47)
4701	228	25.4 (82)	295	6.4 (56)	55	*	230	13.0 (68)	306	7.5 (67)
5050	360	9.7 (21)	496	4.2 (36)	31	*	360	6.9 (26)	500	5.8 (51)
5051	537	8.6 (17)	751	2.4 (21)	37	*	538	6.9 (26)	750	5.2 (47)
5052	594	5.9 (8)	835	2.0 (16)	35	*	594	4.2 (8)	829	3.1 (27)
5055	339	1.8 (2)	429	1.2 (7)	5	*	339	2.4 (4)	429	0.0 (1)
5101	334	6.3 (10)	439	2.7 (24)	21	*	334	9.3 (39)	448	2.2 (14)
5202	144	42.4 (96)	174	12.1 (88)	59	*	144	11.1 (55)	183	8.7 (77)
5601	214	29.9 (90)	266	7.1 (66)	57	*	216	10.2 (47)	284	6.3 (55)
5801	357	27.2 (86)	453	8.8 (74)	103	47.6 (3)	366	12.3 (67)	456	8.3 (72)
5803	3	*	3	*	0	*	3	*	3	*
5804	1	*	1	*	0	*	1	*	1	*
5806	11	*	29	*	4	*	12	*	29	*
6101	14	*	18	*	2	*	14	*	18	*
6102	11	*	13	*	0	*	11	*	13	*
6103	790	14.6 (42)	942	7.2 (69)	107	33.6 (6)	792	13.6 (71)	960	8.0 (68)
6105	1	*	1	*	0	*	1	*	1	*
6301	800	7.0 (12)	991	2.1 (19)	55	*	804	6.0 (17)	999	2.9 (24)
6312	405	2.0 (3)	560	1.1 (6)	11	*	407	4.2 (8)	562	2.8 (21)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 44

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
6316	255	7.8 (13)	355	3.1 (28)	20	*	258	3.5 (6)	357	2.8 (21)
6318	1	*	9	*	0	*	1	*	8	*
6320	148	11.5 (26)	164	0.6 (4)	14	*	148	4.7 (13)	165	3.6 (32)
6325	72	*	96	*	19	*	74	*	97	*
6335	220	13.6 (35)	278	0.7 (5)	21	*	219	11.4 (58)	278	6.5 (56)
6345	45	*	49	*	15	*	45	*	52	*
6360	180	17.2 (49)	238	2.9 (27)	31	*	180	11.7 (60)	241	5.4 (49)
7306	402	18.9 (59)	507	5.3 (48)	70	*	402	11.2 (56)	511	6.1 (54)
7401	426	22.3 (74)	493	9.5 (78)	85	*	431	14.8 (77)	506	8.5 (75)
7601	109	11.9 (29)	133	6.8 (63)	14	*	110	11.8 (61)	138	8.0 (68)
7850	253	10.3 (23)	315	5.4 (50)	29	*	254	5.1 (14)	323	6.8 (60)
7852	2	*	1	*	0	*	2	*	1	*
8101	350	14.3 (38)	486	1.9 (15)	49	*	352	8.0 (35)	492	4.7 (42)
8102	80	*	90	*	5	*	80	*	91	*
8106	103	15.5 (45)	147	6.8 (63)	14	*	103	9.7 (43)	145	4.1 (37)
8150	1	*	1	*	0	*	1	*	1	*
8151	750	14.4 (39)	936	4.5 (38)	92	*	755	5.6 (15)	943	3.8 (35)
8155	2	*	2	*	0	*	2	*	2	*
8201	491	14.5 (40)	664	4.7 (42)	61	*	490	6.5 (22)	666	7.2 (65)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 45

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8202	1,038	2.1 (4)	1,254	0.5 (2)	21	*	1,039	3.9 (7)	1,260	1.2 (3)
8203	80	*	107	5.6 (51)	14	*	81	*	108	16.7 (101)
8250	97	*	124	4.0 (35)	10	*	97	*	124	8.1 (71)
8252	25	*	24	*	3	*	25	*	26	*
8255	362	17.4 (52)	545	5.9 (52)	54	*	362	10.2 (47)	553	9.6 (80)
8257	408	0.2 (1)	480	0.4 (1)	2	*	407	0.7 (1)	479	0.4 (2)
8263	100	10.0 (22)	77	*	11	*	101	9.9 (44)	77	*
8264	188	11.7 (28)	252	1.6 (10)	19	*	188	6.4 (20)	256	2.3 (17)
8267	170	13.5 (34)	254	5.1 (45)	20	*	170	8.8 (37)	261	6.9 (62)
8901	158	13.3 (33)	221	3.6 (33)	19	*	158	5.7 (16)	222	4.5 (40)
8902	261	13.0 (32)	344	2.0 (16)	31	*	262	7.3 (30)	348	2.0 (12)
8903	268	15.7 (47)	374	2.4 (21)	40	*	270	7.0 (29)	374	3.5 (30)
8904	395	18.0 (55)	538	2.2 (20)	61	*	397	9.3 (39)	540	3.7 (33)
8905	357	6.4 (11)	469	1.7 (12)	24	*	359	6.4 (20)	469	5.8 (51)
8909	231	9.5 (20)	316	3.2 (29)	23	*	233	9.0 (38)	319	4.7 (42)
9045	327	2.8 (5)	400	0.5 (2)	9	*	326	4.3 (10)	402	1.2 (3)
9050	191	17.3 (50)	239	3.3 (31)	30	*	193	7.3 (30)	241	1.7 (8)
9051	484	3.5 (6)	591	2.0 (16)	21	*	488	1.2 (2)	592	1.9 (11)
9053	316	17.4 (52)	567	1.8 (13)	61	*	324	10.8 (52)	567	1.8 (9)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2023 Pregnancy Nutrition Surveillance (1)

Page : 46

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
9055	2	*	2	*	0	*	2	*	2	*
9058	265	6.0 (9)	434	1.8 (13)	16	*	267	10.5 (49)	430	2.8 (21)
9066	145	19.3 (65)	134	5.2 (47)	27	*	145	13.1 (69)	135	2.2 (14)
9068	199	8.0 (14)	276	1.4 (8)	14	*	199	6.0 (17)	276	2.9 (24)
9069	356	4.8 (7)	502	3.2 (29)	16	*	357	6.2 (19)	505	6.9 (62)
9070	425	12.5 (30)	616	1.5 (9)	50	*	426	13.6 (71)	619	3.7 (33)
9080	662	14.2 (37)	876	2.7 (24)	86	*	667	12.0 (64)	877	4.4 (38)
9082	17	*	37	*	2	*	18	*	37	*
9083	7	*	6	*	0	*	7	*	6	*
Michigan	33,089	15.8	41,903	5.7	4,925	45.2	33,292	10.2	42,493	6.0
National (Prior Year)	40,927	23.4	49,975	7.8	7,918	31.5	39,948	6.0	53,241	3.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 47

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	46	*	*	47	*	43	*	46	*
0102	98	*	*	100	9.0 (8)	87	*	97	*
0103	123	5.7 (5)	17.1 (102)	124	6.5 (2)	114	4.4 (40)	122	79.5 (52)
0104	39	*	*	39	*	31	*	39	*
0105	1	*	*	1	*	1	*	1	*
0250	33	*	*	30	*	29	*	33	*
0251	23	*	*	24	*	18	*	23	*
0252	32	*	*	32	*	32	*	32	*
0253	43	*	*	45	*	36	*	43	*
0401	119	10.9 (55)	5.9 (47)	117	15.4 (83)	97	*	119	78.2 (57)
0402	82	*	*	83	*	78	*	82	*
0403	30	*	*	29	*	27	*	30	*
0404	27	*	*	28	*	21	*	27	*
0501	112	6.3 (8)	4.5 (24)	121	10.7 (30)	101	1.0 (3)	117	89.7 (3)
0503	15	*	*	16	*	15	*	15	*
0504	83	*	*	86	*	74	*	82	*
0505	45	*	*	53	*	40	*	51	*
0506	45	*	*	48	*	40	*	45	*
0507	18	*	*	19	*	17	*	19	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 48

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	4	*	*	4	*	4	*	4	*
0650	295	9.2 (31)	8.1 (83)	303	9.6 (12)	265	4.2 (35)	299	83.9 (23)
0651	128	15.6 (91)	7.8 (76)	143	14.0 (64)	108	9.3 (89)	128	73.4 (69)
0652	77	*	*	77	*	71	*	77	*
0653	110	5.5 (4)	7.3 (70)	113	11.5 (36)	97	*	110	85.5 (13)
0654	129	9.3 (32)	3.9 (18)	128	10.2 (24)	115	5.2 (54)	128	82.8 (27)
0655	93	*	*	95	*	84	*	93	*
0657	6	*	*	7	*	5	*	6	*
0701	114	4.4 (2)	9.6 (96)	117	12.0 (41)	101	2.0 (7)	116	89.7 (3)
0702	27	*	*	28	*	21	*	27	*
0703	54	*	*	54	*	46	*	54	*
0704	16	*	*	16	*	15	*	16	*
0705	20	*	*	21	*	20	*	20	*
0706	1	*	*	1	*	1	*	1	*
0850	193	12.4 (71)	7.8 (76)	200	15.5 (86)	163	3.7 (26)	198	79.8 (51)
0851	178	9.6 (36)	6.2 (52)	196	15.3 (82)	152	2.0 (7)	183	81.4 (39)
0901	385	9.4 (35)	9.1 (91)	398	10.6 (28)	344	3.8 (27)	384	70.6 (77)
0902	3	*	*	3	*	3	*	3	*
1051	304	7.6 (16)	8.6 (88)	309	9.1 (11)	274	4.0 (31)	308	91.2 (2)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 49

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1101	373	9.7 (37)	6.2 (52)	387	13.4 (57)	322	5.3 (57)	377	68.4 (83)
1102	179	8.9 (28)	10.1 (99)	184	10.3 (26)	159	4.4 (40)	178	75.3 (65)
1106	19	*	*	19	*	16	*	19	*
1250	293	9.9 (41)	9.2 (92)	311	9.0 (8)	263	5.7 (65)	299	81.9 (33)
1251	206	7.8 (18)	7.3 (70)	206	14.6 (72)	173	4.6 (44)	206	81.1 (41)
1252	141	9.9 (41)	8.5 (87)	145	15.9 (91)	119	4.2 (35)	143	81.8 (34)
1253	178	6.2 (7)	9.6 (96)	180	10.0 (19)	155	0.0 (1)	182	80.8 (44)
1350	622	10.9 (55)	6.4 (57)	642	10.0 (19)	557	6.1 (70)	624	84.1 (20)
1352	84	*	*	84	*	67	*	84	*
1401	66	*	*	67	*	61	*	66	*
1402	20	*	*	21	*	19	*	20	*
1403	1	*	*	1	*	1	*	1	*
1501	215	9.3 (32)	5.1 (32)	224	12.5 (51)	187	3.2 (20)	218	84.9 (14)
1503	220	8.2 (21)	4.5 (24)	221	10.0 (19)	196	3.6 (24)	220	79.5 (52)
1504	501	11.4 (60)	6.0 (49)	527	11.0 (34)	444	3.8 (27)	523	83.7 (24)
1506	145	10.3 (47)	8.3 (84)	152	6.6 (3)	133	6.0 (69)	148	79.1 (55)
1507	140	8.6 (25)	7.9 (80)	150	14.7 (76)	118	2.5 (13)	146	88.4 (5)
1509	94	*	*	93	*	82	*	94	*
1511	165	9.1 (30)	7.9 (80)	170	12.4 (50)	144	5.6 (62)	166	76.5 (62)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 50

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1512	108	10.2 (45)	6.5 (61)	110	14.5 (70)	92	*	109	86.2 (9)
1514	193	16.6 (96)	9.3 (94)	196	16.8 (96)	160	4.4 (40)	193	84.5 (17)
1701	138	5.8 (6)	13.0 (101)	143	9.8 (16)	124	2.4 (12)	139	83.5 (25)
1950	152	6.6 (10)	3.9 (18)	156	9.6 (12)	138	2.9 (17)	152	84.9 (14)
1952	41	*	*	39	*	38	*	41	*
1953	93	*	*	95	*	79	*	93	*
1954	100	8.0 (20)	8.0 (82)	102	8.8 (7)	90	*	100	84.0 (22)
1955	38	*	*	40	*	32	*	39	*
1957	99	*	*	103	15.5 (86)	84	*	99	*
1963	25	*	*	26	*	23	*	26	*
2150	158	8.2 (21)	6.3 (55)	166	9.0 (8)	143	2.1 (10)	158	82.3 (32)
2151	68	*	*	69	*	56	*	67	*
2152	6	*	*	6	*	4	*	6	*
2251	98	*	*	96	*	83	*	97	*
2252	50	*	*	53	*	40	*	49	*
2554	9	*	*	9	*	8	*	9	*
2555	2,004	14.0 (85)	5.5 (37)	2,033	14.3 (67)	1,706	7.2 (81)	2,011	66.9 (85)
2556	19	*	*	19	*	17	*	19	*
2563	1	*	*	1	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 51

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2564	1	*	*	1	*	1	*	1	*
3001	311	13.2 (78)	5.1 (32)	320	15.6 (88)	262	4.6 (44)	313	65.5 (87)
3002	176	11.4 (60)	4.0 (20)	180	15.0 (79)	150	2.0 (7)	176	73.3 (71)
3004	99	*	*	102	10.8 (33)	88	*	100	88.0 (6)
3005	46	*	*	46	*	40	*	46	*
3007	2	*	*	2	*	2	*	2	*
3201	131	10.7 (52)	8.4 (86)	134	13.4 (57)	113	0.9 (2)	131	79.4 (54)
3301	1,250	9.8 (39)	6.6 (64)	1,288	12.2 (47)	1,094	2.8 (16)	1,266	83.0 (26)
3304	22	*	*	24	*	20	*	22	*
3305	44	*	*	44	*	42	*	44	*
3306	53	*	*	53	*	42	*	53	*
3308	20	*	*	20	*	18	*	20	*
3312	125	12.8 (75)	3.2 (13)	130	15.4 (83)	106	7.5 (83)	126	85.7 (11)
3314	1	*	*	1	*	1	*	1	*
3550	8	*	*	8	*	7	*	8	*
3701	245	10.2 (45)	6.5 (61)	251	10.0 (19)	221	5.4 (61)	245	86.5 (8)
3850	720	11.5 (63)	7.8 (76)	762	12.1 (44)	630	6.2 (71)	726	80.0 (49)
3901	381	13.4 (79)	5.5 (37)	397	15.1 (80)	321	5.6 (62)	386	73.6 (68)
3950	599	10.4 (50)	7.2 (68)	615	12.0 (41)	526	5.3 (57)	604	80.8 (44)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 52

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4106	671	13.1 (77)	4.3 (23)	685	16.6 (95)	555	6.7 (77)	682	79.0 (56)
4107	773	11.4 (60)	5.0 (29)	801	13.4 (57)	670	5.2 (54)	786	81.6 (36)
4109	218	12.4 (71)	7.8 (76)	222	17.6 (99)	178	3.9 (29)	219	85.8 (10)
4112	180	2.2 (1)	5.6 (40)	180	3.9 (1)	172	1.2 (5)	180	45.6 (101)
4115	985	13.6 (80)	5.8 (44)	1,027	14.0 (64)	849	5.7 (65)	1,011	82.4 (30)
4351	150	7.3 (14)	7.3 (70)	150	11.3 (35)	131	3.1 (18)	150	84.7 (16)
4352	46	*	*	47	*	39	*	46	*
4353	144	11.8 (66)	5.6 (40)	144	14.6 (72)	123	6.5 (73)	145	84.1 (20)
4354	217	9.7 (37)	8.3 (84)	217	12.0 (41)	187	5.3 (57)	218	82.6 (29)
4355	83	*	*	84	*	71	*	83	*
4356	108	6.5 (9)	9.3 (94)	110	8.2 (5)	96	*	111	85.6 (12)
4359	143	9.8 (39)	7.7 (75)	146	12.3 (48)	124	4.0 (31)	143	80.4 (47)
4360	71	*	*	72	*	64	*	72	*
4361	69	*	*	69	*	61	*	69	*
4362	72	*	*	78	*	63	*	72	*
4363	195	10.3 (47)	9.2 (92)	195	13.3 (55)	165	4.2 (35)	197	91.9 (1)
4401	212	9.0 (29)	9.0 (90)	221	11.8 (38)	185	2.7 (15)	212	81.6 (36)
4501	219	5.0 (3)	6.4 (57)	224	8.0 (4)	201	1.5 (6)	220	82.7 (28)
4601	323	10.8 (54)	6.2 (52)	334	13.2 (54)	282	5.0 (52)	323	81.7 (35)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 53

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4701	291	10.7 (52)	6.9 (66)	291	13.7 (61)	248	3.2 (20)	291	84.2 (19)
5050	484	12.6 (74)	5.2 (35)	481	14.3 (67)	402	6.5 (73)	484	75.4 (64)
5051	709	11.6 (64)	6.1 (50)	719	16.1 (93)	583	4.3 (39)	715	76.1 (63)
5052	797	15.8 (92)	4.6 (27)	805	14.5 (70)	674	9.3 (89)	800	72.6 (74)
5055	391	12.0 (67)	4.9 (28)	408	12.3 (48)	343	5.2 (54)	403	71.0 (75)
5101	440	8.6 (25)	6.1 (50)	433	11.8 (38)	377	3.2 (20)	441	78.2 (57)
5202	176	7.4 (15)	6.3 (55)	176	9.7 (15)	157	5.7 (65)	177	80.8 (44)
5601	263	6.8 (11)	10.3 (100)	271	9.6 (12)	237	2.5 (13)	263	81.0 (43)
5801	452	8.6 (25)	5.3 (36)	461	10.0 (19)	407	3.9 (29)	451	73.4 (69)
5803	3	*	*	3	*	3	*	3	*
5804	1	*	*	1	*	1	*	1	*
5806	28	*	*	28	*	20	*	28	*
6101	18	*	*	18	*	17	*	18	*
6102	13	*	*	13	*	13	*	13	*
6103	901	11.3 (59)	5.5 (37)	932	15.1 (80)	760	5.8 (68)	906	74.5 (66)
6105	1	*	*	1	*	1	*	1	*
6301	940	10.6 (51)	5.1 (32)	973	14.3 (67)	802	4.1 (33)	959	74.3 (67)
6312	519	13.7 (82)	4.0 (20)	540	14.8 (77)	436	4.6 (44)	529	82.4 (30)
6316	331	12.1 (69)	7.3 (70)	341	13.8 (62)	283	4.6 (44)	334	81.1 (41)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 54

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6318	8	*	*	8	*	5	*	8	*
6320	154	12.3 (70)	6.5 (61)	163	12.9 (53)	136	6.6 (75)	154	84.4 (18)
6325	93	*	*	95	*	87	*	94	*
6335	266	10.9 (55)	6.4 (57)	273	12.1 (44)	233	4.7 (48)	268	78.0 (59)
6345	52	*	*	51	*	45	*	52	*
6360	222	16.2 (94)	4.5 (24)	232	12.1 (44)	192	9.4 (92)	228	81.6 (36)
7306	484	9.9 (41)	5.8 (44)	494	12.8 (52)	422	3.1 (18)	488	73.0 (73)
7401	478	10.3 (47)	8.6 (88)	491	9.8 (16)	433	4.4 (40)	481	77.1 (60)
7601	129	9.3 (32)	7.0 (67)	135	11.9 (40)	114	3.5 (23)	131	64.1 (88)
7850	306	6.9 (12)	5.6 (40)	313	10.5 (27)	273	2.2 (11)	308	79.9 (50)
7852	1	*	*	1	*	1	*	1	*
8101	433	14.3 (87)	2.1 (1)	473	17.8 (101)	347	6.3 (72)	452	59.3 (91)
8102	85	*	*	84	*	76	*	86	*
8106	103	15.5 (90)	9.7 (98)	143	15.4 (83)	85	*	141	68.8 (82)
8150	1	*	*	1	*	1	*	1	*
8151	875	12.5 (73)	5.9 (47)	905	15.6 (88)	734	5.6 (62)	890	86.9 (7)
8155	2	*	*	2	*	2	*	2	*
8201	637	13.7 (82)	3.6 (17)	656	14.6 (72)	542	6.8 (78)	646	69.8 (79)
8202	1,102	8.5 (24)	3.5 (16)	1,193	9.9 (18)	987	4.2 (35)	1,179	70.6 (77)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 55

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8203	108	11.1 (58)	5.6 (40)	108	10.2 (24)	97	*	108	76.9 (61)
8250	117	7.7 (17)	2.6 (7)	123	10.6 (28)	104	1.0 (3)	119	69.7 (80)
8252	18	*	*	26	*	16	*	25	*
8255	533	12.0 (67)	5.8 (44)	547	14.1 (66)	453	4.9 (51)	537	70.9 (76)
8257	429	12.8 (75)	2.6 (7)	466	10.7 (30)	379	10.8 (95)	444	56.8 (96)
8263	70	*	*	76	*	65	*	72	*
8264	249	11.6 (64)	7.2 (68)	252	13.9 (63)	215	4.7 (48)	250	80.4 (47)
8267	239	7.9 (19)	6.7 (65)	253	8.3 (6)	219	4.1 (33)	245	69.4 (81)
8901	201	15.9 (93)	5.0 (29)	216	19.4 (104)	160	6.9 (80)	205	58.5 (93)
8902	320	17.8 (98)	5.0 (29)	331	16.0 (92)	264	9.5 (93)	325	66.2 (86)
8903	354	14.1 (86)	3.4 (15)	363	16.3 (94)	292	7.9 (85)	353	49.0 (100)
8904	497	18.3 (99)	2.4 (4)	523	20.1 (105)	384	8.9 (87)	503	59.0 (92)
8905	405	10.1 (44)	4.0 (20)	452	13.3 (55)	350	5.1 (53)	438	73.1 (72)
8909	282	19.5 (102)	2.5 (5)	306	18.6 (102)	224	10.3 (94)	292	67.1 (84)
9045	375	6.9 (12)	7.5 (74)	392	11.7 (37)	332	3.6 (24)	383	43.6 (102)
9050	228	13.6 (80)	3.1 (11)	235	14.9 (78)	192	6.8 (78)	231	59.7 (90)
9051	534	8.4 (23)	6.4 (57)	559	10.7 (30)	473	5.3 (57)	562	81.3 (40)
9053	529	13.8 (84)	2.5 (5)	556	14.6 (72)	450	8.4 (86)	533	56.3 (97)
9055	2	*	*	2	*	2	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2023 Pregnancy Nutrition Surveillance (1)

Page : 56

Michigan

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9058	405	16.5 (95)	3.0 (10)	421	17.3 (98)	335	7.5 (83)	410	57.3 (95)
9066	130	19.2 (101)	3.1 (11)	131	17.6 (99)	106	4.7 (48)	129	57.4 (94)
9068	263	17.1 (97)	2.3 (3)	267	13.5 (60)	225	9.3 (89)	263	50.2 (99)
9069	474	14.6 (88)	3.2 (13)	489	16.8 (96)	393	7.4 (82)	477	43.0 (103)
9070	585	18.5 (100)	2.2 (2)	603	19.1 (103)	470	8.9 (87)	587	54.0 (98)
9080	829	15.4 (89)	2.7 (9)	866	15.8 (90)	696	6.6 (75)	846	63.9 (89)
9082	35	*	*	36	*	30	*	35	*
9083	6	*	*	6	*	6	*	6	*
Michigan	39,736	11.4	5.8	41,121	13.2	34,238	5.2	40,322	74.6
National (Prior Year)	50,081	11.0	6.0	51,393	13.2	42,775	5.0	50,583	46.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* National includes previous year's data for all participating States and ITOs.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 57

Michigan

Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0101	123	91.9	0.0	4.9	0.0	0.0	2.4	0.8
0102	377	90.5	0.3	3.2	0.3	0.0	5.6	0.3
0103	371	92.5	0.8	2.7	0.0	0.0	3.8	0.3
0104	127	93.7	0.8	2.4	0.0	0.0	3.1	0.0
0105	4	*	*	*	*	*	*	*
0250	117	78.6	0.0	2.6	6.0	0.0	12.8	0.0
0251	75	*	*	*	*	*	*	*
0252	103	61.2	1.9	1.9	1.9	1.0	32.0	0.0
0253	120	70.0	0.0	2.5	10.8	0.8	15.8	0.0
0401	419	94.3	0.5	2.4	0.2	0.2	2.4	0.0
0402	347	84.4	2.6	2.6	0.0	0.0	10.4	0.0
0403	121	93.4	0.0	3.3	0.8	0.8	1.7	0.0
0404	107	96.3	0.0	0.9	0.0	0.0	2.8	0.0
0501	399	89.5	1.0	3.8	0.3	0.5	4.8	0.3
0503	58	*	*	*	*	*	*	*
0504	319	84.3	0.9	4.4	1.3	1.3	7.8	0.0
0505	196	89.3	0.0	4.1	0.0	0.0	6.6	0.0
0506	173	94.2	0.6	1.2	0.6	0.0	3.5	0.0
0507	71	*	*	*	*	*	*	*
0508	13	*	*	*	*	*	*	*
0650	1,076	77.9	4.3	7.5	3.2	0.5	6.6	0.1
0651	539	91.1	0.9	5.0	0.2	0.6	2.2	0.0
0652	287	94.1	0.7	3.1	0.3	0.0	1.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 58

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0653	378	93.9	0.3	2.4	0.0	0.5	2.6	0.3
0654	417	92.6	0.5	2.9	0.5	0.2	3.1	0.2
0655	315	95.6	0.0	1.3	0.0	0.0	2.9	0.3
0657	35	*	*	*	*	*	*	*
0701	468	87.4	1.7	1.7	1.1	2.8	5.1	0.2
0702	72	*	*	*	*	*	*	*
0703	165	84.8	3.6	1.8	1.2	0.6	7.3	0.6
0704	51	*	*	*	*	*	*	*
0705	66	*	*	*	*	*	*	*
0706	1	*	*	*	*	*	*	*
0850	686	83.2	5.1	8.2	0.0	1.9	1.6	0.0
0851	600	91.0	0.5	5.3	0.5	0.7	1.8	0.2
0901	1,419	80.0	2.5	10.1	0.1	0.4	6.6	0.3
0902	25	*	*	*	*	*	*	*
1051	1,117	81.5	1.3	8.5	2.5	1.1	4.8	0.4
1052	3	*	*	*	*	*	*	*
1101	1,397	35.7	47.7	9.4	0.5	2.3	3.3	1.1
1102	739	72.4	10.4	8.3	0.3	0.9	6.5	1.2
1106	58	*	*	*	*	*	*	*
1250	969	79.9	4.6	12.8	0.2	0.4	1.7	0.4
1251	643	91.9	0.3	3.7	0.3	0.2	3.0	0.6
1252	522	60.3	0.2	35.6	0.0	0.0	3.3	0.6
1253	645	78.1	0.6	11.8	0.0	0.3	8.5	0.6
1350	2,368	59.5	19.6	8.7	0.3	5.0	6.5	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 59

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1351	1	*	*	*	*	*	*	*
1352	354	67.2	20.6	5.6	0.3	0.0	5.1	1.1
1401	208	87.0	0.0	8.2	0.5	0.0	4.3	0.0
1402	88	*	*	*	*	*	*	*
1403	18	*	*	*	*	*	*	*
1501	746	47.3	5.8	40.5	0.1	0.3	3.8	2.3
1503	804	58.8	7.8	22.0	0.2	0.5	7.6	3.0
1504	1,848	37.5	5.5	47.8	0.1	4.1	4.5	0.6
1505	22	*	*	*	*	*	*	*
1506	533	76.7	1.1	16.3	0.0	0.8	4.1	0.9
1507	525	84.0	2.5	8.6	0.2	1.7	2.5	0.6
1509	371	48.5	1.6	44.5	0.0	1.1	2.4	1.9
1511	614	85.3	2.3	7.5	0.2	0.5	3.9	0.3
1512	373	84.7	1.3	7.0	0.3	1.1	5.6	0.0
1514	682	74.5	5.1	13.9	0.1	1.5	4.4	0.4
1701	513	59.3	0.8	2.7	18.3	1.2	17.7	0.0
1950	568	92.4	0.4	6.0	0.0	0.0	0.9	0.4
1952	132	90.9	0.8	6.1	0.0	0.0	2.3	0.0
1953	271	88.6	0.0	9.2	0.0	0.0	2.2	0.0
1954	356	83.7	0.3	11.8	0.0	0.0	3.7	0.6
1955	124	83.1	0.0	12.9	0.0	0.0	3.2	0.8
1957	382	83.8	1.6	10.7	0.0	0.3	3.4	0.3
1963	71	*	*	*	*	*	*	*
2150	553	83.4	0.7	3.8	2.0	0.5	9.4	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 60

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2151	227	81.1	1.8	7.9	0.0	0.0	6.6	2.6
2152	38	*	*	*	*	*	*	*
2251	376	93.9	0.0	2.1	0.0	0.5	2.7	0.8
2252	169	89.9	0.0	4.7	0.0	0.0	3.6	1.8
2554	49	*	*	*	*	*	*	*
2555	7,094	49.9	36.7	6.6	0.0	0.3	6.2	0.3
2556	117	27.4	59.8	2.6	0.0	0.0	10.3	0.0
2557	1	*	*	*	*	*	*	*
2560	1	*	*	*	*	*	*	*
2562	3	*	*	*	*	*	*	*
2563	6	*	*	*	*	*	*	*
2564	3	*	*	*	*	*	*	*
3001	1,159	27.9	43.2	14.8	0.3	0.5	13.1	0.3
3002	304	33.6	38.5	20.1	0.0	0.0	7.9	0.0
3004	301	77.1	0.3	18.3	0.0	0.3	3.7	0.3
3005	160	86.3	0.6	8.1	1.3	0.6	3.1	0.0
3007	7	*	*	*	*	*	*	*
3201	407	91.9	0.5	5.4	0.0	0.2	0.7	1.2
3301	4,300	38.5	30.4	16.2	0.2	6.0	8.5	0.3
3304	100	40.0	23.0	14.0	0.0	19.0	4.0	0.0
3305	158	60.8	12.7	10.1	0.0	5.7	10.8	0.0
3306	206	85.9	2.9	8.3	0.5	1.0	1.0	0.5
3308	94	*	*	*	*	*	*	*
3312	466	42.7	25.8	20.2	0.4	2.4	8.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 61

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3314	1	*	*	*	*	*	*	*
3550	64	*	*	*	*	*	*	*
3551	6	*	*	*	*	*	*	*
3552	1	*	*	*	*	*	*	*
3701	837	88.5	0.2	7.9	0.0	0.1	3.2	0.0
3850	2,696	71.7	13.6	7.2	0.1	0.7	6.6	0.1
3901	1,467	37.4	49.2	6.1	0.1	1.0	5.7	0.5
3902	9	*	*	*	*	*	*	*
3903	2	*	*	*	*	*	*	*
3950	2,001	52.0	23.7	13.5	0.1	0.7	9.0	0.8
3954	1	*	*	*	*	*	*	*
4106	2,174	12.1	35.7	46.7	0.1	1.0	4.0	0.3
4107	2,780	41.4	29.0	20.6	0.4	2.0	6.4	0.2
4109	639	85.9	1.3	8.6	0.2	0.9	2.7	0.5
4112	585	0.9	1.5	96.9	0.0	0.0	0.2	0.5
4115	3,376	29.8	31.7	22.3	0.3	10.7	5.0	0.2
4351	524	86.8	0.6	6.3	0.2	1.0	5.2	0.0
4352	185	73.0	9.2	7.0	0.0	0.0	10.3	0.5
4353	474	56.5	0.4	38.8	0.2	0.0	3.2	0.8
4354	754	87.5	2.8	4.2	0.4	0.3	4.2	0.5
4355	264	85.2	0.4	6.4	1.1	0.0	6.8	0.0
4356	381	77.7	1.3	12.9	0.3	0.5	6.6	0.8
4359	580	60.5	0.7	35.9	0.5	0.2	1.4	0.9
4360	229	87.8	0.4	7.4	0.4	0.0	3.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 62

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4361	244	87.7	1.2	6.6	0.0	0.0	4.5	0.0
4362	267	92.5	0.4	3.7	0.0	0.0	3.0	0.0
4363	641	90.5	1.1	3.7	0.0	0.3	4.2	0.2
4401	730	88.1	1.0	6.3	0.0	0.5	4.0	0.1
4501	750	93.9	0.4	4.1	0.0	0.1	1.3	0.1
4601	1,188	72.1	2.8	19.0	0.2	0.5	4.9	0.6
4701	999	88.4	1.5	4.7	0.1	1.4	2.9	1.0
5050	1,619	43.7	45.3	3.3	0.0	2.0	5.4	0.2
5051	2,574	48.4	38.5	4.2	0.1	3.7	4.6	0.5
5052	2,671	29.4	40.5	3.2	0.2	21.9	4.6	0.2
5055	1,349	76.4	9.9	1.3	0.2	11.3	0.7	0.1
5101	1,456	63.0	14.0	15.9	0.1	1.7	5.2	0.1
5102	5	*	*	*	*	*	*	*
5202	644	87.3	1.4	2.8	2.8	0.0	5.0	0.8
5206	11	*	*	*	*	*	*	*
5601	962	86.2	2.4	6.9	0.0	1.5	3.0	0.1
5801	1,769	80.8	6.4	7.6	0.0	0.1	4.2	0.8
5803	21	*	*	*	*	*	*	*
5804	10	*	*	*	*	*	*	*
5806	57	*	*	*	*	*	*	*
6101	65	*	*	*	*	*	*	*
6102	68	*	*	*	*	*	*	*
6103	3,246	56.0	25.9	10.4	0.4	0.6	6.4	0.3
6105	6	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 63

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
6301	3,378	38.3	30.7	23.8	0.2	2.7	4.0	0.3
6312	1,773	33.6	55.8	5.2	0.1	1.7	3.3	0.2
6316	1,127	59.4	24.5	8.7	0.0	3.1	3.7	0.5
6318	10	*	*	*	*	*	*	*
6320	562	36.8	51.4	3.7	0.2	2.7	5.2	0.0
6325	322	89.4	1.2	5.6	0.0	0.0	3.7	0.0
6335	928	62.3	23.4	4.1	0.0	7.8	2.0	0.4
6345	145	83.4	3.4	4.1	0.0	3.4	5.5	0.0
6360	903	32.9	43.0	17.7	0.0	2.3	3.8	0.3
7306	1,874	46.6	30.9	16.4	0.2	0.4	5.3	0.2
7401	1,777	80.5	5.0	6.9	0.1	0.5	6.6	0.3
7601	505	93.1	0.0	5.0	0.0	0.2	1.2	0.6
7850	1,040	90.7	0.4	4.5	0.0	0.1	3.8	0.5
7852	8	*	*	*	*	*	*	*
8101	1,705	7.6	88.4	1.6	0.0	0.2	1.9	0.2
8102	444	23.4	19.1	53.8	0.0	1.8	1.6	0.2
8105	1	*	*	*	*	*	*	*
8106	550	26.9	51.1	14.2	0.0	0.5	7.1	0.2
8150	7	*	*	*	*	*	*	*
8151	3,223	37.4	37.2	13.4	0.1	3.6	7.8	0.5
8152	7	*	*	*	*	*	*	*
8153	1	*	*	*	*	*	*	*
8155	7	*	*	*	*	*	*	*
8201	2,587	42.9	41.8	5.4	0.2	3.9	5.3	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 64

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8202	3,922	88.9	8.0	1.6	0.0	0.4	0.9	0.2
8203	358	54.5	32.4	7.3	0.0	0.0	5.9	0.0
8250	464	20.7	69.0	3.4	0.2	1.1	5.6	0.0
8252	133	22.6	63.2	6.8	0.0	4.5	3.0	0.0
8255	1,897	53.5	30.4	9.4	0.4	2.6	3.5	0.2
8257	1,627	53.3	9.0	1.0	0.1	35.0	1.5	0.2
8263	389	60.9	17.7	15.7	0.0	1.0	4.6	0.0
8264	795	55.0	31.7	5.8	0.3	2.8	4.4	0.1
8267	707	54.7	18.5	22.2	0.1	0.8	3.3	0.3
8901	794	9.3	82.6	3.1	0.0	2.3	2.5	0.1
8902	1,252	8.7	86.0	2.4	0.0	0.6	2.2	0.1
8903	1,347	6.3	86.0	1.6	0.1	2.1	3.9	0.0
8904	1,935	4.3	90.1	1.9	0.1	1.0	2.4	0.2
8905	1,898	64.3	23.4	7.6	0.2	2.0	2.4	0.1
8909	1,168	5.8	89.1	2.4	0.1	0.3	2.1	0.2
9045	1,310	10.1	21.8	65.5	0.1	0.5	1.4	0.8
9050	882	18.6	59.9	18.6	0.0	0.1	2.8	0.0
9051	1,649	7.0	4.8	87.0	0.0	0.2	0.8	0.3
9053	1,597	7.9	83.5	2.6	0.1	3.0	2.6	0.3
9055	7	*	*	*	*	*	*	*
9058	1,409	15.8	73.4	1.2	0.1	7.2	2.2	0.1
9066	689	2.6	93.5	0.9	0.0	0.7	2.3	0.0
9068	818	2.8	94.1	0.6	0.0	0.5	1.6	0.4
9069	1,657	1.1	96.6	1.4	0.0	0.1	0.7	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 3A**2021 - 2023 Pregnancy Nutrition Surveillance (1)**
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

Page : 65

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9070	1,892	6.4	89.0	1.6	0.1	1.7	1.1	0.2
9072	59	*	*	*	*	*	*	*
9080	2,628	19.1	74.1	2.5	0.2	0.3	3.4	0.3
9082	108	0.9	95.4	1.9	0.9	0.9	0.0	0.0
9083	13	*	*	*	*	*	*	*
9087	11	*	*	*	*	*	*	*
Michigan	142,422	51.2	29.1	11.9	0.3	2.6	4.5	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 66

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	123	0.0	1.6	8.9	56.1	31.7	1.6	0.0
0102	377	0.0	1.3	8.5	59.2	28.1	2.7	0.3
0103	371	0.3	2.2	7.5	61.7	26.4	1.1	0.8
0104	127	0.0	3.9	7.1	60.6	27.6	0.0	0.8
0105	4	*	*	*	*	*	*	*
0250	117	0.0	1.7	8.5	53.8	31.6	2.6	1.7
0251	75	*	*	*	*	*	*	*
0252	103	0.0	1.9	2.9	54.4	38.8	1.9	0.0
0253	120	0.0	2.5	5.0	66.7	23.3	2.5	0.0
0401	419	0.0	1.7	6.2	59.2	32.0	1.0	0.0
0402	347	0.3	2.0	6.9	63.4	26.5	0.6	0.3
0403	121	0.0	2.5	9.9	56.2	28.9	2.5	0.0
0404	107	0.0	1.9	3.7	60.7	29.9	0.9	2.8
0501	399	0.0	1.3	10.3	59.9	26.3	1.8	0.5
0503	58	*	*	*	*	*	*	*
0504	319	0.0	1.3	7.2	56.4	32.6	2.5	0.0
0505	196	0.0	1.0	7.7	62.2	26.0	1.5	1.5
0506	173	0.0	2.3	5.8	69.4	22.0	0.6	0.0
0507	71	*	*	*	*	*	*	*
0508	13	*	*	*	*	*	*	*
0650	1,076	0.1	1.6	6.9	63.6	25.1	2.0	0.7
0651	539	0.4	1.7	9.3	62.3	24.5	0.9	0.9
0652	287	0.0	1.7	9.8	61.3	25.4	1.0	0.7
0653	378	0.0	1.6	4.5	63.8	27.8	0.5	1.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 67

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0654	417	0.0	2.6	6.7	65.7	21.1	1.7	2.2
0655	315	0.0	3.8	5.4	58.4	29.8	1.3	1.3
0657	35	*	*	*	*	*	*	*
0701	468	0.0	1.1	8.3	60.3	28.0	2.4	0.0
0702	72	*	*	*	*	*	*	*
0703	165	0.0	0.0	5.5	57.0	37.6	0.0	0.0
0704	51	*	*	*	*	*	*	*
0705	66	*	*	*	*	*	*	*
0706	1	*	*	*	*	*	*	*
0850	686	0.0	2.0	7.7	56.3	31.0	1.7	1.2
0851	600	0.0	2.2	6.7	64.5	24.0	0.8	1.8
0901	1,419	0.0	2.6	8.1	59.5	28.0	1.1	0.7
0902	25	*	*	*	*	*	*	*
1051	1,117	0.1	1.1	5.6	59.5	31.4	1.8	0.4
1052	3	*	*	*	*	*	*	*
1101	1,397	0.2	2.5	9.9	57.8	25.3	2.4	1.9
1102	739	0.1	2.6	7.4	62.9	25.2	0.8	0.9
1106	58	*	*	*	*	*	*	*
1250	969	0.0	1.7	6.9	62.0	24.8	3.8	0.8
1251	643	0.8	2.5	6.4	64.2	21.8	2.8	1.6
1252	522	0.0	2.9	7.9	63.4	21.5	3.3	1.1
1253	645	0.0	2.8	6.2	62.0	26.0	1.2	1.7
1350	2,368	0.0	2.3	7.6	61.1	25.7	1.9	1.5
1351	1	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 68

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1352	354	0.0	0.8	7.1	63.3	26.0	0.8	2.0
1401	208	1.0	1.0	5.3	56.3	33.2	1.9	1.4
1402	88	*	*	*	*	*	*	*
1403	18	*	*	*	*	*	*	*
1501	746	0.1	2.5	6.0	60.2	26.8	2.5	1.7
1503	804	0.1	2.2	7.7	60.8	25.2	1.7	2.1
1504	1,848	0.1	2.2	6.5	58.5	30.6	1.7	0.4
1505	22	*	*	*	*	*	*	*
1506	533	0.0	0.8	6.0	64.5	24.8	1.7	2.3
1507	525	0.0	0.8	5.7	56.2	34.7	1.9	0.8
1509	371	0.0	1.3	7.3	53.9	33.7	3.2	0.5
1511	614	0.0	2.3	5.4	60.7	30.0	0.7	1.0
1512	373	0.0	1.1	6.2	59.5	30.8	1.9	0.5
1514	682	0.1	1.0	4.5	58.7	32.1	2.9	0.6
1701	513	0.0	1.8	5.1	62.4	28.8	0.8	1.2
1950	568	0.0	3.2	8.1	61.3	25.2	1.1	1.2
1952	132	0.0	1.5	2.3	63.6	29.5	1.5	1.5
1953	271	0.0	1.5	8.5	61.6	25.1	2.2	1.1
1954	356	0.0	2.0	8.1	63.2	25.0	1.4	0.3
1955	124	0.0	2.4	8.1	58.1	28.2	2.4	0.8
1957	382	0.0	1.0	3.7	58.9	34.3	1.3	0.8
1963	71	*	*	*	*	*	*	*
2150	553	0.2	0.4	6.3	62.9	28.2	1.8	0.2
2151	227	0.4	2.6	10.6	61.2	23.8	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 69

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2152	38	*	*	*	*	*	*	*
2251	376	0.3	0.3	5.1	62.2	31.1	1.1	0.0
2252	169	0.0	5.3	5.9	58.6	28.4	1.2	0.6
2554	49	*	*	*	*	*	*	*
2555	7,094	0.1	1.8	6.9	62.0	27.1	1.6	0.5
2556	117	0.0	0.0	2.6	72.6	23.1	1.7	0.0
2557	1	*	*	*	*	*	*	*
2560	1	*	*	*	*	*	*	*
2562	3	*	*	*	*	*	*	*
2563	6	*	*	*	*	*	*	*
2564	3	*	*	*	*	*	*	*
3001	1,159	0.0	1.2	7.9	63.8	23.7	1.2	2.2
3002	304	0.0	1.0	6.6	57.9	28.6	3.0	3.0
3004	301	0.0	1.0	8.6	53.8	34.2	1.3	1.0
3005	160	0.0	0.0	6.9	60.6	28.1	3.1	1.3
3007	7	*	*	*	*	*	*	*
3201	407	0.0	0.7	2.9	60.7	33.7	1.2	0.7
3301	4,300	0.1	2.2	5.7	59.7	28.9	2.2	1.2
3304	100	0.0	0.0	6.0	49.0	39.0	6.0	0.0
3305	158	0.0	0.6	1.9	60.1	32.9	4.4	0.0
3306	206	0.0	1.5	1.9	59.2	32.5	3.9	1.0
3308	94	*	*	*	*	*	*	*
3312	466	0.0	1.5	6.4	57.9	31.3	1.9	0.9
3314	1	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 70

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3550	64	*	*	*	*	*	*	*
3551	6	*	*	*	*	*	*	*
3552	1	*	*	*	*	*	*	*
3701	837	0.2	1.7	6.5	62.8	25.9	1.7	1.2
3850	2,696	0.1	2.6	6.9	61.4	25.7	1.2	2.1
3901	1,467	0.1	2.7	7.8	57.9	27.9	2.3	1.2
3902	9	*	*	*	*	*	*	*
3903	2	*	*	*	*	*	*	*
3950	2,001	0.0	2.1	6.9	60.2	27.5	1.9	1.3
3954	1	*	*	*	*	*	*	*
4106	2,174	0.1	2.2	6.3	58.1	28.2	3.2	1.8
4107	2,780	0.1	1.9	6.0	59.0	29.5	2.4	1.1
4109	639	0.0	1.9	4.2	58.1	33.0	2.3	0.5
4112	585	0.3	4.1	8.4	54.0	28.0	3.9	1.2
4115	3,376	0.1	1.1	4.8	55.6	34.5	3.1	0.7
4351	524	0.0	1.0	5.7	67.9	21.2	1.9	2.3
4352	185	0.0	2.2	7.0	60.0	28.1	1.6	1.1
4353	474	0.0	1.3	7.2	61.0	25.5	3.6	1.5
4354	754	0.0	2.0	7.3	64.3	23.7	2.0	0.7
4355	264	0.4	0.8	7.6	53.8	32.2	3.4	1.9
4356	381	0.0	1.6	7.9	63.3	24.4	0.8	2.1
4359	580	0.2	2.4	5.7	61.4	26.6	2.8	1.0
4360	229	0.0	1.3	7.0	60.7	27.9	1.7	1.3
4361	244	0.0	1.2	11.1	61.9	23.4	2.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 71

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4362	267	0.0	3.0	9.0	56.9	28.1	1.9	1.1
4363	641	0.0	2.2	6.7	63.8	24.6	1.2	1.4
4401	730	0.0	1.5	4.7	62.9	28.2	1.8	1.0
4501	750	0.0	1.9	6.1	65.6	23.6	1.6	1.2
4601	1,188	0.0	1.0	6.5	64.0	26.5	1.2	0.8
4701	999	0.0	1.2	4.9	56.7	34.3	2.3	0.6
5050	1,619	0.1	0.9	3.9	56.8	35.6	2.5	0.2
5051	2,574	0.0	0.7	3.9	59.1	33.5	2.4	0.4
5052	2,671	0.0	1.0	4.1	55.6	36.7	2.0	0.6
5055	1,349	0.0	0.4	1.6	46.8	47.4	3.6	0.1
5101	1,456	0.1	0.8	3.1	58.0	35.6	2.3	0.2
5102	5	*	*	*	*	*	*	*
5202	644	0.0	1.7	4.8	62.7	28.7	2.0	0.0
5206	11	*	*	*	*	*	*	*
5601	962	0.0	0.7	5.9	63.6	28.4	1.1	0.2
5801	1,769	0.0	1.3	6.6	64.4	25.8	1.4	0.5
5803	21	*	*	*	*	*	*	*
5804	10	*	*	*	*	*	*	*
5806	57	*	*	*	*	*	*	*
6101	65	*	*	*	*	*	*	*
6102	68	*	*	*	*	*	*	*
6103	3,246	0.1	2.5	7.7	61.2	25.4	1.4	1.5
6105	6	*	*	*	*	*	*	*
6301	3,378	0.1	1.8	6.0	58.0	31.9	1.9	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 72

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
6312	1,773	0.0	0.6	3.8	49.9	41.8	3.7	0.3
6316	1,127	0.0	0.4	2.9	52.5	39.0	5.0	0.1
6318	10	*	*	*	*	*	*	*
6320	562	0.0	0.2	3.7	56.2	35.6	3.6	0.7
6325	322	0.0	1.2	2.8	59.0	34.8	0.6	1.6
6335	928	0.0	0.4	2.9	52.3	40.8	3.1	0.4
6345	145	0.0	1.4	4.1	63.4	28.3	2.1	0.7
6360	903	0.0	1.6	5.1	60.0	30.7	1.6	1.1
7306	1,874	0.3	2.6	6.4	59.4	29.1	1.7	0.5
7401	1,777	0.0	1.6	7.5	61.5	26.6	1.6	1.2
7601	505	0.0	2.2	6.7	63.8	25.7	1.0	0.6
7850	1,040	0.1	2.4	6.3	64.7	25.0	1.2	0.3
7852	8	*	*	*	*	*	*	*
8101	1,705	0.2	2.2	6.2	54.1	33.8	2.4	1.1
8102	444	0.0	2.0	5.0	59.2	31.5	2.0	0.2
8105	1	*	*	*	*	*	*	*
8106	550	0.0	1.5	4.7	60.9	31.5	1.3	0.2
8150	7	*	*	*	*	*	*	*
8151	3,223	0.0	1.4	5.3	54.6	34.8	3.4	0.6
8152	7	*	*	*	*	*	*	*
8153	1	*	*	*	*	*	*	*
8155	7	*	*	*	*	*	*	*
8201	2,587	0.0	1.3	5.2	56.4	34.5	2.1	0.5
8202	3,922	0.0	0.9	3.3	51.2	38.7	5.4	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 73

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8203	358	0.0	1.1	5.3	58.7	30.4	4.2	0.3
8250	464	0.0	2.4	5.0	53.2	35.8	3.4	0.2
8252	133	0.0	0.8	3.8	50.4	42.9	2.3	0.0
8255	1,897	0.1	1.1	4.2	60.2	32.4	1.9	0.1
8257	1,627	0.0	0.6	2.5	53.7	38.6	4.5	0.1
8263	389	0.0	1.5	5.7	56.6	34.2	2.1	0.0
8264	795	0.1	0.6	3.1	59.2	33.3	3.3	0.3
8267	707	0.0	0.6	4.8	60.1	31.8	2.3	0.4
8901	794	0.1	2.8	5.7	55.5	32.1	2.8	1.0
8902	1,252	0.0	1.4	4.8	51.9	38.1	3.4	0.5
8903	1,347	0.0	2.2	6.0	56.8	31.5	2.4	1.1
8904	1,935	0.3	2.2	6.6	56.8	31.6	1.8	0.7
8905	1,898	0.0	0.7	3.4	55.0	37.0	3.5	0.3
8909	1,168	0.0	1.6	6.8	57.7	30.0	3.4	0.5
9045	1,310	0.2	2.4	6.3	54.8	31.8	3.8	0.8
9050	882	0.1	2.0	5.8	57.5	31.4	2.3	0.9
9051	1,649	0.1	2.1	5.6	55.1	32.6	3.7	0.9
9053	1,597	0.0	2.0	4.8	57.6	31.6	3.1	0.9
9055	7	*	*	*	*	*	*	*
9058	1,409	0.1	1.3	5.8	54.7	34.8	2.7	0.5
9066	689	0.1	2.8	6.0	52.4	36.3	2.3	0.1
9068	818	0.2	3.2	6.8	55.3	29.8	2.4	2.2
9069	1,657	0.2	2.7	5.2	53.5	35.5	2.5	0.4
9070	1,892	0.1	2.9	7.6	54.7	32.0	1.6	1.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 4A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 74

Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9072	59	*	*	*	*	*	*	*
9080	2,628	0.0	1.6	6.3	52.9	35.5	2.8	0.9
9082	108	0.0	1.9	4.6	59.3	33.3	0.9	0.0
9083	13	*	*	*	*	*	*	*
9087	11	*	*	*	*	*	*	*
Michigan	142,422	0.1	1.7	5.9	58.4	30.8	2.3	0.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 75

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	99	*	*	55	*	*	7	*	15	*	112	7.1 (18)	112	14.3 (67)
0102	288	6.3 (127)	62.5 (74)	145	15.9 (33)	56.6 (88)	17	*	37	*	343	9.0 (64)	343	12.8 (37)
0103	305	3.3 (24)	65.9 (114)	157	9.6 (1)	62.4 (112)	27	*	40	*	349	9.2 (70)	350	16.0 (100)
0104	94	*	*	32	*	*	6	*	22	*	116	7.8 (34)	116	14.7 (77)
0105	4	*	*	2	*	*	0	*	0	*	4	*	4	*
0250	99	*	*	59	*	*	7	*	22	*	108	12.0 (128)	108	15.7 (93)
0251	54	*	*	25	*	*	4	*	16	*	67	*	67	*
0252	92	*	*	68	*	*	4	*	8	*	91	*	91	*
0253	118	3.4 (27)	66.9 (120)	87	*	*	11	*	16	*	115	13.0 (134)	115	12.2 (28)
0401	379	5.5 (114)	60.9 (44)	157	15.3 (25)	59.9 (105)	28	*	37	*	385	10.6 (104)	386	17.6 (122)
0402	276	5.1 (103)	60.5 (34)	64	*	*	23	*	34	*	316	11.1 (113)	316	13.3 (49)
0403	108	5.6 (117)	61.1 (50)	54	*	*	5	*	7	*	111	11.7 (122)	111	8.1 (6)
0404	89	*	*	27	*	*	5	*	11	*	98	*	98	*
0501	263	4.2 (70)	56.3 (11)	55	*	*	53	*	80	*	374	8.3 (46)	374	12.3 (31)
0503	39	*	*	10	*	*	4	*	12	*	55	*	55	*
0504	280	1.8 (4)	63.9 (97)	111	20.7 (97)	53.2 (52)	44	*	56	*	305	7.5 (28)	305	13.1 (44)
0505	135	4.4 (80)	54.8 (7)	33	*	*	29	*	39	*	185	8.1 (41)	185	14.6 (75)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 76

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	117	4.3 (75)	47.9 (1)	27	*	*	21	*	32	*	163	6.7 (14)	163	12.3 (31)
0507	33	*	*	10	*	*	5	*	18	*	70	*	70	*
0508	5	*	*	3	*	*	1	*	3	*	13	*	13	*
0650	907	4.5 (84)	63.7 (92)	562	13.5 (11)	57.1 (93)	57	*	118	20.3 (14)	1,006	11.8 (125)	1,007	17.2 (117)
0651	410	5.4 (113)	61.2 (52)	134	16.4 (39)	59.0 (101)	26	*	64	*	493	9.1 (67)	495	14.1 (62)
0652	250	2.8 (13)	64.4 (106)	197	21.3 (99)	47.7 (15)	24	*	28	*	279	8.6 (51)	279	15.8 (94)
0653	333	4.8 (93)	60.7 (38)	227	19.8 (88)	48.0 (17)	31	*	46	*	353	9.9 (88)	355	16.1 (101)
0654	236	5.5 (114)	60.6 (35)	137	15.3 (25)	53.3 (57)	14	*	50	*	388	9.8 (85)	389	15.9 (98)
0655	288	5.6 (117)	58.0 (17)	93	*	*	17	*	15	*	297	11.8 (125)	298	18.8 (128)
0657	20	*	*	9	*	*	0	*	4	*	34	*	34	*
0701	400	3.8 (46)	56.5 (13)	261	17.2 (54)	45.6 (8)	11	*	36	*	418	9.3 (73)	416	17.1 (115)
0702	65	*	*	31	*	*	2	*	6	*	64	*	64	*
0703	135	2.2 (6)	60.7 (38)	80	*	*	11	*	20	*	144	11.8 (125)	144	11.1 (17)
0704	43	*	*	24	*	*	4	*	6	*	42	*	43	*
0705	48	*	*	33	*	*	4	*	8	*	58	*	58	*
0706	1	*	*	1	*	*	0	*	0	*	1	*	1	*
0850	419	4.8 (93)	63.2 (85)	122	21.3 (99)	47.5 (13)	13	*	39	*	631	10.6 (104)	632	15.8 (94)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 77

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0851	402	5.7 (120)	60.7 (38)	158	17.1 (53)	57.0 (92)	17	*	50	*	563	11.2 (115)	566	16.6 (110)
0901	1,058	4.5 (84)	61.0 (47)	584	16.1 (37)	57.4 (95)	33	*	85	*	1,336	8.9 (63)	1,337	15.0 (85)
0902	20	*	*	13	*	*	0	*	1	*	24	*	24	*
1051	896	5.6 (117)	55.5 (8)	630	16.0 (35)	54.3 (69)	49	*	83	*	1,016	7.8 (34)	1,021	12.1 (26)
1052	2	*	*	1	*	*	1	*	0	*	3	*	3	*
1101	1,172	4.4 (80)	63.0 (84)	788	18.5 (76)	52.0 (36)	87	*	159	25.8 (26)	1,247	9.1 (67)	1,249	19.3 (133)
1102	641	5.3 (111)	60.8 (41)	447	16.6 (44)	56.6 (88)	44	*	76	*	646	8.8 (59)	648	16.7 (112)
1106	52	*	*	41	*	*	0	*	7	*	49	*	51	*
1250	381	5.2 (109)	59.8 (28)	136	24.3 (109)	47.8 (16)	63	*	101	38.6 (38)	898	10.0 (92)	899	10.3 (13)
1251	563	4.3 (75)	61.1 (50)	414	15.5 (27)	54.1 (68)	40	*	74	*	587	9.9 (88)	589	8.5 (7)
1252	397	6.0 (125)	63.2 (85)	165	17.6 (60)	53.3 (57)	25	*	54	*	466	10.1 (94)	467	11.8 (23)
1253	552	6.3 (127)	59.8 (28)	235	17.0 (50)	60.0 (107)	48	*	73	*	579	9.8 (85)	579	14.7 (77)
1350	2,048	4.1 (65)	62.1 (68)	1,476	18.1 (73)	53.7 (64)	64	*	78	*	2,192	7.3 (24)	2,193	11.4 (19)
1351	0	*	*	0	*	*	0	*	0	*	1	*	1	*
1352	262	5.3 (111)	64.1 (100)	177	15.8 (32)	59.9 (105)	8	*	12	*	325	5.5 (4)	325	9.2 (10)
1401	170	5.9 (123)	51.2 (2)	98	*	*	9	*	34	*	192	5.2 (2)	192	13.5 (54)
1402	59	*	*	30	*	*	6	*	5	*	77	*	77	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 78

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1403	13	*	*	8	*	*	0	*	2	*	17	*	17	*
1501	582	3.4 (27)	67.5 (123)	245	38.0 (114)	37.6 (2)	95	*	112	12.5 (3)	667	13.0 (134)	668	13.9 (59)
1503	673	5.1 (103)	64.2 (103)	483	20.5 (92)	53.6 (62)	91	*	99	*	741	11.2 (115)	740	15.8 (94)
1504	572	5.1 (103)	56.1 (10)	209	11.5 (5)	52.2 (41)	203	34.5 (3)	320	24.7 (23)	1,684	9.9 (88)	1,685	9.0 (9)
1505	9	*	*	0	*	*	0	*	0	*	19	*	19	*
1506	432	3.5 (32)	62.5 (74)	67	*	*	68	*	110	20.9 (15)	492	12.0 (128)	492	15.4 (90)
1507	364	8.5 (133)	54.7 (6)	57	*	*	37	*	114	42.1 (39)	482	11.2 (115)	483	11.2 (18)
1509	255	3.9 (50)	66.7 (119)	112	26.8 (113)	50.0 (22)	45	*	73	*	330	8.8 (59)	331	7.6 (5)
1511	471	4.0 (57)	64.8 (107)	279	17.9 (68)	52.0 (36)	71	*	135	22.2 (16)	574	11.7 (122)	575	14.6 (75)
1512	249	4.0 (57)	61.4 (53)	125	16.8 (48)	62.4 (112)	34	*	130	34.6 (36)	353	8.8 (59)	354	14.7 (77)
1514	366	3.6 (35)	53.8 (5)	102	25.5 (111)	50.0 (22)	56	*	224	16.5 (6)	643	9.3 (73)	644	15.2 (89)
1701	455	3.1 (19)	60.2 (32)	313	16.6 (44)	54.6 (71)	33	*	72	*	466	12.0 (128)	466	8.6 (8)
1950	501	3.0 (17)	63.3 (87)	349	13.8 (13)	58.7 (100)	12	*	39	*	534	10.7 (107)	535	16.1 (101)
1952	115	1.7 (3)	73.0 (133)	84	*	*	4	*	9	*	126	10.3 (97)	126	19.0 (130)
1953	229	6.6 (130)	59.0 (23)	164	15.9 (33)	50.6 (27)	5	*	32	*	248	10.5 (102)	248	16.5 (109)
1954	322	1.9 (5)	68.9 (127)	196	25.5 (111)	50.5 (25)	7	*	28	*	336	11.0 (112)	336	12.2 (28)
1955	111	2.7 (12)	68.5 (126)	75	*	*	0	*	12	*	113	15.0 (139)	113	15.0 (85)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 79

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1957	336	3.9 (50)	69.0 (128)	215	11.2 (3)	55.3 (77)	12	*	27	*	357	14.8 (138)	358	11.7 (21)
1963	59	*	*	36	*	*	0	*	8	*	65	*	65	*
2150	519	3.1 (19)	62.0 (64)	371	23.2 (107)	46.4 (9)	53	*	113	15.0 (4)	513	10.1 (94)	514	15.4 (90)
2151	213	4.2 (70)	61.0 (47)	168	21.4 (101)	52.4 (46)	23	*	48	*	196	7.1 (18)	196	16.8 (113)
2152	34	*	*	18	*	*	3	*	2	*	32	*	32	*
2251	272	5.1 (103)	66.2 (117)	157	17.8 (65)	52.2 (41)	0	*	7	*	341	3.8 (1)	341	16.1 (101)
2252	89	*	*	24	*	*	0	*	0	*	153	9.8 (85)	153	13.1 (44)
2554	29	*	*	8	*	*	7	*	5	*	45	*	45	*
2555	5,086	4.6 (90)	61.0 (47)	2,411	19.7 (87)	53.0 (51)	121	42.1 (7)	408	28.4 (30)	6,503	9.5 (78)	6,512	17.3 (118)
2556	80	*	*	34	*	*	0	*	2	*	113	8.8 (59)	113	14.2 (64)
2557	1	*	*	0	*	*	0	*	0	*	1	*	1	*
2560	1	*	*	0	*	*	0	*	0	*	1	*	1	*
2562	2	*	*	0	*	*	0	*	0	*	3	*	3	*
2563	3	*	*	3	*	*	1	*	0	*	5	*	5	*
2564	1	*	*	0	*	*	0	*	0	*	2	*	2	*
3001	716	4.3 (75)	67.5 (123)	276	18.8 (82)	45.3 (6)	11	*	44	*	1,096	8.6 (51)	1,095	17.1 (115)
3002	217	5.1 (103)	62.7 (78)	105	18.1 (73)	47.6 (14)	11	*	31	*	288	6.6 (11)	288	14.2 (64)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 80

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3004	168	4.8 (93)	63.7 (92)	61	*	*	3	*	8	*	289	6.6 (11)	289	15.6 (92)
3005	85	*	*	28	*	*	1	*	7	*	152	8.6 (51)	152	11.8 (23)
3007	5	*	*	1	*	*	0	*	0	*	7	*	7	*
3201	345	4.9 (97)	67.0 (121)	221	19.5 (85)	58.4 (99)	42	*	78	*	388	11.1 (113)	388	14.4 (70)
3301	3,509	3.6 (35)	60.9 (44)	2,250	17.6 (60)	55.4 (80)	111	33.3 (2)	135	17.0 (7)	4,094	7.4 (27)	4,098	12.9 (41)
3304	85	*	*	53	*	*	2	*	0	*	96	*	96	*
3305	134	3.7 (38)	59.0 (23)	88	*	*	3	*	1	*	148	9.5 (78)	148	7.4 (4)
3306	180	2.2 (6)	72.2 (132)	114	16.7 (47)	55.3 (77)	4	*	5	*	200	8.0 (40)	200	13.5 (54)
3308	88	*	*	58	*	*	4	*	2	*	82	*	83	*
3312	400	3.8 (46)	61.5 (54)	247	17.4 (57)	55.1 (76)	12	*	9	*	439	8.7 (57)	438	15.8 (94)
3314	1	*	*	0	*	*	0	*	0	*	1	*	1	*
3550	50	*	*	30	*	*	1	*	2	*	61	*	61	*
3551	3	*	*	1	*	*	0	*	0	*	5	*	5	*
3552	1	*	*	1	*	*	0	*	0	*	1	*	1	*
3701	630	4.0 (57)	64.3 (105)	470	20.6 (94)	49.8 (21)	83	*	139	10.1 (2)	778	11.3 (118)	781	19.1 (131)
3850	2,088	4.5 (84)	59.7 (27)	1,382	15.2 (23)	55.7 (82)	69	*	153	19.6 (11)	2,507	9.4 (76)	2,507	14.2 (64)
3901	1,134	4.1 (65)	61.6 (56)	519	15.2 (23)	59.5 (104)	18	*	73	*	1,323	7.6 (29)	1,321	14.5 (73)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 81

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3902	7	*	*	3	*	*	0	*	0	*	9	*	9	*
3903	1	*	*	1	*	*	0	*	0	*	2	*	2	*
3950	1,503	4.2 (70)	62.3 (72)	809	14.7 (18)	55.9 (84)	36	*	102	26.5 (28)	1,849	10.7 (107)	1,854	12.6 (34)
3954	1	*	*	0	*	*	0	*	0	*	1	*	1	*
4106	830	3.4 (27)	61.7 (58)	214	21.5 (102)	43.9 (5)	284	32.4 (1)	509	18.3 (8)	2,030	8.6 (51)	2,036	17.7 (124)
4107	1,727	3.8 (46)	58.9 (21)	594	17.0 (50)	51.9 (35)	367	37.9 (6)	702	22.9 (18)	2,542	8.2 (43)	2,549	19.3 (133)
4109	375	4.0 (57)	58.7 (20)	204	11.3 (4)	56.9 (91)	72	*	211	19.9 (12)	606	7.6 (29)	606	17.5 (121)
4112	568	1.6 (1)	68.0 (125)	476	50.0 (115)	25.2 (1)	5	*	24	*	555	21.3 (141)	555	24.1 (141)
4115	1,607	4.1 (65)	60.8 (41)	503	15.7 (31)	52.1 (38)	418	35.4 (4)	942	26.2 (27)	3,180	9.3 (73)	3,181	17.3 (118)
4351	459	4.1 (65)	60.8 (41)	271	20.7 (97)	50.6 (27)	7	*	25	*	484	12.8 (133)	484	22.9 (140)
4352	160	6.3 (127)	56.9 (15)	117	17.9 (68)	55.6 (81)	2	*	9	*	168	10.7 (107)	168	20.8 (136)
4353	357	3.9 (50)	69.2 (129)	265	20.0 (89)	55.8 (83)	24	*	40	*	428	14.0 (137)	428	12.6 (34)
4354	499	4.4 (80)	63.9 (97)	142	15.5 (27)	52.1 (38)	6	*	37	*	709	10.6 (104)	709	20.9 (137)
4355	180	7.8 (132)	60.6 (35)	110	14.5 (16)	58.2 (98)	5	*	19	*	243	9.5 (78)	242	14.0 (61)
4356	350	2.6 (11)	67.1 (122)	256	17.6 (60)	47.3 (11)	8	*	23	*	350	11.7 (122)	351	17.9 (125)
4359	515	3.3 (24)	66.4 (118)	360	25.3 (110)	46.9 (10)	5	*	29	*	535	9.0 (64)	535	17.6 (122)
4360	211	6.2 (126)	61.6 (56)	149	18.8 (82)	55.0 (74)	11	*	20	*	214	8.4 (47)	214	12.1 (26)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 82

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4361	223	2.2 (6)	62.8 (81)	152	21.7 (103)	53.3 (57)	4	*	15	*	230	7.0 (17)	230	11.7 (21)
4362	235	5.1 (103)	55.7 (9)	171	11.7 (6)	60.2 (108)	8	*	20	*	245	7.8 (34)	246	13.0 (42)
4363	552	5.8 (122)	58.0 (17)	389	19.5 (85)	52.7 (48)	8	*	50	*	604	10.4 (98)	604	14.4 (70)
4401	708	4.9 (97)	60.6 (35)	595	18.7 (78)	51.4 (32)	23	*	47	*	684	6.1 (8)	686	10.3 (13)
4501	571	4.2 (70)	63.6 (91)	330	18.5 (76)	50.0 (22)	37	*	51	*	720	11.3 (118)	720	14.7 (77)
4601	996	3.7 (38)	61.7 (58)	653	15.6 (29)	52.2 (41)	46	*	128	26.6 (29)	1,050	9.7 (84)	1,050	16.4 (108)
4701	834	5.5 (114)	57.4 (16)	490	20.2 (90)	52.7 (48)	38	*	106	9.4 (1)	924	10.0 (92)	926	21.3 (139)
5050	1,544	4.0 (57)	65.9 (114)	1,277	16.9 (49)	52.3 (44)	25	*	221	32.1 (35)	1,514	9.6 (82)	1,515	16.6 (110)
5051	2,398	3.8 (46)	61.9 (63)	1,872	18.0 (72)	54.6 (71)	58	*	344	28.5 (31)	2,374	9.2 (70)	2,373	13.1 (44)
5052	2,511	3.1 (19)	62.0 (64)	2,026	20.5 (92)	47.4 (12)	79	*	273	24.5 (22)	2,536	12.3 (131)	2,534	13.5 (54)
5055	1,075	1.6 (1)	62.6 (77)	814	22.9 (105)	45.3 (6)	20	*	61	*	1,297	13.5 (136)	1,297	7.0 (3)
5101	1,112	3.7 (38)	62.7 (78)	854	12.4 (7)	56.3 (86)	95	*	290	23.4 (19)	1,354	7.9 (38)	1,358	12.4 (33)
5102	3	*	*	1	*	*	0	*	0	*	3	*	3	*
5202	532	4.5 (84)	56.8 (14)	344	17.4 (57)	61.6 (111)	5	*	37	*	590	9.2 (70)	592	12.7 (36)
5206	8	*	*	5	*	*	0	*	1	*	11	*	11	*
5601	711	3.5 (32)	63.7 (92)	328	9.8 (2)	59.1 (102)	51	*	111	23.4 (19)	917	9.9 (88)	915	13.9 (59)
5801	1,648	4.9 (97)	64.9 (110)	1,283	18.9 (84)	53.2 (52)	55	*	128	15.6 (5)	1,615	8.5 (48)	1,617	18.8 (128)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 83

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5803	21	*	*	18	*	*	0	*	0	*	21	*	21	*
5804	10	*	*	7	*	*	0	*	0	*	10	*	10	*
5806	49	*	*	30	*	*	0	*	9	*	54	*	54	*
6101	60	*	*	44	*	*	1	*	1	*	60	*	61	*
6102	58	*	*	47	*	*	2	*	2	*	63	*	63	*
6103	2,903	3.9 (50)	63.5 (89)	2,192	17.8 (65)	53.3 (57)	42	*	52	*	3,003	9.6 (82)	3,008	19.1 (131)
6105	3	*	*	1	*	*	0	*	0	*	5	*	5	*
6301	2,892	4.0 (57)	62.0 (64)	1,931	17.7 (63)	49.5 (19)	93	*	226	25.2 (25)	3,069	5.9 (6)	3,070	13.1 (44)
6312	1,549	4.1 (65)	60.9 (44)	1,010	17.3 (56)	52.4 (46)	62	*	168	38.1 (37)	1,628	8.5 (48)	1,631	10.5 (15)
6316	1,006	4.9 (97)	56.3 (11)	676	16.4 (39)	55.0 (74)	28	*	78	*	1,043	7.9 (38)	1,043	11.6 (20)
6318	8	*	*	5	*	*	1	*	0	*	9	*	9	*
6320	412	3.4 (27)	64.1 (100)	193	16.6 (44)	54.9 (73)	11	*	43	*	501	10.4 (98)	502	17.9 (125)
6325	228	3.9 (50)	60.1 (31)	118	12.7 (8)	60.2 (108)	6	*	31	*	291	8.6 (51)	291	17.9 (125)
6335	687	5.7 (120)	59.2 (25)	343	14.0 (15)	51.6 (33)	40	*	93	*	868	11.6 (120)	869	14.4 (70)
6345	110	7.3 (131)	61.8 (61)	57	*	*	3	*	13	*	129	11.6 (120)	129	16.3 (105)
6360	652	4.3 (75)	58.9 (21)	321	15.0 (22)	51.7 (34)	28	*	69	*	829	7.7 (32)	829	15.1 (87)
7306	1,464	3.1 (19)	64.2 (103)	1,012	17.7 (63)	53.2 (52)	61	*	99	*	1,784	10.4 (98)	1,785	12.8 (37)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 84

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators ^(2,3)

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7401	1,667	4.2 (70)	63.5 (89)	1,306	16.5 (42)	54.4 (70)	81	*	241	31.5 (33)	1,587	9.5 (78)	1,592	15.1 (87)
7601	186	5.9 (123)	58.6 (19)	29	*	*	16	*	58	*	463	8.2 (43)	463	13.4 (52)
7850	889	4.9 (97)	62.8 (81)	711	22.1 (104)	50.8 (29)	22	*	60	*	964	6.8 (15)	966	9.2 (10)
7852	8	*	*	4	*	*	0	*	0	*	6	*	6	*
8101	1,445	3.7 (38)	64.1 (100)	928	14.8 (19)	56.4 (87)	23	*	68	*	1,598	7.8 (34)	1,597	17.3 (118)
8102	346	2.3 (9)	69.4 (131)	184	17.9 (68)	53.3 (57)	2	*	8	*	413	6.8 (15)	413	10.7 (16)
8105	1	*	*	0	*	*	0	*	0	*	0	*	0	*
8106	267	5.2 (109)	59.6 (26)	99	*	*	8	*	27	*	521	7.1 (18)	521	19.8 (135)
8150	6	*	*	2	*	*	0	*	1	*	7	*	7	*
8151	2,515	4.0 (57)	62.0 (64)	1,589	20.6 (94)	49.4 (18)	272	37.5 (5)	505	22.4 (17)	2,898	9.1 (67)	2,899	14.9 (83)
8152	5	*	*	3	*	*	0	*	3	*	7	*	7	*
8153	1	*	*	0	*	*	1	*	0	*	0	*	0	*
8155	5	*	*	1	*	*	0	*	0	*	6	*	6	*
8201	2,010	3.6 (35)	62.1 (68)	1,172	16.4 (39)	56.1 (85)	79	*	300	23.7 (21)	2,276	10.7 (107)	2,281	14.7 (77)
8202	1,423	3.9 (50)	52.4 (4)	731	17.5 (59)	49.5 (19)	74	*	175	18.3 (8)	3,802	7.1 (18)	3,801	3.5 (1)
8203	313	4.8 (93)	63.3 (87)	218	17.9 (68)	50.9 (30)	7	*	54	*	315	10.2 (96)	313	14.1 (62)
8250	369	3.5 (32)	65.6 (113)	216	14.8 (19)	53.2 (52)	10	*	52	*	406	7.1 (18)	407	12.8 (37)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 85

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8252	78	*	*	37	*	*	0	*	2	*	118	18.6 (140)	119	14.3 (67)
8255	1,618	4.5 (84)	62.2 (70)	1,010	18.7 (78)	54.0 (67)	69	*	235	24.7 (23)	1,693	10.5 (102)	1,695	14.3 (67)
8257	916	5.0 (102)	51.3 (3)	578	23.5 (108)	41.0 (3)	51	*	105	20.0 (13)	1,543	8.7 (57)	1,547	4.1 (2)
8263	330	3.0 (17)	60.0 (30)	176	13.1 (10)	57.4 (95)	0	*	27	*	320	8.1 (41)	320	12.8 (37)
8264	667	3.7 (38)	64.8 (107)	460	13.0 (9)	62.4 (112)	28	*	130	19.2 (10)	725	8.6 (51)	724	13.1 (44)
8267	601	4.0 (57)	65.9 (114)	359	20.3 (91)	53.8 (65)	45	*	128	28.9 (32)	647	8.5 (48)	647	16.2 (104)
8901	657	3.7 (38)	64.8 (107)	406	16.5 (42)	53.2 (52)	4	*	19	*	757	8.2 (43)	756	16.3 (105)
8902	945	4.6 (90)	61.7 (58)	519	17.0 (50)	53.8 (65)	18	*	56	*	1,164	7.3 (24)	1,162	13.5 (54)
8903	973	2.8 (13)	65.4 (112)	628	13.5 (11)	61.5 (110)	12	*	30	*	1,251	6.5 (10)	1,258	14.9 (83)
8904	1,583	3.4 (27)	65.2 (111)	903	16.1 (37)	52.9 (50)	23	*	69	*	1,818	7.7 (32)	1,820	16.3 (105)
8905	1,378	3.7 (38)	60.4 (33)	730	17.8 (65)	53.6 (62)	15	*	66	*	1,792	12.3 (131)	1,792	13.3 (49)
8909	939	3.9 (50)	62.8 (81)	545	14.9 (21)	59.4 (103)	14	*	45	*	1,072	7.6 (29)	1,071	14.8 (82)
9045	963	3.3 (24)	69.2 (129)	587	16.0 (35)	52.1 (38)	12	*	37	*	1,230	9.4 (76)	1,229	9.5 (12)
9050	763	4.6 (90)	63.7 (92)	216	13.9 (14)	64.4 (115)	10	*	23	*	828	7.1 (18)	827	13.4 (52)
9051	703	2.8 (13)	63.7 (92)	310	22.9 (105)	41.3 (4)	27	*	95	*	1,562	10.8 (111)	1,562	12.0 (25)
9053	1,426	3.2 (23)	61.5 (54)	1,095	14.6 (17)	57.4 (95)	24	*	41	*	1,522	6.3 (9)	1,522	12.2 (28)
9055	6	*	*	5	*	*	0	*	0	*	6	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 5A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 86

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9058	936	4.3 (75)	62.7 (78)	667	18.3 (75)	52.3 (44)	12	*	46	*	1,339	9.0 (64)	1,340	15.9 (98)
9066	573	4.5 (84)	62.3 (72)	378	20.6 (94)	51.1 (31)	0	*	7	*	657	5.9 (6)	658	21.0 (138)
9068	766	2.3 (9)	64.0 (99)	635	18.7 (78)	57.3 (94)	11	*	20	*	760	5.8 (5)	762	13.0 (42)
9069	1,295	3.7 (38)	62.2 (70)	866	18.7 (78)	50.5 (25)	28	*	55	*	1,566	7.3 (24)	1,567	13.3 (49)
9070	1,687	4.4 (80)	62.5 (74)	1,255	15.6 (29)	56.6 (88)	25	*	65	*	1,768	5.4 (3)	1,771	13.7 (58)
9072	53	*	*	16	*	*	0	*	0	*	47	*	47	*
9080	1,905	2.9 (16)	61.8 (61)	998	17.2 (54)	55.3 (77)	37	*	100	32.0 (34)	2,486	10.4 (98)	2,484	14.5 (73)
9082	88	*	*	48	*	*	0	*	4	*	106	6.6 (11)	106	17.0 (114)
9083	11	*	*	7	*	*	0	*	2	*	12	*	12	*
9087	10	*	*	6	*	*	0	*	0	*	10	*	10	*
Michigan	107,908	4.0	62.0	65,438	17.9	53.1	5,611	33.3	13,095	23.8	131,970	9.1	132,088	14.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy

(ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 87

Michigan

Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	106	88.7 (110)	2.8 (132)	120	30.0 (70)	121	30.6 (44)	85	*
0102	321	90.0 (102)	0.9 (66)	363	32.2 (48)	362	24.9 (80)	273	90.5 (85)
0103	332	92.2 (69)	0.9 (66)	364	36.0 (27)	360	19.7 (113)	281	93.6 (36)
0104	112	93.8 (43)	0.0 (1)	124	41.9 (10)	125	28.8 (55)	99	*
0105	4	*	*	4	*	4	*	4	*
0250	94	*	*	112	41.1 (13)	114	29.8 (47)	89	*
0251	66	*	*	75	*	73	*	58	*
0252	91	*	*	100	42.0 (9)	97	*	76	*
0253	110	92.7 (64)	0.0 (1)	120	45.8 (3)	120	26.7 (69)	102	93.1 (44)
0401	366	94.5 (30)	0.5 (24)	401	26.4 (91)	402	27.6 (64)	285	95.1 (18)
0402	282	87.9 (115)	2.5 (128)	341	32.0 (51)	342	31.6 (39)	281	96.4 (6)
0403	104	89.4 (107)	0.0 (1)	118	39.0 (16)	116	26.7 (69)	93	*
0404	96	*	*	105	28.6 (77)	105	39.0 (18)	86	*
0501	360	96.7 (7)	0.6 (32)	388	28.9 (76)	368	28.3 (57)	291	90.4 (86)
0503	55	*	*	56	*	58	*	40	*
0504	295	93.9 (41)	0.7 (45)	311	30.5 (68)	312	40.1 (14)	240	92.9 (46)
0505	180	95.0 (22)	1.1 (78)	194	23.7 (109)	192	39.6 (15)	145	91.0 (74)
0506	151	91.4 (81)	0.7 (45)	169	32.5 (45)	168	25.0 (79)	122	91.0 (74)
0507	69	*	*	69	*	71	*	42	*
0508	12	*	*	13	*	13	*	5	*
0650	939	92.8 (62)	1.4 (100)	1,053	35.6 (28)	1,042	32.3 (35)	820	92.1 (60)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 88

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0651	462	90.5 (94)	1.7 (113)	532	38.0 (18)	489	28.2 (59)	423	88.7 (110)
0652	258	93.4 (51)	1.6 (109)	280	36.1 (26)	279	24.7 (83)	224	84.8 (129)
0653	314	88.2 (112)	3.8 (136)	374	38.0 (18)	367	32.4 (34)	293	89.4 (100)
0654	314	87.9 (115)	3.2 (134)	411	44.0 (4)	404	16.3 (120)	328	91.8 (69)
0655	283	93.6 (48)	0.7 (45)	309	40.5 (14)	287	35.5 (24)	260	87.7 (118)
0657	31	*	*	35	*	34	*	32	*
0701	370	92.4 (68)	0.8 (56)	461	33.2 (37)	455	40.2 (13)	352	92.9 (46)
0702	58	*	*	69	*	70	*	51	*
0703	144	94.4 (33)	1.4 (100)	163	30.1 (69)	156	44.2 (9)	109	98.2 (1)
0704	42	*	*	50	*	49	*	42	*
0705	56	*	*	65	*	63	*	42	*
0706	1	*	*	1	*	1	*	0	*
0850	619	91.4 (81)	1.3 (94)	676	25.9 (94)	661	31.5 (40)	472	93.6 (36)
0851	551	94.6 (29)	0.4 (19)	589	28.2 (80)	590	37.1 (20)	418	95.5 (13)
0901	1,183	89.9 (104)	2.0 (119)	1,402	38.0 (18)	1,390	26.1 (73)	1,091	91.8 (69)
0902	21	*	*	25	*	24	*	18	*
1051	917	91.3 (85)	2.4 (127)	1,088	29.8 (72)	1,086	29.2 (52)	784	89.9 (95)
1052	3	*	*	2	*	3	*	2	*
1101	1,106	81.5 (138)	1.7 (113)	1,367	27.9 (83)	1,200	22.7 (93)	967	89.6 (98)
1102	552	86.6 (122)	1.8 (115)	727	33.3 (36)	656	23.5 (88)	509	94.7 (24)
1106	49	*	*	56	*	53	*	37	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 89

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1250	838	89.5 (105)	2.0 (119)	942	29.1 (75)	906	37.0 (21)	686	92.7 (51)
1251	572	94.9 (23)	0.3 (17)	622	26.2 (92)	620	46.3 (6)	448	94.2 (27)
1252	362	80.9 (139)	5.5 (140)	508	33.7 (35)	503	23.1 (90)	376	89.1 (105)
1253	456	85.1 (129)	3.9 (138)	621	37.7 (21)	626	23.0 (92)	490	88.8 (109)
1350	2,120	96.2 (13)	0.5 (24)	2,304	32.6 (44)	2,166	32.6 (32)	1,732	92.7 (51)
1351	1	*	*	1	*	1	*	1	*
1352	318	99.4 (1)	0.0 (1)	349	34.4 (31)	333	6.6 (136)	253	92.9 (46)
1401	178	91.6 (76)	1.1 (78)	205	31.2 (62)	202	39.6 (15)	148	93.2 (42)
1402	73	*	*	84	*	78	*	63	*
1403	16	*	*	17	*	18	*	12	*
1501	538	90.0 (102)	1.9 (118)	735	32.9 (41)	696	21.6 (101)	541	85.6 (126)
1503	705	96.6 (9)	0.6 (32)	788	34.0 (33)	753	22.4 (95)	586	81.9 (131)
1504	1,627	96.4 (10)	0.4 (19)	1,810	27.7 (85)	1,756	21.5 (103)	1,235	89.3 (102)
1505	18	*	*	22	*	13	*	22	*
1506	405	93.1 (55)	1.5 (106)	525	30.9 (64)	513	12.1 (130)	391	92.1 (60)
1507	474	96.2 (13)	0.8 (56)	513	31.8 (54)	489	30.5 (46)	339	97.6 (2)
1509	308	91.9 (71)	0.6 (32)	365	31.5 (58)	346	20.2 (110)	275	94.2 (27)
1511	545	91.4 (81)	0.6 (32)	602	32.9 (41)	597	28.6 (56)	445	88.5 (112)
1512	331	95.2 (18)	0.9 (66)	365	21.6 (121)	357	31.1 (42)	212	92.0 (64)
1514	594	94.1 (38)	0.8 (56)	663	24.9 (102)	582	30.6 (44)	408	92.9 (46)
1701	458	96.7 (7)	0.7 (45)	502	34.1 (32)	492	55.7 (3)	390	94.1 (29)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 90

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1950	497	93.8 (43)	0.8 (56)	556	31.3 (61)	487	40.7 (12)	420	96.7 (5)
1952	115	93.9 (41)	0.0 (1)	128	28.1 (81)	111	47.7 (5)	86	*
1953	228	95.2 (18)	0.0 (1)	265	25.7 (97)	236	45.3 (8)	195	96.9 (4)
1954	311	94.5 (30)	0.6 (32)	350	38.3 (17)	296	49.0 (4)	290	95.5 (13)
1955	107	91.6 (76)	0.0 (1)	121	28.1 (81)	104	36.5 (22)	80	*
1957	333	92.8 (62)	0.6 (32)	374	31.6 (57)	296	43.6 (10)	271	94.8 (22)
1963	63	*	*	69	*	59	*	49	*
2150	510	94.3 (36)	0.2 (13)	542	31.7 (56)	535	28.2 (59)	398	95.0 (19)
2151	183	95.1 (21)	1.1 (78)	222	27.0 (87)	169	36.1 (23)	155	91.0 (74)
2152	32	*	*	38	*	37	*	31	*
2251	324	96.3 (12)	1.5 (106)	367	36.8 (23)	363	30.9 (43)	298	95.6 (12)
2252	136	94.9 (23)	2.2 (124)	165	46.7 (1)	155	68.4 (1)	138	91.3 (72)
2554	46	*	*	49	*	38	*	43	*
2555	6,059	90.1 (100)	0.7 (45)	6,954	25.5 (99)	5,800	19.8 (112)	5,081	92.5 (55)
2556	100	87.0 (120)	0.0 (1)	117	19.7 (129)	94	*	97	*
2557	1	*	*	1	*	1	*	1	*
2560	1	*	*	1	*	1	*	1	*
2562	3	*	*	3	*	3	*	3	*
2563	4	*	*	6	*	5	*	6	*
2564	2	*	*	3	*	2	*	3	*
3001	1,035	90.1 (100)	1.3 (94)	1,144	36.8 (23)	1,131	12.3 (128)	924	87.1 (122)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 91

Michigan

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3002	274	93.4 (51)	0.7 (45)	297	32.0 (51)	296	14.5 (125)	234	91.0 (74)
3004	274	94.9 (23)	0.0 (1)	289	33.2 (37)	293	23.9 (85)	213	92.0 (64)
3005	142	91.5 (78)	2.8 (132)	154	39.6 (15)	158	23.4 (89)	132	93.2 (42)
3007	7	*	*	7	*	7	*	6	*
3201	340	91.5 (78)	1.2 (89)	399	24.1 (106)	394	40.9 (11)	275	90.9 (80)
3301	3,568	90.4 (96)	1.1 (78)	4,240	27.5 (86)	4,044	12.6 (127)	3,122	90.0 (94)
3304	82	*	*	98	*	98	*	78	*
3305	127	90.6 (92)	0.8 (56)	156	31.4 (60)	151	14.6 (124)	112	92.0 (64)
3306	177	93.8 (43)	0.6 (32)	202	30.7 (66)	195	14.9 (123)	154	92.2 (59)
3308	69	*	*	93	*	89	*	55	*
3312	374	91.7 (74)	1.1 (78)	456	32.9 (41)	448	13.6 (126)	356	90.4 (86)
3314	1	*	*	1	*	1	*	1	*
3550	56	*	*	62	*	62	*	49	*
3551	5	*	*	6	*	6	*	5	*
3552	1	*	*	1	*	1	*	1	*
3701	752	94.8 (26)	0.7 (45)	820	26.6 (89)	814	22.5 (94)	575	95.8 (8)
3850	2,253	91.7 (74)	1.6 (109)	2,660	37.6 (22)	2,637	23.8 (86)	2,049	89.3 (102)
3901	1,203	92.1 (70)	0.4 (19)	1,442	33.1 (40)	1,412	16.4 (119)	1,081	85.9 (125)
3902	9	*	*	9	*	9	*	9	*
3903	2	*	*	2	*	2	*	1	*
3950	1,735	90.3 (97)	1.2 (89)	1,953	30.6 (67)	1,904	22.0 (98)	1,408	90.6 (83)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 92

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3954	1	*	*	1	*	1	*	1	*
4106	1,808	87.2 (119)	1.8 (115)	2,134	26.1 (93)	2,023	12.3 (128)	1,601	85.0 (127)
4107	2,427	91.8 (72)	1.1 (78)	2,731	23.7 (109)	2,580	29.5 (49)	1,964	92.6 (53)
4109	567	95.2 (18)	1.1 (78)	621	25.0 (100)	612	26.0 (75)	393	94.1 (29)
4112	553	63.5 (140)	0.0 (1)	574	22.1 (120)	117	46.2 (7)	510	90.4 (86)
4115	2,792	93.4 (51)	1.3 (94)	3,311	23.2 (112)	2,435	20.2 (110)	2,354	92.6 (53)
4351	409	89.2 (108)	1.2 (89)	515	43.7 (6)	478	21.3 (104)	404	87.6 (119)
4352	150	83.3 (134)	1.3 (94)	179	41.3 (12)	179	26.8 (68)	144	84.7 (130)
4353	331	86.4 (124)	4.2 (139)	459	36.6 (25)	443	28.0 (62)	344	88.4 (113)
4354	651	96.2 (13)	0.6 (32)	739	46.4 (2)	715	18.9 (115)	579	89.6 (98)
4355	223	87.4 (117)	3.6 (135)	255	31.8 (54)	257	58.8 (2)	200	92.0 (64)
4356	304	91.1 (86)	1.3 (94)	370	33.2 (37)	367	26.4 (71)	285	91.9 (68)
4359	395	85.1 (129)	3.8 (136)	563	44.0 (4)	567	22.0 (98)	458	87.3 (120)
4360	190	94.2 (37)	2.1 (122)	222	31.5 (58)	224	31.3 (41)	177	91.0 (74)
4361	218	93.1 (55)	1.8 (115)	237	43.0 (7)	236	29.2 (52)	189	92.1 (60)
4362	223	90.6 (92)	2.2 (124)	262	34.0 (33)	250	29.6 (48)	201	89.1 (105)
4363	548	90.7 (90)	1.6 (109)	624	31.1 (63)	620	27.9 (63)	460	91.5 (71)
4401	600	90.8 (89)	2.0 (119)	718	32.3 (47)	710	29.4 (50)	527	94.9 (20)
4501	648	92.6 (65)	1.4 (100)	735	32.1 (50)	733	25.5 (77)	545	90.8 (81)
4601	1,006	93.0 (57)	0.2 (13)	1,162	28.6 (77)	1,157	20.3 (109)	839	93.0 (45)
4701	852	91.8 (72)	2.2 (124)	973	29.9 (71)	965	37.2 (19)	692	93.6 (36)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 93

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5050	1,470	94.1 (38)	0.5 (24)	1,575	21.6 (121)	1,488	11.7 (131)	1,054	97.0 (3)
5051	2,260	92.9 (59)	0.8 (56)	2,496	22.8 (117)	2,284	10.4 (132)	1,684	95.2 (17)
5052	2,437	93.0 (57)	0.5 (24)	2,600	21.4 (123)	2,380	9.2 (134)	1,814	94.3 (25)
5055	1,169	98.4 (3)	1.2 (89)	1,326	41.7 (11)	512	6.1 (137)	1,067	95.3 (16)
5101	1,290	92.9 (59)	0.9 (66)	1,416	23.1 (113)	786	39.1 (17)	999	95.7 (9)
5102	3	*	*	5	*	2	*	5	*
5202	535	93.5 (50)	2.1 (122)	624	35.3 (29)	534	28.1 (61)	490	95.7 (9)
5206	7	*	*	10	*	9	*	9	*
5601	859	94.1 (38)	1.4 (100)	946	26.7 (88)	941	24.8 (81)	666	94.1 (29)
5801	1,528	92.6 (65)	0.8 (56)	1,739	27.9 (83)	1,532	27.3 (65)	1,228	93.3 (40)
5803	20	*	*	21	*	19	*	18	*
5804	9	*	*	10	*	10	*	10	*
5806	53	*	*	56	*	47	*	32	*
6101	52	*	*	63	*	62	*	47	*
6102	57	*	*	65	*	62	*	48	*
6103	2,790	90.9 (88)	1.0 (70)	3,177	35.0 (30)	2,955	23.7 (87)	2,419	92.9 (46)
6105	4	*	*	6	*	6	*	4	*
6301	2,896	93.8 (43)	0.8 (56)	3,321	25.6 (98)	3,057	20.5 (107)	2,326	91.0 (74)
6312	1,531	93.6 (48)	0.5 (24)	1,728	25.9 (94)	1,589	22.2 (96)	1,188	90.6 (83)
6316	986	94.7 (27)	1.0 (70)	1,092	24.8 (103)	1,000	27.3 (65)	752	92.4 (57)
6318	7	*	*	9	*	9	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 94

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6320	486	94.7 (27)	1.0 (70)	548	23.0 (116)	491	32.8 (30)	402	95.5 (13)
6325	285	97.2 (5)	0.4 (19)	312	23.1 (113)	309	32.7 (31)	207	93.7 (35)
6335	833	96.4 (10)	0.6 (32)	905	29.5 (73)	727	29.4 (50)	657	94.8 (22)
6345	124	94.4 (33)	0.0 (1)	141	24.1 (106)	132	32.6 (32)	95	*
6360	776	93.7 (47)	0.8 (56)	881	25.0 (100)	795	25.7 (76)	647	94.1 (29)
7306	1,660	90.7 (90)	0.7 (45)	1,849	32.2 (48)	1,811	28.3 (57)	1,377	92.1 (60)
7401	1,269	88.7 (110)	2.6 (129)	1,746	42.4 (8)	1,730	26.1 (73)	1,393	89.4 (100)
7601	452	98.5 (2)	0.2 (13)	496	32.5 (45)	489	32.1 (38)	375	94.1 (29)
7850	926	96.8 (6)	0.6 (32)	1,019	32.0 (51)	266	34.6 (27)	729	85.0 (127)
7852	6	*	*	8	*	3	*	7	*
8101	1,521	83.2 (136)	1.4 (100)	1,663	19.7 (129)	1,413	21.9 (100)	1,102	87.9 (116)
8102	392	89.5 (105)	0.8 (56)	432	22.2 (119)	377	21.0 (106)	329	87.8 (117)
8105	0	*	*	1	*	0	*	1	*
8106	494	86.0 (125)	1.0 (70)	536	15.3 (141)	529	22.1 (97)	355	96.1 (7)
8150	6	*	*	7	*	7	*	6	*
8151	2,744	93.3 (54)	1.0 (70)	3,148	29.4 (74)	3,110	18.2 (116)	2,355	92.4 (57)
8152	6	*	*	7	*	6	*	6	*
8153	0	*	*	1	*	1	*	1	*
8155	6	*	*	7	*	7	*	7	*
8201	2,214	94.4 (33)	0.5 (24)	2,547	19.4 (131)	1,948	35.0 (25)	1,737	88.9 (107)
8202	3,566	98.0 (4)	0.2 (13)	3,829	30.8 (65)	3,448	5.1 (139)	3,030	95.7 (9)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 95

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy ⁽⁴⁾		During Pregnancy ⁽⁵⁾	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8203	304	91.4 (81)	0.7 (45)	355	23.4 (111)	307	32.2 (37)	249	90.4 (86)
8250	401	90.3 (97)	0.5 (24)	461	17.8 (135)	376	27.1 (67)	323	88.9 (107)
8252	112	86.6 (122)	2.7 (131)	130	23.1 (113)	125	24.8 (81)	100	94.0 (34)
8255	1,659	92.9 (59)	0.6 (32)	1,860	17.0 (138)	1,602	32.3 (35)	1,177	90.4 (86)
8257	1,518	96.1 (16)	0.0 (1)	1,596	24.4 (104)	1,160	5.9 (138)	1,242	72.7 (132)
8263	305	95.7 (17)	1.0 (70)	388	26.5 (90)	332	33.1 (29)	374	92.5 (55)
8264	712	94.5 (30)	0.6 (32)	778	16.2 (140)	668	34.9 (26)	495	88.3 (114)
8267	638	91.5 (78)	1.6 (109)	692	14.3 (143)	595	29.2 (52)	449	89.3 (102)
8901	735	84.2 (132)	1.4 (100)	778	19.4 (131)	530	20.4 (108)	506	86.8 (124)
8902	1,134	89.2 (108)	0.4 (19)	1,214	21.3 (124)	961	24.1 (84)	852	89.9 (95)
8903	1,201	83.3 (134)	0.7 (45)	1,317	20.5 (125)	1,029	21.1 (105)	928	87.2 (121)
8904	1,732	83.1 (137)	1.1 (78)	1,887	19.8 (128)	1,546	21.6 (101)	1,258	90.3 (92)
8905	1,698	92.6 (65)	0.7 (45)	1,855	24.0 (108)	1,614	34.4 (28)	1,230	90.4 (86)
8909	1,019	88.0 (114)	0.6 (32)	1,143	20.3 (126)	921	23.1 (90)	796	90.3 (92)
9045	1,178	91.1 (86)	1.1 (78)	1,291	24.2 (105)	566	15.7 (121)	962	94.9 (20)
9050	730	85.5 (126)	0.5 (24)	871	25.9 (94)	805	25.5 (77)	633	88.2 (115)
9051	1,503	90.5 (94)	0.3 (17)	1,601	22.7 (118)	1,331	3.7 (140)	1,232	94.3 (25)
9053	1,497	87.4 (117)	1.1 (78)	1,563	15.0 (142)	1,277	19.0 (114)	880	93.5 (39)
9055	6	*	*	7	*	7	*	3	*
9058	1,284	86.7 (121)	1.3 (94)	1,379	16.9 (139)	1,274	9.9 (133)	753	88.7 (110)
9066	613	84.8 (131)	1.5 (106)	685	28.5 (79)	611	18.0 (118)	641	90.8 (81)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 6A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 96

Michigan

Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9068	738	83.7 (133)	2.6 (129)	803	18.9 (133)	773	15.3 (122)	542	87.1 (122)
9069	1,558	85.2 (127)	1.0 (70)	1,632	17.7 (136)	1,152	8.6 (135)	1,016	93.3 (40)
9070	1,704	85.2 (127)	1.1 (78)	1,863	17.4 (137)	1,659	18.2 (116)	1,185	91.1 (73)
9072	46	*	*	59	*	59	*	59	*
9080	2,367	88.1 (113)	1.2 (89)	2,581	20.1 (127)	1,852	26.3 (72)	1,749	89.8 (97)
9082	102	90.2 (99)	1.0 (70)	107	10.3 (144)	84	*	60	*
9083	12	*	*	12	*	12	*	11	*
9087	10	*	*	11	*	11	*	11	*
Michigan	123,165	91.4	1.0	139,456	27.5	123,767	23.1	100,835	91.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 97

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	85	*	104	18.3 (125)	24	*	85	*	112	7.1 (116)
0102	274	35.0 (121)	331	19.3 (130)	92	*	273	11.7 (99)	347	6.9 (112)
0103	281	34.5 (119)	342	16.4 (116)	92	*	281	21.0 (130)	351	10.0 (131)
0104	101	32.7 (115)	113	16.8 (119)	28	*	99	*	116	4.3 (72)
0105	4	*	3	*	1	*	4	*	4	*
0250	85	*	106	17.9 (121)	31	*	89	*	106	13.2 (138)
0251	58	*	65	*	22	*	58	*	69	*
0252	78	*	90	*	16	*	75	*	92	*
0253	102	47.1 (132)	111	25.2 (138)	45	*	102	15.7 (120)	115	13.0 (137)
0401	285	37.2 (125)	375	18.9 (128)	96	*	285	11.9 (104)	386	4.1 (70)
0402	282	31.9 (110)	307	15.0 (112)	81	*	281	11.7 (99)	316	7.0 (113)
0403	94	*	107	23.4 (136)	34	*	93	*	111	17.1 (142)
0404	87	*	97	*	16	*	86	*	100	5.0 (85)
0501	293	30.4 (105)	373	16.1 (115)	80	*	291	18.6 (127)	378	7.4 (120)
0503	40	*	54	*	8	*	40	*	55	*
0504	243	25.1 (84)	301	13.6 (101)	59	*	238	9.7 (80)	301	4.0 (69)
0505	147	23.8 (80)	183	13.1 (97)	32	*	145	11.7 (99)	185	5.4 (90)
0506	127	43.3 (130)	160	25.6 (139)	48	*	122	16.4 (122)	165	15.2 (141)
0507	44	*	69	*	7	*	42	*	70	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 98

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0508	5	*	13	*	1	*	5	*	13	*
0650	820	22.4 (74)	995	12.6 (92)	179	31.8 (48)	821	9.3 (75)	1,005	5.5 (92)
0651	428	37.6 (127)	485	23.5 (137)	141	24.1 (54)	422	18.5 (126)	493	8.1 (125)
0652	224	29.9 (101)	275	16.7 (118)	64	*	224	20.1 (129)	278	13.3 (139)
0653	296	35.5 (122)	340	18.2 (124)	97	*	293	18.8 (128)	358	11.2 (133)
0654	333	27.0 (94)	379	20.6 (133)	83	*	328	17.7 (124)	389	11.8 (135)
0655	260	37.3 (126)	286	23.1 (135)	91	*	260	17.7 (124)	299	10.0 (131)
0657	32	*	34	*	4	*	32	*	34	*
0701	355	29.6 (99)	419	17.9 (121)	97	*	352	6.8 (53)	422	3.6 (58)
0702	53	*	64	*	14	*	51	*	64	*
0703	108	40.7 (128)	144	22.9 (134)	36	*	109	24.8 (132)	146	9.6 (130)
0704	42	*	42	*	10	*	42	*	43	*
0705	43	*	58	*	16	*	42	*	58	*
0706	0	*	1	*	0	*	0	*	1	*
0850	485	25.4 (85)	636	11.0 (84)	103	43.7 (31)	472	9.1 (73)	636	5.0 (85)
0851	422	23.0 (76)	565	10.4 (79)	94	*	416	11.1 (92)	568	4.4 (75)
0901	1,106	29.4 (96)	1,325	13.7 (103)	304	36.8 (40)	1,091	11.5 (96)	1,338	5.8 (96)
0902	18	*	22	*	7	*	18	*	22	*
1051	809	32.0 (112)	997	13.0 (95)	229	43.2 (32)	784	9.4 (79)	1,027	6.2 (104)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 99

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1052	2	*	3	*	1	*	2	*	3	*
1101	966	18.3 (54)	1,244	7.1 (55)	161	48.4 (18)	966	12.2 (106)	1,246	6.7 (108)
1102	516	30.4 (105)	640	15.6 (114)	140	42.9 (33)	510	15.9 (121)	647	7.1 (116)
1106	38	*	50	*	12	*	37	*	51	*
1250	699	18.7 (56)	886	11.3 (86)	111	25.2 (51)	684	10.1 (85)	900	5.6 (95)
1251	447	30.4 (105)	574	18.5 (126)	124	23.4 (56)	448	11.6 (97)	589	7.6 (121)
1252	378	19.3 (61)	461	9.3 (76)	63	*	376	7.4 (60)	466	5.8 (96)
1253	489	25.8 (87)	574	13.1 (97)	112	35.7 (44)	488	10.2 (86)	579	5.9 (99)
1350	1,754	22.3 (73)	2,152	7.5 (57)	348	44.0 (28)	1,730	4.2 (20)	2,185	3.1 (43)
1351	1	*	1	*	0	*	1	*	1	*
1352	253	20.6 (70)	317	3.8 (27)	48	*	252	3.2 (13)	323	0.9 (5)
1401	150	36.0 (123)	186	12.4 (91)	51	*	149	6.7 (50)	192	6.3 (106)
1402	63	*	77	*	11	*	63	*	78	*
1403	12	*	17	*	4	*	12	*	17	*
1501	545	21.1 (71)	648	6.2 (48)	109	44.0 (28)	541	5.9 (40)	671	4.8 (83)
1503	591	20.0 (67)	740	8.1 (64)	115	33.0 (47)	586	5.5 (36)	743	1.5 (10)
1504	1,255	12.6 (29)	1,681	4.4 (36)	146	47.3 (19)	1,235	2.6 (6)	1,690	1.5 (10)
1505	22	*	19	*	6	*	22	*	19	*
1506	395	19.0 (59)	475	8.0 (62)	67	*	391	4.3 (23)	494	4.7 (80)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 100

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1507	344	32.8 (116)	466	8.4 (67)	104	47.1 (21)	336	10.7 (90)	487	6.2 (104)
1509	277	18.8 (58)	329	9.7 (77)	49	*	275	5.5 (36)	331	3.9 (65)
1511	446	26.7 (91)	563	12.3 (90)	111	37.8 (38)	446	11.7 (99)	576	9.2 (129)
1512	217	15.2 (37)	353	8.5 (69)	34	*	212	8.5 (69)	354	3.4 (50)
1514	428	16.4 (45)	641	5.8 (45)	64	*	404	3.0 (9)	642	1.4 (8)
1701	389	24.4 (83)	467	19.9 (131)	87	*	388	11.1 (92)	471	4.5 (78)
1950	419	33.2 (117)	513	13.5 (100)	132	44.7 (26)	420	14.3 (116)	535	6.7 (108)
1952	88	*	123	8.9 (73)	16	*	86	*	126	7.1 (116)
1953	197	29.4 (96)	239	8.4 (67)	53	*	195	7.2 (57)	247	6.1 (102)
1954	289	32.5 (113)	318	11.9 (89)	93	*	290	11.7 (99)	335	7.8 (122)
1955	82	*	106	13.2 (99)	30	*	80	*	113	7.1 (116)
1957	273	30.0 (102)	353	13.0 (95)	74	*	271	6.6 (48)	359	3.9 (65)
1963	49	*	65	*	14	*	48	*	67	*
2150	408	32.6 (114)	511	20.4 (132)	120	27.5 (50)	396	11.6 (97)	514	6.0 (100)
2151	159	25.8 (87)	194	14.9 (111)	36	*	155	11.0 (91)	196	6.1 (102)
2152	31	*	32	*	13	*	31	*	32	*
2251	302	29.5 (98)	336	14.6 (109)	76	*	297	3.0 (9)	341	2.1 (23)
2252	135	41.5 (129)	152	26.3 (140)	53	*	138	13.0 (112)	154	5.8 (96)
2554	41	*	44	*	13	*	42	*	44	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 101

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2555	5,163	19.8 (66)	6,402	8.7 (71)	896	42.9 (33)	5,083	9.2 (74)	6,494	4.3 (72)
2556	97	*	111	12.6 (92)	25	*	97	*	113	1.8 (14)
2557	1	*	1	*	0	*	1	*	1	*
2560	1	*	1	*	0	*	1	*	1	*
2562	3	*	3	*	0	*	3	*	3	*
2563	6	*	5	*	2	*	6	*	5	*
2564	3	*	2	*	0	*	3	*	2	*
3001	929	23.3 (77)	1,088	8.5 (69)	206	49.0 (16)	924	10.3 (89)	1,097	6.0 (100)
3002	233	19.3 (61)	288	8.0 (62)	42	*	233	17.2 (123)	287	7.0 (113)
3004	219	19.2 (60)	289	5.5 (42)	41	*	213	14.6 (117)	289	3.8 (63)
3005	131	26.0 (89)	152	6.6 (52)	34	*	132	11.4 (95)	152	6.6 (107)
3007	5	*	7	*	2	*	6	*	7	*
3201	283	30.0 (102)	387	15.0 (112)	79	*	275	12.0 (105)	387	4.7 (80)
3301	3,180	17.6 (49)	4,094	6.9 (54)	527	47.2 (20)	3,118	6.1 (45)	4,100	2.7 (35)
3304	79	*	96	*	9	*	78	*	96	*
3305	114	16.7 (46)	148	6.1 (47)	16	*	112	3.6 (15)	149	2.7 (35)
3306	153	19.6 (64)	199	10.1 (78)	30	*	154	3.9 (16)	199	2.5 (32)
3308	60	*	83	*	7	*	55	*	83	*
3312	363	17.1 (48)	437	8.7 (71)	58	*	355	5.6 (39)	439	3.2 (46)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 102

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3314	1	*	1	*	0	*	1	*	1	*
3550	49	*	61	*	28	*	49	*	61	*
3551	5	*	5	*	4	*	5	*	5	*
3552	1	*	1	*	1	*	1	*	1	*
3701	589	27.7 (95)	755	14.6 (109)	147	23.8 (55)	574	12.2 (106)	782	3.8 (63)
3850	2,081	24.2 (82)	2,514	14.0 (105)	460	30.9 (49)	2,049	9.3 (75)	2,510	4.6 (79)
3901	1,095	19.5 (63)	1,313	7.9 (60)	185	36.2 (43)	1,079	5.9 (40)	1,324	2.7 (35)
3902	8	*	8	*	4	*	9	*	9	*
3903	1	*	2	*	0	*	1	*	2	*
3950	1,421	19.6 (64)	1,832	9.2 (75)	259	39.0 (36)	1,406	6.5 (47)	1,855	4.4 (75)
3954	1	*	1	*	1	*	1	*	1	*
4106	1,605	10.4 (17)	2,037	5.9 (46)	147	35.4 (45)	1,594	3.9 (16)	2,039	1.9 (16)
4107	1,999	16.3 (44)	2,533	6.3 (49)	304	49.7 (13)	1,953	5.3 (34)	2,553	2.6 (34)
4109	392	17.6 (49)	605	10.6 (81)	69	*	392	4.1 (18)	607	3.5 (54)
4112	517	4.6 (7)	552	1.6 (6)	18	*	460	8.9 (72)	538	3.5 (54)
4115	2,401	7.8 (11)	3,162	3.4 (20)	193	45.1 (25)	2,343	3.1 (12)	3,175	1.1 (6)
4351	408	31.4 (109)	473	17.8 (120)	115	25.2 (51)	402	10.2 (86)	477	8.0 (124)
4352	139	47.5 (133)	163	29.4 (141)	63	*	143	21.7 (131)	168	13.7 (140)
4353	345	20.3 (68)	427	11.7 (87)	63	*	344	6.4 (46)	427	3.0 (42)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 103

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4354	586	30.2 (104)	654	14.1 (107)	165	20.0 (57)	579	12.8 (110)	708	5.5 (92)
4355	205	33.2 (117)	236	18.6 (127)	58	*	200	13.0 (112)	245	8.6 (127)
4356	285	31.9 (110)	349	14.0 (105)	84	*	284	13.7 (115)	351	11.7 (134)
4359	462	14.5 (33)	535	7.9 (60)	63	*	459	2.8 (8)	532	1.9 (16)
4360	177	20.3 (68)	214	13.6 (101)	33	*	177	11.3 (94)	216	8.3 (126)
4361	192	29.7 (100)	226	18.1 (123)	53	*	189	14.8 (118)	232	8.6 (127)
4362	205	37.1 (124)	236	19.1 (129)	64	*	201	12.9 (111)	248	12.9 (136)
4363	469	22.4 (74)	606	12.9 (94)	94	*	459	12.2 (106)	610	7.9 (123)
4401	534	23.4 (78)	679	14.3 (108)	117	42.7 (35)	527	6.6 (48)	685	2.0 (22)
4501	554	31.2 (108)	718	13.9 (104)	169	49.1 (15)	545	15.2 (119)	721	6.8 (111)
4601	857	23.7 (79)	1,051	10.6 (81)	171	43.9 (30)	839	8.7 (70)	1,053	5.1 (87)
4701	704	24.0 (81)	905	8.2 (65)	153	49.0 (16)	692	7.1 (56)	927	4.1 (70)
5050	1,074	12.4 (27)	1,510	5.5 (42)	120	38.3 (37)	1,053	5.0 (31)	1,513	2.8 (38)
5051	1,728	10.8 (18)	2,369	4.4 (36)	157	45.2 (23)	1,680	4.9 (30)	2,363	2.9 (40)
5052	1,841	9.3 (14)	2,533	3.4 (20)	155	45.2 (23)	1,811	4.1 (18)	2,526	2.1 (23)
5055	1,081	1.9 (2)	1,297	0.8 (3)	18	*	1,067	2.5 (5)	1,297	0.2 (1)
5101	1,013	11.2 (20)	1,346	3.8 (27)	108	60.2 (8)	994	4.8 (28)	1,354	2.1 (23)
5102	5	*	3	*	0	*	5	*	3	*
5202	492	43.5 (131)	565	16.6 (117)	203	37.4 (39)	488	8.4 (67)	603	3.6 (58)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 104

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
5206	10	*	11	*	3	*	9	*	11	*
5601	677	26.9 (93)	888	11.0 (84)	165	34.5 (46)	665	6.0 (44)	916	3.1 (43)
5801	1,239	26.8 (92)	1,588	10.8 (83)	315	44.1 (27)	1,217	9.9 (82)	1,596	5.1 (87)
5803	18	*	21	*	1	*	18	*	21	*
5804	10	*	10	*	3	*	10	*	10	*
5806	31	*	53	*	10	*	32	*	53	*
6101	49	*	60	*	11	*	47	*	61	*
6102	49	*	63	*	11	*	48	*	64	*
6103	2,453	21.6 (72)	2,985	10.4 (79)	483	36.4 (42)	2,422	9.3 (75)	3,030	4.8 (83)
6105	4	*	5	*	0	*	4	*	5	*
6301	2,369	8.0 (12)	3,054	3.5 (24)	180	36.7 (41)	2,319	3.5 (14)	3,069	1.8 (14)
6312	1,217	2.9 (4)	1,633	1.3 (5)	35	*	1,185	2.7 (7)	1,636	1.7 (13)
6316	769	7.7 (10)	1,038	3.8 (27)	57	*	753	2.3 (4)	1,041	1.4 (8)
6318	1	*	9	*	0	*	1	*	8	*
6320	413	11.4 (22)	502	2.0 (8)	38	*	401	2.2 (3)	503	2.2 (28)
6325	211	25.6 (86)	286	9.1 (74)	50	*	207	6.8 (53)	292	5.5 (92)
6335	675	10.2 (16)	866	2.5 (11)	54	*	657	5.2 (33)	868	3.1 (43)
6345	97	*	125	4.0 (31)	26	*	95	*	130	5.4 (90)
6360	659	17.0 (47)	820	5.2 (39)	99	*	643	7.2 (57)	827	3.3 (47)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 105

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7306	1,388	26.0 (89)	1,739	8.3 (66)	342	51.5 (12)	1,375	9.9 (82)	1,787	3.9 (65)
7401	1,404	34.5 (119)	1,572	11.8 (88)	421	46.8 (22)	1,393	13.4 (114)	1,603	6.7 (108)
7601	379	12.7 (30)	456	7.7 (59)	45	*	374	6.7 (50)	463	3.9 (65)
7850	740	14.9 (34)	955	7.5 (57)	103	24.3 (53)	724	4.3 (23)	957	3.3 (47)
7852	7	*	6	*	1	*	7	*	5	*
8101	1,119	15.3 (39)	1,590	3.0 (18)	158	68.4 (4)	1,100	9.3 (75)	1,597	3.4 (50)
8102	337	11.3 (21)	411	3.4 (20)	34	*	327	5.5 (36)	413	3.4 (50)
8105	1	*	0	*	0	*	1	*	0	*
8106	365	18.6 (55)	521	6.5 (50)	55	*	355	5.9 (40)	519	2.3 (30)
8150	6	*	7	*	2	*	6	*	7	*
8151	2,399	15.1 (35)	2,884	5.2 (39)	299	54.2 (10)	2,355	4.4 (26)	2,894	2.1 (23)
8152	6	*	7	*	1	*	6	*	7	*
8153	1	*	0	*	0	*	1	*	0	*
8155	7	*	6	*	1	*	7	*	6	*
8201	1,791	13.5 (32)	2,279	5.3 (41)	204	58.8 (9)	1,734	4.2 (20)	2,282	3.3 (47)
8202	3,074	2.5 (3)	3,794	0.7 (2)	70	*	3,029	3.0 (9)	3,799	0.7 (2)
8203	251	18.7 (56)	312	6.7 (53)	36	*	249	10.0 (84)	314	7.0 (113)
8250	329	15.2 (37)	405	4.0 (31)	38	*	323	4.3 (23)	406	3.7 (61)
8252	106	15.1 (35)	118	4.2 (34)	13	*	100	5.0 (31)	119	0.8 (4)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 106

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators ^(2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes ⁽⁴⁾		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8255	1,206	18.2 (52)	1,691	7.2 (56)	186	49.5 (14)	1,178	6.9 (55)	1,699	4.4 (75)
8257	1,276	0.3 (1)	1,555	0.3 (1)	5	*	1,244	0.8 (1)	1,548	0.7 (2)
8263	379	17.9 (51)	321	6.5 (50)	56	*	373	5.9 (40)	321	3.7 (61)
8264	504	11.1 (19)	721	2.4 (10)	49	*	495	4.4 (26)	723	2.2 (28)
8267	460	15.9 (41)	641	5.1 (38)	59	*	448	6.7 (50)	650	5.2 (89)
8901	519	12.3 (26)	761	4.2 (34)	59	*	508	8.3 (66)	761	3.5 (54)
8902	866	11.7 (25)	1,167	2.5 (11)	94	*	851	5.4 (35)	1,169	2.3 (30)
8903	933	16.0 (42)	1,259	4.0 (31)	130	68.5 (3)	921	7.6 (62)	1,259	2.5 (32)
8904	1,283	16.0 (42)	1,821	2.8 (15)	177	71.8 (2)	1,256	8.4 (67)	1,824	3.6 (58)
8905	1,256	8.4 (13)	1,790	2.8 (15)	101	62.4 (6)	1,225	4.8 (28)	1,791	3.5 (54)
8909	811	10.1 (15)	1,073	3.4 (20)	78	*	795	8.2 (65)	1,073	3.4 (50)
9045	978	4.1 (5)	1,233	0.9 (4)	35	*	956	4.2 (20)	1,233	1.5 (10)
9050	636	15.6 (40)	824	3.8 (27)	95	*	635	9.8 (81)	825	1.9 (16)
9051	1,239	4.3 (6)	1,567	2.1 (9)	56	*	1,233	1.1 (2)	1,567	1.3 (7)
9053	905	12.5 (28)	1,525	1.8 (7)	113	77.0 (1)	881	7.6 (62)	1,523	2.1 (23)
9055	3	*	6	*	0	*	3	*	6	*
9058	767	7.3 (9)	1,341	3.1 (19)	54	*	752	7.4 (60)	1,336	1.9 (16)
9066	659	18.2 (52)	661	5.7 (44)	109	53.2 (11)	641	12.2 (106)	661	2.9 (40)
9068	548	11.5 (23)	770	2.5 (11)	55	*	542	7.2 (57)	771	1.9 (16)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 7A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 107

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9069	1,045	6.4 (8)	1,577	3.5 (24)	65	*	1,015	7.8 (64)	1,580	4.7 (80)
9070	1,203	11.6 (24)	1,771	2.5 (11)	128	64.8 (5)	1,182	10.2 (86)	1,772	2.8 (38)
9072	59	*	47	*	14	*	59	*	47	*
9080	1,757	13.0 (31)	2,487	3.7 (26)	213	61.0 (7)	1,748	8.7 (70)	2,491	4.3 (72)
9082	63	*	106	2.8 (15)	6	*	60	*	106	1.9 (16)
9083	12	*	12	*	0	*	11	*	12	*
9087	11	*	10	*	1	*	11	*	10	*
Michigan	102,357	17.7	131,072	7.4	16,505	43.2	100,645	7.4	132,152	3.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾ Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 108

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	104	5.8 (6)	8.7 (120)	107	13.1 (84)	90	*	104	71.2 (105)
0102	317	11.4 (94)	10.7 (136)	325	11.7 (56)	274	5.1 (82)	315	73.7 (89)
0103	326	7.1 (17)	9.5 (127)	335	8.1 (3)	299	5.7 (98)	326	77.0 (78)
0104	105	8.6 (40)	12.4 (138)	110	18.2 (138)	86	*	105	80.0 (60)
0105	4	*	*	4	*	4	*	4	*
0250	97	*	*	96	*	85	*	96	*
0251	63	*	*	66	*	55	*	63	*
0252	85	*	*	89	*	80	*	85	*
0253	106	8.5 (39)	8.5 (117)	114	16.7 (128)	89	*	106	72.6 (97)
0401	370	10.3 (72)	7.3 (86)	375	12.5 (75)	313	5.1 (82)	370	75.9 (80)
0402	304	9.5 (60)	8.9 (122)	305	10.2 (25)	269	4.5 (64)	304	83.9 (27)
0403	103	10.7 (82)	8.7 (120)	105	9.5 (15)	91	*	105	86.7 (13)
0404	94	*	*	95	*	78	*	94	*
0501	320	6.6 (12)	6.9 (74)	361	7.8 (2)	291	3.1 (30)	351	84.6 (21)
0503	52	*	*	55	*	51	*	53	*
0504	284	9.9 (64)	8.1 (108)	293	11.9 (62)	249	2.4 (13)	285	89.5 (3)
0505	146	10.3 (72)	8.9 (122)	180	11.1 (41)	131	4.6 (68)	172	86.0 (16)
0506	141	9.2 (53)	9.2 (124)	153	8.5 (6)	125	5.6 (96)	148	81.8 (37)
0507	57	*	*	67	*	49	*	65	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 109

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	12	*	*	13	*	11	*	13	*
0650	942	9.3 (54)	8.2 (110)	978	9.5 (15)	852	5.0 (78)	946	81.0 (48)
0651	443	10.6 (80)	7.7 (99)	481	12.3 (72)	387	6.2 (109)	443	72.9 (94)
0652	263	6.8 (15)	6.5 (64)	267	8.2 (4)	237	2.5 (15)	261	76.6 (79)
0653	324	7.7 (25)	6.5 (64)	335	11.6 (52)	286	3.5 (40)	324	81.5 (41)
0654	370	10.5 (76)	5.9 (47)	375	11.5 (48)	324	5.9 (102)	369	80.5 (53)
0655	275	8.0 (29)	7.6 (95)	283	8.8 (9)	248	5.2 (85)	275	79.3 (71)
0657	29	*	*	33	*	23	*	29	*
0701	393	5.3 (4)	8.4 (116)	412	10.4 (30)	352	2.3 (10)	396	90.2 (2)
0702	60	*	*	61	*	54	*	60	*
0703	136	5.9 (7)	5.9 (47)	142	10.6 (32)	121	1.7 (5)	138	72.5 (98)
0704	41	*	*	42	*	33	*	41	*
0705	55	*	*	58	*	49	*	55	*
0706	1	*	*	1	*	1	*	1	*
0850	604	12.6 (111)	7.3 (86)	627	14.5 (106)	515	5.0 (78)	611	83.5 (30)
0851	518	9.3 (54)	6.9 (74)	554	14.3 (105)	446	2.7 (18)	525	83.8 (28)
0901	1,234	10.5 (76)	8.3 (112)	1,298	11.6 (52)	1,090	4.2 (59)	1,236	72.2 (99)
0902	23	*	*	23	*	20	*	23	*
1051	958	8.8 (44)	9.9 (130)	989	11.2 (44)	843	4.0 (53)	967	90.3 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 110

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1052	3	*	*	2	*	1	*	3	*
1101	1,169	12.1 (102)	5.7 (39)	1,206	13.1 (84)	1,004	6.6 (112)	1,173	69.4 (111)
1102	602	7.3 (20)	7.8 (103)	629	9.9 (22)	535	3.7 (44)	603	77.1 (77)
1106	49	*	*	51	*	42	*	49	*
1250	843	9.0 (48)	8.3 (112)	883	9.9 (22)	749	5.9 (102)	852	79.9 (62)
1251	549	10.7 (82)	5.8 (42)	564	15.4 (120)	456	5.0 (78)	548	80.3 (55)
1252	439	8.0 (29)	10.5 (134)	453	11.9 (62)	388	3.1 (30)	442	81.9 (36)
1253	542	6.3 (11)	9.8 (129)	560	11.4 (46)	470	1.5 (1)	546	84.2 (25)
1350	2,047	9.9 (64)	6.4 (61)	2,112	10.5 (31)	1,814	5.3 (88)	2,057	80.0 (60)
1351	1	*	*	1	*	1	*	1	*
1352	307	9.1 (50)	8.1 (108)	315	11.7 (56)	269	1.9 (8)	309	71.5 (102)
1401	183	6.6 (12)	9.3 (126)	186	7.5 (1)	168	4.2 (59)	183	89.1 (7)
1402	69	*	*	74	*	64	*	70	*
1403	16	*	*	16	*	14	*	16	*
1501	603	10.0 (67)	6.1 (53)	645	11.6 (52)	530	4.2 (59)	622	80.2 (56)
1503	691	7.2 (19)	6.9 (74)	707	10.7 (34)	616	2.9 (25)	693	79.8 (63)
1504	1,480	10.5 (76)	7.0 (79)	1,639	13.3 (88)	1,276	3.9 (50)	1,603	81.7 (38)
1505	19	*	*	19	*	17	*	19	*
1506	456	9.4 (57)	7.9 (104)	479	11.1 (41)	402	4.5 (64)	463	75.2 (83)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 111

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1507	430	9.8 (63)	10.0 (132)	464	12.9 (81)	372	2.7 (18)	448	85.7 (17)
1509	301	8.0 (29)	5.6 (37)	313	12.5 (75)	262	2.7 (18)	302	72.8 (96)
1511	541	9.6 (61)	7.2 (85)	557	13.3 (88)	468	3.8 (48)	548	79.4 (69)
1512	338	11.5 (95)	7.7 (99)	345	16.2 (126)	282	3.5 (40)	339	84.4 (23)
1514	592	14.2 (120)	7.6 (95)	627	16.3 (127)	493	3.9 (50)	605	86.4 (15)
1701	435	6.2 (9)	10.6 (135)	456	8.8 (9)	393	2.3 (10)	436	81.2 (46)
1950	504	10.1 (68)	5.2 (30)	517	13.3 (88)	430	2.8 (23)	505	83.2 (31)
1952	120	6.7 (14)	5.8 (42)	122	10.7 (34)	105	1.9 (8)	120	89.2 (5)
1953	233	13.3 (116)	6.0 (51)	244	13.5 (93)	200	6.0 (105)	236	88.1 (10)
1954	310	8.7 (42)	5.5 (35)	320	9.7 (19)	278	2.9 (25)	309	84.5 (22)
1955	106	5.7 (5)	8.5 (117)	110	14.5 (106)	90	*	107	72.9 (94)
1957	340	15.6 (129)	7.4 (91)	347	17.9 (133)	277	4.7 (71)	342	88.0 (11)
1963	59	*	*	59	*	51	*	60	*
2150	472	6.1 (8)	7.6 (95)	495	8.9 (11)	426	1.6 (4)	472	80.1 (59)
2151	190	7.4 (21)	6.3 (59)	193	9.3 (14)	171	4.7 (71)	189	68.8 (114)
2152	30	*	*	32	*	24	*	30	*
2251	330	4.5 (1)	5.8 (42)	332	10.2 (25)	293	1.7 (5)	329	81.5 (41)
2252	141	10.6 (80)	8.5 (117)	148	15.5 (123)	121	1.7 (5)	138	84.1 (26)
2554	45	*	*	45	*	40	*	45	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 112

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2555	6,215	12.9 (113)	5.7 (39)	6,386	13.7 (98)	5,329	6.3 (110)	6,228	65.4 (120)
2556	109	13.8 (118)	3.7 (12)	112	17.9 (133)	90	*	111	45.9 (137)
2557	1	*	*	1	*	1	*	1	*
2560	1	*	*	1	*	1	*	1	*
2562	3	*	*	3	*	2	*	3	*
2563	5	*	*	5	*	5	*	5	*
2564	2	*	*	2	*	2	*	2	*
3001	1,018	15.1 (125)	4.1 (16)	1,056	15.2 (116)	865	6.1 (108)	1,016	61.7 (125)
3002	270	10.7 (82)	4.4 (21)	275	15.3 (117)	229	2.6 (17)	270	69.3 (112)
3004	259	8.9 (46)	7.3 (86)	271	15.1 (112)	217	3.2 (33)	259	77.6 (75)
3005	145	8.3 (35)	12.4 (138)	147	8.2 (4)	133	3.0 (27)	147	79.6 (66)
3007	7	*	*	7	*	5	*	7	*
3201	365	7.4 (21)	9.9 (130)	375	10.7 (34)	325	1.5 (1)	364	80.2 (56)
3301	3,809	11.2 (91)	6.2 (55)	3,968	13.5 (93)	3,289	4.0 (53)	3,847	81.7 (38)
3304	92	*	*	96	*	83	*	92	*
3305	136	7.4 (21)	10.3 (133)	144	11.8 (59)	120	3.3 (34)	137	88.3 (9)
3306	187	7.5 (24)	5.9 (47)	190	12.1 (66)	163	2.5 (15)	186	85.5 (18)
3308	77	*	*	80	*	70	*	77	*
3312	409	10.3 (72)	4.6 (25)	422	12.1 (66)	360	5.6 (96)	411	83.0 (32)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 113

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3314	1	*	*	1	*	1	*	1	*
3550	52	*	*	55	*	43	*	52	*
3551	3	*	*	5	*	3	*	4	*
3552	1	*	*	1	*	1	*	1	*
3701	737	8.4 (38)	7.3 (86)	762	11.2 (44)	650	3.4 (37)	739	87.1 (12)
3850	2,335	11.5 (95)	7.5 (93)	2,435	12.6 (77)	2,038	5.2 (85)	2,344	77.6 (75)
3901	1,223	12.2 (107)	4.3 (20)	1,277	13.2 (87)	1,050	5.3 (88)	1,234	74.4 (86)
3902	5	*	*	8	*	5	*	5	*
3903	2	*	*	2	*	2	*	2	*
3950	1,726	10.9 (87)	8.3 (112)	1,801	12.7 (79)	1,503	4.9 (77)	1,738	79.7 (64)
3954	1	*	*	1	*	1	*	1	*
4106	1,907	13.4 (117)	4.6 (25)	1,984	15.3 (117)	1,607	6.7 (113)	1,927	78.3 (73)
4107	2,394	12.9 (113)	5.5 (35)	2,512	15.1 (112)	2,035	5.7 (98)	2,431	79.5 (68)
4109	569	12.3 (110)	6.9 (74)	582	17.9 (133)	464	3.9 (50)	568	84.7 (20)
4112	529	4.5 (1)	4.2 (18)	529	9.8 (21)	469	2.3 (10)	529	37.4 (140)
4115	2,966	13.1 (115)	6.2 (55)	3,111	14.1 (103)	2,549	5.5 (92)	3,031	81.7 (38)
4351	449	10.9 (87)	8.0 (105)	465	14.2 (104)	384	5.7 (98)	451	86.5 (14)
4352	152	11.8 (100)	6.6 (69)	159	13.8 (100)	125	7.2 (118)	152	79.6 (66)
4353	406	10.1 (68)	7.1 (82)	405	12.1 (66)	353	5.4 (91)	409	80.2 (56)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 114

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4354	664	9.6 (61)	7.7 (99)	679	11.5 (48)	581	4.3 (62)	666	82.0 (35)
4355	219	10.5 (76)	8.2 (110)	233	15.0 (111)	186	3.8 (48)	219	81.3 (45)
4356	328	7.0 (16)	7.6 (95)	334	10.2 (25)	290	3.1 (30)	331	80.4 (54)
4359	490	10.4 (75)	6.7 (71)	499	12.4 (73)	420	4.8 (75)	490	81.0 (48)
4360	194	6.2 (9)	5.7 (39)	208	9.1 (12)	174	3.4 (37)	195	89.2 (5)
4361	203	9.4 (57)	5.4 (33)	217	12.9 (81)	176	2.8 (23)	203	82.8 (33)
4362	232	10.8 (85)	7.3 (86)	242	11.6 (52)	203	3.0 (27)	231	88.7 (8)
4363	575	10.1 (68)	9.6 (128)	583	13.0 (83)	490	3.5 (40)	577	89.4 (4)
4401	639	8.6 (40)	8.0 (105)	662	10.9 (37)	565	4.4 (63)	641	80.8 (50)
4501	676	5.2 (3)	8.0 (105)	702	11.1 (41)	604	1.5 (1)	677	81.2 (46)
4601	992	9.0 (48)	6.4 (61)	1,036	11.4 (46)	879	3.6 (43)	991	79.4 (69)
4701	882	11.5 (95)	6.5 (64)	900	13.9 (101)	756	4.5 (64)	881	83.7 (29)
5050	1,466	12.1 (102)	4.4 (21)	1,470	15.4 (120)	1,218	6.0 (105)	1,464	70.9 (107)
5051	2,232	12.1 (102)	6.0 (51)	2,283	15.6 (124)	1,853	4.6 (68)	2,244	73.6 (92)
5052	2,403	14.8 (124)	4.5 (24)	2,450	14.7 (110)	2,030	8.0 (122)	2,412	70.4 (108)
5055	1,191	10.9 (87)	5.3 (32)	1,230	10.2 (25)	1,069	6.0 (105)	1,213	69.3 (112)
5101	1,313	8.8 (44)	5.2 (30)	1,313	12.2 (70)	1,138	3.0 (27)	1,320	77.7 (74)
5102	3	*	*	3	*	2	*	3	*
5202	576	8.0 (29)	6.9 (74)	580	10.0 (24)	512	4.1 (56)	579	80.7 (51)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 115

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5206	11	*	*	10	*	9	*	11	*
5601	857	8.3 (35)	9.2 (124)	890	10.3 (29)	769	3.3 (34)	859	79.7 (64)
5801	1,532	8.7 (42)	7.0 (79)	1,572	10.9 (37)	1,364	3.7 (44)	1,531	71.3 (104)
5803	20	*	*	20	*	19	*	20	*
5804	10	*	*	10	*	10	*	10	*
5806	51	*	*	51	*	41	*	51	*
6101	57	*	*	60	*	52	*	58	*
6102	63	*	*	64	*	54	*	63	*
6103	2,836	11.7 (98)	6.5 (64)	2,929	13.7 (98)	2,419	5.3 (88)	2,846	70.1 (109)
6105	5	*	*	5	*	4	*	4	*
6301	2,854	12.2 (107)	6.3 (59)	2,992	13.4 (92)	2,458	5.5 (92)	2,900	74.0 (88)
6312	1,515	12.1 (102)	5.1 (29)	1,583	14.5 (106)	1,287	4.1 (56)	1,543	81.4 (43)
6316	960	10.1 (68)	5.6 (37)	1,005	12.1 (66)	840	4.0 (53)	973	81.4 (43)
6318	8	*	*	8	*	5	*	8	*
6320	479	11.3 (93)	6.7 (71)	493	9.7 (19)	430	7.0 (116)	481	82.7 (34)
6325	285	7.7 (25)	11.6 (137)	286	9.1 (12)	255	2.7 (18)	285	80.7 (51)
6335	830	10.8 (85)	6.7 (71)	847	11.8 (59)	724	5.1 (82)	839	75.4 (82)
6345	127	9.4 (57)	7.1 (82)	129	12.4 (73)	110	2.7 (18)	128	84.4 (23)
6360	771	15.6 (129)	4.9 (27)	809	13.1 (84)	660	9.2 (129)	784	75.1 (84)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 116

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7306	1,690	11.9 (101)	6.1 (53)	1,736	13.6 (96)	1,459	4.7 (71)	1,692	71.8 (101)
7401	1,498	9.3 (54)	7.7 (99)	1,542	11.8 (59)	1,321	3.7 (44)	1,501	73.7 (89)
7601	429	9.1 (50)	5.8 (42)	454	11.7 (56)	373	4.6 (68)	441	66.0 (119)
7850	911	7.7 (25)	6.4 (61)	947	10.6 (32)	812	3.4 (37)	915	73.6 (92)
7852	4	*	*	4	*	4	*	4	*
8101	1,457	15.2 (127)	2.4 (2)	1,531	18.0 (136)	1,172	7.9 (121)	1,478	54.9 (133)
8102	380	7.1 (17)	7.4 (91)	398	9.5 (15)	336	4.8 (75)	387	66.9 (117)
8105	0	*	*	0	*	0	*	0	*
8106	442	14.3 (121)	5.9 (47)	507	15.4 (120)	367	6.3 (110)	496	65.1 (123)
8150	6	*	*	7	*	6	*	7	*
8151	2,675	12.2 (107)	6.2 (55)	2,796	13.6 (96)	2,299	5.9 (102)	2,718	85.1 (19)
8152	6	*	*	7	*	6	*	6	*
8153	0	*	*	0	*	0	*	0	*
8155	5	*	*	6	*	5	*	5	*
8201	2,158	12.1 (102)	5.4 (33)	2,251	13.9 (101)	1,862	5.5 (92)	2,173	71.0 (106)
8202	3,373	8.0 (29)	4.1 (16)	3,611	9.5 (15)	3,024	3.7 (44)	3,566	67.5 (115)
8203	303	9.9 (64)	8.3 (112)	309	12.0 (65)	266	4.5 (64)	303	74.6 (85)
8250	384	9.1 (50)	4.2 (18)	401	12.7 (79)	335	2.4 (13)	386	65.3 (121)
8252	98	*	*	115	8.7 (8)	88	*	112	75.9 (80)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance ⁽¹⁾

Page : 117

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators ^(2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8255	1,618	11.7 (98)	6.2 (55)	1,673	12.6 (77)	1,400	5.2 (85)	1,626	70.0 (110)
8257	1,398	14.7 (123)	2.9 (5)	1,509	10.9 (37)	1,233	10.6 (132)	1,441	65.2 (122)
8263	305	8.2 (34)	6.6 (69)	317	11.0 (40)	272	3.3 (34)	305	71.5 (102)
8264	687	11.2 (91)	7.1 (82)	705	12.2 (70)	601	4.7 (71)	690	78.4 (72)
8267	598	7.7 (25)	6.5 (64)	632	8.5 (6)	542	4.1 (56)	610	66.9 (117)
8901	706	16.6 (134)	4.4 (21)	740	17.8 (131)	572	9.1 (127)	709	57.5 (131)
8902	1,092	16.8 (135)	3.9 (14)	1,125	18.7 (140)	874	8.4 (126)	1,103	67.5 (115)
8903	1,164	14.3 (121)	2.8 (4)	1,217	15.9 (125)	967	7.1 (117)	1,158	54.1 (135)
8904	1,694	16.9 (136)	3.0 (7)	1,764	18.4 (139)	1,348	7.6 (119)	1,707	58.2 (130)
8905	1,621	11.0 (90)	5.8 (42)	1,727	13.5 (93)	1,385	5.5 (92)	1,672	72.2 (99)
8909	986	17.3 (137)	3.5 (10)	1,038	18.1 (137)	798	9.1 (127)	997	63.8 (124)
9045	1,171	8.3 (35)	7.5 (93)	1,211	11.5 (48)	1,032	5.0 (78)	1,188	41.2 (139)
9050	782	12.8 (112)	4.9 (27)	809	13.3 (88)	673	6.7 (113)	786	59.7 (127)
9051	1,464	8.9 (46)	7.0 (79)	1,498	11.9 (62)	1,273	5.7 (98)	1,503	73.7 (89)
9053	1,416	16.2 (132)	3.2 (8)	1,487	14.6 (109)	1,195	9.2 (129)	1,423	59.9 (126)
9055	6	*	*	6	*	6	*	5	*
9058	1,224	15.8 (131)	2.9 (5)	1,302	15.1 (112)	1,028	8.0 (122)	1,265	58.7 (128)
9066	620	17.6 (139)	3.9 (14)	648	16.8 (129)	520	8.1 (124)	620	54.4 (134)
9068	735	17.3 (137)	2.6 (3)	754	15.3 (117)	620	9.4 (131)	735	47.9 (136)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Run Date: 6/28/2024

Table 8A

2021 - 2023 Pregnancy Nutrition Surveillance (1)

Page : 118

Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9069	1,485	15.4 (128)	3.6 (11)	1,539	17.8 (131)	1,221	7.8 (120)	1,484	41.8 (138)
9070	1,670	16.4 (133)	3.7 (12)	1,735	17.0 (130)	1,377	8.1 (124)	1,670	56.3 (132)
9072	43	*	*	47	*	36	*	43	*
9080	2,345	15.1 (125)	3.3 (9)	2,440	15.1 (112)	1,975	6.9 (115)	2,397	58.7 (128)
9082	100	14.0 (119)	1.0 (1)	104	11.5 (48)	88	*	101	74.3 (87)
9083	12	*	*	11	*	9	*	12	*
9087	9	*	*	10	*	7	*	9	*
Michigan	123,246	11.5	6.0	128,228	13.2	106,221	5.2	124,658	49.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.