

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2024 Pregnancy Nutrition Surveillance (1)

Page : 1

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
0101	34	*	*	*	*	*	*	*
0102	120	88.3	1.7	5.8	0.0	0.0	4.2	0.0
0103	122	95.1	0.0	1.6	0.0	0.0	3.3	0.0
0104	28	*	*	*	*	*	*	*
0250	43	*	*	*	*	*	*	*
0251	22	*	*	*	*	*	*	*
0252	29	*	*	*	*	*	*	*
0253	29	*	*	*	*	*	*	*
0401	122	96.7	0.0	0.8	0.0	0.0	2.5	0.0
0402	105	77.1	1.0	9.5	0.0	1.9	10.5	0.0
0403	42	*	*	*	*	*	*	*
0404	42	*	*	*	*	*	*	*
0501	139	90.6	1.4	4.3	0.0	0.0	3.6	0.0
0503	15	*	*	*	*	*	*	*
0504	111	85.6	2.7	5.4	1.8	0.9	3.6	0.0
0505	75	*	*	*	*	*	*	*
0506	58	*	*	*	*	*	*	*
0507	20	*	*	*	*	*	*	*
0508	2	*	*	*	*	*	*	*
0650	348	78.7	3.2	9.2	3.4	0.3	3.7	1.4
0651	156	94.2	0.0	3.8	0.0	0.0	1.9	0.0
0652	86	*	*	*	*	*	*	*
0653	136	90.4	0.7	3.7	0.0	0.7	4.4	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2024 Pregnancy Nutrition Surveillance (1)

Page : 2

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
0654	96	*	*	*	*	*	*	*
0655	101	91.1	0.0	5.0	0.0	0.0	4.0	0.0
0657	10	*	*	*	*	*	*	*
0701	134	85.8	5.2	2.2	0.0	3.0	3.7	0.0
0702	18	*	*	*	*	*	*	*
0703	66	*	*	*	*	*	*	*
0704	19	*	*	*	*	*	*	*
0705	31	*	*	*	*	*	*	*
0850	200	84.5	6.0	6.5	0.0	1.0	2.0	0.0
0851	217	90.3	0.9	5.5	0.5	0.9	1.8	0.0
0901	425	80.5	2.8	9.4	0.0	0.7	6.4	0.2
0902	8	*	*	*	*	*	*	*
1051	339	80.2	2.1	10.9	1.5	1.8	3.2	0.3
1052	1	*	*	*	*	*	*	*
1101	386	37.3	42.5	12.4	0.3	2.1	4.9	0.5
1102	240	72.1	10.8	6.7	0.0	1.3	8.3	0.8
1106	10	*	*	*	*	*	*	*
1250	350	76.6	8.3	13.7	0.3	0.0	1.1	0.0
1251	248	94.0	0.4	2.8	0.0	0.0	2.4	0.4
1252	173	55.5	1.2	38.7	0.0	0.6	2.9	1.2
1253	188	75.5	0.5	14.9	0.0	0.0	9.0	0.0
1350	760	55.3	19.3	13.0	0.4	4.2	6.3	1.4
1352	127	71.7	22.8	3.1	0.0	0.0	2.4	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2024 Pregnancy Nutrition Surveillance (1)

Page : 3

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
1401	69	*	*	*	*	*	*	*
1402	28	*	*	*	*	*	*	*
1403	5	*	*	*	*	*	*	*
1501	246	45.5	4.1	45.9	0.0	0.0	3.3	1.2
1503	272	56.3	6.3	27.2	0.0	0.7	8.1	1.5
1504	624	33.3	3.4	55.3	0.2	3.7	2.9	1.3
1506	171	81.9	0.6	12.3	0.0	0.0	4.7	0.6
1507	165	77.0	2.4	17.0	0.0	0.6	2.4	0.6
1509	114	43.9	1.8	48.2	0.9	0.9	2.6	1.8
1511	178	90.4	0.6	3.4	0.0	1.1	4.5	0.0
1512	122	80.3	3.3	9.0	0.8	2.5	2.5	1.6
1514	242	74.0	5.8	16.1	0.0	0.8	2.1	1.2
1701	186	59.1	1.1	1.1	23.7	0.5	14.0	0.5
1950	196	90.3	1.0	4.6	0.0	1.5	2.6	0.0
1952	39	*	*	*	*	*	*	*
1953	80	*	*	*	*	*	*	*
1954	115	86.1	0.0	7.8	0.9	0.0	4.3	0.9
1955	48	*	*	*	*	*	*	*
1957	128	78.1	0.8	14.8	0.0	2.3	3.9	0.0
1963	35	*	*	*	*	*	*	*
2150	166	82.5	0.6	4.2	2.4	1.2	9.0	0.0
2151	75	*	*	*	*	*	*	*
2152	7	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2024 Pregnancy Nutrition Surveillance (1)

Page : 4

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
2251	99	*	*	*	*	*	*	*
2252	53	*	*	*	*	*	*	*
2554	9	*	*	*	*	*	*	*
2555	2,186	48.8	37.6	6.5	0.0	0.4	6.4	0.3
2556	106	29.2	48.1	13.2	0.0	1.9	7.5	0.0
3001	348	31.9	44.0	17.5	0.6	0.6	5.2	0.3
3002	184	39.1	34.8	17.4	0.0	1.1	7.6	0.0
3004	91	*	*	*	*	*	*	*
3005	53	*	*	*	*	*	*	*
3201	145	92.4	0.0	6.2	0.0	0.0	0.7	0.7
3301	1,540	38.0	30.7	19.5	0.1	5.3	5.8	0.6
3304	11	*	*	*	*	*	*	*
3305	26	*	*	*	*	*	*	*
3306	34	*	*	*	*	*	*	*
3308	13	*	*	*	*	*	*	*
3312	91	*	*	*	*	*	*	*
3314	1	*	*	*	*	*	*	*
3550	15	*	*	*	*	*	*	*
3551	4	*	*	*	*	*	*	*
3701	274	85.4	0.4	9.5	0.0	0.0	4.0	0.7
3850	766	69.6	12.8	8.5	0.4	0.5	7.3	0.8
3901	492	34.8	51.4	6.5	0.0	0.8	5.9	0.6
3950	603	49.8	21.7	17.7	0.0	2.5	8.1	0.2

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2024 Pregnancy Nutrition Surveillance (1)

Page : 5

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
4106	910	13.3	29.8	52.5	0.0	0.7	3.6	0.1
4107	733	36.0	26.9	28.2	0.5	1.5	6.7	0.1
4109	280	77.9	2.9	13.2	0.0	1.4	4.6	0.0
4112	149	0.7	0.0	98.7	0.0	0.0	0.7	0.0
4115	1,176	27.0	35.1	24.1	0.2	8.8	4.7	0.2
4351	157	91.7	0.0	5.1	0.0	0.6	2.5	0.0
4352	64	*	*	*	*	*	*	*
4353	138	51.4	0.7	43.5	0.0	0.0	1.4	2.9
4354	223	83.0	4.0	7.2	0.9	0.4	4.0	0.4
4355	90	*	*	*	*	*	*	*
4356	105	78.1	1.0	18.1	1.0	0.0	1.9	0.0
4359	160	66.3	0.6	30.0	0.0	0.0	2.5	0.6
4360	57	*	*	*	*	*	*	*
4361	76	*	*	*	*	*	*	*
4362	75	*	*	*	*	*	*	*
4363	219	87.2	0.9	6.8	0.0	0.0	4.1	0.9
4401	254	84.3	0.8	7.5	0.4	0.8	6.3	0.0
4501	212	91.5	0.0	6.6	0.0	0.0	0.0	1.9
4601	348	71.6	1.7	21.0	0.0	0.3	3.4	2.0
4701	335	84.2	0.6	9.0	0.3	2.4	2.7	0.9
5050	459	47.1	41.4	4.1	0.4	2.2	4.6	0.2
5051	853	49.9	34.7	6.2	0.2	3.4	4.9	0.6
5052	948	26.2	39.2	2.8	0.0	27.4	3.7	0.6

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2024 Pregnancy Nutrition Surveillance (1)

Page : 6

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
5055	419	69.9	14.1	1.4	0.0	14.1	0.5	0.0
5101	420	61.2	10.0	21.9	0.0	3.6	3.3	0.0
5202	203	86.7	2.0	3.9	1.0	0.5	5.4	0.5
5601	287	88.5	2.1	4.9	0.0	0.7	3.8	0.0
5801	541	77.8	6.7	10.5	0.0	0.2	4.1	0.7
5803	30	*	*	*	*	*	*	*
5806	19	*	*	*	*	*	*	*
6101	12	*	*	*	*	*	*	*
6102	12	*	*	*	*	*	*	*
6103	999	57.3	23.6	11.8	0.2	0.5	6.3	0.3
6105	3	*	*	*	*	*	*	*
6301	1,101	31.3	28.6	33.3	0.2	2.5	3.8	0.2
6312	631	30.1	57.2	4.9	0.0	3.3	4.4	0.0
6316	436	55.0	23.6	14.0	0.0	2.8	4.4	0.2
6318	1	*	*	*	*	*	*	*
6320	75	*	*	*	*	*	*	*
6325	95	*	*	*	*	*	*	*
6335	276	53.3	27.5	5.8	0.0	11.2	1.8	0.4
6345	29	*	*	*	*	*	*	*
6360	250	24.0	41.6	30.0	0.4	0.8	2.4	0.8
7306	554	44.9	30.3	17.7	0.4	1.1	5.2	0.4
7401	570	76.7	6.0	8.6	0.0	0.5	7.9	0.4
7601	155	89.7	0.0	7.7	0.0	0.0	2.6	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2024 Pregnancy Nutrition Surveillance (1)

Page : 7

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
7850	367	90.2	0.3	6.3	0.0	0.3	2.7	0.3
8101	519	5.0	89.2	2.3	0.0	0.6	2.9	0.0
8102	125	16.0	18.4	60.0	0.0	3.2	2.4	0.0
8106	184	40.2	39.7	13.6	0.0	1.1	4.3	1.1
8150	2	*	*	*	*	*	*	*
8151	943	35.8	35.1	17.4	0.1	3.6	7.1	0.8
8201	677	40.8	40.6	6.5	0.1	5.3	6.1	0.6
8202	1,185	94.6	3.5	0.8	0.0	0.4	0.4	0.3
8203	122	59.8	27.9	3.3	0.0	0.0	9.0	0.0
8250	90	*	*	*	*	*	*	*
8252	21	*	*	*	*	*	*	*
8255	788	52.5	22.7	15.0	0.4	4.2	4.2	1.0
8257	492	55.7	5.5	0.8	0.0	37.0	1.0	0.0
8264	285	58.9	26.0	7.7	0.0	4.6	2.5	0.4
8267	160	52.5	14.4	27.5	0.0	0.6	5.0	0.0
8901	192	12.5	77.1	4.2	0.0	1.6	4.2	0.5
8902	327	7.0	89.0	1.2	0.0	0.6	2.1	0.0
8903	376	4.0	88.0	2.4	0.0	2.9	2.7	0.0
8904	534	4.7	89.0	2.8	0.2	1.5	1.7	0.2
8905	608	64.3	23.8	7.7	0.0	0.8	2.3	1.0
8909	323	4.6	89.5	3.1	0.0	0.0	2.8	0.0
9045	384	7.8	16.9	74.0	0.0	0.3	0.5	0.5
9050	280	18.9	58.2	20.7	0.0	0.0	2.1	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2024 Pregnancy Nutrition Surveillance (1)

Page : 8

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
9051	767	5.3	2.5	91.1	0.0	0.0	0.5	0.5
9053	444	9.9	77.3	3.2	0.0	7.9	1.4	0.5
9058	607	18.6	68.9	1.5	0.0	8.6	2.0	0.5
9068	279	5.0	93.5	0.7	0.0	0.0	0.7	0.0
9069	426	2.6	95.5	0.7	0.0	0.0	1.2	0.0
9070	570	3.3	92.3	1.8	0.2	1.2	1.2	0.0
9080	1,029	22.4	69.0	4.1	0.0	0.1	3.5	0.9
9082	44	*	*	*	*	*	*	*
9083	3	*	*	*	*	*	*	*
<b>Michigan</b>	<b>44,869</b>	<b>50.2</b>	<b>27.3</b>	<b>14.6</b>	<b>0.4</b>	<b>3.0</b>	<b>4.2</b>	<b>0.5</b>
<b>National (Prior Year)</b>	<b>55,377</b>	<b>48.7</b>	<b>25.3</b>	<b>16.3</b>	<b>1.8</b>	<b>3.1</b>	<b>4.4</b>	<b>0.4</b>

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**WIC PNSS & PedNSS**  
**Table 4A**  
**2024 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Run Date: 4/14/2025

Page : 9

<b>Comparison of Age Distribution (2)</b>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
0101	34	*	*	*	*	*	*	*
0102	120	0.0	0.8	7.5	60.8	22.5	2.5	5.8
0103	122	0.0	3.3	8.2	62.3	23.8	0.8	1.6
0104	28	*	*	*	*	*	*	*
0250	43	*	*	*	*	*	*	*
0251	22	*	*	*	*	*	*	*
0252	29	*	*	*	*	*	*	*
0253	29	*	*	*	*	*	*	*
0401	122	0.0	0.0	9.8	63.9	24.6	1.6	0.0
0402	105	0.0	1.0	5.7	60.0	31.4	0.0	1.9
0403	42	*	*	*	*	*	*	*
0404	42	*	*	*	*	*	*	*
0501	139	0.0	0.0	1.4	76.3	17.3	0.7	4.3
0503	15	*	*	*	*	*	*	*
0504	111	0.0	0.0	5.4	58.6	32.4	2.7	0.9
0505	75	*	*	*	*	*	*	*
0506	58	*	*	*	*	*	*	*
0507	20	*	*	*	*	*	*	*
0508	2	*	*	*	*	*	*	*
0650	348	0.0	2.0	6.3	66.7	21.3	1.4	2.3
0651	156	0.0	1.9	3.8	64.7	23.1	1.9	4.5
0652	86	*	*	*	*	*	*	*
0653	136	0.0	1.5	3.7	62.5	22.8	2.2	7.4
0654	96	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**WIC PNSS & PedNSS**  
**Table 4A**  
**2024 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Run Date: 4/14/2025

Page : 10

<b>Comparison of Age Distribution (2)</b>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
0655	101	0.0	0.0	1.0	68.3	23.8	2.0	5.0
0657	10	*	*	*	*	*	*	*
0701	134	0.0	0.7	6.7	50.0	40.3	0.7	1.5
0702	18	*	*	*	*	*	*	*
0703	66	*	*	*	*	*	*	*
0704	19	*	*	*	*	*	*	*
0705	31	*	*	*	*	*	*	*
0850	200	0.0	1.0	6.0	56.0	32.0	3.0	2.0
0851	217	0.0	1.8	5.1	64.1	24.4	1.8	2.8
0901	425	0.2	1.9	7.1	64.0	24.7	1.4	0.7
0902	8	*	*	*	*	*	*	*
1051	339	0.0	2.4	5.0	53.7	33.6	3.8	1.5
1052	1	*	*	*	*	*	*	*
1101	386	0.3	1.0	3.9	59.3	30.1	1.0	4.4
1102	240	0.0	1.3	6.7	63.3	24.2	2.1	2.5
1106	10	*	*	*	*	*	*	*
1250	350	0.0	1.4	4.6	59.7	27.7	4.9	1.7
1251	248	0.0	2.4	8.1	60.5	21.8	1.2	6.0
1252	173	0.0	2.9	4.0	60.7	22.0	2.3	8.1
1253	188	0.0	1.1	6.9	53.7	31.9	2.1	4.3
1350	760	0.0	1.8	4.3	58.9	29.9	2.2	2.8
1352	127	0.0	1.6	4.7	61.4	24.4	1.6	6.3
1401	69	*	*	*	*	*	*	*
1402	28	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**WIC PNSS & PedNSS**  
**Table 4A**  
**2024 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Run Date: 4/14/2025

Page : 11

<b>Comparison of Age Distribution (2)</b>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
1403	5	*	*	*	*	*	*	*
1501	246	0.0	3.7	6.5	54.1	30.9	2.4	2.4
1503	272	0.0	0.4	4.8	58.8	31.6	1.5	2.9
1504	624	0.3	0.8	6.7	58.2	31.1	2.7	0.2
1506	171	0.0	3.5	6.4	59.1	25.7	4.1	1.2
1507	165	0.0	0.0	4.2	61.2	29.7	0.6	4.2
1509	114	0.0	3.5	4.4	52.6	34.2	0.9	4.4
1511	178	0.0	0.6	7.9	58.4	28.7	3.4	1.1
1512	122	0.0	1.6	6.6	60.7	29.5	1.6	0.0
1514	242	0.0	0.8	3.7	55.0	38.8	1.7	0.0
1701	186	0.5	0.0	5.4	60.8	27.4	2.2	3.8
1950	196	0.0	2.0	6.6	64.8	23.5	1.5	1.5
1952	39	*	*	*	*	*	*	*
1953	80	*	*	*	*	*	*	*
1954	115	0.0	0.9	6.1	60.0	29.6	2.6	0.9
1955	48	*	*	*	*	*	*	*
1957	128	0.0	3.1	8.6	59.4	28.1	0.8	0.0
1963	35	*	*	*	*	*	*	*
2150	166	0.0	1.2	4.8	59.0	28.9	6.0	0.0
2151	75	*	*	*	*	*	*	*
2152	7	*	*	*	*	*	*	*
2251	99	*	*	*	*	*	*	*
2252	53	*	*	*	*	*	*	*
2554	9	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2024 Pregnancy Nutrition Surveillance (1)

Page : 12

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2555	2,186	0.1	2.5	6.3	59.2	28.9	1.7	1.3
2556	106	0.0	1.9	6.6	55.7	33.0	1.9	0.9
3001	348	0.0	0.9	2.0	63.2	25.3	1.4	7.2
3002	184	0.0	1.1	2.7	54.9	30.4	1.1	9.8
3004	91	*	*	*	*	*	*	*
3005	53	*	*	*	*	*	*	*
3201	145	0.0	0.0	6.9	61.4	28.3	1.4	2.1
3301	1,540	0.1	1.9	5.4	57.3	31.8	2.0	1.6
3304	11	*	*	*	*	*	*	*
3305	26	*	*	*	*	*	*	*
3306	34	*	*	*	*	*	*	*
3308	13	*	*	*	*	*	*	*
3312	91	*	*	*	*	*	*	*
3314	1	*	*	*	*	*	*	*
3550	15	*	*	*	*	*	*	*
3551	4	*	*	*	*	*	*	*
3701	274	0.0	3.3	9.9	57.7	26.3	1.1	1.8
3850	766	0.0	1.7	6.0	56.3	28.1	1.7	6.3
3901	492	0.0	2.4	6.5	55.5	29.3	2.8	3.5
3950	603	0.2	1.7	5.6	58.0	29.5	3.5	1.5
4106	910	0.1	3.3	6.0	53.3	30.9	3.2	3.2
4107	733	0.3	2.2	4.8	55.5	32.1	2.6	2.6
4109	280	0.0	1.4	6.1	54.6	34.3	3.2	0.4
4112	149	0.0	1.3	3.4	60.4	28.2	4.7	2.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2024 Pregnancy Nutrition Surveillance (1)

Page : 13

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4115	1,176	0.1	1.3	3.4	52.8	37.2	2.6	2.6
4351	157	0.0	0.6	4.5	60.5	28.7	1.3	4.5
4352	64	*	*	*	*	*	*	*
4353	138	0.0	2.2	6.5	51.4	28.3	0.7	10.9
4354	223	0.0	1.8	8.5	59.2	27.8	2.7	0.0
4355	90	*	*	*	*	*	*	*
4356	105	0.0	0.0	3.8	57.1	34.3	1.0	3.8
4359	160	0.0	2.5	3.1	56.3	33.1	2.5	2.5
4360	57	*	*	*	*	*	*	*
4361	76	*	*	*	*	*	*	*
4362	75	*	*	*	*	*	*	*
4363	219	0.0	0.9	6.4	53.4	32.9	3.2	3.2
4401	254	0.0	0.8	3.5	59.1	28.3	3.1	5.1
4501	212	0.0	0.5	5.2	57.5	28.8	2.8	5.2
4601	348	0.3	1.4	4.0	61.8	27.6	1.4	3.4
4701	335	0.0	1.5	6.3	56.4	33.4	1.8	0.6
5050	459	0.0	1.3	3.5	48.4	41.6	3.3	2.0
5051	853	0.0	0.9	3.3	55.0	36.9	2.9	0.9
5052	948	0.0	1.2	2.0	56.9	37.3	1.9	0.7
5055	419	0.0	0.2	2.4	41.1	51.8	4.3	0.2
5101	420	0.0	0.7	3.3	54.8	39.0	1.4	0.7
5202	203	0.5	1.0	4.9	58.6	32.0	2.0	1.0
5601	287	0.0	1.7	4.5	59.2	30.7	3.1	0.7
5801	541	0.0	1.3	5.2	61.4	28.5	1.5	2.2

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2024 Pregnancy Nutrition Surveillance (1)

Page : 14

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5803	30	*	*	*	*	*	*	*
5806	19	*	*	*	*	*	*	*
6101	12	*	*	*	*	*	*	*
6102	12	*	*	*	*	*	*	*
6103	999	0.1	0.8	3.4	61.0	27.6	2.2	4.9
6105	3	*	*	*	*	*	*	*
6301	1,101	0.1	1.6	4.6	58.7	32.2	2.3	0.5
6312	631	0.0	0.3	2.7	50.9	41.8	3.6	0.6
6316	436	0.0	0.7	2.3	55.0	38.3	3.0	0.7
6318	1	*	*	*	*	*	*	*
6320	75	*	*	*	*	*	*	*
6325	95	*	*	*	*	*	*	*
6335	276	0.4	0.7	3.3	48.2	42.4	4.0	1.1
6345	29	*	*	*	*	*	*	*
6360	250	0.4	0.4	3.2	61.2	29.2	1.2	4.4
7306	554	0.2	2.2	7.2	54.7	33.0	2.5	0.2
7401	570	0.2	1.4	5.3	60.9	26.7	1.9	3.7
7601	155	0.0	3.2	3.9	61.3	29.0	1.3	1.3
7850	367	0.0	1.9	3.8	60.2	31.3	1.6	1.1
8101	519	0.0	3.3	6.2	51.3	34.1	3.5	1.7
8102	125	0.0	2.4	5.6	56.8	31.2	2.4	1.6
8106	184	0.0	1.1	3.3	57.6	35.3	2.7	0.0
8150	2	*	*	*	*	*	*	*
8151	943	0.2	1.1	4.7	52.3	36.8	3.8	1.2

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2024 Pregnancy Nutrition Surveillance (1)

Page : 15

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8201	677	0.1	1.2	3.1	53.6	36.9	4.3	0.7
8202	1,185	0.0	0.4	2.3	51.6	39.7	5.3	0.6
8203	122	0.0	0.8	4.1	57.4	34.4	3.3	0.0
8250	90	*	*	*	*	*	*	*
8252	21	*	*	*	*	*	*	*
8255	788	0.1	1.4	3.7	57.7	34.5	1.9	0.6
8257	492	0.0	0.2	1.6	47.6	45.1	5.1	0.4
8264	285	0.0	0.7	3.5	55.1	35.8	4.2	0.7
8267	160	0.0	1.9	6.9	58.1	30.0	3.1	0.0
8901	192	0.5	2.6	7.8	51.0	34.9	0.5	2.6
8902	327	0.3	1.5	4.0	47.4	40.4	3.7	2.8
8903	376	0.0	2.4	3.5	51.1	34.6	2.9	5.6
8904	534	0.0	1.9	4.3	55.4	35.4	1.9	1.1
8905	608	0.0	0.3	1.8	53.0	41.8	3.0	0.2
8909	323	0.0	1.5	4.0	54.2	36.8	2.8	0.6
9045	384	0.3	2.1	6.8	53.1	33.1	2.9	1.8
9050	280	0.7	1.4	6.4	54.6	32.1	2.5	2.1
9051	767	0.1	1.2	5.1	58.5	30.2	3.4	1.4
9053	444	0.0	2.3	5.4	53.8	35.1	2.3	1.1
9058	607	0.0	2.5	3.6	51.6	37.7	3.0	1.6
9068	279	0.4	1.1	2.2	53.4	37.6	2.5	2.9
9069	426	0.0	2.3	5.6	48.8	38.7	3.3	1.2
9070	570	0.0	2.6	6.0	51.8	36.0	1.9	1.8
9080	1,029	0.1	1.6	4.8	51.0	38.4	2.2	1.9

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**WIC PNSS & PedNSS**  
**Table 4A**  
**2024 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Run Date: 4/14/2025

Page : 16

<b>Comparison of Age Distribution (2)</b>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
9082	44	*	*	*	*	*	*	*
9083	3	*	*	*	*	*	*	*
<b>Michigan</b>	<b>44,870</b>	<b>0.1</b>	<b>1.5</b>	<b>4.8</b>	<b>56.3</b>	<b>32.7</b>	<b>2.6</b>	<b>2.1</b>
<b>National (Prior Year)</b>	<b>55,378</b>	<b>0.1</b>	<b>1.6</b>	<b>5.5</b>	<b>56.6</b>	<b>32.4</b>	<b>2.6</b>	<b>1.2</b>

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2024 Pregnancy Nutrition Surveillance (1)

Page : 17

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	23	*	*	14	*	*	6	*	7	*	31	*	31	*
0102	73	*	*	11	*	*	17	*	26	*	113	5.3 (4)	113	11.5 (25)
0103	78	*	*	18	*	*	16	*	32	*	117	11.1 (74)	117	17.1 (73)
0104	14	*	*	3	*	*	5	*	5	*	25	*	25	*
0250	29	*	*	9	*	*	6	*	12	*	39	*	39	*
0251	15	*	*	2	*	*	3	*	8	*	21	*	21	*
0252	25	*	*	17	*	*	0	*	8	*	26	*	26	*
0253	28	*	*	21	*	*	4	*	10	*	27	*	27	*
0401	108	2.8 (10)	52.8 (5)	67	*	*	15	*	19	*	110	5.5 (8)	111	21.6 (95)
0402	85	*	*	3	*	*	15	*	30	*	97	*	97	*
0403	38	*	*	20	*	*	8	*	10	*	39	*	39	*
0404	23	*	*	3	*	*	7	*	8	*	38	*	38	*
0501	111	0.9 (2)	59.5 (30)	13	*	*	16	*	32	*	122	9.0 (45)	122	10.7 (17)
0503	9	*	*	1	*	*	2	*	6	*	14	*	14	*
0504	104	1.9 (4)	51.9 (4)	40	*	*	17	*	22	*	105	9.5 (53)	105	11.4 (23)
0505	62	*	*	7	*	*	15	*	18	*	71	*	71	*
0506	22	*	*	5	*	*	11	*	9	*	53	*	53	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2024 Pregnancy Nutrition Surveillance (1)

Page : 18

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0507	13	*	*	2	*	*	1	*	6	*	19	*	19	*
0508	2	*	*	2	*	*	0	*	2	*	2	*	2	*
0650	220	3.6 (29)	62.7 (67)	107	15.0 (20)	58.9 (47)	50	*	93	*	318	12.9 (92)	320	17.8 (78)
0651	116	5.2 (67)	61.2 (51)	23	*	*	20	*	35	*	145	11.7 (82)	145	13.1 (37)
0652	74	*	*	44	*	*	10	*	11	*	84	*	84	*
0653	127	7.1 (85)	64.6 (80)	62	*	*	18	*	22	*	131	9.2 (48)	131	18.3 (79)
0654	54	*	*	21	*	*	7	*	36	*	89	*	89	*
0655	96	*	*	12	*	*	7	*	16	*	94	*	95	*
0657	5	*	*	2	*	*	0	*	4	*	9	*	9	*
0701	70	*	*	28	*	*	16	*	44	*	123	11.4 (79)	123	11.4 (23)
0702	10	*	*	1	*	*	4	*	4	*	14	*	14	*
0703	43	*	*	12	*	*	5	*	21	*	58	*	58	*
0704	17	*	*	12	*	*	6	*	2	*	16	*	16	*
0705	10	*	*	4	*	*	7	*	12	*	28	*	28	*
0850	102	3.9 (32)	56.9 (14)	18	*	*	21	*	44	*	182	15.4 (102)	182	18.7 (83)
0851	120	8.3 (91)	54.2 (7)	52	*	*	31	*	63	*	208	7.2 (23)	208	21.6 (95)
0901	354	7.3 (89)	58.2 (20)	240	18.3 (44)	54.2 (21)	67	*	146	27.4 (18)	394	10.7 (70)	395	13.9 (50)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 19

### Michigan

### Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0902	6	*	*	3	*	*	0	*	3	*	8	*	8	*
1051	248	3.2 (18)	60.1 (38)	153	20.3 (52)	49.0 (7)	34	*	60	*	307	8.8 (44)	308	13.3 (39)
1052	0	*	*	0	*	*	0	*	0	*	1	*	1	*
1101	298	6.4 (78)	62.1 (61)	137	24.8 (58)	47.4 (6)	60	*	117	24.8 (14)	342	9.6 (56)	341	19.4 (89)
1102	196	7.1 (85)	56.6 (13)	87	*	*	39	*	87	*	214	7.5 (27)	215	10.2 (15)
1106	9	*	*	1	*	*	1	*	2	*	8	*	8	*
1250	153	4.6 (53)	62.1 (61)	36	*	*	72	*	72	*	327	7.0 (18)	327	7.6 (7)
1251	215	4.7 (58)	59.1 (27)	150	14.7 (18)	60.0 (48)	32	*	59	*	234	14.5 (100)	233	18.5 (81)
1252	133	4.5 (48)	61.7 (58)	53	*	*	24	*	36	*	156	13.5 (95)	156	10.9 (18)
1253	156	7.1 (85)	62.2 (63)	66	*	*	29	*	53	*	174	12.6 (89)	174	14.4 (51)
1350	677	3.0 (15)	63.4 (71)	483	16.8 (37)	56.3 (33)	108	29.6 (2)	103	22.3 (7)	693	8.2 (37)	693	13.4 (42)
1352	108	4.6 (53)	64.8 (82)	64	*	*	19	*	21	*	118	2.5 (1)	118	9.3 (11)
1401	62	*	*	38	*	*	9	*	15	*	66	*	66	*
1402	18	*	*	11	*	*	3	*	6	*	24	*	24	*
1403	4	*	*	3	*	*	0	*	4	*	4	*	4	*
1501	188	3.2 (18)	63.8 (76)	73	*	*	42	*	41	*	220	11.8 (83)	221	10.4 (16)
1503	233	3.4 (23)	66.1 (87)	152	15.1 (21)	50.0 (10)	42	*	48	*	239	12.1 (85)	239	11.3 (21)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2024 Pregnancy Nutrition Surveillance (1)

Page : 20

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1504	185	10.3 (92)	50.8 (3)	45	*	*	88	*	108	15.7 (3)	563	8.5 (42)	563	8.2 (8)
1506	146	7.5 (90)	61.0 (48)	14	*	*	29	*	44	*	156	12.8 (91)	156	12.2 (32)
1507	102	11.8 (94)	45.1 (1)	16	*	*	19	*	45	*	149	8.1 (36)	148	12.2 (32)
1509	70	*	*	20	*	*	13	*	22	*	105	12.4 (87)	105	6.7 (5)
1511	114	2.6 (9)	60.5 (42)	53	*	*	23	*	51	*	166	7.8 (31)	167	17.4 (75)
1512	73	*	*	43	*	*	18	*	44	*	110	5.5 (8)	110	15.5 (59)
1514	135	6.7 (80)	55.6 (11)	17	*	*	30	*	89	*	228	7.5 (27)	228	16.7 (69)
1701	153	3.3 (21)	62.7 (67)	83	*	*	13	*	48	*	163	10.4 (67)	163	16.6 (68)
1950	172	5.2 (67)	59.9 (36)	122	14.8 (19)	57.4 (38)	26	*	64	*	183	10.9 (72)	183	17.5 (76)
1952	31	*	*	22	*	*	1	*	12	*	38	*	38	*
1953	70	*	*	55	*	*	9	*	34	*	75	*	75	*
1954	103	2.9 (11)	64.1 (78)	70	*	*	22	*	37	*	107	15.0 (101)	107	11.2 (20)
1955	43	*	*	28	*	*	8	*	16	*	41	*	41	*
1957	113	5.3 (69)	64.6 (80)	77	*	*	17	*	37	*	121	6.6 (16)	121	13.2 (38)
1963	28	*	*	16	*	*	0	*	5	*	31	*	31	*
2150	159	3.1 (16)	61.0 (48)	109	18.3 (44)	55.0 (27)	28	*	38	*	153	9.2 (48)	154	21.4 (94)
2151	73	*	*	57	*	*	11	*	15	*	64	*	64	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 21

### Michigan

### Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2152	7	*	*	4	*	*	0	*	4	*	6	*	6	*
2251	88	*	*	67	*	*	7	*	16	*	91	*	91	*
2252	30	*	*	5	*	*	3	*	9	*	44	*	43	*
2554	2	*	*	2	*	*	2	*	2	*	9	*	9	*
2555	1,501	6.7 (80)	59.2 (29)	566	20.0 (50)	51.8 (14)	394	42.9 (16)	501	31.5 (28)	2,005	11.3 (76)	2,007	16.3 (64)
2556	93	*	*	84	*	*	3	*	32	*	106	12.3 (86)	106	17.0 (72)
3001	316	3.5 (24)	62.7 (67)	237	21.9 (56)	47.3 (5)	43	*	57	*	321	7.2 (23)	321	13.7 (49)
3002	173	3.5 (24)	70.5 (93)	126	17.5 (42)	51.6 (13)	31	*	36	*	177	5.6 (10)	178	4.5 (3)
3004	74	*	*	59	*	*	9	*	16	*	87	*	87	*
3005	47	*	*	37	*	*	11	*	9	*	52	*	52	*
3201	117	5.1 (63)	65.0 (83)	60	*	*	17	*	38	*	142	7.0 (18)	142	25.4 (101)
3301	1,251	4.3 (42)	58.8 (26)	730	15.5 (23)	56.0 (32)	273	32.6 (6)	271	22.5 (10)	1,459	7.1 (22)	1,459	11.5 (25)
3304	5	*	*	3	*	*	1	*	3	*	10	*	10	*
3305	20	*	*	9	*	*	1	*	3	*	23	*	23	*
3306	31	*	*	15	*	*	1	*	6	*	32	*	33	*
3308	12	*	*	5	*	*	2	*	2	*	12	*	12	*
3312	81	*	*	46	*	*	10	*	8	*	87	*	87	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2024 Pregnancy Nutrition Surveillance (1)

Page : 22

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3314	1	*	*	0	*	*	0	*	0	*	1	*	1	*
3550	8	*	*	4	*	*	0	*	1	*	15	*	15	*
3551	2	*	*	0	*	*	1	*	1	*	4	*	4	*
3701	237	7.2 (88)	59.9 (36)	177	16.4 (34)	61.6 (50)	34	*	66	*	257	7.8 (31)	257	19.1 (86)
3850	608	5.6 (74)	59.5 (30)	384	18.8 (48)	53.9 (18)	80	*	111	33.3 (31)	697	10.3 (64)	699	13.6 (45)
3901	388	3.6 (29)	63.4 (71)	182	9.3 (2)	62.1 (53)	65	*	118	23.7 (12)	440	6.4 (12)	440	13.6 (45)
3950	462	6.1 (75)	60.4 (41)	224	11.6 (6)	63.4 (56)	79	*	132	15.2 (2)	548	9.3 (50)	547	15.4 (57)
4106	317	4.4 (45)	59.6 (33)	89	*	*	144	29.9 (3)	289	23.2 (11)	873	9.0 (45)	874	21.7 (97)
4107	492	3.5 (24)	58.5 (23)	137	10.2 (5)	66.4 (60)	130	32.3 (5)	227	27.8 (20)	666	9.6 (56)	667	23.4 (99)
4109	172	4.1 (36)	58.7 (25)	99	*	*	36	*	89	*	261	8.4 (40)	262	19.8 (91)
4112	147	0.7 (1)	72.1 (94)	132	70.5 (60)	11.4 (1)	28	*	33	*	144	18.8 (103)	144	19.4 (89)
4115	692	4.6 (53)	60.8 (44)	297	20.5 (53)	50.5 (11)	184	39.7 (10)	433	25.4 (16)	1,091	9.3 (50)	1,092	16.1 (62)
4351	139	2.9 (11)	62.6 (65)	100	10.0 (3)	65.0 (59)	16	*	36	*	143	9.8 (59)	144	31.9 (103)
4352	62	*	*	45	*	*	10	*	16	*	58	*	58	*
4353	111	4.5 (48)	64.0 (77)	86	*	*	25	*	29	*	126	10.3 (64)	126	8.7 (9)
4354	176	5.1 (63)	57.4 (15)	58	*	*	14	*	32	*	200	9.5 (53)	201	31.3 (102)
4355	79	*	*	56	*	*	11	*	21	*	82	*	82	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2024 Pregnancy Nutrition Surveillance (1)

Page : 23

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4356	99	*	*	78	*	*	13	*	34	*	99	*	99	*
4359	144	3.5 (24)	65.3 (84)	99	*	*	20	*	50	*	151	11.3 (76)	151	22.5 (98)
4360	53	*	*	41	*	*	5	*	19	*	54	*	54	*
4361	63	*	*	45	*	*	6	*	27	*	67	*	66	*
4362	63	*	*	48	*	*	14	*	24	*	66	*	66	*
4363	201	5.5 (71)	60.2 (39)	164	20.7 (54)	56.7 (35)	35	*	70	*	206	5.3 (4)	206	11.7 (28)
4401	250	4.8 (59)	54.0 (6)	183	12.6 (9)	58.5 (46)	40	*	78	*	238	10.5 (69)	238	11.3 (21)
4501	125	6.4 (78)	60.8 (44)	65	*	*	19	*	50	*	194	10.8 (71)	195	12.8 (34)
4601	266	5.3 (69)	59.8 (34)	144	6.9 (1)	61.8 (52)	38	*	83	*	319	10.0 (61)	319	13.5 (43)
4701	285	6.7 (80)	55.1 (9)	178	17.4 (40)	58.4 (43)	32	*	95	*	310	10.0 (61)	311	24.4 (100)
5050	438	4.6 (53)	61.2 (51)	383	17.0 (38)	53.0 (17)	87	*	179	27.9 (21)	438	11.4 (79)	438	19.2 (87)
5051	790	3.3 (21)	64.3 (79)	545	12.7 (10)	54.5 (23)	147	37.4 (9)	342	30.4 (26)	787	10.2 (63)	788	13.6 (45)
5052	896	4.6 (53)	59.5 (30)	757	19.7 (49)	49.1 (8)	152	45.4 (17)	347	30.0 (25)	900	13.6 (96)	902	13.3 (39)
5055	372	2.2 (6)	61.6 (56)	311	16.1 (31)	54.0 (19)	51	*	86	*	406	12.6 (89)	406	9.4 (12)
5101	377	4.5 (48)	62.9 (70)	303	15.5 (23)	55.1 (28)	89	*	250	26.8 (17)	393	8.4 (40)	393	16.5 (67)
5202	164	6.1 (75)	55.5 (10)	111	18.0 (43)	55.9 (31)	21	*	49	*	185	7.0 (18)	186	16.7 (69)
5601	229	4.4 (45)	67.2 (90)	66	*	*	28	*	49	*	269	9.7 (58)	269	16.7 (69)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2024 Pregnancy Nutrition Surveillance (1)

Page : 24

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5801	515	5.0 (60)	67.2 (90)	383	15.9 (28)	60.1 (49)	80	*	104	18.3 (5)	499	12.4 (87)	499	20.0 (92)
5803	28	*	*	26	*	*	3	*	11	*	27	*	27	*
5806	15	*	*	14	*	*	5	*	7	*	19	*	19	*
6101	11	*	*	6	*	*	3	*	0	*	11	*	10	*
6102	9	*	*	3	*	*	1	*	2	*	11	*	11	*
6103	733	4.5 (48)	62.2 (63)	400	15.5 (23)	54.5 (23)	130	42.3 (15)	67	*	934	10.3 (64)	927	19.3 (88)
6105	3	*	*	1	*	*	1	*	0	*	3	*	3	*
6301	951	4.0 (34)	61.6 (56)	627	16.4 (34)	57.3 (37)	164	40.9 (13)	288	29.5 (24)	1,016	6.4 (12)	1,017	10.9 (18)
6312	516	2.9 (11)	60.3 (40)	265	16.2 (33)	52.5 (16)	119	46.2 (18)	205	33.7 (32)	575	6.4 (12)	575	13.0 (35)
6316	410	4.1 (36)	61.2 (51)	307	20.8 (55)	55.7 (29)	66	*	121	22.3 (7)	388	7.5 (27)	388	14.4 (51)
6318	1	*	*	1	*	*	1	*	0	*	1	*	1	*
6320	53	*	*	17	*	*	13	*	24	*	70	*	70	*
6325	68	*	*	28	*	*	14	*	37	*	86	*	86	*
6335	229	3.1 (16)	58.5 (23)	112	13.4 (13)	49.1 (8)	50	*	73	*	259	10.4 (67)	260	15.4 (57)
6345	24	*	*	9	*	*	4	*	6	*	27	*	27	*
6360	183	5.5 (71)	60.7 (43)	84	*	*	53	*	80	*	227	11.5 (81)	226	16.4 (65)
7306	471	4.2 (39)	63.5 (73)	313	18.5 (47)	56.5 (34)	68	*	56	*	500	11.0 (73)	501	11.6 (27)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2024 Pregnancy Nutrition Surveillance (1)

Page : 25

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7401	523	4.2 (39)	59.8 (34)	423	11.6 (6)	64.5 (58)	70	*	179	22.3 (7)	524	11.3 (76)	522	15.1 (56)
7601	71	*	*	10	*	*	18	*	41	*	141	7.8 (31)	141	6.4 (4)
7850	274	5.5 (71)	61.3 (54)	163	12.9 (11)	57.1 (36)	46	*	79	*	339	8.3 (39)	338	14.5 (53)
8101	399	4.3 (42)	61.4 (55)	244	20.1 (51)	54.9 (26)	87	*	106	40.6 (33)	476	6.7 (17)	475	18.7 (83)
8102	96	*	*	48	*	*	20	*	33	*	116	5.2 (3)	116	12.1 (31)
8106	94	*	*	21	*	*	28	*	74	*	169	9.5 (53)	169	16.0 (61)
8150	2	*	*	1	*	*	0	*	1	*	2	*	2	*
8151	704	5.1 (63)	60.9 (47)	412	14.6 (17)	55.8 (30)	155	40.0 (11)	247	24.3 (13)	853	11.1 (74)	852	16.4 (65)
8201	531	4.3 (42)	62.0 (60)	299	15.4 (22)	57.5 (39)	136	40.4 (12)	354	27.4 (18)	598	12.9 (92)	598	17.2 (74)
8202	420	6.7 (80)	49.0 (2)	184	22.3 (57)	42.4 (4)	193	29.0 (1)	312	20.8 (6)	1,137	7.0 (18)	1,137	2.9 (1)
8203	108	11.1 (93)	58.3 (21)	92	*	*	14	*	57	*	113	14.2 (99)	113	8.8 (10)
8250	82	*	*	44	*	*	27	*	45	*	80	*	80	*
8252	14	*	*	8	*	*	6	*	6	*	19	*	19	*
8255	666	4.5 (48)	60.8 (44)	411	15.8 (27)	52.1 (15)	176	36.4 (8)	364	31.3 (27)	710	13.1 (94)	710	18.3 (79)
8257	291	6.2 (77)	54.6 (8)	201	17.4 (40)	41.3 (3)	109	41.3 (14)	164	32.9 (30)	474	9.1 (47)	474	3.2 (2)
8264	257	2.3 (7)	66.9 (89)	215	13.5 (14)	63.3 (55)	35	*	188	18.1 (4)	273	13.9 (98)	273	11.7 (28)
8267	143	4.2 (39)	58.0 (17)	74	*	*	51	*	67	*	140	13.6 (96)	140	20.0 (92)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2024 Pregnancy Nutrition Surveillance (1)

Page : 26

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8901	137	4.4 (45)	57.7 (16)	89	*	*	31	*	24	*	171	6.4 (12)	171	14.6 (55)
8902	269	4.1 (36)	63.6 (75)	146	17.1 (39)	51.4 (12)	50	*	79	*	299	7.7 (30)	299	13.0 (35)
8903	313	3.5 (24)	61.7 (58)	178	10.1 (4)	63.5 (57)	53	*	72	*	351	8.5 (42)	352	18.8 (85)
8904	394	5.1 (63)	59.1 (27)	221	13.1 (12)	58.4 (43)	81	*	130	43.1 (34)	497	8.2 (37)	498	17.5 (76)
8905	379	3.2 (18)	58.0 (17)	252	15.5 (23)	62.3 (54)	90	*	139	25.2 (15)	561	11.8 (83)	561	13.5 (43)
8909	275	2.9 (11)	66.2 (88)	162	14.2 (16)	61.7 (51)	50	*	58	*	287	7.3 (26)	287	18.5 (81)
9045	329	2.1 (5)	68.7 (92)	255	16.5 (36)	54.1 (20)	62	*	100	11.0 (1)	351	5.4 (7)	351	7.1 (6)
9050	227	1.8 (3)	65.6 (86)	91	*	*	52	*	68	*	264	8.0 (35)	264	13.3 (39)
9051	307	2.3 (7)	63.5 (73)	141	28.4 (59)	39.7 (2)	133	30.1 (4)	244	28.7 (22)	699	9.3 (50)	698	9.5 (13)
9053	404	5.0 (60)	58.4 (22)	320	15.9 (28)	58.4 (43)	87	*	70	*	398	6.0 (11)	398	10.1 (14)
9058	413	6.8 (84)	58.1 (19)	268	12.3 (8)	58.2 (42)	90	*	124	32.3 (29)	571	7.2 (23)	571	11.9 (30)
9068	263	3.8 (31)	65.4 (85)	196	18.4 (46)	54.6 (25)	17	*	35	*	257	7.8 (31)	255	14.5 (53)
9069	373	4.0 (34)	56.3 (12)	264	15.9 (28)	54.2 (21)	76	*	57	*	397	5.3 (4)	396	13.6 (45)
9070	516	5.0 (60)	62.6 (65)	355	16.1 (31)	57.5 (39)	110	47.3 (19)	89	*	528	4.7 (2)	527	16.1 (62)
9080	687	3.9 (32)	61.0 (48)	267	13.9 (15)	57.7 (41)	188	35.1 (7)	226	28.8 (23)	950	9.9 (60)	953	15.9 (60)
9082	33	*	*	16	*	*	13	*	16	*	39	*	39	*
9083	3	*	*	3	*	*	0	*	0	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 27

#### Michigan

#### Comparison of Maternal Health Indicators by Clinic

<b>Michigan</b>	<b>33,778</b>	<b>4.7</b>	<b>60.8</b>	<b>19,602</b>	<b>16.8</b>	<b>55.1</b>	<b>6,900</b>	<b>34.8</b>	<b>11,679</b>	<b>25.4</b>	<b>41,389</b>	<b>9.4</b>	<b>41,399</b>	<b>14.5</b>
<b>National (Prior Year)</b>	<b>42,957</b>	<b>4.2</b>	<b>61.7</b>	<b>27,673</b>	<b>18.8</b>	<b>51.7</b>	<b>6,220</b>	<b>26.9</b>	<b>11,239</b>	<b>22.9</b>	<b>42,973</b>	<b>10.4</b>	<b>42,758</b>	<b>15.6</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2024 Pregnancy Nutrition Surveillance (1)

Page : 28

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
0101	30	*	*	32	*	34	*	24 *
0102	101	94.1 (33)	0.0 (1)	119	34.5 (26)	120	25.8 (39)	100 94.0 (7)
0103	111	95.5 (21)	0.9 (81)	118	33.1 (33)	122	27.9 (30)	90 *
0104	24	*	*	28	*	28	*	23 *
0250	28	*	*	43	*	43	*	35 *
0251	20	*	*	22	*	22	*	15 *
0252	27	*	*	29	*	28	*	20 *
0253	26	*	*	28	*	29	*	22 *
0401	94	*	*	117	25.6 (69)	122	32.0 (20)	90 *
0402	89	*	*	104	30.8 (47)	105	29.5 (26)	78 *
0403	38	*	*	40	*	42	*	32 *
0404	39	*	*	42	*	42	*	34 *
0501	121	95.0 (26)	0.0 (1)	138	32.6 (36)	137	47.4 (3)	109 87.2 (55)
0503	14	*	*	15	*	15	*	10 *
0504	102	93.1 (41)	0.0 (1)	107	29.9 (52)	111	36.9 (15)	85 *
0505	62	*	*	73	*	75	*	53 *
0506	48	*	*	56	*	58	*	51 *
0507	19	*	*	20	*	20	*	9 *
0508	2	*	*	2	*	2	*	1 *
0650	279	90.0 (68)	1.1 (90)	343	36.2 (20)	335	31.9 (21)	264 90.5 (24)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 29

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0651	135	92.6 (48)	0.0 (1)	151	41.1 (6)	143	23.8 (48)	128	88.3 (44)
0652	76	*	*	84	*	86	*	75	*
0653	105	97.1 (12)	0.0 (1)	133	51.9 (2)	134	35.1 (18)	121	82.6 (78)
0654	80	*	*	94	*	96	*	68	*
0655	92	*	*	100	37.0 (16)	99	*	76	*
0657	8	*	*	10	*	10	*	8	*
0701	110	93.6 (36)	0.0 (1)	131	35.1 (24)	129	36.4 (16)	96	*
0702	13	*	*	18	*	18	*	15	*
0703	56	*	*	66	*	63	*	47	*
0704	13	*	*	18	*	19	*	18	*
0705	26	*	*	31	*	31	*	25	*
0850	177	94.4 (29)	0.6 (63)	199	38.7 (9)	200	19.0 (66)	150	92.7 (13)
0851	190	96.3 (16)	0.0 (1)	215	33.5 (32)	214	29.0 (27)	156	93.6 (9)
0901	370	94.1 (33)	0.0 (1)	420	33.1 (33)	423	22.5 (52)	328	87.5 (52)
0902	7	*	*	8	*	8	*	8	*
1051	277	92.8 (42)	0.4 (50)	330	36.7 (18)	328	31.4 (23)	244	87.7 (48)
1052	1	*	*	1	*	1	*	0	*
1101	329	89.4 (70)	0.3 (43)	383	20.4 (85)	365	18.4 (69)	245	89.4 (32)
1102	209	92.8 (42)	0.0 (1)	239	22.6 (80)	234	17.9 (72)	163	95.1 (2)
1106	7	*	*	10	*	10	*	8	*
1250	309	98.1 (5)	0.0 (1)	343	26.8 (62)	350	46.0 (5)	267	83.9 (74)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 6A

2024 Pregnancy Nutrition Surveillance (1)

Page : 30

Michigan

## Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1251	228	96.1 (17)	0.9 (81)	245	37.6 (13)	244	22.5 (52)	187	87.7 (48)
1252	123	90.2 (66)	0.0 (1)	169	37.3 (14)	167	18.6 (68)	135	80.7 (81)
1253	152	88.2 (78)	2.0 (96)	185	28.1 (59)	186	14.5 (90)	131	86.3 (61)
1350	674	96.1 (17)	0.3 (43)	746	31.8 (38)	686	40.4 (10)	576	82.6 (78)
1352	117	99.1 (2)	0.0 (1)	125	30.4 (49)	123	18.7 (67)	97	*
1401	60	*	*	68	*	69	*	57	*
1402	22	*	*	27	*	28	*	16	*
1403	4	*	*	5	*	5	*	3	*
1501	183	84.2 (93)	0.5 (58)	242	24.8 (71)	237	46.4 (4)	189	89.4 (32)
1503	223	98.7 (3)	0.0 (1)	264	39.4 (8)	266	18.0 (71)	206	79.6 (84)
1504	537	96.6 (14)	0.0 (1)	617	31.0 (44)	599	22.4 (54)	435	86.2 (62)
1506	122	91.8 (59)	0.0 (1)	171	35.1 (24)	168	17.3 (76)	132	87.1 (56)
1507	150	98.0 (6)	0.0 (1)	161	34.2 (28)	159	25.2 (42)	121	94.2 (6)
1509	95	*	*	112	29.5 (53)	111	16.2 (81)	86	*
1511	144	88.9 (75)	2.1 (98)	174	35.6 (22)	174	43.1 (7)	130	92.3 (16)
1512	98	*	*	121	26.4 (64)	116	44.0 (6)	81	*
1514	220	96.4 (15)	0.0 (1)	238	30.7 (48)	237	38.8 (12)	155	91.0 (22)
1701	155	98.7 (3)	0.6 (63)	183	47.5 (3)	184	58.2 (2)	146	92.5 (14)
1950	169	95.3 (23)	0.0 (1)	193	29.5 (53)	194	42.8 (8)	146	92.5 (14)
1952	35	*	*	39	*	39	*	21	*
1953	71	*	*	80	*	79	*	51	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 31

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1954	104	95.2 (25)	1.0 (85)	115	31.3 (40)	111	65.8 (1)	83	*
1955	39	*	*	47	*	45	*	39	*
1957	110	97.3 (9)	0.0 (1)	126	30.2 (50)	112	38.4 (13)	95	*
1963	30	*	*	34	*	34	*	35	*
2150	151	90.7 (63)	0.0 (1)	166	33.7 (31)	165	27.9 (30)	133	94.0 (7)
2151	60	*	*	74	*	67	*	54	*
2152	6	*	*	7	*	7	*	6	*
2251	82	*	*	99	*	97	*	76	*
2252	45	*	*	51	*	47	*	40	*
2554	9	*	*	9	*	8	*	7	*
2555	1,807	88.2 (78)	0.6 (63)	2,155	27.7 (60)	2,104	10.6 (97)	1,612	84.0 (73)
2556	96	*	*	101	1.0 (106)	106	6.6 (100)	9	*
3001	286	86.7 (82)	0.0 (1)	341	41.9 (5)	346	14.5 (90)	275	86.5 (60)
3002	175	95.4 (22)	0.0 (1)	184	31.0 (44)	183	14.8 (87)	153	90.2 (28)
3004	74	*	*	88	*	91	*	62	*
3005	47	*	*	53	*	53	*	46	*
3201	120	96.7 (13)	0.8 (76)	143	25.9 (68)	143	39.2 (11)	107	91.6 (19)
3301	1,220	90.5 (64)	0.8 (76)	1,518	26.8 (62)	1,510	14.8 (87)	1,099	85.5 (65)
3304	8	*	*	10	*	11	*	11	*
3305	17	*	*	25	*	25	*	22	*
3306	24	*	*	33	*	34	*	32	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 32

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
3308	9	*	*	13	*	13	*	12 *
3312	75	*	*	90	*	91	*	80 *
3314	1	*	*	1	*	1	*	1 *
3550	14	*	*	15	*	15	*	12 *
3551	4	*	*	4	*	4	*	4 *
3701	247	94.3 (30)	0.0 (1)	269	33.1 (33)	269	21.9 (57)	193 91.7 (17)
3850	645	92.1 (53)	0.6 (63)	754	34.4 (27)	760	21.6 (58)	578 89.8 (31)
3901	399	94.2 (32)	0.3 (43)	484	28.9 (56)	484	18.4 (69)	343 84.5 (72)
3950	486	92.0 (56)	0.6 (63)	588	35.5 (23)	595	23.9 (45)	448 87.7 (48)
4106	770	84.5 (90)	1.0 (85)	899	25.1 (70)	892	11.7 (95)	665 79.8 (82)
4107	634	92.7 (47)	0.0 (1)	729	30.2 (50)	708	27.0 (34)	643 91.3 (21)
4109	251	92.8 (42)	1.6 (95)	274	32.1 (37)	275	24.0 (44)	182 89.0 (37)
4112	142	64.8 (98)	0.0 (1)	148	14.2 (98)	16	*	128 85.2 (68)
4115	928	92.8 (42)	0.6 (63)	1,146	24.3 (73)	743	14.9 (86)	846 85.5 (65)
4351	115	85.2 (86)	0.0 (1)	156	38.5 (11)	143	28.7 (28)	109 88.1 (45)
4352	54	*	*	61	*	62	*	49 *
4353	97	*	*	136	33.8 (29)	130	22.3 (55)	103 78.6 (86)
4354	190	95.3 (23)	0.0 (1)	221	52.0 (1)	220	26.4 (35)	179 88.8 (39)
4355	80	*	*	87	*	90	*	68 *
4356	82	*	*	105	40.0 (7)	104	32.7 (19)	78 *
4359	100	82.0 (96)	2.0 (96)	158	44.3 (4)	155	27.1 (33)	127 86.6 (59)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 33

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
4360	51	*	*	55	*	57	*	43 *
4361	63	*	*	75	*	74	*	55 *
4362	57	*	*	75	*	74	*	63 *
4363	195	88.2 (78)	1.0 (85)	209	29.2 (55)	215	31.2 (24)	159 88.7 (40)
4401	211	92.4 (51)	0.9 (81)	246	31.3 (40)	254	26.4 (35)	187 87.7 (48)
4501	177	90.4 (65)	1.1 (90)	207	36.7 (18)	205	23.9 (45)	153 85.0 (69)
4601	292	92.5 (50)	0.0 (1)	339	26.3 (66)	335	17.6 (74)	236 89.0 (37)
4701	277	92.4 (51)	0.7 (73)	327	31.2 (42)	329	36.2 (17)	239 90.0 (29)
5050	428	91.8 (59)	0.2 (40)	449	18.5 (89)	451	17.1 (78)	301 94.4 (5)
5051	761	93.2 (39)	0.7 (73)	835	17.2 (91)	836	21.3 (59)	538 90.5 (24)
5052	872	90.1 (67)	0.6 (63)	920	17.6 (90)	927	17.3 (76)	596 91.6 (19)
5055	378	99.7 (1)	0.3 (43)	415	38.6 (10)	329	4.6 (102)	321 92.8 (12)
5101	374	92.0 (56)	0.3 (43)	410	12.9 (99)	409	23.7 (49)	275 94.9 (3)
5202	155	94.8 (27)	0.6 (63)	201	37.3 (14)	188	37.2 (14)	152 84.9 (70)
5601	257	97.3 (9)	0.4 (50)	286	31.1 (43)	285	26.3 (37)	217 89.4 (32)
5801	461	93.5 (37)	0.4 (50)	538	30.9 (46)	485	42.1 (9)	406 90.9 (23)
5803	28	*	*	29	*	29	*	12 *
5806	19	*	*	19	*	19	*	7 *
6101	10	*	*	12	*	11	*	9 *
6102	10	*	*	12	*	12	*	11 *
6103	886	93.2 (39)	0.5 (58)	982	23.3 (77)	925	31.5 (22)	640 93.3 (10)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 34

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6105	3	*	*	3	*	3	*	2	*
6301	955	94.6 (28)	0.8 (76)	1,090	26.0 (67)	1,024	12.8 (93)	790	86.1 (63)
6312	513	92.8 (42)	0.4 (50)	620	21.0 (83)	598	19.6 (65)	426	73.2 (89)
6316	355	93.5 (37)	0.6 (63)	421	26.4 (64)	407	19.7 (64)	297	82.5 (80)
6318	1	*	*	1	*	1	*	1	*
6320	69	*	*	75	*	73	*	47	*
6325	83	*	*	92	*	93	*	57	*
6335	242	92.1 (53)	0.8 (76)	273	28.9 (56)	272	25.0 (43)	205	91.7 (17)
6345	26	*	*	28	*	29	*	17	*
6360	214	88.8 (77)	0.0 (1)	247	20.2 (86)	243	23.9 (45)	170	89.4 (32)
7306	451	89.4 (70)	1.1 (90)	549	36.8 (17)	546	23.3 (51)	442	87.3 (54)
7401	410	89.0 (74)	0.2 (40)	559	38.1 (12)	568	23.6 (50)	416	88.7 (40)
7601	133	91.0 (62)	0.0 (1)	154	31.8 (38)	155	11.6 (96)	118	83.1 (76)
7850	280	94.3 (30)	0.4 (50)	363	35.8 (21)	341	22.3 (55)	271	88.6 (42)
8101	450	83.8 (94)	1.3 (93)	508	15.4 (94)	504	16.5 (80)	325	90.5 (24)
8102	113	89.4 (70)	0.0 (1)	119	23.5 (76)	123	17.9 (72)	79	*
8106	166	95.8 (20)	0.0 (1)	181	11.6 (102)	179	16.2 (81)	90	*
8150	2	*	*	2	*	2	*	2	*
8151	809	92.6 (48)	0.5 (58)	928	28.2 (58)	916	25.9 (38)	696	86.9 (57)
8201	586	91.8 (59)	0.0 (1)	663	9.2 (103)	642	28.0 (29)	378	94.7 (4)
8202	1,055	97.5 (8)	0.1 (38)	1,176	33.8 (29)	1,176	13.7 (92)	923	85.6 (64)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 35

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
8203	111	97.3 (9)	0.0 (1)	119	15.1 (96)	121	25.6 (40)	72 *
8250	80	*	*	90	*	88	*	90 *
8252	18	*	*	21	*	21	*	17 *
8255	699	92.0 (56)	1.4 (94)	769	8.7 (104)	770	20.9 (62)	470 95.3 (1)
8257	458	95.9 (19)	0.0 (1)	486	16.0 (93)	470	27.4 (32)	344 73.3 (88)
8264	266	92.1 (53)	0.8 (76)	274	7.7 (105)	281	25.3 (41)	148 88.5 (43)
8267	133	97.7 (7)	0.0 (1)	160	12.5 (101)	158	15.8 (84)	160 90.0 (29)
8901	160	83.8 (94)	0.6 (63)	182	20.9 (84)	179	16.8 (79)	124 87.9 (47)
8902	290	85.2 (86)	1.0 (85)	320	15.0 (97)	319	21.0 (61)	203 89.2 (36)
8903	337	80.1 (97)	0.9 (81)	369	12.7 (100)	368	12.8 (93)	235 83.0 (77)
8904	479	86.6 (84)	1.0 (85)	526	15.2 (95)	516	14.7 (89)	316 85.4 (67)
8905	554	93.7 (35)	0.2 (40)	593	24.5 (72)	588	31.1 (25)	414 79.5 (85)
8909	279	84.6 (89)	0.4 (50)	320	23.8 (75)	313	17.6 (74)	234 83.8 (75)
9045	351	84.3 (91)	0.3 (43)	381	23.9 (74)	320	3.8 (104)	271 90.4 (27)
9050	249	84.7 (88)	0.4 (50)	273	22.3 (81)	278	21.2 (60)	190 75.8 (87)
9051	637	88.9 (75)	0.5 (58)	746	27.1 (61)	606	4.3 (103)	625 93.1 (11)
9053	393	89.3 (73)	0.3 (43)	438	19.6 (87)	414	15.9 (83)	294 87.4 (53)
9058	545	87.2 (81)	0.7 (73)	599	19.5 (88)	576	8.2 (99)	372 79.8 (82)
9068	254	84.3 (91)	0.0 (1)	277	22.7 (79)	267	19.9 (63)	194 88.1 (45)
9069	398	86.7 (82)	0.5 (58)	421	23.0 (78)	417	5.8 (101)	282 58.9 (90)
9070	513	85.8 (85)	0.4 (50)	561	16.9 (92)	558	15.1 (85)	348 84.8 (71)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2024 Pregnancy Nutrition Surveillance (1)

Page : 36

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9080	889	89.5 (69)	0.1 (38)	1,014	21.9 (82)	993	10.4 (98)	680	86.9 (57)
9082	38	*	*	43	*	44	*	27	*
9083	3	*	*	3	*	3	*	3	*
<b>Michigan</b>	<b>38,343</b>	<b>91.5</b>	<b>0.5</b>	<b>44,111</b>	<b>26.9</b>	<b>42,685</b>	<b>21.9</b>	<b>31,855</b>	<b>31855.0</b>
<b>National (Prior Year)</b>	<b>45,177</b>	<b>86.0</b>	<b>1.2</b>	<b>50,643</b>	<b>26.0</b>	<b>44,385</b>	<b>23.0</b>	<b>38,838</b>	<b>91.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 37

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	24	*	31	*	4	*	24	*	31	*
0102	99	*	113	7.1 (80)	19	*	100	22.0 (87)	110	9.1 (90)
0103	90	*	118	8.5 (90)	10	*	90	*	118	9.3 (93)
0104	22	*	25	*	0	*	23	*	24	*
0250	34	*	39	*	12	*	35	*	38	*
0251	15	*	21	*	4	*	15	*	21	*
0252	20	*	26	*	4	*	20	*	27	*
0253	22	*	27	*	6	*	22	*	27	*
0401	89	*	111	8.1 (89)	14	*	90	*	109	9.2 (91)
0402	77	*	97	*	16	*	78	*	97	*
0403	32	*	39	*	2	*	32	*	39	*
0404	34	*	40	*	8	*	34	*	40	*
0501	108	16.7 (73)	122	7.4 (81)	17	*	109	16.5 (72)	126	11.1 (97)
0503	10	*	14	*	1	*	10	*	14	*
0504	85	*	105	7.6 (83)	12	*	85	*	105	4.8 (57)
0505	52	*	71	*	9	*	53	*	71	*
0506	51	*	53	*	12	*	51	*	53	*
0507	9	*	19	*	1	*	9	*	19	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 38

Michigan

## Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0508	1	*	2	*	0	*	1	*	2	*
0650	256	10.5 (46)	319	6.9 (79)	30	*	258	16.3 (69)	317	6.9 (70)
0651	128	20.3 (83)	145	13.8 (102)	25	*	129	21.7 (83)	144	11.8 (98)
0652	75	*	84	*	16	*	75	*	84	*
0653	121	22.3 (86)	131	9.9 (97)	26	*	121	23.1 (88)	130	9.2 (91)
0654	68	*	89	*	12	*	68	*	89	*
0655	76	*	95	*	16	*	77	*	96	*
0657	8	*	9	*	2	*	8	*	9	*
0701	94	*	125	16.0 (103)	22	*	96	*	127	2.4 (19)
0702	15	*	14	*	4	*	15	*	14	*
0703	47	*	58	*	11	*	46	*	57	*
0704	18	*	16	*	2	*	18	*	16	*
0705	25	*	28	*	8	*	25	*	27	*
0850	150	8.0 (28)	183	6.0 (71)	13	*	151	15.2 (66)	186	3.2 (32)
0851	156	9.0 (33)	208	3.4 (41)	12	*	156	12.2 (51)	205	2.4 (19)
0901	327	9.8 (43)	397	5.8 (66)	32	*	326	10.4 (41)	395	7.8 (78)
0902	8	*	8	*	1	*	8	*	8	*
1051	243	19.8 (81)	309	6.1 (72)	42	*	242	12.8 (57)	311	4.5 (53)
1052	0	*	1	*	0	*	0	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 39

Michigan

## Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1101	239	11.3 (50)	343	5.8 (66)	27	*	245	12.2 (51)	342	7.6 (77)
1102	163	16.0 (70)	215	6.5 (74)	26	*	162	12.3 (54)	215	8.8 (86)
1106	8	*	8	*	0	*	8	*	8	*
1250	267	9.7 (41)	327	5.5 (61)	24	*	266	9.4 (33)	327	4.9 (58)
1251	187	20.3 (83)	236	9.7 (95)	35	*	187	20.3 (81)	237	8.9 (87)
1252	135	11.9 (55)	156	5.1 (56)	11	*	136	14.0 (60)	158	3.2 (32)
1253	131	17.6 (76)	174	9.2 (93)	19	*	131	26.7 (90)	174	7.5 (76)
1350	569	11.8 (54)	697	2.6 (30)	70	*	563	12.6 (56)	683	3.7 (42)
1352	93	*	117	5.1 (56)	17	*	96	*	119	1.7 (12)
1401	57	*	66	*	10	*	56	*	65	*
1402	16	*	24	*	2	*	16	*	24	*
1403	3	*	4	*	1	*	3	*	4	*
1501	188	11.2 (49)	222	5.9 (68)	19	*	189	9.5 (35)	223	2.7 (26)
1503	205	14.6 (67)	239	5.9 (68)	23	*	206	10.2 (39)	241	3.3 (36)
1504	432	6.7 (20)	566	2.3 (22)	27	*	437	3.9 (6)	565	1.4 (9)
1506	132	12.9 (60)	156	10.3 (98)	17	*	131	4.6 (10)	156	3.2 (32)
1507	121	19.0 (80)	151	3.3 (39)	19	*	119	16.8 (74)	150	4.0 (46)
1509	86	*	105	1.9 (18)	8	*	86	*	106	5.7 (65)
1511	127	14.2 (66)	167	6.6 (75)	21	*	130	23.8 (89)	167	9.0 (88)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 40

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1512	81	*	110	1.8 (17)	4	*	81	*	111	2.7 (26)
1514	152	7.9 (27)	228	2.6 (30)	14	*	151	4.0 (7)	228	1.8 (13)
1701	144	17.4 (75)	165	10.9 (100)	25	*	146	19.9 (80)	167	9.0 (88)
1950	145	15.2 (68)	183	8.7 (91)	21	*	144	20.8 (82)	183	12.6 (101)
1952	21	*	38	*	2	*	20	*	38	*
1953	51	*	75	*	4	*	51	*	75	*
1954	83	*	106	3.8 (47)	8	*	83	*	108	12.0 (100)
1955	39	*	40	*	7	*	39	*	40	*
1957	95	*	114	3.5 (44)	11	*	95	*	121	7.4 (75)
1963	35	*	31	*	5	*	35	*	31	*
2150	133	26.3 (89)	155	13.5 (101)	31	*	133	15.8 (67)	154	2.6 (22)
2151	53	*	64	*	12	*	53	*	64	*
2152	6	*	6	*	1	*	6	*	6	*
2251	76	*	90	*	16	*	75	*	91	*
2252	38	*	44	*	9	*	40	*	46	*
2554	8	*	9	*	0	*	7	*	9	*
2555	1,612	12.0 (56)	2,010	5.7 (64)	177	32.8 (2)	1,590	14.6 (63)	1,986	7.0 (71)
2556	9	*	106	5.7 (64)	1	*	9	*	106	4.7 (56)
3001	274	11.7 (53)	321	6.2 (73)	29	*	273	14.3 (62)	320	6.6 (68)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 41

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3002	153	7.8 (24)	178	3.9 (48)	12	*	152	21.7 (83)	178	10.1 (94)
3004	62	*	87	*	3	*	62	*	86	*
3005	46	*	52	*	5	*	46	*	52	*
3201	107	16.8 (74)	142	2.8 (34)	17	*	107	18.7 (77)	142	7.0 (71)
3301	1,097	9.3 (37)	1,457	5.5 (61)	102	37.3 (1)	1,096	10.6 (43)	1,460	2.9 (29)
3304	11	*	10	*	0	*	11	*	10	*
3305	21	*	23	*	1	*	21	*	23	*
3306	32	*	33	*	5	*	31	*	33	*
3308	12	*	12	*	2	*	12	*	12	*
3312	80	*	87	*	9	*	80	*	87	*
3314	1	*	1	*	0	*	1	*	1	*
3550	12	*	15	*	5	*	12	*	15	*
3551	4	*	4	*	2	*	4	*	4	*
3701	192	18.8 (78)	257	9.7 (95)	36	*	192	21.9 (86)	258	5.0 (59)
3850	577	13.3 (62)	701	7.4 (81)	68	*	576	17.0 (76)	700	10.1 (94)
3901	343	10.5 (46)	441	5.9 (68)	30	*	340	10.0 (38)	438	3.2 (32)
3950	447	12.3 (58)	550	5.3 (59)	50	*	449	9.4 (33)	550	4.5 (53)
4106	659	7.6 (23)	872	3.7 (46)	56	*	624	5.6 (14)	874	1.1 (7)
4107	636	11.3 (50)	667	3.0 (37)	68	*	626	11.5 (46)	668	3.6 (41)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 42

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4109	176	9.1 (35)	261	4.2 (53)	20	*	181	12.2 (51)	262	4.2 (48)
4112	128	2.3 (6)	144	0.0 (1)	3	*	128	2.3 (4)	144	0.0 (1)
4115	841	4.9 (14)	1,098	1.7 (15)	43	*	825	4.5 (9)	1,091	2.1 (16)
4351	111	25.2 (88)	145	10.3 (98)	25	*	106	16.0 (68)	145	13.1 (102)
4352	49	*	58	*	14	*	49	*	57	*
4353	102	7.8 (24)	126	5.6 (63)	7	*	102	9.8 (37)	126	1.6 (10)
4354	179	21.8 (85)	202	9.4 (94)	35	*	179	21.8 (85)	203	11.8 (98)
4355	69	*	82	*	10	*	69	*	82	*
4356	77	*	99	*	19	*	78	*	99	*
4359	124	11.3 (50)	152	7.9 (88)	16	*	127	7.1 (21)	151	5.3 (62)
4360	43	*	54	*	5	*	43	*	54	*
4361	55	*	67	*	11	*	55	*	67	*
4362	62	*	66	*	9	*	63	*	66	*
4363	160	18.8 (78)	206	8.7 (91)	28	*	160	19.4 (79)	205	8.3 (83)
4401	184	15.2 (68)	238	7.6 (83)	29	*	181	10.5 (42)	238	5.0 (59)
4501	153	22.9 (87)	195	7.7 (85)	33	*	153	16.3 (69)	198	10.6 (96)
4601	236	8.1 (29)	319	5.3 (59)	16	*	236	8.9 (28)	321	3.7 (42)
4701	238	16.0 (70)	310	3.9 (48)	36	*	240	7.5 (23)	311	3.9 (44)
5050	300	9.0 (33)	433	3.0 (37)	25	*	298	11.7 (48)	433	5.3 (62)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 43

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
5051	537	5.4 (16)	788	1.3 (9)	27	*	535	10.7 (44)	785	7.0 (71)
5052	596	4.4 (9)	900	2.3 (22)	25	*	582	9.1 (30)	890	5.1 (61)
5055	321	1.6 (4)	406	0.5 (6)	4	*	319	0.6 (2)	405	0.5 (3)
5101	275	6.5 (19)	393	2.8 (34)	17	*	274	5.5 (13)	393	3.3 (36)
5202	149	20.1 (82)	182	7.7 (85)	32	*	146	6.2 (18)	182	3.3 (36)
5601	216	13.9 (64)	268	3.4 (41)	29	*	214	6.5 (19)	273	4.4 (51)
5801	400	17.8 (77)	488	6.6 (75)	74	*	398	16.6 (73)	498	7.8 (78)
5803	12	*	28	*	1	*	12	*	27	*
5806	7	*	19	*	1	*	7	*	19	*
6101	9	*	12	*	0	*	9	*	12	*
6102	11	*	11	*	1	*	11	*	11	*
6103	639	9.7 (41)	939	7.8 (87)	61	*	640	10.3 (40)	944	4.4 (51)
6105	2	*	3	*	0	*	2	*	3	*
6301	785	4.8 (13)	1,022	1.9 (18)	37	*	775	5.2 (12)	1,016	2.6 (22)
6312	423	2.1 (5)	576	0.7 (7)	8	*	419	6.9 (20)	570	1.6 (10)
6316	297	4.7 (12)	391	1.5 (13)	10	*	293	9.6 (36)	392	7.1 (74)
6318	1	*	1	*	0	*	1	*	1	*
6320	46	*	70	*	3	*	47	*	70	*
6325	57	*	88	*	4	*	56	*	88	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 44

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
6335	205	4.9 (14)	260	1.5 (13)	10	*	204	13.2 (58)	261	8.0 (80)
6345	17	*	26	*	3	*	17	*	27	*
6360	169	7.1 (21)	228	3.9 (48)	12	*	169	18.9 (78)	228	6.6 (68)
7306	440	9.8 (43)	505	5.1 (56)	42	*	442	12.4 (55)	506	5.7 (65)
7401	417	16.1 (72)	524	6.7 (77)	58	*	416	16.3 (69)	521	8.4 (84)
7601	114	14.0 (65)	141	2.1 (20)	19	*	118	8.5 (27)	141	6.4 (67)
7850	267	10.9 (48)	342	6.7 (77)	30	*	270	5.9 (15)	342	5.6 (64)
8101	324	9.3 (37)	476	2.7 (33)	28	*	325	8.0 (25)	476	4.0 (46)
8102	79	*	116	1.7 (15)	3	*	78	*	116	2.6 (22)
8106	90	*	168	3.6 (45)	5	*	90	*	170	0.6 (4)
8150	2	*	2	*	0	*	2	*	2	*
8151	697	8.9 (31)	855	4.4 (54)	56	*	694	7.1 (21)	857	2.7 (26)
8201	372	7.8 (24)	601	3.3 (39)	31	*	377	9.0 (29)	596	4.2 (48)
8202	924	1.4 (3)	1,137	0.3 (4)	13	*	925	3.2 (5)	1,135	1.8 (13)
8203	72	*	114	0.9 (8)	7	*	72	*	114	15.8 (103)
8250	90	*	79	*	7	*	90	*	80	*
8252	17	*	19	*	2	*	17	*	19	*
8255	469	10.4 (45)	716	3.9 (48)	45	*	469	13.2 (58)	719	8.1 (81)
8257	345	0.0 (1)	475	0.2 (2)	0	*	343	1.5 (3)	475	0.6 (4)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 45

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8264	148	9.5 (39)	273	2.2 (21)	14	*	148	4.1 (8)	273	3.3 (36)
8267	160	13.8 (63)	140	2.9 (36)	12	*	159	11.9 (49)	140	8.6 (85)
8901	124	7.3 (22)	171	2.3 (22)	7	*	123	4.9 (11)	171	2.3 (17)
8902	202	4.5 (11)	300	2.3 (22)	9	*	203	5.9 (15)	300	2.3 (17)
8903	234	13.2 (61)	353	3.4 (41)	30	*	233	14.6 (63)	353	4.2 (48)
8904	313	8.6 (30)	501	5.0 (55)	29	*	315	14.0 (60)	501	3.0 (31)
8905	414	4.3 (8)	563	2.5 (29)	12	*	413	15.0 (65)	563	8.2 (82)
8909	234	6.4 (18)	288	1.4 (12)	13	*	230	10.9 (45)	288	3.5 (40)
9045	270	4.4 (9)	353	2.3 (22)	11	*	270	9.3 (31)	354	4.5 (53)
9050	189	12.2 (57)	266	2.6 (30)	22	*	190	11.6 (47)	265	2.6 (22)
9051	625	1.1 (2)	699	0.4 (5)	7	*	626	0.5 (1)	704	0.1 (2)
9053	292	8.9 (31)	398	4.0 (52)	23	*	290	9.3 (31)	393	2.0 (15)
9058	372	5.6 (17)	567	2.3 (22)	18	*	365	7.9 (24)	571	1.2 (8)
9068	194	12.4 (59)	258	2.3 (22)	21	*	193	11.9 (49)	257	3.9 (44)
9069	282	2.8 (7)	397	1.3 (9)	6	*	282	6.0 (17)	399	1.0 (6)
9070	350	9.1 (35)	528	0.2 (2)	26	*	349	16.9 (75)	525	2.9 (29)
9080	680	9.6 (40)	949	1.3 (9)	58	*	681	8.2 (26)	953	2.4 (19)
9082	27	*	39	*	3	*	27	*	39	*
9083	3	*	3	*	1	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2024 Pregnancy Nutrition Surveillance (1)

Page : 46

### Michigan

#### Comparison of Smoking Indicators by Clinic

Michigan	31,723	10.3	41,470	4.5	3,060	41.6	31,605	11.1	41,446	4.9
National (Prior Year)	40,473	22.0	47,284	6.2	7,409	32.6	39,853	9.6	51,116	5.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 47

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	29	*	*	29	*	24	*	29	*
0102	107	6.5 (8)	12.1 (99)	110	10.9 (21)	94	*	106	71.7 (72)
0103	114	9.6 (35)	9.6 (92)	114	12.3 (41)	97	*	114	71.1 (74)
0104	24	*	*	24	*	24	*	24	*
0250	35	*	*	36	*	31	*	35	*
0251	21	*	*	21	*	21	*	21	*
0252	24	*	*	24	*	22	*	24	*
0253	27	*	*	27	*	23	*	27	*
0401	107	2.8 (1)	9.3 (90)	107	11.2 (29)	91	*	107	86.9 (6)
0402	90	*	*	92	*	76	*	90	*
0403	38	*	*	37	*	32	*	38	*
0404	38	*	*	38	*	35	*	38	*
0501	111	8.1 (21)	9.0 (87)	114	7.9 (3)	102	2.9 (15)	111	86.5 (9)
0503	14	*	*	14	*	11	*	14	*
0504	99	*	*	100	15.0 (78)	83	*	99	*
0505	67	*	*	66	*	60	*	68	*
0506	50	*	*	50	*	43	*	50	*
0507	16	*	*	19	*	16	*	17	*
0508	2	*	*	2	*	1	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 48

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	299	8.7 (25)	7.0 (66)	305	10.2 (14)	266	4.5 (42)	299	83.3 (28)
0651	135	9.6 (35)	6.7 (61)	139	7.2 (1)	125	8.8 (81)	135	81.5 (39)
0652	78	*	*	79	*	67	*	78	*
0653	118	5.9 (4)	5.9 (51)	120	9.2 (8)	107	4.7 (45)	117	83.8 (22)
0654	82	*	*	83	*	68	*	82	*
0655	88	*	*	91	*	81	*	88	*
0657	9	*	*	9	*	9	*	9	*
0701	119	6.7 (9)	4.2 (20)	124	10.5 (18)	105	1.0 (2)	121	90.9 (1)
0702	13	*	*	14	*	13	*	13	*
0703	56	*	*	57	*	54	*	56	*
0704	16	*	*	15	*	14	*	16	*
0705	25	*	*	26	*	21	*	24	*
0850	176	11.4 (60)	6.8 (63)	179	12.8 (48)	154	5.2 (55)	177	86.4 (10)
0851	194	8.8 (26)	6.7 (61)	202	14.4 (64)	167	2.4 (10)	196	85.2 (16)
0901	380	12.1 (66)	8.7 (84)	390	14.1 (62)	326	5.2 (55)	382	74.1 (68)
0902	8	*	*	8	*	8	*	8	*
1051	297	7.7 (15)	8.8 (85)	300	11.3 (32)	260	4.2 (38)	299	89.6 (2)
1052	1	*	*	1	*	1	*	1	*
1101	316	11.7 (61)	4.7 (27)	334	13.2 (52)	273	4.4 (40)	317	74.1 (68)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 49

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1102	200	5.0 (2)	5.5 (41)	208	8.2 (4)	184	1.1 (3)	202	74.8 (64)
1106	7	*	*	7	*	7	*	7	*
1250	278	9.0 (30)	6.1 (55)	295	11.9 (37)	239	5.4 (60)	292	76.4 (57)
1251	215	11.2 (58)	6.0 (54)	224	16.1 (86)	178	3.9 (30)	217	87.1 (5)
1252	141	9.9 (41)	7.8 (77)	150	14.0 (60)	120	1.7 (5)	144	84.0 (21)
1253	159	6.9 (11)	12.6 (100)	168	11.9 (37)	138	2.2 (8)	163	82.8 (32)
1350	643	10.6 (49)	7.9 (80)	665	12.2 (40)	561	5.0 (53)	650	80.0 (47)
1352	107	14.0 (81)	11.2 (98)	112	13.4 (55)	93	*	109	77.1 (54)
1401	58	*	*	63	*	51	*	58	*
1402	20	*	*	23	*	19	*	20	*
1403	4	*	*	4	*	3	*	4	*
1501	205	9.8 (39)	9.8 (93)	214	14.5 (67)	172	2.3 (9)	210	81.0 (41)
1503	222	7.7 (15)	7.7 (76)	221	10.9 (21)	193	3.1 (18)	222	80.2 (44)
1504	499	10.0 (43)	7.0 (66)	542	13.7 (58)	434	4.1 (35)	531	83.8 (22)
1506	150	8.0 (20)	7.3 (72)	153	18.3 (94)	122	1.6 (4)	150	83.3 (28)
1507	144	8.3 (22)	5.6 (46)	146	14.4 (64)	121	4.1 (35)	146	83.6 (26)
1509	97	*	*	96	*	82	*	97	*
1511	148	12.2 (67)	4.7 (27)	156	15.4 (83)	124	4.8 (49)	150	84.7 (18)
1512	107	10.3 (45)	4.7 (27)	109	11.0 (25)	95	*	108	86.1 (11)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 50

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1514	211	12.8 (72)	4.3 (22)	219	16.9 (90)	174	1.7 (5)	214	87.4 (4)
1701	146	10.3 (45)	4.8 (31)	152	11.8 (34)	128	4.7 (45)	147	85.0 (17)
1950	173	11.0 (55)	5.8 (48)	177	12.4 (43)	150	5.3 (58)	174	86.8 (7)
1952	34	*	*	37	*	34	*	35	*
1953	70	*	*	73	*	62	*	72	*
1954	100	9.0 (30)	10.0 (94)	104	7.7 (2)	92	*	100	86.0 (12)
1955	38	*	*	37	*	31	*	38	*
1957	112	8.9 (28)	8.9 (86)	118	14.4 (64)	97	*	112	86.6 (8)
1963	29	*	*	28	*	27	*	29	*
2150	144	8.3 (22)	5.6 (46)	149	14.8 (75)	124	3.2 (22)	145	78.6 (50)
2151	62	*	*	63	*	54	*	62	*
2152	6	*	*	6	*	6	*	6	*
2251	89	*	*	90	*	78	*	89	*
2252	40	*	*	43	*	34	*	41	*
2554	9	*	*	9	*	9	*	9	*
2555	1,894	13.0 (74)	5.3 (37)	1,930	13.2 (52)	1,634	7.1 (74)	1,908	65.4 (84)
2556	100	23.0 (100)	3.0 (9)	101	22.8 (102)	74	*	101	62.4 (88)
3001	286	16.8 (92)	3.1 (10)	292	19.2 (98)	228	8.3 (79)	287	61.3 (91)
3002	170	10.0 (43)	2.9 (7)	171	11.1 (27)	151	4.0 (32)	170	67.6 (79)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 51

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3004	83	*	*	83	*	65	*	83	*
3005	50	*	*	50	*	47	*	50	*
3201	131	5.3 (3)	7.6 (75)	134	11.2 (29)	116	2.6 (13)	132	83.3 (28)
3301	1,358	12.4 (68)	6.6 (58)	1,405	13.5 (56)	1,170	5.1 (54)	1,378	83.4 (27)
3304	8	*	*	8	*	7	*	8	*
3305	23	*	*	23	*	20	*	23	*
3306	30	*	*	30	*	25	*	30	*
3308	9	*	*	9	*	8	*	9	*
3312	83	*	*	87	*	70	*	84	*
3314	1	*	*	1	*	1	*	1	*
3550	15	*	*	15	*	13	*	15	*
3551	4	*	*	4	*	3	*	4	*
3701	243	12.8 (72)	8.6 (83)	246	15.0 (78)	204	6.4 (68)	244	85.7 (15)
3850	659	12.6 (69)	5.9 (51)	669	15.7 (85)	553	4.2 (38)	662	81.1 (40)
3901	401	11.7 (61)	6.2 (56)	417	12.5 (45)	348	5.7 (64)	406	72.4 (70)
3950	511	10.8 (51)	7.0 (66)	515	12.6 (46)	441	4.8 (49)	515	77.3 (52)
4106	790	13.5 (79)	4.7 (27)	824	14.8 (75)	671	7.0 (73)	807	77.3 (52)
4107	632	11.1 (56)	4.4 (23)	649	15.4 (83)	534	3.7 (27)	641	81.0 (41)
4109	240	15.0 (86)	9.2 (88)	249	18.1 (93)	195	3.1 (18)	244	81.6 (38)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 52

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4112	132	7.6 (14)	1.5 (2)	137	14.6 (70)	113	3.5 (26)	133	46.6 (101)
4115	986	11.3 (59)	5.5 (41)	1,051	12.7 (47)	855	4.1 (35)	1,040	82.9 (31)
4351	132	6.1 (6)	6.8 (63)	136	14.7 (73)	111	0.9 (1)	133	82.7 (33)
4352	55	*	*	53	*	44	*	55	*
4353	119	10.9 (53)	8.4 (82)	122	9.0 (6)	108	5.6 (63)	120	81.7 (36)
4354	191	8.9 (28)	7.3 (72)	195	9.2 (8)	172	6.4 (68)	193	84.5 (19)
4355	75	*	*	75	*	66	*	75	*
4356	92	*	*	96	*	79	*	93	*
4359	143	7.7 (15)	2.8 (4)	144	11.8 (34)	124	3.2 (22)	145	75.9 (60)
4360	50	*	*	51	*	46	*	51	*
4361	58	*	*	63	*	51	*	60	*
4362	62	*	*	66	*	53	*	63	*
4363	190	8.4 (24)	11.1 (96)	193	14.0 (60)	160	3.1 (18)	192	85.9 (13)
4401	219	6.8 (10)	5.5 (41)	231	9.1 (7)	199	2.5 (11)	221	83.7 (25)
4501	180	7.8 (18)	7.8 (77)	183	10.4 (17)	159	1.9 (7)	180	81.7 (36)
4601	307	6.2 (7)	9.4 (91)	306	9.2 (8)	275	2.9 (15)	306	80.1 (46)
4701	295	10.8 (51)	6.8 (63)	297	16.5 (88)	244	3.3 (24)	297	85.9 (13)
5050	423	10.9 (53)	5.0 (33)	425	11.1 (27)	374	6.1 (66)	425	76.2 (59)
5051	752	9.7 (38)	3.5 (15)	766	12.1 (39)	657	5.5 (62)	760	78.7 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 53

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5052	850	13.3 (78)	4.5 (25)	860	15.1 (82)	710	7.5 (75)	858	75.5 (62)
5055	373	9.9 (41)	4.6 (26)	391	11.0 (25)	330	4.8 (49)	385	64.2 (86)
5101	379	14.0 (81)	6.6 (58)	378	16.7 (89)	310	4.8 (49)	379	80.2 (44)
5202	173	15.0 (86)	5.2 (36)	179	18.4 (95)	141	6.4 (68)	176	87.5 (3)
5601	246	6.9 (11)	10.2 (95)	259	8.9 (5)	224	4.5 (42)	246	68.3 (77)
5801	466	7.9 (19)	7.3 (72)	481	9.6 (13)	417	3.1 (18)	467	71.5 (73)
5803	27	*	*	26	*	24	*	27	*
5806	19	*	*	19	*	16	*	19	*
6101	12	*	*	12	*	11	*	12	*
6102	10	*	*	11	*	8	*	10	*
6103	876	11.1 (56)	5.9 (51)	920	15.0 (78)	742	5.8 (65)	887	71.8 (71)
6105	3	*	*	3	*	2	*	3	*
6301	923	11.8 (63)	6.5 (57)	986	13.0 (51)	799	5.4 (60)	967	75.4 (63)
6312	543	13.1 (76)	5.5 (41)	559	14.5 (67)	456	5.3 (58)	548	78.3 (51)
6316	365	12.6 (69)	5.5 (41)	377	14.6 (70)	309	5.2 (55)	367	80.9 (43)
6318	1	*	*	1	*	0	*	1	*
6320	63	*	*	69	*	57	*	68	*
6325	84	*	*	84	*	70	*	84	*
6335	244	11.9 (64)	5.3 (37)	251	14.3 (63)	208	2.9 (15)	249	79.1 (48)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 54

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6345	27	*	*	26	*	22	*	27	*
6360	211	14.2 (83)	7.1 (69)	218	12.8 (48)	182	8.8 (81)	214	77.1 (54)
7306	479	11.9 (64)	7.1 (69)	487	14.6 (70)	408	3.7 (27)	485	69.9 (75)
7401	500	9.2 (32)	7.8 (77)	506	10.5 (18)	445	4.0 (32)	501	75.8 (61)
7601	135	5.9 (4)	11.1 (96)	137	9.5 (12)	122	2.5 (11)	136	61.0 (92)
7850	323	7.1 (13)	8.0 (81)	330	13.3 (54)	282	2.8 (14)	324	81.8 (35)
8101	417	16.5 (90)	2.4 (3)	456	20.6 (101)	327	8.9 (83)	452	62.2 (89)
8102	92	*	*	108	10.2 (14)	83	*	102	67.6 (79)
8106	158	17.1 (94)	5.1 (34)	166	19.3 (99)	129	4.7 (45)	163	58.9 (96)
8150	2	*	*	2	*	2	*	2	*
8151	797	13.0 (74)	5.8 (48)	818	14.7 (73)	675	3.9 (30)	811	82.6 (34)
8201	550	9.3 (33)	4.9 (32)	580	10.9 (21)	486	3.7 (27)	562	74.7 (65)
8202	1,051	8.8 (26)	3.7 (17)	1,100	9.4 (11)	949	4.4 (40)	1,090	67.7 (78)
8203	111	12.6 (69)	7.2 (71)	110	10.9 (21)	98	*	111	83.8 (22)
8250	76	*	*	78	*	67	*	77	*
8252	16	*	*	17	*	14	*	17	*
8255	654	9.8 (39)	5.8 (48)	683	12.4 (43)	566	4.6 (44)	672	74.7 (65)
8257	387	13.2 (77)	3.1 (10)	427	10.5 (18)	344	8.1 (78)	415	62.7 (87)
8264	261	10.3 (45)	9.2 (88)	265	11.7 (33)	227	4.0 (32)	262	84.4 (20)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 55

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8267	137	9.5 (34)	6.6 (58)	138	12.3 (41)	121	3.3 (24)	136	74.3 (67)
8901	150	22.7 (99)	1.3 (1)	160	16.3 (87)	121	13.2 (90)	159	59.1 (94)
8902	252	16.3 (89)	4.4 (23)	286	13.6 (57)	214	9.8 (86)	275	69.8 (76)
8903	307	17.9 (95)	2.9 (7)	339	18.0 (92)	250	11.2 (89)	323	57.3 (97)
8904	432	16.7 (91)	4.2 (20)	477	18.9 (97)	346	6.9 (72)	455	67.3 (82)
8905	469	10.4 (48)	5.1 (34)	537	11.2 (29)	412	6.1 (66)	527	76.3 (58)
8909	251	14.3 (84)	3.6 (16)	282	14.5 (67)	213	7.5 (75)	266	67.3 (82)
9045	341	10.6 (49)	5.3 (37)	349	12.9 (50)	297	4.7 (45)	344	50.9 (99)
9050	250	16.8 (92)	4.0 (18)	262	17.9 (91)	203	8.9 (83)	256	59.8 (93)
9051	570	9.6 (35)	5.4 (40)	637	11.8 (34)	493	6.5 (71)	643	76.5 (56)
9053	369	13.6 (80)	3.3 (14)	391	10.2 (14)	331	9.7 (85)	371	56.3 (98)
9058	523	18.7 (96)	4.0 (18)	545	15.0 (78)	436	11.0 (88)	531	67.4 (81)
9068	250	19.6 (97)	2.8 (4)	256	14.8 (75)	211	13.7 (91)	251	59.0 (95)
9069	372	21.0 (98)	3.2 (12)	387	19.9 (100)	298	10.4 (87)	375	49.9 (100)
9070	497	15.3 (88)	3.2 (12)	508	18.5 (96)	402	7.5 (75)	503	61.6 (90)
9080	867	14.9 (85)	2.8 (4)	915	13.7 (58)	747	8.3 (79)	900	65.2 (85)
9082	38	*	*	39	*	29	*	38	*
9083	3	*	*	3	*	3	*	3	*
<b>Michigan</b>	<b>38,264</b>	<b>11.5</b>	<b>5.9</b>	<b>39,791</b>	<b>13.4</b>	<b>32,907</b>	<b>5.3</b>	<b>39,073</b>	<b>75.2</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2024 Pregnancy Nutrition Surveillance (1)

Page : 56

### Michigan

### Comparison of Infant Health Indicators by Clinic

<b>National (Prior Year)</b>	<b>47,760</b>	<b>11.0</b>	<b>5.8</b>	<b>49,201</b>	<b>13.1</b>	<b>40,792</b>	<b>5.0</b>	<b>48,458</b>	<b>47.2</b>
------------------------------	---------------	-------------	------------	---------------	-------------	---------------	------------	---------------	-------------

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 57

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
0101	122	90.2	0.0	7.4	0.0	0.0	2.5	0.0
0102	362	89.2	0.6	4.4	0.0	0.0	5.5	0.3
0103	369	94.0	0.5	1.6	0.0	0.0	3.5	0.3
0104	113	93.8	0.9	3.5	0.0	0.0	1.8	0.0
0105	1	*	*	*	*	*	*	*
0250	121	81.8	0.0	3.3	5.8	0.0	9.1	0.0
0251	66	*	*	*	*	*	*	*
0252	104	53.8	2.9	3.8	1.9	1.0	36.5	0.0
0253	113	72.6	0.0	3.5	10.6	1.8	11.5	0.0
0401	399	95.5	0.3	2.0	0.0	0.3	2.0	0.0
0402	327	81.0	3.1	4.9	0.0	0.6	10.4	0.0
0403	120	89.2	0.0	6.7	1.7	0.8	1.7	0.0
0404	111	93.7	0.9	2.7	0.0	0.0	2.7	0.0
0501	401	88.8	1.5	4.5	0.2	0.5	4.2	0.2
0503	51	*	*	*	*	*	*	*
0504	311	85.2	1.6	5.1	1.6	1.0	5.5	0.0
0505	202	91.6	0.5	2.5	0.0	0.0	5.4	0.0
0506	171	90.1	1.2	3.5	0.0	0.6	4.7	0.0
0507	57	*	*	*	*	*	*	*
0508	10	*	*	*	*	*	*	*
0650	1,069	78.8	3.9	7.9	3.6	0.5	4.9	0.6
0651	512	91.4	0.8	5.3	0.2	0.4	2.0	0.0
0652	258	93.8	1.2	3.1	0.8	0.0	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 58

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
0653	383	92.7	0.5	2.9	0.0	0.3	3.4	0.3
0654	375	92.3	0.5	2.9	0.5	0.5	2.9	0.3
0655	312	94.9	0.0	2.2	0.0	0.0	2.9	0.0
0657	26	*	*	*	*	*	*	*
0701	418	86.1	3.3	1.7	0.5	2.9	5.3	0.2
0702	73	*	*	*	*	*	*	*
0703	181	85.6	3.3	2.2	3.3	0.0	4.4	1.1
0704	56	*	*	*	*	*	*	*
0705	83	*	*	*	*	*	*	*
0706	1	*	*	*	*	*	*	*
0850	669	82.1	6.0	7.8	0.0	2.1	2.1	0.0
0851	631	90.6	0.6	5.4	0.5	1.0	1.9	0.0
0901	1,343	80.2	2.8	9.4	0.1	0.5	6.7	0.4
0902	22	*	*	*	*	*	*	*
1051	1,058	80.0	1.5	10.2	1.8	1.4	4.8	0.3
1052	3	*	*	*	*	*	*	*
1101	1,295	37.2	44.3	10.9	0.5	2.3	3.7	1.0
1102	716	70.5	10.5	8.1	0.3	1.3	8.4	1.0
1106	52	*	*	*	*	*	*	*
1250	993	77.6	7.0	13.2	0.3	0.3	1.4	0.1
1251	685	92.3	0.3	3.8	0.3	0.1	3.1	0.1
1252	530	59.8	0.4	35.7	0.0	0.2	3.2	0.8
1253	618	77.3	0.5	13.4	0.0	0.2	7.9	0.6
1350	2,297	58.3	18.8	10.4	0.3	4.7	6.7	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 59

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
1352	354	68.1	22.6	4.5	0.3	0.0	4.2	0.3
1401	204	88.2	0.0	7.4	0.5	0.0	3.9	0.0
1402	79	*	*	*	*	*	*	*
1403	18	*	*	*	*	*	*	*
1501	750	44.8	5.5	43.7	0.0	0.0	4.0	2.0
1503	809	58.2	7.5	23.9	0.2	0.6	7.2	2.3
1504	1,846	35.3	4.7	51.8	0.1	3.8	3.5	0.8
1506	519	78.8	0.4	14.6	0.0	0.4	4.8	1.0
1507	498	81.3	2.0	11.8	0.2	1.4	2.4	0.8
1509	342	47.4	2.0	45.0	0.3	0.9	2.9	1.5
1511	585	86.8	1.5	6.3	0.0	0.7	4.6	0.0
1512	385	82.1	2.3	9.1	0.5	1.3	4.2	0.5
1514	714	75.8	5.5	14.0	0.0	1.1	2.9	0.7
1701	519	59.9	1.0	1.7	20.6	1.0	15.6	0.2
1950	563	92.4	0.4	5.0	0.0	0.5	1.6	0.2
1952	127	92.1	0.8	5.5	0.0	0.0	1.6	0.0
1953	263	87.8	0.4	9.1	0.0	0.8	1.9	0.0
1954	353	84.7	0.0	10.5	0.3	0.0	4.2	0.3
1955	137	86.9	0.0	10.9	0.0	0.0	2.2	0.0
1957	378	81.7	1.6	12.7	0.0	1.1	2.6	0.3
1963	83	*	*	*	*	*	*	*
2150	519	83.6	0.4	4.4	2.3	0.4	8.7	0.2
2151	225	80.0	1.3	8.9	0.0	0.0	6.7	3.1
2152	32	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 60

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2251	334	92.8	0.0	3.6	0.0	0.6	2.4	0.6
2252	175	92.0	0.0	4.0	0.0	0.0	2.3	1.7
2554	39	*	*	*	*	*	*	*
2555	6,909	49.9	36.6	6.7	0.0	0.3	6.2	0.3
2556	162	30.2	50.6	9.9	0.0	1.2	8.0	0.0
2557	1	*	*	*	*	*	*	*
2562	1	*	*	*	*	*	*	*
2563	3	*	*	*	*	*	*	*
2564	3	*	*	*	*	*	*	*
3001	1,116	29.7	40.3	16.0	0.4	0.6	12.6	0.3
3002	434	37.3	36.2	18.4	0.0	0.5	7.6	0.0
3004	297	75.4	0.7	20.2	0.0	0.3	3.0	0.3
3005	163	89.6	0.6	5.5	1.2	0.6	2.5	0.0
3007	3	*	*	*	*	*	*	*
3201	428	91.1	0.5	5.8	0.0	0.2	0.9	1.4
3301	4,373	37.8	30.3	17.7	0.1	6.1	7.6	0.4
3304	83	*	*	*	*	*	*	*
3305	143	62.9	13.3	9.8	0.7	4.2	9.1	0.0
3306	174	83.3	4.6	9.8	0.0	0.6	1.1	0.6
3308	76	*	*	*	*	*	*	*
3312	420	45.7	23.3	20.0	0.7	3.1	7.1	0.0
3314	2	*	*	*	*	*	*	*
3550	53	*	*	*	*	*	*	*
3551	7	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 61

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3701	819	88.0	0.2	8.1	0.0	0.1	3.3	0.2
3850	2,502	71.1	12.7	7.8	0.2	0.6	7.2	0.4
3901	1,396	35.7	50.4	6.4	0.1	1.1	5.9	0.5
3902	2	*	*	*	*	*	*	*
3950	1,942	51.4	22.7	15.0	0.0	1.3	9.0	0.6
3954	1	*	*	*	*	*	*	*
4106	2,379	12.4	32.7	49.7	0.0	0.9	4.0	0.3
4107	2,568	40.3	28.2	22.7	0.4	1.8	6.5	0.1
4109	712	82.3	1.8	10.5	0.1	1.4	3.4	0.4
4112	536	0.7	1.1	97.8	0.0	0.0	0.4	0.0
4115	3,422	29.4	32.7	22.6	0.3	9.8	5.0	0.2
4351	522	88.7	0.6	5.9	0.0	0.8	4.0	0.0
4352	192	72.9	8.3	8.3	0.0	0.0	9.9	0.5
4353	451	53.7	0.7	41.2	0.0	0.0	2.7	1.8
4354	720	85.1	3.6	5.3	0.6	0.4	4.4	0.6
4355	266	83.5	0.8	8.6	1.9	0.0	5.3	0.0
4356	367	79.0	0.8	13.6	0.3	0.3	5.7	0.3
4359	535	62.4	0.7	34.4	0.4	0.0	1.1	0.9
4360	215	89.8	0.5	7.4	0.5	0.0	1.9	0.0
4361	226	88.9	0.9	4.9	0.0	0.0	4.9	0.4
4362	250	92.4	0.4	4.0	0.0	0.0	2.4	0.4
4363	646	89.6	0.9	5.3	0.0	0.2	3.6	0.5
4401	729	86.7	0.8	6.9	0.1	0.5	4.8	0.1
4501	711	93.2	0.4	4.8	0.0	0.1	0.7	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 62

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4601	1,147	72.6	2.2	19.4	0.2	0.5	4.2	1.0
4701	994	86.7	1.2	6.3	0.2	1.9	2.7	0.9
5050	1,582	44.3	43.6	3.8	0.1	2.4	5.6	0.3
5051	2,544	48.3	37.1	5.1	0.1	4.0	4.7	0.5
5052	2,771	28.1	39.0	3.1	0.1	24.8	4.6	0.4
5055	1,330	74.3	10.9	1.2	0.2	12.7	0.7	0.1
5101	1,358	61.4	12.2	18.9	0.1	2.2	5.1	0.1
5102	2	*	*	*	*	*	*	*
5202	625	87.0	1.3	3.2	2.4	0.2	5.1	0.8
5206	7	*	*	*	*	*	*	*
5601	919	86.8	2.4	6.4	0.0	0.9	3.5	0.0
5801	1,714	79.9	6.3	8.9	0.0	0.1	3.9	1.0
5803	40	*	*	*	*	*	*	*
5804	2	*	*	*	*	*	*	*
5806	67	*	*	*	*	*	*	*
6101	58	*	*	*	*	*	*	*
6102	53	*	*	*	*	*	*	*
6103	3,148	55.7	25.9	11.1	0.4	0.6	6.1	0.2
6105	7	*	*	*	*	*	*	*
6301	3,412	35.4	30.4	27.2	0.2	2.7	3.7	0.3
6312	1,814	31.6	56.6	5.4	0.1	2.3	3.9	0.1
6316	1,209	58.4	23.2	11.0	0.0	2.7	4.1	0.5
6318	11	*	*	*	*	*	*	*
6320	451	37.3	50.1	4.7	0.4	2.9	4.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 63

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
6325	311	88.4	1.3	5.5	0.0	0.0	4.8	0.0
6335	906	57.6	25.4	4.9	0.0	9.8	1.8	0.6
6345	135	80.7	3.7	8.1	0.0	2.2	5.2	0.0
6360	768	29.3	41.7	22.9	0.1	2.2	3.4	0.4
7306	1,699	46.3	29.7	17.4	0.3	0.7	5.4	0.2
7401	1,757	78.9	5.3	7.3	0.1	0.5	7.5	0.4
7601	487	91.4	0.0	6.8	0.0	0.0	1.6	0.2
7850	1,076	91.5	0.3	4.7	0.0	0.1	2.9	0.5
7852	5	*	*	*	*	*	*	*
8101	1,645	6.9	88.4	2.0	0.0	0.3	2.2	0.1
8102	373	21.4	20.1	53.6	0.0	2.7	1.9	0.3
8106	538	31.4	46.3	14.1	0.0	0.7	6.9	0.6
8150	4	*	*	*	*	*	*	*
8151	3,087	37.2	35.3	15.8	0.1	3.8	7.2	0.6
8152	5	*	*	*	*	*	*	*
8153	1	*	*	*	*	*	*	*
8155	4	*	*	*	*	*	*	*
8201	2,342	42.4	40.9	6.0	0.2	4.4	5.6	0.5
8202	3,862	91.2	5.9	1.3	0.0	0.4	0.8	0.3
8203	348	57.5	29.6	5.7	0.0	0.0	7.2	0.0
8250	395	23.5	65.1	4.8	0.3	0.5	5.8	0.0
8252	97	*	*	*	*	*	*	*
8255	1,997	52.4	28.1	11.2	0.4	3.3	4.1	0.6
8257	1,547	53.8	7.4	1.0	0.0	36.1	1.5	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 64

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
8263	234	59.0	19.7	15.8	0.0	0.9	4.7	0.0
8264	841	56.2	30.1	6.3	0.2	3.6	3.3	0.2
8267	629	53.4	17.6	23.8	0.0	0.8	4.1	0.2
8901	707	9.1	81.0	4.0	0.0	2.3	3.5	0.1
8902	1,154	7.9	87.1	2.3	0.0	0.7	2.0	0.1
8903	1,253	5.1	87.4	1.8	0.0	2.6	3.1	0.0
8904	1,791	4.7	89.3	2.3	0.1	1.1	2.4	0.1
8905	1,753	66.3	21.4	7.8	0.1	1.7	2.4	0.3
8909	1,067	5.6	88.8	2.4	0.1	0.3	2.8	0.0
9045	1,240	9.0	19.7	68.9	0.1	0.4	1.0	0.9
9050	862	18.0	59.3	20.0	0.0	0.1	2.7	0.0
9051	1,945	6.1	4.0	89.0	0.0	0.1	0.6	0.3
9053	1,560	9.1	80.8	2.8	0.1	4.7	2.3	0.3
9055	7	*	*	*	*	*	*	*
9058	1,546	16.8	72.0	1.3	0.1	7.4	2.1	0.3
9066	368	2.4	92.9	1.1	0.0	1.1	2.4	0.0
9068	820	3.5	93.9	0.9	0.0	0.4	1.2	0.1
9069	1,483	1.4	96.2	1.2	0.0	0.1	1.0	0.1
9070	1,900	6.1	89.5	1.6	0.1	1.6	1.1	0.2
9080	2,839	20.7	71.3	3.2	0.1	0.3	3.7	0.6
9082	106	0.9	92.5	4.7	0.9	0.9	0.0	0.0
9083	12	*	*	*	*	*	*	*
9087	3	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 3A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 65

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic  
Combined 3 Years

Michigan	138,955	50.8	28.2	13.0	0.4	2.8	4.4	0.4
----------	---------	------	------	------	-----	-----	-----	-----

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 66

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	122	0.0	1.6	10.7	57.4	28.7	1.6	0.0
0102	362	0.0	0.8	8.3	58.8	26.8	3.0	2.2
0103	369	0.3	2.2	6.8	62.1	26.3	1.1	1.4
0104	113	0.0	3.5	6.2	57.5	31.0	0.0	1.8
0105	1	*	*	*	*	*	*	*
0250	121	0.0	0.8	8.3	55.4	30.6	2.5	2.5
0251	66	*	*	*	*	*	*	*
0252	104	0.0	1.9	2.9	57.7	35.6	1.9	0.0
0253	113	0.0	2.7	2.7	63.7	30.1	0.9	0.0
0401	399	0.0	1.0	6.8	60.9	30.3	1.0	0.0
0402	327	0.0	1.5	6.1	62.1	29.1	0.3	0.9
0403	120	0.0	2.5	8.3	62.5	25.0	1.7	0.0
0404	111	0.0	0.9	5.4	64.0	26.1	0.9	2.7
0501	401	0.0	0.7	6.2	65.6	23.4	2.0	2.0
0503	51	*	*	*	*	*	*	*
0504	311	0.0	0.3	6.8	56.6	33.8	2.3	0.3
0505	202	0.0	0.5	5.0	62.4	27.2	2.5	2.5
0506	171	0.0	2.3	9.4	62.6	24.0	0.6	1.2
0507	57	*	*	*	*	*	*	*
0508	10	*	*	*	*	*	*	*
0650	1,069	0.1	1.6	6.4	64.3	24.4	1.9	1.4
0651	512	0.2	1.4	6.4	63.9	24.6	1.2	2.3
0652	258	0.0	1.6	8.5	62.4	22.9	1.9	2.7
0653	383	0.0	1.3	4.4	62.9	26.1	1.0	4.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 67

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0654	375	0.0	1.6	5.1	66.4	23.2	0.8	2.9
0655	312	0.0	1.9	4.2	63.1	26.9	1.0	2.9
0657	26	*	*	*	*	*	*	*
0701	418	0.0	0.7	7.7	58.6	30.6	1.9	0.5
0702	73	*	*	*	*	*	*	*
0703	181	0.0	0.0	5.5	58.0	35.4	1.1	0.0
0704	56	*	*	*	*	*	*	*
0705	83	*	*	*	*	*	*	*
0706	1	*	*	*	*	*	*	*
0850	669	0.0	1.6	7.6	55.3	31.7	2.1	1.6
0851	631	0.0	1.9	6.5	64.7	23.3	1.4	2.2
0901	1,343	0.1	2.6	8.6	59.8	26.8	1.1	1.0
0902	22	*	*	*	*	*	*	*
1051	1,058	0.1	1.6	5.2	56.9	32.7	2.6	0.9
1052	3	*	*	*	*	*	*	*
1101	1,295	0.2	1.9	7.9	57.1	27.7	2.2	3.1
1102	716	0.1	1.8	7.7	63.1	24.6	1.3	1.4
1106	52	*	*	*	*	*	*	*
1250	993	0.0	1.5	5.9	61.0	26.1	4.1	1.3
1251	685	0.3	2.6	7.2	63.2	21.0	2.3	3.4
1252	530	0.0	2.6	6.4	62.5	22.1	2.6	3.8
1253	618	0.0	1.8	6.6	58.1	29.0	1.5	3.1
1350	2,297	0.0	2.1	6.4	59.6	27.6	2.2	2.1
1352	354	0.0	1.4	5.9	59.9	27.4	1.1	4.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 68

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1401	204	1.0	1.0	3.9	58.8	29.9	3.4	2.0
1402	79	*	*	*	*	*	*	*
1403	18	*	*	*	*	*	*	*
1501	750	0.1	2.9	5.6	56.1	30.0	2.8	2.4
1503	809	0.0	1.5	5.9	60.9	27.4	1.6	2.6
1504	1,846	0.2	1.8	6.8	57.9	30.9	2.0	0.4
1506	519	0.0	1.5	6.7	60.7	26.2	2.3	2.5
1507	498	0.0	0.4	5.2	57.4	33.7	1.2	2.0
1509	342	0.0	1.8	6.7	52.3	34.8	2.3	2.0
1511	585	0.0	1.7	5.8	61.4	28.5	1.4	1.2
1512	385	0.0	0.8	6.5	60.0	29.9	2.3	0.5
1514	714	0.0	1.3	4.2	55.0	36.8	2.2	0.4
1701	519	0.2	1.0	4.6	62.2	28.3	1.2	2.5
1950	563	0.0	2.5	7.6	63.8	23.4	1.1	1.6
1952	127	0.0	1.6	5.5	59.8	29.9	0.8	2.4
1953	263	0.0	0.0	8.4	65.0	23.6	1.1	1.9
1954	353	0.0	2.0	7.6	60.3	27.5	2.0	0.6
1955	137	0.0	1.5	5.8	63.5	25.5	2.2	1.5
1957	378	0.0	1.9	4.8	60.1	31.5	1.1	0.8
1963	83	*	*	*	*	*	*	*
2150	519	0.0	0.8	5.8	61.1	28.7	3.5	0.2
2151	225	0.4	2.7	9.8	60.9	24.9	0.9	0.4
2152	32	*	*	*	*	*	*	*
2251	334	0.3	0.9	4.5	59.9	32.0	1.5	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 69

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2252	175	0.0	3.4	5.1	60.6	29.1	1.1	0.6
2554	39	*	*	*	*	*	*	*
2555	6,909	0.1	2.0	6.6	60.8	28.1	1.5	0.9
2556	162	0.0	1.2	4.3	61.1	30.9	1.9	0.6
2557	1	*	*	*	*	*	*	*
2562	1	*	*	*	*	*	*	*
2563	3	*	*	*	*	*	*	*
2564	3	*	*	*	*	*	*	*
3001	1,116	0.0	0.6	5.4	63.6	24.7	1.4	4.2
3002	434	0.0	1.2	5.3	55.3	29.7	2.3	6.2
3004	297	0.0	1.0	6.4	52.9	36.7	2.0	1.0
3005	163	0.0	0.0	6.1	57.7	30.7	2.5	3.1
3007	3	*	*	*	*	*	*	*
3201	428	0.0	0.5	4.7	60.7	31.3	1.4	1.4
3301	4,373	0.1	2.0	5.3	58.4	30.5	2.1	1.6
3304	83	*	*	*	*	*	*	*
3305	143	0.0	0.7	2.1	55.9	37.8	3.5	0.0
3306	174	0.0	0.6	3.4	56.9	35.1	2.9	1.1
3308	76	*	*	*	*	*	*	*
3312	420	0.0	1.4	5.5	57.6	31.9	2.1	1.4
3314	2	*	*	*	*	*	*	*
3550	53	*	*	*	*	*	*	*
3551	7	*	*	*	*	*	*	*
3701	819	0.1	2.2	7.7	60.1	26.9	1.5	1.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 70

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3850	2,502	0.1	2.2	6.4	59.4	26.8	1.2	3.9
3901	1,396	0.1	2.4	7.4	57.4	27.7	2.8	2.3
3902	2	*	*	*	*	*	*	*
3950	1,942	0.1	1.8	6.4	59.0	28.6	2.5	1.6
3954	1	*	*	*	*	*	*	*
4106	2,379	0.1	2.5	6.4	55.9	29.3	3.3	2.5
4107	2,568	0.2	1.7	5.3	57.2	31.5	2.5	1.6
4109	712	0.0	1.7	4.8	56.5	33.6	2.9	0.6
4112	536	0.2	2.4	6.3	56.9	28.2	4.1	1.9
4115	3,422	0.1	1.1	4.0	53.9	36.4	3.1	1.4
4351	522	0.0	1.0	5.6	64.6	23.9	1.5	3.4
4352	192	0.5	1.0	5.7	55.7	30.7	1.6	4.7
4353	451	0.0	1.8	6.9	55.9	27.7	3.1	4.7
4354	720	0.0	2.1	7.6	62.4	25.1	2.2	0.6
4355	266	0.0	0.8	9.0	57.5	26.3	1.5	4.9
4356	367	0.0	0.5	7.1	61.0	27.8	0.8	2.7
4359	535	0.2	1.9	4.3	60.9	28.6	2.6	1.5
4360	215	0.0	1.4	5.6	59.1	29.3	1.9	2.8
4361	226	0.0	0.9	10.2	57.1	26.5	4.9	0.4
4362	250	0.0	2.4	8.0	58.4	26.0	2.0	3.2
4363	646	0.0	1.5	6.3	60.4	27.1	2.2	2.5
4401	729	0.0	1.0	4.3	61.9	28.4	1.9	2.6
4501	711	0.0	1.3	5.8	64.0	24.3	2.0	2.7
4601	1,147	0.1	1.3	5.5	62.9	27.1	1.4	1.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 71

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4701	994	0.0	1.2	4.9	56.8	34.3	2.1	0.6
5050	1,582	0.1	1.1	3.8	53.9	37.5	2.8	0.8
5051	2,544	0.0	0.8	3.5	56.8	35.7	2.5	0.7
5052	2,771	0.0	0.9	2.8	55.9	37.6	1.9	0.8
5055	1,330	0.0	0.2	1.7	44.9	49.5	3.6	0.1
5101	1,358	0.0	0.9	3.2	56.6	36.6	2.3	0.4
5102	2	*	*	*	*	*	*	*
5202	625	0.2	1.0	5.4	60.6	30.4	2.1	0.3
5206	7	*	*	*	*	*	*	*
5601	919	0.0	0.9	4.8	62.1	30.1	1.6	0.4
5801	1,714	0.0	1.2	6.2	63.5	26.5	1.5	1.2
5803	40	*	*	*	*	*	*	*
5804	2	*	*	*	*	*	*	*
5806	67	*	*	*	*	*	*	*
6101	58	*	*	*	*	*	*	*
6102	53	*	*	*	*	*	*	*
6103	3,148	0.1	2.0	5.6	61.3	26.3	1.8	2.9
6105	7	*	*	*	*	*	*	*
6301	3,412	0.1	1.6	5.5	58.1	32.1	2.1	0.6
6312	1,814	0.0	0.5	3.5	49.0	42.6	4.0	0.4
6316	1,209	0.0	0.4	2.6	53.3	39.1	4.2	0.3
6318	11	*	*	*	*	*	*	*
6320	451	0.0	0.2	4.0	50.6	39.9	4.0	1.3
6325	311	0.0	1.0	3.5	57.2	32.2	2.6	3.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 72

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
6335	906	0.1	0.7	3.1	50.4	41.7	3.2	0.8
6345	135	0.0	2.2	4.4	63.0	27.4	2.2	0.7
6360	768	0.1	0.5	2.3	60.4	32.6	1.7	2.3
7306	1,699	0.4	2.3	6.3	56.1	32.4	2.2	0.4
7401	1,757	0.1	1.5	6.7	60.6	27.5	1.5	2.1
7601	487	0.0	2.1	6.0	63.9	25.7	1.4	1.0
7850	1,076	0.1	2.0	5.4	63.0	27.4	1.5	0.7
7852	5	*	*	*	*	*	*	*
8101	1,645	0.1	2.4	6.3	52.7	34.4	2.7	1.5
8102	373	0.0	1.9	5.4	57.4	32.4	2.1	0.8
8106	538	0.0	0.9	4.6	57.1	35.1	2.0	0.2
8150	4	*	*	*	*	*	*	*
8151	3,087	0.1	1.4	4.9	53.2	36.1	3.7	0.8
8152	5	*	*	*	*	*	*	*
8153	1	*	*	*	*	*	*	*
8155	4	*	*	*	*	*	*	*
8201	2,342	0.0	1.2	4.3	54.8	35.8	3.0	0.8
8202	3,862	0.0	0.7	3.0	51.0	39.5	5.3	0.4
8203	348	0.0	1.1	4.3	58.0	32.8	3.4	0.3
8250	395	0.0	3.3	6.1	52.4	34.4	3.8	0.0
8252	97	*	*	*	*	*	*	*
8255	1,997	0.1	1.3	3.4	59.4	33.5	2.1	0.4
8257	1,547	0.0	0.5	2.0	52.4	40.3	4.6	0.2
8263	234	0.0	0.4	6.4	59.4	32.1	1.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 4A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 73

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8264	841	0.1	0.8	3.6	57.9	33.5	3.6	0.5
8267	629	0.0	0.8	4.8	57.9	33.5	2.5	0.5
8901	707	0.1	2.8	5.9	53.5	33.9	2.0	1.7
8902	1,154	0.1	1.4	5.2	48.2	40.6	3.5	1.1
8903	1,253	0.0	2.1	4.8	55.4	32.5	2.6	2.6
8904	1,791	0.3	2.3	5.8	56.4	32.4	1.7	0.9
8905	1,753	0.0	0.6	2.9	53.7	39.4	3.3	0.3
8909	1,067	0.0	1.7	5.7	54.9	33.5	3.5	0.7
9045	1,240	0.2	2.1	5.9	52.9	34.3	3.5	1.2
9050	862	0.2	1.6	5.9	55.9	32.5	2.4	1.4
9051	1,945	0.1	1.9	5.2	56.1	31.9	3.5	1.3
9053	1,560	0.0	1.8	4.5	57.1	32.4	3.1	1.2
9055	7	*	*	*	*	*	*	*
9058	1,546	0.1	1.6	4.9	52.3	37.1	3.0	1.0
9066	368	0.0	3.0	5.4	50.5	38.6	2.4	0.0
9068	820	0.4	1.7	4.9	53.8	34.1	2.3	2.8
9069	1,483	0.1	2.3	5.1	51.9	37.1	3.0	0.5
9070	1,900	0.1	2.6	6.8	53.6	33.3	1.9	1.7
9080	2,839	0.1	1.4	5.8	51.5	37.3	2.6	1.4
9082	106	0.0	0.9	6.6	49.1	41.5	1.9	0.0
9083	12	*	*	*	*	*	*	*
9087	3	*	*	*	*	*	*	*
<b>Michigan</b>	<b>138,955</b>	<b>0.1</b>	<b>1.5</b>	<b>5.3</b>	<b>57.3</b>	<b>31.9</b>	<b>2.4</b>	<b>1.4</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 74

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	96	*	*	52	*	*	13	*	20	*	111	11.7 (126)	111	17.1 (109)
0102	262	7.6 (132)	63.0 (87)	96	*	*	34	*	62	*	334	8.1 (43)	334	13.2 (49)
0103	281	5.7 (113)	62.6 (78)	121	9.1 (1)	64.5 (110)	43	*	68	*	354	11.3 (121)	354	16.1 (95)
0104	80	*	*	24	*	*	11	*	26	*	102	6.9 (20)	102	9.8 (14)
0105	1	*	*	0	*	*	0	*	0	*	1	*	1	*
0250	95	*	*	41	*	*	13	*	34	*	110	7.3 (30)	110	15.5 (90)
0251	41	*	*	14	*	*	7	*	24	*	61	*	61	*
0252	91	*	*	66	*	*	4	*	16	*	92	*	92	*
0253	112	5.4 (107)	63.4 (90)	86	*	*	15	*	26	*	109	14.7 (140)	109	8.3 (7)
0401	360	5.0 (89)	56.9 (15)	151	20.5 (96)	52.3 (30)	43	*	56	*	361	8.6 (57)	363	18.2 (119)
0402	259	4.2 (54)	60.6 (48)	39	*	*	38	*	63	*	299	10.0 (86)	299	11.4 (22)
0403	106	6.6 (127)	64.2 (100)	57	*	*	13	*	16	*	111	13.5 (138)	111	14.4 (67)
0404	78	*	*	11	*	*	12	*	19	*	100	10.0 (86)	100	13.0 (45)
0501	301	2.7 (9)	58.1 (21)	46	*	*	58	*	96	*	371	8.4 (54)	371	12.1 (31)
0503	38	*	*	7	*	*	5	*	15	*	48	*	48	*
0504	285	2.1 (5)	59.6 (31)	115	18.3 (75)	58.3 (95)	50	*	59	*	298	8.1 (43)	298	12.4 (33)
0505	158	4.4 (68)	54.4 (6)	30	*	*	35	*	50	*	194	6.7 (16)	194	13.4 (53)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 75

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	102	4.9 (86)	52.0 (2)	19	*	*	31	*	29	*	160	7.5 (35)	160	11.9 (29)
0507	30	*	*	7	*	*	5	*	17	*	55	*	55	*
0508	5	*	*	3	*	*	1	*	4	*	10	*	10	*
0650	824	3.4 (17)	64.8 (108)	488	13.9 (15)	58.8 (97)	104	23.1 (3)	205	19.5 (17)	990	12.4 (130)	992	18.5 (120)
0651	382	5.8 (114)	59.4 (30)	85	*	*	46	*	99	*	470	11.3 (121)	470	15.1 (84)
0652	218	4.1 (49)	66.5 (122)	156	21.2 (103)	51.3 (27)	34	*	38	*	250	9.6 (76)	250	16.0 (94)
0653	342	5.6 (111)	62.0 (67)	196	15.8 (36)	54.6 (55)	47	*	59	*	362	10.2 (93)	363	18.5 (120)
0654	213	6.1 (118)	57.3 (18)	124	20.2 (92)	51.6 (29)	21	*	85	*	351	9.4 (72)	351	17.9 (117)
0655	290	6.6 (127)	57.9 (19)	48	*	*	23	*	30	*	292	9.2 (67)	293	13.3 (50)
0657	14	*	*	9	*	*	0	*	8	*	24	*	24	*
0701	308	4.5 (73)	55.8 (11)	169	16.0 (38)	48.5 (9)	26	*	79	*	376	10.9 (114)	374	15.2 (86)
0702	58	*	*	18	*	*	6	*	10	*	62	*	62	*
0703	135	4.4 (68)	57.0 (17)	60	*	*	14	*	40	*	160	7.5 (35)	160	9.4 (11)
0704	48	*	*	32	*	*	9	*	8	*	48	*	48	*
0705	47	*	*	26	*	*	10	*	20	*	72	*	72	*
0706	1	*	*	1	*	*	0	*	0	*	1	*	1	*
0850	397	4.0 (44)	61.7 (63)	94	*	*	31	*	82	*	612	12.6 (133)	613	16.2 (100)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 76

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0851	398	6.3 (125)	59.8 (37)	149	16.8 (51)	59.7 (103)	47	*	110	17.3 (12)	597	10.2 (93)	597	19.1 (126)
0901	1,014	6.0 (117)	59.7 (34)	570	16.0 (38)	55.4 (68)	100	27.0 (5)	229	25.3 (37)	1,260	9.0 (62)	1,262	15.2 (86)
0902	17	*	*	10	*	*	0	*	4	*	22	*	22	*
1051	823	4.7 (80)	56.9 (15)	540	15.6 (34)	53.7 (43)	80	*	142	25.4 (38)	957	7.4 (34)	962	13.5 (55)
1052	1	*	*	1	*	*	1	*	0	*	3	*	3	*
1101	1,060	5.1 (95)	63.3 (89)	634	20.3 (94)	49.7 (13)	123	42.3 (30)	257	26.1 (42)	1,138	9.3 (70)	1,138	19.1 (126)
1102	609	5.6 (111)	59.8 (37)	370	17.8 (66)	54.9 (58)	75	*	153	15.7 (9)	633	8.2 (48)	635	14.3 (65)
1106	48	*	*	33	*	*	1	*	9	*	45	*	47	*
1250	437	5.3 (103)	61.3 (59)	149	19.5 (87)	50.3 (16)	133	45.1 (36)	170	34.7 (69)	930	8.5 (56)	930	8.9 (10)
1251	618	4.4 (68)	60.2 (41)	462	15.2 (30)	57.1 (85)	71	*	124	22.6 (24)	637	12.6 (133)	637	12.4 (33)
1252	409	6.1 (118)	62.8 (83)	174	14.4 (19)	52.3 (30)	48	*	85	*	478	10.0 (86)	478	11.5 (23)
1253	523	5.9 (116)	60.2 (41)	245	15.1 (28)	57.6 (91)	75	*	121	12.4 (3)	566	9.9 (82)	566	15.0 (81)
1350	2,026	4.1 (49)	62.0 (67)	1,463	18.2 (72)	55.2 (65)	172	32.0 (15)	178	23.0 (29)	2,098	7.9 (40)	2,099	12.6 (40)
1352	294	5.4 (107)	63.6 (94)	191	15.2 (30)	57.6 (91)	27	*	33	*	326	4.3 (1)	326	10.1 (16)
1401	181	6.1 (118)	55.2 (9)	114	15.8 (36)	50.9 (21)	16	*	44	*	192	5.2 (3)	192	16.1 (95)
1402	48	*	*	26	*	*	9	*	9	*	69	*	69	*
1403	13	*	*	9	*	*	0	*	6	*	17	*	17	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 77

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

### Comparison of Maternal Health Indicators <sup>(2,3)</sup>

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1501	574	3.5 (24)	66.2 (121)	242	37.6 (109)	35.5 (2)	115	25.2 (4)	130	13.1 (5)	669	12.3 (129)	672	13.1 (47)
1503	685	4.8 (82)	64.4 (103)	474	19.0 (80)	54.4 (54)	119	22.7 (2)	128	22.7 (25)	736	12.0 (128)	735	15.0 (81)
1504	623	6.6 (127)	55.2 (9)	189	11.6 (3)	54.0 (46)	259	30.9 (13)	349	20.6 (21)	1,685	9.5 (74)	1,686	8.2 (6)
1506	435	4.4 (68)	60.5 (45)	61	*	*	89	*	130	18.5 (15)	476	11.1 (116)	476	15.5 (90)
1507	340	10.6 (134)	54.4 (6)	49	*	*	49	*	130	30.0 (60)	457	10.7 (107)	457	12.3 (32)
1509	221	5.0 (89)	65.2 (116)	89	*	*	52	*	76	*	309	9.7 (79)	309	7.1 (4)
1511	424	4.2 (54)	64.9 (112)	241	17.4 (59)	51.5 (28)	81	*	157	22.9 (28)	546	10.8 (111)	547	14.8 (74)
1512	239	3.8 (33)	64.0 (98)	135	18.5 (78)	59.3 (100)	45	*	144	30.6 (64)	359	9.2 (67)	360	14.4 (67)
1514	401	5.0 (89)	53.9 (4)	87	*	*	79	*	267	12.7 (4)	670	8.2 (48)	671	14.8 (74)
1701	454	3.1 (13)	61.2 (56)	290	17.9 (67)	55.2 (65)	46	*	119	28.6 (54)	464	11.2 (119)	464	11.6 (26)
1950	493	3.7 (29)	62.1 (71)	343	13.4 (10)	59.5 (101)	36	*	103	15.5 (8)	529	10.4 (100)	530	16.4 (102)
1952	110	2.7 (9)	67.3 (128)	76	*	*	3	*	20	*	122	10.7 (107)	122	18.9 (125)
1953	226	5.3 (103)	61.9 (66)	166	14.5 (21)	57.2 (88)	14	*	65	*	244	13.1 (137)	244	14.8 (74)
1954	319	1.6 (2)	66.8 (124)	189	20.1 (91)	54.0 (46)	29	*	65	*	330	11.5 (124)	330	11.2 (20)
1955	123	4.1 (49)	63.4 (90)	80	*	*	8	*	27	*	121	12.4 (130)	121	6.6 (3)
1957	328	4.3 (61)	66.8 (124)	214	12.6 (7)	58.9 (98)	29	*	64	*	353	10.5 (102)	354	11.3 (21)
1963	70	*	*	42	*	*	0	*	13	*	75	*	75	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 78

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2150	486	3.1 (13)	62.3 (75)	338	22.8 (106)	49.7 (13)	69	*	120	17.5 (13)	483	10.4 (100)	485	17.5 (113)
2151	212	4.2 (54)	66.5 (122)	171	17.5 (60)	59.1 (99)	31	*	59	*	194	8.8 (59)	194	18.6 (124)
2152	28	*	*	12	*	*	2	*	6	*	26	*	26	*
2251	277	5.8 (114)	65.3 (118)	184	16.8 (51)	56.0 (74)	7	*	22	*	301	6.0 (5)	301	13.3 (50)
2252	93	*	*	24	*	*	3	*	9	*	154	9.7 (79)	153	11.1 (19)
2554	19	*	*	8	*	*	7	*	6	*	36	*	36	*
2555	4,975	5.0 (89)	60.9 (54)	2,180	19.7 (88)	53.0 (35)	482	43.6 (34)	862	29.8 (58)	6,322	10.2 (93)	6,328	17.0 (108)
2556	134	4.5 (73)	54.5 (8)	99	*	*	3	*	34	*	161	9.9 (82)	161	16.1 (95)
2557	1	*	*	0	*	*	0	*	0	*	1	*	1	*
2562	0	*	*	0	*	*	0	*	0	*	1	*	1	*
2563	0	*	*	0	*	*	0	*	0	*	2	*	2	*
2564	1	*	*	0	*	*	0	*	0	*	2	*	2	*
3001	788	3.9 (40)	64.8 (108)	448	21.0 (101)	46.0 (6)	54	*	101	14.9 (6)	1,048	8.3 (51)	1,047	16.9 (107)
3002	356	4.8 (82)	64.9 (112)	223	17.5 (60)	51.1 (25)	42	*	67	*	414	6.0 (5)	415	9.9 (15)
3004	185	5.4 (107)	67.6 (130)	108	17.6 (65)	50.9 (21)	12	*	24	*	287	7.0 (22)	287	12.5 (38)
3005	97	*	*	61	*	*	12	*	16	*	155	6.5 (12)	155	7.1 (4)
3007	3	*	*	1	*	*	0	*	0	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 79

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3201	349	4.9 (86)	66.8 (124)	192	18.2 (72)	55.7 (71)	55	*	102	23.5 (31)	410	10.0 (86)	410	17.6 (114)
3301	3,545	3.8 (33)	60.0 (39)	2,096	16.1 (41)	55.2 (65)	374	32.6 (16)	404	20.8 (22)	4,153	7.1 (24)	4,156	12.6 (40)
3304	68	*	*	37	*	*	3	*	3	*	79	*	79	*
3305	120	4.2 (54)	60.8 (53)	72	*	*	4	*	4	*	133	5.3 (4)	133	8.3 (7)
3306	155	1.3 (1)	73.5 (134)	82	*	*	5	*	11	*	168	6.5 (12)	169	14.8 (74)
3308	70	*	*	41	*	*	6	*	4	*	68	*	69	*
3312	360	3.9 (40)	59.7 (34)	199	16.6 (49)	57.8 (94)	22	*	17	*	390	7.2 (26)	390	14.9 (79)
3314	2	*	*	0	*	*	0	*	0	*	2	*	2	*
3550	37	*	*	24	*	*	1	*	3	*	51	*	51	*
3551	3	*	*	0	*	*	1	*	1	*	6	*	6	*
3701	670	5.2 (98)	61.5 (61)	498	19.3 (85)	53.4 (38)	100	17.0 (1)	185	10.8 (1)	766	10.7 (107)	766	18.0 (118)
3850	2,066	5.2 (98)	60.5 (45)	1,345	16.1 (41)	56.4 (78)	134	30.6 (11)	228	26.8 (44)	2,303	10.2 (93)	2,304	14.8 (74)
3901	1,105	4.3 (61)	60.7 (49)	518	12.5 (5)	61.2 (106)	83	*	187	27.3 (48)	1,262	7.3 (30)	1,259	14.5 (69)
3902	1	*	*	1	*	*	0	*	0	*	2	*	2	*
3950	1,473	5.2 (98)	60.5 (45)	776	14.6 (22)	57.3 (89)	115	44.3 (35)	232	19.8 (19)	1,781	10.3 (98)	1,784	13.8 (60)
3954	1	*	*	0	*	*	0	*	0	*	1	*	1	*
4106	842	3.9 (40)	60.2 (41)	205	20.0 (90)	49.3 (11)	371	31.3 (14)	674	19.7 (18)	2,239	9.1 (64)	2,244	19.7 (132)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 80

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

### Comparison of Maternal Health Indicators (2,3)

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4107	1,604	3.8 (33)	58.0 (20)	473	14.8 (24)	55.0 (61)	435	34.5 (17)	734	24.4 (32)	2,343	8.7 (58)	2,349	20.3 (134)
4109	410	4.4 (68)	58.3 (23)	241	12.0 (4)	56.4 (78)	99	*	246	21.5 (23)	672	7.7 (38)	673	19.2 (129)
4112	527	1.7 (3)	68.9 (132)	455	56.5 (110)	20.7 (1)	32	*	53	*	512	21.9 (141)	512	24.8 (138)
4115	1,822	4.5 (73)	60.7 (49)	684	18.3 (75)	50.4 (18)	518	38.0 (22)	1,122	25.8 (41)	3,225	9.6 (76)	3,226	17.4 (110)
4351	458	4.1 (49)	61.4 (60)	318	16.0 (38)	56.0 (74)	20	*	60	*	481	12.5 (132)	482	27.4 (141)
4352	176	6.8 (131)	59.1 (28)	125	14.4 (19)	60.0 (104)	12	*	25	*	172	10.5 (102)	172	23.8 (137)
4353	346	3.8 (33)	67.3 (128)	258	20.5 (96)	55.0 (61)	44	*	63	*	407	13.0 (135)	407	11.5 (23)
4354	526	4.8 (82)	62.2 (73)	162	14.8 (24)	56.8 (82)	20	*	66	*	667	10.8 (111)	668	25.1 (139)
4355	224	6.3 (125)	62.9 (86)	150	14.7 (23)	58.7 (96)	16	*	40	*	243	9.5 (74)	242	14.9 (79)
4356	340	2.6 (8)	67.9 (131)	256	18.0 (69)	52.7 (34)	20	*	55	*	345	10.7 (107)	346	18.5 (120)
4359	469	3.4 (17)	65.5 (119)	323	22.9 (107)	48.9 (10)	25	*	77	*	496	9.9 (82)	496	18.5 (120)
4360	201	5.5 (110)	60.7 (49)	144	18.1 (71)	56.9 (84)	14	*	39	*	201	7.0 (22)	201	12.4 (33)
4361	202	4.0 (44)	64.4 (103)	149	22.1 (105)	50.3 (16)	10	*	41	*	210	8.1 (43)	209	12.4 (33)
4362	225	6.2 (123)	56.0 (12)	168	14.3 (17)	54.2 (49)	21	*	44	*	225	6.2 (8)	225	12.4 (33)
4363	578	5.2 (98)	58.8 (25)	423	18.9 (79)	54.8 (56)	43	*	120	17.5 (13)	603	8.1 (43)	603	12.6 (40)
4401	710	5.1 (95)	59.7 (34)	570	17.5 (60)	55.1 (64)	63	*	125	24.8 (34)	688	7.8 (39)	689	10.7 (17)
4501	508	4.9 (86)	62.0 (67)	262	16.8 (51)	55.0 (61)	50	*	96	*	669	11.2 (119)	670	14.6 (71)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 81

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4601	936	4.7 (80)	61.2 (56)	563	13.5 (11)	55.4 (68)	84	*	211	27.0 (46)	1,024	9.8 (81)	1,023	16.2 (100)
4701	809	6.2 (123)	56.5 (13)	468	19.0 (80)	54.3 (52)	69	*	201	11.9 (2)	926	10.6 (105)	928	22.7 (136)
5050	1,516	4.3 (61)	64.1 (99)	1,289	16.5 (48)	52.6 (33)	112	41.1 (27)	400	30.3 (61)	1,480	10.3 (98)	1,480	17.4 (110)
5051	2,355	3.9 (40)	62.6 (78)	1,738	16.4 (45)	53.7 (43)	205	39.0 (26)	686	29.4 (57)	2,351	9.3 (70)	2,351	14.2 (64)
5052	2,603	3.7 (29)	61.0 (55)	2,165	20.8 (100)	47.6 (8)	231	42.4 (31)	620	27.6 (49)	2,638	13.0 (135)	2,638	13.8 (60)
5055	1,197	1.8 (4)	62.1 (71)	1,001	21.3 (104)	47.3 (7)	71	*	147	27.2 (47)	1,281	14.3 (139)	1,281	8.4 (9)
5101	1,117	3.8 (33)	64.2 (100)	872	13.1 (9)	55.8 (72)	171	30.4 (9)	538	25.1 (36)	1,267	7.2 (26)	1,271	13.7 (58)
5102	1	*	*	0	*	*	0	*	0	*	1	*	1	*
5202	506	6.1 (118)	54.0 (5)	326	18.4 (77)	57.7 (93)	24	*	85	*	570	8.8 (59)	571	14.5 (69)
5206	7	*	*	5	*	*	0	*	1	*	7	*	7	*
5601	718	4.0 (44)	64.8 (108)	292	9.9 (2)	59.6 (102)	74	*	156	28.8 (56)	871	10.0 (86)	870	15.4 (89)
5801	1,606	5.2 (98)	65.6 (120)	1,224	19.0 (80)	54.3 (52)	135	29.6 (8)	232	16.8 (11)	1,562	9.9 (82)	1,564	19.4 (130)
5803	38	*	*	33	*	*	3	*	11	*	37	*	37	*
5804	2	*	*	2	*	*	0	*	0	*	2	*	2	*
5806	58	*	*	41	*	*	5	*	16	*	65	*	65	*
6101	53	*	*	38	*	*	4	*	1	*	54	*	53	*
6102	43	*	*	32	*	*	3	*	4	*	50	*	50	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 82

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6103	2,637	4.2 (54)	63.9 (97)	1,789	17.3 (58)	54.2 (49)	172	41.3 (28)	115	30.4 (63)	2,914	9.6 (76)	2,906	20.1 (133)
6105	5	*	*	2	*	*	1	*	0	*	6	*	6	*
6301	2,969	4.1 (49)	61.7 (63)	2,016	16.4 (45)	53.4 (38)	256	41.4 (29)	514	27.6 (49)	3,107	6.1 (7)	3,109	13.0 (45)
6312	1,571	3.5 (24)	61.2 (56)	976	18.0 (69)	50.9 (21)	181	48.6 (39)	373	35.7 (70)	1,667	8.3 (51)	1,670	11.9 (29)
6316	1,109	4.6 (79)	56.8 (14)	791	18.2 (72)	54.9 (58)	94	*	199	24.6 (33)	1,100	8.4 (54)	1,100	12.7 (43)
6318	9	*	*	6	*	*	2	*	0	*	10	*	10	*
6320	340	3.2 (16)	63.2 (88)	146	13.7 (12)	56.8 (82)	23	*	67	*	408	9.1 (64)	409	17.8 (116)
6325	227	5.3 (103)	59.0 (27)	112	14.3 (17)	57.1 (85)	20	*	68	*	283	8.1 (43)	283	17.7 (115)
6335	720	4.3 (61)	59.6 (31)	356	13.8 (13)	51.1 (25)	89	*	166	31.3 (66)	845	11.4 (123)	847	14.6 (71)
6345	106	6.6 (127)	58.5 (24)	51	*	*	7	*	19	*	120	10.8 (111)	120	15.0 (81)
6360	575	5.0 (89)	58.1 (21)	288	14.9 (26)	50.7 (20)	81	*	149	39.6 (74)	710	9.4 (72)	709	16.1 (95)
7306	1,395	3.7 (29)	64.8 (108)	990	19.0 (80)	54.0 (46)	129	43.4 (33)	155	31.6 (67)	1,591	11.1 (116)	1,594	12.9 (44)
7401	1,635	4.2 (54)	62.4 (77)	1,292	15.6 (34)	57.4 (90)	147	36.1 (19)	407	26.8 (44)	1,578	10.5 (102)	1,580	15.1 (84)
7601	213	6.1 (118)	60.1 (40)	26	*	*	33	*	97	*	445	8.8 (59)	445	11.5 (23)
7850	874	5.0 (89)	62.0 (67)	638	19.3 (85)	52.4 (32)	67	*	138	25.4 (38)	998	7.6 (37)	999	11.8 (28)
7852	5	*	*	2	*	*	0	*	0	*	3	*	3	*
8101	1,321	3.6 (27)	62.8 (83)	823	15.2 (30)	56.0 (74)	110	45.5 (37)	174	36.2 (71)	1,523	7.3 (30)	1,521	17.4 (110)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 83

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

### Comparison of Maternal Health Indicators (2,3)

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8102	295	2.4 (6)	67.1 (127)	152	21.1 (102)	49.3 (11)	22	*	41	*	342	6.4 (10)	342	9.6 (13)
8106	253	4.3 (61)	65.2 (116)	72	*	*	36	*	101	22.8 (26)	505	8.3 (51)	505	19.4 (130)
8150	4	*	*	1	*	*	0	*	2	*	4	*	4	*
8151	2,360	4.5 (73)	61.7 (63)	1,459	20.2 (92)	50.4 (18)	393	38.4 (25)	701	22.8 (26)	2,772	9.2 (67)	2,772	15.3 (88)
8152	4	*	*	2	*	*	0	*	2	*	5	*	5	*
8153	1	*	*	0	*	*	1	*	0	*	0	*	0	*
8155	3	*	*	0	*	*	0	*	0	*	3	*	3	*
8201	1,879	4.2 (54)	62.6 (78)	1,041	17.1 (56)	54.8 (56)	215	42.8 (32)	650	25.7 (40)	2,061	11.0 (115)	2,064	16.6 (103)
8202	1,259	4.8 (82)	50.8 (1)	569	20.4 (95)	45.2 (5)	267	27.3 (6)	487	19.9 (20)	3,724	7.1 (24)	3,723	3.3 (1)
8203	308	7.8 (133)	62.7 (82)	236	19.9 (89)	53.0 (35)	21	*	110	16.4 (10)	314	11.5 (124)	313	13.7 (58)
8250	328	3.4 (17)	64.3 (102)	189	16.4 (45)	53.4 (38)	37	*	97	*	349	7.2 (26)	350	13.4 (53)
8252	52	*	*	29	*	*	6	*	8	*	89	*	90	*
8255	1,701	4.5 (73)	62.3 (75)	1,032	17.1 (56)	53.4 (38)	245	37.1 (20)	599	28.7 (55)	1,794	11.1 (116)	1,796	16.8 (106)
8257	990	5.3 (103)	52.6 (3)	659	20.6 (98)	41.9 (4)	160	37.5 (21)	269	27.9 (52)	1,467	9.1 (64)	1,471	4.1 (2)
8263	197	2.5 (7)	58.9 (26)	95	*	*	0	*	27	*	188	6.4 (10)	188	13.3 (50)
8264	734	3.7 (29)	64.6 (105)	547	13.0 (8)	62.9 (109)	63	*	318	18.6 (16)	782	10.6 (105)	782	12.5 (38)
8267	553	3.4 (17)	64.9 (112)	306	19.0 (80)	55.9 (73)	96	*	195	30.3 (61)	575	10.1 (91)	575	19.1 (126)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 84

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8901	565	3.5 (24)	63.4 (90)	350	15.1 (28)	54.9 (58)	35	*	43	*	666	8.0 (42)	664	16.1 (95)
8902	938	4.3 (61)	62.6 (78)	514	17.9 (67)	53.1 (37)	68	*	135	34.1 (68)	1,067	6.7 (16)	1,065	14.1 (62)
8903	987	3.1 (13)	64.6 (105)	601	12.5 (5)	62.6 (108)	65	*	102	40.2 (75)	1,168	6.8 (19)	1,173	15.9 (93)
8904	1,423	4.0 (44)	63.5 (93)	831	15.2 (30)	54.2 (49)	104	51.0 (40)	199	38.2 (73)	1,676	7.3 (30)	1,678	16.6 (103)
8905	1,250	3.4 (17)	59.2 (29)	700	16.9 (55)	57.1 (85)	105	30.5 (10)	205	24.9 (35)	1,634	11.9 (127)	1,634	13.1 (47)
8909	886	3.8 (33)	63.8 (96)	530	13.8 (13)	60.9 (105)	64	*	103	28.2 (53)	975	6.7 (16)	974	16.6 (103)
9045	971	3.4 (17)	69.3 (133)	652	16.6 (49)	53.5 (42)	74	*	137	15.3 (7)	1,143	8.2 (48)	1,142	9.4 (11)
9050	733	3.8 (33)	63.7 (95)	243	14.0 (16)	62.1 (107)	62	*	91	*	810	7.2 (26)	809	13.5 (55)
9051	862	2.8 (11)	64.7 (107)	398	25.6 (108)	40.2 (3)	160	30.6 (11)	339	27.7 (51)	1,819	10.1 (91)	1,818	11.6 (26)
9053	1,407	3.4 (17)	59.6 (31)	1,126	14.9 (26)	56.7 (81)	110	38.2 (23)	111	26.1 (42)	1,469	6.2 (8)	1,468	11.0 (18)
9055	6	*	*	5	*	*	0	*	0	*	6	*	6	*
9058	1,040	5.1 (95)	60.7 (49)	727	16.2 (44)	53.8 (45)	102	38.2 (23)	170	31.2 (65)	1,462	7.9 (40)	1,462	14.7 (73)
9066	323	4.3 (61)	62.8 (83)	217	20.7 (99)	49.8 (15)	0	*	7	*	349	6.6 (15)	349	21.8 (135)
9068	773	3.0 (12)	64.9 (112)	627	17.5 (60)	56.6 (80)	28	*	55	*	759	6.5 (12)	758	14.1 (62)
9069	1,236	4.0 (44)	60.4 (44)	878	17.5 (60)	51.0 (24)	104	28.8 (7)	111	23.4 (30)	1,393	6.9 (20)	1,393	13.5 (55)
9070	1,711	4.5 (73)	62.2 (73)	1,276	16.1 (41)	55.5 (70)	135	47.4 (38)	153	36.6 (72)	1,767	5.1 (2)	1,769	14.3 (65)
9080	2,077	3.6 (27)	61.6 (62)	974	16.8 (51)	56.1 (77)	225	35.6 (18)	325	29.8 (58)	2,659	10.2 (93)	2,662	15.6 (92)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 5A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 85

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9082	84	*	*	40	*	*	13	*	20	*	100	9.0 (62)	100	26.0 (140)
9083	12	*	*	10	*	*	0	*	2	*	11	*	11	*
9087	2	*	*	1	*	*	0	*	0	*	3	*	3	*
Michigan	106,070	4.3	61.6	63,239	17.6	53.7	11,903	34.3	23,406	24.6	128,518	9.2	128,592	14.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 86

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	105	90.5 (97)	1.0 (120)	118	30.5 (69)	122	32.0 (38)	89	*
0102	310	91.3 (87)	0.3 (40)	351	31.9 (55)	357	24.4 (79)	280	91.4 (47)
0103	339	93.8 (46)	0.3 (40)	360	35.6 (36)	365	23.6 (85)	280	93.2 (31)
0104	98	*	*	111	39.6 (14)	113	31.9 (40)	90	*
0105	1	*	*	1	*	1	*	1	*
0250	91	*	*	117	40.2 (12)	121	25.6 (71)	94	*
0251	60	*	*	66	*	66	*	47	*
0252	93	*	*	103	42.7 (7)	101	51.5 (5)	77	*
0253	103	91.3 (87)	0.0 (1)	112	40.2 (12)	113	26.5 (67)	89	*
0401	333	95.5 (26)	0.0 (1)	381	24.4 (106)	389	29.3 (53)	269	94.8 (13)
0402	272	89.3 (111)	1.1 (126)	324	29.6 (77)	326	31.9 (40)	259	95.4 (7)
0403	103	92.2 (76)	0.0 (1)	115	37.4 (23)	116	33.6 (33)	92	*
0404	99	*	*	110	31.8 (57)	110	35.5 (26)	90	*
0501	358	96.1 (14)	0.0 (1)	396	29.5 (79)	377	36.9 (19)	299	88.0 (102)
0503	48	*	*	49	*	51	*	34	*
0504	290	93.1 (59)	0.3 (40)	301	30.6 (67)	307	38.4 (18)	236	93.6 (26)
0505	178	94.9 (33)	0.6 (78)	199	22.6 (111)	200	36.5 (21)	146	93.8 (21)
0506	146	91.1 (90)	0.7 (92)	167	35.9 (30)	171	30.4 (45)	139	89.2 (80)
0507	55	*	*	55	*	57	*	29	*
0508	10	*	*	10	*	10	*	5	*
0650	909	92.8 (64)	0.6 (78)	1,045	35.9 (30)	1,031	31.4 (43)	824	91.3 (50)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 87

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0651	440	91.4 (86)	0.7 (92)	502	38.4 (19)	463	26.8 (64)	403	88.3 (96)
0652	228	96.5 (9)	0.4 (53)	251	39.4 (15)	255	27.1 (62)	203	87.2 (110)
0653	308	94.5 (38)	1.3 (131)	379	43.5 (5)	377	35.8 (24)	321	86.9 (113)
0654	287	90.6 (96)	0.7 (92)	371	42.0 (8)	370	17.6 (116)	292	91.4 (47)
0655	282	92.9 (63)	0.4 (53)	307	38.4 (19)	292	38.7 (17)	250	88.8 (88)
0657	20	*	*	26	*	26	*	23	*
0701	327	93.9 (45)	0.3 (40)	412	34.5 (41)	406	39.4 (14)	310	89.4 (76)
0702	55	*	*	70	*	71	*	57	*
0703	158	95.6 (23)	0.0 (1)	179	33.0 (50)	174	39.7 (12)	127	96.1 (2)
0704	44	*	*	55	*	55	*	51	*
0705	69	*	*	82	*	81	*	58	*
0706	1	*	*	1	*	1	*	0	*
0850	599	92.7 (67)	0.8 (105)	661	31.0 (65)	662	26.6 (66)	476	93.7 (22)
0851	574	95.8 (18)	0.0 (1)	620	30.2 (73)	624	34.6 (28)	453	94.0 (18)
0901	1,131	92.6 (70)	0.7 (92)	1,327	36.7 (24)	1,330	25.3 (72)	1,042	89.6 (69)
0902	18	*	*	22	*	22	*	18	*
1051	864	92.4 (73)	0.8 (105)	1,032	31.3 (62)	1,029	30.0 (49)	750	88.5 (92)
1052	3	*	*	2	*	3	*	1	*
1101	1,014	84.5 (130)	0.9 (113)	1,277	27.4 (92)	1,122	22.5 (94)	862	89.1 (83)
1102	556	89.6 (107)	0.4 (53)	705	30.5 (69)	646	21.7 (96)	500	94.8 (13)
1106	45	*	*	52	*	49	*	36	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 88

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1250	871	93.8 (46)	0.1 (28)	970	27.5 (91)	979	38.9 (16)	724	89.6 (69)
1251	623	95.7 (21)	0.5 (68)	665	30.1 (74)	675	39.6 (13)	487	90.8 (56)
1252	360	88.3 (116)	0.6 (78)	514	35.8 (33)	513	22.8 (91)	391	87.0 (112)
1253	468	88.9 (114)	1.1 (126)	599	34.1 (42)	606	20.6 (102)	458	88.4 (94)
1350	2,042	96.1 (14)	0.3 (40)	2,244	33.4 (47)	2,078	35.6 (25)	1,717	89.3 (78)
1352	321	99.1 (2)	0.0 (1)	349	33.8 (44)	343	12.0 (133)	261	83.9 (125)
1401	180	88.3 (116)	1.7 (134)	203	32.0 (54)	201	36.8 (20)	149	92.6 (36)
1402	67	*	*	77	*	75	*	54	*
1403	16	*	*	18	*	18	*	12	*
1501	544	89.2 (112)	0.7 (92)	738	28.7 (85)	714	32.4 (36)	547	87.9 (103)
1503	692	98.1 (3)	0.0 (1)	789	34.7 (40)	774	22.0 (95)	594	83.0 (127)
1504	1,622	96.7 (6)	0.2 (32)	1,814	28.1 (87)	1,781	20.4 (105)	1,251	89.5 (74)
1506	389	94.3 (41)	0.0 (1)	514	29.6 (77)	504	14.7 (125)	380	90.8 (56)
1507	457	96.7 (6)	0.4 (53)	487	33.5 (46)	473	30.4 (45)	336	96.4 (1)
1509	281	92.2 (76)	0.0 (1)	338	29.9 (75)	329	17.3 (118)	252	93.7 (22)
1511	507	89.9 (103)	0.6 (78)	573	33.2 (48)	572	34.4 (29)	421	88.6 (89)
1512	333	94.0 (43)	0.6 (78)	378	22.0 (114)	370	34.1 (30)	230	93.0 (33)
1514	627	95.2 (28)	0.5 (68)	700	28.6 (86)	656	34.0 (31)	449	91.3 (50)
1701	456	97.4 (5)	0.4 (53)	510	37.6 (22)	505	60.6 (2)	394	93.7 (22)
1950	486	94.7 (36)	0.0 (1)	555	30.3 (72)	518	45.6 (8)	415	95.2 (8)
1952	112	93.8 (46)	0.0 (1)	123	29.3 (81)	113	51.3 (6)	77	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 89

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1953	226	96.5 (9)	0.0 (1)	258	22.5 (113)	241	44.8 (9)	182	95.6 (5)
1954	311	95.2 (28)	0.6 (78)	350	38.6 (17)	318	59.1 (3)	283	95.1 (10)
1955	114	89.5 (110)	0.0 (1)	134	29.1 (82)	121	46.3 (7)	95	*
1957	324	94.1 (42)	0.0 (1)	373	31.9 (55)	311	43.4 (10)	278	92.8 (34)
1963	72	*	*	81	*	77	*	68	*
2150	479	92.7 (67)	0.2 (32)	512	33.6 (45)	511	29.2 (54)	386	94.0 (18)
2151	181	94.5 (38)	0.0 (1)	221	31.2 (64)	171	40.4 (11)	152	91.4 (47)
2152	26	*	*	32	*	32	*	25	*
2251	280	96.4 (12)	0.4 (53)	326	38.0 (21)	326	32.2 (37)	266	95.1 (10)
2252	143	96.5 (9)	0.7 (92)	170	47.1 (2)	158	62.7 (1)	141	90.8 (56)
2554	36	*	*	39	*	30	*	32	*
2555	5,786	89.6 (107)	0.4 (53)	6,779	26.8 (94)	6,030	17.6 (116)	5,022	89.5 (74)
2556	142	80.3 (136)	0.7 (92)	157	7.6 (143)	151	11.9 (134)	54	*
2557	1	*	*	1	*	1	*	1	*
2562	1	*	*	1	*	1	*	1	*
2563	1	*	*	3	*	2	*	3	*
2564	2	*	*	3	*	2	*	3	*
3001	971	89.9 (103)	0.2 (32)	1,100	39.2 (16)	1,103	12.4 (132)	887	86.5 (115)
3002	404	95.3 (27)	0.2 (32)	428	31.3 (62)	431	14.2 (127)	349	90.5 (59)
3004	263	95.8 (18)	0.4 (53)	286	35.7 (34)	294	20.1 (108)	207	89.9 (65)
3005	144	94.4 (40)	0.7 (92)	159	32.1 (52)	162	19.1 (111)	135	90.4 (61)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 90

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3007	3	*	*	3	*	3	*	3	*
3201	350	94.0 (43)	0.9 (113)	420	25.0 (102)	419	39.4 (14)	301	91.0 (54)
3301	3,521	89.9 (103)	0.8 (105)	4,309	27.8 (88)	4,240	14.3 (126)	3,181	88.3 (96)
3304	64	*	*	80	*	81	*	72	*
3305	110	90.0 (102)	0.9 (113)	141	36.2 (27)	139	18.0 (115)	111	90.1 (62)
3306	146	93.2 (57)	0.0 (1)	170	35.9 (30)	171	17.0 (121)	141	94.3 (16)
3308	55	*	*	76	*	74	*	51	*
3312	331	92.4 (73)	0.0 (1)	414	35.5 (37)	414	15.0 (124)	335	89.6 (69)
3314	2	*	*	2	*	2	*	2	*
3550	46	*	*	52	*	53	*	42	*
3551	6	*	*	7	*	7	*	6	*
3701	734	95.1 (31)	0.0 (1)	805	29.4 (80)	808	23.1 (87)	577	94.3 (16)
3850	2,083	92.2 (76)	0.5 (68)	2,470	36.3 (26)	2,479	22.9 (90)	1,908	89.0 (87)
3901	1,135	92.6 (70)	0.3 (40)	1,379	31.8 (57)	1,369	17.1 (120)	1,011	84.1 (124)
3902	2	*	*	2	*	2	*	2	*
3950	1,643	90.5 (97)	0.5 (68)	1,898	31.7 (61)	1,886	22.6 (93)	1,398	89.6 (69)
3954	1	*	*	1	*	1	*	1	*
4106	1,961	86.1 (126)	1.3 (131)	2,340	26.6 (95)	2,295	12.5 (131)	1,756	82.1 (129)
4107	2,222	92.2 (76)	0.4 (53)	2,530	27.3 (93)	2,421	30.0 (49)	1,976	92.0 (42)
4109	644	94.9 (33)	1.2 (130)	695	26.3 (99)	693	24.7 (77)	443	91.6 (46)
4112	510	70.4 (137)	0.0 (1)	529	18.3 (132)	29	*	461	88.3 (96)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 91

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4115	2,767	93.2 (57)	0.7 (92)	3,351	25.0 (102)	2,366	19.4 (109)	2,445	89.4 (76)
4351	400	87.8 (119)	1.0 (120)	514	43.0 (6)	480	26.0 (70)	390	87.9 (103)
4352	154	88.3 (116)	0.0 (1)	186	40.9 (10)	187	23.0 (89)	148	87.8 (107)
4353	307	90.9 (94)	1.0 (120)	441	38.5 (18)	431	26.7 (65)	339	86.4 (117)
4354	619	95.8 (18)	0.2 (32)	708	52.5 (1)	693	23.8 (83)	571	89.1 (83)
4355	224	90.2 (99)	1.8 (135)	257	36.2 (27)	261	36.4 (23)	204	88.2 (99)
4356	293	91.8 (83)	0.7 (92)	356	36.0 (29)	360	27.2 (61)	272	89.7 (67)
4359	343	86.0 (127)	2.0 (137)	520	44.6 (3)	525	25.3 (72)	418	86.6 (114)
4360	178	91.6 (85)	1.1 (126)	209	33.0 (50)	212	30.2 (48)	164	92.7 (35)
4361	197	94.9 (33)	1.5 (133)	220	44.1 (4)	223	29.1 (55)	178	87.1 (111)
4362	198	92.4 (73)	1.0 (120)	247	33.2 (48)	243	28.0 (58)	198	82.8 (128)
4363	564	91.1 (90)	0.7 (92)	622	29.9 (75)	635	29.6 (52)	463	91.1 (53)
4401	597	91.0 (92)	0.8 (105)	713	31.8 (57)	723	28.1 (57)	536	92.4 (37)
4501	600	93.7 (50)	0.3 (40)	694	34.9 (39)	700	23.9 (82)	527	88.6 (89)
4601	960	92.8 (64)	0.2 (32)	1,122	29.1 (82)	1,121	18.4 (114)	802	91.9 (43)
4701	842	93.5 (53)	0.8 (105)	968	30.7 (66)	976	36.5 (21)	695	91.8 (45)
5050	1,433	93.3 (56)	0.5 (68)	1,540	20.8 (119)	1,525	12.7 (130)	1,037	96.0 (4)
5051	2,229	92.6 (70)	0.9 (113)	2,475	21.5 (116)	2,438	13.5 (129)	1,660	93.7 (22)
5052	2,532	91.7 (84)	0.4 (53)	2,694	19.7 (126)	2,633	11.7 (135)	1,801	93.3 (29)
5055	1,164	99.7 (1)	0.1 (28)	1,314	41.1 (9)	506	6.3 (140)	1,044	95.2 (8)
5101	1,213	92.7 (67)	0.2 (32)	1,313	18.6 (128)	946	32.0 (38)	905	95.5 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 92

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5102	1	*	*	2	*	1	*	2	*
5202	499	95.6 (23)	0.6 (78)	607	35.7 (34)	551	32.8 (34)	459	92.2 (39)
5206	4	*	*	7	*	6	*	7	*
5601	820	95.9 (16)	0.6 (78)	909	29.0 (84)	913	25.3 (72)	665	91.9 (43)
5801	1,450	93.0 (60)	0.6 (78)	1,692	30.4 (71)	1,450	31.7 (42)	1,237	92.2 (39)
5803	37	*	*	39	*	37	*	21	*
5804	2	*	*	2	*	2	*	2	*
5806	64	*	*	66	*	58	*	31	*
6101	48	*	*	57	*	57	*	43	*
6102	48	*	*	51	*	53	*	37	*
6103	2,718	92.2 (76)	0.4 (53)	3,093	31.8 (57)	2,991	25.2 (75)	2,264	93.4 (27)
6105	5	*	*	7	*	7	*	5	*
6301	2,912	94.6 (37)	0.5 (68)	3,365	26.5 (96)	3,138	20.4 (105)	2,393	89.1 (83)
6312	1,523	93.0 (60)	0.2 (32)	1,777	24.6 (105)	1,683	21.0 (100)	1,216	84.8 (123)
6316	1,017	95.2 (28)	0.6 (78)	1,168	26.4 (98)	1,115	24.4 (79)	812	88.4 (94)
6318	8	*	*	10	*	10	*	2	*
6320	396	93.4 (54)	0.0 (1)	440	23.0 (110)	433	33.7 (32)	311	96.1 (2)
6325	276	96.4 (12)	0.4 (53)	302	20.9 (117)	303	35.0 (27)	193	92.2 (39)
6335	808	95.0 (32)	0.5 (68)	885	27.7 (89)	849	27.8 (59)	642	94.5 (15)
6345	115	91.3 (87)	0.0 (1)	131	26.0 (100)	131	29.8 (51)	87	*
6360	664	91.9 (82)	0.5 (68)	750	21.6 (115)	725	24.6 (78)	527	93.2 (31)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 93

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
7306	1,456	90.7 (95)	0.4 (53)	1,678	34.1 (42)	1,668	26.5 (67)	1,270	90.5 (59)
7401	1,236	89.6 (107)	1.1 (126)	1,723	40.4 (11)	1,737	25.2 (75)	1,335	90.0 (64)
7601	426	96.7 (6)	0.0 (1)	482	35.3 (38)	484	20.2 (107)	372	90.1 (62)
7850	910	95.9 (16)	0.5 (68)	1,056	36.5 (25)	523	27.3 (60)	790	85.7 (119)
7852	3	*	*	5	*	3	*	5	*
8101	1,440	83.8 (133)	1.0 (120)	1,610	18.1 (133)	1,418	20.5 (103)	1,047	87.9 (103)
8102	330	89.7 (106)	0.3 (40)	360	25.3 (101)	333	21.3 (98)	272	87.5 (109)
8106	482	90.2 (99)	0.6 (78)	527	15.6 (136)	531	20.5 (103)	335	94.9 (12)
8150	4	*	*	4	*	4	*	4	*
8151	2,624	93.0 (60)	0.3 (40)	3,030	27.6 (90)	3,002	21.6 (97)	2,259	89.8 (66)
8152	5	*	*	5	*	5	*	5	*
8153	0	*	*	1	*	1	*	1	*
8155	3	*	*	4	*	4	*	4	*
8201	1,995	93.4 (54)	0.3 (40)	2,304	15.5 (137)	1,956	31.2 (44)	1,499	89.2 (80)
8202	3,444	98.1 (3)	0.1 (28)	3,787	32.1 (52)	3,773	7.1 (137)	3,019	93.4 (27)
8203	305	92.8 (64)	0.3 (40)	345	20.9 (117)	332	28.3 (56)	222	91.0 (54)
8250	345	91.0 (92)	0.3 (40)	394	15.2 (138)	338	24.0 (81)	303	89.1 (83)
8252	85	*	*	96	*	95	*	76	*
8255	1,765	92.2 (76)	1.0 (120)	1,956	13.3 (141)	1,877	26.3 (69)	1,206	92.3 (38)
8257	1,436	95.7 (21)	0.0 (1)	1,524	20.8 (119)	1,259	14.1 (128)	1,152	70.8 (130)
8263	180	95.6 (23)	0.0 (1)	234	26.5 (96)	216	26.9 (63)	234	89.7 (67)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 94

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8264	766	93.6 (51)	0.5 (68)	821	13.6 (140)	776	30.3 (47)	502	86.3 (118)
8267	561	93.8 (46)	0.9 (113)	619	13.7 (139)	574	23.7 (84)	453	88.1 (100)
8901	636	83.3 (134)	0.8 (105)	687	20.4 (123)	551	17.2 (119)	466	89.3 (78)
8902	1,037	87.3 (121)	0.6 (78)	1,118	19.8 (125)	998	21.1 (99)	761	87.9 (103)
8903	1,114	82.6 (135)	0.6 (78)	1,228	18.5 (130)	1,001	19.1 (111)	842	85.5 (120)
8904	1,600	83.9 (132)	0.8 (105)	1,756	18.6 (128)	1,531	19.3 (110)	1,144	88.5 (92)
8905	1,581	93.6 (51)	0.1 (28)	1,711	24.9 (104)	1,577	32.5 (35)	1,179	86.5 (115)
8909	924	87.2 (122)	0.3 (40)	1,049	20.8 (119)	869	23.1 (87)	729	88.1 (100)
9045	1,107	89.0 (113)	0.7 (92)	1,223	24.2 (107)	653	6.9 (138)	904	94.0 (18)
9050	734	84.9 (129)	0.4 (53)	851	23.7 (108)	821	23.4 (86)	609	85.1 (121)
9051	1,703	90.1 (101)	0.4 (53)	1,888	23.6 (109)	1,508	3.3 (141)	1,508	93.3 (29)
9053	1,451	87.7 (120)	0.9 (113)	1,533	17.2 (135)	1,302	18.9 (113)	910	91.2 (52)
9055	6	*	*	7	*	7	*	3	*
9058	1,399	87.1 (123)	0.9 (113)	1,519	18.5 (130)	1,435	9.5 (136)	885	85.1 (121)
9066	321	86.6 (124)	0.0 (1)	366	30.6 (67)	322	22.7 (92)	367	89.6 (69)
9068	749	84.0 (131)	1.9 (136)	808	19.9 (124)	788	17.0 (121)	552	87.7 (108)
9069	1,389	86.3 (125)	0.8 (105)	1,462	18.7 (127)	1,203	6.7 (139)	944	83.8 (126)
9070	1,704	86.0 (127)	0.6 (78)	1,870	17.3 (134)	1,756	16.4 (123)	1,190	89.2 (80)
9080	2,523	88.8 (115)	0.7 (92)	2,799	20.6 (122)	2,283	20.7 (101)	1,898	88.6 (89)
9082	96	*	*	104	9.6 (142)	87	*	59	*
9083	11	*	*	12	*	12	*	12	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 6A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 95

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
9087	3	*	*	3	*	3	*	3	*
Michigan	119,166	91.7	0.5	136,281	27.5	125,882	22.8	98,807	89.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 96

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	88	*	103	16.5 (129)	24	*	89	*	111	6.3 (83)
0102	278	28.1 (113)	322	11.8 (113)	77	*	280	14.6 (97)	334	9.0 (110)
0103	280	29.6 (121)	348	12.4 (117)	78	*	280	23.9 (127)	356	11.0 (121)
0104	89	*	99	*	19	*	90	*	101	7.9 (103)
0105	1	*	0	*	0	*	1	*	1	*
0250	88	*	108	14.8 (125)	32	*	94	*	109	18.3 (139)
0251	47	*	59	*	18	*	47	*	63	*
0252	77	*	91	*	16	*	77	*	94	*
0253	89	*	106	21.7 (139)	36	*	89	*	109	17.4 (138)
0401	264	28.8 (116)	352	13.9 (123)	72	*	269	14.9 (101)	361	6.6 (85)
0402	257	25.7 (104)	290	10.3 (98)	62	*	259	16.6 (112)	299	8.0 (104)
0403	92	*	108	15.7 (128)	24	*	92	*	111	19.8 (140)
0404	90	*	102	10.8 (104)	17	*	90	*	104	11.5 (124)
0501	296	23.0 (96)	370	10.3 (98)	64	*	299	17.7 (115)	379	10.0 (115)
0503	34	*	47	*	6	*	34	*	48	*
0504	234	21.4 (86)	294	11.9 (114)	51	*	234	11.1 (76)	295	5.1 (65)
0505	145	21.4 (86)	192	8.9 (90)	31	*	146	15.8 (104)	194	5.7 (78)
0506	139	36.7 (128)	157	21.0 (137)	47	*	139	25.2 (130)	162	21.0 (141)
0507	29	*	54	*	4	*	29	*	55	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 97

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0508	5	*	10	*	1	*	5	*	10	*
0650	806	18.5 (75)	979	10.4 (101)	154	27.9 (41)	820	12.3 (81)	988	7.6 (98)
0651	401	32.4 (125)	461	19.5 (133)	119	28.6 (40)	403	22.3 (126)	469	11.3 (122)
0652	202	20.8 (85)	246	11.0 (106)	41	*	203	24.1 (128)	249	17.3 (137)
0653	320	32.2 (124)	349	14.0 (124)	98	*	321	21.8 (124)	365	12.6 (129)
0654	291	25.8 (105)	342	20.5 (136)	69	*	292	19.5 (120)	351	15.4 (136)
0655	250	34.4 (127)	281	21.4 (138)	79	*	251	20.7 (123)	295	14.2 (134)
0657	23	*	24	*	4	*	23	*	24	*
0701	307	27.7 (112)	379	17.4 (131)	81	*	310	8.1 (51)	384	3.6 (40)
0702	57	*	62	*	14	*	57	*	62	*
0703	126	37.3 (129)	158	19.6 (134)	38	*	126	22.2 (125)	159	6.9 (92)
0704	51	*	47	*	10	*	51	*	48	*
0705	58	*	72	*	21	*	58	*	71	*
0706	0	*	1	*	0	*	0	*	1	*
0850	475	15.6 (59)	617	7.8 (79)	70	*	477	12.4 (83)	620	5.2 (66)
0851	448	12.9 (41)	597	6.5 (66)	58	*	451	12.9 (86)	595	4.4 (57)
0901	1,040	21.4 (86)	1,252	9.6 (94)	215	37.7 (33)	1,040	12.3 (81)	1,258	7.6 (98)
0902	18	*	21	*	5	*	18	*	21	*
1051	746	26.4 (106)	945	11.2 (108)	175	38.3 (32)	748	12.4 (83)	971	7.3 (96)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 98

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1052	1	*	3	*	1	*	1	*	3	*
1101	847	15.2 (53)	1,137	6.0 (59)	120	47.5 (17)	861	13.6 (90)	1,136	7.8 (102)
1102	495	25.1 (100)	628	12.3 (116)	118	39.8 (28)	500	16.0 (107)	635	9.9 (114)
1106	36	*	46	*	10	*	36	*	47	*
1250	723	14.5 (48)	918	8.5 (86)	96	*	723	9.8 (71)	930	6.7 (88)
1251	483	25.5 (103)	625	13.8 (122)	114	23.7 (43)	487	15.8 (104)	641	9.7 (112)
1252	390	15.9 (61)	473	7.2 (75)	52	*	392	9.7 (70)	479	5.4 (73)
1253	453	21.9 (90)	560	11.6 (110)	88	*	456	16.2 (110)	568	7.6 (98)
1350	1,706	17.5 (71)	2,088	5.0 (51)	273	47.6 (16)	1,704	7.5 (41)	2,085	4.1 (52)
1352	257	19.8 (81)	318	4.1 (39)	51	*	260	2.7 (4)	325	1.5 (5)
1401	150	27.3 (110)	187	10.7 (103)	39	*	149	7.4 (38)	191	6.8 (91)
1402	53	*	69	*	9	*	54	*	70	*
1403	12	*	17	*	4	*	12	*	17	*
1501	545	16.9 (66)	661	6.1 (60)	87	*	547	8.6 (56)	676	5.2 (66)
1503	591	19.0 (79)	735	7.3 (76)	103	34.0 (35)	594	7.6 (45)	740	2.4 (17)
1504	1,245	9.5 (23)	1,684	3.0 (29)	113	47.8 (15)	1,252	3.4 (6)	1,691	1.7 (9)
1506	380	17.1 (69)	463	8.4 (85)	59	*	379	4.5 (12)	478	5.2 (66)
1507	331	26.9 (109)	444	5.9 (57)	83	*	332	13.9 (91)	463	6.7 (88)
1509	252	14.3 (47)	308	5.2 (52)	34	*	252	5.2 (17)	310	5.2 (66)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 99

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1511	416	21.9 (90)	539	10.4 (101)	88	*	422	18.0 (118)	547	11.7 (126)
1512	227	10.6 (28)	360	6.7 (68)	27	*	230	9.6 (68)	362	3.6 (40)
1514	445	15.5 (56)	668	4.6 (44)	66	*	444	4.1 (9)	672	1.5 (5)
1701	388	20.6 (84)	468	16.9 (130)	76	*	393	15.5 (103)	473	6.6 (85)
1950	413	23.7 (98)	514	8.6 (87)	94	*	413	16.5 (111)	530	10.6 (118)
1952	77	*	119	4.2 (40)	10	*	76	*	122	9.8 (113)
1953	182	22.5 (93)	236	6.8 (70)	38	*	182	9.3 (64)	243	8.6 (107)
1954	280	25.4 (102)	312	6.7 (68)	69	*	283	14.1 (93)	331	11.5 (124)
1955	95	*	114	9.6 (94)	26	*	95	*	120	10.0 (115)
1957	278	23.4 (97)	342	9.1 (91)	57	*	278	10.8 (75)	354	5.6 (76)
1963	68	*	75	*	16	*	67	*	76	*
2150	385	29.4 (120)	483	17.4 (131)	105	22.9 (44)	384	13.5 (89)	485	6.6 (85)
2151	150	25.3 (101)	192	12.5 (118)	34	*	151	14.6 (97)	194	7.2 (95)
2152	25	*	26	*	6	*	25	*	26	*
2251	266	27.4 (111)	296	15.2 (126)	63	*	264	5.3 (19)	302	3.0 (28)
2252	137	29.9 (122)	153	20.3 (135)	39	*	141	17.7 (115)	157	10.8 (120)
2554	31	*	35	*	6	*	32	*	36	*
2555	5,019	16.7 (65)	6,246	6.9 (72)	754	41.2 (25)	5,003	11.2 (77)	6,296	5.9 (79)
2556	54	*	160	6.3 (63)	11	*	54	*	161	3.7 (42)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 100

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2557	1	*	1	*	0	*	1	*	1	*
2562	1	*	1	*	0	*	1	*	1	*
2563	3	*	2	*	1	*	3	*	2	*
2564	3	*	2	*	0	*	3	*	2	*
3001	885	17.2 (70)	1,042	6.8 (70)	141	47.5 (17)	885	11.2 (77)	1,048	6.7 (88)
3002	348	12.1 (33)	415	5.8 (56)	39	*	347	19.6 (121)	414	8.9 (109)
3004	207	13.0 (42)	287	4.9 (50)	27	*	207	8.7 (59)	286	3.5 (37)
3005	134	19.4 (80)	155	7.7 (78)	25	*	135	14.1 (93)	155	12.3 (128)
3007	3	*	3	*	0	*	3	*	3	*
3201	301	21.6 (89)	409	8.1 (82)	62	*	301	14.6 (97)	409	5.9 (79)
3301	3,175	15.3 (54)	4,153	6.2 (62)	472	45.1 (20)	3,176	8.6 (56)	4,161	2.9 (23)
3304	72	*	79	*	7	*	72	*	79	*
3305	110	13.6 (45)	132	6.1 (60)	14	*	110	5.5 (23)	133	3.8 (45)
3306	140	17.9 (72)	169	5.9 (57)	26	*	140	6.4 (31)	169	2.4 (17)
3308	51	*	69	*	8	*	51	*	69	*
3312	336	17.0 (68)	388	8.8 (89)	52	*	334	7.8 (48)	390	3.8 (45)
3314	2	*	2	*	0	*	2	*	2	*
3550	42	*	51	*	23	*	42	*	51	*
3551	6	*	6	*	3	*	6	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 101

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3701	572	22.6 (94)	742	11.6 (110)	123	22.8 (45)	575	17.2 (114)	768	5.3 (71)
3850	1,907	18.9 (78)	2,312	10.3 (98)	324	31.2 (39)	1,906	12.9 (86)	2,308	7.3 (96)
3901	1,005	15.5 (56)	1,253	6.6 (67)	139	32.4 (38)	1,007	7.7 (46)	1,261	3.7 (42)
3902	2	*	2	*	0	*	2	*	2	*
3950	1,391	16.6 (64)	1,767	7.1 (74)	213	38.5 (31)	1,397	8.2 (54)	1,788	5.3 (71)
3954	1	*	1	*	1	*	1	*	1	*
4106	1,743	9.0 (20)	2,243	4.7 (46)	149	39.6 (29)	1,709	5.3 (19)	2,248	2.0 (11)
4107	1,954	15.0 (51)	2,331	4.8 (47)	288	52.8 (8)	1,952	7.3 (36)	2,353	3.3 (34)
4109	428	11.7 (31)	671	7.0 (73)	60	*	442	7.5 (41)	675	4.0 (50)
4112	468	5.1 (8)	509	1.0 (4)	18	*	450	6.9 (32)	510	2.9 (23)
4115	2,433	7.0 (14)	3,221	2.9 (26)	177	46.9 (19)	2,420	4.2 (10)	3,221	1.6 (7)
4351	392	28.6 (115)	473	13.5 (120)	103	33.0 (36)	385	14.0 (92)	476	11.3 (122)
4352	145	42.8 (130)	166	27.1 (140)	56	*	147	20.4 (122)	171	12.9 (132)
4353	338	16.9 (66)	406	9.4 (93)	49	*	338	7.4 (38)	406	3.0 (28)
4354	570	26.7 (108)	623	10.8 (104)	143	25.2 (42)	570	16.1 (108)	670	8.8 (108)
4355	205	28.8 (116)	236	15.3 (127)	50	*	205	16.1 (108)	244	10.7 (119)
4356	270	30.4 (123)	344	13.7 (121)	79	*	271	14.4 (96)	347	13.5 (133)
4359	415	12.3 (35)	497	6.4 (64)	51	*	418	3.6 (7)	493	3.4 (36)
4360	164	16.5 (63)	201	11.9 (114)	24	*	164	17.7 (115)	203	11.8 (127)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 102

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators <sup>(2,3)</sup>										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes <sup>(4)</sup>	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4361	178	29.2 (119)	206	11.7 (112)	45	*	178	24.2 (129)	212	12.7 (130)
4362	197	28.9 (118)	216	13.0 (119)	53	*	198	19.2 (119)	227	15.0 (135)
4363	464	19.8 (81)	605	10.2 (97)	84	*	464	15.3 (102)	608	10.2 (117)
4401	526	18.8 (76)	682	11.0 (106)	100	42.0 (23)	530	7.5 (41)	689	3.5 (37)
4501	526	28.3 (114)	668	11.2 (108)	144	51.4 (10)	527	15.9 (106)	674	9.3 (111)
4601	800	18.1 (73)	1,022	8.1 (82)	125	40.0 (27)	802	9.4 (66)	1,026	4.7 (61)
4701	692	20.5 (83)	908	5.7 (55)	132	50.0 (12)	696	8.3 (55)	931	5.0 (63)
5050	1,036	9.7 (24)	1,471	3.9 (37)	91	*	1,033	7.4 (38)	1,473	4.2 (53)
5051	1,657	8.3 (15)	2,347	2.9 (26)	119	50.4 (11)	1,654	7.1 (35)	2,341	4.7 (61)
5052	1,798	6.8 (13)	2,634	2.7 (21)	121	43.8 (21)	1,784	5.8 (26)	2,619	3.5 (37)
5055	1,044	1.4 (2)	1,281	0.7 (3)	13	*	1,042	1.4 (3)	1,280	0.2 (1)
5101	901	8.5 (16)	1,260	3.2 (31)	75	*	901	6.1 (29)	1,267	2.8 (21)
5102	2	*	1	*	0	*	2	*	1	*
5202	457	32.6 (126)	548	10.0 (96)	144	39.6 (29)	454	7.7 (46)	575	4.5 (59)
5206	7	*	7	*	2	*	7	*	7	*
5601	660	22.9 (95)	844	7.9 (81)	141	36.9 (34)	661	7.3 (36)	876	4.2 (53)
5801	1,209	24.2 (99)	1,521	8.7 (88)	292	41.4 (24)	1,218	12.4 (83)	1,542	6.3 (83)
5803	21	*	38	*	1	*	21	*	37	*
5804	2	*	2	*	1	*	2	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 103

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
5806	30	*	64	*	8	*	31	*	64	*
6101	43	*	54	*	4	*	43	*	55	*
6102	37	*	51	*	5	*	37	*	51	*
6103	2,261	15.6 (59)	2,900	8.1 (82)	333	32.7 (37)	2,265	10.5 (74)	2,948	5.9 (79)
6105	5	*	6	*	0	*	5	*	6	*
6301	2,377	6.4 (11)	3,101	2.7 (21)	148	40.5 (26)	2,370	4.2 (10)	3,108	2.2 (16)
6312	1,210	2.5 (4)	1,674	1.1 (6)	32	*	1,207	4.6 (13)	1,671	2.0 (11)
6316	811	6.3 (10)	1,098	2.8 (25)	47	*	809	5.1 (16)	1,104	3.7 (42)
6318	2	*	10	*	0	*	2	*	9	*
6320	309	11.0 (30)	408	1.2 (7)	28	*	310	4.8 (15)	409	3.9 (48)
6325	191	22.0 (92)	281	7.8 (79)	39	*	192	14.1 (93)	287	7.7 (101)
6335	642	9.0 (20)	844	1.8 (9)	46	*	641	8.9 (62)	847	5.5 (74)
6345	87	*	115	2.6 (18)	23	*	87	*	121	8.3 (106)
6360	524	15.3 (54)	703	4.8 (47)	76	*	523	12.0 (80)	712	5.5 (74)
7306	1,267	18.8 (76)	1,581	7.4 (77)	225	48.4 (14)	1,269	10.3 (73)	1,599	5.0 (63)
7401	1,331	26.4 (106)	1,568	9.2 (92)	308	43.8 (21)	1,335	14.7 (100)	1,590	8.1 (105)
7601	366	12.8 (39)	438	4.8 (47)	49	*	371	7.5 (41)	445	5.6 (76)
7850	786	10.3 (25)	990	6.4 (64)	83	*	789	3.9 (8)	997	4.6 (60)
7852	5	*	3	*	0	*	5	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 104

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8101	1,042	13.5 (44)	1,515	2.6 (18)	132	71.2 (3)	1,045	8.0 (50)	1,522	4.0 (50)
8102	271	8.9 (19)	340	2.1 (12)	22	*	271	5.2 (17)	341	3.2 (32)
8106	335	13.1 (43)	504	5.2 (52)	41	*	335	5.4 (21)	504	1.6 (7)
8150	4	*	4	*	0	*	4	*	4	*
8151	2,253	12.5 (37)	2,762	4.5 (43)	239	51.5 (9)	2,257	5.6 (25)	2,776	2.7 (20)
8152	5	*	5	*	1	*	5	*	5	*
8153	1	*	0	*	0	*	1	*	0	*
8155	4	*	3	*	0	*	4	*	3	*
8201	1,490	12.1 (33)	2,065	4.3 (42)	159	57.9 (7)	1,496	5.9 (27)	2,061	4.4 (57)
8202	3,019	2.1 (3)	3,716	0.6 (2)	60	*	3,021	3.1 (5)	3,720	1.1 (3)
8203	221	14.5 (48)	313	4.2 (40)	26	*	222	17.1 (113)	314	12.7 (130)
8250	304	11.8 (32)	347	3.7 (35)	28	*	303	6.9 (32)	349	5.2 (66)
8252	76	*	88	*	10	*	76	*	90	*
8255	1,205	14.6 (50)	1,799	5.4 (54)	156	48.7 (13)	1,206	9.6 (68)	1,809	6.9 (92)
8257	1,154	0.3 (1)	1,479	0.2 (1)	3	*	1,149	1.1 (2)	1,473	0.6 (2)
8263	232	16.4 (62)	188	2.7 (21)	33	*	234	6.0 (28)	188	4.3 (56)
8264	501	8.8 (17)	778	1.9 (11)	42	*	502	4.6 (13)	781	2.9 (23)
8267	452	12.8 (39)	570	3.7 (35)	43	*	451	8.6 (56)	578	6.9 (92)
8901	465	9.2 (22)	666	3.2 (31)	39	*	465	6.9 (32)	667	3.3 (34)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 105

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8902	757	10.4 (26)	1,069	2.2 (14)	76	*	760	5.4 (21)	1,070	2.1 (14)
8903	839	15.5 (56)	1,176	3.3 (34)	117	71.8 (2)	835	8.1 (51)	1,176	2.9 (23)
8904	1,138	13.7 (46)	1,678	3.2 (31)	142	68.3 (4)	1,143	9.5 (67)	1,682	3.0 (28)
8905	1,177	6.7 (12)	1,635	2.4 (16)	70	*	1,178	8.7 (59)	1,636	5.9 (79)
8909	725	8.8 (17)	975	2.9 (26)	60	*	724	9.3 (64)	978	3.9 (48)
9045	903	3.7 (6)	1,147	1.0 (4)	30	*	898	5.5 (23)	1,150	2.8 (21)
9050	602	15.1 (52)	809	3.1 (30)	90	*	611	8.7 (59)	812	1.7 (9)
9051	1,504	2.7 (5)	1,823	1.6 (8)	44	*	1,509	1.0 (1)	1,828	1.1 (3)
9053	899	12.5 (37)	1,470	2.1 (12)	113	77.9 (1)	907	8.9 (62)	1,463	2.1 (14)
9055	3	*	6	*	0	*	3	*	6	*
9058	881	6.0 (9)	1,459	2.5 (17)	51	*	878	8.1 (51)	1,459	2.0 (11)
9066	367	18.3 (74)	350	4.6 (44)	66	*	368	11.4 (79)	350	2.6 (19)
9068	552	10.7 (29)	769	2.3 (15)	54	*	551	7.8 (48)	769	2.9 (23)
9069	942	4.9 (7)	1,398	2.7 (21)	43	*	943	6.2 (30)	1,403	3.8 (45)
9070	1,188	10.4 (26)	1,769	1.8 (9)	111	64.9 (5)	1,190	12.9 (86)	1,770	3.2 (32)
9080	1,891	12.4 (36)	2,659	2.6 (18)	217	64.1 (6)	1,899	10.1 (72)	2,666	4.2 (53)
9082	58	*	100	4.0 (38)	5	*	59	*	100	3.0 (28)
9083	12	*	11	*	1	*	12	*	11	*
9087	3	*	3	*	0	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 7A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 106

### Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Michigan	98,366	14.6	127,800	5.8	13,377	43.1	98,459	9.1	128,740	4.8
----------	--------	------	---------	-----	--------	------	--------	-----	---------	-----

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 107

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	106	8.5 (38)	9.4 (126)	107	15.0 (111)	89	*	106	71.7 (109)
0102	314	9.6 (62)	10.8 (135)	318	10.7 (29)	273	4.4 (63)	312	73.4 (102)
0103	337	6.8 (15)	11.3 (136)	344	9.6 (13)	303	3.3 (32)	337	76.6 (91)
0104	95	*	*	96	*	80	*	95	*
0105	1	*	*	1	*	1	*	1	*
0250	98	*	*	98	*	87	*	97	*
0251	59	*	*	60	*	51	*	59	*
0252	89	*	*	89	*	84	*	89	*
0253	104	9.6 (62)	6.7 (77)	109	13.8 (95)	89	*	104	73.1 (104)
0401	350	8.3 (37)	7.1 (89)	351	12.0 (56)	297	4.4 (63)	351	82.6 (49)
0402	285	9.5 (59)	10.2 (132)	289	10.7 (29)	252	3.2 (28)	285	80.7 (67)
0403	104	8.7 (43)	7.7 (103)	103	11.7 (52)	89	*	106	84.0 (32)
0404	98	*	*	99	*	85	*	98	*
0501	325	7.7 (23)	6.2 (55)	358	7.3 (3)	300	3.7 (42)	346	86.4 (19)
0503	46	*	*	48	*	43	*	47	*
0504	279	10.8 (83)	7.9 (109)	285	14.0 (97)	238	2.1 (13)	280	89.3 (8)
0505	163	5.5 (4)	9.2 (124)	185	8.6 (5)	147	1.4 (6)	181	91.2 (2)
0506	143	7.0 (16)	7.7 (103)	150	8.7 (7)	128	4.7 (72)	148	81.8 (55)
0507	46	*	*	54	*	42	*	51	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 108

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	10	*	*	10	*	9	*	10	*
0650	926	8.6 (41)	7.6 (99)	953	9.8 (17)	831	4.6 (69)	930	83.8 (34)
0651	426	10.3 (73)	7.7 (103)	452	10.2 (23)	380	6.8 (112)	426	77.9 (81)
0652	237	5.1 (2)	6.3 (58)	237	8.9 (9)	213	0.9 (2)	235	83.8 (34)
0653	331	6.6 (13)	6.3 (58)	342	9.9 (20)	298	3.7 (42)	330	85.2 (24)
0654	334	11.1 (90)	6.0 (47)	338	12.4 (62)	293	5.1 (89)	333	81.1 (61)
0655	272	7.4 (19)	5.5 (36)	280	8.2 (4)	249	4.8 (77)	272	80.5 (71)
0657	21	*	*	23	*	18	*	21	*
0701	359	5.6 (6)	7.0 (88)	372	12.1 (58)	315	1.3 (4)	365	90.7 (3)
0702	57	*	*	59	*	51	*	57	*
0703	152	6.6 (13)	7.2 (93)	156	9.0 (10)	138	2.9 (23)	152	75.0 (95)
0704	47	*	*	47	*	40	*	47	*
0705	66	*	*	70	*	57	*	65	*
0706	1	*	*	1	*	1	*	1	*
0850	593	12.1 (103)	6.9 (83)	609	14.4 (102)	508	5.1 (89)	599	83.6 (40)
0851	551	9.3 (54)	6.5 (70)	583	14.8 (109)	472	2.5 (19)	558	83.3 (45)
0901	1,179	11.4 (94)	8.3 (117)	1,225	13.0 (76)	1,023	4.6 (69)	1,182	73.9 (98)
0902	21	*	*	21	*	19	*	21	*
1051	918	8.5 (38)	8.2 (115)	936	11.5 (49)	803	4.1 (58)	927	90.5 (4)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 109

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1052	3	*	*	2	*	2	*	3	*
1101	1,060	12.1 (103)	5.6 (38)	1,105	13.5 (91)	910	6.4 (109)	1,064	72.4 (107)
1102	591	6.4 (11)	7.1 (89)	613	10.0 (22)	527	2.5 (19)	593	76.9 (89)
1106	46	*	*	46	*	41	*	46	*
1250	837	8.8 (44)	7.6 (99)	887	10.3 (25)	739	5.4 (100)	859	79.5 (76)
1251	591	11.0 (88)	5.8 (42)	606	15.5 (118)	489	4.9 (82)	592	83.6 (40)
1252	445	9.2 (53)	9.2 (124)	463	12.7 (70)	388	3.4 (34)	451	83.8 (34)
1253	532	7.1 (18)	9.8 (130)	548	12.0 (56)	457	1.3 (4)	540	83.7 (38)
1350	1,959	10.2 (71)	6.9 (83)	2,019	11.0 (39)	1,727	5.2 (94)	1,974	81.7 (56)
1352	303	12.2 (105)	9.6 (129)	312	14.4 (102)	258	1.9 (10)	306	73.5 (101)
1401	180	7.8 (26)	11.7 (137)	188	9.6 (13)	164	3.7 (42)	180	87.2 (14)
1402	61	*	*	66	*	56	*	62	*
1403	16	*	*	16	*	14	*	16	*
1501	610	10.0 (69)	7.4 (96)	645	12.6 (68)	527	4.0 (53)	625	81.9 (54)
1503	685	7.0 (16)	6.1 (51)	691	10.9 (37)	604	2.6 (21)	687	81.1 (61)
1504	1,515	10.5 (77)	6.3 (58)	1,631	13.5 (91)	1,310	3.8 (48)	1,595	83.4 (44)
1506	450	9.8 (65)	7.6 (99)	466	12.7 (70)	389	4.9 (82)	453	79.7 (73)
1507	425	9.9 (67)	7.3 (95)	442	14.3 (100)	362	3.3 (32)	435	87.6 (11)
1509	284	8.1 (31)	6.3 (58)	289	12.8 (72)	245	3.7 (42)	286	76.9 (89)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 110

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1511	506	10.7 (79)	6.3 (58)	522	13.0 (76)	438	5.0 (88)	510	80.6 (70)
1512	347	10.7 (79)	6.3 (58)	352	14.5 (104)	296	3.0 (25)	349	84.8 (25)
1514	628	14.2 (119)	6.8 (80)	649	15.9 (123)	522	3.6 (38)	637	87.4 (13)
1701	434	7.8 (26)	8.8 (121)	449	10.7 (29)	385	3.1 (27)	437	83.5 (43)
1950	499	8.6 (41)	4.4 (21)	512	12.3 (60)	433	3.5 (35)	501	85.4 (23)
1952	115	4.3 (1)	6.1 (51)	117	6.8 (1)	105	1.9 (10)	116	92.2 (1)
1953	230	9.6 (62)	6.1 (51)	238	10.9 (37)	204	4.4 (63)	232	85.8 (21)
1954	307	9.1 (51)	7.2 (93)	316	9.2 (12)	277	4.7 (72)	306	85.9 (20)
1955	112	8.9 (47)	6.3 (58)	114	16.7 (126)	92	*	113	79.6 (75)
1957	332	12.0 (102)	7.8 (106)	345	15.7 (122)	281	4.3 (61)	333	87.1 (15)
1963	70	*	*	69	*	63	*	71	*
2150	448	7.4 (19)	6.0 (47)	464	11.0 (39)	398	2.3 (14)	449	81.1 (61)
2151	187	5.9 (9)	5.9 (45)	191	11.0 (39)	165	1.8 (9)	186	77.4 (85)
2152	25	*	*	26	*	22	*	25	*
2251	293	5.5 (4)	4.8 (29)	293	10.2 (23)	259	1.9 (10)	292	83.9 (33)
2252	141	11.3 (92)	8.5 (120)	149	18.1 (134)	117	1.7 (8)	139	88.5 (10)
2554	36	*	*	36	*	34	*	36	*
2555	6,045	13.3 (114)	5.9 (45)	6,156	13.5 (91)	5,188	6.8 (112)	6,070	67.2 (117)
2556	153	19.6 (137)	3.9 (16)	155	20.0 (137)	119	9.2 (127)	155	59.4 (128)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 111

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2557	1	*	*	1	*	1	*	1	*
2562	1	*	*	1	*	0	*	1	*
2563	2	*	*	2	*	2	*	2	*
2564	2	*	*	2	*	2	*	2	*
3001	961	14.6 (120)	3.9 (16)	994	15.6 (119)	810	6.5 (110)	961	64.3 (121)
3002	393	10.4 (75)	3.8 (14)	398	13.1 (81)	342	2.9 (23)	393	70.7 (111)
3004	266	9.0 (49)	8.3 (117)	274	15.0 (111)	224	4.9 (82)	267	78.7 (78)
3005	148	9.5 (59)	8.8 (121)	150	8.7 (7)	135	3.7 (42)	149	81.2 (60)
3007	3	*	*	3	*	3	*	3	*
3201	385	8.1 (31)	9.1 (123)	391	12.3 (60)	336	2.4 (16)	386	82.1 (53)
3301	3,884	11.7 (97)	6.5 (70)	4,018	13.4 (88)	3,350	4.2 (59)	3,930	83.6 (40)
3304	74	*	*	77	*	68	*	74	*
3305	122	9.8 (65)	10.7 (134)	129	12.4 (62)	107	4.7 (72)	123	90.2 (6)
3306	155	5.8 (7)	7.1 (89)	157	15.3 (116)	130	0.8 (1)	155	86.5 (18)
3308	63	*	*	63	*	57	*	63	*
3312	363	9.4 (58)	5.2 (34)	376	13.0 (76)	319	6.3 (107)	365	87.1 (15)
3314	2	*	*	2	*	2	*	2	*
3550	45	*	*	48	*	38	*	45	*
3551	5	*	*	6	*	4	*	5	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

### 2022 - 2024 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 112

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3701	725	10.3 (73)	6.6 (74)	742	12.5 (66)	631	4.8 (77)	727	87.6 (11)
3850	2,155	12.2 (105)	7.1 (89)	2,231	13.9 (96)	1,851	5.1 (89)	2,165	81.0 (66)
3901	1,168	12.2 (105)	5.2 (34)	1,212	13.4 (88)	1,005	5.6 (101)	1,179	74.2 (97)
3902	1	*	*	1	*	1	*	1	*
3950	1,685	10.8 (83)	7.4 (96)	1,721	12.5 (66)	1,468	4.8 (77)	1,700	80.3 (72)
3954	1	*	*	1	*	1	*	1	*
4106	2,090	13.5 (115)	4.4 (21)	2,159	15.6 (119)	1,760	6.8 (112)	2,120	78.5 (79)
4107	2,223	12.3 (109)	4.8 (29)	2,305	15.2 (115)	1,881	5.1 (89)	2,260	81.1 (61)
4109	624	13.6 (116)	7.9 (109)	642	17.4 (127)	511	3.5 (35)	628	84.7 (27)
4112	482	5.2 (3)	3.7 (12)	490	10.8 (35)	427	2.3 (14)	483	43.9 (137)
4115	2,979	12.5 (111)	6.0 (47)	3,131	13.2 (82)	2,581	5.2 (94)	3,083	83.3 (45)
4351	450	10.0 (69)	6.9 (83)	459	14.6 (105)	380	4.2 (59)	452	86.7 (17)
4352	157	7.6 (22)	6.4 (68)	161	12.4 (62)	132	4.5 (68)	157	82.8 (48)
4353	387	10.9 (86)	6.7 (77)	391	11.5 (49)	342	5.8 (104)	389	80.7 (67)
4354	627	9.1 (51)	6.5 (70)	636	11.3 (45)	549	4.9 (82)	631	83.8 (34)
4355	227	8.8 (44)	6.6 (74)	232	13.4 (88)	197	5.1 (89)	227	83.7 (38)
4356	319	8.2 (36)	10.0 (131)	327	10.7 (29)	281	3.6 (38)	323	84.5 (28)
4359	458	9.0 (49)	6.3 (58)	467	11.8 (54)	397	4.0 (53)	460	81.3 (59)
4360	187	6.4 (11)	8.0 (112)	193	9.8 (17)	167	2.4 (16)	189	90.5 (4)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 113

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4361	194	7.7 (23)	4.6 (25)	200	11.0 (39)	171	1.2 (3)	196	84.2 (30)
4362	213	10.8 (83)	8.0 (112)	224	12.9 (74)	186	2.7 (22)	213	89.2 (9)
4363	571	9.5 (59)	9.5 (128)	574	12.9 (74)	485	3.9 (51)	575	90.1 (7)
4401	634	8.8 (44)	8.2 (115)	665	10.7 (29)	565	3.7 (42)	638	82.6 (49)
4501	633	5.8 (7)	7.6 (99)	647	9.7 (15)	570	1.6 (7)	634	82.6 (49)
4601	972	7.9 (28)	7.4 (96)	996	10.8 (35)	864	3.2 (28)	970	81.4 (57)
4701	886	11.2 (91)	6.3 (58)	895	15.3 (116)	745	3.8 (48)	888	84.2 (30)
5050	1,429	12.3 (109)	4.6 (25)	1,430	13.2 (82)	1,219	6.8 (112)	1,431	75.3 (94)
5051	2,219	11.7 (97)	4.7 (28)	2,263	15.0 (111)	1,858	4.9 (82)	2,237	77.8 (83)
5052	2,504	14.7 (121)	4.6 (25)	2,535	15.0 (111)	2,106	8.4 (123)	2,520	73.9 (98)
5055	1,183	10.7 (79)	4.8 (29)	1,225	10.4 (26)	1,056	5.9 (105)	1,211	69.9 (113)
5101	1,227	10.4 (75)	5.6 (38)	1,221	13.3 (85)	1,045	3.6 (38)	1,232	79.7 (73)
5102	1	*	*	1	*	1	*	1	*
5202	549	10.2 (71)	4.9 (32)	555	13.3 (85)	472	4.9 (82)	554	84.8 (25)
5206	7	*	*	7	*	6	*	7	*
5601	805	8.1 (31)	9.4 (126)	842	9.9 (20)	725	4.0 (53)	807	77.2 (87)
5801	1,479	8.1 (31)	6.9 (83)	1,514	10.6 (28)	1,317	3.2 (28)	1,478	72.3 (108)
5803	36	*	*	35	*	33	*	36	*
5804	2	*	*	2	*	2	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 114

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5806	62	*	*	62	*	51	*	62	*
6101	54	*	*	55	*	50	*	54	*
6102	49	*	*	51	*	43	*	49	*
6103	2,753	11.9 (100)	6.0 (47)	2,861	14.6 (105)	2,332	5.9 (105)	2,771	73.6 (100)
6105	6	*	*	6	*	5	*	5	*
6301	2,884	11.5 (95)	6.2 (55)	3,029	13.5 (91)	2,486	5.2 (94)	2,961	75.7 (93)
6312	1,569	12.5 (111)	5.0 (33)	1,625	14.3 (100)	1,330	4.8 (77)	1,594	81.1 (61)
6316	1,026	11.5 (95)	5.8 (42)	1,064	13.0 (76)	885	4.6 (69)	1,037	81.4 (57)
6318	9	*	*	9	*	5	*	9	*
6320	387	12.7 (113)	6.2 (55)	400	9.0 (10)	349	8.3 (122)	394	84.3 (29)
6325	279	7.9 (28)	10.4 (133)	279	10.4 (26)	247	2.4 (16)	280	80.7 (67)
6335	809	11.0 (88)	6.7 (77)	823	12.4 (62)	702	4.4 (63)	819	76.3 (92)
6345	118	9.3 (54)	6.8 (80)	119	12.6 (68)	101	3.0 (25)	119	83.2 (47)
6360	657	15.4 (124)	5.6 (38)	686	12.8 (72)	567	9.7 (131)	670	77.8 (83)
7306	1,509	11.3 (92)	6.4 (68)	1,545	14.0 (97)	1,297	3.9 (51)	1,521	72.5 (106)
7401	1,507	9.3 (54)	7.8 (106)	1,537	11.0 (39)	1,340	3.8 (48)	1,510	77.2 (87)
7601	420	8.1 (31)	7.9 (109)	435	11.3 (45)	371	3.5 (35)	428	67.1 (118)
7850	943	7.5 (21)	6.9 (83)	973	11.7 (52)	833	3.2 (28)	947	78.4 (80)
7852	2	*	*	2	*	2	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 115

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8101	1,373	15.1 (123)	2.5 (1)	1,468	17.7 (130)	1,111	7.7 (118)	1,431	59.9 (126)
8102	300	7.7 (23)	8.0 (112)	324	8.6 (5)	269	4.8 (77)	316	70.6 (112)
8106	427	15.5 (126)	6.6 (74)	492	16.3 (125)	353	5.7 (103)	484	67.1 (118)
8150	3	*	*	4	*	3	*	4	*
8151	2,577	12.2 (105)	6.1 (51)	2,669	14.0 (97)	2,203	5.2 (94)	2,626	85.5 (22)
8152	4	*	*	5	*	4	*	4	*
8153	0	*	*	0	*	0	*	0	*
8155	3	*	*	3	*	3	*	3	*
8201	1,944	11.9 (100)	4.5 (24)	2,023	13.3 (85)	1,683	5.2 (94)	1,968	73.1 (104)
8202	3,327	8.5 (38)	3.6 (11)	3,538	9.7 (15)	2,980	4.0 (53)	3,508	69.4 (114)
8203	306	11.8 (99)	7.8 (106)	308	11.4 (47)	270	5.6 (101)	306	78.8 (77)
8250	334	10.5 (77)	3.9 (16)	346	10.7 (29)	297	4.0 (53)	337	68.0 (115)
8252	72	*	*	88	*	65	*	86	*
8255	1,701	10.9 (86)	5.8 (42)	1,760	13.0 (76)	1,466	4.7 (72)	1,723	73.2 (103)
8257	1,287	13.8 (118)	3.0 (4)	1,405	11.1 (44)	1,134	9.3 (128)	1,345	62.0 (124)
8263	180	6.1 (10)	5.6 (38)	186	7.0 (2)	167	3.6 (38)	181	77.3 (86)
8264	747	10.7 (79)	8.3 (117)	762	12.2 (59)	653	4.4 (63)	751	82.2 (52)
8267	541	8.9 (47)	6.5 (70)	562	9.8 (17)	486	4.3 (61)	550	71.1 (110)
8901	611	17.5 (134)	3.8 (14)	645	17.8 (132)	492	9.8 (132)	624	59.8 (127)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 116

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8902	975	16.5 (130)	4.2 (20)	1,023	17.7 (130)	787	8.6 (125)	1,004	67.9 (116)
8903	1,074	15.5 (126)	3.0 (4)	1,138	16.1 (124)	893	8.7 (126)	1,084	55.4 (133)
8904	1,532	16.8 (132)	3.7 (12)	1,629	19.0 (136)	1,216	7.9 (121)	1,562	61.1 (125)
8905	1,432	9.9 (67)	5.5 (36)	1,571	11.8 (54)	1,250	5.3 (99)	1,528	74.7 (96)
8909	877	16.5 (130)	3.1 (7)	946	17.5 (129)	711	8.4 (123)	902	66.7 (120)
9045	1,094	8.0 (30)	6.8 (80)	1,127	11.4 (47)	965	4.7 (72)	1,109	44.2 (136)
9050	773	13.6 (116)	4.4 (21)	800	14.6 (105)	656	7.5 (116)	783	59.4 (128)
9051	1,612	9.3 (54)	6.3 (58)	1,708	11.5 (49)	1,405	6.3 (107)	1,719	77.9 (81)
9053	1,370	15.4 (124)	3.0 (4)	1,436	13.2 (82)	1,178	9.5 (130)	1,379	58.2 (130)
9055	6	*	*	6	*	6	*	5	*
9058	1,341	17.4 (133)	3.4 (9)	1,407	15.6 (119)	1,118	9.3 (128)	1,371	62.2 (123)
9066	335	17.6 (136)	3.9 (16)	344	17.4 (127)	276	6.5 (110)	334	57.8 (131)
9068	741	17.5 (134)	2.8 (2)	755	14.7 (108)	628	10.4 (133)	742	52.6 (134)
9069	1,322	16.3 (129)	3.1 (7)	1,363	18.1 (134)	1,083	7.8 (119)	1,327	45.9 (135)
9070	1,671	15.9 (128)	3.4 (9)	1,723	18.0 (133)	1,360	7.8 (119)	1,678	57.4 (132)
9080	2,478	15.0 (122)	2.9 (3)	2,592	14.8 (109)	2,101	7.5 (116)	2,547	62.9 (122)
9082	95	*	*	98	*	75	*	96	*
9083	11	*	*	11	*	9	*	11	*
9087	3	*	*	3	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 4/14/2025

## Table 8A

2022 - 2024 Pregnancy Nutrition Surveillance (1)

Page : 117

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Michigan	119,947	11.5	5.9	124,372	13.2	103,291	5.3	121,765	45.9
----------	---------	------	-----	---------	------	---------	-----	---------	------

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.