

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2022 Pregnancy Nutrition Surveillance (1)

Page : 1

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
0101	36	*	*	*	*	*	*	*
0102	124	87.1	0.0	4.8	0.0	0.0	8.1	0.0
0103	113	96.5	0.9	0.9	0.0	0.0	1.8	0.0
0104	42	*	*	*	*	*	*	*
0250	38	*	*	*	*	*	*	*
0251	16	*	*	*	*	*	*	*
0252	39	*	*	*	*	*	*	*
0253	37	*	*	*	*	*	*	*
0401	146	94.5	0.0	2.7	0.0	0.7	2.1	0.0
0402	130	82.3	3.8	3.1	0.0	0.0	10.8	0.0
0403	41	*	*	*	*	*	*	*
0404	36	*	*	*	*	*	*	*
0501	131	89.3	1.5	3.8	0.0	0.0	4.6	0.8
0503	18	*	*	*	*	*	*	*
0504	104	88.5	1.9	3.8	1.9	0.0	3.8	0.0
0505	72	*	*	*	*	*	*	*
0506	61	*	*	*	*	*	*	*
0507	16	*	*	*	*	*	*	*
0508	4	*	*	*	*	*	*	*
0650	376	79.0	4.5	7.7	4.5	0.8	3.5	0.0
0651	194	89.7	1.5	6.7	0.0	0.5	1.5	0.0
0652	89	*	*	*	*	*	*	*
0653	124	96.0	0.0	0.8	0.0	0.0	3.2	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2022 Pregnancy Nutrition Surveillance (1)

Page : 2

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
0654	142	91.5	0.0	3.5	1.4	0.0	3.5	0.0
0655	103	97.1	0.0	0.0	0.0	0.0	2.9	0.0
0657	9	*	*	*	*	*	*	*
0701	147	83.7	2.7	2.7	0.0	1.4	8.8	0.7
0702	22	*	*	*	*	*	*	*
0703	52	*	*	*	*	*	*	*
0704	16	*	*	*	*	*	*	*
0705	24	*	*	*	*	*	*	*
0850	248	81.0	6.9	8.5	0.0	2.8	0.8	0.0
0851	198	92.4	0.5	4.5	0.5	0.0	2.0	0.0
0901	484	81.6	2.3	9.1	0.2	0.2	6.2	0.4
0902	11	*	*	*	*	*	*	*
1051	366	80.9	1.1	9.3	1.6	0.8	5.7	0.5
1052	2	*	*	*	*	*	*	*
1101	454	37.2	44.1	10.6	0.9	1.8	4.0	1.5
1102	259	73.7	8.9	6.9	0.0	1.5	8.5	0.4
1106	21	*	*	*	*	*	*	*
1250	303	82.5	5.0	9.2	0.3	1.0	1.7	0.3
1251	200	90.5	0.0	5.0	1.0	0.0	3.5	0.0
1252	193	61.1	0.0	35.8	0.0	0.0	2.1	1.0
1253	224	77.2	0.0	13.8	0.0	0.4	7.6	0.9
1350	799	61.1	19.6	7.6	0.1	4.4	6.9	0.3
1352	131	67.2	21.4	6.1	0.8	0.0	4.6	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2022 Pregnancy Nutrition Surveillance (1)

Page : 3

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
1401	60	*	*	*	*	*	*	*
1402	25	*	*	*	*	*	*	*
1403	11	*	*	*	*	*	*	*
1501	241	46.1	5.0	41.5	0.0	0.0	5.8	1.7
1503	279	60.6	8.6	21.5	0.4	0.4	6.5	2.2
1504	618	36.9	5.5	49.7	0.0	2.9	4.5	0.5
1506	175	80.0	0.6	13.1	0.0	0.6	4.0	1.7
1507	165	86.7	2.4	5.5	0.0	2.4	3.0	0.0
1509	119	47.1	2.5	47.1	0.0	0.0	2.5	0.8
1511	218	85.3	2.3	6.4	0.0	0.5	5.5	0.0
1512	144	85.4	1.4	8.3	0.0	0.0	4.9	0.0
1514	260	76.9	6.2	13.1	0.0	1.2	1.9	0.8
1701	172	59.9	0.6	1.2	20.9	0.6	16.9	0.0
1950	195	93.3	0.0	5.1	0.0	0.0	1.0	0.5
1952	44	*	*	*	*	*	*	*
1953	79	*	*	*	*	*	*	*
1954	121	84.3	0.0	9.9	0.0	0.0	5.8	0.0
1955	44	*	*	*	*	*	*	*
1957	134	85.1	0.0	10.4	0.0	0.0	3.7	0.7
1963	18	*	*	*	*	*	*	*
2150	170	85.9	0.6	5.3	1.8	0.0	5.9	0.6
2151	73	*	*	*	*	*	*	*
2152	16	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2022 Pregnancy Nutrition Surveillance (1)

Page : 4

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
2251	121	95.0	0.0	2.5	0.0	1.7	0.8	0.0
2252	63	*	*	*	*	*	*	*
2554	19	*	*	*	*	*	*	*
2555	2,410	52.2	34.9	6.7	0.0	0.2	5.6	0.3
2556	35	*	*	*	*	*	*	*
2557	1	*	*	*	*	*	*	*
2562	1	*	*	*	*	*	*	*
2563	2	*	*	*	*	*	*	*
2564	2	*	*	*	*	*	*	*
3001	407	29.7	32.9	17.9	0.0	0.2	18.9	0.2
3002	51	*	*	*	*	*	*	*
3004	94	*	*	*	*	*	*	*
3005	56	*	*	*	*	*	*	*
3007	1	*	*	*	*	*	*	*
3201	135	89.6	0.7	5.9	0.0	0.7	1.5	1.5
3301	1,424	38.1	29.4	16.9	0.1	6.8	8.6	0.2
3304	48	*	*	*	*	*	*	*
3305	66	*	*	*	*	*	*	*
3306	78	*	*	*	*	*	*	*
3308	38	*	*	*	*	*	*	*
3312	176	45.5	26.7	19.9	0.0	2.3	5.7	0.0
3550	28	*	*	*	*	*	*	*
3551	3	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2022 Pregnancy Nutrition Surveillance (1)

Page : 5

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
3701	276	90.2	0.0	6.2	0.0	0.0	3.6	0.0
3850	887	72.7	12.1	7.4	0.1	0.5	7.1	0.1
3901	455	37.6	51.4	4.8	0.0	0.9	5.1	0.2
3902	2	*	*	*	*	*	*	*
3950	640	54.1	23.3	12.5	0.0	1.1	8.3	0.8
3954	1	*	*	*	*	*	*	*
4106	717	12.0	34.9	47.8	0.0	1.0	4.0	0.3
4107	953	43.0	28.4	19.8	0.4	1.8	6.3	0.2
4109	191	88.5	0.0	7.9	0.0	1.0	2.1	0.5
4112	186	1.1	1.1	97.3	0.0	0.0	0.5	0.0
4115	1,136	29.7	31.8	22.2	0.4	10.9	4.7	0.4
4351	195	88.2	1.0	6.2	0.0	0.5	4.1	0.0
4352	70	*	*	*	*	*	*	*
4353	149	48.3	0.0	45.0	0.0	0.0	4.0	2.7
4354	246	85.0	4.9	3.3	0.8	0.4	5.3	0.4
4355	79	*	*	*	*	*	*	*
4356	134	84.3	0.0	9.7	0.0	0.0	6.0	0.0
4359	206	64.1	0.5	33.0	0.5	0.0	1.0	1.0
4360	80	*	*	*	*	*	*	*
4361	76	*	*	*	*	*	*	*
4362	92	*	*	*	*	*	*	*
4363	208	89.4	1.4	3.4	0.0	0.0	5.3	0.5
4401	231	87.4	1.3	6.1	0.0	0.9	4.3	0.0

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2022 Pregnancy Nutrition Surveillance (1)

Page : 6

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
4501	251	92.8	0.8	4.8	0.0	0.0	1.6	0.0
4601	408	72.1	2.2	18.9	0.5	0.7	5.1	0.5
4701	338	88.8	2.1	3.0	0.0	1.5	3.8	0.9
5050	576	41.0	46.2	3.8	0.0	3.3	5.6	0.2
5051	869	46.1	40.0	4.9	0.0	3.7	4.7	0.5
5052	931	31.4	38.5	3.3	0.2	21.1	5.5	0.1
5055	466	76.4	10.1	1.3	0.0	11.2	0.9	0.2
5101	449	64.8	13.4	15.4	0.2	1.1	5.1	0.0
5102	2	*	*	*	*	*	*	*
5202	223	87.0	0.9	2.7	3.6	0.0	4.5	1.3
5206	7	*	*	*	*	*	*	*
5601	327	84.7	2.8	7.6	0.0	0.6	4.3	0.0
5801	640	82.5	6.7	7.0	0.0	0.2	2.8	0.8
5803	7	*	*	*	*	*	*	*
5804	1	*	*	*	*	*	*	*
5806	16	*	*	*	*	*	*	*
6101	28	*	*	*	*	*	*	*
6102	27	*	*	*	*	*	*	*
6103	1,109	55.9	26.2	11.2	0.5	0.5	5.7	0.1
6105	3	*	*	*	*	*	*	*
6301	1,183	37.9	32.8	22.9	0.3	2.9	3.0	0.3
6312	583	32.9	56.6	4.8	0.0	1.5	3.8	0.2
6316	385	59.7	23.9	9.4	0.0	2.6	4.2	0.3

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2022 Pregnancy Nutrition Surveillance (1)

Page : 7

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
6320	191	31.9	56.0	4.7	0.5	1.6	5.2	0.0
6325	106	88.7	0.0	6.6	0.0	0.0	4.7	0.0
6335	322	60.9	24.5	4.0	0.0	8.7	1.6	0.3
6345	46	*	*	*	*	*	*	*
6360	256	34.8	40.2	19.9	0.0	1.2	3.5	0.4
7306	606	46.4	30.0	16.0	0.5	0.5	6.6	0.0
7401	629	80.1	5.2	5.1	0.0	0.5	8.6	0.5
7601	180	92.8	0.0	5.6	0.0	0.0	1.7	0.0
7850	363	92.8	0.3	3.9	0.0	0.0	2.8	0.3
7852	3	*	*	*	*	*	*	*
8101	591	9.0	86.3	1.9	0.0	0.2	2.5	0.2
8102	149	26.2	22.1	47.7	0.0	2.0	1.3	0.7
8106	197	29.4	47.7	14.2	0.0	1.0	7.6	0.0
8150	1	*	*	*	*	*	*	*
8151	1,100	38.5	37.2	12.8	0.0	3.5	7.4	0.5
8152	5	*	*	*	*	*	*	*
8153	1	*	*	*	*	*	*	*
8155	2	*	*	*	*	*	*	*
8201	897	44.8	40.1	5.2	0.2	3.7	5.6	0.2
8202	1,370	88.0	8.6	1.8	0.1	0.4	1.0	0.1
8203	103	61.2	32.0	4.9	0.0	0.0	1.9	0.0
8250	168	16.7	71.4	5.4	0.0	1.2	5.4	0.0
8252	48	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2022 Pregnancy Nutrition Surveillance (1)

Page : 8

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
8255	592	55.2	31.4	6.9	0.0	2.4	3.9	0.2
8257	543	52.3	9.6	1.3	0.0	34.8	1.7	0.4
8263	133	53.4	25.6	15.0	0.0	1.5	4.5	0.0
8264	270	54.8	31.9	6.3	0.4	2.6	3.7	0.4
8267	189	51.9	21.2	23.3	0.0	1.1	2.6	0.0
8901	283	9.2	80.2	4.9	0.0	2.5	3.2	0.0
8902	443	9.0	84.2	4.1	0.0	0.7	2.0	0.0
8903	478	5.4	86.6	1.7	0.0	2.1	4.2	0.0
8904	673	3.7	90.6	1.8	0.0	0.9	3.0	0.0
8905	635	66.1	21.3	8.0	0.0	2.5	2.0	0.0
8909	396	5.1	89.1	1.8	0.3	0.8	3.0	0.0
9045	419	7.4	20.5	69.5	0.2	0.5	1.4	0.5
9050	320	16.9	60.0	20.6	0.0	0.0	2.5	0.0
9051	547	6.8	5.1	86.7	0.0	0.0	1.3	0.2
9053	525	7.2	84.6	2.1	0.2	3.0	2.5	0.4
9055	4	*	*	*	*	*	*	*
9058	479	17.3	71.2	1.5	0.2	7.1	2.7	0.0
9066	223	1.8	94.2	0.4	0.0	1.8	1.8	0.0
9068	243	1.6	96.3	0.4	0.0	0.8	0.8	0.0
9069	518	1.2	97.1	0.6	0.0	0.0	1.0	0.2
9070	664	7.8	87.8	1.4	0.0	1.8	1.1	0.2
9080	871	19.9	71.3	2.4	0.1	0.8	4.8	0.7
9082	24	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2022 Pregnancy Nutrition Surveillance (1)

Page : 9

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9083	2	*	*	*	*	*	*	*
9087	3	*	*	*	*	*	*	*
Michigan	47,919	51.7	28.7	11.8	0.4	2.6	4.6	0.3
National (Prior Year)	66,073	48.2	27.8	15.1	1.9	2.8	4.0	0.3

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2022 Pregnancy Nutrition Surveillance (1)

Page : 10

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	36	*	*	*	*	*	*	*
0102	124	0.0	1.6	9.7	54.8	31.5	2.4	0.0
0103	113	0.0	1.8	10.6	60.2	24.8	1.8	0.9
0104	42	*	*	*	*	*	*	*
0250	38	*	*	*	*	*	*	*
0251	16	*	*	*	*	*	*	*
0252	39	*	*	*	*	*	*	*
0253	37	*	*	*	*	*	*	*
0401	146	0.0	2.1	7.5	57.5	32.9	0.0	0.0
0402	130	0.0	1.5	6.9	63.8	26.9	0.8	0.0
0403	41	*	*	*	*	*	*	*
0404	36	*	*	*	*	*	*	*
0501	131	0.0	0.8	7.6	60.3	29.0	2.3	0.0
0503	18	*	*	*	*	*	*	*
0504	104	0.0	1.0	3.8	54.8	37.5	2.9	0.0
0505	72	*	*	*	*	*	*	*
0506	61	*	*	*	*	*	*	*
0507	16	*	*	*	*	*	*	*
0508	4	*	*	*	*	*	*	*
0650	376	0.3	1.6	4.8	63.8	26.9	1.9	0.8
0651	194	0.5	1.0	8.2	64.9	24.2	0.5	0.5
0652	89	*	*	*	*	*	*	*
0653	124	0.0	0.8	6.5	64.5	27.4	0.0	0.8
0654	142	0.0	1.4	4.9	66.2	26.1	0.7	0.7

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2022 Pregnancy Nutrition Surveillance (1)

Page : 11

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0655	103	0.0	4.9	4.9	63.1	24.3	1.0	1.9
0657	9	*	*	*	*	*	*	*
0701	147	0.0	0.7	10.2	62.6	24.5	2.0	0.0
0702	22	*	*	*	*	*	*	*
0703	52	*	*	*	*	*	*	*
0704	16	*	*	*	*	*	*	*
0705	24	*	*	*	*	*	*	*
0850	248	0.0	2.8	10.1	54.8	28.2	2.4	1.6
0851	198	0.0	1.5	7.1	68.7	19.7	1.0	2.0
0901	484	0.0	3.1	9.5	57.4	27.7	1.0	1.2
0902	11	*	*	*	*	*	*	*
1051	366	0.3	1.6	5.2	55.7	35.2	1.9	0.0
1052	2	*	*	*	*	*	*	*
1101	454	0.0	2.4	10.8	50.9	31.1	2.6	2.2
1102	259	0.0	1.9	8.5	64.1	23.2	1.2	1.2
1106	21	*	*	*	*	*	*	*
1250	303	0.0	1.0	7.9	61.1	26.1	3.3	0.7
1251	200	0.5	4.0	7.0	59.0	24.5	4.0	1.0
1252	193	0.0	2.6	9.8	63.7	19.7	3.6	0.5
1253	224	0.0	1.3	7.1	60.3	29.5	1.3	0.4
1350	799	0.0	3.0	7.9	59.8	26.5	1.8	1.0
1352	131	0.0	1.5	8.4	60.3	28.2	0.0	1.5
1401	60	*	*	*	*	*	*	*
1402	25	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**WIC PNSS & PedNSS**  
**Table 4A**  
**2022 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Run Date: 5/29/2024

Page : 12

<b>Comparison of Age Distribution (2)</b>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
1403	11	*	*	*	*	*	*	*
1501	241	0.0	2.9	7.1	57.3	29.5	0.8	2.5
1503	279	0.0	2.5	6.5	63.4	25.4	0.4	1.8
1504	618	0.2	2.9	7.8	59.4	28.2	1.0	0.6
1506	175	0.0	0.6	5.7	61.7	28.6	1.1	2.3
1507	165	0.0	0.6	4.8	57.6	34.5	1.8	0.6
1509	119	0.0	0.8	6.7	47.9	39.5	5.0	0.0
1511	218	0.0	1.4	3.2	71.6	22.5	0.5	0.9
1512	144	0.0	0.7	9.7	53.5	31.3	4.2	0.7
1514	260	0.0	1.9	5.4	55.4	34.6	2.3	0.4
1701	172	0.0	1.7	5.2	61.6	30.2	0.6	0.6
1950	195	0.0	3.6	8.7	69.2	16.9	0.5	1.0
1952	44	*	*	*	*	*	*	*
1953	79	*	*	*	*	*	*	*
1954	121	0.0	2.5	10.7	62.0	21.5	2.5	0.8
1955	44	*	*	*	*	*	*	*
1957	134	0.0	2.2	3.7	56.7	35.1	1.5	0.7
1963	18	*	*	*	*	*	*	*
2150	170	0.0	0.0	7.6	61.8	28.8	1.8	0.0
2151	73	*	*	*	*	*	*	*
2152	16	*	*	*	*	*	*	*
2251	121	0.8	0.0	5.0	60.3	32.2	1.7	0.0
2252	63	*	*	*	*	*	*	*
2554	19	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2022 Pregnancy Nutrition Surveillance (1)

Page : 13

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2555	2,410	0.2	1.7	6.7	62.3	27.1	1.5	0.5
2556	35	*	*	*	*	*	*	*
2557	1	*	*	*	*	*	*	*
2562	1	*	*	*	*	*	*	*
2563	2	*	*	*	*	*	*	*
2564	2	*	*	*	*	*	*	*
3001	407	0.0	0.5	9.3	61.7	24.6	2.0	2.0
3002	51	*	*	*	*	*	*	*
3004	94	*	*	*	*	*	*	*
3005	56	*	*	*	*	*	*	*
3007	1	*	*	*	*	*	*	*
3201	135	0.0	0.7	5.2	62.2	30.4	1.5	0.0
3301	1,424	0.1	2.1	5.7	61.3	27.7	2.1	1.1
3304	48	*	*	*	*	*	*	*
3305	66	*	*	*	*	*	*	*
3306	78	*	*	*	*	*	*	*
3308	38	*	*	*	*	*	*	*
3312	176	0.0	0.6	4.5	58.5	35.2	1.1	0.0
3550	28	*	*	*	*	*	*	*
3551	3	*	*	*	*	*	*	*
3701	276	0.4	1.4	5.4	63.4	26.8	1.4	1.1
3850	887	0.2	2.4	6.8	61.7	26.5	1.2	1.2
3901	455	0.2	2.2	7.7	57.8	27.9	2.9	1.3
3902	2	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**WIC PNSS & PedNSS**  
**Table 4A**  
**2022 Pregnancy Nutrition Surveillance (1)**  
**Michigan**  
**Comparison of Age Distribution by Clinic**

Run Date: 5/29/2024

Page : 14

<b>Comparison of Age Distribution (2)</b>								
<b>Clinic</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
3950	640	0.0	1.9	8.8	58.8	27.7	2.3	0.6
3954	1	*	*	*	*	*	*	*
4106	717	0.1	1.8	7.0	57.2	27.6	3.8	2.5
4107	953	0.1	1.3	5.6	59.2	30.6	2.6	0.6
4109	191	0.0	2.1	5.8	60.2	28.8	2.6	0.5
4112	186	0.5	2.2	7.0	57.0	28.5	3.2	1.6
4115	1,136	0.2	0.9	4.7	55.4	34.9	3.3	0.8
4351	195	0.0	1.5	5.6	67.2	23.1	0.5	2.1
4352	70	*	*	*	*	*	*	*
4353	149	0.0	2.7	7.4	63.1	20.1	3.4	3.4
4354	246	0.0	2.0	8.5	64.2	23.6	1.6	0.0
4355	79	*	*	*	*	*	*	*
4356	134	0.0	0.7	8.2	66.4	22.4	0.7	1.5
4359	206	0.0	1.9	4.9	63.1	26.7	1.9	1.5
4360	80	*	*	*	*	*	*	*
4361	76	*	*	*	*	*	*	*
4362	92	*	*	*	*	*	*	*
4363	208	0.0	2.4	6.7	63.5	23.6	1.9	1.9
4401	231	0.0	0.9	4.3	64.5	27.7	1.7	0.9
4501	251	0.0	2.0	6.0	65.7	23.9	2.0	0.4
4601	408	0.0	1.0	5.4	63.7	27.5	1.0	1.5
4701	338	0.0	0.9	3.8	58.6	33.7	2.1	0.9
5050	576	0.2	0.9	4.9	55.4	35.4	3.0	0.3
5051	869	0.0	0.8	3.9	60.3	31.9	2.8	0.3

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2022 Pregnancy Nutrition Surveillance (1)

Page : 15

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5052	931	0.0	0.9	2.9	54.5	39.7	1.8	0.2
5055	466	0.0	0.4	1.1	48.7	46.4	3.4	0.0
5101	449	0.0	0.7	3.6	59.0	34.3	2.2	0.2
5102	2	*	*	*	*	*	*	*
5202	223	0.0	0.9	4.9	62.8	29.1	2.2	0.0
5206	7	*	*	*	*	*	*	*
5601	327	0.0	0.6	4.3	66.4	27.8	0.9	0.0
5801	640	0.0	1.3	6.6	65.3	25.0	1.4	0.5
5803	7	*	*	*	*	*	*	*
5804	1	*	*	*	*	*	*	*
5806	16	*	*	*	*	*	*	*
6101	28	*	*	*	*	*	*	*
6102	27	*	*	*	*	*	*	*
6103	1,109	0.0	2.8	7.7	61.0	26.1	1.5	0.8
6105	3	*	*	*	*	*	*	*
6301	1,183	0.0	1.4	5.7	58.9	31.5	2.1	0.3
6312	583	0.0	0.3	3.1	49.6	42.9	3.9	0.2
6316	385	0.0	0.3	3.1	54.0	37.9	4.4	0.3
6320	191	0.0	0.0	4.2	56.5	36.6	2.1	0.5
6325	106	0.0	0.9	4.7	51.9	39.6	1.9	0.9
6335	322	0.0	0.3	2.2	54.7	38.5	3.7	0.6
6345	46	*	*	*	*	*	*	*
6360	256	0.0	0.4	2.0	61.3	34.4	1.2	0.8
7306	606	0.5	2.1	6.1	58.7	30.7	1.3	0.5

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2022 Pregnancy Nutrition Surveillance (1)

Page : 16

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
7401	629	0.0	1.3	9.2	59.6	27.8	1.1	1.0
7601	180	0.0	1.7	6.1	66.1	25.6	0.6	0.0
7850	363	0.3	2.2	6.1	64.7	25.3	1.4	0.0
7852	3	*	*	*	*	*	*	*
8101	591	0.3	1.9	5.2	55.0	35.0	1.4	1.2
8102	149	0.0	1.3	7.4	61.1	28.2	1.3	0.7
8106	197	0.0	1.0	4.6	59.4	34.0	0.5	0.5
8150	1	*	*	*	*	*	*	*
8151	1,100	0.0	1.5	5.1	55.5	34.3	3.2	0.4
8152	5	*	*	*	*	*	*	*
8153	1	*	*	*	*	*	*	*
8155	2	*	*	*	*	*	*	*
8201	897	0.0	1.4	4.3	57.0	34.6	2.0	0.7
8202	1,370	0.1	0.8	3.6	52.3	37.4	5.5	0.3
8203	103	0.0	1.0	2.9	55.3	36.9	3.9	0.0
8250	168	0.0	1.8	4.8	54.2	34.5	4.8	0.0
8252	48	*	*	*	*	*	*	*
8255	592	0.2	1.5	2.9	62.3	31.4	1.5	0.2
8257	543	0.0	0.7	2.8	55.6	37.6	3.1	0.2
8263	133	0.0	0.0	4.5	59.4	34.6	1.5	0.0
8264	270	0.4	1.5	3.3	57.4	33.3	3.7	0.4
8267	189	0.0	0.0	3.7	64.0	29.6	2.1	0.5
8901	283	0.0	3.5	4.9	58.3	31.1	1.8	0.4
8902	443	0.0	1.8	6.3	50.8	37.9	2.9	0.2

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2022 Pregnancy Nutrition Surveillance (1)

Page : 17

## Michigan

### Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8903	478	0.0	1.7	5.9	61.1	28.0	2.9	0.4
8904	673	0.1	3.3	6.2	57.7	30.3	1.6	0.7
8905	635	0.0	0.8	3.8	52.8	38.1	4.3	0.3
8909	396	0.0	1.8	6.1	56.6	30.6	4.5	0.5
9045	419	0.2	1.9	5.5	52.0	35.1	4.5	0.7
9050	320	0.0	1.3	5.6	60.3	30.0	1.9	0.9
9051	547	0.2	2.6	4.8	56.5	32.0	3.5	0.5
9053	525	0.0	1.5	4.4	58.3	30.9	4.2	0.8
9055	4	*	*	*	*	*	*	*
9058	479	0.2	1.5	6.5	54.1	33.8	3.3	0.6
9066	223	0.0	4.0	5.8	53.8	34.1	2.2	0.0
9068	243	0.8	1.2	7.0	55.1	30.9	2.9	2.1
9069	518	0.2	2.3	6.6	52.3	35.7	2.5	0.4
9070	664	0.2	3.3	7.4	55.1	30.9	2.0	1.2
9080	871	0.1	1.3	6.2	51.1	37.4	2.6	1.3
9082	24	*	*	*	*	*	*	*
9083	2	*	*	*	*	*	*	*
9087	3	*	*	*	*	*	*	*
<b>Michigan</b>	<b>47,924</b>	<b>0.1</b>	<b>1.6</b>	<b>5.9</b>	<b>58.5</b>	<b>30.8</b>	<b>2.3</b>	<b>0.7</b>
<b>National (Prior Year)</b>	<b>66,076</b>	<b>0.1</b>	<b>2.1</b>	<b>6.2</b>	<b>58.9</b>	<b>30.1</b>	<b>2.4</b>	<b>0.3</b>

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 18

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	32	*	*	18	*	*	2	*	1	*	32	*	32	*
0102	101	6.9 (98)	64.4 (72)	51	*	*	5	*	11	*	116	9.5 (63)	115	14.8 (56)
0103	99	*	*	53	*	*	10	*	7	*	108	7.4 (27)	108	18.5 (94)
0104	32	*	*	10	*	*	3	*	6	*	36	*	36	*
0250	34	*	*	20	*	*	4	*	10	*	36	*	36	*
0251	8	*	*	5	*	*	2	*	3	*	15	*	15	*
0252	33	*	*	23	*	*	4	*	2	*	34	*	34	*
0253	37	*	*	29	*	*	3	*	8	*	37	*	37	*
0401	132	6.8 (97)	56.1 (8)	48	*	*	9	*	9	*	131	9.2 (59)	131	13.0 (35)
0402	105	3.8 (39)	61.0 (31)	28	*	*	8	*	12	*	115	15.7 (108)	115	11.3 (18)
0403	36	*	*	18	*	*	0	*	2	*	39	*	39	*
0404	26	*	*	3	*	*	0	*	3	*	34	*	34	*
0501	88	*	*	17	*	*	22	*	32	*	123	8.9 (53)	123	13.0 (35)
0503	14	*	*	4	*	*	2	*	5	*	18	*	18	*
0504	91	*	*	36	*	*	17	*	16	*	103	6.8 (20)	103	9.7 (13)
0505	52	*	*	7	*	*	12	*	12	*	69	*	69	*
0506	45	*	*	9	*	*	9	*	8	*	57	*	57	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 19

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0507	6	*	*	2	*	*	2	*	8	*	16	*	16	*
0508	2	*	*	1	*	*	1	*	2	*	4	*	4	*
0650	331	2.7 (14)	65.9 (85)	262	14.5 (12)	58.4 (60)	9	*	19	*	355	12.1 (95)	355	19.7 (102)
0651	143	4.2 (51)	60.1 (22)	37	*	*	6	*	20	*	177	9.6 (66)	177	14.7 (54)
0652	79	*	*	60	*	*	8	*	15	*	88	*	88	*
0653	109	2.8 (18)	60.6 (27)	66	*	*	15	*	16	*	117	6.8 (20)	117	19.7 (102)
0654	53	*	*	28	*	*	3	*	7	*	133	9.8 (71)	133	12.0 (23)
0655	92	*	*	21	*	*	3	*	4	*	99	*	99	*
0657	3	*	*	2	*	*	0	*	1	*	8	*	8	*
0701	123	5.7 (90)	57.7 (10)	79	*	*	2	*	2	*	134	12.7 (100)	134	15.7 (68)
0702	21	*	*	15	*	*	0	*	2	*	19	*	19	*
0703	44	*	*	29	*	*	3	*	7	*	48	*	48	*
0704	12	*	*	9	*	*	2	*	3	*	16	*	16	*
0705	21	*	*	16	*	*	0	*	0	*	23	*	23	*
0850	139	4.3 (56)	60.4 (26)	43	*	*	2	*	0	*	230	13.9 (104)	231	14.7 (54)
0851	131	5.3 (81)	64.1 (69)	55	*	*	4	*	2	*	187	13.4 (103)	188	18.6 (96)
0901	344	6.7 (94)	58.7 (14)	156	11.5 (3)	54.5 (48)	10	*	17	*	460	8.7 (49)	460	14.8 (56)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 20

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0902	8	*	*	4	*	*	0	*	0	*	11	*	11	*
1051	297	5.4 (85)	54.2 (6)	210	12.4 (6)	56.2 (55)	9	*	18	*	339	6.8 (20)	340	13.8 (45)
1052	1	*	*	1	*	*	1	*	0	*	2	*	2	*
1101	396	4.0 (45)	61.4 (34)	271	19.6 (41)	48.7 (11)	17	*	49	*	399	9.5 (63)	399	18.3 (93)
1102	230	4.3 (56)	63.5 (59)	162	16.7 (23)	56.2 (55)	11	*	23	*	226	8.8 (51)	227	17.6 (87)
1106	21	*	*	18	*	*	0	*	0	*	20	*	20	*
1250	135	5.9 (92)	59.3 (16)	58	*	*	12	*	27	*	285	9.5 (63)	285	9.8 (14)
1251	184	4.9 (71)	59.8 (19)	135	15.6 (15)	53.3 (36)	11	*	26	*	188	5.9 (7)	188	6.4 (4)
1252	149	6.7 (94)	65.1 (79)	60	*	*	1	*	10	*	172	8.1 (36)	172	11.6 (20)
1253	186	5.4 (85)	61.8 (40)	83	*	*	11	*	14	*	202	10.9 (84)	203	16.3 (78)
1350	681	4.1 (50)	63.6 (61)	493	19.9 (44)	53.3 (36)	12	*	5	*	737	5.8 (5)	738	11.5 (19)
1352	101	5.0 (73)	62.4 (49)	73	*	*	0	*	1	*	121	4.1 (2)	121	9.1 (11)
1401	54	*	*	27	*	*	3	*	15	*	59	*	59	*
1402	17	*	*	8	*	*	2	*	2	*	22	*	22	*
1403	7	*	*	4	*	*	0	*	2	*	11	*	11	*
1501	185	3.2 (25)	70.3 (97)	82	*	*	36	*	46	*	215	11.6 (94)	216	13.4 (41)
1503	235	5.1 (75)	63.8 (65)	164	22.0 (56)	55.5 (53)	37	*	39	*	262	12.6 (98)	262	19.8 (104)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 21

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1504	207	5.3 (81)	62.8 (54)	78	*	*	92	*	121	26.4 (3)	573	9.8 (71)	573	7.9 (9)
1506	147	2.7 (14)	63.9 (66)	21	*	*	23	*	40	*	164	11.0 (86)	164	16.5 (80)
1507	123	11.4 (100)	59.3 (16)	14	*	*	11	*	42	*	149	8.7 (49)	150	12.7 (32)
1509	79	*	*	38	*	*	23	*	28	*	104	9.6 (66)	104	7.7 (8)
1511	156	6.4 (93)	67.3 (90)	90	*	*	31	*	62	*	202	10.4 (79)	202	14.4 (48)
1512	92	*	*	52	*	*	15	*	50	*	137	10.2 (74)	138	18.1 (92)
1514	151	5.3 (81)	51.7 (1)	37	*	*	27	*	92	*	238	8.8 (51)	239	12.1 (24)
1701	160	3.8 (39)	58.1 (12)	107	26.2 (61)	50.5 (20)	16	*	24	*	157	11.5 (93)	157	7.0 (5)
1950	165	3.0 (22)	61.8 (40)	116	11.2 (2)	60.3 (62)	0	*	9	*	184	10.9 (84)	185	15.1 (63)
1952	39	*	*	28	*	*	0	*	1	*	42	*	42	*
1953	66	*	*	45	*	*	1	*	3	*	72	*	72	*
1954	106	0.0 (1)	67.9 (92)	58	*	*	0	*	2	*	116	12.1 (95)	116	12.1 (24)
1955	41	*	*	26	*	*	0	*	1	*	38	*	38	*
1957	113	4.4 (60)	66.4 (87)	71	*	*	5	*	3	*	127	11.0 (86)	127	11.0 (16)
1963	14	*	*	8	*	*	0	*	0	*	17	*	17	*
2150	156	5.1 (75)	64.1 (69)	106	26.4 (62)	42.5 (3)	16	*	38	*	158	11.4 (91)	158	16.5 (80)
2151	64	*	*	52	*	*	8	*	21	*	59	*	59	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 22

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2152	13	*	*	4	*	*	2	*	1	*	14	*	14	*
2251	89	*	*	47	*	*	0	*	0	*	110	2.7 (1)	110	16.4 (79)
2252	29	*	*	8	*	*	0	*	0	*	56	*	56	*
2554	12	*	*	6	*	*	4	*	3	*	18	*	18	*
2555	1,727	3.4 (27)	62.9 (55)	871	19.9 (44)	54.0 (40)	6	*	14	*	2,237	9.7 (69)	2,239	17.6 (87)
2556	25	*	*	10	*	*	0	*	0	*	35	*	35	*
2557	1	*	*	0	*	*	0	*	0	*	1	*	1	*
2562	0	*	*	0	*	*	0	*	0	*	1	*	1	*
2563	0	*	*	0	*	*	0	*	0	*	1	*	1	*
2564	0	*	*	0	*	*	0	*	0	*	1	*	1	*
3001	220	3.6 (34)	64.5 (74)	57	*	*	0	*	0	*	393	8.4 (43)	393	17.6 (87)
3002	24	*	*	9	*	*	0	*	1	*	49	*	49	*
3004	45	*	*	15	*	*	0	*	0	*	91	*	91	*
3005	18	*	*	7	*	*	0	*	1	*	54	*	54	*
3007	1	*	*	0	*	*	0	*	0	*	1	*	1	*
3201	110	2.7 (14)	70.9 (98)	78	*	*	18	*	25	*	126	14.3 (106)	126	15.9 (74)
3301	1,136	3.4 (27)	60.3 (23)	650	16.2 (20)	51.7 (30)	19	*	17	*	1,363	7.2 (24)	1,363	13.1 (38)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 23

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3304	40	*	*	22	*	*	1	*	0	*	45	*	45	*
3305	56	*	*	34	*	*	0	*	0	*	64	*	64	*
3306	70	*	*	37	*	*	2	*	1	*	76	*	76	*
3308	35	*	*	21	*	*	0	*	0	*	35	*	35	*
3312	145	4.8 (68)	61.4 (34)	76	*	*	6	*	0	*	164	6.7 (18)	165	13.3 (39)
3550	21	*	*	14	*	*	0	*	0	*	28	*	28	*
3551	1	*	*	0	*	*	0	*	0	*	2	*	2	*
3701	203	5.4 (85)	63.1 (56)	146	23.3 (57)	47.3 (8)	23	*	42	*	253	14.2 (105)	253	19.0 (98)
3850	740	4.2 (51)	61.1 (32)	478	15.7 (17)	56.5 (57)	19	*	23	*	823	9.7 (69)	820	15.7 (68)
3901	358	5.6 (89)	59.2 (15)	158	17.1 (26)	55.1 (52)	0	*	2	*	413	8.2 (39)	411	14.4 (48)
3902	1	*	*	1	*	*	0	*	0	*	2	*	2	*
3950	467	4.7 (66)	61.5 (37)	262	17.2 (27)	51.1 (25)	0	*	2	*	604	11.4 (91)	606	12.5 (30)
3954	1	*	*	0	*	*	0	*	0	*	1	*	1	*
4106	250	2.8 (18)	60.8 (28)	51	*	*	118	30.5 (1)	165	20.0 (1)	662	9.1 (58)	664	16.7 (84)
4107	551	4.0 (45)	55.4 (7)	156	15.4 (14)	49.4 (15)	147	36.7 (2)	215	26.0 (2)	867	9.0 (54)	869	16.1 (77)
4109	107	4.7 (66)	56.1 (8)	71	*	*	33	*	68	*	181	4.4 (3)	182	18.7 (97)
4112	179	2.2 (8)	67.6 (91)	158	45.6 (64)	25.9 (1)	1	*	4	*	182	21.4 (110)	182	26.9 (110)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 24

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4115	563	3.4 (27)	59.7 (18)	199	15.6 (15)	54.3 (45)	179	39.1 (3)	299	26.8 (4)	1,077	10.5 (80)	1,076	19.0 (98)
4351	171	3.5 (31)	62.0 (43)	125	21.6 (54)	51.2 (27)	0	*	4	*	180	12.8 (101)	180	22.8 (109)
4352	61	*	*	39	*	*	1	*	2	*	64	*	64	*
4353	113	1.8 (6)	71.7 (99)	84	*	*	9	*	8	*	130	16.9 (109)	130	12.3 (28)
4354	158	2.5 (10)	65.8 (84)	61	*	*	0	*	3	*	236	8.1 (36)	236	21.6 (108)
4355	55	*	*	37	*	*	0	*	0	*	73	*	72	*
4356	124	1.6 (4)	70.2 (96)	86	*	*	1	*	2	*	128	10.2 (74)	128	18.0 (91)
4359	175	2.9 (20)	65.1 (79)	115	25.2 (60)	49.6 (16)	0	*	1	*	189	10.6 (81)	189	11.6 (20)
4360	76	*	*	49	*	*	2	*	1	*	72	*	72	*
4361	70	*	*	48	*	*	1	*	3	*	71	*	71	*
4362	84	*	*	58	*	*	3	*	1	*	81	*	81	*
4363	176	5.7 (90)	60.8 (28)	121	19.0 (38)	51.2 (27)	2	*	7	*	193	9.3 (60)	193	14.5 (52)
4401	219	5.0 (73)	64.4 (72)	184	26.6 (63)	52.7 (35)	0	*	4	*	222	5.9 (7)	222	7.2 (6)
4501	187	4.3 (56)	61.5 (37)	91	*	*	6	*	12	*	243	12.8 (101)	243	18.5 (94)
4601	354	4.0 (45)	64.7 (76)	231	16.5 (22)	51.1 (25)	0	*	16	*	362	8.3 (40)	361	17.7 (90)
4701	267	7.1 (99)	57.7 (10)	143	19.6 (41)	49.0 (14)	13	*	28	*	312	10.3 (77)	312	21.5 (107)
5050	549	4.2 (51)	61.4 (34)	470	17.4 (28)	50.2 (18)	0	*	4	*	542	10.3 (77)	542	14.9 (59)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 25

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5051	807	4.0 (45)	61.1 (32)	641	20.0 (47)	51.8 (32)	1	*	3	*	809	8.5 (45)	809	13.0 (35)
5052	870	3.7 (37)	62.4 (49)	726	20.7 (50)	46.8 (7)	0	*	1	*	903	12.6 (98)	903	13.3 (39)
5055	414	1.0 (3)	62.3 (46)	343	24.8 (59)	44.6 (6)	0	*	0	*	446	14.6 (107)	446	5.2 (2)
5101	334	2.1 (7)	68.0 (94)	271	12.2 (5)	55.0 (51)	32	*	55	*	426	7.5 (29)	429	12.1 (24)
5102	1	*	*	0	*	*	0	*	0	*	1	*	1	*
5202	180	6.7 (94)	53.9 (4)	116	20.7 (50)	56.9 (58)	1	*	1	*	205	12.2 (97)	203	15.8 (72)
5206	7	*	*	5	*	*	0	*	1	*	7	*	7	*
5601	247	4.9 (71)	61.9 (42)	120	10.8 (1)	54.2 (43)	16	*	18	*	318	11.3 (90)	318	14.8 (56)
5801	589	5.3 (81)	64.2 (71)	452	19.9 (44)	52.4 (33)	17	*	15	*	588	8.5 (45)	588	16.5 (80)
5803	7	*	*	6	*	*	0	*	0	*	7	*	7	*
5804	1	*	*	1	*	*	0	*	0	*	1	*	1	*
5806	14	*	*	8	*	*	0	*	2	*	16	*	16	*
6101	26	*	*	19	*	*	0	*	0	*	25	*	25	*
6102	23	*	*	20	*	*	1	*	0	*	27	*	27	*
6103	1,008	3.8 (39)	65.3 (81)	749	18.0 (30)	54.1 (42)	7	*	6	*	1,030	8.6 (47)	1,030	20.3 (105)
6105	1	*	*	0	*	*	0	*	0	*	2	*	2	*
6301	1,042	3.9 (43)	62.4 (49)	717	16.7 (23)	50.3 (19)	4	*	24	*	1,090	5.4 (4)	1,091	15.5 (67)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 26

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6312	520	3.1 (24)	64.0 (67)	354	18.4 (33)	49.7 (17)	3	*	18	*	532	9.0 (54)	535	10.1 (15)
6316	352	4.8 (68)	54.0 (5)	238	17.6 (29)	52.5 (34)	4	*	9	*	355	7.9 (34)	355	12.7 (32)
6320	139	2.9 (20)	66.2 (86)	64	*	*	0	*	9	*	173	6.4 (15)	174	16.7 (84)
6325	77	*	*	41	*	*	0	*	6	*	102	6.9 (23)	102	19.6 (101)
6335	248	5.2 (79)	65.3 (81)	139	15.8 (19)	54.0 (40)	10	*	27	*	310	11.0 (86)	309	14.9 (59)
6345	34	*	*	18	*	*	0	*	3	*	42	*	42	*
6360	186	4.8 (68)	58.6 (13)	95	*	*	7	*	18	*	241	8.3 (40)	242	15.3 (65)
7306	454	2.6 (12)	64.8 (77)	305	18.7 (36)	50.8 (24)	1	*	29	*	580	10.7 (82)	582	12.2 (27)
7401	584	4.6 (65)	63.7 (64)	464	19.6 (41)	51.3 (29)	33	*	80	*	559	9.8 (71)	561	16.0 (75)
7601	47	*	*	6	*	*	4	*	14	*	166	9.0 (54)	166	15.1 (63)
7850	310	4.2 (51)	62.3 (46)	248	21.8 (55)	48.8 (13)	0	*	2	*	336	6.3 (13)	336	9.2 (12)
7852	3	*	*	1	*	*	0	*	0	*	2	*	2	*
8101	474	3.0 (22)	62.7 (53)	303	14.5 (12)	54.8 (49)	0	*	1	*	556	6.7 (18)	555	16.9 (86)
8102	115	3.5 (31)	63.5 (59)	58	*	*	0	*	0	*	135	5.9 (7)	135	7.4 (7)
8106	89	*	*	23	*	*	0	*	0	*	189	7.4 (27)	189	19.0 (98)
8150	1	*	*	0	*	*	0	*	0	*	1	*	1	*
8151	848	4.2 (51)	61.6 (39)	524	24.4 (58)	43.5 (4)	100	44.0 (4)	185	28.1 (5)	977	8.6 (47)	978	14.5 (52)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 27

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8152	4	*	*	2	*	*	0	*	2	*	5	*	5	*
8153	1	*	*	0	*	*	1	*	0	*	0	*	0	*
8155	1	*	*	0	*	*	0	*	0	*	1	*	1	*
8201	719	3.2 (25)	64.5 (74)	422	18.5 (34)	53.6 (39)	0	*	8	*	798	9.6 (66)	798	14.9 (59)
8202	532	3.6 (34)	53.8 (3)	275	18.9 (37)	47.3 (8)	0	*	0	*	1,326	8.0 (35)	1,325	3.5 (1)
8203	85	*	*	54	*	*	0	*	1	*	92	*	92	*
8250	135	3.7 (37)	66.7 (88)	87	*	*	0	*	2	*	145	7.6 (31)	146	14.4 (48)
8252	28	*	*	15	*	*	0	*	0	*	44	*	45	*
8255	483	5.2 (79)	60.9 (30)	327	21.1 (53)	50.5 (20)	0	*	1	*	535	9.0 (54)	535	15.0 (62)
8257	332	5.4 (85)	52.1 (2)	202	20.3 (48)	44.1 (5)	0	*	2	*	518	9.3 (60)	519	5.4 (3)
8263	116	2.6 (12)	60.3 (23)	62	*	*	0	*	0	*	111	8.1 (36)	111	14.4 (48)
8264	224	4.0 (45)	62.1 (45)	144	11.8 (4)	63.9 (64)	0	*	1	*	253	7.5 (29)	253	13.8 (45)
8267	157	0.6 (2)	70.1 (95)	96	*	*	0	*	1	*	177	10.7 (82)	177	15.8 (72)
8901	228	4.4 (60)	63.6 (61)	147	14.3 (11)	54.4 (46)	0	*	0	*	275	6.5 (16)	273	13.6 (43)
8902	358	3.9 (43)	62.3 (46)	213	19.2 (39)	54.9 (50)	1	*	0	*	421	7.6 (31)	419	13.6 (43)
8903	365	2.7 (14)	67.9 (92)	230	13.9 (9)	60.9 (63)	0	*	0	*	443	6.3 (13)	447	15.7 (68)
8904	555	3.8 (39)	66.8 (89)	344	15.7 (17)	51.7 (30)	0	*	2	*	640	7.3 (26)	641	15.3 (65)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2022 Pregnancy Nutrition Surveillance (1)

Page : 28

Michigan

## Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8905	448	4.5 (63)	60.0 (20)	238	18.5 (34)	54.2 (43)	0	*	0	*	605	11.2 (89)	605	11.2 (17)
8909	315	4.4 (60)	63.2 (57)	178	12.9 (7)	60.1 (61)	0	*	2	*	370	5.9 (7)	369	15.7 (68)
9045	308	3.6 (34)	72.4 (100)	184	14.1 (10)	53.3 (36)	0	*	0	*	392	8.4 (43)	392	8.7 (10)
9050	287	4.5 (63)	62.4 (49)	83	*	*	0	*	0	*	307	7.2 (24)	307	14.0 (47)
9051	250	1.6 (4)	64.8 (77)	119	21.0 (52)	42.0 (2)	0	*	0	*	530	9.4 (62)	530	12.8 (34)
9053	463	2.2 (8)	60.3 (23)	359	13.6 (8)	57.1 (59)	0	*	0	*	504	6.2 (12)	503	12.5 (30)
9055	4	*	*	4	*	*	0	*	0	*	4	*	4	*
9058	297	3.4 (27)	64.0 (67)	212	19.3 (40)	50.5 (20)	0	*	0	*	458	7.6 (31)	457	16.0 (75)
9066	198	5.1 (75)	63.6 (61)	133	20.3 (48)	47.4 (10)	0	*	0	*	214	6.5 (16)	215	20.9 (106)
9068	226	3.5 (31)	65.5 (83)	187	18.2 (31)	55.6 (54)	0	*	0	*	233	6.0 (11)	233	12.4 (29)
9069	397	4.3 (56)	62.0 (43)	269	16.4 (21)	48.7 (11)	0	*	0	*	494	8.3 (40)	495	13.5 (42)
9070	587	5.1 (75)	60.0 (20)	451	16.9 (25)	50.6 (23)	0	*	0	*	623	5.8 (5)	624	11.9 (22)
9080	647	2.5 (10)	63.2 (57)	349	18.3 (32)	54.4 (46)	0	*	1	*	833	10.2 (74)	833	16.6 (83)
9082	19	*	*	11	*	*	0	*	0	*	24	*	24	*
9083	2	*	*	2	*	*	0	*	0	*	2	*	2	*
9087	2	*	*	1	*	*	0	*	0	*	3	*	3	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 5A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 29

#### Michigan

#### Comparison of Maternal Health Indicators by Clinic

<b>Michigan</b>	<b>36,308</b>	<b>4.0</b>	<b>62.2</b>	<b>22,132</b>	<b>18.5</b>	<b>51.9</b>	<b>1,472</b>	<b>31.5</b>	<b>2,621</b>	<b>25.6</b>	<b>44,703</b>	<b>9.0</b>	<b>44,728</b>	<b>14.3</b>
<b>National (Prior Year)</b>	<b>48,187</b>	<b>3.7</b>	<b>62.1</b>	<b>33,280</b>	<b>19.3</b>	<b>50.7</b>	<b>2,408</b>	<b>26.4</b>	<b>2,474</b>	<b>22.5</b>	<b>48,523</b>	<b>10.6</b>	<b>48,164</b>	<b>13.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2022 Pregnancy Nutrition Surveillance (1)

Page : 30

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
0101	28	*	*	36	*	36	*	30 *
0102	114	91.2 (68)	0.9 (82)	116	31.9 (32)	122	19.7 (80)	90 *
0103	106	92.5 (53)	0.0 (1)	111	36.9 (15)	112	18.8 (85)	85 *
0104	35	*	*	40	*	42	*	34 *
0250	31	*	*	38	*	38	*	27 *
0251	15	*	*	16	*	16	*	12 *
0252	34	*	*	38	*	38	*	29 *
0253	35	*	*	37	*	37	*	26 *
0401	123	96.7 (12)	0.0 (1)	140	27.1 (54)	137	24.8 (57)	98 *
0402	108	89.8 (80)	0.9 (82)	128	26.6 (59)	129	36.4 (16)	102 97.1 (5)
0403	33	*	*	39	*	40	*	31 *
0404	31	*	*	36	*	36	*	34 *
0501	117	95.7 (18)	0.0 (1)	128	23.4 (75)	121	30.6 (29)	88 *
0503	18	*	*	16	*	18	*	11 *
0504	100	92.0 (61)	0.0 (1)	98	*	102	48.0 (3)	72 *
0505	66	*	*	72	*	71	*	57 *
0506	53	*	*	59	*	61	*	46 *
0507	16	*	*	15	*	16	*	7 *
0508	4	*	*	4	*	4	*	1 *
0650	328	94.5 (31)	0.3 (43)	365	39.2 (11)	364	32.7 (20)	296 91.9 (50)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 31

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0651	165	90.9 (70)	0.6 (65)	193	37.8 (13)	188	30.9 (28)	155	89.7 (74)
0652	79	*	*	85	*	86	*	60	*
0653	103	92.2 (57)	1.9 (106)	124	39.5 (9)	123	36.6 (15)	100	91.0 (60)
0654	104	93.3 (41)	1.9 (106)	141	41.1 (6)	139	14.4 (92)	112	91.1 (59)
0655	98	*	*	100	37.0 (14)	86	*	83	*
0657	7	*	*	9	*	9	*	8	*
0701	114	93.0 (47)	0.9 (82)	145	34.5 (22)	143	37.8 (11)	112	90.2 (70)
0702	17	*	*	21	*	21	*	15	*
0703	48	*	*	51	*	51	*	35	*
0704	15	*	*	16	*	16	*	14	*
0705	22	*	*	23	*	24	*	14	*
0850	224	92.9 (50)	0.4 (51)	243	27.6 (48)	244	31.1 (27)	177	93.8 (32)
0851	185	93.5 (38)	0.0 (1)	190	25.8 (64)	194	37.1 (13)	144	95.8 (14)
0901	397	91.9 (63)	1.0 (88)	477	35.8 (19)	479	28.4 (38)	375	92.5 (43)
0902	8	*	*	11	*	11	*	9	*
1051	301	93.0 (47)	0.7 (68)	357	28.3 (44)	357	32.5 (21)	243	90.5 (68)
1052	2	*	*	1	*	2	*	1	*
1101	357	82.9 (107)	0.6 (65)	444	35.1 (21)	426	24.6 (58)	320	87.8 (84)
1102	190	85.8 (98)	0.5 (57)	253	32.4 (29)	248	24.2 (60)	187	96.3 (10)
1106	19	*	*	21	*	21	*	14	*
1250	272	90.8 (71)	0.0 (1)	293	26.3 (63)	291	24.4 (59)	212	92.9 (39)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 32

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1251	186	96.2 (16)	0.5 (57)	190	20.5 (88)	198	73.2 (1)	127	96.1 (12)
1252	128	86.7 (95)	0.8 (75)	184	33.2 (27)	186	19.9 (79)	137	91.2 (57)
1253	162	87.7 (92)	1.2 (98)	215	36.7 (16)	216	26.4 (47)	172	91.3 (55)
1350	715	96.9 (11)	0.3 (43)	772	32.3 (30)	772	35.9 (18)	577	92.5 (43)
1352	118	98.3 (4)	0.0 (1)	130	33.8 (25)	128	8.6 (99)	90	*
1401	56	*	*	60	*	58	*	37	*
1402	22	*	*	24	*	22	*	18	*
1403	10	*	*	11	*	11	*	8	*
1501	174	88.5 (88)	1.1 (92)	237	26.6 (59)	229	8.3 (101)	164	87.2 (87)
1503	249	98.4 (3)	0.0 (1)	272	33.1 (28)	269	24.2 (60)	202	88.6 (80)
1504	553	95.7 (18)	0.4 (51)	606	26.6 (59)	606	18.3 (86)	408	90.7 (65)
1506	133	95.5 (22)	0.0 (1)	174	27.0 (56)	167	10.2 (97)	124	93.5 (34)
1507	151	94.0 (34)	1.3 (100)	163	30.1 (40)	158	29.1 (36)	102	98.0 (2)
1509	97	*	*	118	28.0 (45)	114	21.9 (71)	88	*
1511	190	91.6 (65)	0.0 (1)	213	31.9 (32)	216	26.9 (45)	150	82.0 (96)
1512	131	95.4 (25)	0.8 (75)	140	17.1 (102)	139	27.3 (42)	77	*
1514	212	94.3 (33)	0.9 (82)	254	27.2 (52)	212	29.2 (35)	159	93.1 (35)
1701	158	94.9 (26)	0.0 (1)	168	23.8 (72)	162	61.7 (2)	128	94.5 (26)
1950	165	93.3 (41)	0.0 (1)	191	27.7 (46)	163	36.2 (17)	137	97.1 (5)
1952	38	*	*	43	*	32	*	26	*
1953	63	*	*	76	*	66	*	55	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2022 Pregnancy Nutrition Surveillance (1)

Page : 33

Michigan

## Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1954	105	92.4 (55)	1.0 (88)	119	40.3 (7)	102	42.2 (6)	102	97.1 (5)
1955	33	*	*	43	*	33	*	31	*
1957	114	92.1 (59)	0.0 (1)	131	31.3 (35)	109	44.0 (5)	98	*
1963	15	*	*	17	*	14	*	14	*
2150	159	93.1 (45)	0.6 (65)	165	32.1 (31)	166	25.9 (51)	117	94.0 (31)
2151	55	*	*	72	*	55	*	41	*
2152	14	*	*	16	*	16	*	11	*
2251	104	98.1 (5)	0.0 (1)	118	39.0 (12)	118	19.5 (81)	98	*
2252	50	*	*	60	*	58	*	50	*
2554	18	*	*	19	*	14	*	15	*
2555	2,077	90.6 (75)	0.4 (51)	2,351	24.0 (71)	1,959	20.5 (75)	1,707	92.9 (39)
2556	30	*	*	35	*	28	*	28	*
2557	1	*	*	1	*	1	*	1	*
2562	1	*	*	1	*	1	*	1	*
2563	1	*	*	2	*	1	*	2	*
2564	1	*	*	2	*	1	*	2	*
3001	362	89.5 (81)	0.3 (43)	404	33.9 (24)	401	10.2 (97)	318	85.2 (93)
3002	47	*	*	48	*	51	*	36	*
3004	87	*	*	89	*	92	*	62	*
3005	50	*	*	54	*	55	*	46	*
3007	1	*	*	1	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 34

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3201	113	95.6 (21)	0.9 (82)	132	23.5 (74)	131	41.2 (7)	84	*
3301	1,192	90.2 (78)	0.9 (82)	1,410	26.5 (62)	1,345	13.3 (93)	1,031	90.4 (69)
3304	39	*	*	47	*	47	*	41	*
3305	56	*	*	65	*	65	*	46	*
3306	66	*	*	77	*	75	*	57	*
3308	33	*	*	38	*	37	*	22	*
3312	140	91.4 (67)	0.0 (1)	173	27.2 (52)	172	12.8 (94)	134	91.0 (60)
3550	24	*	*	27	*	28	*	22	*
3551	2	*	*	3	*	3	*	2	*
3701	244	95.5 (22)	0.0 (1)	271	27.3 (51)	273	27.1 (44)	191	95.3 (19)
3850	721	91.5 (66)	0.1 (35)	871	35.9 (18)	880	25.9 (51)	677	87.7 (85)
3901	373	92.0 (61)	0.0 (1)	449	34.3 (23)	448	15.8 (88)	339	84.4 (94)
3902	2	*	*	2	*	2	*	2	*
3950	562	88.8 (84)	0.5 (57)	620	30.0 (41)	615	20.5 (75)	448	92.0 (49)
3954	1	*	*	1	*	1	*	1	*
4106	572	88.1 (90)	1.4 (102)	708	27.0 (56)	690	14.6 (91)	535	83.2 (95)
4107	830	90.8 (71)	0.5 (57)	932	27.4 (50)	907	26.2 (49)	681	92.2 (48)
4109	174	94.8 (27)	1.7 (105)	185	21.6 (84)	185	22.2 (70)	108	94.4 (27)
4112	181	68.0 (108)	0.0 (1)	182	19.8 (91)	10	*	160	92.5 (43)
4115	951	92.3 (56)	0.8 (75)	1,108	22.7 (78)	904	21.2 (73)	766	91.4 (54)
4351	152	88.8 (84)	1.3 (100)	190	39.5 (9)	191	27.2 (43)	141	91.5 (52)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 35

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
4352	55	*	*	69	*	69	*	54 *
4353	95	*	*	144	43.8 (3)	141	26.2 (49)	110 88.2 (83)
4354	214	96.7 (12)	0.0 (1)	242	49.2 (1)	236	19.5 (81)	189 91.5 (52)
4355	65	*	*	77	*	75	*	63 *
4356	116	88.8 (84)	0.0 (1)	129	24.8 (67)	132	28.0 (39)	96 *
4359	140	89.3 (82)	0.7 (68)	195	43.1 (4)	204	27.5 (41)	158 89.9 (72)
4360	62	*	*	78	*	77	*	61 *
4361	68	*	*	75	*	75	*	65 *
4362	73	*	*	89	*	89	*	66 *
4363	180	90.6 (75)	1.1 (92)	202	27.7 (46)	205	31.7 (24)	142 93.0 (37)
4401	194	88.1 (90)	1.5 (104)	225	31.6 (34)	227	28.6 (37)	156 95.5 (18)
4501	224	95.5 (22)	0.0 (1)	244	27.5 (49)	250	31.2 (26)	173 89.6 (76)
4601	354	93.8 (35)	0.3 (43)	396	29.8 (42)	402	16.9 (87)	276 94.6 (25)
4701	284	95.8 (17)	0.7 (68)	326	29.8 (42)	332	38.9 (8)	226 94.7 (24)
5050	520	93.8 (35)	0.4 (51)	561	22.6 (79)	553	6.3 (104)	376 98.1 (1)
5051	767	93.4 (39)	0.4 (51)	840	21.2 (86)	810	7.2 (103)	584 95.0 (20)
5052	858	93.8 (35)	0.2 (39)	905	20.7 (87)	872	5.3 (105)	610 94.4 (27)
5055	407	99.8 (2)	0.0 (1)	458	44.5 (2)	121	5.0 (106)	384 95.8 (14)
5101	411	92.5 (53)	0.2 (39)	435	21.4 (85)	228	44.3 (4)	295 96.3 (10)
5102	1	*	*	2	*	1	*	2 *
5202	179	97.2 (8)	0.0 (1)	215	36.7 (16)	193	36.8 (14)	164 95.7 (17)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 36

### Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
5206	4	*	*	7	*	6	*	7 *
5601	300	94.7 (30)	0.7 (68)	320	30.9 (37)	326	23.3 (64)	231 94.8 (22)
5801	538	93.1 (45)	0.7 (68)	630	30.2 (39)	597	22.6 (69)	455 93.0 (37)
5803	7	*	*	7	*	6	*	6 *
5804	1	*	*	1	*	1	*	1 *
5806	16	*	*	16	*	16	*	12 *
6101	23	*	*	27	*	28	*	20 *
6102	25	*	*	25	*	27	*	15 *
6103	940	92.7 (52)	0.3 (43)	1,086	35.7 (20)	1,060	23.5 (62)	832 92.3 (47)
6105	1	*	*	3	*	3	*	2 *
6301	1,037	94.5 (31)	0.2 (39)	1,160	24.6 (69)	1,130	22.9 (67)	795 90.6 (66)
6312	500	93.4 (39)	0.0 (1)	567	25.0 (66)	557	23.5 (62)	381 92.7 (42)
6316	338	96.4 (15)	0.0 (1)	372	24.7 (68)	374	25.9 (51)	259 93.1 (35)
6320	169	91.7 (64)	0.0 (1)	180	16.1 (106)	179	37.4 (12)	115 97.4 (3)
6325	101	98.0 (6)	0.0 (1)	102	15.7 (109)	102	32.4 (22)	62 *
6335	294	96.6 (14)	0.3 (43)	308	25.6 (65)	277	30.0 (32)	217 97.2 (4)
6345	40	*	*	44	*	45	*	25 *
6360	219	92.2 (57)	0.5 (57)	246	23.2 (76)	233	25.8 (54)	176 94.3 (29)
7306	538	92.8 (51)	0.0 (1)	597	31.3 (35)	593	29.3 (34)	426 93.7 (33)
7401	436	90.4 (77)	1.4 (102)	614	42.3 (5)	617	27.6 (40)	488 90.6 (66)
7601	161	100.0 (1)	0.0 (1)	176	39.8 (8)	178	32.0 (23)	143 96.5 (9)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 6A

2022 Pregnancy Nutrition Surveillance (1)

Page : 37

Michigan

## Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)								
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number % (rank)
7850	326	97.2 (8)	0.3 (43)	353	33.7 (26)	44	*	264 85.6 (91)
7852	2	*	*	3	*	1	*	3 *
8101	525	83.8 (104)	0.8 (75)	578	18.0 (96)	486	22.8 (68)	369 85.4 (92)
8102	127	92.1 (59)	0.0 (1)	147	24.5 (70)	133	21.1 (74)	113 89.4 (77)
8106	184	90.2 (78)	0.5 (57)	192	16.1 (106)	196	23.0 (66)	142 95.8 (14)
8150	1	*	*	1	*	1	*	1 *
8151	927	93.2 (44)	0.1 (35)	1,079	27.1 (54)	1,072	19.1 (83)	808 92.8 (41)
8152	5	*	*	5	*	5	*	5 *
8153	0	*	*	1	*	1	*	1 *
8155	1	*	*	2	*	2	*	2 *
8201	764	94.8 (27)	0.1 (35)	883	19.8 (91)	704	38.5 (9)	629 87.1 (88)
8202	1,206	97.8 (7)	0.1 (35)	1,331	30.4 (38)	1,320	3.6 (108)	1,057 97.1 (5)
8203	89	*	*	103	20.4 (89)	95	*	69 *
8250	145	86.2 (96)	0.7 (68)	167	13.2 (111)	135	26.7 (46)	116 89.7 (74)
8252	45	*	*	47	*	47	*	34 *
8255	526	93.0 (47)	0.8 (75)	579	17.6 (99)	540	33.0 (19)	373 91.2 (57)
8257	506	95.7 (18)	0.0 (1)	531	21.7 (82)	402	8.5 (100)	399 69.7 (97)
8263	108	97.2 (8)	0.0 (1)	133	21.8 (81)	122	30.3 (30)	133 92.5 (43)
8264	250	94.8 (27)	0.8 (75)	263	14.8 (110)	235	37.9 (10)	166 87.3 (86)
8267	174	90.8 (71)	1.1 (92)	183	19.7 (93)	163	30.1 (31)	123 86.2 (90)
8901	261	85.4 (100)	0.8 (75)	277	19.1 (94)	168	25.0 (56)	184 90.8 (64)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2022 Pregnancy Nutrition Surveillance (1)

Page : 38

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8902	414	88.9 (83)	0.5 (57)	427	17.8 (98)	380	20.0 (78)	295	89.2 (78)
8903	417	84.4 (102)	0.7 (68)	469	21.7 (82)	363	21.5 (72)	337	88.4 (81)
8904	607	83.4 (106)	1.0 (88)	660	17.6 (99)	532	23.3 (64)	431	91.0 (60)
8905	580	93.3 (41)	0.2 (39)	617	23.7 (73)	551	31.6 (25)	407	90.2 (70)
8909	349	87.4 (93)	0.3 (43)	389	18.0 (96)	315	26.3 (48)	261	91.6 (51)
9045	379	90.8 (71)	1.1 (92)	411	22.1 (80)	170	7.6 (102)	306	96.1 (12)
9050	279	83.9 (103)	0.4 (51)	318	23.0 (77)	298	25.8 (54)	227	89.9 (72)
9051	511	91.0 (69)	0.0 (1)	526	20.0 (90)	468	4.5 (107)	395	94.2 (30)
9053	493	87.4 (93)	1.0 (88)	512	15.8 (108)	413	19.1 (83)	293	94.9 (21)
9055	4	*	*	4	*	4	*	1	*
9058	435	86.2 (96)	1.1 (92)	468	16.2 (105)	450	11.3 (95)	246	87.0 (89)
9066	193	85.5 (99)	0.0 (1)	222	27.0 (56)	207	20.3 (77)	222	91.0 (60)
9068	228	83.8 (104)	2.2 (108)	238	18.1 (95)	234	15.0 (89)	159	88.7 (79)
9069	492	84.6 (101)	1.2 (98)	509	16.5 (103)	364	10.7 (96)	305	94.8 (22)
9070	596	88.4 (89)	0.5 (57)	653	16.5 (103)	592	14.9 (90)	416	91.3 (55)
9080	798	88.6 (87)	1.1 (92)	850	17.5 (101)	601	29.8 (33)	550	88.4 (81)
9082	23	*	*	24	*	20	*	14	*
9083	2	*	*	2	*	2	*	2	*
9087	3	*	*	3	*	3	*	3	*
<b>Michigan</b>	<b>41,530</b>	<b>91.7</b>	<b>0.5</b>	<b>46,804</b>	<b>26.8</b>	<b>42,624</b>	<b>23.1</b>	<b>33,621</b>	<b>33621.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2022 Pregnancy Nutrition Surveillance (1)

Page : 39

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic

National (Prior Year)	53,284	81.8	3.2	52,379	24.5	48,576	23.1	44,162	92.4
-----------------------	--------	------	-----	--------	------	--------	------	--------	------

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2022 Pregnancy Nutrition Surveillance (1)

Page : 40

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	30	*	30	*	8	*	30	*	32	*
0102	90	*	113	16.8 (98)	31	*	90	*	117	9.4 (106)
0103	85	*	105	18.1 (101)	35	*	85	*	109	5.5 (86)
0104	34	*	33	*	8	*	34	*	36	*
0250	27	*	35	*	6	*	27	*	36	*
0251	12	*	15	*	4	*	12	*	15	*
0252	29	*	34	*	6	*	29	*	34	*
0253	26	*	36	*	12	*	26	*	37	*
0401	97	*	126	19.0 (102)	36	*	98	*	131	4.6 (78)
0402	101	31.7 (87)	112	14.3 (90)	29	*	102	10.8 (86)	115	8.7 (105)
0403	31	*	37	*	10	*	31	*	39	*
0404	34	*	34	*	5	*	34	*	34	*
0501	87	*	124	16.9 (99)	28	*	88	*	127	7.9 (102)
0503	11	*	18	*	2	*	11	*	18	*
0504	72	*	102	17.6 (100)	20	*	71	*	100	5.0 (79)
0505	57	*	69	*	14	*	57	*	69	*
0506	46	*	56	*	18	*	46	*	58	*
0507	7	*	16	*	2	*	7	*	16	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2022 Pregnancy Nutrition Surveillance (1)

Page : 41

Michigan

## Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0508	1	*	4	*	1	*	1	*	4	*
0650	295	22.7 (68)	351	12.5 (79)	65	*	296	7.8 (68)	354	5.4 (83)
0651	154	39.6 (96)	173	24.3 (109)	55	*	154	15.6 (97)	176	8.0 (103)
0652	60	*	87	*	9	*	60	*	88	*
0653	100	38.0 (94)	114	20.2 (104)	36	*	100	15.0 (96)	117	12.0 (111)
0654	111	27.9 (80)	133	23.3 (108)	30	*	112	12.5 (89)	133	3.8 (63)
0655	83	*	91	*	38	*	83	*	100	10.0 (108)
0657	8	*	8	*	2	*	8	*	8	*
0701	112	29.5 (84)	134	19.4 (103)	31	*	112	7.1 (62)	134	6.0 (91)
0702	15	*	19	*	1	*	15	*	19	*
0703	35	*	48	*	11	*	35	*	48	*
0704	14	*	16	*	3	*	14	*	16	*
0705	14	*	23	*	8	*	14	*	23	*
0850	176	22.2 (66)	231	7.8 (59)	38	*	177	10.2 (83)	231	2.6 (36)
0851	143	17.5 (48)	188	10.6 (73)	25	*	143	14.7 (95)	188	5.3 (82)
0901	374	28.3 (82)	451	10.2 (67)	102	42.2 (7)	375	9.1 (76)	458	5.9 (90)
0902	9	*	10	*	4	*	9	*	10	*
1051	242	33.5 (91)	332	15.1 (93)	71	*	243	6.6 (61)	341	6.7 (97)
1052	1	*	2	*	1	*	1	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2022 Pregnancy Nutrition Surveillance (1)

Page : 42

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1101	319	14.7 (40)	398	5.8 (44)	41	*	319	9.1 (76)	398	5.8 (89)
1102	186	33.3 (89)	222	16.7 (97)	56	*	187	14.4 (94)	227	10.6 (110)
1106	14	*	20	*	4	*	14	*	20	*
1250	212	19.3 (50)	282	13.5 (85)	37	*	212	4.7 (46)	285	4.2 (72)
1251	127	25.2 (74)	187	21.4 (105)	28	*	127	4.7 (46)	188	7.4 (99)
1252	137	19.7 (54)	170	9.4 (65)	25	*	137	5.1 (50)	171	3.5 (60)
1253	171	25.7 (75)	201	13.4 (84)	38	*	171	10.5 (85)	204	5.4 (83)
1350	576	21.7 (63)	729	6.6 (49)	103	43.7 (5)	577	2.8 (17)	736	3.8 (63)
1352	90	*	117	6.0 (45)	16	*	90	*	120	0.8 (4)
1401	38	*	58	*	9	*	38	*	59	*
1402	18	*	22	*	4	*	18	*	23	*
1403	8	*	11	*	3	*	8	*	11	*
1501	163	19.6 (53)	212	5.7 (42)	30	*	164	4.3 (34)	216	4.2 (72)
1503	201	21.4 (62)	261	7.3 (55)	43	*	202	2.0 (12)	262	1.9 (23)
1504	406	13.1 (32)	571	4.2 (34)	52	*	407	2.7 (16)	572	1.6 (18)
1506	124	19.4 (52)	159	5.7 (42)	19	*	124	1.6 (8)	164	2.4 (34)
1507	101	36.6 (93)	145	11.0 (74)	34	*	100	13.0 (91)	153	3.9 (68)
1509	88	*	105	10.5 (71)	12	*	88	*	104	2.9 (46)
1511	148	27.7 (78)	197	14.7 (92)	38	*	150	12.7 (90)	202	8.4 (104)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2022 Pregnancy Nutrition Surveillance (1)

Page : 43

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1512	77	*	139	12.2 (78)	11	*	77	*	139	3.6 (62)
1514	158	13.3 (33)	239	6.3 (47)	19	*	158	3.8 (28)	240	1.3 (12)
1701	128	29.7 (86)	159	21.4 (105)	34	*	127	11.8 (88)	158	3.2 (52)
1950	137	32.8 (88)	176	9.7 (66)	43	*	137	10.9 (87)	185	7.6 (101)
1952	26	*	41	*	4	*	26	*	42	*
1953	55	*	70	*	16	*	55	*	72	*
1954	99	*	110	7.3 (55)	27	*	102	5.9 (55)	116	6.0 (91)
1955	31	*	33	*	11	*	31	*	38	*
1957	98	*	123	14.6 (91)	32	*	98	*	127	1.6 (18)
1963	14	*	17	*	2	*	14	*	17	*
2150	117	33.3 (89)	159	21.4 (105)	37	*	116	7.8 (68)	158	6.3 (95)
2151	40	*	59	*	15	*	41	*	59	*
2152	11	*	14	*	3	*	11	*	14	*
2251	98	*	107	15.9 (95)	24	*	97	*	110	2.7 (40)
2252	50	*	56	*	16	*	50	*	57	*
2554	15	*	17	*	2	*	15	*	18	*
2555	1,706	20.0 (56)	2,211	8.1 (60)	309	43.4 (6)	1,708	7.3 (64)	2,239	4.0 (69)
2556	28	*	34	*	10	*	28	*	35	*
2557	1	*	1	*	0	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2022 Pregnancy Nutrition Surveillance (1)

Page : 44

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2562	1	*	1	*	0	*	1	*	1	*
2563	2	*	1	*	1	*	2	*	1	*
2564	2	*	1	*	0	*	2	*	1	*
3001	318	20.1 (57)	386	8.8 (62)	63	*	318	8.8 (74)	392	3.8 (63)
3002	36	*	49	*	6	*	36	*	48	*
3004	62	*	91	*	11	*	62	*	91	*
3005	46	*	54	*	13	*	46	*	54	*
3007	1	*	1	*	0	*	1	*	1	*
3201	84	*	125	13.6 (87)	20	*	84	*	126	7.1 (98)
3301	1,029	19.8 (55)	1,360	6.8 (50)	197	49.7 (1)	1,029	4.5 (39)	1,364	3.2 (52)
3304	41	*	45	*	5	*	41	*	45	*
3305	46	*	63	*	8	*	46	*	64	*
3306	56	*	76	*	13	*	57	*	76	*
3308	22	*	35	*	3	*	22	*	35	*
3312	134	24.6 (71)	163	15.3 (94)	30	*	133	4.5 (39)	164	6.1 (94)
3550	22	*	28	*	15	*	22	*	28	*
3551	2	*	2	*	1	*	2	*	2	*
3701	189	24.9 (73)	248	13.7 (89)	42	*	190	13.2 (93)	254	4.3 (74)
3850	677	23.0 (70)	823	12.5 (79)	141	30.5 (9)	677	8.7 (72)	824	4.1 (71)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2022 Pregnancy Nutrition Surveillance (1)

Page : 45

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3901	339	22.1 (65)	409	7.3 (55)	61	*	338	3.8 (28)	412	2.9 (46)
3902	2	*	2	*	0	*	2	*	2	*
3950	445	19.3 (50)	602	9.3 (63)	83	*	447	4.5 (39)	607	4.4 (75)
3954	1	*	1	*	1	*	1	*	1	*
4106	534	10.5 (22)	667	6.0 (45)	47	*	529	3.2 (21)	667	2.7 (40)
4107	680	15.6 (41)	869	6.4 (48)	99	*	680	4.0 (30)	878	1.8 (22)
4109	106	17.0 (46)	183	10.4 (69)	18	*	108	0.9 (2)	184	3.8 (63)
4112	164	3.7 (5)	180	1.1 (5)	5	*	147	4.8 (48)	180	5.0 (79)
4115	767	8.3 (14)	1,070	3.3 (19)	63	*	764	3.7 (26)	1,075	1.3 (12)
4351	141	27.7 (78)	178	13.5 (85)	36	*	141	7.1 (62)	182	6.6 (96)
4352	54	*	61	*	27	*	54	*	64	*
4353	110	22.7 (68)	130	8.5 (61)	21	*	110	4.5 (39)	130	0.8 (4)
4354	188	29.3 (83)	211	11.8 (77)	53	*	189	10.1 (82)	235	3.4 (55)
4355	63	*	72	*	14	*	63	*	73	*
4356	96	*	129	16.3 (96)	32	*	96	*	129	10.1 (109)
4359	158	14.6 (39)	189	6.9 (51)	22	*	158	1.9 (11)	189	2.1 (26)
4360	61	*	73	*	8	*	61	*	74	*
4361	65	*	70	*	13	*	65	*	73	*
4362	66	*	79	*	19	*	66	*	83	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2022 Pregnancy Nutrition Surveillance (1)

Page : 46

## Michigan

### Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators <small>(2,3)</small>										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes <small>(4)</small>	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4363	142	21.8 (64)	198	12.6 (81)	28	*	142	6.3 (57)	198	9.6 (107)
4401	156	22.4 (67)	221	13.6 (87)	34	*	156	3.2 (21)	222	0.9 (8)
4501	173	29.5 (84)	242	12.8 (83)	48	*	173	9.8 (79)	244	5.7 (88)
4601	275	26.5 (76)	362	10.5 (71)	62	*	276	9.8 (79)	361	5.0 (79)
4701	226	20.4 (58)	303	6.9 (51)	41	*	226	4.4 (37)	314	3.8 (63)
5050	376	10.4 (20)	542	4.2 (34)	35	*	375	4.3 (34)	540	1.9 (23)
5051	583	10.6 (23)	808	5.1 (39)	55	*	581	4.1 (31)	806	2.1 (26)
5052	608	10.0 (19)	899	3.6 (27)	61	*	608	4.1 (31)	900	2.3 (32)
5055	384	1.0 (2)	446	0.4 (2)	4	*	384	1.3 (3)	446	0.2 (1)
5101	292	13.0 (30)	428	4.0 (29)	37	*	293	3.1 (20)	426	2.8 (42)
5102	2	*	1	*	0	*	2	*	1	*
5202	164	35.4 (92)	192	10.4 (69)	53	*	164	6.1 (56)	210	1.9 (23)
5206	7	*	7	*	2	*	7	*	7	*
5601	230	24.8 (72)	310	12.6 (81)	55	*	231	5.2 (51)	319	2.2 (30)
5801	452	27.4 (77)	580	10.3 (68)	115	44.3 (4)	454	8.8 (74)	588	3.4 (55)
5803	6	*	7	*	0	*	6	*	7	*
5804	1	*	1	*	1	*	1	*	1	*
5806	12	*	16	*	3	*	12	*	16	*
6101	20	*	24	*	2	*	20	*	25	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

## 2022 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic

Page : 47

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
6102	15	*	27	*	4	*	15	*	27	*
6103	832	21.2 (60)	1,019	9.3 (63)	165	33.9 (8)	833	7.7 (67)	1,044	5.4 (83)
6105	2	*	2	*	0	*	2	*	2	*
6301	792	7.4 (13)	1,088	4.0 (29)	56	*	791	1.5 (6)	1,093	1.1 (11)
6312	382	3.4 (4)	538	1.5 (8)	13	*	381	2.4 (15)	539	1.7 (20)
6316	259	6.6 (9)	352	4.0 (29)	17	*	258	1.6 (8)	355	0.8 (4)
6320	115	13.0 (30)	174	1.1 (5)	11	*	115	1.7 (10)	174	2.3 (32)
6325	62	*	97	*	16	*	62	*	102	2.9 (46)
6335	217	8.3 (14)	306	2.9 (16)	15	*	218	2.3 (13)	308	2.6 (36)
6345	25	*	40	*	5	*	25	*	42	*
6360	175	21.1 (59)	237	7.6 (58)	33	*	174	5.7 (53)	243	4.5 (76)
7306	425	28.0 (81)	569	11.2 (75)	113	48.7 (2)	425	7.3 (64)	582	3.4 (55)
7401	488	38.7 (95)	551	11.3 (76)	165	46.7 (3)	488	13.1 (92)	563	7.5 (100)
7601	143	12.6 (28)	164	5.5 (41)	16	*	143	3.5 (25)	166	3.0 (50)
7850	266	9.8 (18)	333	6.9 (51)	24	*	265	0.8 (1)	332	1.5 (15)
7852	3	*	2	*	0	*	3	*	1	*
8101	368	16.6 (44)	553	3.1 (18)	55	*	368	8.2 (71)	554	3.4 (55)
8102	112	13.4 (34)	134	2.2 (11)	14	*	113	5.3 (52)	134	4.5 (76)
8106	142	16.2 (43)	189	5.3 (40)	22	*	142	1.4 (5)	189	0.5 (3)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2022 Pregnancy Nutrition Surveillance (1)

Page : 48

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8150	1	*	1	*	0	*	1	*	1	*
8151	806	13.9 (38)	971	4.5 (37)	91	*	808	4.5 (39)	976	1.5 (15)
8152	5	*	5	*	1	*	5	*	5	*
8153	1	*	0	*	0	*	1	*	0	*
8155	2	*	1	*	0	*	2	*	1	*
8201	627	12.8 (29)	800	4.8 (38)	67	*	629	3.7 (26)	799	2.1 (26)
8202	1,057	2.7 (3)	1,325	1.1 (5)	26	*	1,057	2.3 (13)	1,325	0.4 (2)
8203	69	*	92	*	5	*	69	*	92	*
8250	117	13.7 (36)	144	3.5 (25)	11	*	116	4.3 (34)	145	3.4 (55)
8252	34	*	45	*	5	*	34	*	45	*
8255	374	17.1 (47)	538	7.1 (54)	57	*	375	4.5 (39)	537	2.6 (36)
8257	401	0.5 (1)	524	0.0 (1)	1	*	399	1.3 (3)	519	0.8 (4)
8263	132	21.2 (60)	111	1.8 (9)	22	*	133	3.0 (18)	111	0.9 (8)
8264	165	4.8 (8)	253	2.0 (10)	9	*	166	3.0 (18)	252	3.2 (52)
8267	122	10.7 (24)	176	2.3 (12)	11	*	122	4.1 (31)	177	5.6 (87)
8901	183	7.1 (12)	274	3.3 (19)	13	*	184	9.2 (78)	274	2.9 (46)
8902	294	12.2 (27)	425	2.4 (13)	36	*	295	3.4 (24)	422	2.1 (26)
8903	337	16.9 (45)	449	4.0 (29)	47	*	332	4.5 (39)	449	1.3 (12)
8904	430	13.5 (35)	639	2.5 (14)	52	*	431	6.5 (59)	641	2.5 (35)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2022 Pregnancy Nutrition Surveillance (1)

Page : 49

Michigan

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8905	406	9.4 (17)	603	3.0 (17)	34	*	406	4.4 (37)	604	4.0 (69)
8909	260	10.4 (20)	371	3.8 (28)	24	*	261	8.0 (70)	371	3.5 (60)
9045	306	3.9 (6)	394	0.5 (3)	10	*	302	3.3 (23)	394	2.8 (42)
9050	222	15.8 (42)	304	3.3 (19)	38	*	228	7.5 (66)	306	1.0 (10)
9051	395	4.3 (7)	533	2.8 (15)	16	*	395	1.5 (6)	532	1.5 (15)
9053	291	10.7 (24)	505	1.0 (4)	29	*	293	6.5 (59)	503	2.6 (36)
9055	1	*	4	*	0	*	1	*	4	*
9058	244	6.6 (9)	458	3.5 (25)	17	*	246	5.7 (53)	458	2.2 (30)
9066	222	17.6 (49)	216	4.2 (34)	39	*	223	10.3 (84)	215	2.8 (42)
9068	159	11.9 (26)	235	3.4 (22)	19	*	159	5.0 (49)	236	1.7 (20)
9069	304	6.9 (11)	499	3.4 (22)	21	*	304	6.3 (57)	499	2.8 (42)
9070	413	9.2 (16)	625	3.4 (22)	35	*	415	8.7 (72)	626	3.0 (50)
9080	549	13.8 (37)	834	4.1 (33)	73	*	551	10.0 (81)	836	6.0 (91)
9082	14	*	24	*	0	*	14	*	24	*
9083	2	*	2	*	0	*	2	*	2	*
9087	3	*	3	*	0	*	3	*	3	*
Michigan	33,554	17.5	44,427	7.2	5,392	42.2	33,562	6.0	44,801	3.5
National (Prior Year)	49,474	24.1	56,944	9.7	9,950	32.6	46,000	5.6	60,042	1.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 50

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	31	*	*	31	*	22	*	31	*
0102	109	11.9 (67)	8.3 (96)	108	12.0 (47)	92	*	109	78.0 (63)
0103	100	5.0 (2)	6.0 (49)	106	10.4 (22)	92	*	101	79.2 (57)
0104	32	*	*	33	*	25	*	32	*
0250	30	*	*	32	*	27	*	29	*
0251	15	*	*	15	*	12	*	15	*
0252	33	*	*	33	*	30	*	33	*
0253	34	*	*	37	*	30	*	34	*
0401	124	10.5 (53)	6.5 (63)	127	9.4 (10)	109	7.3 (79)	125	83.2 (31)
0402	113	8.8 (31)	9.7 (103)	114	12.3 (52)	98	*	113	83.2 (31)
0403	36	*	*	37	*	30	*	38	*
0404	33	*	*	33	*	29	*	33	*
0501	102	8.8 (31)	4.9 (34)	123	3.3 (1)	97	*	118	83.1 (34)
0503	17	*	*	18	*	17	*	18	*
0504	97	*	*	99	*	81	*	99	*
0505	51	*	*	66	*	47	*	62	*
0506	48	*	*	52	*	45	*	53	*
0507	12	*	*	16	*	9	*	15	*
0508	4	*	*	4	*	4	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 51

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0650	332	8.1 (22)	7.5 (85)	345	9.6 (13)	300	5.0 (51)	332	84.0 (26)
0651	163	6.7 (10)	8.6 (101)	170	9.4 (10)	147	3.4 (21)	163	78.5 (60)
0652	82	*	*	81	*	75	*	80	*
0653	103	8.7 (29)	5.8 (44)	109	9.2 (9)	94	*	103	86.4 (15)
0654	123	10.6 (54)	7.3 (80)	127	11.0 (30)	110	5.5 (58)	123	83.7 (27)
0655	91	*	*	94	*	84	*	91	*
0657	6	*	*	7	*	4	*	6	*
0701	126	5.6 (7)	7.1 (76)	131	13.7 (68)	109	0.9 (1)	128	91.4 (3)
0702	17	*	*	17	*	17	*	17	*
0703	42	*	*	45	*	38	*	42	*
0704	15	*	*	16	*	11	*	15	*
0705	21	*	*	23	*	16	*	21	*
0850	224	12.5 (73)	6.3 (58)	230	14.8 (87)	191	6.3 (68)	224	84.8 (20)
0851	179	9.5 (40)	6.7 (68)	185	14.6 (84)	153	3.3 (19)	179	83.2 (31)
0901	414	12.6 (75)	7.2 (79)	437	14.2 (79)	353	4.8 (45)	416	76.9 (65)
0902	10	*	*	10	*	8	*	10	*
1051	317	10.1 (46)	7.3 (80)	327	14.1 (76)	269	4.1 (28)	320	90.6 (6)
1052	2	*	*	1	*	1	*	2	*
1101	371	14.8 (94)	5.7 (42)	384	13.8 (70)	315	9.2 (95)	370	74.9 (70)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 52

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1102	212	5.7 (8)	6.1 (52)	221	11.3 (37)	184	2.2 (9)	213	80.3 (53)
1106	20	*	*	20	*	18	*	20	*
1250	266	7.5 (16)	7.5 (85)	281	10.0 (18)	237	5.1 (54)	268	80.2 (54)
1251	170	14.7 (91)	3.5 (10)	176	15.9 (95)	138	6.5 (70)	169	82.2 (42)
1252	163	8.0 (19)	11.0 (106)	168	8.9 (7)	149	4.0 (26)	164	85.4 (18)
1253	195	8.2 (24)	7.7 (89)	200	14.0 (74)	164	1.8 (5)	195	87.2 (11)
1350	694	9.1 (36)	6.3 (58)	712	11.0 (30)	609	4.6 (35)	700	81.0 (49)
1352	112	8.9 (33)	10.7 (105)	116	11.2 (34)	98	*	113	67.3 (90)
1401	56	*	*	58	*	52	*	56	*
1402	21	*	*	22	*	18	*	22	*
1403	11	*	*	11	*	10	*	11	*
1501	190	11.1 (60)	7.4 (83)	207	10.6 (26)	168	6.5 (70)	197	79.7 (55)
1503	243	5.3 (4)	6.2 (54)	249	11.6 (41)	215	1.4 (2)	245	83.3 (29)
1504	515	10.1 (46)	5.8 (44)	562	15.8 (94)	432	3.5 (23)	541	82.6 (37)
1506	155	11.0 (59)	7.1 (76)	161	13.0 (62)	134	6.7 (73)	155	76.8 (66)
1507	141	12.8 (76)	8.5 (99)	146	13.7 (68)	123	3.3 (19)	143	90.9 (4)
1509	93	*	*	100	13.0 (62)	81	*	95	*
1511	193	10.9 (58)	6.2 (54)	196	11.7 (43)	170	4.7 (38)	194	80.9 (50)
1512	132	11.4 (65)	7.6 (87)	133	17.3 (102)	109	4.6 (35)	132	82.6 (37)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 53

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1514	224	13.4 (83)	7.1 (76)	234	14.1 (76)	188	4.8 (45)	230	90.0 (7)
1701	150	7.3 (14)	8.7 (102)	154	10.4 (22)	133	2.3 (10)	151	82.1 (44)
1950	174	8.0 (19)	3.4 (8)	179	14.5 (82)	145	2.1 (8)	175	84.6 (22)
1952	40	*	*	41	*	33	*	40	*
1953	67	*	*	70	*	63	*	67	*
1954	107	10.3 (50)	3.7 (16)	110	10.9 (28)	95	*	106	87.7 (10)
1955	36	*	*	37	*	29	*	36	*
1957	121	16.5 (105)	8.3 (96)	124	16.9 (101)	100	8.0 (89)	122	86.9 (12)
1963	16	*	*	15	*	13	*	16	*
2150	146	5.5 (5)	6.2 (54)	149	9.4 (10)	131	1.5 (4)	146	82.2 (42)
2151	57	*	*	59	*	55	*	57	*
2152	13	*	*	14	*	12	*	13	*
2251	106	3.8 (1)	3.8 (18)	107	6.5 (3)	98	*	106	84.9 (19)
2252	51	*	*	53	*	43	*	49	*
2554	18	*	*	18	*	17	*	18	*
2555	2,147	12.9 (77)	6.8 (71)	2,193	13.1 (64)	1,848	6.2 (67)	2,151	69.2 (88)
2556	34	*	*	35	*	28	*	35	*
2557	1	*	*	1	*	1	*	1	*
2562	1	*	*	1	*	0	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 54

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2563	1	*	*	1	*	1	*	1	*
2564	1	*	*	1	*	1	*	1	*
3001	364	14.0 (87)	3.3 (7)	382	12.8 (59)	320	6.9 (74)	361	65.7 (94)
3002	47	*	*	47	*	41	*	47	*
3004	84	*	*	89	*	71	*	84	*
3005	52	*	*	54	*	48	*	53	*
3007	1	*	*	1	*	1	*	1	*
3201	123	8.1 (22)	11.4 (107)	123	12.2 (51)	107	3.7 (25)	123	83.7 (27)
3301	1,276	12.9 (77)	6.3 (58)	1,325	14.6 (84)	1,086	4.5 (33)	1,286	84.4 (24)
3304	44	*	*	45	*	41	*	44	*
3305	55	*	*	62	*	45	*	56	*
3306	72	*	*	74	*	63	*	72	*
3308	34	*	*	34	*	31	*	34	*
3312	155	9.0 (34)	4.5 (28)	159	8.8 (6)	143	7.0 (76)	155	88.4 (8)
3550	22	*	*	25	*	18	*	22	*
3551	1	*	*	2	*	1	*	1	*
3701	237	8.0 (19)	4.6 (30)	245	12.7 (58)	206	2.4 (11)	238	90.8 (5)
3850	776	12.4 (72)	7.6 (87)	800	14.0 (74)	668	4.8 (45)	777	81.7 (46)
3901	386	11.4 (65)	3.9 (20)	398	12.6 (55)	336	5.4 (55)	387	76.7 (67)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 55

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3902	1	*	*	1	*	1	*	1	*
3950	575	11.3 (61)	8.0 (91)	591	12.9 (60)	501	4.2 (30)	581	82.4 (39)
3954	1	*	*	1	*	1	*	1	*
4106	629	14.0 (87)	4.3 (24)	650	15.5 (92)	534	6.6 (72)	631	79.4 (56)
4107	818	14.2 (89)	4.9 (34)	855	16.8 (99)	677	6.1 (65)	833	80.8 (51)
4109	166	13.3 (82)	6.0 (49)	171	16.4 (97)	138	3.6 (24)	165	87.9 (9)
4112	170	6.5 (9)	3.5 (10)	173	15.0 (89)	142	2.8 (13)	170	40.0 (107)
4115	1,008	12.5 (73)	6.6 (67)	1,053	12.9 (60)	877	5.7 (60)	1,032	84.7 (21)
4351	168	15.5 (98)	6.5 (63)	173	17.3 (102)	138	8.0 (89)	169	91.7 (2)
4352	56	*	*	61	*	49	*	56	*
4353	124	9.7 (41)	6.5 (63)	125	10.4 (22)	111	5.4 (55)	124	75.8 (69)
4354	219	8.7 (29)	4.1 (21)	224	12.5 (54)	190	3.2 (18)	220	84.5 (23)
4355	69	*	*	73	*	60	*	69	*
4356	119	7.6 (17)	6.7 (68)	121	9.9 (16)	106	2.8 (13)	119	85.7 (17)
4359	172	9.3 (38)	8.1 (93)	177	11.3 (37)	149	4.7 (38)	172	86.6 (13)
4360	66	*	*	70	*	57	*	66	*
4361	67	*	*	68	*	59	*	67	*
4362	79	*	*	80	*	70	*	78	*
4363	186	9.7 (41)	8.1 (93)	186	11.3 (37)	160	4.4 (31)	186	92.5 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 56

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4401	203	10.8 (56)	10.3 (104)	213	11.3 (37)	181	6.1 (65)	205	82.4 (39)
4501	234	5.1 (3)	8.5 (99)	240	10.8 (27)	210	1.4 (2)	234	83.3 (29)
4601	342	6.7 (10)	6.7 (68)	356	10.1 (20)	307	2.0 (6)	341	82.4 (39)
4701	300	12.0 (69)	5.3 (38)	307	15.6 (93)	253	4.7 (38)	300	82.7 (36)
5050	522	13.2 (80)	3.8 (18)	524	13.9 (72)	443	7.7 (84)	522	74.5 (74)
5051	758	13.9 (84)	4.7 (33)	778	16.8 (99)	618	4.9 (48)	762	78.6 (59)
5052	857	15.2 (97)	4.6 (30)	870	15.2 (90)	722	8.4 (92)	862	73.5 (80)
5055	419	10.3 (50)	5.0 (37)	426	8.2 (5)	383	7.3 (79)	423	74.2 (78)
5101	408	9.1 (36)	4.2 (22)	410	11.7 (43)	358	3.1 (17)	412	80.8 (51)
5102	1	*	*	1	*	1	*	1	*
5202	200	8.5 (26)	3.5 (10)	200	12.0 (47)	174	2.9 (15)	201	86.1 (16)
5206	7	*	*	7	*	6	*	7	*
5601	296	10.1 (46)	8.1 (93)	312	10.9 (28)	264	4.9 (48)	298	81.2 (48)
5801	561	7.8 (18)	7.8 (90)	572	12.1 (49)	493	2.6 (12)	560	72.0 (85)
5803	6	*	*	6	*	6	*	6	*
5804	1	*	*	1	*	1	*	1	*
5806	15	*	*	15	*	15	*	15	*
6101	24	*	*	25	*	22	*	24	*
6102	26	*	*	27	*	22	*	26	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 8A

2022 Pregnancy Nutrition Surveillance (1)

Page : 57

Michigan

## Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6103	976	13.2 (80)	6.4 (61)	1,009	13.8 (70)	830	6.0 (63)	978	74.3 (77)
6105	2	*	*	2	*	2	*	1	*
6301	1,021	12.1 (70)	7.0 (72)	1,070	13.4 (65)	885	6.0 (63)	1,035	77.2 (64)
6312	507	10.7 (55)	5.3 (38)	526	13.5 (66)	438	4.6 (35)	517	82.8 (35)
6316	330	9.7 (41)	4.5 (28)	346	10.4 (22)	293	4.1 (28)	336	82.1 (44)
6320	170	12.9 (77)	6.5 (63)	168	5.4 (2)	156	10.9 (98)	172	84.3 (25)
6325	102	6.9 (12)	14.7 (108)	100	10.0 (18)	90	*	102	78.4 (61)
6335	299	10.4 (52)	8.0 (91)	299	11.0 (30)	261	5.4 (55)	302	72.5 (84)
6345	39	*	*	42	*	34	*	40	*
6360	224	15.6 (99)	5.4 (40)	236	13.6 (67)	193	10.9 (98)	228	74.6 (71)
7306	546	11.9 (67)	6.2 (54)	564	14.5 (82)	467	4.9 (48)	548	74.5 (74)
7401	529	8.5 (26)	7.0 (72)	540	12.6 (55)	462	3.0 (16)	528	78.4 (61)
7601	156	9.0 (34)	5.8 (44)	163	12.3 (52)	135	4.4 (31)	161	74.5 (74)
7850	314	8.6 (28)	7.0 (72)	330	11.2 (34)	278	4.7 (38)	315	73.3 (82)
7852	1	*	*	1	*	1	*	1	*
8101	523	14.7 (91)	2.9 (2)	539	15.2 (90)	437	8.0 (89)	527	58.4 (101)
8102	123	7.3 (14)	5.7 (42)	132	9.1 (8)	110	4.5 (33)	128	73.4 (81)
8106	166	13.9 (84)	6.0 (49)	183	14.2 (79)	139	5.0 (51)	180	73.3 (82)
8150	0	*	*	1	*	0	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 58

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8151	905	11.3 (61)	6.4 (61)	946	11.8 (45)	794	5.9 (62)	925	86.6 (13)
8152	4	*	*	5	*	4	*	4	*
8153	0	*	*	0	*	0	*	0	*
8155	1	*	*	1	*	1	*	1	*
8201	757	12.3 (71)	4.9 (34)	787	14.1 (76)	655	5.0 (51)	760	74.6 (71)
8202	1,174	8.3 (25)	3.7 (16)	1,245	9.7 (15)	1,044	3.4 (21)	1,239	69.9 (87)
8203	87	*	*	90	*	75	*	87	*
8250	141	11.3 (61)	4.3 (24)	145	10.3 (21)	126	4.0 (26)	141	66.0 (92)
8252	38	*	*	45	*	35	*	44	*
8255	514	11.3 (61)	5.8 (44)	530	12.6 (55)	447	4.7 (38)	514	73.7 (79)
8257	471	15.1 (96)	3.2 (5)	512	11.9 (46)	411	9.0 (94)	486	66.3 (91)
8263	110	5.5 (5)	5.5 (41)	110	7.3 (4)	102	2.0 (6)	109	78.9 (58)
8264	237	10.1 (46)	8.4 (98)	245	11.0 (30)	211	4.7 (38)	239	81.6 (47)
8267	165	9.7 (41)	6.1 (52)	171	9.9 (16)	146	5.5 (58)	169	71.0 (86)
8901	260	15.8 (103)	4.2 (22)	269	17.5 (105)	211	10.0 (96)	260	61.2 (96)
8902	403	15.6 (99)	3.5 (10)	406	21.9 (109)	309	7.1 (78)	404	68.1 (89)
8903	413	15.0 (95)	2.7 (1)	436	14.4 (81)	351	7.7 (84)	408	59.3 (98)
8904	603	15.6 (99)	4.3 (24)	629	18.1 (106)	486	7.8 (87)	604	58.3 (102)
8905	558	9.3 (38)	7.0 (72)	582	11.2 (34)	488	4.7 (38)	563	74.6 (71)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2022 Pregnancy Nutrition Surveillance (1)

Page : 59

### Michigan

### Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8909	344	15.7 (102)	3.2 (5)	358	19.0 (108)	274	7.7 (84)	344	66.0 (92)
9045	378	6.9 (12)	7.4 (83)	386	9.6 (13)	336	5.7 (60)	382	38.7 (108)
9050	295	10.8 (56)	5.8 (44)	303	11.6 (41)	261	6.9 (74)	296	58.8 (100)
9051	508	9.8 (45)	7.3 (80)	512	12.1 (49)	439	7.3 (79)	514	75.9 (68)
9053	472	18.6 (108)	3.4 (8)	489	13.9 (72)	397	10.6 (97)	475	61.7 (95)
9055	4	*	*	4	*	4	*	4	*
9058	413	16.7 (107)	3.1 (4)	441	14.7 (86)	347	8.9 (93)	430	60.5 (97)
9066	205	16.6 (106)	4.4 (27)	213	17.4 (104)	170	7.6 (83)	205	58.0 (103)
9068	228	15.8 (103)	3.5 (10)	232	15.9 (95)	192	7.8 (87)	228	48.2 (105)
9069	476	14.5 (90)	2.9 (2)	487	18.1 (106)	392	6.4 (69)	475	45.7 (106)
9070	589	13.9 (84)	4.6 (30)	612	16.5 (98)	488	7.0 (76)	588	57.3 (104)
9080	782	14.7 (91)	3.5 (10)	811	14.9 (88)	658	7.4 (82)	801	59.2 (99)
9082	22	*	*	23	*	16	*	23	*
9083	2	*	*	2	*	0	*	2	*
9087	3	*	*	3	*	3	*	3	*
<b>Michigan</b>	<b>41,947</b>	<b>11.5</b>	<b>5.9</b>	<b>43,460</b>	<b>13.2</b>	<b>36,146</b>	<b>5.3</b>	<b>42,370</b>	<b>75.5</b>
<b>National (Prior Year)</b>	<b>57,663</b>	<b>10.8</b>	<b>6.5</b>	<b>58,093</b>	<b>13.0</b>	<b>48,228</b>	<b>4.7</b>	<b>58,053</b>	<b>55.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 60

## Michigan

### Comparison of Racial and Ethnic Distribution by Clinic Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
0101	111	93.7	0.0	1.8	0.0	0.0	3.6	0.9
0102	410	90.2	0.5	2.4	0.5	0.0	6.3	0.0
0103	358	93.6	0.6	2.2	0.0	0.0	3.4	0.3
0104	123	95.1	0.0	2.4	0.0	0.0	2.4	0.0
0105	14	*	*	*	*	*	*	*
0250	116	79.3	0.9	1.7	5.2	0.0	12.9	0.0
0251	76	*	*	*	*	*	*	*
0252	97	*	*	*	*	*	*	*
0253	119	70.6	0.0	0.8	11.8	0.0	16.8	0.0
0401	441	94.6	0.5	2.0	0.2	0.2	2.5	0.0
0402	373	85.5	1.6	3.2	0.3	0.0	9.4	0.0
0403	137	94.2	0.7	2.2	1.5	0.0	0.7	0.7
0404	116	92.2	0.9	0.9	0.9	0.0	5.2	0.0
0501	427	93.4	0.5	2.3	0.0	0.2	3.3	0.2
0503	74	*	*	*	*	*	*	*
0504	358	86.6	1.7	3.1	1.4	0.6	6.7	0.0
0505	230	90.0	0.0	3.9	0.0	0.0	6.1	0.0
0506	169	92.9	0.0	1.2	0.6	0.6	4.7	0.0
0507	70	*	*	*	*	*	*	*
0508	19	*	*	*	*	*	*	*
0650	1,074	78.2	4.4	6.7	3.9	0.7	6.1	0.1
0651	559	92.7	0.7	4.1	0.0	0.4	2.1	0.0
0652	300	94.0	0.0	2.7	0.3	0.0	3.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 61

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
0653	407	94.8	0.0	2.2	0.0	0.5	2.5	0.0
0654	410	91.2	1.2	3.4	0.5	0.0	3.7	0.0
0655	308	95.8	0.0	1.0	0.0	0.3	2.6	0.3
0657	47	*	*	*	*	*	*	*
0701	490	87.8	2.0	2.4	0.6	2.0	4.7	0.4
0702	68	*	*	*	*	*	*	*
0703	151	85.4	2.6	2.0	0.7	1.3	7.3	0.7
0704	50	*	*	*	*	*	*	*
0705	69	*	*	*	*	*	*	*
0706	3	*	*	*	*	*	*	*
0850	701	83.3	5.0	9.4	0.0	1.6	0.6	0.1
0851	600	92.5	0.3	4.8	0.3	0.3	1.5	0.2
0901	1,489	80.3	2.1	10.5	0.1	0.2	6.5	0.3
0902	34	*	*	*	*	*	*	*
1051	1,138	83.6	1.0	7.6	2.4	0.7	4.2	0.5
1052	3	*	*	*	*	*	*	*
1101	1,467	35.0	49.4	8.9	0.3	1.7	3.8	0.9
1102	801	75.0	10.4	7.4	0.1	0.9	5.6	0.6
1106	66	*	*	*	*	*	*	*
1250	974	83.9	2.1	10.7	0.1	0.7	1.6	0.9
1251	622	93.1	0.2	3.2	0.5	0.0	2.4	0.6
1252	525	60.2	0.2	35.4	0.0	0.0	3.6	0.6
1253	691	77.4	0.9	12.0	0.0	0.3	9.0	0.4
1350	2,393	58.3	20.4	8.3	0.4	5.6	6.8	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 62

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
1351	28	*	*	*	*	*	*	*
1352	402	70.1	18.4	5.2	0.2	0.0	5.2	0.7
1401	197	89.8	0.5	4.6	1.0	0.0	4.1	0.0
1402	102	58.8	0.0	20.6	8.8	0.0	11.8	0.0
1403	21	*	*	*	*	*	*	*
1501	745	48.9	5.0	40.4	0.1	0.3	3.5	1.9
1503	832	60.2	7.8	21.4	0.4	0.2	7.9	2.0
1504	1,907	37.1	5.5	47.7	0.1	4.0	5.0	0.6
1505	131	76.3	6.1	13.7	0.0	0.0	3.1	0.8
1506	551	80.0	1.5	14.0	0.0	0.7	2.7	1.1
1507	528	84.1	2.5	7.6	0.0	1.3	4.0	0.6
1508	6	*	*	*	*	*	*	*
1509	384	47.1	1.0	46.9	0.0	0.5	2.9	1.6
1511	613	86.8	2.6	6.2	0.3	0.5	3.3	0.3
1512	365	86.0	0.8	6.3	0.0	0.8	6.0	0.0
1514	587	75.6	5.1	12.8	0.3	1.4	4.3	0.5
1701	543	57.5	0.7	2.0	19.0	0.7	20.1	0.0
1950	609	91.1	0.3	6.4	0.0	0.2	1.6	0.3
1952	140	94.3	0.7	3.6	0.0	0.0	0.7	0.7
1953	249	93.2	0.0	5.6	0.0	0.0	1.2	0.0
1954	391	84.7	0.3	11.3	0.3	0.0	3.1	0.5
1955	113	85.0	0.0	12.4	0.0	0.0	1.8	0.9
1957	381	83.7	1.0	10.2	0.0	0.0	4.5	0.5
1963	57	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 3A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 63

Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
1964	1	*	*	*	*	*	*	*
2150	582	84.5	0.7	3.3	1.5	0.5	9.1	0.3
2151	210	79.0	2.4	8.6	0.0	0.0	5.7	4.3
2152	33	*	*	*	*	*	*	*
2251	391	95.1	0.0	2.0	0.0	0.5	1.8	0.5
2252	168	90.5	0.0	3.0	0.6	0.0	4.2	1.8
2554	49	*	*	*	*	*	*	*
2555	7,177	49.4	37.6	6.3	0.0	0.3	6.1	0.3
2556	184	26.6	60.9	3.3	0.5	0.0	8.7	0.0
2557	2	*	*	*	*	*	*	*
2558	1	*	*	*	*	*	*	*
2560	3	*	*	*	*	*	*	*
2561	1	*	*	*	*	*	*	*
2562	6	*	*	*	*	*	*	*
2563	13	*	*	*	*	*	*	*
2564	17	*	*	*	*	*	*	*
3001	1,181	25.2	48.5	15.5	0.1	0.4	10.1	0.2
3002	197	25.4	44.7	23.4	0.0	0.0	6.6	0.0
3004	260	70.0	0.4	25.8	0.0	0.4	3.5	0.0
3005	153	87.6	0.0	6.5	0.7	2.0	3.3	0.0
3007	9	*	*	*	*	*	*	*
3201	404	91.8	0.5	5.7	0.0	0.2	1.2	0.5
3203	1	*	*	*	*	*	*	*
3301	4,277	38.1	30.7	16.3	0.1	5.9	8.7	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 64

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
3304	103	46.6	17.5	12.6	0.0	16.5	6.8	0.0
3305	157	60.5	12.1	12.7	0.0	5.7	8.9	0.0
3306	234	86.8	3.0	7.7	0.4	0.9	0.9	0.4
3308	114	90.4	0.9	6.1	0.0	0.0	2.6	0.0
3312	426	38.3	26.8	25.4	0.5	1.9	7.3	0.0
3550	80	*	*	*	*	*	*	*
3551	10	*	*	*	*	*	*	*
3552	14	*	*	*	*	*	*	*
3701	876	88.6	0.2	7.8	0.0	0.0	3.4	0.0
3850	2,830	71.4	14.0	6.5	0.1	0.6	7.2	0.1
3901	1,528	36.8	50.7	6.0	0.0	0.9	5.4	0.3
3902	42	*	*	*	*	*	*	*
3903	7	*	*	*	*	*	*	*
3950	2,021	52.5	24.1	12.9	0.2	1.0	8.5	0.7
3954	6	*	*	*	*	*	*	*
4106	2,160	12.0	39.3	43.3	0.2	0.8	4.3	0.1
4107	2,880	41.1	28.8	21.0	0.3	2.0	6.5	0.2
4109	643	84.4	1.6	9.5	0.0	0.5	3.9	0.2
4110	2	*	*	*	*	*	*	*
4112	557	1.1	1.1	96.8	0.2	0.2	0.2	0.5
4115	3,417	29.9	31.6	21.9	0.3	11.6	4.6	0.3
4351	532	88.2	0.8	5.8	0.2	0.6	4.3	0.2
4352	180	74.4	10.6	4.4	0.0	0.0	10.0	0.6
4353	457	55.8	0.4	38.7	0.4	0.0	3.7	0.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 65

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
4354	734	86.8	3.7	3.1	0.5	0.5	4.9	0.4
4355	264	84.5	0.8	5.7	1.9	0.0	7.2	0.0
4356	390	81.0	1.3	10.8	0.3	0.8	5.4	0.5
4358	23	*	*	*	*	*	*	*
4359	593	63.2	0.5	33.2	0.5	0.2	1.3	1.0
4360	218	87.6	0.5	7.3	0.0	0.0	4.6	0.0
4361	255	88.6	1.2	6.3	0.0	0.0	3.9	0.0
4362	279	90.3	0.4	4.7	0.0	0.4	3.9	0.0
4363	680	90.1	0.9	2.8	0.0	0.1	5.7	0.3
4401	758	88.9	1.1	5.7	0.0	0.7	3.7	0.0
4501	779	93.2	0.4	4.7	0.0	0.1	1.5	0.0
4601	1,237	71.7	2.7	19.7	0.2	0.4	4.9	0.5
4701	998	90.0	1.7	3.4	0.0	1.0	3.1	0.8
5050	1,576	45.3	44.2	3.1	0.0	2.0	5.1	0.3
5051	2,681	49.2	38.5	4.3	0.0	2.9	4.7	0.4
5052	2,718	32.4	40.9	3.1	0.3	18.9	4.2	0.2
5055	1,308	79.3	8.9	1.1	0.1	9.9	0.7	0.1
5101	1,508	66.5	13.7	12.9	0.3	1.7	4.9	0.0
5102	17	*	*	*	*	*	*	*
5202	676	86.4	1.5	3.1	2.5	0.4	5.5	0.6
5206	16	*	*	*	*	*	*	*
5601	957	86.7	2.1	6.0	0.0	1.8	3.3	0.1
5801	1,847	81.5	7.0	6.4	0.1	0.2	4.1	0.8
5803	52	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 66

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
5804	17	*	*	*	*	*	*	*
5806	45	*	*	*	*	*	*	*
6101	57	*	*	*	*	*	*	*
6102	86	*	*	*	*	*	*	*
6103	3,333	54.5	27.5	10.0	0.4	0.5	6.8	0.3
6105	9	*	*	*	*	*	*	*
6106	31	*	*	*	*	*	*	*
6301	3,372	38.9	31.2	22.8	0.2	2.3	4.4	0.2
6312	1,762	34.8	55.1	4.1	0.1	1.9	3.9	0.1
6316	1,100	61.0	23.6	8.0	0.1	3.3	3.7	0.3
6318	22	*	*	*	*	*	*	*
6320	490	35.3	53.5	3.9	0.4	1.8	5.1	0.0
6325	315	90.5	0.6	5.1	0.0	0.0	3.5	0.3
6335	931	66.8	21.2	3.1	0.0	6.7	2.1	0.1
6340	1	*	*	*	*	*	*	*
6345	115	82.6	3.5	4.3	0.0	4.3	5.2	0.0
6360	975	32.9	45.7	15.7	0.0	1.4	3.9	0.3
7306	2,065	45.6	31.9	15.4	0.1	0.4	6.2	0.3
7401	1,903	80.5	5.8	6.0	0.0	0.5	7.0	0.3
7601	558	93.4	0.0	4.5	0.2	0.4	1.1	0.5
7850	1,025	89.2	0.4	6.3	0.0	0.1	3.8	0.2
7852	15	*	*	*	*	*	*	*
8101	1,727	6.9	88.9	1.2	0.0	0.3	2.5	0.2
8102	485	20.2	17.5	58.1	0.0	1.4	2.5	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 67

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution <sup>(2)</sup>								
Clinic	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
8105	235	31.9	63.4	2.6	0.4	0.4	0.9	0.4
8106	574	27.2	52.1	13.2	0.2	0.9	6.3	0.2
8150	7	*	*	*	*	*	*	*
8151	3,165	37.7	39.7	11.0	0.1	3.1	7.9	0.4
8152	24	*	*	*	*	*	*	*
8153	5	*	*	*	*	*	*	*
8155	10	*	*	*	*	*	*	*
8157	1	*	*	*	*	*	*	*
8201	2,775	44.2	40.5	4.9	0.2	3.8	5.9	0.4
8202	3,986	85.7	11.0	1.7	0.0	0.4	1.0	0.2
8203	375	58.1	31.7	5.6	0.3	0.5	3.7	0.0
8250	589	20.9	68.6	4.4	0.2	1.0	4.9	0.0
8252	163	12.3	79.1	4.3	0.0	2.5	1.8	0.0
8255	2,059	54.8	30.3	9.0	0.2	2.4	3.2	0.1
8257	1,801	53.3	10.8	0.8	0.2	33.4	1.4	0.2
8263	527	62.2	18.4	13.5	0.0	1.5	4.4	0.0
8264	790	58.1	30.1	5.6	0.1	2.0	3.9	0.1
8267	634	54.3	19.6	22.7	0.2	0.6	2.5	0.2
8901	870	8.3	84.7	2.5	0.0	1.5	2.9	0.1
8902	1,347	8.2	86.5	2.8	0.0	0.3	2.0	0.1
8903	1,463	5.6	87.1	2.1	0.1	1.3	3.8	0.1
8904	2,171	3.6	90.9	1.8	0.1	0.9	2.4	0.3
8905	1,887	62.9	25.3	7.7	0.1	1.6	2.2	0.1
8909	1,094	5.3	90.0	2.0	0.1	0.3	2.1	0.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 3A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 68

## Michigan

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
9045	1,371	9.6	25.4	62.6	0.1	0.4	1.5	0.3
9050	954	17.1	62.7	17.3	0.0	0.0	2.9	0.0
9051	1,515	7.8	5.0	85.3	0.0	0.1	1.5	0.3
9053	1,386	6.8	86.2	2.2	0.1	2.2	2.3	0.2
9055	7	*	*	*	*	*	*	*
9058	1,277	16.9	71.3	1.5	0.1	7.9	2.2	0.1
9066	984	2.2	94.2	0.5	0.0	0.7	2.3	0.0
9068	850	2.2	95.5	0.2	0.0	0.6	1.2	0.2
9069	1,858	1.3	97.2	1.0	0.0	0.1	0.4	0.1
9070	1,643	5.8	89.8	1.5	0.1	1.5	1.1	0.1
9072	386	2.8	93.3	0.8	0.3	0.8	2.1	0.0
9080	2,520	19.0	75.0	2.2	0.1	0.4	3.0	0.3
9082	110	1.8	94.5	1.8	0.0	0.9	0.9	0.0
9083	7	*	*	*	*	*	*	*
9084	5	*	*	*	*	*	*	*
9087	15	*	*	*	*	*	*	*
<b>Michigan</b>	<b>146,297</b>	<b>51.2</b>	<b>29.9</b>	<b>11.3</b>	<b>0.4</b>	<b>2.4</b>	<b>4.5</b>	<b>0.3</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 69

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0101	111	0.0	1.8	3.6	64.0	27.0	3.6	0.0
0102	410	0.0	2.7	8.5	59.8	27.1	2.0	0.0
0103	358	0.0	3.1	8.7	62.8	24.0	1.1	0.3
0104	123	0.0	4.9	8.9	65.0	20.3	0.8	0.0
0105	14	*	*	*	*	*	*	*
0250	116	0.0	4.3	9.5	51.7	31.0	2.6	0.9
0251	76	*	*	*	*	*	*	*
0252	97	*	*	*	*	*	*	*
0253	119	0.0	0.8	5.9	67.2	24.4	1.7	0.0
0401	441	0.0	2.0	8.6	59.9	28.8	0.7	0.0
0402	373	0.3	1.6	9.4	62.5	24.9	1.3	0.0
0403	137	0.0	3.6	8.8	58.4	27.0	2.2	0.0
0404	116	0.0	0.9	4.3	64.7	29.3	0.9	0.0
0501	427	0.2	1.2	8.4	63.7	24.8	1.6	0.0
0503	74	*	*	*	*	*	*	*
0504	358	0.0	2.2	4.5	58.7	31.6	3.1	0.0
0505	230	0.0	0.9	9.1	66.5	22.2	0.9	0.4
0506	169	0.0	3.6	6.5	68.6	20.7	0.6	0.0
0507	70	*	*	*	*	*	*	*
0508	19	*	*	*	*	*	*	*
0650	1,074	0.1	2.2	6.8	64.4	24.5	1.6	0.4
0651	559	0.5	3.2	8.9	62.6	23.6	0.9	0.2
0652	300	0.0	2.7	9.7	65.3	22.0	0.3	0.0
0653	407	0.0	1.5	5.7	64.1	27.3	1.0	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 70

## Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
0654	410	0.0	2.7	7.8	62.2	24.9	1.5	1.0
0655	308	0.0	3.9	7.8	58.1	27.9	1.6	0.6
0657	47	*	*	*	*	*	*	*
0701	490	0.0	1.4	8.0	58.4	30.4	1.8	0.0
0702	68	*	*	*	*	*	*	*
0703	151	0.0	0.0	7.9	53.0	39.1	0.0	0.0
0704	50	*	*	*	*	*	*	*
0705	69	*	*	*	*	*	*	*
0706	3	*	*	*	*	*	*	*
0850	701	0.0	2.4	9.1	58.3	27.1	2.0	1.0
0851	600	0.0	2.0	6.7	66.5	22.5	1.2	1.2
0901	1,489	0.0	2.5	7.9	61.1	26.9	1.2	0.4
0902	34	*	*	*	*	*	*	*
1051	1,138	0.2	1.6	6.5	58.2	31.4	1.9	0.3
1052	3	*	*	*	*	*	*	*
1101	1,467	0.1	3.7	9.6	57.7	24.7	2.0	2.0
1102	801	0.0	2.4	8.9	62.3	24.1	1.4	1.0
1106	66	*	*	*	*	*	*	*
1250	974	0.0	1.8	8.6	62.1	24.1	3.0	0.3
1251	622	0.6	2.9	7.7	63.2	22.3	2.3	1.0
1252	525	0.2	2.7	9.1	63.4	21.3	3.0	0.2
1253	691	0.0	2.2	7.2	65.8	23.6	1.0	0.1
1350	2,393	0.0	3.0	7.4	62.6	24.7	1.6	0.8
1351	28	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 71

## Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
1352	402	0.0	1.2	7.7	65.2	23.9	1.2	0.7
1401	197	1.0	1.5	5.6	61.9	28.4	1.0	0.5
1402	102	0.0	1.0	1.0	50.0	44.1	2.9	1.0
1403	21	*	*	*	*	*	*	*
1501	745	0.0	2.0	7.7	61.6	25.8	1.5	1.5
1503	832	0.2	3.1	9.0	60.8	24.9	0.8	1.1
1504	1,907	0.2	2.4	6.9	60.8	27.6	1.8	0.3
1505	131	0.0	0.8	4.6	70.2	21.4	2.3	0.8
1506	551	0.0	0.9	5.4	64.1	27.2	1.5	0.9
1507	528	0.0	1.1	5.5	59.3	31.1	2.7	0.4
1508	6	*	*	*	*	*	*	*
1509	384	0.0	1.6	6.8	55.7	31.8	3.9	0.3
1511	613	0.0	2.1	6.7	62.8	27.1	0.8	0.5
1512	365	0.0	1.9	7.1	56.7	31.0	3.0	0.3
1514	587	0.2	0.9	4.8	61.3	29.6	2.7	0.5
1701	543	0.0	1.8	6.8	62.6	27.3	1.3	0.2
1950	609	0.0	3.3	10.5	62.7	21.8	1.0	0.7
1952	140	0.0	0.0	4.3	69.3	24.3	1.4	0.7
1953	249	0.0	2.8	8.4	64.7	21.7	1.6	0.8
1954	391	0.3	2.3	7.9	65.0	22.5	1.5	0.5
1955	113	0.0	2.7	10.6	61.1	25.7	0.0	0.0
1957	381	0.0	1.6	4.7	55.9	36.2	1.3	0.3
1963	57	*	*	*	*	*	*	*
1964	1	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 72

## Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2150	582	0.2	0.9	6.9	60.8	30.4	0.9	0.0
2151	210	0.5	2.9	9.5	63.3	23.3	0.5	0.0
2152	33	*	*	*	*	*	*	*
2251	391	0.3	1.3	5.6	61.9	29.9	1.0	0.0
2252	168	0.0	4.8	4.8	64.3	25.6	0.6	0.0
2554	49	*	*	*	*	*	*	*
2555	7,177	0.2	2.1	7.3	63.1	25.5	1.6	0.3
2556	184	0.0	0.5	4.9	70.7	22.3	1.6	0.0
2557	2	*	*	*	*	*	*	*
2558	1	*	*	*	*	*	*	*
2560	3	*	*	*	*	*	*	*
2561	1	*	*	*	*	*	*	*
2562	6	*	*	*	*	*	*	*
2563	13	*	*	*	*	*	*	*
2564	17	*	*	*	*	*	*	*
3001	1,181	0.0	1.9	9.2	63.3	22.6	1.6	1.4
3002	197	0.0	3.6	4.6	65.5	23.9	2.5	0.0
3004	260	0.0	0.4	7.7	59.2	31.2	1.2	0.4
3005	153	0.0	0.7	8.5	65.4	22.9	2.0	0.7
3007	9	*	*	*	*	*	*	*
3201	404	0.0	0.7	5.2	60.1	32.4	1.5	0.0
3203	1	*	*	*	*	*	*	*
3301	4,277	0.1	2.6	6.9	61.2	26.7	1.9	0.5
3304	103	0.0	0.0	5.8	52.4	38.8	2.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 73

## Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
3305	157	0.0	0.0	4.5	59.2	33.8	2.5	0.0
3306	234	0.4	2.6	6.8	53.0	32.9	3.4	0.9
3308	114	0.0	0.0	4.4	65.8	28.1	1.8	0.0
3312	426	0.0	1.2	6.3	62.0	28.9	1.6	0.0
3550	80	*	*	*	*	*	*	*
3551	10	*	*	*	*	*	*	*
3552	14	*	*	*	*	*	*	*
3701	876	0.2	2.2	7.2	65.3	23.3	1.3	0.6
3850	2,830	0.1	2.7	7.2	62.6	25.3	1.3	0.8
3901	1,528	0.2	3.1	7.7	57.9	28.2	2.2	0.8
3902	42	*	*	*	*	*	*	*
3903	7	*	*	*	*	*	*	*
3950	2,021	0.0	2.6	7.7	61.1	26.1	1.9	0.5
3954	6	*	*	*	*	*	*	*
4106	2,160	0.0	2.3	7.4	58.4	27.3	3.0	1.6
4107	2,880	0.1	2.2	6.8	59.5	28.9	1.9	0.5
4109	643	0.0	2.0	6.2	61.4	28.6	1.6	0.2
4110	2	*	*	*	*	*	*	*
4112	557	0.5	4.3	8.6	55.3	27.6	2.9	0.7
4115	3,417	0.1	1.2	5.5	57.0	33.1	2.6	0.6
4351	532	0.0	1.1	6.6	68.8	21.2	1.3	0.9
4352	180	0.0	2.8	7.2	62.2	26.7	1.1	0.0
4353	457	0.0	2.2	7.2	65.4	20.4	3.5	1.3
4354	734	0.0	1.9	7.1	65.3	24.0	1.6	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 74

## Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
4355	264	0.4	0.8	7.6	53.8	33.0	3.4	1.1
4356	390	0.0	2.6	7.4	66.2	21.8	1.0	1.0
4358	23	*	*	*	*	*	*	*
4359	593	0.0	2.5	5.9	61.0	26.8	2.7	1.0
4360	218	0.0	0.9	8.7	61.5	27.1	1.4	0.5
4361	255	0.0	2.0	11.0	62.7	22.4	1.6	0.4
4362	279	0.0	2.5	10.4	59.1	26.2	0.7	1.1
4363	680	0.0	2.5	8.4	65.0	22.4	1.2	0.6
4401	758	0.0	2.1	4.9	63.1	27.7	1.7	0.5
4501	779	0.0	2.4	6.2	64.2	24.8	2.1	0.4
4601	1,237	0.1	1.1	6.8	62.9	27.4	1.0	0.7
4701	998	0.0	1.1	5.1	58.6	32.7	2.0	0.5
5050	1,576	0.1	1.0	4.4	57.3	34.6	2.5	0.1
5051	2,681	0.0	1.0	4.3	59.8	32.2	2.6	0.1
5052	2,718	0.0	1.3	4.2	56.3	35.9	2.1	0.1
5055	1,308	0.0	0.4	1.4	49.0	45.3	3.9	0.1
5101	1,508	0.1	0.9	3.7	59.1	33.8	2.4	0.1
5102	17	*	*	*	*	*	*	*
5202	676	0.1	1.5	4.0	62.1	29.6	2.7	0.0
5206	16	*	*	*	*	*	*	*
5601	957	0.0	1.3	5.9	64.4	27.4	1.1	0.0
5801	1,847	0.0	1.8	6.9	63.7	25.7	1.6	0.3
5803	52	*	*	*	*	*	*	*
5804	17	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 75

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
5806	45	*	*	*	*	*	*	*
6101	57	*	*	*	*	*	*	*
6102	86	*	*	*	*	*	*	*
6103	3,333	0.1	2.9	9.0	61.2	24.7	1.4	0.6
6105	9	*	*	*	*	*	*	*
6106	31	*	*	*	*	*	*	*
6301	3,372	0.0	2.2	6.0	59.6	30.0	2.1	0.1
6312	1,762	0.0	0.5	3.0	52.4	40.9	3.1	0.1
6316	1,100	0.0	0.5	2.9	55.7	36.8	4.0	0.1
6318	22	*	*	*	*	*	*	*
6320	490	0.0	0.6	2.9	59.8	33.5	2.9	0.4
6325	315	0.0	1.3	5.1	58.1	34.0	1.0	0.6
6335	931	0.0	0.1	2.3	53.9	39.4	4.1	0.2
6340	1	*	*	*	*	*	*	*
6345	115	0.0	0.0	3.5	60.9	33.0	2.6	0.0
6360	975	0.0	2.2	7.0	59.4	28.7	1.7	1.0
7306	2,065	0.2	2.8	7.5	61.6	26.5	1.2	0.3
7401	1,903	0.1	1.8	8.2	63.1	24.6	1.6	0.7
7601	558	0.0	2.2	6.3	65.6	25.4	0.5	0.0
7850	1,025	0.1	2.6	6.5	66.0	23.9	0.9	0.0
7852	15	*	*	*	*	*	*	*
8101	1,727	0.2	2.2	6.5	56.7	31.7	2.0	0.8
8102	485	0.0	2.5	5.6	60.0	29.7	2.1	0.2
8105	235	0.0	2.6	5.1	53.2	37.4	1.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 76

## Michigan

### Comparison of Age Distribution by Clinic

### Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
8106	574	0.0	1.7	4.5	64.6	27.7	1.2	0.2
8150	7	*	*	*	*	*	*	*
8151	3,165	0.0	1.4	5.5	56.3	33.7	2.7	0.3
8152	24	*	*	*	*	*	*	*
8153	5	*	*	*	*	*	*	*
8155	10	*	*	*	*	*	*	*
8157	1	*	*	*	*	*	*	*
8201	2,775	0.0	1.5	5.0	58.0	33.3	1.9	0.3
8202	3,986	0.0	1.0	3.4	53.4	36.4	5.2	0.6
8203	375	0.0	0.8	5.1	56.5	34.1	3.5	0.0
8250	589	0.0	1.9	4.9	56.4	32.9	3.7	0.2
8252	163	0.0	1.2	6.1	58.9	31.9	1.8	0.0
8255	2,059	0.1	1.4	5.6	61.3	29.9	1.7	0.0
8257	1,801	0.0	0.6	3.1	56.0	35.8	4.4	0.1
8263	527	0.0	0.9	4.7	59.8	32.4	2.1	0.0
8264	790	0.1	1.0	2.5	59.6	33.5	3.0	0.1
8267	634	0.0	1.7	5.2	63.2	27.3	2.4	0.2
8901	870	0.3	3.3	6.7	57.9	28.7	2.6	0.3
8902	1,347	0.1	1.5	5.0	56.6	34.1	2.4	0.2
8903	1,463	0.0	2.9	7.5	59.7	27.2	2.3	0.4
8904	2,171	0.2	2.7	6.5	59.0	29.4	1.8	0.4
8905	1,887	0.0	1.0	3.6	54.4	36.6	4.2	0.3
8909	1,094	0.0	1.6	7.0	59.0	28.8	3.4	0.3
9045	1,371	0.2	2.6	6.7	54.9	31.0	4.2	0.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 4A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 77

## Michigan

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
9050	954	0.1	1.9	6.1	58.5	31.2	1.7	0.5
9051	1,515	0.1	2.0	6.5	55.6	31.4	4.2	0.2
9053	1,386	0.0	2.5	5.7	57.0	31.3	3.0	0.6
9055	7	*	*	*	*	*	*	*
9058	1,277	0.1	1.5	6.3	56.4	32.3	3.0	0.5
9066	984	0.3	2.9	6.3	58.1	29.9	2.2	0.2
9068	850	0.4	2.8	6.6	58.1	28.2	2.8	1.1
9069	1,858	0.2	3.3	6.1	53.9	33.7	2.4	0.3
9070	1,643	0.1	3.2	7.2	56.8	30.0	2.1	0.5
9072	386	0.5	2.3	11.1	57.3	26.7	1.8	0.3
9080	2,520	0.0	1.7	6.1	54.7	34.3	2.5	0.6
9082	110	0.0	1.8	2.7	60.9	31.8	2.7	0.0
9083	7	*	*	*	*	*	*	*
9084	5	*	*	*	*	*	*	*
9087	15	*	*	*	*	*	*	*
<b>Michigan</b>	<b>146,297</b>	<b>0.1</b>	<b>2.0</b>	<b>6.3</b>	<b>59.6</b>	<b>29.4</b>	<b>2.2</b>	<b>0.4</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 78

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	86	*	*	47	*	*	9	*	4	*	100	9.0 (75)	100	13.0 (59)
0102	292	5.8 (128)	62.7 (87)	135	17.0 (45)	54.1 (74)	17	*	13	*	372	8.3 (56)	371	12.7 (53)
0103	305	3.6 (47)	65.6 (124)	145	13.8 (7)	57.2 (102)	15	*	14	*	326	5.5 (6)	327	16.2 (112)
0104	87	*	*	28	*	*	5	*	8	*	112	7.1 (27)	113	13.3 (68)
0105	10	*	*	3	*	*	0	*	0	*	14	*	14	*
0250	104	5.8 (128)	56.7 (14)	73	*	*	5	*	11	*	105	14.3 (141)	105	16.2 (112)
0251	56	*	*	29	*	*	2	*	3	*	68	*	68	*
0252	89	*	*	69	*	*	8	*	3	*	87	*	87	*
0253	116	2.6 (16)	65.5 (122)	90	*	*	3	*	8	*	112	9.8 (99)	112	12.5 (51)
0401	397	6.0 (132)	59.9 (39)	185	17.3 (50)	56.8 (98)	20	*	11	*	397	9.8 (99)	398	16.8 (121)
0402	310	4.2 (80)	60.0 (40)	86	*	*	11	*	13	*	335	12.2 (132)	335	11.6 (36)
0403	125	5.6 (124)	62.4 (75)	67	*	*	4	*	4	*	126	12.7 (138)	126	11.1 (28)
0404	101	3.0 (25)	62.4 (75)	53	*	*	3	*	3	*	110	10.9 (119)	110	16.4 (120)
0501	262	4.2 (80)	56.1 (10)	70	*	*	43	*	51	*	396	8.8 (69)	396	12.4 (49)
0503	55	*	*	18	*	*	6	*	8	*	70	*	70	*
0504	310	2.9 (23)	64.2 (110)	132	17.4 (52)	50.0 (26)	31	*	35	*	340	7.1 (27)	340	12.9 (57)
0505	154	3.9 (62)	57.8 (21)	32	*	*	28	*	21	*	213	6.6 (20)	213	16.9 (123)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 79

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	120	5.0 (112)	55.0 (6)	27	*	*	14	*	21	*	156	7.1 (27)	156	10.9 (26)
0507	38	*	*	11	*	*	3	*	16	*	70	*	70	*
0508	12	*	*	7	*	*	1	*	3	*	19	*	19	*
0650	932	4.7 (97)	62.2 (71)	622	14.5 (13)	54.2 (78)	32	*	36	*	997	10.7 (113)	998	14.5 (88)
0651	452	3.8 (58)	61.9 (67)	210	17.6 (54)	53.8 (70)	16	*	27	*	513	7.6 (41)	515	12.0 (42)
0652	258	2.7 (18)	66.3 (127)	202	16.8 (42)	53.0 (62)	14	*	18	*	292	8.9 (71)	293	17.1 (126)
0653	373	3.5 (36)	62.7 (87)	274	14.2 (11)	52.9 (61)	23	*	29	*	381	8.7 (67)	383	16.2 (112)
0654	241	4.1 (74)	61.4 (59)	135	17.8 (56)	52.6 (52)	13	*	9	*	382	7.6 (41)	383	11.5 (34)
0655	285	4.2 (80)	61.1 (53)	157	18.5 (70)	54.1 (74)	11	*	5	*	284	10.6 (109)	285	19.3 (137)
0657	26	*	*	11	*	*	2	*	1	*	45	*	45	*
0701	426	3.3 (30)	57.3 (19)	303	18.2 (66)	45.5 (7)	10	*	4	*	439	9.3 (79)	439	14.8 (96)
0702	67	*	*	52	*	*	0	*	5	*	63	*	63	*
0703	132	0.8 (2)	59.1 (29)	100	20.0 (88)	52.0 (43)	9	*	8	*	137	15.3 (144)	137	10.2 (18)
0704	44	*	*	28	*	*	3	*	3	*	44	*	45	*
0705	60	*	*	46	*	*	4	*	0	*	65	*	65	*
0706	3	*	*	2	*	*	0	*	0	*	3	*	3	*
0850	403	6.2 (134)	59.8 (35)	151	18.5 (70)	47.0 (10)	16	*	7	*	649	11.7 (128)	650	16.3 (116)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 80

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

### Comparison of Maternal Health Indicators <sup>(2,3)</sup>

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0851	338	5.9 (130)	59.8 (35)	131	12.2 (3)	57.3 (103)	23	*	8	*	559	12.5 (135)	563	15.8 (109)
0901	1,162	4.2 (80)	61.4 (59)	625	16.2 (34)	55.5 (88)	38	*	22	*	1,388	8.1 (53)	1,388	13.1 (63)
0902	26	*	*	13	*	*	0	*	1	*	30	*	30	*
1051	931	5.5 (122)	53.8 (5)	693	15.9 (30)	52.1 (45)	33	*	21	*	1,048	7.3 (34)	1,052	10.8 (25)
1052	2	*	*	1	*	*	1	*	0	*	3	*	3	*
1101	1,261	3.6 (47)	62.5 (81)	888	18.7 (74)	53.9 (73)	87	*	101	27.7 (8)	1,317	9.3 (79)	1,318	19.5 (140)
1102	709	5.1 (116)	60.6 (47)	481	15.2 (20)	57.6 (104)	43	*	53	*	684	8.6 (66)	686	17.2 (127)
1106	57	*	*	46	*	*	4	*	2	*	55	*	55	*
1250	404	4.7 (97)	61.1 (53)	153	24.8 (112)	47.7 (14)	28	*	33	*	894	10.5 (108)	895	9.7 (14)
1251	523	4.8 (100)	62.7 (87)	371	15.4 (24)	50.9 (36)	16	*	41	*	576	5.9 (9)	577	6.1 (5)
1252	403	5.2 (119)	64.5 (113)	142	19.7 (85)	50.0 (26)	10	*	16	*	465	9.7 (95)	466	10.7 (21)
1253	593	5.6 (124)	60.2 (44)	232	20.7 (98)	52.6 (52)	21	*	26	*	619	10.7 (113)	619	13.4 (69)
1350	1,954	3.9 (62)	61.5 (62)	1,328	18.8 (75)	52.1 (45)	46	*	17	*	2,201	7.2 (33)	2,203	10.9 (26)
1351	20	*	*	9	*	*	3	*	0	*	23	*	23	*
1352	246	4.9 (107)	63.4 (98)	154	18.8 (75)	59.7 (111)	3	*	2	*	368	4.3 (2)	368	6.5 (6)
1401	157	3.8 (58)	50.3 (1)	77	*	*	9	*	21	*	184	5.4 (4)	184	15.2 (102)
1402	83	*	*	43	*	*	2	*	4	*	90	*	90	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 81

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1403	14	*	*	8	*	*	0	*	2	*	20	*	20	*
1501	593	2.7 (18)	69.1 (133)	250	34.8 (115)	38.8 (2)	80	*	83	*	668	12.6 (136)	668	11.7 (37)
1503	696	4.5 (90)	62.6 (84)	490	19.0 (79)	52.0 (43)	69	*	69	*	776	10.1 (104)	777	15.7 (107)
1504	626	6.1 (133)	57.0 (16)	272	14.3 (12)	52.2 (47)	171	35.1 (2)	241	27.4 (7)	1,726	9.6 (89)	1,727	8.8 (11)
1505	74	*	*	13	*	*	4	*	6	*	115	4.3 (2)	115	7.8 (7)
1506	436	3.9 (62)	64.7 (117)	60	*	*	42	*	74	*	505	12.9 (139)	506	13.4 (69)
1507	357	6.7 (137)	56.6 (13)	57	*	*	29	*	78	*	479	9.2 (78)	480	9.2 (12)
1508	4	*	*	3	*	*	0	*	1	*	4	*	4	*
1509	271	5.2 (119)	64.6 (114)	143	19.6 (84)	53.8 (70)	36	*	60	*	337	8.3 (56)	338	8.6 (10)
1511	474	4.4 (87)	62.9 (92)	264	19.7 (85)	50.0 (26)	60	*	103	23.3 (4)	574	11.3 (122)	575	14.6 (89)
1512	258	4.3 (84)	62.0 (68)	103	11.7 (2)	67.0 (116)	29	*	88	*	342	6.4 (16)	343	16.3 (116)
1514	297	3.7 (55)	51.9 (2)	92	*	*	43	*	154	20.8 (2)	552	9.1 (77)	553	15.7 (107)
1701	486	3.3 (30)	60.3 (45)	337	18.1 (65)	52.5 (51)	20	*	29	*	492	10.8 (117)	494	8.3 (9)
1950	537	2.8 (20)	62.4 (75)	342	14.6 (14)	56.7 (96)	12	*	12	*	567	11.8 (130)	568	14.4 (85)
1952	126	1.6 (6)	69.8 (134)	86	*	*	8	*	4	*	134	9.0 (75)	134	14.9 (98)
1953	214	6.5 (136)	58.4 (24)	142	15.5 (25)	52.8 (56)	7	*	7	*	225	9.3 (79)	225	17.8 (134)
1954	352	1.7 (7)	69.0 (132)	228	25.0 (113)	49.6 (22)	9	*	5	*	367	15.0 (143)	367	14.7 (92)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 82

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1955	106	2.8 (20)	62.3 (72)	70	*	*	1	*	3	*	103	12.6 (136)	103	15.5 (105)
1957	336	4.5 (90)	69.9 (135)	206	14.1 (10)	48.5 (18)	12	*	3	*	356	16.3 (145)	356	14.0 (77)
1963	45	*	*	27	*	*	0	*	0	*	52	*	53	*
1964	1	*	*	1	*	*	0	*	0	*	1	*	1	*
2150	546	3.5 (36)	62.6 (84)	398	21.6 (102)	47.2 (12)	39	*	72	*	532	10.7 (113)	532	14.3 (83)
2151	191	4.7 (97)	55.5 (8)	150	18.0 (61)	55.3 (87)	12	*	25	*	173	6.4 (16)	173	17.3 (128)
2152	30	*	*	16	*	*	3	*	1	*	29	*	30	*
2251	268	5.6 (124)	57.1 (18)	123	20.3 (96)	50.4 (32)	9	*	5	*	355	2.8 (1)	356	16.3 (116)
2252	94	*	*	23	*	*	4	*	1	*	148	10.1 (104)	148	11.5 (34)
2554	34	*	*	15	*	*	8	*	4	*	47	*	47	*
2555	4,979	4.3 (84)	60.9 (51)	2,391	18.8 (75)	52.8 (56)	183	40.4 (5)	96	*	6,580	8.9 (71)	6,591	16.8 (121)
2556	144	3.5 (36)	59.7 (33)	64	*	*	3	*	4	*	174	8.0 (49)	174	15.5 (105)
2557	2	*	*	1	*	*	0	*	0	*	2	*	2	*
2558	1	*	*	0	*	*	0	*	0	*	0	*	0	*
2560	2	*	*	0	*	*	0	*	0	*	2	*	2	*
2561	1	*	*	1	*	*	1	*	0	*	1	*	1	*
2562	2	*	*	0	*	*	0	*	0	*	6	*	6	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 83

Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2563	6	*	*	4	*	*	1	*	1	*	10	*	10	*
2564	8	*	*	1	*	*	1	*	1	*	12	*	12	*
3001	812	3.9 (62)	65.5 (122)	287	19.9 (87)	43.2 (4)	17	*	21	*	1,126	8.0 (49)	1,126	14.2 (81)
3002	140	1.4 (3)	76.4 (139)	50	*	*	4	*	7	*	184	7.6 (41)	184	16.3 (116)
3004	144	2.8 (20)	64.6 (114)	41	*	*	1	*	1	*	244	9.4 (82)	244	14.8 (96)
3005	97	*	*	30	*	*	4	*	3	*	146	11.6 (125)	146	10.3 (20)
3007	6	*	*	0	*	*	0	*	0	*	8	*	8	*
3201	350	4.3 (84)	68.6 (131)	259	20.1 (91)	56.0 (95)	32	*	42	*	387	11.4 (123)	388	14.7 (92)
3203	1	*	*	0	*	*	0	*	0	*	1	*	1	*
3301	3,459	3.6 (47)	61.6 (63)	2,420	19.0 (79)	54.3 (80)	116	36.2 (3)	36	*	4,069	7.4 (38)	4,070	12.9 (57)
3304	82	*	*	56	*	*	1	*	1	*	96	*	96	*
3305	127	5.5 (122)	56.7 (14)	89	*	*	4	*	0	*	150	10.7 (113)	150	10.7 (21)
3306	199	3.0 (25)	65.3 (120)	146	16.4 (36)	56.8 (98)	5	*	1	*	226	6.2 (11)	226	11.1 (28)
3308	105	4.8 (100)	60.0 (40)	77	*	*	3	*	0	*	103	6.8 (22)	103	14.6 (89)
3312	358	3.9 (62)	62.6 (84)	237	21.9 (104)	51.9 (41)	10	*	0	*	402	9.7 (95)	402	14.7 (92)
3550	64	*	*	30	*	*	1	*	0	*	75	*	75	*
3551	5	*	*	1	*	*	0	*	0	*	8	*	8	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 84

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3552	14	*	*	5	*	*	1	*	0	*	12	*	12	*
3701	631	5.1 (116)	63.7 (104)	462	20.1 (91)	50.9 (36)	50	*	66	*	806	11.0 (120)	809	19.3 (137)
3850	2,052	4.0 (68)	60.0 (40)	1,362	15.3 (22)	54.6 (81)	70	*	68	*	2,638	8.8 (69)	2,638	13.0 (59)
3901	1,166	3.5 (36)	61.6 (63)	506	17.0 (45)	55.7 (91)	24	*	11	*	1,358	8.4 (59)	1,357	12.8 (55)
3902	33	*	*	15	*	*	3	*	0	*	31	*	31	*
3903	6	*	*	3	*	*	0	*	1	*	7	*	7	*
3950	1,538	4.1 (74)	62.4 (75)	838	15.0 (18)	53.2 (65)	29	*	15	*	1,835	10.6 (109)	1,838	11.8 (39)
3954	5	*	*	4	*	*	0	*	0	*	6	*	6	*
4106	964	3.2 (27)	62.1 (69)	318	21.7 (103)	43.4 (5)	227	33.9 (1)	312	20.8 (2)	2,007	7.9 (47)	2,011	15.0 (101)
4107	1,863	4.0 (68)	58.3 (23)	687	16.4 (36)	52.8 (56)	261	43.7 (6)	432	24.8 (5)	2,610	8.0 (49)	2,615	17.0 (124)
4109	380	3.2 (27)	57.9 (22)	228	13.2 (4)	57.9 (106)	56	*	127	20.5 (1)	600	6.7 (21)	601	17.0 (124)
4110	1	*	*	1	*	*	0	*	0	*	2	*	2	*
4112	539	1.5 (5)	68.1 (130)	412	50.7 (116)	24.3 (1)	15	*	10	*	527	18.8 (146)	527	20.1 (144)
4115	1,614	3.8 (58)	60.1 (43)	450	16.9 (43)	52.7 (55)	331	36.3 (4)	577	26.7 (6)	3,163	9.6 (89)	3,164	16.0 (110)
4351	477	3.6 (47)	62.5 (81)	284	22.9 (107)	46.8 (9)	12	*	10	*	488	11.9 (131)	488	20.1 (144)
4352	159	5.0 (112)	58.5 (26)	116	20.7 (98)	50.0 (26)	4	*	2	*	166	10.2 (106)	166	19.3 (137)
4353	351	2.6 (16)	70.1 (136)	264	19.3 (83)	54.9 (83)	19	*	19	*	405	14.3 (141)	405	14.6 (89)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 85

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4354	470	3.6 (47)	65.7 (125)	155	21.9 (104)	47.1 (11)	19	*	11	*	691	7.5 (39)	691	19.7 (142)
4355	136	4.4 (87)	58.8 (28)	74	*	*	4	*	3	*	245	10.6 (109)	245	11.4 (32)
4356	360	1.4 (3)	64.4 (112)	266	16.2 (34)	51.9 (41)	6	*	7	*	359	11.4 (123)	359	17.8 (134)
4358	15	*	*	14	*	*	0	*	0	*	22	*	22	*
4359	518	3.7 (55)	65.4 (121)	368	23.6 (108)	47.3 (13)	4	*	12	*	547	9.9 (102)	547	14.4 (85)
4360	205	4.9 (107)	62.4 (75)	139	27.3 (114)	49.6 (22)	5	*	2	*	196	7.1 (27)	196	12.2 (47)
4361	226	0.4 (1)	59.7 (33)	129	24.0 (110)	49.6 (22)	3	*	5	*	235	8.5 (61)	235	11.9 (41)
4362	231	4.8 (100)	55.0 (6)	160	13.8 (7)	58.1 (108)	6	*	6	*	257	8.9 (71)	258	10.1 (17)
4363	569	5.6 (124)	58.5 (26)	400	20.0 (88)	50.8 (35)	17	*	12	*	642	10.6 (109)	642	13.2 (64)
4401	730	4.5 (90)	59.2 (31)	615	20.0 (88)	51.7 (40)	12	*	9	*	703	5.4 (4)	704	9.8 (15)
4501	595	4.0 (68)	63.5 (99)	394	20.8 (100)	49.5 (21)	28	*	21	*	743	11.6 (125)	744	14.9 (98)
4601	959	3.4 (33)	62.3 (72)	638	15.5 (25)	50.6 (33)	27	*	24	*	1,105	9.4 (82)	1,106	15.4 (104)
4701	890	4.9 (107)	57.5 (20)	583	18.0 (61)	52.3 (49)	31	*	36	*	916	9.6 (89)	917	20.0 (143)
5050	1,498	4.1 (74)	63.5 (99)	1,253	17.1 (48)	53.1 (63)	26	*	10	*	1,479	9.5 (88)	1,480	14.3 (83)
5051	2,488	3.4 (33)	61.8 (66)	2,010	18.0 (61)	54.7 (82)	53	*	20	*	2,468	8.7 (67)	2,469	10.7 (21)
5052	2,544	3.5 (36)	61.4 (59)	2,072	20.2 (93)	48.6 (20)	43	*	16	*	2,582	11.6 (125)	2,582	12.2 (47)
5055	812	1.7 (7)	60.8 (48)	501	23.6 (108)	46.5 (8)	16	*	6	*	1,260	13.0 (140)	1,260	4.6 (4)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 86

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

### Comparison of Maternal Health Indicators (2,3)

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5101	1,133	4.1 (74)	60.5 (46)	800	13.3 (5)	55.9 (94)	82	*	72	*	1,402	9.7 (95)	1,405	12.6 (52)
5102	14	*	*	8	*	*	3	*	0	*	15	*	15	*
5202	589	4.4 (87)	59.1 (29)	399	17.8 (56)	57.9 (106)	18	*	11	*	615	9.8 (99)	617	13.0 (59)
5206	12	*	*	9	*	*	0	*	1	*	16	*	16	*
5601	696	3.6 (47)	62.9 (92)	238	10.5 (1)	55.0 (84)	36	*	26	*	901	10.8 (117)	900	12.1 (44)
5801	1,679	5.0 (112)	63.3 (96)	1,287	17.1 (48)	54.2 (78)	52	*	27	*	1,688	8.5 (61)	1,689	17.5 (130)
5803	50	*	*	42	*	*	3	*	2	*	48	*	48	*
5804	17	*	*	12	*	*	1	*	0	*	16	*	16	*
5806	31	*	*	18	*	*	0	*	2	*	43	*	43	*
6101	54	*	*	40	*	*	0	*	0	*	51	*	52	*
6102	78	*	*	64	*	*	1	*	0	*	82	*	82	*
6103	3,006	3.5 (36)	64.3 (111)	2,319	17.9 (58)	52.2 (47)	46	*	18	*	3,089	9.6 (89)	3,091	18.7 (136)
6105	5	*	*	3	*	*	0	*	0	*	8	*	8	*
6106	28	*	*	19	*	*	0	*	2	*	26	*	26	*
6301	2,813	3.6 (47)	62.1 (69)	1,751	18.0 (61)	48.5 (18)	76	*	44	*	3,069	5.7 (7)	3,069	13.0 (59)
6312	1,528	3.7 (55)	61.1 (53)	965	17.0 (45)	53.1 (63)	42	*	29	*	1,593	6.5 (18)	1,596	9.8 (15)
6316	970	4.9 (107)	56.5 (12)	614	17.9 (58)	54.1 (74)	26	*	14	*	1,010	6.9 (24)	1,010	11.1 (28)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 87

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6318	7	*	*	0	*	*	2	*	0	*	21	*	21	*
6320	339	3.5 (36)	63.7 (104)	165	15.2 (20)	55.8 (92)	8	*	10	*	436	9.6 (89)	437	16.0 (110)
6325	221	5.9 (130)	55.7 (9)	113	13.3 (5)	62.8 (114)	7	*	9	*	281	8.5 (61)	281	17.4 (129)
6335	597	6.4 (135)	59.6 (32)	288	14.9 (16)	50.7 (34)	31	*	33	*	888	12.2 (132)	887	13.2 (64)
6340	1	*	*	0	*	*	0	*	0	*	0	*	0	*
6345	81	*	*	47	*	*	0	*	3	*	105	12.4 (134)	105	16.2 (112)
6360	651	4.0 (68)	58.4 (24)	292	16.1 (33)	54.1 (74)	20	*	20	*	887	7.3 (34)	888	14.0 (77)
7306	1,554	2.9 (23)	62.3 (72)	972	15.8 (28)	52.6 (52)	48	*	41	*	1,935	9.4 (82)	1,937	11.4 (32)
7401	1,790	4.1 (74)	61.7 (65)	1,447	17.7 (55)	53.8 (70)	80	*	102	34.3 (10)	1,708	8.3 (56)	1,712	14.7 (92)
7601	123	7.3 (139)	56.1 (10)	29	*	*	14	*	21	*	496	7.7 (44)	496	12.1 (44)
7850	897	4.6 (94)	63.5 (99)	724	21.0 (101)	50.3 (31)	13	*	8	*	933	7.1 (27)	934	9.6 (13)
7852	14	*	*	10	*	*	0	*	0	*	13	*	13	*
8101	1,480	3.2 (27)	64.1 (109)	1,010	18.3 (68)	52.8 (56)	40	*	11	*	1,628	7.7 (44)	1,627	17.5 (130)
8102	341	2.3 (13)	68.0 (129)	176	16.5 (39)	56.8 (98)	8	*	3	*	451	7.3 (34)	452	10.2 (18)
8105	187	4.8 (100)	71.1 (138)	85	*	*	31	*	3	*	189	6.9 (24)	189	4.2 (2)
8106	277	5.4 (121)	57.0 (16)	113	18.6 (72)	55.8 (92)	11	*	2	*	542	6.3 (12)	542	17.7 (133)
8150	6	*	*	2	*	*	0	*	0	*	7	*	7	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 88

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8151	2,463	3.9 (62)	61.3 (57)	1,544	20.5 (97)	48.1 (16)	175	44.0 (7)	254	29.1 (9)	2,863	9.6 (89)	2,864	14.4 (85)
8152	19	*	*	11	*	*	2	*	3	*	22	*	22	*
8153	5	*	*	1	*	*	2	*	0	*	3	*	3	*
8155	6	*	*	2	*	*	0	*	1	*	8	*	8	*
8157	1	*	*	1	*	*	0	*	0	*	1	*	1	*
8201	1,963	3.5 (36)	61.2 (56)	1,209	15.6 (27)	55.6 (90)	39	*	31	*	2,474	8.9 (71)	2,474	11.7 (37)
8202	1,813	4.6 (94)	52.3 (3)	938	18.9 (78)	48.1 (16)	74	*	23	*	3,856	6.3 (12)	3,854	3.1 (1)
8203	313	5.1 (116)	63.6 (103)	204	16.7 (41)	51.5 (38)	12	*	5	*	329	9.7 (95)	328	14.0 (77)
8250	440	4.8 (100)	62.7 (87)	263	16.0 (31)	53.2 (65)	17	*	8	*	508	7.5 (39)	508	12.0 (42)
8252	101	2.0 (10)	67.3 (128)	40	*	*	3	*	2	*	143	11.2 (121)	144	13.9 (76)
8255	1,658	5.0 (112)	60.8 (48)	1,119	18.3 (68)	53.3 (67)	64	*	18	*	1,822	8.5 (61)	1,822	13.8 (75)
8257	797	4.9 (107)	53.3 (4)	413	24.0 (110)	41.4 (3)	23	*	13	*	1,715	8.4 (59)	1,717	4.5 (3)
8263	438	3.4 (33)	59.8 (35)	260	13.8 (7)	61.9 (113)	16	*	3	*	442	10.0 (103)	441	11.8 (39)
8264	585	3.6 (47)	64.6 (114)	385	15.3 (22)	59.2 (110)	19	*	6	*	708	7.3 (34)	707	13.2 (64)
8267	463	4.1 (74)	65.0 (118)	307	18.2 (66)	57.0 (101)	12	*	5	*	581	8.1 (53)	579	11.2 (31)
8901	716	3.5 (36)	63.5 (99)	456	17.3 (50)	53.3 (67)	21	*	6	*	825	6.3 (12)	824	13.7 (73)
8902	883	4.0 (68)	60.8 (48)	463	18.6 (72)	52.3 (49)	32	*	5	*	1,254	7.8 (46)	1,252	12.4 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 89

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8903	1,026	2.2 (11)	65.9 (126)	608	14.8 (15)	60.0 (112)	22	*	5	*	1,347	6.8 (22)	1,353	17.5 (130)
8904	1,722	3.5 (36)	65.2 (119)	987	16.6 (40)	53.4 (69)	63	*	9	*	2,023	8.1 (53)	2,027	14.9 (98)
8905	1,235	4.0 (68)	59.8 (35)	630	19.2 (82)	51.6 (39)	31	*	7	*	1,781	11.7 (128)	1,783	12.1 (44)
8909	825	4.6 (94)	63.3 (96)	432	16.0 (31)	58.6 (109)	14	*	3	*	1,009	8.0 (49)	1,008	14.1 (80)
9045	942	2.2 (11)	70.7 (137)	531	16.4 (36)	48.0 (15)	26	*	6	*	1,293	9.4 (82)	1,293	8.1 (8)
9050	805	4.8 (100)	62.4 (75)	207	15.0 (18)	63.8 (115)	17	*	4	*	901	7.0 (26)	901	12.8 (55)
9051	706	1.8 (9)	63.7 (104)	312	20.2 (93)	44.6 (6)	23	*	2	*	1,437	10.2 (106)	1,437	10.7 (21)
9053	1,228	3.3 (30)	63.7 (104)	873	14.9 (16)	57.7 (105)	20	*	4	*	1,310	6.5 (18)	1,310	13.2 (64)
9055	7	*	*	4	*	*	0	*	0	*	7	*	7	*
9058	775	4.5 (90)	62.8 (91)	504	19.0 (79)	52.8 (56)	22	*	2	*	1,207	9.4 (82)	1,207	15.2 (102)
9066	736	3.8 (58)	61.0 (52)	421	21.9 (104)	49.9 (25)	25	*	5	*	938	6.3 (12)	943	20.5 (146)
9068	786	2.4 (14)	64.0 (108)	600	20.2 (93)	55.5 (88)	17	*	3	*	803	6.1 (10)	805	13.4 (69)
9069	1,361	3.5 (36)	62.5 (81)	800	17.4 (52)	50.0 (26)	55	*	5	*	1,774	7.1 (27)	1,775	12.7 (53)
9070	1,451	4.8 (100)	61.3 (57)	1,012	15.8 (28)	55.0 (84)	36	*	4	*	1,532	5.8 (8)	1,535	13.5 (72)
9072	298	6.7 (137)	63.1 (94)	134	17.9 (58)	56.7 (96)	25	*	4	*	341	7.9 (47)	341	19.6 (141)
9080	1,652	2.5 (15)	63.1 (94)	871	16.9 (43)	55.1 (86)	60	*	10	*	2,397	9.4 (82)	2,395	13.7 (73)
9082	90	*	*	49	*	*	1	*	0	*	106	8.5 (61)	106	14.2 (81)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 5A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 90

## Michigan

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)														
Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9083	4	*	*	2	*	*	0	*	0	*	7	*	7	*
9084	4	*	*	3	*	*	1	*	1	*	4	*	4	*
9087	14	*	*	8	*	*	0	*	0	*	13	*	13	*
Michigan	108,849	3.9	61.7	65,508	18.1	52.5	4,914	36.3	4,945	26.1	135,141	8.8	135,252	13.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy

(ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 91

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	95	*	*	110	26.4 (99)	104	30.8 (40)	81	*
0102	361	87.8 (93)	3.9 (100)	392	35.5 (34)	386	22.8 (87)	311	88.4 (110)
0103	315	87.3 (96)	5.7 (114)	354	33.3 (48)	339	18.6 (110)	273	90.5 (78)
0104	111	91.9 (45)	2.7 (66)	120	42.5 (7)	118	25.4 (65)	99	*
0105	14	*	*	14	*	14	*	14	*
0250	93	*	*	114	48.2 (1)	101	34.7 (25)	89	*
0251	66	*	*	75	*	68	*	59	*
0252	87	*	*	94	*	90	*	73	*
0253	110	89.1 (81)	2.7 (66)	119	45.4 (2)	114	22.8 (87)	99	*
0401	381	89.8 (65)	6.3 (120)	429	34.0 (45)	410	24.4 (73)	331	95.8 (7)
0402	315	81.9 (129)	7.6 (128)	364	32.7 (54)	352	31.8 (34)	294	94.6 (19)
0403	121	87.6 (95)	0.8 (7)	134	32.8 (52)	128	24.2 (74)	103	97.1 (2)
0404	107	98.1 (2)	0.0 (1)	115	29.6 (73)	110	44.5 (5)	94	*
0501	389	95.9 (10)	1.0 (12)	417	29.5 (75)	383	29.2 (48)	305	91.5 (59)
0503	70	*	*	70	*	65	*	50	*
0504	332	90.4 (58)	2.7 (66)	349	28.4 (85)	339	40.4 (14)	257	93.0 (43)
0505	214	91.1 (52)	4.7 (109)	227	28.2 (89)	221	35.3 (23)	182	90.7 (74)
0506	149	87.9 (91)	3.4 (87)	165	29.7 (70)	161	26.1 (62)	121	87.6 (116)
0507	69	*	*	69	*	66	*	43	*
0508	18	*	*	19	*	18	*	10	*
0650	946	89.9 (61)	4.0 (103)	1,058	36.5 (27)	1,008	31.9 (33)	831	91.5 (59)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 92

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0651	493	87.8 (93)	4.3 (105)	555	40.7 (12)	507	28.4 (50)	456	89.3 (102)
0652	276	89.9 (61)	3.3 (85)	294	34.4 (40)	281	23.1 (85)	229	84.3 (131)
0653	359	77.4 (138)	14.2 (142)	403	41.2 (11)	388	32.2 (32)	315	89.2 (105)
0654	334	85.9 (106)	8.4 (130)	402	44.5 (4)	388	20.4 (102)	322	87.6 (116)
0655	277	92.1 (44)	2.2 (48)	300	38.7 (18)	266	27.8 (52)	242	89.3 (102)
0657	44	*	*	47	*	45	*	40	*
0701	405	86.7 (99)	5.9 (117)	481	32.8 (52)	455	38.0 (18)	369	93.8 (30)
0702	61	*	*	66	*	62	*	44	*
0703	135	94.1 (18)	1.5 (24)	150	28.7 (83)	138	39.9 (15)	96	*
0704	44	*	*	49	*	49	*	42	*
0705	63	*	*	67	*	66	*	53	*
0706	3	*	*	3	*	3	*	3	*
0850	634	89.4 (77)	2.2 (48)	691	27.2 (93)	654	31.2 (37)	497	93.2 (40)
0851	547	92.5 (42)	1.6 (27)	588	26.9 (95)	564	32.4 (30)	415	96.9 (3)
0901	1,282	83.2 (122)	8.7 (131)	1,473	38.3 (19)	1,423	22.5 (91)	1,167	90.1 (86)
0902	27	*	*	34	*	33	*	25	*
1051	977	86.2 (103)	7.3 (125)	1,114	29.0 (80)	1,067	29.3 (46)	776	90.5 (78)
1052	3	*	*	2	*	3	*	2	*
1101	1,230	78.0 (137)	4.6 (108)	1,432	29.2 (78)	1,316	19.8 (106)	1,041	88.0 (113)
1102	621	82.3 (127)	6.8 (123)	785	33.5 (46)	741	24.2 (74)	568	95.4 (10)
1106	52	*	*	64	*	61	*	44	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 93

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1250	808	85.3 (109)	5.6 (112)	944	29.7 (70)	858	29.5 (45)	706	89.7 (94)
1251	565	90.8 (55)	3.7 (97)	606	28.9 (81)	574	43.7 (6)	439	94.5 (22)
1252	397	73.3 (141)	11.6 (139)	512	34.4 (40)	497	21.7 (94)	393	89.6 (97)
1253	521	74.7 (140)	13.1 (141)	671	37.0 (26)	642	21.7 (94)	529	90.0 (89)
1350	2,126	95.8 (11)	1.5 (24)	2,333	31.8 (61)	2,066	32.3 (31)	1,711	89.0 (106)
1351	23	*	*	26	*	27	*	27	*
1352	362	97.0 (4)	1.7 (32)	395	32.9 (50)	373	4.8 (144)	287	90.9 (68)
1401	173	87.9 (91)	5.8 (115)	193	30.6 (67)	186	43.0 (7)	142	90.8 (70)
1402	85	*	*	97	*	84	*	70	*
1403	19	*	*	20	*	21	*	14	*
1501	582	84.4 (117)	6.5 (121)	735	32.5 (55)	616	12.5 (128)	547	84.5 (130)
1503	747	92.9 (34)	3.3 (85)	817	38.2 (21)	757	21.5 (96)	619	81.9 (134)
1504	1,686	94.0 (22)	1.7 (32)	1,872	29.3 (77)	1,757	23.8 (80)	1,310	87.5 (118)
1505	113	85.0 (111)	10.6 (137)	129	42.6 (6)	101	22.8 (87)	130	90.0 (89)
1506	441	84.8 (115)	9.5 (135)	543	35.7 (32)	514	9.1 (138)	415	93.3 (36)
1507	468	93.8 (25)	2.8 (73)	519	31.0 (64)	480	26.3 (59)	349	95.4 (10)
1508	4	*	*	6	*	5	*	6	*
1509	327	91.1 (52)	2.1 (45)	379	34.6 (38)	347	21.9 (93)	291	93.1 (41)
1511	549	93.4 (27)	0.9 (8)	602	34.6 (38)	585	22.9 (86)	453	88.5 (109)
1512	328	96.0 (8)	1.2 (18)	359	22.3 (117)	333	31.5 (35)	209	93.3 (36)
1514	512	94.1 (18)	1.6 (27)	568	22.0 (120)	446	31.2 (37)	313	93.9 (29)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 94

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1701	487	96.1 (7)	1.4 (23)	529	36.5 (27)	505	46.5 (4)	427	89.7 (94)
1950	538	89.8 (65)	4.3 (105)	594	31.6 (62)	502	29.9 (42)	445	96.0 (5)
1952	126	89.7 (68)	5.6 (112)	138	26.8 (96)	113	34.5 (26)	94	*
1953	211	90.0 (60)	3.8 (99)	243	29.6 (73)	213	37.1 (21)	183	97.3 (1)
1954	347	90.2 (59)	3.5 (91)	384	37.5 (22)	320	33.1 (28)	317	94.6 (19)
1955	97	*	*	110	35.5 (34)	90	*	80	*
1957	336	90.8 (55)	1.8 (37)	373	30.0 (69)	284	41.2 (10)	273	94.5 (22)
1963	52	*	*	55	*	40	*	39	*
1964	1	*	*	1	*	1	*	1	*
2150	531	93.4 (27)	0.9 (8)	572	32.2 (57)	547	26.5 (57)	431	94.9 (15)
2151	161	90.7 (57)	3.7 (97)	206	26.7 (97)	161	30.4 (41)	129	93.0 (43)
2152	30	*	*	32	*	30	*	23	*
2251	347	94.2 (15)	3.2 (82)	387	35.7 (32)	371	25.6 (63)	305	93.1 (41)
2252	139	94.2 (15)	3.6 (94)	163	45.4 (2)	152	60.5 (2)	135	90.4 (81)
2554	48	*	*	49	*	37	*	39	*
2555	6,293	89.2 (80)	1.7 (32)	7,047	25.2 (104)	5,571	14.6 (123)	5,096	91.3 (63)
2556	163	84.7 (116)	2.5 (60)	183	24.6 (107)	124	14.5 (124)	142	89.4 (99)
2557	2	*	*	2	*	1	*	1	*
2558	0	*	*	1	*	0	*	1	*
2560	2	*	*	3	*	3	*	3	*
2561	1	*	*	1	*	1	*	1	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 95

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2562	6	*	*	6	*	4	*	6	*
2563	10	*	*	13	*	7	*	8	*
2564	12	*	*	17	*	14	*	14	*
3001	1,073	85.6 (107)	3.9 (100)	1,168	34.1 (43)	1,108	12.3 (129)	934	86.0 (124)
3002	174	89.7 (68)	1.7 (32)	193	39.9 (15)	185	13.0 (127)	157	90.4 (81)
3004	236	92.8 (38)	2.1 (45)	249	32.9 (50)	247	23.9 (79)	181	89.0 (106)
3005	139	89.9 (61)	2.9 (77)	149	38.9 (17)	150	20.7 (99)	129	94.6 (19)
3007	8	*	*	9	*	8	*	6	*
3201	365	86.8 (98)	3.0 (78)	398	24.6 (107)	368	36.4 (22)	265	90.9 (68)
3203	1	*	*	1	*	1	*	1	*
3301	3,757	89.8 (65)	3.2 (82)	4,232	26.7 (97)	3,871	9.9 (135)	3,111	89.9 (92)
3304	88	*	*	102	39.2 (16)	101	13.9 (125)	77	*
3305	137	89.1 (81)	3.6 (94)	155	28.4 (85)	146	10.3 (132)	104	90.4 (81)
3306	207	92.3 (43)	2.4 (58)	232	28.0 (90)	208	14.9 (121)	167	91.0 (66)
3308	98	*	*	112	24.1 (111)	102	18.6 (110)	62	*
3312	359	89.7 (68)	3.6 (94)	418	32.5 (55)	397	10.3 (132)	325	90.8 (70)
3550	71	*	*	77	*	75	*	63	*
3551	8	*	*	10	*	10	*	9	*
3552	12	*	*	14	*	14	*	14	*
3701	795	92.8 (38)	3.0 (78)	862	28.4 (85)	820	22.8 (87)	601	95.5 (9)
3850	2,443	88.1 (88)	4.8 (110)	2,785	37.5 (22)	2,620	23.6 (81)	2,168	88.3 (111)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 96

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3901	1,282	89.9 (61)	2.0 (41)	1,496	35.0 (37)	1,419	15.8 (118)	1,151	85.6 (125)
3902	31	*	*	42	*	40	*	41	*
3903	7	*	*	7	*	7	*	5	*
3950	1,754	88.1 (88)	3.2 (82)	1,978	31.1 (63)	1,842	20.3 (103)	1,417	90.1 (86)
3954	6	*	*	6	*	6	*	6	*
4106	1,864	87.3 (96)	2.8 (73)	2,132	25.3 (102)	1,911	11.6 (130)	1,584	85.0 (129)
4107	2,542	89.4 (77)	2.8 (73)	2,835	22.9 (114)	2,602	26.4 (58)	1,994	91.7 (58)
4109	569	88.2 (86)	6.2 (119)	628	28.8 (82)	595	23.4 (84)	422	94.8 (17)
4110	2	*	*	2	*	2	*	0	*
4112	526	52.5 (142)	0.0 (1)	546	23.4 (112)	269	56.5 (3)	492	92.5 (50)
4115	2,913	91.5 (49)	3.1 (80)	3,355	21.7 (123)	2,636	18.4 (113)	2,363	92.8 (47)
4351	437	89.5 (75)	2.7 (66)	521	41.5 (9)	385	25.5 (64)	410	90.7 (74)
4352	153	75.2 (139)	7.2 (124)	175	38.3 (19)	169	25.4 (65)	139	84.2 (132)
4353	340	79.4 (133)	11.8 (140)	441	37.4 (24)	420	26.2 (61)	330	85.2 (128)
4354	648	95.1 (13)	2.3 (54)	724	40.3 (14)	669	15.2 (119)	564	91.5 (59)
4355	236	85.2 (110)	8.1 (129)	258	34.1 (43)	244	76.6 (1)	199	91.5 (59)
4356	334	86.2 (103)	5.1 (111)	381	33.3 (48)	363	27.3 (54)	291	92.4 (52)
4358	22	*	*	23	*	23	*	23	*
4359	454	78.9 (134)	10.1 (136)	577	40.6 (13)	547	22.1 (92)	455	89.9 (92)
4360	182	89.6 (72)	6.0 (118)	210	34.3 (42)	207	31.4 (36)	170	91.8 (56)
4361	229	91.7 (47)	2.2 (48)	252	41.3 (10)	243	30.9 (39)	201	91.0 (66)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 97

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4362	245	83.7 (120)	9.4 (134)	271	36.2 (31)	255	29.8 (43)	208	89.4 (99)
4363	597	83.1 (124)	8.7 (131)	665	33.4 (47)	626	28.4 (50)	491	90.8 (70)
4401	656	84.9 (113)	7.3 (125)	742	30.2 (68)	703	26.7 (56)	532	94.7 (18)
4501	708	86.3 (102)	7.3 (125)	768	31.9 (60)	741	29.6 (44)	559	92.5 (50)
4601	1,084	92.9 (34)	1.3 (21)	1,214	28.5 (84)	1,153	20.7 (99)	876	93.0 (43)
4701	866	89.6 (72)	4.5 (107)	974	28.3 (88)	904	38.9 (17)	681	94.4 (24)
5050	1,448	93.9 (23)	0.7 (5)	1,542	21.6 (126)	1,313	9.7 (137)	1,043	96.9 (3)
5051	2,404	93.4 (27)	1.0 (12)	2,609	21.9 (121)	2,172	10.1 (134)	1,777	94.9 (15)
5052	2,508	92.9 (34)	1.6 (27)	2,653	22.8 (116)	2,178	8.3 (139)	1,893	94.2 (26)
5055	1,181	92.9 (34)	6.5 (121)	1,282	44.5 (4)	808	5.6 (143)	1,058	93.0 (43)
5101	1,355	91.9 (45)	2.5 (60)	1,479	24.8 (106)	695	40.7 (13)	1,043	95.2 (14)
5102	15	*	*	17	*	12	*	15	*
5202	571	91.6 (48)	3.9 (100)	662	36.4 (29)	530	32.8 (29)	526	92.6 (48)
5206	12	*	*	14	*	10	*	9	*
5601	864	92.6 (41)	2.0 (41)	941	29.4 (76)	912	25.4 (65)	677	95.3 (12)
5801	1,623	89.6 (72)	3.4 (87)	1,817	28.0 (90)	1,701	24.8 (72)	1,294	92.0 (54)
5803	48	*	*	52	*	51	*	43	*
5804	15	*	*	17	*	16	*	13	*
5806	42	*	*	45	*	40	*	34	*
6101	46	*	*	54	*	54	*	42	*
6102	74	*	*	83	*	74	*	63	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 98

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6103	2,909	89.7 (68)	2.7 (66)	3,255	35.5 (34)	2,764	24.2 (74)	2,462	90.0 (89)
6105	7	*	*	9	*	9	*	5	*
6106	26	*	*	31	*	29	*	31	*
6301	2,971	91.0 (54)	2.8 (73)	3,320	25.0 (105)	2,947	16.5 (115)	2,326	90.1 (86)
6312	1,544	93.4 (27)	1.6 (27)	1,725	24.6 (107)	1,495	23.5 (83)	1,194	91.1 (64)
6316	989	93.6 (26)	1.7 (32)	1,077	24.4 (110)	955	25.3 (70)	747	93.7 (33)
6318	21	*	*	22	*	14	*	7	*
6320	423	93.9 (23)	1.9 (38)	474	19.4 (133)	368	35.3 (23)	320	95.3 (12)
6325	277	96.0 (8)	0.7 (5)	306	22.9 (114)	282	27.7 (53)	197	95.9 (6)
6335	855	94.2 (15)	3.4 (87)	909	32.0 (59)	509	28.9 (49)	678	93.8 (30)
6340	0	*	*	1	*	1	*	1	*
6345	102	94.1 (18)	2.0 (41)	112	16.1 (144)	99	*	69	*
6360	833	91.4 (50)	2.3 (54)	951	26.1 (100)	744	27.2 (55)	703	92.6 (48)
7306	1,847	88.2 (86)	3.1 (80)	2,042	32.1 (58)	1,913	24.9 (71)	1,525	90.8 (70)
7401	1,495	81.0 (131)	10.8 (138)	1,877	41.9 (8)	1,757	25.4 (65)	1,510	87.9 (114)
7601	489	94.9 (14)	2.5 (60)	548	36.3 (30)	526	43.0 (7)	432	94.4 (24)
7850	911	93.2 (32)	4.1 (104)	1,007	31.0 (64)	353	26.3 (59)	728	85.3 (127)
7852	13	*	*	15	*	7	*	12	*
8101	1,566	82.1 (128)	2.0 (41)	1,674	20.1 (128)	1,283	20.1 (105)	1,108	87.0 (121)
8102	433	86.4 (100)	2.5 (60)	475	22.1 (119)	384	19.5 (107)	344	86.3 (123)
8105	191	98.4 (1)	1.0 (12)	234	37.2 (25)	231	18.6 (110)	234	92.3 (53)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 99

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup>									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8106	530	83.8 (118)	1.9 (38)	562	16.0 (145)	511	20.9 (98)	365	93.4 (35)
8150	6	*	*	7	*	7	*	6	*
8151	2,758	92.8 (38)	2.2 (48)	3,094	29.7 (70)	2,939	16.1 (116)	2,290	93.8 (30)
8152	21	*	*	24	*	14	*	23	*
8153	3	*	*	4	*	4	*	4	*
8155	8	*	*	10	*	9	*	10	*
8157	1	*	*	1	*	1	*	1	*
8201	2,425	94.1 (18)	1.1 (16)	2,736	19.7 (130)	1,935	40.9 (12)	1,807	90.3 (85)
8202	3,675	96.7 (6)	1.2 (18)	3,902	29.1 (79)	2,967	5.7 (142)	3,020	93.3 (36)
8203	322	93.2 (32)	2.2 (48)	370	21.9 (121)	292	38.0 (18)	277	94.2 (26)
8250	506	88.9 (83)	1.6 (27)	581	17.2 (143)	437	29.3 (46)	413	88.6 (108)
8252	139	79.9 (132)	5.8 (115)	157	21.7 (123)	146	19.2 (109)	119	93.3 (36)
8255	1,788	93.4 (27)	0.6 (4)	2,025	18.7 (136)	1,501	39.0 (16)	1,353	91.1 (64)
8257	1,683	96.9 (5)	0.3 (3)	1,756	25.7 (101)	1,240	8.1 (140)	1,344	79.4 (135)
8263	432	97.2 (3)	0.9 (8)	526	22.2 (118)	354	41.0 (11)	446	95.7 (8)
8264	701	95.6 (12)	0.9 (8)	775	18.7 (136)	610	37.5 (20)	502	91.8 (56)
8267	571	89.5 (75)	2.6 (65)	622	18.0 (141)	498	41.8 (9)	428	90.7 (74)
8901	808	84.9 (113)	2.4 (58)	857	18.4 (138)	467	24.0 (78)	549	84.2 (132)
8902	1,236	88.1 (88)	1.2 (18)	1,317	19.7 (130)	903	23.6 (81)	911	90.7 (74)
8903	1,300	81.1 (130)	1.9 (38)	1,429	21.7 (123)	1,101	19.3 (108)	999	85.6 (125)
8904	1,964	82.6 (126)	2.2 (48)	2,124	19.0 (135)	1,520	20.7 (99)	1,406	89.3 (102)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 6A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 100

## Michigan

### Comparison of Maternal Behavioral Indicators by Clinic Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8905	1,703	91.3 (51)	1.5 (24)	1,844	23.4 (112)	1,539	33.5 (27)	1,233	89.4 (99)
8909	978	86.4 (100)	1.1 (16)	1,071	19.2 (134)	862	20.3 (103)	727	90.4 (81)
9045	1,263	88.9 (83)	3.4 (87)	1,352	27.1 (94)	657	21.2 (97)	1,013	91.9 (55)
9050	826	83.8 (118)	2.3 (54)	943	25.3 (102)	823	25.4 (65)	682	87.2 (120)
9051	1,403	88.6 (85)	2.1 (45)	1,466	27.8 (92)	1,338	8.0 (141)	1,146	93.6 (34)
9053	1,286	86.2 (103)	1.3 (21)	1,355	15.4 (146)	1,090	16.0 (117)	755	94.2 (26)
9055	7	*	*	6	*	6	*	2	*
9058	1,166	85.0 (111)	2.7 (66)	1,252	18.2 (140)	1,105	9.8 (136)	701	87.3 (119)
9066	894	78.7 (136)	3.5 (91)	976	21.2 (127)	785	13.2 (126)	706	89.7 (94)
9068	782	83.1 (124)	3.5 (91)	835	20.0 (129)	741	14.7 (122)	574	87.8 (115)
9069	1,771	83.2 (122)	2.3 (54)	1,833	17.9 (142)	1,127	10.6 (131)	1,092	90.5 (78)
9070	1,496	83.6 (121)	2.7 (66)	1,617	18.3 (139)	1,330	17.1 (114)	1,023	89.6 (97)
9072	339	78.8 (135)	8.8 (133)	384	30.7 (66)	374	15.0 (120)	385	86.8 (122)
9080	2,310	85.6 (107)	2.5 (60)	2,465	19.6 (132)	1,723	24.1 (77)	1,666	88.1 (112)
9082	103	89.3 (79)	1.0 (12)	110	8.2 (147)	96	*	67	*
9083	7	*	*	6	*	5	*	4	*
9084	4	*	*	5	*	5	*	5	*
9087	13	*	*	15	*	15	*	15	*
<b>Michigan</b>	<b>129,039</b>	<b>89.2</b>	<b>3.1</b>	<b>143,475</b>	<b>27.7</b>	<b>121,256</b>	<b>22.2</b>	<b>103,878</b>	<b>90.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

### 2020 - 2022 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Smoking Indicators by Clinic Combined 3 Years

Page : 101

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0101	86	*	98	*	25	*	81	*	100	4.0 (109)
0102	324	38.0 (126)	370	25.7 (139)	109	21.1 (67)	311	14.1 (123)	375	5.1 (128)
0103	286	35.3 (119)	323	22.0 (129)	87	*	273	19.4 (134)	328	6.1 (141)
0104	104	28.8 (88)	110	18.2 (114)	24	*	99	*	113	4.4 (116)
0105	14	*	14	*	6	*	14	*	14	*
0250	92	*	103	20.4 (123)	28	*	89	*	102	4.9 (126)
0251	61	*	68	*	20	*	59	*	68	*
0252	78	*	87	*	18	*	72	*	87	*
0253	102	46.1 (138)	110	26.4 (140)	40	*	99	*	112	8.9 (144)
0401	346	41.6 (136)	393	22.6 (133)	119	33.6 (52)	331	12.1 (112)	397	2.5 (66)
0402	305	35.1 (117)	331	17.8 (112)	95	*	294	9.5 (82)	335	5.1 (128)
0403	111	42.3 (137)	122	23.8 (136)	39	*	103	15.5 (127)	126	5.6 (135)
0404	103	22.3 (66)	109	16.5 (103)	21	*	94	*	110	3.6 (99)
0501	327	35.8 (122)	398	22.1 (130)	102	31.4 (56)	305	19.3 (133)	401	4.7 (123)
0503	52	*	70	*	12	*	50	*	70	*
0504	290	32.1 (103)	339	15.9 (98)	76	*	256	6.3 (56)	336	2.4 (61)
0505	194	28.9 (90)	213	16.0 (99)	49	*	181	10.5 (96)	213	4.2 (110)
0506	130	41.5 (135)	155	25.2 (138)	44	*	121	14.0 (121)	157	8.9 (144)
0507	47	*	70	*	12	*	43	*	70	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 102

### Michigan

#### Comparison of Smoking Indicators by Clinic

#### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
0508	11	*	19	*	4	*	10	*	19	*
0650	883	26.3 (78)	995	14.7 (93)	205	32.2 (54)	830	7.7 (69)	996	2.9 (76)
0651	474	40.5 (133)	511	27.8 (142)	166	24.7 (63)	455	16.0 (129)	513	3.9 (106)
0652	243	35.4 (120)	292	21.9 (128)	79	*	229	17.5 (132)	293	10.2 (146)
0653	340	35.9 (123)	380	21.8 (127)	104	22.1 (66)	315	16.2 (130)	384	6.5 (142)
0654	340	30.3 (95)	382	20.2 (122)	93	*	322	14.6 (125)	383	4.2 (110)
0655	255	41.2 (134)	277	26.7 (141)	89	*	242	17.4 (131)	286	5.2 (130)
0657	43	*	45	*	14	*	40	*	45	*
0701	388	32.5 (107)	439	20.0 (121)	109	37.6 (44)	369	6.0 (52)	439	3.0 (81)
0702	50	*	63	*	10	*	44	*	63	*
0703	107	38.3 (127)	139	23.7 (135)	34	*	96	*	139	5.8 (138)
0704	43	*	45	*	10	*	42	*	45	*
0705	54	*	65	*	23	*	53	*	65	*
0706	3	*	3	*	1	*	3	*	3	*
0850	546	33.7 (113)	651	14.0 (89)	155	47.1 (18)	497	7.4 (67)	651	2.5 (66)
0851	464	32.5 (107)	562	15.3 (96)	125	48.8 (15)	413	10.7 (100)	564	2.7 (73)
0901	1,229	34.7 (116)	1,384	16.8 (105)	383	35.2 (48)	1,167	10.2 (92)	1,396	4.2 (110)
0902	29	*	28	*	9	*	25	*	28	*
1051	873	35.4 (120)	1,041	14.6 (92)	256	45.3 (21)	776	6.8 (62)	1,056	3.9 (106)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 103

### Michigan

#### Comparison of Smoking Indicators by Clinic

#### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1052	2	*	3	*	1	*	2	*	3	*
1101	1,125	21.7 (63)	1,315	9.5 (59)	198	45.5 (20)	1,040	11.0 (104)	1,317	4.4 (116)
1102	620	34.5 (114)	679	17.1 (107)	163	40.5 (37)	568	12.9 (117)	685	4.5 (120)
1106	50	*	55	*	9	*	44	*	55	*
1250	767	23.2 (69)	892	13.5 (83)	144	27.8 (60)	703	8.1 (74)	897	1.9 (44)
1251	467	35.1 (117)	576	24.1 (137)	140	30.0 (59)	439	9.6 (83)	576	3.6 (99)
1252	414	22.7 (67)	464	11.6 (75)	79	*	393	7.1 (64)	466	3.4 (95)
1253	557	31.1 (99)	619	15.2 (95)	149	36.9 (46)	528	8.5 (79)	617	3.2 (87)
1350	1,841	25.2 (75)	2,165	10.0 (65)	380	37.4 (45)	1,716	4.3 (30)	2,195	2.0 (49)
1351	27	*	22	*	9	*	27	*	23	*
1352	309	20.7 (55)	362	7.7 (50)	60	*	285	2.5 (11)	367	0.3 (2)
1401	154	37.0 (125)	182	18.7 (116)	52	*	143	11.9 (111)	184	6.0 (140)
1402	78	*	90	*	18	*	70	*	91	*
1403	16	*	20	*	5	*	14	*	20	*
1501	577	20.1 (53)	655	6.6 (44)	101	42.6 (31)	542	3.3 (15)	669	1.9 (44)
1503	657	21.5 (61)	776	9.9 (64)	134	32.1 (55)	619	3.6 (19)	777	0.9 (10)
1504	1,434	14.8 (31)	1,727	6.1 (38)	175	48.0 (16)	1,311	2.3 (9)	1,730	0.9 (10)
1505	130	23.1 (68)	115	9.6 (61)	26	*	125	4.0 (25)	115	5.2 (130)
1506	441	20.9 (57)	497	10.1 (66)	75	*	415	4.6 (35)	506	2.0 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 104

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
1507	385	33.5 (112)	474	10.1 (66)	106	43.4 (27)	343	10.2 (92)	482	3.5 (98)
1508	6	*	4	*	0	*	6	*	4	*
1509	307	18.9 (50)	338	12.7 (79)	52	*	291	4.5 (33)	338	2.4 (61)
1511	477	32.1 (103)	566	15.4 (97)	135	42.2 (34)	453	7.3 (66)	576	4.3 (114)
1512	241	17.4 (45)	342	9.4 (57)	33	*	209	5.3 (42)	342	2.3 (57)
1514	391	12.3 (21)	553	6.3 (41)	36	*	306	2.0 (7)	548	1.3 (20)
1701	453	30.7 (97)	495	21.6 (125)	122	15.6 (68)	423	10.6 (98)	495	3.2 (87)
1950	478	39.1 (129)	552	19.0 (117)	161	38.5 (41)	445	15.7 (128)	567	4.8 (125)
1952	101	23.8 (70)	133	13.5 (83)	20	*	94	*	134	3.0 (81)
1953	195	33.3 (110)	222	13.5 (83)	54	*	183	7.7 (69)	225	1.8 (43)
1954	328	31.1 (99)	359	16.2 (101)	96	*	318	10.7 (100)	366	2.5 (66)
1955	87	*	97	*	31	*	80	*	103	3.9 (106)
1957	282	32.3 (105)	352	14.8 (94)	87	*	273	3.7 (22)	357	1.4 (27)
1963	42	*	54	*	5	*	39	*	55	*
1964	1	*	1	*	0	*	1	*	1	*
2150	465	38.9 (128)	533	23.1 (134)	149	30.2 (58)	430	10.5 (96)	532	3.0 (81)
2151	147	33.3 (110)	173	22.5 (132)	38	*	129	10.9 (103)	173	2.3 (57)
2152	25	*	30	*	11	*	23	*	30	*
2251	330	30.9 (98)	352	17.9 (113)	87	*	304	3.3 (15)	355	1.4 (27)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 105

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
2252	138	47.8 (139)	148	31.1 (143)	56	*	134	14.2 (124)	148	4.7 (123)
2554	41	*	46	*	14	*	38	*	46	*
2555	5,508	21.4 (59)	6,539	10.7 (71)	984	40.4 (38)	5,091	8.0 (73)	6,582	3.0 (81)
2556	150	27.3 (84)	174	14.4 (91)	39	*	142	5.6 (46)	177	1.1 (13)
2557	1	*	2	*	0	*	1	*	2	*
2558	1	*	0	*	0	*	1	*	0	*
2560	3	*	2	*	0	*	3	*	2	*
2561	1	*	1	*	1	*	1	*	1	*
2562	6	*	6	*	1	*	6	*	6	*
2563	9	*	10	*	2	*	8	*	10	*
2564	16	*	12	*	4	*	14	*	12	*
3001	970	27.3 (84)	1,116	10.5 (69)	248	50.0 (14)	934	10.6 (98)	1,125	4.2 (110)
3002	160	26.3 (78)	184	13.0 (81)	38	*	156	13.5 (119)	182	3.3 (94)
3004	201	20.4 (54)	243	7.0 (49)	34	*	181	13.8 (120)	244	4.5 (120)
3005	135	29.6 (91)	146	10.3 (68)	39	*	129	11.6 (109)	146	5.5 (134)
3007	6	*	8	*	2	*	6	*	8	*
3201	301	34.6 (115)	387	19.6 (120)	93	*	265	10.2 (92)	388	5.2 (130)
3203	1	*	1	*	0	*	1	*	1	*
3301	3,354	17.5 (46)	4,065	8.3 (53)	519	43.5 (25)	3,101	4.1 (28)	4,069	2.4 (61)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 106

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
3304	85	*	96	*	8	*	77	*	96	*
3305	117	14.5 (28)	150	6.7 (47)	13	*	104	1.0 (4)	150	1.3 (20)
3306	178	20.8 (56)	225	9.8 (63)	35	*	168	3.6 (19)	224	1.3 (20)
3308	81	*	104	9.6 (61)	12	*	62	*	104	0.0 (1)
3312	341	21.4 (59)	401	10.7 (71)	66	*	324	3.7 (22)	401	3.2 (87)
3550	64	*	76	*	36	*	63	*	76	*
3551	9	*	8	*	6	*	9	*	8	*
3552	14	*	12	*	9	*	14	*	12	*
3701	669	28.8 (88)	801	17.5 (109)	163	27.0 (61)	600	8.2 (75)	812	2.5 (66)
3850	2,289	29.8 (94)	2,643	17.7 (110)	610	30.3 (57)	2,168	7.6 (68)	2,643	2.4 (61)
3901	1,219	22.2 (65)	1,353	9.2 (56)	223	37.7 (43)	1,146	5.3 (42)	1,353	1.7 (41)
3902	40	*	30	*	8	*	41	*	31	*
3903	6	*	7	*	0	*	5	*	7	*
3950	1,541	21.6 (62)	1,830	10.7 (71)	279	38.4 (42)	1,414	5.0 (39)	1,836	2.3 (57)
3954	6	*	6	*	2	*	6	*	6	*
4106	1,671	13.0 (23)	2,014	6.4 (43)	177	40.7 (36)	1,573	2.6 (13)	2,013	1.1 (13)
4107	2,217	17.8 (48)	2,617	8.3 (53)	307	47.2 (17)	1,988	5.7 (49)	2,623	1.4 (27)
4109	463	24.0 (71)	602	12.8 (80)	93	*	421	4.0 (25)	601	2.0 (49)
4110	2	*	2	*	0	*	0	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 107

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
4112	503	3.2 (4)	528	1.1 (4)	14	*	402	8.5 (79)	507	3.2 (87)
4115	2,603	9.3 (9)	3,149	3.7 (17)	216	43.5 (25)	2,342	2.2 (8)	3,153	0.8 (8)
4351	432	29.6 (91)	485	17.3 (108)	111	27.0 (61)	410	6.8 (62)	490	3.7 (103)
4352	143	51.0 (140)	164	32.9 (144)	72	*	139	21.6 (135)	166	7.8 (143)
4353	355	24.5 (73)	405	11.9 (76)	70	*	329	6.7 (61)	405	2.0 (49)
4354	588	31.3 (101)	656	17.7 (110)	164	23.2 (65)	565	11.3 (106)	688	2.5 (66)
4355	216	32.4 (106)	245	21.6 (125)	64	*	199	9.0 (81)	247	4.5 (120)
4356	309	35.9 (123)	360	19.4 (118)	97	*	291	12.4 (113)	359	5.6 (135)
4358	23	*	22	*	1	*	23	*	22	*
4359	491	17.1 (42)	546	9.0 (55)	79	*	456	4.8 (38)	547	1.3 (20)
4360	178	24.7 (74)	197	18.3 (115)	36	*	170	10.0 (89)	197	3.6 (99)
4361	215	28.4 (87)	234	20.5 (124)	56	*	200	12.5 (114)	236	3.4 (95)
4362	220	40.0 (131)	255	22.4 (131)	72	*	207	14.0 (121)	260	5.8 (138)
4363	529	25.9 (76)	648	16.7 (104)	117	23.9 (64)	490	11.4 (107)	647	5.4 (133)
4401	584	29.6 (91)	703	16.2 (101)	139	43.2 (29)	532	6.0 (52)	703	1.3 (20)
4501	607	32.9 (109)	746	16.8 (105)	179	41.3 (35)	559	11.8 (110)	748	3.2 (87)
4601	953	28.1 (86)	1,109	12.4 (77)	214	44.4 (22)	876	9.8 (87)	1,108	4.3 (114)
4701	766	26.5 (80)	905	13.0 (81)	167	40.1 (39)	683	4.7 (37)	916	1.9 (44)
5050	1,171	16.7 (40)	1,479	6.8 (48)	166	39.8 (40)	1,042	6.1 (55)	1,478	1.2 (19)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 108

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
5051	2,010	13.6 (27)	2,467	5.8 (34)	217	42.4 (32)	1,773	3.9 (24)	2,461	1.7 (41)
5052	2,059	13.5 (26)	2,579	5.2 (33)	228	44.3 (23)	1,887	4.5 (33)	2,578	1.4 (27)
5055	1,125	2.0 (2)	1,260	0.9 (2)	18	*	1,058	3.2 (14)	1,260	0.5 (3)
5101	1,159	14.8 (31)	1,402	6.3 (41)	152	52.6 (12)	1,038	4.1 (28)	1,403	1.5 (35)
5102	17	*	15	*	2	*	15	*	15	*
5202	563	40.0 (131)	603	19.4 (118)	210	35.2 (48)	523	8.2 (75)	632	1.4 (27)
5206	12	*	16	*	3	*	9	*	16	*
5601	736	27.0 (83)	890	13.5 (83)	172	36.6 (47)	677	5.0 (39)	897	1.3 (20)
5801	1,421	30.5 (96)	1,684	13.7 (88)	361	42.9 (30)	1,293	10.2 (92)	1,689	4.4 (116)
5803	49	*	48	*	9	*	43	*	48	*
5804	16	*	16	*	4	*	13	*	16	*
5806	37	*	43	*	14	*	34	*	43	*
6101	45	*	51	*	12	*	42	*	52	*
6102	65	*	81	*	17	*	63	*	82	*
6103	2,618	26.6 (81)	3,077	14.3 (90)	601	33.9 (51)	2,468	8.4 (77)	3,108	3.2 (87)
6105	7	*	8	*	0	*	5	*	8	*
6106	31	*	25	*	8	*	31	*	26	*
6301	2,555	9.4 (10)	3,064	5.0 (31)	209	34.4 (50)	2,321	3.6 (19)	3,071	1.1 (13)
6312	1,342	4.2 (5)	1,598	1.3 (5)	42	*	1,192	2.3 (9)	1,598	0.8 (8)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 109

### Michigan

#### Comparison of Smoking Indicators by Clinic

#### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
6316	847	10.3 (15)	1,009	5.1 (32)	73	*	747	2.5 (11)	1,010	0.6 (5)
6318	14	*	21	*	1	*	7	*	21	*
6320	367	10.6 (16)	438	2.5 (8)	29	*	320	0.6 (1)	437	1.1 (13)
6325	236	26.7 (82)	275	13.5 (83)	48	*	196	4.6 (35)	279	2.2 (54)
6335	750	8.1 (8)	884	3.5 (15)	49	*	679	1.9 (6)	886	1.1 (13)
6340	1	*	0	*	0	*	1	*	0	*
6345	81	*	103	2.9 (11)	14	*	69	*	105	2.9 (76)
6360	771	16.2 (37)	880	6.6 (44)	104	44.2 (24)	698	6.0 (52)	886	1.6 (38)
7306	1,619	31.8 (102)	1,896	11.0 (74)	453	52.8 (11)	1,523	10.1 (91)	1,939	3.1 (85)
7401	1,592	39.8 (130)	1,698	16.1 (100)	533	43.3 (28)	1,510	13.2 (118)	1,715	5.0 (127)
7601	459	15.7 (35)	494	10.5 (69)	58	*	430	5.8 (50)	496	1.6 (38)
7850	788	26.0 (77)	934	12.4 (77)	167	32.3 (53)	722	7.1 (64)	927	3.6 (99)
7852	12	*	13	*	3	*	12	*	12	*
8101	1,199	15.4 (34)	1,625	4.1 (21)	160	60.6 (4)	1,105	9.6 (83)	1,625	2.8 (74)
8102	369	12.2 (20)	451	4.0 (20)	38	*	340	5.6 (46)	453	2.9 (76)
8105	234	9.8 (12)	190	3.2 (12)	18	*	233	0.9 (3)	189	0.5 (3)
8106	409	21.0 (58)	543	8.1 (52)	68	*	365	5.5 (44)	543	1.3 (20)
8150	6	*	7	*	2	*	6	*	7	*
8151	2,494	16.2 (37)	2,850	6.0 (37)	328	51.5 (13)	2,287	4.3 (30)	2,857	1.1 (13)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 110

## Michigan

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
8152	23	*	22	*	7	*	23	*	22	*
8153	4	*	3	*	1	*	4	*	3	*
8155	10	*	8	*	2	*	10	*	8	*
8157	1	*	1	*	0	*	1	*	1	*
8201	2,093	14.7 (30)	2,473	6.1 (38)	245	55.5 (8)	1,801	4.0 (25)	2,479	1.5 (35)
8202	3,262	3.1 (3)	3,855	1.0 (3)	87	*	3,018	3.4 (17)	3,855	0.7 (7)
8203	295	19.3 (51)	327	9.5 (59)	39	*	277	6.5 (58)	329	1.5 (35)
8250	454	17.6 (47)	505	4.6 (26)	53	*	413	6.5 (58)	508	1.4 (27)
8252	129	17.1 (42)	145	4.8 (29)	18	*	119	8.4 (77)	144	2.1 (53)
8255	1,520	19.5 (52)	1,820	9.4 (57)	224	42.4 (32)	1,358	6.4 (57)	1,822	1.9 (44)
8257	1,489	0.5 (1)	1,725	0.5 (1)	6	*	1,347	0.7 (2)	1,717	1.0 (12)
8263	498	21.9 (64)	443	7.7 (50)	75	*	444	4.3 (30)	442	1.4 (27)
8264	577	12.0 (19)	708	3.5 (15)	56	*	501	3.4 (17)	706	1.4 (27)
8267	472	15.3 (33)	578	6.6 (44)	53	*	428	5.1 (41)	581	2.9 (76)
8901	602	13.0 (23)	829	6.2 (40)	70	*	551	9.6 (83)	826	3.4 (95)
8902	988	11.9 (18)	1,261	3.3 (13)	103	63.1 (2)	912	5.9 (51)	1,259	2.8 (74)
8903	1,079	15.8 (36)	1,353	4.7 (28)	133	55.6 (7)	995	10.7 (100)	1,354	2.2 (54)
8904	1,563	16.2 (37)	2,035	4.3 (22)	204	63.7 (1)	1,406	9.7 (86)	2,036	3.8 (104)
8905	1,348	10.2 (13)	1,780	3.3 (13)	122	60.7 (3)	1,222	5.5 (44)	1,780	2.6 (72)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

Table 7A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 111

Michigan

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8909	798	12.7 (22)	1,013	3.9 (19)	87	*	727	10.0 (89)	1,010	3.2 (87)
9045	1,087	7.3 (7)	1,298	2.4 (6)	67	*	1,008	5.6 (46)	1,294	1.6 (38)
9050	718	16.7 (40)	897	4.8 (29)	119	55.5 (8)	684	12.7 (115)	895	2.3 (57)
9051	1,202	7.2 (6)	1,441	2.5 (8)	78	*	1,147	1.7 (5)	1,440	0.6 (5)
9053	895	10.2 (13)	1,313	2.4 (6)	79	*	754	7.7 (69)	1,311	2.2 (54)
9055	3	*	7	*	0	*	2	*	7	*
9058	774	9.7 (11)	1,209	4.3 (22)	65	*	694	6.6 (60)	1,208	1.9 (44)
9066	809	17.8 (48)	946	5.8 (34)	115	53.9 (10)	706	12.7 (115)	944	3.1 (85)
9068	618	17.2 (44)	807	5.9 (36)	89	*	575	11.0 (104)	808	2.5 (66)
9069	1,291	10.9 (17)	1,787	4.6 (26)	112	46.4 (19)	1,088	11.4 (107)	1,785	4.4 (116)
9070	1,124	13.3 (25)	1,541	3.8 (18)	127	60.6 (4)	1,013	9.9 (88)	1,536	2.4 (61)
9072	385	24.2 (72)	341	4.4 (24)	82	*	386	14.8 (126)	341	2.9 (76)
9080	1,789	14.6 (29)	2,397	4.5 (25)	231	58.4 (6)	1,665	7.7 (69)	2,400	3.8 (104)
9082	82	*	106	2.8 (10)	6	*	68	*	106	5.7 (137)
9083	5	*	7	*	0	*	4	*	7	*
9084	5	*	4	*	1	*	5	*	4	*
9087	15	*	13	*	1	*	15	*	13	*
Michigan	112,918	20.2	134,825	9.3	19,273	41.2	103,605	7.0	135,290	2.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 112

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0101	93	*	*	96	*	82	*	93	*
0102	343	11.7 (99)	9.3 (124)	350	11.4 (55)	297	6.7 (119)	342	70.5 (88)
0103	304	7.6 (27)	7.2 (82)	315	8.6 (6)	278	5.8 (108)	305	71.5 (85)
0104	99	*	*	107	14.0 (108)	86	*	99	*
0105	12	*	*	13	*	11	*	12	*
0250	95	*	*	96	*	82	*	94	*
0251	64	*	*	66	*	57	*	64	*
0252	81	*	*	85	*	75	*	81	*
0253	105	7.6 (27)	7.6 (96)	111	16.2 (129)	89	*	105	71.4 (86)
0401	379	7.4 (25)	7.9 (100)	387	10.1 (27)	334	3.6 (42)	379	74.4 (76)
0402	324	8.6 (44)	8.0 (103)	324	11.1 (49)	282	4.3 (67)	324	81.5 (27)
0403	117	8.5 (42)	10.3 (133)	119	10.1 (27)	103	4.9 (81)	119	81.5 (27)
0404	106	9.4 (61)	11.3 (139)	106	15.1 (123)	89	*	106	77.4 (57)
0501	344	9.6 (67)	8.7 (114)	389	8.7 (10)	307	5.2 (90)	381	77.7 (54)
0503	67	*	*	68	*	62	*	69	*
0504	315	9.5 (64)	6.7 (68)	326	11.0 (46)	276	2.9 (18)	321	90.7 (1)
0505	179	11.7 (99)	7.3 (87)	204	12.3 (74)	157	5.7 (104)	199	83.9 (19)
0506	135	9.6 (67)	6.7 (68)	144	7.6 (2)	121	5.0 (87)	142	76.8 (62)
0507	56	*	*	67	*	47	*	63	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 113

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
0508	18	*	*	19	*	16	*	19	*
0650	944	9.5 (64)	9.0 (118)	978	10.4 (33)	847	5.4 (97)	945	76.9 (61)
0651	474	7.2 (21)	7.8 (98)	503	9.9 (24)	428	4.2 (65)	473	72.3 (82)
0652	276	8.0 (32)	7.2 (82)	280	8.6 (6)	248	3.2 (27)	274	76.6 (63)
0653	345	8.4 (41)	7.5 (92)	356	11.0 (46)	306	3.3 (29)	346	80.6 (32)
0654	356	11.0 (89)	7.0 (76)	364	11.5 (59)	311	6.4 (114)	357	77.3 (58)
0655	263	6.5 (9)	7.2 (82)	269	8.6 (6)	235	3.4 (33)	263	75.3 (70)
0657	41	*	*	44	*	32	*	41	*
0701	410	6.8 (13)	9.5 (130)	432	9.3 (17)	370	3.5 (39)	412	88.6 (4)
0702	60	*	*	60	*	58	*	60	*
0703	125	8.0 (32)	9.6 (131)	133	9.8 (23)	113	2.7 (12)	127	74.8 (74)
0704	43	*	*	44	*	33	*	43	*
0705	62	*	*	64	*	52	*	62	*
0706	3	*	*	3	*	3	*	3	*
0850	618	11.8 (104)	7.1 (78)	642	13.1 (87)	535	4.9 (81)	624	85.4 (13)
0851	522	9.6 (67)	6.7 (68)	552	12.7 (79)	456	3.7 (48)	528	83.3 (21)
0901	1,289	10.9 (87)	8.5 (110)	1,354	12.6 (78)	1,127	4.3 (67)	1,297	70.1 (90)
0902	28	*	*	28	*	25	*	28	*
1051	976	8.9 (49)	9.4 (126)	1,011	11.2 (50)	860	3.8 (53)	983	87.1 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 114

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1052	3	*	*	2	*	1	*	3	*
1101	1,243	11.7 (99)	5.1 (29)	1,273	12.1 (71)	1,075	6.6 (118)	1,244	69.5 (95)
1102	639	6.1 (6)	6.6 (66)	664	8.1 (3)	575	3.5 (39)	641	75.5 (69)
1106	53	*	*	55	*	45	*	53	*
1250	851	8.7 (46)	7.9 (100)	878	9.7 (22)	754	5.3 (94)	857	76.5 (65)
1251	540	11.1 (91)	6.3 (55)	557	13.8 (103)	459	4.8 (77)	540	79.1 (45)
1252	438	7.1 (18)	10.7 (136)	449	9.6 (21)	395	2.8 (15)	441	83.7 (20)
1253	580	5.7 (5)	9.1 (122)	603	11.3 (52)	508	1.8 (5)	582	86.1 (8)
1350	2,081	9.8 (76)	6.5 (62)	2,132	10.7 (36)	1,839	4.9 (81)	2,092	76.1 (67)
1351	21	*	*	20	*	19	*	21	*
1352	343	9.3 (56)	9.3 (124)	356	9.3 (17)	308	3.9 (56)	350	66.6 (104)
1401	169	8.3 (39)	8.3 (109)	175	9.1 (14)	153	5.2 (90)	169	85.2 (15)
1402	84	*	*	88	*	73	*	85	*
1403	19	*	*	20	*	16	*	19	*
1501	601	9.7 (71)	8.0 (103)	646	10.7 (36)	532	4.3 (67)	628	75.2 (72)
1503	724	6.6 (12)	8.1 (106)	742	11.5 (59)	639	3.1 (23)	727	77.7 (54)
1504	1,403	9.3 (56)	7.1 (78)	1,691	13.4 (93)	1,198	3.6 (42)	1,657	80.5 (33)
1505	106	3.8 (1)	11.3 (139)	110	14.5 (116)	92	*	108	78.7 (47)
1506	473	8.5 (42)	6.8 (72)	490	13.3 (91)	409	3.2 (27)	477	73.6 (79)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 115

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
1507	433	9.7 (71)	10.6 (134)	463	11.9 (66)	381	3.7 (48)	445	85.2 (15)
1508	4	*	*	4	*	3	*	4	*
1509	305	7.9 (31)	6.2 (52)	323	13.6 (98)	263	1.9 (6)	307	69.1 (97)
1511	539	9.3 (56)	6.5 (62)	558	12.7 (79)	468	3.4 (33)	550	80.2 (35)
1512	327	10.7 (83)	6.7 (68)	335	15.5 (125)	276	3.6 (42)	327	78.9 (46)
1514	505	14.1 (121)	7.1 (78)	540	17.2 (134)	415	3.4 (33)	521	86.9 (7)
1701	457	7.2 (21)	10.9 (138)	473	10.1 (27)	403	2.7 (12)	457	75.3 (70)
1950	534	11.2 (94)	6.0 (48)	549	14.0 (108)	450	3.1 (23)	536	82.5 (24)
1952	125	6.4 (7)	4.8 (26)	130	11.5 (59)	110	0.9 (2)	125	90.4 (2)
1953	214	12.1 (109)	6.1 (49)	221	12.7 (79)	185	4.9 (81)	217	88.9 (3)
1954	346	8.1 (35)	5.2 (31)	353	10.2 (31)	308	1.6 (3)	345	81.4 (29)
1955	97	*	*	99	*	83	*	97	*
1957	341	16.7 (138)	9.4 (126)	347	17.6 (137)	279	5.7 (104)	344	85.5 (11)
1963	48	*	*	48	*	42	*	48	*
1964	1	*	*	1	*	1	*	1	*
2150	489	5.1 (4)	8.6 (112)	516	9.1 (14)	441	1.6 (3)	490	80.2 (35)
2151	166	7.2 (21)	9.0 (118)	170	6.5 (1)	156	4.5 (73)	166	59.6 (121)
2152	28	*	*	29	*	23	*	28	*
2251	345	6.4 (7)	5.8 (45)	350	11.4 (55)	305	2.3 (7)	346	77.5 (56)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance <sup>(1)</sup>

Page : 116

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
2252	138	8.7 (46)	8.7 (114)	141	11.3 (52)	123	2.4 (9)	136	80.1 (37)
2554	45	*	*	46	*	40	*	45	*
2555	6,294	11.9 (107)	5.4 (35)	6,470	13.1 (87)	5,431	5.7 (104)	6,304	61.8 (115)
2556	168	15.5 (132)	4.2 (20)	175	20.6 (143)	133	6.0 (112)	170	47.6 (137)
2557	2	*	*	2	*	1	*	2	*
2558	0	*	*	0	*	0	*	0	*
2560	2	*	*	2	*	2	*	2	*
2561	1	*	*	1	*	1	*	1	*
2562	6	*	*	6	*	4	*	6	*
2563	10	*	*	10	*	10	*	10	*
2564	12	*	*	12	*	12	*	12	*
3001	1,055	15.2 (129)	3.8 (12)	1,089	14.5 (116)	903	6.5 (117)	1,051	59.5 (122)
3002	175	9.1 (52)	5.1 (29)	177	13.6 (98)	152	3.9 (56)	175	59.4 (123)
3004	217	11.1 (91)	5.5 (39)	229	17.5 (136)	176	4.0 (59)	218	69.3 (96)
3005	139	5.0 (3)	10.8 (137)	141	8.5 (5)	128	2.3 (7)	141	75.9 (68)
3007	8	*	*	8	*	6	*	8	*
3201	369	7.3 (24)	10.6 (134)	382	10.7 (36)	331	3.0 (20)	368	79.3 (41)
3203	0	*	*	0	*	0	*	0	*
3301	3,794	11.8 (104)	5.7 (41)	3,951	13.9 (106)	3,262	4.6 (76)	3,822	78.4 (51)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 117

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
3304	92	*	*	94	*	82	*	92	*
3305	139	9.4 (61)	9.4 (126)	148	13.5 (95)	120	3.3 (29)	141	85.8 (10)
3306	217	6.9 (14)	6.9 (73)	222	9.9 (24)	195	2.6 (10)	216	79.2 (43)
3308	97	*	*	100	11.0 (46)	86	*	96	*
3312	381	9.7 (71)	6.3 (55)	392	11.7 (64)	336	4.5 (73)	383	78.3 (52)
3550	66	*	*	69	*	54	*	66	*
3551	6	*	*	8	*	6	*	7	*
3552	11	*	*	12	*	10	*	11	*
3701	769	7.4 (25)	8.5 (110)	792	10.7 (36)	681	2.8 (15)	770	86.0 (9)
3850	2,487	10.5 (80)	8.0 (103)	2,563	12.2 (72)	2,189	4.8 (77)	2,493	73.9 (78)
3901	1,261	10.7 (83)	4.1 (18)	1,314	12.2 (72)	1,093	4.8 (77)	1,274	69.9 (91)
3902	27	*	*	30	*	23	*	27	*
3903	7	*	*	7	*	6	*	7	*
3950	1,696	10.8 (85)	8.2 (108)	1,788	12.0 (67)	1,485	4.0 (59)	1,715	75.1 (73)
3954	6	*	*	6	*	6	*	6	*
4106	1,880	13.4 (117)	4.7 (25)	1,962	14.7 (119)	1,602	6.4 (114)	1,897	76.6 (63)
4107	2,448	13.6 (118)	5.7 (41)	2,576	16.1 (128)	2,053	5.8 (108)	2,487	77.8 (53)
4109	565	11.3 (97)	6.4 (60)	581	17.0 (132)	468	3.4 (33)	563	82.4 (25)
4110	2	*	*	2	*	0	*	2	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

### 2020 - 2022 Pregnancy Nutrition Surveillance (1) Michigan Comparison of Infant Health Indicators by Clinic Combined 3 Years

Page : 118

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
4112	496	6.5 (9)	4.6 (23)	499	12.8 (82)	425	3.3 (29)	497	28.4 (141)
4115	2,959	13.0 (115)	6.1 (49)	3,106	14.3 (113)	2,534	5.3 (94)	3,011	80.1 (37)
4351	454	11.2 (94)	7.7 (97)	474	14.3 (113)	390	5.9 (110)	456	85.5 (11)
4352	152	12.5 (111)	7.9 (100)	160	13.1 (87)	127	5.5 (100)	152	77.0 (59)
4353	382	9.4 (61)	7.3 (87)	381	12.9 (84)	326	4.3 (67)	386	78.5 (49)
4354	652	9.0 (50)	7.2 (82)	669	10.5 (35)	581	3.6 (42)	654	79.2 (43)
4355	217	11.1 (91)	9.2 (123)	232	14.7 (119)	184	4.3 (67)	219	74.0 (77)
4356	334	9.0 (50)	6.9 (73)	341	12.0 (67)	290	3.8 (53)	334	76.3 (66)
4358	20	*	*	21	*	20	*	20	*
4359	501	8.6 (44)	7.0 (76)	509	10.8 (40)	438	3.9 (56)	502	79.3 (41)
4360	168	7.1 (18)	6.5 (62)	185	8.6 (6)	149	4.0 (59)	168	87.5 (5)
4361	195	10.8 (85)	8.7 (114)	225	13.8 (103)	169	3.6 (42)	196	84.2 (18)
4362	247	9.7 (71)	8.9 (117)	250	10.0 (26)	218	2.8 (15)	246	84.6 (17)
4363	607	9.2 (54)	9.7 (132)	615	12.8 (82)	519	2.7 (12)	608	85.4 (13)
4401	666	9.2 (54)	8.6 (112)	679	11.6 (63)	582	5.2 (90)	668	80.4 (34)
4501	705	6.5 (9)	8.1 (106)	729	11.8 (65)	625	2.6 (10)	705	78.6 (48)
4601	1,054	6.9 (14)	7.3 (87)	1,095	8.9 (12)	959	2.9 (18)	1,053	73.6 (79)
4701	874	11.0 (89)	6.2 (52)	896	13.5 (95)	755	4.1 (63)	872	83.3 (21)
5050	1,431	11.9 (107)	4.8 (26)	1,442	15.7 (126)	1,193	5.2 (90)	1,430	65.4 (106)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 119

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low		High	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
5051	2,342	11.7 (99)	5.3 (32)	2,393	14.0 (108)	1,984	4.5 (73)	2,350	68.6 (100)
5052	2,458	14.1 (121)	4.6 (23)	2,503	14.8 (121)	2,069	6.9 (121)	2,468	65.4 (106)
5055	1,150	9.5 (64)	5.3 (32)	1,182	8.2 (4)	1,056	5.6 (103)	1,165	62.5 (113)
5101	1,355	8.7 (46)	5.7 (41)	1,365	11.3 (52)	1,193	3.0 (20)	1,362	74.7 (75)
5102	15	*	*	15	*	13	*	15	*
5202	603	7.1 (18)	7.5 (92)	606	9.1 (14)	542	3.1 (23)	605	83.0 (23)
5206	16	*	*	14	*	13	*	16	*
5601	855	8.2 (37)	9.0 (118)	883	10.2 (31)	767	3.1 (23)	857	78.5 (49)
5801	1,607	8.0 (32)	7.5 (92)	1,645	11.2 (50)	1,423	3.0 (20)	1,607	68.8 (99)
5803	47	*	*	47	*	42	*	47	*
5804	15	*	*	16	*	14	*	15	*
5806	40	*	*	41	*	36	*	40	*
6101	48	*	*	50	*	41	*	49	*
6102	81	*	*	82	*	71	*	81	*
6103	2,928	11.6 (98)	6.3 (55)	2,991	13.0 (86)	2,508	5.4 (97)	2,936	66.7 (103)
6105	8	*	*	8	*	7	*	7	*
6106	25	*	*	26	*	22	*	25	*
6301	2,859	12.9 (114)	6.5 (62)	3,005	13.2 (90)	2,461	5.9 (110)	2,904	72.0 (84)
6312	1,481	11.2 (94)	5.6 (40)	1,560	13.6 (98)	1,276	3.7 (48)	1,508	79.6 (40)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 120

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
6316	927	9.7 (71)	5.4 (35)	983	12.5 (77)	809	3.7 (48)	943	80.0 (39)
6318	18	*	*	20	*	17	*	19	*
6320	415	10.6 (82)	6.3 (55)	425	10.8 (40)	367	6.3 (113)	419	81.9 (26)
6325	275	9.1 (52)	12.0 (141)	276	10.1 (27)	244	3.3 (29)	274	77.0 (59)
6335	845	10.5 (80)	5.9 (46)	854	12.4 (76)	732	4.9 (81)	852	73.4 (81)
6340	0	*	*	0	*	0	*	0	*
6345	100	7.0 (17)	9.0 (118)	105	11.4 (55)	88	*	101	81.2 (30)
6360	839	14.3 (123)	5.4 (35)	870	14.0 (108)	712	8.1 (128)	848	71.1 (87)
7306	1,831	12.5 (111)	6.2 (52)	1,894	13.6 (98)	1,582	5.0 (87)	1,837	68.4 (101)
7401	1,614	9.3 (56)	7.1 (78)	1,655	12.3 (74)	1,414	3.6 (42)	1,614	69.1 (97)
7601	466	7.7 (29)	6.4 (60)	489	10.8 (40)	409	4.2 (65)	478	60.7 (118)
7850	884	8.1 (35)	7.5 (92)	921	10.9 (45)	785	3.7 (48)	887	69.7 (93)
7852	11	*	*	11	*	10	*	11	*
8101	1,509	15.9 (135)	2.5 (2)	1,547	18.4 (141)	1,199	8.5 (132)	1,516	49.7 (136)
8102	419	6.9 (14)	7.2 (82)	442	10.4 (33)	370	3.8 (53)	427	60.4 (119)
8105	148	4.1 (2)	4.1 (18)	189	9.0 (13)	138	0.7 (1)	181	54.1 (132)
8106	499	14.0 (119)	5.4 (35)	531	16.0 (127)	416	4.8 (77)	518	62.2 (114)
8150	6	*	*	7	*	6	*	7	*
8151	2,664	12.2 (110)	5.7 (41)	2,772	13.8 (103)	2,280	5.3 (94)	2,701	81.2 (30)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 121

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8152	20	*	*	22	*	20	*	20	*
8153	3	*	*	2	*	2	*	3	*
8155	7	*	*	8	*	6	*	7	*
8157	1	*	*	1	*	0	*	1	*
8201	2,328	11.8 (104)	5.9 (46)	2,440	13.4 (93)	2,016	5.4 (97)	2,354	65.4 (106)
8202	3,464	8.2 (37)	4.5 (22)	3,685	9.3 (17)	3,113	4.0 (59)	3,636	59.1 (124)
8203	310	10.3 (79)	9.4 (126)	320	14.4 (115)	262	3.4 (33)	314	70.4 (89)
8250	484	12.8 (113)	5.0 (28)	498	13.5 (95)	415	5.5 (100)	488	57.6 (126)
8252	127	10.2 (77)	3.1 (5)	138	8.7 (10)	113	4.4 (72)	136	69.9 (91)
8255	1,730	11.7 (99)	6.1 (49)	1,796	11.5 (59)	1,521	5.5 (100)	1,751	64.5 (111)
8257	1,530	14.9 (127)	2.7 (3)	1,669	10.8 (40)	1,345	9.7 (135)	1,603	65.1 (109)
8263	422	8.3 (39)	7.3 (87)	436	10.8 (40)	378	3.4 (33)	423	64.8 (110)
8264	664	10.2 (77)	6.6 (66)	685	12.0 (67)	583	4.1 (63)	668	72.3 (82)
8267	539	7.8 (30)	7.4 (91)	566	9.4 (20)	484	3.5 (39)	549	61.0 (117)
8901	775	16.9 (140)	3.4 (8)	809	17.8 (140)	631	9.7 (135)	772	52.3 (134)
8902	1,151	15.6 (133)	3.2 (6)	1,222	18.8 (142)	930	7.0 (122)	1,194	63.7 (112)
8903	1,236	14.5 (124)	3.2 (6)	1,304	16.6 (131)	1,016	7.4 (126)	1,233	54.2 (131)
8904	1,913	15.9 (135)	3.7 (11)	1,971	17.7 (138)	1,543	7.1 (123)	1,923	54.4 (129)
8905	1,607	10.9 (87)	6.3 (55)	1,718	13.9 (106)	1,363	5.1 (89)	1,678	69.7 (93)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

Run Date: 5/29/2024

## Table 8A

2020 - 2022 Pregnancy Nutrition Surveillance (1)

Page : 122

## Michigan

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)									
Clinic	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
8909	940	15.2 (129)	3.9 (16)	975	16.3 (130)	779	8.1 (128)	948	60.3 (120)
9045	1,237	9.3 (56)	7.8 (98)	1,274	12.0 (67)	1,080	4.9 (81)	1,248	42.6 (139)
9050	829	13.3 (116)	5.3 (32)	880	14.0 (108)	708	6.8 (120)	847	56.6 (127)
9051	1,370	9.6 (67)	6.9 (73)	1,376	13.3 (91)	1,164	5.7 (104)	1,384	68.4 (101)
9053	1,223	16.6 (137)	3.8 (12)	1,277	13.7 (102)	1,039	9.3 (133)	1,228	61.3 (116)
9055	6	*	*	6	*	5	*	6	*
9058	1,101	15.2 (129)	2.7 (3)	1,176	12.9 (84)	946	8.4 (131)	1,151	59.0 (125)
9066	880	17.0 (141)	4.0 (17)	926	17.0 (132)	736	8.3 (130)	887	51.6 (135)
9068	773	16.7 (138)	3.6 (10)	790	15.1 (123)	653	9.3 (133)	773	44.2 (138)
9069	1,680	15.8 (134)	3.5 (9)	1,749	17.4 (135)	1,386	7.8 (127)	1,678	38.8 (140)
9070	1,452	14.5 (124)	4.4 (21)	1,508	14.9 (122)	1,224	7.3 (125)	1,451	56.5 (128)
9072	318	15.1 (128)	3.8 (12)	339	17.7 (138)	267	6.4 (114)	324	54.3 (130)
9080	2,226	14.7 (126)	3.8 (12)	2,339	14.5 (116)	1,882	7.1 (123)	2,307	53.1 (133)
9082	100	14.0 (119)	2.0 (1)	105	11.4 (55)	88	*	101	66.3 (105)
9083	7	*	*	6	*	4	*	7	*
9084	4	*	*	4	*	4	*	4	*
9087	12	*	*	13	*	10	*	12	*
<b>Michigan</b>	<b>126,335</b>	<b>11.2</b>	<b>6.1</b>	<b>131,479</b>	<b>13.0</b>	<b>109,108</b>	<b>5.1</b>	<b>127,943</b>	<b>52.7</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.