

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0101	188	93.6	0.0	2.7	0.5	0.0	3.2	0.0
0102	658	91.8	0.5	2.9	0.2	0.0	4.7	0.0
0103	653	91.6	0.0	2.9	0.0	0.0	5.5	0.0
0104	225	89.8	0.0	3.6	0.0	0.4	6.2	0.0
0250	258	80.2	0.0	0.8	1.2	0.0	17.8	0.0
0251	172	82.6	0.0	0.6	3.5	0.6	12.8	0.0
0252	198	41.4	0.5	2.0	1.0	0.0	55.1	0.0
0253	287	66.9	0.3	1.0	4.9	0.7	26.1	0.0
0401	771	91.3	0.5	2.1	0.0	0.1	6.0	0.0
0402	585	82.2	0.5	1.9	0.7	0.3	14.4	0.0
0403	205	95.6	0.5	2.4	0.0	0.0	1.5	0.0
0404	239	88.7	0.0	2.9	0.0	0.0	8.4	0.0
0501	845	92.2	0.5	2.7	0.1	0.2	4.3	0.0
0502	3	*	*	*	*	*	*	*
0503	252	88.1	0.0	5.2	0.4	0.0	6.3	0.0
0504	836	84.0	0.6	2.8	1.7	1.2	9.7	0.1
0505	505	85.9	0.2	4.2	0.4	0.4	8.9	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0506	329	93.6	0.0	2.1	0.0	0.0	4.0	0.3
0507	217	85.7	0.0	2.8	0.0	0.0	11.5	0.0
0650	1,529	80.0	3.7	4.9	1.4	0.9	9.0	0.1
0651	996	86.5	0.1	6.5	0.2	0.6	5.9	0.1
0652	475	92.2	0.0	3.2	0.0	0.0	4.6	0.0
0653	732	96.3	0.0	2.2	0.1	0.0	1.4	0.0
0654	662	92.6	1.1	1.5	0.2	0.0	4.7	0.0
0655	614	89.6	0.0	3.3	0.2	0.0	7.0	0.0
0657	120	95.0	0.0	0.8	0.0	0.0	4.2	0.0
0701	726	86.2	1.1	3.7	0.0	2.3	6.6	0.0
0702	149	85.9	0.0	0.7	0.7	0.0	12.8	0.0
0703	336	89.0	0.0	3.0	0.9	0.0	6.8	0.3
0704	135	91.9	0.0	2.2	0.7	0.0	4.4	0.7
0705	204	89.7	0.0	1.0	0.0	0.0	9.3	0.0
0850	1,706	78.2	4.3	8.5	0.0	0.9	8.0	0.1
0851	1,416	90.3	0.6	4.4	0.0	0.1	4.4	0.1
0901	2,657	69.5	2.0	16.4	0.0	0.1	11.9	0.1

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0902	128	89.8	0.0	4.7	0.0	0.0	5.5	0.0
1051	2,077	85.5	0.8	6.3	0.5	0.2	6.6	0.1
1101	2,753	32.4	54.4	6.1	0.1	0.8	6.2	0.1
1102	1,308	68.1	11.2	6.0	0.3	0.7	13.5	0.2
1106	166	74.1	0.0	15.1	0.6	0.0	10.2	0.0
1250	1,707	78.6	0.2	16.6	0.1	0.0	4.5	0.0
1251	1,360	89.4	0.2	5.1	0.1	0.1	5.1	0.0
1252	462	48.7	0.4	47.6	0.0	0.0	3.2	0.0
1253	1,968	72.3	4.7	14.5	0.1	0.2	8.3	0.0
1350	3,923	48.2	19.7	12.0	0.1	5.3	14.7	0.1
1351	248	87.1	0.8	4.4	0.0	1.2	6.5	0.0
1352	707	53.7	23.8	9.2	0.0	0.3	13.0	0.0
1401	347	85.6	0.3	6.3	0.0	0.0	7.5	0.3
1402	258	49.6	0.8	34.9	6.2	0.0	8.5	0.0
1403	55	*	*	*	*	*	*	*
1404	17	*	*	*	*	*	*	*
1501	1,743	42.7	6.0	44.7	0.2	0.0	5.7	0.6

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
1503	1,945	52.6	6.2	31.7	0.2	0.6	8.0	0.7
1504	3,814	38.9	4.0	48.6	0.0	2.9	5.5	0.1
1505	1,330	82.5	1.5	11.1	0.2	0.4	4.4	0.0
1506	1,119	76.2	1.3	17.1	0.1	0.0	5.2	0.2
1507	1,063	84.1	1.1	7.1	0.0	0.8	6.8	0.0
1508	150	0.0	0.0	100.0	0.0	0.0	0.0	0.0
1509	997	50.5	2.7	43.9	0.0	0.1	2.7	0.1
1511	1,177	86.9	0.9	6.5	0.1	0.3	5.4	0.0
1512	682	80.8	0.3	9.7	0.1	0.0	9.1	0.0
1701	1,160	48.4	0.5	3.5	14.8	0.4	32.2	0.1
1950	953	90.1	0.0	6.0	0.0	0.2	3.7	0.0
1952	248	88.3	0.0	5.2	0.4	0.4	5.6	0.0
1953	504	85.1	0.8	9.7	0.2	0.0	4.2	0.0
1954	962	81.3	0.1	16.1	0.0	0.0	2.5	0.0
1955	251	73.7	0.4	23.5	0.0	0.4	2.0	0.0
1957	1,070	80.8	2.0	12.1	0.1	0.4	4.7	0.0
2150	1,122	82.4	0.1	3.4	0.9	0.0	13.0	0.2

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
2151	406	88.4	0.2	4.4	0.2	0.0	6.7	0.0
2251	698	86.2	0.1	3.9	0.0	0.4	9.2	0.1
2252	336	94.0	0.0	1.8	0.0	0.0	4.2	0.0
2551	1	*	*	*	*	*	*	*
2552	19	*	*	*	*	*	*	*
2554	257	78.6	7.8	5.1	0.0	0.0	8.6	0.0
2555	10,909	44.9	39.2	6.2	0.1	0.3	9.4	0.0
3001	1,625	10.4	67.3	17.7	0.0	0.4	4.2	0.1
3002	452	16.8	46.5	31.2	0.0	0.0	5.5	0.0
3004	270	0.0	0.0	100.0	0.0	0.0	0.0	0.0
3007	54	*	*	*	*	*	*	*
3009	15	*	*	*	*	*	*	*
3201	884	91.5	0.3	5.2	0.2	0.0	2.7	0.0
3301	5,298	30.8	28.6	21.7	0.0	4.9	13.9	0.1
3302	1	*	*	*	*	*	*	*
3304	243	43.2	14.4	11.1	0.4	24.7	6.2	0.0
3305	257	57.6	12.1	14.4	0.4	3.5	12.1	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
3306	516	83.5	3.5	5.6	0.0	0.2	7.2	0.0
3308	124	86.3	0.8	7.3	0.8	0.0	4.8	0.0
3309	71	*	*	*	*	*	*	*
3312	765	22.1	32.3	27.1	0.0	2.7	15.7	0.1
3550	140	0.0	0.0	2.9	92.1	0.0	5.0	0.0
3551	51	*	*	*	*	*	*	*
3552	82	*	*	*	*	*	*	*
3701	1,870	83.4	0.3	11.9	0.1	0.1	4.3	0.0
3850	5,061	62.6	12.4	9.1	0.1	0.3	15.5	0.0
3950	2,990	56.8	16.9	11.9	0.0	0.8	13.5	0.1
3952	2	*	*	*	*	*	*	*
3954	101	68.3	5.0	9.9	0.0	5.9	10.9	0.0
4001	2,019	22.1	57.3	9.2	0.0	0.4	10.8	0.0
4002	1,828	41.8	24.7	18.9	0.1	0.4	13.9	0.0
4103	208	7.7	66.8	5.8	0.0	8.7	11.1	0.0
4106	3,890	10.9	36.0	44.9	0.0	0.4	7.7	0.1
4107	5,619	40.8	20.5	23.3	0.1	1.1	14.0	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
4109	1,925	81.9	1.2	9.7	0.0	0.2	6.9	0.1
4110	2	*	*	*	*	*	*	*
4112	989	0.1	0.2	99.6	0.0	0.1	0.0	0.0
4113	1,252	4.5	20.0	71.8	0.0	0.6	3.2	0.0
4120	3,380	43.5	7.3	35.3	0.0	2.6	11.3	0.1
4160	3,062	30.6	30.5	17.6	0.1	11.0	10.3	0.0
4351	934	87.3	0.4	8.2	0.0	0.0	4.1	0.0
4352	322	67.1	10.9	5.0	0.0	0.0	17.1	0.0
4353	840	60.5	0.1	34.5	0.0	0.0	4.9	0.0
4354	1,207	83.1	4.0	3.6	0.5	0.4	8.1	0.3
4355	521	75.4	0.2	10.2	0.0	0.0	14.2	0.0
4356	841	78.1	0.2	15.3	0.0	0.0	6.3	0.0
4358	301	7.3	0.0	91.7	0.0	0.0	1.0	0.0
4359	996	65.4	0.1	32.4	0.0	0.0	2.1	0.0
4360	359	90.5	0.0	8.4	0.0	0.0	1.1	0.0
4361	363	92.6	0.0	0.8	0.0	0.0	6.6	0.0
4362	528	91.3	0.4	3.2	0.0	0.2	4.9	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
4363	1,431	90.4	0.2	4.1	0.0	0.1	4.8	0.3
4401	2,012	84.6	0.4	7.4	0.0	0.0	7.4	0.1
4501	1,564	86.6	1.0	8.2	0.1	0.0	4.2	0.0
4601	2,253	67.2	1.3	24.2	0.0	0.0	7.1	0.0
4701	2,086	88.8	0.6	5.0	0.1	0.5	5.0	0.0
5050	2,474	56.8	26.7	3.2	0.1	1.9	11.3	0.0
5051	3,823	51.6	30.0	5.0	0.1	1.8	11.4	0.0
5052	3,674	49.9	24.9	3.5	0.0	11.4	10.3	0.0
5055	784	90.8	2.8	2.6	0.0	3.1	0.6	0.1
5101	3,238	67.3	9.0	12.9	0.0	1.5	9.1	0.1
5202	983	82.5	0.5	3.8	0.4	0.1	12.6	0.1
5203	224	75.4	1.3	8.0	0.0	0.4	14.7	0.0
5204	149	76.5	0.0	2.7	0.0	0.0	20.8	0.0
5601	2,031	87.8	1.0	3.7	0.1	0.3	6.7	0.2
5801	2,888	78.4	5.0	7.7	0.1	0.1	8.7	0.1
5803	359	87.7	0.6	6.1	0.3	0.6	4.5	0.3
5804	133	83.5	0.0	10.5	0.0	0.0	5.3	0.8

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
6101	130	88.5	0.0	7.7	0.0	0.0	3.8	0.0
6102	447	78.7	2.0	10.7	0.0	0.0	8.5	0.0
6103	5,583	52.2	22.8	12.9	0.2	0.1	11.8	0.0
6104	672	25.9	52.1	10.9	0.0	0.3	10.9	0.0
6105	107	79.4	1.9	15.0	0.0	0.0	3.7	0.0
6301	6,464	39.7	26.0	24.4	0.1	2.3	7.5	0.0
6312	4,021	40.0	48.2	3.0	0.1	2.0	6.6	0.1
6316	2,417	68.3	15.7	8.0	0.1	2.2	5.7	0.0
6320	367	61.0	25.9	5.4	0.3	2.7	4.6	0.0
6325	279	91.0	0.7	4.3	0.0	0.0	3.9	0.0
6330	902	18.7	54.8	17.7	0.0	0.6	8.2	0.0
6335	1,017	82.3	6.7	2.1	0.1	3.0	5.8	0.0
6340	170	27.6	22.9	42.9	0.0	2.9	3.5	0.0
6345	45	*	*	*	*	*	*	*
6350	404	49.3	35.1	3.5	0.0	0.5	11.6	0.0
6877	57	*	*	*	*	*	*	*
6895	29	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
7304	363	87.9	0.8	7.2	0.0	0.0	4.1	0.0
7306	4,015	40.7	32.5	18.4	0.0	0.4	7.9	0.1
7308	280	88.2	1.1	6.4	0.0	0.0	4.3	0.0
7310	55	*	*	*	*	*	*	*
7401	3,685	74.7	4.7	8.1	0.1	0.2	12.3	0.1
7601	1,052	88.3	0.0	8.3	0.0	0.0	3.4	0.0
7850	1,768	85.4	0.3	7.5	0.0	0.2	6.4	0.1
7852	167	96.4	1.2	1.2	0.0	0.0	1.2	0.0
8101	1,541	9.3	85.1	1.4	0.1	0.8	3.3	0.0
8102	643	7.0	3.0	88.8	0.0	0.0	1.2	0.0
8105	557	41.1	41.3	5.0	0.2	0.9	11.5	0.0
8106	485	20.4	59.0	9.1	0.0	0.0	11.3	0.2
8150	164	45.1	20.1	17.7	0.0	7.9	9.1	0.0
8151	4,742	32.4	38.4	12.8	0.2	3.4	12.7	0.1
8152	308	26.6	51.0	5.8	0.0	0.3	16.2	0.0
8153	52	*	*	*	*	*	*	*
8154	89	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
8155	168	48.8	24.4	8.9	0.6	10.7	6.5	0.0
8156	4	*	*	*	*	*	*	*
8157	33	*	*	*	*	*	*	*
8201	3,682	49.4	22.8	6.4	0.1	1.6	19.5	0.1
8202	5,298	86.8	6.9	3.8	0.0	0.7	1.7	0.1
8203	505	59.4	26.7	4.2	0.0	1.0	8.7	0.0
8250	1,291	23.2	61.5	5.3	0.2	1.2	8.7	0.0
8252	162	0.0	96.3	0.0	0.0	0.0	3.7	0.0
8255	3,942	56.2	26.5	8.9	0.1	1.8	6.6	0.0
8257	714	50.8	17.2	0.7	0.0	29.0	2.2	0.0
8258	1,516	62.7	19.7	6.7	0.0	1.2	9.6	0.1
8260	1,374	46.5	26.9	2.0	0.0	22.3	2.3	0.0
8901	1,750	0.7	96.0	0.5	0.1	0.5	2.2	0.0
8902	1,339	3.1	93.4	0.8	0.0	0.2	2.5	0.0
8903	1,845	1.1	96.5	0.4	0.1	0.3	1.7	0.0
8904	1,889	2.8	86.3	0.6	0.2	6.4	3.7	0.1
8905	4,673	48.2	26.2	20.7	0.0	0.6	4.2	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
9010	171	3.5	94.7	1.2	0.0	0.0	0.6	0.0
9012	96	*	*	*	*	*	*	*
9013	130	0.0	95.4	0.8	0.0	1.5	1.5	0.8
9015	54	*	*	*	*	*	*	*
9016	296	0.3	95.6	0.7	0.0	0.3	3.0	0.0
9017	148	1.4	95.3	1.4	0.0	0.0	2.0	0.0
9018	277	0.4	94.6	1.8	0.0	0.0	3.2	0.0
9019	83	*	*	*	*	*	*	*
9021	19	*	*	*	*	*	*	*
9041	7	*	*	*	*	*	*	*
9045	2,551	6.6	32.7	58.3	0.0	0.0	2.2	0.1
9050	2,453	11.4	53.7	32.2	0.0	0.3	2.4	0.0
9051	2,580	6.4	9.6	81.7	0.0	0.0	2.4	0.0
9053	3,292	0.9	95.6	1.1	0.0	0.7	1.6	0.0
9055	134	6.0	90.3	3.0	0.0	0.0	0.7	0.0
9058	1,283	55.9	34.9	1.0	0.1	6.4	1.7	0.0
9066	2,741	1.1	96.3	0.7	0.0	0.1	1.8	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
9068	1,851	1.1	94.6	0.9	0.3	0.3	2.9	0.0
9069	4,042	0.8	94.5	1.0	0.0	0.0	3.6	0.0
9070	2,379	1.8	85.0	0.4	0.1	9.1	3.5	0.0
9072	2,244	1.2	95.0	0.3	0.0	1.1	2.4	0.0
9074	61	*	*	*	*	*	*	*
9080	3,880	25.3	69.2	2.5	0.0	0.1	2.9	0.0
Michigan	257,024	50.4	26.2	13.7	0.3	1.6	7.8	0.1
Nation (Prior Year)	562,982	51.4	21.4	18.0	0.5	2.3	6.4	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Page: 14
Run Date: 04/03/2014

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0101	188	28.2	25.0	52.1
0102	658	39.8	23.6	40.9
0103	653	34.5	23.1	45.6
0104	225	34.7	23.6	44.9
0250	258	28.3	24.0	53.9
0251	172	29.1	22.7	52.3
0252	198	18.2	29.3	57.6
0253	287	30.3	20.6	50.5
0401	771	29.3	26.3	47.6
0402	585	33.8	20.7	47.9
0403	205	32.2	23.4	48.3
0404	239	31.4	22.6	49.4
0501	845	30.9	24.5	47.7
0502	3	*	*	*
0503	252	34.5	21.0	46.4
0504	836	34.9	23.9	43.7
0505	505	35.8	22.0	44.8
0506	329	28.3	20.1	54.4
0507	217	23.0	24.9	54.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0650	1,529	29.9	25.7	47.7
0651	996	28.4	27.8	48.3
0652	475	25.7	29.5	48.0
0653	732	23.6	26.0	54.9
0654	662	27.9	25.5	49.8
0655	614	29.3	27.9	45.8
0657	120	31.7	22.5	48.3
0701	726	32.5	23.1	46.0
0702	149	30.2	24.2	49.0
0703	336	30.7	22.9	49.7
0704	135	21.5	24.4	54.1
0705	204	27.5	30.4	47.1
0850	1,706	33.9	25.1	44.4
0851	1,416	31.6	24.4	47.0
0901	2,657	29.4	27.3	47.0
0902	128	18.8	29.7	57.8
1051	2,077	31.1	27.8	43.9
1101	2,753	30.9	26.1	46.6
1102	1,308	34.5	26.5	42.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1106	166	27.1	29.5	45.8
1250	1,707	28.1	25.5	48.1
1251	1,360	34.2	22.5	46.3
1252	462	26.6	26.4	51.3
1253	1,968	33.5	21.2	47.4
1350	3,923	33.5	22.4	47.2
1351	248	34.3	19.0	50.0
1352	707	28.1	26.9	48.7
1401	347	29.1	24.5	48.4
1402	258	25.6	21.7	54.7
1403	55	*	*	*
1404	17	*	*	*
1501	1,743	28.9	24.7	50.7
1503	1,945	29.6	25.2	48.9
1504	3,814	28.7	26.2	48.3
1505	1,330	27.9	25.4	48.9
1506	1,119	29.5	23.9	50.0
1507	1,063	29.4	26.5	46.8
1508	150	18.0	30.7	54.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Page: 17
Run Date: 04/03/2014

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1509	997	27.1	23.6	52.1
1511	1,177	33.1	24.0	45.4
1512	682	28.9	24.3	50.0
1701	1,160	30.2	24.9	48.1
1950	953	34.5	26.0	43.0
1952	248	38.3	21.8	43.5
1953	504	31.5	26.6	45.8
1954	962	35.6	20.7	46.5
1955	251	29.9	25.5	48.2
1957	1,070	33.4	26.3	43.5
2150	1,122	30.7	25.4	46.5
2151	406	30.0	27.6	47.3
2251	698	33.4	24.5	45.7
2252	336	27.4	21.7	53.9
2551	1	*	*	*
2552	19	*	*	*
2554	257	26.8	19.5	54.9
2555	10,909	38.7	23.2	40.4
3001	1,625	30.1	23.5	49.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
3002	452	31.6	19.9	51.1
3004	270	21.9	25.2	56.7
3007	54	*	*	*
3009	15	*	*	*
3201	884	29.8	24.4	50.2
3301	5,298	35.9	23.8	42.6
3302	1	*	*	*
3304	243	28.8	23.0	50.6
3305	257	31.9	28.0	44.0
3306	516	32.2	23.4	47.3
3308	124	40.3	14.5	49.2
3309	71	*	*	*
3312	765	28.6	23.8	50.6
3550	140	34.3	22.9	43.6
3551	51	*	*	*
3552	82	*	*	*
3701	1,870	30.3	26.8	46.6
3850	5,061	31.6	24.5	46.8
3950	2,990	31.2	23.4	47.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Page: 19
Run Date: 04/03/2014

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
3952	2	*	*	*
3954	101	21.8	25.7	56.4
4001	2,019	32.0	23.2	47.7
4002	1,828	32.1	23.5	47.6
4103	208	15.4	26.0	63.0
4106	3,890	29.7	24.1	49.6
4107	5,619	33.0	24.0	46.0
4109	1,925	29.8	23.2	51.0
4110	2	*	*	*
4112	989	28.0	21.6	54.8
4113	1,252	35.3	20.4	47.0
4120	3,380	33.9	23.3	46.1
4160	3,062	34.1	24.2	44.9
4351	934	26.6	24.8	50.9
4352	322	29.2	18.0	55.0
4353	840	26.1	25.8	51.5
4354	1,207	32.4	24.9	45.5
4355	521	32.2	23.2	47.2
4356	841	30.2	23.5	48.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Page: 20
Run Date: 04/03/2014

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
4358	301	23.6	21.3	58.1
4359	996	26.3	24.2	51.8
4360	359	34.5	19.5	47.1
4361	363	26.4	26.7	50.1
4362	528	29.0	26.7	47.0
4363	1,431	31.3	24.9	47.3
4401	2,012	32.0	27.0	43.5
4501	1,564	30.3	24.7	48.9
4601	2,253	34.4	25.4	42.5
4701	2,086	33.2	27.0	42.0
5050	2,474	34.8	25.5	43.0
5051	3,823	35.2	24.7	42.5
5052	3,674	35.3	25.4	42.0
5055	784	36.4	18.1	48.2
5101	3,238	32.7	27.2	43.3
5202	983	31.3	25.7	45.2
5203	224	26.8	20.5	54.0
5204	149	27.5	26.2	47.7
5601	2,031	31.6	23.8	48.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
5801	2,888	35.1	24.4	44.1
5803	359	25.3	28.4	51.0
5804	133	20.3	27.8	56.4
6101	130	29.2	24.6	48.5
6102	447	24.4	25.1	53.2
6103	5,583	30.0	24.8	48.1
6104	672	40.0	25.4	36.6
6105	107	19.6	27.1	57.9
6301	6,464	39.8	21.9	40.9
6312	4,021	40.5	22.8	39.3
6316	2,417	40.6	22.1	39.8
6320	367	29.2	17.7	55.6
6325	279	24.4	21.5	55.9
6330	902	37.5	16.3	47.8
6335	1,017	29.9	19.9	53.7
6340	170	28.2	21.8	52.9
6345	45	*	*	*
6350	404	33.4	19.3	50.0
6877	57	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Page: 22
Run Date: 04/03/2014

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
6895	29	*	*	*
7304	363	32.2	21.8	49.9
7306	4,015	33.8	23.4	45.4
7308	280	34.6	24.3	43.6
7310	55	*	*	*
7401	3,685	31.3	26.5	45.5
7601	1,052	30.5	24.4	48.1
7850	1,768	32.5	25.8	46.1
7852	167	26.9	26.3	52.1
8101	1,541	55.3	17.3	29.0
8102	643	28.0	18.7	56.1
8105	557	45.6	18.0	38.1
8106	485	41.4	19.8	40.8
8150	164	25.6	26.8	49.4
8151	4,742	33.5	24.4	45.2
8152	308	52.6	22.1	27.3
8153	52	*	*	*
8154	89	*	*	*
8155	168	25.6	23.8	52.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Page: 23
Run Date: 04/03/2014

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months	12 - 23 Months	24 - 59 Months
		%	%	%
8156	4	*	*	*
8157	33	*	*	*
8201	3,682	40.1	25.4	36.7
8202	5,298	32.4	22.7	47.7
8203	505	37.6	20.0	45.5
8250	1,291	34.4	30.7	37.4
8252	162	34.0	23.5	44.4
8255	3,942	40.6	24.3	37.5
8257	714	30.5	21.7	50.4
8258	1,516	34.8	24.1	43.5
8260	1,374	29.2	28.5	45.4
8901	1,750	33.5	23.0	46.0
8902	1,339	37.0	24.3	41.5
8903	1,845	32.4	23.7	46.3
8904	1,889	36.0	24.5	42.6
8905	4,673	31.9	21.2	49.8
9010	171	100.0	0.0	0.0
9012	96	*	*	*
9013	130	27.7	25.4	48.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
9015	54	*	*	*
9016	296	32.8	24.0	44.9
9017	148	34.5	21.6	45.3
9018	277	27.8	22.4	53.8
9019	83	*	*	*
9021	19	*	*	*
9041	7	*	*	*
9045	2,551	29.2	25.2	49.5
9050	2,453	32.7	21.2	50.0
9051	2,580	34.1	18.8	49.6
9053	3,292	40.1	22.7	39.9
9055	134	99.3	0.0	0.7
9058	1,283	27.5	23.1	52.8
9066	2,741	34.8	22.4	46.0
9068	1,851	33.8	23.3	46.6
9069	4,042	40.8	22.2	39.5
9070	2,379	32.4	25.3	46.2
9072	2,244	36.2	20.6	45.4
9074	61	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Page: 25
Run Date: 04/03/2014

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
9080	3,880	30.1	24.1	48.1
Michigan	257,024	33.4	21.0	45.6
Nation (Prior Year)	562,982	29.9	21.8	48.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0101	34	*	*	186	3.2	2.7	8.6	98	*	*	127	10.2
0102	162	9.3	10.5	645	7.1	2.5	12.4	262	14.9	16.8	414	12.8
0103	140	2.9	8.6	638	4.4	1.6	10.2	283	23.7	14.8	431	12.8
0104	56	*	*	225	4.4	2.2	8.9	101	14.9	11.9	135	11.1
0250	49	*	*	248	4.8	3.6	10.1	129	14.0	19.4	172	8.1
0251	37	*	*	167	3.6	0.0	10.8	88	*	*	112	1.8
0252	32	*	*	191	6.3	1.0	16.2	108	14.8	25.9	141	8.5
0253	56	*	*	283	1.4	11.0	5.3	141	8.5	5.7	189	9.0
0401	170	8.2	10.6	756	6.7	1.5	13.4	355	13.2	18.3	471	3.4
0402	122	6.6	6.6	575	4.9	1.9	11.1	272	18.0	14.7	323	9.9
0403	51	*	*	200	3.5	1.5	15.0	94	*	*	119	1.7
0404	53	*	*	234	2.1	1.7	11.1	113	21.2	18.6	135	11.1
0501	142	5.6	9.2	834	4.7	4.0	11.4	395	19.2	13.9	560	11.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0502	0	*	*	3	*	*	*	2	*	*	1	*
0503	53	*	*	248	4.0	6.0	9.7	114	15.8	9.6	163	6.7
0504	159	6.3	9.4	817	5.4	1.5	10.4	354	21.2	14.1	545	7.5
0505	127	4.7	8.7	486	5.3	3.7	7.6	218	17.9	10.1	307	9.1
0506	54	*	*	321	5.0	3.1	9.7	173	17.3	11.0	222	5.9
0507	27	*	*	212	6.1	2.4	12.7	113	23.9	17.7	160	17.5
0650	346	6.6	8.1	1,499	4.4	2.7	10.1	706	20.1	15.2	1,032	5.5
0651	208	4.3	3.8	960	5.0	2.9	11.1	449	16.7	15.8	664	9.9
0652	86	*	*	467	9.4	2.1	12.6	220	18.6	13.6	305	11.5
0653	135	2.2	6.7	711	3.7	1.5	11.8	382	17.0	16.2	504	9.1
0654	146	8.9	13.0	642	4.7	2.5	8.3	314	19.1	13.4	464	12.5
0655	133	7.5	12.0	596	6.7	3.0	9.6	266	16.2	15.0	408	3.9
0657	24	*	*	117	6.0	0.9	12.0	56	*	*	85	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0701	159	5.0	15.7	710	4.9	2.1	8.6	319	20.7	9.7	425	5.4
0702	32	*	*	144	7.6	2.8	6.9	69	*	*	88	*
0703	66	*	*	329	3.6	2.7	14.6	160	18.1	19.4	218	2.8
0704	24	*	*	133	6.0	1.5	12.0	71	*	*	94	*
0705	32	*	*	199	6.5	0.5	10.1	93	*	*	131	1.5
0850	395	9.6	9.1	1,658	5.2	2.9	11.0	721	17.9	15.4	1,093	5.4
0851	318	11.0	7.5	1,373	4.4	3.9	9.3	638	16.3	12.5	939	6.7
0901	562	10.5	9.6	2,601	6.5	1.8	10.9	1,206	17.6	15.2	1,760	6.1
0902	10	*	*	127	5.5	4.7	9.4	74	*	*	104	20.2
1051	502	7.0	13.1	1,976	4.8	5.4	9.8	881	18.4	13.3	1,248	11.7
1101	666	8.3	7.7	2,638	6.7	2.7	10.4	1,203	16.6	13.1	1,791	14.2
1102	332	10.5	6.6	1,267	5.9	1.3	10.5	532	15.4	15.0	804	7.2
1106	33	*	*	159	6.3	0.6	10.1	71	*	*	118	5.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1250	333	6.9	10.2	1,658	6.9	2.1	13.9	790	15.9	15.8	1,136	5.8
1251	309	6.8	9.4	1,322	6.4	2.7	10.1	604	16.9	11.6	889	4.5
1252	72	*	*	438	8.2	1.4	13.2	221	16.3	19.0	339	8.6
1253	436	8.5	10.3	1,897	6.9	2.8	10.8	884	16.7	13.2	1,295	11.0
1350	915	9.4	8.1	3,821	7.2	2.9	13.2	1,790	17.4	16.3	2,569	15.4
1351	62	*	*	238	6.3	3.4	11.8	118	16.9	16.1	158	11.4
1352	132	11.4	5.3	688	6.0	2.8	11.0	329	22.8	12.8	464	18.1
1401	72	*	*	334	4.5	5.1	12.6	162	23.5	16.0	231	5.6
1402	43	*	*	245	4.9	2.9	12.2	135	17.8	17.0	155	21.3
1403	8	*	*	53	*	*	*	32	*	*	35	*
1404	4	*	*	16	*	*	*	8	*	*	13	*
1501	370	9.5	11.4	1,693	4.3	2.4	13.6	844	18.8	18.5	1,180	30.6
1503	438	9.4	8.7	1,891	4.2	2.3	11.8	908	17.7	16.1	1,304	20.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1504	813	8.0	6.5	3,710	4.5	3.2	10.5	1,772	16.8	13.5	2,492	30.1
1505	243	8.6	16.0	1,293	4.9	3.1	9.3	623	14.8	10.3	894	18.2
1506	243	7.0	8.6	1,090	4.2	3.7	10.6	532	16.2	13.2	760	21.2
1507	213	5.2	8.9	1,034	3.9	3.8	7.0	477	17.2	8.2	699	32.9
1508	20	*	*	149	4.0	0.7	16.1	82	*	*	112	25.0
1509	187	8.0	12.3	967	4.4	3.1	11.5	494	14.8	16.2	710	23.1
1511	261	5.4	10.7	1,142	3.5	3.3	10.1	507	19.3	15.4	764	25.3
1512	133	7.5	15.8	666	4.8	2.7	9.9	333	12.6	9.9	463	29.6
1701	237	5.9	10.5	1,109	5.0	2.3	13.2	520	18.3	14.8	729	8.2
1950	220	10.0	7.7	939	6.7	1.8	10.2	399	16.8	13.3	609	6.1
1952	57	*	*	245	2.9	2.4	11.0	105	23.8	15.2	144	3.5
1953	108	11.1	5.6	492	8.7	2.2	13.6	222	11.7	12.6	330	3.9
1954	210	4.8	7.6	935	4.6	2.4	12.0	425	15.3	16.7	626	4.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1955	47	*	*	246	6.1	1.2	11.4	118	12.7	14.4	173	6.4
1957	242	7.9	7.4	1,039	5.1	4.5	10.7	442	17.6	15.8	720	4.4
2150	248	5.6	14.1	1,096	7.3	0.8	12.0	499	15.2	17.4	667	2.8
2151	91	*	*	403	4.5	1.0	9.7	189	12.7	14.8	277	1.4
2251	154	14.9	4.5	675	5.9	0.9	9.9	302	16.9	13.9	459	7.6
2252	65	*	*	322	6.5	2.2	12.4	168	24.4	18.5	233	9.0
2551	0	*	*	1	*	*	*	1	*	*	1	*
2552	10	*	*	19	*	*	*	3	*	*	5	*
2554	48	*	*	255	4.3	3.9	12.9	139	14.4	13.7	161	8.7
2555	2,835	11.1	5.7	10,666	6.7	2.9	10.6	4,257	15.4	13.4	6,697	16.9
3001	353	13.0	4.8	1,561	6.5	2.2	12.1	777	19.9	12.2	1,099	7.6
3002	97	*	*	425	4.2	1.4	15.8	220	18.6	12.7	306	12.1
3004	43	*	*	252	4.8	0.8	19.0	141	16.3	27.7	188	4.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
3007	10	*	*	54	*	*	*	34	*	*	39	*
3009	1	*	*	10	*	*	*	8	*	*	12	*
3201	169	7.7	13.6	871	6.4	2.8	14.5	435	15.4	16.1	594	6.7
3301	1,262	10.4	6.3	5,180	6.0	3.5	11.1	2,176	15.5	15.0	3,443	12.7
3302	0	*	*	1	*	*	*	1	*	*	1	*
3304	45	*	*	240	2.9	3.8	11.3	120	14.2	11.7	171	14.0
3305	58	*	*	253	2.8	3.6	8.3	113	9.7	9.7	168	8.3
3306	109	6.4	8.3	508	3.0	2.0	11.2	238	16.8	14.3	346	4.6
3308	35	*	*	121	7.4	1.7	7.4	58	*	*	70	*
3309	12	*	*	70	*	*	*	37	*	*	48	*
3312	145	14.5	8.3	746	6.3	1.3	13.9	377	19.1	17.8	521	11.9
3550	32	*	*	137	9.5	4.4	18.2	58	*	*	80	*
3551	9	*	*	50	*	*	*	22	*	*	26	*

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
3552	19	*	*	79	*	*	*	37	*	*	44	*
3701	409	9.3	7.8	1,816	4.3	3.2	9.3	836	16.4	13.4	1,248	8.2
3850	1,123	10.7	6.9	4,839	5.0	3.6	9.8	2,245	17.6	14.2	3,200	16.3
3950	712	7.7	7.7	2,897	4.5	3.7	10.6	1,372	18.8	14.1	1,886	23.4
3952	0	*	*	2	*	*	*	1	*	*	1	*
3954	13	*	*	99	*	*	*	56	*	*	70	*
4001	425	9.4	4.9	1,944	6.2	3.5	11.4	931	16.3	12.7	1,391	33.3
4002	390	7.7	6.4	1,765	5.1	2.5	11.6	836	15.6	14.0	1,255	19.8
4103	24	*	*	204	3.4	7.8	11.8	127	17.3	15.7	141	29.1
4106	820	11.7	5.6	3,776	7.3	1.7	10.9	1,852	17.3	14.5	2,669	17.5
4107	1,276	9.9	9.0	5,456	6.3	1.5	9.9	2,481	17.5	12.0	3,744	15.5
4109	414	8.2	11.6	1,867	4.8	3.1	7.3	931	15.8	10.8	1,302	4.9
4110	0	*	*	2	*	*	*	2	*	*	2	*

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
4112	185	6.5	7.6	952	5.8	4.6	17.5	514	18.1	22.8	773	7.2
4113	311	8.0	6.1	1,225	3.7	2.2	11.9	567	16.0	20.3	810	8.5
4120	749	7.2	10.0	3,290	5.9	1.2	11.4	1,493	15.5	12.4	2,206	8.3
4160	662	8.8	6.5	2,963	6.1	2.1	9.4	1,307	17.8	11.9	2,024	14.4
4351	174	10.9	6.9	912	7.0	2.0	9.4	462	20.6	12.1	658	10.6
4352	69	*	*	304	6.6	1.0	11.8	164	17.1	14.6	208	15.4
4353	159	10.1	10.7	797	5.5	2.9	13.8	401	21.2	19.5	577	15.3
4354	271	7.0	10.7	1,190	6.0	1.6	9.7	535	18.7	12.7	803	13.8
4355	132	8.3	9.8	511	4.9	3.5	9.4	240	15.0	15.4	329	14.3
4356	193	7.8	5.2	778	5.5	1.9	12.1	376	17.8	17.6	547	21.4
4358	51	*	*	281	9.6	1.1	24.2	160	21.9	33.8	216	23.6
4359	202	9.4	8.4	969	6.0	1.2	13.9	501	18.0	15.4	650	14.3
4360	86	*	*	347	4.6	2.0	8.6	162	22.2	11.7	233	17.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
4361	74	*	*	350	6.0	2.0	10.3	174	20.7	12.1	249	18.5
4362	129	9.3	8.5	517	7.4	1.0	11.6	241	15.4	16.6	335	13.4
4363	300	7.7	8.7	1,393	5.6	1.7	9.6	648	17.7	14.7	965	18.7
4401	467	4.7	10.3	1,971	6.0	2.5	10.5	853	16.3	13.7	1,290	10.2
4501	326	8.0	6.7	1,521	6.4	1.5	9.1	734	14.7	12.9	1,047	7.3
4601	548	8.2	7.3	2,197	6.3	2.5	12.8	919	19.9	18.0	1,190	8.7
4701	488	9.0	8.8	2,028	5.9	1.5	10.0	831	15.9	15.0	1,340	6.3
5050	592	10.8	5.4	2,412	6.1	1.7	11.4	1,020	13.9	13.6	1,559	17.3
5051	957	10.6	7.2	3,726	6.9	1.2	12.6	1,555	14.9	13.8	2,382	17.2
5052	928	9.5	6.4	3,580	6.1	1.5	11.8	1,477	15.6	14.8	2,314	17.1
5055	206	8.7	7.3	762	4.2	1.3	12.5	364	15.9	14.8	501	29.9
5101	761	9.2	5.5	3,163	7.5	1.8	11.2	1,348	17.8	15.9	2,063	11.3
5202	224	9.8	7.6	910	6.3	3.5	12.7	407	17.0	16.7	531	5.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
5203	51	*	*	201	8.0	0.5	9.5	110	16.4	12.7	109	4.6
5204	35	*	*	137	6.6	2.9	8.8	66	*	*	76	*
5601	420	7.9	7.1	1,988	4.2	1.6	11.5	943	16.8	13.7	1,284	11.7
5801	721	8.7	8.9	2,796	6.4	3.2	9.6	1,205	18.2	14.0	1,809	8.3
5803	51	*	*	357	5.3	0.8	8.7	182	13.7	11.5	252	7.9
5804	15	*	*	129	6.2	3.1	12.4	71	*	*	100	5.0
6101	28	*	*	126	7.1	2.4	11.9	60	*	*	84	*
6102	85	*	*	435	7.1	3.4	10.8	229	13.1	12.2	296	11.8
6103	1,268	9.5	6.2	5,418	5.1	3.5	9.5	2,582	18.5	13.4	3,656	19.0
6104	179	11.7	3.4	658	6.5	4.7	8.1	244	16.8	9.4	384	25.5
6105	12	*	*	103	3.9	2.9	8.7	59	*	*	75	*
6301	1,573	10.9	6.2	6,301	6.4	1.8	9.2	2,545	16.1	13.6	3,933	12.9
6312	985	9.8	8.1	3,940	6.0	2.9	7.7	1,527	15.1	9.2	2,513	17.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
6316	634	7.4	9.1	2,370	4.9	1.9	9.9	930	16.0	13.5	1,410	11.3
6320	66	*	*	346	4.0	4.9	8.7	188	12.2	13.3	239	11.3
6325	46	*	*	272	7.4	4.4	11.8	150	21.3	14.7	189	12.7
6330	231	15.6	5.6	864	6.0	3.6	8.0	417	15.8	11.5	545	18.2
6335	168	9.5	9.5	975	3.7	3.7	9.1	514	13.0	12.1	698	13.8
6340	22	*	*	168	3.0	1.8	10.7	89	*	*	122	12.3
6345	12	*	*	43	*	*	*	19	*	*	27	*
6350	90	*	*	389	4.1	5.7	12.3	193	10.9	16.6	259	8.9
6877	5	*	*	56	*	*	*	27	*	*	34	*
6895	1	*	*	29	*	*	*	17	*	*	26	*
7304	74	*	*	353	5.7	1.4	12.5	173	19.7	16.2	251	5.2
7306	949	10.1	7.0	3,906	5.6	1.6	13.4	1,760	15.5	17.6	2,678	14.7
7308	66	*	*	271	5.9	1.8	12.2	119	16.0	12.6	190	7.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
7310	11	*	*	53	*	*	*	19	*	*	29	*
7401	835	9.7	6.9	3,592	6.0	2.0	8.7	1,606	15.6	13.4	2,494	4.3
7601	222	8.6	8.1	1,031	4.0	2.7	9.7	489	19.8	14.3	726	5.4
7850	415	7.5	8.9	1,708	4.5	4.4	11.4	772	15.7	17.2	1,177	6.3
7852	25	*	*	158	1.9	3.2	7.6	80	*	*	126	5.6
8101	571	16.1	4.2	1,485	7.4	6.8	10.4	431	11.4	13.5	757	28.5
8102	120	6.7	10.0	610	5.7	7.0	18.9	329	17.9	25.2	451	16.0
8105	191	12.6	5.8	531	9.0	4.0	10.4	201	18.9	12.4	302	19.2
8106	152	11.2	5.3	471	7.6	1.7	10.4	194	18.0	13.4	277	22.4
8150	29	*	*	158	5.1	2.5	4.4	76	*	*	114	17.5
8151	1,085	9.8	6.1	4,627	5.6	2.3	10.0	2,062	16.7	13.9	3,157	18.4
8152	103	11.7	8.7	305	8.5	3.9	10.5	82	*	*	170	30.0
8153	9	*	*	52	*	*	*	24	*	*	36	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
8154	10	*	*	88	*	*	*	51	*	*	49	*
8155	24	*	*	166	2.4	1.8	12.7	87	*	*	123	24.4
8156	0	*	*	4	*	*	*	1	*	*	2	*
8157	1	*	*	32	*	*	*	28	*	*	29	*
8201	1,030	10.1	6.8	3,392	6.5	3.1	10.7	1,223	15.7	12.9	1,816	9.1
8202	1,097	7.8	6.7	5,198	4.5	2.3	10.1	2,446	13.7	11.7	3,498	26.9
8203	110	5.5	6.4	500	5.6	1.8	11.4	225	20.4	12.0	286	12.6
8250	343	12.8	4.4	1,258	5.3	3.7	8.1	465	10.3	9.2	763	27.3
8252	34	*	*	147	13.6	4.1	12.2	68	*	*	101	38.6
8255	1,058	9.8	7.5	3,846	7.7	1.5	10.1	1,415	18.1	11.2	2,083	9.5
8257	138	8.7	3.6	688	5.2	5.4	11.2	341	13.2	13.2	443	21.9
8258	354	8.8	7.3	1,480	5.1	2.3	11.9	635	21.9	16.2	913	18.0
8260	281	12.5	3.6	1,317	5.5	2.6	9.0	604	15.6	10.8	952	16.4

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
8901	371	11.6	3.0	1,709	5.8	5.3	9.8	786	15.8	11.5	1,148	30.2
8902	299	16.1	3.3	1,299	5.2	6.7	7.9	543	12.3	9.8	827	33.4
8903	357	12.6	2.5	1,792	6.4	3.9	12.2	827	16.0	15.2	1,219	35.5
8904	414	10.4	4.1	1,825	3.8	5.7	9.4	779	14.1	12.7	1,195	28.8
8905	976	8.2	6.7	4,501	4.4	6.1	11.3	2,205	16.1	13.5	3,149	20.8
9010	171	12.3	4.7	150	6.0	15.3	1.3	0	*	*	0	*
9012	20	*	*	95	*	*	*	54	*	*	58	*
9013	19	*	*	129	3.9	4.7	7.8	63	*	*	72	*
9015	12	*	*	52	*	*	*	27	*	*	30	*
9016	54	*	*	286	6.3	2.4	9.1	130	25.4	8.5	184	32.6
9017	30	*	*	140	5.7	3.6	12.1	65	*	*	83	*
9018	47	*	*	270	7.0	3.3	10.0	145	20.7	13.8	45	*
9019	16	*	*	81	*	*	*	46	*	*	45	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
9021	2	*	*	17	*	*	*	8	*	*	12	*
9041	5	*	*	7	*	*	*	2	*	*	1	*
9045	537	9.1	7.6	2,424	5.1	1.5	14.2	1,179	17.5	18.1	1,720	20.3
9050	514	12.6	6.0	2,362	5.9	3.5	11.2	1,166	16.0	14.3	1,632	16.1
9051	579	7.8	8.1	2,367	7.5	1.4	19.1	1,133	20.3	22.1	1,683	27.4
9053	894	14.8	3.1	3,172	6.2	3.4	8.2	1,254	14.8	10.7	2,040	27.1
9055	133	9.0	4.5	120	8.3	13.3	7.5	1	*	*	1	*
9058	223	7.2	2.7	1,252	4.6	2.9	9.1	659	14.4	12.3	895	19.0
9066	666	11.6	3.0	2,677	5.5	5.3	7.3	1,227	10.3	8.2	1,707	13.5
9068	430	14.0	2.8	1,775	7.5	9.6	8.6	833	7.2	7.2	1,206	26.5
9069	1,155	12.8	3.9	3,915	6.2	3.8	8.1	1,539	14.7	10.9	2,418	26.0
9070	549	12.4	4.0	2,305	7.6	9.2	9.7	1,073	7.6	6.6	1,392	21.1
9072	542	13.1	3.1	1,994	7.5	3.7	8.6	884	13.9	11.3	1,155	19.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
9074	0	*	*	59	*	*	*	25	*	*	31	*
9080	755	12.8	4.1	3,798	5.3	2.2	8.7	1,819	15.7	12.1	2,734	37.6
Michigan	58,707	9.7	7.1	249,036	5.8	2.9	10.7	112,234	16.4	13.8	164,326	16.1
Nation (Prior Year)	131,697	9.1	6.8	550,807	6.2	2.9	10.7	235,427	16.6	14.6	377,330	13.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
0101	36	*	26	*	25	*	20	*	25	*	103	88.3	186	5.4
0102	167	62.9	128	21.1	109	9.2	93	*	113	8.0	287	83.3	658	6.8
0103	142	59.9	118	22.9	102	8.8	87	*	103	6.8	314	87.6	651	6.6
0104	56	*	29	*	28	*	31	*	28	*	108	85.2	224	11.2
0250	49	*	52	*	38	*	33	*	45	*	144	86.8	258	5.0
0251	37	*	30	*	16	*	28	*	29	*	98	*	172	7.0
0252	32	*	19	*	30	*	7	*	12	*	120	85.0	198	4.0
0253	56	*	42	*	38	*	30	*	35	*	155	85.2	285	14.0
0401	170	55.9	123	19.5	136	12.5	76	*	85	*	388	92.8	771	4.8
0402	121	62.8	112	25.9	87	*	67	*	79	*	298	85.6	581	5.7
0403	51	*	29	*	26	*	25	*	23	*	105	89.5	204	6.9
0404	53	*	39	*	33	*	23	*	25	*	125	81.6	239	12.1
0501	175	70.9	166	21.1	144	15.3	115	21.7	154	12.3	425	90.4	840	17.5
0502	0	*	0	*	0	*	0	*	0	*	2	*	3	*
0503	52	*	52	*	25	*	38	*	48	*	129	89.1	251	3.2
0504	197	79.7	143	28.7	135	14.1	100	38.0	114	16.7	394	92.4	834	4.9
0505	137	77.4	69	*	67	*	64	*	59	*	238	90.3	503	7.8
0506	66	*	52	*	50	*	39	*	46	*	188	88.3	328	16.5

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
0507	29	*	38	*	36	*	26	*	32	*	125	93.6	217	8.3
0650	346	77.2	274	26.3	274	13.5	173	30.6	231	15.6	775	84.5	1,525	6.2
0651	208	62.0	178	16.3	192	5.7	103	18.4	125	4.0	515	93.2	993	9.8
0652	86	*	72	*	90	*	41	*	53	*	249	92.4	473	7.4
0653	135	76.3	100	22.0	110	12.7	65	*	79	*	428	90.7	728	14.0
0654	146	78.1	93	*	121	10.7	69	*	78	*	348	93.4	661	9.7
0655	133	72.9	135	17.8	121	10.7	68	*	106	12.3	295	90.2	613	6.9
0657	24	*	18	*	21	*	11	*	11	*	59	*	120	11.7
0701	159	79.9	117	35.0	96	*	87	*	101	10.9	363	97.2	725	1.4
0702	32	*	26	*	20	*	12	*	17	*	76	*	148	4.7
0703	66	*	57	*	50	*	37	*	49	*	178	92.1	335	7.8
0704	24	*	14	*	25	*	5	*	11	*	78	*	133	11.3
0705	32	*	35	*	33	*	21	*	32	*	107	97.2	204	3.9
0850	400	73.0	315	21.3	279	9.0	206	21.4	273	11.7	807	87.4	1,700	4.6
0851	320	67.2	270	22.6	212	9.9	162	24.1	228	14.9	724	89.8	1,414	5.0
0901	568	51.1	472	16.3	503	8.0	250	20.4	319	3.4	1,333	96.4	2,645	6.2
0902	10	*	21	*	18	*	12	*	18	*	80	*	128	11.7
1051	503	83.1	367	33.8	395	20.0	209	39.2	315	15.6	971	92.6	2,065	3.4

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
1101	669	50.7	393	17.3	454	9.0	187	21.9	248	5.6	1,337	78.0	2,739	6.4
1102	334	54.2	244	14.8	228	9.6	135	19.3	168	4.2	598	83.1	1,304	6.2
1106	33	*	26	*	33	*	9	*	16	*	79	*	166	7.2
1250	335	68.4	288	19.4	293	13.7	189	26.5	242	10.7	881	78.1	1,703	6.4
1251	310	78.1	214	26.2	202	9.9	165	28.5	180	19.4	669	88.2	1,351	7.5
1252	72	*	101	24.8	87	*	59	*	81	*	247	89.1	460	3.3
1253	438	74.2	299	21.1	272	11.0	231	22.9	253	13.8	984	92.6	1,955	5.1
1350	918	58.5	643	14.5	559	7.5	381	16.3	410	5.9	1,972	89.9	3,881	4.1
1351	62	*	38	*	35	*	36	*	34	*	130	88.5	246	2.8
1352	132	51.5	95	*	102	9.8	61	*	80	*	364	80.2	707	8.3
1401	72	*	47	*	51	*	28	*	35	*	183	88.5	347	6.6
1402	44	*	36	*	36	*	22	*	34	*	146	95.2	254	4.3
1403	8	*	4	*	5	*	2	*	3	*	35	*	55	*
1404	4	*	3	*	4	*	1	*	3	*	9	*	17	*
1501	374	71.4	247	24.3	245	18.8	195	27.2	205	11.2	922	85.8	1,726	5.0
1503	444	68.7	299	24.4	289	13.1	218	25.2	251	12.4	1,017	90.8	1,930	3.7
1504	837	71.6	516	24.6	667	12.1	264	27.3	375	10.4	1,929	87.7	3,786	2.7
1505	266	77.1	193	38.9	221	19.0	128	36.7	163	20.9	693	89.5	1,317	2.7

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
1506	244	72.5	183	27.9	178	11.8	142	25.4	153	14.4	597	94.1	1,118	3.5
1507	226	80.1	168	25.6	205	13.2	124	29.0	145	17.9	521	87.9	1,055	5.9
1508	20	*	15	*	21	*	6	*	12	*	86	*	150	0.0
1509	193	69.9	140	25.7	152	11.2	97	*	111	10.8	561	85.0	995	7.5
1511	263	74.9	225	16.0	205	10.7	148	18.9	193	4.1	569	92.1	1,168	3.4
1512	136	79.4	104	23.1	112	7.1	65	*	76	*	357	94.4	681	4.3
1701	238	70.2	195	29.2	184	15.8	113	42.5	131	27.5	587	86.5	1,156	8.1
1950	221	73.8	162	22.8	131	8.4	122	29.5	146	11.0	445	89.2	948	8.6
1952	57	*	44	*	34	*	30	*	39	*	111	82.0	247	7.7
1953	108	73.1	69	*	66	*	53	*	65	*	249	81.9	504	6.9
1954	211	73.5	185	22.7	146	12.3	131	26.0	145	11.0	474	89.5	959	7.2
1955	47	*	47	*	47	*	25	*	38	*	126	86.5	251	7.2
1957	249	80.7	196	24.5	181	17.1	145	19.3	166	7.2	488	87.5	1,057	3.5
2150	248	69.8	197	18.3	197	9.6	133	15.8	165	4.8	564	91.1	1,114	4.9
2151	91	*	65	*	67	*	50	*	57	*	204	91.2	401	5.5
2251	157	64.3	143	25.2	126	16.7	93	*	116	9.5	334	93.4	697	3.2
2252	65	*	52	*	49	*	27	*	43	*	189	85.2	336	6.5
2551	0	*	0	*	0	*	0	*	0	*	1	*	1	*

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breastfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breastfeed At Least 6 Months	Number	%	Number	%
2552	10	*	8	*	2	*	5	*	5	*	3	*	19	*
2554	49	*	35	*	37	*	24	*	23	*	147	91.8	255	8.2
2555	2,849	55.2	2,185	11.7	1,660	5.8	1,246	12.0	1,583	5.1	4,655	78.4	10,739	6.7
3001	354	57.1	267	11.6	204	4.4	177	7.3	222	3.2	862	79.9	1,614	9.0
3002	97	*	65	*	44	*	45	*	58	*	244	97.1	445	7.0
3004	44	*	33	*	33	*	21	*	31	*	162	74.7	270	2.2
3007	10	*	4	*	3	*	1	*	2	*	37	*	54	*
3009	2	*	3	*	2	*	1	*	2	*	8	*	15	*
3201	168	59.5	129	16.3	123	10.6	76	*	86	*	469	98.1	884	5.9
3301	1,285	71.1	1,014	21.9	902	11.4	571	15.6	741	6.3	2,405	82.8	5,245	4.9
3302	0	*	0	*	0	*	0	*	0	*	1	*	1	*
3304	46	*	38	*	33	*	22	*	32	*	131	85.5	241	0.8
3305	58	*	41	*	44	*	21	*	35	*	119	96.6	254	0.8
3306	111	72.1	88	*	81	*	57	*	74	*	259	93.1	509	4.1
3308	35	*	21	*	15	*	15	*	16	*	61	*	124	2.4
3309	12	*	9	*	13	*	3	*	7	*	40	*	70	*
3312	148	70.3	111	14.4	98	*	76	*	88	*	406	79.6	760	4.5
3550	32	*	26	*	23	*	24	*	23	*	66	*	138	2.9

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
3551	9	*	6	*	7	*	7	*	6	*	25	*	51	*
3552	19	*	8	*	9	*	3	*	8	*	44	*	80	*
3701	412	73.3	343	18.7	339	7.1	228	22.4	253	10.7	941	79.6	1,866	8.9
3850	1,131	63.6	943	17.9	833	9.4	547	19.6	750	8.8	2,515	84.8	5,038	7.1
3950	716	70.3	446	30.7	450	14.0	312	22.4	355	11.3	1,514	85.2	2,958	3.5
3952	0	*	0	*	0	*	0	*	0	*	2	*	2	*
3954	13	*	15	*	16	*	4	*	9	*	60	*	101	5.0
4001	433	62.6	337	15.4	304	8.6	220	20.0	237	10.1	1,029	64.8	2,010	7.1
4002	391	61.1	260	25.8	239	11.3	192	21.9	215	11.2	934	77.0	1,827	6.1
4103	25	*	21	*	31	*	11	*	14	*	139	77.0	207	7.2
4106	824	67.4	672	19.8	607	7.4	388	13.7	474	9.5	2,026	85.6	3,865	3.2
4107	1,278	67.7	1,027	20.4	935	9.6	593	23.6	764	13.0	2,756	88.2	5,566	2.8
4109	417	67.6	328	22.3	307	11.4	214	24.8	246	12.6	1,030	94.2	1,911	4.8
4110	0	*	0	*	0	*	0	*	0	*	2	*	2	*
4112	185	29.2	118	18.6	114	13.2	89	*	91	*	581	96.6	988	3.5
4113	311	71.1	196	17.9	185	13.5	152	8.6	154	5.8	614	84.0	1,250	1.8
4120	750	70.4	662	20.1	517	7.7	425	17.6	522	8.4	1,652	88.8	3,338	1.5
4160	672	71.9	637	27.5	484	15.5	369	24.1	470	16.2	1,479	89.5	3,034	2.0

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
4351	177	70.6	128	19.5	142	12.7	95	*	109	16.5	512	97.7	934	9.5
4352	69	*	47	*	44	*	35	*	37	*	182	82.4	320	15.9
4353	159	72.3	114	15.8	121	12.4	84	*	87	*	460	83.9	835	6.5
4354	271	73.4	188	23.4	204	12.7	141	31.9	147	21.8	584	93.8	1,200	4.1
4355	133	75.2	68	*	81	*	58	*	47	*	256	90.2	518	6.0
4356	195	74.4	127	21.3	151	10.6	100	19.0	90	*	436	92.4	840	7.0
4358	51	*	39	*	44	*	25	*	33	*	183	84.7	299	1.3
4359	201	77.6	125	32.0	165	15.8	80	*	106	7.5	553	79.9	991	4.7
4360	86	*	66	*	53	*	50	*	64	*	182	86.8	357	9.0
4361	76	*	49	*	57	*	25	*	40	*	196	91.3	362	5.8
4362	130	80.8	73	*	76	*	36	*	60	*	264	88.6	526	7.6
4363	300	77.3	245	20.0	244	11.1	175	14.9	228	5.3	720	89.2	1,424	5.8
4401	467	66.4	363	25.6	352	11.4	239	20.5	317	8.8	937	92.4	1,993	4.2
4501	329	66.9	268	17.5	235	6.4	149	22.1	196	7.1	816	89.7	1,550	5.4
4601	553	70.3	443	21.4	399	12.8	322	15.8	397	7.6	1,015	79.6	2,189	5.3
4701	487	73.1	388	26.0	401	13.2	226	26.1	297	12.5	935	90.6	2,085	3.3
5050	595	58.7	398	13.3	407	8.1	260	12.7	312	5.1	1,130	81.2	2,443	3.5
5051	963	60.5	596	16.9	620	6.0	422	12.8	471	6.8	1,725	79.0	3,789	3.7

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
5052	932	57.3	549	13.5	607	6.6	362	12.2	430	5.1	1,641	78.9	3,632	3.1
5055	222	38.7	132	14.4	103	8.7	64	*	78	*	393	65.9	783	3.3
5101	778	67.9	565	19.5	590	9.8	385	16.6	462	7.1	1,484	87.1	3,220	2.5
5202	225	76.4	147	25.9	161	14.9	107	19.6	132	11.4	480	88.3	977	3.9
5203	52	*	22	*	33	*	14	*	17	*	124	89.5	223	4.9
5204	35	*	24	*	26	*	14	*	21	*	77	*	145	2.8
5601	420	80.0	339	24.5	329	11.2	293	25.9	296	12.2	1,031	91.6	1,996	4.2
5801	729	66.1	555	15.1	490	8.2	350	14.0	391	7.7	1,340	82.5	2,873	7.0
5803	51	*	55	*	66	*	25	*	42	*	195	90.8	355	4.2
5804	15	*	13	*	24	*	8	*	10	*	80	*	133	3.0
6101	28	*	16	*	21	*	14	*	13	*	67	*	130	13.8
6102	86	*	54	*	73	*	55	*	49	*	249	88.8	446	6.3
6103	1,272	56.0	830	13.0	897	6.7	560	13.4	579	5.7	2,813	86.8	5,563	6.7
6104	182	43.4	152	6.6	139	4.3	83	*	102	1.0	259	79.2	668	9.6
6105	12	*	14	*	17	*	11	*	14	*	64	*	107	4.7
6301	1,581	54.0	1,108	15.3	821	8.5	861	10.3	942	5.8	2,793	82.3	6,418	2.8
6312	989	66.2	800	22.9	568	14.3	572	16.4	676	10.2	1,679	79.5	3,978	2.8
6316	634	62.6	463	22.7	341	9.4	328	19.5	400	10.3	1,033	82.2	2,396	2.3

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
6320	69	*	62	*	40	*	40	*	53	*	210	95.7	364	2.5
6325	47	*	26	*	29	*	22	*	20	*	165	89.7	277	5.1
6330	233	54.5	149	10.7	91	*	95	*	98	*	443	77.7	891	6.5
6335	179	60.9	146	26.7	108	11.1	121	19.0	127	10.2	584	94.7	1,002	0.8
6340	22	*	25	*	24	*	17	*	22	*	96	*	166	1.2
6345	12	*	7	*	3	*	9	*	7	*	23	*	44	*
6350	96	*	43	*	42	*	35	*	35	*	214	90.2	402	2.2
6877	5	*	10	*	9	*	5	*	9	*	28	*	55	*
6895	1	*	3	*	9	*	0	*	2	*	17	*	29	*
7304	75	*	49	*	45	*	42	*	28	*	195	89.2	361	7.8
7306	951	62.6	605	14.2	600	5.3	480	12.7	417	5.8	1,951	80.9	3,959	6.5
7308	66	*	48	*	37	*	31	*	27	*	129	86.0	272	5.5
7310	11	*	14	*	8	*	3	*	6	*	21	*	53	*
7401	837	56.8	702	13.7	640	7.5	383	16.7	559	5.9	1,795	87.6	3,652	7.7
7601	222	68.0	163	22.7	151	6.0	116	15.5	142	8.5	530	88.5	1,050	6.2
7850	417	69.8	353	23.2	327	7.0	210	17.6	252	4.8	858	95.1	1,759	6.3
7852	25	*	26	*	32	*	21	*	20	*	92	*	166	5.4
8101	572	46.2	345	11.0	217	6.0	283	4.2	301	2.3	463	74.7	1,530	7.2

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
8102	120	37.5	70	*	55	*	58	*	63	*	379	79.7	642	4.7
8105	192	66.1	82	*	71	*	73	*	72	*	216	85.2	556	5.6
8106	153	49.0	63	*	50	*	76	*	55	*	207	59.9	485	4.9
8150	29	*	20	*	27	*	13	*	18	*	89	*	162	4.9
8151	1,091	77.6	849	31.7	786	18.4	601	17.3	729	11.4	2,282	83.3	4,707	4.7
8152	103	78.6	83	*	64	*	57	*	63	*	88	*	304	7.9
8153	9	*	9	*	13	*	6	*	8	*	24	*	52	*
8154	10	*	9	*	10	*	8	*	9	*	52	*	86	*
8155	24	*	27	*	24	*	22	*	25	*	95	*	166	4.2
8156	0	*	1	*	0	*	0	*	1	*	1	*	4	*
8157	1	*	2	*	2	*	2	*	2	*	30	*	33	*
8201	1,047	55.4	714	18.6	597	8.4	456	15.1	586	6.8	1,458	86.0	3,663	3.4
8202	1,147	57.1	834	32.3	749	9.6	526	5.7	675	4.3	2,680	82.4	5,251	3.0
8203	114	61.4	91	*	60	*	66	*	76	*	233	89.3	503	5.4
8250	345	43.2	239	12.6	280	5.0	92	*	95	*	511	92.8	1,269	3.3
8252	34	*	30	*	28	*	22	*	27	*	73	*	160	3.1
8255	1,075	52.2	778	15.9	626	7.8	524	11.1	702	6.3	1,554	82.9	3,933	4.7
8257	139	66.2	103	30.1	104	13.5	57	*	77	*	375	93.3	703	1.7

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breastsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
8258	359	53.2	232	13.8	227	7.5	144	11.8	189	4.8	694	85.4	1,511	5.1
8260	286	75.2	233	29.2	259	16.6	162	4.3	212	2.4	682	74.8	1,373	3.2
8901	372	33.9	238	9.7	207	1.9	182	4.9	179	2.2	858	63.6	1,749	11.1
8902	300	38.0	234	9.8	195	4.1	164	4.9	179	1.1	586	73.0	1,337	5.9
8903	357	32.5	224	6.3	197	3.0	198	2.5	194	0.5	915	71.4	1,836	9.4
8904	419	47.5	313	16.6	250	6.8	230	5.7	242	4.5	857	73.6	1,884	9.2
8905	982	41.3	566	15.4	492	5.1	455	3.3	481	2.7	2,477	81.0	4,662	5.3
9010	171	32.7	21	*	0	*	24	*	14	*	0	*	170	8.8
9012	20	*	7	*	9	*	4	*	7	*	59	*	96	*
9013	19	*	24	*	25	*	12	*	19	*	70	*	130	13.8
9015	12	*	6	*	9	*	6	*	6	*	28	*	54	*
9016	54	*	46	*	46	*	27	*	37	*	145	63.4	293	10.2
9017	30	*	18	*	17	*	9	*	15	*	72	*	146	11.6
9018	47	*	37	*	42	*	20	*	24	*	156	64.1	275	8.0
9019	16	*	11	*	12	*	7	*	8	*	49	*	83	*
9021	2	*	4	*	3	*	0	*	2	*	10	*	18	*
9041	5	*	0	*	0	*	1	*	0	*	2	*	7	*
9045	538	47.8	428	8.9	432	3.0	210	3.8	234	2.6	1,344	76.2	2,540	5.0

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breastsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
9050	537	42.1	363	12.1	279	5.7	230	6.5	264	2.7	1,303	65.0	2,437	8.2
9051	594	55.1	353	17.8	257	8.6	279	3.6	317	2.8	1,346	77.4	2,578	2.5
9053	897	37.1	600	11.2	494	6.5	397	10.1	441	6.8	1,361	81.3	3,289	16.9
9055	132	12.9	5	*	0	*	5	*	5	*	1	*	134	7.5
9058	231	63.2	196	25.5	181	8.8	143	7.7	178	2.8	725	97.7	1,281	4.7
9066	667	35.8	363	8.5	339	1.8	296	4.7	289	4.5	1,328	73.3	2,729	8.8
9068	431	39.4	286	11.2	230	3.9	209	11.5	229	4.8	917	70.0	1,848	15.7
9069	1,159	47.7	671	13.9	561	4.1	526	4.9	585	2.1	1,679	69.5	4,032	9.1
9070	558	44.4	293	12.3	293	2.0	192	12.5	210	5.7	1,187	62.3	2,360	8.1
9072	558	41.6	310	5.5	285	2.5	195	2.6	199	1.5	1,029	65.6	2,228	12.1
9074	0	*	9	*	5	*	0	*	9	*	28	*	60	*
9080	757	51.3	598	23.1	627	6.7	337	6.5	385	2.3	1,973	78.8	3,823	4.1
Michigan	59,352	60.9	42,948	19.1	39,324	9.5	28,358	16.3	33,631	7.9	124,279	83.7	255,219	5.6
Nation (Prior Year)	132,549	64.2	71,014	24.4	76,220	11.5	59,502	16.4	68,534	8.8	262,129	81.3	552,622	13.7

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 55
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)
Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0101	113	7.1	10.6	593	5.4	2.9	8.3	293	16.7	10.6	375	13.3
0102	476	6.7	12.2	1,927	7.5	3.5	12.7	801	17.2	16.9	1,133	13.1
0103	426	5.6	8.2	1,914	5.9	2.1	12.7	843	20.6	17.1	1,181	11.5
0104	144	4.2	7.6	638	4.9	2.5	9.1	273	16.1	11.4	373	7.5
0250	158	11.4	10.1	769	4.6	4.0	10.9	365	16.4	18.9	495	10.7
0251	111	8.1	4.5	564	3.0	0.5	12.8	272	17.3	17.6	337	3.9
0252	112	8.0	7.1	637	5.5	1.4	14.0	332	22.3	22.6	396	8.8
0253	169	7.1	8.9	841	2.0	9.9	5.2	396	9.6	6.1	535	8.4
0401	544	7.9	11.4	2,401	5.9	1.9	11.2	1,074	14.3	16.5	1,367	4.0
0402	380	5.5	5.8	1,821	4.8	2.3	11.7	852	15.5	15.4	1,036	11.2
0403	142	6.3	7.0	661	4.1	1.5	14.4	325	16.9	20.9	372	1.3
0404	145	9.7	9.0	709	3.4	2.5	11.1	339	16.8	16.8	393	10.4
0501	530	8.9	7.9	2,610	4.6	3.2	10.9	1,179	17.6	13.7	1,604	8.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 56
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
0502	1	*	*	14	*	*	*	7	*	*	8	*
0503	178	7.3	7.9	809	5.2	5.8	9.1	344	11.9	10.2	491	9.2
0504	528	7.4	9.8	2,597	5.4	2.0	11.9	1,138	18.9	15.1	1,594	6.9
0505	399	5.5	8.5	1,703	5.7	4.2	8.9	763	17.7	12.3	983	6.0
0506	172	5.2	8.7	991	5.1	4.5	10.6	528	16.9	12.7	652	6.9
0507	37	*	*	299	5.4	1.7	12.4	162	23.5	16.7	233	16.7
0650	1,059	7.6	7.0	4,671	6.0	2.6	10.4	2,095	18.9	16.0	3,107	6.3
0651	667	6.6	5.7	3,021	5.9	2.7	11.0	1,384	15.5	15.7	1,991	11.2
0652	293	8.2	11.6	1,548	8.2	2.1	11.8	768	17.4	14.8	993	8.3
0653	434	6.7	8.8	2,229	5.1	1.7	11.3	1,101	18.5	15.4	1,501	10.4
0654	414	7.2	11.6	1,961	4.6	1.8	9.0	940	18.2	13.9	1,371	15.9
0655	407	7.1	9.3	1,757	7.1	2.5	9.4	768	16.1	14.6	1,145	3.4
0657	69	*	*	375	4.8	1.3	12.0	194	21.6	18.6	267	15.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 57
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0701	457	6.1	10.9	2,065	5.9	1.7	7.9	923	14.5	10.2	1,127	7.6
0702	95	*	*	443	6.1	2.0	7.7	201	19.9	12.9	240	7.5
0703	230	6.1	8.3	1,051	4.9	2.3	11.1	486	20.6	17.9	611	2.8
0704	75	*	*	418	7.2	1.4	12.4	224	25.0	14.3	249	2.8
0705	141	7.8	9.9	766	6.1	1.2	9.0	366	17.5	9.6	408	2.2
0850	1,202	7.1	7.9	5,151	5.3	3.1	11.0	2,242	18.9	15.9	3,251	5.7
0851	928	8.3	8.4	4,299	4.7	3.3	9.1	2,069	16.1	11.7	2,883	5.8
0901	1,798	9.0	9.3	8,106	6.5	1.7	10.1	3,602	17.0	14.4	5,046	10.1
0902	70	*	*	420	9.3	3.6	10.0	203	17.2	14.8	287	19.9
1051	1,581	8.1	10.7	6,167	5.0	6.4	9.7	2,730	17.8	12.9	3,521	12.4
1101	2,035	10.5	7.2	8,321	7.0	2.2	10.9	3,674	16.4	14.5	5,548	12.7
1102	1,022	8.4	8.2	3,993	5.4	1.2	10.8	1,670	17.0	15.1	2,515	6.6
1106	115	6.1	7.8	512	4.9	0.6	9.2	219	19.2	10.5	331	5.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 58
Run Date: 04/02/2014

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1250	1,049	7.1	9.2	5,215	6.8	2.1	13.6	2,500	16.9	15.1	3,462	4.8
1251	915	7.8	9.3	4,147	6.3	2.0	9.9	1,918	16.5	13.3	2,767	3.1
1252	72	*	*	439	8.2	1.4	13.4	222	16.2	19.4	340	8.5
1253	1,505	8.0	9.2	6,734	7.4	3.5	11.7	3,153	18.5	14.7	4,529	11.3
1350	2,628	8.6	7.8	11,381	6.7	2.8	13.5	5,228	18.4	16.7	7,542	15.8
1351	148	8.8	14.2	700	6.3	1.9	12.9	340	15.3	16.2	477	13.6
1352	433	8.8	6.9	2,089	6.0	2.9	11.4	1,007	18.7	14.3	1,392	20.8
1401	222	10.4	8.6	1,109	5.0	5.3	12.8	534	17.6	16.3	664	4.7
1402	153	5.2	13.7	804	4.6	2.6	12.8	408	20.8	14.5	519	16.2
1403	27	*	*	148	9.5	7.4	14.9	78	*	*	93	*
1404	11	*	*	73	*	*	*	38	*	*	62	*
1501	1,121	7.7	9.6	5,478	4.3	2.5	14.3	2,736	18.1	17.2	3,817	24.8
1503	1,225	8.4	8.2	6,008	4.8	2.9	11.7	2,912	16.7	15.2	4,073	28.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 59
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1504	2,483	7.1	6.9	11,608	4.3	3.1	10.7	5,573	16.3	14.1	7,743	23.1
1505	841	8.1	11.9	4,269	5.3	3.0	10.4	2,120	14.7	10.8	2,850	20.8
1506	675	7.9	8.6	3,204	5.1	3.7	11.0	1,545	17.9	13.1	2,183	21.3
1507	642	7.2	9.8	3,138	3.8	3.5	8.0	1,499	18.2	10.0	2,071	31.7
1508	99	*	*	569	2.5	1.4	15.5	290	24.5	21.4	399	21.1
1509	579	9.7	9.8	3,144	4.1	3.2	12.2	1,606	17.1	16.2	2,257	21.8
1511	809	5.8	9.6	3,427	3.4	2.8	10.2	1,535	18.5	14.6	2,196	23.5
1512	384	8.9	10.7	1,967	4.5	2.6	11.2	966	15.2	12.7	1,337	32.4
1701	758	5.0	12.0	3,500	5.1	2.8	12.8	1,572	17.4	15.6	2,155	6.1
1950	727	8.4	7.0	2,998	7.2	2.0	10.9	1,221	18.1	15.7	1,915	6.1
1952	160	5.6	8.1	745	4.2	2.4	10.9	338	18.6	14.8	478	5.4
1953	310	6.5	6.5	1,546	7.0	1.9	12.4	721	13.7	15.7	1,023	5.6
1954	599	6.8	6.7	2,901	6.2	2.5	11.4	1,277	14.1	15.6	1,851	5.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 60

Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1955	141	6.4	9.9	771	5.4	1.7	13.0	379	12.7	16.4	515	6.4
1957	650	8.2	8.0	2,857	5.7	4.1	10.9	1,208	16.7	14.8	1,924	4.5
1959	44	*	*	261	5.7	3.1	13.0	127	10.2	17.3	163	3.1
2150	703	7.0	11.2	3,333	6.8	1.2	12.5	1,497	16.1	17.2	1,926	3.3
2151	276	8.3	7.6	1,232	5.0	1.5	10.1	556	16.9	14.7	762	3.0
2251	470	9.8	6.4	2,041	5.6	1.3	10.1	932	16.7	14.5	1,254	7.1
2252	201	7.5	4.0	1,030	6.4	2.0	12.8	501	24.6	18.0	691	7.5
2551	722	12.2	3.6	2,874	6.4	3.3	11.7	1,178	17.2	13.4	1,519	20.1
2552	16	*	*	28	*	*	*	4	*	*	6	*
2554	135	7.4	3.7	702	6.0	2.6	16.1	353	16.4	14.2	420	9.8
2555	7,740	11.0	5.4	29,991	6.8	3.3	10.5	11,800	15.2	13.4	17,563	17.9
3001	1,011	13.0	4.9	4,761	6.2	2.3	14.1	2,332	18.7	14.9	3,096	8.2
3002	268	9.3	7.1	1,320	5.2	1.5	14.3	661	19.8	13.0	877	17.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 61
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
3003	2	*	*	13	*	*	*	7	*	*	10	*
3004	137	8.0	2.2	825	4.6	1.2	20.6	453	20.5	28.9	563	7.6
3007	22	*	*	167	1.8	3.0	17.4	103	17.5	19.4	127	17.3
3009	8	*	*	39	*	*	*	24	*	*	37	*
3010	2	*	*	3	*	*	*	2	*	*	2	*
3201	592	9.6	11.5	2,704	6.6	3.7	13.1	1,214	14.9	15.8	1,730	9.0
3301	4,000	9.1	7.2	15,966	5.7	3.8	11.3	6,687	16.2	14.8	10,248	12.2
3302	185	8.1	8.6	995	5.2	1.9	12.7	498	20.7	17.9	684	14.5
3304	145	4.8	14.5	754	3.6	3.6	9.3	378	15.1	12.7	517	10.8
3305	170	4.7	11.2	788	3.3	3.7	8.5	373	14.7	9.9	517	6.8
3306	301	6.3	11.0	1,442	3.8	2.6	9.2	667	18.1	10.2	949	8.5
3308	84	*	*	352	4.8	1.4	9.9	159	17.0	11.3	220	8.2
3309	48	*	*	252	2.4	0.8	10.7	123	22.0	13.0	171	4.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 62
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
3312	207	11.6	7.2	1,078	6.2	1.9	13.5	537	19.9	18.4	782	11.8
3550	94	*	*	440	6.6	3.4	20.5	211	19.9	21.3	234	20.1
3551	24	*	*	131	0.0	3.1	17.6	67	*	*	64	*
3552	56	*	*	245	5.7	1.2	24.9	116	17.2	21.6	128	27.3
3701	1,234	7.5	7.1	5,388	5.0	3.6	8.9	2,427	20.9	12.3	3,630	8.2
3805	2,336	9.3	8.5	9,960	4.6	4.4	9.1	4,512	17.2	14.7	5,265	13.1
3850	1,123	10.7	6.9	4,839	5.0	3.6	9.8	2,245	17.6	14.2	3,200	16.3
3950	2,058	7.2	8.7	8,919	4.3	4.4	10.7	4,054	19.1	13.6	5,689	24.0
3952	0	*	*	7	*	*	*	6	*	*	5	*
3953	1	*	*	7	*	*	*	6	*	*	5	*
3954	45	*	*	324	5.9	2.5	10.2	182	18.1	11.0	229	15.7
4001	1,294	12.1	4.9	5,813	5.7	4.4	11.5	2,716	15.8	14.0	3,976	35.7
4002	1,172	9.3	6.0	5,345	5.4	3.8	12.4	2,494	16.1	15.5	3,692	24.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 63
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
4103	97	*	*	623	4.0	4.8	11.1	352	15.9	13.1	417	24.7
4106	2,189	11.6	5.8	9,894	7.1	1.8	11.3	4,682	16.6	15.0	6,709	17.6
4107	3,492	10.0	8.3	15,280	6.6	1.7	10.1	6,872	16.9	12.6	10,170	15.3
4109	1,199	7.3	10.2	5,918	4.9	2.7	7.5	2,932	15.8	11.4	3,976	6.8
4110	563	8.7	7.3	3,134	5.6	1.5	12.4	1,601	17.7	18.6	2,069	10.9
4112	585	5.0	8.2	3,097	6.2	4.2	17.3	1,667	18.2	22.4	2,517	11.7
4113	804	5.3	8.1	3,531	3.7	2.1	13.2	1,711	17.3	21.1	2,305	7.9
4120	2,218	8.0	9.2	10,103	6.8	1.5	11.5	4,636	16.3	13.4	6,643	11.3
4160	1,948	9.4	7.4	8,768	7.0	1.9	10.4	3,856	16.7	12.6	5,627	13.2
4351	573	8.6	9.2	2,907	6.2	2.0	10.3	1,398	19.5	13.7	1,848	13.1
4352	178	11.8	5.1	920	5.7	1.7	11.4	491	20.2	15.3	616	15.1
4353	467	10.3	9.9	2,446	6.2	3.4	12.1	1,186	20.4	16.8	1,636	14.4
4354	820	5.9	10.7	3,623	5.3	1.7	9.6	1,567	19.3	13.3	2,251	10.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 64

Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
4355	341	7.3	9.4	1,544	5.4	2.5	11.5	725	18.1	18.8	972	13.2
4356	571	7.7	6.1	2,586	6.2	1.7	11.9	1,194	20.9	16.8	1,660	17.5
4358	159	8.2	8.2	959	6.8	0.8	22.9	524	23.5	33.4	671	20.0
4359	570	7.2	10.4	2,861	5.7	1.7	14.3	1,477	19.6	16.0	1,840	14.3
4360	216	6.5	6.9	948	3.9	2.2	7.8	452	19.0	10.6	583	14.2
4361	225	9.3	5.8	1,065	6.7	1.6	10.8	495	22.0	16.4	688	14.0
4362	338	8.3	8.9	1,574	7.1	1.5	10.3	732	14.5	15.2	985	12.2
4363	952	6.9	9.2	4,253	5.1	1.7	9.0	1,918	17.7	14.4	2,697	16.4
4401	1,432	7.7	8.7	5,925	6.1	2.8	10.4	2,543	17.2	14.5	3,812	8.8
4501	1,000	6.8	7.1	4,820	5.9	1.8	8.8	2,240	14.2	12.2	3,089	8.5
4502	1	*	*	2	*	*	*	0	*	*	0	*
4591	0	*	*	1	*	*	*	1	*	*	1	*
4601	1,680	7.9	6.6	6,722	5.7	2.8	11.6	2,748	19.1	15.8	3,545	7.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 65
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)
Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
4701	1,500	6.9	9.8	6,033	5.6	1.8	9.6	2,492	17.0	15.2	3,887	6.4
5050	1,854	11.8	5.7	7,193	6.0	2.4	11.4	2,875	14.1	12.7	4,577	15.7
5051	2,989	10.3	6.2	11,260	6.5	2.3	11.7	4,478	13.4	12.3	6,930	16.6
5052	2,966	9.6	5.7	10,998	5.7	1.9	12.0	4,311	16.2	15.2	6,710	15.1
5055	206	8.7	7.3	762	4.2	1.3	12.5	364	15.9	14.8	501	29.9
5101	2,421	8.6	6.6	9,740	6.2	2.4	10.7	3,953	17.7	15.3	5,824	9.6
5202	687	8.9	7.4	2,733	6.5	2.9	12.7	1,156	18.0	18.3	1,473	7.6
5203	156	8.3	10.9	718	6.3	2.1	11.0	358	15.6	14.8	394	11.7
5204	105	11.4	8.6	482	5.2	2.1	9.5	217	21.2	13.8	264	9.8
5601	1,307	8.6	8.6	5,953	6.0	1.8	13.1	2,665	16.7	14.3	3,391	9.8
5801	2,109	9.3	7.8	8,373	7.4	2.1	10.5	3,468	17.8	14.9	5,366	8.0
5803	224	8.9	8.0	1,222	5.8	2.1	9.7	552	14.1	11.6	818	9.0
5804	61	*	*	488	4.5	3.3	9.4	242	16.5	10.7	375	6.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 66

Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
6101	85	*	*	409	7.6	2.9	10.3	200	20.5	9.0	273	12.8
6102	218	5.5	7.3	1,217	6.3	4.1	10.3	635	14.0	12.8	822	13.5
6103	3,829	9.3	6.1	16,827	4.7	3.7	9.6	7,929	17.5	12.7	11,223	21.2
6104	651	11.8	5.1	1,844	4.8	5.8	8.8	679	16.8	14.1	993	19.2
6105	47	*	*	346	3.5	2.0	8.1	189	18.5	9.5	249	11.6
6106	19	*	*	39	*	*	*	16	*	*	15	*
6301	5,170	10.2	6.1	20,017	6.6	2.0	9.1	7,889	15.7	14.0	11,845	13.1
6312	3,144	10.1	6.9	12,174	7.3	2.7	8.3	4,664	15.3	11.3	7,303	18.1
6316	1,885	9.0	7.5	6,989	5.3	2.0	9.1	2,595	14.4	13.4	4,033	11.1
6320	210	9.0	11.4	1,165	3.9	4.3	8.8	602	14.6	12.3	741	8.5
6325	141	4.3	5.0	789	6.7	3.4	10.4	389	21.1	14.9	503	7.8
6330	489	16.0	5.5	2,059	7.1	3.5	9.0	962	14.8	10.2	1,284	19.2
6335	524	7.6	9.0	2,910	3.5	3.8	8.5	1,433	14.3	11.7	1,919	10.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 67
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
6340	104	8.7	4.8	603	4.5	2.0	7.3	304	14.8	12.5	392	8.2
6345	25	*	*	147	6.1	2.0	16.3	69	*	*	91	*
6350	267	12.0	9.0	1,255	5.2	5.0	11.7	550	12.7	16.5	788	12.9
6877	5	*	*	56	*	*	*	27	*	*	34	*
6895	1	*	*	29	*	*	*	17	*	*	26	*
7304	223	3.6	8.5	1,081	4.7	2.5	13.3	518	21.0	17.2	702	3.1
7306	2,817	10.5	7.7	11,727	5.1	2.7	12.1	5,176	15.6	16.6	7,502	14.2
7308	188	8.5	11.2	874	4.5	2.6	11.6	374	18.7	12.8	553	7.8
7310	54	*	*	196	6.1	3.6	11.2	76	*	*	107	3.7
7401	2,696	8.6	6.8	11,531	6.5	2.1	8.4	4,971	15.6	13.2	7,123	6.1
7601	735	7.1	6.4	3,281	4.2	3.2	8.0	1,431	15.9	11.9	2,133	6.3
7850	1,197	7.7	6.7	5,145	4.3	5.3	10.7	2,372	17.3	16.6	3,439	4.1
7852	101	5.9	12.9	462	2.2	5.2	9.1	208	14.9	15.9	323	4.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 68
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
8101	1,024	18.7	4.3	2,279	7.5	8.5	9.6	670	10.9	12.8	1,129	29.8
8102	146	5.5	9.6	811	6.2	6.4	19.1	446	20.2	24.4	620	15.5
8105	303	10.9	6.6	778	9.1	3.5	11.2	279	18.6	13.3	414	16.9
8106	218	11.5	3.7	674	6.8	1.8	11.6	296	16.2	13.9	418	24.6
8150	58	*	*	354	4.2	2.8	6.8	177	15.8	9.6	243	18.9
8151	3,228	9.7	5.9	13,284	6.0	2.2	10.3	5,569	16.8	14.1	8,665	19.8
8152	385	12.5	6.0	1,129	8.0	5.5	10.6	305	15.4	17.4	609	38.3
8153	27	*	*	137	5.1	0.7	13.9	69	*	*	94	*
8154	242	9.1	7.0	1,100	3.5	3.6	6.3	517	14.9	8.3	647	5.9
8155	75	*	*	435	2.5	2.3	10.8	234	14.1	13.2	309	21.0
8156	10	*	*	79	*	*	*	46	*	*	59	*
8157	10	*	*	117	0.9	1.7	7.7	81	*	*	93	*
8201	3,004	9.5	6.9	9,431	6.9	2.5	10.1	2,810	17.2	12.8	4,796	6.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 69
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
8202	3,427	7.4	7.0	15,288	4.8	1.9	10.6	7,112	16.5	12.6	9,943	27.3
8203	411	8.8	6.8	1,569	7.2	1.8	9.2	575	20.7	9.9	846	14.1
8250	1,143	12.2	5.6	3,817	6.3	3.3	7.9	1,217	12.3	10.0	2,036	22.0
8252	81	*	*	302	13.6	5.6	10.9	150	18.0	18.0	202	37.6
8255	3,332	9.4	8.1	10,975	7.8	1.3	10.3	3,445	19.7	12.4	5,747	12.6
8257	417	11.3	5.0	2,192	5.4	4.7	9.9	1,000	12.1	14.0	1,333	13.6
8258	1,176	9.5	6.1	5,227	6.8	1.8	12.5	2,031	18.7	14.9	2,984	13.9
8260	852	10.9	3.4	3,542	5.4	3.1	7.8	1,589	14.7	10.3	2,393	18.6
8901	1,261	13.3	3.3	5,094	5.8	5.7	9.3	2,190	14.9	11.8	3,320	36.0
8902	925	12.5	3.1	3,584	5.7	7.2	8.8	1,474	11.9	10.9	2,251	33.8
8903	1,356	13.3	3.2	5,822	6.6	5.1	12.1	2,542	15.9	14.2	3,795	38.6
8904	1,360	11.9	5.4	5,253	4.1	6.8	8.8	2,175	14.1	11.3	3,328	33.7
8905	3,024	7.9	6.3	14,079	4.7	7.0	12.3	6,528	16.6	14.4	9,752	24.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 70
Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
9004	1	*	*	1	*	*	*	0	*	*	0	*
9010	177	14.1	4.5	153	5.9	15.0	1.3	0	*	*	0	*
9012	48	*	*	275	2.9	4.4	8.4	146	19.9	8.9	87	*
9013	97	*	*	420	5.7	4.0	7.9	177	14.1	7.3	138	35.5
9015	37	*	*	166	7.8	4.8	16.9	89	*	*	79	*
9016	192	16.1	2.1	764	6.8	3.3	9.0	324	22.5	8.6	419	41.1
9017	76	*	*	362	8.8	1.9	15.5	174	26.4	15.5	125	23.2
9018	147	20.4	3.4	630	8.9	4.0	9.5	280	19.6	15.0	60	*
9019	44	*	*	228	8.3	3.9	11.0	117	12.8	13.7	55	*
9021	87	*	*	338	7.1	3.8	14.5	146	25.3	13.0	44	*
9040	0	*	*	1	*	*	*	1	*	*	0	*
9041	41	*	*	373	5.1	4.3	11.0	210	17.1	11.9	103	32.0
9045	1,651	10.7	7.2	7,324	5.7	1.4	13.9	3,412	17.8	17.6	4,724	21.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Page: 71

Run Date: 04/02/2014

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
9050	1,612	11.1	6.1	7,577	6.4	3.1	12.7	3,437	15.2	15.9	4,990	16.5
9051	1,600	7.8	7.9	7,846	6.1	1.3	16.7	3,687	18.8	21.1	5,520	18.4
9053	2,571	14.7	3.7	9,125	7.0	3.2	8.6	3,475	15.7	12.1	5,670	24.9
9055	147	8.2	4.8	132	8.3	12.9	6.8	2	*	*	2	*
9058	722	8.9	4.4	3,865	3.8	2.9	8.7	1,922	13.3	10.7	2,676	21.6
9066	1,964	13.1	4.1	8,177	5.6	6.5	7.0	3,414	7.9	7.4	5,203	12.2
9068	1,423	12.5	3.0	5,679	6.5	9.9	7.1	2,404	7.2	5.3	3,692	27.2
9069	2,686	12.6	4.6	9,276	7.0	3.3	8.9	3,306	14.3	12.3	5,549	23.8
9070	1,795	13.7	2.8	7,332	8.3	8.0	11.0	3,104	7.3	6.5	4,431	23.9
9072	1,616	12.7	3.2	6,012	7.7	3.1	9.6	2,357	14.8	11.1	3,656	14.6
9074	754	13.4	4.1	3,341	7.8	3.0	9.4	1,349	13.6	11.5	2,100	15.5
9080	2,674	12.5	4.2	11,879	5.6	2.5	8.6	5,401	14.7	12.4	7,879	33.6
Michigan	177,968	9.5	7.0	757,067	5.9	3.0	10.7	331,475	16.4	14.0	477,486	15.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
0101	118	71.2	85	*	82	*	64	*	82	*	320	86.3	599	7.0
0102	494	66.4	360	17.2	296	8.1	267	18.7	331	6.0	903	84.7	1,950	9.0
0103	433	61.0	341	18.2	297	7.7	254	17.7	315	7.0	964	84.4	1,938	10.4
0104	146	63.0	101	15.8	95	*	81	*	97	*	307	79.2	638	15.5
0250	156	67.9	125	24.0	120	10.8	87	*	113	9.7	406	86.0	785	7.6
0251	111	66.7	92	*	65	*	70	*	88	*	303	87.5	571	6.8
0252	112	72.3	78	*	92	*	42	*	71	*	385	84.2	650	8.2
0253	169	59.8	127	16.5	107	10.3	101	16.8	119	7.6	450	86.7	846	13.2
0401	544	52.8	399	18.5	404	11.6	291	24.4	355	10.4	1,218	91.7	2,428	5.6
0402	379	64.9	302	20.9	269	10.0	213	26.3	260	7.7	965	85.4	1,845	7.5
0403	142	67.6	96	*	92	*	79	*	88	*	377	84.1	676	8.0
0404	145	49.0	100	17.0	112	10.7	78	*	86	*	381	82.7	718	18.5
0501	583	72.6	449	27.8	405	12.1	324	27.5	412	13.8	1,323	90.2	2,633	14.4
0502	1	*	2	*	3	*	0	*	2	*	8	*	14	*
0503	179	79.9	149	32.9	126	14.3	101	33.7	140	12.1	397	90.2	819	4.8
0504	597	79.6	453	33.3	376	17.0	332	33.4	399	14.8	1,301	91.6	2,643	5.6
0505	421	77.4	269	31.6	240	16.3	231	37.2	251	12.7	878	89.1	1,754	5.8

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
0506	190	80.0	151	16.6	146	11.6	101	23.8	138	7.2	582	86.4	1,006	14.2
0507	39	*	48	*	50	*	36	*	42	*	185	91.4	304	7.6
0650	1,059	73.5	822	24.5	797	11.8	577	23.1	748	11.4	2,366	85.2	4,734	7.5
0651	668	59.0	491	16.7	535	7.7	357	14.0	430	3.5	1,584	89.6	3,076	12.5
0652	293	59.0	222	16.7	245	3.7	149	18.1	187	5.9	864	87.8	1,570	10.5
0653	435	69.4	331	24.5	353	11.9	224	25.0	296	13.9	1,270	89.8	2,266	14.9
0654	413	69.2	297	22.6	335	11.3	218	23.9	272	11.0	1,064	92.1	2,005	12.6
0655	407	72.7	329	20.1	299	10.7	214	20.1	292	10.6	889	90.3	1,799	6.6
0657	69	*	46	*	57	*	38	*	39	*	211	96.2	380	17.6
0701	457	81.4	333	37.8	296	16.9	261	21.5	308	11.7	1,057	95.9	2,091	0.8
0702	95	*	74	*	67	*	45	*	61	*	230	80.4	450	3.3
0703	230	62.6	181	23.2	131	9.9	121	24.8	169	11.2	552	90.6	1,068	7.2
0704	75	*	45	*	56	*	30	*	42	*	253	86.6	424	11.1
0705	141	74.5	116	21.6	97	*	80	*	110	8.2	419	94.7	779	2.2
0850	1,213	72.0	957	21.1	863	9.5	664	19.3	872	10.1	2,498	87.0	5,241	4.6
0851	932	66.2	786	20.2	682	8.9	518	22.0	718	11.4	2,304	88.4	4,417	4.9
0901	1,814	56.1	1,383	14.6	1,383	6.6	1,009	15.0	1,207	3.6	4,094	94.2	8,214	7.1

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
0902	71	*	61	*	59	*	46	*	54	*	230	88.3	425	9.9
1051	1,582	81.4	1,033	37.6	1,026	18.4	661	35.6	938	14.9	3,067	91.8	6,383	3.4
1101	2,042	47.7	1,263	15.8	1,329	8.1	846	14.3	1,095	5.2	4,156	74.7	8,531	7.8
1102	1,025	52.7	681	17.3	673	10.3	476	15.1	590	6.9	1,891	81.8	4,078	9.0
1106	116	69.8	89	*	85	*	62	*	77	*	249	87.1	523	7.3
1250	1,056	61.6	840	20.0	769	11.6	603	19.4	772	10.1	2,765	82.2	5,303	7.3
1251	922	72.5	655	23.8	571	10.9	522	22.2	593	13.3	2,128	89.9	4,201	6.8
1252	72	*	101	24.7	87	*	59	*	81	*	248	89.1	461	3.3
1253	1,518	67.3	1,075	21.7	948	9.9	838	16.7	994	11.8	3,534	87.6	6,873	4.8
1350	2,639	55.7	1,860	14.6	1,683	6.4	1,335	11.2	1,538	4.6	5,765	89.4	11,543	4.4
1351	149	76.5	116	20.7	102	13.7	89	*	110	5.5	386	94.0	716	3.1
1352	436	52.3	297	10.1	303	7.3	222	9.0	278	3.2	1,120	86.6	2,139	9.2
1401	223	69.1	150	25.3	157	10.2	101	21.8	131	9.9	597	88.9	1,132	4.5
1402	153	83.7	114	41.2	116	24.1	80	*	107	17.8	454	87.4	830	5.8
1403	27	*	17	*	21	*	15	*	16	*	89	*	152	5.9
1404	11	*	13	*	11	*	8	*	13	*	42	*	76	*
1501	1,147	70.4	802	24.9	813	13.9	635	23.3	710	9.3	3,000	85.2	5,571	5.9

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
1503	1,264	67.2	967	22.3	907	12.5	709	17.6	857	9.9	3,250	87.1	6,098	5.5
1504	2,586	71.9	1,629	27.3	1,888	11.7	1,044	23.0	1,344	9.1	6,064	86.0	11,793	3.3
1505	879	76.9	618	36.9	646	16.9	445	29.4	554	17.1	2,331	91.3	4,333	2.1
1506	678	72.3	507	23.7	510	12.2	394	21.6	452	11.1	1,715	93.0	3,253	4.3
1507	674	74.3	475	26.9	527	14.0	361	28.3	444	14.6	1,635	89.1	3,190	4.2
1508	103	76.7	71	*	94	*	38	*	55	*	318	93.4	590	0.5
1509	594	67.2	457	27.6	479	11.1	313	16.9	400	8.5	1,779	84.3	3,204	6.6
1511	819	72.6	652	18.1	528	9.3	490	20.2	607	4.6	1,713	90.3	3,472	3.9
1512	392	68.9	311	23.5	283	10.2	222	22.5	276	13.0	1,061	94.2	2,003	3.7
1701	757	67.9	567	28.7	503	14.1	427	33.3	476	20.2	1,784	86.8	3,571	7.7
1950	727	71.4	521	20.7	423	8.5	424	20.0	490	9.2	1,422	88.9	3,040	8.8
1952	160	71.3	115	19.1	104	13.5	80	*	106	8.5	376	87.0	755	7.7
1953	309	69.6	235	19.6	212	6.6	173	19.1	224	9.4	819	85.6	1,565	7.2
1954	623	68.9	528	22.3	421	10.2	389	22.4	481	11.0	1,489	89.5	2,963	6.9
1955	143	71.3	118	27.1	105	18.1	79	*	100	11.0	426	86.4	780	10.3
1957	656	77.7	550	23.8	461	15.2	425	21.6	506	7.5	1,359	86.9	2,896	3.9
1959	47	*	37	*	28	*	26	*	34	*	147	85.7	262	5.7

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
2150	703	67.1	574	19.7	533	9.0	417	17.0	526	6.5	1,739	92.3	3,366	5.9
2151	276	62.7	202	16.3	190	9.5	163	17.2	188	5.3	620	89.8	1,227	6.4
2251	474	65.2	378	24.3	329	13.1	265	26.8	344	9.9	1,044	89.5	2,082	4.4
2252	201	62.7	150	18.0	144	13.9	104	26.0	136	9.6	581	88.1	1,053	6.0
2551	724	36.7	467	8.8	375	2.9	326	5.8	455	1.8	1,288	72.7	2,945	9.5
2552	16	*	9	*	3	*	6	*	6	*	4	*	28	*
2554	136	41.2	101	9.9	92	*	80	*	87	*	401	90.3	702	12.0
2555	7,761	52.4	6,141	12.4	4,509	5.9	4,416	9.6	5,364	4.1	13,204	78.8	30,306	6.5
3001	1,010	53.2	746	8.0	629	3.7	537	5.2	691	1.6	2,654	78.6	4,906	11.1
3002	269	53.5	194	5.2	163	3.1	136	2.9	184	2.2	746	94.0	1,361	6.0
3003	2	*	1	*	1	*	0	*	1	*	7	*	13	*
3004	139	78.4	100	28.0	99	*	74	*	96	*	541	71.7	871	0.9
3007	22	*	15	*	18	*	9	*	13	*	118	85.6	173	0.6
3009	9	*	7	*	6	*	4	*	6	*	28	*	53	*
3010	2	*	1	*	0	*	1	*	1	*	2	*	4	*
3201	592	55.6	426	16.4	388	8.2	323	18.6	378	5.3	1,358	95.4	2,742	6.1
3301	4,044	67.3	2,939	19.8	2,517	10.1	2,012	11.4	2,545	5.2	7,379	80.4	16,159	5.8

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
3302	185	60.0	163	17.8	144	9.0	106	4.7	154	3.2	543	79.0	1,016	3.8
3304	146	92.5	107	64.5	104	44.2	66	*	90	*	429	80.4	763	0.8
3305	170	78.2	114	28.9	123	17.9	78	*	102	8.8	407	89.2	795	1.8
3306	305	72.1	245	18.4	217	9.7	177	13.6	226	4.4	733	92.1	1,447	5.3
3308	85	*	56	*	59	*	42	*	51	*	172	88.4	355	3.1
3309	48	*	39	*	37	*	19	*	36	*	140	85.7	254	5.9
3312	212	67.0	180	12.2	176	9.7	127	9.4	153	6.5	590	79.2	1,099	4.1
3550	94	*	70	*	57	*	60	*	61	*	240	90.4	446	6.1
3551	24	*	18	*	13	*	21	*	18	*	75	*	132	7.6
3552	56	*	31	*	38	*	21	*	30	*	134	89.6	247	2.0
3701	1,239	68.0	950	18.5	921	9.2	695	19.4	829	11.0	2,775	88.4	5,489	7.6
3805	2,344	60.3	1,898	14.4	1,544	7.8	1,339	12.6	1,804	5.7	5,017	83.8	10,224	7.5
3850	1,131	63.6	943	17.9	833	9.4	547	19.6	750	8.8	2,515	84.8	5,038	7.1
3950	2,073	66.8	1,468	28.1	1,356	14.2	1,102	21.8	1,335	10.6	4,551	84.3	9,076	4.9
3952	0	*	0	*	0	*	0	*	0	*	7	*	7	*
3953	1	*	0	*	0	*	1	*	0	*	6	*	7	*
3954	45	*	38	*	43	*	21	*	32	*	205	87.8	327	2.8

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
4001	1,337	60.2	935	15.7	838	7.6	704	15.2	791	7.6	3,039	65.8	5,986	7.9
4002	1,179	62.8	840	22.0	730	11.0	686	16.2	767	6.3	2,808	73.3	5,475	6.9
4103	98	*	74	*	87	*	58	*	66	*	386	77.7	628	7.5
4106	2,201	64.4	1,691	19.3	1,498	8.1	1,188	10.4	1,419	6.5	5,211	80.6	10,048	4.4
4107	3,503	66.7	2,849	21.9	2,409	10.0	1,981	19.5	2,473	11.4	7,704	87.6	15,539	3.3
4109	1,207	66.3	965	21.8	899	12.5	689	21.0	835	11.6	3,230	91.5	6,005	4.7
4110	567	63.8	428	21.7	404	9.9	266	6.4	362	2.8	1,736	79.3	3,191	5.0
4112	586	35.8	404	19.3	384	14.3	331	5.1	372	1.9	1,869	96.7	3,168	5.2
4113	804	68.8	494	19.0	442	12.4	430	7.4	440	3.9	1,878	83.2	3,580	1.1
4120	2,235	66.4	1,846	19.6	1,485	9.5	1,292	14.2	1,632	7.5	5,166	86.9	10,234	2.0
4160	1,975	69.0	1,686	27.1	1,336	15.3	1,156	21.1	1,420	14.7	4,329	86.6	8,911	2.3
4351	578	67.1	421	20.2	416	10.8	327	18.7	394	12.2	1,592	91.1	2,960	10.5
4352	179	66.5	121	16.5	117	8.5	89	*	106	6.6	556	79.9	957	18.1
4353	469	67.6	386	19.9	388	11.6	286	15.4	354	9.6	1,366	77.7	2,530	9.0
4354	821	69.8	617	20.9	592	9.1	513	23.6	569	14.4	1,782	95.2	3,662	5.2
4355	342	70.5	230	22.2	241	8.3	178	21.3	204	2.9	823	89.1	1,592	7.9
4356	574	69.7	417	23.7	437	12.6	325	21.2	370	7.3	1,393	90.7	2,690	6.8

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
4358	159	78.6	124	32.3	142	25.4	97	*	115	2.6	604	89.7	997	1.0
4359	570	76.8	353	28.0	416	14.7	253	16.2	322	5.9	1,666	79.8	2,926	6.4
4360	216	78.2	164	20.1	139	13.7	131	13.7	159	6.9	511	87.7	963	11.1
4361	228	73.2	150	25.3	166	9.0	89	*	140	11.4	579	86.5	1,092	7.3
4362	340	79.4	228	27.2	240	14.2	131	24.4	211	10.0	834	89.6	1,595	8.9
4363	951	75.0	743	18.7	649	8.9	557	15.1	717	5.3	2,172	87.9	4,328	7.6
4401	1,433	65.5	1,065	21.3	974	10.5	787	15.1	988	7.7	2,868	92.7	6,042	4.9
4501	1,004	57.7	792	16.7	686	7.3	556	14.7	702	4.8	2,561	88.1	4,878	6.7
4502	1	*	1	*	0	*	1	*	1	*	1	*	2	*
4591	0	*	0	*	0	*	0	*	0	*	1	*	1	*
4601	1,716	69.2	1,260	21.5	1,087	10.7	933	15.9	1,163	7.3	3,136	76.3	6,574	5.1
4701	1,498	69.8	1,047	25.9	1,037	11.3	745	21.2	932	10.9	2,820	90.7	6,121	4.4
5050	1,861	55.0	1,257	13.0	1,236	6.6	954	9.7	1,130	4.4	3,246	81.4	7,371	4.7
5051	2,992	56.1	1,969	15.9	1,863	6.6	1,592	10.6	1,772	6.2	5,121	79.1	11,592	4.5
5052	2,971	53.3	1,932	12.9	1,844	6.3	1,507	7.9	1,735	4.7	4,920	78.2	11,269	4.2
5055	222	38.7	132	14.4	103	8.7	64	*	78	*	393	65.9	783	3.3
5101	2,458	66.7	1,782	19.9	1,689	10.3	1,365	13.5	1,604	5.9	4,466	85.9	9,862	3.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
5202	689	71.6	488	24.2	464	13.1	369	20.3	461	9.3	1,354	88.3	2,872	3.7
5203	157	65.0	102	23.5	125	11.2	77	*	97	*	408	86.8	758	5.4
5204	105	73.3	78	*	87	*	55	*	72	*	259	90.0	503	4.2
5601	1,306	73.9	995	26.6	911	9.8	830	25.4	921	11.3	3,023	89.1	5,958	4.6
5801	2,120	62.7	1,606	15.0	1,391	7.6	1,172	13.2	1,405	6.4	3,951	82.1	8,521	7.4
5803	224	66.5	205	22.4	205	10.2	147	23.1	188	10.1	639	87.5	1,227	5.9
5804	61	*	78	*	85	*	51	*	74	*	280	83.6	494	3.8
6101	85	*	50	*	58	*	43	*	44	*	235	89.8	424	12.7
6102	219	68.0	159	28.9	181	15.5	143	25.2	145	9.0	713	88.8	1,243	6.6
6103	3,839	54.4	2,523	14.8	2,550	6.3	1,955	13.6	2,190	5.3	8,686	86.0	17,141	6.7
6104	654	43.0	339	9.4	288	3.8	265	9.4	279	2.2	729	77.2	1,882	12.2
6105	47	*	46	*	54	*	33	*	45	*	210	90.0	353	7.1
6106	19	*	6	*	1	*	6	*	6	*	15	*	39	*
6301	5,205	54.0	3,604	17.1	2,730	8.4	3,011	10.2	3,360	5.4	8,901	81.9	20,340	3.7
6312	3,167	64.2	2,408	24.4	1,737	13.0	1,881	14.9	2,231	8.3	5,264	78.3	12,318	3.1
6316	1,886	59.9	1,361	20.2	1,058	9.6	1,078	14.7	1,261	7.4	2,966	82.0	7,076	2.6
6320	215	56.7	178	26.4	141	18.4	121	17.4	165	13.3	687	92.6	1,198	1.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
6325	143	60.8	109	21.1	98	*	87	*	98	*	440	87.3	803	5.6
6330	503	49.3	333	10.8	253	3.6	245	7.3	275	3.3	1,046	77.6	2,116	7.7
6335	558	53.2	439	22.6	377	9.0	364	12.9	414	7.5	1,638	93.2	2,967	1.0
6340	108	50.0	89	*	81	*	65	*	84	*	341	95.9	607	1.5
6345	25	*	27	*	18	*	22	*	26	*	77	*	149	1.3
6350	283	51.2	178	20.2	178	6.2	158	8.9	166	4.8	617	90.6	1,282	2.9
6877	5	*	10	*	9	*	5	*	9	*	28	*	55	*
6895	1	*	3	*	9	*	0	*	2	*	17	*	29	*
7304	225	54.7	157	15.9	147	4.1	112	8.0	122	5.7	591	88.7	1,101	9.9
7306	2,827	53.4	1,769	16.2	1,677	4.7	1,449	11.0	1,460	5.9	5,870	80.4	11,898	7.9
7308	185	58.9	143	21.0	132	8.3	101	20.8	105	10.5	435	86.0	875	7.0
7310	54	*	36	*	33	*	21	*	27	*	84	*	202	16.8
7401	2,696	56.5	2,177	13.8	1,880	6.5	1,457	12.5	1,954	4.9	5,694	86.8	11,709	9.0
7601	735	63.9	582	21.0	459	8.5	427	17.1	548	6.2	1,617	85.7	3,297	7.8
7850	1,201	66.8	928	21.0	834	7.9	660	13.5	783	3.4	2,650	92.0	5,268	8.9
7852	101	75.2	74	*	76	*	65	*	68	*	243	96.7	477	6.5
8101	1,027	45.8	476	11.8	297	5.7	443	5.4	424	2.1	735	72.5	2,377	7.1

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
8102	146	39.0	97	*	80	*	81	*	89	*	514	79.4	845	4.6
8105	307	64.2	119	17.6	98	*	115	14.8	105	5.7	307	84.7	805	4.7
8106	219	47.9	94	*	76	*	103	8.7	84	*	315	63.2	701	5.1
8150	58	*	48	*	50	*	39	*	46	*	207	90.8	359	3.1
8151	3,250	71.5	2,463	32.0	2,172	17.0	1,864	17.7	2,264	11.7	6,304	82.1	13,471	6.8
8152	390	67.7	266	14.7	202	6.9	198	12.1	236	5.9	347	86.5	1,147	10.0
8153	27	*	22	*	25	*	16	*	18	*	76	*	139	10.1
8154	242	76.9	161	59.6	148	42.6	145	45.5	157	38.9	579	95.7	1,077	1.3
8155	75	*	61	*	56	*	51	*	57	*	263	90.1	438	6.6
8156	10	*	12	*	10	*	6	*	11	*	51	*	80	*
8157	10	*	14	*	11	*	8	*	12	*	89	*	119	5.9
8201	3,046	56.2	2,050	17.6	1,755	7.7	1,477	11.6	1,875	5.5	3,361	85.0	9,782	4.0
8202	3,572	60.1	2,306	29.1	2,054	9.8	1,786	5.5	2,097	3.6	7,925	79.6	15,382	4.1
8203	418	54.3	310	18.4	249	10.8	237	13.9	285	4.9	640	83.6	1,577	6.0
8250	1,145	45.7	821	13.3	724	7.0	490	9.6	604	5.8	1,391	91.2	3,868	4.4
8252	82	*	46	*	41	*	38	*	39	*	165	60.6	325	5.8
8255	3,363	51.4	2,242	15.0	1,837	6.7	1,700	10.4	2,141	5.0	3,980	82.2	11,123	5.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
8257	475	52.6	345	26.1	333	13.2	261	2.7	309	2.3	1,153	93.1	2,235	1.0
8258	1,252	48.5	950	13.2	813	5.4	727	8.9	892	4.4	2,318	85.8	5,300	6.1
8260	866	72.9	589	31.6	590	20.5	471	6.6	558	2.7	1,813	73.3	3,651	4.0
8901	1,264	32.0	840	9.5	641	2.2	694	4.6	747	2.3	2,452	66.3	5,246	10.9
8902	929	37.6	639	11.4	514	4.3	530	5.5	567	2.3	1,631	69.1	3,688	8.6
8903	1,362	26.0	919	8.2	749	2.9	778	2.3	838	1.2	2,854	69.5	5,973	10.5
8904	1,370	43.4	929	14.3	728	6.5	755	5.2	821	2.9	2,442	72.8	5,394	9.6
8905	3,039	39.6	2,062	12.5	1,808	4.9	1,636	2.7	1,881	1.9	7,556	77.2	14,445	6.5
9004	1	*	0	*	0	*	0	*	0	*	0	*	1	*
9010	177	32.2	21	*	0	*	24	*	14	*	0	*	176	8.5
9012	49	*	37	*	32	*	27	*	37	*	166	77.1	283	9.5
9013	98	*	72	*	71	*	55	*	65	*	198	62.6	425	12.5
9015	37	*	18	*	23	*	18	*	18	*	98	*	174	18.4
9016	192	48.4	131	5.3	93	*	111	2.7	122	0.8	367	61.3	779	14.5
9017	77	*	49	*	46	*	29	*	43	*	193	72.5	373	9.9
9018	147	51.7	100	8.0	89	*	77	*	85	*	312	60.6	638	7.8
9019	44	*	31	*	32	*	20	*	26	*	126	65.9	231	14.7

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2010 - 2012 Pediatric Nutrition Surveillance (1)
Michigan

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
9021	87	*	51	*	36	*	43	*	47	*	168	76.2	340	8.5
9040	0	*	0	*	0	*	0	*	0	*	1	*	1	*
9041	41	*	27	*	40	*	20	*	26	*	236	60.2	382	12.3
9045	1,675	45.4	1,195	9.9	1,147	3.9	859	3.1	948	1.9	3,929	70.8	7,568	6.4
9050	1,642	37.3	1,250	13.5	1,061	5.3	917	3.9	1,120	1.8	3,934	64.0	7,731	8.0
9051	1,669	53.3	1,173	19.3	1,033	7.8	859	3.8	1,043	2.9	4,315	77.7	8,145	2.8
9053	2,598	41.5	1,828	11.3	1,358	6.0	1,394	6.2	1,650	4.2	3,896	76.1	9,356	19.3
9055	147	15.0	6	*	0	*	6	*	6	*	2	*	150	7.3
9058	751	63.4	594	19.7	564	6.9	463	3.7	558	1.8	2,191	87.2	3,925	6.5
9066	1,974	35.6	1,434	5.6	1,201	1.7	1,167	3.0	1,353	2.1	3,826	71.4	8,279	8.6
9068	1,428	39.8	982	9.9	833	3.7	789	6.0	919	3.5	2,725	70.7	5,826	14.4
9069	2,705	50.5	1,858	11.2	1,417	4.4	1,481	4.9	1,756	1.9	3,744	70.3	9,478	9.8
9070	1,823	38.0	1,183	9.3	1,105	2.6	872	7.5	1,029	2.9	3,476	61.0	7,460	9.3
9072	1,638	40.6	1,100	8.2	909	3.2	881	4.8	973	2.0	2,727	64.0	6,306	10.8
9074	763	50.8	666	11.3	489	5.1	521	4.4	657	1.4	1,542	68.7	3,408	10.2
9080	2,684	51.2	1,929	21.9	1,807	7.1	1,443	5.3	1,637	2.3	5,988	71.8	12,029	6.0
Michigan	179,867	58.8	129,914	18.8	114,932	9.0	96,818	13.5	116,216	6.6	373,467	82.6	771,257	6.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.