

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0101	158	90.5	0.0	5.7	0.0	0.0	3.8	0.0
0102	689	88.2	0.1	3.9	0.4	0.0	7.3	0.0
0103	608	88.5	0.2	4.4	0.0	0.0	6.9	0.0
0104	223	93.3	0.0	3.6	0.0	0.0	3.1	0.0
0250	204	87.3	0.0	0.0	2.9	0.0	9.8	0.0
0251	135	87.4	1.5	0.0	5.2	0.0	5.9	0.0
0252	199	32.7	0.5	1.0	4.0	0.0	61.8	0.0
0253	232	72.4	1.3	0.0	9.5	1.7	15.1	0.0
0401	731	91.8	0.0	2.6	0.0	0.0	5.6	0.0
0402	554	80.9	0.5	3.4	0.2	0.5	14.4	0.0
0403	176	94.3	0.6	1.1	0.6	0.0	2.8	0.6
0404	207	84.5	1.4	3.9	0.0	1.0	9.2	0.0
0501	792	94.7	0.1	1.1	0.3	0.6	3.2	0.0
0502	2	*	*	*	*	*	*	*
0503	193	86.5	0.0	6.7	0.5	0.0	6.2	0.0
0504	721	81.6	2.4	3.3	9.3	1.2	2.2	0.0
0505	427	83.8	0.2	6.1	0.7	1.2	8.0	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0506	312	92.0	0.0	2.2	0.0	0.0	5.8	0.0
0507	262	88.2	0.0	2.3	0.4	0.0	9.2	0.0
0508	100	78.0	0.0	2.0	9.0	0.0	11.0	0.0
0509	10	*	*	*	*	*	*	*
0650	1,500	78.5	3.9	4.7	1.5	1.6	9.7	0.1
0651	894	87.9	0.0	6.2	0.0	0.2	5.5	0.2
0652	493	90.1	0.6	3.0	0.0	0.2	6.1	0.0
0653	742	95.3	0.4	1.6	0.4	0.0	2.3	0.0
0654	566	92.9	0.7	1.8	0.0	0.0	4.6	0.0
0655	538	88.5	0.0	3.7	0.0	0.4	7.4	0.0
0657	101	89.1	1.0	5.0	0.0	0.0	5.0	0.0
0701	639	85.3	0.3	1.3	0.0	3.6	9.4	0.2
0702	134	87.3	0.0	0.0	0.0	0.0	12.7	0.0
0703	358	88.3	0.8	3.9	0.6	0.3	6.1	0.0
0704	87	*	*	*	*	*	*	*
0705	189	91.5	0.0	1.1	0.0	0.0	7.4	0.0
0850	1,478	80.2	2.9	9.7	0.0	0.7	6.4	0.1

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
0851	1,270	93.3	0.6	3.5	0.0	0.0	2.6	0.1
0901	2,492	70.8	2.1	14.8	0.0	0.2	12.0	0.1
0902	74	*	*	*	*	*	*	*
1051	1,894	84.1	1.0	7.5	0.4	0.2	6.7	0.3
1101	2,533	33.6	51.2	7.9	0.2	0.5	6.0	0.7
1102	1,256	66.1	13.5	8.4	0.2	1.0	10.4	0.4
1106	179	81.0	1.1	12.3	0.0	0.6	5.0	0.0
1250	1,508	80.6	0.6	13.8	0.0	0.1	4.7	0.1
1251	1,322	85.6	0.2	7.5	0.1	0.1	6.2	0.4
1252	724	49.7	0.8	43.5	0.0	0.0	5.5	0.4
1253	1,490	74.5	6.0	10.8	0.0	0.0	8.7	0.1
1350	3,706	47.4	19.4	10.9	0.2	7.3	14.9	0.1
1351	255	87.8	1.6	4.7	0.8	0.4	4.7	0.0
1352	590	52.4	28.0	8.3	0.0	0.2	11.2	0.0
1401	361	86.7	0.3	9.1	0.3	0.0	3.3	0.3
1402	241	45.2	0.8	29.0	9.5	0.4	14.1	0.8
1403	31	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
1404	6	*	*	*	*	*	*	*
1501	1,483	42.8	5.9	42.9	0.1	0.1	5.9	2.4
1503	1,860	49.6	6.6	31.9	0.1	0.3	9.4	2.1
1504	3,407	39.6	4.7	47.7	0.0	2.9	4.9	0.2
1505	1,136	79.8	1.5	13.7	0.1	0.2	4.6	0.1
1506	1,040	74.6	0.5	19.5	0.0	0.1	4.4	0.9
1507	884	80.8	1.8	9.0	0.2	1.5	6.4	0.2
1508	141	0.7	0.0	93.6	0.0	0.0	0.0	5.7
1509	858	46.7	2.7	45.6	0.1	0.0	2.7	2.2
1511	1,080	90.8	1.3	3.3	0.0	0.1	4.4	0.1
1512	637	81.6	0.6	8.3	0.3	0.0	8.9	0.2
1701	1,095	45.8	0.3	4.0	17.4	0.6	31.7	0.1
1950	919	88.9	0.0	7.2	0.2	0.0	3.4	0.3
1952	174	90.2	0.0	5.2	0.0	0.0	4.6	0.0
1953	504	81.2	0.2	11.5	0.2	0.2	6.7	0.0
1954	957	77.3	0.3	18.2	0.0	0.0	4.2	0.0
1955	189	70.4	0.0	24.3	0.0	0.0	4.8	0.5

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
1957	931	77.1	0.5	15.7	0.0	0.4	6.2	0.0
2150	1,097	81.0	0.0	3.7	0.1	0.1	15.0	0.0
2151	378	85.2	0.0	4.2	1.1	0.0	9.5	0.0
2251	652	89.1	0.2	3.5	0.0	0.0	6.7	0.5
2252	300	95.0	0.0	2.0	0.7	0.0	2.3	0.0
2554	211	83.4	5.7	5.7	0.0	0.0	5.2	0.0
2555	11,081	41.6	37.6	7.1	0.0	0.3	13.2	0.1
3001	1,533	11.2	62.5	20.0	0.1	0.7	5.6	0.0
3002	460	14.3	48.3	30.0	0.0	0.0	7.4	0.0
3004	219	4.1	0.0	95.9	0.0	0.0	0.0	0.0
3007	33	*	*	*	*	*	*	*
3009	15	*	*	*	*	*	*	*
3201	732	88.8	0.3	6.8	0.0	0.0	4.1	0.0
3301	5,100	27.1	29.4	21.4	0.0	6.4	15.6	0.0
3304	240	35.0	12.9	10.0	0.0	29.6	12.5	0.0
3305	325	58.8	11.1	12.9	0.0	4.6	12.6	0.0
3306	539	79.6	2.4	8.9	0.0	0.0	9.1	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
3308	151	81.5	0.0	11.9	0.7	0.0	6.0	0.0
3309	70	*	*	*	*	*	*	*
3312	770	23.2	30.8	27.4	0.1	1.9	16.4	0.1
3550	129	0.0	0.0	4.7	89.1	0.0	6.2	0.0
3551	40	*	*	*	*	*	*	*
3552	76	*	*	*	*	*	*	*
3701	1,665	82.2	0.4	11.7	0.0	0.2	5.5	0.0
3850	4,711	62.1	13.3	8.4	0.1	0.3	15.7	0.0
3950	2,737	54.8	18.2	11.8	0.0	0.8	14.4	0.0
3954	21	*	*	*	*	*	*	*
4001	1,934	22.2	56.4	8.2	0.2	1.4	11.5	0.1
4002	1,705	42.6	24.5	18.4	0.1	0.4	14.0	0.0
4003	3	*	*	*	*	*	*	*
4004	1	*	*	*	*	*	*	*
4103	106	3.8	76.4	12.3	0.0	1.9	5.7	0.0
4106	3,252	10.5	38.4	42.4	0.0	0.6	7.9	0.2
4107	5,155	38.6	22.5	22.4	0.1	1.7	14.6	0.1

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
4109	1,678	80.5	1.2	8.2	0.0	0.1	10.0	0.1
4112	806	0.1	0.4	98.5	0.0	0.6	0.4	0.0
4113	1,213	5.7	12.4	77.4	0.0	2.1	2.4	0.0
4115	5,571	34.1	21.8	22.8	0.0	11.0	10.2	0.1
4351	892	85.0	0.2	7.5	0.1	0.1	6.6	0.4
4352	293	62.1	12.6	8.2	0.0	0.0	17.1	0.0
4353	686	58.6	0.1	35.4	0.3	0.0	4.8	0.7
4354	1,142	83.9	4.6	3.9	0.0	0.2	7.3	0.3
4355	489	74.0	1.0	8.6	1.0	0.0	15.1	0.2
4356	749	77.0	0.7	14.8	0.1	0.1	7.2	0.0
4358	261	16.1	0.0	81.2	0.0	0.0	1.5	1.1
4359	803	65.3	0.5	31.9	0.0	0.0	2.0	0.4
4360	387	91.7	0.0	7.0	0.0	0.0	1.3	0.0
4361	345	91.6	0.3	1.7	0.0	0.0	6.4	0.0
4362	410	92.4	0.0	4.6	0.0	0.0	2.9	0.0
4363	1,349	91.5	0.1	4.2	0.1	0.1	4.1	0.0
4401	1,852	83.9	1.0	8.9	0.1	0.1	5.9	0.1

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
4501	1,500	87.4	0.4	7.9	0.3	0.1	3.9	0.1
4601	2,224	60.5	1.9	27.7	0.0	0.1	9.7	0.2
4602	8	*	*	*	*	*	*	*
4701	1,824	88.5	0.9	4.8	0.1	0.6	5.0	0.1
5050	2,331	51.7	30.2	4.2	0.0	0.9	12.8	0.1
5051	3,304	50.7	32.1	4.8	0.0	1.6	10.7	0.2
5052	3,265	42.4	29.9	3.0	0.0	13.7	10.8	0.2
5055	1,846	95.2	1.1	0.5	0.0	2.8	0.3	0.1
5101	2,979	66.5	9.4	12.0	0.0	1.7	10.3	0.0
5202	1,024	78.9	0.4	4.8	0.8	0.8	14.3	0.1
5203	119	76.5	0.0	0.8	0.8	0.0	21.8	0.0
5204	97	*	*	*	*	*	*	*
5601	1,786	90.0	1.2	3.6	0.0	0.3	4.7	0.2
5801	2,515	75.0	6.1	8.5	0.0	0.2	10.2	0.1
5803	344	85.2	0.6	9.0	0.3	0.3	4.7	0.0
5804	98	*	*	*	*	*	*	*
6101	92	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
6102	285	81.4	1.1	8.1	0.0	0.0	9.5	0.0
6103	5,696	48.9	24.7	13.1	0.1	0.2	12.9	0.0
6104	113	24.8	51.3	12.4	0.0	1.8	9.7	0.0
6105	64	*	*	*	*	*	*	*
6301	5,686	37.8	27.1	23.7	0.0	2.5	8.8	0.1
6312	3,443	38.5	48.2	3.9	0.1	1.6	7.6	0.0
6316	1,886	67.1	15.2	9.2	0.1	1.9	6.5	0.0
6320	236	68.6	18.2	3.0	0.0	1.7	8.5	0.0
6325	242	90.9	1.2	2.9	0.0	1.2	3.7	0.0
6330	1,292	17.5	55.9	18.2	0.2	1.2	7.0	0.2
6335	1,286	75.7	12.4	2.5	0.0	3.1	6.2	0.1
6340	149	20.1	12.8	59.7	0.0	3.4	4.0	0.0
6345	39	*	*	*	*	*	*	*
6350	33	*	*	*	*	*	*	*
7304	250	86.0	1.2	10.8	0.0	0.0	2.0	0.0
7306	3,532	36.2	30.7	23.3	0.1	0.3	9.4	0.1
7308	234	74.8	1.7	15.0	0.0	0.0	8.5	0.0

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
7310	57	*	*	*	*	*	*	*
7401	3,313	70.9	5.5	9.7	0.0	0.2	13.5	0.2
7601	887	86.5	0.6	8.7	0.0	0.0	4.1	0.2
7850	1,774	84.1	0.3	7.5	0.0	0.2	7.9	0.1
7852	113	92.9	0.0	1.8	0.0	0.0	5.3	0.0
8101	2,374	5.0	89.3	0.8	0.0	0.7	4.0	0.1
8102	986	8.7	5.2	83.1	0.0	0.0	2.9	0.1
8105	2,231	34.9	55.9	3.4	0.2	0.8	4.5	0.3
8106	691	17.2	59.5	11.6	0.3	0.3	10.7	0.4
8150	149	43.0	20.1	23.5	0.0	11.4	2.0	0.0
8151	4,680	30.0	37.5	13.0	0.2	4.2	15.0	0.2
8152	230	27.4	45.7	10.0	0.0	0.9	16.1	0.0
8153	46	*	*	*	*	*	*	*
8155	140	50.7	22.9	10.0	0.7	6.4	9.3	0.0
8157	50	*	*	*	*	*	*	*
8201	3,709	47.9	30.9	6.3	0.1	2.9	11.6	0.3
8202	5,695	89.0	6.6	2.9	0.0	0.2	1.3	0.1

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
8203	672	59.7	24.0	5.2	0.0	1.0	10.1	0.0
8250	1,001	23.7	59.1	6.7	0.1	1.1	9.3	0.0
8252	173	2.9	92.5	0.0	0.0	0.0	4.6	0.0
8255	4,016	48.7	28.7	9.4	0.1	2.6	10.3	0.2
8257	1,689	52.8	11.6	1.6	0.2	32.1	1.5	0.2
8263	1,186	60.8	11.6	16.5	0.1	0.8	10.1	0.1
8264	707	48.1	36.4	6.6	0.0	1.0	7.9	0.0
8265	152	59.9	11.8	19.1	0.0	0.0	9.2	0.0
8267	16	*	*	*	*	*	*	*
8901	1,658	0.8	96.0	0.9	0.1	0.4	1.7	0.0
8902	1,554	2.6	92.5	1.2	0.0	0.1	3.6	0.0
8903	1,704	0.8	95.2	0.7	0.0	0.6	2.7	0.1
8904	1,933	2.3	90.5	0.9	0.2	2.8	3.2	0.1
8905	2,680	53.4	26.6	14.8	0.0	0.6	4.6	0.0
8908	9	*	*	*	*	*	*	*
9012	63	*	*	*	*	*	*	*
9013	46	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
9016	324	0.3	96.9	1.9	0.0	0.0	0.9	0.0
9017	94	*	*	*	*	*	*	*
9018	254	0.8	92.5	2.8	0.0	0.0	3.9	0.0
9041	30	*	*	*	*	*	*	*
9045	3,527	4.6	42.1	50.8	0.0	0.1	2.4	0.1
9050	1,631	9.6	59.3	27.1	0.0	0.3	3.7	0.1
9051	2,007	3.7	7.6	85.4	0.0	0.1	3.1	0.1
9053	2,714	2.1	93.6	0.7	0.3	1.8	1.6	0.0
9055	713	1.4	94.5	1.7	0.0	1.0	1.4	0.0
9058	1,603	42.0	40.4	1.9	0.1	13.7	1.9	0.0
9066	1,798	1.7	96.3	0.9	0.0	0.1	0.9	0.1
9068	1,633	1.3	95.5	0.6	0.0	0.2	2.1	0.1
9069	3,881	0.5	96.7	0.6	0.0	0.1	1.9	0.1
9070	1,925	3.0	86.1	0.5	0.1	7.4	3.0	0.0
9072	2,272	0.9	94.5	0.9	0.1	1.0	2.6	0.0
9080	3,472	22.2	72.1	2.4	0.0	0.1	3.1	0.1
9081	177	23.7	41.8	23.7	1.1	0.0	9.0	0.6

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 4A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Michigan	240,590	49.0	27.1	13.5	0.3	1.9	8.1	0.2
Nation (Prior Year)	911,219	48.5	22.9	14.8	0.7	3.5	9.1	0.5

(1) Reporting period is January 1 through December 31

(2) Analyses based on one record per child

(3) Excludes records with unknown data errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0101	158	35.4	27.2	41.1
0102	689	40.5	19.4	42.8
0103	608	34.2	21.1	49.0
0104	223	42.6	17.0	44.4
0250	204	30.9	24.0	51.0
0251	135	32.6	25.9	50.4
0252	199	30.7	25.6	47.7
0253	232	34.5	21.6	50.9
0401	731	34.9	23.4	45.0
0402	554	40.1	23.1	40.4
0403	176	35.8	25.6	43.8
0404	207	32.9	19.8	50.7
0501	792	38.8	20.3	43.4
0502	2	*	*	*
0503	193	31.6	24.9	46.6
0504	721	39.3	22.7	42.4
0505	427	35.1	19.7	48.0
0506	312	33.7	26.0	44.9
0507	262	26.0	16.4	60.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
0508	100	16.0	27.0	62.0
0509	10	*	*	*
0650	1,500	36.9	25.1	41.8
0651	894	34.3	23.9	46.5
0652	493	36.9	21.5	44.0
0653	742	28.6	27.5	48.2
0654	566	29.9	23.1	51.9
0655	538	32.7	24.3	47.6
0657	101	26.7	23.8	51.5
0701	639	36.3	20.2	46.3
0702	134	32.8	24.6	46.3
0703	358	39.7	20.4	44.1
0704	87	*	*	*
0705	189	33.3	19.0	50.3
0850	1,478	38.4	21.7	44.3
0851	1,270	36.1	20.9	47.2
0901	2,492	32.9	24.6	46.1
0902	74	*	*	*
1051	1,894	34.8	24.9	43.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
1101	2,533	39.7	21.0	43.5
1102	1,256	40.5	21.3	41.8
1106	179	38.5	25.7	40.2
1250	1,508	32.4	24.9	46.2
1251	1,322	36.9	22.2	44.3
1252	724	31.5	24.9	48.3
1253	1,490	37.2	20.6	45.0
1350	3,706	36.2	22.2	45.4
1351	255	37.3	20.4	45.5
1352	590	30.7	22.4	50.8
1401	361	37.4	24.7	39.1
1402	241	36.9	21.6	43.2
1403	31	*	*	*
1404	6	*	*	*
1501	1,483	31.9	23.0	49.2
1503	1,860	33.4	23.8	46.5
1504	3,407	27.5	28.6	47.3
1505	1,136	30.5	24.0	47.6
1506	1,040	33.3	23.3	48.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months	12 - 23 Months	24 - 59 Months
		%	%	%
1507	884	33.6	23.1	45.9
1508	141	22.0	22.0	59.6
1509	858	29.4	21.8	52.0
1511	1,080	35.5	22.6	45.6
1512	637	32.7	21.7	48.7
1701	1,095	36.9	21.0	45.3
1950	919	37.4	21.1	44.7
1952	174	36.2	20.7	47.1
1953	504	34.5	20.4	47.2
1954	957	39.5	20.3	42.6
1955	189	27.0	20.1	55.0
1957	931	37.6	18.7	46.1
2150	1,097	36.2	22.6	44.4
2151	378	36.8	24.3	43.7
2251	652	35.1	23.0	46.3
2252	300	31.3	22.3	50.7
2554	211	23.2	28.0	53.1
2555	11,081	41.5	20.4	41.1
3001	1,533	35.4	19.7	49.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
3002	460	31.5	19.6	52.4
3004	219	31.1	20.5	51.6
3007	33	*	*	*
3009	15	*	*	*
3201	732	37.8	20.1	44.0
3301	5,100	39.9	21.5	42.0
3304	240	32.5	20.8	48.8
3305	325	36.3	19.4	46.8
3306	539	33.8	21.9	48.2
3308	151	33.8	21.2	49.0
3309	70	*	*	*
3312	770	31.9	20.4	50.4
3550	129	31.8	19.4	53.5
3551	40	*	*	*
3552	76	*	*	*
3701	1,665	35.6	24.6	44.3
3850	4,711	35.6	21.6	46.3
3950	2,737	40.9	19.7	41.6
3954	21	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
4001	1,934	36.0	19.6	47.4
4002	1,705	35.0	21.2	46.5
4003	3	*	*	*
4004	1	*	*	*
4103	106	24.5	26.4	52.8
4106	3,252	32.5	21.8	49.3
4107	5,155	39.4	20.4	43.3
4109	1,678	35.3	23.1	46.1
4112	806	34.0	20.7	50.4
4113	1,213	41.5	17.1	45.1
4115	5,571	36.9	20.6	45.7
4351	892	32.8	23.0	47.1
4352	293	31.7	27.0	44.7
4353	686	34.7	22.3	45.9
4354	1,142	33.8	22.8	47.3
4355	489	34.4	19.8	48.5
4356	749	35.9	21.4	45.9
4358	261	25.7	19.9	58.2
4359	803	29.6	22.0	52.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
4360	387	31.5	23.8	49.4
4361	345	39.1	24.1	39.7
4362	410	35.1	23.7	45.1
4363	1,349	34.8	22.8	46.4
4401	1,852	35.0	24.2	43.3
4501	1,500	34.5	21.2	48.1
4601	2,224	39.5	21.9	41.9
4602	8	*	*	*
4701	1,824	36.3	22.7	45.2
5050	2,331	41.4	22.9	39.0
5051	3,304	41.7	22.6	38.7
5052	3,265	41.6	21.7	40.2
5055	1,846	34.0	21.6	47.5
5101	2,979	38.5	23.1	41.2
5202	1,024	35.5	25.8	41.4
5203	119	20.2	29.4	54.6
5204	97	*	*	*
5601	1,786	39.7	18.6	44.5
5801	2,515	37.6	22.3	43.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
5803	344	28.8	23.3	53.2
5804	98	*	*	*
6101	92	*	*	*
6102	285	28.4	22.8	54.4
6103	5,696	35.7	22.6	45.0
6104	113	50.4	25.7	23.9
6105	64	*	*	*
6301	5,686	42.4	20.1	40.3
6312	3,443	41.2	19.6	41.4
6316	1,886	38.7	20.6	43.8
6320	236	33.9	20.3	48.7
6325	242	44.2	19.8	38.0
6330	1,292	42.4	17.6	41.7
6335	1,286	37.4	17.5	48.4
6340	149	34.2	24.2	46.3
6345	39	*	*	*
6350	33	*	*	*
7304	250	36.4	20.4	46.0
7306	3,532	39.7	20.2	43.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
7308	234	37.6	17.9	45.3
7310	57	*	*	*
7401	3,313	36.0	24.4	43.7
7601	887	40.2	18.7	45.3
7850	1,774	36.3	21.7	45.5
7852	113	35.4	12.4	54.9
8101	2,374	42.9	19.1	40.9
8102	986	32.6	18.4	52.4
8105	2,231	46.1	20.1	36.6
8106	691	38.1	21.0	42.4
8150	149	26.8	17.4	59.1
8151	4,680	35.2	23.3	44.9
8152	230	41.7	20.9	40.9
8153	46	*	*	*
8155	140	24.3	17.1	61.4
8157	50	*	*	*
8201	3,709	42.7	20.7	39.5
8202	5,695	35.8	18.5	48.3
8203	672	39.3	24.9	39.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
8250	1,001	37.6	23.9	40.5
8252	173	37.6	16.8	48.6
8255	4,016	44.4	20.1	39.0
8257	1,689	38.1	18.3	45.7
8263	1,186	41.7	22.9	37.7
8264	707	48.8	16.4	37.1
8265	152	59.2	6.6	34.9
8267	16	*	*	*
8901	1,658	35.6	17.2	49.4
8902	1,554	40.7	20.9	41.0
8903	1,704	36.9	17.7	47.4
8904	1,933	38.9	18.2	45.1
8905	2,680	34.1	22.1	47.3
8908	9	*	*	*
9012	63	*	*	*
9013	46	*	*	*
9016	324	32.4	17.3	53.7
9017	94	*	*	*
9018	254	28.0	20.9	55.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Age Distribution				
Clinic	Number Included in Analysis	0 - 11 Months	12 - 23 Months	24 - 59 Months
		%	%	%
9041	30	*	*	*
9045	3,527	48.8	14.5	39.7
9050	1,631	36.0	19.9	48.4
9051	2,007	34.7	20.7	47.3
9053	2,714	37.5	23.4	42.4
9055	713	100.0	0.0	0.0
9058	1,603	26.7	22.5	53.9
9066	1,798	39.2	20.4	43.5
9068	1,633	39.9	19.1	43.5
9069	3,881	42.2	20.9	39.8
9070	1,925	38.4	19.8	44.4
9072	2,272	43.0	19.5	40.4
9080	3,472	32.2	20.3	49.7
9081	177	55.4	15.3	31.1
Michigan	240,590	37.6	18.0	44.3
Nation (Prior Year)	911,219	37.3	19.1	43.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0101	31	*	*	149	8.1	2.0	10.1	60	*	*	100	16.0
0102	159	7.5	8.8	672	6.7	4.0	7.9	282	14.5	11.7	431	21.6
0103	113	5.3	6.2	592	7.4	0.7	14.4	289	17.6	15.2	423	26.0
0104	60	*	*	215	9.3	5.6	7.0	95	*	*	143	18.2
0250	37	*	*	195	4.1	1.5	6.7	96	*	*	140	9.3
0251	27	*	*	134	4.5	3.7	6.0	67	*	*	75	*
0252	44	*	*	193	4.1	2.1	10.9	92	*	*	137	5.1
0253	46	*	*	229	1.7	9.2	2.2	116	3.4	2.6	145	15.9
0401	168	8.3	8.3	720	7.6	2.1	12.9	323	18.6	14.2	450	11.6
0402	153	9.8	8.5	539	8.3	3.0	11.1	211	13.3	18.5	315	12.4
0403	43	*	*	170	4.1	2.9	8.8	74	*	*	115	12.2
0404	39	*	*	203	3.4	3.0	5.4	101	14.9	8.9	125	16.8
0501	191	7.9	11.0	780	6.2	4.7	11.2	339	19.5	16.8	489	7.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0502	0	*	*	2	*	*	*	1	*	*	2	*
0503	39	*	*	187	7.5	7.0	9.6	86	*	*	122	9.0
0504	163	7.4	6.7	701	7.1	4.1	11.4	294	19.0	13.9	459	10.2
0505	86	*	*	408	5.1	4.9	10.3	202	20.8	13.9	290	4.8
0506	50	*	*	302	5.6	3.3	9.9	132	18.9	12.9	218	5.5
0507	27	*	*	260	6.5	1.5	10.4	158	15.2	12.7	195	7.2
0508	10	*	*	99	*	*	*	61	*	*	83	*
0509	4	*	*	9	*	*	*	5	*	*	6	*
0650	360	8.9	10.6	1,464	7.0	3.1	10.5	601	17.1	16.3	939	10.8
0651	191	6.8	6.3	865	6.4	1.4	10.9	399	17.0	15.3	601	10.3
0652	109	4.6	8.3	475	6.7	1.3	10.3	206	19.4	17.0	294	7.1
0653	146	7.5	7.5	726	4.3	2.9	10.7	345	12.8	14.2	497	8.2
0654	113	5.3	10.6	547	5.9	2.7	9.7	278	13.3	14.4	369	8.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0655	104	9.6	8.7	520	6.3	3.7	8.5	242	17.8	13.6	375	9.6
0657	18	*	*	100	6.0	4.0	8.0	52	*	*	66	*
0701	153	8.5	9.8	615	6.8	3.1	5.5	282	21.3	9.2	428	7.7
0702	33	*	*	125	6.4	4.0	6.4	53	*	*	82	*
0703	84	*	*	350	5.1	6.0	8.6	152	20.4	9.9	211	7.6
0704	25	*	*	86	*	*	*	40	*	*	56	*
0705	43	*	*	183	6.6	2.7	9.8	90	*	*	117	7.7
0850	333	6.6	6.9	1,440	6.2	3.9	12.6	626	16.9	17.7	932	6.2
0851	268	10.4	8.6	1,219	5.3	3.7	7.9	569	18.6	12.5	825	8.6
0901	510	10.0	9.8	2,431	7.9	2.2	11.2	1,102	15.7	15.6	1,687	7.7
0902	13	*	*	73	*	*	*	35	*	*	53	*
1051	470	6.8	8.7	1,788	4.1	6.4	9.5	798	17.5	13.2	1,110	17.1
1101	642	10.9	6.4	2,455	5.6	5.3	9.1	1,049	15.7	14.0	1,459	14.1

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1102	324	8.6	8.3	1,220	5.2	3.8	9.4	501	16.6	13.6	740	7.8
1106	43	*	*	176	5.7	3.4	11.9	69	*	*	106	6.6
1250	332	8.1	9.6	1,463	7.5	1.8	15.1	663	17.5	17.2	949	3.0
1251	279	9.0	7.9	1,291	8.0	1.3	11.3	562	16.7	13.7	828	3.0
1252	137	8.0	11.7	708	6.6	2.0	12.7	334	19.8	16.8	498	7.4
1253	334	7.2	6.9	1,460	5.5	2.5	9.2	651	14.4	12.0	963	11.0
1350	825	8.6	7.0	3,602	7.6	2.9	13.7	1,617	17.4	15.4	2,378	22.9
1351	54	*	*	248	6.5	2.0	13.7	110	15.5	18.2	159	23.3
1352	115	15.7	3.5	564	5.7	2.8	14.4	289	19.7	20.1	407	22.6
1401	86	*	*	351	5.7	8.0	10.3	134	15.7	14.2	230	9.1
1402	57	*	*	234	5.1	5.1	10.3	100	22.0	10.0	151	9.9
1403	5	*	*	30	*	*	*	16	*	*	23	*
1404	1	*	*	6	*	*	*	4	*	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1501	305	6.6	7.9	1,431	3.2	3.8	12.6	685	19.3	18.4	988	16.6
1503	397	7.3	11.1	1,824	5.3	2.4	10.9	827	18.7	14.3	1,230	16.3
1504	733	7.9	7.4	3,317	4.7	2.3	11.1	1,553	16.4	15.4	2,294	34.4
1505	218	5.5	11.5	1,112	6.7	2.3	8.5	525	17.9	9.9	772	12.4
1506	208	5.3	8.7	1,006	6.7	2.5	11.8	474	18.8	14.1	688	20.6
1507	188	4.3	9.0	851	4.5	2.7	8.3	384	14.1	9.9	571	18.0
1508	16	*	*	132	5.3	1.5	16.7	79	*	*	103	26.2
1509	157	7.0	5.7	828	3.5	4.5	11.2	423	17.7	16.1	586	27.5
1511	250	8.4	11.2	1,045	4.3	2.4	9.7	470	19.4	17.0	692	16.9
1512	111	9.9	9.0	617	5.8	2.3	11.8	293	18.4	15.0	406	13.5
1701	234	5.6	8.5	1,057	7.2	3.0	12.5	473	18.2	13.7	667	18.7
1950	213	8.5	7.5	883	7.1	1.4	9.3	380	16.6	14.2	604	5.8
1952	33	*	*	170	6.5	0.0	8.8	80	*	*	126	4.8

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1953	103	11.7	6.8	491	7.7	1.4	11.4	232	22.0	9.5	340	3.8
1954	223	14.8	9.4	930	6.1	3.2	8.7	392	15.3	13.3	590	3.4
1955	31	*	*	184	5.4	4.3	15.8	98	*	*	133	2.3
1957	195	6.2	7.2	902	6.0	1.4	10.6	408	17.4	12.7	574	2.8
2150	255	6.7	10.6	1,076	5.2	2.2	9.0	470	20.0	10.6	682	2.2
2151	84	*	*	374	6.4	3.2	13.6	160	16.9	17.5	221	3.6
2251	133	11.3	9.8	637	8.3	1.6	8.3	291	17.2	10.3	416	9.6
2252	55	*	*	294	7.1	0.7	14.6	148	16.2	18.9	199	8.0
2554	39	*	*	205	3.4	1.5	11.2	109	14.7	16.5	142	12.7
2555	2,759	12.4	5.1	10,753	9.6	2.1	10.8	4,409	15.8	12.5	6,831	23.8
3001	334	11.7	5.4	1,475	7.8	2.0	12.4	733	18.8	13.0	1,033	2.2
3002	89	*	*	446	7.4	1.3	11.9	233	21.0	14.2	316	7.3
3004	38	*	*	206	2.9	1.0	22.3	109	22.0	32.1	161	7.5

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
3007	5	*	*	31	*	*	*	18	*	*	25	*
3009	0	*	*	8	*	*	*	7	*	*	15	*
3201	164	7.3	11.0	714	6.6	2.7	13.3	310	18.1	15.2	469	7.9
3301	1,308	10.2	5.5	4,973	6.5	3.6	9.5	2,059	15.2	13.9	3,192	13.9
3304	42	*	*	230	3.5	1.3	10.9	112	11.6	10.7	162	6.2
3305	65	*	*	319	4.1	1.3	8.2	147	12.9	6.1	216	6.5
3306	89	*	*	534	6.2	1.3	8.2	254	20.5	5.5	383	5.2
3308	30	*	*	147	4.8	0.0	10.2	71	*	*	105	4.8
3309	15	*	*	68	*	*	*	27	*	*	45	*
3312	144	10.4	5.6	750	5.3	3.2	11.7	373	18.5	15.8	540	12.8
3550	22	*	*	126	6.3	0.0	20.6	66	*	*	75	*
3551	10	*	*	39	*	*	*	20	*	*	19	*
3552	19	*	*	73	*	*	*	35	*	*	41	*

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
3701	375	6.7	8.5	1,628	4.3	4.9	10.8	709	16.2	16.1	1,066	8.9
3850	1,024	10.4	7.4	4,565	5.8	3.8	9.7	2,084	19.5	15.7	3,123	20.0
3950	666	10.1	9.9	2,614	5.8	3.6	10.0	1,083	17.3	13.2	1,720	18.1
3954	2	*	*	20	*	*	*	14	*	*	15	*
4001	419	10.3	6.9	1,874	4.6	3.2	10.1	882	15.8	13.3	1,315	31.9
4002	347	11.8	8.9	1,654	7.9	2.0	12.6	767	17.5	14.2	1,148	17.3
4003	1	*	*	2	*	*	*	0	*	*	2	*
4004	0	*	*	1	*	*	*	0	*	*	1	*
4103	17	*	*	101	3.0	5.9	6.9	52	*	*	49	*
4106	622	9.3	6.3	3,173	8.1	1.6	10.6	1,540	17.2	12.1	2,175	14.2
4107	1,189	12.4	8.2	4,994	8.1	2.1	9.9	2,133	15.7	12.0	3,270	13.1
4109	351	10.5	10.8	1,629	5.7	2.0	8.2	741	14.7	11.3	1,091	7.6
4112	187	8.6	11.2	775	5.4	3.6	18.3	384	18.2	25.5	580	13.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
4113	305	6.9	7.9	1,186	7.8	1.9	12.8	528	17.8	17.2	739	3.5
4115	1,147	10.0	7.7	5,370	7.2	2.3	10.7	2,443	15.8	12.2	3,594	8.8
4351	181	7.2	9.4	866	6.4	1.7	9.6	402	15.9	13.9	572	22.6
4352	69	*	*	277	8.7	1.8	9.7	122	17.2	14.8	195	14.4
4353	155	5.8	11.0	660	8.9	2.6	13.6	303	19.1	18.2	437	14.9
4354	243	8.2	7.8	1,105	7.1	1.7	10.6	512	20.7	16.6	758	13.7
4355	107	6.5	5.6	467	6.9	3.2	10.9	219	18.3	18.3	299	10.7
4356	173	8.7	9.2	707	8.9	1.0	11.6	316	21.5	16.8	483	15.9
4358	47	*	*	251	10.0	0.8	21.1	144	20.8	26.4	194	17.5
4359	162	8.0	14.8	779	6.4	1.7	14.6	404	18.8	16.3	564	14.9
4360	70	*	*	374	4.5	2.1	8.3	179	24.0	12.3	255	9.4
4361	96	*	*	342	12.0	2.3	12.9	135	25.9	17.8	218	10.1
4362	97	*	*	401	8.7	0.5	10.7	179	16.8	15.6	259	18.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
4363	293	4.8	9.6	1,319	5.2	1.4	8.4	599	18.7	12.2	866	12.8
4401	396	9.1	9.1	1,810	6.5	2.4	9.0	769	18.9	15.2	1,225	10.4
4501	315	8.3	5.1	1,453	7.3	2.4	10.3	682	17.0	14.8	1,017	7.6
4601	552	8.3	8.5	2,152	5.8	2.6	9.3	875	20.0	15.8	1,260	9.4
4602	3	*	*	8	*	*	*	5	*	*	5	*
4701	419	7.4	8.1	1,776	6.9	1.0	9.0	784	16.5	12.2	1,189	6.3
5050	616	7.6	7.1	2,274	6.9	1.3	10.6	867	15.2	13.7	1,402	16.9
5051	885	13.1	6.1	3,232	7.6	1.3	12.0	1,240	12.7	14.1	1,946	16.6
5052	880	13.3	5.6	3,186	8.1	2.3	9.7	1,268	14.0	11.9	1,893	14.3
5055	222	7.2	5.0	1,809	2.8	2.2	10.9	841	19.1	15.3	1,271	35.6
5101	707	9.3	9.1	2,904	6.0	2.0	9.0	1,177	16.9	14.7	1,862	20.0
5202	265	8.7	10.2	971	8.2	2.6	11.2	394	18.5	16.0	524	7.4
5203	15	*	*	111	7.2	0.9	16.2	58	*	*	68	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
5204	17	*	*	93	*	*	*	48	*	*	56	*
5601	410	9.3	11.5	1,744	8.0	1.3	12.0	759	19.2	16.6	1,063	7.0
5801	618	8.6	8.3	2,433	7.6	3.2	10.1	1,030	15.7	16.2	1,548	10.5
5803	52	*	*	334	7.8	2.1	12.0	172	13.4	15.1	241	11.2
5804	9	*	*	96	*	*	*	52	*	*	80	*
6101	13	*	*	90	*	*	*	47	*	*	62	*
6102	45	*	*	273	4.4	3.3	8.8	148	12.8	10.8	195	12.3
6103	1,362	12.5	6.5	5,516	6.9	2.9	10.5	2,487	16.2	12.8	3,537	17.7
6104	25	*	*	111	3.6	2.7	9.9	27	*	*	56	*
6105	9	*	*	60	*	*	*	32	*	*	38	*
6301	1,388	12.6	6.1	5,525	9.7	1.6	10.2	2,226	16.2	12.9	3,405	13.9
6312	788	9.6	7.1	3,366	8.7	2.3	8.1	1,370	15.7	10.0	2,141	18.6
6316	427	9.4	4.9	1,849	5.3	1.9	9.7	794	15.4	13.9	1,221	8.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
6320	52	*	*	229	4.8	3.1	8.7	108	8.3	13.0	154	9.7
6325	62	*	*	236	6.4	4.2	8.1	88	*	*	151	11.9
6330	314	11.8	4.8	1,258	7.2	3.7	6.4	522	15.1	10.0	827	21.6
6335	279	10.0	5.7	1,236	3.6	4.3	9.2	582	10.7	12.4	875	15.1
6340	22	*	*	147	4.1	2.0	9.5	67	*	*	109	23.9
6345	8	*	*	36	*	*	*	21	*	*	30	*
6350	5	*	*	31	*	*	*	13	*	*	23	*
7304	53	*	*	245	3.3	6.1	10.6	111	18.9	18.0	154	9.7
7306	859	12.1	6.4	3,414	7.1	3.0	12.3	1,464	16.7	15.8	2,221	16.8
7308	47	*	*	228	3.9	2.6	8.8	102	24.5	12.7	160	11.3
7310	18	*	*	55	*	*	*	24	*	*	26	*
7401	729	8.9	6.4	3,206	5.6	1.9	8.9	1,366	18.4	15.8	2,178	7.7
7601	207	6.8	7.2	859	6.5	1.0	7.9	378	20.1	13.2	568	13.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
7850	401	7.5	10.2	1,722	5.5	4.1	9.1	769	18.7	15.2	1,179	13.7
7852	19	*	*	112	3.6	4.5	10.7	61	*	*	80	*
8101	606	16.7	3.5	2,286	7.0	5.5	7.7	942	13.1	10.4	1,418	38.0
8102	185	5.9	9.7	950	6.0	4.0	17.8	490	21.0	23.9	648	27.2
8105	594	10.6	6.7	2,169	5.0	4.7	8.4	785	14.1	9.2	1,259	18.6
8106	169	14.2	4.1	667	9.3	0.6	10.8	280	18.9	16.1	412	15.3
8150	15	*	*	140	3.6	3.6	10.0	78	*	*	110	20.0
8151	1,052	12.2	7.3	4,542	7.0	2.1	9.9	1,998	16.2	13.8	3,049	17.7
8152	60	*	*	226	8.8	1.3	9.3	93	*	*	145	24.8
8153	5	*	*	43	*	*	*	27	*	*	34	*
8155	17	*	*	137	8.8	2.9	8.0	84	*	*	103	26.2
8157	9	*	*	48	*	*	*	29	*	*	34	*
8201	950	10.0	5.7	3,631	7.7	2.0	10.7	1,418	15.3	14.0	2,282	16.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)						Anemia Low Hb/Hct ⁽⁷⁾		
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
8202	1,086	8.4	4.7	5,587	4.9	2.3	8.1	2,663	12.7	10.4	3,733	19.4
8203	169	12.4	3.0	654	6.9	1.4	8.1	256	14.8	8.2	404	6.4
8250	192	7.8	9.4	987	6.5	2.8	10.6	399	13.8	12.8	615	14.0
8252	47	*	*	160	9.4	1.9	6.9	80	*	*	105	16.2
8255	1,098	10.2	9.2	3,920	6.7	2.2	7.8	1,499	18.1	10.9	2,272	16.2
8257	403	12.7	3.0	1,646	6.0	3.5	9.4	739	14.6	15.0	1,034	19.1
8263	277	9.4	6.5	1,162	7.0	1.4	8.3	429	19.1	10.7	705	12.2
8264	263	8.7	7.2	689	8.0	1.3	10.6	254	14.2	9.4	360	15.3
8265	83	*	*	150	6.7	1.3	7.3	53	*	*	62	*
8267	14	*	*	16	*	*	*	2	*	*	2	*
8901	350	17.1	1.4	1,598	7.6	6.2	7.6	785	14.9	9.6	1,105	35.8
8902	342	12.6	2.9	1,503	6.3	6.6	8.0	615	8.9	10.9	1,009	38.0
8903	352	12.2	3.1	1,641	6.9	3.6	10.0	781	17.0	13.7	1,110	38.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
8904	414	13.5	5.1	1,851	5.2	5.6	8.1	836	10.9	9.7	1,224	42.8
8905	488	7.6	5.7	2,582	6.4	4.6	13.6	1,210	15.4	15.0	1,823	28.0
8908	4	*	*	6	*	*	*	4	*	*	5	*
9012	11	*	*	63	*	*	*	35	*	*	38	*
9013	6	*	*	45	*	*	*	24	*	*	13	*
9016	56	*	*	315	8.9	3.5	11.4	169	17.8	11.2	191	46.1
9017	11	*	*	87	*	*	*	52	*	*	55	*
9018	41	*	*	245	4.1	6.5	9.8	139	20.1	10.8	38	*
9041	3	*	*	30	*	*	*	18	*	*	3	*
9045	1,376	15.4	4.8	3,335	5.0	6.0	11.8	1,315	14.6	18.7	1,839	15.0
9050	327	15.9	4.0	1,578	7.2	2.4	9.5	763	13.8	13.8	1,031	28.8
9051	368	8.2	5.4	1,904	8.2	1.9	17.6	869	17.7	20.7	1,265	22.1
9053	501	17.0	3.6	2,636	7.5	3.0	6.3	1,108	14.2	9.1	1,801	26.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2015 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Run Date: 05/02/2016

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	% <2500 g	% >4000 g	Number	%	%	%	Number	% Overweight	% Obese	Number	%
9055	710	9.7	2.5	669	7.6	15.7	2.8	0	*	*	0	*
9058	251	16.3	3.2	1,556	6.2	3.9	9.4	834	12.7	11.5	1,128	20.0
9066	399	15.5	2.5	1,729	7.1	2.7	8.9	749	15.0	12.4	1,071	16.6
9068	316	12.7	3.5	1,576	5.2	4.5	7.1	683	15.2	8.1	1,039	19.8
9069	888	15.5	2.7	3,767	7.4	3.3	8.5	1,491	12.2	11.9	2,416	26.9
9070	346	15.6	3.2	1,777	6.6	4.2	8.0	813	10.8	10.7	1,181	45.0
9072	570	15.4	3.3	2,195	7.8	2.6	7.7	878	14.5	12.2	1,316	31.7
9080	725	15.2	4.6	3,330	6.6	3.2	7.9	1,680	14.6	10.0	2,255	28.0
9081	43	*	*	172	12.2	6.4	9.3	53	*	*	67	*
Michigan	55,089	10.5	6.8	233,217	6.7	2.9	10.2	102,221	16.3	13.6	151,524	17.2
Nation (Prior Year)	210,775	9.8	7.3	877,334	6.8	2.8	10.5	381,637	16.2	13.6	527,531	14.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan**

Run Date: 05/03/2016

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)**

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
0101	31	*	29	*	19	*	22	*	29	*	75	*	157	3.8
0102	160	52.5	142	16.9	119	11.8	125	14.4	140	12.9	325	86.5	678	5.5
0103	113	68.1	107	22.4	104	9.6	84	*	100	10.0	323	85.4	604	6.3
0104	60	*	55	*	38	*	40	*	53	*	106	81.1	223	9.0
0250	37	*	42	*	47	*	27	*	40	*	108	89.8	204	3.9
0251	27	*	23	*	18	*	21	*	23	*	71	*	135	9.6
0252	44	*	25	*	30	*	14	*	24	*	102	72.5	199	5.5
0253	48	*	44	*	41	*	42	*	44	*	125	81.6	231	9.1
0401	170	54.1	117	26.5	126	11.9	123	22.0	114	16.7	359	95.8	728	2.2
0402	153	74.5	92	*	98	*	82	*	90	*	243	88.9	551	5.4
0403	43	*	29	*	29	*	28	*	29	*	84	*	172	12.8
0404	39	*	35	*	42	*	26	*	35	*	110	95.5	207	9.2
0501	204	71.1	148	21.6	136	10.3	128	14.8	145	8.3	373	89.5	789	5.4
0502	0	*	1	*	1	*	0	*	1	*	1	*	2	*
0503	39	*	32	*	27	*	26	*	31	*	103	90.3	192	4.2
0504	167	82.6	127	42.5	119	22.7	105	41.0	122	22.1	344	89.8	719	5.6
0505	96	*	75	*	68	*	66	*	75	*	220	89.1	427	5.6
0506	63	*	62	*	60	*	46	*	62	*	150	82.7	311	8.7

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breastsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
0507	28	*	44	*	40	*	29	*	43	*	166	89.8	261	8.0
0508	10	*	12	*	17	*	8	*	11	*	67	*	100	2.0
0509	4	*	1	*	0	*	1	*	1	*	6	*	10	*
0650	360	72.2	297	23.9	286	10.8	249	20.9	284	12.0	688	91.4	1,495	3.4
0651	191	67.5	165	18.2	184	11.4	142	18.3	164	10.4	439	91.3	888	8.7
0652	110	70.0	83	*	89	*	75	*	80	*	242	90.9	488	9.6
0653	147	76.2	136	19.9	131	12.2	108	24.1	133	12.0	396	95.2	739	7.3
0654	113	76.1	92	*	84	*	83	*	91	*	319	92.2	564	6.2
0655	104	70.2	116	20.7	106	12.3	86	*	113	7.1	270	90.7	533	7.3
0657	19	*	11	*	13	*	13	*	11	*	56	*	101	5.0
0701	153	81.7	116	40.5	95	*	98	*	112	17.0	317	92.4	633	1.9
0702	33	*	24	*	18	*	21	*	23	*	67	*	133	4.5
0703	85	*	64	*	51	*	59	*	64	*	170	85.3	357	3.9
0704	25	*	7	*	13	*	10	*	7	*	41	*	87	*
0705	44	*	29	*	32	*	25	*	28	*	101	86.1	188	4.3
0850	353	77.9	325	25.8	224	10.3	255	22.4	310	12.6	715	86.0	1,476	3.1
0851	274	75.2	245	26.5	203	12.8	172	27.9	232	16.4	643	93.5	1,265	2.8
0901	520	63.3	413	20.3	455	11.4	391	16.4	400	5.0	1,245	96.5	2,486	3.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breastsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
0902	13	*	13	*	12	*	11	*	12	*	38	*	74	*
1051	470	83.4	286	33.2	346	20.2	202	34.7	274	13.1	885	88.4	1,883	2.9
1101	641	60.4	425	17.9	392	12.0	399	14.8	405	7.9	1,170	79.9	2,500	5.3
1102	328	68.3	247	27.1	196	11.7	221	28.1	242	14.9	547	84.5	1,238	4.4
1106	43	*	34	*	31	*	30	*	33	*	79	*	179	5.0
1250	331	81.0	227	21.1	261	12.3	209	16.7	219	10.5	755	83.7	1,502	4.7
1251	281	75.8	232	17.2	196	11.2	190	21.6	228	11.4	647	88.3	1,315	4.9
1252	137	85.4	115	22.6	116	13.8	94	*	97	*	372	82.8	720	2.4
1253	337	75.7	286	20.6	228	9.2	225	18.2	257	7.4	717	86.9	1,477	3.2
1350	833	70.1	644	16.9	635	9.6	536	14.7	576	5.9	1,795	90.8	3,673	3.2
1351	54	*	44	*	34	*	33	*	39	*	128	98.4	252	1.2
1352	115	58.3	105	14.3	113	6.2	74	*	100	8.0	321	96.3	588	0.7
1401	87	*	70	*	58	*	50	*	69	*	159	88.7	361	7.5
1402	57	*	41	*	39	*	33	*	40	*	113	83.2	240	1.7
1403	5	*	2	*	2	*	3	*	2	*	18	*	30	*
1404	1	*	1	*	1	*	1	*	1	*	4	*	6	*
1501	313	68.4	261	16.9	225	7.6	240	17.5	252	6.3	776	94.6	1,463	1.4
1503	399	76.7	316	25.0	331	14.8	292	20.2	299	13.4	936	94.4	1,849	1.2

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breastsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
1504	754	74.5	400	22.8	620	11.8	221	20.4	351	10.0	1,670	87.8	3,353	1.8
1505	226	80.5	163	31.9	214	25.2	148	25.7	142	16.9	563	95.2	1,117	1.8
1506	210	74.3	181	24.3	167	15.6	159	27.0	179	11.2	555	97.5	1,036	1.2
1507	193	76.2	139	30.2	162	24.7	131	26.0	130	20.8	424	85.8	873	4.2
1508	20	*	26	*	24	*	11	*	20	*	82	*	134	0.0
1509	165	75.2	136	27.2	137	11.7	103	23.3	117	12.8	448	87.1	843	3.7
1511	252	81.3	181	27.6	194	12.9	149	26.8	168	10.7	517	91.3	1,068	2.8
1512	115	74.8	112	23.2	124	9.7	101	25.7	107	12.1	331	92.7	624	2.4
1701	237	68.8	205	29.3	187	20.3	165	27.9	198	19.7	529	92.2	1,078	13.5
1950	213	83.1	169	21.9	137	14.6	134	21.6	159	11.3	441	87.3	911	6.8
1952	33	*	27	*	32	*	21	*	26	*	90	*	172	4.1
1953	103	81.6	87	*	75	*	67	*	81	*	251	81.7	498	8.4
1954	224	83.5	180	32.2	137	14.6	160	25.0	173	15.0	440	90.7	936	3.8
1955	31	*	29	*	23	*	25	*	29	*	111	90.1	185	3.8
1957	196	85.2	185	27.6	126	15.9	146	24.0	167	13.8	454	88.3	911	2.4
2150	258	74.8	212	20.8	218	9.6	196	20.9	208	11.1	522	91.2	1,090	1.9
2151	84	*	82	*	76	*	69	*	82	*	178	95.5	371	5.9
2251	133	69.9	108	24.1	103	16.5	84	*	104	15.4	336	92.0	647	1.9

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
2252	55	*	47	*	50	*	44	*	47	*	161	87.6	300	6.0
2554	39	*	26	*	40	*	19	*	26	*	116	90.5	210	3.8
2555	2,777	60.1	2,288	13.5	1,827	8.1	1,780	11.9	2,002	5.6	4,872	80.2	10,912	4.3
3001	334	63.2	248	9.7	216	6.9	211	5.2	244	0.8	806	84.5	1,519	6.8
3002	89	*	69	*	66	*	56	*	69	*	261	93.9	455	3.3
3004	38	*	31	*	35	*	29	*	31	*	119	81.5	218	1.4
3007	5	*	7	*	5	*	5	*	7	*	20	*	33	*
3009	0	*	2	*	3	*	0	*	1	*	10	*	15	*
3201	164	64.0	111	23.4	103	14.6	111	18.0	110	9.1	343	89.8	730	4.9
3301	1,312	74.6	948	23.8	839	15.0	677	15.7	861	7.4	2,282	85.4	5,037	4.3
3304	45	*	41	*	44	*	26	*	39	*	125	86.4	235	1.7
3305	66	*	59	*	51	*	42	*	57	*	165	92.1	319	1.6
3306	93	*	101	28.7	85	*	71	*	96	*	286	92.7	531	3.4
3308	30	*	25	*	20	*	19	*	24	*	80	*	151	3.3
3309	16	*	18	*	10	*	10	*	18	*	34	*	69	*
3312	145	73.1	124	19.4	118	6.8	90	*	116	5.2	408	82.8	760	3.8
3550	22	*	17	*	21	*	15	*	16	*	75	*	128	2.3
3551	10	*	4	*	3	*	7	*	4	*	23	*	39	*

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
3552	19	*	12	*	13	*	11	*	12	*	38	*	75	*
3701	378	81.5	333	21.6	312	11.5	255	22.4	327	7.6	801	85.8	1,655	5.1
3850	1,029	70.4	1,027	16.7	861	9.4	703	15.1	973	6.3	2,370	80.9	4,700	5.8
3950	674	69.3	579	21.9	443	13.5	467	21.0	547	10.1	1,230	86.2	2,705	2.3
3954	2	*	3	*	0	*	3	*	3	*	15	*	21	*
4001	422	59.5	340	17.9	292	8.6	297	16.8	330	6.1	985	64.1	1,917	5.6
4002	350	60.3	301	16.3	266	12.4	242	10.7	294	5.4	852	83.6	1,688	2.9
4003	1	*	0	*	1	*	0	*	0	*	1	*	3	*
4004	0	*	0	*	1	*	0	*	0	*	0	*	1	*
4103	17	*	15	*	15	*	16	*	15	*	60	*	106	10.4
4106	627	69.9	581	13.1	539	8.2	456	9.9	549	5.5	1,713	87.7	3,205	4.3
4107	1,196	69.6	1,084	21.1	861	13.0	842	16.5	1,017	11.4	2,392	86.5	5,071	2.0
4109	358	81.8	328	24.1	261	9.6	280	17.5	318	8.2	819	88.6	1,648	2.2
4112	187	41.2	136	9.6	136	8.1	127	3.9	136	2.9	431	97.9	805	1.6
4113	305	81.3	227	17.6	187	9.6	205	6.3	214	4.7	571	91.9	1,202	0.9
4115	1,156	78.9	1,171	24.6	901	16.9	854	18.7	1,072	10.6	2,734	87.5	5,490	1.2
4351	182	79.1	137	16.8	171	14.6	135	20.0	133	11.3	449	93.5	887	4.5
4352	69	*	39	*	49	*	39	*	38	*	149	80.5	293	11.9

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
4353	154	72.1	114	25.4	120	9.2	100	18.0	110	12.7	338	83.7	685	4.1
4354	243	74.9	192	20.3	214	13.6	161	24.2	188	13.3	572	94.4	1,140	3.5
4355	107	78.5	79	*	81	*	66	*	79	*	247	91.9	484	4.8
4356	172	75.6	139	28.1	149	16.1	124	19.4	134	11.9	367	89.9	746	4.7
4358	47	*	30	*	38	*	25	*	29	*	162	87.0	260	1.5
4359	162	72.8	128	27.3	139	11.5	95	*	120	14.2	444	87.4	795	1.9
4360	70	*	67	*	87	*	45	*	66	*	198	87.9	385	7.0
4361	96	*	70	*	72	*	52	*	69	*	150	89.3	345	3.8
4362	97	*	72	*	65	*	60	*	69	*	208	91.8	408	4.9
4363	294	75.2	252	20.2	248	13.7	209	15.3	252	6.0	675	88.9	1,340	4.1
4401	396	70.5	349	27.8	352	13.9	296	19.9	344	10.8	865	88.3	1,826	1.5
4501	315	74.3	273	19.4	262	12.6	232	11.6	262	3.4	770	93.9	1,492	5.5
4601	557	77.6	433	22.6	384	10.4	377	19.9	411	8.0	1,005	90.5	2,202	2.6
4602	3	*	0	*	0	*	0	*	0	*	5	*	8	*
4701	421	79.6	346	24.0	338	15.7	289	23.2	340	12.9	874	92.9	1,819	1.5
5050	618	60.5	454	15.9	438	8.9	401	11.5	429	4.9	962	87.7	2,296	2.5
5051	886	63.1	621	20.3	577	8.8	568	15.1	587	8.7	1,378	86.0	3,267	2.6
5052	884	58.4	647	12.8	565	7.1	578	9.5	615	5.2	1,406	86.3	3,226	2.2

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan**

Run Date: 05/03/2016

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)**

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
5055	388	36.9	287	5.6	312	1.9	227	0.9	271	0.7	965	57.7	1,843	1.6
5101	710	70.3	619	20.0	580	13.4	450	14.9	566	7.2	1,319	82.2	2,956	3.5
5202	266	78.9	178	32.0	193	20.7	146	29.5	165	14.5	456	89.7	1,019	3.2
5203	15	*	14	*	20	*	6	*	14	*	72	*	119	5.0
5204	17	*	8	*	12	*	6	*	7	*	58	*	97	*
5601	410	77.6	357	22.1	266	11.3	357	24.6	347	10.7	846	91.1	1,746	3.3
5801	618	61.3	462	13.4	408	8.3	420	13.3	452	6.4	1,192	89.5	2,465	5.2
5803	52	*	56	*	59	*	38	*	54	*	197	95.9	340	2.4
5804	9	*	11	*	11	*	9	*	11	*	63	*	95	*
6101	13	*	17	*	17	*	10	*	13	*	51	*	91	*
6102	45	*	39	*	50	*	38	*	36	*	161	90.7	283	4.6
6103	1,362	63.1	874	14.5	938	7.5	914	12.6	810	5.9	2,722	84.8	5,662	6.1
6104	25	*	37	*	37	*	22	*	34	*	30	*	112	4.5
6105	9	*	10	*	12	*	7	*	9	*	39	*	64	*
6301	1,414	61.9	1,159	18.6	788	8.9	961	13.0	1,140	7.8	2,457	87.9	5,592	2.2
6312	793	67.7	713	21.7	494	14.4	561	15.5	694	9.5	1,539	87.1	3,406	1.3
6316	429	65.0	397	21.2	298	12.1	315	16.8	384	9.4	891	82.3	1,879	8.9
6320	53	*	41	*	23	*	35	*	38	*	124	93.5	232	0.4

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breastsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
6325	63	*	47	*	29	*	48	*	46	*	99	*	242	2.9
6330	323	65.9	250	11.6	167	6.6	218	9.6	238	5.0	568	90.0	1,286	5.0
6335	282	59.2	242	14.9	170	7.6	187	5.9	234	3.4	673	94.9	1,274	0.6
6340	23	*	28	*	25	*	24	*	28	*	73	*	149	2.0
6345	9	*	6	*	4	*	6	*	6	*	23	*	39	*
6350	5	*	5	*	6	*	3	*	5	*	14	*	33	*
7304	53	*	47	*	28	*	45	*	40	*	126	88.9	248	2.0
7306	861	58.1	648	13.6	570	5.8	530	9.1	584	5.1	1,600	86.8	3,446	4.9
7308	47	*	44	*	45	*	27	*	32	*	106	88.7	231	1.3
7310	18	*	13	*	4	*	9	*	9	*	26	*	53	*
7401	730	59.2	725	18.6	661	8.2	546	15.8	697	6.2	1,595	87.5	3,291	7.0
7601	208	64.4	170	15.9	124	8.9	149	14.1	165	6.1	417	90.9	881	5.8
7850	401	68.8	361	21.3	305	9.2	257	10.1	338	5.6	849	91.6	1,750	3.5
7852	19	*	24	*	12	*	19	*	20	*	50	*	110	0.9
8101	626	49.7	479	8.1	341	6.5	437	4.1	450	2.0	1,034	68.7	2,358	5.9
8102	186	38.2	144	6.9	117	4.3	133	2.3	140	0.7	566	77.0	981	2.9
8105	595	51.4	454	16.5	334	7.2	409	10.8	448	6.7	904	79.4	2,175	2.6
8106	176	52.3	108	9.3	128	2.3	118	6.8	104	1.9	305	60.0	691	4.5

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breastsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
8150	15	*	28	*	20	*	23	*	28	*	96	*	148	0.0
8151	1,057	81.4	850	32.0	777	22.1	694	20.7	818	14.3	2,265	88.8	4,591	1.6
8152	60	*	47	*	40	*	38	*	44	*	101	84.2	224	3.6
8153	5	*	7	*	7	*	4	*	7	*	30	*	46	*
8155	17	*	17	*	22	*	14	*	16	*	89	*	139	0.0
8157	10	*	9	*	5	*	9	*	8	*	31	*	50	*
8201	962	62.6	743	15.3	587	10.4	619	10.2	723	5.5	1,591	90.0	3,659	3.3
8202	1,259	40.9	935	14.4	781	8.1	814	4.9	891	4.2	2,985	83.7	5,644	2.6
8203	171	70.8	116	17.2	133	6.8	66	*	105	5.7	281	91.8	667	5.2
8250	192	53.6	247	11.7	175	9.7	191	6.8	241	4.1	438	84.5	967	4.0
8252	49	*	25	*	15	*	28	*	24	*	83	*	171	4.7
8255	1,101	61.7	876	14.7	623	7.9	786	10.8	845	4.7	1,693	84.9	3,948	2.9
8257	421	77.2	261	33.7	259	20.8	204	3.9	234	4.3	818	75.4	1,666	0.7
8263	278	57.9	271	10.7	266	5.6	198	10.6	266	3.0	483	87.6	1,169	2.0
8264	265	66.8	153	15.7	80	*	145	9.0	148	2.7	282	83.3	704	3.0
8265	83	*	24	*	6	*	31	*	24	*	54	*	152	1.3
8267	14	*	1	*	0	*	1	*	1	*	2	*	16	*
8901	364	47.0	269	7.4	211	1.9	249	3.6	257	0.4	882	71.4	1,652	7.7

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
8902	355	57.7	282	13.1	259	5.0	266	9.0	273	5.5	682	73.8	1,543	6.4
8903	364	37.1	267	8.2	221	1.8	275	2.9	261	1.9	853	69.9	1,684	7.1
8904	441	53.5	345	11.3	249	6.0	323	5.3	334	2.7	924	69.6	1,918	6.7
8905	498	56.2	443	13.5	360	6.9	353	8.2	417	6.0	1,393	71.8	2,674	3.0
8908	4	*	0	*	0	*	0	*	0	*	5	*	9	*
9012	11	*	6	*	8	*	8	*	6	*	37	*	63	*
9013	6	*	4	*	7	*	3	*	4	*	24	*	46	*
9016	58	*	53	*	38	*	43	*	53	*	186	66.1	324	7.1
9017	11	*	14	*	13	*	12	*	14	*	56	*	91	*
9018	41	*	38	*	32	*	34	*	38	*	152	75.0	253	4.3
9041	3	*	3	*	5	*	2	*	3	*	20	*	30	*
9045	1,381	35.6	658	8.5	374	4.0	693	3.5	640	2.0	1,492	82.1	3,506	5.3
9050	328	65.9	274	16.1	222	5.9	233	6.0	259	3.1	790	78.4	1,631	7.2
9051	384	74.2	371	14.3	287	7.7	269	3.3	362	2.2	1,041	69.1	1,988	2.6
9053	502	49.2	515	14.2	426	8.0	463	7.3	506	5.5	1,258	72.7	2,705	17.1
9055	712	32.3	141	10.6	0	*	288	5.2	134	3.0	0	*	713	2.2
9058	265	64.5	196	15.8	250	12.4	177	7.3	188	3.2	927	92.6	1,601	1.6
9066	395	36.5	326	4.3	282	4.6	277	2.2	323	1.5	818	82.3	1,785	5.8

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 7A
2015 Pediatric Nutrition Surveillance (1)
Michigan**

Run Date: 05/03/2016

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)**

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
9068	318	45.9	322	12.1	273	4.8	292	7.9	316	4.4	754	79.2	1,616	9.3
9069	889	50.7	771	9.3	619	4.7	699	3.7	760	2.1	1,652	74.7	3,855	8.4
9070	351	46.4	377	11.1	318	5.0	316	4.4	363	1.7	917	56.3	1,905	11.7
9072	569	42.7	456	5.9	327	1.8	405	4.0	446	1.8	987	61.5	2,250	9.5
9080	728	59.8	508	11.6	501	6.2	447	5.8	486	2.9	1,835	70.1	3,457	5.2
9081	43	*	42	*	33	*	27	*	41	*	56	*	177	0.0
Michigan	55,956	64.3	44,452	18.4	38,816	10.8	37,300	13.8	42,242	7.3	114,298	84.5	238,131	4.2
Nation (Prior Year)	207,356	66.8	100,563	26.5	83,118	13.7	70,540	14.5	82,222	8.6	283,232	83.6	605,284	7.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾
Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0101	115	8.7	5.2	510	4.9	3.3	8.4	220	18.2	10.0	341	10.9
0102	494	5.1	8.7	2,006	6.5	3.7	11.3	823	18.3	14.0	1,323	17.0
0103	400	5.3	7.0	1,894	6.2	1.6	13.2	875	18.2	15.8	1,362	18.7
0104	142	12.7	10.6	625	7.8	4.0	11.0	281	19.2	16.0	425	16.5
0250	135	10.4	9.6	669	4.0	1.5	8.4	334	19.8	12.6	470	10.2
0251	94	*	*	485	3.9	2.1	8.9	234	23.9	12.4	298	13.8
0252	133	8.3	5.3	592	3.2	1.5	10.6	282	14.2	13.5	406	6.2
0253	149	7.4	8.7	749	1.1	9.6	3.3	365	8.2	3.8	486	13.6
0401	492	8.3	9.3	2,197	7.0	1.8	13.4	992	18.1	16.6	1,431	10.4
0402	405	8.4	7.7	1,622	6.2	2.3	11.7	681	17.5	17.5	981	14.3
0403	121	9.1	7.4	517	5.2	1.5	12.4	228	14.9	18.4	349	16.6
0404	124	7.3	8.1	611	3.6	2.1	9.2	288	15.6	14.6	399	17.8
0501	549	8.0	9.3	2,340	4.9	4.8	11.4	1,041	19.1	15.3	1,528	8.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
0502	4	*	*	16	*	*	*	5	*	*	10	*
0503	134	6.0	11.2	669	7.0	4.8	10.0	322	18.0	12.4	459	6.3
0504	478	6.7	8.8	2,153	6.5	2.7	11.3	904	18.1	13.9	1,473	10.0
0505	300	8.3	9.7	1,329	6.2	3.8	9.4	628	19.7	13.2	931	8.8
0506	168	5.4	9.5	944	5.7	4.3	9.4	420	19.3	12.4	656	6.4
0507	89	*	*	753	5.4	1.9	10.8	405	16.8	13.3	578	10.7
0508	27	*	*	236	7.2	2.1	17.4	135	9.6	21.5	185	2.7
0509	4	*	*	9	*	*	*	5	*	*	6	*
0650	1,095	7.7	9.5	4,442	6.2	2.8	10.8	1,869	18.3	17.7	2,946	8.0
0651	599	9.0	7.0	2,775	6.3	2.2	11.2	1,305	16.9	16.1	1,949	11.8
0652	304	6.6	7.6	1,375	6.8	2.1	11.4	639	19.1	15.6	904	9.7
0653	450	6.9	9.6	2,164	5.0	2.3	10.9	1,055	15.6	13.8	1,505	8.8
0654	355	6.5	11.0	1,760	5.4	2.6	9.4	853	14.5	13.6	1,234	10.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
0655	350	7.4	8.6	1,623	5.6	2.8	8.6	743	18.8	12.9	1,137	6.0
0657	60	*	*	326	7.7	2.8	8.9	169	23.1	13.0	230	11.7
0701	425	6.4	8.0	1,950	6.4	2.5	6.8	906	18.9	8.7	1,419	10.5
0702	89	*	*	400	7.5	2.5	9.3	183	15.8	12.6	271	7.7
0703	230	6.1	9.1	1,017	4.0	4.2	9.8	473	16.7	13.1	632	8.2
0704	61	*	*	288	4.2	2.1	8.7	156	21.2	9.6	201	3.0
0705	118	5.9	9.3	546	5.3	2.2	9.5	280	14.3	11.8	359	6.1
0850	1,062	7.1	7.9	4,469	5.6	3.4	11.4	1,909	18.6	15.2	2,915	6.5
0851	796	7.0	7.5	3,748	4.8	3.0	9.3	1,751	17.8	13.0	2,595	7.7
0901	1,597	8.5	9.2	7,443	7.4	1.8	11.5	3,346	17.0	16.0	5,235	7.8
0902	45	*	*	276	4.0	2.9	8.0	157	13.4	9.6	197	15.7
1051	1,514	8.4	10.2	5,702	4.2	5.4	9.7	2,515	17.8	12.9	3,627	16.1
1101	1,974	10.5	5.9	7,767	5.1	4.5	9.1	3,367	15.8	13.4	5,015	14.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾

Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1102	1,001	8.0	8.9	3,868	4.9	2.6	10.1	1,637	16.5	13.1	2,481	7.1
1106	138	6.5	8.7	549	5.8	2.0	9.5	235	27.2	11.1	346	6.6
1250	1,031	8.1	8.2	4,666	7.1	2.0	14.0	2,183	17.9	16.4	3,129	4.1
1251	972	7.8	7.4	4,082	7.3	1.8	11.3	1,731	18.8	14.0	2,599	3.2
1252	453	9.3	8.2	2,118	6.1	2.2	13.3	972	19.4	18.3	1,480	7.6
1253	1,103	9.0	6.7	4,492	6.6	2.2	10.5	1,950	16.8	13.9	2,989	9.0
1350	2,596	8.9	6.6	11,040	7.3	2.5	13.7	4,932	17.6	15.9	7,496	19.2
1351	145	7.6	9.0	745	6.6	2.8	11.9	344	14.5	15.4	527	13.7
1352	358	10.3	5.6	1,745	6.5	2.1	13.6	840	19.8	17.3	1,243	16.7
1401	274	7.7	9.5	1,068	4.4	5.6	10.2	433	15.5	16.4	718	8.4
1402	166	7.2	12.7	734	4.8	4.8	10.5	343	18.1	12.8	486	9.7
1403	14	*	*	111	2.7	2.7	15.3	68	*	*	80	*
1404	5	*	*	39	*	*	*	20	*	*	30	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/03/2016

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1501	933	6.1	8.8	4,665	3.6	2.6	13.5	2,271	18.4	18.3	3,362	16.6
1503	1,239	7.3	8.6	5,719	4.7	2.0	10.7	2,634	18.6	14.8	3,942	16.0
1504	2,370	7.6	8.1	10,464	4.7	2.5	11.9	4,872	17.1	15.2	7,307	36.2
1505	732	7.5	10.0	3,653	5.0	2.6	7.9	1,690	15.2	9.3	2,565	25.8
1506	643	6.8	8.6	3,153	6.5	2.4	12.5	1,482	18.6	15.0	2,224	19.4
1507	615	6.8	8.1	2,810	4.3	2.9	8.6	1,247	14.2	10.8	1,942	24.6
1508	71	*	*	469	4.7	0.9	18.1	273	22.7	22.7	366	18.9
1509	461	8.2	8.2	2,578	3.6	3.3	12.7	1,347	16.6	17.3	1,895	23.4
1511	770	7.8	11.8	3,272	4.1	2.4	9.8	1,482	18.1	15.4	2,178	21.0
1512	345	9.0	7.8	1,842	4.9	2.8	11.0	892	19.2	12.8	1,268	18.3
1701	748	4.7	11.1	3,315	5.6	2.5	12.8	1,523	17.1	14.1	2,119	15.2
1950	628	6.4	6.5	2,688	6.9	1.2	10.0	1,166	19.4	15.3	1,855	6.7
1952	127	8.7	8.7	579	5.5	0.5	10.0	238	16.8	14.3	398	5.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/03/2016

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
1953	334	9.0	6.3	1,571	7.1	1.2	11.7	711	18.8	13.8	1,113	2.8
1954	653	10.4	10.4	2,834	5.5	2.3	10.6	1,194	15.7	14.6	1,844	4.3
1955	101	8.9	6.9	599	5.3	2.7	13.7	297	19.2	16.2	440	2.7
1957	585	6.2	7.5	2,823	6.4	1.6	11.1	1,251	19.3	12.5	1,885	2.9
2150	768	7.7	12.1	3,223	5.3	1.7	10.1	1,387	17.6	13.3	2,139	2.6
2151	273	6.2	8.4	1,130	6.0	1.6	10.8	475	17.7	14.9	709	2.8
2251	464	10.1	9.9	2,030	7.2	1.5	9.7	895	16.0	12.0	1,366	7.9
2252	186	10.8	4.8	929	4.8	1.5	12.5	470	19.8	14.9	657	5.6
2554	117	6.8	6.8	593	4.4	2.2	11.5	320	17.2	15.6	422	7.8
2555	8,580	12.2	5.2	32,722	8.3	2.0	10.7	13,163	16.1	13.2	20,970	21.5
3001	995	12.7	4.5	4,522	7.7	1.7	12.0	2,202	18.1	13.2	3,196	5.0
3002	267	10.9	7.1	1,353	6.0	1.8	14.3	698	20.8	15.5	981	16.1
3004	129	1.6	3.1	657	4.3	0.9	21.6	356	25.3	28.4	501	5.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
3007	17	*	*	116	5.2	1.7	16.4	63	*	*	89	*
3009	1	*	*	24	*	*	*	20	*	*	31	*
3201	488	7.8	9.8	2,330	6.4	2.1	14.2	1,094	16.9	18.2	1,619	6.9
3301	3,996	9.6	6.6	15,349	6.0	2.5	9.8	6,354	16.1	13.0	10,061	13.2
3302	0	*	*	2	*	*	*	2	*	*	1	*
3304	148	10.1	12.2	737	3.5	1.1	10.7	343	14.6	11.7	512	9.2
3305	200	6.0	9.0	919	5.0	2.1	9.0	412	14.1	8.7	624	7.7
3306	323	7.7	10.5	1,633	5.4	0.9	8.6	753	19.1	8.8	1,151	6.4
3308	95	*	*	408	8.6	0.7	10.5	156	15.4	11.5	279	4.7
3309	46	*	*	200	7.0	0.5	14.5	85	*	*	143	4.9
3312	418	7.9	4.1	2,277	6.8	2.1	13.8	1,137	15.4	18.6	1,642	11.2
3550	73	*	*	400	4.8	1.5	20.5	188	26.6	26.6	259	26.3
3551	18	*	*	127	3.1	3.9	28.3	72	*	*	75	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
3552	53	*	*	236	4.7	4.7	20.3	109	22.9	26.6	133	24.1
3701	1,171	6.7	7.1	5,052	4.5	3.7	9.9	2,241	16.3	15.1	3,465	8.3
3850	3,203	8.4	7.7	14,132	4.9	4.0	9.7	6,491	19.8	15.3	9,600	20.9
3950	2,131	9.8	9.4	8,293	5.8	3.3	10.7	3,473	17.6	13.4	5,468	24.7
3954	11	*	*	136	0.0	0.0	11.8	88	*	*	105	20.0
4001	1,292	10.2	5.9	5,776	4.9	3.0	10.7	2,719	15.8	13.2	4,106	35.2
4002	1,093	10.3	7.4	5,075	7.1	1.7	13.6	2,358	17.1	15.2	3,582	17.3
4003	1	*	*	2	*	*	*	0	*	*	2	*
4004	0	*	*	1	*	*	*	0	*	*	1	*
4103	69	*	*	369	5.1	8.1	6.8	201	11.9	8.0	199	34.7
4106	1,982	11.4	5.8	10,101	7.7	1.6	11.7	4,876	17.0	14.3	7,063	15.1
4107	3,563	10.9	7.8	15,263	7.5	1.6	10.4	6,652	16.5	12.6	10,241	13.2
4109	1,097	10.3	8.6	5,175	5.3	2.2	8.4	2,361	14.5	11.4	3,503	7.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾

Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
4110	0	*	*	2	*	*	*	2	*	*	2	*
4112	489	5.1	8.8	2,404	6.7	3.2	18.6	1,250	19.0	22.5	1,922	9.5
4113	827	8.5	7.3	3,469	6.0	2.2	12.3	1,529	20.2	18.8	2,297	4.4
4115	1,716	9.7	7.5	6,550	7.4	2.2	10.4	2,899	15.7	12.3	4,248	9.1
4120	1,215	8.8	8.2	5,895	6.0	1.5	10.5	2,653	15.3	12.6	4,049	7.6
4160	1,102	9.3	7.8	5,446	6.5	2.4	10.4	2,443	16.6	13.0	3,920	11.8
4351	560	6.1	8.8	2,667	5.3	1.9	10.3	1,300	17.3	14.2	1,846	19.0
4352	193	8.8	7.8	890	7.0	1.5	10.7	416	18.5	17.3	638	13.3
4353	458	9.0	7.9	2,136	7.3	2.4	13.5	1,044	20.1	18.5	1,525	13.2
4354	752	8.9	8.4	3,419	6.1	1.3	9.7	1,530	19.9	14.3	2,372	11.7
4355	304	5.6	8.2	1,417	5.4	3.2	10.0	645	17.7	18.0	990	12.3
4356	515	8.2	9.1	2,264	7.6	1.6	11.4	1,053	20.2	16.1	1,574	18.0
4358	131	11.5	6.9	751	9.5	1.5	22.1	408	25.7	27.5	583	22.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/03/2016

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
4359	502	6.4	12.2	2,478	6.4	1.5	14.5	1,234	17.4	16.9	1,744	13.4
4360	236	6.8	10.2	1,105	4.2	1.7	7.2	506	21.7	12.3	755	11.3
4361	254	9.8	7.1	1,016	9.0	1.1	13.1	435	23.9	15.2	687	11.4
4362	294	7.1	8.8	1,331	7.4	0.8	11.5	595	20.2	17.8	911	16.7
4363	959	6.6	8.9	4,120	5.4	1.3	8.9	1,843	19.1	13.3	2,817	12.7
4401	1,264	7.7	8.6	5,612	5.9	2.1	9.6	2,397	19.4	14.8	3,841	11.6
4501	992	7.7	6.9	4,442	6.8	2.3	9.9	2,028	17.6	14.3	3,092	6.6
4601	1,701	8.1	7.6	6,460	6.5	2.2	11.5	2,620	18.3	17.2	3,724	10.5
4602	3	*	*	8	*	*	*	5	*	*	5	*
4701	1,355	8.3	9.4	5,616	6.3	1.4	9.2	2,399	17.1	13.2	3,793	6.4
5050	1,959	10.0	6.8	7,175	6.8	1.5	10.3	2,812	16.6	11.8	4,516	18.7
5051	2,795	11.2	6.5	10,126	6.7	1.3	11.6	3,936	14.3	14.1	6,299	19.1
5052	2,642	10.9	6.1	10,012	6.7	1.9	10.8	4,090	15.2	14.2	6,396	16.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
5055	839	8.6	5.1	4,763	3.2	2.1	12.1	2,235	17.4	15.7	3,405	32.6
5101	2,217	9.1	8.7	8,857	6.0	1.8	10.2	3,611	18.2	15.4	5,784	15.7
5202	684	6.7	9.6	2,628	6.8	2.1	10.7	1,126	19.8	13.6	1,488	7.3
5203	87	*	*	451	5.8	1.3	15.3	215	15.3	16.7	281	7.8
5204	85	*	*	400	7.5	2.0	12.8	194	12.9	17.5	248	6.0
5601	1,174	9.3	9.3	5,456	7.0	1.3	11.6	2,509	18.4	15.5	3,500	8.7
5801	1,884	7.8	8.3	7,804	7.6	2.4	10.9	3,337	16.3	15.0	5,090	8.3
5803	175	7.4	9.1	1,021	6.7	1.3	11.0	535	14.8	14.4	757	9.0
5804	35	*	*	305	5.2	3.0	9.8	161	14.9	8.1	232	6.5
6101	54	*	*	324	5.6	1.5	11.4	156	16.0	12.8	243	14.4
6102	200	7.5	10.5	1,051	4.9	2.7	8.8	530	15.3	11.3	764	10.6
6103	3,888	11.4	6.6	16,392	6.0	2.6	10.5	7,538	16.1	13.1	11,126	17.6
6104	330	16.4	4.5	1,181	6.8	2.6	8.7	428	15.0	11.0	738	22.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/03/2016

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
6105	42	*	*	244	8.2	1.6	13.1	132	21.2	15.2	178	7.9
6301	4,448	11.4	6.1	17,475	8.2	1.7	10.0	7,021	16.0	13.1	11,043	16.0
6312	2,543	11.2	6.3	10,447	7.9	2.2	8.2	4,161	15.5	9.8	6,871	18.4
6316	1,388	9.5	6.6	6,007	5.3	1.6	10.1	2,537	15.6	13.8	3,933	10.5
6320	155	14.8	5.2	815	5.8	3.7	8.2	387	12.4	9.8	582	12.0
6325	158	8.9	9.5	677	6.1	2.5	8.7	308	16.9	12.0	446	10.5
6330	868	13.7	3.8	3,543	6.3	3.4	8.0	1,558	15.7	10.2	2,331	19.6
6335	642	8.3	6.9	2,981	3.5	3.9	8.8	1,476	11.1	12.9	2,158	13.3
6340	79	*	*	429	3.7	3.0	9.3	219	14.6	12.3	315	15.6
6345	19	*	*	111	3.6	3.6	11.7	59	*	*	90	*
6350	143	8.4	11.9	782	3.5	3.6	10.4	366	14.8	16.4	570	15.1
6877	1	*	*	14	*	*	*	7	*	*	11	*
7304	176	10.8	9.1	843	5.7	3.2	13.5	396	21.2	19.7	596	8.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Run Date: 05/03/2016

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾												
Clinic	Birth Weight ⁽⁴⁾			Height and Weight ^(5,6)							Anemia Low Hb/Hct ⁽⁷⁾	
	Low	High		Short Stature	Under- Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
7306	2,627	10.0	7.1	10,852	6.4	2.2	12.6	4,694	15.9	15.5	7,339	17.6
7308	173	6.4	11.6	806	5.3	2.6	10.9	364	18.7	12.4	582	13.7
7310	35	*	*	149	2.7	2.7	9.4	66	*	*	94	*
7401	2,443	7.8	7.7	10,213	5.3	1.9	9.5	4,418	17.5	15.9	7,041	8.2
7601	651	8.9	8.1	2,696	5.3	1.8	9.0	1,174	21.0	14.4	1,804	8.8
7850	1,252	7.3	8.9	5,255	5.2	3.9	9.4	2,305	18.5	14.8	3,588	13.0
7852	67	*	*	380	2.6	2.9	8.9	202	16.8	12.9	278	9.4
8101	1,766	17.5	3.4	6,187	6.7	5.3	8.8	2,283	13.1	11.2	3,870	33.8
8102	544	6.4	10.1	2,653	5.4	3.8	17.7	1,350	19.9	23.3	1,915	26.7
8105	1,598	12.0	6.2	5,235	5.6	4.5	9.2	1,816	14.9	11.2	2,723	30.4
8106	457	14.0	4.2	1,803	7.2	1.4	12.1	808	19.9	15.7	1,215	20.6
8150	60	*	*	411	3.4	3.2	8.5	222	13.1	13.1	318	21.1
8151	3,265	10.8	7.1	14,103	6.2	1.9	10.6	6,249	17.3	14.7	9,792	17.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
8152	199	11.6	6.0	780	6.5	1.4	11.7	301	19.6	13.0	521	26.1
8153	19	*	*	155	7.7	0.0	19.4	92	*	*	122	17.2
8155	62	*	*	436	3.9	2.5	12.2	257	9.7	14.4	342	25.1
8157	21	*	*	138	2.9	5.1	5.8	94	*	*	101	28.7
8201	3,030	9.5	6.7	11,351	6.8	1.9	11.8	4,313	18.3	15.8	6,712	15.3
8202	3,356	7.8	5.2	16,278	4.6	2.1	9.4	7,739	12.9	11.1	11,191	21.5
8203	452	9.1	5.5	1,713	5.1	1.2	8.6	688	15.7	10.0	995	13.4
8250	920	10.0	6.2	3,805	6.8	2.4	10.5	1,584	14.2	12.7	2,254	22.0
8252	128	12.5	2.3	546	8.6	3.1	8.1	280	14.3	11.1	390	28.2
8255	3,211	9.3	7.9	12,001	6.3	1.7	9.3	4,576	17.0	11.9	7,132	14.4
8257	893	11.4	3.2	3,861	4.8	4.3	10.6	1,791	14.3	16.0	2,524	19.8
8258	465	11.8	8.0	1,780	6.2	2.7	10.4	729	19.5	16.5	1,007	18.7
8260	143	8.4	3.5	934	4.1	2.0	11.1	478	16.7	15.3	687	15.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Combined 3 Years

Comparison of Growth and Anemia Indicators ⁽³⁾

Clinic	Birth Weight (4)			Height and Weight (5,6)						Anemia Low Hb/Hct (7)		
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	%	%	Number	%
8263	738	9.9	7.0	2,469	7.4	1.5	9.6	899	16.6	12.3	1,434	14.7
8264	275	8.4	7.3	707	7.8	1.3	10.5	257	14.0	9.3	366	15.3
8265	83	*	*	150	6.7	1.3	7.3	53	*	*	62	*
8267	14	*	*	16	*	*	*	2	*	*	2	*
8901	970	13.7	2.3	4,695	6.3	5.1	8.6	2,261	14.7	10.2	3,312	38.3
8902	1,052	14.1	3.4	4,336	6.4	5.9	8.2	1,768	10.5	10.6	2,845	36.6
8903	1,084	12.5	4.0	4,892	6.6	3.7	11.0	2,313	16.1	14.3	3,332	39.6
8904	1,138	11.7	4.1	5,328	4.3	5.0	8.9	2,451	12.3	11.1	3,662	38.6
8905	2,009	8.5	6.5	9,666	6.0	4.4	13.1	4,545	15.8	15.0	6,698	28.1
8908	4	*	*	6	*	*	*	4	*	*	5	*
9012	35	*	*	248	2.0	6.9	7.3	146	15.8	8.2	151	31.1
9013	28	*	*	199	3.5	4.5	4.5	105	13.3	3.8	65	*
9015	6	*	*	49	*	*	*	24	*	*	28	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
9016	193	17.6	4.1	1,017	6.6	2.9	10.1	534	18.7	10.3	622	42.3
9017	39	*	*	288	6.9	2.4	8.3	158	18.4	13.9	186	32.8
9018	116	13.8	3.4	773	3.5	4.4	9.4	420	17.1	13.6	112	30.4
9019	8	*	*	75	*	*	*	42	*	*	22	*
9021	16	*	*	94	*	*	*	50	*	*	51	*
9041	15	*	*	96	*	*	*	52	*	*	10	*
9045	3,446	12.8	5.1	9,394	5.1	5.5	12.7	3,799	16.8	18.6	5,524	16.9
9050	1,233	11.9	4.7	5,634	6.0	2.3	10.3	2,578	16.8	14.0	3,784	22.1
9051	1,278	8.1	7.4	6,319	7.7	1.4	19.6	2,920	19.7	22.8	4,194	22.9
9053	2,377	14.6	3.5	9,375	6.5	3.2	6.8	3,700	15.5	10.2	6,167	26.0
9055	1,102	9.0	3.5	1,024	7.0	14.6	2.7	2	*	*	4	*
9058	878	13.1	4.8	4,928	4.9	3.1	9.2	2,528	14.4	11.0	3,562	18.3
9066	1,406	13.9	3.6	6,039	6.4	2.9	9.4	2,655	14.4	12.1	4,006	16.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3)												
Clinic	Birth Weight (4)			Height and Weight (5,6)							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature	Under-Weight	Obese	>= 2 Yrs Overweight and obese					
	Number	%	%	Number	%	%	%	Number	% Overweight	% Obese	Number	%
9068	1,232	13.7	3.7	5,134	6.1	3.5	8.6	2,228	13.3	10.1	3,416	25.1
9069	2,867	14.1	3.9	11,345	7.0	3.5	8.3	4,479	13.2	11.6	7,293	23.5
9070	1,399	15.3	3.4	6,110	6.0	4.3	8.6	2,754	11.7	11.7	4,087	37.9
9072	1,841	13.0	3.6	6,981	6.8	2.8	7.9	2,861	14.0	10.9	4,439	32.3
9080	2,143	13.3	4.3	10,320	5.6	2.9	9.4	5,153	14.7	11.1	7,470	28.0
9081	165	13.9	6.1	399	10.5	6.5	11.8	119	16.0	15.1	146	20.5
Michigan	171,231	10.0	6.9	723,225	6.2	2.6	10.6	317,700	16.6	14.0	480,706	17.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.

(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
0101	115	75.7	94	*	80	*	77	*	89	*	250	90.4	519	5.8
0102	495	59.0	395	16.7	338	11.2	343	18.1	379	8.2	943	84.1	2,036	5.4
0103	400	68.8	360	19.4	301	9.0	289	18.0	343	6.4	982	83.8	1,932	6.1
0104	143	72.7	132	22.7	105	10.5	96	*	120	6.7	317	81.4	642	8.3
0250	135	71.1	140	19.3	131	10.7	102	20.6	134	6.0	375	87.2	694	6.2
0251	94	*	71	*	90	*	69	*	71	*	255	85.9	490	9.0
0252	134	74.6	81	*	107	12.1	52	*	77	*	305	79.0	602	6.3
0253	151	57.0	126	14.3	121	7.4	120	18.3	125	8.8	404	77.0	754	13.1
0401	494	54.9	403	18.9	395	8.6	389	18.0	393	11.2	1,118	90.7	2,233	3.8
0402	407	71.7	299	22.1	298	13.4	252	21.8	284	9.5	772	80.8	1,648	6.8
0403	120	64.2	98	*	102	2.9	87	*	92	*	266	91.4	528	13.6
0404	124	52.4	107	21.5	113	7.1	88	*	103	14.6	324	88.9	621	13.4
0501	583	72.4	405	23.7	394	12.2	352	19.0	399	13.3	1,145	87.2	2,366	6.6
0502	4	*	6	*	6	*	3	*	5	*	5	*	16	*
0503	136	87.5	119	42.0	101	22.8	90	*	112	22.3	363	87.3	679	4.0
0504	506	81.4	389	36.5	376	21.5	338	36.7	375	19.5	1,051	87.5	2,196	4.9
0505	319	77.7	223	22.9	202	13.9	197	26.4	215	8.4	702	87.2	1,382	5.6

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
0506	209	72.2	190	15.8	170	10.6	141	22.7	183	7.7	480	80.2	972	10.6
0507	95	*	118	23.7	137	19.0	93	*	118	5.9	443	89.8	768	6.4
0508	29	*	35	*	41	*	31	*	35	*	147	86.4	238	2.5
0509	4	*	1	*	0	*	1	*	1	*	6	*	10	*
0650	1,102	74.4	880	22.8	836	13.5	700	20.7	842	11.9	2,110	86.1	4,523	4.7
0651	600	68.7	511	16.6	532	7.9	432	16.4	491	6.1	1,470	87.8	2,838	9.5
0652	306	73.2	226	16.4	239	7.1	205	17.6	213	6.6	723	87.6	1,405	8.3
0653	453	78.4	360	21.9	386	16.3	263	25.1	337	15.1	1,179	94.4	2,195	10.7
0654	355	78.3	295	24.4	319	12.5	269	20.4	293	8.9	966	90.1	1,796	7.0
0655	350	71.7	340	20.3	298	11.7	242	16.5	323	8.4	843	85.4	1,657	6.3
0657	61	*	41	*	50	*	40	*	40	*	187	90.9	329	11.9
0701	428	80.8	362	41.7	294	18.4	288	31.3	333	20.1	1,020	91.2	2,002	1.5
0702	88	*	74	*	64	*	64	*	72	*	216	85.6	415	3.9
0703	225	72.0	177	24.3	129	16.3	150	21.3	170	12.4	523	83.4	1,036	5.0
0704	61	*	33	*	38	*	29	*	33	*	170	87.6	293	6.5
0705	119	81.5	90	*	67	*	73	*	84	*	313	88.2	562	3.6
0850	1,093	78.7	985	23.6	777	11.2	734	19.3	918	9.9	2,146	83.6	4,558	3.5

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
0851	807	75.7	743	23.4	642	10.6	520	24.0	663	13.1	1,986	88.1	3,866	3.9
0901	1,622	60.7	1,330	16.3	1,396	8.9	1,192	15.5	1,270	4.4	3,809	92.3	7,604	4.6
0902	45	*	40	*	35	*	36	*	41	*	167	90.4	281	7.8
1051	1,517	84.0	1,013	30.9	1,061	19.2	766	29.4	968	12.6	2,789	87.7	5,976	3.2
1101	1,982	57.2	1,408	14.8	1,289	9.1	1,163	14.2	1,297	6.0	3,766	74.9	7,957	6.0
1102	1,008	62.4	793	20.2	653	9.3	652	21.5	739	9.7	1,819	80.2	3,941	4.9
1106	138	73.9	109	30.3	89	*	91	*	103	13.6	265	83.8	562	5.3
1250	1,036	76.7	797	23.5	839	12.0	701	20.4	753	11.2	2,472	80.3	4,798	5.1
1251	981	77.1	772	22.0	628	13.4	657	21.6	742	13.2	1,977	82.4	4,153	6.6
1252	454	81.7	382	22.5	352	13.1	309	16.2	336	6.8	1,100	81.3	2,165	3.0
1253	1,111	77.0	832	22.2	702	10.8	673	16.3	732	6.1	2,192	83.1	4,572	4.0
1350	2,611	67.0	1,992	14.7	1,984	8.8	1,574	11.7	1,721	5.4	5,475	86.6	11,235	3.9
1351	147	73.5	137	27.7	120	16.7	110	17.3	127	3.9	397	93.5	760	2.2
1352	361	58.4	299	9.4	321	4.7	224	12.1	266	4.9	949	90.1	1,803	2.5
1401	274	82.5	212	25.5	169	18.9	151	17.9	202	9.4	494	84.2	1,103	7.3
1402	166	90.4	126	42.9	107	29.0	108	30.6	120	15.0	385	82.9	747	1.7
1403	14	*	11	*	9	*	9	*	11	*	75	*	114	7.9

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
1404	5	*	8	*	4	*	6	*	7	*	20	*	38	*
1501	959	69.3	805	20.4	814	12.3	680	16.3	755	6.9	2,531	90.8	4,746	2.3
1503	1,247	73.5	1,012	25.5	1,020	15.1	890	21.9	944	13.7	2,963	89.6	5,773	2.1
1504	2,440	75.6	1,263	23.0	1,896	11.0	736	21.7	1,038	9.4	5,340	84.6	10,641	1.8
1505	787	81.3	613	31.3	674	19.4	513	27.7	534	15.7	1,875	90.3	3,692	1.8
1506	654	72.2	562	21.5	583	15.4	487	23.4	530	9.1	1,699	93.0	3,239	1.9
1507	640	79.2	479	28.6	518	17.0	415	28.0	450	15.8	1,390	87.1	2,869	3.3
1508	78	*	54	*	73	*	24	*	40	*	284	92.6	475	0.0
1509	491	72.3	391	25.1	434	10.4	261	14.6	291	9.3	1,443	84.9	2,624	3.9
1511	775	80.9	598	24.2	564	13.5	510	22.2	544	9.4	1,621	90.0	3,329	2.7
1512	377	76.1	270	23.0	326	11.7	196	26.5	244	9.4	1,001	88.5	1,879	3.3
1701	755	66.8	615	27.6	499	16.0	517	27.5	591	19.3	1,733	85.1	3,402	13.7
1950	632	79.9	521	23.8	402	14.9	410	23.7	482	11.8	1,347	86.1	2,761	7.6
1952	127	75.6	110	18.2	103	8.7	88	*	103	9.7	273	85.3	587	5.1
1953	336	78.3	298	25.5	229	10.5	232	27.2	268	12.7	793	84.6	1,600	8.3
1954	653	82.4	560	26.1	436	14.7	492	20.7	543	10.1	1,357	87.9	2,854	4.8
1955	101	70.3	100	27.0	90	*	77	*	95	*	348	88.2	608	5.1

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
1957	596	83.2	566	27.2	405	16.5	417	22.1	522	11.7	1,384	86.4	2,845	2.9
2150	771	73.7	606	17.8	639	9.9	556	18.0	584	9.1	1,593	86.2	3,271	3.9
2151	274	68.2	225	18.7	222	11.7	195	20.0	214	10.7	531	91.0	1,143	4.7
2251	465	72.5	393	25.4	353	15.6	306	27.5	371	14.0	1,021	89.0	2,071	2.1
2252	186	71.5	134	20.9	149	14.8	135	24.4	132	8.3	518	87.3	952	5.8
2554	118	35.6	72	*	103	6.8	45	*	66	*	342	86.8	605	5.8
2555	8,612	57.1	6,941	13.8	5,488	7.2	5,332	11.5	5,997	5.6	14,458	77.2	33,025	4.9
3001	998	59.4	748	8.6	709	5.5	661	4.7	720	1.8	2,470	82.6	4,646	6.9
3002	268	51.9	225	5.3	189	3.2	188	6.9	219	0.9	782	94.5	1,381	3.0
3004	129	73.6	97	*	97	*	84	*	96	*	398	75.4	687	0.7
3007	16	*	16	*	17	*	14	*	16	*	71	*	117	0.0
3009	1	*	5	*	5	*	2	*	4	*	23	*	33	*
3201	491	68.6	393	20.6	304	11.8	348	17.0	375	6.7	1,204	84.6	2,364	5.2
3301	4,019	75.8	2,906	22.8	2,626	13.8	2,161	13.8	2,617	6.3	7,090	82.1	15,518	4.1
3302	0	*	0	*	0	*	0	*	0	*	2	*	2	*
3304	155	91.0	129	62.0	131	38.9	91	*	117	21.4	389	85.1	748	0.7
3305	203	80.8	173	34.7	147	17.0	127	18.9	161	8.7	456	90.8	925	1.5

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
3306	330	77.6	287	27.2	268	19.0	218	18.3	263	5.7	843	92.5	1,641	3.5
3308	96	*	81	*	73	*	59	*	72	*	178	89.9	413	3.4
3309	47	*	42	*	31	*	27	*	42	*	103	88.3	204	3.9
3312	423	71.6	379	18.7	362	9.1	280	11.8	339	3.8	1,250	83.1	2,295	3.4
3550	73	*	56	*	72	*	51	*	55	*	214	85.5	406	1.5
3551	18	*	15	*	15	*	15	*	14	*	81	*	127	1.6
3552	53	*	39	*	42	*	35	*	37	*	122	88.5	239	4.6
3701	1,178	75.5	1,015	20.4	980	11.7	846	19.6	994	9.5	2,515	81.4	5,154	6.1
3850	3,218	68.1	3,083	16.5	2,606	8.9	2,104	14.7	2,777	6.2	7,290	81.7	14,523	5.9
3950	2,151	69.0	1,719	23.7	1,402	12.4	1,365	20.3	1,573	9.6	3,946	85.6	8,536	2.6
3954	11	*	15	*	15	*	11	*	12	*	97	*	141	0.7
4001	1,312	60.6	1,028	17.7	877	9.6	857	13.8	957	6.4	3,021	61.3	5,904	6.1
4002	1,101	63.4	877	20.2	758	12.3	792	16.7	851	8.1	2,661	78.6	5,192	4.4
4003	1	*	0	*	1	*	0	*	0	*	1	*	3	*
4004	0	*	0	*	1	*	0	*	0	*	0	*	1	*
4103	69	*	49	*	45	*	45	*	50	*	229	84.3	382	7.6
4106	2,005	67.4	1,906	14.0	1,691	8.5	1,462	8.3	1,768	6.0	5,467	84.7	10,232	4.0

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
4107	3,580	70.3	3,086	21.1	2,424	12.0	2,413	17.5	2,826	12.3	7,524	84.4	15,453	2.3
4109	1,117	73.8	1,034	23.1	866	11.4	853	18.9	975	10.2	2,638	85.9	5,215	3.0
4110	0	*	0	*	0	*	0	*	0	*	2	*	2	*
4112	489	33.5	404	12.9	401	7.7	332	5.4	377	4.2	1,426	93.2	2,474	2.0
4113	827	82.2	602	18.6	557	11.0	557	6.5	572	5.1	1,744	86.2	3,534	1.4
4115	1,730	77.9	1,396	24.4	1,048	16.3	1,013	20.0	1,272	10.8	3,203	87.3	6,737	1.2
4120	1,224	73.0	1,248	17.6	994	10.3	955	12.6	1,111	6.8	2,972	86.2	6,002	1.8
4160	1,111	70.7	1,082	28.4	863	17.6	819	20.5	983	13.5	2,798	83.2	5,641	1.6
4351	556	77.3	413	17.9	469	13.2	382	20.4	398	12.1	1,434	90.2	2,718	4.8
4352	193	71.5	138	20.3	166	7.2	134	14.9	134	10.4	493	79.9	932	13.4
4353	458	73.6	344	19.5	368	10.1	291	16.5	326	7.4	1,172	84.6	2,210	4.6
4354	757	70.7	640	23.1	647	12.2	552	25.4	624	16.7	1,724	92.1	3,500	3.0
4355	304	75.3	252	23.0	285	10.5	194	20.6	241	5.8	760	91.6	1,484	5.3
4356	517	73.5	388	26.5	440	13.4	343	21.9	376	9.6	1,212	90.8	2,373	4.9
4358	133	85.0	122	26.2	131	16.0	100	7.0	119	3.4	466	84.8	778	1.7
4359	501	75.2	382	28.3	430	14.0	288	17.4	353	11.9	1,377	79.7	2,522	3.3
4360	237	84.0	210	24.8	222	15.3	177	18.6	208	6.3	568	85.2	1,125	7.4

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
4361	254	80.3	192	20.8	195	10.8	161	12.4	189	5.8	482	86.5	1,034	4.0
4362	294	84.7	234	30.3	264	18.2	196	18.4	228	7.5	689	89.0	1,358	6.9
4363	959	78.8	781	20.5	799	12.5	659	15.3	762	5.8	2,093	85.0	4,200	5.4
4401	1,263	70.2	1,066	23.5	1,052	12.2	895	22.0	1,016	10.2	2,702	88.5	5,680	2.6
4501	994	72.5	790	18.4	771	12.5	675	13.0	752	4.3	2,325	90.4	4,536	4.7
4601	1,710	74.6	1,348	19.6	1,177	9.8	1,120	18.2	1,282	6.8	2,980	81.2	6,520	4.1
4602	3	*	0	*	0	*	0	*	0	*	5	*	8	*
4701	1,360	77.7	1,097	21.6	1,053	12.8	872	21.1	1,048	10.1	2,679	91.3	5,742	2.2
5050	1,961	60.5	1,372	15.3	1,335	7.9	1,222	11.5	1,267	6.1	3,098	84.0	7,225	2.9
5051	2,799	63.6	1,941	16.8	1,844	7.9	1,718	11.9	1,834	7.1	4,324	81.6	10,238	2.8
5052	2,650	56.7	1,877	12.7	1,788	6.9	1,666	8.0	1,785	4.3	4,542	81.5	10,124	2.5
5055	1,055	29.7	839	11.2	783	4.7	687	1.7	777	1.8	2,559	53.8	4,842	3.0
5101	2,231	66.9	1,811	22.4	1,726	11.4	1,451	15.5	1,683	8.4	4,053	82.4	9,020	3.0
5202	689	78.2	481	29.5	504	17.3	388	26.8	452	11.5	1,305	88.4	2,787	3.4
5203	89	*	65	*	95	*	41	*	59	*	260	90.0	488	5.9
5204	85	*	55	*	90	*	42	*	52	*	229	87.3	437	3.4
5601	1,175	77.7	1,089	24.5	835	14.7	1,028	24.7	1,064	10.6	2,800	88.5	5,481	3.7

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
5801	1,884	63.3	1,439	14.3	1,391	8.6	1,271	12.6	1,361	6.4	3,788	85.3	7,901	5.8
5803	174	63.2	166	31.9	171	15.2	119	23.5	158	14.6	595	91.4	1,042	3.2
5804	35	*	48	*	54	*	35	*	45	*	183	92.9	304	3.9
6101	54	*	54	*	62	*	41	*	48	*	194	88.1	342	9.9
6102	200	76.5	150	27.3	188	16.0	138	23.2	139	8.6	596	89.1	1,076	4.6
6103	3,903	61.2	2,472	14.2	2,823	7.2	2,533	11.8	2,349	5.5	8,320	84.4	16,774	6.2
6104	333	51.4	204	9.3	225	3.1	197	5.6	200	2.5	495	71.1	1,215	8.1
6105	42	*	37	*	43	*	37	*	36	*	148	85.8	254	6.3
6301	4,506	59.2	3,679	17.2	2,667	7.2	3,100	12.0	3,559	6.6	7,743	84.2	17,694	2.2
6312	2,557	66.0	2,307	22.8	1,648	14.6	1,856	17.7	2,208	10.7	4,667	83.2	10,570	1.7
6316	1,391	66.0	1,275	20.4	996	9.8	1,020	13.7	1,204	8.6	2,847	81.6	6,071	5.6
6320	161	65.8	146	24.7	109	19.3	109	11.0	131	6.9	433	93.8	829	1.1
6325	162	75.9	129	19.4	79	*	119	14.3	125	10.4	340	92.9	692	4.2
6330	883	55.6	695	11.2	445	4.5	590	6.6	651	3.8	1,700	82.1	3,625	5.2
6335	665	52.2	509	18.5	372	9.7	436	10.3	493	5.1	1,678	93.6	3,082	0.8
6340	82	*	69	*	54	*	61	*	62	*	240	92.9	442	2.0
6345	20	*	20	*	17	*	16	*	20	*	65	*	116	0.0

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				>=2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
6350	146	47.3	147	18.4	104	8.7	125	13.6	141	8.5	414	90.3	799	1.5
6877	1	*	3	*	1	*	1	*	3	*	8	*	14	*
7304	177	59.9	137	10.2	120	5.0	111	8.1	118	8.5	435	88.5	859	3.0
7306	2,641	63.9	1,892	13.8	1,708	6.6	1,649	9.6	1,710	6.0	5,165	82.5	10,938	5.4
7308	174	71.3	137	18.2	133	9.0	92	*	92	*	378	87.8	807	2.2
7310	36	*	33	*	19	*	20	*	24	*	70	*	145	7.6
7401	2,443	58.9	2,245	14.9	1,947	6.0	1,663	13.3	2,097	6.2	5,118	88.4	10,435	7.2
7601	652	64.6	549	16.0	402	7.7	466	15.5	529	6.6	1,297	86.2	2,736	5.9
7850	1,252	72.2	1,113	22.3	943	12.2	823	11.9	1,009	5.2	2,516	87.4	5,336	4.0
7852	67	*	61	*	49	*	45	*	55	*	196	91.8	376	1.3
8101	1,805	46.5	1,317	8.6	986	4.6	1,276	4.2	1,269	2.8	2,573	69.6	6,374	6.6
8102	550	38.7	403	6.9	355	3.4	383	1.6	396	0.8	1,551	73.3	2,756	2.6
8105	1,597	49.0	1,087	15.4	810	7.0	1,031	9.7	1,096	6.6	2,043	82.1	5,262	2.3
8106	469	47.3	304	8.9	338	3.6	337	6.2	302	3.0	875	66.3	1,909	3.7
8150	60	*	62	*	62	*	57	*	60	*	260	92.7	425	0.9
8151	3,281	79.6	2,720	30.5	2,493	18.2	2,182	20.2	2,563	11.7	7,056	86.5	14,279	2.2
8152	199	77.9	174	25.3	146	7.5	131	16.0	162	7.4	345	79.7	790	3.4

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
8153	19	*	17	*	22	*	10	*	16	*	105	90.5	160	1.3
8155	63	*	54	*	67	*	42	*	51	*	279	93.2	439	0.7
8157	22	*	19	*	13	*	20	*	18	*	97	*	139	2.9
8201	3,095	60.3	2,567	15.9	1,943	8.2	2,097	10.1	2,410	5.4	4,837	85.6	11,487	3.5
8202	3,624	45.0	2,625	21.4	2,236	10.1	2,245	4.9	2,456	4.1	8,671	80.6	16,404	2.9
8203	470	61.9	356	19.1	308	9.4	251	14.7	321	7.2	746	90.3	1,731	4.4
8250	927	49.4	815	10.3	591	6.9	690	6.7	776	4.1	1,730	80.2	3,781	3.8
8252	130	60.8	82	*	78	*	79	*	87	*	305	70.8	588	4.9
8255	3,216	57.9	2,694	14.0	2,075	6.9	2,362	9.7	2,569	4.9	5,159	82.1	12,123	3.5
8257	942	75.8	654	30.7	581	18.2	524	3.4	583	3.1	1,967	78.1	3,915	0.9
8258	519	53.9	334	12.0	229	9.2	300	6.0	329	3.6	804	79.2	1,814	3.7
8260	145	68.3	116	25.9	136	13.2	81	*	104	3.8	547	73.3	998	1.7
8263	744	58.2	608	13.8	470	6.8	508	9.8	595	5.4	993	84.4	2,492	2.6
8264	277	66.8	154	15.6	83	*	145	9.0	149	2.7	285	82.8	722	3.0
8265	83	*	24	*	6	*	31	*	24	*	54	*	152	1.3
8267	14	*	1	*	0	*	1	*	1	*	2	*	16	*
8901	994	43.0	787	5.6	677	1.9	726	2.9	745	0.7	2,527	66.0	4,835	8.7

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%		
8902	1,082	48.5	838	10.0	693	3.5	820	6.6	818	4.0	1,980	70.6	4,461	6.7
8903	1,096	37.9	780	8.6	678	2.1	793	3.2	771	1.8	2,543	69.1	5,029	6.7
8904	1,180	48.1	957	10.1	780	5.1	859	5.5	920	3.5	2,721	71.1	5,499	7.2
8905	2,028	48.8	1,664	11.6	1,368	5.0	1,431	6.2	1,615	4.3	5,154	72.4	9,977	3.4
8908	4	*	0	*	0	*	0	*	0	*	5	*	9	*
9012	35	*	38	*	36	*	28	*	34	*	157	61.1	254	12.2
9013	28	*	24	*	36	*	21	*	25	*	114	60.5	201	8.5
9015	5	*	6	*	7	*	5	*	6	*	29	*	50	*
9016	195	42.1	162	3.7	145	2.1	143	1.4	154	0.0	586	63.8	1,043	8.8
9017	38	*	54	*	38	*	39	*	51	*	175	64.0	300	10.3
9018	116	62.9	131	14.5	104	5.8	104	9.6	124	3.2	465	68.4	791	6.3
9019	8	*	8	*	8	*	7	*	7	*	47	*	76	*
9021	16	*	15	*	13	*	10	*	15	*	52	*	94	*
9041	16	*	14	*	21	*	10	*	14	*	55	*	98	*
9045	3,462	35.4	1,690	6.8	1,194	2.2	1,647	3.7	1,634	1.7	4,386	77.5	9,807	6.1
9050	1,253	60.7	987	14.3	806	4.6	875	4.5	944	2.5	2,769	71.7	5,818	6.3
9051	1,329	59.1	1,170	13.7	946	7.3	915	2.4	1,114	1.5	3,430	72.2	6,674	3.0

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2013 - 2015 Pediatric Nutrition Surveillance (1)
Michigan

Run Date: 05/03/2016

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)
Children Aged < 5 Years (3)
Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4)														
Clinic	Breastfeeding (5)				Exclusive Breastfeeding (6)				≥2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfeed	Number	% Breastfeed At Least 6 Months	Number	% Breatsfeed At Least 12 Months	Number	% Breastfeed At Least 3 Months	Number	% Breatfeed At Least 6 Months	Number	%	Number	%
9053	2,381	49.9	1,883	13.9	1,418	6.1	1,784	7.2	1,878	4.5	4,162	71.0	9,633	16.6
9055	1,104	31.3	174	8.6	0	*	371	5.1	171	2.3	3	*	1,109	4.1
9058	916	59.6	750	21.6	864	13.2	674	5.9	726	3.3	2,797	90.6	5,063	2.2
9066	1,402	39.9	1,075	7.0	890	3.4	965	5.5	1,066	3.0	2,909	74.0	6,199	8.5
9068	1,235	40.2	936	11.2	734	5.0	934	8.1	921	4.3	2,480	69.0	5,278	13.9
9069	2,874	53.0	2,388	10.3	1,830	4.0	2,196	5.0	2,349	2.6	4,954	72.8	11,566	8.8
9070	1,409	43.8	1,130	9.6	937	4.4	1,029	4.2	1,086	2.8	3,070	60.7	6,351	11.6
9072	1,844	41.5	1,387	8.0	1,043	2.5	1,338	3.9	1,394	2.2	3,170	55.3	7,116	14.2
9080	2,156	53.6	1,685	15.0	1,711	6.3	1,449	5.7	1,622	2.4	5,631	72.6	10,659	4.5
9081	165	28.5	92	*	51	*	92	*	88	*	129	78.3	418	4.8
	123	51.2	29	*	50	*	33	*	28	*	95	*	261	33.7
Michigan	173,348	63.1	136,177	18.2	120,019	10.0	113,976	13.4	127,736	7.0	355,550	81.9	737,734	4.7

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.