

Public Health Assessment

Final Release

**Kalamazoo River/Enbridge Spill: Evaluation of Kalamazoo River surface water
and fish after a crude oil release
(Calhoun and Kalamazoo Counties, Michigan)**

**Prepared by the
Michigan Department of Community Health**

JUNE 2, 2014

**Prepared under a Cooperative Agreement with the
U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Agency for Toxic Substances and Disease Registry
Division of Community Health Investigations
Atlanta, Georgia 30333**

THE ATSDR PUBLIC HEALTH ASSESSMENT: A NOTE OF EXPLANATION

This Public Health Assessment was prepared by ATSDR's Cooperative Agreement Partner pursuant to the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA or Superfund) section 104 (i)(6) (42 U.S.C. 9604 (i)(6)), and in accordance with our implementing regulations (42 C.F.R. Part 90). In preparing this document, ATSDR's Cooperative Agreement Partner has collected relevant health data, environmental data, and community health concerns from the Environmental Protection Agency (EPA), state and local health and environmental agencies, the community, and potentially responsible parties, where appropriate.

In addition, this document has previously been provided to EPA and the affected states in an initial release, as required by CERCLA section 104 (i)(6)(H) for their information and review. The revised document was released for a 60-day public comment period. Subsequent to the public comment period, ATSDR's Cooperative Agreement Partner addressed all public comments and revised or appended the document as appropriate. The public health assessment has now been reissued. This concludes the public health assessment process for this site, unless additional information is obtained by ATSDR's Cooperative Agreement Partner which, in the agency's opinion, indicates a need to revise or append the conclusions previously issued.

Use of trade names is for identification only and does not constitute endorsement by the U.S. Department of Health and Human Services. Additional copies of this report are available from:

National Technical Information Service, Springfield, Virginia
(703) 605-6000

You May Contact ATSDR Toll Free at
1-800-CDC-INFO

or

Visit our Home Page at: <http://www.atsdr.cdc.gov>

PUBLIC HEALTH ASSESSMENT

Kalamazoo River/Enbridge Spill: Evaluation of Kalamazoo River surface water
and fish after a crude oil release
(Calhoun and Kalamazoo Counties, Michigan)

Prepared by:

Michigan Department of Community Health
Under Cooperative Agreement with the
U.S. Department of Health and Human Services
Agency for Toxic Substances and Disease Registry

Foreword

The Michigan Department of Community Health (MDCH) conducted this evaluation under a cooperative agreement with federal Agency for Toxic Substances and Disease Registry (ATSDR). ATSDR conducts public health activities (assessments/consultations, advisories, education) at sites of environmental contamination. The purpose of this document is to identify potentially harmful exposures and recommends actions that would minimize those exposures. This is not a regulatory document and does not evaluate or confirm compliance with laws. This is a publicly available document and is provided to the appropriate regulatory agencies for their consideration.

The following steps are necessary to conduct public health assessments/consultations:

- Evaluating exposure: MDCH toxicologists begin by reviewing available information about environmental conditions at the site: how much contamination is present, where it is found on the site, and how people might be exposed to it. This process requires the measurement of chemicals in air, water, soil, or animals. Usually, MDCH does not collect its own environmental sampling data. We rely on information provided by the Michigan Department of Environmental Quality (MDEQ), U.S. Environmental Protection Agency (EPA), and other government agencies, businesses, and the general public.
- Evaluating health effects: If there is evidence that people are being exposed – or could be exposed – to hazardous substances, MDCH toxicologists then determine whether that exposure could be harmful to human health, using existing scientific information. The report focuses on public health – the health impact on the community as a whole.
- Developing recommendations: In its report, MDCH outlines conclusions regarding any potential health threat posed by a site, and offers recommendations for reducing or eliminating human exposure to contaminants. If there is an immediate health threat, MDCH will issue a public health advisory warning people of the danger, and will work with the appropriate agencies to resolve the problem.
- Soliciting community input: The evaluation process is interactive. MDCH solicits and considers information from various government agencies, parties responsible for the site, and the community. If you have any questions or comments about this report, we encourage you to contact us.

Please write to: Toxicology and Response Section
 Division of Environmental Health
 Michigan Department of Community Health
 PO Box 30195
 Lansing, MI 48909

Or call us at: 1-800-648-6942 (toll free)

For more information, please visit:

www.michigan.gov/mdch-toxics

Table of Contents

Summary.....	7
Purpose and Health Issues	7
Background	8
Discussion.....	10
Environmental Contamination	10
Surface water	10
Fish from Marshall Pond, Ceresco Impoundment, and Morrow Lake	17
Exposure Pathways Analysis	21
Toxicological Evaluation	23
Non-oil-related Chemicals	24
1,2,3-Trichloropropane	24
trans-1,4-Dichloro-2-butene.....	24
Analytes with No Screening Levels.....	25
Oil-related Chemicals	25
Non-oil-related Chemicals	26
Community Health Concerns.....	26
Children’s Health Considerations	27
Conclusions.....	27
Recommendations	28
Public Health Action Plan	28
Report Preparation.....	29
References.....	30

List of Tables

Table 1: Chemicals detected in the Michigan Department of Environmental Quality’s (MDEQ) analysis of the crude oil were:	10
Table 2: Levels of analytes (in microgram per liter [µg/L]) with no screening levels in surface water samples collected January 2011 through April 2011.....	13
Table 3: Levels of analytes (in microgram per liter [µg/L]) over or with no screening levels in surface water samples collected May 2011 through August 2011.	14
Table 4: Levels of analytes (in microgram per liter [µg/L]) over or with no screening levels in surface water samples collected September 2011 through December 2011.....	15
Table 5: Levels of analytes (in microgram per liter [µg/L]) with reporting limits over or with no screening levels in surface water samples collected January 2011 through April 17, 2012. No analytes were detected in these samples.	16
Table 6: Fish collected from the locations in the Kalamazoo River in 2010 and 2011.....	17

Table 7: Range of PAHs (in parts per billion [ppb]) in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.	19
Table 8: Minimum, Maximum, and 95% Upper Confidence Limit of the mean (UCL) in parts per million (ppm) of polychlorinated biphenyls (PCBs), in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.	20
Table 9: Minimum, Maximum, and 95% Upper Confidence Limit of the mean (UCL) in parts per million (ppm) of mercury, in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.	20
Table 10: Exposure pathway for residents and visitors recreationally using the areas of the Kalamazoo River and Morrow Lake (Calhoun and Kalamazoo Counties), Michigan, impacted by the July 2010 Enbridge pipeline release of heavy crude oil.	22
Table A-1: Variables used to calculate the site-specific surface water screening levels using the EPA's online screening level calculator for recreational use (http://epa-prgs.ornl.gov/cgi-bin/chemicals/csl_search).	A-2
Table A-2: Generic and site-specific surface water screening levels in either micrograms per liter (µg/L) or milligrams per liter (mg/L).	A-2
Table A-3: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between July to September 2010.	A-9
Table A-4: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between October to December 2010.	A-18
Table A-5: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between January through April 2011.	A-26
Table A-6: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between May to Aug 2011.	A-27
Table A-7: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between September to December 2011.	A-31
Table A-8: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between January to April 17, 2012. (No analytes were detected.)	A-35
Table B-1: Preliminary screening levels (in parts per billion) for polycyclic aromatic hydrocarbons (PAHs), nickel, and vanadium.	B-1
Table B-2: Range (minimum and maximum in parts per billion [ppb]) of polycyclic aromatic hydrocarbons (PAHs), nickel, and vanadium in fish collected in 2010 and 2011 from Marshall Pond.	B-2
Table B-3: Range (minimum and maximum in parts per billion [ppb]) of polycyclic aromatic hydrocarbons (PAHs), nickel, and vanadium in fish collected in 2010 and 2011 from Ceresco Impoundment.	B-4
Table B-4: Range (minimum and maximum in parts per billion [ppb]) of polycyclic aromatic hydrocarbons (PAHs), nickel, and vanadium in fish collected in 2010 and 2011 from Morrow Lake.	B-5
Table B-5: Minimum, Maximum, and 95% Upper Confidence Limit of the mean (UCL) in parts per million (ppm) of polychlorinated biphenyls (PCBs), in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.	B-6

Table B-6: Minimum, Maximum, and 95% Upper Confidence Limit of the mean (UCL) in parts per million (ppm) of mercury, in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.	B-6
--	-----

List of Figures

Figure 1: Overview of the areas of Talmadge Creek and the Kalamazoo River impacted by the July 2010 oil spill (Calhoun and Kalamazoo Counties, Michigan).The lines indicate the area impacted by the July 2010 oil spill and the fish indicate locations where fish were collected (one location upstream of the oil spill: Marshall Pond, and two locations in the area impacted by the oil spill: Ceresco Impoundment and Morrow Lake).	9
--	---

List of Appendices

Appendix A : Levels of chemicals in surface water	A-1
Appendix B : Levels of chemicals in fish filets	B-1

Acronyms and Abbreviations

ATSDR	Agency for Toxic Substances and Disease Registry
BaP	benzo(a)pyrene
BW	Body weight
CR	Consumption rate
CSF	Cancer slope factor
DL	Detection limit
DRO	diesel range organics
EPA	United States Environmental Protection Agency
FDA	United States Food and Drug Administration
GRO	gasoline range organics
kg	kilogram
MDCH	Michigan Department of Community Health
MDEQ	Michigan Department of Environmental Quality
mg/L	milligrams per liter
NA	not available
NLM	United States National Library of Medicine
NRC	National Response Center
NS	no samples
ORO	oil range organics
PAHs	polycyclic aromatic hydrocarbons
PCBs	polychlorinated biphenyls
PHA	Public Health Assessment
ppb	Parts per billion
ppm	Parts per million
RDWC	Residential Drinking Water Criteria
RfD	reference dose
RL	Risk level
UCL	upper confidence limit on the mean
µg/L	micrograms per liter

Summary

Kalamazoo River and Morrow Lake surface water sampling began in July 2010. The sampling was in response to an oil spill from a pipeline operated by Enbridge Energy Partners, LLP (Enbridge). The Michigan Department of Community Health (MDCH) issued an advisory against swimming or fishing in the oil-impacted stretch of the Kalamazoo River and Morrow Lake. The local health departments (in Calhoun and Kalamazoo counties) banned recreational use of the river to prevent people's exposure. This Public Health Assessment (PHA) evaluates chemical levels in surface water and fish. The surface water data were collected between July 2010 and April 2012 from the Kalamazoo River and Morrow Lake. Fish were collected from the Kalamazoo River and Morrow Lake in 2010 and 2011.

MDCH has reached the following conclusions about chemical levels in Kalamazoo River and Morrow Lake surface water and fish:

1. *Chemical levels found in surface water are not expected to cause long-term harm to people's health. People may have temporary health effects, such as skin irritation, from contact with oil sheen or tar globules in the water.* Oil-related and non-oil related chemicals were measured in the surface water. Only a very few of these chemicals were detected above health-protective screening levels in surface water samples. These chemicals were mostly polycyclic aromatic hydrocarbons. People are not expected to be exposed to levels that would cause long-term health concerns. However, if people have contact with these chemicals or the oil sheen and tar globules in the river, they may develop temporary health effects, such as skin irritation.

Next steps: MDCH will continue to review new surface water data.

2. *Oil-related chemical levels found in fish from the Kalamazoo River and Morrow Lake will not harm people's health. However, MDCH has issued fish consumption guidelines because of levels of mercury and polychlorinated biphenyls (PCBs) levels found in the filets.* Fish were collected from Marshall Pond, Ceresco Impoundment, and Morrow Lake. Marshall Pond is upstream of the oil spill. Chemical levels in Marshall Pond fish filets were similar to levels found in Ceresco Impoundment and Morrow Lake fish. Two chemicals not related to the oil spill, mercury and PCBs, were found in the fish filets. MDCH recommends that people limit the amount of certain Kalamazoo River and Morrow Lake fish they eat. (Guidelines listed in this PHA are listed in the 2014-2015 Eat Safe Fish Guide [formerly the Michigan Fish Advisory].) Current guidelines can be found at: www.michigan.gov/eatsafefish.

Next steps: MDCH will evaluate new fish data and issue fish consumption guidelines as needed.

Purpose and Health Issues

The purpose of this Public Health Assessment (PHA) is to determine if people are or may be exposed to chemicals in the Kalamazoo River or Morrow Lake surface water or fish and if any

potential exposure is harmful to people's health. Within days of the oil spill in July 2010, surface water samples were collected from the Kalamazoo River and Morrow Lake. Surface water samples were collected for the rest of 2010, 2011, and 2012 (and continues as of November 2, 2012). Fish from the Kalamazoo River (Marshall Pond and Ceresco Impoundment) and Morrow Lake were collected in 2010 and 2011. Ecological assessment, such as determining effects to fish, is beyond the scope of this PHA. This PHA does not evaluate or confirm regulatory compliance, but determines if any potentially harmful human exposures are occurring or may occur in the future.

This PHA was available for public comment from September 30 to November 14, 2013. MDCH did not receive any comments.

Background

In July 2010, more than 800,000 gallons of crude oil flowed into a wetland area near Talmadge Creek and into the creek, which is a tributary of the Kalamazoo River. The oil was from a 30-inch pipeline near the city of Marshall, Calhoun County, Michigan operated by Enbridge Energy Partners, LLP (Enbridge). Enbridge reported the spill to the National Response Center (NRC), which notified the United States Environmental Protection Agency (EPA), among other agencies.

The EPA was the lead agency for response to this spill and on July 27, the Federal On-Scene Coordinator and Incident Commander issued the EPA Removal Order. Using guidelines of the Incident Command System¹, a Unified Command was established later that week. Members of the Unified Command included federal, state and local agencies, along with Enbridge representatives. At the request of the EPA Incident Commander, MDCH staff deployed to the Command Center to provide public health support.

The spilled oil was eventually contained at Morrow Lake, which was more than 37 miles downstream from the spill (MDEQ 2010a). See Figure 1 for the extent of the oil spill. At the time of the spill, Talmadge Creek and the Kalamazoo River were between 25- and 50-year flood levels due to rain that had fallen during the previous days. Because the river and creek were at high water levels, oil flowed into overbank areas, wetlands, and floodplains.

In July 2010, Calhoun County Public Health and Kalamazoo County Health and Community Services Department issued a ban on recreational river use. MDCH issued a swimming and fishing advisory and a "do not eat" guideline for fish in the river.² In June 2012, a majority of the river was re-opened for recreational use by the county health departments and MDCH lifted the advisory and guideline. MDCH recommended that people follow the pre-spill fish consumption guidelines for the Kalamazoo River and Morrow Lake.

¹ The Incident Command System is a management system for incidents of all sizes and types. This system is used when one agency is responding to an incident and can be scaled up for when many agencies are responding to an incident.

² The MDCH guideline extended from downstream (west) of I-69 on the Kalamazoo River to the west end of Morrow Lake.

Figure 1: Overview of the areas of Talmadge Creek and the Kalamazoo River impacted by the July 2010 oil spill (Calhoun and Kalamazoo Counties, Michigan). The lines indicate the area impacted by the July 2010 oil spill and the fish indicate locations where fish were collected (one location upstream of the oil spill: Marshall Pond, and two locations in the area impacted by the oil spill: Ceresco Impoundment and Morrow Lake).



Discussion

This PHA evaluates two ways people may be exposed to chemicals. One from contact with the Talmadge Creek, Kalamazoo River, and Morrow Lake surface water and the second from eating fish caught from the Kalamazoo River or Morrow Lake. (Talmadge Creek is not fishable.)

Along with the chemicals detected in the Michigan Department of Environmental Quality's (MDEQ) analysis of the oil (MDEQ 2010b), surface water samples were tested for a variety of other chemicals. Some may have been in the oil, but some are naturally occurring or may have been previously present in the surface water. MDCH tested fish filets for chemicals found in the oil, as well as chemicals that were previously found in fish from the Kalamazoo River and Morrow Lake before the oil spill.³ Table 1 contains the chemicals identified in the MDEQ's analysis.

Table 1: Chemicals detected in the Michigan Department of Environmental Quality's (MDEQ) analysis of the crude oil were:

<u>Inorganic chemicals (metals)⁴</u>	<u>Organic chemicals</u>
Beryllium	1,2,3-Trimethylbenzene
Iron	1,2,4-Trimethylbenzene
Mercury	1,3,5-Trimethylbenzene
Molybdenum	2-Methylnaphthalene
Nickel	Benzene
Titanium	Cyclohexane
Vanadium	Ethylbenzene
	Isopropylbenzene
	Naphthalene
	n-Propylbenzene
	Phenanthrene
	p-Isopropyltoluene
	sec-Butylbenzene
	Toluene
	Xylenes, Total

Environmental Contamination

Surface water

Data presented in this PHA were collected from July 2010 to April 2012. These data were obtained from the EPA's Scribe data base. In 2010, MDCH advised people to avoid contact with the oil and the local health departments banned recreational river use. Levels of chemicals

³ The fish were tested for chemicals previously found to ensure that the presence of the oil or any cleanup activities (changes to the river or lake) did not increase the chemical levels in the fish.

⁴ Aluminum, arsenic, barium, cadmium, chromium, cobalt, copper, lead, lithium, manganese, and zinc were not detected in samples of the crude oil (MDEQ 2010b).

measured in surface water samples collected in 2010 are provided in Appendix A (Tables A-2 through A-8). These data are not evaluated in this PHA because the oil floating on Talmadge Creek, the Kalamazoo River, and Morrow Lake was removed in the cleanup conducted in 2010. The 2011 data are evaluated here as it provides information about chemical levels across four seasons of river and lake conditions (temperature, water levels, etc.). Data from 2012 provide information about conditions within a few months prior to the Kalamazoo River and Morrow Lake opening.

Surface water sampling results were compared to health-based screening levels. These screening levels are expected to be protective for everybody, including children. People's actual exposure would most likely be lower than the exposure used in the screening levels. If the levels in the surface water were below the screening level, no further evaluation was necessary. If chemical levels were above the screening level, those chemicals were evaluated further. All screening levels are listed Table A-1 and described in Appendix A.

Samples were collected throughout the impacted area of the Kalamazoo River and Morrow Lake. Surface water was collected from more than 30 locations throughout the impacted area in 2011. From January to April 17, 2012, surface water samples were collected from 27 locations throughout the area impacted by the oil spill. Data was grouped into four month intervals: January to April 2011, May to August 2011, September to December 2011, and January to April 17, 2012. (Data from 2010 is in Appendix A.)

Table 2 presents the data from the surface water samples collected from January 2011 to April 2011. Screening levels are not available for diesel range organics (DRO), gasoline range organics (GRO), oil range organics (ORO), p-isopropyltoluene, and titanium. These chemicals were not detected in most samples collected. As multiple laboratories and methods were used to analyze for these chemicals, reporting limits were presented when the chemical was not detected. These analytes will be discussed in the "Analytes with No Screening Levels" section. No analytes were over available screening levels.

Results for the surface water samples collected between May and August 2011 are presented in Table 3. DRO, GRO, methyl iodide, ORO, p-isopropyltoluene, and titanium have no screening levels and will be discussed in the "Analytes with No Screening Levels" section. Of the remaining analytes listed in Table 3, only benzo(a)pyrene was detected in surface water over the screening level of 2 µg/L. All other chemicals in Table 3 were not detected, but had reporting limits over the screening level (reporting limits were presented as minimum and maximum in the "Not detected" column). Reporting limits are the amount of a chemical that can accurately be detected and its concentration reliably measured. Analytes may be present in the sample below the reporting limit, but will be discussed if the largest reporting limit is over the screening level.

Acenaphthylene was not detected in any of the surface water samples collected between May to August 2011; however, the reporting limit was above the screening level for two out of 503 samples. The average of these samples is 2.6 µg/L.⁵ This average is well below the screening level of 52 µg/L.

⁵ All values (detected and non-detected) were included in the calculation. This value was calculated with EPA's ProUCL 4.1 software.

Indeno(1,2,3-c,d)pyrene was detected in only one of the 503 samples collected between May and August 2011. Only reporting limits from two of the samples were above the screening level. The amount detected, 2.1 µg/L, was below the screening level of 20 µg/L. The average level of indeno(1,2,3-c,d)pyrene was 2.2 µg/L,⁶ which is under the screening level. Acenaphthylene and indeno(1,2,3-c,d)pyrene will not be discussed further. Benzo(a)pyrene, dibenz(a,h)anthracene, 1,2,3-trichloropropane, and trans-1,4-dichloro-2-butene will be discussed below.

Table 4 presents the analytes over or with no screening levels in surface water samples collected between September and December 2011. DRO, GRO, methyl iodide, ORO, and p-isopropyltoluene will be discussed in the “Analytes with No Screening Levels” section. Of the remaining analytes, only benzo(a)pyrene was detected in surface water samples. All other analytes had reporting limits that were over the screening levels.

For the 631 surface water samples collected between May and December 2011 (Tables 3 and 4), benzo(a)pyrene was only detected in three samples. Only two of the detected samples were over the screening level of 2 µg/L. The average level of benzo(a)pyrene in the surface water was about 2 µg/L.⁷ Benzo(a)pyrene will be discussed further in the “Exposure Pathways” section.

For the 631 surface water samples collected between May and December 2011 (Tables 3 and 4), dibenz(a,h)anthracene was not detected in any of the samples. For the samples where dibenz(a,h)anthracene was not detected, 131 samples had reporting limits over the screening level of 2 µg/L. The average level for all the samples tested for dibenz(a,h)anthracene was 2.2 µg/L.⁸ This analyte will be discussed further in the “Exposure Pathways” section.

No analytes were detected in surface water samples collected from January to April 17, 2012 (Table 5). Methyl iodide and p-isopropyl toluene will be discussed in the “Analytes with No Screening Levels” section. The reporting limits were over the screening levels for two analytes, 1,2,3-trichloropropane and trans-1,4-dichloro-2-butene. These analytes will be discussed in the “Exposure Pathways” section.

⁶ All values (detected and non-detected) were included in the calculation. This value was calculated with EPA’s ProUCL 4.1 software. The EPA’s ProUCL software can be found at: <http://www.epa.gov/osp/hstl/tsc/software.htm#Download>.

⁷ If all values (detected and non-detected) were included in the calculation, the average was 1.9 µg/L. If the reporting limits were identified as non-detects, the average was 2.2 µg/L. These values were calculated with EPA’s ProUCL 4.1 software.

⁸ The average was calculated, using the EPA’s ProUCL 4.1 software, using the reporting limits because dibenz(a,h)anthracene was not detected in any of the samples.

Table 2: Levels of analytes (in microgram per liter [$\mu\text{g/L}$]) with no screening levels in surface water samples collected January 2011 through April 2011.

Analyte	Total number of samples	Detected			Not detected ^a		
		Number of samples	Minimum level	Maximum level	Number of samples	Minimum reporting limit	Maximum reporting level
Diesel Range Organics (DRO)	337	0	ND ^b	ND	337	100	100
Gasoline Range Organics (GRO)	334	0	ND	ND	334	200	200
Oil Range Organics (ORO)	70	67	48	220	3	100	100
p-Isopropyltoluene	337	0	ND	ND	337	1	2
Titanium	656	325	0.55	14	331	1.0	1.0

a = These chemicals were not detected. The minimum and maximum levels were the reporting limits for the chemicals.

b = These chemicals were not detected in the samples (ND = not detected).

Table 3: Levels of analytes (in microgram per liter [$\mu\text{g/L}$]) over or with no screening levels in surface water samples collected May 2011 through August 2011.

Analyte	Screening level	Total number of samples	Detected				Not detected ^a			
			Number of samples	Minimum level	Maximum level	Samples over screening level	Number of samples	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,2,3-Trichloropropane	0.07	9	0	ND ^b	ND	NA	9	1	1	9
Acenaphthylene	52	503	0	ND	ND	NA	503	1	100	2
Benzo(a)pyrene	2	503	2	2.1	3.4	2	501	1	20	259
Dibenz(a,h)anthracene	2	503	0	NA	NA	NA	503	1	40	130
Diesel Range Organics (DRO)	NA	537	3	7,300	40,000	NA	534	100	1,000	NA
Gasoline Range Organics (GRO)	NA	537	0	ND	ND	NA	537	200	200	NA
Indeno(1,2,3-c,d)pyrene	20	503	1	2.1	2.1	0	502	1	40	2
Methyl iodide	NA	9	0	ND	ND	NA	9	1	1	NA
Oil Range Organics (ORO)	NA	537	7	440	170,000	NA	530	100	100	NA
p-Isopropyltoluene	NA	537	1	0.41	0.41	NA	536	2	5	NA
Titanium	NA	526	526	1.2	49	NA	0	NA	NA	NA
trans-1,4-Dichloro-2-butene	0.0012	9	0	ND	ND	NA	9	5	5	9

Bold values are above the screening levels.

a = These chemicals were not detected. The minimum and maximum levels were the reporting limits for the chemicals.

b = "NA" indicates that no screening levels were available or no values were available.

Table 4: Levels of analytes (in microgram per liter [$\mu\text{g/L}$]) over or with no screening levels in surface water samples collected September 2011 through December 2011.

Analyte	Screening level	Total number of samples	Detected				Not detected ^a			
			Number of samples	Minimum level	Maximum level	Samples over screening level	Number of samples	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,2,3-Trichloropropane	0.07	128	0	ND ^b	ND	NA	128	1	1	128
Benzo(a)pyrene	2	128	1	1	1	0	127	1	5	1
Dibenz(a,h)anthracene	2	128	0	ND	ND	NA	128	2	5	1
Diesel Range Organics (DRO)	NA	38	1	140	140	NA	37	100	100	NA
Gasoline Range Organics (GRO)	NA	8	0	ND	ND	NA	8	200	200	NA
Methyl iodide	NA	128	0	ND	ND	NA	128	1	1	NA
Oil Range Organics (ORO)	NA	38	5	260	710	NA	33	100	100	NA
p-Isopropyltoluene	NA	128	1	9.5	9.5	NA	127	5	5	NA
trans-1,4-Dichloro-2-butene	0.0012	128	0	ND	ND	NA	128	1	1	128

Bold values are above the screening levels.

a = These chemicals were not detected. The minimum and maximum levels were the reporting limits for the chemicals.

b = “NA” indicates that no screening levels were available or no values were available.

Table 5: Levels of analytes (in microgram per liter [$\mu\text{g/L}$]) with reporting limits over or with no screening levels in surface water samples collected January 2011 through April 17, 2012. No analytes were detected in these samples.

Analyte	Screening level	Not detected ^a			
		Number of samples	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,2,3-Trichloropropane	0.07	24	1	1	24
Methyl iodide	NA ^b	24	1	1	NA
p-Isopropyltoluene	NA	24	5	5	NA
trans-1,4-Dichloro-2-butene	0.0012	24	1	1	24

Bold values are above the screening levels.

a = These chemicals were not detected. The minimum and maximum levels were the reporting limits for the chemicals.

b = "NA" indicates that no screening levels were available or no values were available.

Fish from Marshall Pond, Ceresco Impoundment, and Morrow Lake

Fish were collected from three locations in 2010 and 2011. One location, Marshall Pond, is upstream of the spill and provides background levels of oil-related chemicals. Two locations, Ceresco Impoundment and Morrow Lake, were within the stretch of the Kalamazoo River impacted by the crude oil spill. Table 6 presents the fish species collected in 2010 and 2011 from these three locations. Fish filets were analyzed for nickel, vanadium, 21 individual polycyclic aromatic hydrocarbons (PAHs), mercury, and 83 polychlorinated biphenyl (PCB) congeners.⁹ Nickel, vanadium, and a few of the PAHs were found in samples of the oil that spilled. Mercury and PCBs were measured to ensure that the oil spill, or the resulting clean-up activities, did not cause changes to chemical levels in the fish.

Table 6: Fish collected from the locations in the Kalamazoo River in 2010 and 2011.

Location	Collected in 2010 (number of fish)	Collected in 2011 (number of fish)
Marshall Pond	Carp (10) and Smallmouth Bass (10)	Carp (10), Largemouth Bass (10), Rock Bass (10), and Smallmouth Bass (10)
Ceresco Impoundment	Carp (10) and Rock Bass (10)	Carp (10), Rock Bass (5), Smallmouth Bass (10)
Morrow Lake	Bluegill (6 ^a), Carp (10), and Rock Bass (8)	Bluegill (3 ^b) and Carp (10)

a = Ten bluegill were collected in 2010 from Morrow Lake. The 6 bluegill filet samples include 3 composite samples (one composed of 3 fish filets, one composed of 2 fish filets, and one composed of 2 fish filets) and filets from three individual fish.

b = Ten bluegill were collected in 2011 from Morrow Lake. The 3 samples include 2 composite samples (one composed of 6 fish filets and one composed of 3 fish filets) and one individual filet.

MDCH has established fish consumption screening levels for mercury and PCBs. As no fish consumption screening levels were available for PAHs, nickel, and vanadium, MDCH calculated preliminary fish consumption screening levels using the assumption that a person would eat 16 *MI Servings* per month, which is the least restrictive consumption guideline category.¹⁰ The screening level for benzo(a)pyrene and the carcinogenic PAHs grouped with benzo(a)pyrene, called BaP equivalent values,¹¹ was 0.9 ppb. For the other PAHs, nickel, and vanadium the screening level was 2,700 ppb.

Nickel and vanadium were not detected in any of the filets from fish collected in either 2010 or 2011 from Marshall Pond, Ceresco Impoundment, or Morrow Lake. The detection limit for nickel was 600 parts per billion (ppb). The detection limit for vanadium was 200 ppb. As there was no difference in the nickel and vanadium levels in fish collected from the three locations and

⁹ Some other organochlorine chemicals are measured when PCBs are measured, but no fish consumption guidelines are issued due to these chemicals.

¹⁰ MDCH's standard exposure assumptions and equations for calculating fish consumption guidelines are available at: www.michigan.gov/eatsafefish under "Reports and Science." The number of ounces in a *MI Serving* changes with body weight. For example, a *MI Serving* is 8 ounces for an 80 kilogram (kg) person, 4 ounces for a 40 kg person, and 2 ounces for a 20 kg person.

¹¹ Along with benzo(a)pyrene, six other PAHs (benzo(b)fluoranthene, benzo(a)anthracene, benzo(k)fluoranthene, chrysene, dibenz(a,h)anthracene, and indeno(1,2,3-cd)pyrene) are included in the benzo(a)pyrene (BaP) equivalent values.

the levels were not above the screening level, no fish consumption guidelines are necessary for nickel and vanadium. Appendix B contains additional discussion.

Fish filets were tested for 21 PAHs. Seven of the PAHs can be evaluated as a group by relating the amount present to an equivalent amount of benzo(a)pyrene (BaP equivalent value). These seven are benzo(a)pyrene, benzo(b)fluoranthene, benzo(a)anthracene, benzo(k)fluoranthene, chrysene, dibenz(a,h)anthracene, and indeno(1,2,3-cd)pyrene. See Appendix B for further details. BaP equivalent values are presented in Table 7. (See Table B-1, -2, and -3 in Appendix B for individual chemical levels.) For most of the samples, the PAHs included in the BaP equivalent value were not detected. These PAHs were only detected in carp and rock bass from Marshall Pond; carp from Ceresco Impoundment; and bluegill from Morrow Lake. The maximum BaP equivalent value was 1.26 ppb, from rock bass collected in Marshall Pond (a location upstream of the spill). While the maximum level is over the screening level of 0.8 ppb, an upper-bound estimate of the average concentration, called the 95% upper confidence limit of the mean (UCL), for all the rock bass collected from Marshall Pond (0.7 ppb) is not. No fish consumption guidelines are necessary.

The other fourteen PAHs were evaluated individually (1-methylnaphthalene, 1-methylphenanthrene, 2,6-dimethylnaphthalene, 2-methylnaphthalene, acenaphthene, acenaphthylene, anthracene, benzo(e)pyrene, benzo(g,h,i)perylene, fluoranthene, fluorene, naphthalene, phenanthrene, and pyrene). The highest levels of these PAHs were found in carp collected from Ceresco Impoundment in 2010 (naphthalene at 16.1 ppb) and Marshall Pond in 2010 (acenaphthene at 14.91 ppb). These levels are well below the screening level of 2,300 ppb.

No fish consumption guidelines are necessary due to the PAHs levels. Appendix B contains additional discussion.

Table 7: Range of PAHs (in parts per billion [ppb]) in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.

Fish	Range (minimum and maximum) of PAHs (in ppb)					
	Marshall Pond		Ceresco Impoundment		Morrow Lake	
	BaP equivalent value ^a	Other PAHs ^b	BaP equivalent value ^a	Other PAHs ^b	BaP equivalent value ^a	Other PAHs ^b
Bluegill	NS ^c	NS	NS	NS	0.24-0.81 ^e	0.2 (DL) ^d - 3.23
Carp	0.46 ^e	0.2 (DL) - 14.91	0.46-0.67 ^e	0.2 (DL) - 16.1	0.46	0.2 (DL) - 6.58
Largemouth Bass	0.46	0.2 (DL) - 2.92	NS	NS	NS	NS
Smallmouth Bass	0.46	0.2 (DL) - 4.67	0.46	0.2 (DL) - 6.96	NS	NS
Rock Bass	0.2-1.26 ^e	0.2 (DL) - 8.99	0.46	0.2 (DL) - 7.5	0.46	0.2 (DL) - 3.37

a = The individual PAHs included in the BaP equivalent values are benzo(a)pyrene, benzo(b)fluoranthene, benzo(a)anthracene, benzo(k)fluoranthene, chrysene, dibenz(a,h)anthracene, and indeno(1,2,3-cd)pyrene. See Appendix B for a description of the BaP equivalent values. Unless otherwise noted, these PAHs were not detected.

b = These PAHs are 1-methylnaphthalene, 1-methylphenanthrene, 2,6-dimethylnaphthalene, 2-methylnaphthalene, acenaphthene, acenaphthylene, anthracene, benzo(e)pyrene, benzo(g,h,i)perylene, fluoranthene, fluorene, naphthalene, phenanthrene, and pyrene.

c = "NS" indicates that the fish were not sampled.

d = The value was at the detection limit (DL).

e = At least one of the PAHs was detected.

Tables 8 and 9 present the PCB and mercury levels, respectively, in fish collected from Marshall Pond, Ceresco Impoundment, and Morrow Lake. Based on the levels (95% UCLs) of PCBs and mercury measured in the fish filets, fish consumption guidelines are needed for certain fish species. The fish consumption guidelines are located in the discussion following Table 9.

Table 8: Minimum, Maximum, and 95% Upper Confidence Limit of the mean (UCL) in parts per million (ppm) of polychlorinated biphenyls (PCBs), in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.

Fish	Levels of PCBs (in ppm)								
	Marshall Pond			Ceresco Impoundment			Morrow Lake		
	2010	2011	95% UCL ^a	2010	2011	95% UCL ^a	2010	2011	95% UCL ^a
Bluegill	NS ^b	NS	NA ^c	NS	NS	NA	0.0054-0.0162	0.0019-0.0074	0.01
Carp	0.001-0.0908	0.0031-0.0826	0.04	0.0139-0.2448	0.0035-0.066	0.12	0.0519-0.3041	0.0697-0.8292	0.55
Largemouth Bass	NS	0.001-0.0126	0.01	NS	NS	NA	NS	NS	NA
Smallmouth Bass	0.0016-0.1646	0.001-0.24	0.04	NS	0.001-0.0172	0.04 ^d	NS	NS	0.24 ^d
Rock Bass	NS	0.001-0.0174	0.01	0.0012-0.0252	0.0019-0.0041	0.01	0.0021-0.237	NS	0.24 ^d

a = The 95% upper confidence limit on the mean (UCL) calculation includes all available data sets.

b = “NS” (no samples) indicates that the fish were not collected.

c = A 95% UCL was not available (NA) because no fish were collected.

d = The 95% UCL includes data from fish collected in 2006 or may have been calculated only with data from fish collected in 2006.

Table 9: Minimum, Maximum, and 95% Upper Confidence Limit of the mean (UCL) in parts per million (ppm) of mercury, in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.

Fish	Levels of mercury (in ppm)								
	Marshall Pond			Ceresco Impoundment			Morrow Lake		
	2010	2011	95% UCL ^a	2010	2011	95% UCL ^a	2010	2011	95% UCL ^a
Bluegill	NS ^b	NS	NA ^c	NS	NS	NA	0.025-0.055	0.022-0.063	0.05
Carp	0.041-0.23	0.067-0.28	0.15	0.074-0.21	0.061-0.16	0.13	0.059-0.65	0.082-0.37	0.27
Largemouth Bass	NS	0.12-0.45	0.27	NS	NS	NA	NS	NS	NA
Smallmouth Bass	0.064-0.14	0.068-0.24	0.16	NS	0.064-0.38	1.0 ^d	NS	NS	0.59 ^d
Rock Bass	NS	0.054-0.13	0.1	0.052-0.085	0.034-0.061	0.07	0.075-0.14	NS	0.59 ^d

a = The 95% upper confidence limit on the mean (UCL) calculation includes all available data sets.

b = “NS” (no samples) indicates that the fish were not collected.

c = A 95% UCL was not available (NA) because no fish were collected.

d = The 95% UCL includes data from fish collected in 2006 or may have been calculated only with data from fish collected in 2006.

MDCH issued fish consumption guidelines based on PCB and mercury levels for Kalamazoo River and Morrow Lake fish. The levels of PCBs and mercury were similar to those seen before the oil spill. Based on the levels of nickel, vanadium, and PAHs, no fish consumption guidelines are necessary for these chemicals.

- For the Kalamazoo River, in Marshall Pond, people should eat no more than 8 *MI Servings* per month (*MI Servings*/month) of rock bass; 4 *MI Servings*/month of smallmouth bass, largemouth bass less than 18 inches, or carp; and 2 *MI Servings*/month of largemouth bass greater than 18 inches. A *MI Serving* is 8 ounces for an adult (80 kilograms [kgs]) and 2-4 ounces for children (20-40 kgs). See the Eat Safe Fish Guide for more information (www.michigan.gov/eatsafefish).
- For the Kalamazoo River, from the Marshall Pond Dam to Ceresco Impoundment, people should eat no more than 8 *MI Servings*/month of rock bass greater than 8 inches; 4 *MI Servings*/month of smallmouth bass less than 18 inches; 2 *MI Servings*/month of smallmouth bass greater than 18 inches; and 1 meal/month of carp.
- Kalamazoo River from the Ceresco Dam to Morrow Dam, including Morrow Lake, people should eat no more than 16 *MI Servings*/month of bluegill; 8 *MI Servings*/month of rock bass; 6 *MI Servings*/year of smallmouth bass; and healthy adults should limit their consumption (to 1-2 *MI Servings*/year) of carp. See the Eat Safe Fish Guide for more information.

The above guidelines are from the 2014-2015 Eat Safe Fish Guide (formerly the Michigan Fish Advisory). Check the Eat Safe Fish Guide for the most current fish consumption guidelines (www.michigan.gov/eatsafefish).

Exposure Pathways Analysis

There are five things to consider when deciding if a person may be exposed to a chemical, also known as an *exposure pathway*: (1) where is the chemical coming from (source), (2) what in a person's environment has been contaminated (environmental medium), (3) is there a way a person might come into contact with the chemical (exposure point), (4) how they might come into contact with the chemical (exposure route), and (5) who might be exposed to it (exposed population). An exposure pathway is complete if it is expected or there is proof that all five elements are present. It is considered either a potential or an incomplete pathway if there is a lower probability of exposure or there is no evidence that at least one of the elements above are, have been, or will be present. Table 10 summarizes the exposure pathway for residents and visitors recreationally using the areas of the Kalamazoo River and Morrow Lake (Calhoun and Kalamazoo Counties), Michigan, impacted by the July 2010 Enbridge pipeline release of crude oil.

Table 10: Exposure pathway for residents and visitors recreationally using the areas of the Kalamazoo River and Morrow Lake (Calhoun and Kalamazoo Counties), Michigan, impacted by the July 2010 Enbridge pipeline release of heavy crude oil.

Source	Environmental Medium	Exposure Point	Exposure Route	Exposed Population	Time Frame	Exposure
Heavy crude oil (crude oil-related chemicals)	Surface water	Surface water	Incidental ingestion and dermal contact	Residents and visitors using the river and lake	Past Present Future	Incomplete ^a Completed ^b Potential ^c
Heavy crude oil (crude oil-related chemicals)	Surface water and sediment	Fish	Ingestion	Residents and visitors who eat fish	Past Present Future	Incomplete Completed Potential
Other sources or contaminated areas (non-oil-related chemicals)	Surface water	Surface water	Incidental ingestion and dermal contact	Residents and visitors using the river and lake	Past Present Future	Incomplete Completed Potential
Other sources or contaminated areas (non-oil-related chemicals)	Surface water and sediment	Fish	Ingestion	Residents and visitors who eat fish	Past Present Future	Incomplete Completed Potential

a = Incomplete indicates that all five elements of the exposure were or are not present.

b = Completed indicates that all five elements of the exposure pathway are either expected to occur or occurring.

c = Potential indicates that all five elements of the exposure pathway may have occurred in the past or may occur in the future.

People using the Kalamazoo River or Morrow Lake for recreational activities, such as canoeing, fishing, or swimming, may encounter oil sheen on the water or vegetation. Although this oil sheen may be present in a few millimeters thick layer on top of the water, very few chemicals were found in the surface water in samples collected in 2011 and 2012.

Only one analyte, benzo(a)pyrene, was detected above the surface water screening levels from samples collected between January 2011 and April 17, 2012. Benzo(a)pyrene was detected in two out of 503 surface water samples collected between May and August 2011. The average benzo(a)pyrene value is approximately equivalent to the screening level of 2 µg/L. While people may occasionally encounter benzo(a)pyrene in the water, the levels present are not expected to be harmful to people's health.

Dibenz(a,h)anthracene was not detected in any of the 631 samples collected from May to December 2011. Although 131 samples had reporting limits over the screening level of 2 µg/L, the average (2.2 µg/L) was just slightly over the screening level. Dibenz(a,h)anthracene may sometimes be present in the surface water in amounts over the screening level. People are not expected to encounter amounts that are consistently over the screening level. The amounts that could be present (those above the screening level, but below the reporting limit) are not expected to be harmful to people's health.

Higher levels of benzo(a)pyrene or dibenz(a,h)anthracene may be present in areas with oil sheen (both sheen from remaining oil from the July 2010 oil spill and sheen from other sources). In general, people should minimize their contact with oil sheen as skin contact can cause temporary skin irritation. Avoid areas with sheen, if possible. If people get sheen or tar globules on their skin, wash with soap and water as soon as possible.

Some analytes had reporting limits consistently above the surface water screening levels. These were 1,2,3-trichloropropane and trans-1,4-dichloro-2-butene. For 1,2,3-trichloropropane, all 161 surface water samples collected between May 2011 and April 17, 2012 had reporting limits over the screening level (0.07 µg/L). The reporting limit for all samples was 1 µg/L. Since this chemical is not expected to be present in the water, one-half of the reporting limit can be used to compare to the screening level. Since one-half of the reporting limit, 0.5 µg/L, is still above the screening level, 1,2,3-trichloropropane will be discussed in the “Toxicological Evaluation” section.

All 161 surface water samples, collected between May 2011 and April 17, 2012, tested for trans-1,4-dichloro-2-butene were above the screening level (0.0012 µg/L). The highest reporting limit was 5 µg/L. As half of that limit, 2.5 µg/L is still above the screening level, trans-1,4-dichloro-2-butene will be discussed in the “Toxicological Evaluation” section.

Some analytes did not have any screening levels (DRO, GRO, ORO, p-isopropyltoluene, titanium, and methyl iodide). These will be discussed in the “Analytes with No Screening Levels” section.

People fishing in the Kalamazoo River, including Marshall Pond and Ceresco Impoundment, or Morrow Lake should limit their consumption of certain fish because of mercury and PCB levels. Nickel and vanadium were not detected in any of the fish filets and only low levels of the PAHs measured were found. The levels of PAHs present in the fish filets are not a health concern and do not result in any fish consumption guidelines.

Toxicological Evaluation

No chemicals identified in the oil were present in the surface water or fish filet samples at levels that would be expected to cause more than temporary skin irritation with contact.¹²

The mercury and PCB levels present in some fish species were high enough for MDCH to issue fish consumption guidelines. Following the fish consumption guidelines will limit people’s exposure to these chemicals so that no health effects will be expected for any individuals, including children. The fish consumption guidelines can be found after Table 9 (on page 21). Check www.michigan.gov/eatsafefish for the most current guidelines.

¹² Titanium and p-isopropyltoluene were detected in surface water samples, but there are no screening levels for these two analytes. They will be discussed in the “Analytes with No Screening Levels” section.

Non-oil-related Chemicals

1,2,3-Trichloropropane

One hundred and sixty-one surface water samples were tested for 1,2,3-trichloropropane between May 2011 and April 17, 2012. This chemical was not detected in any of the samples, however, the reporting limit (1 µg/L, the lowest level that could accurately be measured) for all samples was over the screening level of 0.07 µg/L.

Regular swimmers would be expected to ingest about 0.38 µg¹³ of 1,2,3-trichloropropane over the course of a year. This amount is lower than the amount that people may be exposed to through their diet. FDA food survey values indicate that levels between 3.53 and 6.12 part per billion (or µg/kg), with an average of 4.21ppb, were found in nine food items tested, with the highest levels in sandwich cookies (Heikes et al. 1995). The total amount a person may ingest over 150 days swimming is about equal to 2 servings of sandwich cookies.¹⁴ People are expected to be exposed to this chemical through foods, drinking water, and outside air (NLM 2009c). The levels that could be present in the water are not expected to add much to people's typical exposure. People's health would not be harmed from this exposure.¹⁵

trans-1,4-Dichloro-2-butene

This chemical, trans-1,4-dichloro-2-butene, was measured in 161 surface water samples collected between May 2011 and April 17, 2012. It was not detected in any of the samples. The reporting limit (the lowest levels that could accurately be measured) was 1 µg/L for 152 of the samples and 5 µg/L for nine of the samples. Both of these reporting limits were over the screening level of 0.0012 µg/L.¹⁶ This screening level is based on inhalation of the chemical. If this chemical is even present in the surface water, it is expected to volatilize and be degraded in the air within about a week. It will also degrade in the water within a few days (NLM 2009b). Because it was not detected, it degrades fairly quickly in the air and water, and no locations along the river seem to be contaminated with this chemical,¹⁷ people are expected to be exposed to very little or no trans-1,4-dichloro-2-butene in surface water. The only exposure expected is to the background levels that may be present here and in other similar locations. This exposure is not expected to harm people's health.

¹³ As this chemical is not expected to be present, one-half of the reporting limit (0.5 µg/L) was used to calculate this value. Regular swimmers would be expected to accidentally swallow about 0.75 L of water over 150 days. This would result in swallowing a maximum of 0.38 µg of 1,2,3-trichloropropane over those 150 days. The amount that would be dermally absorbed would add only an insignificant amount of 1,2,3-trichloropropane to the amount ingested (less than 0.01 µg).

¹⁴ If a serving of sandwich cookies is 34 grams (1.2 ounces), the amount of 1,2,3-trichloropropane is 0.204 µg/serving (6.12 µg/30 servings in a kilogram [kg]). Two servings is approximately 0.4 µg of 1,2,3-trichloropropane.

¹⁵ Adults and children would ingest less (0.000032-0.00025 µg/kg-day) than the reference dose of 4 µg/kg-day from accidentally swallowing river or lake water while swimming. If people did swallow this amount, an estimated cancer risk would be less than 1 in 100,000, which means out of 100,000 people exposed less than one additional person could develop cancer.

¹⁶ As trans-1,4-dichloro-2-butene is not expected to be present in the surface water, one-half of the reporting limit would be 0.5 or 2.5 µg/L. Both are above the screening level of 0.0012 µg/L.

¹⁷ This information is from the list of chemicals found in nearby Part 201 sites (Enbridge 2010).

Analytes with No Screening Levels

Oil-related Chemicals

DRO (C10-C20)

DRO (diesel range organics) are chemicals with 10 to 20 carbons in the structure. This is a generic designation for many specific chemicals present in petroleum products and may measure chemicals that are from plant or bacterial sources. Of the 912 samples collected in 2011, DRO was detected in four samples (less than 0.5% of the total samples). The maximum level detected was 40,000 µg/L. DRO data are most useful in identifying whether petroleum products are present in the samples, not to assess any potential health concerns. The results from the surface water samples indicate that chemicals that can be measured with this analysis were very rarely detected in surface water in 2011.

GRO (C6-C10)

GRO (gasoline range organics) are chemicals with six to ten carbons in the structure. Similar to DRO, this is a generic designation. GRO was not detected in any of the 879 surface water samples analyzed and the reporting limit was 200 µg/L. GRO data are most useful in identifying whether petroleum products are present in the samples, not to assess any potential health concerns. The results from the surface water samples do not indicate that GRO were present in surface water, above 200 µg/L, in 2011.

ORO (C20-C34)

ORO (oil range organics) are chemicals with 20 to 34 carbons in the structure. As with DRO and GRO, any hydrocarbon with the correct number of carbons will be measured, which include chemicals from plant or bacterial sources. OROs were detected in 79 out of the 748 surface water samples analyzed (about 10% of the total samples). The detections ranged from 48 to 170,000 µg/L. ORO data are most useful in identifying whether petroleum products are present in the samples, not to assess any potential health concerns. The results from the surface water samples indicate that petroleum products, or chemicals from plant or bacterial sources, may have been present in surface water in 2011.

p-Isopropyltoluene

Over 1,000 surface water samples (1,026) were analyzed for p-isopropyltoluene, also called p-cymene, from January 2011 to April 17, 2012. This chemical was only detected in two, at 0.41 and 9.5 µg/L, of the samples.¹⁸ The U.S. Food and Drug Administration (FDA) allows this chemical to be added to foods as a flavoring (NLM 2010a). Exposure to the levels present in surface water is not a health concern.

Titanium

People ingest titanium in food, such as milled grains, butter, corn oil, and lettuce (IPCS 1982). There is not enough information available on titanium to determine levels that could cause harm to people's health. However, titanium is used in a variety of medical devices, such as knee or hip joint replacements and dental implants and there is no evidence of toxicity from these implants

¹⁸ These two samples represent 0.2% of the total samples tested from January 2011 to April 17, 2012.

(IPCS 1982). One form of titanium, titanium dioxide, is FDA-approved for use as a color additive in food (NLM 2009a). Titanium was tested in 1,182 surface water samples collected between January and August 2011. The range of titanium detected in the samples was 0.55 to 49 µg/L. People's bodies will not absorb a majority of the titanium in the water. The highest amount that regular swimmers may accidentally drink totals 40 µg of titanium.¹⁹ This total (for the entire year) would only be about 10% of a person's daily titanium intake from the average amount in the typical U.S. diet (NLM 2002). People's exposure to titanium from accidental swallows of water is not expected to add much to the typical daily exposure and is not expected to harm people's health.

Non-oil-related Chemicals

Methyl iodide

One hundred and sixty-one surface water samples were tested for methyl iodide between May 2011 and April 17, 2012. This chemical was not detected in any of the samples. The reporting limit (the lowest level that could accurately be measured) was 1 µg/L.

People are exposed to small amounts of methyl iodide in the air and in ocean seafood (NLM 2010b). Based on the reporting limits for the surface water samples, the maximum amount of methyl iodide that a person would accidentally swallow over the course of 150 days is 0.38 µg.²⁰ This is about ten times lower than the amount that a person could have from one meal of ocean fish.²¹ The amount of methyl iodide that a person could accidentally swallow would not harm people's health.

Community Health Concerns

Individuals have concerns about watering their gardens and/or lawns with water from the river. These concerns are, in part, due to on-going oil recovery activities, such as surface boom on the river or submerged oil removal. People are primarily concerned about chemicals dissolved in the water, but are also concerned about sheen or tar globules being sprayed onto the edible and non-edible parts of plants (Steve Noble, MDEQ, personal communication 2012). People's health will not be harmed from direct contact with chemicals dissolved in the water. Any sheen or tar globules in the water may cause temporary skin irritation while people are irrigating.

The chemicals present, or possibly present, in the water may accumulate in the plants. However, this accumulation is smaller than the amount of the chemicals that would be expected to remain on the surface of the plant from splashed soil/water droplets (Samsoe-Petersen et al. 2002, Fismes et al. 2002). The potential oil sheen or tar globules remaining from being sprayed on gardens could irritate people's skin if they are tending the garden without gloves. Produce with visible oil sheen or tar globules should not be eaten. Individuals may want to consider alternate

¹⁹ This is if people swim for 0.1 hours daily for 150 days (15 hours) and accidentally swallow 0.05 L of water per hour of swimming (0.75 L of water). If the water always had 49 µg titanium/L, people would swallow 0.040 µg of titanium over the 150 days.

²⁰ Since the chemical was never detected and is not expected to be present, one-half of the reporting limit was used (0.5 µg/L). A person swimming each day for 150 days would accidentally swallow 0.75 L of river or lake water. People would swallow approximately 0.38 µg of methyl iodide over those 150 days (0.5 µg/L x 0.75 L = 0.38 µg).

²¹ The median level of methyl iodide in five species of fish was 17 µg/kg (NLM 2010b). From one 8 ounce meal (0.227 kg), people would ingest 3.9 µg of methyl iodide (17 µg/kg x 0.227 kg = 3.9 µg).

watering methods, such as drip irrigation (root vegetables may still have this material on the surface) or an alternate water source.

Children's Health Considerations

Children do more hand-to-mouth behaviors than adults, which may increase the exposure to any chemicals present. Children may be at greater risk than adults when exposed to certain hazardous substances. A child's lower body weight and higher intake rate result in a greater dose of hazardous chemicals compared to their weight. If toxic exposure levels are high enough during critical growth stages, the developing body systems of children may be damaged. Screening levels used in this PHA are protective for children as well as adults.

Oil sheen or tar globules may be attractive for children to touch. This could result in temporary skin irritation. If children get oil sheen or tar globules on their skin, it should be washed off as soon as possible with soap and water. Accidentally swallowing sheen or a tar globule may cause gastrointestinal upset.

Conclusions

Chemical levels found in surface water are not expected to cause long-term harm to people's health. People may have temporary health effects, such as skin irritation, from contact with oil sheen or tar globules in the water. Oil-related and non-oil related chemicals were measured in the surface water. Only a very few of these chemicals were detected above health-protective screening levels in surface water samples. These chemicals were mostly polycyclic aromatic hydrocarbons. People are not expected to be exposed to chemical levels that would cause long-term health concerns. However, if people have contact with these chemicals or the oil sheen and globules in the river, they may develop temporary health effects, such as skin irritation.

Oil-related chemical levels found in fish from the Kalamazoo River and Morrow Lake will not harm people's health. However, MDCH has issued fish consumption guidelines because of mercury and polychlorinated biphenyls (PCBs) levels found in the filets. Fish were collected from Marshall Pond, Ceresco Impoundment, and Morrow Lake. Marshall Pond is upstream of the oil spill. Chemical levels in Marshall Pond fish filets were similar to levels found in Ceresco Impoundment and Morrow Lake fish. Two chemicals not related to the oil spill, mercury and PCBs, were found in the fish filets. MDCH recommends that people limit the amount of certain Kalamazoo River and Morrow Lake fish they eat. (Guidelines listed in this PHA are listed in the 2014-2015 Eat Safe Fish Guide.) Current guidelines can be found at: www.michigan.gov/eatsafefish.

Recommendations

- Avoid contact with oil sheen, if possible. If people do have direct contact with the sheen, wash skin with soap and water.
- Follow the fish consumption guidelines in the Eat Safe Fish Guide (www.michigan.gov/eatsafefish).

Public Health Action Plan

- MDCH will remain available as needed for health concerns related to the Kalamazoo River/Enbridge Spill.
- MDCH will update the Eat Safe Fish Guide as needed.

If individuals want additional information or have health concerns, please contact MDCH's Division of Environmental Health at 1-800-648-6942.

Report Preparation

This Public Health Assessment was prepared by the Michigan Department of Community Health under a cooperative agreement with the federal Agency for Toxic Substances and Disease Registry (ATSDR). It is in accordance with the approved agency methods, policies, procedures existing at the date of publication. Editorial review was completed by the cooperative agreement partner. ATSDR has reviewed this document and concurs with its findings based on the information presented. ATSDR's approval of this document has been captured in an electronic database, and the approving agency reviewers are listed below.

Author

Jennifer Gray, Ph.D.
Toxicologist

ATSDR Reviewers

Trent LeCoulre
Technical Project Officer
Division of Community Health Investigations

References

Agency for Toxic Substances and Disease Registry (ATSDR). Guidance on including child health issues in Division of Health Assessment and Consultation documents. July 2, 1998.

Enbridge Energy Partners, LLP (Enbridge). 2010 October. Enbridge Line 6B MP 608 Pipeline Release, Marshall, Michigan - Evaluation of Potential Impact of Released Oil on Groundwater used for Drinking Water (Hydrogeological Evaluation Report). Appendix A: Part 201 Sites of Environmental Contamination (FOIA)

http://www.epa.gov/enbridgespill/pdfs/enbridge_appendixa_part201.pdf.

Fismes, J., Perrin-Ganier, C., Empereur-Bissonnet, P., et al. 2002. Soil-to-root transfer and translocation of polycyclic aromatic hydrocarbons by vegetables grown on industrial contaminated soils. *J Environ Qual* 31(5): 1649-1656.

Heikes, D.L., Jensen, S.R., and Fleming-Jones, M.E. 1995. Purge and Trap Extraction with GC-MS Determination of Volatile Organic Compounds in Table-Ready Foods. *J Agric Food Chem* 43(11): 2869-2875.

International Programme on Chemical Safety (IPCS). Titanium (EHC 24, 1982). [updated 1982; accessed November]. Available from: <http://www.inchem.org/documents/ehc/ehc/ehc24.htm>.

Michigan Department of Environmental Quality (MDEQ). 2010a November. Administrative Consent Order and Partial Settlement Agreement. http://www.michigan.gov/documents/deq/deq-water-enbridge-DNRE-Admin_Order_350688_7.pdf.

Michigan Department of Environmental Quality (MDEQ). 2010b September. Laboratory data sheets – oil samples collected on August 6, 2010.

Samsoe-Petersen, L., Larsen, E. H., Larsen, P. B., et al. 2002. Uptake of trace elements and PAHs by fruit and vegetables from contaminated soils. *Environ Sci Technol* 36(14): 3057-3063.

U.S. Environmental Protection Agency (EPA). 1993. Provisional guidance for quantitative risk assessment of polycyclic aromatic hydrocarbons. EAP/600/R-93/089. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=49732#Download>.

U.S. National Library of Medicine (NLM). Hazardous Substance Data Bank: Titanium Compounds CASRN: No CAS RN. [updated August 2002; accessed February 2012]. Available from: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search>.

U.S. National Library of Medicine (NLM). Hazardous Substance Data Bank: Titanium Dioxide CASRN: 13463-67-7. [updated April 2009a; accessed February 2012]. Available from: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search>.

U.S. National Library of Medicine (NLM). Hazardous Substance Data Bank: 1,4-Dichloro-trans-2-butene CASRN: 110-57-6. [updated April 2009b; accessed October 2012]. Available from: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search>.

U.S. National Library of Medicine (NLM). Hazardous Substance Data Bank: 1,2,3-Trichloropropane CASRN: 96-18-4. [updated December 2009c; accessed October 2012]. Available from: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search>.

U.S. National Library of Medicine (NLM). Hazardous Substance Data Bank: p-Cymene CASRN: 99-87-6. [updated June 2010a; accessed November 2011]. Available from: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search>.

U.S. National Library of Medicine (NLM). Hazardous Substance Data Bank: Methyl Iodide CASRN: 74-88-4. [updated September 2010b; accessed October 2012]. Available from: <http://toxnet.nlm.nih.gov/cgi-bin/sis/search>.

Appendix A: Levels of chemicals in surface water

Analyte levels were compared to generic and site-specific screening levels. Since the Kalamazoo River, including Morrow Lake, is not used as a source for drinking water (it is surface water used for recreational purposes) and drinking water analytical methods were not used to measure the analytes, drinking water screening levels were not selected first.

Generic surface water screening levels were selected based on the hierarchy below.

1. First selected were the EPA's Regional Screening Levels for tapwater. These screening levels include ingestion of the water and dermal exposure (skin contact) typical of bathing for both adults and children. These screening levels account for a greater exposure than what people will have recreationally using the Kalamazoo River.
2. If the above was not available, an ATSDR drinking water comparison value was used. This value only accounts for ingestion. All values selected were protective for children and adults. This comparison value accounts for a greater exposure than people will have recreationally using the Kalamazoo River.
3. If none of the above was available, MDEQ's Rule 57 human non-drinking water value was used. There are two sets of values, one protective for the development of cancer and the other protective for the development of non-cancer effects. The lower of the two was selected. These values are protective for incidental ingestion of water, such as an accidental swallow of water, and accumulation of chemicals in the fish. These values will account for a greater exposure than people are expected to have as they assume people are swimming one hour a day for the recreational season (123 days).
4. If none of the above were available the MDEQ Residential Drinking Water Criteria (RDWC) was selected. These are the chemical levels that would not cause health effects if adults or children drank the water daily for 30 years. This criterion will account for a greater exposure than people are expected to have from recreational use of the Kalamazoo River.

If analytes were higher than the generic screening levels (described above), site-specific screening levels were calculated. These were calculated using the EPA's online screening level calculator (http://epa-prgs.ornl.gov/cgi-bin/chemicals/csl_search). This calculator uses chemical-specific default values and user-provided exposure parameters to determine screening levels for surface water. All available default values were used. Other variables were entered as presented in Table A-1. These variables represent a daily wading-type exposure (about one-third of an adult's body and a little less than half of a child's body [about 40%] covered in water) with ingestion of 3 to 5 milliliters of surface water daily for 150 days (5 months covering the May through October recreational season). Table A-2 lists both the generic and site-specific screening levels.

Table A-1: Variables used to calculate the site-specific surface water screening levels using the EPA's online screening level calculator for recreational use (http://epa-prgs.ornl.gov/cgi-bin/chemicals/csl_search).

Variable	Age range			
	0-2 years of age	2-6 years of age	6-16 years of age	16-30 years of age
Body weight (in kilograms)	10	15	50	80
Exposure duration (years)	2	4	10	14
Exposure frequency (days/year)	150	150	150	150
Exposure time (hours/event)	0.06	0.1	0.1	0.1
Events (events/day)	1	1	1	1
Water intake rate (liters/hour)	0.05	0.05	0.05	0.05
Skin surface area (square centimeters)	2,670	2,670	5,800	5,800

Table A-2: Generic and site-specific surface water screening levels in either micrograms per liter (µg/L) or milligrams per liter (mg/L).

Analyte	Units	Generic screening level ^a	Site-specific screening level ^b
1,1,1,2-Tetrachloroethane	µg/L	0.5	140
1,1,1-Trichloroethane	µg/L	7,500	ND ^c
1,1,2,2-Tetrachloroethane	µg/L	0.066	40
1,1,2-Trichloroethane	µg/L	0.24	200
1,1,2-Trichlorotrifluoroethane (Freon 113)	µg/L	53,000	ND
1,1'-Biphenyl	µg/L	0.83	ND
1,1-Dichloroethane	µg/L	2.4	ND
1,1-Dichloroethene	µg/L	260	ND
1,2,3-Trichlorobenzene	µg/L	5.2	ND
1,2,3-Trichloropropane	µg/L	0.00065	0.07
1,2,3-Trimethylbenzene	µg/L	10	ND
1,2,4,5-Tetrachlorobenzene	µg/L	1.2	30
1,2,4-Trichlorobenzene	µg/L	0.99	30
1,2,4-Trimethylbenzene	µg/L	15	ND
1,2-Dibromo-3-chloropropane	µg/L	0.0032	2
1,2-Dibromoethane	µg/L	0.0065	7
1,2-Dichlorobenzene	µg/L	280	ND
1,2-Dichloroethane	µg/L	0.15	170
1,2-Dichloropropane	µg/L	0.38	260
1,3,5-Trimethylbenzene	µg/L	87	ND
1,3-Dichlorobenzene	µg/L	200 ^d	ND
1,4-Dichlorobenzene	µg/L	0.42	290
1,4-Dioxane	µg/L	0.67	400

Table A-2 continued			
Analyte	Units	Generic screening level ^a	Site-specific screening level ^b
2,3,4,6-Tetrachlorophenol	µg/L	170	ND
2,4,5-T	µg/L	120	ND
2,4,5-TP (Silvex)	µg/L	84	ND
2,4,5-Trichlorophenol	µg/L	890	ND
2,4,6-Trichlorophenol	µg/L	3.5	130
2,4-D	µg/L	130	ND
2,4-Dichlorophenol	µg/L	35	ND
2,4-Dimethylphenol	µg/L	270	ND
2,4-Dinitrophenol	µg/L	30	7,500
2,4-Dinitrotoluene	µg/L	30	ND
2,6-Dinitrotoluene	µg/L	15	ND
2-Butanone (Methyl ethyl ketone)	µg/L	4,900	ND
2-Chloronaphthalene	µg/L	550	ND
2-Chlorophenol	µg/L	71	ND
2-Hexanone (Methyl butyl ketone)	µg/L	34	17,800
2-Methylnaphthalene	µg/L	27	780
2-Methylphenol	µg/L	720	ND
2-Nitroaniline	µg/L	150	ND
2-Nitrophenol	µg/L	20 ^d	ND
3- & 4-Methylphenol	µg/L	1,400	ND
3,3'-Dichlorobenzidine	µg/L	0.11	6
3-Nitroaniline	µg/L	NA ^f	ND
4,4'-DDD	µg/L	0.28	ND
4,4'-DDT	µg/L	0.2	ND
4,6-Dinitro-2-methylphenol	µg/L	1.2	210
4-Bromophenyl phenyl ether	µg/L	NA	ND
4-Chloro-3-methylphenol	µg/L	1,100	ND
4-Chloroaniline	µg/L	0.32	60
4-Chlorophenyl phenyl ether	µg/L	NA	ND
4-Methyl-2-pentanone	µg/L	1,000	ND
4-Methylphenol	µg/L	1,400	ND
4-Nitroaniline	µg/L	3.3	10
4-Nitrophenol	µg/L	60 ^g	ND
Acenaphthene	µg/L	400	ND
Acenaphthylene	µg/L	52 ^e	ND
Acetone	µg/L	12,000	ND
Acetonitrile	µg/L	130	ND
Acetophenone	µg/L	1,500	ND
Acrylonitrile	µg/L	0.045	60

Table A-2 continued			
Analyte	Units	Generic screening level ^a	Site-specific screening level ^b
Aldrin	µg/L	0.00021	0.004
alpha-BHC	µg/L	0.0062	0.2
alpha-Chlordane	µg/L	NA	ND
Aluminum	mg/L	16	ND
Anthracene	µg/L	1,300	ND
Antimony	mg/L	0.006	ND
Arsenic	mg/L	0.0047	0.03
Atrazine	µg/L	0.26	ND
Barium	mg/L	2.9	ND
Benzaldehyde	µg/L	1,500	ND
Benzene	µg/L	0.39	120
Benzo(a)anthracene	µg/L	0.029	20
Benzo(a)pyrene	µg/L	0.0029	2
Benzo(b)fluoranthene	µg/L	0.029	20
Benzo(g,h,i)perylene	µg/L	1 ^e	53,000
Benzo(k)fluoranthene	µg/L	0.29	170
Beryllium	mg/L	0.016	ND
beta-BHC	µg/L	0.022	ND
Bis(2-chloroethoxy)methane	µg/L	47	ND
Bis(2-chloroethyl)ether	µg/L	0.012	20
Bis(2-chloroisopropyl)ether	µg/L	0.31	ND
Bis(2-ethylhexyl)phthalate	µg/L	0.071	1
Bromobenzene	µg/L	54	ND
Bromochloromethane	µg/L	83	ND
Bromodichloromethane	µg/L	0.12	190
Bromoform	µg/L	7.9	ND
Bromomethane	µg/L	7	ND
Butyl benzyl phthalate	µg/L	14	ND
Cadmium	mg/L	0.0069	ND
Calcium	mg/L	NA	ND
Caprolactam	µg/L	7,700	ND
Carbazole	µg/L	41 ^h	ND
Carbon disulfide	µg/L	720	ND
Carbon tetrachloride	µg/L	0.39	60
Chlordane, technical	µg/L	0.027	ND
Chloride	mg/L	250 ^e	ND
Chlorobenzene	µg/L	72	ND
Chloroethane	µg/L	190	ND
Chloroform	µg/L	0.19	320

Table A-2 continued			
Analyte	Units	Generic screening level ^a	Site-specific screening level ^b
Chloromethane	µg/L	190	ND
Chromium	mg/L	0.00031	0.003
Chrysene	µg/L	2.9	1,700
cis-1,2-Dichloroethene	µg/L	28	ND
cis-1,3-Dichloropropene	µg/L	0.41	90
Cobalt	mg/L	0.0047	ND
Copper	mg/L	0.62	ND
Cyclohexane	µg/L	13,000	ND
delta-BHC	µg/L	NA	ND
Dibenz(a,h)anthracene	µg/L	0.0029	2
Dibenzofuran	µg/L	5.8	160
Dibromochloromethane	µg/L	0.15	150
Dibromomethane	µg/L	7.9	ND
Dichlorodifluoromethane (Freon 12)	µg/L	190	ND
Dieldrin	µg/L	0.0015	0.04
Diethyl ether	µg/L	3,100	ND
Diethyl phthalate	µg/L	11,000	ND
Di-isopropyl ether (DIPE)	µg/L	1,500	ND
Dimethoate	µg/L	3.1	1,300
Dimethyl phthalate	µg/L	73,000 ^e	ND
Di-n-butyl phthalate	µg/L	670	ND
Di-n-octyl phthalate	µg/L	4,000 ^d	ND
Disulfoton	µg/L	0.38	20
DRO (C10-C20)	µg/L	NA	ND
Endosulfan I	µg/L	NA	ND
Endosulfan II	µg/L	NA	ND
Endosulfan sulfate	µg/L	NA	ND
Endrin	µg/L	1.7	ND
Endrin aldehyde	µg/L	NA	ND
Endrin ketone	µg/L	NA	ND
Ethylbenzene	µg/L	1.3	170
Ethyl-tert-butyl ether (ETBE)	µg/L	130,000 ^h	ND
Famphur	µg/L	NA	ND
Fluoranthene	µg/L	630	ND
Fluorene	µg/L	220	ND
gamma-BHC (Lindane)	µg/L	0.036	ND
gamma-Chlordane	µg/L	NA	ND
GRO (C6-C10)	µg/L	NA	ND
Heptachlor	µg/L	0.0018	0.03

Table A-2 continued			
Analyte	Units	Generic screening level ^a	Site-specific screening level ^b
Heptachlor epoxide	µg/L	0.0033	0.1
Hexachlorobenzene	µg/L	0.042	30
Hexachlorobutadiene	µg/L	0.26	6
Hexachlorocyclopentadiene	µg/L	22	500
Hexachloroethane	µg/L	0.79	20
Indeno(1,2,3-c,d)pyrene	µg/L	0.029	20
Iron	mg/L	11	5,000
Isophorone	µg/L	67	ND
Isopropylbenzene (Cumene)	µg/L	390	ND
Lead	mg/L	0.19 ^h	ND
m- & p-Xylenes	µg/L	190	ND
Magnesium	mg/L	400 ^e	ND
Manganese	mg/L	0.32	ND
Mercury	mg/L	0.0016	ND
Methoxychlor	µg/L	27	ND
Methyl acetate	µg/L	16,000	ND
Methyl iodide	µg/L	NA	ND
Methyl parathion	µg/L	3.4	ND
Methyl tert-butyl ether	µg/L	12	ND
Methylcyclohexane	µg/L	NA	ND
Methylene chloride	µg/L	9.9	ND
Molybdenum	mg/L	0.078	ND
Naphthalene	µg/L	6.1	8,200
n-Butylbenzene	µg/L	780	ND
Nickel	mg/L	0.3	ND
Nitrate	mg/L	25	ND
Nitrite	mg/L	1.6	ND
Nitrobenzene	µg/L	0.12	5,000
n-Nitrosodi-n-propylamine	µg/L	0.0093	3
n-Nitrosodiphenylamine	µg/L	10	ND
n-Propylbenzene	µg/L	530	ND
o,o,o-Triethylphosphorothioate	µg/L	NA	ND
ORO (C20-C34)	µg/L	NA	ND
o-Xylene	µg/L	190	ND
Parathion	µg/L	65	ND
PCB-1016	µg/L	0.96	650
PCB-1221	µg/L	0.0043	0.2
PCB-1232	µg/L	0.0043	0.2
PCB-1242	µg/L	0.034	20

Table A-2 continued			
Analyte	Units	Generic screening level ^a	Site-specific screening level ^b
PCB-1248	µg/L	0.034	20
PCB-1254	µg/L	0.034	20
PCB-1260	µg/L	0.034	20
PCB-1262	µg/L	NA	ND
PCB-1268	µg/L	NA	ND
Pentachlorophenol	µg/L	0.17	110
Phenanthrene	µg/L	1.4 ⁱ	53,000
Phenol	µg/L	4,500	ND
Phorate	µg/L	2.3	140
Phosphorus, Total	mg/L	0.031	0.14
p-Isopropyltoluene	µg/L	NA	ND
Polychlorinated biphenyls, Total	µg/L	0.17	110
Potassium	mg/L	NA	ND
Pyrene	µg/L	87	1,900
sec-Butylbenzene	µg/L	80 ^e	ND
Selenium	mg/L	0.078	ND
Silver	mg/L	0.071	ND
Sodium	mg/L	120 ^e	ND
Styrene	µg/L	1,100	ND
Sulfate	mg/L	250 ^e	ND
Sulfide	mg/L	NA	ND
Sulfotep	µg/L	5.3	270
t-Amyl methyl ether (TAME)	µg/L	190 ^e	ND
t-Butyl alcohol	µg/L	1,500	ND
tert-Butylbenzene	µg/L	80 ^e	ND
Tetrachloroethene	µg/L	9.7	ND
Tetraethyl pyrophosphate (TEPP)	µg/L	NA	ND
Tetrahydrofuran	µg/L	3,200	ND
Thallium	mg/L	0.016	ND
Thionazin	µg/L	NA	ND
Titanium	µg/L	NA	ND
Toluene	µg/L	860	ND
Toxaphene	µg/L	0.013	0.3
trans-1,2- Dichloroethene	µg/L	86	ND
trans-1,3-Dichloropropene	µg/L	0.41	90
trans-1,4-Dichloro-2-butene	µg/L	0.0012	ND
Trichloroethene	µg/L	0.44	40
Trichlorofluoromethane (Freon 11)	µg/L	1,100	ND
Vanadium	mg/L	0.078	4

Table A-2 continued			
Analyte	Units	Generic screening level ^a	Site-specific screening level ^b
Vinyl acetate	µg/L	410	ND
Vinyl chloride	µg/L	0.015	1
Xylenes, total	µg/L	190	ND
Zinc	mg/L	4.7	ND

a = Unless otherwise noted, the generic screening levels are the EPA's Regional Screening Levels for tapwater.

b = Site-specific screening levels were calculated based the variables in Table A-1 and the EPA's online screening level calculator for recreational use (http://epa-prgs.ornl.gov/cgi-bin/chemicals/csl_search).

c = A site-specific screening levels was not determined (ND).

d = This is the ATSDR's child intermediate child Environmental Media Evaluation Guide.

e = This is the MDEQ's Part 201 Residential Drinking Water Criterion.

f = No generic screening levels were available (NA).

g = This is one of ATSDR's Comparison Values (EPA's Lifetime Health Advisory).

h = This is the MDEQ's Rule 57 human non-drink value.

i = This is the MDEQ's Rule 57 Final Chronic Value, which is protective of human exposure.

Table A-3: Levels of analytes (in either microgram per liter [$\mu\text{g/L}$] or milligrams per liter [mg/L]) in surface water samples collected between July to September 2010.

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,1,1,2-Tetrachloroethane	$\mu\text{g/L}$	140	1398	0	ND ^a	ND	NA ^b	1398	1	1	0
1,1,1-Trichloroethane	$\mu\text{g/L}$	7,500	1669	0	ND	ND	NA	1669	1	1	0
1,1,2,2-Tetrachloroethane	$\mu\text{g/L}$	40	1669	0	ND	ND	NA	1669	1	1	0
1,1,2-Trichloroethane	$\mu\text{g/L}$	200	1669	0	ND	ND	NA	1669	1	1	0
1,1,2-Trichlorotrifluoroethane (Freon 113)	$\mu\text{g/L}$	53,000	1669	0	ND	ND	NA	1669	1	1	0
1,1'-Biphenyl	$\mu\text{g/L}$	0.83	273	0	ND	ND	NA	273	0.5	0.59	0
1,1-Dichloroethane	$\mu\text{g/L}$	2.4	1669	0	ND	ND	NA	1669	1	1	0
1,1-Dichloroethene	$\mu\text{g/L}$	260	1669	2	0.25	0.35	0	1667	1	1	0
1,2,3-Trichlorobenzene	$\mu\text{g/L}$	5.2	271	0	ND	ND	NA	271	1	1	0
1,2,3-Trichloropropane	$\mu\text{g/L}$	0.07	1398	0	ND	ND	NA	1398	1	1	1398
1,2,4,5-Tetrachlorobenzene	$\mu\text{g/L}$	30	273	0	ND	ND	NA	273	2	2.4	0
1,2,4-Trichlorobenzene	$\mu\text{g/L}$	30	3067	1	0.3	0.3	0	3066	1	14	0
1,2,4-Trimethylbenzene	$\mu\text{g/L}$	15	1686	32	0.15	50	2	1654	1	2	0
1,2-Dibromo-3-chloropropane	$\mu\text{g/L}$	2	1669	0	ND	ND	NA	1669	1	2	0
1,2-Dibromoethane	$\mu\text{g/L}$	7	1669	0	ND	ND	NA	1669	1	1	0
1,2-Dichlorobenzene	$\mu\text{g/L}$	280	3067	2	0.32	0.35	0	3065	1	14	0
1,2-Dichloroethane	$\mu\text{g/L}$	170	1669	0	ND	ND	NA	1669	1	1	0
1,2-Dichloropropane	$\mu\text{g/L}$	260	1669	0	ND	ND	NA	1669	1	2	0
1,3,5-Trimethylbenzene	$\mu\text{g/L}$	87	1686	11	0.13	50	0	1675	1	2	0

Table A-3 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,3-Dichlorobenzene	µg/L	200	3067	2	0.34	0.35	0	3065	1	14	0
1,4-Dichlorobenzene	µg/L	290	3067	2	0.34	0.38	0	3065	1	14	0
1,4-Dioxane	µg/L	400	271	0	ND	ND	NA	271	25	25	0
2,3,4,6-Tetrachlorophenol	µg/L	170	273	0	ND	ND	NA	273	5	5.9	0
2,4,5-T	µg/L	120	5	0	ND	ND	NA	5	1	1	0
2,4,5-TP (Silvex)	µg/L	84	5	0	ND	ND	NA	5	2	2	0
2,4,5-Trichlorophenol	µg/L	890	1671	0	ND	ND	NA	1671	0.5	14	0
2,4,6-Trichlorophenol	µg/L	130	1671	0	ND	ND	NA	1671	0.5	14	0
2,4-D	µg/L	130	5	0	ND	ND	NA	5	2	2	0
2,4-Dichlorophenol	µg/L	35	1671	0	ND	ND	NA	1671	0.5	29	0
2,4-Dimethylphenol	µg/L	270	1671	0	ND	ND	NA	1671	1	14	0
2,4-Dinitrophenol	µg/L	7,500	1671	0	ND	ND	NA	1671	5	57	0
2,4-Dinitrotoluene	µg/L	30	1671	0	ND	ND	NA	1671	0.5	14	0
2,6-Dinitrotoluene	µg/L	15	1671	0	ND	ND	NA	1671	0.47	14	0
2-Butanone (Methyl Ethyl Ketone)	µg/L	4,900	1669	7	0.69	7.9	0	1662	5	25	0
2-Chloronaphthalene	µg/L	550	1785	0	ND	ND	NA	1785	0.5	14	0
2-Chlorophenol	µg/L	71	1670	0	ND	ND	NA	1670	0.5	14	0
2-Hexanone (Methyl Butyl Ketone)	µg/L	17,800	1669	0	ND	ND	NA	1669	5	50	0
2-Methylnaphthalene	µg/L	780	3643	18	0.031	63	0	3625	0.47	14	0
2-Methylphenol	µg/L	720	1647	0	ND	ND	NA	1647	0.5	14	0
2-Nitroaniline	µg/L	150	1671	0	ND	ND	NA	1671	1	57	0
2-Nitrophenol	µg/L	20	1671	0	ND	ND	NA	1671	0.5	14	0
3- & 4- Methylphenol	µg/L	1,400	610	0	ND	ND	NA	610	1.9	2.8	0

Table A-3 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
3,3'-Dichlorobenzidine	µg/L	6	1671	0	ND	ND	NA	1671	4	57	731
3-Nitroaniline	µg/L	NA	1671	0	ND	ND	NA	1671	1	57	NA
4,4'-DDD	µg/L	0.28	10	0	ND	ND	NA	10	0.02	0.029	0
4,4'-DDT	µg/L	0.2	5	0	ND	ND	NA	5	0.02	0.029	0
4,6-Dinitro-2-methylphenol	µg/L	210	1671	0	ND	ND	NA	1671	0.5	57	0
4-Bromophenyl phenyl ether	µg/L	NA	1671	0	ND	ND	NA	1671	0.5	14	NA
4-Chloro-3-methylphenol	µg/L	1,100	1671	6	0.19	0.21	0	1665	0.5	14	0
4-Chloroaniline	µg/L	60	1671	0	ND	ND	NA	1671	1	57	0
4-Chlorophenyl phenyl ether	µg/L	NA	1671	0	ND	ND	NA	1671	0.5	14	NA
4-Methyl-2-pentanone (MIBK)	µg/L	1,000	1669	0	ND	ND	NA	1669	5	50	0
4-Methylphenol	µg/L	1,400	1037	1	0.45	0.45	0	1036	0.5	14	0
4-Nitroaniline	µg/L	10	1671	0	ND	ND	NA	1671	1	57	1292
4-Nitrophenol	µg/L	60	1671	0	ND	ND	NA	1671	5	57	0
Acenaphthene	µg/L	400	1957	4	0.04	0.4	0	1953	0.5	14	0
Acenaphthylene	µg/L	52	1957	2	0.08	0.13	0	1955	0.5	14	0
Acetone	µg/L	12,000	1669	168	0.36	64	0	1501	0.67	50	0
Acetophenone	µg/L	1,500	273	1	0.09	0.09	0	272	0.5	1	0
Acrylonitrile	µg/L	60	788	0	ND	ND	NA	788	1	2	0
Aldrin	µg/L	0.004	5	0	ND	ND	NA	5	0.01	0.014	5
alpha-BHC	µg/L	0.2	5	0	ND	ND	NA	5	0.01	0.014	0
alpha-Chlordane	µg/L	NA	5	0	ND	ND	NA	5	0.02	0.029	NA
Aluminum	mg/L	16	488	430	0.013	7.27	0	58	0.016	0.1	0
Anthracene	µg/L	1,300	1957	5	0.04	0.22	0	1952	0.5	14	0

Table A-3 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Antimony	mg/L	0.006	1425	273	0.00013	0.0046	0	1152	0.00013	0.005	0
Arsenic	mg/L	0.03	1425	1299	0.00041	0.03	0	126	0.001	0.005	0
Atrazine	µg/L	0.26	273	0	ND	ND	NA	273	0.5	1	273
Barium	mg/L	2.9	1425	1242	0.0065	0.66	0	183	0.0098	0.1	0
Benzaldehyde	µg/L	1,500	273	2	0.5	0.51	0	271	0.5	1	0
Benzene	µg/L	120	1957	34	0.18	49	0	1923	1	1	0
Benzo(a)anthracene	µg/L	20	1957	28	0.03	25	1	1929	0.19	14	0
Benzo(a)pyrene	µg/L	2	1957	44	0.03	38	2	1913	0.19	14	713
Benzo(b)fluoranthene	µg/L	20	1957	28	0.03	2.5	0	1929	0.19	14	0
Benzo(g,h,i)perylene	µg/L	53,000	1957	17	0.061	2.2	0	1940	0.5	14	0
Benzo(k)fluoranthene	µg/L	170	1957	26	0.051	0.54	0	1931	0.19	14	0
Beryllium	mg/L	0.016	1425	92	4.50E-05	0.00058	0	1333	4.40E-05	0.002	0
beta-BHC	µg/L	0.022	5	0	ND	ND	NA	5	0.01	0.014	0
Bis(2-chloroethoxy)methane	µg/L	47	1671	0	ND	ND	NA	1671	0.5	14	0
Bis(2-chloroethyl)ether	µg/L	20	1671	1	0.22	0.22	0	1670	0.5	14	0
Bis(2-chloroisopropyl)ether	µg/L	0.31	1671	0	ND	ND	NA	1671	0.5	14	1671
Bis(2-ethylhexyl)phthalate	µg/L	1	1671	115	0.25	47	30	1556	0.5	14	940
Bromochloromethane	µg/L	83	1669	0	ND	ND	NA	1669	1	1	0
Bromodichloromethane	µg/L	190	1669	14	0.15	5.3	0	1655	1	1	0
Bromoform	µg/L	7.9	1669	1	0.64	0.64	0	1668	1	1	0
Bromomethane	µg/L	7	1669	1	0.47	0.47	0	1668	1	5	0
Butyl benzyl phthalate	µg/L	14	1671	10	0.061	1.1	0	1661	1	14	0
Cadmium	mg/L	0.0069	1425	61	0.0001	0.0052	0	1364	0.00024	0.001	0
Calcium	mg/L	NA	717	661	19.8	79,000	NA	56	62.6	88.1	NA

Table A-3 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Caprolactam	µg/L	7,700	243	0	ND	ND	NA	243	1	1.2	0
Carbazole	µg/L	41	1637	0	ND	ND	NA	1637	0.5	29	0
Carbon disulfide	µg/L	720	1669	31	0.19	1.8	0	1638	2.5	5	0
Carbon tetrachloride	µg/L	60	1669	0	ND	ND	NA	1669	1	1	0
Chlordane, Technical	µg/L	0.5	5	0	ND	ND	NA	5	0.5	0.71	1
Chlorobenzene	µg/L	72	1669	0	ND	ND	NA	1669	1	1	0
Chloroethane	µg/L	190	1669	0	ND	ND	NA	1669	1	5	0
Chloroform	µg/L	320	1669	40	0.16	6.8	0	1629	1	1	0
Chloromethane	µg/L	190	1669	10	0.32	1.4	0	1659	1	5	0
Chromium	mg/L	0.003	1425	715	3.90E-05	0.11	25	710	4.10E-05	0.01	366
Chrysene	µg/L	1,700	1957	24	0.03	28	1	1933	0.47	14	0
cis-1,2-Dichloroethene	µg/L	28	1669	0	ND	ND	NA	1669	1	1	0
cis-1,3-Dichloropropene	µg/L	90	1669	1	0.66	0.66	0	1668	1	1	0
Cobalt	mg/L	2	1425	1177	5.90E-05	0.0041	0	248	0.00015	0.02	0
Copper	mg/L	283	1425	908	0.0003	0.668	0	517	0.00037	0.005	0
Cyclohexane	µg/L	13,000	271	0	ND	ND	NA	271	5	5	0
delta-BHC	µg/L	NA	5	0	ND	ND	NA	5	0.01	0.014	NA
Dibenzo(a,h)anthracene	µg/L	2	1957	11	0.1	1	0	1946	0.28	14	730
Dibenzofuran	µg/L	160	1671	0	ND	ND	NA	1671	0.5	14	0
Dibromochloromethane	µg/L	150	1669	6	0.51	3.7	0	1663	1	5	0
Dibromomethane	µg/L	7.9	1398	0	ND	ND	NA	1398	1	5	0
Dichlorodifluoromethane (Freon 12)	µg/L	190	1669	0	ND	ND	NA	1669	1	5	0
Dieldrin	µg/L	0.04	5	0	ND	ND	NA	5	0.02	0.029	0

Table A-3 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Diethyl ether	µg/L	3,100	1398	14	0.33	2.5	0	1384	1	10	0
Diethyl phthalate	µg/L	11,000	1671	27	0.25	1.5	0	1644	0.5	57	0
Dimethoate	µg/L	1,300	5	0	ND	ND	NA	5	10	10	0
Dimethyl phthalate	µg/L	73,000	1671	0	ND	ND	NA	1671	0.5	57	0
Di-n-butyl phthalate	µg/L	670	1671	17	0.27	0.88	0	1654	1	14	0
Di-n-octyl phthalate	µg/L	4,000	1671	6	0.39	1.4	0	1665	0.5	14	0
Disulfoton	µg/L	20	5	0	ND	ND	NA	5	5	5	0
DRO (C10-C20)	µg/L	NA	1517	106	15	7,300	NA	1411	100	2,500	NA
DRO (C10-C28)	µg/L	NA	159	11	65	390	NA	148	200	300	NA
DRO (C10-C38)	µg/L	NA	286	61	0.24	27,000	NA	225	100	122	NA
Endosulfan I	µg/L	NA	5	0	ND	ND	NA	5	0.02	0.029	NA
Endosulfan II	µg/L	NA	5	0	ND	ND	NA	5	0.02	0.029	NA
Endosulfan sulfate	µg/L	NA	5	0	ND	ND	NA	5	0.02	0.029	NA
Endrin	µg/L	1.7	5	0	ND	ND	NA	5	0.02	0.029	0
Endrin aldehyde	µg/L	NA	5	0	ND	ND	NA	5	0.02	0.029	NA
Endrin ketone	µg/L	NA	5	0	ND	ND	NA	5	0.02	0.029	NA
Ethylbenzene	µg/L	170	1957	24	0.16	43	0	1933	1	1	0
Famphur	µg/L	NA	5	0	ND	ND	NA	5	20	20	NA
Fluoranthene	µg/L	630	1957	45	0.041	2.8	0	1912	0.5	14	0
Fluorene	µg/L	220	1957	3	0.07	1.1	0	1954	0.5	14	0
gamma-BHC (Lindane)	µg/L	0.036	5	0	ND	ND	NA	5	0.01	0.014	0
gamma-Chlordane	µg/L	NA	5	0	ND	ND	NA	5	0.02	0.029	NA
GRO (C5-C10)	µg/L	NA	320	5	130	540	NA	315	100	200	NA
GRO (C6-C10)	µg/L	NA	1632	43	11	85	NA	1589	50	200	NA

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Heptachlor	µg/L	0.03	5	0	ND	ND	NA	5	0.01	0.014	0
Heptachlor epoxide	µg/L	0.09	5	0	ND	ND	NA	5	0.01	0.014	0
Hexachlorobenzene	µg/L	30	1676	1	0.23	0.23	0	1675	0.01	14	0
Hexachlorobutadiene	µg/L	6	1671	0	ND	ND	NA	1671	0.5	14	22
Hexachlorocyclopentadiene	µg/L	500	1671	0	ND	ND	NA	1671	0.5	57	0
Hexachloroethane	µg/L	20	2459	0	ND	ND	NA	2459	0.5	14	0
Indeno(1,2,3-cd)pyrene	µg/L	20	1957	20	0.061	0.97	0	1937	0.19	14	0
Iron	mg/L	5,000	1425	1402	0.0304	50	0	23	0.076	0.2	0
Isophorone	µg/L	67	1671	0	ND	ND	NA	1671	0.5	14	0
Isopropylbenzene (Cumene)	µg/L	390	1669	1	0.49	0.49	0	1668	1	5	0
Lead	mg/L	0.19	1425	1098	4.40E-05	0.247	2	327	4.40E-05	0.003	0
m- & p-Xylene	µg/L	190	1957	57	0.32	93	0	1900	1	2	0
Magnesium	mg/L	400	717	704	6.3	27	0	13	19.4	25.4	0
Manganese	mg/L	80	1425	1409	0.018	5.3	0	16	0.026	0.097	0
Mercury	mg/L	0.0016	1425	31	9.00E-06	0.00043	0	1394	0.0002	0.0002	0
Methoxychlor	µg/L	27	5	0	ND	ND	NA	5	0.04	0.057	0
Methyl Acetate	µg/L	16,000	271	0	ND	ND	NA	271	5	5	0
Methyl iodide	µg/L	NA	1398	0	ND	ND	NA	1398	1	5	NA
Methyl parathion	µg/L	400	5	0	ND	ND	NA	5	5	5	0
Methyl tert-butyl ether	µg/L	12	1669	0	ND	ND	NA	1669	1	5	0
Methylcyclohexane	µg/L	NA	271	0	ND	ND	NA	271	5	5	NA
Methylene chloride	µg/L	9.9	1669	12	0.9	3.6	0	1657	1	5	0
Molybdenum	mg/L	0.078	937	370	0.00033	0.0023	0	567	5.10E-05	0.0057	0
Naphthalene	µg/L	8,200	3529	23	0.031	57	0	3506	0.5	14	0

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Nickel	mg/L	0.3	1638	1056	0.00013	0.072	0	582	0.00015	0.02	0
Nitrate	mg/L	25	5	5	1.8	2	0	NA	NA	NA	NA
Nitrite	mg/L	1.6	5	0	ND	ND	NA	5	0.02	0.02	0
Nitrobenzene	µg/L	5,000	1671	0	ND	ND	NA	1671	0.5	14	0
n-Nitrosodi-n-propylamine	µg/L	3	1671	2	1.1	7.9	1	1669	0.47	14	1046
n-Nitrosodiphenylamine	µg/L	680	1671	0	ND	ND	NA	1671	0.5	14	0
n-Propylbenzene	µg/L	530	1398	2	0.15	0.16	0	1396	1	1	0
^{o,o,o} -Triethylphosphorothioate	µg/L	NA	5	0	ND	ND	NA	5	10	10	NA
ORO (C20-C34)	µg/L	NA	1398	101	46	2,700	NA	1297	100	5,000	NA
ORO (C28-C36)	µg/L	NA	261	34	85	8,500	NA	227	200	410	NA
o-Xylene	µg/L	190	1957	31	0.15	48	0	1926	1	1	0
Parathion	µg/L	65	10	0	ND	ND	NA	10	10	10	0
PCB-1016	µg/L	650	1663	0	ND	ND	NA	1663	0.12	2.8	NA
PCB-1221	µg/L	0.2	1663	0	ND	ND	NA	1663	0.12	2.8	672
PCB-1232	µg/L	0.2	1663	0	ND	ND	NA	1663	0.12	2.8	672
PCB-1242	µg/L	20	1663	0	ND	ND	NA	1663	0.12	2.8	0
PCB-1248	µg/L	20	1663	14	0.064	9.3	0	1649	0.12	0.73	0
PCB-1254	µg/L	20	1663	5	0.03	0.1	0	1658	0.12	2.8	0
PCB-1260	µg/L	20	1663	1	0.46	0.46	0	1662	0.12	2.8	0
PCB-1262	µg/L	NA	271	0	ND	ND	NA	271	0.2	0.23	NA
PCB-1268	µg/L	NA	271	0	ND	ND	NA	271	0.2	0.23	NA
Pentachlorophenol	µg/L	110	1671	0	ND	ND	NA	1671	0.5	57	0
Phenanthrene	µg/L	53,000	1957	32	0.03	2.4	0	1925	0.5	14	0

Table A-3 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Phenol	µg/L	4,500	1671	0	ND	ND	NA	1671	0.5	14	0
Phorate	µg/L	140	5	0	ND	ND	NA	5	5	5	0
Phosphorus, Total	mg/L	0.14	5	3	0.033	0.044	0	2	0.05	0.05	0
Polychlorinated biphenyls, Total	µg/L	110	351	15	0.03	9.3	0	336	0.12	0.73	0
Potassium	mg/L	NA	488	380	1	90.3	NA	108	0.622	10	NA
Pyrene	µg/L	87	1957	72	0.03	27	0	1885	0.5	14	0
Selenium	mg/L	0.078	1425	112	0.00042	0.0013	0	1313	0.00081	0.005	0
Silver	mg/L	0.071	1425	86	1.30E-05	0.003	0	1339	1.30E-05	0.0005	0
Sodium	mg/L	120	488	488	1.7	76.3	0	NA	NA	NA	NA
Styrene	µg/L	1,100	1669	1	0.5	0.5	0	1668	1	1	0
Sulfate	mg/L	250	5	5	36	49	0	NA	NA	NA	NA
Sulfide	mg/L	NA	6	0	ND	ND	NA	6	1	1	NA
Sulfotepp (Tetraethyl Dithiopyrophosphate)	µg/L	270	5	0	ND	ND	NA	5	10	10	0
Tetrachloroethene	µg/L	9.7	1669	0	ND	ND	NA	1669	1	2	0
Tetraethyl pyrophosphate (TEPP)	µg/L	NA	5	0	ND	ND	NA	5	10	10	NA
Thallium	mg/L	0.016	1425	88	0.00011	0.0014	0	1337	0.00012	0.005	0
Thionazin	µg/L	NA	5	0	ND	ND	NA	5	5	5	NA
Toluene	µg/L	860	1957	173	0.14	46	0	1784	1	1	0
Toxaphene	µg/L	0.3	5	0	ND	ND	NA	5	2	2.9	5
trans-1,2-Dichloroethene	µg/L	86	1669	0	ND	ND	NA	1669	1	1	0
trans-1,3-Dichloropropene	µg/L	90	1669	0	ND	ND	NA	1669	1	1	0
trans-1,4-Dichloro-2-butene	µg/L	0.0012	1398	0	ND	ND	NA	1398	1	10	1398

Table A-3 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Trichloroethene	µg/L	40	1669	0	ND	ND	NA	1669	1	1	0
Trichlorofluoromethane (Freon 11)	µg/L	1,100	1669	0	ND	ND	NA	1669	1	1	0
Vanadium	mg/L	4	1638	815	0.00013	0.14	0	823	0.00013	0.01	0
Vinyl acetate	µg/L	410	1398	0	ND	ND	NA	1398	1	100	0
Vinyl chloride	µg/L	1	1669	0	ND	ND	NA	1669	1	1	0
Xylenes, Total	µg/L	190	1513	41	0.32	4.3	0	1472	1	3	0
Zinc	mg/L	4.7	1425	791	0.0013	0.726	0	634	0.0013	0.05	0

a = These chemicals were not detected (ND).

b= No values were available (NA).

Table A-4: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between October to December 2010.

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,1,1,2-Tetrachloroethane	µg/L	140	162	0	ND ^a	ND	NA ^b	162	1	1	0
1,1,1-Trichloroethane	µg/L	7,500	162	0	ND	ND	NA	162	1	1	0
1,1,2,2-Tetrachloroethane	µg/L	40	162	0	ND	ND	NA	162	1	1	0
1,1,2-Trichloroethane	µg/L	200	162	0	ND	ND	NA	162	1	1	0
1,1,2-Trichlorotrifluoroethane (Freon 113)	µg/L	53,000	162	0	ND	ND	NA	162	1	1	0

Table A-4 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,1-Dichloroethane	µg/L	2.4	162	0	ND	ND	NA	162	1	1	0
1,1-Dichloroethene	µg/L	260	162	0	ND	ND	NA	162	1	1	0
1,2,3-Trichloropropane	µg/L	0.07	162	0	ND	ND	NA	162	1	1	162
1,2,3-Trimethylbenzene	µg/L	10	215	0	ND	ND	NA	215	1	1	0
1,2,4-Trichlorobenzene	µg/L	30	324	0	ND	ND	NA	324	1	10	0
1,2,4-Trimethylbenzene	µg/L	15	377	7	0.22	1.9	0	370	1	1	0
1,2-Dibromo-3-chloropropane	µg/L	2	162	0	ND	ND	NA	162	1	1	0
1,2-Dibromoethane	µg/L	7	162	0	ND	ND	NA	162	1	1	0
1,2-Dichlorobenzene	µg/L	280	324	0	ND	ND	NA	324	1	10	0
1,2-Dichloroethane	µg/L	170	162	0	ND	ND	NA	162	1	1	0
1,2-Dichloropropane	µg/L	260	162	0	ND	ND	NA	162	2	2	0
1,3,5-Trimethylbenzene	µg/L	87	377	5	0.11	0.48	0	372	1	1	0
1,3-Dichlorobenzene	µg/L	200	324	0	ND	ND	NA	324	2	10	0
1,4-Dichlorobenzene	µg/L	290	324	0	ND	ND	NA	324	2	10	0
2,4,5-Trichlorophenol	µg/L	890	162	0	ND	ND	NA	162	5	10	0
2,4,6-Trichlorophenol	µg/L	130	162	0	ND	ND	NA	162	5	10	0
2,4-Dichlorophenol	µg/L	35	162	0	ND	ND	NA	162	10	20	0
2,4-Dimethylphenol	µg/L	270	162	0	ND	ND	NA	162	5	10	0
2,4-Dinitrophenol	µg/L	7,500	162	0	ND	ND	NA	162	20	40	0
2,4-Dinitrotoluene	µg/L	30	162	0	ND	ND	NA	162	5	10	0
2,6-Dinitrotoluene	µg/L	15	162	0	ND	ND	NA	162	5	10	0
2-Butanone (Methy Ethyl Ketone)	µg/L	4,900	162	0	ND	ND	NA	162	5	5	0
2-Chloronaphthalene	µg/L	550	162	0	ND	ND	NA	162	5	10	0

Table A-4 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
2-Chlorophenol	µg/L	71	162	0	ND	ND	NA	162	5	10	0
2-Hexanone (Methyl Butyl Ketone)	µg/L	34	162	0	ND	ND	NA	162	5	5	0
2-Methylnaphthalene	µg/L	27	754	4	0.9	6.4	0	750	1	10	0
2-Methylphenol	µg/L	720	162	0	ND	ND	NA	162	5	10	0
2-Nitroaniline	µg/L	150	162	0	ND	ND	NA	162	20	40	0
2-Nitrophenol	µg/L	20	162	0	ND	ND	NA	162	5	10	0
3,3'-Dichlorobenzidine	µg/L	6	162	0	ND	ND	NA	162	20	40	162
3-Nitroaniline	µg/L	NA	162	0	ND	ND	NA	162	20	40	NA
4,6-Dinitro-2-methylphenol	µg/L	210	162	0	ND	ND	NA	162	20	40	0
4-Bromophenyl phenyl ether	µg/L	NA	162	0	ND	ND	NA	162	5	10	NA
4-Chloro-3-methylphenol	µg/L	1,100	162	0	ND	ND	NA	162	5	10	0
4-Chloroaniline	µg/L	60	162	0	ND	ND	NA	162	20	40	0
4-Chlorophenyl phenyl ether	µg/L	NA	162	0	ND	ND	NA	162	5	10	NA
4-Methyl-2-pentanone	µg/L	1,000	162	0	ND	ND	NA	162	5	5	0
4-Methylphenol	µg/L	1,400	162	0	ND	ND	NA	162	5	10	0
4-Nitroaniline	µg/L	10	162	0	ND	ND	NA	162	20	40	162
4-Nitrophenol	µg/L	60	162	0	ND	ND	NA	162	20	40	0
Acenaphthene	µg/L	400	162	0	ND	ND	NA	162	5	10	0
Acenaphthylene	µg/L	52	162	0	ND	ND	NA	162	5	10	0
Acetone	µg/L	12,000	162	3	0.7	3.4	0	159	0.39	20	0
Acrylonitrile	µg/L	60	162	0	ND	ND	NA	162	1	1	0
Anthracene	µg/L	1,300	162	0	ND	ND	NA	162	5	10	0
Antimony	mg/L	0.006	162	90	0.0001	0.0045	0	72	0.0002	0.002	0

Table A-4 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Arsenic	mg/L	0.03	162	156	0.0006	0.0017	0	6	0.005	0.005	0
Barium	mg/L	2.9	162	162	0.026	0.086	0	NA	NA	NA	NA
Benzene	µg/L	120	377	7	0.28	1.6	0	370	1	1	0
Benzo(a)anthracene	µg/L	20	162	0	ND	ND	NA	162	5	10	0
Benzo(a)pyrene	µg/L	2	162	0	ND	ND	NA	162	5	10	162
Benzo(b)fluoranthene	µg/L	20	162	0	ND	ND	NA	162	5	10	0
Benzo(g,h,i)perylene	µg/L	53,000	162	0	ND	ND	NA	162	5	10	0
Benzo(k)fluoranthene	µg/L	170	162	0	ND	ND	NA	162	5	10	0
Beryllium	mg/L	0.016	377	37	5.00E-05	0.0007	0	340	5.00E-05	0.001	0
Bis(2-chloroethoxy)methane	µg/L	47	162	0	ND	ND	NA	162	5	10	0
Bis(2-chloroethyl) Ether	µg/L	20	162	0	ND	ND	NA	162	5	10	0
Bis(2-chloroisopropyl) Ether	µg/L	0.31	162	0	ND	ND	NA	162	5	10	162
Bis(2-ethylhexyl) Phthalate	µg/L	1	162	2	0.41	0.81	0	160	5	10	160
Bromochloromethane	µg/L	83	162	0	ND	ND	NA	162	1	1	0
Bromodichloromethane	µg/L	190	162	0	ND	ND	NA	162	1	1	0
Bromoform	µg/L	7.9	162	0	ND	ND	NA	162	1	1	0
Bromomethane	µg/L	7	162	0	ND	ND	NA	162	1	1	0
Butyl Benzyl Phthalate	µg/L	14	162	0	ND	ND	NA	162	5	10	0
Cadmium	mg/L	0.0069	162	1	0.0001	0.0001	0	161	0.001	0.001	0
Calcium	mg/L	NA	8	8	74	88	NA	NA	NA	NA	NA
Carbazole	µg/L	41	162	0	ND	ND	NA	162	10	20	0
Carbon disulfide	µg/L	720	162	0	ND	ND	NA	162	2.5	2.5	0
Carbon tetrachloride	µg/L	60	162	0	ND	ND	NA	162	1	1	0
Chloride (as Cl)	mg/L	250	8	8	36	50	0	NA	NA	NA	NA

Table A-4 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Chlorobenzene	µg/L	72	162	0	ND	ND	NA	162	1	1	0
Chloroethane	µg/L	190	162	0	ND	ND	NA	162	1	1	0
Chloroform	µg/L	320	162	0	ND	ND	NA	162	1	1	0
Chloromethane	µg/L	190	162	0	ND	ND	NA	162	1	1	0
Chromium	mg/L	0.003	162	125	7.00E-05	0.0023	0	37	0.0001	0.005	3
Chrysene	µg/L	1,700	162	0	ND	ND	NA	162	5	10	0
cis-1,2-Dichloroethene	µg/L	28	162	0	ND	ND	NA	162	1	1	0
cis-1,3-Dichloropropene	µg/L	90	162	0	ND	ND	NA	162	1	1	0
Cobalt	mg/L	0.0047	162	142	0.0001	0.0006	0	20	0.0003	0.0004	0
Copper	mg/L	0.62	162	135	0.0004	0.019	0	27	0.001	0.005	0
Cyclohexane	µg/L	13,000	215	0	ND	ND	NA	215	1	1	0
Dibenz(a,h)anthracene	µg/L	2	162	0	ND	ND	NA	162	5	10	162
Dibenzofuran	µg/L	160	162	0	ND	ND	NA	162	5	10	0
Dibromochloromethane	µg/L	150	162	0	ND	ND	NA	162	1	1	0
Dibromomethane	µg/L	7.9	162	0	ND	ND	NA	162	1	1	0
Dichlorodifluoromethane (Freon 12)	µg/L	190	162	0	ND	ND	NA	162	1	1	0
Diethyl ether	µg/L	3,100	162	0	ND	ND	NA	162	10	10	0
Diethyl Phthalate	µg/L	11,000	162	11	0.25	0.72	0	151	20	40	0
Dimethyl Phthalate	µg/L	73,000	162	0	ND	ND	NA	162	20	40	0
Di-n-butyl Phthalate	µg/L	670	162	0	ND	ND	NA	162	5	10	0
Di-n-octyl Phthalate	µg/L	4,000	162	0	ND	ND	NA	162	5	10	0
DRO (C10-C20)	µg/L	NA	377	0	ND	ND	NA	377	100	200	NA
Ethylbenzene	µg/L	170	377	6	0.22	0.97	0	371	1	1	0

Table A-4 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Fluoranthene	µg/L	630	162	0	ND	ND	NA	162	5	10	0
Fluorene	µg/L	220	162	0	ND	ND	NA	162	5	10	0
GRO (C6-C10)	µg/L	NA	377	0	ND	ND	NA	377	200	200	NA
Hexachlorobenzene	µg/L	30	162	0	ND	ND	NA	162	5	10	0
Hexachlorobutadiene	µg/L	6	162	0	ND	ND	NA	162	5	10	1
Hexachlorocyclopentadiene	µg/L	500	162	0	ND	ND	NA	162	20	40	0
Hexachloroethane	µg/L	20	324	0	ND	ND	NA	324	1	10	0
Indeno(1,2,3-cd)pyrene	µg/L	20	162	0	ND	ND	NA	162	5	10	0
Iron	mg/L	11	385	384	0.14	3.3	0	1	0.2	0.2	0
Isophorone	µg/L	67	162	0	ND	ND	NA	162	5	10	0
Isopropylbenzene (Cumene)	µg/L	390	377	0	ND	ND	NA	377	1	1	0
Lead	mg/L	0.19	162	114	4.00E-05	0.0022	0	48	5.00E-05	0.003	0
m- & p-Xylene	µg/L	190	377	12	0.36	3.6	0	365	1	2	0
Magnesium	mg/L	400	8	8	21	26	0	NA	NA	NA	NA
Manganese	mg/L	0.32	162	162	0.014	0.22	0	NA	NA	NA	NA
Mercury	mg/L	0.0016	377	22	9.00E-06	0.0001	0	355	1.00E-05	0.0002	0
Methyl iodide	µg/L	NA	162	0	ND	ND	NA	162	5	5	NA
Methyl tert-Butyl Ether	µg/L	12	162	0	ND	ND	NA	162	5	5	0
Methylene chloride	µg/L	9.9	162	0	ND	ND	NA	162	5	5	0
Molybdenum	mg/L	0.078	377	161	8.00E-05	0.037	0	216	1.00E-04	0.005	0
Naphthalene	µg/L	8,200	754	4	0.35	2	0	750	1	10	0
Nickel	mg/L	0.3	377	296	0.0001	0.0061	0	81	0.0002	0.005	0
Nitrobenzene	µg/L	5,000	162	0	ND	ND	NA	162	5	10	0
N-Nitrosodi-n-propylamine	µg/L	3	162	0	ND	ND	NA	162	5	10	162

Table A-4 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
N-Nitrosodiphenylamine	µg/L	10	162	0	ND	ND	NA	162	5	10	0
N-Propylbenzene	µg/L	530	377	1	0.36	0.36	0	376	1	1	0
ORO (C20-C34)	µg/L	NA	377	0	ND	ND	NA	377	100	200	NA
o-Xylene	µg/L	190	377	7	0.23	1.6	0	370	1	1	0
PCB-1016	µg/L	650	162	0	ND	ND	NA	162	0.4	1	0
PCB-1221	µg/L	0.2	162	0	ND	ND	NA	162	0.4	1	162
PCB-1232	µg/L	0.2	162	0	ND	ND	NA	162	0.4	1	162
PCB-1242	µg/L	20	162	0	ND	ND	NA	162	0.4	1	0
PCB-1248	µg/L	20	162	0	ND	ND	NA	162	0.4	1	0
PCB-1254	µg/L	20	162	0	ND	ND	NA	162	0.4	1	0
PCB-1260	µg/L	20	162	0	ND	ND	NA	162	0.4	1	0
Pentachlorophenol	µg/L	110	162	0	ND	ND	NA	162	20	40	0
Phenanthrene	µg/L	53,000	377	1	0.88	0.88	0	376	1	10	0
Phenol	µg/L	4,500	162	0	ND	ND	NA	162	5	10	0
p-Isopropyltoluene	µg/L	NA	215	0	ND	ND	NA	215	1	1	NA
Potassium	mg/L	NA	8	0	ND	ND	NA	8	10	10	NA
Pyrene	µg/L	87	162	0	ND	ND	NA	162	5	10	0
sec-Butylbenzene	µg/L	80	215	0	ND	ND	NA	215	1	1	0
Selenium	mg/L	0.078	162	22	0.0006	0.0012	0	140	0.0007	0.005	0
Silver	mg/L	0.071	162	26	1.00E-05	7.00E-05	0	136	1.00E-05	0.0002	0
Sodium	mg/L	120	8	8	16	23	0	NA	NA	NA	NA
Styrene	µg/L	1,100	162	0	ND	ND	NA	162	1	1	0
Sulfate	mg/L	250	8	8	34	38	0	NA	NA	NA	NA
Tetrachloroethene	µg/L	9.7	162	0	ND	ND	NA	162	2	2	0

Table A-4 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Thallium	mg/L	0.016	162	8	0.0001	0.0004	0	154	0.0001	0.002	0
Titanium	mg/L	NA	215	202	0.0005	0.02	NA	13	0.0007	0.001	NA
Toluene	µg/L	860	377	32	0.14	4.7	0	345	1	1	0
trans-1,2-Dichloroethene	µg/L	86	162	0	ND	ND	NA	162	1	1	0
trans-1,3-Dichloropropene	µg/L	90	162	0	ND	ND	NA	162	1	1	0
trans-1,4-Dichloro-2-butene	µg/L	0.0012	162	0	ND	ND	NA	162	5	5	162
Trichloroethene	µg/L	40	162	0	ND	ND	NA	162	1	1	0
Trichlorofluoromethane (Freon 11)	µg/L	1,100	162	0	ND	ND	NA	162	1	1	0
Vanadium	mg/L	0.078	377	263	0.0001	0.0058	0	114	0.0002	0.004	0
Vinyl acetate	µg/L	410	162	0	ND	ND	NA	162	1	1	0
Vinyl chloride	µg/L	1	162	0	ND	ND	NA	162	1	1	0
Xylenes, Total	µg/L	190	377	7	0.69	5.2	0	370	2	3	0
Zinc	mg/L	4.7	162	125	0.0013	0.036	0	37	0.0016	0.01	0

a = These chemicals were not detected (ND).

b= No values were available (NA).

Table A-5: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between January through April 2011.

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,2,3- Trimethylbenzene	µg/L	10	334	0	ND ^a	ND	NA ^b	334	1	1	0
1,2,4- Trimethylbenzene	µg/L	15	338	335	0.27	0.93	0	3	1	1	0
1,3,5- Trimethylbenzene	µg/L	87	675	673	0.12	0.24	0	2	1	1	0
2-Methylnaphthalene	µg/L	27	337	336	4.3	4.3	0	1	1	10	0
Benzene	µg/L	120	304	303	0.37	0.37	0	1	1	1	0
Beryllium	mg/L	0.016	371	337	0.00003	0.00021	0	34	0.000052	0.001	0
Cyclohexane	µg/L	13,000	336	0	ND	ND	NA	336	1	5	0
DRO (C10-C20)	µg/L	NA	337	0	ND	ND	NA	337	100	100	NA
Ethylbenzene	µg/L	1.3	338	337	0.23	0.23	0	1	1	1	0
GRO (C6-C10)	µg/L	NA	334	0	ND	ND	NA	334	200	200	NA
Iron	mg/L	11	330	330	0.17	2.8	0	0	NA	NA	NA
Isopropylbenzene (Cumene)	µg/L	390	337	0	ND	ND	NA	337	1	1	0
m- & p-Xylenes	µg/L	190	341	334	0.31	1.8	0	7	1	2	0
Mercury	mg/L	0.0016	30	27	0.000037	0.000039	0	3	0.0002	0.0002	0
Molybdenum	mg/L	0.078	981	671	0.00042	0.02	0	310	0.00083	0.001	0
Naphthalene	µg/L	6.1	72	69	0.35	1	0	3	1	5	0
Nickel	mg/L	0.3	605	337	0.00015	0.0067	0	268	0.0011	0.005	0
n-Propylbenzene	µg/L	530	336	0	ND	ND	NA	336	1	1	0
ORO (C20-C34)	µg/L	NA	70	67	48	220	NA	3	100	100	NA
o-Xylene	µg/L	190	338	337	0.38	0.38	0	1	1	1	0
Phenanthrene	µg/L	1.4	337	0	ND	ND	NA	337	1	1	0
p-Isopropyltoluene	µg/L	NA	337	0	ND	ND	NA	337	1	2	NA

Table A-5 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
sec-Butylbenzene	µg/L	80	6	0	ND	ND	NA	6	1	2	0
Titanium	mg/L	NA	656	325	0.00055	0.014	NA	331	0.001	0.001	NA
Toluene	µg/L	860	349	337	0.14	0.79	0	12	1	1	0
Vanadium	mg/L	0.078	601	331	0.00013	0.0018	0	270	0.004	0.004	0
Xylenes, total	µg/L	190	337	6	0.31	1.8	0	331	2	3	0

a = These chemicals were not detected (ND).

b= No values were available (NA).

Table A-6: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between May to Aug 2011.

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,1,1,2-Tetrachloroethane	µg/L	140	9	0	ND ^a	ND	NA ^b	9	1	1	0
1,1,1-Trichloroethane	µg/L	7,500	9	0	ND	ND	NA	9	1	1	0
1,1,2,2-Tetrachloroethane	µg/L	40	9	0	ND	ND	NA	9	1	1	0
1,1,2-Trichloroethane	µg/L	200	9	0	ND	ND	NA	9	1	1	0
1,1,2-Trichlorotrifluoroethane (Freon 113)	µg/L	53,000	9	0	ND	ND	NA	9	1	1	0
1,1-Dichloroethane	µg/L	2.4	9	0	ND	ND	NA	9	1	1	0
1,1-Dichloroethene	µg/L	260	9	0	ND	ND	NA	9	1	1	0
1,2,3-Trichlorobenzene	µg/L	5.2	9	0	ND	ND	NA	9	5	5	0
1,2,3-Trichloropropane	µg/L	0.07	9	0	ND	ND	NA	9	1	1	9
1,2,3-Trimethylbenzene	µg/L	10	528	0	ND	ND	NA	528	1	5	0
1,2,4-Trichlorobenzene	µg/L	30	9	0	ND	ND	NA	9	5	5	0

Table A-6 continued

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,2,4-Trimethylbenzene	µg/L	15	537	2	0.19	2.6	0	535	1	1	0
1,2-Dibromo-3-chloropropane	µg/L	2	9	0	ND	ND	NA	9	1	1	0
1,2-Dibromoethane	µg/L	7	9	0	ND	ND	NA	9	1	1	0
1,2-Dichlorobenzene	µg/L	280	9	0	ND	ND	NA	9	1	1	0
1,2-Dichloroethane	µg/L	170	9	0	ND	ND	NA	9	1	1	0
1,2-Dichloropropane	µg/L	260	9	0	ND	ND	NA	9	1	1	0
1,3,5-Trimethylbenzene	µg/L	87	537	1	0.12	0.12	0	536	1	1	0
1,3-Dichlorobenzene	µg/L	200	9	0	ND	ND	NA	9	1	1	0
1,4-Dichlorobenzene	µg/L	290	9	0	ND	ND	NA	9	1	1	0
2-Butanone (Methyl ethyl ketone)	µg/L	4,900	9	0	ND	ND	NA	9	25	25	0
2-Hexanone (Methyl butyl ketone)	µg/L	17,800	9	0	ND	ND	NA	9	50	50	0
2-Methylnaphthalene	µg/L	780	1074	1	3.8	3.8	0	1073	1	100	0
4-Methyl-2-pentanone	µg/L	1,000	9	0	ND	ND	NA	9	50	50	0
Acenaphthene	µg/L	400	503	0	ND	ND	NA	503	1	100	0
Acenaphthylene	µg/L	52	503	0	ND	ND	NA	503	1	100	2
Acetone	µg/L	12,000	9	0	ND	ND	NA	9	50	50	0
Acrylonitrile	µg/L	130	9	0	ND	ND	NA	9	2	2	0
Anthracene	µg/L	1,300	503	0	ND	ND	NA	503	1	100	0
Benzene	µg/L	120	537	1	1.4	1.4	0	536	1	1	0
Benzo(a)anthracene	µg/L	20	503	2	3.8	8.3	0	501	1	20	0
Benzo(a)pyrene	µg/L	2	503	2	2.1	3.4	2	501	1	20	259
Benzo(b)fluoranthene	µg/L	20	503	2	6.5	15	0	501	1	20	0
Benzo(g,h,i)perylene	µg/L	53,000	503	2	1.2	2	0	501	1	20	0
Benzo(k)fluoranthene	µg/L	170	503	3	0.29	11	0	500	1	20	0
Beryllium	mg/L	0.016	535	59	0.00003	0.00014	0	476	0.001	0.002	0
Bromochloromethane	µg/L	83	9	0	ND	ND	NA	9	1	1	0
Bromodichloromethane	µg/L	190	9	0	ND	ND	NA	9	1	1	0
Bromoform	µg/L	7.9	9	0	ND	ND	NA	9	1	1	0
Bromomethane	µg/L	7	9	0	ND	ND	NA	9	5	5	0
Carbon disulfide	µg/L	720	9	0	ND	ND	NA	9	5	5	0

Table A-6 continued

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Carbon tetrachloride	µg/L	60	9	0	ND	ND	NA	9	1	1	0
Chlorobenzene	µg/L	72	9	0	ND	ND	NA	9	1	1	0
Chloroethane	µg/L	190	9	0	ND	ND	NA	9	5	5	0
Chloroform	µg/L	320	9	1	1.3	1.3	0	8	1	1	0
Chloromethane	µg/L	190	9	0	ND	ND	NA	9	5	5	0
Chrysene	µg/L	1,700	503	2	16	30	0	501	1	20	0
cis-1,2-Dichloroethene	µg/L	28	9	0	ND	ND	NA	9	1	1	0
cis-1,3-Dichloropropene	µg/L	90	9	0	ND	ND	NA	9	1	1	0
Cyclohexane	µg/L	13,000	537	0	ND	ND	NA	537	5	5	0
Dibenz(a,h)anthracene	µg/L	2	503	0	ND	ND	NA	503	1	40	130
Dibromochloromethane	µg/L	150	9	0	ND	ND	NA	9	5	5	0
Dichlorodifluoromethane (Freon 12)	µg/L	190	9	0	ND	ND	NA	9	5	5	0
Diethyl ether	µg/L	3,100	9	0	ND	ND	NA	9	10	10	0
Di-isopropyl ether (DIPE)	µg/L	1,500	9	0	ND	ND	NA	9	5	5	0
DRO (C10-C20)	µg/L	NA	537	3	7,300	40,000	NA	534	100	1,000	NA
Ethylbenzene	µg/L	170	537	2	0.7	1.8	0	535	1	1	0
Ethyl-tert-butyl ether (ETBE)	µg/L	130,000	9	0	ND	ND	NA	9	5	5	0
Fluoranthene	µg/L	630	503	2	29	86	0	501	1	20	0
Fluorene	µg/L	220	503	0	ND	ND	NA	503	1	100	0
GRO (C6-C10)	µg/L	NA	537	0	ND	ND	NA	537	200	200	NA
Hexachloroethane	µg/L	20	9	0	ND	ND	NA	9	5	5	0
Indeno(1,2,3-c,d)pyrene	µg/L	20	503	1	2.1	2.1	0	502	1	40	2
Iron	mg/L	5,000	526	526	0.12	27	0	0	NA	NA	NA
Isopropylbenzene (Cumene)	µg/L	390	537	0	ND	ND	NA	537	1	5	0
m- & p-Xylenes	µg/L	190	537	4	0.37	2.3	0	533	2	2	0
Mercury	mg/L	0.0016	526	16	0.000016	0.000059	0	510	0.0002	0.0002	0
Methyl iodide	µg/L	NA	9	0	ND	ND	NA	9	1	1	NA
Methyl tert-butyl ether	µg/L	12	9	0	ND	ND	NA	9	5	5	0
Methylene chloride	µg/L	9.9	9	0	ND	ND	NA	9	5	5	0
Molybdenum	mg/L	0.078	535	511	0.00083	0.0041	0	24	0.001	0.05	0
Naphthalene	µg/L	8,200	1074	8	0.48	1.5	0	1066	1	100	0

Table A-6 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
n-Butylbenzene	µg/L	780	9	0	ND	ND	NA	9	1	1	0
Nickel	mg/L	0.3	535	436	0.00021	0.15	0	99	0.005	0.02	0
Nitrate	mg/L	25	74	74	0.27	1.6	0	0	NA	NA	NA
Nitrite	mg/L	1.6	74	19	0.02	0.039	0	55	0.02	0.02	0
n-Propylbenzene	µg/L	530	537	0	ND	ND	NA	537	1	1	0
ORO (C20-C34)	µg/L	NA	537	7	440	170,000	NA	530	100	100	NA
o-Xylene	µg/L	190	537	2	0.16	2.3	0	535	1	1	0
Phenanthrene	µg/L	53,000	537	2	9.6	27	0	535	1	40	0
p-Isopropyltoluene	µg/L	NA	537	1	0.41	0.41	NA	536	2	5	NA
Pyrene	µg/L	1,900	503	2	18	54	0	501	1	100	0
sec-Butylbenzene	µg/L	80	537	0	ND	ND	NA	537	1	2	0
Styrene	µg/L	1,100	9	0	ND	ND	NA	9	1	1	0
t-Amyl methyl ether (TAME)	µg/L	190	9	0	ND	ND	NA	9	5	5	0
t-Butyl alcohol	µg/L	1,500	9	0	ND	ND	NA	9	50	50	0
tert-Butylbenzene	µg/L	80	9	0	ND	ND	NA	9	2	2	0
Tetrachloroethene	µg/L	9.7	9	0	ND	ND	NA	9	1	1	0
Tetrahydrofuran	µg/L	3,200	9	0	ND	ND	NA	9	90	90	0
Titanium	mg/L	NA	526	526	0.0012	0.049	NA	0	NA	NA	NA
Toluene	µg/L	860	537	25	0.14	5.4	0	512	1	1	0
trans-1,2- Dichloroethene	µg/L	86	9	0	ND	ND	NA	9	1	1	0
trans-1,3-Dichloropropene	µg/L	90	9	0	ND	ND	NA	9	1	1	0
trans-1,4-Dichloro-2-butene	µg/L	0.0012	9	0	ND	ND	NA	9	5	5	9
Trichloroethene	µg/L	40	9	0	ND	ND	NA	9	1	1	0
Trichlorofluoromethane (Freon 11)	µg/L	1,100	9	0	ND	ND	NA	9	1	1	0
Vanadium	mg/L	0.078	535	475	0.00015	0.14	0	60	0.004	0.004	0
Vinyl chloride	µg/L	1	9	0	ND	ND	NA	9	1	1	0
Xylenes, total	µg/L	190	528	4	0.53	3.2	0	524	2	3	0

a = These chemicals were not detected (ND).

b= No values were available (NA).

Table A-7: Levels of analytes (in either microgram per liter [$\mu\text{g/L}$] or milligrams per liter [mg/L]) in surface water samples collected between September to December 2011.

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,1,1,2-Tetrachloroethane	$\mu\text{g/L}$	140	128	0	ND ^a	ND	NA ^b	128	1	1	0
1,1,1-Trichloroethane	$\mu\text{g/L}$	7,500	128	0	ND	ND	NA	128	1	1	0
1,1,2,2-Tetrachloroethane	$\mu\text{g/L}$	40	128	0	ND	ND	NA	128	1	1	0
1,1,2-Trichloroethane	$\mu\text{g/L}$	200	128	0	ND	ND	NA	128	1	1	0
1,1,2-Trichlorotrifluoroethane (Freon 113)	$\mu\text{g/L}$	53,000	8	0	ND	ND	NA	8	1	1	0
1,1-Dichloroethane	$\mu\text{g/L}$	2.4	128	0	ND	ND	NA	128	1	1	0
1,1-Dichloroethene	$\mu\text{g/L}$	260	128	0	ND	ND	NA	128	1	1	0
1,2,3-Trichlorobenzene	$\mu\text{g/L}$	5.2	128	0	ND	ND	NA	128	5	5	0
1,2,3-Trichloropropane	$\mu\text{g/L}$	0.07	128	0	ND	ND	NA	128	1	1	128
1,2,3-Trimethylbenzene	$\mu\text{g/L}$	10	120	0	ND	ND	NA	120	5	5	0
1,2,4-Trichlorobenzene	$\mu\text{g/L}$	30	128	0	ND	ND	NA	128	5	5	0
1,2,4-Trimethylbenzene	$\mu\text{g/L}$	15	128	0	ND	ND	NA	128	1	1	0
1,2-Dibromo-3-chloropropane	$\mu\text{g/L}$	2	128	0	ND	ND	NA	128	1	1	0
1,2-Dibromoethane	$\mu\text{g/L}$	7	128	0	ND	ND	NA	128	1	1	0
1,2-Dichlorobenzene	$\mu\text{g/L}$	280	128	0	ND	ND	NA	128	1	1	0
1,2-Dichloroethane	$\mu\text{g/L}$	170	128	0	ND	ND	NA	128	1	1	0
1,2-Dichloropropane	$\mu\text{g/L}$	260	128	0	ND	ND	NA	128	1	1	0
1,3-Dichlorobenzene	$\mu\text{g/L}$	200	128	0	ND	ND	NA	128	1	1	0
1,3,5-Trimethylbenzene	$\mu\text{g/L}$	87	128	0	ND	ND	NA	128	1	1	0
1,4-Dichlorobenzene	$\mu\text{g/L}$	290	128	0	ND	ND	NA	128	1	1	0
2-Butanone (Methyl ethyl ketone)	$\mu\text{g/L}$	4,900	128	0	ND	ND	NA	128	25	25	0
2-Hexanone (Methyl butyl ketone)	$\mu\text{g/L}$	17,800	128	0	ND	ND	NA	128	50	50	0
2-Methylnaphthalene	$\mu\text{g/L}$	27	136	0	ND	ND	NA	136	5	5	0
4-Methyl-2-pentanone	$\mu\text{g/L}$	1,000	128	0	ND	ND	NA	128	50	50	0
Acenaphthene	$\mu\text{g/L}$	400	128	0	ND	ND	NA	128	5	5	0

Table A-7 continued

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Acenaphthylene	µg/L	52	128	0	ND	ND	NA	128	5	5	0
Acetone	µg/L	12,000	128	0	ND	ND	NA	128	50	50	0
Acetonitrile	µg/L	130	120	0	ND	ND	NA	120	50	50	0
Acrylonitrile	µg/L	60	8	0	ND	ND	NA	8	2	2	0
Anthracene	µg/L	1,300	128	0	ND	ND	NA	128	5	5	0
Benzene	µg/L	120	128	0	ND	ND	NA	128	1	1	0
Benzo(a)anthracene	µg/L	20	128	0	ND	ND	NA	128	1	5	0
Benzo(a)pyrene	µg/L	2	128	1	1	1	0	127	1	5	1
Benzo(b)fluoranthene	µg/L	20	128	1	1.1	1.1	0	127	1	5	0
Benzo(g,h,i)perylene	µg/L	53,000	128	0	ND	ND	NA	128	1	5	0
Benzo(k)fluoranthene	µg/L	170	128	1	1.3	1.3	0	127	1	5	0
Beryllium	mg/L	0.016	128	0	ND	ND	NA	128	0.001	0.002	0
Bromobenzene	µg/L	54	120	0	ND	ND	NA	120	1	1	0
Bromochloromethane	µg/L	83	128	0	ND	ND	NA	128	1	1	0
Bromodichloromethane	µg/L	190	128	0	ND	ND	NA	128	1	1	0
Bromoform	µg/L	7.9	128	0	ND	ND	NA	128	1	1	0
Bromomethane	µg/L	7	128	0	ND	ND	NA	128	5	5	0
Carbon disulfide	µg/L	720	128	0	ND	ND	NA	128	5	5	0
Carbon tetrachloride	µg/L	60	128	0	ND	ND	NA	128	1	1	0
Chlorobenzene	µg/L	72	128	0	ND	ND	NA	128	1	1	0
Chloroethane	µg/L	190	128	0	ND	ND	NA	128	5	5	0
Chloroform	µg/L	320	128	0	ND	ND	NA	128	1	1	0
Chloromethane	µg/L	190	128	0	ND	ND	NA	128	5	5	0
Chrysene	µg/L	1,700	128	0	ND	ND	NA	128	1	5	0
cis-1,2-Dichloroethene	µg/L	28	128	0	ND	ND	NA	128	1	1	0
cis-1,3-Dichloropropene	µg/L	90	128	0	ND	ND	NA	128	1	1	0
Cyclohexane	µg/L	13,000	128	0	ND	ND	NA	128	5	5	0
Dibenz(a,h)anthracene	µg/L	2	128	0	ND	ND	NA	128	2	5	1
Dibromochloromethane	µg/L	150	128	0	ND	ND	NA	128	5	5	0
Dibromomethane	µg/L	7.9	120	0	ND	ND	NA	120	5	5	0
Dichlorodifluoromethane (Freon 12)	µg/L	190	128	0	ND	ND	NA	128	5	5	0

Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Diethyl ether	µg/L	3,100	128	0	ND	ND	NA	128	10	10	0
Di-isopropyl ether (DIPE)	µg/L	1,500	128	0	ND	ND	NA	128	5	5	0
DRO (C10-C20)	µg/L	NA	38	1	140	140	NA	37	100	100	NA
Ethylbenzene	µg/L	170	128	0	ND	ND	NA	128	1	1	0
Ethyl-tert-butyl ether (ETBE)	µg/L	130,000	128	0	ND	ND	NA	128	5	5	0
Fluoranthene	µg/L	630	128	1	1.8	1.8	0	127	1	5	0
Fluorene	µg/L	220	128	0	ND	ND	NA	128	5	5	0
GRO (C6-C10)	µg/L	NA	8	0	ND	ND	NA	8	200	200	NA
Hexachloroethane	µg/L	20	128	0	ND	ND	NA	128	5	5	0
Indeno(1,2,3-c,d)pyrene	µg/L	20	128	0	ND	ND	NA	128	2	5	0
Isopropylbenzene (Cumene)	µg/L	390	128	0	ND	ND	NA	128	5	5	0
m- & p-Xylenes	µg/L	190	128	0	ND	ND	NA	128	2	2	0
Mercury	mg/L	0.0016	30	0	ND	ND	NA	30	0.0002	0.0002	0
Methyl iodide	µg/L	NA	128	0	ND	ND	NA	128	1	1	NA
Methyl tert-butyl ether	µg/L	12	128	0	ND	ND	NA	128	5	5	0
Methylene chloride	µg/L	9.9	128	0	ND	ND	NA	128	5	5	0
Molybdenum	mg/L	0.078	128	0	ND	ND	NA	128	0.005	0.05	0
Naphthalene	µg/L	6.1	136	0	ND	ND	NA	136	5	5	0
n-Butylbenzene	µg/L	780	128	0	ND	ND	NA	128	1	1	0
Nickel	mg/L	0.3	8	0	ND	ND	NA	8	0.02	0.02	0
Nitrate	mg/L	25	13	13	0.67	1.4	0	0	NA	NA	NA
Nitrite	mg/L	1.6	13	0	ND	ND	NA	13	0.02	0.02	0
n-Propylbenzene	µg/L	530	128	0	ND	ND	NA	128	1	1	0
ORO (C20-C34)	µg/L	NA	38	5	260	710	NA	33	100	100	NA
o-Xylene	µg/L	190	128	0	ND	ND	NA	128	1	1	0
Phenanthrene	µg/L	53,000	128	0	ND	ND	NA	128	2	5	0
p-Isopropyltoluene	µg/L	NA	128	1	9.5	9.5	NA	127	5	5	NA
Pyrene	µg/L	87	128	0	ND	ND	NA	128	5	5	0
sec-Butylbenzene	µg/L	80	128	0	ND	ND	NA	128	1	1	0
Styrene	µg/L	1,100	128	0	ND	ND	NA	128	1	1	0

Table A-7 continued											
Analyte	Units	Screening level	Total number of samples	Detected				Not detected			
				Number of samples detected	Minimum level	Maximum level	Samples over screening level	Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
t-Amyl methyl ether (TAME)	µg/L	190	128	0	ND	ND	NA	128	5	5	0
t-Butyl alcohol	µg/L	1,500	128	0	ND	ND	NA	128	50	50	0
tert-Butylbenzene	µg/L	80	128	0	ND	ND	NA	128	1	1	0
Tetrachloroethene	µg/L	9.7	128	0	ND	ND	NA	128	1	1	0
Tetrahydrofuran	µg/L	3,200	128	0	ND	ND	NA	128	90	90	0
Toluene	µg/L	860	128	0	ND	ND	NA	128	1	1	0
trans-1,2- Dichloroethene	µg/L	86	128	0	ND	ND	NA	128	1	1	0
trans-1,3-Dichloropropene	µg/L	90	128	0	ND	ND	NA	128	1	1	0
trans-1,4-Dichloro-2-butene	µg/L	0.0012	128	0	ND	ND	NA	128	1	1	128
Trichloroethene	µg/L	40	128	0	ND	ND	NA	128	1	1	0
Trichlorofluoromethane (Freon 11)	µg/L	1,100	128	0	ND	ND	NA	128	1	1	0
Vanadium	mg/L	0.078	128	2	0.0048	0.0094	0	126	0.004	0.005	0
Vinyl chloride	µg/L	1	128	0	ND	ND	NA	128	1	1	0

a = These chemicals were not detected (ND).

b= No values were available (NA).

Table A-8: Levels of analytes (in either microgram per liter [µg/L] or milligrams per liter [mg/L]) in surface water samples collected between January to April 17, 2012. (No analytes were detected.)

Analyte	Units	Screening level	Not detected			
			Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
1,1,1,2-Tetrachloroethane	µg/L	140	24	1	1	0
1,1,1-Trichloroethane	µg/L	7,500	24	1	1	0
1,1,2,2-Tetrachloroethane	µg/L	40	24	1	1	0
1,1,2-Trichloroethane	µg/L	20	24	1	1	0
1,1-Dichloroethane	µg/L	2.4	24	1	1	0
1,1-Dichloroethene	µg/L	260	24	1	1	0
1,2,3-Trichlorobenzene	µg/L	5.2	24	5	5	0
1,2,3-Trichloropropane	µg/L	0.07	24	1	1	24
1,2,3-Trimethylbenzene	µg/L	10	24	5	5	0
1,2,4-Trichlorobenzene	µg/L	30	24	5	5	0
1,2,4-Trimethylbenzene	µg/L	15	24	1	1	0
1,2-Dibromo-3-chloropropane	µg/L	2	24	1	1	0
1,2-Dibromoethane	µg/L	7	24	1	1	0
1,2-Dichlorobenzene	µg/L	280	24	1	1	0
1,2-Dichloroethane	µg/L	170	24	1	1	0
1,2-Dichloropropane	µg/L	260	24	1	1	0
1,3,5-Trimethylbenzene	µg/L	87	24	1	1	0
1,3-Dichlorobenzene	µg/L	200	24	1	1	0
1,4-Dichlorobenzene	µg/L	290	24	1	1	0
2-Butanone (Methyl ethyl ketone)	µg/L	4,900	24	25	25	0
2-Hexanone (Methyl butyl ketone)	µg/L	17,800	24	50	50	0
2-Methylnaphthalene	µg/L	27	24	5	5	0
4-Methyl-2-pentanone	µg/L	1,000	24	50	50	0
Acenaphthene	µg/L	400	84	5	5	0

Table A-8 continued						
Analyte	Units	Screening level	Not detected			
			Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Acenaphthylene	µg/L	52	84	5	5	0
Acetone	µg/L	12,000	24	50	50	0
Acetonitrile	µg/L	130	24	50	50	0
Anthracene	µg/L	1,300	84	5	5	0
Benzene	µg/L	120	24	1	1	0
Benzo(a)anthracene	µg/L	20	84	1	1	0
Benzo(a)pyrene	µg/L	2	84	1	1	0
Benzo(b)fluoranthene	µg/L	20	84	1	1	0
Benzo(g,h,i)perylene	µg/L	1	84	1	1	0
Benzo(k)fluoranthene	µg/L	168	84	1	1	0
Beryllium	mg/L	0.016	83	0.001	0.001	0
Bromobenzene	µg/L	54	24	1	1	0
Bromochloromethane	µg/L	83	24	1	1	0
Bromodichloromethane	µg/L	190	24	1	1	0
Bromoform	µg/L	7.9	24	1	1	0
Bromomethane	µg/L	7	24	5	5	0
Carbon disulfide	µg/L	720	24	5	5	0
Carbon tetrachloride	µg/L	60	24	1	1	0
Chlorobenzene	µg/L	72	24	1	1	0
Chloroethane	µg/L	190	24	5	5	0
Chloroform	µg/L	320	24	1	1	0
Chloromethane	µg/L	190	24	5	5	0
Chrysene	µg/L	2.9	84	1	1	0
cis-1,2-Dichloroethene	µg/L	28	24	1	1	0
cis-1,3-Dichloropropene	µg/L	90	24	1	1	0

Table A-8 continued						
Analyte	Units	Screening level	Not detected			
			Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Cyclohexane	µg/L	13,000	24	5	5	0
Dibenz(a,h)anthracene	µg/L	2	84	2	2	0
Dibromochloromethane	µg/L	150	24	5	5	0
Dibromomethane	µg/L	7.9	24	5	5	0
Dichlorodifluoromethane (Freon 12)	µg/L	190	24	5	5	0
Diethyl ether	µg/L	3,100	24	10	10	0
Di-isopropyl ether (DIPE)	µg/L	1,500	24	5	5	0
Ethylbenzene	µg/L	1.3	24	1	1	0
Ethyl-tert-butyl ether (ETBE)	µg/L	130,000	24	5	5	0
Fluoranthene	µg/L	630	84	1	1	0
Fluorene	µg/L	220	84	5	5	0
Hexachloroethane	µg/L	20	24	5	5	0
Indeno(1,2,3-c,d)pyrene	µg/L	20	84	2	2	0
Isopropylbenzene (Cumene)	µg/L	390	24	5	5	0
m- & p-Xylenes	µg/L	190	24	2	2	0
Methyl iodide	µg/L	NA ^a	24	1	1	NA
Methyl tert-butyl ether	µg/L	12	24	5	5	0
Methylene chloride	µg/L	9.9	24	5	5	0
Molybdenum	mg/L	0.078	83	0.05	0.05	0
Naphthalene	µg/L	6.1	24	5	5	0
n-Butylbenzene	µg/L	780	24	1	1	0
n-Propylbenzene	µg/L	530	24	1	1	0
o-Xylene	µg/L	190	24	1	1	0
Phenanthrene	µg/L	53,000	84	2	2	0
p-Isopropyltoluene	µg/L	NA	24	5	5	NA

Table A-8 continued						
Analyte	Units	Screening level	Not detected			
			Number of samples not-detected	Minimum reporting limit	Maximum reporting limit	Samples over screening level
Pyrene	µg/L	87	84	5	5	0
sec-Butylbenzene	µg/L	80	24	1	1	0
Styrene	µg/L	1,100	24	1	1	0
t-Amyl methyl ether (TAME)	µg/L	190	24	5	5	0
t-Butyl alcohol	µg/L	3,900	24	50	50	0
tert-Butylbenzene	µg/L	80	24	1	1	0
Tetrachloroethene	µg/L	9.7	24	1	1	0
Tetrahydrofuran	µg/L	3,200	24	90	90	0
Toluene	µg/L	860	24	1	1	0
trans-1,2- Dichloroethene	µg/L	86	24	1	1	0
trans-1,3-Dichloropropene	µg/L	90	24	1	1	0
trans-1,4-Dichloro-2-butene	µg/L	0.0012	24	1	1	24
Trichloroethene	µg/L	40	24	1	1	0
Trichlorofluoromethane (Freon 11)	µg/L	1,100	24	1	1	0
Vanadium	mg/L	0.078	83	0.004	0.004	0
Vinyl chloride	µg/L	1	24	1	1	0

a = No value was available.

Appendix B: Levels of chemicals in fish filets

Fish consumption guidelines were previously issued for fish from the Kalamazoo River (Ceresco Impoundment) and Morrow Lake (listed as part of the Kalamazoo River – from Battle Creek to Morrow Pond Dam) due to PCBs and mercury. Guidelines will continue to be issued for these chemicals, using the fish consumption screening levels established by MDCH for the Michigan Fish Consumption Advisory Program, because of the chemical levels present in the fish filets.

Since there are no screening levels for polycyclic aromatic hydrocarbons (PAHs), nickel, and vanadium, preliminary screening levels were used to evaluate the levels present in the fish filets. Preliminary screening levels are used to evaluate chemicals without established fish consumption screening levels and to determine whether screening levels should be created. Screening level are calculated using standard meal size (0.227 kilograms [kg] per serving [one half pound]) and body weight (80 kg) for a meal frequency of 16 *MI Servings*/month. Sixteen *MI Servings*/month is the least restrictive category issued in the fish consumption guidelines.

The equation used to calculate the screening level protective for non-cancer effects is:

$$\text{non-cancer screening} = (\text{RfD} \cdot \text{BW}) / \text{CR}$$

Where:

- Reference Dose (RfD) = 0.004 mg/kg/day (for 2-methylnaphthlene, which is the lowest RfD for the non-BaP PAHs, nickel, and vanadium)
- Body weight (BW) = 80 kg
- Consumption rate (CR) = 0.12 g/day (16 *MI Servings*/month)

The equation used to calculate the screening level protective for development of cancer is:

$$\text{cancer screening level} = ([\text{RL} / \text{CSF}] \cdot \text{BW}) / \text{CR}$$

Where:

- Risk level (RL) = 0.00001
- Cancer Slope Factor (CSF) = 7.3 (mg/kg/day)⁻¹ for benzo(a)pyrene (BaP)
- Body weight (BW) = 80 kg
- Consumption rate (CR) = 0.12 g/day (16 *MI Servings*/month)

Table B-1: Preliminary screening levels (in parts per billion [ppb]) for polycyclic aromatic hydrocarbons (PAHs), nickel, and vanadium.

Chemical	Preliminary screening level in ppb
Benzo(a)pyrene and select PAHs (BaP equivalent value)	0.9
Other PAHs, nickel, and vanadium	2,700

The maximum chemicals levels in the fish filets were compared to the preliminary screening level. If the maximums were below the screening level, no further evaluation was necessary. If the maximum was above the screening level the 95% upper confidence limit of the mean (UCL)

for that chemical was calculated. If the 95% UCL was below the screening level, no further evaluation was necessary. The 95% UCL was used because waterbody-specific fish guidelines are set based on the 95% UCL for a specific chemical.

Table B-2, B-3, and B-4 presents the PAH, nickel, and vanadium levels in Marshall Pond, Ceresco Impoundment, and Morrow Lake, respectively. Benzo(a)pyrene (BaP) equivalent values are included in the tables. The BaP equivalent value is the amount of BaP and PAHs that act in the same manner in people's bodies. The other PAHs are benzo(b)fluoranthene, benzo(a)anthracene, benzo(k)fluoranthene, chrysene, dibenz(a,h)anthracene, and indeno(1,2,3-cd)pyrene. The concentration of these PAHs present in the fish filet are multiplied by a relative potency factor (EPA 1993). The BaP equivalent value is the sum of all seven values (concentration x relative potency factor).

Table B-2: Range (minimum and maximum in parts per billion [ppb]) of polycyclic aromatic hydrocarbons (PAHs), nickel, and vanadium in fish collected in 2010 and 2011 from Marshall Pond.

Chemical	Collected in 2010		Collected in 2011			
	Carp	Smallmouth Bass	Carp	Largemouth Bass	Rock Bass	Smallmouth Bass
1-Methylnaphthalene	0.31-2.23	0.33-0.53	0.48-2.05	0.5-1.83	0.87-1.64	1.08-2.63
1-Methylphenanthrene	0.2 (DL) ^a -0.54	0.2 (DL)	0.2 (DL)-2.15	0.48-1	0.26-1.36	0.27-0.67
2,6-Dimethylnaphthalene	0.57-2.01	0.65-0.98	0.2 (DL)	0.29-0.61	0.25-0.4	0.35-0.9
2-Methylnaphthalene	0.2 (DL)-2.63	0.28-0.61	0.36-1.8	0.34-1.01	0.53-1.09	0.57-1.46
Acenaphthene	0.24-14.91	0.85-2.64	0.94-7.04	0.54-1.07	0.34-0.95	0.63-2.18
Acenaphthylene	0.2 (DL)-0.73	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Anthracene	0.2 (DL)	0.26-0.66	0.2 (DL)-0.74	0.2 (DL)-0.33	0.2 (DL)-2.12	0.2 (DL)-0.54
Benzo(a)pyrene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.81	0.2 (DL)
Benzo(b)fluoranthene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.85	0.2 (DL)
Benzo(a)anthracene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)-1.68	0.2 (DL)
Benzo(e)pyrene	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.92	0.2 (DL)
Benzo(g,h,i)perylene	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.35	0.2 (DL)
Benzo(k)fluoranthene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Chrysene ^b	0.2 (DL)-0.31	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)-1.74	0.2 (DL)
Dibenz(a,h)anthracene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Fluoranthene	0.2 (DL)-4.71	0.48-1.36	0.37-4.21	0.4-0.83	0.5-8.19	0.62-1.22
Fluorene	0.26-8.27	0.76-1.89	0.48-4.19	0.47-0.85	0.33-1.19	0.68-1.59

Table B-2 continued						
Chemical	Collected in 2010		Collected in 2011			
	Carp	Smallmouth Bass	Carp	Largemouth Bass	Rock Bass	Smallmouth Bass
Indeno(1,2,3-cd)pyrene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Naphthalene	0.84-4.86	1.02-1.65	1.1-3.28	0.97-2.92	1.48-2.19	1.41-2.83
Phenanthrene	0.4-13.26	1.53-3.84	1.12-7.72	1.64-2.88	0.86-8.99	2.05-4.67
Pyrene	0.2 (DL)-2.16	0.2 (DL)-0.34	0.2 (DL)-1.86	0.2 (DL)-0.21	0.61-5.74	0.38-0.6
BaP equivalents ^c	0.46-0.46	0.46-0.46	0.46-0.46	0.46-0.46	0.20-1.26	0.46-0.46
Nickel	0.6 (DL)	0.6 (DL)	0.6 (DL)	0.6 (DL)	0.6 (DL)	0.6 (DL)
Vanadium	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)

a = DL indicated detection limit

b = PAHs included in the BaP equivalent values

c = Sum of chemical relative to benzo(a)pyrene toxicity indicated with “b” note. (BaP equivalent = sum of [concentration* relative potency]. Relative potencies were from EPA [1993].)

Only the maximum level of BaP equivalents was over the preliminary screening level for rock bass, collected in 2011, from Marshall Pond. However, the 95% UCL (0.7 ppb) was not over the screening level of 0.9 ppb. No further evaluation of Marshall Pond fish was needed.

Table B-3: Range (minimum and maximum in parts per billion [ppb]) of polycyclic aromatic hydrocarbons (PAHs), nickel, and vanadium in fish collected in 2010 and 2011 from Ceresco Impoundment.

Chemical	Collected in 2010		Collected in 2011		
	Carp	Rock Bass	Carp	Rock Bass	Smallmouth Bass
1-Methylnaphthalene	0.59-12.8	0.87-7.5	1.15-3.13	1.22-2.26	1.15-2.73
1-Methylphenanthrene	0.2 (DL) ^a -0.81	0.2 (DL)	0.2 (DL)	0.21-2.41	0.2 (DL)-1.24
2,6-Dimethylnaphthalene	0.63-1.97	0.62-1.5	NA	0.89-0.89	0.44-0.95
2-Methylnaphthalene	0.6-8.1	0.45-3.91	0.96-3.14	0.79-1.47	0.84-1.55
Acenaphthene	1-8.35	0.33-0.84	1.45-8.02	0.8-1.12	0.61-1.83
Acenaphthylene	0.2 (DL)-1.7	0.2 (DL)-0.26	NA	0.2 (DL)	0.2 (DL)
Anthracene	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.23	0.2 (DL)-0.36
Benzo(a)pyrene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.36	0.2 (DL)	0.2 (DL)
Benzo(b)fluoranthene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Benzo(a)anthracene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Benzo(e)pyrene	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.4	0.2 (DL)	0.2 (DL)
Benzo(g,h,i)perylene	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.31	0.2 (DL)	0.2 (DL)
Benzo(k)fluoranthene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Chrysene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.34	0.2 (DL)
Dibenz(a,h)anthracene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.29	0.2 (DL)	0.2 (DL)
Fluoranthene	0.3-1.49	0.2 (DL)-0.37	0.75-3.42	0.52-1.18	0.41-2.25
Fluorene	0.66-3.41	0.41-0.64	1.12-3.31	0.63-0.96	0.57-1.55
Indeno(1,2,3-cd)pyrene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Naphthalene	1.85-16.1	1.73-7.3	2.24-6.1	3.22-3.83	3.09-6.96
Phenanthrene	0.59-3.07	0.43-0.78	1.14-4.48	1.74-3.42	1.35-4.94
Pyrene	0.2 (DL)-0.81	0.2 (DL)	0.46-1.77	0.32-0.7	0.21-0.92
BaP equivalents ^c	0.46-0.46	0.46-0.46	0.46-0.67	0.46-0.46	0.46-0.46
Nickel	600 (DL)	600 (DL)	600 (DL)	600 (DL)	600 (DL)
Vanadium	200 (DL)	200 (DL)	200 (DL)	200 (DL)	200 (DL)

a = DL indicated detection limit

b = PAHs included in the BaP equivalent values

c = Sum of chemical relative to benzo(a)pyrene toxicity indicated with “b” note. (BaP equivalent = sum of [concentration_x * relative potency_x]. Relative potencies were from EPA [1993].)

None of the maximum chemical levels measured in fish filets collected from Ceresco Impoundment were over the preliminary screening levels. No further evaluation is needed.

Table B-4: Range (minimum and maximum in parts per billion [ppb]) of polycyclic aromatic hydrocarbons (PAHs), nickel, and vanadium in fish collected in 2010 and 2011 from Morrow Lake.

Chemical	Collected in 2010			Collected in 2011	
	Bluegill	Carp	Rock Bass	Bluegill	Carp
1-Methylnaphthalene	0.56-1.2	1.4-2.91	0.84-1.55	1.11-1.74	0.63-1.95
1-Methylphenanthrene	0.2 (DL)-0.37	0.2 (DL)-0.82	0.2 (DL)-0.85	0.2 (DL)	0.2 (DL)
2,6-Dimethylnaphthalene	0.53-1.3	0.2 (DL)-0.72	0.2 (DL)-0.78	0.48-0.48	0.53-0.53
2-Methylnaphthalene	0.34-0.75	0.82-2.39	0.41-0.79	0.6-0.94	0.38-1.49
Acenaphthene	0.2 (DL)	0.2 (DL)-0.21	0.2 (DL)	0.33-0.69	0.37-2.26
Acenaphthylene	0.2 (DL)	0.2 (DL)-0.23	0.2 (DL)	0.2 (DL)	0.2 (DL)
Anthracene	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Benzo(a)pyrene ^b	0.2 (DL)-0.48	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Benzo(b)fluoranthene ^b	0.2 (DL)-0.55	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Benzo(a)anthracene ^b	0.2 (DL)-0.38	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Benzo(e)pyrene	0.2 (DL)-0.56	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.25	0.2 (DL)
Benzo(g,h,i)perylene	0.2 (DL)-0.31	0.2 (DL)	0.2 (DL)	0.2 (DL)-0.4	0.2 (DL)
Benzo(k)fluoranthene ^b	0.2 (DL)-0.25	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Chrysene ^b	0.2 (DL)-0.7	0.2 (DL)-0.44	0.2 (DL)	0.2 (DL)	0.2 (DL)
Dibenz(a,h)anthracene ^b	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Fluoranthene	0.2 (DL)-1.93	0.2 (DL)-2.16	0.2 (DL)-0.4	0.76-1.3	0.58-2.84
Fluorene	0.36-0.36	0.2 (DL)-0.25	0.2 (DL)	0.47-0.7	NA
Indeno(1,2,3-cd)pyrene ^b	0.2 (DL)-0.36	0.2 (DL)	0.2 (DL)	0.2 (DL)	0.2 (DL)
Naphthalene	1.87-3.23	2.78-6.58	1.77-3.37	1.7-3	0.96-2.74
Phenanthrene	0.38-1.65	0.31-1.38	0.34-0.92	1.28-2.02	0.81-2.85
Pyrene	0.2-1.28	0.2 (DL)-0.78	0.2 (DL)	0.52-0.52	0.51-1.46
BaP equivalents ^c	0.46-0.81	0.46-0.46	0.46-0.46	0.24-0.46	0.46-0.46
Nickel	600 (DL)	600 (DL)	600 (DL)	600 (DL)	600 (DL)
Vanadium	200 (DL)	200 (DL)	200 (DL)	200 (DL)	200 (DL)

a = DL indicated detection limit

b = PAHs included in the BaP equivalent values

c = Sum of chemical relative to benzo(a)pyrene toxicity indicated with “b” note. (BaP equivalent = sum of [concentration* relative potency]. Relative potencies were from EPA [1993].)

None of the maximum chemical levels measured in fish filets collected from Morrow Lake were over the preliminary screening levels. No further evaluation is needed.

Tables B-5 and B-6 present the PCB and mercury, respectively, in fish collected from Marshall Pond, Ceresco Impoundment, and Morrow Lake. MDCH issued fish consumption guidelines for certain fish species due to the presence of PCBs and mercury in the fish filets. The fish consumption guidelines are listed below the tables.

Table B-5: Minimum, Maximum, and 95% Upper Confidence Limit of the mean (UCL) in parts per million (ppm) of polychlorinated biphenyls (PCBs), in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.

Fish	Levels of PCBs (in ppm)								
	Marshall Pond			Ceresco Impoundment			Morrow Lake		
	2010	2011	95% UCL ^a	2010	2011	95% UCL	2010	2011	95% UCL ^a
Bluegill	NS ^b	NS	NA ^c	NS	NS	NA	0.0054-0.0162	0.0019-0.0074	0.01
Carp	0.001-0.0908	0.0031-0.0826	0.04	0.0139-0.2448	0.0035-0.066	0.12	0.0519-0.3041	0.0697-0.8292	0.55
Largemouth Bass	NS	0.001-0.0126	0.01	NS	NS	NA	NS	NS	NA
Smallmouth Bass	0.0016-0.1646	0.001-0.24	0.04	NS	0.001-0.0172	0.04 ^d	NS	NS	0.24 ^d
Rock Bass	NS	0.001-0.0174	0.01	0.0012-0.0252	0.0019-0.0041	0.01	0.0021-0.237	NS	0.24 ^d

a = The 95% upper confidence limit on the mean (UCL) calculation includes all available data sets.

b = No samples were collected (NS).

c = The 95% UCL could not be calculated and was not available (NA).

d = These values include fish collected in 2006 or may only be from fish collected in 2006.

Table B-6: Minimum, Maximum, and 95% Upper Confidence Limit of the mean (UCL) in parts per million (ppm) of mercury, in fish collected in 2010 and 2011 from Marshall Pond, Ceresco Impoundment, and Morrow Lake.

Fish	Levels of mercury (in ppm)								
	Marshall Pond			Ceresco Impoundment			Morrow Lake		
	2010	2011	95% UCL ^a	2010	2011	95% UCL	2010	2011	95% UCL ^a
Bluegill	NS ^b	NS	NA ^c	NS	NS	NA	0.025-0.055	0.022-0.063	0.05
Carp	0.041-0.23	0.067-0.28	0.15	0.074-0.21	0.061-0.16	0.13	0.059-0.65	0.082-0.37	0.27
Largemouth Bass	NS	0.12-0.45	0.27	NS	NS	NA	NS	NS	NA
Smallmouth Bass	0.064-0.14	0.068-0.24	0.16	NS	0.064-0.38	1.0 ^d	NS	NS	0.59 ^d
Rock Bass	NS	0.054-0.13	0.1	0.052-0.085	0.034-0.061	0.07	0.075-0.14	NS	0.59 ^d

a = The 95% upper confidence limit on the mean (UCL) calculation includes all available data sets.

b = No samples were collected (NS).

c = The 95% UCL could not be calculated and was not available (NA).

d = These values include fish collected in 2006 or may only be from fish collected in 2006.

Recommendations for fish consumption guidelines based on PCB and mercury levels in Kalamazoo River and Morrow Lake fish are below (2014-2015 Eat Safe Fish Guide [formerly the Michigan Fish Advisory]). Fish consumption guidelines are not needed for nickel, vanadium, and PAHs.

- For the Kalamazoo River, in Marshall Pond, people should eat no more than 8 *MI Servings* per month (*MI Servings*/month) of rock bass; 4 *MI Servings*/month of smallmouth bass, largemouth bass less than 18 inches, or carp; and 2 *MI Servings*/month of largemouth bass greater than 18 inches. The typical meal size is 8 ounces for adults and 2-4 ounces for children. See the Eat Safe Fish Guide for more information (www.michigan.gov/eatsafefish).
- For the Kalamazoo River, from the Marshall Pond Dam to Ceresco Impoundment, people should eat no more than 8 *MI Servings*/month of rock bass greater than 8 inches; 4 *MI Servings*/month of smallmouth bass less than 18 inches; 2 *MI Servings*/month of smallmouth bass greater than 18 inches; and 1 meal/month of carp.
- Kalamazoo River from the Ceresco Dam to Morrow Dam, including Morrow Lake, people should eat no more than 16 *MI Servings*/month of bluegill; 8 *MI Servings*/month of rock bass; 6 *MI Servings*/year of smallmouth bass; and healthy adults should limit their consumption (to 1-2 *MI Servings* per year) of carp. See the Eat Safe Fish Guide for more information.

Check the Eat Safe Fish Guide for the most current fish guidelines (www.michigan.gov/eatsafefish).