

**Estimates for Selected Risk Factors and Health Indicators  
by Michigan Economic Development Collaborative Region**

**State of Michigan**

**Michigan Behavioral Risk Factor Surveillance System  
(Michigan BRFSS)**

Chronic Disease Epidemiology Unit  
Surveillance and Program Evaluation Section  
Lifecourse Epidemiology and Genomics Division  
Bureau of Disease Control, Prevention, and Epidemiology  
Michigan Department of Community Health

## Table of Contents

Introduction .....	3
Map of Economic Development Collaborative Regions.....	4
Health Status Tables	
Table 1: General Health Status .....	5
Table 2: Health Status on at Least 14 Days in the Past Month.....	6
Table 3: Mean Number of Days of Poor Physical Health, Poor Mental Health, and Activity Limitations.....	7
Table 4: Disability .....	8
Table 5: Weight Status .....	9
Table 6: Health Care Access.....	10
Table 7: Medicaid and Medicare Coverage.....	11
Risk Behaviors Tables	
Table 8: No Leisure-Time Physical Activity .....	12
Table 9: Aerobic Physical Activity and Muscle Strengthening .....	13
Table 10: Adequate Physical Activity .....	14
Table 11: Median Fruit and Vegetable Intake Per Day .....	15
Table 12: Adequate Fruit and Vegetable Consumption .....	16
Table 13: Cigarette Smoking .....	17
Table 14: Current Smokers Who Attempted to Quit.....	18
Table 15: Smokeless Tobacco .....	19
Table 16: Alcohol Consumption.....	20
Table 17: Mean Number of Binge Drinking Occassions During the Past Month Among Binge Drinkers .....	21
Table 18: Seatbelt Use .....	22
Table 19: Hypertension Awareness and Medication Use .....	23
Clinical Preventative Practices	
Table 20: Cholesterol Screening and Awareness .....	24
Table 21: No Routine Checkup in Past Year.....	25
Table 22: Immunizations Among Adults Aged 65 Years and Older .....	26
Table 23: HIV Testing Among Adults Aged 18 - 64 Years .....	27
Chronic Conditions	
Table 24: Asthma Among Adults .....	28
Table 25: Asthma Among Children.....	29
Table 26: Chronic Obstructive Pulmonary Disease (COPD).....	30
Table 27: Arthritis.....	31
Table 28: Cardiovascular Disease.....	32
Table 29: Cancer .....	33
Table 30: Diabetes .....	34
Table 31: Kidney Disease.....	35
Table 32: Depression .....	36
Table 33: Vision Impairment.....	37

## Introduction

The Michigan Behavioral Risk Factor Survey (BRFS) is an annual, statewide telephone survey of Michigan adults aged 18 years and older that is conducted through a collaborative effort among the Division of Behavioral Surveillance (DBS) of the Centers for Disease Control and Prevention (CDC), the Michigan State University Institute for Public Policy and Social Research (IPPSR), and the Michigan Department of Community Health. Michigan BRFS data contribute to the national [Behavioral Risk Factor Surveillance System](#) that is managed by the DBS at the CDC.

Due to increases in the utilization of cell phone communication, 2011 marks the first year in which the Michigan BRFS collected data from both land line and cell phone interviews. In 2011, the total sample size of completed interviews for the core survey was 11,049 (land line = 9,308; cell phone = 1,741). Due to variability in response rates to individual questions and differences in specific sub-populations of interest for particular questions, not all estimates are based on a total sample size of 11,049. As a result, item-by-item sample sizes and population descriptions are included within each table. Furthermore, only comparisons between estimates with non-overlapping 95% confidence intervals should be considered significantly different.

For the 2011 Michigan BRFS, the sample of land line telephone numbers that were utilized for data collection was selected using a list-assisted, random-digit-dialed methodology with a disproportionate stratification based on phone bank density, and whether or not the phone numbers were directory listed. In addition, census tracts with a high population density of African Americans were oversampled within the land line portion of the survey. The sample of cell phone numbers used within the 2011 Michigan BRFS data collection process was randomly selected from dedicated cellular telephone banks sorted on the basis of area code and exchange within the State of Michigan.

To allow for the incorporation of cell phone data into the final Michigan BRFS dataset and to improve the accuracy of prevalence estimates based on Michigan BRFS data, a new weighting methodology, known as iterative proportional fitting or raking, was introduced in 2011. Estimates based on this new methodology were weighted to adjust for the probabilities of selection and a raking adjustment factor that adjusted for the distribution of Michigan adults by telephone source (land line vs. cell phone), detailed race/ethnicity, education level, marital status, age by gender, gender by race/ethnicity, age by race/ethnicity, and renter/owner status.

In 2012, Michigan BRFS data from the 2011 survey year were analyzed by Michigan Economic Development Collaborative (MEDC) region. A map identifying each of Michigan's ten economic development collaborative regions is included on the following page. Population-based prevalence estimates and asymmetric confidence intervals were calculated for indicators of health status, health risk behaviors, clinical preventive practices, and chronic conditions among the adult population in Michigan. Respondents who refused to answer a question were removed from the denominator during the analysis of such questions. Those who responded "Don't Know/Not Sure" were also excluded from the denominator unless indicated otherwise. All analyses were performed in SAS-Callable SUDAAN®, a statistical computing that is designed for complex sample surveys.

### Important Disclaimer for 2011

**Significant changes have been made to the BRFS methodology in 2011. As a result, the estimates provided within the following tables should not be compared to Michigan BRFS estimates from previous years. For more information regarding these changes, please visit the following website: <http://www.cdc.gov/surveillancepractice/reports/brfss/brfss.html>.**

If you have any questions regarding any of the estimates within the following tables, please contact Chris Fussman at MDCH, by phone, (517) 335-8144, or by email, [MIBRFSS@michigan.gov](mailto:MIBRFSS@michigan.gov).



<b>Table 1: Health Status by MEDC Region 2011 Michigan BRFSS</b>		
	<b>General Health, Fair or Poor<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>17.2</b>	<b>(16.2-18.3)</b>
<b>MEDC Region</b>		
Upper Peninsula	16.7	(13.5-20.3)
Northwest	15.0	(11.1-20.0)
Northeast	24.3	(18.2-31.8)
West Central	15.2	(12.8-17.9)
Bay	15.7	(12.6-19.3)
Southwest	23.3	(19.2-28.0)
Central	14.8	(10.9-19.9)
East Central	20.9	(16.7-25.8)
South Central	13.2	(10.4-16.7)
Southeast	17.2	(15.5-19.1)

<sup>a</sup> Among all adults, the proportion who reported that their health, in general, was either fair or poor. (N = 11,037)

<b>Table 2: Health Status on at Least 14 Days in the Past Month by MEDC Region 2011 Michigan BRFS</b>						
	<b>Poor Physical Health<sup>a</sup></b>		<b>Poor Mental Health<sup>b</sup></b>		<b>Activity Limitation<sup>c</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>	<b>%</b>	<b>95% Confidence Interval</b>	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>13.1</b>	<b>(12.2-14.1)</b>	<b>13.1</b>	<b>(12.1-14.2)</b>	<b>9.4</b>	<b>(8.5-10.2)</b>
<b>MEDC Region</b>						
Upper Peninsula	13.4	(10.5-17.0)	11.1	(8.3-14.9)	8.9	(6.6-11.8)
Northwest	10.3	(7.2-14.5)	10.4	(6.9-15.2)	6.6	(4.2-10.2)
Northeast	16.0	(11.1-22.5)	12.9	(8.7-18.8)	12.0	(8.1-17.6)
West Central	12.4	(10.2-14.8)	13.3	(10.9-16.2)	8.5	(6.7-10.7)
Bay	12.0	(9.4-15.2)	15.7	(12.3-19.8)	10.0	(7.5-13.2)
Southwest	13.4	(10.2-17.4)	11.6	(8.8-15.2)	8.9	(6.4-12.2)
Central	17.4	(12.9-23.0)	12.6	(8.9-17.6)	10.5	(7.0-15.6)
East Central	15.7	(12.1-20.2)	15.8	(11.7-21.0)	11.7	(8.7-15.5)
South Central	10.3	(8.0-13.2)	11.3	(8.8-14.4)	9.0	(6.9-11.7)
Southeast	13.2	(11.6-15.0)	13.4	(11.6-15.4)	9.3	(7.9-11.0)

<sup>a</sup> Among all adults, the proportion who reported 14 or more days of poor physical health, which includes physical illness and injury, during the past 30 days. (N = 10,866)  
<sup>b</sup> Among all adults, the proportion who reported 14 or more days of poor mental health, which includes stress, depression, and problems with emotions, during the past 30 days. (N = 10,915)  
<sup>c</sup> Among all adults, the proportion who reported 14 or more days in the past 30 days in which either poor physical health or poor mental health kept respondents from doing their usual activities, such as self-care, work, and recreation. (N = 10,970)

**Table 3: Mean Number of Days of Poor Physical Health, Poor Mental Health, and Activity Limitations by MEDC Region  
2011 Michigan BRFS**

	Poor Physical Health <sup>a</sup>		Poor Mental Health <sup>b</sup>		Activity Limitation <sup>c</sup>	
	Mean # of days	95% Confidence Interval	Mean # of days	95% Confidence Interval	Mean # of days	95% Confidence Interval
<b>Michigan Total</b>	<b>4.1</b>	<b>(3.9-4.4)</b>	<b>4.2</b>	<b>(4.0-4.5)</b>	<b>2.9</b>	<b>(2.7-3.1)</b>
<b>MEDC Region</b>						
Upper Peninsula	4.2	(3.4-4.9)	3.6	(2.9-4.4)	2.6	(2.0-3.2)
Northwest	3.4	(2.4-4.3)	3.3	(2.3-4.3)	2.0	(1.3-2.7)
Northeast	4.8	(3.3-6.3)	4.4	(3.0-5.8)	3.7	(2.4-5.0)
West Central	3.8	(3.3-4.4)	4.1	(3.5-4.8)	2.6	(2.1-3.1)
Bay	3.8	(3.1-4.5)	4.9	(4.0-5.9)	3.1	(2.3-3.8)
Southwest	4.2	(3.3-5.1)	3.7	(2.9-4.5)	3.0	(2.2-3.8)
Central	5.4	(4.2-6.7)	4.1	(3.1-5.0)	3.4	(2.3-4.5)
East Central	4.8	(3.8-5.8)	4.7	(3.6-5.9)	3.3	(2.4-4.1)
South Central	3.6	(2.9-4.3)	3.6	(2.9-4.3)	2.6	(2.1-3.2)
Southeast	4.1	(3.7-4.6)	4.5	(4.0-5.0)	2.9	(2.5-3.3)

<sup>a</sup> Among all adults, the mean number of days during the past 30 days in which physical health, including illness and injury, was not good. (N = 10,866)

<sup>b</sup> Among all adults, the mean number of days during the past 30 days in which mental health, including stress, depression, and problems with emotions, was not good. (N = 10,915)

<sup>c</sup> Among all adults, the mean number of days during the past 30 days in which poor physical health or poor mental health kept respondent from doing their usual activities, such as self-care, work, and recreation. (N = 10,970)

**Table 4: Disability by MEDC Region  
2011 Michigan BRFS**

	Total Disability <sup>a</sup>		Any Activity Limitation <sup>b</sup>		Used Special Equipment <sup>c</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>28.2</b>	<b>(27.0-29.5)</b>	<b>26.4</b>	<b>(25.2-27.7)</b>	<b>8.4</b>	<b>(7.7-9.2)</b>
<b>MEDC Region</b>						
Upper Peninsula	30.8	(26.5-35.4)	29.4	(25.3-33.9)	8.3	(6.1-11.1)
Northwest	26.5	(21.5-32.2)	25.0	(20.1-30.7)	8.1	(5.4-12.0)
Northeast	33.7	(26.9-41.3)	32.7	(25.9-40.2)	10.1	(6.4-15.6)
West Central	27.6	(24.5-31.0)	25.9	(22.8-29.2)	7.2	(5.8-8.9)
Bay	27.0	(23.0-31.5)	24.8	(21.0-29.2)	8.0	(6.0-10.7)
Southwest	30.4	(26.1-35.2)	28.5	(24.2-33.2)	8.9	(6.5-12.0)
Central	33.8	(28.3-39.9)	30.7	(25.4-36.6)	13.4	(9.5-18.7)
East Central	30.6	(25.9-35.7)	28.4	(23.8-33.4)	9.2	(6.7-12.4)
South Central	26.9	(23.0-31.2)	25.8	(21.9-30.1)	6.6	(4.8-9.1)
Southeast	26.8	(24.6-29.0)	25.1	(23.0-27.3)	8.4	(7.2-9.8)

<sup>a</sup> Among all adults, the proportion who reported being limited in any activities because of physical, mental, or emotional problems, or reported that they required use of special equipment (such as a cane, a wheelchair, a special bed, or a special telephone) due to a health problem. (N = 10,691)

<sup>b</sup> Among all adults, the proportion who reported being limited in any activities because of physical, mental, or emotional problems. (N = 10,688)

<sup>c</sup> Among all adults, the proportion who reported that they required use of special equipment (such as a can, a wheelchair, a special bed, or a special telephone) due to a health problem. (N = 10,718)



**Table 5: Weight Status by MEDC Region  
2011 Michigan BRFSS**

	Obese <sup>a</sup>		Overweight <sup>b</sup>		Normal Weight <sup>c</sup>		Underweight <sup>d</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>31.3</b>	<b>(30.0-32.6)</b>	<b>34.2</b>	<b>(32.8-35.5)</b>	<b>33.0</b>	<b>(31.6-34.4)</b>	<b>1.6</b>	<b>(1.2-2.1)</b>
<b>MEDC Region</b>								
Upper Peninsula	33.5	(29.0-38.4)	35.5	(30.9-40.4)	29.4	(25.1-34.2)	1.5	(0.6-3.5)
Northwest	28.2	(22.9-34.2)	34.8	(28.9-41.3)	35.9	(29.9-42.4)	1.0	(0.3-3.1)
Northeast	37.2	(29.7-45.4)	32.7	(25.6-40.7)	29.3	(22.2-37.6)	0.8	(0.2-2.8)
West Central	30.4	(27.1-33.8)	33.8	(30.5-37.4)	34.4	(30.8-38.2)	1.4	(0.6-2.8)
Bay	36.9	(32.2-41.9)	34.7	(30.4-39.4)	27.4	(23.4-31.9)	0.9	(0.3-2.5)
Southwest	39.2	(34.2-44.4)	35.1	(30.3-40.2)	24.7	(20.7-29.3)	1.0	(0.4-2.5)
Central	33.4	(27.7-39.6)	29.9	(24.7-35.7)	34.7	(29.2-40.6)	2.1	(0.8-5.4)
East Central	32.9	(28.0-38.2)	36.2	(31.1-41.7)	28.4	(23.5-33.8)	2.5	(0.9-7.0)
South Central	27.2	(23.2-31.5)	36.8	(32.3-41.5)	33.9	(29.5-38.5)	2.2	(1.1-4.4)
Southeast	28.9	(26.7-31.2)	33.1	(30.8-35.5)	36.3	(33.8-38.9)	1.7	(1.1-2.6)

Note: BMI, body mass index, is defined as weight (in kilograms) divided by height (in meters) squared [weight in kg/(height in meters)<sup>2</sup>]. Weight and height were self-reported. Pregnant women were excluded. (N = 10,579)

<sup>a</sup> Among all adults, the proportion of respondents whose BMI was greater than or equal to 30.0.

<sup>b</sup> Among all adults, the proportion of respondents whose BMI was greater than or equal to 25.0, but less than 30.0.

<sup>c</sup> Among all adults, the proportion of respondents whose BMI was greater than or equal to 18.5, but less than 25.0.

<sup>d</sup> Among all adults, the proportion of respondents whose BMI was less than 18.5.

**Table 6: Health Care Access by MEDC Region  
2011 Michigan BRFSS**

	No Health Care Coverage <sup>a</sup> (Among 18 - 64 year olds)		No Personal Health Care Provider <sup>b</sup>		No Health Care Access During Past 12 Months Due to Cost <sup>c</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>18.3</b>	<b>(17.0-19.6)</b>	<b>15.5</b>	<b>(14.4-16.7)</b>	<b>16.5</b>	<b>(15.4-17.6)</b>
<b>MEDC Region</b>						
Upper Peninsula	21.0	(16.2-26.8)	10.9	(8.1-14.4)	15.7	(12.2-19.9)
Northwest	21.0	(15.1-28.4)	12.7	(9.2-17.3)	14.4	(10.3-19.7)
Northeast	14.2	(8.9-22.0)	14.6	(9.0-22.7)	18.0	(12.7-24.9)
West Central	15.8	(13.1-19.0)	13.4	(10.9-16.5)	16.6	(14.0-19.5)
Bay	21.1	(16.6-26.4)	15.0	(11.8-19.0)	18.5	(14.8-22.8)
Southwest	20.8	(16.1-26.5)	17.8	(14.0-22.5)	15.3	(12.0-19.4)
Central	12.9	(8.6-19.0)	16.0	(11.6-21.5)	13.2	(9.5-18.0)
East Central	14.7	(10.8-19.8)	15.6	(11.7-20.4)	17.9	(13.8-22.9)
South Central	15.9	(12.3-20.3)	14.8	(11.6-18.6)	14.7	(11.6-18.5)
Southeast	19.9	(17.6-22.4)	16.4	(14.4-18.6)	17.0	(15.1-19.0)

<sup>a</sup> Among adults aged 18-64 years, the proportion who reported having no health care coverage, including health insurance, prepaid plans such as HMOs, or government plans, such as Medicare. (N = 7,367)

<sup>b</sup> Among all adults, the proportion who reported that they did not have anyone that they thought of as their personal doctor or health care provider. (N = 11,007)

<sup>c</sup> Among all adults, the proportion who reported that in the past 12 months, they could not see a doctor when they needed to due to the cost. (N = 11,034)

**Table 7: Medicaid and Medicare Coverage by MEDC Region  
2011 Michigan BRFS**

	Medicaid Only <sup>a</sup>		Medicare Only <sup>b</sup>		Both Medicaid and Medicare <sup>c</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>6.8</b>	<b>(6.1-7.7)</b>	<b>20.1</b>	<b>(19.2-21.1)</b>	<b>3.3</b>	<b>(2.9-3.8)</b>
<b>MEDC Region</b>						
Upper Peninsula	4.9	(3.0-7.8)	24.5	(21.2-28.2)	4.1	(2.6-6.5)
Northwest	5.4	(2.7-10.4)	25.2	(20.6-30.5)	2.1	(1.1-4.0)
Northeast	10.4	(6.1-17.1)	25.5	(19.6-32.4)	4.5	(2.3-8.7)
West Central	7.0	(5.1-9.5)	19.5	(17.0-22.3)	2.9	(1.9-4.3)
Bay	8.1	(5.7-11.4)	21.7	(18.4-25.4)	3.1	(2.0-4.8)
Southwest	7.9	(5.5-11.0)	20.1	(16.8-23.8)	4.0	(2.5-6.1)
Central	5.9	(3.3-10.5)	17.4	(13.6-22.0)	2.3	(1.1-4.6)
East Central	7.1	(4.2-11.6)	19.5	(15.9-23.6)	3.6	(2.2-5.7)
South Central	3.6	(2.0-6.3)	20.2	(17.2-23.7)	2.6	(1.5-4.4)
Southeast	7.3	(6.0-8.9)	19.2	(17.5-20.9)	3.8	(3.0-4.7)

<sup>a</sup> Among all adults, the proportion who reported that they currently have Medicaid insurance coverage, but not Medicare. (N = 10,833)  
<sup>b</sup> Among all adults, the proportion who reported that they currently have Medicare insurance coverage, but not Medicaid. (N = 10,833)  
<sup>c</sup> Among all adults, the proportion who reported that they currently have Medicaid and Medicare insurance coverage. (N = 10,833)

<b>Table 8: No Leisure-Time Physical Activity by MEDC Region 2011 Michigan BRFS</b>		
	<b>No Leisure-Time Physical Activity<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>23.6</b>	<b>(22.4-24.8)</b>
<b>MEDC Region</b>		
Upper Peninsula	23.1	(19.2-27.6)
Northwest	26.2	(20.7-32.6)
Northeast	22.2	(16.7-28.8)
West Central	22.2	(19.4-25.2)
Bay	25.6	(21.7-29.9)
Southwest	25.1	(21.1-29.5)
Central	26.7	(21.4-32.7)
East Central	24.5	(20.1-29.6)
South Central	20.7	(17.1-24.8)
Southeast	23.1	(21.0-25.3)

<sup>a</sup> Among all adults, the proportion who reported not participating in any leisure-time physical activities or exercises such as running, calisthenics, golf, gardening, or walking during the past month. (N = 10,772)

**Table 9: Aerobic Physical Activity and Muscle Strengthening by MEDC Region  
2011 Michigan BRFSS**

	Adequate Aerobic Physical Activity (≥ 150 minutes / week) <sup>a</sup>		Adequate Muscle Strengthening (≥ 2 days / week) <sup>b</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>53.5</b>	<b>(52.1-55.0)</b>	<b>28.8</b>	<b>(27.5-30.1)</b>
<b>MEDC Region</b>				
Upper Peninsula	54.1	(49.1-59.0)	24.5	(20.4-29.0)
Northwest	57.0	(50.4-63.3)	29.2	(23.5-35.5)
Northeast	59.6	(51.6-67.1)	21.8	(15.8-29.2)
West Central	56.3	(52.5-59.9)	28.9	(25.5-32.6)
Bay	50.7	(45.9-55.6)	23.5	(19.7-27.8)
Southwest	54.1	(48.9-59.1)	24.2	(20.1-28.9)
Central	54.5	(48.3-60.6)	36.5	(30.7-42.7)
East Central	52.8	(47.1-58.4)	29.0	(24.0-34.6)
South Central	57.8	(53.0-62.5)	28.3	(24.3-32.6)
Southeast	50.8	(48.2-53.4)	30.7	(28.4-33.1)

<sup>a</sup> Among all adults, the proportion who reported that they do either moderate physical activities for at least 150 minutes per week, vigorous physical activities for at least 75 minutes per week, or an equivalent combination of moderate and vigorous physical activities. (N = 10,446)  
<sup>b</sup> Among all adults, the proportion who reported that they do muscle strengthening activities on two or more days per week. (N = 10,670)

<b>Table 10: Adequate Physical Activity by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Adequate Physical Activity<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>19.7</b>	<b>(18.6-20.9)</b>
<b>MEDC Region</b>		
Upper Peninsula	17.4	(14.0-21.5)
Northwest	21.2	(16.4-26.9)
Northeast	18.0	(12.5-25.2)
West Central	20.3	(17.3-23.5)
Bay	15.4	(12.3-19.3)
Southwest	17.3	(13.8-21.4)
Central	25.3	(20.2-31.1)
East Central	17.4	(13.6-22.1)
South Central	21.4	(17.8-25.5)
Southeast	20.5	(18.5-22.7)

<sup>a</sup> Among all adults, the proportion who reported that they do either moderate physical activities for at least 150 minutes per week, vigorous physical activities for at least 75 minutes per week, or an equivalent combination of moderate and vigorous physical activities **and** also participate in muscle strengthening activities on two or more days per week. (N = 10,375)

<b>Table 11: Median Fruit and Vegetable Intake Per Day by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Median Fruit Intake (times/day)<sup>a</sup></b>	<b>Median Vegetable Intake (times/day)<sup>b</sup></b>
	<b>%</b>	<b>%</b>
<b>Michigan Total</b>	<b>1.1</b>	<b>1.6</b>
<b>MEDC Region</b>		
Upper Peninsula	1.2	1.5
Northwest	1.1	1.7
Northeast	1.1	1.6
West Central	1.0	1.5
Bay	1.0	1.4
Southwest	1.0	1.6
Central	1.0	1.5
East Central	1.1	1.5
South Central	1.1	1.6
Southeast	1.1	1.6
<sup>a</sup> Among all adults, the median number of times per day during the past 30 days in which 100% pure fruit juices or fresh, frozen, or canned fruit were consumed. (N = 10,842) <sup>b</sup> Among all adults, the median number of times per day during the past 30 days in which cooked or canned beans, dark green vegetables, orange-colored vegetables, or other vegetables were consumed. (N = 10,822)		

<b>Table 12: Adequate Fruit and Vegetable Consumption by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Fruits and Vegetables (≥ 5 times / day)<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>17.8</b>	<b>(16.8-19.0)</b>
<b>MEDC Region</b>		
Upper Peninsula	13.3	(10.7-16.4)
Northwest	15.6	(11.7-20.4)
Northeast	20.2	(14.2-27.9)
West Central	14.9	(12.4-17.7)
Bay	15.3	(12.2-19.2)
Southwest	18.2	(14.6-22.5)
Central	19.2	(14.7-24.5)
East Central	17.1	(13.3-21.6)
South Central	20.0	(16.4-24.2)
Southeast	19.6	(17.7-21.7)

<sup>a</sup> Among all adults, the proportion whose total reported consumption of fruits (including juice) and vegetables was five or more times per day. (N = 10,813)



**Table 13: Cigarette Smoking by MEDC Region  
2011 Michigan BRFSS**

	Current Smoking <sup>a</sup>		Former Smoking <sup>b</sup>		Never Smoked	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>23.3</b>	<b>(22.0-24.6)</b>	<b>25.7</b>	<b>(24.5-26.8)</b>	<b>51.0</b>	<b>(49.6-52.4)</b>
<b>MEDC Region</b>						
Upper Peninsula	23.0	(18.7-27.9)	28.1	(24.4-32.1)	49.0	(44.1-53.8)
Northwest	21.0	(15.7-27.6)	29.9	(24.6-35.9)	49.0	(42.7-55.4)
Northeast	18.9	(13.2-26.2)	34.9	(27.7-42.9)	46.2	(38.3-54.3)
West Central	21.4	(18.3-24.7)	25.5	(22.5-28.7)	53.2	(49.5-56.8)
Bay	23.0	(19.1-27.3)	24.7	(20.9-28.9)	52.4	(47.6-57.1)
Southwest	25.7	(21.2-30.7)	26.3	(22.4-30.8)	47.9	(43.0-52.9)
Central	19.3	(14.7-24.9)	25.2	(20.6-30.5)	55.5	(49.4-61.4)
East Central	29.3	(24.1-35.1)	22.2	(18.3-26.6)	48.5	(43.0-54.0)
South Central	21.0	(17.1-25.4)	23.5	(20.0-27.3)	55.6	(50.9-60.2)
Southeast	24.0	(21.8-26.5)	26.0	(24.1-28.1)	49.9	(47.4-52.4)

<sup>a</sup> Among all adults, the proportion who reported that they had ever smoked at least 100 cigarettes (5 packs) in their life and that they smoke cigarettes now, either every day or on some days. (N = 11,002)

<sup>b</sup> Among all adults, the proportion who reported that they had ever smoked at least 100 cigarettes (5 packs) in their life, but they do not smoke cigarettes now. (N = 11,002)

<b>Table 14: Current Smokers Who Attempted to Quit by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Tried to Quit Smoking One Day or Longer in Past Year<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>62.4</b>	<b>(59.2-65.5)</b>
<b>MEDC Region</b>		
Upper Peninsula	56.3	(44.3-67.6)
Northwest	61.3	(44.7-75.6)
Northeast	82.2	(67.5-91.2)
West Central	51.5	(42.9-60.0)
Bay	66.0	(56.0-74.8)
Southwest	61.4	(50.1-71.5)
Central	62.0	(46.8-75.1)
East Central	63.7	(52.7-73.4)
South Central	61.3	(49.7-71.7)
Southeast	66.6	(61.1-71.8)

<sup>a</sup> Among current smokers, the proportion who reported that during the past 12 months, they had tried to quit smoking for one day or longer. (N = 1,996)

<b>Table 15: Smokeless Tobacco Use by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Current Smokeless Tobacco Use<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>4.4</b>	<b>(3.8-5.1)</b>
<b>MEDC Region</b>		
Upper Peninsula	5.4	(3.2-8.9)
Northwest	6.5	(3.6-11.5)
Northeast	7.4	(4.1-13.0)
West Central	4.7	(3.3-6.6)
Bay	5.0	(3.2-7.7)
Southwest	4.3	(2.3-7.8)
Central	4.3	(2.1-8.7)
East Central	5.9	(3.6-9.6)
South Central	4.8	(3.0-7.8)
Southeast	3.1	(2.2-4.3)

<sup>a</sup> Among all adults, the proportion who reported that they currently use chewing tobacco, snuff or snus, either every day or on some days. (N = 11,042)

<b>Table 16: Alcohol Consumption by MEDC Region 2011 Michigan BRFSS</b>				
	<b>Heavy Drinking<sup>a</sup></b>		<b>Binge Drinking<sup>b</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>7.0</b>	<b>(6.3-7.9)</b>	<b>19.7</b>	<b>(18.5-20.9)</b>
<b>MEDC Region</b>				
Upper Peninsula	8.2	(5.9-11.1)	24.0	(19.9-28.8)
Northwest	9.9	(6.9-14.0)	21.3	(16.5-27.1)
Northeast	6.0	(3.4-10.3)	16.3	(11.2-23.2)
West Central	8.1	(5.9-10.9)	21.2	(18.1-24.8)
Bay	9.1	(6.7-12.2)	22.2	(18.4-26.5)
Southwest	4.6	(3.0-7.0)	19.0	(15.0-23.7)
Central	6.6	(4.1-10.5)	19.1	(14.7-24.4)
East Central	6.2	(4.0-9.5)	18.2	(14.2-23.1)
South Central	7.8	(5.6-10.6)	19.0	(15.5-23.1)
Southeast	6.5	(5.3-7.9)	18.7	(16.6-20.9)

<sup>a</sup> Among all adults, the proportion who reported consuming an average of more than two alcoholic drinks per day for men or more than one per day for women in the previous month. (N = 10,584)

<sup>b</sup> Among all adults, the proportion who reported consuming five or more drinks per occasion (for men) or 4 or more drinks per occasion (for women) at least once in the previous month. (N = 10,587)

<b>Table 17: Mean Number of Binge Drinking Occasions during the Past Month among Binge Drinkers by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Number of Binge Drinking Occasions<sup>a</sup></b>	
	<b>Mean</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>4.5</b>	<b>(4.1-4.9)</b>
<b>MEDC Region</b>		
Upper Peninsula	4.3	(2.9-5.6)
Northwest	4.7	(3.2-6.2)
Northeast	5.4	(2.5-8.3)
West Central	4.1	(3.2-5.0)
Bay	4.4	(3.6-5.2)
Southwest	3.7	(2.7-4.8)
Central	4.8	(3.1-6.5)
East Central	4.9	(3.0-6.8)
South Central	5.2	(3.6-6.8)
Southeast	4.5	(3.7-5.2)

<sup>a</sup> Among adults who binge drank in the past month, the mean number of times during which five or more alcoholic beverages (for men) or four or more (for women) were consumed during the past month. (N = 1,636)

<b>Table 18: Seatbelt Use by MEDC Region 2011 Michigan BRFS</b>		
	<b>Always Uses a Seatbelt<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>88.7</b>	<b>(87.7-89.7)</b>
<b>MEDC Region</b>		
Upper Peninsula	87.4	(83.6-90.4)
Northwest	87.2	(82.2-90.9)
Northeast	91.7	(87.0-94.9)
West Central	88.8	(85.9-91.2)
Bay	89.5	(85.9-92.3)
Southwest	85.8	(81.0-89.6)
Central	88.9	(83.7-92.6)
East Central	87.2	(82.3-90.9)
South Central	91.2	(88.0-93.6)
Southeast	89.0	(87.2-90.7)

<sup>a</sup> Among all adults, the proportion who reported always using a seatbelt when driving or riding in a car. (N = 10,701)

**Table 19: Hypertension Awareness and Medication Use by MEDC Region  
2011 Michigan BRFS**

	Ever Told HBP <sup>a</sup>		Taking BP Medication <sup>b</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>34.2</b>	<b>(32.9-35.4)</b>	<b>76.5</b>	<b>(74.4-78.5)</b>
<b>MEDC Region</b>				
Upper Peninsula	32.2	(28.3-36.4)	83.1	(76.1-88.3)
Northwest	36.6	(31.0-42.5)	82.4	(74.2-88.4)
Northeast	45.5	(37.8-53.5)	74.3	(62.8-83.2)
West Central	32.0	(28.8-35.3)	76.8	(70.8-81.8)
Bay	38.5	(34.0-43.1)	76.7	(69.4-82.6)
Southwest	35.6	(31.1-40.3)	72.3	(63.5-79.7)
Central	33.4	(28.0-39.2)	77.3	(67.6-84.8)
East Central	37.3	(32.3-42.7)	71.7	(62.6-79.4)
South Central	30.7	(26.9-34.9)	75.4	(67.9-81.6)
Southeast	33.7	(31.5-36.0)	78.2	(74.6-81.4)

<sup>a</sup> Among all adults, the proportion who reported that they were ever told by a doctor that they had high blood pressure (HBP). Women who had HBP only during pregnancy and adults who were borderline hypertensive were considered to not have been diagnosed. (N = 11,014)

<sup>b</sup> Among adults who were ever told that they had HBP, the proportion that reported they were currently taking blood pressure (BP) medicines for their HBP. (N = 4,656)

**Table 20: Cholesterol Screening and Awareness by MEDC Region  
2011 Michigan BRFS**

	Cholesterol Ever Checked <sup>a</sup>		Checked Within Past Five Years <sup>b</sup>		Ever Told High Cholesterol <sup>c</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>81.1</b>	<b>(79.7-82.4)</b>	<b>76.7</b>	<b>(75.3-78.1)</b>	<b>41.8</b>	<b>(40.4-43.3)</b>
<b>MEDC Region</b>						
Upper Peninsula	85.0	(80.3-88.7)	79.8	(74.9-83.9)	43.5	(38.7-48.4)
Northwest	83.2	(77.2-87.9)	78.2	(71.9-83.5)	41.9	(35.5-48.6)
Northeast	85.0	(76.0-91.1)	81.8	(72.7-88.3)	48.9	(40.7-57.1)
West Central	80.4	(76.6-83.7)	76.1	(72.2-79.6)	39.6	(36.0-43.4)
Bay	77.2	(72.6-81.3)	72.0	(67.1-76.4)	45.1	(40.0-50.3)
Southwest	81.6	(76.6-85.8)	76.9	(71.7-81.4)	38.4	(33.5-43.5)
Central	78.2	(71.8-83.4)	74.7	(68.3-80.1)	41.6	(35.5-48.0)
East Central	79.4	(73.6-84.2)	73.9	(68.0-79.1)	42.9	(37.3-48.7)
South Central	83.3	(78.6-87.1)	78.5	(73.7-82.7)	41.1	(36.4-45.9)
Southeast	82.1	(79.7-84.3)	78.3	(75.8-80.5)	42.4	(39.8-45.0)

<sup>a</sup> Among all adults, the proportion who reported ever having had their blood cholesterol checked. (N = 10,875)

<sup>b</sup> Among all adults, the proportion who reported that they have had their blood cholesterol checked within the past five years. (N = 10,742)

<sup>c</sup> Among adults who ever had their blood cholesterol checked, the proportion who reported that a doctor, nurse, or other health professional had told them that their cholesterol was high. (N = 9,622)



<b>Table 21: No Routine Checkup in Past Year by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Had No Routine Checkup in Past Year<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>33.5</b>	<b>(32.2-34.9)</b>
<b>MEDC Region</b>		
Upper Peninsula	28.6	(24.3-33.3)
Northwest	30.1	(24.4-36.5)
Northeast	29.3	(22.5-37.1)
West Central	34.2	(30.7-37.9)
Bay	38.6	(34.0-43.5)
Southwest	37.6	(32.8-42.6)
Central	35.3	(29.5-41.6)
East Central	34.4	(29.3-40.0)
South Central	33.1	(28.7-37.8)
Southeast	31.4	(29.0-33.9)

<sup>a</sup> Among all adults, the proportion who reported that they did not have a routine checkup in the past year. (N = 10,935)

**Table 22: Immunizations among Adults Aged 65 Years and Older by MEDC Region  
2011 Michigan BRFS**

	Had Flu Vaccine in Past Year <sup>a</sup>		Ever Had Pneumonia Vaccine <sup>b</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>58.0</b>	<b>(55.5-60.4)</b>	<b>67.1</b>	<b>(64.7-69.4)</b>
<b>MEDC Region</b>				
Upper Peninsula	62.3	(55.3-68.8)	69.4	(62.3-75.7)
Northwest	70.1	(60.0-78.5)	79.3	(70.5-86.0)
Northeast	60.7	(47.4-72.6)	63.8	(49.7-75.8)
West Central	60.5	(54.0-66.7)	68.3	(61.8-74.1)
Bay	57.8	(49.1-66.0)	65.3	(56.6-73.0)
Southwest	56.9	(48.2-65.2)	69.0	(60.7-76.3)
Central	55.0	(42.2-67.2)	67.5	(53.6-78.8)
East Central	58.9	(48.8-68.3)	61.4	(50.9-70.9)
South Central	60.8	(52.5-68.4)	64.6	(56.2-72.1)
Southeast	53.5	(49.2-57.7)	66.2	(61.9-70.2)

<sup>a</sup> Among adults aged 65 years and older, the proportion who reported that they had a flu vaccine, either by an injection in the arm or sprayed in the nose during the past 12 months. (N = 3,449)

<sup>b</sup> Among adults aged 65 years and older, the proportion who reported that they ever had a pneumococcal vaccine. (N = 3,357)

<b>Table 23: HIV Testing among Adults Aged 18 - 64 Years by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Ever Had an HIV Test<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>41.3</b>	<b>(39.6-43.0)</b>
<b>MEDC Region</b>		
Upper Peninsula	34.8	(29.3-40.9)
Northwest	39.2	(31.5-47.4)
Northeast	35.0	(26.4-44.7)
West Central	37.9	(33.8-42.2)
Bay	38.9	(33.5-44.6)
Southwest	44.3	(38.4-50.4)
Central	43.6	(36.8-50.6)
East Central	39.7	(33.5-46.2)
South Central	38.6	(33.4-44.2)
Southeast	45.0	(42.1-48.0)

<sup>a</sup> Among adults aged 18 - 64 years, the proportion who reported that they ever had been tested for HIV, apart from tests that were part of a blood donation. Don't know was considered a valid response to this question. (N = 6,992)

<b>Table 24: Asthma among Adults by MEDC Region 2011 Michigan BRFSS</b>				
	<b>Lifetime Asthma Prevalence<sup>a</sup></b>		<b>Current Asthma Prevalence<sup>b</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>14.8</b>	<b>(13.8-15.9)</b>	<b>9.9</b>	<b>(9.1-10.8)</b>
<b>MEDC Region</b>				
Upper Peninsula	16.9	(13.3-21.2)	10.7	(8.2-13.9)
Northwest	16.5	(11.8-22.6)	11.3	(7.3-17.0)
Northeast	14.4	(9.9-20.5)	10.6	(6.8-16.1)
West Central	13.1	(10.8-15.8)	7.9	(6.1-10.1)
Bay	11.3	(8.8-14.4)	7.5	(5.5-10.1)
Southwest	15.1	(11.6-19.4)	11.1	(8.1-15.0)
Central	15.3	(11.5-20.0)	10.6	(7.4-15.0)
East Central	14.2	(11.0-18.1)	9.3	(6.8-12.5)
South Central	16.6	(13.1-20.7)	10.2	(7.5-13.7)
Southeast	15.6	(13.8-17.5)	10.7	(9.2-12.4)
<sup>a</sup> Among all adults, the proportion who reported that they were ever told by a doctor, nurse, or other health care professional that they had asthma. (N = 11,012) <sup>b</sup> Among all adults, the proportion who reported that they still had asthma. (N = 10,978)				

<b>Table 25: Asthma among Children by MEDC Region 2011 Michigan BRFSS</b>				
	<b>Lifetime Asthma Prevalence<sup>a</sup></b>		<b>Current Asthma Prevalence<sup>b</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>14.5</b>	<b>(12.9-16.3)</b>	<b>10.0</b>	<b>(8.6-11.6)</b>
<b>MEDC Region</b>				
Upper Peninsula	11.1	(7.5-16.3)	8.5	(5.3-13.4)
Northwest	15.9	(9.3-25.9)	11.3	(5.6-21.5)
Northeast	8.5	(3.8-17.9)	6.1	(2.2-16.3)
West Central	12.3	(9.1-16.5)	8.7	(6.0-12.6)
Bay	20.7	(14.8-28.1)	15.1	(10.2-21.8)
Southwest	11.6	(6.7-19.1)	7.3	(3.6-14.2)
Central	13.4	(7.8-21.9)	11.1	(6.1-19.2)
East Central	11.0	(6.1-19.2)	8.7	(4.4-16.6)
South Central	18.9	(12.2-28.0)	12.6	(7.1-21.4)
Southeast	15.7	(12.9-18.9)	10.0	(7.8-12.7)
<sup>a</sup> Estimated proportion of Michigan children aged 0-17 years who were ever told by a doctor, nurse or other health professional that they had asthma, using proxy information from the adult respondent. (N = 2,594). <sup>b</sup> Estimated proportion of Michigan children aged 0-17 years who still have asthma, using proxy information from the adult respondent. (N = 2,589)				

<b>Table 26: Chronic Obstructive Pulmonary Disease (COPD) by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Ever Told COPD, Emphysema or Chronic Bronchitis<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>8.0</b>	<b>(7.3-8.7)</b>
<b>MEDC Region</b>		
Upper Peninsula	9.2	(6.6-12.6)
Northwest	5.8	(3.6-9.3)
Northeast	12.3	(7.7-18.9)
West Central	6.8	(5.4-8.6)
Bay	8.7	(6.4-11.6)
Southwest	8.1	(5.9-11.0)
Central	5.4	(3.6-8.1)
East Central	9.6	(6.9-13.4)
South Central	9.1	(6.7-12.1)
Southeast	7.9	(6.8-9.3)

<sup>a</sup> Among all adults, the proportion who reported ever being told by a doctor that they had chronic obstructive pulmonary disease (COPD), emphysema or chronic bronchitis. (N = 10,987)

<b>Table 27: Arthritis by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Ever Told Arthritis<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>31.0</b>	<b>(29.8-32.2)</b>
<b>MEDC Region</b>		
Upper Peninsula	36.0	(31.7-40.5)
Northwest	31.8	(26.5-37.5)
Northeast	37.4	(30.4-45.0)
West Central	28.0	(25.0-31.1)
Bay	34.9	(30.6-39.5)
Southwest	35.3	(30.8-40.0)
Central	32.5	(27.3-38.3)
East Central	30.9	(26.4-35.9)
South Central	28.3	(24.5-32.5)
Southeast	29.9	(27.8-32.1)

<sup>a</sup> Among all adults, the proportion who reported ever being told by a doctor that they had some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia. (N = 10,974)

**Table 28: Cardiovascular Disease by MEDC Region  
2011 Michigan BRFS**

	Ever Told Heart Attack <sup>a</sup>		Had Told Angina or Coronary Heart Disease <sup>b</sup>		Ever Told Stroke <sup>c</sup>		Ever Told Any Cardiovascular Disease <sup>d</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>5.3</b>	<b>(4.8-5.9)</b>	<b>5.0</b>	<b>(4.4-5.5)</b>	<b>3.3</b>	<b>(2.9-3.8)</b>	<b>9.5</b>	<b>(8.8-10.3)</b>
<b>MEDC Region</b>								
Upper Peninsula	5.7	(4.3-7.5)	5.4	(4.0-7.4)	2.8	(2.0-4.1)	10.3	(8.2-12.7)
Northwest	6.0	(3.7-9.5)	4.4	(2.8-6.9)	3.8	(2.2-6.6)	11.5	(8.3-15.8)
Northeast	8.2	(5.2-12.6)	7.0	(4.3-11.0)	4.5	(2.3-8.5)	12.5	(8.7-17.7)
West Central	4.9	(3.7-6.5)	4.5	(3.3-6.1)	2.6	(1.7-3.9)	8.4	(6.7-10.4)
Bay	6.4	(4.5-8.9)	5.5	(3.9-7.7)	3.2	(2.0-5.0)	10.0	(7.7-12.8)
Southwest	5.4	(3.4-8.3)	6.6	(4.4-9.6)	3.1	(1.9-5.1)	11.0	(8.3-14.5)
Central	4.1	(2.2-7.5)	4.2	(2.4-7.3)	4.1	(2.3-7.4)	8.2	(5.6-12.0)
East Central	5.3	(3.4-7.9)	5.8	(3.9-8.4)	3.3	(2.0-5.3)	10.6	(8.0-13.9)
South Central	4.1	(2.8-5.9)	4.9	(3.4-7.2)	3.4	(2.3-5.0)	8.5	(6.5-11.1)
Southeast	5.5	(4.5-6.6)	4.5	(3.7-5.5)	3.4	(2.6-4.4)	9.4	(8.2-10.8)

Among all adults, the proportion who had ever been told by a doctor that...

<sup>a</sup> they had a heart attack or myocardial infarction (N = 10,997);

<sup>b</sup> they had angina or coronary heart disease (N = 10,963);

<sup>c</sup> they had a stroke. (N = 11,018)

<sup>d</sup> they had a heart attack, coronary heart disease or a stroke. (N = 10,947)



**Table 29: Cancer by MEDC Region  
2011 Michigan BRFSS**

	Ever Told Skin Cancer <sup>a</sup>		Ever Told Any Other Types of Cancer <sup>b</sup>		Ever Told Cancer <sup>c</sup>	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
<b>Michigan Total</b>	<b>5.6</b>	<b>(5.1-6.1)</b>	<b>7.3</b>	<b>(6.7-7.9)</b>	<b>11.8</b>	<b>(11.1-12.6)</b>
<b>MEDC Region</b>						
Upper Peninsula	5.1	(3.7-7.1)	7.2	(5.7-9.1)	11.2	(9.0-13.7)
Northwest	7.4	(5.3-10.2)	8.3	(5.9-11.7)	13.7	(10.5-17.6)
Northeast	9.4	(6.1-14.3)	8.7	(5.7-13.0)	16.3	(11.8-22.1)
West Central	5.2	(4.1-6.7)	7.5	(6.0-9.3)	11.8	(9.9-13.9)
Bay	7.3	(5.3-10.1)	7.0	(5.3-9.2)	13.0	(10.4-16.2)
Southwest	8.2	(6.3-10.6)	8.0	(6.0-10.5)	14.7	(12.0-17.9)
Central	4.3	(2.6-7.1)	7.8	(5.4-11.2)	10.5	(7.7-14.1)
East Central	4.6	(3.1-6.6)	8.6	(6.1-12.0)	12.7	(9.7-16.4)
South Central	5.1	(3.6-7.2)	7.5	(5.6-10.0)	11.9	(9.5-14.9)
Southeast	5.0	(4.2-5.9)	6.5	(5.5-7.6)	10.6	(9.4-12.0)

<sup>a</sup> Among all adults, the proportion who reported ever being told by a doctor that they had skin cancer. (N = 11,023)

<sup>b</sup> Among all adults, the proportion who reported ever being told by a doctor that they had a form of cancer other than skin cancer. (N = 11,020)

<sup>c</sup> Among all adults, the proportion who reported ever being told by a doctor that they had skin cancer or any other type of cancer. (N = 11,008)

<b>Table 30: Diabetes by MEDC Region 2011 Michigan BRFS</b>		
	<b>Ever Told Diabetes<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>10.0</b>	<b>(9.3-10.7)</b>
<b>MEDC Region</b>		
Upper Peninsula	10.0	(8.1-12.2)
Northwest	10.4	(7.3-14.7)
Northeast	17.2	(12.3-23.5)
West Central	9.8	(8.1-11.9)
Bay	10.2	(7.9-13.1)
Southwest	9.4	(7.2-12.1)
Central	9.0	(6.3-12.7)
East Central	10.6	(8.0-14.1)
South Central	7.7	(5.8-10.1)
Southeast	10.1	(8.9-11.5)
<sup>a</sup> Among all adults, the proportion who reported that they were ever told by a doctor that they had diabetes. Adults told they have prediabetes and women who had diabetes only during pregnancy were classified as not having been diagnosed. (N = 11,032)		

<b>Table 31: Kidney Disease by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Ever Told Kidney Disease<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>3.0</b>	<b>(2.6-3.5)</b>
<b>MEDC Region</b>		
Upper Peninsula	4.8	(2.8-8.0)
Northwest	3.4	(1.8-6.4)
Northeast	4.8	(2.4-9.3)
West Central	2.3	(1.6-3.3)
Bay	3.0	(1.9-4.8)
Southwest	2.6	(1.6-4.2)
Central	3.5	(1.9-6.4)
East Central	3.0	(1.9-5.0)
South Central	2.6	(1.7-4.0)
Southeast	3.0	(2.3-4.0)

<sup>a</sup> Among all adults, the proportion who reported ever being told by a doctor that they had kidney disease. (N = 11,003)

<b>Table 32: Depression by MEDC Region 2011 Michigan BRFSS</b>		
	<b>Ever Told Depression<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>20.6</b>	<b>(19.5-21.8)</b>
<b>MEDC Region</b>		
Upper Peninsula	22.3	(18.3-26.9)
Northwest	20.9	(16.1-26.7)
Northeast	20.7	(15.4-27.2)
West Central	23.7	(20.8-27.0)
Bay	22.3	(18.5-26.6)
Southwest	21.3	(17.5-25.7)
Central	21.8	(17.0-27.5)
East Central	19.9	(15.7-24.9)
South Central	18.9	(15.5-22.8)
Southeast	19.2	(17.2-21.4)

<sup>a</sup> Among all adults, the proportion who reported ever being told by a doctor that they had a depressive disorder including depression, major depression, dysthymia, or minor depression. (N = 10,989)

<b>Table 33: Visual Impairment by MEDC Region 2011 Michigan BRFS</b>		
	<b>Ever Told Non-Correctable Visual Impairment<sup>a</sup></b>	
	<b>%</b>	<b>95% Confidence Interval</b>
<b>Michigan Total</b>	<b>19.2</b>	<b>(18.2-20.3)</b>
<b>MEDC Region</b>		
Upper Peninsula	24.3	(20.3-28.7)
Northwest	24.5	(19.4-30.4)
Northeast	17.6	(12.8-23.7)
West Central	16.5	(14.2-19.2)
Bay	19.7	(16.2-23.7)
Southwest	20.8	(17.0-25.3)
Central	19.3	(15.4-24.1)
East Central	17.3	(13.8-21.4)
South Central	17.2	(14.2-20.7)
Southeast	19.8	(18.1-21.8)

<sup>a</sup> Among all adults, the proportion who reported ever being told by a doctor that they had a visual impairment in one or both eyes, even when wearing glasses. (N = 10,880)