

**Health Indicators and Risk Estimates by
Community Health Assessment Regions
&
Local Health Departments**

State of Michigan

Selected Tables

**Michigan Behavioral Risk Factor Surveillance System
(Michigan BRFSS)**

2017 - 2019

Lifecourse Epidemiology and Genomics Division
Bureau of Epidemiology and Population Health
Michigan Department of Health and Human Services

Table of Contents

Table of Contents	2
Introduction	3
Community Health Assessment and Improvement Regions	4
Local Health Departments.....	5
Table 1: Health Status.....	6
Table 2: Poor Physical Health on at Least 14 Days in the Past Month	8
Table 3: Poor Mental Health on at Least 14 Days in the Past Month	10
Table 4: Activity Limitation on at Least 14 Days in the Past Month	12
Table 5: Disability.....	14
Table 6: Weight Status.....	16
Table 7: No Health Care Coverage Among Those Aged 18-64 Years.....	18
Table 8: Health Care Access	20
Table 9: No Leisure Time Physical Activity	22
Table 10: Adequate Physical Activity	24
Table 11: Cigarette Smoking.....	26
Table 12: Alcohol Consumption	28
Table 13: High Blood Pressure	30
Table 14: Cholesterol Screening and Awareness.....	32
Table 15: No Routine Checkup in Past Year	34
Table 16: Immunizations Among Adults Aged 65 Years and Older	36
Table 17: HIV Testing Among Adults Aged 18-64 Years.....	38
Table 18: Asthma	40
Table 19: Chronic Obstructive Pulmonary Disease (COPD)	42
Table 20: Arthritis	44
Table 21: Heart Attack	46
Table 22: Angina or Coronary Heart Disease	48
Table 23: Stroke	50
Table 24: Cardiovascular Disease	52
Table 25: Cancer.....	54
Table 26: Diabetes	56
Table 27: Kidney Disease	58
Table 28: Depression	60

Introduction

The Michigan Behavioral Risk Factor Survey (MiBRFS) is an annual, statewide telephone survey of Michigan adults aged 18 years and older that is conducted through a collaborative effort among the Population Health Surveillance Branch (PHSB) of the Centers for Disease Control and Prevention (CDC), the Michigan State University Institute for Public Policy and Social Research (IPPSR), and the Michigan Department of Health and Human Services. Michigan BRFS data contribute to the national [Behavioral Risk Factor Surveillance System](#) that is managed by the PHSB at the CDC.

State-specific, population-based prevalence estimates (and asymmetric 95% confidence intervals) of health risk behaviors, preventive health practices, and chronic conditions are calculated using annual data sets (see Annual Tables at the [MiBRFSS website \[http://www.michigan.gov/brfs\]](#)). However, region-specific and local health department-specific prevalence rates are usually computed using BRFS data combined across years.

For this report, data from the 2017-2019 Michigan BRFS were combined. The total sample sizes by year were: 10,889 in 2017, 10,322 in 2018, and 10,518 in 2019. Not all questions were included annually in the survey. The table subtitles indicate which years were used for that particular analysis.

For the 2017 through 2019 Michigan Behavioral Risk Factor Surveys, the sample of land line telephone numbers that were utilized for data collection was selected using a list-assisted, random-digit-dialed methodology with a disproportionate stratification based on phone bank density, and whether or not the phone numbers were directory listed. The sample of cell phone numbers used within each survey year was randomly selected from dedicated cellular telephone banks sorted on the basis of area code and exchange within the State of Michigan.


Iterative proportional fitting or raking was the weighting methodology used to make the 2017-2019 Michigan BRFS data representative of the current Michigan adult population. Estimates based on this methodology were weighted to adjust for the probabilities of selection and a raking adjustment factor that adjusted for the distribution of Michigan adults by telephone source (land line vs. cell phone), detailed race/ethnicity, education level, marital status, age by gender, gender by race/ethnicity, age by race/ethnicity, and renter/owner status. No additional weighting factors were computed at the regional or local health department (LHD) level, thus the weights used might not always provide accurate regional and/or LHD estimates, particularly when regional and/or LHD demographic distributions vary greatly from that of the state.

Population-based prevalence estimates and asymmetric 95% confidence intervals were calculated for indicators of health status, health risk behaviors, clinical preventive practices, and chronic conditions among the overall adult population in Michigan, as well as at the regional and local health department level. All analyses were performed in SAS-Callable SUDAAN®, a statistical computing program that was designed for complex sample surveys. In an effort to provide the most reliable information possible, prevalence estimates are suppressed if they are based on a denominator of less than 50 respondents or have a relative standard error of greater than 30%. Comparisons between estimates with non-overlapping 95% confidence intervals should be considered significantly different.

Due to the BRFSS methodology changes that took place in 2011, the estimates provided within the following tables should not be compared to Michigan BRFSS estimates from years prior to 2011.

If you have any questions regarding any of the estimates within the following tables, please contact Yan Tian at tiany@michigan.gov.

Community Health Assessment and Improvement Regions State of Michigan



Local Health Departments
State of Michigan




Table 1: Health Status Michigan BRFSS, 2017-2019 Combined			
Geographic Area	General Health, Fair or Poor ^a		
	N	%	95% Confidence Interval
Michigan Total	31,669	18.9	(18.3-19.5)
Region 1	12,890	18.9	(18.0-19.8)
Livingston	1,540	12.6	(10.4-15.3)
Macomb	1,886	19.1	(17.0-21.4)
Monroe	363	18.4	(14.4-23.3)
Oakland	3,189	14.9	(13.4-16.6)
St. Clair	381	23.9	(19.0-29.6)
Washtenaw	1,474	13.0	(10.8-15.6)
City of Detroit	1,402	31.9	(29.0-35.0)
Wayne exc. Detroit	2,582	18.2	(16.5-20.1)
Region 2	2,157	21.4	(19.2-23.8)
Genesee	1,624	22.3	(19.7-25.2)
Lapeer	249	22.1	(16.8-28.6)
Shiawassee	264	15.8	(10.5-23.1)
Region 3	1,764	18.2	(15.9-20.9)
Jackson	1,404	18.2	(15.9-20.8)
Lenawee	296	18.6	(13.9-24.6)
Region 4	1,755	16.9	(14.8-19.3)
Barry-Eaton	454	19.5	(15.3-24.5)
Mid-Michigan ^b	523	19.0	(15.2-23.5)
Ingham	953	15.5	(12.7-18.8)
Region 5	1,488	18.8	(16.3-21.6)
Branch-Hillsdale-St. Joseph	434	22.3	(17.4-28.1)
Calhoun	369	18.8	(13.9-24.9)
Kalamazoo	662	16.6	(13.2-20.6)
Region 6	670	19.2	(15.8-23.1)
Van Buren-Cass	282	22.3	(16.9-28.7)
Berrien	333	16.5	(12.3-21.7)
Region 7	3,090	15.1	(13.6-16.7)
Allegan	350	17.9	(13.7-23.0)
Ionia	188	18.3	(12.4-26.1)
Kent	1,625	15.0	(13.0-17.2)
Mid-Michigan ^b	523	19.0	(15.2-23.5)
Ottawa	724	11.2	(8.5-14.5)

Table 1 Cont'd			
Geographic Area	General Health, Fair or Poor^a		
	N	%	95% Confidence Interval
Michigan Total	31,669	18.9	(18.3-19.5)
Region 8	2,605	23.8	(21.3-26.5)
District #10	1,085	25.1	(21.5-29.0)
Muskegon	1,416	23.2	(20.0-26.7)
Region 9	1,781	17.4	(15.1-19.9)
District #2	210	18.0	(12.4-25.3)
District #4	264	20.6	(14.5-28.5)
Northwest Michigan	545	19.9	(15.6-24.9)
Benzie-Leelanau	254	14.6	(9.9-21.1)
Grand Traverse	451	14.2	(10.5-19.0)
Region 10	1,162	21.0	(18.0-24.3)
Bay	258	22.5	(15.9-30.8)
Huron	117	17.1	(10.8-26.0)
Saginaw	471	21.6	(17.0-26.9)
Sanilac	150	21.2	(14.8-29.5)
Tuscola	153	19.3	(13.3-27.1)
Region 11	959	21.4	(18.1-25.1)
Central Michigan	548	24.5	(20.1-29.5)
Midland	231	19.4	(13.3-27.3)
Region 12	813	20.1	(17.1-23.6)
Luce-Mackinac-Alger-Schoolcraft	104	24.0	(15.4-35.3)
Western Upper Peninsula	192	15.7	(10.6-22.7)
Delta-Menominee	145	22.5	(15.9-31.0)
Chippewa	114	20.5	(13.2-30.3)
Dickinson-Iron	69	24.9	(14.3-39.9)
Marquette	155	24.0	(15.4-35.3)

^a The proportion of adults who reported that their health, in general, was either fair or poor.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

Table 2: Poor Physical Health on at Least 14 Days in the Past Month
Michigan BRFSS, 2017-2019 Combined

Geographic Area	Poor Physical Health^a		
	N	%	95% Confidence Interval
Michigan Total	31,201	14.3	(13.8-14.9)
Region 1			
Livingston	1,522	9.5	(7.6-11.7)
Macomb	1,864	14.3	(12.5-16.4)
Monroe	355	15.3	(11.6-20.0)
Oakland	3,149	12.1	(10.8-13.6)
St. Clair	378	17.9	(13.9-22.7)
Washtenaw	1,458	9.2	(7.4-11.4)
City of Detroit	1,371	19.4	(17.0-22.2)
Wayne exc. Detroit	2,548	13.6	(12.1-15.2)
Region 2			
Genesee	1,602	17.6	(15.4-20.2)
Lapeer	244	17.6	(12.7-24.0)
Shiawassee	261	16.5	(10.8-24.4)
Region 3			
Jackson	1,387	13.3	(11.4-15.6)
Lenawee	290	13.2	(9.3-18.2)
Region 4			
Barry-Eaton	444	14.0	(10.6-18.3)
Mid-Michigan ^b	513	15.6	(12.1-19.8)
Ingham	938	13.8	(11.3-16.8)
Region 5			
Branch-Hillsdale-St. Joseph	428	16.7	(12.3-22.2)
Calhoun	365	16.3	(11.7-22.3)
Kalamazoo	649	13.2	(10.3-16.6)
Region 6			
Van Buren-Cass	277	16.2	(11.7-22.1)
Berrien	327	13.5	(9.3-19.2)
Region 7			
Allegan	341	16.8	(12.4-22.3)
Ionia	185	13.2	(8.7-19.5)
Kent	1,604	13.0	(11.1-15.2)
Mid-Michigan ^b	513	15.6	(12.1-19.8)
Ottawa	718	9.8	(7.5-12.7)

Table 2 Cont'd			
Geographic Area	Poor Physical Health ^a		
	N	%	95% Confidence Interval
Michigan Total	31,201	14.3	(13.8-14.9)
Region 8	2,564	17.0	(15.0-19.2)
District #10	1,069	17.9	(15.0-21.2)
Muskegon	1,392	18.1	(15.5-21.0)
Region 9	1,765	14.5	(12.4-16.9)
District #2	206	12.2	(7.8-18.5)
District #4	263	17.0	(11.4-24.5)
Northwest Michigan	538	18.2	(14.1-23.3)
Benzie-Leelanau	255	13.9	(9.4-20.0)
Grand Traverse	446	10.6	(7.7-14.3)
Region 10	1,136	14.4	(12.2-17.0)
Bay	253	10.7	(7.3-15.5)
Huron	-- ^c	-- ^c	-- ^c
Saginaw	460	16.2	(12.6-20.6)
Sanilac	148	19.0	(12.8-27.2)
Tuscola	146	13.8	(8.7-21.1)
Region 11	944	13.8	(11.2-16.8)
Central Michigan	538	17.3	(13.6-21.7)
Midland	226	9.1	(5.7-14.2)
Region 12	802	17.3	(14.3-20.7)
Luce-Mackinac-Alger-Schoolcraft	101	17.3	(10.0-28.2)
Western Upper Peninsula	190	12.4	(7.8-19.0)
Delta-Menominee	146	20.0	(13.5-28.6)
Chippewa	113	22.8	(14.1-34.8)
Dickinson-Iron	-- ^c	-- ^c	-- ^c
Marquette	151	20.4	(13.8-29.1)

^a The proportion of adults who reported 14 or more days, out of the previous 30, on which their physical health was not good, which includes physical illness and injury.
^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.
^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 3: Poor Mental Health on at Least 14 Days in the Past Month Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Poor Mental Health^a		
	N	%	95% Confidence Interval
Michigan Total	31,268	14.7	(14.2-15.3)
Region 1			
Livingston	1,523	12.9	(10.5-15.7)
Macomb	1,868	14.0	(12.1-16.0)
Monroe	360	14.1	(10.3-19.0)
Oakland	3,148	12.8	(11.3-14.4)
St. Clair	380	14.9	(11.1-19.7)
Washtenaw	1,458	16.5	(13.9-19.5)
City of Detroit	1,380	17.5	(15.1-20.2)
Wayne exc. Detroit	2,561	15.3	(13.6-17.2)
Region 2	2,125	16.4	(14.4-18.6)
Genesee	1,598	17.1	(14.8-19.7)
Lapeer	246	14.3	(9.6-20.7)
Shiawassee	261	14.2	(9.0-21.5)
Region 3	1,736	15.5	(13.1-18.2)
Jackson	1,379	15.4	(13.1-18.1)
Lenawee	293	15.0	(10.3-21.3)
Region 4	1,729	14.2	(12.3-16.4)
Barry-Eaton	447	10.9	(7.9-14.9)
Mid-Michigan ^b	519	11.9	(8.9-15.7)
Ingham	938	17.1	(14.2-20.4)
Region 5	1,467	16.8	(14.4-19.6)
Branch-Hillsdale-St. Joseph	424	16.6	(12.0-22.6)
Calhoun	369	14.6	(10.6-19.9)
Kalamazoo	652	18.1	(14.6-22.4)
Region 6	659	17.6	(14.1-21.7)
Van Buren-Cass	275	14.5	(10.0-20.6)
Berrien	330	18.8	(13.9-24.9)
Region 7	3,053	13.2	(11.7-14.8)
Allegan	344	18.1	(13.0-24.4)
Ionia	185	19.5	(13.3-27.8)
Kent	1,609	11.5	(9.8-13.6)
Mid-Michigan ^b	519	11.9	(8.9-15.7)
Ottawa	713	13.8	(10.6-17.6)

Table 3 Cont'd			
Geographic Area	Poor Mental Health ^a		
	N	%	95% Confidence Interval
Michigan Total	31,268	14.7	(14.2-15.3)
Region 8	2,571	15.9	(13.9-18.0)
District #10	1,071	14.3	(11.8-17.3)
Muskegon	1,397	17.6	(14.7-20.8)
Region 9	1,765	12.9	(10.9-15.2)
District #2	210	14.4	(9.4-21.5)
District #4	261	10.8	(6.8-16.7)
Northwest Michigan	541	14.3	(10.4-19.3)
Benzie-Leelanau	251	10.4	(6.5-16.3)
Grand Traverse	445	13.3	(9.8-17.8)
Region 10	1,145	16.4	(13.6-19.6)
Bay	256	24.0	(17.0-32.9)
Huron	-- ^c	-- ^c	-- ^c
Saginaw	460	16.6	(12.7-21.4)
Sanilac	150	12.4	(7.6-19.8)
Tuscola	150	7.5	(4.1-13.1)
Region 11	950	15.3	(12.4-18.9)
Central Michigan	542	17.4	(13.4-22.3)
Midland	-- ^c	-- ^c	-- ^c
Region 12	802	12.5	(10.0-15.5)
Luce-Mackinac-Alger-Schoolcraft	-- ^c	-- ^c	-- ^c
Western Upper Peninsula	190	8.3	(5.1-13.3)
Delta-Menominee	145	16.9	(11.0-25.0)
Chippewa	111	16.8	(9.7-27.4)
Dickinson-Iron	-- ^c	-- ^c	-- ^c
Marquette	151	12.0	(7.0-19.7)

^a The proportion of adults who reported 14 or more days, out of the previous 30, on which their mental health was not good, which includes stress, depression, and problems with emotions.
^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.
^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 4: Activity Limitation on at Least 14 Days in the Past Month Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Activity Limitation^a		
	N	%	95% Confidence Interval
Michigan Total	31,434	10.3	(9.9-10.8)
Region 1	12,799	10.3	(9.6-11.0)
Livingston	1,531	6.4	(5.0-8.3)
Macomb	1,872	10.4	(8.8-12.3)
Monroe	356	12.2	(8.6-17.0)
Oakland	3,174	8.7	(7.5-10.0)
St. Clair	382	15.4	(11.4-20.5)
Washtenaw	1,463	7.4	(5.7-9.6)
City of Detroit	1,384	15.2	(13.0-17.7)
Wayne exc. Detroit	2,564	10.1	(8.7-11.6)
Region 2	2,139	12.2	(10.5-14.2)
Genesee	1,609	12.4	(10.5-14.7)
Lapeer	246	10.0	(6.5-15.2)
Shiawassee	264	12.6	(7.5-20.5)
Region 3	1,754	10.3	(8.4-12.6)
Jackson	1,397	9.6	(7.8-11.6)
Lenawee	293	12.7	(8.6-18.3)
Region 4	1,741	10.3	(8.6-12.3)
Barry-Eaton	449	11.9	(8.5-16.4)
Mid-Michigan ^b	521	10.8	(8.0-14.5)
Ingham	946	9.3	(7.3-11.8)
Region 5	1,476	10.1	(8.2-12.4)
Branch-Hillsdale-St. Joseph	429	13.7	(9.6-19.2)
Calhoun	365	11.6	(7.9-16.7)
Kalamazoo	659	7.1	(5.0-9.9)
Region 6	666	10.4	(7.8-13.7)
Van Buren-Cass	280	9.5	(6.1-14.6)
Berrien	331	12.5	(8.6-17.8)
Region 7	3,070	8.3	(7.1-9.6)
Allegan	345	11.2	(7.5-16.4)
Ionia	188	8.5	(4.9-14.6)
Kent	1,619	8.0	(6.5-9.7)
Mid-Michigan ^b	521	10.8	(8.0-14.5)
Ottawa	715	7.0	(4.8-10.2)

Table 4 Cont'd			
Geographic Area	Activity Limitation ^a		
	N	%	95% Confidence Interval
Michigan Total	31,434	10.3	(9.9-10.8)
Region 8	2,583	11.9	(10.2-13.7)
District #10	1,074	11.2	(9.0-13.9)
Muskegon	1,405	13.5	(11.2-16.2)
Region 9	1,773	10.0	(8.1-12.2)
District #2	207	8.9	(5.2-14.7)
District #4	265	12.5	(7.5-20.3)
Northwest Michigan	541	12.1	(8.7-16.8)
Benzie-Leelanau	254	10.2	(6.4-15.8)
Grand Traverse	449	6.6	(4.5-9.8)
Region 10	1,150	10.6	(8.7-12.9)
Bay	255	12.8	(8.4-19.0)
Huron	117	7.6	(3.9-14.3)
Saginaw	464	9.4	(6.8-12.8)
Sanilac	149	12.7	(7.6-20.5)
Tuscola	152	9.8	(5.7-16.4)
Region 11	957	10.1	(8.1-12.6)
Central Michigan	546	11.6	(8.8-15.2)
Midland	230	8.0	(4.7-13.3)
Region 12	804	12.5	(9.9-15.8)
Luce-Mackinac-Alger-Schoolcraft	-- ^c	-- ^c	-- ^c
Western Upper Peninsula	-- ^c	-- ^c	-- ^c
Delta-Menominee	145	14.6	(9.2-22.2)
Chippewa	114	17.1	(10.3-26.9)
Dickinson-Iron	-- ^c	-- ^c	-- ^c
Marquette	152	10.9	(6.3-18.2)

^a The proportion of adults who reported 14 or more days, out of the previous 30, on which either poor physical health or poor mental health kept them from doing their usual activities, such as self-care, work, and recreation.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 5: Disability Michigan BRFSS, 2017 and 2018 Combined			
Geographic Area	Disability^a		
	N	%	95% Confidence Interval
Michigan Total	20,100	27.3	(26.6-28.1)
Region 1	8,319	27.1	(25.9-28.3)
Livingston	986	20.8	(17.3-24.7)
Macomb	1,187	30.4	(27.3-33.6)
Monroe	244	28.4	(22.5-35.2)
Oakland	2,070	21.2	(19.2-23.4)
St. Clair	259	32.4	(26.2-39.3)
Washtenaw	936	22.8	(19.5-26.4)
City of Detroit	942	35.1	(31.6-38.8)
Wayne exc. Detroit	1,690	28.3	(25.9-31.0)
Region 2	1,366	29.5	(26.6-32.5)
Genesee	1,003	30.0	(26.6-33.6)
Lapeer	175	31.5	(24.3-39.7)
Shiawassee	188	25.5	(18.4-34.2)
Region 3	1,067	27.6	(24.2-31.2)
Jackson	875	28.8	(25.3-32.6)
Lenawee	192	25.5	(18.9-33.4)
Region 4	1,136	23.5	(20.6-26.7)
Barry-Eaton	311	25.0	(19.7-31.3)
Mid-Michigan ^b	338	23.1	(18.4-28.6)
Ingham	606	23.8	(19.8-28.3)
Region 5	975	29.6	(26.1-33.3)
Branch-Hillsdale-St. Joseph	286	28.8	(22.7-35.7)
Calhoun	254	31.0	(24.4-38.6)
Kalamazoo	435	29.2	(24.2-34.8)
Region 6	412	32.3	(27.2-37.9)
Van Buren-Cass	185	31.8	(24.8-39.6)
Berrien	227	32.7	(25.6-40.7)
Region 7	2,016	21.6	(19.6-23.7)
Allegan	227	29.2	(22.8-36.5)
Ionia	119	20.9	(13.7-30.5)
Kent	1,087	20.4	(17.9-23.3)
Mid-Michigan ^b	338	23.1	(18.4-28.6)
Ottawa	464	19.4	(15.4-24.0)

Table 5 Cont'd			
Geographic Area	Disability ^a		
	N	%	95% Confidence Interval
Michigan Total	20,100	27.3	(26.6-28.1)
Region 8			
District #10	1,610	32.3	(29.1-35.5)
	710	32.7	(28.4-37.4)
Muskegon	900	31.4	(27.5-35.6)
Region 9			
District #2	1,124	26.9	(23.7-30.4)
	142	31.5	(23.2-41.2)
District #4	178	27.3	(20.1-35.9)
Northwest Michigan	351	25.7	(20.6-31.6)
Benzie-Leelanau	158	26.2	(17.8-36.8)
Grand Traverse	295	25.2	(18.8-32.9)
Region 10			
Bay	751	29.2	(25.0-33.7)
Huron	173	33.8	(24.2-45.0)
Saginaw	79	25.7	(16.2-38.3)
Sanilac	292	28.1	(22.0-35.1)
Tuscola	110	28.1	(19.5-38.7)
	97	27.1	(18.5-37.7)
Region 11			
Central Michigan	525	31.5	(26.8-36.6)
Midland	367	33.3	(27.4-39.7)
	158	26.6	(19.6-34.9)
Region 12			
Luce-Mackinac-Alger-Schoolcraft	504	33.4	(28.7-38.5)
Western Upper Peninsula	67	43.1	(29.6-57.6)
Delta-Menominee	129	26.8	(17.5-38.7)
Chippewa	90	35.7	(25.7-47.2)
Dickinson-Iron	73	37.7	(25.7-51.4)
Marquette	-- ^c	-- ^c	-- ^c
	100	36.5	(26.2-48.1)

^a The proportion of adults who reported being limited in any activities because of physical, mental, or emotional problems, or reported that they required use of special equipment (such as a cane, a wheelchair, a special bed, or a special telephone) due to a health problem.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 6: Weight Status
Michigan BRFSS, 2017-2019 Combined

Geographic Area	Sample Size	Obese ^a		Overweight ^b		Healthy Weight ^c	
		N	%	95% Confidence Interval	%	95% Confidence Interval	%
Michigan Total	29,575	33.8	(33.1-34.5)	34.6	(33.9-35.3)	29.8	(29.1-30.5)
Region 1							
Livingston	1,458	30.4	(27.1-34.0)	40.4	(36.7-44.2)	28.1	(24.9-31.6)
Macomb	1,760	31.7	(29.2-34.3)	37.0	(34.3-39.7)	30.1	(27.6-32.8)
Monroe	337	38.4	(32.4-44.9)	39.1	(33.2-45.4)	21.9	(17.2-27.5)
Oakland	2,957	27.5	(25.6-29.6)	37.3	(35.2-39.5)	33.0	(31.0-35.1)
St. Clair	366	35.2	(29.7-41.2)	35.5	(29.9-41.6)	28.4	(22.7-34.8)
Washtenaw	1,414	27.9	(24.8-31.3)	32.5	(29.4-35.8)	37.7	(34.3-41.2)
City of Detroit	1,292	41.0	(37.8-44.3)	28.2	(25.3-31.3)	28.5	(25.6-31.6)
Wayne exc. Detroit	2,436	34.3	(32.0-36.6)	35.2	(32.9-37.6)	28.7	(26.6-31.0)
Region 2	1,993	37.7	(35.1-40.4)	35.3	(32.7-38.0)	25.1	(22.8-27.5)
Genesee	1,490	36.7	(33.7-39.8)	35.7	(32.7-38.9)	25.5	(22.8-28.4)
Lapeer	239	39.4	(32.6-46.8)	30.1	(23.9-37.1)	28.1	(22.2-35.0)
Shiawassee	244	42.0	(34.3-50.1)	36.9	(29.7-44.8)	20.0	(14.7-26.8)
Region 3	1,628	35.4	(32.3-38.8)	36.3	(33.0-39.6)	26.8	(23.9-29.9)
Jackson	1,301	34.4	(31.4-37.6)	35.7	(32.4-39.1)	28.8	(25.7-32.0)
Lenawee	271	38.5	(31.7-45.8)	39.2	(32.3-46.6)	20.5	(15.2-26.9)
Region 4	1,660	34.0	(31.2-36.9)	33.4	(30.6-36.4)	30.7	(28.0-33.6)
Barry-Eaton	424	33.1	(27.7-38.9)	40.9	(35.1-46.9)	25.4	(20.6-30.8)
Mid-Michigan ^d	492	44.7	(39.3-50.1)	24.7	(20.5-29.4)	27.7	(23.0-33.1)
Ingham	906	31.6	(27.9-35.6)	32.2	(28.3-36.3)	34.2	(30.3-38.3)
Region 5	1,413	34.5	(31.4-37.7)	34.1	(31.1-37.2)	29.4	(26.4-32.5)
Branch-Hillsdale-St. Joseph	410	39.1	(33.2-45.4)	34.5	(29.1-40.4)	25.2	(20.1-31.1)
Calhoun	351	36.0	(29.7-42.9)	38.8	(32.5-45.4)	23.2	(18.1-29.4)
Kalamazoo	629	29.8	(25.7-34.3)	32.0	(27.7-36.6)	35.5	(30.9-40.3)
Region 6	633	33.6	(29.2-38.3)	32.0	(27.7-36.7)	32.1	(27.9-36.6)
Van Buren-Cass	258	35.5	(28.9-42.6)	29.4	(23.4-36.2)	33.6	(27.2-40.6)
Berrien	322	32.8	(26.7-39.5)	33.7	(27.4-40.7)	30.5	(24.7-36.9)
Region 7	2,885	32.9	(30.8-35.1)	33.5	(31.5-35.7)	32.1	(30.0-34.3)
Allegan	332	35.0	(28.6-42.0)	36.6	(30.5-43.2)	26.2	(21.0-32.2)
Ionia	171	39.2	(31.0-48.0)	32.6	(24.8-41.4)	26.9	(19.8-35.4)
Kent	1,520	32.5	(29.6-35.5)	33.1	(30.3-36.0)	33.5	(30.6-36.5)
Mid-Michigan ^d	492	44.7	(39.3-50.1)	24.7	(20.5-29.4)	27.7	(23.0-33.1)
Ottawa	674	26.6	(22.7-31.0)	35.9	(31.5-40.7)	35.4	(30.6-40.5)

Table 6 Cont'd

Geographic Area	Sample Size	Obese^a		Overweight^b		Healthy Weight^c	
		N	%	95% Confidence Interval	%	95% Confidence Interval	%
Michigan Total	29,575	33.8	(33.1-34.5)	34.6	(33.9-35.3)	29.8	(29.1-30.5)
Region 8	2,458	37.8	(35.0-40.7)	33.3	(30.7-36.0)	27.3	(24.8-30.0)
District #10	1,037	38.8	(34.8-43.0)	33.9	(30.2-37.8)	25.7	(22.3-29.5)
Muskegon	1,320	36.9	(33.3-40.6)	33.8	(30.4-37.3)	28.1	(24.8-31.7)
Region 9	1,681	32.2	(29.3-35.2)	35.1	(32.2-38.1)	30.4	(27.5-33.4)
District #2	196	38.6	(30.4-47.6)	34.8	(26.7-43.8)	25.4	(18.5-33.8)
District #4	247	37.6	(30.2-45.7)	27.7	(21.5-34.8)	31.1	(23.3-40.1)
Northwest Michigan	516	30.5	(25.3-36.2)	39.4	(34.1-45.0)	26.9	(22.4-32.0)
Benzie-Leelanau	242	34.5	(27.4-42.4)	31.3	(24.8-38.6)	30.5	(23.7-38.4)
Grand Traverse	425	26.5	(21.6-32.0)	37.4	(31.9-43.3)	35.7	(29.9-42.0)
Region 10	1,086	39.2	(35.7-42.9)	32.8	(29.5-36.4)	26.1	(22.8-29.7)
Bay	242	32.2	(25.6-39.6)	34.2	(26.8-42.6)	30.7	(23.6-38.9)
Huron	108	40.3	(30.1-51.4)	38.2	(27.9-49.6)	20.7	(13.3-30.9)
Saginaw	435	43.1	(37.3-49.0)	30.5	(25.6-35.9)	25.4	(20.3-31.3)
Sanilac	139	39.4	(30.4-49.2)	34.4	(25.3-44.9)	24.0	(17.1-32.5)
Tuscola	151	38.2	(29.7-47.4)	35.1	(27.1-44.1)	23.0	(15.6-32.7)
Region 11	895	38.5	(34.4-42.7)	31.9	(28.2-35.8)	28.4	(24.6-32.5)
Central Michigan	518	40.0	(34.7-45.6)	32.5	(27.6-37.8)	26.1	(21.2-31.6)
Midland	218	40.9	(33.1-49.1)	28.6	(22.1-36.0)	28.7	(21.5-37.1)
Region 12	765	34.9	(30.9-39.0)	35.8	(31.7-40.0)	26.7	(23.1-30.7)
Luce-Mackinac-Alger-Schoolcraft	98	35.4	(24.9-47.7)	30.7	(20.7-42.8)	28.4	(18.1-41.6)
Western Upper Peninsula	183	31.9	(24.3-40.7)	37.7	(29.3-46.9)	27.5	(20.2-36.2)
Delta-Menominee	141	39.5	(30.5-49.4)	38.1	(29.3-47.9)	22.0	(15.2-30.7)
Chippewa	105	43.9	(32.6-55.8)	32.9	(22.4-45.4)	22.9	(14.7-33.8)
Dickinson-Iron	63	28.4	(17.7-42.3)	44.0	(30.0-59.1)	22.5	(12.0-38.1)
Marquette	144	33.5	(24.7-43.6)	33.9	(25.6-43.3)	29.4	(21.3-39.1)

Note: BMI, body mass index, is defined as weight (in kilograms) divided by height (in meters) squared [weight in kg/(height in meters)²]. Weight and height were self-reported. Pregnant women were excluded.

^a The proportion of adults whose BMI was greater than or equal to 30.0.

^b The proportion of adults whose BMI was greater than or equal to 25.0, but less than 30.0.

^c The proportion of adults whose BMI was greater than or equal to 18.5, but less than 25.0.

^d The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

Table 7: No Health Care Coverage Among Those Aged 18-64 Years Michigan BRFSS, 2017-2019 Combined			
Geographic Area	No Health Care Coverage^a		
	N	%	95% Confidence Interval
Michigan Total	20,331	10.0	(9.4-10.5)
Region 1			
Livingston	964	7.8	(5.6-10.7)
Macomb	1,200	9.0	(7.2-11.3)
Monroe	245	5.3	(3.0-9.3)
Oakland	2,071	5.9	(4.7-7.3)
St. Clair	260	7.9	(4.5-13.4)
Washtenaw	934	7.1	(5.3-9.5)
City of Detroit	988	14.6	(12.1-17.5)
Wayne exc. Detroit	1,709	7.5	(6.0-9.2)
Region 2			
Genesee	1,063	10.6	(8.3-13.5)
Lapeer	166	12.6	(7.4-20.5)
Shiawassee	-- ^c	-- ^c	-- ^c
Region 3			
Jackson	874	10.2	(7.7-13.4)
Lenawee	189	15.7	(9.9-24.0)
Region 4			
Barry-Eaton	272	11.8	(7.3-18.4)
Mid-Michigan ^b	326	11.6	(7.9-16.8)
Ingham	671	9.7	(7.2-13.1)
Region 5			
Branch-Hillsdale-St. Joseph	261	14.9	(10.3-21.0)
Calhoun	227	11.3	(7.0-17.7)
Kalamazoo	435	10.2	(7.3-14.1)
Region 6			
Van Buren-Cass	170	8.0	(4.7-13.5)
Berrien	205	13.4	(8.4-20.7)
Region 7			
Allegan	218	10.3	(6.3-16.4)
Ionia	129	19.2	(10.9-31.8)
Kent	1,093	9.2	(7.1-11.7)
Mid-Michigan ^b	326	11.6	(7.9-16.8)
Ottawa	476	8.9	(6.0-13.0)

Table 7 Cont'd

Geographic Area	No Health Care Coverage^a		
	N	%	95% Confidence Interval
Michigan Total	20,331	10.0	(9.4-10.5)
Region 8			
District #10	1,628	12.7	(10.4-15.4)
Muskegon	636	15.8	(12.2-20.1)
	913	10.1	(7.5-13.4)
Region 9			
District #2	1,047	10.9	(8.8-13.5)
District #4	124	15.0	(8.7-24.6)
Northwest Michigan	-- ^c	-- ^c	-- ^c
Benzie-Leelanau	310	10.9	(7.2-16.2)
Grand Traverse	127	11.3	(6.6-18.7)
	277	8.7	(5.7-13.2)
Region 10			
Bay	714	11.1	(8.5-14.4)
Huron	166	10.1	(5.7-17.4)
Saginaw	-- ^c	-- ^c	-- ^c
Sanilac	301	11.2	(7.3-16.8)
Tuscola	99	17.1	(9.2-29.7)
	-- ^c	-- ^c	-- ^c
Region 11			
Central Michigan	619	12.1	(8.7-16.7)
Midland	328	13.8	(9.1-20.5)
	-- ^c	-- ^c	-- ^c
Region 12			
Luce-Mackinac-Alger-Schoolcraft	509	12.8	(9.5-16.9)
Western Upper Peninsula	-- ^c	-- ^c	-- ^c
Delta-Menominee	115	16.9	(9.7-27.7)
Chippewa	91	14.8	(8.2-25.2)
Dickinson-Iron	-- ^c	-- ^c	-- ^c
Marquette	-- ^c	-- ^c	-- ^c

^a Among adults aged 18-64 years, the proportion who reported having no health care coverage, including health insurance, prepaid plans such as HMOs, or government plans, such as Medicare.
^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.
^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 8: Health Care Access
Michigan BRFSS, 2017-2019 Combined

Geographic Area	No Personal Health Care Provider^a			No Health Care Access During Past 12 Months Due to Cost^b		
	N	%	95% Confidence Interval	N	%	95% Confidence Interval
Michigan Total	31,539	14.8	(14.3-15.4)	31,653	11.6	(11.2-12.1)
Region 1						
Livingston	1,532	13.3	(10.7-16.5)	1,540	7.7	(6.1-9.8)
Macomb	1,878	12.9	(11.0-15.1)	1,883	13.3	(11.4-15.4)
Monroe	360	13.0	(8.9-18.7)	363	7.2	(4.8-10.5)
Oakland	3,184	12.7	(11.2-14.3)	3,188	10.5	(9.1-12.0)
St. Clair	382	5.4	(2.9-9.6)	383	9.6	(6.6-13.9)
Washtenaw	1,468	14.3	(11.9-17.1)	1,474	9.8	(7.8-12.1)
City of Detroit	1,394	19.6	(17.1-22.5)	1,401	18.5	(16.1-21.3)
Wayne exc. Detroit	2,583	13.5	(11.8-15.4)	2,586	12.2	(10.7-13.9)
Region 2						
Genesee	1,617	13.4	(11.2-15.9)	1,625	11.9	(9.8-14.2)
Lapeer	248	15.0	(10.0-21.9)	250	14.3	(9.6-20.8)
Shiawassee	263	10.5	(6.0-17.7)	264	10.5	(6.9-15.6)
Region 3						
Jackson	1,394	16.4	(13.7-19.4)	1,406	10.9	(8.9-13.2)
Lenawee	293	19.0	(13.5-25.9)	294	13.3	(8.9-19.2)
Region 4						
Barry-Eaton	452	14.8	(10.6-20.3)	453	7.0	(4.6-10.5)
Mid-Michigan ^c	520	13.6	(10.0-18.2)	523	10.2	(7.5-13.7)
Ingham	947	20.3	(17.0-24.1)	952	11.4	(8.9-14.5)
Region 5						
Branch-Hillsdale-St. Joseph	434	19.1	(14.6-24.6)	433	11.6	(7.9-16.7)
Calhoun	367	14.6	(10.6-19.9)	367	9.3	(6.2-13.8)
Kalamazoo	658	17.5	(14.1-21.7)	662	12.4	(9.4-16.1)
Region 6						
Van Buren-Cass	279	13.5	(9.2-19.2)	280	13.0	(9.0-18.6)
Berrien	330	21.3	(15.5-28.5)	332	10.2	(6.8-15.1)

Table 8 Cont'd

Geographic Area	No Personal Health Care Provider^a			No Health Care Access During Past 12 Months Due to Cost^b		
	N	%	95% Confidence Interval	N	%	95% Confidence Interval
Michigan Total	31,539	14.8	(14.3-15.4)	31,653	11.6	(11.2-12.1)
Region 7						
Allegan	3,074	11.3	(9.7-13.0)	3,082	11.5	(10.1-13.1)
Ionia	347	10.5	(6.9-15.6)	348	12.0	(8.3-17.1)
Kent	185	14.2	(7.5-25.4)	-- ^d	-- ^d	-- ^d
Mid-Michigan ^c	1,618	12.4	(10.3-14.7)	1,620	12.2	(10.3-14.4)
Ottawa	520	13.6	(10.0-18.2)	523	10.2	(7.5-13.7)
Ottawa	722	7.8	(5.0-12.2)	723	11.1	(7.9-15.3)
Region 8	2,594	14.6	(12.5-17.0)	2,602	13.8	(11.9-15.9)
District #10	1,083	15.2	(12.3-18.6)	1,084	14.5	(11.8-17.7)
Muskegon	1,409	10.8	(8.4-13.8)	1,416	14.1	(11.6-17.1)
Region 9	1,776	15.9	(13.8-18.3)	1,784	10.2	(8.4-12.3)
District #2	208	21.6	(15.1-30.0)	210	10.4	(6.2-17.0)
District #4	265	14.3	(9.8-20.3)	266	8.4	(4.7-14.5)
Northwest Michigan	545	14.8	(11.2-19.4)	545	12.2	(8.7-17.0)
Benzie-Leelanau	252	13.0	(8.4-19.6)	255	10.1	(6.5-15.5)
Grand Traverse	449	11.7	(8.6-15.9)	451	9.7	(6.7-13.8)
Region 10	1,154	16.6	(13.6-20.0)	1,160	9.9	(7.8-12.5)
Bay	257	16.4	(10.3-25.1)	259	11.7	(6.9-19.0)
Huron	116	20.9	(13.4-31.1)	-- ^d	-- ^d	-- ^d
Saginaw	466	17.9	(13.2-23.7)	469	10.2	(7.0-14.6)
Sanilac	149	11.9	(7.3-18.9)	149	11.4	(7.0-18.0)
Tuscola	153	11.2	(6.6-18.6)	-- ^d	-- ^d	-- ^d
Region 11	958	17.8	(14.4-21.7)	962	8.6	(6.7-10.9)
Central Michigan	548	15.7	(11.6-21.0)	550	8.6	(6.3-11.6)
Midland	230	11.1	(7.0-17.3)	231	8.3	(4.9-13.7)
Region 12	811	13.3	(10.7-16.4)	811	10.8	(8.4-13.7)
Luce-Mackinac-Alger-Schoolcraft	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d
Western Upper Peninsula	190	17.3	(11.6-25.2)	191	8.4	(4.8-14.5)
Delta-Menominee	146	12.6	(7.6-20.3)	144	10.8	(6.4-17.7)
Chippewa	114	11.7	(6.5-20.2)	-- ^d	-- ^d	-- ^d
Dickinson-Iron	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d
Marquette	-- ^d	-- ^d	-- ^d	155	11.0	(6.1-18.8)

^a The proportion of adults who reported that they did not have anyone that they thought of as their personal doctor or health care provider.

^b The proportion of adult s who reported that in the past 12 months there was a time when they needed to see a doctor but could not due to the cost.

^c The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^d Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 9: No Leisure Time Physical Activity Michigan BRFSS, 2017-2019 Combined			
Geographic Area	No Leisure Time Physical Activity^a		
	N	%	95% Confidence Interval
Michigan Total	30,795	25.5	(24.8-26.1)
Region 1			
Livingston	1,505	20.7	(17.9-23.8)
Macomb	1,825	27.1	(24.7-29.7)
Monroe	352	25.6	(20.5-31.5)
Oakland	3,100	20.2	(18.5-22.0)
St. Clair	378	26.8	(21.5-32.9)
Washtenaw	1,447	17.2	(14.7-20.1)
City of Detroit	1,321	35.4	(32.4-38.7)
Wayne exc. Detroit	2,518	24.8	(22.7-26.9)
Region 2	2,098	29.1	(26.7-31.6)
Genesee	1,574	30.4	(27.6-33.5)
Lapeer	246	24.2	(18.5-30.9)
Shiawassee	259	27.8	(21.6-35.1)
Region 3	1,720	27.9	(24.9-31.1)
Jackson	1,370	29.0	(25.9-32.3)
Lenawee	287	27.0	(21.3-33.7)
Region 4	1,714	24.1	(21.6-26.9)
Barry-Eaton	440	26.2	(21.3-31.9)
Mid-Michigan ^b	516	25.5	(21.2-30.3)
Ingham	931	22.9	(19.5-26.6)
Region 5	1,458	28.7	(25.8-31.8)
Branch-Hillsdale-St. Joseph	428	31.7	(26.3-37.6)
Calhoun	364	33.3	(27.1-40.0)
Kalamazoo	644	24.1	(20.2-28.5)
Region 6	650	24.4	(20.6-28.5)
Van Buren-Cass	272	28.6	(22.6-35.4)
Berrien	324	21.6	(16.9-27.3)
Region 7	3,018	21.9	(20.1-23.8)
Allegan	341	27.2	(21.9-33.2)
Ionia	185	26.3	(19.5-34.6)
Kent	1,581	20.8	(18.3-23.4)
Mid-Michigan ^b	516	25.5	(21.2-30.3)
Ottawa	710	19.2	(15.8-23.2)

Table 9 Cont'd			
Geographic Area	No Leisure Time Physical Activity^a		
	N	%	95% Confidence Interval
Michigan Total	30,795	25.5	(24.8-26.1)
Region 8			
District #10	2,541	29.1	(26.5-31.8)
Muskegon	1,068	31.4	(27.6-35.5)
	1,371	26.4	(23.5-29.5)
Region 9			
District #2	1,749	27.5	(24.7-30.5)
District #4	204	35.1	(27.4-43.5)
Northwest Michigan	256	33.5	(25.9-42.0)
Benzie-Leelanau	535	26.8	(21.7-32.5)
Grand Traverse	251	25.6	(19.0-33.6)
	446	20.5	(16.3-25.4)
Region 10			
Bay	1,123	28.7	(25.4-32.2)
Huron	252	25.0	(18.8-32.4)
Saginaw	116	28.3	(19.9-38.6)
Sanilac	449	30.4	(25.0-36.3)
Tuscola	145	30.8	(23.0-40.0)
	151	27.3	(19.9-36.1)
Region 11			
Central Michigan	929	26.5	(22.8-30.4)
Midland	535	29.0	(24.2-34.4)
	223	22.7	(16.6-30.3)
Region 12			
Luce-Mackinac-Alger-Schoolcraft	791	25.8	(22.3-29.7)
Western Upper Peninsula	100	28.3	(18.8-40.3)
Delta-Menominee	190	22.1	(16.0-29.6)
Chippewa	140	37.0	(27.8-47.3)
Dickinson-Iron	113	32.9	(22.8-44.8)
Marquette	68	27.1	(16.7-40.8)
	151	15.4	(10.1-22.7)

^a The proportion of adults who reported not participating in any leisure time physical activities or exercises such as running, calisthenics, golf, gardening, or walking during the past month.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

Table 10: Adequate Physical Activity Michigan BRFSS, 2017 and 2019 Combined			
Geographic Area	Adequate Physical Activity^a		
	N	%	95% Confidence Interval
Michigan Total	19,301	21.3	(20.5-22.0)
Region 1			
Livingston	944	22.9	(19.3-26.9)
Macomb	1,191	18.6	(16.1-21.4)
Monroe	218	16.8	(11.6-23.7)
Oakland	1,979	25.0	(22.7-27.4)
St. Clair	221	20.6	(15.0-27.7)
Washtenaw	929	21.8	(18.5-25.4)
City of Detroit	797	18.1	(15.1-21.7)
Wayne exc. Detroit	1,645	24.1	(21.6-26.9)
Region 2	1,292	19.1	(16.6-21.9)
Genesee	989	18.3	(15.5-21.5)
Lapeer	144	22.4	(15.6-31.0)
Shiawassee	147	19.8	(12.8-29.4)
Region 3	1,090	20.3	(17.1-23.8)
Jackson	879	19.4	(16.3-22.9)
Lenawee	175	22.4	(16.1-30.3)
Region 4	1,081	23.2	(20.1-26.6)
Barry-Eaton	265	23.8	(17.8-31.2)
Mid-Michigan ^b	340	21.0	(15.9-27.1)
Ingham	586	23.3	(19.2-28.0)
Region 5	898	21.2	(18.1-24.7)
Branch-Hillsdale-St. Joseph	267	17.2	(12.5-23.3)
Calhoun	211	23.4	(16.4-32.1)
Kalamazoo	411	23.3	(18.8-28.5)
Region 6	415	22.6	(17.5-28.5)
Van Buren-Cass	177	19.4	(13.5-27.0)
Berrien	205	26.0	(18.2-35.8)
Region 7	1,892	21.8	(19.5-24.2)
Allegan	215	18.3	(12.6-25.8)
Ionia	117	17.8	(11.2-27.1)
Kent	984	23.6	(20.3-27.1)
Mid-Michigan ^b	340	21.0	(15.9-27.1)
Ottawa	448	21.4	(17.0-26.6)

Table 10 Cont'd			
Geographic Area	Adequate Physical Activity ^a		
	N	%	95% Confidence Interval
Michigan Total	19,301	21.3	(20.5-22.0)
Region 8			
District #10	1,565	18.7	(16.1-21.7)
District #10	662	16.6	(13.3-20.6)
Muskegon	842	21.9	(17.5-27.1)
Region 9			
District #2	1,088	20.6	(17.6-23.8)
District #2	129	13.0	(7.8-21.0)
District #4	155	16.1	(10.4-24.3)
Northwest Michigan	323	22.6	(17.2-29.1)
Benzie-Leelanau	163	20.5	(14.3-28.5)
Grand Traverse	285	27.2	(20.4-35.1)
Region 10			
Bay	707	19.4	(16.0-23.3)
Huron	161	16.9	(10.9-25.4)
Saginaw	-- ^c	-- ^c	-- ^c
Sanilac	289	20.2	(15.0-26.7)
Tuscola	81	24.3	(15.0-37.0)
Tuscola	99	22.8	(14.6-33.9)
Region 11			
Central Michigan	558	18.8	(14.9-23.5)
Midland	331	18.5	(13.7-24.5)
Midland	150	21.3	(14.5-30.2)
Region 12			
Luce-Mackinac-Alger-Schoolcraft	494	20.8	(16.9-25.3)
Western Upper Peninsula	55	25.5	(13.6-42.7)
Delta-Menominee	113	18.3	(11.3-28.4)
Chippewa	85	17.9	(10.4-29.0)
Dickinson-Iron	75	20.5	(12.1-32.7)
Marquette	-- ^c	-- ^c	-- ^c
Marquette	97	24.5	(15.2-37.0)

^a The proportion who reported that they do either moderate physical activities for at least 150 minutes per week, vigorous physical activities for at least 75 minutes per week, or an equivalent combination of moderate and vigorous physical activities and also participate in muscle strengthening activities on two or more days per week.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 11: Cigarette Smoking
Michigan BRFSS, 2017-2019 Combined

Geographic Area	Sample Size	Current Smoking ^a		Former Smoking ^b		Never Smoked	
		N	%	95% Confidence Interval	%	95% Confidence Interval	%
Michigan Total	30,159	19.0	(18.4-19.6)	26.6	(26.0-27.2)	54.4	(53.7-55.2)
Region 1	12,460	18.0	(17.1-18.9)	25.8	(24.9-26.8)	56.2	(55.0-57.3)
Livingston	1,496	12.9	(10.6-15.4)	35.7	(32.0-39.5)	51.5	(47.7-55.2)
Macomb	1,820	20.8	(18.5-23.2)	27.3	(25.0-29.7)	51.9	(49.2-54.7)
Monroe	348	22.4	(17.4-28.4)	26.8	(21.8-32.3)	50.8	(44.6-57.0)
Oakland	3,089	12.4	(10.9-14.0)	28.1	(26.2-30.1)	59.5	(57.4-61.6)
St. Clair	371	25.9	(20.6-32.0)	30.1	(24.9-35.8)	44.0	(38.1-50.1)
Washtenaw	1,437	10.7	(8.5-13.3)	20.1	(17.7-22.7)	69.2	(66.0-72.3)
City of Detroit	1,324	27.1	(24.3-30.2)	17.8	(15.4-20.3)	55.1	(51.9-58.4)
Wayne exc. Detroit	2,508	19.0	(17.1-21.0)	26.5	(24.5-28.6)	54.5	(52.2-56.9)
Region 2	2,081	23.1	(20.7-25.7)	29.4	(27.0-31.8)	47.5	(44.8-50.2)
Genesee	1,559	24.7	(21.9-27.7)	27.8	(25.2-30.6)	47.5	(44.4-50.6)
Lapeer	247	20.7	(15.1-27.7)	34.9	(28.5-42.0)	44.4	(37.5-51.5)
Shiawassee	256	18.0	(12.1-25.9)	30.8	(24.2-38.2)	51.3	(43.4-59.1)
Region 3	1,710	21.9	(19.0-25.0)	25.2	(22.6-28.0)	52.9	(49.6-56.3)
Jackson	1,358	22.7	(20.0-25.7)	28.3	(25.4-31.4)	49.0	(45.6-52.3)
Lenawee	289	24.0	(17.9-31.4)	23.3	(18.2-29.2)	52.7	(45.7-59.7)
Region 4	1,710	16.6	(14.4-19.1)	25.0	(22.6-27.5)	58.4	(55.4-61.3)
Barry-Eaton	442	16.5	(12.7-21.1)	27.4	(22.6-32.8)	56.1	(50.3-61.7)
Mid-Michigan ^c	512	21.4	(17.0-26.5)	25.6	(21.4-30.3)	53.0	(47.7-58.3)
Ingham	927	15.6	(12.7-19.1)	24.5	(21.3-28.0)	59.8	(55.8-63.8)
Region 5	1,447	21.7	(19.0-24.8)	27.2	(24.4-30.1)	51.1	(47.9-54.4)
Branch-Hillsdale-St. Joseph	423	29.8	(24.2-36.2)	26.2	(21.6-31.4)	44.0	(38.1-50.0)
Calhoun	365	19.6	(14.6-25.7)	33.5	(27.4-40.2)	46.9	(40.4-53.5)
Kalamazoo	638	17.7	(14.2-21.9)	23.7	(19.9-27.9)	58.6	(53.8-63.2)
Region 6	646	19.8	(16.1-24.0)	26.4	(22.7-30.5)	53.8	(49.1-58.4)
Van Buren-Cass	274	19.4	(14.3-25.7)	33.6	(27.5-40.5)	46.9	(40.2-53.8)
Berrien	317	19.5	(14.5-25.8)	22.5	(17.7-28.0)	58.0	(51.2-64.6)
Region 7	3,000	16.9	(15.1-18.8)	25.6	(23.8-27.5)	57.5	(55.3-59.7)
Allegan	341	16.5	(12.2-21.9)	26.3	(21.0-32.4)	57.2	(50.6-63.6)
Ionia	182	23.0	(16.2-31.5)	21.0	(15.2-28.4)	56.0	(46.9-64.7)
Kent	1,572	16.2	(13.9-18.7)	26.1	(23.6-28.7)	57.8	(54.8-60.7)
Mid-Michigan ^c	512	21.4	(17.0-26.5)	25.6	(21.4-30.3)	53.0	(47.7-58.3)
Ottawa	707	15.0	(11.1-19.9)	24.2	(20.5-28.3)	60.8	(55.8-65.5)

Table 11 Cont'd

Geographic Area	Sample Size	Current Smoking ^a		Former Smoking ^b		Never Smoked	
		N	%	95% Confidence Interval	%	95% Confidence Interval	%
Michigan Total	30,159	19.0	(18.4-19.6)	26.6	(26.0-27.2)	54.4	(53.7-55.2)
Region 8	2,527	21.8	(19.5-24.3)	30.2	(27.7-32.7)	48.0	(45.2-50.9)
District #10	1,062	21.5	(18.4-25.0)	32.4	(28.9-36.2)	46.1	(42.0-50.2)
Muskegon	1,364	25.1	(21.7-29.0)	28.4	(25.4-31.5)	46.5	(42.9-50.2)
Region 9	1,736	21.4	(18.8-24.2)	31.1	(28.4-34.0)	47.5	(44.4-50.6)
District #2	204	27.9	(20.9-36.2)	27.6	(20.8-35.6)	44.5	(36.1-53.3)
District #4	254	26.7	(19.4-35.4)	37.1	(29.6-45.2)	36.3	(29.2-44.0)
Northwest Michigan	530	18.6	(14.6-23.4)	31.6	(26.6-37.1)	49.8	(44.3-55.3)
Benzie-Leelanau	249	12.3	(8.2-18.0)	33.4	(26.6-40.9)	54.3	(46.6-61.9)
Grand Traverse	442	21.3	(16.8-26.8)	31.4	(26.4-36.8)	47.3	(41.4-53.2)
Region 10	1,119	20.8	(17.7-24.3)	28.0	(24.9-31.3)	51.2	(47.5-55.0)
Bay	254	21.6	(15.1-29.8)	26.6	(20.7-33.5)	51.8	(43.7-59.9)
Huron	115	15.0	(8.5-25.2)	38.7	(28.9-49.5)	46.3	(35.9-57.0)
Saginaw	446	23.0	(18.0-28.9)	28.6	(23.6-34.1)	48.5	(42.6-54.4)
Sanilac	143	18.6	(12.5-26.9)	21.4	(14.9-29.6)	60.0	(50.4-68.9)
Tuscola	149	16.8	(11.2-24.6)	28.2	(20.7-37.2)	54.9	(45.6-63.9)
Region 11	926	21.2	(17.7-25.1)	26.3	(22.9-29.9)	52.5	(48.3-56.7)
Central Michigan	534	23.6	(19.1-28.9)	29.6	(25.0-34.7)	46.7	(41.3-52.3)
Midland	222	17.0	(10.8-25.7)	26.1	(19.9-33.4)	56.9	(48.6-64.8)
Region 12	797	20.4	(17.1-24.1)	29.6	(26.0-33.5)	50.0	(45.8-54.2)
Luce-Mackinac-Alger-Schoolcraft	100	19.5	(11.4-31.3)	31.9	(21.7-44.1)	48.7	(36.8-60.7)
Western Upper Peninsula	190	23.7	(17.1-32.0)	25.0	(18.6-32.7)	51.3	(42.5-60.0)
Delta-Menominee	143	17.4	(11.5-25.3)	28.9	(21.1-38.2)	53.7	(44.2-63.0)
Chippewa	113	24.5	(15.3-36.8)	31.8	(22.6-42.6)	43.7	(32.9-55.2)
Dickinson-Iron	68	24.9	(14.2-39.9)	36.7	(24.1-51.4)	38.4	(26.1-52.4)
Marquette	152	17.4	(11.4-25.8)	31.4	(23.4-40.7)	51.2	(41.7-60.5)

^a The proportion of adults who reported that they had ever smoked at least 100 cigarettes (five packs) in their life and that they smoke cigarettes now, either every day or on some days.

^b The proportion of adults who reported that they had ever smoked at least 100 cigarettes (five packs) in their life, but they do not smoke cigarettes now.

^c The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

**Table 12: Alcohol Consumption
Michigan BRFSS, 2017-2019 Combined**

Geographic Area	Sample Size	Any Alcohol in Past Month^a		Heavy Drinking^b		Binge Drinking^c	
		N	%	95% Confidence Interval	%	95% Confidence Interval	%
Michigan Total	30,498	56.8	(56.0-57.5)	6.5	(6.1-6.8)	18.1	(17.5-18.7)
Region 1	12,388	58.6	(57.5-59.7)	6.3	(5.7-6.8)	18.5	(17.6-19.4)
Livingston	1,488	63.7	(60.0-67.2)	7.7	(6.0-10.0)	19.5	(16.6-22.9)
Macomb	1,815	58.7	(56.0-61.4)	6.4	(5.1-8.0)	19.4	(17.2-21.8)
Monroe	348	55.2	(49.0-61.3)	9.7	(6.4-14.3)	18.9	(14.5-24.3)
Oakland	3,070	62.8	(60.7-64.9)	6.0	(5.0-7.1)	19.1	(17.4-21.0)
St. Clair	369	54.9	(48.7-60.9)	6.9	(4.3-10.7)	18.9	(14.5-24.3)
Washtenaw	1,432	62.5	(58.9-65.9)	6.5	(5.1-8.3)	18.4	(15.8-21.4)
City of Detroit	1,311	49.0	(45.7-52.3)	5.5	(4.2-7.3)	16.3	(14.0-18.9)
Wayne exc. Detroit	2,490	57.6	(55.2-59.9)	5.9	(4.8-7.3)	17.8	(15.9-19.9)
Region 2	2,067	51.8	(49.1-54.5)	4.5	(3.5-5.7)	14.8	(12.9-16.9)
Genesee	1,549	49.4	(46.3-52.5)	4.8	(3.6-6.3)	14.8	(12.6-17.4)
Lapeer	246	59.9	(52.7-66.7)	-- ^e	-- ^e	17.4	(12.3-23.9)
Shiawassee	253	55.0	(46.9-62.9)	-- ^e	-- ^e	12.5	(8.1-18.8)
Region 3	1,704	56.6	(53.3-59.9)	6.6	(4.9-8.7)	17.0	(14.5-19.9)
Jackson	1,352	52.3	(48.9-55.6)	5.0	(3.9-6.5)	14.7	(12.5-17.1)
Lenawee	288	63.1	(56.3-69.4)	7.9	(4.7-13.2)	19.9	(14.6-26.6)
Region 4	1,710	55.6	(52.6-58.6)	6.3	(5.0-7.9)	19.2	(16.9-21.8)
Barry-Eaton	439	49.2	(43.4-55.1)	7.3	(4.9-10.7)	16.6	(12.6-21.4)
Mid-Michigan ^d	513	57.1	(51.7-62.2)	8.7	(6.1-12.3)	20.3	(16.0-25.4)
Ingham	929	58.1	(54.0-62.2)	5.6	(4.0-7.7)	20.6	(17.4-24.3)
Region 5	1,444	53.9	(50.6-57.1)	6.3	(4.9-8.0)	17.8	(15.4-20.4)
Branch-Hillsdale-St. Joseph	422	46.9	(41.0-52.9)	6.3	(3.9-10.0)	15.5	(11.6-20.6)
Calhoun	363	44.3	(37.9-51.0)	5.0	(2.8-9.0)	12.0	(8.3-17.1)
Kalamazoo	636	64.3	(59.5-68.8)	6.9	(4.9-9.5)	22.2	(18.4-26.5)
Region 6	647	52.4	(47.6-57.1)	5.2	(3.6-7.6)	15.2	(12.0-19.0)
Van Buren-Cass	272	53.4	(46.5-60.2)	6.3	(3.6-10.9)	14.3	(10.1-20.0)
Berrien	321	50.4	(43.5-57.3)	4.0	(2.1-7.3)	15.5	(10.9-21.5)
Region 7	2,992	60.3	(58.1-62.5)	7.4	(6.4-8.6)	18.7	(17.0-20.5)
Allegan	340	56.6	(50.1-62.9)	7.9	(5.0-12.2)	12.9	(9.1-18.0)
Ionia	182	48.1	(39.1-57.2)	-- ^e	-- ^e	16.8	(10.9-25.2)
Kent	1,567	62.1	(59.1-65.0)	6.6	(5.3-8.1)	19.4	(17.1-21.9)
Mid-Michigan ^d	513	57.1	(51.7-62.2)	8.7	(6.1-12.3)	20.3	(16.0-25.4)
Ottawa	705	63.0	(58.1-67.6)	9.1	(6.8-12.0)	19.7	(16.0-24.0)

Table 12 Cont'd

Geographic Area	Sample Size	Any Alcohol in Past Month ^a		Heavy Drinking ^b		Binge Drinking ^c	
		N	%	95% Confidence Interval	%	95% Confidence Interval	%
Michigan Total	30,498	56.8	(56.0-57.5)	6.5	(6.1-6.8)	18.1	(17.5-18.7)
Region 8	2,518	50.5	(47.6-53.4)	5.4	(4.4-6.7)	16.1	(14.0-18.3)
District #10	1,057	46.9	(42.9-51.0)	4.7	(3.4-6.3)	14.4	(11.9-17.4)
Muskegon	1,359	54.7	(51.1-58.4)	7.1	(5.2-9.7)	20.1	(16.7-24.0)
Region 9	1,731	59.2	(56.2-62.2)	8.7	(7.2-10.6)	18.7	(16.4-21.2)
District #2	201	62.2	(53.7-69.9)	10.6	(6.3-17.3)	23.5	(16.9-31.8)
District #4	256	53.4	(45.2-61.5)	7.8	(4.4-13.4)	16.8	(11.8-23.4)
Northwest Michigan	527	56.1	(50.4-61.5)	7.6	(5.2-11.0)	16.8	(13.0-21.6)
Benzie-Leelanau	250	60.4	(52.4-67.8)	7.9	(4.8-12.8)	15.8	(11.0-22.1)
Grand Traverse	440	64.7	(58.7-70.2)	10.3	(7.3-14.5)	19.4	(15.0-24.8)
Region 10	1,117	52.8	(49.0-56.5)	5.5	(4.1-7.3)	18.0	(15.3-21.0)
Bay	254	53.1	(44.7-61.2)	--e	--e	16.5	(11.6-23.0)
Huron	113	62.5	(51.8-72.0)	--e	--e	14.2	(8.5-22.9)
Saginaw	446	52.4	(46.5-58.3)	6.5	(4.3-9.9)	18.7	(14.5-23.7)
Sanilac	141	49.8	(40.1-59.6)	--e	--e	17.9	(11.5-26.8)
Tuscola	150	48.8	(39.6-58.0)	--e	--e	21.0	(14.1-30.1)
Region 11	922	53.9	(49.7-58.1)	7.8	(5.8-10.4)	18.6	(15.5-22.2)
Central Michigan	531	48.8	(43.3-54.3)	7.9	(5.4-11.3)	17.7	(14.0-22.2)
Midland	223	57.1	(49.0-64.8)	5.8	(3.3-10.1)	16.5	(10.3-25.4)
Region 12	793	56.8	(52.6-60.9)	9.7	(7.4-12.8)	21.0	(17.5-24.9)
Luce-Mackinac-Alger-Schoolcraft	101	40.4	(29.6-52.1)	--e	--e	14.3	(8.0-24.4)
Western Upper Peninsula	188	58.5	(49.6-66.8)	9.6	(5.2-16.9)	22.3	(14.8-32.2)
Delta-Menominee	142	64.0	(54.5-72.4)	--e	--e	16.3	(10.1-25.1)
Chippewa	112	54.9	(43.5-65.7)	--e	--e	20.7	(12.2-33.1)
Dickinson-Iron	68	57.5	(42.8-71.0)	--e	--e	26.4	(16.0-40.2)
Marquette	152	56.3	(46.6-65.5)	11.9	(7.2-19.1)	24.4	(17.1-33.5)

^a Among all adults, the proportion who reported consuming at least one drink of any alcoholic beverage in the previous month.

^b Among all adults, the proportion who reported consuming an average of more than two alcoholic drinks per day for men or more than one per day for women in the previous month.

^c Among all adults, the proportion who reported consuming five or more drinks per occasion (for men) or 4 or more drinks per occasion (for women) at least once in the previous month.

^d The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^e Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 13: High Blood Pressure Michigan BRFSS, 2017 and 2019 Combined			
Geographic Area	Ever Told HBP ^a		
	N	%	95% Confidence Interval
Michigan Total	21,341	34.9	(34.1-35.8)
Region 1	8,786	34.4	(33.2-35.7)
Livingston	1,017	33.3	(29.4-37.5)
Macomb	1,330	35.3	(32.4-38.4)
Monroe	246	36.6	(30.2-43.6)
Oakland	2,172	32.1	(29.8-34.5)
St. Clair	240	34.3	(27.7-41.5)
Washtenaw	986	23.7	(20.6-27.1)
City of Detroit	937	44.9	(41.0-48.8)
Wayne exc. Detroit	1,810	34.1	(31.6-36.8)
Region 2	1,427	41.4	(38.2-44.6)
Genesee	1,097	40.5	(36.9-44.1)
Lapeer	154	46.4	(37.6-55.4)
Shiawassee	162	41.0	(31.4-51.4)
Region 3	1,199	35.9	(32.1-39.8)
Jackson	964	36.7	(33.0-40.6)
Lenawee	197	39.2	(31.4-47.6)
Region 4	1,188	34.9	(31.6-38.4)
Barry-Eaton	294	44.4	(37.5-51.6)
Mid-Michigan ^b	370	34.7	(29.1-40.7)
Ingham	643	31.0	(26.7-35.8)
Region 5	985	33.7	(30.1-37.4)
Branch-Hillsdale-St. Joseph	287	30.9	(24.6-38.1)
Calhoun	229	40.7	(32.8-49.1)
Kalamazoo	457	32.8	(27.9-38.1)
Region 6	453	34.9	(29.9-40.2)
Van Buren-Cass	194	39.6	(32.0-47.7)
Berrien	223	34.4	(27.3-42.3)
Region 7	2,070	32.2	(29.8-34.7)
Allegan	234	37.4	(30.4-45.0)
Ionia	127	42.7	(31.9-54.2)
Kent	1,082	31.4	(28.2-34.8)
Mid-Michigan ^b	370	34.7	(29.1-40.7)
Ottawa	490	27.4	(22.8-32.4)

Table 13 Cont'd			
Geographic Area	Ever Told HBP ^a		
	N	%	95% Confidence Interval
Michigan Total	21,341	34.9	(34.1-35.8)
Region 8	1,727	36.2	(33.0-39.5)
District #10	720	34.7	(30.4-39.3)
Muskegon	937	38.0	(33.6-42.6)
Region 9	1,190	35.5	(31.9-39.2)
District #2	140	42.3	(32.5-52.8)
District #4	172	38.2	(29.0-48.2)
Northwest Michigan	364	34.8	(28.7-41.4)
Benzie-Leelanau	179	38.1	(29.8-47.1)
Grand Traverse	301	30.9	(24.8-37.6)
Region 10	783	37.7	(33.6-42.0)
Bay	175	30.0	(22.6-38.6)
Huron	71	36.8	(25.5-49.7)
Saginaw	331	40.7	(34.1-47.7)
Sanilac	88	42.6	(31.6-54.4)
Tuscola	107	39.5	(29.6-50.5)
Region 11	626	37.7	(33.1-42.5)
Central Michigan	374	38.6	(32.9-44.7)
Midland	162	39.9	(31.3-49.2)
Region 12	555	37.3	(32.8-42.0)
Luce-Mackinac-Alger-Schoolcraft	63	35.0	(22.9-49.3)
Western Upper Peninsula	121	41.8	(32.1-52.1)
Delta-Menominee	98	39.0	(28.5-50.6)
Chippewa	81	25.6	(17.0-36.7)
Dickinson-Iron	-- ^c	-- ^c	-- ^c
Marquette	110	38.8	(28.8-49.8)

^a Among all adults, the proportion who reported that they were ever told by a doctor that they had high blood pressure (HBP). Women who had HBP only during pregnancy and adults who were borderline hypertensive were considered to not have been diagnosed.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 14: Cholesterol Screening and Awareness
Michigan BRFSS, 2017 and 2019 Combined

Geographic Area	Cholesterol Ever Checked ^a			Ever Told High Cholesterol ^b		
	N	%	95% Confidence Interval	N	%	95% Confidence Interval
Michigan Total	20,119	93.6	(93.1-94.1)	19,742	33.6	(32.8-34.5)
Region 1						
Livingston	8,340	94.7	(93.9-95.3)	8,215	33.9	(32.6-35.1)
Macomb	969	93.4	(90.7-95.4)	945	35.5	(31.3-40.0)
Monroe	1,267	94.0	(91.7-95.6)	1,242	35.9	(32.9-39.1)
Oakland	-- ^d	-- ^d	-- ^d	226	30.9	(24.8-37.9)
St. Clair	2,076	96.8	(95.6-97.7)	2,069	35.0	(32.5-37.5)
Washtenaw	-- ^d	-- ^d	-- ^d	217	39.6	(32.2-47.5)
City of Detroit	931	91.1	(87.8-93.5)	913	26.6	(23.3-30.2)
Wayne exc. Detroit	882	94.6	(92.3-96.3)	859	32.6	(28.9-36.5)
	1,719	94.8	(93.2-96.1)	1,701	33.7	(31.1-36.4)
Region 2						
Genesee	1,341	93.5	(91.2-95.2)	1,314	36.3	(33.2-39.6)
Lapeer	1,030	93.9	(91.4-95.8)	1,012	35.9	(32.4-39.6)
Shiawassee	-- ^d	-- ^d	-- ^d	141	40.6	(31.9-50.0)
	-- ^d	-- ^d	-- ^d	148	33.1	(24.2-43.4)
Region 3						
Jackson	1,125	93.3	(90.3-95.4)	1,102	34.5	(30.8-38.5)
Lenawee	905	92.6	(89.5-94.9)	888	36.3	(32.6-40.3)
	-- ^d	-- ^d	-- ^d	178	36.2	(28.4-44.8)
Region 4						
Barry-Eaton	1,098	91.4	(88.6-93.6)	1,073	33.1	(29.8-36.5)
Mid-Michigan ^c	-- ^d	-- ^d	-- ^d	276	40.1	(33.5-47.2)
Ingham	-- ^d	-- ^d	-- ^d	349	28.7	(23.5-34.6)
	585	89.3	(85.4-92.2)	562	31.6	(27.1-36.5)
Region 5						
Branch-Hillsdale-St. Joseph	929	90.2	(86.8-92.8)	897	34.8	(31.2-38.7)
Calhoun	-- ^d	-- ^d	-- ^d	265	37.7	(30.7-45.1)
Kalamazoo	-- ^d	-- ^d	-- ^d	211	35.2	(27.7-43.4)
	426	88.4	(83.1-92.2)	409	33.5	(28.3-39.1)
Region 6						
Van Buren-Cass	426	94.3	(90.4-96.7)	422	35.5	(30.3-40.9)
Berrien	-- ^d	-- ^d	-- ^d	184	39.9	(32.1-48.3)
	-- ^d	-- ^d	-- ^d	206	32.4	(25.6-40.1)

Table 14 Cont'd

Geographic Area	Cholesterol Ever Checked^a			Ever Told High Cholesterol^b		
	N	%	95% Confidence Interval	N	%	95% Confidence Interval
Michigan Total	20,119	93.6	(93.1-94.1)	19,742	33.6	(32.8-34.5)
Region 7						
Allegan	1,944	95.3	(93.8-96.4)	1,933	28.0	(25.7-30.4)
Ionia	--d	--d	--d	216	31.0	(24.3-38.5)
Kent	--d	--d	--d	118	29.8	(21.1-40.3)
Mid-Michigan ^c	1,008	96.1	(94.1-97.4)	1,011	27.9	(24.8-31.2)
Ottawa	--d	--d	--d	349	28.7	(23.5-34.6)
	459	93.9	(89.9-96.3)	456	25.4	(21.1-30.2)
Region 8	1,629	92.0	(89.7-93.8)	1,582	34.8	(31.6-38.2)
District #10	683	92.0	(88.5-94.5)	651	33.2	(28.8-37.9)
Muskegon	885	91.6	(88.1-94.2)	866	38.4	(34.3-42.7)
Region 9						
District #2	1,119	91.5	(88.8-93.5)	1,090	34.9	(31.3-38.7)
District #4	--d	--d	--d	124	29.3	(20.0-40.7)
Northwest Michigan	--d	--d	--d	157	37.9	(28.7-48.1)
Benzie-Leelanau	337	90.4	(84.0-94.4)	332	41.1	(34.4-48.2)
Grand Traverse	--d	--d	--d	168	34.9	(26.7-44.0)
	--d	--d	--d	279	30.4	(24.5-37.1)
Region 10	749	93.5	(90.4-95.7)	729	35.7	(31.5-40.1)
Bay	--d	--d	--d	161	35.9	(26.5-46.6)
Huron	--d	--d	--d	68	39.2	(27.1-52.9)
Saginaw	--d	--d	--d	312	33.1	(27.0-39.9)
Sanilac	--d	--d	--d	77	40.4	(28.9-52.9)
Tuscola	--d	--d	--d	101	40.2	(29.9-51.5)
Region 11	588	94.4	(91.2-96.5)	581	36.2	(31.4-41.3)
Central Michigan	351	94.5	(90.5-96.9)	349	37.6	(31.6-44.1)
Midland	--d	--d	--d	153	37.7	(29.0-47.3)
Region 12						
Luce-Mackinac-Alger-Schoolcraft	516	90.2	(85.8-93.4)	503	37.2	(32.5-42.2)
Western Upper Peninsula	--d	--d	--d	61	33.8	(21.5-48.8)
Delta-Menominee	--d	--d	--d	110	38.8	(28.9-49.6)
Chippewa	--d	--d	--d	89	43.9	(32.5-55.9)
Dickinson-Iron	--d	--d	--d	68	30.7	(20.2-43.7)
Marquette	--d	--d	--d	--d	--d	--d
	--d	--d	--d	100	42.5	(31.3-54.5)

^a Among all adults, the proportion who reported ever having had their blood cholesterol checked.

^b Among adults who ever had their blood cholesterol checked, the proportion who reported that a doctor, nurse, or other health professional had told them that their cholesterol was high.

^c The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^d Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 15: No Routine Checkup in Past Year Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Had No Routine Checkup in Past Year^a		
	N	%	95% Confidence Interval
Michigan Total	31,384	22.3	(21.6-22.9)
Region 1	12,788	21.3	(20.4-22.3)
Livingston	1,523	20.8	(17.9-24.1)
Macomb	1,868	23.5	(21.2-26.1)
Monroe	357	21.0	(16.3-26.6)
Oakland	3,162	21.9	(20.1-23.8)
St. Clair	382	22.3	(17.2-28.3)
Washtenaw	1,470	24.8	(22.0-28.0)
City of Detroit	1,394	18.1	(15.6-20.9)
Wayne exc. Detroit	2,562	19.3	(17.4-21.3)
Region 2	2,147	20.0	(17.9-22.4)
Genesee	1,620	18.9	(16.5-21.5)
Lapeer	246	25.2	(19.2-32.2)
Shiawassee	261	21.1	(15.0-29.0)
Region 3	1,749	26.5	(23.5-29.8)
Jackson	1,393	23.8	(21.0-26.9)
Lenawee	292	28.7	(22.4-35.8)
Region 4	1,735	23.9	(21.4-26.7)
Barry-Eaton	448	21.8	(17.0-27.5)
Mid-Michigan ^b	521	21.5	(17.2-26.5)
Ingham	941	25.8	(22.3-29.6)
Region 5	1,475	24.2	(21.4-27.1)
Branch-Hillsdale-St. Joseph	432	26.2	(21.2-31.9)
Calhoun	367	21.1	(16.2-27.1)
Kalamazoo	654	24.6	(20.6-29.2)
Region 6	666	24.4	(20.6-28.8)
Van Buren-Cass	281	22.5	(17.3-28.7)
Berrien	330	25.0	(19.5-31.5)
Region 7	3,063	22.1	(20.2-24.1)
Allegan	344	21.3	(16.5-27.0)
Ionia	186	25.2	(17.0-35.6)
Kent	1,613	22.3	(19.8-24.9)
Mid-Michigan ^b	521	21.5	(17.2-26.5)
Ottawa	717	21.7	(17.7-26.3)

Table 15 Cont'd			
Geographic Area	Had No Routine Checkup in Past Year ^a		
	N	%	95% Confidence Interval
Michigan Total	31,384	22.3	(21.6-22.9)
Region 8			
District #10	2,577	21.7	(19.4-24.1)
Muskegon	1,070	24.0	(20.8-27.6)
	1,405	18.8	(16.0-21.9)
Region 9	1,762	22.1	(19.5-24.8)
District #2	206	22.7	(15.8-31.4)
District #4	261	20.2	(14.3-27.8)
Northwest Michigan	539	21.9	(17.6-26.8)
Benzie-Leelanau	252	23.7	(17.4-31.4)
Grand Traverse	450	18.5	(14.4-23.4)
Region 10	1,153	24.9	(21.6-28.6)
Bay	258	27.7	(20.3-36.6)
Huron	116	26.1	(17.6-36.7)
Saginaw	465	25.4	(20.3-31.3)
Sanilac	150	23.6	(15.9-33.5)
Tuscola	151	16.6	(10.7-24.9)
Region 11	948	25.3	(21.6-29.4)
Central Michigan	541	24.9	(20.0-30.5)
Midland	228	20.5	(14.0-29.0)
Region 12	805	23.5	(20.0-27.4)
Luce-Mackinac-Alger-Schoolcraft	104	20.8	(12.3-33.0)
Western Upper Peninsula	190	22.1	(15.4-30.7)
Delta-Menominee	143	24.7	(16.7-34.9)
Chippewa	113	27.8	(18.6-39.5)
Dickinson-Iron	68	18.2	(8.4-34.8)
Marquette	154	25.2	(18.1-34.1)

^a The proportion of adults who reported that they did not have a routine checkup in the past year.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

Table 16: Immunizations Among Adults Aged 65 Years and Older Michigan BRFSS, 2017-2019 Combined						
Geographic Area	Had Flu Vaccine in Past Year ^a			Ever Had Pneumonia Vaccine ^b		
	N	%	95% Confidence Interval	N	%	95% Confidence Interval
Michigan Total	10,443	57.6	(56.3-58.8)	10,182	74.1	(73.0-75.3)
Region 1						
Livingston	542	61.0	(54.9-66.8)	530	79.6	(74.2-84.2)
Macomb	615	61.7	(57.0-66.3)	598	78.5	(74.2-82.2)
Monroe	111	50.0	(39.3-60.8)	110	75.9	(64.7-84.4)
Oakland	1,013	60.5	(56.8-64.1)	977	75.2	(71.8-78.3)
St. Clair	113	54.4	(43.6-64.8)	111	75.0	(64.9-83.0)
Washtenaw	515	62.7	(57.3-67.8)	498	79.3	(74.6-83.4)
City of Detroit	362	47.7	(41.3-54.2)	350	60.7	(54.0-67.0)
Wayne exc. Detroit	804	58.9	(54.5-63.1)	780	73.6	(69.2-77.6)
Region 2						
Genesee	511	57.8	(52.6-62.9)	499	68.3	(63.1-73.1)
Lapeer	80	49.2	(37.2-61.2)	79	58.6	(45.8-70.3)
Shiawassee	96	61.9	(50.1-72.4)	96	79.7	(69.6-87.0)
Region 3						
Jackson	492	53.3	(47.9-58.7)	488	75.5	(70.4-80.0)
Lenawee	102	46.9	(36.0-58.1)	101	75.8	(64.9-84.1)
Region 4						
Barry-Eaton	170	54.5	(45.8-63.0)	164	75.7	(67.5-82.4)
Mid-Michigan ^c	185	66.5	(58.3-73.8)	182	80.5	(73.5-86.0)
Ingham	266	66.2	(58.8-72.8)	262	77.9	(70.6-83.7)
Region 5						
Branch-Hillsdale-St. Joseph	164	52.3	(43.4-61.1)	161	65.6	(56.3-73.9)
Calhoun	135	54.7	(44.3-64.7)	133	66.9	(56.2-76.1)
Kalamazoo	211	59.3	(50.6-67.5)	205	79.5	(70.0-86.5)
Region 6						
Van Buren-Cass	105	66.3	(54.2-76.6)	103	79.4	(68.9-87.1)
Berrien	120	52.2	(41.7-62.6)	118	75.6	(65.7-83.4)

Table 16 Cont'd

Geographic Area	Had Flu Vaccine in Past Year^a			Ever Had Pneumonia Vaccine^b		
	N	%	95% Confidence Interval	N	%	95% Confidence Interval
Michigan Total	10,443	57.6	(56.3-58.8)	10,182	74.1	(73.0-75.3)
Region 7						
Allegan	977	56.8	(53.1-60.5)	952	78.0	(74.7-81.0)
Ionia	122	50.3	(39.9-60.7)	121	75.0	(65.9-82.4)
Kent	55	53.8	(38.1-68.8)	54	67.0	(51.0-79.8)
Mid-Michigan ^c	498	59.0	(53.7-64.0)	482	78.3	(73.3-82.6)
Ottawa	185	66.5	(58.3-73.8)	182	80.5	(73.5-86.0)
	236	57.4	(50.1-64.4)	229	83.4	(77.1-88.1)
Region 8						
District #10	936	59.7	(55.5-63.7)	916	76.1	(72.0-79.8)
Muskegon	436	57.9	(52.1-63.4)	427	75.7	(70.0-80.7)
	476	62.2	(56.6-67.6)	465	76.0	(70.4-80.8)
Region 9						
District #2	697	58.5	(53.4-63.5)	677	73.8	(68.9-78.2)
District #4	79	62.4	(48.7-74.4)	78	76.9	(63.4-86.4)
Northwest Michigan	107	48.5	(35.7-61.5)	101	62.7	(48.6-74.9)
Benzie-Leelanau	219	58.9	(49.0-68.2)	213	77.1	(69.3-83.3)
Grand Traverse	125	59.0	(48.6-68.8)	122	74.1	(63.7-82.3)
	167	66.2	(56.7-74.6)	163	79.2	(70.7-85.7)
Region 10						
Bay	412	57.2	(50.9-63.2)	403	72.7	(67.2-77.6)
Huron	87	69.6	(58.2-79.1)	84	77.8	(66.2-86.3)
Saginaw	56	63.3	(48.2-76.1)	56	71.1	(56.9-82.1)
Sanilac	154	48.3	(37.9-58.8)	151	70.6	(60.8-78.8)
Tuscola	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d
	69	54.9	(41.4-67.7)	67	66.7	(52.2-78.6)
Region 11						
Central Michigan	317	52.1	(45.1-59.0)	306	78.7	(73.1-83.4)
Midland	204	48.1	(39.5-56.8)	197	77.9	(70.8-83.6)
	88	60.8	(48.3-72.0)	85	77.1	(65.3-85.7)
Region 12						
Luce-Mackinac-Alger-Schoolcraft	285	51.2	(44.2-58.1)	281	65.3	(58.1-71.9)
Western Upper Peninsula	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d
Delta-Menominee	74	42.0	(29.8-55.2)	73	66.7	(52.8-78.2)
Chippewa	51	44.3	(28.4-61.4)	51	56.4	(39.4-72.0)
Dickinson-Iron	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d
Marquette	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d
	60	65.8	(50.0-78.7)	57	72.3	(55.6-84.5)

^a Among adults aged 65 years and older, the proportion who reported that they had a flu vaccine, either by an injection in the arm or sprayed in the nose during the past 12 months.

^b Among adults aged 65 years and older, the proportion who reported that they ever had a pneumococcal vaccine.

^c The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^d Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 17: HIV Testing Among Adults Aged 18-64 Years Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Had an HIV Test ^a		
	N	%	95% Confidence Interval
Michigan Total	18,393	46.3	(45.4-47.2)
Region 1	7,612	50.2	(48.8-51.5)
Livingston	877	43.4	(38.8-48.1)
Macomb	1,089	44.6	(41.3-48.1)
Monroe	221	42.4	(35.2-50.0)
Oakland	1,861	48.2	(45.5-50.9)
St. Clair	231	44.0	(36.7-51.7)
Washtenaw	868	49.7	(45.5-53.9)
City of Detroit	861	74.0	(70.3-77.3)
Wayne exc. Detroit	1,558	46.1	(43.2-49.1)
Region 2	1,275	49.2	(45.9-52.6)
Genesee	949	52.5	(48.6-56.4)
Lapeer	162	39.8	(31.6-48.6)
Shiawassee	152	43.0	(33.3-53.3)
Region 3	1,009	45.9	(41.7-50.1)
Jackson	790	50.3	(46.1-54.6)
Lenawee	171	39.1	(30.8-48.1)
Region 4	1,067	44.4	(40.8-48.1)
Barry-Eaton	253	50.5	(43.1-57.8)
Mid-Michigan ^b	300	41.2	(34.8-47.9)
Ingham	614	42.5	(37.7-47.4)
Region 5	848	46.6	(42.5-50.7)
Branch-Hillsdale-St. Joseph	238	43.6	(36.1-51.5)
Calhoun	209	49.1	(40.7-57.5)
Kalamazoo	387	46.7	(40.9-52.7)
Region 6	388	45.8	(39.9-51.7)
Van Buren-Cass	154	42.4	(33.9-51.4)
Berrien	189	47.4	(38.9-56.1)
Region 7	1,883	42.3	(39.6-45.0)
Allegan	205	42.3	(34.4-50.6)
Ionia	118	38.0	(28.4-48.7)
Kent	1,007	45.0	(41.5-48.7)
Mid-Michigan ^b	300	41.2	(34.8-47.9)
Ottawa	434	36.8	(31.3-42.6)

Table 17 Cont'd			
Geographic Area	Ever Had an HIV Test ^a		
	N	%	95% Confidence Interval
Michigan Total	18,393	46.3	(45.4-47.2)
Region 8			
District #10	1,446	40.8	(37.2-44.4)
Muskegon	571	37.5	(32.5-42.8)
	810	47.2	(42.6-51.9)
Region 9			
District #2	948	39.0	(35.3-42.8)
District #4	112	34.9	(25.2-46.0)
Northwest Michigan	130	35.0	(25.9-45.4)
Benzie-Leelanau	279	39.9	(33.2-47.0)
Grand Traverse	115	40.2	(30.3-50.9)
	259	42.5	(35.4-49.8)
Region 10			
Bay	649	42.4	(37.8-47.3)
Huron	150	44.3	(34.1-55.0)
Saginaw	55	21.6	(11.5-37.0)
Sanilac	272	44.1	(37.0-51.4)
Tuscola	90	44.5	(33.2-56.4)
	74	43.0	(31.1-55.8)
Region 11			
Central Michigan	552	37.2	(32.3-42.4)
Midland	295	39.0	(32.3-46.1)
	127	30.5	(22.0-40.5)
Region 12			
Luce-Mackinac-Alger-Schoolcraft	461	40.1	(35.0-45.5)
Western Upper Peninsula	53	42.0	(27.5-58.0)
Delta-Menominee	110	34.2	(24.7-45.0)
Chippewa	83	38.7	(27.8-51.0)
Dickinson-Iron	69	49.9	(35.6-64.2)
Marquette	-- ^c	-- ^c	-- ^c
	80	36.7	(25.5-49.5)

Note: "Don't know" was considered a valid response.

^a Among adults aged 18 - 64 years, the proportion who reported that they ever had been tested for HIV, apart from tests that were part of a blood donation.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 18: Asthma
Michigan BRFSS, 2017-2019 Combined

Geographic Area	Ever Told Have Asthma ^a			Still Have Asthma ^b		
	N	%	95% Confidence Interval	N	%	95% Confidence Interval
Michigan Total	31,602	16.1	(15.6-16.7)	31,441	11.1	(10.6-11.6)
Region 1						
Livingston	12,863	16.3	(15.5-17.1)	12,806	11.3	(10.6-12.0)
Macomb	1,536	13.5	(11.2-16.1)	1,531	8.3	(6.8-10.1)
Monroe	1,881	17.1	(15.1-19.3)	1,869	11.2	(9.5-13.2)
Oakland	363	15.7	(11.9-20.6)	363	10.0	(7.0-14.1)
St. Clair	3,181	14.3	(12.9-15.9)	3,169	9.3	(8.2-10.7)
Washtenaw	383	23.1	(18.1-29.0)	377	17.9	(13.2-23.7)
City of Detroit	1,472	15.4	(13.2-18.0)	1,464	9.8	(8.0-12.1)
Wayne exc. Detroit	1,398	20.6	(18.2-23.3)	1,394	16.2	(14.0-18.7)
	2,577	15.5	(13.9-17.3)	2,567	11.2	(9.8-12.7)
Region 2						
Genesee	2,158	17.3	(15.2-19.5)	2,143	12.8	(11.0-14.8)
Lapeer	1,626	18.0	(15.7-20.5)	1,617	12.8	(10.9-15.0)
Shiawassee	250	13.0	(9.0-18.6)	247	11.5	(7.6-17.1)
	262	18.1	(12.0-26.5)	259	13.8	(8.2-22.3)
Region 3						
Jackson	1,762	17.1	(14.7-19.7)	1,750	11.6	(9.7-13.9)
Lenawee	1,402	16.5	(14.2-19.0)	1,395	11.9	(9.9-14.2)
	296	19.1	(14.1-25.3)	293	13.1	(8.9-18.9)
Region 4						
Barry-Eaton	1,750	17.0	(14.9-19.4)	1,742	10.9	(9.2-12.8)
Mid-Michigan ^c	453	16.5	(12.5-21.5)	452	10.0	(7.0-14.0)
Ingham	523	14.6	(11.1-18.9)	523	10.3	(7.4-14.0)
	949	18.5	(15.5-21.9)	942	12.3	(9.9-15.2)
Region 5						
Branch-Hillsdale-St. Joseph	1,486	18.2	(15.8-20.9)	1,480	11.6	(9.7-13.8)
Calhoun	432	17.0	(12.8-22.3)	432	10.4	(7.3-14.6)
Kalamazoo	368	18.5	(14.0-23.9)	364	12.5	(8.8-17.5)
	663	18.5	(14.9-22.7)	661	11.4	(8.7-14.8)
Region 6						
Van Buren-Cass	668	13.6	(10.7-17.0)	662	7.9	(5.7-10.9)
Berrien	280	10.9	(7.3-15.8)	275	5.4	(3.0-9.7)
	333	13.4	(9.8-18.1)	333	8.0	(5.4-11.9)
Region 7						
Allegan	3,079	14.6	(13.1-16.3)	3,067	9.8	(8.6-11.2)
Ionia	348	15.9	(11.5-21.5)	345	10.5	(7.1-15.3)
Kent	187	11.3	(7.0-17.8)	187	7.4	(4.4-12.3)
Mid-Michigan ^c	1,622	15.1	(13.1-17.3)	1,616	10.4	(8.7-12.4)
Ottawa	523	14.6	(11.1-18.9)	523	10.3	(7.4-14.0)
	719	13.6	(10.3-17.9)	716	7.8	(5.8-10.5)

Table 18 Cont'd

Geographic Area	Ever Told Have Asthma^a			Still Have Asthma^b		
	N	%	95% Confidence Interval	N	%	95% Confidence Interval
Michigan Total	31,602	16.1	(15.6-16.7)	31,441	11.1	(10.6-11.6)
Region 8						
District #10	2,598	15.4	(13.3-17.8)	2,582	10.5	(8.6-12.7)
Muskegon	1,082	15.2	(12.2-18.8)	1,074	10.5	(7.8-14.0)
	1,413	15.9	(13.4-18.8)	1,406	10.1	(8.2-12.5)
Region 9						
District #2	1,780	14.8	(12.7-17.2)	1,774	10.1	(8.3-12.2)
District #4	210	16.3	(10.7-24.0)	209	13.1	(8.0-20.8)
Northwest Michigan	265	13.7	(9.2-19.9)	265	8.2	(4.9-13.2)
Benzie-Leelanau	546	15.7	(12.0-20.3)	543	12.2	(8.9-16.5)
Grand Traverse	253	11.3	(7.1-17.4)	-- ^d	-- ^d	-- ^d
	449	15.1	(11.3-19.9)	447	9.2	(6.1-13.6)
Region 10						
Bay	1,160	17.8	(14.8-21.2)	1,152	12.5	(9.8-15.8)
Huron	257	22.3	(15.1-31.7)	255	16.7	(9.9-26.6)
Saginaw	116	16.1	(9.6-25.8)	-- ^d	-- ^d	-- ^d
Sanilac	471	18.3	(13.9-23.6)	469	12.6	(8.8-17.7)
Tuscola	150	17.6	(11.5-26.1)	146	12.3	(7.2-20.1)
	153	7.7	(4.2-13.5)	-- ^d	-- ^d	-- ^d
Region 11						
Central Michigan	961	15.6	(12.8-18.9)	957	11.5	(9.0-14.5)
Midland	550	13.9	(10.5-18.3)	547	11.6	(8.3-15.9)
	230	18.9	(13.4-26.0)	229	13.1	(8.8-19.2)
Region 12						
Luce-Mackinac-Alger-Schoolcraft	809	16.1	(13.0-19.8)	804	13.1	(10.2-16.7)
Western Upper Peninsula	104	13.4	(7.6-22.6)	-- ^d	-- ^d	-- ^d
Delta-Menominee	191	16.4	(9.8-26.2)	-- ^d	-- ^d	-- ^d
Chippewa	144	16.7	(10.0-26.5)	-- ^d	-- ^d	-- ^d
Dickinson-Iron	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d	-- ^d
Marquette	68	23.9	(12.7-40.3)	-- ^d	-- ^d	-- ^d
	154	18.2	(12.1-26.4)	152	16.3	(10.5-24.6)

^a The proportion of adults who reported that they were ever told by a doctor, nurse, or other health care professional that they had asthma.

^b Among all adults, the proportion who reported that they still had asthma.

^c The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^d Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 19: Chronic Obstructive Pulmonary Disease (COPD) Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told COPD, Emphysema or Chronic Bronchitis^a		
	N	%	95% Confidence Interval
Michigan Total	31,546	8.6	(8.2-9.0)
Region 1	12,842	8.3	(7.7-8.9)
Livingston	1,534	6.5	(5.0-8.4)
Macomb	1,875	8.5	(7.2-10.1)
Monroe	360	10.6	(7.6-14.5)
Oakland	3,175	6.5	(5.5-7.6)
St. Clair	384	14.1	(10.2-19.2)
Washtenaw	1,474	4.0	(3.0-5.4)
City of Detroit	1,393	11.9	(10.0-14.1)
Wayne exc. Detroit	2,575	9.1	(7.9-10.4)
Region 2	2,152	12.8	(11.1-14.7)
Genesee	1,620	13.8	(11.8-16.2)
Lapeer	248	12.4	(8.2-18.2)
Shiawassee	264	8.1	(4.7-13.5)
Region 3	1,759	10.3	(8.5-12.4)
Jackson	1,400	10.4	(8.6-12.5)
Lenawee	295	11.4	(7.6-16.7)
Region 4	1,750	6.9	(5.7-8.4)
Barry-Eaton	452	8.8	(6.2-12.4)
Mid-Michigan ^b	522	8.1	(6.0-11.1)
Ingham	951	5.9	(4.4-7.8)
Region 5	1,480	9.7	(7.8-11.9)
Branch-Hillsdale-St. Joseph	433	10.3	(7.3-14.4)
Calhoun	365	13.2	(9.0-19.1)
Kalamazoo	659	7.4	(5.1-10.7)
Region 6	668	9.1	(6.8-12.0)
Van Buren-Cass	281	12.2	(8.2-17.8)
Berrien	333	7.8	(5.1-11.8)
Region 7	3,085	6.2	(5.3-7.2)
Allegan	349	9.9	(7.0-13.8)
Ionia	188	9.0	(5.2-15.0)
Kent	1,621	5.5	(4.4-6.8)
Mid-Michigan ^b	522	8.1	(6.0-11.1)
Ottawa	724	4.4	(3.1-6.2)

Table 19 Cont'd			
Geographic Area	Ever Told COPD, Emphysema or Chronic Bronchitis ^a		
	N	%	95% Confidence Interval
Michigan Total	31,546	8.6	(8.2-9.0)
Region 8			
District #10	2,592	10.2	(8.7-11.9)
Muskegon	1,079	12.1	(9.9-14.6)
	1,409	7.6	(6.1-9.3)
Region 9	1,776	10.6	(8.7-12.9)
District #2	208	19.4	(12.9-28.2)
District #4	264	13.1	(8.4-19.9)
Northwest Michigan	546	9.1	(6.6-12.4)
Benzie-Leelanau	255	6.1	(3.5-10.5)
Grand Traverse	446	7.4	(5.0-10.9)
Region 10	1,149	8.7	(6.9-10.9)
Bay	254	7.3	(4.7-11.2)
Huron	-- ^c	-- ^c	-- ^c
Saginaw	468	9.0	(6.1-13.1)
Sanilac	150	13.2	(7.9-21.2)
Tuscola	150	9.4	(5.5-15.5)
Region 11	959	8.5	(6.5-11.1)
Central Michigan	549	11.1	(8.1-15.0)
Midland	231	5.7	(3.1-10.0)
Region 12	804	8.4	(6.5-10.7)
Luce-Mackinac-Alger-Schoolcraft	-- ^c	-- ^c	-- ^c
Western Upper Peninsula	187	7.5	(4.2-12.9)
Delta-Menominee	-- ^c	-- ^c	-- ^c
Chippewa	-- ^c	-- ^c	-- ^c
Dickinson-Iron	66	18.1	(9.8-31.0)
Marquette	153	10.3	(6.2-16.6)

^a Among all adults, the proportion who reported ever being told by a doctor that they had chronic obstructive pulmonary disease (COPD), emphysema or chronic bronchitis.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 20: Arthritis Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told Arthritis^a		
	N	%	95% Confidence Interval
Michigan Total	31,544	31.1	(30.5-31.8)
Region 1			
Livingston	1,538	27.6	(24.6-30.9)
Macomb	1,871	32.0	(29.6-34.5)
Monroe	362	35.6	(30.1-41.4)
Oakland	3,176	28.3	(26.5-30.2)
St. Clair	383	36.9	(31.4-42.7)
Washtenaw	1,468	22.2	(19.8-24.8)
City of Detroit	1,399	32.5	(29.7-35.5)
Wayne exc. Detroit	2,572	31.0	(28.9-33.1)
Region 2	2,148	38.0	(35.6-40.6)
Genesee	1,617	37.4	(34.6-40.3)
Lapeer	249	40.3	(33.8-47.2)
Shiawassee	262	39.3	(31.9-47.1)
Region 3	1,758	33.6	(30.6-36.8)
Jackson	1,401	33.6	(30.6-36.7)
Lenawee	293	37.1	(30.7-44.0)
Region 4	1,746	30.7	(28.2-33.4)
Barry-Eaton	454	32.7	(27.8-38.0)
Mid-Michigan ^b	522	35.9	(31.2-41.0)
Ingham	946	28.0	(24.5-31.7)
Region 5	1,485	31.6	(28.8-34.5)
Branch-Hillsdale-St. Joseph	434	37.9	(32.4-43.7)
Calhoun	369	34.8	(28.9-41.2)
Kalamazoo	659	25.8	(22.2-29.9)
Region 6	665	33.4	(29.2-37.8)
Van Buren-Cass	278	36.4	(30.1-43.2)
Berrien	333	31.8	(26.1-38.1)
Region 7	3,074	26.9	(25.1-28.7)
Allegan	346	31.7	(26.3-37.7)
Ionia	188	35.0	(27.3-43.6)
Kent	1,614	25.7	(23.3-28.3)
Mid-Michigan ^b	522	35.9	(31.2-41.0)
Ottawa	722	22.3	(19.2-25.9)

Table 20 Cont'd			
Geographic Area	Ever Told Arthritis ^a		
	N	%	95% Confidence Interval
Michigan Total	31,544	31.1	(30.5-31.8)
Region 8			
District #10	2,595	36.1	(33.6-38.7)
Muskegon	1,081	37.4	(33.7-41.1)
	1,410	37.4	(34.2-40.8)
Region 9	1,778	34.6	(31.8-37.5)
District #2	209	41.8	(33.8-50.2)
District #4	264	37.6	(30.5-45.3)
Northwest Michigan	546	36.8	(31.6-42.3)
Benzie-Leelanau	252	30.1	(24.0-37.0)
Grand Traverse	450	31.1	(26.1-36.6)
Region 10	1,155	33.0	(29.7-36.4)
Bay	258	36.3	(29.0-44.4)
Huron	115	19.4	(13.1-27.6)
Saginaw	468	31.4	(26.5-36.7)
Sanilac	148	39.9	(31.0-49.5)
Tuscola	153	37.4	(29.2-46.4)
Region 11	959	30.5	(27.1-34.1)
Central Michigan	548	34.9	(30.2-39.9)
Midland	230	33.8	(27.1-41.2)
Region 12	810	36.2	(32.4-40.2)
Luce-Mackinac-Alger-Schoolcraft	103	40.0	(29.0-52.1)
Western Upper Peninsula	192	34.9	(27.3-43.3)
Delta-Menominee	146	36.4	(28.1-45.5)
Chippewa	113	34.9	(25.5-45.7)
Dickinson-Iron	68	44.8	(31.5-58.8)
Marquette	154	40.6	(31.8-50.1)

^a Among all adults, the proportion who reported ever being told by a health care professional that they had some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates

Table 21: Heart Attack Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told Heart Attack ^a		
	N	%	95% Confidence Interval
Michigan Total	31,548	5.0	(4.8-5.3)
Region 1	12,853	5.0	(4.5-5.4)
Livingston	1,535	3.8	(2.9-5.1)
Macomb	1,879	5.6	(4.5-6.9)
Monroe	362	7.4	(4.9-11.1)
Oakland	3,186	4.1	(3.4-5.0)
St. Clair	384	6.1	(4.0-9.2)
Washtenaw	1,470	3.1	(2.3-4.2)
City of Detroit	1,391	5.1	(3.9-6.6)
Wayne exc. Detroit	2,573	5.8	(4.9-7.0)
Region 2	2,142	5.7	(4.6-7.0)
Genesee	1,611	5.5	(4.3-6.9)
Lapeer	249	9.1	(5.5-14.6)
Shiawassee	262	3.9	(2.2-6.7)
Region 3	1,761	5.9	(4.7-7.4)
Jackson	1,402	5.8	(4.7-7.3)
Lenawee	295	7.1	(4.5-11.0)
Region 4	1,753	3.9	(3.1-5.0)
Barry-Eaton	453	3.8	(2.4-5.8)
Mid-Michigan ^b	524	5.5	(3.7-8.2)
Ingham	952	3.2	(2.2-4.7)
Region 5	1,482	5.9	(4.6-7.5)
Branch-Hillsdale-St. Joseph	434	7.5	(4.9-11.3)
Calhoun	366	5.6	(3.4-9.1)
Kalamazoo	659	5.0	(3.4-7.3)
Region 6	665	5.2	(3.6-7.4)
Van Buren-Cass	281	8.5	(5.5-12.8)
Berrien	-- ^c	-- ^c	-- ^c
Region 7	3,077	3.1	(2.5-3.8)
Allegan	349	3.3	(2.0-5.5)
Ionia	-- ^c	-- ^c	-- ^c
Kent	1,619	3.0	(2.2-4.2)
Mid-Michigan ^b	524	5.5	(3.7-8.2)
Ottawa	719	2.3	(1.5-3.8)

Table 21 Cont'd

Geographic Area	Ever Told Heart Attack^a		
	N	%	95% Confidence Interval
Michigan Total	31,548	5.0	(4.8-5.3)
Region 8			
District #10	2,591	5.9	(4.8-7.2)
Muskegon	1,076	7.5	(5.8-9.6)
	1,411	4.4	(3.4-5.7)
Region 9	1,777	5.4	(4.3-6.8)
District #2	-- ^c	-- ^c	-- ^c
District #4	266	6.7	(4.4-9.9)
Northwest Michigan	544	6.2	(4.1-9.1)
Benzie-Leelanau	-- ^c	-- ^c	-- ^c
Grand Traverse	449	5.1	(3.2-7.9)
Region 10	1,151	6.4	(4.9-8.3)
Bay	257	4.5	(2.6-7.7)
Huron	-- ^c	-- ^c	-- ^c
Saginaw	467	7.4	(4.9-11.2)
Sanilac	147	8.7	(4.8-15.4)
Tuscola	-- ^c	-- ^c	-- ^c
Region 11	959	6.2	(4.5-8.6)
Central Michigan	550	7.2	(4.8-10.7)
Midland	230	6.6	(3.7-11.4)
Region 12	809	7.0	(5.2-9.2)
Luce-Mackinac-Alger-Schoolcraft	-- ^c	-- ^c	-- ^c
Western Upper Peninsula	-- ^c	-- ^c	-- ^c
Delta-Menominee	-- ^c	-- ^c	-- ^c
Chippewa	-- ^c	-- ^c	-- ^c
Dickinson-Iron	69	18.9	(10.5-31.5)
Marquette	-- ^c	-- ^c	-- ^c

^a Among all adults, the proportion who had ever been told by a doctor that they had a heart attack or myocardial infarction.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 22: Angina or Coronary Heart Disease Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told Angina or Coronary Heart Disease ^a		
	N	%	95% Confidence Interval
Michigan Total	31,441	4.9	(4.6-5.2)
Region 1	12,806	4.9	(4.5-5.3)
Livingston	1,532	4.4	(3.2-5.9)
Macomb	1,872	5.4	(4.5-6.6)
Monroe	356	4.3	(2.5-7.3)
Oakland	3,167	4.7	(3.9-5.6)
St. Clair	384	5.9	(4.0-8.7)
Washtenaw	1,464	3.6	(2.7-4.7)
City of Detroit	1,388	5.2	(4.0-6.7)
Wayne exc. Detroit	2,571	4.9	(4.1-5.8)
Region 2	2,142	5.6	(4.6-6.7)
Genesee	1,614	5.7	(4.6-7.0)
Lapeer	-- ^c	-- ^c	-- ^c
Shiawassee	262	4.4	(2.6-7.6)
Region 3	1,745	4.7	(3.7-5.9)
Jackson	1,390	5.5	(4.4-6.9)
Lenawee	291	4.6	(2.7-7.7)
Region 4	1,743	4.3	(3.4-5.4)
Barry-Eaton	451	4.5	(3.0-6.6)
Mid-Michigan ^b	519	4.9	(3.3-7.3)
Ingham	947	3.4	(2.3-4.9)
Region 5	1,471	4.9	(3.9-6.3)
Branch-Hillsdale-St. Joseph	428	6.2	(4.3-8.9)
Calhoun	365	4.8	(3.0-7.7)
Kalamazoo	656	4.2	(2.8-6.4)
Region 6	662	5.7	(4.1-7.9)
Van Buren-Cass	277	9.7	(6.4-14.4)
Berrien	330	3.5	(2.0-5.9)
Region 7	3,071	3.0	(2.4-3.8)
Allegan	-- ^c	-- ^c	-- ^c
Ionia	-- ^c	-- ^c	-- ^c
Kent	1,620	3.3	(2.4-4.4)
Mid-Michigan ^b	519	4.9	(3.3-7.3)
Ottawa	716	2.2	(1.4-3.4)

Table 22 Cont'd			
Geographic Area	Ever Told Angina or Coronary Heart Disease ^a		
	N	%	95% Confidence Interval
Michigan Total	31,441	4.9	(4.6-5.2)
Region 8	2,590	5.0	(4.1-6.1)
District #10	1,081	5.2	(4.0-6.7)
Muskegon	1,408	4.9	(3.9-6.2)
Region 9	1,769	5.7	(4.6-7.0)
District #2	207	7.3	(4.4-11.9)
District #4	265	6.4	(4.3-9.4)
Northwest Michigan	541	4.6	(3.1-6.7)
Benzie-Leelanau	251	6.9	(3.8-12.1)
Grand Traverse	448	5.6	(3.5-8.9)
Region 10	1,149	6.6	(5.1-8.5)
Bay	255	5.8	(3.5-9.3)
Huron	115	8.7	(4.9-15.0)
Saginaw	468	6.5	(4.1-10.1)
Sanilac	-- ^c	-- ^c	-- ^c
Tuscola	151	7.0	(3.9-12.2)
Region 11	956	6.1	(4.4-8.5)
Central Michigan	547	6.1	(4.0-9.1)
Midland	228	7.4	(4.4-12.1)
Region 12	809	6.9	(5.1-9.2)
Luce-Mackinac-Alger-Schoolcraft	-- ^c	-- ^c	-- ^c
Western Upper Peninsula	-- ^c	-- ^c	-- ^c
Delta-Menominee	-- ^c	-- ^c	-- ^c
Chippewa	-- ^c	-- ^c	-- ^c
Dickinson-Iron	-- ^c	-- ^c	-- ^c
Marquette	155	9.3	(5.1-16.3)

^a Among all adults, the proportion who had ever been told by a doctor that they had angina or coronary heart disease.
^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.
^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 23: Stroke Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told Stroke^a		
	N	%	95% Confidence Interval
Michigan Total	31,636	3.6	(3.3-3.8)
Region 1			
Livingston	1,537	2.0	(1.3-2.9)
Macomb	1,883	4.1	(3.2-5.3)
Monroe	-- ^c	-- ^c	-- ^c
Oakland	3,187	2.9	(2.3-3.7)
St. Clair	383	4.9	(3.0-7.7)
Washtenaw	1,473	3.1	(2.1-4.6)
City of Detroit	1,397	5.9	(4.7-7.4)
Wayne exc. Detroit	2,585	3.6	(2.8-4.6)
Region 2	2,154	3.7	(2.8-4.9)
Genesee	1,622	4.2	(3.1-5.7)
Lapeer	-- ^c	-- ^c	-- ^c
Shiawassee	-- ^c	-- ^c	-- ^c
Region 3	1,763	3.8	(2.8-5.1)
Jackson	1,405	4.0	(3.1-5.2)
Lenawee	-- ^c	-- ^c	-- ^c
Region 4	1,755	3.2	(2.4-4.3)
Barry-Eaton	454	3.0	(1.8-4.8)
Mid-Michigan ^b	525	4.3	(2.5-7.2)
Ingham	952	2.5	(1.6-3.8)
Region 5	1,487	3.8	(2.8-5.2)
Branch-Hillsdale-St. Joseph	434	4.9	(2.8-8.4)
Calhoun	-- ^c	-- ^c	-- ^c
Kalamazoo	662	3.3	(2.1-5.2)
Region 6	669	3.3	(2.2-5.1)
Van Buren-Cass	282	4.6	(2.6-8.0)
Berrien	-- ^c	-- ^c	-- ^c
Region 7	3,081	2.7	(2.1-3.4)
Allegan	-- ^c	-- ^c	-- ^c
Ionia	-- ^c	-- ^c	-- ^c
Kent	1,623	2.9	(2.1-4.0)
Mid-Michigan ^b	525	4.3	(2.5-7.2)
Ottawa	719	2.8	(1.8-4.4)

Table 23 Cont'd			
Geographic Area	Ever Told Stroke ^a		
	N	%	95% Confidence Interval
Michigan Total	31,636	3.6	(3.3-3.8)
Region 8	2,602	3.5	(2.8-4.4)
District #10	1,082	3.8	(2.7-5.3)
Muskegon	1,416	3.3	(2.5-4.4)
Region 9	1,785	3.8	(2.8-5.3)
District #2	-- ^c	-- ^c	-- ^c
District #4	-- ^c	-- ^c	-- ^c
Northwest Michigan	547	3.3	(1.8-6.0)
Benzie-Leelanau	-- ^c	-- ^c	-- ^c
Grand Traverse	450	2.2	(1.2-3.8)
Region 10	1,158	3.7	(2.5-5.4)
Bay	-- ^c	-- ^c	-- ^c
Huron	-- ^c	-- ^c	-- ^c
Saginaw	469	5.0	(2.8-8.7)
Sanilac	-- ^c	-- ^c	-- ^c
Tuscola	-- ^c	-- ^c	-- ^c
Region 11	960	4.0	(2.8-5.6)
Central Michigan	549	3.6	(2.4-5.3)
Midland	-- ^c	-- ^c	-- ^c
Region 12	813	3.4	(2.4-4.9)
Luce-Mackinac-Alger-Schoolcraft	-- ^c	-- ^c	-- ^c
Western Upper Peninsula	-- ^c	-- ^c	-- ^c
Delta-Menominee	-- ^c	-- ^c	-- ^c
Chippewa	-- ^c	-- ^c	-- ^c
Dickinson-Iron	-- ^c	-- ^c	-- ^c
Marquette	-- ^c	-- ^c	-- ^c

^a Among all adults, the proportion who had ever been told by a doctor that they had a stroke.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 24: Cardiovascular Disease Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told Any Cardiovascular Disease^a		
	N	%	95% Confidence Interval
Michigan Total	31,382	9.9	(9.5-10.3)
Region 1	12,780	10.0	(9.3-10.6)
Livingston	1,534	7.9	(6.4-9.7)
Macomb	1,866	11.2	(9.7-12.9)
Monroe	355	9.7	(6.7-13.7)
Oakland	3,162	8.6	(7.6-9.8)
St. Clair	383	10.9	(8.1-14.7)
Washtenaw	1,462	7.1	(5.7-8.9)
City of Detroit	1,383	12.1	(10.2-14.2)
Wayne exc. Detroit	2,563	10.7	(9.4-12.1)
Region 2	2,133	10.7	(9.3-12.3)
Genesee	1,603	11.0	(9.3-12.9)
Lapeer	249	11.9	(7.9-17.5)
Shiawassee	262	8.5	(5.5-12.8)
Region 3	1,748	9.7	(8.2-11.4)
Jackson	1,392	10.8	(9.2-12.6)
Lenawee	293	10.0	(6.9-14.3)
Region 4	1,746	8.5	(7.2-10.1)
Barry-Eaton	452	8.1	(6.0-10.8)
Mid-Michigan ^b	521	10.8	(8.1-14.4)
Ingham	948	7.1	(5.5-9.2)
Region 5	1,470	10.8	(9.1-12.8)
Branch-Hillsdale-St. Joseph	430	13.8	(10.4-18.1)
Calhoun	363	10.4	(7.2-14.7)
Kalamazoo	654	8.9	(6.7-11.7)
Region 6	662	10.5	(8.3-13.3)
Van Buren-Cass	280	16.1	(11.9-21.4)
Berrien	327	7.3	(4.8-11.1)
Region 7	3,063	6.7	(5.8-7.8)
Allegan	349	7.4	(5.1-10.6)
Ionia	186	8.4	(4.9-14.0)
Kent	1,614	6.6	(5.3-8.2)
Mid-Michigan ^b	521	10.8	(8.1-14.4)
Ottawa	711	6.1	(4.5-8.1)

Table 24 Cont'd			
Geographic Area	Ever Told Any Cardiovascular Disease ^a		
	N	%	95% Confidence Interval
Michigan Total	31,382	9.9	(9.5-10.3)
Region 8			
District #10	2,585	11.2	(9.7-12.9)
Muskegon	1,076	13.3	(11.1-15.8)
	1,407	8.8	(7.4-10.5)
Region 9			
District #2	1,767	11.2	(9.5-13.1)
District #4	206	13.1	(8.5-19.8)
Northwest Michigan	266	13.5	(9.6-18.8)
Benzie-Leelanau	539	10.9	(8.1-14.6)
Grand Traverse	253	10.8	(6.9-16.5)
	446	10.0	(7.2-13.8)
Region 10			
Bay	1,144	11.3	(9.3-13.7)
Huron	254	8.3	(5.5-12.4)
Saginaw	116	15.1	(9.6-23.0)
Sanilac	464	12.2	(8.9-16.6)
Tuscola	146	13.0	(8.1-20.1)
	151	9.9	(6.1-15.7)
Region 11			
Central Michigan	955	11.3	(9.0-14.1)
Midland	547	12.1	(9.2-15.8)
	228	11.5	(7.6-17.0)
Region 12			
Luce-Mackinac-Alger-Schoolcraft	805	12.3	(10.0-15.1)
Western Upper Peninsula	103	16.3	(9.9-25.4)
Delta-Menominee	190	8.3	(5.1-13.2)
Chippewa	143	12.7	(7.7-20.2)
Dickinson-Iron	114	12.2	(6.5-21.5)
Marquette	67	24.8	(14.6-38.7)
	154	12.4	(7.7-19.4)

^a Among all adults, the proportion who had ever been told by a doctor that they had a heart attack, coronary heart disease or a stroke.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

Table 25: Cancer Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told Cancer ^a		
	N	%	95% Confidence Interval
Michigan Total	31,609	13.0	(12.6-13.4)
Region 1			
Livingston	1,536	14.6	(12.4-17.2)
Macomb	1,881	13.0	(11.4-14.7)
Monroe	363	17.8	(13.9-22.4)
Oakland	3,185	13.3	(12.1-14.6)
St. Clair	383	17.8	(14.1-22.4)
Washtenaw	1,474	12.3	(10.6-14.4)
City of Detroit	1,402	9.4	(7.8-11.3)
Wayne exc. Detroit	2,580	13.1	(11.7-14.6)
Region 2			
Genesee	1,623	11.6	(10.0-13.3)
Lapeer	247	18.2	(13.5-24.0)
Shiawassee	263	12.8	(9.0-17.9)
Region 3			
Jackson	1,402	14.1	(12.2-16.1)
Lenawee	294	13.1	(9.5-17.9)
Region 4			
Barry-Eaton	453	11.5	(9.0-14.7)
Mid-Michigan ^b	525	12.2	(9.6-15.5)
Ingham	951	11.4	(9.4-13.8)
Region 5			
Branch-Hillsdale-St. Joseph	434	13.4	(10.3-17.4)
Calhoun	369	11.9	(8.7-16.0)
Kalamazoo	658	9.4	(7.4-11.8)
Region 6			
Van Buren-Cass	279	13.6	(9.8-18.5)
Berrien	331	12.4	(9.3-16.2)
Region 7			
Allegan	350	15.0	(11.6-19.2)
Ionia	187	14.0	(9.5-20.2)
Kent	1,621	11.6	(10.2-13.3)
Mid-Michigan ^b	525	12.2	(9.6-15.5)
Ottawa	720	13.7	(11.3-16.5)

Table 25 Cont'd			
Geographic Area	Ever Told Cancer ^a		
	N	%	95% Confidence Interval
Michigan Total	31,609	13.0	(12.6-13.4)
Region 8			
District #10	2,603	15.7	(14.0-17.7)
Muskegon	1,084	19.2	(16.4-22.4)
	1,415	12.0	(10.4-14.0)
Region 9			
District #2	1,780	15.8	(13.8-18.1)
District #4	210	22.3	(16.4-29.6)
Northwest Michigan	265	12.1	(7.5-18.9)
Benzie-Leelanau	543	14.6	(11.4-18.4)
Grand Traverse	255	17.2	(12.3-23.4)
	450	17.4	(13.7-21.9)
Region 10			
Bay	1,154	14.2	(12.0-16.6)
Huron	259	13.1	(8.9-19.0)
Saginaw	115	13.8	(8.2-22.4)
Sanilac	467	13.0	(10.1-16.7)
Tuscola	148	17.7	(10.9-27.3)
	152	17.7	(12.0-25.2)
Region 11			
Central Michigan	959	14.1	(11.6-16.9)
Midland	548	15.4	(12.1-19.4)
	230	13.3	(9.5-18.4)
Region 12			
Luce-Mackinac-Alger-Schoolcraft	810	14.8	(12.2-17.7)
Western Upper Peninsula	103	21.6	(13.3-33.0)
Delta-Menominee	191	11.3	(7.2-17.2)
Chippewa	146	15.6	(10.6-22.3)
Dickinson-Iron	114	11.5	(6.4-19.9)
Marquette	68	17.7	(10.0-29.5)
	154	15.8	(9.8-24.4)

^a Among all adults, the proportion who reported ever being told by a doctor that they had skin cancer or any other type of cancer.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

Table 26: Diabetes Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told Diabetes ^a		
	N	%	95% Confidence Interval
Michigan Total	31,684	11.3	(10.9-11.7)
Region 1	12,897	11.1	(10.5-11.8)
Livingston	1,541	9.5	(7.6-11.9)
Macomb	1,881	11.6	(10.1-13.3)
Monroe	363	10.2	(7.3-14.1)
Oakland	3,193	10.1	(8.9-11.3)
St. Clair	384	8.6	(6.1-12.1)
Washtenaw	1,476	7.9	(6.4-9.6)
City of Detroit	1,403	14.2	(12.2-16.4)
Wayne exc. Detroit	2,583	12.1	(10.7-13.6)
Region 2	2,157	12.1	(10.6-13.8)
Genesee	1,625	13.9	(12.0-16.1)
Lapeer	250	7.7	(4.8-11.9)
Shiawassee	262	6.9	(4.2-11.2)
Region 3	1,765	10.6	(9.0-12.4)
Jackson	1,405	12.0	(10.2-14.0)
Lenawee	296	9.8	(6.8-13.9)
Region 4	1,754	10.7	(9.1-12.5)
Barry-Eaton	453	11.9	(8.7-16.0)
Mid-Michigan ^b	524	10.9	(8.3-14.1)
Ingham	953	9.7	(7.7-12.2)
Region 5	1,487	13.1	(11.2-15.3)
Branch-Hillsdale-St. Joseph	434	12.6	(9.5-16.7)
Calhoun	370	17.3	(12.7-23.1)
Kalamazoo	660	11.3	(8.9-14.3)
Region 6	670	10.9	(8.6-13.8)
Van Buren-Cass	282	14.4	(10.3-19.7)
Berrien	333	9.3	(6.4-13.2)
Region 7	3,091	10.3	(9.1-11.6)
Allegan	349	12.4	(8.8-17.1)
Ionia	188	11.8	(7.7-17.6)
Kent	1,626	10.5	(8.9-12.4)
Mid-Michigan ^b	524	10.9	(8.3-14.1)
Ottawa	724	8.6	(6.5-11.3)

Table 26 Cont'd			
Geographic Area	Ever Told Diabetes ^a		
	N	%	95% Confidence Interval
Michigan Total	31,684	11.3	(10.9-11.7)
Region 8			
District #10	2,606	13.2	(11.6-15.0)
Muskegon	1,086	14.3	(12.0-17.0)
	1,416	12.4	(10.5-14.6)
Region 9	1,785	10.4	(8.7-12.4)
District #2	210	13.7	(9.2-20.0)
District #4	266	14.1	(9.7-20.1)
Northwest Michigan	546	8.2	(5.2-12.6)
Benzie-Leelanau	255	11.0	(7.3-16.2)
Grand Traverse	451	8.8	(6.0-12.7)
Region 10	1,161	12.9	(10.8-15.4)
Bay	259	10.2	(6.8-15.0)
Huron	116	11.4	(6.7-18.9)
Saginaw	470	14.5	(11.0-18.9)
Sanilac	150	14.6	(9.2-22.5)
Tuscola	153	12.2	(7.9-18.3)
Region 11	962	11.5	(9.2-14.4)
Central Michigan	550	13.4	(10.2-17.4)
Midland	231	11.0	(7.1-16.6)
Region 12	814	12.0	(9.9-14.6)
Luce-Mackinac-Alger-Schoolcraft	-- ^c	-- ^c	-- ^c
Western Upper Peninsula	192	13.8	(9.2-20.0)
Delta-Menominee	146	14.2	(9.2-21.3)
Chippewa	114	13.5	(7.9-22.0)
Dickinson-Iron	-- ^c	-- ^c	-- ^c
Marquette	155	10.0	(6.0-16.2)

^a The proportion of adults who reported that they were ever told by a doctor that they have diabetes. Women who had diabetes only during pregnancy and adults who were diagnosed with pre-diabetes were considered to not have been diagnosed with diabetes.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.

^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 27: Kidney Disease Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told Kidney Disease ^a		
	N	%	95% Confidence Interval
Michigan Total	31,629	3.5	(3.3-3.8)
Region 1	12,871	3.6	(3.2-4.0)
Livingston	1,536	2.1	(1.4-3.1)
Macomb	1,880	3.2	(2.4-4.1)
Monroe	361	4.3	(2.4-7.5)
Oakland	3,189	3.3	(2.7-4.1)
St. Clair	382	4.8	(2.7-8.5)
Washtenaw	1,472	3.1	(2.1-4.6)
City of Detroit	1,400	4.5	(3.4-6.0)
Wayne exc. Detroit	2,578	3.9	(3.1-4.8)
Region 2	2,157	3.6	(2.8-4.7)
Genesee	1,623	3.6	(2.7-4.8)
Lapeer	-- ^c	-- ^c	-- ^c
Shiawassee	-- ^c	-- ^c	-- ^c
Region 3	1,767	3.5	(2.6-4.6)
Jackson	1,407	3.6	(2.7-4.9)
Lenawee	296	3.3	(1.8-6.0)
Region 4	1,750	2.6	(1.9-3.5)
Barry-Eaton	452	3.2	(2.0-5.1)
Mid-Michigan ^b	524	3.1	(1.8-5.4)
Ingham	950	2.6	(1.7-3.9)
Region 5	1,480	5.0	(3.7-6.6)
Branch-Hillsdale-St. Joseph	433	5.7	(3.5-9.1)
Calhoun	365	6.8	(3.8-12.0)
Kalamazoo	659	3.5	(2.4-5.2)
Region 6	665	3.4	(2.2-5.2)
Van Buren-Cass	-- ^c	-- ^c	-- ^c
Berrien	-- ^c	-- ^c	-- ^c
Region 7	3,088	3.0	(2.4-3.7)
Allegan	350	2.9	(1.7-5.0)
Ionia	-- ^c	-- ^c	-- ^c
Kent	1,624	3.0	(2.2-4.1)
Mid-Michigan ^b	524	3.1	(1.8-5.4)
Ottawa	723	2.4	(1.4-4.1)

Table 27 Cont'd			
Geographic Area	Ever Told Kidney Disease ^a		
	N	%	95% Confidence Interval
Michigan Total	31,629	3.5	(3.3-3.8)
Region 8			
District #10	2,600	3.7	(2.9-4.7)
Muskegon	1,081	4.0	(2.8-5.5)
Muskegon	1,415	3.8	(2.8-5.1)
Region 9	1,782	3.4	(2.4-4.7)
District #2	-- ^c	-- ^c	-- ^c
District #4	-- ^c	-- ^c	-- ^c
Northwest Michigan	545	1.9	(1.1-3.3)
Benzie-Leelanau	-- ^c	-- ^c	-- ^c
Grand Traverse	-- ^c	-- ^c	-- ^c
Region 10	1,160	3.4	(2.3-5.0)
Bay	-- ^c	-- ^c	-- ^c
Huron	-- ^c	-- ^c	-- ^c
Saginaw	-- ^c	-- ^c	-- ^c
Sanilac	-- ^c	-- ^c	-- ^c
Tuscola	-- ^c	-- ^c	-- ^c
Region 11	959	4.1	(2.8-5.8)
Central Michigan	548	3.9	(2.5-6.2)
Midland	-- ^c	-- ^c	-- ^c
Region 12	813	4.6	(3.3-6.4)
Luce-Mackinac-Alger-Schoolcraft	-- ^c	-- ^c	-- ^c
Western Upper Peninsula	-- ^c	-- ^c	-- ^c
Delta-Menominee	-- ^c	-- ^c	-- ^c
Chippewa	-- ^c	-- ^c	-- ^c
Dickinson-Iron	-- ^c	-- ^c	-- ^c
Marquette	-- ^c	-- ^c	-- ^c

^a Among all adults, the proportion who reported ever being told by a doctor that they had kidney disease.
^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.
^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 28: Depression Michigan BRFSS, 2017-2019 Combined			
Geographic Area	Ever Told Depression ^a		
	N	%	95% Confidence Interval
Michigan Total	31,555	22.7	(22.1-23.3)
Region 1	12,840	21.0	(20.1-21.9)
Livingston	1,536	20.5	(17.7-23.7)
Macomb	1,879	20.2	(18.1-22.5)
Monroe	361	25.1	(20.2-30.8)
Oakland	3,174	18.6	(17.0-20.4)
St. Clair	380	24.5	(19.6-30.2)
Washtenaw	1,465	24.5	(21.6-27.6)
City of Detroit	1,396	21.1	(18.6-23.7)
Wayne exc. Detroit	2,576	22.2	(20.3-24.2)
Region 2	2,152	23.1	(20.9-25.4)
Genesee	1,619	23.7	(21.2-26.5)
Lapeer	249	22.3	(16.8-28.9)
Shiawassee	264	20.7	(14.8-28.3)
Region 3	1,761	24.1	(21.5-27.0)
Jackson	1,402	26.3	(23.5-29.3)
Lenawee	295	22.3	(17.0-28.6)
Region 4	1,746	23.5	(21.2-26.1)
Barry-Eaton	453	21.4	(17.1-26.4)
Mid-Michigan ^b	524	21.0	(17.0-25.5)
Ingham	945	26.1	(22.7-29.7)
Region 5	1,485	27.3	(24.4-30.4)
Branch-Hillsdale-St. Joseph	433	26.5	(21.2-32.5)
Calhoun	370	26.0	(20.3-32.5)
Kalamazoo	660	28.4	(24.3-33.0)
Region 6	668	23.6	(19.8-27.9)
Van Buren-Cass	281	24.3	(18.9-30.7)
Berrien	332	21.9	(16.8-28.0)
Region 7	3,075	24.2	(22.3-26.2)
Allegan	349	26.7	(21.1-33.1)
Ionia	188	30.9	(22.6-40.7)
Kent	1,615	23.7	(21.2-26.3)
Mid-Michigan ^b	524	21.0	(17.0-25.5)
Ottawa	719	22.7	(18.5-27.5)

Table 28 Cont'd			
Geographic Area	Ever Told Depression ^a		
	N	%	95% Confidence Interval
Michigan Total	31,555	22.7	(22.1-23.3)
Region 8			
District #10	2,595	28.3	(25.7-30.9)
Muskegon	1,081	26.9	(23.3-30.8)
	1,410	31.7	(28.3-35.3)
Region 9	1,779	23.8	(21.3-26.5)
District #2	210	21.5	(15.5-28.9)
District #4	263	19.5	(14.0-26.4)
Northwest Michigan	544	25.8	(21.2-31.0)
Benzie-Leelanau	255	20.8	(15.5-27.4)
Grand Traverse	450	28.6	(23.4-34.4)
Region 10	1,157	24.4	(21.2-27.8)
Bay	258	32.9	(25.0-41.9)
Huron	115	11.6	(6.7-19.3)
Saginaw	468	23.6	(19.1-28.7)
Sanilac	150	22.4	(15.4-31.2)
Tuscola	153	20.5	(14.3-28.4)
Region 11	958	21.8	(18.6-25.3)
Central Michigan	550	23.9	(19.7-28.7)
Midland	228	16.8	(11.8-23.5)
Region 12	807	22.4	(19.2-26.1)
Luce-Mackinac-Alger-Schoolcraft	103	27.3	(17.4-40.1)
Western Upper Peninsula	192	16.8	(11.7-23.6)
Delta-Menominee	143	20.9	(14.4-29.2)
Chippewa	114	25.5	(16.8-36.7)
Dickinson-Iron	67	27.0	(15.9-42.0)
Marquette	154	27.5	(20.0-36.5)

^a Among all adults, the proportion who reported ever being told by a doctor that they had a depressive disorder including depression, major depression, dysthymia, or minor depression.

^b The Mid-Michigan District Health Department consists of Clinton, Gratiot, and Montcalm. Clinton and Gratiot are Region 4 counties, while Montcalm is a Region 7 county. All three counties were included in the Mid-Michigan estimate, but only the respective counties were included in the regional estimates.