## COVID-19 Vaccine Resource Guide

COVID-19 Vaccine	Pfizer (mRNA)¹: Orange Cap	Pfizer (mRNA)²: Purple Cap	Pfizer (mRNA)³: Gray Cap DO NOT DILUTE	Moderna (mRNA) <sup>4</sup> (Note: Booster Dose Volume)*	J&J/ Janssen <sup>5</sup> Not Preferred <sup>6</sup> (mRNA vaccines are preferred for primary, and booster doses)
Primary Series for Immunocompetent: Age Indication	5 through 11 years	12 years and older	12 years and older	18 years and older	18 years and older
Number of Primary Doses	2 doses	2 doses	2 doses	2 doses	1 dose
Interval Between Doses	21 days	3 weeks (8 weeks optimal for some persons) <sup>7</sup>	3 weeks (8 weeks optimal for some persons) <sup>7</sup>	4 weeks (8 weeks optimal for some persons) <sup>7</sup>	Not Applicable
Dose Volume	0.2mL	0.3mL	0.3mL	0.5mL	0.5mL

## Please See Page 2 For Guidance Specific to Moderately or Severely Immunocompromised and Booster Doses

Preparation/dosage:  (Discard diluent vial after one use, even if diluent remains)	1.3mL diluent/ 0.2mL dose	1.8mL diluent/ 0.3mL dose	DO NOT DILUTE/ 0.3mL dose	DO NOT DILUTE/ 0.5mL dose OR 0.25mL dose for booster only	DO NOT DILUTE/ 0.5mL dose
Storage Requirements:  DO NOT REFREEZE VACCINE	Ultra-Cold Freezer: Until expiration date <sup>1</sup>	Ultra-Cold Freezer: expiry date on label + 3 months	Ultra-Cold Freezer: Until expiration date <sup>3</sup>	DO NOT STORE IN ULTRA- COLD FREEZER	DO NOT STORE IN ULTRA-COLD FREEZER
	DO NOT STORE IN A STANDARD FREEZER	Standard Freezer: 2 weeks	DO NOT STORE IN A STANDARD FREEZER	Standard Freezer: Until expiration date <sup>8</sup> (QR code)	DO NOT STORE IN A STANDARD FREEZER
See below for temperature ranges <sup>†</sup>	Refrigerator: 10 weeks	Refrigerator: 31 days	Refrigerator: 10 weeks	Refrigerator: 30 days	Refrigerator: Until expiration date <sup>9</sup> (QR code)
Thaw Time	30 minutes-4 hours	30 minutes-3 hours	30 minutes-6 hours	Refrigerator-2.5 -3 hours Room Temp-1 hour	DO NOT FREEZE
Room Temp <b>Unpunctured</b>	12 hours	2 hours	12 hours	24 hours	12 hours
Room Temp <b>Punctured</b> (Room temp in syringe)	12 hours	6 hours	12 hours	12 hours (no more than 20 punctures per vial)	2 hours

<sup>†</sup>Ultra-Cold Freezer: -90°C to -60°C (-130°F to -76°F) Standard Freezer: -25°C to -15°C (-13°F to 5°F) Refrigerator: 2°C to 8°C (35°F to 46°F)

COVID-19 Vaccine	Pfizer (mRNA)¹: Orange Cap	Pfizer (mRNA)²: Purple Cap	Pfizer (mRNA)³: Gray Cap DO NOT DILUTE	Moderna (mRNA) <sup>4</sup> (Note: Booster Dose Volume)*	J&J/ Janssen <sup>5</sup> Not Preferred <sup>6</sup> (mRNA vaccines are preferred for primary, and booster doses)
Primary Series for Moderately or Severely Immunocompromised <sup>10</sup> : Age indication	5 through 11 years	12 years and older	12 years and older	18 years and older	18 years and older <sup>11</sup>
Number of Primary Doses	3 doses	3 doses	3 doses	3 doses	2 doses
Interval between 1 <sup>st</sup> & 2 <sup>nd</sup> Dose	21 days	21 days	21 days	28 days	28 days- mRNA only
Interval between 2 <sup>nd</sup> & 3 <sup>rd</sup> Dose	28 days	28 days	28 days	28 days	N/A
Vaccine Dose Volume	0.2mL	0.3mL	0.3mL	0.5mL	0.3mL-Pfizer 0.5mL-Moderna
Booster Dose: Age Eligibility for Booster dose <sup>12</sup>	Not Eligible	12 years and older with a complete Primary Series	12 years and older with a complete Primary Series	18 years and older with a complete Primary Series	18 years and older with a complete Primary Series (mRNA vaccines preferred) <sup>6</sup>
Interval between Primary and Booster for Immunocompetent	Not Eligible	5 months or greater after the second dose	5 months or greater after the second dose	5 months or greater after the second dose	2 months or greater after the first dose
Interval between Primary and Booster for Immunocompromised	Not Eligible	3 months or greater after the third dose	3 months or greater after the third dose	3 months or greater after the third dose	2 months or greater after the second dose
Dose Volume for Booster	Not Eligible	0.3mL	0.3mL	0.25mL*	mRNA Preferred 0.5mL-J&J

<sup>&</sup>lt;sup>1</sup>Fact Sheet for Healthcare Providers Administering Vaccine - Pfizer-BioNTech COVID-19 Vaccine for 5 - 11 Years of Age-orange cap must dilute (fda.gov)

<sup>&</sup>lt;sup>2</sup>Fact Sheet for Healthcare Providers Administering Vaccine - Pfizer-BioNTech COVID-19 Vaccine for 12 and older - purple cap must dilute (fda.gov)

<sup>&</sup>lt;sup>3</sup>Fact Sheet for Healthcare Providers Administering Vaccine - Pfizer-BioNTech COVID-19 Vaccine for 12 and older - gray cap do not dilute (fda.gov)

<sup>&</sup>lt;sup>4</sup>Moderna COVID-19 Vaccine Fact Sheet for Healthcare Providers Administering Vaccine (fda.gov)

<sup>&</sup>lt;sup>5</sup>Janssen COVID-19 Vaccine - EUA Fact Sheet for Healthcare Providers Administering Vaccine (janssenlabels.com)

<sup>&</sup>lt;sup>6</sup>Johnson & Johnson's Janssen COVID-19 Vaccine Overview and Safety | CDC

<sup>&</sup>lt;sup>7</sup>Interim Clinical Considerations Primary Series Vaccination Schedule —An 8-week interval may be optimal for **some** people ages 12 years and older, especially for males ages 12 to 39 years. Follow hyperlink for further guidance on who is recommended to follow the shorter intervals.

<sup>&</sup>lt;sup>8</sup>Vaccination Provider Overview | Moderna COVID-19 Vaccine (EUA) (modernatx.com)

<sup>&</sup>lt;sup>9</sup>Janssen COVID-19 Vaccine Expiry Checker (vaxcheck.jnj-19)

<sup>&</sup>lt;sup>10</sup>COVID-19 Vaccines for Moderately or Severely Immunocompromised People | CDC

<sup>&</sup>lt;sup>11</sup>Appendix B. Guidance for People who are Moderately or Severely Immunocompromised and Vaccinated with Janssen COVID-19 Vaccine

<sup>12</sup> COVID-19 Vaccine Booster Shots | CDC